

Dendrochronologically dated pine stumps document phase wise bog expansion at a northwest German site between c. 6700 BC and c. 3400 BC

Inke E. M. Achterberg¹, Jan Eckstein, Bernhard Birkholz, Andreas Bauerochse², Hanns Hubert Leuschner¹

¹Department for Palynology and Climate Dynamics, University of Göttingen, 37073 Göttingen, Germany

²Lower Saxony State Service of Cultural Heritage, 30175 Hannover, Germany

Correspondence to: Inke E. M. Achterberg (iachter@gwdg.de)

Abstract. The investigated northwest German mire site at *Tote Moor* is densely covered with subfossil pine stumps (*Pinus sylvestris* L.) from the fen-bog transition. This facilitates the spatio-temporal reconstruction of mire development, which is based on 212 in situ tree stumps in the case study presented here. Six dendrochronologically dated site chronologies together cover 2345 years between 6703 BC and 3403 BC. The gaps in between are 6 to 550 years long. Additionally, a floating chronology of 309 years, containing 30 trees, was radiocarbon dated to the beginning of the 7th millennium cal. BC. Peat-stratigraphical survey was carried out additionally, and elevations a.s.l. were determined at several locations.

Tree dying-off phases, which indicate water level rise at the site, mostly in context of the local fen-bog transition, are evident for c. 6600 – 6450 BC, c. 6350 – 5750 BC, c. 5300 – 4900 BC, c. 4700 – 4550 BC, c. 3900 – 3850 BC, c. 3700 – 3600 BC, c. 3500 – 3450 BC and c. 3400 BC. The spatial distribution of the dated in situ trees illustrates the phase wise expansion of raised bog over fen peat at the site. The documented bog expansion pulses likely correspond to climatic wet sifts.

1 Introduction

Raised bog development shaped the northwest German lowland during the Holocene, as eventually about a third of the area had been covered by mires (fens and bogs) (Metzler, 2004). The development of these mires on the underlying nutrient poor glacial deposits was largely determined by climatic variations (Ellenberg, 1996). The area was particularly characterized by large lowland raised bogs, which grow better under humid and cool conditions (Behre, 2008). The expansion of these raised bogs is evident since the 7th millennium BC, with a strong increase between 5100 and 3600 BC (Eckstein et al., 2011, Petzelberger et al., 1999). The peat in the area contains a valuable environmental record of the past millennia.

The raised bogs were treeless in their central parts, while most of the surrounding region was wooded. Even the fens were wooded in large parts, often forming alder (*Alnus*) swamp forests or carrying other tree species. Often at the margins of a raised bog a swamp (lagg) develops, fed by run-off water from the bog mixing with ground water. In contrast to the otherwise generally treeless raised bog (Ellenberg, 1996), the more drained margin of the raised bog towards the lagg, is

commonly wooded by pine (*Pinus sylvestris* L.). The expansion of the raised bogs (which can be expected to occur during moist periods) often causes severe growth conditions for bog trees and consequently widespread dying-off phases. The remains of the affected trees are preserved under *Sphagnum* peat in the process. Stratigraphically the pine stumps mostly mark the local fen-bog transition. When in situ stumps are dated, they document the raised bog expansion.

5 Due to peat mining and the uncovering of such stump layers areal dendrochronological investigation became possible. The trees dated absolutely by dendrochronology deliver the finest temporal reconstruction of bog expansion and local bog formation respectively, particularly by their dying-off dates (Eckstein et al., 2009, 2010, Edvardsson et al., 2014, 2016, Krapiec et al. 2016, Leuschner et al., 2002, 2007). Eckstein et al. (2009, 2010, 2011) have shown, that the tree dying-off occurs in phases, which are often synchronous in different mires. They were able to show that the tree dying-off is mostly
10 due to hydrological changes, while fire and storm have been found to be of little relevance on the investigated sites. Particularly the observed synchrony of pine establishment and dying-off phases in different mires shows, that the local mire development documented by the trees is strongly linked to climatic variations (Eckstein et al., 2009, 2010, 2011, Edvardsson et al., 2012a, 2012b).

The phase-wise advance of a raised bog in northwest Germany is particularly well documented in the mire *Tote Moor* at the
15 site *TOMO-south* (Fig. 1). At the site, tree remains from one stratigraphical stump-layer (Fig. 2) turned out to originate from four millennia, documenting the bog progress across some 500 m.

At site *TOMO_south*, the spatial as well as the temporal distribution of many preserved trees has been obtained. The present study aims to connect areal changes to the temporal patterns observed.

2 Material and Methods

20 The *Tote Moor* mire complex is located north of lake *Steinhuder Meer* near Hannover. The undulating relief below the mire consist of sand, and is likely to have held several small ponds and isolated mires before the expanding mire complex connected them. The whole depression of the lake is based on the same nutrient-poor glaciofluvial deposits (mostly sand) that characterize the *geest* region.

25 On the site *TOMO_south* many in situ tree stumps, and also in situ stems and ex situ stumps (pulled at ditch-digging and left at ditch-side) were sampled as radial slices using a chain saw. The tree remains were documented by using a feature table (regarding their growth, size, conservation condition etc.), photographic pictures and GPS coordinates. On some tree stumps, the root depth and shape was investigated. Later, the samples were dried, reduced by circular saw, and frozen. Then, suitable measuring radii were surface-cut by scalpel, contrasted by rubbed-in chalk dust and measured on an *Aniol* motorized
30 measuring device with CATRAS measuring software to a precision of 0.01 mm. The tree ring-widths series were cross dated using mainly the V-program-set (a.o. SYNCH2) by Thomas Riemer (Riemer, 1994 and unpublished work) and the *TSAPWin* program by Rinntech (Rinn, 2003). The Lower Saxony chronologies of oak and pine were used as a base for dating. Part of

the Lower Saxony Pine Chronology is also a product of this work. For the floating chronology segment three radiocarbon dates, which were determined at the Leibniz Institute for Applied Geophysics (LIAG) in Hanover, were wiggle matched on base of the IntCal13 calibration curve (Reimer et al., 2013) using the *OxCal 4.2* online software (Bronk Ramsey et al., 2001, Bronk Ramsey, 2009).

5 In situ finds were later separated from ex situ finds. For the spatio-temporal reconstruction of raised bog development (Fig. 5), stumps with rootplates partially dragged upward, which retained part of their root system within the grown peat and thus their location, were used along with the finds that were in situ in the stricter sense. Up to 96 trees have possibly been moved in such a way. They were added to the 116 trees which were considered 'in situ' in a stricter sense. In total 210 tree stumps with dendrochronological datings and 2 radiocarbon dated tree stumps are included in the reconstruction and displayed (Fig. 10 5 and 6). All 212 are referred to as 'in situ' in the following. All maps were created using ArcGIS 9 and 10 (Esri) mapping software. Figures were prepared using CoralDRAW software.

In addition, the peat-stratigraphy of 56 cores at the site was investigated macroscopically. Elevations a.s.l. were determined by stadia survey for 36 of the peat cores taken for this study, and for 63 on behalf of ASB-Humus peat mining company, who kindly shared their data. The elevation a.s.l. of the mineral base (sand) beneath the peat is depicted as a regularized 15 spline interpolation on base of the total 99 elevation measurements (Fig. 6).

Dendrochronological dates are given in years BC, radiocarbon dates in years cal. BC. Labelled time spans, like chronology segments or gaps, always include the years named.

3 Results

At the site *TOMO_south*, a rather levelled field left by peat mining revealed the remains of an apparent pine forest (Fig. 2). 20 A closer look revealed, that what had appeared to be one continuous tree layer actually did contain neighbouring stumps grown on slightly different elevation levels (Fig. 3).

Of the 700 tree stumps sampled at this site most were pine (*Pinus sylvestris*), only 10 oak (*Quercus spec.*) stumps were sighted and sampled. The pine remains often had retained bark in the lower parts, often being preserved to bark edge or showing only minimal decay at the outermost rings.

25 3.1 Temporal distribution of the trees

Only after many pine stumps had been sampled (a first sampling of about 70 trees did not provide good cross-dating results), dendrochronological dating was successful for 477 trees. An additional floating chronology segment of 30 trees was radiocarbon dated. The trees at the seemingly homogeneous site in fact originated from various centuries (Fig. 4, Table 1). They grew around 7000 to 6700 +/-80 cal. BC (floating chronology segment) and between 6703 and 3403 BC 30 (dendrochronological calendar dates). The floating chronology segment covers 309 years, the chronology segments with dendrochronological calendar dates cover 2345 years.

The majority of the trees (443) originates from the 7th to 5th millennia BC. Much less trees (34) represent the 4th millennium BC. Tree occurrence was not scattered over time, but is found to cluster in at least eight groups (chronology segments C14 and A1 – F, Fig. 4). There are also periods without pine trees preserved at the site: while the two shortest of the five gaps between the site chronologies (6 and 11 years in length) belong to the poorly replicated 4th millennium BC and are likely due to sampling design (which, in this case, is the placing of the site borderline through peat mining), the longer gaps between the well replicated chronologies likely reflect the actual dynamic at the site.

There are periods where tree establishment and tree **dying-off events** appear rather scattered (e.g. 6000 – 5800 BC or 5230 – 5120 BC), and intervals where the wooded phases display rather clustered tree establishment and tree dying-off. Particularly clear are the dying-off events around 6315 BC, 5060 BC and 3838 BC (Fig. 4).

Rejuvenation pulses, where several trees germinate within a few years, are found within several segments. This can be seen clearly in segments C14, A2 (repeatedly) and B. Less clearly segments C and E depict the same, while segment F lacks replication and no such pattern is found in segments A1 and D (which is also poorly replicated). A closer look reveals, that all rejuvenation pulses within segments are preceded by a time of limited or even missing rejuvenation. The rejuvenation pulses hence seem to mark the end of periods with limited tree establishment. In several cases the phases of limited tree establishment are also phases of frequent tree dying-off. This is particularly clear for the germination pulse beginning with the dying-off event around 6315 BC.

3.2 Reconstructed bog expansion

Mapping of the dated in situ trees generally shows groups of contemporaneous trees growing together. The oldest trees, from the early 7th millennium BC, cluster in the south-east of the plot, while the youngest, those of the 4th millennium BC, are found to the north-west, with the rest arranged in between. This clearly shows the general direction of raised bog expansion from **south-west to north-east** across the plot, even though there are some discontinuities within the site. Firstly, the groups of trees are not arranged in orderly stripes (Fig. 5), but form tongues and islands, often in correspondence to the dynamics of the mineral base (sand) beneath (Fig. 6). On sandy elevations trees persisted much longer, while the surrounding mire had long become a treeless raised bog (Fig. 7). Secondly, there are sometimes individual trees from one time interspersed into a group from another.

The mineral base is lowest to the south-east, where the oldest trees were found. Those trees document the oldest part of raised-bog in the plot.

In Fig. 5 the raised bog advance has been reconstructed according to the dated in situ tree remains. The interpolated areas (Fig. 5) were set in a 'no tree growth after' approach. That means: In places, where trees from different chronology segments occur intertwined, the place was assigned to the younger specimen.

This was done in an attempt to depict the advance of the raised bog, which, in its final state, is mostly treeless (Ellenberg, 1996).

3.3 Peat-stratigraphy

The peat stratigraphy is shown for 31 cores of the site (Fig. 8), displayed in elevational relation. In total, elevation a.s.l. has been determined for 36 of the 56 cores which were peat-stratigraphically investigated at the site. Please note, that some cores have been taken at the edge of the peat excavation field, where peat was left standing more than 1.5 m higher than the level of the tree stump layer.

The general order above the glacial sand deposits is the following: brown moss peat on bottom, wood rich peat above, followed by *Sphagnum* peat on top. The pine tree remains are found at the fen-bog-transition. This picture is rather continuous over the site, with differences mostly restricted to layer thickness. The brown moss peat is mostly weakly humified, often features *Menyanthes* seeds (*Menyanthes trifolia*) in the middle or upper part, and silt near the bottom.

The wood rich peat varies, mostly being humified stronger, but containing in different spots and layers various portions of *Betula* (bark), *Alnus* (wood), *Pinus* (bark and cone) and charcoal.

A few spots feature an *Eriophorum* layer (*Eriophorum vaginatum*) below the *Sphagnum* peat. More cores show some *Eriophorum* mixed in with the lower *Sphagnum* peat or the upper wood rich peat.

The *Sphagnum* peat, where distinguishable, consists of *Sphagnum* section *Acutifolia* peat, *Sphagnum* section *Cuspidata* peat and *Sphagnum-Carex* peat.

A few of the cores (two of the ones shown) feature highly humified fen peat in the lower part, which is likely to be brown moss peat in a decomposed state. In agreement with this assumption, the brown moss peat of most cores is more humified at its base.

3.4 Root morphology

There is one main feature, dividing the pines into two groups by their roots: the first type of root systems (type 1) spreads out horizontally, without any downward pointing roots. The central root at these trees has either died off at a length of about 10 cm or less, or is not traceable at all. The second type of root systems (type 2), however, displays downward growing roots, most of them with a pronounced central root that has grown vertically downwards. Most often, these reach 20 to 40cm below the root plate.

These two types clearly show up in the ex situ material pulled from the ditches. Roots below the stem were investigated also for 18 in situ stumps (Fig. 9), primarily to clarify if their roots reached the mineral soil below the peat. This was not the case with the examined specimen.

4 Discussion

4.1 The preserved trees

During dry periods, with consequently lowered bog water tables, pine forestation can temporally also occur on the bog. Such growth of large pines within raised bog (rooting in *Sphagnum* peat) was described i.a. by Overbeck (1954) for a mire in northern Germany, by Moir et al. (2010) for such findings in Scotland, by Edvardsson et al. (2014) in southern Sweden, or in general terms by Ellenberg (1996).

However, this is not the case at *TOMO_south*, where the pines occur at the fen-bog transition and not within *Sphagnum* peat layers. As confirmed by peat stratigraphy, the tree remains in *TOMO_south* represent a persistent pine forestation at the raised bog margin. Pine forestation at raised bog margins is a very typical occurrence. They colonize marginal parts of the bog, where the ascending bog surface is well drained compared to more central parts (Ellenberg, 1996, Overbeck, 1975).

The death and conservation of the trees however, appears to be closely connected to the expanding of the raised bog and the rise of the corresponding water table. This has been found evident on base of abundant upward growing roots late in the trees life, the drastically narrowing rings near the bark, peat stratigraphical context and the state of conservation at various comparable sites investigated in northwest Germany and southern Sweden (Eckstein et al., 2009, 2010, Edvardsson et al., 2012b, 2014, Leuschner et al., 2002, 2007). Therefore the death of the trees and their conservation under *Sphagnum* peat dates raised bog expansion, and using in situ stumps adds location to the event. Seeing the continuously dense (spatial) cover of the site with tree stumps (Fig. 2), all significant and lasting raised bog expansion should be documented by embedded trees.

The tree data displays phases of cumulative dying-off on the one hand and gaps between site chronologies on the other. This documents phases of the bog expanding, and phases without significant expansion at the site.

The phases of bog expansion should generally relate to relatively humid climate phases, as the bogs are dependent on rain water (Ellenberg, 1996). Raised bog formation can be favoured by a preceding lowering of water table through the disconnection of the fen peat surface from the lowered ground water and its nutrient supply (Hughes 2004, Tahvanainen, 2011, von Bülow, 1935). At site *TOMO_south* this combination of a drier phase with lowered water tables followed by a phase of swift raised bog expansion may have occurred repeatedly. A lowered water table is documented by numerous tree stumps with vertical roots (type 2). These roots often reach 20 to 40 cm below the root plate, and include a thick central root going straight down (Fig. 9). Root system type 1, however, being spread out flat and without any pronounced vertical root, is also common at the site. Such roots indicate a higher water table, limiting rooting space at the time of growth. This shows, that not all of the trees embedded by the expanding raised bog grew under drier conditions with significantly lowered water tables.

In general, tree growth on a mire can also contribute to a lower water table, as transpiration is being increased (Moir et al. 2010, Limpens et al. 2014). This effect can be reversed however, where shading in high density stands reduces evaporation even more and thus results in increased surface wetness (Limpens et al. 2014).

The area around the bog appears to have been wooded continuously, as pollen records from the bog itself (Shumilovskikh et al. 2015, Schlütz unpublished) and the regional landscape history (Behre 2008) suggest. The trees from these surrounding mineral soils would not be preserved, however.

4.2 Dying-off phases

Most indicative for periods of lateral bog growth are those of tree dying-off. As there are seven dendrochronological dated chronology segments, there are at least seven such periods (Fig. 4). Particularly for the chronology segments A2 and B, phases of accelerated and decelerated bog advance appear to be depicted in the trees. Even though this could partially also relate to the irregular sampling pattern, some prominent short-term events of cumulative tree dying-off, like those around 6315 BC, 5060 BC and 3838 BC, are clear. The periods in which the trees died off are: 6614 – 6436 BC, 6365 – 5733 BC, 5308 – 4902 BC, 4712 – 4537 BC, 3895 – 3838 BC, 3691 – 3614 BC, 3496 – 3473 BC and 3407 – 3403 BC (Fig. 4, Table 1). These are interpreted as phases of raised bog expansion, which in turn imply moist phases. How the trees are successively affected by the local raised bog development is also illustrated in the individual tree ring series, as shown for segment F (Fig. 10).

The floating chronology segment C14 shows subsequent dying-off approximately. The trees of segment C14 are less well preserved. This might indicate a mire environment at the time with higher microbial activity, possibly with less continuous water logging of the mire surface.

4.3 Rejuvenation pulses and suppression

The rejuvenation discussed here is limited to trees, which grew up to form a number of rings sufficient for dendrochronological dating.

In the course of dying-off phases, rejuvenation is often missing. When all trees of stand are killed by a fatal influence, young trees germinated shortly prior to the event would not occur in the dendrochronological record due to insufficient tree ring numbers for dating. This can not be more than part of an explanation however, since the dying-off phases are in many cases stretched longer than the lengths of time required of an tree ring sequence for dating.

The moist conditions relating to the dying-off phases however, would likely also suppress successful tree establishment.

Holmgren et al. (2015) found *Pinus sylvestris* recruitment on bogs to be more successful on drier sites (hummocks) with lower water tables. They found natural seedling establishment to occur on both, moist (lawns) and drier microsites

(hummocks) equally. Young trees however were much less abundant on the moister sites (lawns), and adult trees were only found on the drier sites (hummocks) with lower water tables (Holmgren et al. 2015).

Several times in the record, numerous germinations occur synchronous. These are mostly found following a phase (mostly of one to a few decades) of suppressed rejuvenation. Hence, they most likely mark the end of the unfavourable conditions which suppressed rejuvenation. Drier conditions on the peat surface are likely a factor. Furthermore, the large numbers of simultaneous seedling establishment can be explained by the lack of shading undergrowth as a preceding generation of young trees is missing.

Tree rejuvenation events (germination phases), which took place simultaneously in several different peat bogs, sometimes coincide with dying-off events, and more often follow directly after them. Eckstein et al. (2010, 2012) and Leuschner et al. (2007) have referred to this as Germination-Dying-Off-phases (GDOs). Even though a spatial pilot study (Stenzel, 2013) did not find the newly established trees directly in the area shaded by those trees which had just died off, the mechanism causing the coherence is still suspected to be driven by competition for light. Other trees and shrub (e.g. *Betula* or young pine trees) would have suffered from the same influence which killed the large pine trees and thereby created an opening in canopy. This could explain the coherence or close succession of dying-off and germination phases.

Zackrisson et al. (1995) also take **seed production** into account to explain rejuvenation pulses in pine populations. Conditions stressful to the trees can enhance seed production. On the contrary, favourable climate conditions may also influence seed production. Enhanced seed production under stressful conditions supports rejuvenation when the older tree generation is about to die off. The germination and establishment of these seedlings however, depends on the opening of canopy created by the tree dying-off, and on otherwise suitable conditions at the site. In how far seed production might be reflected in tree establishment at the given site is therefore unclear.

4.4 Chronology-gaps

The site chronologies of *TOMO_south* are interrupted by 5 gaps, ranging between 550 and 6 years in length (Table 1). The two very short gaps (6 and 11 years respectively) are rather insignificant, especially, since the respective period is not well replicated. These gaps are likely to result from sampling design, as the neighbouring chronology segments are poorly replicated and the according trees are found to the rim of the investigated area. It is well possible, that more trees represent that period, but were located just outside the margin of the site to the north-west. Both short gaps belong to the 4th millennium BC {3837 – 3832 BC (6 years) and 4613-3603 BC (11 years)} (Table 1), when trees only grew at the site on sandy elevations, forming wooded 'islands' in the mire (Fig. 5, 6 and 7).

The three longer gaps are more meaningful. The two very long gaps are 5732 – 5405 BC (328 years) and 4536 – 3987 BC (550 years), the gap of intermediate length (64 years) is 4901 – 4836 BC. All three are framed by well-replicated site chronology segments, with the respective trees found well within the site. Given the dense sampling of the site, it can be assumed, that the large gaps actually represent periods with no or very few pine trees being embedded at the site.

Other studies have attributed chronology gaps in raised bogs to a lack of tree growth due to high surface wetness. This makes much sense for the tree layers well within *Sphagnum* peat as described e.g. in Scotland (Moir et al., 2010) and south Sweden (Edvardsson et al., 2014). At those sites, tree growth was only possible on the raised bog during drier phases.

In case of the present study however, the stratigraphic position of the tree layer is at the fen-bog transition. Apparently, at 5 *TOMO_south* the expanding raised bog embedded trees grown at its rim. The site was thoroughly covered with pine stumps. These were representatively sampled and dated. Taken together, this suggest, that the large chronology gaps at *TOMO_south* represent periods of (near) stagnation of the lateral raised bog growth. Therefore the gaps are not interpreted as periods of particularly high surface wetness in this case, but quite the contrary. During the periods of site chronology gaps the raised bog at *TOMO_south* appears not to have expanded significantly. There are however, tree dying-off phases documented at 10 other bogs in northwest Germany within the time of these gaps. In particular, in the 550 year gap at *TOMO_south* (4536 – 3987 BC), there are two pronounced dying-off phases (c. 4490 – 4370 BC and 4230 – 4170 BC) within and one at the end of the gap (c. 4010 – 3960 BC) in other bogs (Achterberg et al. 2015). Therefore, these phases of apparent bog growth stagnation at *TOMO_south* can not be related to climatic dry phases directly.

4.5 Spatial distribution

15 The mapping of the dated in situ trees at the site (Fig. 5) show the raised bog expansion from south-east to north-west. The mineral base is lowest in the south-east of the plot and highest to the south-west of the plot, as well as in the centre of the site (Fig. 6). This direction is parallel to the nearest lake shore.

There are clear spatial clusters of trees from the same periods, which usually border to patches of the preceding and following chronology segments. This is coherent with the picture of the advancing raised bog successively embedding the 20 tree stumps.

The oldest parts (with trees only from the first chronology segments) are found to the south-east, where the mineral base is low. The trees from more recent centuries were found in places where the mineral base ascends, like the sandy elevation near the centre of the plot. Apparently wooded islands within the raised bog persisted for a long time (Fig. 5, 6 and 7). The distribution of trees from different chronology segments in the plot can mostly be related to the dynamic relief of the mineral 25 base (Fig. 6).

At some places however, there are also trees from different epochs interspersed (Fig. 5). In the mapping of raised bog expansion (Fig. 5) these places are assigned to the last tree dying-off they document, for the advancing central part of the bog should have been generally treeless (Ellenberg, 1996). Re-establishment of trees in places where trees had already been embedded before may have been favoured by drier conditions, or simply relate to the dynamic relief, with the root plates of 30 previous generations additionally serving as 'stepping stones' for the new trees to grow on.

4.6 Peat stratigraphy

Thick and mostly weakly humified layers of brown moss peat were found at the bottom of *TOMO_south* peat cores. It was the base for extended tree growth, which then itself deposited thick layers of wood rich peat. Were basal swamp forest peat is less humified, the brown moss can still be seen intertwined. Likewise intertwined is the first *Sphagnum* growth into the top layers of pine rich peat. This depicts a succession to ombrotrophy, with one plant community creating the habitat for the next.

There are fine mineral materials found interspersed in several peat cores and layers. These may have been washed into the moss by temporary flooding or, particularly in the older deposits, also might have been blown into the mire.

4.7 Climatic comparisons

The following alignments were largely limited to records, which have a certain precision of dating (dendrochronological, varve, etc.), are located with some level of proximity (European studies), regard the time frame covered by *TOMO_south* trees and also describe a hydrological signal. For the type of proxy used in the studies referred to and their location, please see Table 2. The temporal overview given in the following text is also displayed in Figure 11 A and B.

We interpret the **dying-off phases** as times of lateral mire expansion. Bog expansion again would require a certain humidity, which may be more dependent on a reduced evapotranspiration than the actual precipitation alone. This would mean the dying-off phases in the periods c. 6600 – 6450 BC, 6350 – 5750 BC, 5300 – 4900 BC, 4700 – 4550 BC, 3900 – 3850 BC, 3700 – 3600 BC, 3500 – 3400 BC indicate more humid phases. In turn, the **gaps** between the site chronologies and the periods of rather undisturbed tree growth are interpreted as phases of stagnation of lateral raised bog expansion. These may be related to climatic conditions unfavourable for mire growths, possibly involving drier periods. This would apply to the phases c. 6450 – 6350 BC, 5750 – 5300 BC, 4900 – 4700 BC, 4550 – 3900 BC, 3850 – 3700 BC, and 3600 – 3500 BC.

As mentioned before, the dying-off phases observed at *TOMO_south* show much synchronism with dying-off phases observed at other mire sites in northwest Germany (Eckstein et al., 2009, 2010, 2011), which emphasises their climatic context. In addition to the temporal placing of the *TOMO_south* record (terminating 3400 BC) and its location in the northwest German lowland, where settlements were established later than on the richer soils of the adjacent hills, this synchrony makes it appear unlikely, that anthropogenic clearance activity in the catchment area might have had detectable influence on the bogs hydrology.

The replication of the northwest German dendrochronological pine record, even though it is influenced by sampling patterns and the accessibility of trees depending on peat mining, also reflects climate. This shows in the European similarities of chronology replication (Edvardsson et al. 2016). And it is also illustrated by the comparison with other climate-related records (Fig. 11 A). The number of dates for lowered and raised lake levels in Mid-Europe (Magny et al. 2004) displays

similarities with the replication of the northwest German pine chronology. The northwest German pine declines (die-off phases visible in replication) within the long *TOMO_south* gap (c. 4500 – 4000 BC) fit phases of higher lake levels in the data of Magny et al. (2004) particularly well (Fig. 11 A). The replication at *TOMO_south* is largely in tune with the over-all pine replication record of northwest Germany, except for northwest German phases of pine growth which are not represented at *TOMO_south*, dating c. 4900-4800 BC and c. 4500 – 4000 BC (which is a long gap between site chronologies at *TOMO_south*).

The **8.2 ka event** cooling phase, according to Thomas et al. (2007), began around c. 6300 cal. BC. Kobashi et al. (2007) date its beginning later, to c. 6225 cal. BC, while Veski et al. (2004) observe the time of maximum cooling for 6250 – 6150 varve years BC (Fig. 11 B). Veski et al. (2004) also observe an abrupt end of the 8.2 cool period for 6080 varve years BC.

Dendrochronological data in proposed context with the event includes a phase of pine establishment in Irish bogs from 6210 BC on (Torbensohn et al., 2015), a phase of poor growth and regeneration of west German oaks from the river Main sediments c. 6270 – 6000 BC (Spurk et al., 2002) and a gap in the bog oak chronology 6177 – 6060 BC (Achterberg et al., 2016). A pronounced glacier advance at Mont Miné in the Swiss Alps was dated to c. 6175 BC using dendrochronology (Nicolussi & Schlüchter 2012). The replication of the Eastern Alpine Conifer Chronology (Nicolussi et al. 2009) drops after 6250 BC for about 200 years. The replication of the Northwest German Pine Chronology (Achterberg et al. 2016) is also declining at the time, but does not reach particularly low levels.

The dates mentioned above correspond well in general. Particularly good agreement is found, for example, between the beginning of the gap in the Northwest German Bog Oak Chronology 6177 BC (Achterberg et al. 2016) and the glacier advance dated by Nicolussi and Schlüchter (2012) to c. 6175 BC, as well as between the end of said gap in the Northwest German Bog Oak Chronology 6060 BC (Achterberg et al. 2016) and the end of the cool phase observed by Veski et al. (2004) for 6080 varve BC. The phases of unfavourable growth conditions observed at river Main from c. 6270 BC on (Spurk et al. 2002) and in the Alps from c. 6250 BC on (Nicolussi et al. 2009) are in equally close agreement.

At site *TOMO_south*, a phase of frequent tree dying-off from 6250 BC to 6157 BC matches the time of maximum cooling (6250 – 6150 varve years BC) as described by Veski et al. (2004) and the beginning of the replication decline from c. 6250 BC on in the Eastern Alpine Conifer Chronology (Nicolussi et al. 2009). The trees of *TOMO_south* display a strong dying-off pulse at c. 6315 BC, which may relate to the event as Thomas et al. (2007) date it. However, the 8.2k event is not showing as clearly in the pine record of *TOMO-south* as it is in other records.

Magny et al. (2004) describe several phases of high lake levels in mid-Europe, four of which are within the time frame covered at *TOMO_south* (In the following these are referred to as first to fourth phase, disregarding all which are outside of the time frame discussed here). The first of these (c. 6350 – 6100 cal. BC) is also within the dying-off phase of segment A2 (6365 – 5733 BC). The beginning of the two phases (high lake levels in Magny et al. 2004 and dying-off at *TOMO_south*) are in close temporal accord. The second phase of high lake levels, described by Magny et al. (2004) for c. 5600 – 5300

cal. BC does not fit the data of *TOMO_south*. The onset of the event is contemporaneous to a site chronology gap at *TOMO_south*, which is taken to indicate dry conditions rather than wet ones, and its end is contemporaneous to the beginning of a dying-off phase at *TOMO_south* (segment B, 5308 BC), which should indicate the beginning of a more humid period rather than its end.

5 The data of Schmidt et al. (2004) seems to be contradicting the record of *TOMO_south* at that time as well. They show data from 5600 BC on, which displays more or less humid conditions until about 5420 BC. This is within a chronology gap at *TOMO_south*, with the gap end (5404 BC) closely meeting the end of the relatively humid conditions observed in the data of Schmidt et al.(2004). The subsequent dry phase displayed by Schmidt et al. (2004) for c. 5400 – 5350 BC fits the indications from *TOMO_south* better, where the pines begin establishment contemporary to the dry phase beginning, and start dying off
10 only after the dry phases end, in 5308 BC. The two wet phases that follow according to Schmidt et al. (2004) (c. 5320 – 5000 BC and c. 4950 – 4900 BC) fit the dying-off phase of segment B at *TOMO_south* (5308 – 4902 BC), which also indicates humid conditions. The beginning of the above dying-off phase is temporally close to the beginning of the first of the two mentioned wet phases, and the end of the dying-off phase is contemporary to the end of the second. This is a very close agreement of the indications of the two dendrochronological records. The interjacent dry phase (c. 5000 – 4950 BC)
15 documented by Schmidt et al. (2004) is not reflected in *TOMO_south*.

Gunnarson et al. (2003) describe drier conditions for c. 4900-4800 BC. At *TOMO_south* a site chronology gap (4901 – 4838 BC) begins at the same time. The gap and the following phase of tree establishment comply with drier conditions.

More humid conditions are evident from pollen and peat data composed in context of a trackway (Bauerochse 2003), which is dendrochronologically dated (construction and maintenance 4629 – 4545 BC) (Bauerochse et al. 2012, Achterberg et al.
20 2015). The palaeo-botanical indications for increased humidity described by Bauerochse (2003) can thus be aligned to the dying-off phase of site chronology segment C (4712 – 4537 BC).

The third phase of high lake levels described by Magny et al.'s (2004) (c. 4400 – 3950 cal. BC) does not coincide with indications for increased humidity at *TOMO_south*. It begins within a long site chronology gap (4536 – 3987 BC) and ends before the beginning of the next dying-off phase. There are however, die-off phases of northwest German pines from other
25 bogs which date to c. 4490 – 4370 BC, 4230 – 4170 BC and c. 4010 – 3960 BC (Achterberg et al. 2015), contemporary to that phase of high lake levels described by Magny et al. (2004) for c. 4400 – 3950 cal. BC (Fig. 11 A and B). Turney et al. (2006) on the other hand state drier conditions for Ireland around c. 4250 BC, also dating within the *TOMO_south* site chronology gap 4536 – 3987 BC, which is in agreement with our interpretation of the *TOMO_south* data.

Palynological indications for increased humidity (Bauerochse 2003) are temporally anchored to after 3701 BC by a
30 dendrochronologically dated trackway (Bauerochse et al. 2012, Achterberg et al. 2015). This is within the dying-off phase of segment E at *TOMO_south* (3691 – 3614 BC), and thereby in agreement with its climatic indication. Around the same time Arbogast et al. (2006) as well identify climatic deterioration, a shift towards cooler and possibly more humid conditions for c. 3700 – 3250 BC. Magny et al. (2004) date the beginning of their fourth phase of high lake levels to c. 3700 cal. BC as well

(c. 3700 – 3250 cal. BC). Despite the low replication of segment E, these coherences make the indication of its dying-off phase appear quite valid.

The wet phase observed by Gunnarson et al. (2003) for c. 3600 – 3400 BC covers about the same time as segment F. This is not exactly analogue. However, both dying-off pulses of segment F thereby are within the wet phase documented for Sweden (Gunnarson et al. 2003), with the end of the wet phase meeting the end of the last dying-off pulse. This makes for an intermediate level of agreement.

Dreslerova (2012) points out a pronounced shift towards wetter, cooler and a more variable climate around c. 3550 cal. BC, reviewing numerous European studies of climate proxy for the Holocene. The *TOMO_south* dying-off phase 3496 – 3473 BC (segment F) may well relate to this event, which Dreslerova (2012) describes to be the beginning of a significant climatic phase.

The tree ring chronologies of *TOMO_south* do not cover the end of the Holocene Thermal Maximum (HTM), which occurred around c. 2350 cal. BC with a shift to cooler conditions (Seppä et al. 2005).

The trees of *TOMO_south* record water table change at one mire. The water table changes are largely driven by precipitation in the catchment area, but also affected by transpiration. This hydrological signal has a higher temporal resolution due to its local nature than an over-regional record (such as e.g. Achterberg et al. 2015) would. Even though the climatic drivers are the same in both cases, regional variations of rainfall etc are smoothed out in an over-regional record at the cost of precision, which would in turn be retained in a case study as this.

Even though the dating of the trees and their reactions to environmental change is precise to the calendar year, that does not necessarily apply to the dating of the related climatic causes. The water table rise observed in mires can be lagged behind the climatic shifts by months or even years (Edvardsson et al. 2016).

5 Conclusions

The trees, stratigraphically located at the fen-bog-transition, are viewed to stem from the former bog margin, being embedded by the expanding raised bogs *Sphagnum* peat. The tree dates document a raised bog expansion at the site for 6614 – 6436 BC (c. 250y), 6365 – 5733 BC (c. 750y) , 5308 – 4902 BC (c. 500y), 4712 – 4537 BC (c. 300y), 3895 – 3838 BC (c. 150y), 3691 – 3614 BC (c. 200y), 3496 – 3473 BC (c. 25y) and c. 3400 BC (4y). These phases of lateral raised bog growth likely occurred in periods of rather humid climate.

The shorter gaps between the later site chronologies are viewed insignificant, since they belong to poorly replicated periods documented at the sites margin. The three longer gaps between the earlier site chronology segments however are framed by well replicated sections represented in the central area of the site. These are interpreted to represent periods without significant (and lasting) raised bog advance, since the site is throughout covered with stumps and these were densely sampled. These chronology gaps relating to phases of apparent bog stagnation are 5733 – 5308 BC (328 y), 4902 – 4712 BC

(64y) and 4537 – 3895 BC (550y). There are however, dying-off phases recorded at other northwest German bogs within these periods. These periods therefore do not appear to have been throughout dry phases in the region.

The distribution of the tree stumps of various ages across the site supports the picture of subsequent bog advance embedding the tree stumps. The bog expanded from the south-east towards north-west according to the dating of the trees, the direction of bog growth largely reflecting the elevation of the mineral base below the peat. Rises of the sandy ground formed wooded islands within the bog, being successively covered by *Sphagnum* peat with much delay.

6 Acknowledgements

Thanks goes to Torsten Struck for support and generous help in field work, Jens Will for a first language check-up and Kerry-Ann Williams for a native speaker proofreading on short notice. We would particularly like to express our gratitude to ASB-Humus, especially Mr. Thuernau, for kindly allowing us to work on their peat extraction grounds, their helpful support of our research and for sharing their elevation data. Our sincerest gratitude goes to the German research society DFG for funding the work this manuscript is based on (projects LE 1805/2 and HA 4438/1). We also thank the reviewers and editor.

References

- Achterberg, I., Bauerochse, A., Giesecke, T., Metzler, A., Leuschner, H. (2015): Contemporaneousness of Trackway construction and environmental change: a dendrochronological study in Northwest-German mires. IANSA 6 (1), 19-29.
- Achterberg, I., Frechen, M., Bauerochse, A., Eckstein, J., Leuschner, H.H.: The Goettingen tree-ring chronologies of peat-preserved oaks and pines from northwest Germany. ZDGG – German Journal of Geosciences, DOI: [10.1127/zdgg/2016/0042](https://doi.org/10.1127/zdgg/2016/0042), 2016.
- Arbogast, R. M., Jacomet, S., Magny, M.: The significance of climate fluctuations for lake level changes and shifts in subsistence economy during the Late Neolithic (4300–2400 BC) in Central Europe. Vegetation History and Archaeobotany 15/4, 403–418, 2006.
- Bauerochse, A.: Environmental change and its influence on trackway construction and settlement. In: Bauerochse, A., Hassmann, H. (eds): Peatlands, archaeological sites – archives of nature, nature conservation, wise use. Proceedings of the Peatland Conference 2002 in Hannover, Germany, Rahden/Westf., 66-76, 2003.
- Bauerochse, A., Leuschner, H.H., Metzler, A.: Das Campemoor im Neolithikum. Jahrbuch für das Oldenburger Münsterland 2012, 61. Jahrgang, Heimatbund für das Oldenburger Münsterland, 2012.
- Behre, K.-E.: Landschaftsgeschichte Norddeutschlands- Umwelt und Siedlung von der Steinzeit bis zur Gegenwart. 308 p., Wachholtz Verlag Neumünster 2008.
- Bronk Ramsey, C.: Bayesian analysis of radiocarbon dates. Radiocarbon, 51 (1), 337-360, 2009.

- Bronk Ramsey, C., van der Plicht, J., Weninger, B.: 'Wiggle matching' radiocarbon dates. *Radiocarbon*, 43 (2a), 381-389, 2001.
- Dreslerova, D.: Human Response to Potential Robust Climate Change in the Territory of Bohemia (the Czech Republic). *IANSIA Interdisciplinaria Archaeologica – Natural Sciences in Archaeology* 3 (1), 43-55, 2012.
- 5 Eckstein, J., Leuschner, H.H., Bauerochse, A., Sass-Klaassen, U.: Subfossil bog-pine horizons document climate and ecosystem changes during the Mid- Holocene. *Dendrochronologia* 27, 129-146, 2009.
- Eckstein, J., Leuschner, H.H., Giesecke, T., Schumilovskikh, L., Bauerochse, A.: Dendroecological investigations at Venner Moor (northwest Germany) document climate-driven woodland dynamics and mire development in the period 2450–2050 BC. *The Holocene* 20, 2 , 231-244, 2010.
- 10 Eckstein, J., Leuschner, H.H., Bauerochse, A.: Mid-Holocene pine woodland phases and mire development – significance of dendrochronological data from subfossil trees from northwest Germany. *Journal of Vegetation Science* 22, 781-794, 2011.
- Edvardsson, J., Leuschner, H.H., Linderholm, H., Hammarlund, D.: South Swedish bog pines as indicators of Mid-Holocene climate variability. *Dendrochronologia* 30, 93-103, 2012a.
- 15 Edvardsson, J., Linderson, H., Rundgren, M., Hammarlund, D.: Holocene peatland development and hydrological variability inferred from bog-pine dendrochronology and peat stratigraphy – a case study from southern Sweden. *Journal of Quaternary Science* 27 (6), 553-563, 2012b.
- Edvardsson, J., Poska, A., van der Putten, N., Rundgren, M., Linderson, H., Hammarlund, D.: Late-Holocene expansion of a south Swedish peatland and its impact on marginal ecosystems: evidence from dendrochronology, peat stratigraphy and paleobotanical data. *The Holocene* 24 (4), 466-476, 2014.
- 20 Edvardsson, J., Stoffel, M., Corona, C., Bragazza, L., Leuschner, H.H., Charman, D. J., Helama, S.: Subfossil peatland trees as proxies for Holocene palaeohydrology and palaeoclimate. *Earth-Science Reviews* 163, 118-140, 2016.
- Ellenberg, H.: *Vegetation Mitteleuropas mit den Alpen*. 1096 p., Ulmer Stuttgart 1996.
- Gunnarson, B., Bogmark, A., Wastergård, S.: Holocene humidity fluctuations in Sweden inferred from dendrochronology and peat stratigraphy. *Boreas* 32, 347-360, 2003.
- 25 Holmgren, M., Lin, C.-Y., Murillo, J., Nieuwenhuis, A., Penninkhof, J., Sanders, N., van Bart, T., van Veen, H., Vasander, H., Vollebregt, M., Limpens, J.: Positive shrub-tree interactions facilitate woody encroachment in boreal peatlands. *Journal of Ecology*, 103, 58-66, 2015.doi: 10.1111/1365-2745.12331
- Hughes P., Barber, K.: Contrasting pathway to ombrotrophy in three raised bogs in Ireland and Cumbria, England. *The Holocene* 14 (1), 65-77, 2004.
- 30 Kobashi, T., Severinghaus, J.P., Brook, E.J., Barnola, J.-M., and Grachev, A.M.: Precise timing and characterization of abrupt climate change 8200 years ago from air trapped in polar ice. *Quaternary Science Reviews*, v. 26, 1212–1222, 2007.

- Krapiec, M., Margielewski, W., Korzenń, K., Szychowska-Krapiec, E., Łajczak, A.: Late Holocene palaeoclimate variability: The significance of bog pine dendrochronology related to peat stratigraphy. The Puścizna Wielka raised bog case study (Orawa – Nowy Targ Basin, Polish Inner Carpathians). *Quaternary Science Reviews*, 148, 192-208, 2016.
- 5 Leuschner, H.H., Sass-Klaassen, U., Jansma, E., Baillie, M.G.L., Spurk, M.: Subfossil European bog oaks: population dynamics and long-term growth depressions as indicators of changes in the Holocene hydro-regime and climate. *The Holocene* 12, 695–706, 2002.
- Leuschner, H.H., Bauerochse, A., Metzler, A.: Environmental change, bog history and human impact around 2900 B.C. in NW Germany – preliminary results from a dendroecological study of a sub-fossil pine woodland at Campemoor, Dümmer Basin. *Vegetation History and Archaeobotany* 16: 183-195, 2007.
- 10 Limpens, J., Holmgren, M., Jacobs, C., Van de Zee, S., Karofeld, E., Berendse, F.: How does tree density affect water loss of peatlands? A mesocosm Experiment. *PLOS One* 9 (3), 1-11, 2014.
- Magny, M.: Holocene climate variability as reflected by mid-European lake-level fluctuations and its probable impact on prehistoric human settlements. *Quaternary International* 113, 65-79, 2004.
- 15 Meier-Uhlherr, R., Schulz, C., Luthardt, V.: *Steckbriefe Moorsubstrate. 2., unveränd. Aufl., HNE Eberswalde (Hrsg.), Berlin 2015.*
- Metzler, A.: Moorarchäologie in Niedersachsen. In: Fansa, M.; Both, F. (Eds.): *Archäologie, Land, Niedersachsen: 25 Jahre Denkmalschutzgesetz – 400.000 Jahre Geschichte. Archäologische Mitteilungen aus Nordwestdeutschland, Beiheft 43; Landesmuseum Natur und Mensch Oldenburg, 471-475, 2004.*
- 20 Moir, A.K., Leroy, S., Brown, D., Collins, P.: Dendrochronological evidence for a lower water-table on peatland around 3200-3000 BC from subfossil pine in northern Scotland. *The Holocene* 20(6), 931-942, 2010.
- Nicolussi, K., Kaufmann, M., Melvin, T. M., van der Plicht, J., Schießling, P., Thurner, A.: A 9111 year long conifer tree-ring chronology for the European Alps: a base for environmental and climatic investigations. *The Holocene* 19(6), 909-920, 2009.
- 25 Nicolussi, K. & Schlüchter, C.: The 8.2 ka event – Calendar-dated glacier response in the Alps. *Geology* 40 (9), 819-822, 2012. doi:10.1130/G32406.1
- Overbeck, F.: *Das Große Moor bei Gifhorn – im Wechsel hygrokliner und xerokliner Phasen der nordwestdeutschen Hochmoorentwicklung. Niedersächsisches Amt für Landesplanung und Statistik, Veröffentlichungen Reihe A I, Band 41, Walter Dorn Verlag, Bremen-Horn, 1952.*
- 30 Overbeck, F.: *Botanisch-geologische Moorkunde - unter besonderer Berücksichtigung der Moore Nordwestdeutschlands als Quellen zur Vegetations-, Klima- und Siedlungsgeschichte. Karl Wachtelholz Verlag Neumünster, 1975.*
- Petzelberger, B., Behre, K.-E., Geyh, M.: Beginning and spread of raised bogs Northwest Germany – first results of a new project. *Telma* 29, 21-38, 1999.

- Riemer T.: Über die Varianz von Jahrringbreiten – Statistische Methoden für die Auswertung der jährlichen Dickenzuwächse von Bäumen unter sich ändernden Lebensbedingungen. *Berichte des Forschungszentrums Waldökosysteme, Reihe A* 121:1–375, 1994.
- 5 Reimer, P., Baillie, M., Bard, E., Bayliss, A., Beck, J., Blackwell, P., Bronk Ramsey, C., Grootes, P., Guilderson, T., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T., Hoffmann, D., Hogg, A., Hughen, K., Kaiser, K., Kromer, B., Manning, S., Niu, M., Reimer, R., Richards, D., Scott, E., Southon, J., Staff, R., Turney, C., van der Plicht, J.: IntCal13 and Marine 13 radiocarbon age calibration curves, 0–50,000 years cal BP. *Radiocarbon*, Vol. 55 (4), 1869–1887, 2013.
- 10 Rinn, F.: TSAP-Win: Time Series Analysis and Presentation for Dendrochronology and related Applications. Version 0.55 User reference. Heidelberg, Germany, 2003.
- Schmidt, B., Gruhle, W., Rück, O.: Klimaextreme in Bandkeramischer Zeit (5300 bis 5000 v. Chr.). Interpretation dendrochronologischer und archäologischer Befunde. *Archäologisches Korrespondenzblatt* 34/3, 303–308, 2004.
- Schweingruber, F.: *Tree Rings – Basics and Application of Dendrochronology*. 276 S. D. Reidel Publishing Company, Dordrecht, Holland, 1988.
- 15 Seppä, H., Hammarlund, D., Antonsson, K.: Low-frequency changes in temperature and effective humidity during the Holocene in south-central Sweden: implications for atmospheric and oceanic forcings of climate. *Climate Dynamics* 25, 285–297, 2005. DOI 10.1007/s00382-005-0024-5
- Shumilovskikh, L., Schlütz, F., Achterberg, I., Kvitkina, A., Bauerochse, A., Leuschner, H.H.: Pollen as a nutrient source in Holocene ombrotrophic bogs. *Review of Palaeobotany and Palynology* 221, 171–178, 2015.
- 20 Spurk, M., Leuschner, H., Baillie, M., Briffa, K., Friedrich, M.: Depositional frequency of German subfossil oaks: climatically induced fluctuations in the Holocene. *The Holocene* 12, 6, 707–715, 2002.
- Stenzel, V.: *Dendroökologische Untersuchungen zum räumlich-zeitlichen Wachstum eines subfossilen Kiefernwaldes im Toten Moor (bei Neustadt am Rübenberge in Niedersachsen)*. Bachelorarbeit, Abteilung Palynology und Klimadynamik, Albrecht-von-Haller-Institut für Pflanzenwissenschaften, Biologische Fakultät, Universität
25 Göttingen, 2013.
- Tahvanainen, T.: Abrupt ombrotrophication of a boreal aapa mire triggered by hydrological disturbance in the catchment. *Journal of Ecology* 99, 404–415, 2011.
- Thomas, E., Wolff, E., Mulvaney, R., Steffensen, J., Johnsen, S., Arrowsmith, C., White, J., Vaughn, B., and Popp, T.: The 8.2 ka event from Greenland ice cores. *Quaternary Science Reviews*, v. 26, 70–81, 2007.
- 30 Torbenson, M., Plunkett, G., Brown, D., Pilcher, J., Leuschner, H.H.: Asynchrony in key Holocene chronologies: Evidence from Irish bog pines. *Geology* 43, 799–802, 2015.
- Turney, C., Baillie, M., Palmer, J.: Holocene climatic change and past Irish societal response. *Journal of Archaeological Science* 33, 34–38, 2006.

Veski, S., Seppä, H., Ojala, A.: Cold event at 8200 yr B.P. recorded in annually laminated lake sediments in eastern Europe. *Geology* 32 (8), 681-684, 2004.

von Bülow: Die Bedeutung der Versumpfung für die Entstehung von Hochmooren in Deutschland. *Geologische Rundschau* 26 (4), 277-284, 1935.

- 5 Zackrisson, O., Nilsson, M.-C., Steijlen, I., Hörnberg, G.: Regeneration pulses and climate-vegetation interactions in nonpyrogenic boreal Scots pine stands. *Journal of Ecology* 83, 469-483, 1995.

Table 1: Chronology cover. For this table, the measured rings were used only, estimations of missing rings to pith or bark were not added. Only the trees with dendrochronological calendar dates are listed below.

Chronology segment name	A		B	C	D	E	F	total	
	A1	A2							
Color	Dark violet	Blue	Pink	Red	Orange	Yellow	Green		
Chronology cover [y BC]	6703 – 5733		5404 – 4902	4837 – 4537	3986 – 3838	3831 – 3614	3602 – 3403	Dispersed over 3301 years	
	6703 – 6436	6483 – 5733							
Tree dying-off [y BC]	6614 – 6436	6365 – 5733	5308 – 4902	4712 – 4537	3895 – 3838	3691 – 3614	3496 – 3473 3407 – 3403	1482 years	
Chronology segment length [y]	971		504	301	149	218	202	2345 years	
	268	752							
Chronology gap [y]		-	328	64	550	6	11	-	959 years
Replication [trees]	254		66	123	10	19	5	477 trees	
	53	201							



5 **Figure 1:** Location of the study site *TOMO_south*, indicated by a star.



Figure 2: View on site *TOMO_south*. The upper layers of peat have been removed. Numerous tree-stumps are protruding. Photo: Inke Achterberg.




Figure 3: Tree stumps at site *TOMO_south* grown on two elevation levels. The upper level trees are sometimes supported by their dead successors. Photo: Inke Achterberg.




Figure 4: Temporal distribution of pine trees at site *TOMO_south*. On top the chronology segments C14 and A1-F are indicated. The gaps between the site chronology segments (white) are labelled with their duration. The time covered by the site chronologies of *TOMO_south* is underlain in grey, for the floating chronology segment C14 striped in grey-white. The coloured horizontal lines indicate the lifespans of the individual dated trees from the site (measured rings coloured, estimated missing rings black). The trees of each chronology segment (Table 1) are shown in a different colour (A1: dark violet, A2: blue, B: pink, C: red, D: orange, E: yellow, F: green), the floating chronology segment C14 in grey. The colours are used accordingly in the spatial mapping (Fig. 5 and 6). The oaks from the site are displayed in black, below the pines. In [A] the trees are sorted by dying-off date. The three most prominent events of cumulative tree dying-off are highlighted by a dashed blue vertical line and labelled by year BC. Below, the periods from first to last tree dying-off (last measured ring) are indicated, labelled by years BC. The lighter grey backgrounds indicate when the trees of a chronology segment start to die off. In [B] the trees are sorted by germination date. Here, the lighter grey backgrounds indicate periods with limited tree establishment.





Figure 5: Reconstructed raised bog advance in terms of 'no more tree growth'. The coloured dots indicate in situ tree remains, their colour indicates the chronology segments they belong to. The hachures indicate areas with lac of dated in situ samples. The coloured areas show, where NO MORE trees grew at a certain period. In this sense, the oldest section is where only trees of the floating, radiocarbon dated segment C14 (beginning of the 7th millennium cal. BC) were found, which is coloured grey. The dark violet areas show, were no trees younger than the first group of trees with dendrochronological calendar dates, A1 (early 7th millennium BC), were found. Blue areas feature only trees from the late 7th and early 6th millennium BC (A2) and older. The pink area delivered trees from the second half of the 6th millennium BC and the beginning of the 5th (B). The red area shows were no trees grew after the first half of the 5th millennium BC (C). The orange indicates were, after that, only trees from 4000-3800 BC (D) were found. The yellow area still featured trees in the first half of the 4th millennium BC (E), and the green in the second half of the 4th millennium BC (F). The colours with the according dates are given to the right.



5

Figure 6: Elevations a.s.l. of the mineral base (sand) below the peat. Regularized spline interpolation of 99 measurements. The elevations are interpolated using a regularized spline. The actual measurements range from 36,42 to 38,16 m a.s.l.; thereby, the first and the last elevation class (green, below 35,50 m a.s.l., and white, above 38,51 m a.s.l.) are products of extrapolation only. The coring points are indicated by triangles. The dated in situ trees are shown as circles, for each chronology segment (Table 1) in a different colour (C14: grey, A1: dark violet, A2: blue, B: pink, C: red, D: orange, E: yellow, F: green).



Figure 7: Pine growth at a bog site in Finland. View across a stretch of treeless bog. In front are small trees, thinning out where the peat deepens. In the back large trees form a wooded 'island' where the mineral ground ascends. Photo: Inke Achterberg.




Figure 8: Peat stratigraphy. Six transects are shown with the peat stratigraphy and elevation a.s.l. of the cores. In the sky view, site *TOMO_south* is outlined in pink. The locations of the displayed peat cores are indicated by black triangles, yellow lines connecting the transects. Peat stratigraphy for cores without elevation measurement is not shown. Please note, that cores D8, F1 and F2 were taken at the border to the plot, where peat was left standing significantly higher than within the plot where the tree layer was exposed. Underlying satellite image: ©2008 Google Earth, image ©2009GeoContent, ©2009 Tele Atlas.

5



Figure 9: Strong downward roots below a tree stump *TOMO_south* (tree life 5396-5230 BC, chronology segment B). Photo: Inke Achterberg.





Figure 10: Tree ring widths curves of chronology segment F. The mean value of the tree ring widths is indicated by a horizontal line for each curve, the area below mean is filled black to highlight growth depressions. One sample is not shown, because it is missing several rings to bark and thereby the record of the final years of that trees life. All trees show narrow ring widths prior to their deaths. These growth depressions are not reflected equally in the surviving trees. This illustrates the locality of the calamity.

5



5 **Figure 11 A:** Comparison of the replication of *TOMO_south* northwest German pine chronologies with other climate-related records. From top to bottom: oak deposition at river Main (Spurk et al. 2002); scores of higher and lower lake levels in Mid-Europe (Magny et al. 2004); Replication of the northwest German pine chronology including the pines of *TOMO_south*, which are highlighted red.




Figure 11 B: Comparison of the phases observed at *TOMO_south* with those of other studies.

Table 2: Comparisons with other palaeoenvironmental studies.

Publication	Proxy	Location	Time	Indication	<i>TOMO_south</i>	fit
Thomas et al. (2007)	Ice cores chemistry and stable isotope	Greenland	Begin c. 6300 cal. BC	Cooling	Dying-off phase c. 6350 – 5750 BC; strong dying-off pulse at c. 6315 BC	+
Spurk et al. (2002)	Dendrochronological	West Germany	c. 6270 – 6000 BC	Poor growth and regeneration conditions	Frequent tree dying-off from 6250 BC to 6157 BC	+
Kobashi et al. (2007)	Ice core (GISP2) methane and nitrogen isotopes	Greenland	Begin c. 6225 cal. BC	Cooling	Frequent tree dying-off from 6250 BC to 6157 BC	+
Torbensoen et al. (2015)	Dendrochronological; pine establishment at three bog sites	Ireland	From 6210 BC on		Frequent tree dying-off from 6250 BC to 6157 BC	+
Veski et al. (2004)	Varve	Estonia	6250 – 6150 varve years BC	Maximum cooling	Frequent tree dying-off from 6250 BC to 6157 BC	+
Nicolussi et al. (2009)	Dendrochronological	Alps	c. 6250 – 6050 BC	Low replication	Frequent tree dying-off from 6250 BC to 6157 BC	+
Veski et al. (2004)	Varve	Estonia	6080 varve years BC	Abrupt end of the 8.2k cooling period		
Achterberg et al. (2016)	Dendrochronological	Northwest	6177 – 6060 BC	Gap in bog oak		

		Germany		chronology		
Nicolussi & Schlüchter (2012)	Dendrochronological	Alps	c. 6175 BC	Glacier advance	Frequent tree dying-off from 6250 BC to 6157 BC	+
Magny (2004)	Radiocarbon, dendrochronological and archaeological dates	France and Switzerland	c. 6350 - 6100 cal. BC	High lake levels (I)	Start is beginning of the dying-off phase of segment A2 at <i>TOMO_south</i> (6365 BC)	+
Magny (2004)	Radiocarbon, dendrochronological and archaeological dates	France and Switzerland	c. 5600 - 5300 cal. BC	High lake levels (II)	End coincides with beginning of dying-off phase (5308 - 4902 BC)	+/-
Schmidt et al. (2004)	Dendrochronological	West Germany	(5600) - 5420 BC	More or less humid	Long gap (5732 - 5405 BC)	-
Schmidt et al. (2004)	Dendrochronological	West Germany	c. 5410 - 5330 BC	Dry phase	Tree establishment and growth (no dying-off)	+
Schmidt et al. (2004)	Dendrochronological	West Germany	c. 5320 - 5000 BC	Wet phase	Beginning at the same time as dying-off phase c. 5300 – 4900 BC begins	+
Schmidt et al. (2004)	Dendrochronological	West Germany	c. 5000 - 4950 BC	Dry phase	(Not reflected clearly in the data of <i>TOMO_south</i>)	-
Schmidt et al. (2004)	Dendrochronological	West Germany	c. 4950 - 4900 BC	Wet phase	End at the same time as dying-off phase c. 5300 – 4900 BC ends	+
Gunnarson et al. (2003)	Dendrochronological	Sweden	c. 4900 - 4800 BC	Drier conditions	Found accordingly for <i>TOMO_south</i> (c. 4900 - 4700 BC)	+
Bauerohse 2003	Palynological implications, context of dendro-dated find	NW Germany	Context of find dated to 4629 - 4545 BC	More humid conditions	dying-off phase 4712 - 4537 BC	+
Magny (2004)	Radiocarbon, dendrochronological and archaeological dates	France and Switzerland	c. 4400 - 3950 cal. BC	High lake levels (III)	Gap between the site chronology segments (c. 4500 - 3900 BC)	-
Turney et al. (2006)	Dendrochronological	Ireland	c. 4250 BC	Drier conditions	Within the site chronology gap 4536 - 3987 BC	+
Spurk et al. (2002)	Dendrochronological	West Germany	c. 4160 - 3870 BC	Reduced activity of the river Main	Gap 4536 - 3987 BC, which might point towards drier conditions as well. dying-off phase at <i>TOMO_south</i> from 3895 BC on, 25 years prior to the end of the period identified by Spurk et al. (2002).	+/-
Bauerohse (2003)	Palynological implications, context of dendro-dated find	NW Germany	after 3701 BC	More humid conditions	dying-off phase at <i>TOMO_south</i> from 3691 BC to 3614 BC	+
Arbogast et al. (2006)	Dendrochronological dates from archaeological layers	Swiss Alps, French and German	3700 - 3250 BC	Lake level rise; climatic deterioration, cool and possibly humid conditions	Beginning dying-off phase segment E (3691 BC), (covers dying-off phases segments E and F)	+
Magny (2004)	Radiocarbon, dendrochronological and archaeological dates	France and Switzerland	c. 3700 - 3250 cal. BC, maximum 3350 - 3250 cal. BC	High lake levels (IV)	Start is beginning of dying-off phase segment E (3691 BC)	+
Dreslerova (2012)	Multi-proxy (review)	Europe (multi site)	c. 3550 cal. BC	Pronounced shift towards wetter, cooler and a more variable climate	Shift from stagnating bog growth to a phase of bog expansion found for c. 3500 BC	+
Gunnarson et al. (2003)	Dendrochronological	Sweden	c. 3600 – 3400 BC	Wet phase	Contemporary to cover of segment F (3602-3403 BC), including two dying-off pulses	+/-

Magny & Haas (2004)	Dendrochronological date, pollen and archaeological	Switzerland	c. 3370 BC	Lake level rise	Last trees at <i>TOMO_south</i> died off 35 years - previously
---------------------	--	-------------	------------	-----------------	---