Review 1

Interactive comment on "Periodic input of dust over the Eastern Carpathians during the Holocene linked with Saharan desertification and human impact" by Jack Longman et al. Anonymous Referee #1 Received and published: 24 February 2017

The manuscript presents profiles of major elements and paleoecological indicators from an ombrotrophic peat bog in Romania. The paleoclimate records are associated to mineral dust and the discussion is focused around possible interpretations that try to disentangle local from distal signals. The authors interpret the history of the site in function of a superposition of changes on diverse spatial scales, from that of local hydrology to the large scale / hemispheric patterns derived from Greenland ice cores or North Atlantic marine sediment records. The topic is of interest to the paleoclimate community. The work appears well structured and its presentation in the manuscript is generally clear. I think that a better discussion of uncertainties would improve the manuscript.

We thank the reviewer for their kind words, and constructive review.

General comment

My comment about improving the discussion of uncertainties is articulated in two parts, that have to do with (1) the quantification of the dust flux and (2) the attribution of potential sources.

(1) For the purpose of estimating the dust flux, only the concentrations of Ti from ICPOES are used, although a semi-quantitative comparison is carried out against three major elements counts from XRF. Other studies trying to estimate dust from peat bog records discussed the uncertainties related to this issue (e.g. Marx et al., 2009; Kylander et al., 2016). Please discuss more in detail these aspects, and if possible provide some estimates of the uncertainties.

This is a good point, and I order to address the reviewer's comments, we have included a summary outlining the uncertainties within the updated manuscript, lines 209-219:

"It must be noted here that using Ti alone in dust flux calculations does not allow for reconstruction of all minerals related to dust deposition. Ti, which is lithogenic and conservative, is a major component in soil dust, particularly within clay minerals (Shotyk et al., 2002), but may not be associated with other dust-forming minerals, including phosphates, plagioclase and silicates (Kylander et al., 2016), although our records of K and Si may help indicate changes in deposition rates of these minerals (See Mayewski and Maasch, 2006). As a result, we are unable to infer specific mineral-related changes in the composition of dust. However, Ti alone will record changes in the intensity of deposition of the main dust-forming minerals (Sharifi et al., 2015; Shotyk et al., 2002), and variations in K and Si (particularly with local K- and Si-rich dacites a possible dust source) may further indicate the influx of minerals which are not associated with Ti. Such an approach has been applied successfully to studies of changing dust influx (e.g. Allan et al., 2013; Sapkota et al., 2007; Sharifi et al., 2015), with each study able to identify periods of high and low dust deposition from Tiderived dust flux alone."

(2) The attribution of potential sources is based on a simple analysis of correlations between major elements counts, as discussed by the authors, and is also supported by the interpretation of the evidence from indicators such as testate amoeba and pollen. Nonetheless, it would only be by looking at dust size distributions at the same time, that one could derive more firm conclusions about distal versus local contributions to the dust budget (e.g. Mahowald et al., 2014). If possible, include this kind of data, otherwise please discuss this aspect in the text.

We attempted to address this point via grain size analysis on a Malvern Mastersizer 2000. Unfortunately, due to limited sample material, and the very high organic matter content, obscuration values on the laser particle counter were too low for reliable results to be presented. As Kylander et al. (2016) explain, this is a recurring issue with dust reconstruction within peat cores.

The manuscript has been updated to reflect these attempts between lines 164-169:

2.10 Grain Size

"In an effort to indicate distal versus local inputs to the bog via the dust particle size, grain size analysis was attempted using a Malvern Mastersizer 2000. Unfortunately, as also observed in previous studies (Kylander et al., 2016) due to the lack of available sample material, and low minerogenic matter (and correspondingly high organic matter) present in the samples, satisfactory obscuration values were not achieved for most analyses."

Specific comments

p. 3, 103-106. Please describe how the cores were packed and stored in the phase between recovery and analyses.

Adjustment made to lines 105-107:

"Upon recovery, the material was wrapped in clingfilm, transported to the laboratory, described, imaged, and subjected to further analyses. The core was stored at 3°C"

p. 4, 120-121. Later in the text (line 179) you also mention different detection limits for the different elements. Please provide all this kind of information in the same place in the text, and try to mark it visually in the plots.

The value of 50cps is not a detection limit, it is merely guidance for the interpretation of the results. Indeed, studies have been presented displaying variations not related to noise in Mg analysis, where typical cps values are as low as 5-10 (Dulski et al., 2015). With our data it is clear some fluctuations below 50cps are not just noise, as they are echoed in all three elements, and even within the ICP-OES analysis.

Regardless, since we have now normalised the data with respect to the scattering effect of organic material, this sentence no longer applies to our data and has been removed.

Figures 3 & 4. I think a slight confusion can arise because of the way some of the records are organized between these two figures. For instance, it would be useful to see in Figure 3 the Ti concentration profile from ICP-OES along with the major elements counts from XRF, rather than directly the dust flux which weights in the sedimentation rate and bulk density profiles. On the other hand it would be nice to have all the factors defining the dust profile in the same figure, i.e. Ti concentration profile from ICP-OES, sedimentation rate and bulk density profiles.

Figure 3 has been updated, with all the factors defining the dust flux; raw ICP-OES values, density and sedimentation rate, displayed on one figure. As such, density has been removed from Figure 4.

p. 4, 126. What is the depth span of each sample? i.e. was the full core analyzes, or just portions of it?

Lines 127-128 *have been updated to indicate each sample was* 1*cm*³ *and that the entire core was analysed:*

"ICP-OES analysis was carried out on 105 samples of 1cm3 of sediment, roughly every 10 cm, through the entire core."

p. 5, 160. Harmonize with what you say at line 118. p. 5, 175. What do you mean by significant? Did you apply some statistical test? Otherwise perhaps change with "visible".

The designation "significant" has now been removed. Additionally, we now clarify the reasoning behind the denotation of dust events in lines 183-187:

"Such zones are identified as an increase in two or more of the elements above the background deposition (K > 0.001, Si > 0.001 and Ti > 0.004, see dashed line on figure 3). These intervals are further discussed as reflecting major dust deposition events, and are referenced in the remainder of the text using the denotation D01-D10 (Fig. 3). Two exceptions, at the base of the core, close to the transition from lake to bog, and the last 1000 years, due to high noise, are not highlighted."

p. 6, 190. Please discuss the uncertainties in estimating the dust flux.

These have been outlined earlier in the response, and the discussion has been added to the text (lines 209-219)

p. 6, 208-211. It is not clear at this point what is the contribution of this kind of analysis to the work.

Since we do not use the magnetic susceptibility values within our interpretation, these data have been removed from the manuscript, and from figure 4.

p. 7, 238. The dust "Dn" events are selected based on the occurrences of at least one of the elements form the XRF scan, so it is a bit weird to go through the text until this point with some apparent inconsistency between the discussion of peaks D0 to D3, which are not evident and sometimes in anti-phase the what you call dust record. Please either change your definition of dust event (D) or harmonize the text.

In light of comments from reviewer 2, we have removed D0 from our interpretation of dust events, since it may be simply related to minerotrophy. For events D1-D3, as outlined, we have identified them via one of the elements increasing. We agree some confusion may arise with the interpretation of the 'dust record', which is generally based on the Ti-derived dust flux. As such, we have amended the manuscript on lines 366 & 408, replacing "dust record", with "Ti-derived dust record"

When we are considering the results of more than simply the dust flux, we retain the use of the term "dust record".

p. 7, 244-245. As you discuss below, there is not a correlation, so please rephrase with something like "we compare the timing of the identified dust prepositional events with periods ..."

Adjustment made to lines 265-266:

"Firstly, it is noteworthy that five of the identified dust depositional events may be compared to periods of Rapid Climate Change (RCC)"

p. 7, 260. "Appear to indicate": maybe it would be better "are consistent with", in relation to my comment about the missing information on dust particle size distributions. Please include this kind of data if possible, otherwise add a discussion paragraph about this issue.

"Appear to indicate" has been replaced with "are consistent with" on line 284.

As outlined earlier in this response, and in the manuscript lines 166-169, particle size variations could not be obtained from this highly organic record.

Figure 5. Check the units of the upper curves (the two cores from McGee et al., 2013): I think you reported the values with the wrong units, which I think are g/cm2/kyr in the original publication, so there is a factor ten difference. In fact, on this scale you have the same dust flux, if not lower, in the North African plume and Belgium or Romania, which would be weird.

This has been checked, and the reviewer is correct-Figure 5 has been revised accordingly

p. 8, 266-268. It is not very clear what is the relation between these two studies, please rephrase.

To clarify these studies, their relationship, and locations, the manuscript has been adapted in lines 289-292:

"Saharan dust in Atlantic marine cores strongly increased at this time, with a 140% rise roughly at 5500 cal yr BP observed on the western Saharan margin (Adkins et al., 2006) with another study indicating a rise by a factor of 5 by 4900 cal yr BP in a selection of similarly-located sites (McGee et al., 2013)."

p. 8, 279-289. Again, particle size distributions would help clarify these issues.

We tend to agree particle size data would help the interpretation, but unfortunately we have been unable to attain it.

p. 8, 291-298. Earlier in the text you mentioned the different mobility of these major elements, and how the similarity of their profiles supports the ombrotrophic nature of the peat bog. Please clarify how this is coherent with your analysis here, which is instead based on the difference between the same elements.

These comparisons are not made with regards to any discussion on ombrotrophy. We maintain the similarity of the records, as evidenced by the strong correlations is indicative of the ombrotrophy of the site. However, since the elemental records are not perfectly correlated, and there are periods within which one of the elements is more enriched relative to others, we believe this approach is valid for distinguishing the geochemical signature of the events.

p. 9, 310-313. It would be interesting to compare Type 2 events with the background signature of non-D periods.

This is an interesting idea, and we have implemented it, with results indicating background peat deposition is similar in makeup to Type 2 events- further hinting at the local nature of such deposition. The manuscript has been updated to include these data in lines 339-342:

"The local nature of such deposition is emphasised by the similarity the depositional signal to background values; the elemental composition outside of dust events. For all data points not considered to be related to dust (or the minerotrophic lowermost section), the Ti-K regression is low (r2= 0.1513) with a gradient of 0.0863."

p. 9, 314. "Fig. 8" should probably be "Fig. 6", please check.

The reviewer is correct; this has been corrected on line 343

p. 12, 429. Was the data interpolated somehow before performing wavelet analysis? What is the pace / temporal resolution of the time series fed to the wavelet analysis software?

Before performing wavelet analysis, we used Gaussian interpolation with a timestep of 4 years and a window of 12 years.

This is included in lines 161-163:

"For this analysis, the lithogenic normalised elemental data from ITRAX measurements (Ti, K, and Si) was interpolated to equal time steps of four years using a Gaussian window of 12 years."

References

S.K. Marx, H.A. McGowan, B.S. Kamber, Long-range dust transport from eastern Australia: a proxy for Holocene aridity and ENSO-type climate variability, Earth Planet. Sci. Lett., 282 (2009), pp. 167–177.

Malin E. Kylander, Antonio Martínez-Cortizas, Richard Bindler, Sarah L. Greenwood, Carl-Magnus Mörth, Sebastien Rauch, Potentials and problems of building detailed dust records using peat archives: An example from Store Mosse (the "Great Bog"), Sweden, Geochimica et Cosmochimica Acta, Volume 190, 1 October 2016, Pages 156-174, ISSN 0016-7037, http://dx.doi.org/10.1016/j.gca.2016.06.028.

Mahowald, N., S. Albani, J. F. Kok, S. Engelstaedter, R. Scanza, D. S. Ward, and M. G. Flanner (2014). The size distribution of desert dust aerosols and its impact on the Earth system. Aeolian Research, 15, 53-71, doi: 10.1016/j.aeolia.2013.09.002.

Review 2

Interactive comment on "Periodic input of dust over the Eastern Carpathians during the Holocene linked with Saharan desertification and human impact" by Jack Longman et al. N. Fagel (Referee) nathalie.fagel@ulg.ac.be Received and published: 1 March 2017

The ms presents a very interesting and complete dataset on Holocene dust reconstruction from East European peat record. The ms is well organized with clear aims. The text is too the point. The approach is quite innovative. See the attached pdf with several remarks and/or suggestions on the text and the figures. For my point of view, two mains points should be completed.

First on the methodological approach I suggest to present the calibration between ITRAX and ICP-OES data to better constrain the proposed approach. I will also discuss the potential influence of porosity/humidity changes in the measured peat section on the number of ITRAX counts. Indeed elemental ITRAX data are often normalized, for instance by the number of measured counts in order to take into changes in peat density/porosity but also on surface irregularities. I would explain why this approach is not useful.

We thank Dr. Fagel for the comprehensive review of our work. We fully agree, and have addressed both comments in the revised manuscript. Firstly, we now present the correlation, and significance of said correlation, between ICP-OES and ITRAX.

Secondly, we have normalised our ITRAX data (by incoherent + coherent scattering) with respect to the scattering effect of density, porosity and organic matter changes. As such, all data is now presented as normalised counts per second, and all corresponding further analysis has been performed on such data. Please see the comments below for exact location of added text, and adapted figures.

Second I would complete the text about the relationships between the identified dust events, the wet/dry, cold/warm conditions and the related figures.

We have also addressed these issues; see below for text additions, and figure adaptations

Hoping the review will be helpful to improve this interesting manuscript. Sincerely, Nathalie Fagel Please also note the supplement to this comment: <u>http://www.clim-past-discuss.net/cp-2017-6/cp-2017-6-RC2-supplement.pdf</u>

Specific comments from attached Supplement:

Line 26 unclear older or younger than 6,1 kyr?

"Since" 6100 yr BP; manuscript has been adapted to reflect this

Lines 125-126 For my pint of view those data must be more exploited in the ms. Why do you not present the calibration between ITRAX and ICP data?

This is a valid point and we have performed such measurements. Due to the differences in resolution of the data sets, we have performed a Gaussian interpolation at 100 yr steps, with a window of 300 years in order to put both records on the same timescales. The correlation (Pearson's R, n=105) of this is 0.2649, with a p-value of 1.416×10^{-8} , indicating a significant relationship between the two methods. The apparent weakness of the correlation is likely a function of the early half of the record, where low values are observed in both methods, and so the correlation is simply noise, and so we do not expect a perfect correlation. We have attached a graph to display this here:



Such correlation data has also been added to the manuscript on lines 203-206:

"To facilitate comparison, we bring both records on the same timescale using a Gaussian interpolation with 100 year time steps and a 300 year window. Pearson's r=0.2649, with a p-value of <0.001, indicative of a significant correlation (See SI 4)."

Lines 164-165 You should give the age of the lithological transition. I would suggest to add a schematic lithological column in regard with Fig 3.

The age of the lithological transition has been added to the manuscript on line 173: "transition from a wetland into a bog, at approximately 10,330 yr BP"

A lithological description, and full core image was included in the supplementary information: SI1 & SI2

Lines 175-176 (1) The identification of the D events must be precise. It is not clear why you merge some peaks into one D event (e.g., D1, D8).

Dust events were identified based on the increase in two or more of the ITRAX-derived element concentration profiles (see caption on Fig. 3). For each element, a cutoff point was identified, with K=0.001, Si=0.001 and Ti=0.004. There are two exceptions, the very earliest section of the core, close to the minerotrophic-ombrotrophic transition, and in the most recent 1000 years, where the noise is higher. Some dust events do indeed appear to merge peaks, this is because we believe they are related to the same event, as they are very close to each other chronologically. An explanation of the denotation of dust events is now included on lines 183-187:

"Such zones are identified as an increase in two or more of the elements above the background deposition (K > 0.001, Si > 0.001 and Ti > 0.004, see dashed line on figure 3). These intervals are further discussed as reflecting major dust deposition events, and are referenced in the remainder of the text using the denotation D01-D10 (Fig. 3). Two exceptions, at the base of the core, close to the transition from lake to bog, and the last 1000 years, due to high noise, are not highlighted."

(2) What is the age uncertainty for each interval according to your age model? It is important to precise since some events cover less than 200 years.

The age model uncertainties range from approximately 20 years at the uppermost sections of the core, to around 150 years by the base of the core (See Fig. 2 and Table 1). As such, events lasting 200 years should be discernible at this resolution.

This has been added to the manuscript on lines 175-176: "Age model uncertainties range from 20 years in the uppermost sections to 150 years at the base of the core."

(3) On figure 3, I would suggest to add a secondary scale for dust flux in order to check for smaller peaks.

We have added such a figure to the supplementary information- see SI 3.

Line 192 You must give more arguments to support this assumption. Do you calculate the correlation coefficient between the two datasets? I would add the figure, even as supplementary data. See also notes on fig. 3 and 4.

Such data has been added to lines 203-206. See above for our description of the process, and a simple graph of said comparison (also added to the manuscript as SI 4).

Line 204 add a synthetic lithological column on Fig; 4 to indicate the transition

See SI 2 for lithological column

Line 225 text unclear.

Text has been altered on lines 246-247:

"These persist only for the period in which each element is enriched, with such cycles particularly evident within the last 6000 years."

Line 230 Text unclear. Indeed the similar profiles in the three elements emphasize that there is no post-depositional elemental mobility: all elements behave as conservative in the studied peat. I would reorganize the paragraph.

We have inserted the following into line 254:

" indicating the conservative behaviour of such elements in the studied peat."

Lines 247-248 For your record I would suggest to describe deeper the relationships between air temperature and/or humidity and dust flux. In the Misten dust record most dust peaks coincide with cold events.

We agree such a discussion improves the manuscript, and so we have adapted lines 268-276 to reflect this:

"For example D8, between 3450-2800 cal yr BP falls Europe-wide cold period (Wanner et al., 2011). Such cold-related dust deposition has been observed previously in western Europe. However, within Mohos such a compassion may not be drawn for the majority of dust events. For example, event D9 (860-650 cal yr BP) occurs during the Medieval Climate Anomaly, a period of generally higher European temperatures (Mann et al., 2009) but also one of intense human impact on the environment through deforestation and agriculture (Arnaud et al., 2016; Kaplan et al., 2009). Furthermore, such events within the Misten record (Allan et al., 2013) were also linked to low humidity, whereas the Mohos TA (Fig. 4) record indicates locally wet conditions. This suggests that dust depositional events in this region are a result of a complex interplay of environmental conditions in the dust source areas, rather than simply reflecting locally warm or cold, or even wet or dry periods."

Lines 260-261 also observed in other bog records (misten, EGR)

We have revised the text on lines 284-285 to reflect this:

"a major shift in the controls of dust production and deposition at this time, a shift observed in peatderived dust records from Western Europe (Allan et al., 2013; Le Roux et al., 2012)."

Line 299 it could reflect the minerotrophic stage of the peat, with no meaining on dust flux.

Type 1 deposition only occurs within D10- the most recent dust event, not D0, the event close to the minerotrophic stage of peat.

We agree however, that D0 may therefore not be a dust event, but related to minerotrophy, and the transition to ombrotrophy, and so we now do not discuss it as a dust event.

Line 344 not clear with double (.

Text adapted line 372: "(Gradient of samples pre-10,500 yr BP = 0.7429, D10= 1.0637)."

Lines 353-356 D10 is older or very close to the minerotrophic/ombrotrophic transition. I would not give to much interpretation in term of dust flux for this first D event. The peat at that time may be affected by surface runoff.

As mentioned above, we agree with the reviewer's interpretation regarding D0, and so we do not discuss it as a dust event.

Line 373 you must first remove signal from volcanic eruption observed in EGR record (see Le Roux et al., 2012).

Text has been adapted to reflect this in lines 401-403:

"with a large dust flux peaks identified in Switzerland between 9000-8400 cal yr BP with other volcanic eruption-related peaks (See Fig. 5), when there is little evidence of dust input into Mohos."

Lines 414 and 418 Fig .7 - But there are still several meter variations in water table: it is significant, no? Moreover most D events during this interval coincide with shallower water table. Such wetter conditions may limit local erosion and favor distal input. We observe the same relationship for the Misten bog (see Allan et al., 2013).

The DWT values are in cm, something which has now been indicated on the figures. As such, the variations in this period only indicate changes of roughly 2-5 cm change at most. The second point regarding such wet conditions favouring distal deposition is valid, however, and so we have adapted the text accordingly on lines 448-449:

"Such wetter conditions also limit local drought-related erosion, and so may be further evidence of distal dust input at this time (Allan et al., 2013)."

Lines 427-428 you may add that the local input erase any distal input in that case.

The text has been adapted lines 458-459:

"local dust mobilisation, with K-rich dust present at this time, with local input potentially erasing some distal signals."

Line 436 The time interval is quite short to evidence cyclcity > 1 kyr. i would suggest to keep "millenial cyclicity".

Text has been adapted in lines 467 and 470 to reflect this

Lines 460-463 I would suggest to rewrite point 2. Here it overlaps with the point 3. I would say that two intervals are identified wit different major dust control: 1) dust input over the older interval reflects more local conditions; 2) distal input for the younger interval.

In order to improve the clarity, we have re-written this point, on lines 492-494:

"The two intervals before and after this shift are indicative of a change in major dust controls. For the period prior to 6100 yr BP, dust input is reflective of more local controls, whilst the most recent 6100 yr BP of deposition may be linked to more distal forcings."

Comments on Figures

Figure 3: any influence of density changes in the measured counts? see note for figure 4.

As mentioned above, we have normalised all ITRAX data (relative to scattering) to take into account density shifts.

Figure 4: Do you calculate ash content after 950° calcination? It could also be a good indicator for mineral matter in your peat.

Unfortunately, we did not measure LOI after 950°, as we were primarily interested in organic matter, not CaCo₃ content.

Since ITRAX was performed on the fresh core section how do you take into account the density and/or porosity changes in the measured counts? I suggest to discuss this point in the ms. What could be the influence of the average 20% variation of the bulk density on the elemental counts? How could you normalize each ITRAX elemental profile?

As outlined earlier, in order to circumvent these issues, we have normalised to the scattering values of the ITRAX, after Kylander et al., 2011.

What is the correlation between ITRAX counts and ICP measurements?

The correlation has been calculated, and is presented in lines 203-206:

"To facilitate comparison, we bring both records on the same timescale using a Gaussian interpolation with 100 year time steps and a 300 year window. Pearson's r=0.2649, with a p-value of <0.001, indicative of a significant correlation (See SI 4)"

Figure 5: I would suggest to improve this figure. Add the dust events as grey bars as identified in the two ms of Allan et al. 2013 and Le Roux et al. 2012. Some volcanic events are reported in EGR record: they have to be indicated.

In our opinion adding the dust events as indicated by the other two manuscripts clutters the figure. The data is still presented, and so it is hopefully still clear when those studies indicate increased dust. The volcanic events have been added.

Figure 6: add arrows on the vertical scale trend to indicate wetter or dryer local conditions as identified by the reconstructed water table.

These have been added to all figures displaying the TA record.

Table 2: ? add digitsAll issues relating to digits here have been resolved

Additional Reference

Dulski, P., Brauer, A., Mangili, C., 2015. Combined μ-XRF and Microfacies Techniques for Lake Sediment Analyses, in: Croudace, I.W., Rothwell, R.G. (Eds.), Micro-XRF Studies of Sediment Cores. Springer, Dordrecht, pp. 325–349.

Periodic input of dust over the Eastern Carpathians during the Holocene linked with Saharan desertification and human impact

Jack Longman¹, Daniel Veres², Vasile Ersek¹, Ulrich Salzmann¹, Katalin Hubay³, Marc Bormann⁴, Volker Wennrich⁵, Frank Schäbitz⁴

¹ Department of Geography, Northumbria University, Newcastle-Upon-Tyne, United Kingdom

² Romanian Academy, Institute of Speleology, Clinicilor 5, Cluj-Napoca, Romania

³ Hungarian Academy of Science - Institute for Nuclear Research, Hertelendi Laboratory of Environmental Studies, H-4026 Debrecen, Bem ter 18/C, Hungary

⁴ Institute of Geography Education, University of Cologne, 50931 Köln, Germany

⁵ Institute of Geology and Mineralogy, University of Cologne, 50674 Köln, Germany

Correspondence to: Jack Longman (jack.longman@northumbria.ac.uk) and Daniel Veres (daniel.veres@ubbcluj.ro)

Abstract. Reconstructions of dust flux have been used to produce valuable global records of changes in atmospheric circulation and aridity. These studies have highlighted the importance of atmospheric dust in marine and terrestrial biogeochemistry and nutrient cycling. By investigating a 10,800-year long paleoclimate archive from the Eastern Carpathians (Romania) we present the first peat record of changing dust deposition over the Holocene for the Carpathian-Balkan region. Using qualitative (XRF core scanning) and quantitative (ICP-OES) measurements of lithogenic (Fe, K, Si, Ti) elements, we identify 10 periods of major dust deposition between: 9500-9200, 8400-8100, 7720-7250, 6350-5950, 5450-5050, 4130-3770, 3450-2850, 2000-1450, 800-620, and 60 cal yr BP to present. In addition, we used testate amoeba assemblages preserved within the peat to infer local palaeohydroclimate conditions. Our record highlights several discrepancies between eastern and western European dust depositional records, and the impact of highly complex hydrological regimes in the Carpathian region. Since 6100 cal yr BP, we find that the geochemical indicators of dust flux become uncoupled from the local hydrology. This coincides with the appearance of millennial-scale cycles in the dust input and changes in geochemical composition of dust. We suggest this is indicative of a shift in dust provenance from local/regional (likely loess-related) to distal (Saharan) sources, which <u>coincide</u> with the end of the African Humid Period and the onset of Saharan desertification.

1 Introduction

Atmospheric dust plays a major role in oceanic and lacustrine biogeochemistry and productivity (Jickells, 2005) by providing macronutrients to these systems (Mahowald et al., 2010). Furthermore, climatically dust plays a role in forcing precipitation (Ramanathan, 2001; Yoshioka et al., 2007) and in moderating incoming solar radiation. As such, reconstructions of past dust flux are <u>an</u> important <u>tool</u> to understand Holocene climate variability, biogeochemical cycles, and the planet's feedback to future changes in atmospheric dust loading.

The link between atmospheric circulation patterns and dust input has been studied intensively (Allan et al., 2013; Kylander et al., 2013a; Marx et al., 2009; Le Roux et al., 2012) with clear evidence of climate variations linked with the dust cycle (Goudie and Middleton, 2006). Generally, dust is produced in arid zones (Grousset and Biscaye, 2005) and may be transported thousands of miles before deposition (Grousset et al., 2003). In addition,

dust input into the atmosphere can increase significantly during droughts (e.g., Miao et al., 2007; Notaro et al., 2015; Sharifi et al., 2015). As such, fluctuations in dust loading may be indicative of both regional drying and long-distance transport (Le Roux et al., 2012).

Hydroclimatic fluctuations had a significant effect on the development of civilisations throughout the Holocene (Brooks, 2006; deMenocal, 2001; Sharifi et al., 2015), especially on those which relied heavily on agriculture and pastoralism, as was the case in the Carpathian-Balkan region (Schumacher et al., 2016). To understand the impact hydroclimatic changes had on the population of an area of such importance to European history, high-resolution palaeoclimate and palaeohydrological records are needed. This is especially important in the Carpathian region, given the extensive loess cover in the area (Marković et al., 2015) - a fundamental factor in sustaining high agricultural production. Additionally, the sensitivity of loess to moisture availability and water stress during dry periods may turn this region and other surrounding loess belts into major dust sources (Kok et al., 2014; Rousseau et al., 2014; Sweeney and Mason, 2013). This is particularly true under semi-arid (Edri et al., 2016), or agriculturally-altered conditions (Korcz et al., 2009), as is the case with the major dust fields of Eastern Eurasia (Buggle et al., 2009; Smalley et al., 2011; Újvári et al., 2012). Thus, the dust influx into the Carpathian-Balkan region should be extremely sensitive to relatively small changes in precipitation rates. This hydroclimatic sensitivity is enhanced due to the fact that the Carpathians and the surrounding lowlands are located at a confluence of three major atmospheric systems; the North Atlantic, the Mediterranean and the Siberian High (Obreht et al., 2016). Indeed, research appears to indicate the climate in Romania is controlled, at least in part, by North Atlantic Oscillation (NAO) fluctuations (Bojariu and Giorgi, 2005; Bojariu and Paliu, 2001), but it is yet unclear how this relationship evolved in the past-(Haliuc et al., 2017).

Multi-proxy and high resolution studies of palaeoenvironmental changes in the region are still scarce, with most focusing on reconstructing past vegetation changes (e.g., Feurdean et al., 2012). More recently, testate amoeba (Schnitchen et al., 2006; Feurdean et al., 2015), pollen and diatoms (Magyari et al., 2009, 2013, Buczko et al., 2013), and macrofossils (Gałka et al., 2016) have been utilised to elucidate the history of hydroclimatic variability in the region. What is evident from these studies is the high inter-site variability, with clear disagreements on timing and extent of wet and dry periods within a relatively small spatial distribution (e.g., two spatially close sites displaying differing precipitation trends as reported in Feurdean et al., 2008). It is possible that this variability reflects only site-related (including chronological) uncertainties, or is an indicator of the impact of location at the contact of several climatic zones (Obreht et al., 2016). To determine this, the impact of different modes of atmospheric (and moisture) circulation patterns and their imprint within paleoclimate archives must be investigated through better regional coverage following high-resolution multi-proxy approaches (e.g., Longman et al., in review.); Haliuc et al., 2017).

Our research provides a record of periodic dry and/or dusty periods in Eastern Europe as indicated by reconstructed dust input, using an ombrotrophic bog from the Romanian Carpathians (Fig.1). As the only source of clastic material deposited within ombrotrophic bogs is via atmospheric loading, such records have been used convincingly as archives of dust deposition over the Holocene in Western Europe and Australia (Allan et al., 2013; Kylander et al., 2013a; Marx et al., 2009, 2010; Le Roux et al., 2012). To produce records of dust and/or hydroclimate variability, both inorganic (Allan et al., 2013; Ross-Barraclough and Shotyk, 2003; Shotyk, 2002)

and organic (Booth et al., 2005; Lamentowicz et al., 2008; Morris et al., 2015; Swindles et al., 2010) proxies may be utilised (see Chambers et al. 2012 for a review).

Here we present the first record of dust input over the Carpathian Mountains, documenting changes in dust flux, source, and intensity of deposition using the downcore lithogenic element concentrations from the Mohos ombrotrophic bog profile. The record covers 10,800 years of deposition over 9.5 m of peat, providing a valueablevaluable high-resolution record for this region. Our research utilises both organic and inorganic proxies, with a high-resolution geochemical record of lithogenic elements (Ti, Si, and K), presented alongside the bog surface wetness as reconstructed using testate amoeba to understand dust source changes and the link between regional and extra regional hydroclimate variability and dust.

2 Materials and Methods

2.1 Geographical Setting

The Mohos peat bog (25°55' E; 46°05' N; 1050 m altitude, Fig.1) is located in the Eastern Carpathians, Romania, in the Ciomadul volcanic massif (Fig. 1). The *Sphagnum*-dominated bog covers some 80 hectares, and occupies an infilled volcanic crater. There is no riverine inflow, which means that inorganic material deposited within the bog is almost exclusively derived via direct atmospheric transport. The climate is temperate continental, with average annual temperatures of 15°C and precipitation of 800 mm (Kristó, 1995). Surrounding vegetation is typical of this altitude in the Carpathians (Cristea, 1993), the bog being located at the upper limit of the beech forest, with spruce also found on surrounding slopes. Vegetation on the bog itself is diverse, with common occurrences of *Pinus sylvestris, Alnus glutinosa,* and *Betula pubescens,* alongside various *Salix* species (Pop, 1960; Tanțău et al., 2003).

The Mohos crater is related to volcanic activity from the Ciomadul volcano, which last erupted roughly 29.6 cal kyr BP in the neighbouring younger crater currently occupied by the Lake St Ana (Harangi et al., 2010; Karátson et al., 2016; Magyari et al., 2014; Wulf et al., 2016). The surrounding geology is dominated by andesites and dacites, occasionally capped by pyroclastic deposits and a thick soil cover.

2.2 Coring

A Russian peat corer was used to recover a 950-cm long peat sequence from the middle part of Mohos bog. The material consists mainly of *Sphagnum* peat and lacustrine sediments in the lowermost part. <u>Upon recovery, the material was wrapped in clingfilm, transported to the laboratory, described, imaged, and subjected to further analyses. At both institutions, the The core was stored at 3°C.</u>

2.3 Sedimentological Parameters

Loss on ignition (LOI) was performed on ~1g (exactly 1cm^3) of wet peat, sampled at 2-em2cm resolution. The peat was dried overnight at 105°C prior to ignition at 550°C for four hours. Weight loss after this combustion was used to calculate combusted organic material, followed by further combustion at 950°C for two hours to calculate total carbon content following carbonate removal (Heiri et al., 2001). The dry bulk density was determined from the known volume and the dry weight prior combustion.

2.4 Micro-XRF and MSCL Core Scanning

Non-destructive X-Ray fluorescence (XRF) analysis was performed using an ITRAX core scanner equipped with a Si-drift chamber detector (Croudace et al., 2006) at the University of Cologne (Institute of Geology and

Mineralogy). The analytical resolution employed a 2-mm step size and 20 s counting time using a Cr X-ray tube set to 30 kV and 30 mA. The method allows for a wide range of elements to be analysed, from which we have selected Ti, K, and Si for further interpretation. To allow <u>afor</u> better visibility, all XRF data sets were smoothed using a 9-point running average. Due to the methodological nature of XRF core scanning, the data are presented as <u>raw</u>-counts per second (cps) and are therefore considered as semi-quantitative. To ensure the impact of sedimentological variables, including density, high organic matter₇ and water content, is taken into account, the raw cps values have been normalised with respect to total (incoherent + coherent) scattering (Kylander et al., 2011, 2013b)_{7.}

2.5 ICP-OES

To perform quantitative analysis of elements to allow inference of past dust flux, as well as to validate the ITRAX data, ICP-OES analysis was carried out on 105 samples of 1cm3 of sediment-collected over 1 cm depth interval, and at, roughly 100 year resolutionevery 10 cm, through the entire core. These samples were dried at 105°C overnight before homogenising using a pestle and mortar and then subjected to a mixed acid (HNO₃: HCI: HF) total digestion (adapted from Krachler et al., 2002), for 40 minutes in a MARS accelerated reaction system. The solution was then analysed using a Perkin Elmer Optima 8000 ICP-OES system at Northumbria University. To monitor a potential instrumental drift, internal standard (1ppm Sc) was added to all samples, and analysed alongside Ti. In addition, two Certified Reference Materials (CRMs) were digested and analysed throughout the runs (Montana soil 2711 and NIMT/UOE/FM/001). Recoveries for both CRMs were good for Ti, with average values of 85% and 79% respectively. Blanks with negligible Ti contamination were run alongside the samples and CRMs.

2.6 Calculating Dust Flux

The dust flux delivered to an ombrotrophic bog via atmospheric loading may be calculated using the concentration of a lithogenic element, such as Ti (Allan et al., 2013). Using the averaged occurrence of Ti in the upper continental crust (UCC values from Wedepohl, 1995), the density of the peat as well as the peat accumulation rate (PAR), the following formula may be used:

$$Dust \ Flux \ (g \ m^{-2} yr^{-1}) = \left(\frac{[\text{Ti}]_{sample}}{[Ti]_{UCC}}\right) \times density \times PAR \times 10000 (Eq. 1)$$

2.7 Palaeoecological indicators

A total of 44 samples of roughly 1cm³ each were sampled along the peat profile for testate amoeba analysis. The bulk samples were disaggregated and sieved according to Booth et al. (2010), prior to mounting in water on slides. Two tablets of Lycopodium spores of known value were added toprior to disaggregation to allow for calculation of test density. For each sample at least 150 tests were counted, with identification of taxa following Charman et al. (2000). For interpretation, two methods of determining wet and dry local depositional environments based on changes in testate amoeba assemblages were used. Firstly, a transfer function (Schnitchen et al., 2006) already applied to Carpathian bogs was used to reconstruct past variations in the depth of the water table (DWT). Secondly, the main taxa were grouped into their affinity to wet or dry conditions according to Charman et al. (2000) and plotted as a function of percentage.

2.8 Chronology

The age model for the Mohos peat record is based on 16 radiocarbon dates on bulk peat (collected over less than 1 cm depth interval per sample) consisting only of *Sphagnum* moss remains (Table 1). These analyses were performed via EnvironMICADAS accelerator mass spectrometry (AMS) at the Hertelendi Laboratory of Environmental Studies (HEKAL), Debrecen, Hungary₂ using the methodology outlined in Molnár et al. (2013). The ¹⁴C ages were converted into calendar years using the IntCal13 calibration curve (Reimer et al., 2013) and an age-depth model (see Fig. 2) was generated using Bacon (Blaauw and Christen, 2011).

2.9 Wavelet Analysis

Continuous Morlet wavelet transform was used to identify non-stationary cyclicities in the data (Grinsted et al., 2004; Torrence and Compo, 1998). For this analysis, the lithogenic normalised elemental data from ITRAX measurements (Ti, K, and Si) was interpolated to equal time steps of four years using a Gaussian window of 12 years.

2.10 Grain Size

To indicate particle size distribution throughout the record, grain size analysis was attempted using a Malvern Mastersizer 2000. However, as also noted in other studies (Kylander et al., 2016) due to the very low minerogenic matter (and correspondingly high organic matter) present in all samples, satisfactory obscuration values were not achieved for the vast majority of analyses.

In an effort to indicate distal versus local inputs to the bog via the dust particle size, grain size analysis was attempted using a Malvern Mastersizer 2000. Unfortunately, as also observed in previous studies (Kylander et al., 2016) due to the lack of available sample material, and low minerogenic matter (and correspondingly high organic matter) present in the samples, satisfactory obscuration values were not achieved for most analyses.

3 Results

3.1 Age Model and Lithology

The Mohos peat profile is 950 cm long, and reaches the transition to the underlying basal limnic clay (Tanţău et al., 2003). Between 950-890 cm the record is composed of organic detritus (gytţia) and *Carex* peat deposited prior to the transition from a wetland into a bog, at roughly 10,330 yr BP. From 890 cm upwards, the core is primarily *Sphagnum*-dominated peat. The age-depth model indicates the Mohos peat record covers almost 10,800 years of deposition, with the uppermost layer (growing moss) of the peat dating to 2014. Age model uncertainties range from 20 years in the uppermost sections to 150 years at the base of the core. Thus, the resolution for ITRAX data average ~5yr/sample and for ICP-OES is roughly 100 yr/sample, respectively. The testate amoeba resolution is roughly 200 yr/sample. In the following, all quoted ages are given in calibrated years before present (cal yr BP).

3.2 Dust indicators

3.2.1 Ti, K, and Si

Similar trends for the lithogenic elements Ti, and Si, and the mobile element K, are visible in the record (Fig. 3), with 10 main zones of higher counts above typical background values present. Such zones are identified as an increase in two or more of the elements above the background deposition (K > 0.001, Si > 0.001, and Ti > 0.004, see dashed line on figure-Fig.3). These intervals are further discussed as reflecting major dust deposition events, and are referenced in the remainder of the text using the denotation D01-D10 (Fig. 3). Two exceptions are not

highlighted. Firstly, at the base of the core, close to the transition from lake sediments to bog. Secondly, a very brief enrichment between D9, and D10, related the last 1000 years, due to high variation in the scattering values at this time.noise, are not highlighted. The lithogenic, and therefore soil and rock derived Ti and Si have previously been used as proxies for dust input (e.g., Allan et al., 2013; Sharifi et al., 2015), whilst K covaries with Si $(R^2=0.9945)$ and so controlling factors on their deposition must be <u>nearly the same similar</u>. For these elements, the periods with inferred non-dust deposition are characterised by values approaching the detection limit (150, 15_{τ} and 40 cps, respectively). A short period of very high values for all elements (10,000, 1300 and 8000 cps, respectively) is observed between 10,800-10,500 cal yr BP (not shown on diagram), reflecting the deposition of clastic sediments within the transition from lake to bog at the onset of the Holocene. Zones of elevated values (D1-D5), with average cps values roughly Ti=300, $Si=30_7$ and K=100 and persisting for several centuries each, occur sporadically throughout the next 6000 years of the record, between 9500-9200, 8400-8100, 7720-7250, 6350-59007 and 5450-5050 cal yr BP (Fig. 3). Similarly long periods, but with much higher element counts (Ti=800, Si=60 and K=200 cps) occur between 4130-3770, 3450-2850, and 2000-1450 cal yr BP (D6-8). Two final, short (roughly 100-year duration) but relatively large peaks (D9-10) may be seen in the last 1000 years, between 800-620 cal yr BP (with values Ti=300, Si=40 and K=100 cps) and 60 cal yr BP to present (Ti=300, Si=80 and K=400 cps, respectively).

3.2.2 Dust Flux

Using the quantitative ICP-OES values of Ti (in ppm), and equation 1, the dust flux can be calculated (Fig. 3). The ICP-OES Ti record shows very good correlation with the Ti data derived through ITRAX analysis. After To facilitate comparison, we bring both records on the same timescale using a Gaussian interpolation atwith 100 year time_steps using and a 300 year Gaussian window, R^2 . Pearson's r=0.2649, and with a p-value of 1.416×10^{-8} , indicating<0.001, indicative of a significant correlation. (See SI 4). This further implies indicates the reliability of the XRF core scanning method even for such highly organic sediments (as already suggested by Poto et al., 2014), and validates its usage as toolproxy for deriving dust flux (Fig. 3).

It must be noted here that using Ti alone in dust flux calculations may not allow for reconstruction of all forms of dust deposition. Ti, which is lithogenic and conservative, is a major component in soil dust, particularly within elay minerals (Shotyk et al., 2002). On the other hand Ti may not be associated with other dust forming minerals, including phosphates, plagioclase, and silicates (Kylander et al., 2016), and thus, may help indicate changes in deposition rates of particles bearing these elements. As a result of these limitations, at this stage of research we refrain from inferring changes in mineral composition of dust within the core, and the results should be considered only indicative of general dust deposition. Such an approach has been utilised in a number of previous studies (e.g., Allan et al., 2013; Sapkota et al., 2007; Sharifi et al., 2015).does not allow for reconstruction of all minerals related to dust deposition. Ti, which is lithogenic and conservative, is a major component in soil dust, particularly within clay minerals (Shotyk et al., 2002), but may not be associated with other dust-forming minerals, including phosphates, plagioclase and silicates (Kylander et al., 2016), although our records of K and Si may help indicate changes in the composition of dust. However, Ti alone will record changes in the intensity of deposition of the main dust-forming minerals (Shotyk et al., 2002), and variations in K and Si (particularly with local K- and Si-rich dacites a possible dust source) may further indicate the influx

of minerals which are not associated with Ti. Such an approach has been applied successfully to studies of changing dust influx (e.g. Allan et al., 2013; Sapkota et al., 2007; Sharifi et al., 2015), with each study able to identify periods of high and low dust deposition from Ti-derived dust flux alone.

The Ti-derived dust flux for most the record is below 1 g m⁻²yr⁻¹, but with seven periods of dust deposition clearly identifiable for the last 6100 years, and several smaller fluctuations prior to that (mainly visible in the elemental data). The main peaks are similar in their timing to the ITRAX Ti trend, with three large peaks (dust flux >1.5 g m⁻²yr⁻¹) located between 5400-5050, 2100-1450, and 800-620 cal yr BP, respectively (Fig. 3). Smaller peaks are present (dust flux 0-5-1 g m⁻²yr⁻¹) at 6100-6000, 4150-3770, and 3500-2850 cal yr BP, respectively.

3.3 Density and Loss-on-Ignition (LOI)

Density values are relatively stable throughout the core, with all samples ranging between 0.06-0.1 g-cm⁻³/cm³. This trend is different from the organic matter values, which typically oscillate around 90-100% over the entire record. The very base of the record is however an exception, denoting the gradual transition from limnic clays to the peat reaching, with organic matter values of 80-90% between 10,800-10,000 cal yr BP. Very occasional intervals with lower organic matter content (roughly 85%) may be observed at 5400, 4100-3900, 3300-3200, 1900-1800₇ and 900-800 cal yr BP, respectively (Fig. 4).

3.5 Testate Amoeba

Two methods of clarifying the paleoclimate signal derived through investigating testate amoeba assemblages have been used (Charman et al., 2000; Schnitchen et al., 2006), with both indicating similar hydroclimatic trends. Reconstructions of depth-to-water table (DWT) values indicate three main trends within the record. The first, encompasses the time period between 10,800-7000 cal yr BP, and is characterised by highly fluctuating values, with four very dry periods (DWT ~20 cm) at 10,800-10,200, 9000-8800, 8600-7600, and 7400-6600 cal yr BP interspersed by wetter (DWT 15 cm) conditions (Fig. 4). After 7000 cal yr BP, values are much more stable, with DWT of <u>15-em15cm</u> until the final zone, the last 100 years, where DWT rises to <u>20 em20cm</u>. These fluctuations are in line with those seen in the wet/dry indicator species.

3.6 Wavelet Analysis

The wavelet analysis of K, Si₇ and Ti show significant (above the 95% confidence threshold) periodicities between 1000-2000 years within the past 6000 years (Fig. 8). Prior to this, there appears to be no major cyclicity in the ITRAX data. Within periods, which display raised ITRAX counts, shorter frequency (50-200 year) cycles are seen. These persist only for the period in which each element is enriched, with such cycles particularly evident within the last 6000 years.

4 Discussion

4.1 Peat ombrotrophy

The relative intensities of the lithogenic elements analysed via ITRAX covary throughout the record (Fig. 3), despite their varying post-depositional mobility (Francus et al., 2009; Kylander et al., 2011). For example, the largely immobile Ti shows a very high correlation with that of redox sensitive Fe ($R^2 = 0.962$) and mobile K ($R^2 = 0.970$). This indicates the downcore distribution of these elements is mostly unaffected by post-depositional mobilisation via groundwater leaching and/or organic activity as documented in other studies (e.g., Novak et al., 2011; Rothwell et al., 2010), indicating the conservative behaviour of such elements in the studied peat. This,

alongside the low clastic content (average organic matter of 91%), low density, and domination of *Sphagnum* organic detritus, indicates the ombrotrophic nature of Mohos bog throughout time and validate the use of this record to reconstruct dust fluxes for the last ca. 10,000 years (Fig. 3).

4.2 The Dust Record

The record of inferred lithogenic (dust) input as indicated by Ti, K_{7} and/or Si documents 10 well-constrained periods of major and abrupt dust deposition (denoted D0-D10), with further small, short-term fluctuations (Fig. 3). The dust influx onto the Mohos peat was accompanied by decreases in organic matter (OM) as indicated from the LOI profile, and higher density values (Fig. 4), particularly over the intervals covered by events D5-D10. The major dust deposition events lasted from a few decades to centuries (Fig. 3).

Firstly, it is noteworthy that five of the identified dust depositional events may be compared to periods of Rapid Climate Change (RCC) as outlined by Mayewski et al. (2004) from the Greenland GISP2 record (Fig. 5). However, despite apparent hemispheric-scale influences, the dust events identified within Mohos record have little correlation to reconstructed European paleoclimate changes during Holocene. For example D8, between 3450 2850 cal yr BP falls within a cold period (Wanner et al., 2011). Such cold related dust deposition has been observed previously in western Europe (Allan et al., 2013). However, within Mohos such a comparison may not be drawn with the majority of dust events. D9, for instance, between 800 620 cal yr BP is within the Medieval Climate Anomaly, a period of generally higher European temperatures (Mann et al., 2009) but also one of intense human impact on the environment thorough deforestation and agriculture (Arnaud et al., 2016; Kaplan et al., 2009). Furthermore, such events within the Misten record (Allan et al., 2013) were also linked to low humidity, whereas the Mohos TA (See Fig. 4) record indicates locally wet conditions. This suggests the dust depositional events in this region are a result of a complex interplay of environmental conditions in the dust source areas, rather than simply reflectingFor example D8, between 3450-2800 cal yr BP falls Europe-wide cold period (Wanner et al., 2011). Such cold-related dust deposition has been observed previously in western Europe. However, within Mohos such a compassion may not be drawn for the majority of dust events. For example, event D9 (860-650 cal yr BP) occurs during the Medieval Climate Anomaly, a period of generally higher European temperatures (Mann et al., 2009) but also one of intense human impact on the environment through deforestation and agriculture (Arnaud et al., 2016; Kaplan et al., 2009). Furthermore, such events within the Misten record (Allan et al., 2013) were also linked to low humidity, whereas the Mohos TA (Fig. 4) record indicates locally wet conditions. This suggests that dust depositional events in this region are a result of a complex interplay of environmental conditions in the dust source areas, rather than simply reflecting locally warm or cold, or even wet or dry periods.

In addition to the North Atlantic, the impact of both the Mediterranean and the intertropical convergence zone (ITCZ) atmospheric systems influencing the Mohos dust record are apparent, including major climate changes in North Africa. D4 for example occurs within the chronological span of the 5900 cal yr BP event, a major cooling and drying period- (Bond et al., 2001; Cremaschi and Zerboni, 2009; Shanahan et al., 2015). Increased dust influx is also recorded around 5300 cal yr BP (D5, Fig. 3) which roughly correlates with the end of the African Humid Period and onset of Saharan desertification (DemenocaldeMenocal et al., 2000). The lack of dust flux perturbations prior to 6100 yr BP, and their prevalence thereafter at Mohos are consistent with a major shift in the controls of dust production and deposition at this time, a change observed in peat-derived dust records from Western Europe (Allan et al., 2013; Le Roux et al., 2012). The desertification of the Sahara around this time was

the largest variation in dust production in the northern hemisphere (see McGee et al., 2013; deMenocal et al., 2000).

Within our record, this initial dust flux increase was followed by a period of reduced dust loading, prior to a rapid, and apparently major (highest dust flux values in the record prior to the most recent two millennia) event at 5400-5000 cal yr BP. Regionally, Saharan dust in Atlantic marine cores strongly increased at this time, with a 140% rise roughly at 5500 cal yr BP observed on the westernWestern Saharan margin (Adkins et al., 2006) with another study indicating a rise by a factor of 5 by 4900 cal yr BP in a selection of similarly-located sites (McGee et al., 2013). Furthermore, evidence from marine cores across the Mediterranean indicate decreasing Nile output and increasing dust fluxes into the Eastern Mediterranean at this time (Box et al., 2011; Revel et al., 2010). The correlation of these data to the Mohos record appears indicative of the region-wide impact of North African desertification. It is noteworthy, as seen in Fig. 5, that the release of dust from the Sahara correlates well with increasing frequency and intensity of dust fluxes at Mohos after 6000 cal yr BP, with all major (dust flux >0.5 g m⁻²yr⁻¹) Ti-derived dust flux peaks occurring after this time (Fig. 3). This period is the first indication of the impact the Mediterranean climate and movement of the ITCZ has had on the Carpathian-Balkan region (as simulated by Egerer et al., (2016) and Boos and Korty, (2016). Indeed, intermittent intrusions of Saharan dust over the Carpathian area have been well documented both through direct observations (Labzovskii et al., 2014; Varga et al., 2013), and through provenance studies of past Saharan dust contribution within interglacial soils in the region (Varga et al., 2016).

In addition to Saharan desertification, it is likely that early agriculture in the Carpathian-Balkan region has contributed towards the increase in dust flux values at this time. It is known that advanced agriculture-based societies inhabited the Carpathian area in the mid-Holocene (Carozza et al., 2012), with evidence of farming seen in a number of pollen records (see Schumacher et al., 2016 for a compilation), including in Mohos itself at the end of the Chalcolithic period (Tanțău et al., 2003). Since agriculture and soil erosion may be linked, it is possible events D4 and D5 could also reflect to some extent dust input related to land disturbance by human activities, on a regional scale. However, such evidence for agriculture, particularly in the proximity of Mohos is limited to a few *Plantago* and cereal pollen (Tanțău et al., 2003), whilst the majority of pollen studies in Romania at this time indicate no significant agricultural indicators (e.g., Magyari et al., 2010; Schumacher et al., 2016; Tanțău et al., 2014). As such, it seems unlikely agricultural activity is behind such a large change in the dust deposition record from Mohos.

4.3 Geochemical evidence for a dust provenance shift at 6100-6000 cal yr BP?

To better understand the nature of the shift in dust flux after 6100-6000 cal yr BP, a simple approach to disentangling the geochemical makeup of the reconstructed dust load is discussed below. Figure 6 displays the clustering of the lithogenic elements Ti and K (and Si, due to the similarity in the Si and K records) during dust events D1-D10. The data appear to show three main types of dust (and presumably sources), one with high values for both Ti and K (Type 1), one with relatively high values for K (Type 2), and one with relatively high Ti compared to K (Type 3). The values for Ti-K correlation, average Ti, and average K (in <u>normalised cps</u>) are listed in Table 2. Generally, the periods of no enrichment, and low K and Ti, do not show any correlation, indicative of natural background and instrumental detection limits.

Type 1 deposition occurs only in D10, and is characterised by Ti-K gradient of nearly 1, indicating similar values for both elements throughout the period, and a dust rich in both K and Ti. Type 2 deposition occurs in several the dust events, particularly in D1-2, D4-5, and D7 (Fig. 8). The K enrichment, which characterises these events is, evidenced by the Ti-K gradients <1 and low (even negative in the case of D2) correlations between the two elements. -Finally, Type 3 events (D3, D6, and D8-9) are characterised by an increased Ti-K gradient, generally, around 0.2. The average Ti values during these events and the Ti-derived dust flux, are generally highest in these periods (Table 2). These groupings would indicate similar dust sources within grouped events, and may aid in identifying provenance.

Type 2 events typically occur in the older part of the record, except D7 (3400-3000 cal yr BP_{ia} Fig. 8). Such events are not visible in the Ti-derived dust flux values, indicative of the reduced impact of Ti-bearing dust particles deposited within the corresponding periods. The local rocks consist of K-rich dacites and pyroclastics (Szakács et al., 2015), with relatively low Ti concentrations and enriched in K (Vinkler et al., 2007). Therefore, the likely source of particlesparticulates deposited during these dust events is local or regional, with nearby (or even distal) loess and loess-like deposits as another potential source, since loess sediments in south-eastern Europe are generally depleted in Ti (Buggle et al., 2008). The local nature of such deposition is emphasised by the similarity of type 2 deposits the depositional signal to background values; the elemental composition outside of dust events. For all data points not considered to be related to dust (or the minerotrophic lowermost section), the Ti-K correlationregression is low ($R^2r2= 0.1513$) with a gradient of 0.0863.

Type 3 events, conversely, appear Ti-enriched (Fig._6), with contribution from a source away from the low-Ti dust of south-<u>easteastern</u> European loess fields. These events typically occur after 6100 cal yr BP (Fig. 3). With the periodic influence of the Mediterranean air masses in the region (Apostol, 2008; Bojariu and Paliu, 2001), Saharan dust must be considered as a potential source area, since it appears to play a major role in dust input into Europe today (e.g., Athanasopoulou et al., 2016). Geochemically, Saharan dust is typically Ti-enriched (Nicolás et al., 2008). In particular, the Bodélé depression, the single-largest dust source in the Sahara, exhibits extremely high Ti/Al and Ti enrichment (Bristow et al., 2010; Moreno et al., 2006). Since-the Ti enrichment does not show any regional trends, it is no use for determining exact source areas within the Sahara (Scheuvens et al., 2013), but the presence of Ti-enriched dust appears to reflect a signal of Saharan influence. Consequently, events of Type 3 may be considered to reflect, at least to a large extent, contribution of Saharan dust. Finally, the single Type 1 event may be attributable to a mixing of both local (resulting in high K) and distal sources (resulting in high Ti); sources, evidence for Saharan input and local soil erosion/deflation.

Previous work has indicated the input of Saharan dust in Eastern Europe, with evidence of such a source seen in Carpathian loess (Újvári et al., 2012; Varga et al., 2013) and soil-forming dust (Varga et al., 2016). Additionally, recent atmospheric satellite imagery has further confirmed the extent of Saharan dust outbreaks and depositional events over central-eastern Europe (Varga et al., 2013). However, the lack of long-term dust reconstructions in the region has so far precluded understanding of changing dust sources over the Holocene.

Previous studies across Europe indicate the complex input of dust from various sources over the mid-to-late Holocene (e.g., Veron et al., 2014), but pertinently to our findings at Mohos, many examples exhibit a major shift in dust sources at roughly 5000-7000 cal yr BP. In Belgium, Nd isotopes indicate a local source of dust from the

input of European loess prior to₁ and Saharan dust after 6500 cal yr BP (Allan et al., 2013). This is echoed by data from Le Roux et al. (2012) that implies a major shift in the Nd isotopic composition at 6000 cal yr BP, moving from a local to a mixed source, but with clear Saharan overprinting. The transition identified within the Mohos Ti-derived dust record at 6100-6000 cal yr BP, therefore, appears to echo the appearance of a Saharan dust element within other European bog-based dust reconstructions. However, it appears that input of Saharan dust was not limited to the onset of North Africa desertification, as indicated by input of likely Saharan derived dust within Mohos event D3 already by 7800-7200 cal yr BP. Further, even after 6100 cal yr BP, local sources still played a significant role, with D7 showing clear local or regional (e.g., loess-derived) signal.

D10 is interesting in that it appears to indicate even more K-rich dust sources. The D10 values are similar in compositional gradient to the lake sediments deposited prior to the onset of peat formation in the early Holocene (Gradient of samples pre-10,500 yr BP = 0.7429, D10= 1.0637). Since the surrounding dacites and pyroclastics are K-rich (Vinkler et al., 2007), and the sediment composition prior to peat formation reflects the natural signal of erosion into the lake, it is reasonable to assume this period is indicative of local slope erosion. This is potentially due to the decline of the local forest and agricultural intensification, identified in the most recent sections of the Mohos pollen record (Tanțău et al., 2003). It is sensible to assume the local deforestation (visible around the Mohos bog as meadows for hay harvesting) has caused local soil erosion and increased dust production from very proximal sources (Mulitza et al., 2010). This is a clear sign of the persistent human impact at local to regional scale during the early Holocene (Giosan et al., 2012; Schumacher et al., 2016) that is also mirrored in the nearby Lake St Ana record (Magyari et al., 2009). As indicated by regional studies (e.g., Labzovskii et al., 2014; Varga et al., 2013; Vukmirović et al., 2004)₇ high levels of Ti indicate that the Saharan input does not cease through this period, but that it is rather-matched by high-K local sources. The apparent higher water table of the Mohos bog as implied by the TA record and the increased Ti contents rather point towards an increasing Saharan influence rather than a major local dust source.

4.4 Correlation to other European dust records

Comparison to similar dust records from peat cores in Western Europe (Allan et al., 2013; Le Roux et al., 2012), and Atlantic margin sediments (McGee et al., 2013) reveals some interesting trends visible in all these records (Fig. 5), indicating comparable continent-wide controls on past dust flux. Specifically, the major dust event as seen at 5400-5000 cal yr BP in Mohos, and subsequent increase in number and intensity of dust events is comparable with an intensification of dust deposition over Europe after 6000 cal yr BP (Le Roux et al., 2012), with concurrent increases in dust flux in the Mid-Holocene documented in Belgium (Allan et al., 2013). The authors suggest a cool period as the cause of this dust increase (Wanner et al., 2011). In addition to the reconstructed cool environments in Western Europe, this period is characterised by increased dust production in the Sahara (McGee et al., 2013), which is also likely to have played a role in the increasing dust flux over Europe. After 5000 cal yr BP, it appears Mohos and central-western European records show a more concurrent trend, with comparable dust peaks in the Swiss record (Le Roux et al., 2012) between 4100-3800, 3600-3050, 850-600, and 75 cal yr BP also present in Mohos, and a similar dust peak at 3200-2800 cal yr BP identified in another bog record from Bohemia (Veron et al., 2014).

Despite some similarities between the records, there is also significant variability, highlighting the difference between climatic controls in western and central Europe and those in south-eastern Europe. The disconnection

between Mohos and other records is particularly clear for the early Holocene, with a large dust flux peak identified in Switzerland between 9000-8400 cal yr BP, and other volcanic eruption-related dust (seeSee Fig. 5), when there is little evidence of dust input into Mohos. This discrepancy could be indicative of the east-west (Davis et al., 2003; Mauri et al., 2015; Roberts et al., 2012) and north-south (Magny et al., 2013) hydroclimatic gradients in Europe throughout the Holocene. As other studies indicate, south-eastern Europe was mostly disconnected (in terms of both precipitation and temperature) from the rest of Europe in the early-Mid Holocene (Davis et al., 2003; Drăguşin et al., 2014), clearly indicated by the trend in the Mohos Ti-derived dust record. Since the Sahara had not undergone significant desertification by this time, no clear correlation with western records may be made, hinting at more local source for the early to mid-Holocene, which are not present in the Ti-derived dust record at Mohos, are more likely related to local fluctuations in moisture availability, and Si and K rich soil dust.

4.5 Palaeoecological Proxy Record

To further investigate the difference between local and regional palaeoclimate signals within Mohos, and to reconstruct the local hydroclimate conditions throughout the record, we use the fossil assemblages of testate amoeba (TA). These data, alongside comparisons to existing Carpathian-Balkan and Mediterranean hydroclimate reconstructions (Fig. 7), may be used to further investigate the theory of a distal (most likely Saharan) source for dust after 6100 cal yr BP. The earliest section in the TA record (10,800-6400 cal yr BP) is characterised by fluctuating dry/wet periods, indicative of large shifts in the local hydroclimatic environment (Fig. 4). The earliest identified dry period (10,800-10,000 cal yr BP) is linked to the shift away from a lacustrine to a palustrine environment as a result of local drying. Three subsequent dry periods may be identified in the TA record (9300-8800, 8500-8100, and 7800-7000 cal yr BP, all of which are also identifiable in the geochemical dust record (D1-D3) via peaks in K and Si. Between 10,200-7450 cal yr BP, dust flux at Mohos was low. Dust events during this time are mainly present in the K and Si records (Fig. 3), or in OM and density parameters (Fig. 4).

The first period of elevated dust proxies at roughly 10,300 cal yr BP (D0) correlates well with the 10,200 cal yr BP oscillation (Rasmussen et al., 2007), previously linked to a drop in water levels at nearby Sf. Ana Lake (Korponai et al., 2011; Magyari et al., 2012, 2014). High Difflugia pulex and Trigonopyxis arcula values during D1 as indicator taxa for dry conditions (Allan et al., 2013; Charman et al., 2000) appear to confirm local drying, observed across much of the Mediterranean (Berger et al., 2016; Buczkó et al., 2013; Magyari et al., 2013; Fig. 7). The D2 and D3 events may also be observed in both the TA record and the geochemical dust record, with D2 attributable to the 8200 cal yr BP event (Bond et al., 2001), a paleoclimatic event already identified in other local hydroclimate reconstructions (Buczkó et al., 2013; Magyari et al., 2013; Schnitchen et al., 2006). The transition to the next wet period at 8000 cal yr BP also mirrors the dust record, with a deeper water table occurring during the dust-free conditions between D2 and D3. This is prior to the bog undergoing dry conditions between 7800-7000 cal yr BP, roughly in line with D3, drying which has previously been observed in Romania (Gałka et al., 2016; Magyari et al., 2009; Fig 7). Due to the covariance between geochemical and palaeoecological proxies at this time, and the correlation to other local reconstructions, the early Holocene section of the record indicates a close linkage of local hydroclimate and dust input. These dust events, therefore, are likely the signal of remobilised material (Edri et al., 2016), from proximal or distal sources (including perhaps from loess-derived sediments, at the foot of Ciomadul volcano) as the climate locally appears to become more arid.

Between 6600-1200 cal yr BP, the TA indicate a shift to prolonged wet conditions, with only minor fluctuations and no clear correlation to the geochemically-derived dust record, and so the dust events appear unrelated to local drying within this time period (Fig. 7). Such wetter conditions also limit local <u>drought-related</u> erosion, and so may be further evidence of distal dust input at this time (Allan et al., 2013). Furthermore, this is indicative of a decoupling of the dust record from local climate reconstructions, with dry phases common throughout the Mid-Late Holocene in other Romanian sites (e.g., Magyari et al., 2009; Schnitchen et al., 2006; Fig. 7), and a distal dust source.

In the last millennium, there are two major dust events, with the first, D9, occurring between 850-650 cal yr BP. This episode falls within the late Medieval Warm Period, and could be related to human activity in the local area, as pollen from the Mohos bog indicates strong evidence for agriculture at roughly the same time (Tanţău et al., 2003). This may be seen in the intensity of the dust deposition at this time (dust flux >3 g m⁻²yr⁻¹). D10, from 75 cal yr BP to present is certainly linked to such human influences, with the TA record echoing local studies, which display anthropogenically-altered conditions and intensive agriculture (Buczkó et al., 2013; Diaconu et al., 2016; Giosan et al., 2012; Magyari et al., 2009, 2013; Morellón et al., 2016; Schnitchen et al., 2006; Fig. 7). This appears to validate the geochemical approach used earlier, as intensive farming is likely to result in local dust mobilisation, with K-rich dust present at this time, and local input potentially erasing some distal signals. This does not preclude Saharan input, however, as the dust is also Ti-rich.

4.6 Periodicity

To further understand the nature of the reconstructed dust events, cyclicity within the geochemical record was investigated using wavelet analysis (Fig. 8). The main elements of interest (Ti, Si₇ and K) have no apparent cyclicity in the first half of the record (10,800-6000 cal yr BP) when there is low spectral power at all periods. In contrast, the last 6000 years display clear centennial and <u>millennial</u>-scale cycles. A number of other studies have identified cyclicity shifts at this time (Fletcher et al., 2013; Jiménez-Espejo et al., 2014; Morley et al., 2014), related to North Atlantic variability, but so far mainly in western Mediterranean records. From 6000 cal yr BP onwards, the geochemical record at Mohos preserves two main cyclicities; one <u>millennial</u> cycle (at ~1200-2000 years) and the second at ~ 600-800 years (Fig. 8). A 715-775 year cycle has been determined as a harmonic of Bond event-related dry periods, present in other Northern hemisphere records (Springer et al., 2008) and in central Africa (Russell et al., 2003). The <u>millennial</u> scale cycle, in contrast, is within the envelope of a 1750 year cycle observed within the western Mediterranean, in pollen (Fletcher et al., 2013) and Saharan dust (Debret et al., 2007; Jiménez-Espejo et al., 2014), which is attributed to changes in North Atlantic circulation.

Within the dust deposition events (Fig. 3), there is an overprinting of high-frequency cyclicity in the Ti record, especially within the last 5000 years (Fig. 8). These are particularly clear at 4200, 3400, and 1800 cal yr BP, but lower-power cyclicities may be seen in most dust deposition events. These are generally 100-200 years in length, and only last the extent of the dust outbreak. Cycles with lower than 140-year periodicities are possibly reflecting mainly background noise (Turner et al., 2016), but those longer in duration may be indicative of climatically forced fluctuations within drought events affecting the dust source areas. This suggests the reconstructed dust deposition events based on the Mohos record were not characterised by constant deposition of dust, but by periodic dust pulses. These short cycles could reflect solar forcing, with comparable 200-year cycles observed in humification profiles from peats (Swindles et al., 2012), sediments in the Baltic Sea (Yu, 2003), Pacific Ocean

(Poore et al., 2004), and in North American peatland isotope records (Nichols and Huang, 2012). In many cases, such cycles have been linked to lower solar activity periods, low temperatures and increased precipitation oscillations, related to the De Vries/Suess 200 year cycle (Lüdecke et al., 2015). In the case of Mohos, these fluctuations may have manifested themselves as shifts in dust deposition, and could indicate the persistent effect solar dynamics has on all facets of climate system.

5 Conclusions

- The first record of Holocene drought and dust input in a bog from Eastern Europe documents ten periods of high dust<u>loading</u>: 9500-9100, 8400-8100, 7720-7250, 6150-5900, 5450-5050, 4130-3770, 3450-2850, 2100-1450, 800-620, and 60 cal yr BP to present.
- A major intensification in the number, and severity (as indicated by dust flux values) of dust events is
 observed after 6100 cal yr BP. The two intervals before and after this shift are indicative of an alteration
 in major dust controls. For the period prior to 6100 yr BP, dust input is reflective of more local controls,
 whilst the most recent 6100 yr BP of deposition may be linked to more distal forcings.
- The timing of the major shift at 6100 cal yr BP is possibly related to the end of the African Humid Period, and the establishment of the Sahara Desert, pointing to significantly greater Saharan input within the regional dust loading after this time. This is corroborated by changes in cyclicity attributable to Saharan dust outbreaks, and a shift toward Ti-rich dust (a signal of Saharan rock and sediment) deposited onto the Mohos peat. Our data is the first such indication of the impact Saharan dust has had across Eastern Europe, in line with enhanced deposition of dust across the Mediterranean region. A tentative dust provenance analysis based on a simple geochemical approach to disentangle the composition of the dust has been applied to confirm this, with three main types of deposition documented, indicating the interplay between local/regional (mainly loess-derived) and Saharan dust sources over the Holocene.
- The most recent dust event, between 75 cal yr BP and today is geochemically indicative mainly of local erosion. This may be linked to the increasing human impact through deforestation, agriculture and recently tourism, and associated soil erosion, indicating a shift in the controls on drought and dust in the region.

Author contribution. J. Longman, D.-Veres, V.-Ersek, and U.-Salzmann designed the research, interpreted the results and wrote the paper. D.-Veres, M.-Bormann, and F.-Schäbitz performed the fieldwork. J.-Longman performed the ICP-OES, testate amoeba and statistical analysis. V.-Ersek performed the wavelet analysis. M. Bormann, F. Schäbitz, and V.-Wennrich performed performed the ITRAX and MSCL-analysis. K.-Hubay performed the ¹⁴C dating. All authors approved the content of the paper.

Competing interests. The authors declare that they have no conflict of interest

Acknowledgements. We would like to thank Northumbria University for J.Longman's studentship. This is a contribution to the project PN–II-ID-PCE-2012-4-0530 "Millennial-scale geochemical records of anthropogenic impact and natural climate change in the Romanian Carpathians" and to the CRC806 "Our way to Europe" hosted at University of Cologne, Bonn & Aachen (subproject B2) granted by the DFG (German Research Foundation).

References

Adkins, J., DeMenocal, P. and Eshel, G.: The "African humid period" and the record of marine upwelling from excess 230Th in Ocean Drilling Program Hole 658C, Paleoceanography, 21(4), 1–14, doi:10.1029/2005PA001200, 2006.

Allan, M., Le Roux, G., Piotrowska, N., Beghin, J., Javaux, E., Court-Picon, M., Mattielli, N., Verheyden, S. and Fagel, N.: Mid- and late Holocene dust deposition in western Europe: The Misten peat bog (Hautes Fagnes – Belgium), Clim. Past, 9(5), 2285–2298, doi:10.5194/cp-9-2285-2013, 2013.

Apostol, L.: The Mediterranean cyclones: the role in ensuring water resources and their potential of climatic risk, in the east of Romania, Present Environ. Sustain. Dev., 2, 143–163, 2008.

Arnaud, F., Poulenard, J., Giguet-Covex, C., Wilhelm, B., Révillon, S., Jenny, J.-P., Revel, M., Enters, D., Bajard, M., Fouinat, L., Doyen, E., Simonneau, A., Pignol, C., Chapron, E., Vannière, B. and Sabatier, P.: Erosion under climate and human pressures: An alpine lake sediment perspective, Quat. Sci. Rev., 152, 1–18, doi:10.1016/j.quascirev.2016.09.018, 2016.

Athanasopoulou, E., Protonotariou, A., Papangelis, G., Tombrou, M., Mihalopoulos, N. and Gerasopoulos, E.: Long-range transport of Saharan dust and chemical transformations over the Eastern Mediterranean, Atmos. Environ., 140, 592–604, doi:10.1016/j.atmosenv.2016.06.041, 2016.

Berger, J. F., Lespez, L., Kuzucuoğlu, C., Glais, A., Hourani, F., Barra, A. and Guilaine, J.: Interactions between climate change and human activities during the Early to Mid Holocene in the East Mediterranean basins, Clim. Past Discuss., (February), 1–42, doi:10.5194/cp-2016-4, 2016.

Blaauw, M. and Christen, J. A.: Flexible paleoclimate age-depth models using an autoregressive gamma process, Bayesian Anal., 6(3), 457–474, doi:10.1214/ba/1339616472, 2011.

Bojariu, R. and Giorgi, F.: The North Atlantic Oscillation signal in a regional climate simulation for the European region, Tellus, 57A(4), 641–653, doi:10.1111/j.1600-0870.2005.00122.x, 2005.

Bojariu, R. and Paliu, D.-M.: North Atlantic Oscillation Projection on Romanian Climate Fluctuations in the Cold Season, in Detecting and Modelling Regional Climate Change and Associated Impacts, edited by M. B. India and D. L. Bonillo, pp. 345–356, Springer Berlin Heidelberg, Berlin Heidelburg., 2001.

Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I. and Bonani, G.: Persistent solar influence on North Atlantic climate during the Holocene., Science, 294(5549), 2130–2136, doi:10.1126/science.1065680, 2001.

Booth, R. K., Jackson, S. T., Forman, S. L., Kutzbach, J. E., Bettis, E. a., Kreig, J. and Wright, D. K.: A severe centennial-scale drought in midcontinental North America 4200 years ago and apparent global linkages, The Holocene, 15, 321–328, doi:10.1191/0959683605hl825ft, 2005.

Booth, R. K., Lamentowicz, M. and Charman, D. J.: Preparation and analysis of testate amoebae in peatland palaeoenvironmental studies, Mires Peat, 7(2), 1–7, 2010.

Box, M. R., Krom, M. D., Cliff, R. A., Bar-Matthews, M., Almogi-Labin, A., Ayalon, A. and Paterne, M.: Response of the Nile and its catchment to millennial-scale climatic change since the LGM from Sr isotopes and major elements of East Mediterranean sediments, Quat. Sci. Rev., 30(3), 431–442, doi:10.1016/j.quascirev.2010.12.005, 2011.

Bristow, C. S., Hudson-Edwards, K. A. and Chappell, A.: Fertilizing the Amazon and equatorial Atlantic with West African dust, Geophys. Res. Lett., 37(14), L14807, doi:10.1029/2010GL043486, 2010.

Brooks, N.: Cultural responses to aridity in the Middle Holocene and increased social complexity, Quat. Int., 151(1), 29–49, doi:10.1016/j.quaint.2006.01.013, 2006.

Buczkó, K., Magyari, E. K., Braun, M. and Bálint, M.: Diatom-inferred lateglacial and Holocene climatic

variability in the South Carpathian Mountains (Romania), Quat. Int., 293, 123–135, doi:10.1016/j.quaint.2012.04.042, 2013.

Buggle, B., Glaser, B., Zöller, L., Hambach, U., Marković, S., Glaser, I. and Gerasimenko, N.: Geochemical characterization and origin of Southeastern and Eastern European loesses (Serbia, Romania, Ukraine), Quat. Sci. Rev., 27(9), 1058–1075, doi:10.1016/j.quascirev.2008.01.018, 2008.

Buggle, B., Hambach, U., Glaser, B., Gerasimenko, N., Marković, S., Glaser, I. and Zöller, L.: Stratigraphy, and spatial and temporal paleoclimatic trends in Southeastern/Eastern European loess-paleosol sequences, Quat. Int., 196(1–2), 86–106, doi:10.1016/j.quaint.2008.07.013, 2009.

Carozza, J.-M., Micu, C., Mihail, F. and Carozza, L.: Landscape change and archaeological settlements in the lower Danube valley and delta from early Neolithic to Chalcolithic time: A review, Quat. Int., 261, 21–31, doi:10.1016/j.quaint.2010.07.017, 2012.

Chambers, F. M., Booth, R. K., De Vleeschouwer, F., Lamentowicz, M., Le Roux, G., Mauquoy, D., Nichols, J. E. and van Geel, B.: Development and refinement of proxy-climate indicators from peats, Quat. Int., 268, 21–33, doi:10.1016/j.quaint.2011.04.039, 2012.

Charman, D. J., Hendon, D. and Woodland, W.: The identification of testate amoebae (Protozoa: Rhizopoda) in peats, QRA Techni., Quaternary Research Association, London., 2000.

Cremaschi, M. and Zerboni, A.: Early to Middle Holocene landscape exploitation in a drying environment: Two case studies compared from the central Sahara (SW Fezzan, Libya), Comptes Rendus - Geosci., 341(8–9), 689–702, doi:10.1016/j.crte.2009.05.001, 2009.

Cristea, G., Cuna, S. M., Farcas, S., Tantau, I., Dordai, E. and Magdas, D. A.: Carbon isotope composition as indicator for climatic changes during the middle and late Holocene in a peat bog from Maramures Mountains (Romania), The Holocene, 24, 15–23, doi:10.1177/0959683613512166, 2013.

Cristea, V.: Fitosociologie și vegetația României, Babes-Bolyai University Press, Cluj Napoca., 1993.

Croudace, I. W., Rindby, A. and Rothwell, R. G.: ITRAX: description and evaluation of a new multi-function X-ray core scanner, Geol. Soc. London, 267, 51–63, doi:10.1144/GSL.SP.2006.267.01.04, 2006.

Davis, B. A. S., Brewer, S., Stevenson, A. C. and Guiot, J.: The temperature of Europe during the Holocene reconstructed from pollen data, Quat. Sci. Rev., 22(15–17), 1701–1716, doi:10.1016/S0277-3791(03)00173-2, 2003.

Debret, M., Bout-Roumazeilles, V., Grousset, F., Desmet, M., McManus, J. F., Massei, N., Sebag, D., Petit, J.-R., Copard, Y. and Trentesaux, A.: The origin of the 1500-year climate cycles in Holocene North-Atlantic records, Clim. Past Discuss., 3(2), 679–692, doi:10.5194/cpd-3-679-2007, 2007.

deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L. and Yarusinsky, M.: Abrupt onset and termination of the African Humid Period: Rapid climate responses to gradual insolation forcing, <u>in</u>Quat. Sci. Rev., <u>vol.</u>19 (1–5), <u>pp.</u>347–361, <u>doi:-10.1016/S0277-3791(99)00081-5,...</u>2000.

deMenocal, P. B.: Cultural Responses to Climate Change Duringclimate change during the Latelate Holocene, Science, 292(5517), 667–673, doi:-10.1126/science.1059287, 2001...

Diaconu, A.-C., Grindean, R., Panait, A. and Tanțău, I.: Late Holocene palaeohydrological changes in a *Sphagnum* peat bog from NW Romania based on testate amoebae, Stud. UBB Geol., 60(1), doi: <u>http://dx.doi.org/</u>10.5038/1937-8602.60.1.1285, 2016.

Drăgușin, V., Staubwasser, M., Hoffmann, D. L., Ersek, V., Onac, B. P. and Veres, D.: Constraining Holocene hydrological changes in the Carpathian–Balkan region using speleothem δ 18O and pollen-based temperature reconstructions, Clim. Past, 10(4), 1363–1380, doi:10.5194/cp-10-1363-2014, 2014.

Edri, A., Dody, A., Tanner, S., Swet, N. and Katra, I.: Variations in dust-related PM10 emission from an arid

land due to surface composition and topsoil disturbance, Arab. J. Geosci., 9(12), 607, doi:10.1007/s12517-016-2651-z, 2016.

Egerer, S., Claussen, M., Reick, C. and Stanelle, T.: The link between marine sediment records and changes in Holocene Saharan landscape: Simulating the dust cycle, Clim. Past, 12(4), 1009–1027, doi:10.5194/cp-12-1009-2016, 2016.

Feurdean, A., Klotz, S., Mosbrugger, V. and Wohlfarth, B.: Pollen-based quantitative reconstructions of Holocene climate variability in NW Romania, Palaeogeogr. Palaeoclimatol. Palaeoecol., 260(3–4), 494–504, doi:10.1016/j.palaeo.2007.12.014, 2008.

Feurdean, A., Tămaş, T., Tanțău, I. and Fărcaş, S.: Elevational variation in regional vegetation responses to lateglacial climate changes in the Carpathians, J. Biogeogr., 39(2), 258–271, doi:10.1111/j.1365-2699.2011.02605.x, 2012.

Feurdean, A., Galka, M., Kuske, E., Tantau, I., Lamentowicz, M., Florescu, G., Liakka, J., Hutchinson, S. M., Mulch, A. and Hickler, T.: Last Millennium hydro-climate variability in Central-Eastern Europe (Northern Carpathians, Romania), The Holocene, 25(7), 1179–1192, doi:10.1177/0959683615580197, 2015.

Fletcher, W. J., Debret, M. and Goni, M. F. S.: Mid-Holocene emergence of a low-frequency millennial oscillation in western Mediterranean climate: Implications for past dynamics of the North Atlantic atmospheric westerlies, The Holocene, 23(2), 153–166, doi:10.1177/0959683612460783, 2013.

Francus, P., Lamb, H., Nakagawa, T., Marshall, M., Brown, E. and Members, S. 2006 P.: The potential of high-resolution X-ray fluorescence core scanning : Applications in paleolimnology, PAGES news, 17(3), 93–95, 2009.

Gałka, M., Tanțău, I., Ersek, V. and Feurdean, A.: A 9000year record of cyclic vegetation changes identified in a montane peatland deposit located in the Eastern Carpathians (Central-Eastern Europe): Autogenic succession or regional climatic influences?, Palaeogeogr. Palaeoclimatol. Palaeoecol., 449, 52–61, doi:10.1016/j.palaeo.2016.02.007, 2016.

Giosan, L., Coolen, M. J. L., Kaplan, J. O., Constantinescu, S., Filip, F., Filipova-Marinova, M., Kettner, A. J. and Thom, N.: Early Anthropogenic Transformation of the Danube-Black Sea System, Sci. Rep., 2, 1–6, doi:10.1038/srep00582, 2012.

Goudie, A. S. and Middleton, N. J.: Desert Dust in the Global System, Springer Berlin Heidelberg, Berlin & Heidelburg., 2006.

Grinsted, A., Moore, J. C. and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, Nonlinear Process. Geophys., 11(5/6), 561–566, doi:10.5194/npg-11-561-2004, 2004.

Grousset, F. E. and Biscaye, P. E.: Tracing dust sources and transport patterns using Sr, Nd and Pb isotopes, Chem. Geol., 222(3), 149–167, doi:10.1016/j.chemgeo.2005.05.006, 2005.

Grousset, F. E., Ginoux, P., Bory, A. and Biscaye, P. E.: Case study of a Chinese dust plume reaching the French Alps, Geophys. Res. Lett., 30(6), 1277, doi:10.1029/2002GL016833, 2003.

Haliuc, A., Veres, D., Brauer, A., Hubay, K., Hutchinson, S.M., Begy, R. and Braun, M.: Palaeohydrological changes during the mid and late Holocene in the Carpathian area, central-eastern Europe, Glob. and Planet. Change, 152, 99-114, http://dx.doi.org/10.1016/j.gloplacha.2017.02.010, 2017.

Harangi, S., Molnar, M., Vinkler, A. P., Kiss, B., Jull, A. J. T. and Leonard, A. G.: Radiocarbon Dating of the Last Volcanic Eruptions of Ciomadul Volcano, Southeast Carpathians, Eastern-Central Europe, Radiocarbon, 52(3), 1498–1507, doi:dx.doi.org/10.2458/azu_js_rc.52.3648, 2010.

Heiri, O., Lotter, A. F. and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: Reproducibility and comparability of results, J. Paleolimnol., 25, 101–110, doi:10.1023/A:1008119611481, 2001.

Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global Iron Connections Between Desert Dust, Ocean Biogeochemistry, and Climate, Science, 308, 67-71, doi: 10.1126/science.1105959, 2005.

Jiménez-Espejo, F. J., García-Alix, A., Jiménez-Moreno, G., Rodrigo-Gámiz, M., Anderson, R. S., Rodríguez-Tovar, F. J., Martínez-Ruiz, F., Giralt, S., Delgado Huertas, A. and Pardo-Igúzquiza, E.: Saharan aeolian input and effective humidity variations over western Europe during the Holocene from a high altitude record, Chem. Geol., 374, 1–12, doi:10.1016/j.chemgeo.2014.03.001, 2014.

Kaplan, J. O., Krumhardt, K. M. and Zimmermann, N.: The prehistoric and preindustrial deforestation of Europe, Quat. Sci. Rev., 28(27), 3016–3034, doi:10.1016/j.quascirev.2009.09.028, 2009.

Karátson, D., Wulf, S., Veres, D., Magyari, E. K., Gertisser, R., Timar-Gabor, A., Novothny, Telbisz, T., Szalai, Z., Anechitei-Deacu, V., Appelt, O., Bormann, M., Jánosi, C., Hubay, K. and Schäbitz, F.: The latest explosive eruptions of Ciomadul (Csomád) volcano, East Carpathians - A tephrostratigraphic approach for the 51-29 ka BP time interval, J. Volcanol. Geotherm. Res., 319, 29–51, doi:10.1016/j.jvolgeores.2016.03.005, 2016.

Kok, J. F., Mahowald, N. M., Fratini, G., Gillies, J. A., Ishizuka, M., Leys, J. F. and Mikami, M.: An improved dust emission model – Part 1: Model description and comparison against measurements, Atmos. Chem. Phys, 14, 13023–13041, doi:10.5194/acp-14-13023-2014, 2014.

Korcz, M., Fudała, J. and Kliś, C.: Estimation of wind blown dust emissions in Europe and its vicinity, Atmos. Environ., 43(7), 1410–1420, doi:10.1016/j.atmosenv.2008.05.027, 2009.

Korponai, J., Magyari, E. K., Buczkó, K., Iepure, S., Namiotko, T., Czakó, D., Kövér, C. and Braun, M.: Cladocera response to Late Glacial to Early Holocene climate change in a South Carpathian mountain lake, Hydrobiologia, 676(1), 223–235, doi:10.1007/s10750-011-0881-3, 2011.

Krachler, M., Mohl, C., Emons, H. and Shotyk, W.: Influence of digestion procedures on the determination of rare earth elements in peat and plant samples by USN-ICP-MS, J. Anal. At. Spectrom., 17(8), 844–851, doi:10.1039/b200780k, 2002.

Kristó, A.: A Csomád hegycsoport. A Szent-Anna tó természetvédelmi területe (The Nature Reserve of Lake Saint Ana)., Kristó András emlékére (In Rememb. András Kristó). Balat. Akadémia Könyvek, 13, 38–45, 1995.

Kylander, M. E., Ampel, L., Wohlfarth, B. and Veres, D.: High-resolution X-ray fluorescence core scanning analysis of Les Echets (France) sedimentary sequence: new insights from chemical proxies, J. Quat. Sci., 26(1), 109–117, doi:10.1002/jqs.1438, 2011.

Kylander, M. E., Bindler, R., Cortizas, A. M., Gallagher, K., Mörth, C. M. and Rauch, S.: A novel geochemical approach to paleorecords of dust deposition and effective humidity: 8500 years of peat accumulation at Store Mosse (the "Great Bog"), Sweden, Quat. Sci. Rev., 69, 69–82, doi:10.1016/j.quascirev.2013.02.010, 2013a.

Kylander, M. E., Klaminder, J., Wohlfarth, B. and Löwemark, L.: Geochemical responses to paleoclimatic changes in southern Sweden since the late glacial: the Hässeldala Port lake sediment record, J. Paleolimnol., 50(1), 57–70, doi:10.1007/s10933-013-9704-z, 2013b.

Kylander, M. E., Martínez-Cortizas, A., Bindler, R., Greenwood, S. L., Mörth, C.-M. and Rauch, S.: Potentials and problems of building detailed dust records using peat archives: An example from Store Mosse (the "Great Bog"), Sweden, Geochim. Cosmochim. Acta, 190, 156–174, doi:10.1016/j.gca.2016.06.028, 2016.

Labzovskii, L., Toanca, F. and Nicolae, D.: Determination of Saharan dust properties over Bucharest, Romania. Part 2: Study cases analysis, Rom. J. Phys., 59(9–10), 1097–1108, 2014.

Lamentowicz, M., Cedro, A., Gałka, M., Goslar, T., Miotk-Szpiganowicz, G., Mitchell, E. A. D. and Pawlyta, J.: Last millennium palaeoenvironmental changes from a Baltic bog (Poland) inferred from stable isotopes, pollen, plant macrofossils and testate amoebae, Palaeogeogr. Palaeoclimatol. Palaeoecol., 265(1), 93–106, doi:10.1016/j.palaeo.2008.04.023, 2008.

Longman, J., Ersek, V., Veres, D. and Salzmann, U.: Detrital events and hydroclimate variability in the Romanian Carpathians during the Mid-to-Late Holocene, Quat. Sci. Rev., under review.

Lüdecke, H.-J., Weiss, C. O. and Hempelmann, A.: Paleoclimate forcing by the solar De Vries/Suess cycle, Clim. Past Discuss., 11(1), 279–305, doi:10.5194/cpd-11-279-2015, 2015.

Magny, M.: Holocene climate variability as reflected by mid-European lake-level fluctuations and its probable impact on prehistoric human settlements, Quat. Int., 113(1), 65–79, doi:10.1016/S1040-6182(03)00080-6, 2004.

Magny, M., Combourieu-Nebout, N., De Beaulieu, J. L., Bout-Roumazeilles, V., Colombaroli, D., Desprat, S., Francke, A., Joannin, S., Ortu, E. and Peyron, O.: Geoscientific Instrumentation Methods and Data Systems North–south palaeohydrological contrasts in the central Mediterranean during the Holocene: tentative synthesis and working hypotheses, Clim. Past, 9, 2043–2071, doi:10.5194/cp-9-2043-2013, 2013.

Magyari, E., Buczkó, K., Jakab, G., Braun, M., Pál, Z., Karátson, D., and Pap, I.: Palaeolimnology of the last crater lake in the Eastern Carpathian Mountains: a multiproxy study of Holocene hydrological changes, Hydrobiologia, 631, 29-63, doi: 10.1007/s10750-009-9801-1, 2009.

Magyari, E. K., Chapman, J. C., Passmore, D. G., Allen, J. R. M., Huntley, J. P. and Huntley, B.: Holocene persistence of wooded steppe in the Great Hungarian Plain, J. Biogeogr., 37, 915–935, doi:10.1111/j.1365-2699.2009.02261.x, 2010.

Magyari, E. K., Jakab, G., Bálint, M., Kern, Z., Buczkó, K. and Braun, M.: Rapid vegetation response to Lateglacial and early Holocene climatic fluctuation in the South Carpathian Mountains (Romania), Quat. Sci. Rev., 35, 116–130, doi:10.1016/j.quascirev.2012.01.006, 2012.

Magyari, E. K., Demény, A., Buczkó, K., Kern, Z., Vennemann, T., Fórizs, I., Vincze, I., Braun, M., Kovács, J. I., Udvardi, B. and Veres, D.: A 13,600-year diatom oxygen isotope record from the South Carpathians (Romania): Reflection of winter conditions and possible links with North Atlantic circulation changes, Quat. Int., 293, 136–149, doi:10.1016/j.quaint.2012.05.042, 2013.

Magyari, E. K., Veres, D., Wennrich, V., Wagner, B., Braun, M., Jakab, G., Karátson, D., Pál, Z., Ferenczy, G., St-Onge, G., Rethemeyer, J., Francois, J.-P., von Reumont, F. and Schäbitz, F.: Vegetation and environmental responses to climate forcing during the Last Glacial Maximum and deglaciation in the East Carpathians: attenuated response to maximum cooling and increased biomass burning, Quat. Sci. Rev., 106, 278–298, doi:10.1016/j.quascirev.2014.09.015, 2014.

Mahowald, N. M., Kloster, S., Engelstaedter, S., Moore, J. K., Mukhopadhyay, S., Mcconnell, J. R., Albani, S., Doney, S. C., Bhattacharya, A., Curran, M. A. J., Flanner, M. G., Hoffman, F. M., Lawrence, D. M., Lindsay, K., Mayewski, P. A., Neff, J., Rothenberg, D., Thomas, E., Thornton, P. E. and Zender, C. S.: Observed 20th century desert dust variability: impact on climate and biogeochemistry, Atmos. Chem. Phys. Atmos. Chem. Phys., 10, 10875–10893, doi:10.5194/acp-10-10875-2010, 2010.

Mann, M. E., Zhang, Z., Rutherford, S., Bradley, R. S., Hughes, M. K., Shindell, D., Ammann, C., Faluvegi, G., and Ni, F.: Global Signatures and Dynamical Origins of the Little Ice Age and Medieval Climate Anomaly, Science, 326(5957), 1256-1260, doi: 10.1126/science.1177303, 2009.

Marković, S. B., Stevens, T., Kukla, G. J., Hambach, U., Fitzsimmons, K. E., Gibbard, P., Buggle, B., Zech, M., Guo, Z., Hao, Q., Wu, H., O'Hara Dhand, K., Smalley, I. J., Újvári, G., Sümegi, P., Timar-Gabor, A., Veres, D., Sirocko, F., Vasiljević, D. A., Jary, Z., Svensson, A., Jović, V., Lehmkuhl, F., Kovács, J. and Svirčev, Z.: Danube loess stratigraphy — Towards a pan-European loess stratigraphic model, Earth–Sci. Rev., 148, 228–258, doi:10.1016/j.earscirev.2015.06.005, 2015.

Marx, S. K., McGowan, H. A. and Kamber, B. S.: Long-range dust transport from eastern Australia: A proxy for Holocene aridity and ENSO-type climate variability, Earth Planet. Sci. Lett., 282(1–4), 167–177, doi:10.1016/j.epsl.2009.03.013, 2009.

Marx, S. K., Kamber, B. S., McGowan, H. a. and Zawadzki, A.: Atmospheric pollutants in alpine peat bogs record a detailed chronology of industrial and agricultural development on the Australian continent, Environ. Pollut., 158(5), 1615–1628, doi:10.1016/j.envpol.2009.12.009, 2010.

Mauri, A., Davis, B. A. S., Collins, P. M. and Kaplan, J. O.: The climate of Europe during the Holocene: a gridded pollen-based reconstruction and its multi-proxy evaluation, Quat. Sci. Rev., 112, 109–127, doi:10.1016/j.quascirev.2015.01.013, 2015.

Mayewski, P. a, Rohling, E., Curtstager, J., Karlén, W., Maasch, K., Davidmeeker, L., Meyerson, E., Gasse, F., Vankreveld, S. and Holmgren, K.: Holocene climate variability, Quat. Res., 62(3), 243–255, doi:10.1016/j.yqres.2004.07.001, 2004.

McGee, D., DeMenocal, P. B., Winckler, G., Stuut, J. B. W. and Bradtmiller, L. I.: The magnitude, timing and abruptness of changes in North African dust deposition over the last 20,000 yr, Earth Planet. Sci. Lett., 371–372, 163–176, doi:10.1016/j.epsl.2013.03.054, 2013.

Miao, X., Mason, J. A., Swinehart, J. B., Loope, D. B., Hanson, P. R., Goble, R. J. and Liu, X.: A 10,000 year record of dune activity, dust storms, and severe drought in the central Great Plains, Geology, 35(2), 119, doi:10.1130/G23133A.1, 2007.

Molnár, M., Rinyu, L., Veres, M., Seiler, M., Wacker, L. and Synal, H.-A.: EnvironMICADAS : a mini 14C AMS with enhanced gas ion source, Radiocarbon, 55(2), 338–344, doi:10.2458/azu_js_rc.55.16331, 2013.

Morellón, M., Anselmetti, F. S., Ariztegui, D., Brushulli, B., Sinopoli, G., Wagner, B., Sadori, L., Gilli, A. and Pambuku, A.: Human–climate interactions in the central Mediterranean region during the last millennia: The laminated record of Lake Butrint (Albania), Quat. Sci. Rev., 136, 134–152, doi:10.1016/j.quascirev.2015.10.043, 2016.

Moreno, T., Querol, X., Castillo, S., Alastuey, A., Cuevas, E., Herrmann, L., Mounkaila, M., Elvira, J. and Gibbons, W.: Geochemical variations in aeolian mineral particles from the Sahara–Sahel Dust Corridor, Chemosphere, 65(2), 261–270, doi:10.1016/j.chemosphere.2006.02.052, 2006.

Morley, A., Rosenthal, Y. and DeMenocal, P.: Ocean-atmosphere climate shift during the mid-to-late Holocene transition, Earth Planet. Sci. Lett., 388, 18–26, doi:10.1016/j.epsl.2013.11.039, 2014.

Morris, P. J., Baird, A. J., Young, D. M. and Swindles, G. T.: Untangling climate signals from autogenic changes in long-term peatland development, Geophys. Res. Lett., 42(24), 10,788-10,797, doi:10.1002/2015GL066824, 2015.

Mulitza, S., Heslop, D., Pittauerova, D., Fischer, H. W., Meyer, I., Stuut, J.-B., Zabel, M., Mollenhauer, G., Collins, J. A., Kuhnert, H. and Schulz, M.: Increase in African dust flux at the onset of commercial agriculture in the Sahel region, Nature, 466(7303), 226–228, doi:10.1038/nature09213, 2010.

Nichols, J. E. and Huang, Y.: Hydroclimate of the northeastern United States is highly sensitive to solar forcing, Geophys. Res. Lett., 39(4), L04707, doi:10.1029/2011GL050720, 2012.

Nicolás, J., Chiari, M., Crespo, J., Orellana, I. G., Lucarelli, F., Nava, S., Pastor, C. and Yubero, E.: Quantification of Saharan and local dust impact in an arid Mediterranean area by the positive matrix factorization (PMF) technique, Atmos. Environ., 42(39), 8872–8882, doi:10.1016/j.atmosenv.2008.09.018, 2008.

Notaro, M., Yu, Y. and Kalashnikova, O. V.: Regime shift in Arabian dust activity, triggered by persistent Fertile Crescent drought, J. Geophys. Res. Atmos., 120(19), 10,229-10,249, doi:10.1002/2015JD023855, 2015.

Novak, M., Zemanova, L., Voldrichova, P., Stepanova, M., Adamova, M., Pacherova, P., Komarek, A., Krachler, M. and Prechova, E.: Experimental Evidence for Mobility/Immobility of Metals in Peat, Environ. Sci. Technol., 45(17), 7180–7187, doi:10.1021/es201086v, 2011.

Obreht, I., Zeeden, C., Hambach, U., Veres, D., Marković, S. b., Bösken, J., Svirčev, Z., Bačević, N., Gavrilov, M. B. and Lehmkuhl, F.: Tracing the influence of Mediterranean climate on Southeastern Europe during the past 350,000 years, Sci. Rep., 6, 36334, doi:10.1038/srep36334, 2016.

Poore, R. Z., Quinn, T. M. and Verardo, S.: Century-scale movement of the Atlantic Intertropical Convergence Zone linked to solar variability, Geophys. Res. Lett., 31(12), L12214, doi:10.1029/2004GL019940, 2004.

Pop, E.: Mlaștinile de turbă din Republica Populară Română (Peat bogs from Romania), Editura Academiei Republicii Populare Române, Bucharest., 1960.

Poto, L., Gabrieli, J., Crowhurst, S., Agostinelli, C., Spolaor, A., Cairns, W. R. L., Cozzi, G. and Barbante, C.: Cross calibration between XRF and ICP-MS for high spatial resolution analysis of ombrotrophic peat cores for palaeoclimatic studies, Anal. Bioanal. Chem., 407, 379–385, doi:10.1007/s00216-014-8289-3, 2014.

Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Aerosols, Climate, and the Hydrological Cycle, Science, 294, 2119-2124, doi: 10.1126/science.1064034, 2001.

Rasmussen, S. O., Vinther, B. M., Clausen, H. B. and Andersen, K. K.: Early Holocene climate oscillations recorded in three Greenland ice cores, Quat. Sci. Rev., 26(15–16), 1907–1914, doi:10.1016/j.quascirev.2007.06.015, 2007.

Reimer, P., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R. A., Turney, C. S. M. and van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP, Radiocarbon, 55(4), 1869–1887, doi:10.2458/azu_js_rc.55.16947, 2013.

Revel, M., Ducassou, E., Grousset, F. E., Bernasconi, S. M., Migeon, S., Revillon, S., Mascle, J., Murat, A., Zaragosi, S. and Bosch, D.: 100,000 Years of African monsoon variability recorded in sediments of the Nile margin, Quat. Sci. Rev., 29(11), 1342–1362, doi:10.1016/j.quascirev.2010.02.006, 2010.

Roberts, N., Moreno, A., Valero-Garcés, B. L., Corella, J. P., Jones, M., Allcock, S., Woodbridge, J., Morellón, M., Luterbacher, J., Xoplaki, E. and Türkeş, M.: Palaeolimnological evidence for an east-west climate see-saw in the Mediterranean since AD 900, Glob. Planet. Change, 84–85, 23–34, doi:10.1016/j.gloplacha.2011.11.002, 2012.

Ross-Barraclough, F. and Shotyk, W.: Millennial-scale records of atmospheric mercury deposition obtained from ombrotrophic and minerotrophic peatlands in the Swiss Jura Mountains, Environ. Sci. Technol., 37(2), 235–244, doi:10.1021/es0201496, 2003.

Rothwell, J. J., Taylor, K. G., Chenery, S. R. N., Cundy, A. B., Evans, M. G. and Allott, T. E. H.: Storage and behavior of As, Sb, Pb, and Cu in ombrotrophic peat bogs under contrasting water table conditions, Environ. Sci. Technol., 44(22), 8497–8502, doi:10.1021/es101150w, 2010.

Rousseau, D. D., Chauvel, C., Sima, A., Hatté, C., Lagroix, F., Antoine, P., Balkanski, Y., Fuchs, M., Mellett, C., Kageyama, M., Ramstein, G. and Lang, A.: European glacial dust deposits: Geochemical constraints on atmospheric dust cycle modeling, Geophys. Res. Lett., 41(21), 7666–7674, doi:10.1002/2014GL061382, 2014.

Le Roux, G., Fagel, N., De Vleeschouwer, F., Krachler, M., Debaille, V., Stille, P., Mattielli, N., van der Knaap, W. O., van Leeuwen, J. F. N. and Shotyk, W.: Volcano- and climate-driven changes in atmospheric dust sources and fluxes since the Late Glacial in Central Europe, Geology, 40(4), 335–338, doi:10.1130/g32586.1, 2012.

Russell, J. M., Johnson, T. C., and Talbot, M. R.: A 725 yr cycle in the climate of central Africa during the late Holocene, Geology, 31(8), 677-680, doi: 10.1130/g19449.1, 2003.

-Sapkota, A., Cheburkin, A. K., Bonani, G. and Shotyk, W.: Six millennia of atmospheric dust deposition in southern South America (Isla Navarino, Chile), The Holocene, 17(5), 561–572, doi:10.1177/0959683607078981, 2007.

Scheuvens, D., Schütz, L., Kandler, K., Ebert, M. and Weinbruch, S.: Bulk composition of northern African dust and its source sediments — A compilation, Earth-Science Rev., 116, 170–194, doi:10.1016/j.earscirev.2012.08.005, 2013.

Schnitchen, C., Charman, D. J., Magyari, E., Braun, M., Grigorszky, I., Tóthmérész, B., Molnár, M. and Szántó, Z.: Reconstructing hydrological variability from testate amoebae analysis in Carpathian peatlands, J. Paleolimnol., 36(1), 1–17, doi:10.1007/s10933-006-0001-y, 2006.

Schumacher, M., Schier, W. and Schütt, B.: Mid-Holocene vegetation development and herding-related interferences in the Carpathian region, Quat. Int., 415, 253–267, doi:10.1016/j.quaint.2015.09.074, 2016.

Shanahan, T. M., McKay, N. P., Hughen, K. A., Overpeck, J. T., Otto-Bliesner, B., Heil, C. W., King, J., Scholz, C. A. and Peck, J.: The time-transgressive termination of the African Humid Period, Nat. Geosci., 8(2), 140–144, doi:10.1038/ngeo2329, 2015.

Sharifi, A., Pourmand, A., Canuel, E. A., Ferer-Tyler, E., Peterson, L. C., Aichner, B., Feakins, S. J., Daryaee, T., Djamali, M., Beni, A. N., Lahijani, H. A. K. and Swart, P. K.: Abrupt climate variability since the last deglaciation based on a high-resolution, multi-proxy peat record from NW Iran: The hand that rocked the Cradle of Civilization?, Quat. Sci. Rev., 123, 215–230, doi:10.1016/j.quascirev.2015.07.006, 2015.

Shotyk, W.: The chronology of anthropogenic, atmospheric Pb deposition recorded by peat cores in three minerogenic peat deposits from Switzerland, Sci. Total Environ., 292, 19–31, doi:10.1016/S0048-9697(02)00030-X, 2002.

Shotyk, W., Krachler, M., Martinez-Cortizas, A., Cheburkin, A. K. and Emons, H.: A peat bog record of natural, pre-anthropogenic enrichments of trace elements in atmospheric aerosols since 12 370 14C yr BP, and their variation with Holocene climate change, Earth Planet. Sci. Lett., 199(1–2), 21–37, doi:10.1016/S0012-821X(02)00553-8, 2002.

Smalley, I., Marković, S. B. and Svirčev, Z.: Loess is [almost totally formed by] the accumulation of dust, Quat. Int., 240(1–2), 4–11, doi:10.1016/j.quaint.2010.07.011, 2011.

Springer, G. S., Rowe, H. D., Hardt, B., Edwards, R. L. and Cheng, H.: Solar forcing of Holocene droughts in a stalagmite record from West Virginia in east-central North America, Geophys. Res. Lett., 35(17), L17703, doi:10.1029/2008GL034971, 2008.

Sweeney, M. R. and Mason, J. A.: Mechanisms of dust emission from Pleistocene loess deposits, Nebraska, USA, J. Geophys. Res. Earth Surf., 118(3), 1460–1471, doi:10.1002/jgrf.20101, 2013.

Swindles, G. T., Blundell, A., Roe, H. M. and Hall, V. A.: A 4500-year proxy climate record from peatlands in the North of Ireland: the identification of widespread summer "drought phases"?, Quat. Sci. Rev., 29(13), 1577–1589, doi:10.1016/j.quascirev.2009.01.003, 2010.

Swindles, G. T., Patterson, R. T., Roe, H. M. and Galloway, J. M.: Evaluating periodicities in peat-based climate proxy records, Quat. Sci. Rev., 41, 94–103, doi:10.1016/j.quascirev.2012.03.003, 2012.

Szakács, A., Seghedi, I., Pécskay, Z. and Mirea, V.: Eruptive history of a low-frequency and low-output rate Pleistocene volcano, Ciomadul, South Harghita Mts., Romania, Bull. Volcanol., 77(2), 12, doi:10.1007/s00445-014-0894-7, 2015.

Tanțău, I., Reille, M., De Beaulieu, J. L., Farcas, S., Goslar, T. and Paterne, M.: Vegetation history in the Eastern Romanian Carpathians: Pollen analysis of two sequences from the Mohos crater, Veg. Hist. Archaeobot., 12(2), 113–125, doi:10.1007/s00334-003-0015-6, 2003.

Tanțău, I., Feurdean, A., De Beaulieu, J. L., Reille, M. and Fărcaş, S.: Vegetation sensitivity to climate changes and human impact in the Harghita Mountains (Eastern Romanian Carpathians) over the past 15 000 years, J. Quat. Sci., 29(2), 141–152, doi:10.1002/jqs.2688, 2014.

Torrence, C., and Compo, G. P.: A Practical Guide to Wavelet Analysis, Bulletin of the American Meteorological Society, 79(1), 61-78, doi: 10.1175/1520-0477(1998)079<0061:apgtwa>2.0.co;2, 1998.

-Újvári, G., Varga, A., Ramos, F. C., Kovács, J., Németh, T. and Stevens, T.: Evaluating the use of clay mineralogy, Sr–Nd isotopes and zircon U–Pb ages in tracking dust provenance: An example from loess of the Carpathian Basin, Chem. Geol., 304, 83–96, doi:10.1016/j.chemgeo.2012.02.007, 2012.

Varga, G., Kovács, J. and Újvári, G.: Analysis of Saharan dust intrusions into the Carpathian Basin (Central Europe) over the period of 1979–2011, Glob. Planet. Change, 100, 333–342, doi:10.1016/j.gloplacha.2012.11.007, 2013.

Varga, G., Cserháti, C., Kovács, J. and Szalai, Z.: Saharan dust deposition in the Carpathian Basin and its possible effects on interglacial soil formation, Aeolian Res., 22, 1–12, doi:10.1016/j.aeolia.2016.05.004, 2016.

Veron, A., Novak, M., Brizova, E. and Stepanova, M.: Environmental imprints of climate changes and anthropogenic activities in the Ore Mountains of Bohemia (Central Europe) since 13 cal. kyr BP, The Holocene, 24(8), 919–931, doi:10.1177/0959683614534746, 2014.

Vinkler, A. P., Harangi, S., Ntaflos, T. and Szakács, A.: A Csornád vulkán (Keleti-Kárpátok) horzsaköveinek kőzettani és geokémiai vizsgálata - petrogenetikai következtetések, Földtani Közlöny_, 137(1), 103–128, 2007.

Vukmirović, Z., Unkašević, M., Lazić, L., Tošić, I., Rajšić, S. and Tasić, M.: Analysis of the Saharan dust regional transport, Meteorol. Atmos. Phys., 85(4), 265–273, doi:10.1007/s00703-003-0010-6, 2004.

Wanner, H., Solomina, O., Grosjean, M., Ritz, S. P. and Jetel, M.: Structure and origin of Holocene cold events, Quat. Sci. Rev., 30(21–22), 3109–3123, doi:10.1016/j.quascirev.2011.07.010, 2011.

Wedepohl, K. H.: The composition of the continental crust, Geochim. Cosmochim. Acta, 59(7), 1217–1232, doi:10.1016/0016-7037(95)00038-2, 1995.

Wulf, S., Fedorowicz, S., Veres, D., Łanczont, M., Karátson, D., Gertisser, R., Bormann, M., Magyari, E., Appelt, O., Hambach, U. and Gozhyk, P. F.: The "Roxolany Tephra" (Ukraine) – new evidence for an origin from Ciomadul volcano, East Carpathians, J. Quat. Sci., 31(6), 565–576, doi:10.1002/jqs.2879, 2016.

Yoshioka, M., Mahowald, N. M., Conley, A. J., Collins, W. D., Fillmore, D. W., Zender, C. S. and Coleman, D. B.: Impact of desert dust radiative forcing on sahel precipitation: Relative importance of dust compared to sea surface temperature variations, vegetation changes, and greenhouse gas warming, J. Clim., 20(8), 1445–1467, doi:10.1175/JCLI4056.1, 2007.

Yu, S.-Y.: Centennial-scale cycles in middle Holocene sea level along the southeastern Swedish Baltic coast, Geol. Soc. Am. Bull., 115(11), 1404, doi:10.1130/B25217.1, 2003.



Figure 1: 1.A: Figure 1: 1.A: Map of the Carpathian-Balkan region indicating location of Mohos peat bog (red star), in the South-Eastern Carpathian Mountains. Predominant wind directions relating to air circulation patterns in the area are indicated by black arrows. Major Saharan dust source areas are indicated in yellow (Scheuvens et al., 2013) and local loess fields (including loess-derived alluvium) in green (Marković et al., 2015). 1.B: Map of Mohos and neighbouring Lake Sf Ana, from Google Earth 6.1.7601.1 (June 10th 2016). Harghita County, Romania, 46°05' N ; 25°55' E, Eye alt 3.06 km, CNAS/Astrium, DigitalGlobe 2016. http://www.google.com/earth/index.html (Accessed January 23rd 2017). Coring location within white box. 1.C: Photo of Mohos bog at the coring location with the crater rim visible in the distance.



Figure 2: Age-depth model of Mohos peat record, as determined via Bacon (Blaauw and Christen, 2011). Upper left graph indicates Markov Chain Monte Carlo iterations. Also on the upper panel are prior (green line) and posterior (grey histogram) distributions for the accumulation rate (middle) and memory (right). For the lower panel, calibrated radiocarbon ages are in blue. The age-depth model is outlined in grey, with darker grey indicating more likely calendar ages. Grey stippled lines show 95% confidence intervals, and the red curve indicates the single 'best' model used in this work.



Figure 3: ITRAX-<u>derived data of</u> lithogenic <u>element contents</u> (K, Si, and Ti) <u>concentration</u> throughout the Mohos peat record, with all data smoothed using a 9-point moving average to eliminate noise. Alongside, dust flux as reconstructed from Ti concentration values (also displayed), and sedimentation rate is presented. Dust events (D0-D10), as identified from increases in at least two of the lithogenic elements under discussion, are highlighted in brown, and labelled. <u>Dashed lines on ITRAX data indicate the enrichment above which a dust event is denoted.</u>



Figure 4: Comparison of Ti-derived dust flux record with wet and dry TA indicator species % values, reconstructed Depth to Water Table (DWT) and organic matter (as indicated by Loss on Ignition). Vertical bars as in Fig. 3.



Figure 5: Comparison of dust flux values as reconstructed from Mohos peat bog with similar records. Two Western African dust flux records (GC 68 and 66) from marine cores (McGee et al., 2013), are presented alongside bog-based records from Misten bog in Belgium (Allan et al., 2013) and Etang de la Gruyere in Switzerland (Le Roux et al., 2012) respectively. Indicated on these records are volcanic events as identified by the authors (brown triangles). These are presented alongside the dust flux record from Mohos (lower panel). Also shown, in brown, are periods of Rapid Climate Change derived from Greenland Ice cores (Mayewski et al., 2004). Verticle bars as in Fig. 3.



Figure 6: Correlation graphs and gradients of normalised Ti versus normalised K for each of the dust events (D1-D10)



Figure 7: Comparison of dust events and bog wetness as reconstructed from the Mohos record, to regional hydroclimate reconstructions. Data presented via green bars is drought/dry/low lake periods from the following publications. A: (Magny, 2004), B: (Cristea et al., 2013), C: (Galka et al., 2016), D: (Magyari et al., 2013), E: (Buczkó et al., 2013), F: (Magyari et al., 2009), G: (Schnitchen et al., 2006). These are presented alongside the Mohos testate amoeba-derived Depth to Water Table record, and Ti-derived dust flux.



Figure 8: Spectral analysis of Mohos ITRAX geochemical data for A: <u>K</u>, B: Si, C: <u>Ti</u>. Areas outlined in black are significant at the 95% confidence level. Shaded area indicates the cone of influence, outside of which results may be unreliable.

Lab ID	Depth	¹⁴ C age (yr BP $\pm 1\sigma$)	Calibrated age (cal yr BP $\pm 2\sigma$)	Dated material	
DeA-8343	50	37±18	37-65	bulk peat	
DeA-8344	100	838±19	700-785	bulk peat	
DeA-10111	150	1174±28	1049-1179	bulk peat	
DeA-10112	175	1471±26	1309-1399	bulk peat	
DeA-8345	200	2022±21	1921-2007	bulk peat	
DeA-10137	225	2155±27	2048-2305	bulk peat	
DeA-10138	280	2530±28	2495-2744	bulk peat	
DeA-8346	300	3112±23	3249-3383	bulk peat	
DeA-10139	350	4110±31	4523-4713	bulk peat	
DeA-10140	380	4641±54	5282-5484	bulk peat	
DeA-8347	400	4638±26	5372-5463	bulk peat	
DeA-10141	500	5949±36	6677-6861	bulk peat	
DeA-10142	600	6989±43	7785-7867	bulk peat	
DeA-8348	700	7909±33	8600-8793	bulk peat	
DeA-10143	800	8687±45	9539-9778	bulk peat	
DeA-8349	900	9273±36	10369-10571	bulk peat	

Table 1: Radiocarbon dates used to build the age model for Mohos peat record.

Table 2: Ti-K correlation (\mathbb{R}^2) , alongside average cps for K and Ti for each of the dust events as identified within the Mohos core.

Dust Event	D1	D2	D3	D4	D5	D6	D7	D8	D9	D10
Ti-K Correlation (R ²)	0.072	0.111	0.314	0.162	0.296	0.809	0.248	0.758	0.671	0.645
Average Ti (Normalised cps)	0.0015	0.0015	0.0022	0.0018	0.0026	0.0061	0.0048	0.0044	0.0031	0.0052
Average K (Normalised cps)	0.0006	0.0006	0.0006	0.0007	0.0009	0.0018	0.0016	0.0013	0.0011	0.0064