

Reply to the Referees' comments

The authors are thankful to all the three referees for their critical and constructive comments which helped in improving the manuscript. The necessary changes in view of the comments made by the referee have been incorporated in the manuscript. Here, we provide replies to the comments. In view of the comments and suggestions made by the reviewers, we have changed the title of the manuscript from “Strengthening of the Somali upwelling during the Holocene and its impact on southwest monsoon rainfall” to “Variations of the Somali upwelling since 18.5 ka BP and its relationship with southwest monsoon rainfall”. The manuscript presently focuses on variation of the biogenic silica during the last 18.5 ka and possible implications.

Referee #1

We thank the referee for his recommendation and positive view on our manuscript.

Referee #2

Comment: Line 11: 18.5-15 ka is referred to as the glacial maximum; Line 12: the BA is referred to as post-glacial. The latter term is misleading as the Holocene starts at 11.7 ka.

Reply: The Bølling-Allerød (15-12.9 ka BP) is generally considered to be a warming phase after the last glacial maximum. We did not refer to the B/A event as post glacial one, but the upwelling signal shows an increase during the B/A. This increase in upwelling signal after last glacial maximum is referred here as post glacial intensification of Southwest monsoon. The sentence has been rephrased in the revised manuscript.

Comment: Lines 38-39: the introduction to biogenic silica is too short here. But 4.1 would fit in here.

Reply: The introduction to biogenic silica has been modified by bringing the discussion part as per reviewer suggestion.

Comment: Lines 75ff: repeated in the end of chapter 4.1; see comments on lines 156-162

Reply: The repeated sentences were deleted in the section 4.1 (now part of introduction) from line 158 to 160 (now 80 to 83).

Comment: *The first chapter (4.1) is a very useful part of an introduction because it outlines the problem of silica preservation and thus the suitability of this proxy as a productivity indicator. As own data are not discussed this is certainly not part of a discussion.*

Reply: We agree with the referee that the chapter 4.1 (now part of introduction) does not contain own data, however it contains the synthesis of the previous study in the study area and gives a complete idea about the proxy. As advised, this chapter has been moved to introduction to give a more comprehensive details on biogenic silica.

Comment: *The last lines 156-162 of chapter 4.1 repeat what has already been written in the methods. However, I find this part still very problematic and I am still not sure if I understand it correctly. I guess that the authors used the age model of Tiwari et al. (2010) which is of course what should be done. Second, I understand that the authors used an average sedimentation rate to calculate the biogenic opal accumulation rates. This cannot be done! As Fig. 4 shows and as the authors also state there are large variations of the sedimentation rates with lower rates during the Holocene and higher during the glacial section. If an average rate of the entire core is used these differences are concealed. I still don't understand why this is done and I think that it is not justified to do it. The authors could compare % biogenic Si with accumulation rates and discuss whether the differences are related to winnowing/focusing, they could also discuss problems with accumulation rates but they cannot blur these differences by using an average sedimentation rate.*

Reply: The repeated lines in the chapter 4.1 (now in introduction) have been deleted as per reviewer's suggestion. The average sedimentation rate has been used to overcome the problems of sediment redistribution if at all present in the study area. However, we had no intentions to hide the use of average sedimentation rate and thus have shown the age-depth model in figure 4 and discussed about the validity of its use in the present study (lines 125-127). This is based on the similar method used in a referred study by deMenocal et al.,2000 in east Atlantic sediment core to understand the west African climate.

Comment: *The discussion of the applicability of different temperature proxies (4.2.) is certainly dispensable in this paper as it does not produce own SST data. At least until line 185 the whole section could be deleted.*

Reply: The chapter 4.2 (now 3.1) was added during the first revision of this manuscript as per reviewers' comment on the comparison of biogenic silica flux and SST. We agree with the reviewer that the introduction of different SST proxies in chapter 4.2 is very extensive. Major part of chapter 4.2 have been now deleted in the corrected manuscript as suggested.

Comment: *The next extensive discussion of published data (4.3, lines 198-234) is more important to the data interpretation as the authors discuss how Arabian Sea SST (modulated by upwelling) and moisture transport to the Indian Subcontinent are related. This discussion would need to be focused and shortened. From line 229 to 234 it is much better.*

Reply: We have tried to discuss this at length given the complex nature of relation between the Somali upwelling and Indian rainfall and also to give a fair view on the uncertainties around this relationship.

Comment: *Results and discussion are mixed and the points are not very clearly presented. The results section could therefore be deleted.*

Reply: Result section has been deleted in the revised manuscript as per referee's suggestion.

Referee #3

Comment: *The major part of the paper and discussion (Line 198 - 273) deals with the time period between 18.5 to 11.7 ka BP including Last Glacial Period (18.5 to 15 ka BP) and Deglacial period (15-11.7 ka BP), while only one portions (Line 275 to 310) deals with the Holocene. But paper title gives an impression that entire work is on Holocene time period. I will suggest to make changes in the title accordingly like "Variations of the Somali upwelling since 18.5 ka BP and its relationship with southwest monsoon rainfall".*

Reply: We accept the reviewer's concern and changed the manuscript title to "Variations of the Somali upwelling since 18.5 ka BP and its relationship with southwest monsoon rainfall", as per the suggestion.

Comment: *It is still not clear that, how average sedimentation rate is different from the sedimentation rate given in the Tiwari et al., 2010.*

Reply: Age-depth model presented in Tiwari et al., 2010 was based on the age control points shown in figure 4 and their sedimentation rate was calculated using the variation in two adjacent age control points. However the average sedimentation rate method considers only two dates (minimum and maximum) in the sediment core and give a constant sedimentation rate for the whole core, which is mainly used to overcome the problems associated with sediment redistribution.

Comment: *In Figure 5, 6, 7 the Biogenic silica flux shows increasing trend from ~17 to 13.5, ~11.7 to 9.7, ~8.2 to 5, and ~4 to 2 ka BP and sharp decrease between ~13.5 and 11.7, ~9.7 and 8.2 and ~5 and 4 ka BP. What processes is attributed to sharp decrease in upwelling*

during these periods like 13.5 to 11.7 attributed to Younger Dryas. The Mg/Ca SST is annual SST measured by Anand et al., 2008, while anti-phase relationship between SST and upwelling is seasonal phenomenon (Fig.3) remain for a quarter of year as suggested by the authors. It means Mg/Ca SST shows major glacial/interglacial variations in the SST, than, why authors are trying to co-relate with the upwelling proxy?

Reply: The short term variations in the biogenic silica flux were discussed in the chapter 3.2. We have discussed the possible reasons for the variation in the flux. SST is physically connected with upwelling, and an important proxy for reconstruction of upwelling history. The figure 1 (now figure 1) shows the modern data set with the SST values being measured directly. But the Mg/Ca proxy is based on the foraminifera abundance which may or may not preserve the seasonal SST signal based on the productivity dynamics in a region. For example, if the foram abundance was influenced by seasonal changes in the hydrography then the Mg/Ca of foram would give seasonal SST rather than annual. The reason behind the comparison of biogenic silica with SST, was to show the suitability of biogenic silica flux over SST as upwelling proxy in our study area.

Comment: *Various marine as well as land based proxies suggests huge variations in southwest monsoon during the Holocene. The overall biogenic flux remains between 6 to 4 g.m-2.y-1. On close look (Figure 10), the author may observe low southwest monsoon during low upwelling (i.e., 4 g.m-2.y-1). Hence, line 312 Somali upwelling possibly had a negative impact on southwest monsoon rainfall throughout the Holocene is over exaggeration as authors have not done one to one correlation between upwelling and southwest monsoon record.*

Reply: We agree with the referee that this study does not present one to one correlation between the Somali upwelling and southwest monsoon rainfall. The overall trends have been therefore compared and interpreted accordingly. Though there are points of anti-correlation, the trend shows a negative impact of Somali upwelling on southwest monsoon rainfall baring the beginning of Holocene. The line 368 has been modified as “It is observed that baring the beginning, the Somali upwelling had a negative impact on southwest monsoon rainfall over south-western India and Oman during Holocene”.

Comment: *Line 124 – 126 is repetition of line 107 -117*

Reply: Result section has been deleted in the revised manuscript.

Comment: *Degree symbol is missing at line 178.*

Reply: This part has also been deleted in the revised manuscript as per referee #2 suggestions.

Variations of the Somali upwelling since 18.5 ka BP and its relationship with southwest monsoon rainfall

~~Strengthening of the Somali upwelling during the Holocene and its impact on southwest monsoon rainfall~~

5 Balaji D.^{1,2}, Ravi Bhushan², L. S. Chamyal¹

¹Department of Geology, The Maharaja Sayajirao University of Baroda, India

²Geoscience Division, Physical Research Laboratory, Ahmedabad, India

Correspondence to: Balaji D. (balaji.d86@gmail.com)

Abstract. ~~The history of the~~ Somali upwelling ~~history for during~~ the last 18.5 ka has been reconstructed using biogenic silica
10 fluxes estimated from a sediment core retrieved from the western Arabian Sea. Surface winds along the east African coast
during southwest monsoon causes the Somali upwelling and therefore the intensity of this upwelling has been related to the
southwest monsoonal variability. ~~Variations in biogenic silica fluxes~~ Biogenic silica flux variation suggests periodic
weakening and strengthening of the Somali upwelling. ~~A w~~Weakened upwelling during the last glacial period (18.5-15 ka
15 BP) and strengthened upwelling during the Bølling-Allerød (15-12.9 ka BP) suggest ~~post-glacial~~ onset of the southwest
monsoon. Whereas the Younger Dryas (12.9-11.7 ka BP) is marked by reduced upwelling strength, intensification of the
Somali upwelling at the beginning of the Holocene and ~~a further~~ decline at 8 ka BP ~~have been observed~~. Increase in
upwelling strength recorded since 8 ka BP suggest ~~strengthening of the~~ southwest monsoon ~~strengthening~~ during the latter
part of the Holocene. ~~These~~ Comparison of upwelling variations ~~when compared~~ with the southwest monsoon precipitation
20 rainfall ~~is observed~~ at the beginning of Holocene. The observed shift ~~is has been~~ attributed to the variation in the southwest
monsoon strength due to the latitudinal shift of the Intertropical Convergence Zone (ITCZ) associated with changes in
moisture sources.

1 Introduction

~~The greater part~~ A large fraction of the world's population inhabits the tropical region, where climate is mainly ~~controlled~~
25 ~~driven~~ by monsoon rainfall. Understanding the causes of past ~~climatic~~ changes thus plays a critical role in deciphering past,
present and future monsoon variability. ~~The economy of~~ India, ~~which is being~~ a tropical country that contains a significant
part of the world's population, ~~its economy is largely~~ dependent ~~to a large extent~~ on the southwest monsoon (SWM) rainfall;
hence, slight changes in SWM rainfall can lead to immense societal impacts. Several attempts have been made to identify the
factors responsible for SWM rainfall variations. SWM rainfall variability is correlated with several global phenomena, such

30 as ENSO (Goswami et al., 1999; Annamalai and Liu, 2005), Atlantic Sea Surface Temperature (SST; Goswami et al., 2006; Yadav, 2016), Eurasian snow cover (Hahn and Shukla, 1976; Pant and Rupa Kumar, 1997; Bamzai and Shukla, 1999), the pre-monsoon 500 hPa ridge (Mooley et al., 1986), the Indo-Pacific warm pool (Parthasarathy et al., 1988; Parthasarathy et al., 1991), the Pacific decadal oscillation (Krishnan and Sugi, 2003), and the Atlantic multi-decadal oscillation (Krishnamurthy and Krishnamurthy, 2015). In addition to these factors that influence SWM, the Indian Ocean Warm Pool (IOWP) has been identified as the prominent source of moisture for the SWM rainfall (Ninomiya and Kobayashi, 1999; Gimeno et al., 2010). During its maxima, the IOWP extends throughout the northern Indian Ocean during the pre-monsoon period (April), and it almost reduces to half during the SWM (Izumo et al., 2008). The extent of the IOWP is primarily affected controlled by the Somali upwelling and partly by the latent heat flux increase in the Arabian Sea during the SWM season (Izumo et al., 2008). The Both Somali upwelling, as well as SWM rainfall, are caused by the SWM winds during boreal summer.

Upwelling of deep water during the SWM brings nutrients up to the photic zone, enhancing surface productivity in the western Arabian Sea. Paleoproductivity variations in the coastal regions off Somalia and Oman have been extensively studied to understand past changes in SWM-related upwelling (Sirocko et al., 1993; Naidu and Malmgren, 1996; Gupta et al., 2003; Tiwari et al., 2010). Most of these previous studies were based on the foraminifera shell abundance or its chemistry. The foraminifera production in the study area persists throughout the year with maximum production during monsoon seasons. Siliceous productivity in the study area being restricted to the southwest monsoon season can serve as a better proxy for upwelling. However, The variations in siliceous productivity in the western Arabian Sea, which have thus has direct implications for to the strength of upwelling strength during in the past, have not been understood. The siliceous productivity can be studied using the biogenic silica flux in the marine sediments. The present study thus aims to understand past variations in siliceous productivity in the Somali upwelling region, as well as paleo upwelling strength and its relationship with the southwest monsoon rainfall, using a sediment core retrieved from the western Arabian Sea (Fig.1).

The surface waters of the world ocean are mostly deficient in bioavailable silica (Hurd, 1973), which is a major nutrient for siliceous productivity. Apart from the Southern Ocean, high siliceous productivity can be observed in the major upwelling regions, where upwelled nutrient-rich water causes high primary production (Koning et al., 2001). The ocean is under saturated with respect to silica, and thus biogenic silica flux in sediments is a function of its export flux, which is controlled by its production at the surface and dissolution in the water column as well at the sediment water interface (Hurd, 1989; Broecker and Peng, 1982). Thus, a Using biogenic silica as a proxy for the study of paleo-upwelling requires understanding of its production and burial efficiency. Sediment trap studies from the western Arabian Sea indicated that biogenic silica flux mimics the SWM upwelling (Fig. 1; Nair, 2000; Haake et al., 1993). Studies of sediment trap data and surface sediments (Koning et al., 1997; Koning et al., 2001) of the Somali basin provides better estimates of the burial efficiency of biogenic silica i.e. ratio between diatom abundance in at surface to its concentration in the sediment, in of the western Arabian Sea. Only 6.8–8.7 % of diatom (biogenic silica) productivity is preserved in the sediments of the Somali basin; the rest is remineralised in the water column and at the sediment water interface (Koning et al., 2001). One of the major findings from

65 sediment traps in the Somali basin by Koning et al., (2001) is the selective preservation of upwelling-indicating diatoms in the sediments of the Somali region. This is linked to the silicification of diatom frustules; most pre- and post-upwelling produced diatoms are weakly silicified, enhancing their dissolution in the water column and leading to their low preservation in sediments. Better preservation of upwelling indicating diatoms may also be linked to the increased downward supply due to high surface production. Nutrient availability (Si:N) and concentration of dissolved iron can also affect diatom silicification that leads to variation in preservation (Hutchins and Bruland, 1998). In general, it is noted that high silicate
70 concentration along with micro-nutrient depletion leads to more silicified and faster sinking diatoms (Hutchins and Bruland, 1998), ~~which is a~~ most plausible scenario during the late phase of SWM upwelling. If burial efficiency (BE) is the primary controller of biogenic silica flux variation, ratio of low flux to low BE should have been similar to the ratio of high flux to high BE. However, in the present record using the modern high and low BE values (Koning et al., 2001), the ratio of high flux to high BE (top) is almost three times more than the ratio of low flux to low BE (bottom), thereby indicating the absence
75 of preservation effect in the biogenic silica flux.

Apart from biogenic silica production and preservation efficiency, sediment redistribution can also influence the biogenic silica flux. However, considering the sediment core location and average sedimentation rate, it is likely that the influence of sediment focusing/winnowing on the flux record is minimal. The location of the sediment core is far from the continental slope (Fig. 2) and not directly influenced by coastal currents or fluvial systems that would lead to redistribution of the
80 sediment flux. ~~The chronology of the sediment record is based on the published age model given by Tiwari et al., (2010). The model, however, shows sizeable variations in the sedimentation rate. Because the calculated sedimentation rate is a function of the selected sampling depths (which may vary), average rate of sedimentation has been considered in estimating the fluxes of biogenic silica.~~ The high surface production of biogenic silica during SWM upwelling (Fig. 1; Haake et al., 1993; Koning et al., 1997), together with the increased burial efficiency of upwelling-indicating diatoms (biogenic silica) in
85 the western Arabian Sea sediments (Koning et al., 2001), makes biogenic silica flux as a potential proxy for SWM-related upwelling in the study area.

The present study thus aims to understand past variations in siliceous productivity in the Somali upwelling region, as well as paleo-upwelling strength and its relationship with the southwest monsoon rainfall, using a sediment core retrieved from the western Arabian Sea (Fig. 2).

90 1.1 Modern Oceanography and Productivity

The surface water circulation in the western Arabian Sea is controlled by seasonal changes in atmospheric wind pattern related with annual migration of ITCZ (Wyrтки, 1973). During boreal winter, ITCZ ~~stays~~ ~~is~~ ~~located~~ south of equator and shifts to north during boreal summer. This northward shift of ITCZ during southwest monsoon (SWM, June-September) season drives the southern hemisphere eastern trade winds across equator that turns clockwise and becomes southwest winds
95 (Findlater, 1977; Fig. 3). These southwest winds ~~help to form the result in~~ Somali current along the east African coast towards north. The Somali current is generally associated with near shore upwelling and eddies such as southern gyre, great

whirl and Socotra eddy (Schott et al., 1990; Beal and Chereskin, 2003; Schott et al., 2009). These eddies induce intense upwelling which ~~pumps out the~~ brings low temperature and nutrient rich subsurface water to the surface along the east coast of Africa (Young and Kindle, 1994).

100 Productivity in the western Arabian Sea ~~reflects the~~ demonstrates seasonal changes in surface ocean characteristics (Qasim, 1977; Brock et al., 1991). More than half of the annual productivity in the western Arabian Sea occurs during southwest monsoon due to intense upwelling (Haake et al., 1993). Total flux (biogenic + dust) ~~also~~ peaks at the same time when productivity is at its maximum, indicating that SWM not only causes high productivity in the western Arabian Sea but also contributes to high dust flux (Sirocko and Lange, 1991; Haake et al., 1993). Bhushan et al., 2003 observed that ~~the~~
105 concentration of nitrate and phosphate increased at the bottom of the mixed layer at the core location, whereas, significant increase in silicate concentration occurs only at ~~the~~ depths of thermocline. During the onset of SWM upwelling, nitrate and phosphate rich waters surfaced more compared to silicate due to the upwelling of shallow waters. In presence of high nitrate and phosphate along with the micronutrients (derived by dust flux), the calcareous primary producers dominated the surface productivity. Sediment trap studies in the western Arabian Sea recorded high biogenic carbonate flux at the onset of SWM
110 upwelling (Haake et al., 1993). During the late phase of SWM upwelling, the surfacing of deeper waters increased silicate concentration in surface waters. High silicate content and depletion of micro-nutrients to sustain excessive nutrient utilization results in siliceous productivity and subsequent biogenic silica flux to the sediments (Fig. 1; Koning et al., 2001). Initially, this upwelled deep water surfaced at Somali coastal upwelling zone and transported towards the mouth of Gulf of Aden (core location) through the Socotra channel i.e. between the Socotra Island and Somalia (Young and Kindle, 1994).

115 2 Material and methods

A sediment core SS4018 was collected off the Horn of Africa (north of Socotra island), from the western Arabian Sea (13° 12.80' N, 53° 15.40' E; water depth 2830 m; core length 130 cm; Fig. 2) during FORV Sagar Sampada cruise SS-164 in 1998. Sub-sampling ~~of the core was carried out~~ at 2-cm intervals. ~~was carried out~~. The age-depth model (Fig. 4) as well as the calcareous and organic productivity proxies of this sediment core have been presented elsewhere (Tiwari et al., 2010).
120 Dry bulk density (DBD) was computed using an empirical equation based on the calcium carbonate concentrations (Clemens et al., 1987). The flux rate was estimated using an average sedimentation rate computed based on the age-depth model given by Tiwari et al., (2010). ~~However, the published ages of the individual samples were considered~~. The sedimentation rate at the core site as given by Tiwari et al., (2010) is variable, the lowest being 3.5 and highest at 22.7 cm.ka-1. Since the age model depends on the sample selection criteria and may change according to depth of age control points, an average
125 sedimentation rate was computed for the entire core. ~~The actual sedimentation at SS4018 sediment core site possibly much more complex than the age model presented in Tiwari et al (2010), so we computed the average sedimentation rates for the whole core. This average age model is only used to understand large-scale changes in biogenic silica concentration.~~

The biogenic silica concentration was measured in each sample using the method described by Carter and Colman (1994). Dried homogenized samples weighing 50 mg were placed in centrifuge tubes. Five milliliters of 10 % H₂O₂ was added to each sample at room temperature, and the samples were stored for 2 hours to remove organic matter. Five milliliters of 1N HCl was added to each tube. After acid treatment, ~~sample was washed thoroughly with 20 ml of distilled water~~ ~~was added~~, and the samples were centrifuged for 15 minutes. ~~Sample tubes~~ were kept in an oven for drying after removal of the supernatant. Thirty milliliters of 2 M Na₂CO₃ was added to each sample ~~tube~~, and ~~the tubes~~ were kept in a shaker bath at 95^o C for 5 hours. After 5 hours, the samples were centrifuged for 5 minutes, and 3 ml of hot supernatant was pipetted out of each sample and added to exactly 30 ml of distilled water in pre-cleaned sample tubes. The solution was acidified by adding 0.9 ml of concentrated HNO₃. Sample tubes were sealed after effervescence. Silicon and Aluminium concentrations were measured in these samples using ICP-AES (Jobin-Yvon, Model 38S ~~at Physical Research Laboratory, Ahmedabad~~). The silicon concentrations were then corrected for clay mineral dissolution by using the formula given by Carter and Colman (1994) (Eqn 1):

$$\Delta Si = Si - (Al * 1.93) \quad (1)$$

Where, ΔSi is the corrected silicon concentration, Si and Al are the measured concentrations of silicon and aluminium in the sample, and 1.93 is the Si to Al ratio in smectite. Smectite is an abundant clay mineral in the northern Arabian Sea (Sirocko et al., 1991). Biogenic silica concentrations were calculated using the formula given below (Eqn 2):

$$Biogenic\ silica = \Delta Si * K \quad (2)$$

Where, K is a constant that equals 2.4, which accounts for the ~10 % water content in biogenic silica (Mortlock and Frolich, 1989). Overall, the error associated with the biogenic silica measurement is less than 5 % based on repeat measurements. The biogenic silica flux is calculated by multiplying the biogenic silica fraction by the sedimentation rate (SR) and the dry bulk density (DBD) (Eqn 3),

$$B.Si.flux(g.m^{-2}.y^{-1}) = B.Si * SR(m.y^{-1}) * DBD(g.m^{-3}) \quad (3)$$

The uncertainties associated with the biogenic silica concentration (B.Si) is estimated from the error in Aluminium and Silicon concentration based on ~~repeat~~ measurements of ~~repeat and~~ standard material. The maximum error in biogenic silica concentration is within 5%. Dry bulk density (DBD) is calculated from CaCO₃ concentration using an empirical equation suggested by Clemens et al., 1987. The standard uncertainty in DBD calculation is 0.091 g/cm³. The uncertainty in average sedimentation rate (SR) is 0.12 cm/ky. ~~Finally~~ The uncertainty associated with biogenic silica flux (B.Si flux) is propagated using the ~~below~~ equation below,

$$\sigma_{B.Si\ flux} = B.Si\ flux * \sqrt{\{(\sigma_{B.Si}/B.Si)^2 + (\sigma_{DBD}/DBD)^2 + (\sigma_{SR}/SR)^2\}} \quad (4)$$

Where, prefix “ σ ” stands for uncertainty. Uncertainty in biogenic silica concentration is below 5%. But the uncertainty in flux are up to 15%. This increase in uncertainty is due to the high standard error associated with empirical derivation of Dry bulk density.

160 **3Results**

Biogenic silica concentrations varied from 3 % to 15.2 % during the last 18.5 ka (Table 1), with the lowest concentrations (3–4 %) being observed at the bottom of the core between 18.5 ka and 16 ka BP. Subsequently, it increased continuously up to 13.5 ka BP (9 %) and decreased to 6.5 % at 13 ka BP. No significant variations in biogenic silica concentrations were observed between 13 ka and 11.3 ka BP. After 11.3 ka BP, biogenic silica increased from 6.5 % to 10.5 % and remained stable for a period of almost 1000 years until 10 ka BP. The biogenic silica concentration increased from 10.5 % to 12 % during 10–9.5 ka BP, and it decreased subsequently to 8.5 % at 8 ka BP. From 8 ka to 7.3 ka BP, it changed from 8.5 % to 9.5 % with a peak at 7.7 ka BP (13.3 %), further it remained stable until 6 ka BP. From 6 ka to 5 ka BP, the biogenic silica concentration increased from 10 % to 12.5 % and then decreased at 4 ka BP. After 4 ka BP, a continuous increase to a maximum value of 15 % biogenic silica at 1.5 ka BP and a subsequent decrease to 12 % at 1 ka BP were observed. Biogenic silica concentrations showed no variations during the last 1 ka BP.

The variations in biogenic silica fluxes show an overall increasing trend from 18.5 ka to present (Fig.5). The fluxes varied between 1.4 to 6.8 $\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$. The minimum flux ($<2\text{ g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$) was observed at the bottom of the sediment core, i.e., between 18.5 and 16 ka BP, and the maximum flux at 2 ka BP. Five distinct peaks in biogenic silica flux during the last 18.5 ka BP were observed between 14–13 ka, 11–10 ka, 10–9 ka, 6–4 ka and 2.5–1 ka BP. Uncertainties in biogenic silica concentrations and fluxes are below 5 % and up to 15 %, respectively. The increase in uncertainty in the flux is due to the high standard error associated with the empirical derivation of the dry bulk densities.

3 Result and Discussion**3.1 Biogenic silica flux as an upwelling proxy**

The surface waters of the world ocean are mostly deficient in bioavailable silica (Hurd, 1973), which is a major nutrient for siliceous productivity. Apart from the Southern Ocean, high siliceous productivity can be observed in the major upwelling regions, where upwelled nutrient rich water causes high primary production (Koning et al., 2001). The ocean is under saturated with respect to silica, and thus biogenic silica flux in sediments is a function of its export flux, which is controlled by its production at the surface and dissolution in the water column as well at the sediment water interface (Hurd, 1989; Broecker and Peng, 1982). Thus, using biogenic silica as a proxy for the study of paleo upwelling requires understanding of its production and burial efficiency. Sediment trap studies from the western Arabian Sea indicated that biogenic silica flux mimics the SWM upwelling (Fig. 3; Nair, 2000; Haake et al., 1993). Studies of sediment trap data and surface sediments (Koning et al., 1997; Koning et al., 2001) of the Somali basin provides better estimates of the burial efficiency of biogenic silica i.e. ratio between diatom abundance in at surface to its concentration in the sediment, in of the western Arabian Sea. Only 6.8–8.7 % of diatom (biogenic silica) productivity is preserved in the sediments of the Somali basin; the rest is remineralised in the water column and at the sediment water interface (Koning et al., 2001). One of the major findings from

195 sediment traps in the Somali basin by Koning et al., (2001) is the selective preservation of upwelling-indicating diatoms in the sediments of the Somali region. This is linked to the silicification of diatom frustules; most pre- and post-upwelling produced diatoms are weakly silicified, enhancing their dissolution in the water column and leading to their low preservation in sediments. Better preservation of upwelling-indicating diatoms may also be linked to the increased downward supply due to high surface production. Nutrient availability (Si:N) and concentration of dissolved iron can also affect diatom silicification that leads to variation in preservation (Hutchins and Bruland, 1998). In general, it is noted that high silicate concentration along with micro-nutrient depletion leads to more silicified and faster-sinking diatoms (Hutchins and Bruland, 1998), which is a most plausible scenario during the late phase of SWM upwelling. If burial efficiency (BE) is the primary controller of biogenic silica flux variation, ratio of low flux to low BE should have been similar to the ratio of high flux to high BE. However, in the present record using the modern high and low BE values (Koning et al., 2001), the ratio of high flux to high BE (top) is almost three times more than the ratio of low flux to low BE (bottom), thereby indicating the absence of preservation effect in the biogenic silica flux.

200 Apart from biogenic silica production and preservation efficiency, sediment redistribution can also influence the biogenic silica flux. However, considering the sediment core location and average sedimentation rate, it is likely that the influence of sediment focusing/winnowing on the flux record is minimal. The location of the sediment core is far from the continental slope (Fig. 1) and not directly influenced by coastal currents or fluvial systems that would lead to redistribution of the sediment flux. The chronology of the sediment record is based on the published age model given by Tiwari et al., (2010). The model, however, shows sizeable variations in the sedimentation rate. Because the calculated sedimentation rate is a function of the selected sampling depths (which may vary), average rate of sedimentation has been considered in estimating the fluxes of biogenic silica. The high surface production of biogenic silica during SWM upwelling (Fig. 3; Haake et al., 1993; Koning et al., 1997), together with the increased burial efficiency of upwelling-indicating diatoms (biogenic silica) in the western Arabian Sea sediments (Koning et al., 2001), makes biogenic silica flux as a potential proxy for SWM related upwelling in the study area.

3.1 Biogenic silica flux vs SST

215 Similar to biogenic silica flux, the paleo-SST ~~reconstructions~~ can serve as proxy for upwelling, as upwelling increases siliceous productivity with reduction in SST. However, the inverse relation between siliceous productivity and SST is valid only during the SWM season and not even on annual scale. ~~Mostly the SST reconstructions are based on biomarker (TEX₈₆ and U^K₃₇; Brassell et al., 1986; Schouten et al., 2002) or from planktonic foraminifera shell chemistry ($\delta^{18}\text{O}$ and Mg/Ca; Emiliani, 1955; Chave, 1954). However, the annual mean signal of SST or its reconstruction on seasonal scale depends on~~ the signature of the productivity proxies for the region. The $\delta^{18}\text{O}$ value of foraminifera shell is not only dependent on the SST, but also gets modulated with the preservation of shell, carbonate ion concentration, salinity and $\delta^{18}\text{O}$ (ice volume) of the original seawater (Lea, 2003), which makes the reconstruction rather complex. The SST reconstruction based on Mg/Ca ratio is affected by species dependency and dissolution (Lea, 2003). In most of the cases, it is foraminifera based SST

225 which preserves better signature of annual mean signal due to the foraminifera production throughout the year (Conan et al.,
 2000; Dahl and Oppo, 2006), but the possibility of preserving seasonal signal depends on local hydrography and
 productivity. While, the biomarker based SST records may preserve seasonal signal, because they are mainly produced
 during monsoon season (Huguet et al., 2006). However, there are other limitations with the biomarker based SST records.
 Major limitation for the application of alkenone $U^{K^2}_{37}$ in low latitude regions is that it saturates around 28°C (Prah
 and Wakeham, 1987). While, the TEX_{86} based SST are valid in the range of 5 to 30°C (Kim et al., 2008), with better estimates
 230 above 15°C (Kim et al., 2010). However, there are numerous evidence suggesting TEX_{86} records sub-surface temperature
 rather than SST (Hertzberg et al., 2016; Lee et al., 2008; Lopes dos Santos et al., 2010; Seki et al., 2012; Wuchter et al., 2006).
 The nutrient availability and variation in productivity may also influence the TEX_{86} temperature (Hertzberg et al., 2016).
 Comparing the SST reconstructed using $U^{K^2}_{37}$ and TEX_{86} from a sediment core in the western Arabian Sea, Huguet et al.,
 2006 suggested that the $U^{K^2}_{37}$ SST are in phase with northern hemisphere dynamics during NE monsoon, while, TEX_{86} SST
 235 are controlled by SW monsoon. All SST proxies tend to record annual mean signal with varying fraction of seasonal signal.
 Glacial boundary conditions have strong influence on the annual mean SST in the Arabian Sea irrespective of monsoon
 upwelling (Broccoli, 2000; Dahl and Oppo, 2006). While, the biogenic silica flux has been controlled by the SWM
 upwelling signal due to the fact that since it is produced during the southwest monsoon season and preserves upwelling
 signal. Hence, the biogenic silica flux can be identified as a better proxy than SST to understand SWM upwelling in the
 240 study area. The variations in biogenic silica fluxes during the 18.5 ka are shown in Fig. 5 and the data is presented in table 1.
 Comparison of biogenic silica flux with other paleo-SST records (Anand et al., 2008; Sahar et al., 2007; Huguet et al., 2006)
 from nearby locations are shown in figure 6 and 8. There is No definitive relation can be observed between biogenic silica
 flux and SST records on temporal scale. Also, The SST using different proxies show inconsistent changes in the studied time
 span, the TEX_{86} SST is always higher than Mg/Ca SST (Fig. 6&8). A general observation is that both biogenic silica flux
 245 and SST were low during 18.5 to 15 ka BP, later showing an anti-correlation (Fig. 6). The anti-correlation is strong between
 biogenic silica flux and TEX_{86} SST during 15 to 11.7 ka BP (Fig. 6). However, during the last 11.7 ka the Mg/Ca based SST
 shows a strong anti-correlation with biogenic silica flux record, indicating variation in influence of on the seasonal signal
 for of different proxies (Fig. 8). Temporal relation between biogenic silica flux and SST is discussed in the following
 section.

250 3.2 Somali upwelling strength versus southwest monsoon rainfall

The western Arabian Sea SSTs during SWM are directly related to upwelling strength (enhanced upwelling results in
 lower SSTs and vice versa; Fig. 1). Previous studies have shown that the northern Indian ocean and in particular the Arabian
 Sea to be an important source of moisture for SWM rainfall over India (Ninomiya and Kobayashi, 1999; Gimeno et al.,
 2010). There are several various kind of relationship observed between SST, moisture and SWM rainfall. First order relation
 255 would be positive, i.e. reduced SWM winds would cause reduced upwelling as well as reduced rainfall and vice versa.
 However, the relation between Arabian Sea SST and SWM rainfall is complicated due to the fact that SST modulates the

moisture availability as well as the meridional temperature gradient (Levine and Turner, 2012). Modelling study by Shukla (1975) showed that the cold Arabian Sea SST during SWM tend to reduce the SWM rainfall through reduced moisture transport. However, Webster et al (1999) and Clark et al (2000) showed that the SWM rainfall has stronger connection with winter and spring SSTs rather than summer, and suggested a delayed influence of SST on rainfall. A modelling study by Arpe et al (1998) demonstrated that warmer northern Indian Ocean leads to increased SWM rainfall over India through enhanced evaporation and moisture supply, while indicating the strong influence of pacific SST anomalies on monsoon. It was also suggested that Arabian Sea SST modulates the impact of ENSO on monsoon precipitation (Arpe et al., 1998; Lavine and Turner, 2012). An observational study by Vecchi and Harrison (2004) detected a strong positive correlation between western Arabian SSTs and SWM rainfall over the Western Ghats Mountains in India from 1982 to 2001. Overall, it has been suggested that any isolated cooling of the Arabian Sea will reduce SWM rainfall through reduced moisture supply, in absence of other large-scale forcing (Lavine and Turner, 2012). An observational and modelling study by Izumo et al (2008) signals ~~the~~ causes for the variations in western Arabian Sea SSTs and ~~their~~ its influence on SWM rainfall over the Western Ghats. According to Izumo et al (2008), increased Somali upwelling during the late spring reduces the westward extension of the IOWP during summer, which decreases moisture availability to the air mass that delivers rainfall to the western part of the Indian sub-continent. Though, the upwelling-rainfall connection is not fully understood and difficult to model, the observations suggest an anti-correlation between Somali upwelling (western Arabian Sea SST during SWM) and SWM rainfall. Both Somali upwelling and SWM rainfall ~~were~~ being caused by southwest monsoonal winds during SWM season, hence their anti-correlation indicates a negative feed-back within the system.

Did this anti-correlation exist between the Somali upwelling and SWM rainfall in the geological past? To answer this question, we need to investigate the record of paleo-upwelling in the Somali region and paleo-rainfall in the western part of India and adjoining areas. There is no continuous terrestrial record of paleo-rainfall covering the last 18.5 ka from the Western Ghats, but there are several paleoclimatic records based on marine sediment cores from the eastern Arabian Sea. The biogenic silica flux temporal variability is compared (Fig. 7&10) with paleo-rainfall record ($\delta^{18}\text{O}_w\text{IVF}$ by Anand et al., 2008)) from the eastern Arabian Sea and a speleothem record from Oman (Fleitmann et al., 2003). The $\delta^{18}\text{O}_w$ IVF is the Ice Volume Free (IVF) oxygen isotopic composition of seawater based on the $\delta^{18}\text{O}$ of *G.ruber*. Anand et al. (2008) showed that the reconstructed $\delta^{18}\text{O}_w$ IVF during the last 19 ka from a sediment core (SK-17) in the eastern Arabian Sea was mainly controlled by the SWM rainfall in the Western Ghats. The Qunf speleothem record from Oman (Fleitmann et al., 2003) had been widely used as an indicator for SWM variation. The location of Qunf speleothem is very close to the present study area i.e. downwind side to the present study area during SWM season. If the SWM was the reason for the rainfall in southern Oman, then western Arabian Sea must be the ~~source~~ source of moisture. Though, there are no observational study on the relation between upwelling strength and rainfall in Oman, a comparison is made to give a preliminary assessment. Since records are from different regions and have irregular temporal resolution, only long-term trends have been examined.

3.2.1 Last Glacial Period (18.5–15 ka BP)

290 The biogenic silica flux record does not show any distinct variation between the previously identified Heinrich event 1 and the Last Glacial Maximum (LGM; Clark et al., 2009), the period between 18.5 and 15 ka thus is considered here as the Last Glacial Period (LGP). Both biogenic silica concentrations (3–5 %) and fluxes (~2 g.m-2.y-1) were lowest during the LGP (Fig. 5), similar to earlier findings of low productivity during glacial periods from the western Arabian Sea (Burckle, 1989; Sirocko et al., 1991; Sirocko et al., 2000; Ivanochko et al., 2005; Tiwari et al., 2010). Based on the modern pattern of
295 biogenic silica productivity and its burial efficiency in the western Arabian Sea, the observed low fluxes of biogenic silica indicate that the Somali upwelling was very weak during the LGP. However, the lowest SSTs recorded in the last 18.5 ka in the Somali basin during the LGP (Huguet et al., 2006; Saher et al., 2007; Anand et al., 2008) are related to basin wide cooling and not connected with upwelling strength (Fig. 6). Dahl and Oppo, 2006 observed a reduction in Arabian Sea SST of 2–4° C during LGP. Thus, it is unlikely that the IOWP (SST>28° C) formed during LGP. In absence of IOWP, any
300 relation between the Somali upwelling and rainfall is unexpected. Paleo-rainfall record from the eastern Arabian Sea shows high $\delta^{18}\text{O}_w$ IVF values indicative of reduced freshwater flux and rainfall during this period (Fig. 7). Based on weak upwelling in the western Arabian Sea and reduced fresh water influx to the eastern Arabian Sea, it can be concluded that the SWM was weak/absent during the LGP.

3.2.2 Deglacial Period (15–11.7 ka BP)

305 The Deglacial Period (DP) is a connecting phase between two entirely different climatic periods, the LGP and the Holocene. The DP is basically a composite of two millennial scale events between 15–12.9 ka and 12.9–11.7 ka BP. This period occupied by these two events nearly coincides with well-known climatic events, specifically the Bølling-Allerød (B/A) and the Younger Dryas (YD) event. The beginning of the B/A is marked by an abrupt increase in biogenic silica flux (Fig. 6a) is attributed to the effect of northern limit of southwest monsoon, attained at the study site with subsequent increase in Somali
310 upwelling strength. This is further supported by Zr/Hf in two independent sediment cores near our core site, which shows an increasing flux of wind borne dust from the Horn of Africa (an indicator of the SWM) at the onset of the B/A (Sirocko et al., 2000; Isaji et al., 2015). The reduction in TEX₈₆ SST during the B/A in the Somali basin (Huguet et al., 2006) also suggests increased upwelling (Fig. 5), however the Mg/Ca SST does not show this change (Fig. 6). The inconsistency between the two SST records (TEX₈₆ and Mg/Ca) could be related to the control on seasonal production of the proxy material.
315 The $\delta^{18}\text{O}_w$ IVF record from core SK-17 (Anand et al., 2008) shows depleted values, indicating higher influx of fresh water from the Western Ghats caused by high SWM rainfall, during the B/A (Fig. 7c). The positive correlation between Somali upwelling (high biogenic silica flux) and SWM rainfall in the Western Ghats (high fresh water influx to the eastern Arabian Sea) during the B/A contrasts with the present-day scenario as observed by Vecchi and Harrison (2004). Presently, the moisture source for the SWM rainfall is the Arabian Sea and the central Indian Ocean (IOWP), which is affected by SWM
320 upwelling (Izumo et al., 2008). If the central Indian Ocean ~~were~~ was the source of moisture for SWM rainfall during the

B/A, then the observed co-variation is possible. Thus, it is proposed that the moisture source for SWM rainfall over Western Ghats during the B/A event was different from the modern source. The other possibility, that rainfall in south-western India was enhanced due to a strong NE monsoon during the B/A is unlikely because the siliceous productivity in the western Arabian Sea related to the NE monsoon has not been reported (Koning et al., 1997; Ramaswamy and Gaye, 2006). In contrast to the B/A, the upwelling in the western Arabian Sea was weak during the YD, as revealed by the low biogenic silica fluxes and high SSTs (Fig. 6; Hugué et al., 2006). This is in agreement with the previous studies from Arabian Sea which showed decreased productivity during YD due to reduction in SWM (Altabet et al., 2002; Ivanochko et al., 2005). Furthermore, the high $\delta^{18}\text{O}_{\text{wIVF}}$ in the eastern Arabian Sea (Anand et al., 2008) were caused by low freshwater influx points to weak SWM rainfall (Fig.7).

330 3.2.3 Holocene (11.7–0 ka BP)

The beginning of the Holocene is marked by an abrupt increase in biogenic silica flux (Fig. 8a). This sudden increase in biogenic silica fluxes between 11.7 ka and 9 ka BP could have been caused by be due to the intensification of the SWM (extended season) resulting in northward shift of the ITCZ, following the peak in Northern Hemisphere solar insolation (Fleitmann et al., 2007). Somali basin SST records (Saher et al., 2007; Hugué et al., 2006; Anand et al., 2008) also shows a marginal decrease at the onset of the Holocene, but not up to the levels seen during the B/A (Fig.8). The TEX_{86} SST shows more variation than Mg/Ca at the beginning of Holocene, however, the Mg/Ca SST shows a mirror image to the biogenic silica flux pattern during Holocene, indicating dominance of seasonal signal in Mg/Ca SST during this period (Fig. 8b). Based on the stable isotopic composition of organic carbon and nitrogen in the 4018 core, Tiwari et al (2010) ~~too~~ also suggested an increase in productivity during Holocene and attributed it to the strengthening of Somali upwelling. The synchronous changes in the biogenic silica flux with biogenic silica/carbonate ratio (Fig. 9) indicates a change in dominant plankton community (carbonaceous to siliceous) due to increased upwelling, as suggested by Tiwari et al. (2010). The $\delta^{18}\text{O}_{\text{wIVF}}$ (Anand et al., 2008) display values like those of the YD during the early Holocene (Fig. 10c), indicating reduced rainfall (lower fresh water influx) over the Western Ghats. This anti-correlation between Somali upwelling and SWM rainfall over south-western India during the early Holocene (11.7 ka to 9 ka BP), marks the establishment of the modern-day climate system. The increased Somali upwelling in the western Arabian Sea during the early Holocene (11.7 to 8 ka BP; Fig. 10a) might could have reduced the IOWP expanse during the SWM season, thereby resulting in lower moisture availability and subsequent reduced rainfall over the Western Ghats. Oman speleothem record also shows decreased precipitation during early Holocene (Fig. 10b) and supports this interpretation of low moisture availability.

At ~8 ka BP (Fig. 10a), Somali upwelling strength decreased compared to the early Holocene, but persisted above YD and B/A levels, indicating the presence of ~~the~~ SWM with reduced wind strengths relative to the early Holocene. The SST record also shows an increased value at ~8 ka BP (Fig. 8b) indicating reduction in Somali upwelling. This reduction in upwelling at 8 ka BP can support the westward extension of the IOWP during the SWM season, thereby increasing both moisture availability over the Arabian Sea and rainfall over the Western Ghats. The $\delta^{18}\text{O}_{\text{wIVF}}$ value decreased at 8 ka BP, indicating

an increase in fresh water influx from the Western Ghats (Anand et al., 2008) due to increased SWM rainfall (Fig. 10c).
355 Oman speleothem record too shows decreased $\delta^{18}\text{O}$ value at 8 ka BP (Fig. 10b) suggesting increased SWM rainfall.
Somali upwelling had a gradual increase during the last 8 ka with minor positive excursion at around 5 and 2 ka BP (Fig. 10a). The increase in SWM induced Somali upwelling during the last 8 ka contrasts the idea that SWM followed the northern hemisphere insolation during Holocene (Gupta et al., 2003; Fleitmann et al., 2003 and references therein). However, this interpretation is well in agreement with other studies from the Arabian Sea which shows that SWM did
360 slightly increase during Holocene (Agnihotri et al., 2003; Tiwari et al., 2010). These short-term increase in Somali upwelling at 5 and 2 ka BP can also be observed with reduction in the Mg/Ca SST record (Fig. 8b). Oman speleothem record shows an increase in $\delta^{18}\text{O}$ during the last 8 ka suggesting reduction in SWM rainfall (Fig. 10b). The hiatus in Oman speleothem record at 2 ka BP coincides with the strengthened Somali upwelling, however, it is difficult to explain (Fig. 10a & b). The SK-17 record (Anand et al., 2008) shows slight increase in the $\delta^{18}\text{O}_{\text{wIVF}}$ of surface waters during the last 8 ka (Fig. 10c) indicating
365 reduction in SWM rainfall. The opposite trend in upwelling and rainfall record during the last 8 ka indicates the negative impact of Somali upwelling on SWM rainfall through changing the area of IOWP and moisture availability. However, the short-term variations in upwelling are not observed in the eastern Arabian Sea rainfall record.

It is observed that baring the beginning, the Somali upwelling had a negative impact on southwest monsoon rainfall over south-western India and Oman during Holocene. ~~The Somali upwelling possibly had a negative impact on southwest monsoon rainfall over south western India and Oman throughout the Holocene.~~
370 This finding would have implications in context of the modelling study by deCastro et al. (2016), which shows that Somali upwelling would increase during the twenty-first century.

4 Conclusions

The present study demonstrates the use of biogenic silica flux as a proxy for the temporal variations in the strength of the
375 Somali upwelling during the last 18.5 ka. Some of the salient findings of the present study are summarized below:

1. The Somali upwelling was weak during the LGP coeval with the weak southwest monsoon.
2. The post-glacial onset of the southwest monsoon was marked by an increase in the strength of the Somali upwelling at 15 ka BP, with the eastern Arabian Sea records showing increased southwest monsoon rainfall.
3. The Somali upwelling was weak between 12.9 and 11.7 ka BP, indicating another phase of weak southwest
380 monsoon like that of the LGP. Overall, records of the Somali upwelling and southwest monsoon rainfall exhibit positive correlations between 18.5 and 11.7 ka BP.
4. A Shift from positive to negative correlation between the Somali upwelling strength and southwest monsoon rainfall occurred at 11.7 ka BP at the beginning of the Holocene, which marks the establishment of modern day climate system.

- 385 5. Enhanced Somali upwelling during the last 11.7 ka BP, except for a decline at 8 ka BP, had a negative impact on southwest monsoon rainfall.
6. Both, latitudinal shifts in the Intertropical Convergence Zone (ITCZ) and changes in moisture source region act as causative factors for the reversal in the relationship between upwelling and southwest monsoon rainfall.
7. Future observational and modelling studies on southwest monsoon rainfall reconstruction and prediction should
390 incorporate variations in the moisture source region.

Acknowledgments

Balaji D. is thankful to the Council of Scientific and Industrial Research (CSIR) of India for providing support through a CSIR-NET-Senior Research Fellowship. Ravi Bhushan thanks Director of the PRL for research grant support. We thank Dr. Alpa Sridhar for critical comments and suggestions on the manuscript. We thank Professor Eduardo Zorita (editor) and three
395 ~~the two~~ anonymous reviewers for their constructive comments which helped us in improving the manuscript.

References

- Agnihotri R, Bhattacharya SK, Sarin MM, Somayajulu BLK.: Changes in the surface productivity and subsurface denitrification during the Holocene: a multiproxy study from the eastern Arabian Sea. *The Holocene* 13: 701–713, 2003.
- Altabet, M. A., Higginson, M. J., & Murray, D. W.: The effect of millennial-scale changes in Arabian Sea denitrification on
400 atmospheric CO₂. *Nature*, 415(6868), 159, 2002.
- Anand, P., Kroon, D., Singh, A. D., Ganeshram, R. S., Ganssen, G., and Elderfield, H.: Coupled sea surface temperature–seawater δ¹⁸O reconstructions in the Arabian Sea at the millennial scale for the last 35 ka, *Paleoceanography*, 23, 2008.
- Annamalai, H., and Liu, P.: Response of the Asian summer monsoon to changes in El Niño properties, *Quarterly Journal of the Royal Meteorological Society*, 131, 805-831, 2005.
- 405 Arpe K, Duřmenil L, Giorgetta MA.: Variability of the Indian monsoon in the ECHAM3 model: sensitivity to sea surface temperature, soil moisture, and the stratospheric quasi-biennial oscillation. *J Clim* 11:1837–1858, 1998.
- Beal, L. M., and Chereskin, T. K.: The volume transport of the Somali Current during the 1995 southwest monsoon, *Deep Sea Research Part II: Topical Studies in Oceanography*, 50, 2077-2089, 2003.
- Berrisford, P., Kållberg, P., Kobayashi, S., Dee, D., Uppala, S., Simmons, A., Poli, P., and Sato, H.: Atmospheric
410 conservation properties in ERA-Interim, *Quarterly Journal of the Royal Meteorological Society*, 137, 1381-1399, 2011.
- Bhushan, R., Dutta, K., Mulson, S., Povinec, P. P., & Somayajulu, B. L. K.: Distribution of natural and man-made radionuclides during the reoccupation of GEOSECS stations 413 and 416 in the Arabian Sea: temporal changes. *Deep Sea Research Part II: Topical Studies in Oceanography*, 50(17-21), 2777-2784, 2003.

- Brassell, S. C., G. Eglinton, I. T. Marlowe, U. Pflaumann, and M. Sarnthein.: Molecular stratigraphy: A new tool for climatic
415 assessment, *Nature*, 320, 129–133, 1986.
- Broccoli, A. J.: Tropical cooling at the Last Glacial Maximum: An atmosphere–mixed layer ocean model simulation. *Journal
of Climate*, 13(5), 951-976, 2000.
- Burckle, L. H.: Distribution of diatoms in sediments of the northern Indian Ocean: Relationship to physical oceanography,
Marine Micropaleontology, 15, 53-65, 1989.
- 420 Carter, S. J., and Colman, S. M.: Biogenic silica in Lake Baikal sediments: results from 1990–1992 American cores, *Journal
of Great Lakes Research*, 20, 751-760, 1994.
- Chave, K. E.: Aspects of the biogeochemistry of magnesium. 1. Calcareous marine organisms, *J. Geol.*, 62, 266–283, 1954.
- Clark CO, Cole JE, Webster PJ.: Indian Ocean SST and Indian Summer Rainfall: predictive relationships and their decadal
variability. *J Clim* 13:2503–2519, 2000.
- 425 Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica, J.X., Hostetler, S.W. and McCabe,
A.M.: The last glacial maximum. *Science*, 325(5941), pp.710-714, 2009.
- Clemens, S. C., Prell, W. L., and Howard, W. R.: Retrospective dry bulk density estimates from southeast Indian Ocean
sediments—comparison of water loss and chloride-ion methods, *Marine geology*, 76, 57-69, 1987.
- Conan, S. H., & Brummer, G. J. A.: Fluxes of planktic foraminifera in response to monsoonal upwelling on the Somalia
430 Basin margin. *Deep Sea Research Part II: Topical Studies in Oceanography*, 47(9-11), 2207-2227, 2000.
- Dahl, K. A., and Oppo, D. W.: Sea surface temperature pattern reconstructions in the Arabian Sea, *Paleoceanography*, 21,
2006.
- Emiliani, C.: Pleistocene temperatures, *J. Geol*, 63, 538–578, 1955.
- Findlater, J.: Observational aspects of the low-level cross-equatorial jet stream of the western Indian Ocean, in: *Monsoon
435 Dynamics*, Springer, 1251-1262, 1978.
- Fleitmann, D., Burns, S. J., Mudelsee, M., Neff, U., Kramers, J., Mangini, A., & Matter, A.: Holocene forcing of the Indian
monsoon recorded in a stalagmite from southern Oman. *Science*, 300(5626), 1737-1739, 2003.
- Fleitmann, D., Burns, S. J., Mangini, A., Mudelsee, M., Kramers, J., Villa, I., Neff, U., Al-Subbary, A. A., Buettner, A., and
Hippler, D.: Holocene ITCZ and Indian monsoon dynamics recorded in stalagmites from Oman and Yemen (Socotra),
440 *Quaternary Science Reviews*, 26, 170-188, 2007.
- Gimeno L, Drumond A, Nieto R, Trigo RM, Stohl A.: On the origin of continental precipitation. *Geophys Res Lett*
37:L13804, 2010.
- Goswami, B., Krishnamurthy, V., and Annmalai, H.: A broad-scale circulation index for the interannual variability of the
Indian summer monsoon, *Quarterly Journal of the Royal Meteorological Society*, 125, 611-633, 1999.
- 445 Goswami, B., Madhusoodanan, M., Neema, C., and Sengupta, D.: A physical mechanism for North Atlantic SST influence
on the Indian summer monsoon, *Geophysical Research Letters*, 33, 2006.

- Govil, P., and Naidu, P. D.: Evaporation-precipitation changes in the eastern Arabian Sea for the last 68 ka: Implications on monsoon variability, *Paleoceanography*, 25, 2010.
- 450 Gupta, A. K., Anderson, D. M., and Overpeck, J. T.: Abrupt changes in the Asian southwest monsoon during the Holocene and their links to the North Atlantic Ocean, *Nature*, 421, 354-357, 2003.
- Haake, B., Ittekkot, V., Rixen, T., Ramaswamy, V., Nair, R., and Curry, W.: Seasonality and interannual variability of particle fluxes to the deep Arabian Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 40, 1323-1344, 1993.
- Hahn, D. G., and Shukla, J.: An apparent relationship between Eurasian snow cover and Indian monsoon rainfall, *Journal of the Atmospheric Sciences*, 33, 2461-2462, 1976.
- 455 Hertzberg, J. E., Schmidt, M. W., Bianchi, T. S., Smith, R. W., Shields, M. R., & Marcantonio, F.: Comparison of eastern tropical Pacific TEX₈₆ and Globigerinoides ruber Mg/Ca derived sea surface temperatures: Insights from the Holocene and Last Glacial Maximum. *Earth and Planetary Science Letters*, 434, 320-332, 2016.
- Huguet, C., Kim, J. H., Sinninghe-Damsté, J. S., and Schouten, S.: Reconstruction of sea surface temperature variations in the Arabian Sea over the last 23 kyr using organic proxies (TEX₈₆ and U_{37K'}), *Paleoceanography*, 21, 2006.
- 460 Hurd, D. C.: Interactions of biogenic opal, sediment and seawater in the Central Equatorial Pacific, *Geochimica et Cosmochimica Acta*, 37, 2257-2266, 1973.
- Isaji, Y., Kawahata, H., Ohkouchi, N., Ogawa, N. O., Murayama, M., Inoue, K., and Tamaki, K.: Varying responses to Indian monsoons during the past 220 kyr recorded in deep-sea sediments in inner and outer regions of the Gulf of Aden, *Journal of Geophysical Research: Oceans*, 120, 7253-7270, 2015.
- 465 Ivanochko, T. S., Ganeshram, R. S., Brummer, G.-J. A., Ganssen, G., Jung, S. J., Moreton, S. G., and Kroon, D.: Variations in tropical convection as an amplifier of global climate change at the millennial scale, *Earth and Planetary Science Letters*, 235, 302-314, 2005.
- Izumo, T., Montégut, C. B., Luo, J.-J., Behera, S. K., Masson, S., and Yamagata, T.: The role of the western Arabian Sea upwelling in Indian monsoon rainfall variability, *Journal of Climate*, 21, 5603-5623, 2008.
- 470 Kim, J.-H., Schouten, S., Hopmans, E., Donner, B., Sinninghe-Damsté, J.S.: Global sediment core-top calibration of the TEX₈₆ paleothermometer in the ocean. *Geochim. Cosmochim. Acta* 72, 1154–1173, 2008.
- Kim, J.-H., van der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., Koc, N., Hopmans, E., Sinninghe-Damsté, J.S.: New indices and calibrations derived from the distribution of crenarchaeal isoprenoid tetraether lipids: implications for past sea surface temperature reconstructions. *Geochim. Cosmochim. Acta* 74, 4639–4654, 2010.
- 475 Koning, E., Brummer, G.-J., Van Raaphorst, W., Van Bennekom, J., Helder, W., and Van Iperen, J.: Settling, dissolution and burial of biogenic silica in the sediments off Somalia (northwestern Indian Ocean), *Deep Sea Research Part II: Topical Studies in Oceanography*, 44, 1341-1360, 1997.
- Koning, E., Van Iperen, J., Van Raaphorst, W., Helder, W., Brummer, G.-J., and Van Weering, T.: Selective preservation of upwelling-indicating diatoms in sediments off Somalia, NW Indian Ocean, *Deep Sea Research Part I: Oceanographic*
- 480 *Research Papers*, 48, 2473-2495, 2001.

- Krishnamurthy, L., and Krishnamurthy, V.: Teleconnections of Indian monsoon rainfall with AMO and Atlantic tripole, *Climate Dynamics*, 46, 2269-2285, 2016.
- Krishnan, R., and Sugi, M.: Pacific decadal oscillation and variability of the Indian summer monsoon rainfall, *Climate Dynamics*, 21, 233-242, 2003.
- 485 Lea, D. W.: Elemental and isotopic proxies of past ocean temperatures, in *Treatise on Geochemistry*, vol. 6, *The Ocean and Marine Geochemistry*, edited by H. D. Holland and K. K. Turekian, pp. 365–390, Elsevier, New York, 2003.
- Lee, K.-E., Kim, J.-H., Wilke, I., Helmke, P., Schouten, S.: A study of the alkenone, TEX86, and planktonic foraminifera in the Benguela Upwelling System: implications for past sea surface temperature estimates. *Geochem. Geo-phys. Geosyst.* 9, Q10019, 2008.
- 490 Levine RC, Turner AG.: Dependence of Indian monsoon rainfall on moisture fluxes across the Arabian Sea and the impact of coupled model sea surface temperature biases. *ClimDyn* 38:2167–2190, 2012.
- Lopes dos Santos, R., Prange, M., Castañeda, I.S., Schefuß, E., Mulitza, S., Schulz, M., Niedermeyer, E.M., Sinningh-Samsté, J.S., Schouten, S.: Glacial-interglacial variability in Atlantic meridional overturning circulation and thermocline adjustments in the tropical North Atlantic. *Earth Planet. Sci. Lett.* 300, 407–414, 2010.
- 495 M., d., Sousa, M., Santos, F., Dias, J., and Gómez-Gesteira, M.: How will Somali coastal upwelling evolve under future warming scenarios?, *Scientific Reports*, 6, 2016.
- Mooley, D., Parthasarathy, B., and Pant, G.: Relationship between Indian Summer Monsoon Rainfall and Location of the Ridge at the 500-mb Level along 75° E, *Journal of climate and applied meteorology*, 25, 633-640, 1986.
- Mortlock, R. A., and Froelich, P. N.: A simple method for the rapid determination of biogenic opal in pelagic marine
500 sediments, *Deep Sea Research Part A. Oceanographic Research Papers*, 36, 1415-1426, 1989.
- Naidu, P. D., and Malmgren, B. A.: A high-resolution record of late Quaternary upwelling along the Oman Margin, Arabian Sea based on planktonic foraminifera, *Paleoceanography*, 11, 129-140, 1996.
- Ninomiya, K., and Kobayashi, C.: Precipitation and Moisture Balance of the Asian Summer Monsoon in 1991, *Journal of the Meteorological Society of Japan. Ser. II*, 77, 77-99, 1999.
- 505 Pant, G. B., and Kumar, K. R.: *Climates of south Asia*, John Wiley & Sons, 1997.
- Parthasarathy, B., Diaz, H., and Eischeid, J.: Prediction of all-India summer monsoon rainfall with regional and large-scale parameters, *Journal of Geophysical Research: Atmospheres*, 93, 5341-5350, 1988.
- Parthasarathy, B., Kumar, K. R., and Munot, A.: Evidence of secular variations in Indian monsoon rainfall–circulation relationships, *Journal of Climate*, 4, 927-938, 1991.
- 510 Prahl, F.G., Wakeham, S.G.: Calibration of unsaturation patterns in long-chain ketone compositions for paleotemperature assessment. *Nature* 330, 367–369, 1987.
- Qasim, S.: Biological productivity of the Indian Ocean, *Indian Journal of Marine Sciences*, 6, 16, 1977.
- R., S.: *Ocean Data View*, 2016.

- Ramaswamy, V., and Gaye, B.: Regional variations in the fluxes of foraminifera carbonate, coccolithophorid carbonate and biogenic opal in the northern Indian Ocean, Deep Sea Research Part I: Oceanographic Research Papers, 53, 271-293, 2006.
- Saher, M., Jung, S., Elderfield, H., Greaves, M., and Kroon, D.: Sea surface temperatures of the western Arabian Sea during the last deglaciation, *Paleoceanography*, 22, 2007.
- Schott, F., Swallow, J. C., and Fieux, M.: The Somali Current at the equator: annual cycle of currents and transports in the upper 1000 m and connection to neighbouring latitudes, Deep Sea Research Part A. Oceanographic Research Papers, 37, 1825-1848, 1990.
- Schott, F. A., Xie, S. P., and McCreary, J. P.: Indian Ocean circulation and climate variability, *Reviews of Geophysics*, 47, 2009.
- Schouten, S., E. C. Hopmans, E. Schefuss, and J. S. Sinningh-Damsté': Distributional variations in marine crenarchaeotal membrane lipids: A new tool for reconstructing ancient sea water temperatures?, *Earth Planet. Sci. Lett.*, 204, 265– 274, 2002.
- Seki, O., Schmidt, D., Schouten, S., Hopmans, E., Sinningh-Damsté, J., Pancost, R.: Paleoceanographic changes in the eastern Equatorial Pacific over the last 10 Myr. *Paleoceanography* 27, PA3224, 2012.
- Shukla J.: Effect of Arabian Sea-surface temperature anomaly on Indian summer monsoon: a numerical experiment with the GFDL model. *J AtmosSci* 32(3):503–511, 1975.
- Sirocko, F., and Lange, H.: Clay-mineral accumulation rates in the Arabian Sea during the late Quaternary, *Marine Geology*, 97, 105-119, 1991.
- Sirocko, F., Sarnthein, M., Lange, H., and Erlenkeuser, H.: Atmospheric summer circulation and coastal upwelling in the Arabian Sea during the Holocene and the last glaciation, *Quaternary Research*, 36, 72-93, 1991.
- Sirocko, F., Sarnthein, M., Erlenkeuser, H., Lange, H., Arnold, M., and Duplessy, J.: Century-scale events in monsoonal climate over the past 24,000 years, *Nature*, 364, 322-324, 1993.
- Sirocko, F., Garbe-Schönberg, D., and Devey, C.: Processes controlling trace element geochemistry of Arabian Sea sediments during the last 25,000 years, *Global and Planetary Change*, 26, 217-303, 2000.
- Tiwari, M., Ramesh, R., Bhushan, R., Sheshshayee, M. S., Somayajulu, B. L., Jull, A., and Burr, G. S.: Did the Indo-Asian summer monsoon decrease during the Holocene following insolation?, *Journal of Quaternary Science*, 25, 1179-1188, 2010.
- Vecchi, G. A., and Harrison, D.: Interannual Indian rainfall variability and Indian Ocean sea surface temperature anomalies.. In *Earth Climate: The Ocean-Atmosphere Interaction*, C. Wang, S. P. Xie & J.A. Carton (eds.), American Geophysical Union., *Geophysical Monograph*, 147, 247-259, 2004.
- Webster PJ, Moore AM, Loschnigg JP, Leben RR.: Coupled ocean-atmosphere dynamics in the Indian Ocean during 1997–98. *Nature* 401:356–360, 1999.

Wuchter, C., Schouten, S., Wakeham, S.G., Sinninghe-Damsté, J.S.: Archaeal tetraether membrane lipid fluxes in the northeastern Pacific and the Arabian Sea: implications for TEX₈₆ paleothermometry. *Paleoceanography* 21, PA4208, 2006.

550 Wyrski, K.: Physical oceanography of the Indian Ocean, in: *The biology of the Indian Ocean*, Springer, 18-36, 1973.


Yadav, R. K.: On the relationship between east equatorial Atlantic SST and ISM through Eurasian wave, *Climate Dynamics*, 48, 281-295, 2017.

Young, D. K., and Kindle, J. C.: Physical processes affecting availability of dissolved silicate for diatom production in the Arabian Sea, *Journal of Geophysical Research: Oceans*, 99, 22619-22632, 1994.


555

560

565



570 **Figure 31.** Modern oceanography of western Arabian Sea. Synchronous change in upwelling intensity and biogenic silica flux clearly indicate that the siliceous productivity in western Arabian Sea is controlled by SWM upwelling. Upwelling strength data is used from Nair, 2000 and Opal flux from Haake et al., 2003.



575 **Figure 24:** Location of sediment core SS-4018 (filled star) in the Arabian Sea. Also shown are the sites discussed in the paper: NIOP-929 (Saher et al., 2007), NIOP-905 (Huguet et al., 2006), SK-17 (Anand et al., 2008), Qunf cave (Fleitmann et al., 2007).




Figure 32: Location of our sediment core SS4018 in the western Arabian Sea. Also shown are the sites discussed in the manuscript. Bottom figures shows the seasonal changes in Wind speed, Precipitation and SST during southwest (SWM) and northeast monsoon (NEM). ECMWF-ERA-Interim data (Berrisford et al., 2011) used and the image obtained using Climate Reanalyzer (<http://cci-reanalyzer.org>), Climate Change Institute, University of Maine, USA.

580




Figure 4: Age depth model of the sediment core SS-4018 (adopted from Tiwari et al., 2010). The error bars marks one sigma uncertainty in calibrated age.

585







Figure 5: Temporal variation of Biogenic silica flux with two sigma uncertainty in sediment core SS4018. Filled triangles at the bottom of the plot marks the age-control points with one sigma uncertainty.



590 **Figure 6. Comparison of biogenic silica flux with SST records from western Arabian Sea for pre-Holocene time (18.5-11.7 ka BP).** a) Biogenic silica flux, b) Mg/Ca based SST from NIOP-905 core (Anand et al., 2008), c) Mg/Ca based SST from NIOP-929 core (Saher et al., 2007), d) TEX_{86} SST from NIOP-905 core (Huguet et al., 2006).



595 **Figure 7. Comparison of biogenic silica flux with rainfall record from eastern Arabian Sea for pre-Holocene time (18.5-11.7 ka BP). (a) Biogenic silica flux (present study), (b) $\delta^{18}\text{O}_w$ IVF (Anand et al., 2008).**



600 **Figure 8: Comparison of Somali upwelling with western Arabian Sea SST records. a) Biogenic silica flux, b) Mg/Ca based SST from NIOP-905 core (Anand et al., 2008), c) Mg/Ca based SST from NIOP-929 core (Saher et al., 2007), d) TEX_{86} SST from NIOP-905 core (Huguet et al., 2006). Grey arrow indicate the trend of proxy records during the last 8 ka.**





Figure 9: Comparison of biogenic silica flux with silica to carbonate ratio in 4018 sediment core. Synchronous changes in both parameters indicate the dominance of biogenic silica flux on the ratio.



605

Figure 10: Comparison of Somali upwelling with southwest monsoon rainfall records during Holocene. (a) Biogenic silica flux (present study), (b) Oman speleothem record, (c) $\delta^{18}\text{O}_w$ IVF data from eastern Arabian Sea. Grey arrow indicate the trend of proxy record during the last 8 ka.

610

Age (ky)	B. Si (%)	σ B.Si	B.Si flux (g/m ² /y)	σ B.Si flux	Age (ky)	B. Si (%)	σ B.Si	B.Si flux (g/m ² /y)	σ B.Si flux
0.00	12.58	0.22	5.51	0.79	12.23	6.41	0.11	3.29	0.41
0.54	12.58	0.25	5.44	0.80	12.54	6.49	0.16	3.36	0.42
1.09	12.58	0.28	5.05	0.80	12.86	6.49	0.20	3.40	0.42
1.63	15.20	0.20	6.62	0.96	12.95	7.76	0.13	4.03	0.49
2.17	14.81	0.14	6.80	0.93	13.04	7.37	0.18	3.76	0.47
2.72	13.06	0.05	5.70	0.82	13.13	7.24	0.09	3.77	0.46
3.26	11.70	0.21	5.34	0.74	13.22	7.74	0.13	3.99	0.49
3.81	10.23	0.17	4.66	0.65	13.47	9.06	0.21	4.68	0.58
4.35	11.79	0.10	5.33	0.74	13.72	8.23	0.16	4.27	0.52
4.90	12.49	0.13	5.78	0.79	13.97	6.83	0.11	3.56	0.43
5.44	11.64	0.22	5.59	0.74	14.22	6.83	0.22	3.52	0.44
5.81	10.52	0.17	5.19	0.67	14.47	5.41	0.10	2.84	0.34
6.18	9.74	0.04	4.74	0.61	14.72	6.11	0.16	3.16	0.39
6.55	10.52	0.29	5.21	0.68	14.89	5.74	0.09	3.04	0.36
7.29	9.59	0.12	4.76	0.61	15.06	5.49	0.19	2.64	0.36
7.93	9.59	0.15	4.71	0.61	15.24	5.24	0.14	2.59	0.34
8.20	8.31	0.19	4.23	0.53	15.41	5.07	0.15	2.27	0.32
9.02	9.35	0.10	4.73	0.59	15.58	5.08	0.04	2.21	0.32
9.29	11.88	0.09	6.03	0.75	15.83	4.94	0.08	2.24	0.31
9.56	12.28	0.19	6.17	0.78	16.09	4.53	0.17	2.07	0.29
9.84	11.57	0.14	5.72	0.73	16.34	3.66	0.11	1.65	0.23
10.12	10.43	0.21	5.26	0.66	16.85	3.54	0.13	1.63	0.23
10.29	10.57	0.10	5.47	0.67	16.94	4.75	0.20	2.22	0.31
10.46	10.82	0.14	5.58	0.68	17.03	4.17	0.15	1.92	0.27
10.63	10.59	0.20	5.38	0.67	17.11	4.04	0.17	1.89	0.27
10.80	10.45	0.11	5.27	0.66	17.20	4.17	0.08	2.02	0.26
10.97	9.47	0.12	4.89	0.60	17.29	3.05	0.12	1.44	0.20
11.28	7.04	0.17	3.59	0.45	17.87	3.52	0.09	1.68	0.22
11.60	6.59	0.09	3.48	0.42	18.44	3.25	0.11	1.57	0.21
11.91	6.92	0.17	3.45	0.44					

615 Table 1: Biogenic silica concentration and flux data.