1	"Temperature and mineral dust variability recorded in two low accumulation Alpine
2	ice cores over the last millennium" by Pascal Bohleber et al.
3	- Response to reviews -
4	
5	Please note:
6	• All line numbers in "Changes to manuscript" refer to the new version (if not
7	noted otherwise)
8	• Changes in the corresponding pdf are highlighted in red
9	• Author's responses to the referee's comments are in blue
10	• All new references can be found in the new manuscript
11	
12	Introductory remark:
13	We thank both referees for their very thorough reviews and we appreciate the helpful
14	suggestions and comments. After careful consideration, especially of points commonly
15	raised by both reviewers, we determined the need to clarify our basic line of argument. For
16	this purpose, we would like to emphasize the following key points:
17	1) We aim to distinguish throughout the paper two separate signal components of the
18	Ca2+ record: <i>1. Episodic spikes</i> , typically two orders of magnitude above
19	background levels, and <i>2. Long-term trends</i> of the decadal-scale average Ca2+
20	concentration. Both components are evaluated separately. At CG, mineral
21	background aerosol levels are generally low and the Ca2+ record is dominated by
22	inputs of Saharan dust (e.g. Wagenbach et al. 1996). In this sense, the already
23	established link between Ca2+ and Saharan dust concerns signal component 1. The
24	potential new link between Ca2+ and temperature is evaluated for signal
25	component 2.
26	2) Regarding 1., we do not intend to make quantitative inferences regarding mineral
27	dust concentrations of individual events but aim to estimate their frequency of
28	occurrence at CG. For this purpose we build on what has already been demonstrated
29	in previous studies, namely that Ca2+ combined with an alkalinity measure is in fact
30	a sensitive and appropriate tool to identify Saharan dust layers at CG (Wagenbach et
31	al., 1996).
32	3) Regarding 2., we respond to the intriguing present situation at CG where we face i)
33	fundamental shortcomings in making quantitative use of the stable water isotope

34		thermometer (Bohleber et al. 2013) and ii) the already known co-variation between
35		trends in Ca2+ and delta O-18 (Wagenbach et al. 1996, Wagenbach and Geis, 1989)
36		as well as delta 0-18 and instrumental temperature (Bohleber et al. 2013). This
37		raises the question to what extent a relationship exists between temperature and
38		Ca2+ trends, and if this may serve as a potential substitute for quantitative
39		temperature reconstruction at CG.
40	4)	While we explore the suggested relation between Ca2+ trends and temperature, we
41		strongly emphasize that it is not our intention to introduce a new ice core
42		temperature proxy. We evaluate the Ca2+ trends solely regarding their <i>site-specific</i>
43		temperature connection. This is an analogue approach as pursued for NH4+ in the
44		Bolivian Andes (Kellerhals et al. 2010).
45	5)	We also emphasize that we by no means disregard the influence of snow deposition
46		and post-depositional effects. In fact, the main goal in using the semi-quantitative
47		snow deposition model (section 2.2) is to demonstrate that post-depositional
48		influence may not be disregarded when evaluating the temperature coupling to
49		Ca2+-trends.
50	In orde	er to eliminate the apparent ambiguities in the original version and in order to make
51	our lin	e of argument more clear we have made the following major changes to the
52	manus	cript. We feel that by means of these changes the most important issues raised by the
53	review	ers have been properly addressed and the clarity of the paper has been substantially
54	improv	red. Detailed responses to the referees' comments are given separately for referee #1
55	and #2	, respectively.
56	Chang	es to manuscript:
57	•	We have clarified the abstract and the conclusions according the above points. We
58		now present two additional tables and two additional figures as supporting
59		evidence in the appendix / as supplementary material.
60	•	Page 4 Lines 29ff.: We added a clear statement regarding the separate treatment of
61		Ca2+-spikes and long-term variability in this study.
62	•	We have split up the previous section 2.2. as follows:
63		• Page 3 Lines 16ff.: We combined with the original section 2.1 the
64		fundamental description of snow preservation at CG and its consequences
65		for interpreting the isotope and mineral dust proxies. This also includes the

66	basic reasoning for expecting a temperature-related imprint in the long-
67	term Ca2+-variability.
68	• Since we feel like it has diverted the attention from our main line of
69	argument, we have moved the details of the semi-quantitative treatment of
70	snow deposition to the supplementary material in the appendix.
71	• Page 18, Line 4ff.: We now refer to the semi-quantitative analysis at a later
72	point in the manuscript. The discussion of potential causes of the observed
73	Ca2+-temperature co-variation is now presented within 5. Results and
74	Discussion. We believe this makes it easier to follow for the reader, since the
75	results have been presented at that point.
76	• Page 18, Line 21ff: We have included a clear statement regarding the site-specific
77	nature of the observed Ca2+-temperature connection.
78	
79	Response to anonymous referee #1
80	This paper presents an exceptional data set composed by stable isotopes and various
81	proxies of aeolian dust deposition as measured in two Alpine cores from Colle Gnifetti
82	(European Alps) during the last 1000 years. The quality of the data presented appears
83	robust and, in general, the time scale adopted seems reliable. However, while this data set
84	does deserve publication, the interpretation and use of the results obtained is often weak,
85	speculative and potentially misleading. I believe the discussion section of the journal
86	Climate of the Past represents a suitable venue to possibly sort out several important
87	interpretative issues before the possible final publication.
88	We thank the referee for the detailed comments. As outlined in the introductory remark, we
89	feel that there is an apparent ambiguity regarding the interpretation of the results, which
90	we have now tried to remove and believe that by this means the main issues raised by the
91	reviewer can be addressed properly.
92	
93	Regardless the source of the dust deposited, the correlation between Ca2+ (as dust proxy)
94	and atmospheric temperature may be really linked to a poorly known post depositional
95	process. Dust in surface snow may indeed facilitate a metamorphic process that could
96	consolidate and conserve the snow on site, for instance via a reduced albedo (larger
97	amounts of solar radiation absorbed) at times with more frequent clear sky and higher
98	atmospheric temperatures. In my view the authors should really try to better describe this

99 possible process.

100	We fully agree that a post-depositional influence must be taken into account, and the
101	mechanism described by the reviewer is entirely consistent with our general view
102	regarding the post-depositional effect of dust-rich surface snow. We have added a more
103	detailed description of this process. In addition we now provide supporting evidence for the
104	influence of dust on snow consolidation. We have included a comparison between Ca2+ and
105	the high-resolution density profile to show directly that in general dust-rich snow layers
106	correspond with confined layers of enhanced density (with respect to the surrounding
107	layers). This supports the concept of dust-rich layers featuring faster snow consolidation
108	(Figure A2).
109	Changes to manuscript:
110	• Changes according to splitting the original section 2.2 as outlined in the initial
111	remarks (see above).
112	• Page 18, Line 4ff.: We have included in this discussion our supporting evidence from
113	the high-resolution density data. We also now include references for each potential
114	driver for the Ca2+-temperature co-variation.
115	
116	Considering the poor seasonality of intense dust events of Saharan origin, at this time I do
117	not see how their semi-quantitative model can be useful to support the possible occurrence
118	of out of phase dust (wet) deposition and post-depositional consolidation. Most
119	importantly, the out of phase dust/snow deposition and the possible post depositional
120	process linked to atmospheric temperature is not sufficiently supported by data. In this
121	respect results from past snow pit studies might be illuminating and should be extensively
122	presented.
123	As outlined in the introductory remark, the main purpose of the semi-quantitative model is
124	to demonstrate that snow deposition can have a non-negligible influence on the long-term
125	variability of Ca2+ and thus must be considered as a potential contribution to the Ca2+-
126	temperature coupling. We have clarified this view accordingly and now also include
127	additional information in the text regarding the following points:
128	• Snow pit data presented in earlier studies (Wagenbach and Geis, 1989) show that
129	local maxima of stable isotope and dust-proxies coincide. This is consistent with
130	what we find in the uppermost section of our core, providing a near-seasonal
131	resolution (Figure A1). Analogue to the treatment of the stable isotope signal in

132	Wagenbach et al. (2012), it is thus justified to assume a potential phase-lag between
133	accumulation and Ca2+ maxima. We also note that the principal function of the
134	phase lag is to mimic the seasonally biased sub-sampling. Qualitatively similar
135	results are found when using a different approach to model this process, e.g. by
136	following Fisher and Koerner (1988).
137	Changes to manuscript: Include Figure A1 as additional supporting evidence in the
138	appendix / supplementary material
139	• The connection between atmospheric temperature and snow consolidation is well
140	established, e.g. temperature being used as parameter in various firn-densification
141	models (also consider Figure 27 in Fauve et al. (2002)). A faster snow consolidation
142	driven by higher summer temperature corresponds in the model to an increase in
143	a_r and a decrease in t_phi. We have clarified this accordingly.
144	Changes to manuscript: Added respective references to discussion on page 18, lines
145	4ff. Added text and clarified the influence of temperature via snow consolidation in the
146	semi-quantitative model (appendix A1).
147	• We also now explicitly point out that the seasonality in Ca2+ is not dominated by the
148	seasonal occurrence in Saharan dust events, but rather the result of increased
149	vertical mixing of air masses in summer vs. overall clean winter snow conditions (cf.
150	Figure A2).
151	Changes to manuscript: Added text and references on page 4, lines 17ff.
152	
153	While the occurrence of a Saharan dust fallout via wet deposition is very well established at
154	Colle Gnifetti, the interannual variability of this kind of Saharan events is very high in the
155	Alps and is mostly related to periods when atmospheric advection dominates (late fall,
156	winter, early spring). During summer, however, atmospheric vertical convection rules in the
157	Alps and dry transport and deposition of dust of local/Alpine origin is extremely likely and
158	probably intense also on Colle Gnifetti. Source reconstruction of dust entrapped in ice cores
159	is very complex even when sophisticated multiple proxies are used (e.g Sr and Nd isotopes).
160	Ca2+, even when combined with ECM and dust size, cannot be considered a specific proxy of
161	Saharan dust events and cannot discriminate with sufficient confidence between different
162	kinds of sources (e.g. Saharan vs. local) and atmospheric circulation (e.g. meridional flows
163	vs. vertical convection).

164 We agree that vertical atmospheric convection likely rules the seasonal cycle of Ca2+. The

165 sophisticated isotopic fingerprinting of individual dust layers is certainly a worthwhile 166 future target (and we appreciate the suggestion), although requiring substantial analytical 167 capabilities as well as a significant portion of the ice core. For our main target being 168 quantifying the frequency of occurrence of Saharan dust events, the task was to identify the 169 specific imprint of Saharan dust in presence of events of increased impurity load by 170 enhanced vertical mixing. Here we were able to rely on what has already been shown in the 171 two earlier studies investigating dust-related records at CG (i.e. Wagenbach et al. (1996) 172 and Wagenbach and Geis (1988)). These studies have clearly demonstrated that 173 investigating "whether the size distribution parameters of the Saharan dust • 174 deposited differ significantly from those of dust deposits related to more regional 175 source areas or even background aerosol" revealed that "the volume size 176 distribution of the two Saharan dust events is indeed significantly shifted towards 177 larger particles" (Wagenbach and Geis, 1988, and cf. Figure 7 therein). The authors 178 also conclude that "it seems reasonable to attribute individual differences in the size 179 distribution of the Saharan dust deposited on CG to different transport times which 180 in turn means different transport velocities and/or different lengths of the 181 trajectories". 182 "that the Ca2+ spikes associated with the strongly alkaline snow layers in recent firn • 183 at Colle Gnifetti are most likely due to the mobilisation of calcerous Saharan soil 184 material which, following long range transport to the Alps, is mainly deposited there 185 by precipitation scavenging" (Wagenbach et al. 1996). The authors compile data 186 from various CG snow pits and ice cores, finding that "This compilation suggests that 187 the analyses of ionic Ca2+ in connection with reliable alkalinity measurements 188 appear to be a very sensitive and specific tool to identify Saharan dust influenced 189 snow layers in high depth resolution". 190 We now make additional clear reference to these specific earlier findings to point out that it 191 is not the scope of our work to present a refined detection of Saharan dust in CG ice cores. 192 Instead, we intend to make use of already established tools, in combination with our new 193 ice core chronology comprising the last millennium at sufficient confidence. 194 Changes to manuscript: Page 4, Lines 26-28: Added text and reference accordingly. 195 196 Remarkably, if Ca2+ concentration is assumed to really trace dust from a specific area (e.g. 197 Sahara desert) this would severely prevent its general use as paleothermometer as this

198	parameter would strongly depend on environmental conditions at the source that are not
199	directly related to atmospheric temperature. For instance a Ca2+ based paleo thermometer
200	might be severely biased by different soil conditions at the source, due for instance to
201	changes in atmospheric circulation and precipitation, not temperature (e.g. green Sahara
202	during the middle-early Holocene). The use of Ca2+ to construct a paleothermometer could
203	thus depend on very site specific post-depositional processes and on changing
204	environmental conditions at the source of dust over time. Thus it is fundamental to greatly
205	caution about its general use in space (drilling site dependence) and time (it may work only
206	during certain times). In conclusion, while the relation between Ca2+ and temperature is
207	interesting, its extended use cannot provide unambiguous novel knowledge about past
208	atmospheric temperature.
209	Here we refer again to our initial comment regarding the separate treatment of Ca2+ spikes
210	(Saharan dust proxy together with an alkalinity measure) and the overall trend component
211	of the Ca2+ signal. Only the latter is explored with respect to a potential coupling to
212	temperature, and we are now being more explicit regarding this distinction. We are also
213	clearly stating that, at best, Ca2+ is a site-specific temperature proxy. Regarding its
214	performance to reconstruct temperature in time, please see the next comment below.
215	Changes to manuscript:
216	• Page 4, Lines 29ff.: Added statement regarding distinction of signal components in
217	Ca2+.
218	• Page 18, Lines 21ff.: Added statement regarding site-specific role of Ca2+.
219	
220	When compared to Ca2+, the use of stable isotopes as a proxy of atmospheric temperature
221	is better justified by the physics and is less sensitive to source effects due to moisture
222	changes. It is remarkable that the correlation of stable isotopes and instrumental
223	atmospheric temperature is very good during the last 150 years when the recorded
224	instrumental temperatures are most robust. It appears that the real problem of the possible
225	paleo-thermometer based on stable isotopes is the "excessive" and spatially variable
226	sensitivity (1.4-2.3 per mill/C) when compared to what one would expect (0.65 per mill/C).
227	For this reason the authors decide not to attempt a quantitative temperature reconstruction
228	based on stable isotopes. However, while this decision may be justified, I believe that an
229	extended discussion of the possibly biased paleo-temperatures obtained by means of this
230	presumably too sensitive/variable paleo-thermometer would be very interesting. This

231 could show for instance the inconsistency of the temperatures obtained, even considering

physical processes such as the amplification of atmospheric temperature anomalies with

the elevation.

234 We believe that an adequate discussion of the lags and leads of the peculiar high

- isotope/temperature sensitivity is beyond the scope of this work and would likely result in
- an entirely different paper. Ideally, such an evaluation would need to take into account
- regional isotope-climate modeling. We take the reviewers comment as an encouragement to
- 238 pursue such a dedicated assessment in a future study. The agreement between the isotope
- and temperature trends within the last 150 years has been discussed in an earlier study,
- $240 \qquad also showing that if calibrated with the last 100 years, isotope levels at CG suggest a warmer$
- 241 "early instrumental period" by about 0.4 deg C (Bohleber et al. 2013). Following this
- 242 comment we now include this point in our discussion.
- 243 Remarkably, however, if considering the Ca2+ trends against instrumental temperature, we
- find an agreement of similar quality as for the isotopes, but now persistent over the full
- instrumental 250-year period (cf. Figure 7). Although the exact reason for a potential non-
- stationary isotope-temperature sensitivity remains elusive, this finding suggests that the
- 247 potential Ca2+-temperature relationship may not be affected to the same degree. Moreover,
- $248 \qquad \text{if using the Ca2+-temperature for a tentative calibration, the reconstructed temperature} \\$
- 249 variability is consistent with the latest summer temperature reconstruction based on other
- archives (the Luterbacher et al. (2016) record). The main point here is that given the
- agreement with instrumental data and other temperature reconstructions, we find no
- evidence of potential non-stationary behavior in the Ca2+-temperature correspondence.
- 253 We have added text in order to be more clear about this interpretation.
- 254 Changes to manuscript: Page 19, Lines 13ff.: Added text to clarify the interpretation of the255 comparison with the other proxy reconstruction.
- 256
- 257 A likely reliability of the timescale obtained for the two cores from Colle Gnifetti is
- suggested by the good correlation of Ca2+ with an independent, well dated, past summer
- temperature record obtained from tree rings. However, the reliability of this time scale may
- be just due to the use of recent absolute time horizons (during the last century) and 14C
- 261 measurements (although the significant reference, PhD thesis of Hoffmann 2016, cannot
- 262 provide an accessible and peer reviewed methodological support). Counting of annual
- 263 layers is not convincing in the deep part of core KCC. In particular the concept of "group of

264 peaks" of Ca2+ to identify a single annual layer seems extremely arbitrary, and, at this time, 265 not supported in the paper by any snow pit observation. In addition a possibly larger 266 deflation of lighter snow during the colder summers of the Little Ice Age may have removed 267 more annual layers that expected. In this way, it is very possible that the interval period 268 between 150 and 600 BP remains unconstrained and prone to larger uncertainties. 269 We do not use the comparison with the summer temperature reconstruction to show the 270 reliability of our time scale. This would require comparing an already demonstrated 271 temperature signal reconstructed from the ice core. Our argument goes in the opposite 272 direction: Having constructed a reliable time scale, we look for consistency between the 273 potential Ca2+-based temperature variability and an established reconstruction from 274 another archive. 275 We believe we have demonstrated the reliability of our time scale for the following reasons: 276 Annual layer counting is a well-established and widely used tool for ice core dating. 277 However, at CG the employment of this tool was so far limited by rapid layer thinning 278 beyond the resolution of most melting techniques. Based on the agreement between CFA 279 and LA-ICP-MS, we were able to make extended use of annual layer counting to build our 280 chronology. While the identification of additional absolute dating horizons, such as volcanic 281 eruptions, would of course be desirable, this is at present not possible for CG. However, we 282 have taken great care to account for the resulting uncertainty in "unconstrained" annual 283 laver counting (section 4.4). The resulting age-scale is backed by 14C markers, an approach 284 already employed successfully at CG in previous studies (e.g. Jenk et al. 2009). The PhD 285 thesis by Helene Hoffmann has been reviewed as part of the process in obtaining her PhD at 286 Heidelberg University. It is fully available online and we have now included the link in the 287 respective reference. In addition, we are now able to reference her according publication 288 which has been accepted for publication in *Radiocarbon* in the meantime (Hoffmann et al. 289 2017). 290 We thank the referee for bringing our attention to provide additional justification for the 291 concept of "grouped peaks" in the LA-ICP-MS Ca signal. For this purpose we have prepared 292 a supplementary figure showing the uppermost 5 m of the delta O-18 and Ca2+ stratigraphy 293 in our core. Here, multiple sub-seasonal peaks in Ca2+ are clearly present to back up this 294 concept. In addition, we again refer to the fact that based on its ultra-high resolution and 295 non-destructive technique, LA-ICP-MS can map the spatial heterogenity of impurities even

within a single annual layer.

297	Changes to manuscript:
298	• Page 19, Lines 13ff.: Clarified regarding the comparison with the other proxy
299	reconstruction not being intended for dating verification.
300	• Added reference for Hoffmann et al. (2017a,b)
301	Added Figure A1 as supporting evidence for "grouped peak" concept
302	
303	While Colle Gnifetti is a very well know ice core drilling site and many glaciological studies
304	have been performed, this paper fails to offer the necessary comparison with existing data
305	set available from other ice cores (and snow pit samples) obtained at the same site. In
306	particular several studies of particulate and aerosol deposition were performed (e.g.
307	Thevenon, JGR 2009) and should be carefully compared to check the consistency (or not) of
308	the new findings with the previous results.
309	We already compare our results to the findings of Thevenon et al. (2009) on page 18 line 5
310	(original manuscript). However, a more detailed comparison is hampered given the
311	difference in depth resolution and dating uncertainty between the two studies. We would
312	certainly welcome a specific suggestion by the referee regarding the further comparison of
313	our results.
314	Changes to manuscript: Page 17, Lines 7ff.: Added some more detail regarding the
315	comparison with Thevenon et al. (2009)
316	
317	
318	Specific comments:
319	
320	P1 L1: "Among ice core drilling sites in the European Alps, the Colle Gnifetti (CG) glacier
321	saddle is the only one to offer climate records back to at least 1000 years"
322	
323	There is now in the Eastern Alps a new ice core climate record that goes back almost 7000
324	years (Gabrielli et al. The Cryosphere 10, 2779–2797, 2016; Gabrielli et al. 19th EGU
325	General Assembly, EGU2017, proceedings from the conference held 23-28 April, 2017 in
326	Vienna, Austria., p.9932).
327	Thank you for pointing this out. We have changed our wording in order to be more specific.
328	Colle Gnifetti is the only non-temperate site (Ortles in the Eastern Alps is partially
000	

329 temperate).

330	Changes to manuscript: Page 1, Line 1 and Page 2, Line 5: Changed wording accordingly.
331	
332	P1 L8: "A high and potentially non-stationary isotope/temperature sensitivity limits the
333	quantitative use of the stable isotope variability thus far".
334	
335	This statement is not discussed sufficiently within the text.
336	In this case the statement primarily serves as background and motivation (referring to what
337	is already known at CG).
338	Changes to manuscript: Page 1, Lines 7ff.: We have changed the text accordingly to make
339	this more clear.
340	
341	P1 L15: "the medieval climate period around 1100–1200 AD stands out through an
342	increased occurrence of dust events, potentially resulting from a relative increase in
343	meridional flow and dry conditions over the Mediterranean".
344	
345	While the frequency of the dust horizons is reproducible in the two cores, they cannot be
346	linked to individual dust events of Saharan origin that cannot be unambiguously
347	distinguished from the occurrence of local past summer surfaces marked by dust
348	accumulation.
349	As pointed out in our response above, we are making use of a tool developed in an earlier
350	study and demonstrated to be suitable to identify Saharan dust events in CG ice cores
351	(Wagenbach et al. 1996). Following the discussion with the referee, however, we now
352	include a statement that more sophisticated methods based on isotopic fingerprinting exist
353	today and may be used in the future to test and refine our findings.
354	Changes to manuscript: Page 20, Lines 15ff. Added text with a respective statement.
355	
356	P2 L5: "Colle Gnifetti (CG) in spite of its limited glacier depth – stands out as the only site
357	where net snow accumulation is low enough to provide records over the last millennium
358	and potentially beyond at a reasonable time resolution".
359	
360	Again, this is not correct, please see Gabrielli et al. The Cryosphere 10, 2779–2797,
361	2016.
362	Thanks- we changed our wording accordingly (see above).

364	P4 L5 "A single deposition event typically lasts less than a few days (Sodemann et al., 2005;
365	Schwikowski et al., 1995). The associated warm air temperature and the substantially
366	lowered snow albedo both support surface snow consolidation and partly protect the dust
367	layer from wind erosion."
368	
369	As long as air temperature is below the freezing level (as during snow events), this cannot
370	be a factor facilitating snow consolidation.
371	As discussed in our response above, temperature does play an important role in snow
372	consolidation.
373	Changes to manuscript: Page 18, Lines 4ff.: We have added a specific remark and an
374	according references regarding this point.
375	
376	P4-10 "Therefore, the Ca2+ record of the CG ice cores is primarily related to mineral
377	dust and dominated by Saharan dust related spikes".
378	
379	This conclusion is unsupported by more recent data detailing more specific proxies of
380	Saharan dust.
381	As indicated in the text, this statement refers to findings already published by two previous
382	studies (Wagenbach et al. 1996, Wagenbach and Geis 1988). If the referee would like to
383	suggest a specific study that is providing new (in particular refuting) evidence, we would
384	certainly consider this in our discussion.
385	
386	P4-27 "For instance, warm summers feature increased vertical mixing and hence a higher
387	atmospheric impurity load, and in addition, entail faster fresh snow consolidation. This may
388	lead to an increased relative amount of impurity-rich summer snow deposition."
389	
390	This is a very reasonable and, unlike the meridional Saharan advection, a more regular
391	process in the Alps. I'm not sure why the authors do not consider and discuss it further
392	within the text.
393	From the discussion with the referee we learned that we have not emphasized this point
394	strongly enough, although we believe it is a very important process in this context. In fact
395	part of this process is what we explore with the semi-quantitative model exercise. We have

clarified this and come back to it in the respective parts of the Discussion.
Changes to manuscript:
• Page 3, Lines 18ff.: Clarified and added text.
• Page 18, Lines 4ff.: Included a detailed discussion of all involved processes.
P5-L7 "Here we follow the model of Wagenbach et al. (2012), which assumes sinusoidal
cycles for the precipitation-borne signal $S(t)$ and the surface accumulation pattern $A(t)$, and
a phase-lag t' between S(t) and A(t)."
While accumulation and delta 180 seasonal patterns are accepted at Colle Gnifetti, it is
much less so for a disturbed seasonal signal like dust (Ca2+). The authors need to present
data supporting how the sinusoidal assumption is justified for Ca+2 deposition.
Paragraph 2.2 In general I do not find that this paragraph well written or even necessary. It
is not clearly explained what are the main motivations and conclusions of the conceptual
model. At this time I'm also not sure how it supports the rational of the interpretation. Does
the model show that a phase lag between deposition and consolidation explain some recent
observations? If so, the phenomenology of these processes needs to be supported,
displaying existing data (e.g. snow pits) that could complement the conceptual discussion
performed by means of this model.
As outlined in the initial comment and after careful consideration of the referee's comment
we have decided to break up the original paragraph as not to divert from our main line of
argument. Being now used in the Discussion the scope of the model consideration becomes
more clear, i.e. to semi-quantitatively demonstrate that the influence of snow preservation
may not be disregarded when evaluating the long-term variability of Ca2+ and that it must
be considered as a potential process introducing the coupling to temperature.
Albeit the sinusoidal pattern is of course an idealization, we have now included additional
evidence for the seasonal pattern and link between delta O-18 and Ca2+ (Figure A1).
Changes to manuscript:
• Page 18, Lines 4ff.: Moved and rewrote this section (originally part of 2.2) to clarify
• Additional supporting figures in the appendix (Figure A1, A2)
P7-L2 "The threshold (4.0 μm) was chosen such that it corresponds to the expected median

429 particle diameter of Saharan dust particles at CG (Wagenbach and Geis, 1989)."

430 431 Is this threshold necessary AND sufficient to discriminate between Saharan and non-432 Saharan dust? I do not think so. The reference reported is pretty old and in the meantime 433 many tools have been developed to characterize dust sources. In my view this threshold is 434 just indicative but not strictly discriminant. 435 Although the reference is old we are not aware that its findings have been refuted by newer 436 studies. As said before, we certainly agree that more sophisticated tools may potentially 437 allow for a precise fingerprinting of the individual dust sources, although requiring 438 substantial analytical effort. 439 We intended to use the already established discrimination method based on the median 440 particle diameter and set our threshold according to the previous study. That said, the 441 results are qualitatively independent with respect to the exact choice of the threshold. 442 Notably this includes the outstanding feature in CPP during the medieval period. 443 Only the sensitivity of the CPP to changes in the particle size distribution (PSD) is 444 dependent on the threshold. Choosing the threshold to be the median value of the normal 445 PSD means that the CPP is 50% for "regular" dust and is sensitive to changes in the PSD. 446 Changes to manuscript: Page 20, Line 7-8: We added a statement to point out that the 447 outstanding feature during the medieval period is not a result of the choice of threshold. 448 449 3.2.1 Radiocarbon analysis. A table detailing all the results obtained by analyzing the 6+5 450 samples from KCC and KCI needs to be reported, including the linked uncertainties. 451 We have added a table as requested and would also like to point out that we were able to 452 include an additional radiocarbon measurement for KCI, in clear support of the age scale. 453 **Changes to manuscript:** We included a respective table in the appendix, Table A2. 454 455 4 Ice core dating. A Table summarizing all the time horizons used in KCC and KCI needs to 456 be reported. Another table indicating different kind of annual layer counting in different 457 sections of the two cores would be also very useful. 458 **Changes to manuscript:** We included a respective table in the appendix, Table A1. 459 460 P9-L9 "(BP, referring to the drilling year of the respective ice core if not otherwise noted)". 461

462	This could be very confusing. I strongly suggest to use a different notation.
463	We have generally changed this notation to either year AD or stating the precise year, e.g.
464	year b2005 or year b2013 in order to be more precise.
465	
466	P10-L6 "The groups of peaks are separated by a comparatively stable signal of low Ca
467	concentrations. The latter is interpreted as resulting from the varying degree of winter
468	snow being included in the record otherwise dominated by summer snow. "
469	
470	This "group of peaks" sounds very suspicious and arbitrary. This idea needs to be supported
471	with additional evidences from snow pit studies or from a comparison with the seasonality
472	of stable isotopes at depths where annual layers are still distinguishable.
473	We thank the reviewer for bringing this to our attention. As requested we provide
474	additional support of this concept in a supplementary Figure A1.
475	
476	P10-L8 "Accordingly, the grouped peaks correspond to sub-annual snow deposition events
477	of elevated Ca concentration during the summer period".
478	
479	How can multiple wet dust deposition events be distinguished from the formation of one or
480	more summer surfaces of accumulated dust?
481	We do not see how, based on the LA-ICP-MS Ca signal alone, one could distinguish wet and
482	dry deposition. However, the comparison with the CFA-based counting in depth intervals
483	where CFA clearly identifies the annual layer signal (Figure 4 b)) shows that the "grouped
484	peaks" are not a result of multiple, very closely spaced, summer surfaces. It would be hard
485	to imagine a depositional behavior that produces such a "grouped peak" pattern at the
486	observed regularity. Considering the additional Figure A1, it appears much more plausible
487	to assign this to sub-seasonal structure resolved by LA-ICP-MS.
488	
489	The different age depth relationships displayed in Fig. 5 for KCC and KCI between 150 years
490	BP and 700-800 years BP needs to be carefully discussed. In fact, within this interval both
491	cores do not depend on absolute time horizons.
492	As discussed above, unfortunately there is no means to identify additional horizons in the
493	respective time period. We have accounted for the respective uncertainty in annual layer
494	counting accordingly. Nonetheless, we have added more text to point out this circumstance.

495 **Changes to manuscript:** Page 10, Line 32-34: Added a respective statement to the text. 496 497 Caption Fig. 4b The year to year correspondence between Ca annual layers determined 498 by LA and CFA should be indicated drawing lines connecting the star symbols. At the 499 moment no clear one-to-one link is apparent. 500 Although the overall pattern between the CFA signal and the general baseline in LA-ICP-MS 501 is highly similar, we are cautious about necessarily assigning a one-to-one correspondence 502 between peaks. This is not least due to the possibility of a slight remaining offset between 503 the two depth scales. We have added text to fully explain this in the caption. It was our 504 intention to demonstrate that for a given depth interval, the number of years counted in 505 both signals (CFA and LA-ICP-MS) is consistent within uncertainty. 506 **Changes to manuscript:** Added text to the caption of what is now Figure 3. 507 508 P12 L10 "The frequency of occurrence in these total snow loss events is, however, 509 extremely hard to quantify. Counting annual layers in between the above mentioned (dust) 510 horizons within the last century, reveals an offset of typically only one to two years as 511 compared to the known age of the horizons." The last century is not very much 512 representative of colder periods (Little Ice Age, LIA) where snow drift could have been far 513 more important, possibly eroding more annual layers. Notably the LIA is also the time when 514 no absolute time horizons are available for the time scale that depends entirely on counting 515 annual layers. 516 Agreed, but there is probably not much that can be changed about this until additional 517 absolute horizons are discovered. To reiterate, we have employed a dedicated approach to 518 quantify our counting uncertainty. 519 520 P13 L13 "(e.g. note the distinct isotope minima around 1360 AD)." This is an important note 521 when considering the companion paper by More et al. 2017 in Geohealth as this time 522 corresponds almost exactly with the time of the Black Death. Could this isotope minima 523 have been a large winter snow accumulation event? In this sense the implications for the 524 interpretation of the linked Pb record could be very important. 525 We point out that the two papers were not designed to be companion papers, e.g. as stated 526 in the manuscript there are minor differences in the used age scale. Although the 1360 AD 527 isotope minima could be in principle be connected to a higher percentage of winter snow

528	preservation, the signal in the impurity species is less outstanding. We thank the referee for
529	noting this and take this comment as encouragement for further investigation.
530	
531	P14 L14 "However, this is the first time that the correlation holds to this extent also for the
532	comparatively old core sections".
533	
534	As far as I know, this has been also observed at least in the Ortles ice cores (Gabrielli
535	et al. The Cryosphere 10, 2779–2797, 2016).
536	This statement refers primarily to the CG ice cores, and we have changed the wording
537	accordingly. That said, to our knowledge Gabrielli et al. (2016) show a comparison of the
538	three Ortles ice cores on a depth scale, not as time series.
539	Changes to manuscript: Page 12, Line 17: "However, this is the first time that the
540	correlation holds to this extent also for the comparatively old core sections of CG cores"
541	
542	P15 L5 "Stack of the two stable isotope records (calculated as their simple average)".
543	
544	Please, mention the temporal step used to calculate averages.
545	Changed accordingly.
546	Changes to manuscript: Page 13, Line 32: "Stack of the two stable isotope records
547	(calculated as their simple average, at nominal annual resolution)"
548	
549	P15 L15 "substantially higher sensitivity values for KCI than KCC, revealing 2.3 vs. 1.4 per
550	mill/C, respectively".
551	
552	This is surprising considering the striking similarities of the two stable isotope profiles (Fig.
553	6). Could you provide an explanation?
554	We do not have a full explanation for this phenomenon, adding to the enigmatic nature of
555	the isotope sensitivity at CG. However, the central difference between KCI and KCC is the
556	especially low net accumulation at KCI. This entails an even stricter confinement to
557	sampling mainly the summer season precipitation. This points towards the depositional
558	bias to play a role in explaining the high sensitivity values.
559	
560	P15 L16 "changes in snow preservation are expected to bias sensitivity (cf. the conceptual

561	consideration in section	2.2).	"
-----	--------------------------	-------	---

In this case, the conceptual model presented should be useful to quantify and perhapscorrect this bias.

565 This statement refers to the reasoning of behind the previous comment (see above). While

using the model to correct for the bias is an interesting suggestion, it would certainly

require information on the past variability in snow deposition at each site, which is not

available and generally very hard to quantify at CG. We decided to reword this paragraph toclarify our reasoning.

570 Changes to manuscript: Page 14, Line 10ff.: Reworded the paragraph accordingly.

571

572 P15 L17 "This is consistent with the sensitivity difference among KCI and KCC, since an even

573 more strict confinement towards sampling the high summer season can be expected for the 574 lower accumulation KCI."

575

576 Agree, but from just a look of the two records, KCI and KCC show very similar absolute

577 stable isotopes values during the potential calibration time (Fig. 6).

578 The difference is in fact small compared to the absolute values (e.g. for the last 100 years -

579 13.48 vs -13.71 per mil for KCC and KCI, respectively). However, it also affects the

580 magnitude of long term trends, e.g. the recent increase in isotope values over the last 100

581 years (e.g. based on linear regression 0.20 vs. 0.25 per mil / decade for KCC and KCI,

respectively). We thank the reviewer for pointing this out and have included a respectiveremark in the text.

584 Changes to manuscript: Page 14, Line 10ff.: Reworded the statement accordingly.

585

586 P17 L3 "While the particle signal alone is not sufficient for differentiating these events,

587 Saharan dust layers in CG ice cores can be reliably identified based on the analyses of Ca2+,

588 supplemented by alkalinity measurements and, in principle, particle size distribution

589 (Wagenbach et al., 1996)."

590

591 This information is not sufficient to discriminate between a Saharan dust event and a past

592 summer surface formed by dry dust accumulation. While Saharan dust events may well

593 have these characteristics, also summer dust layers formed during prolonged dry periods

594	could have the same or similar characteristics. In addition, a higher coarse particle
595	percentage may be more indicative of local dust rather than long-range transported dust.
596	In order to avoid redundancies we would like to refer to our previous responses presented
597	above, and references to the earlier studies.
598	
599	P18 L5 "This is in broad agreement with periods of enhanced Saharan dust deposition
600	reported by Thevenon et al. (2009) obtained from elemental analysis in a CG ice core."
601	
602	This presumed broad agreement should be demonstrated in detail. At this time the high
603	frequency of dust events in the described periods is both consistent with more frequent
604	Saharan events and the formation of dust enriched past summer surfaces.
605	Accepting the limitations due to the different depth resolution in both records, we made an
606	attempt to compare our findings with the results by Thevenon et al. (2009) in a little more
607	detail.
608	Changes to manuscript: Page 17, Lines 7 ff.: Added text.
609	
610	P19 L25 "The connection of the distinct increase in coarse particles with enhanced dust
611	event frequency indicates an increase in direct transport of Saharan dust the (as opposed to
612	indirect advection with longer pathway and thus stronger decrease in coarse particles)."
613	
614	This is, at best, consistent (not indicative) with an increase in direct transport of Saharan
615	dust. In fact this observation is also compatible with other scenarios (e.g. higher occurrence
616	of summer surfaces marked by dust, increase in the intensity of the summer vertical
617	transport of dust of Alpine origin).
618	Based on our response presented above regarding the distinct difference in particle sizes of
619	Saharan dust (Wagenbach et al. 1996, Wagenbach and Geis 1988), in our view this
620	observation is not compatible with the mentioned scenarios. However, we have slightly
621	changed the wording of this statement following the referees suggestion.
622	Changes to manuscript: Page 20, Line 11: "the connection of the distinct increase in coarse
623	particles with enhanced dust event frequency rather <i>suggests</i> an increase in direct
624	transport of Saharan dust"
625	

626 P19 L27 "Notably, this view is consistent with increased deuterium excess, which would be

627 exp	pected from wa	arm and dry air	masses collecting	g moisture over the	e Mediterranean."
---------	----------------	-----------------	-------------------	---------------------	-------------------

- 628
- 629 This may be an important note that needs to be expanded and adequately referenced.
- 630 Finding increased values of deuterium excess supports the view of a relative increase in
- 631 Saharan dust advection, which we have made more clear. We have also added an according
- 632 reference as suggested.
- 633 **Changes to manuscript:** Page 20, Line 15ff.: Changed accordingly.
- 634
- 635 P20 L12 "increased meridional transport favoring direct Saharan dust advection over the
- 636 Mediterranean is consistent with a NAO+ dominated MCA, lasting 1100 to about 1300 AD,
- 637 as proposed by Trouet et al. (2009). Although NAO is mainly a winter signal (thus not a
- 638 direct concern to CG summer representative ice core signals)"
- 639
- 640 As mentioned, Saharan dust advection is just one of the possibilities and a NAO signal
- 641 consistent with this hypothesis is really a weak reasoning, especially considering that a NAO
- 642 winter signal can have little impact on the summer biased cores from Colle Gnifetti. Please,
- 643 consider to entirely remove this section (lines 10-17).
- 644 Following the reviewers comment and after additional consideration, we have decided to645 remove this section.
- 646
- 647 Conclusions: conclusions needs to be tuned down accordingly to the main observations
- 648 performed.
- 649 Changed accordingly.
- 650 Changes to manuscript: Page 20, Line 25ff.: Reworded the second part of the conclusions.651
- 652 P21-L12 "The intrinsic contribution of snow preservation may bias the isotope temperature653 sensitivity".
- 055 Selisiti
- 654
- 655 This is interesting but it has not been shown and adequately discussed within the text.
- 656 We have removed this statement from the conclusions.
- 657

1	"Temperature and mineral dust variability recorded in two low accumulation Alpine
2	ice cores over the last millennium" by Pascal Bohleber et al.
3	- Response to reviews -
4	
5	Please note:
6	• All line numbers in "Changes to manuscript" refer to the new version (if not
7	noted otherwise)
8	• Changes in the corresponding pdf are highlighted in red
9	• Author's responses to the referee's comments are in blue
10	• All new references can be found in the new manuscript
11	
12	Introductory remark:
13	We thank both referees for their very thorough reviews and we appreciate the helpful
14	suggestions and comments. After careful consideration, especially of points commonly
15	raised by both reviewers, we determined the need to clarify our basic line of argument. For
16	this purpose, we would like to emphasize the following key points:
17	1) We aim to distinguish throughout the paper two separate signal components of the
18	Ca2+ record: <i>1. Episodic spikes</i> , typically two orders of magnitude above
19	background levels, and <i>2. Long-term trends</i> of the decadal-scale average Ca2+
20	concentration. Both components are evaluated separately. At CG, mineral
21	background aerosol levels are generally low and the Ca2+ record is dominated by
22	inputs of Saharan dust (e.g. Wagenbach et al. 1996). In this sense, the already
23	established link between Ca2+ and Saharan dust concerns signal component 1. The
24	potential new link between Ca2+ and temperature is evaluated for signal
25	component 2.
26	2) Regarding 1., we do not intend to make quantitative inferences regarding mineral
27	dust concentrations of individual events but aim to estimate their frequency of
28	occurrence at CG. For this purpose we build on what has already been demonstrated
29	in previous studies, namely that Ca2+ combined with an alkalinity measure is in fact
30	a sensitive and appropriate tool to identify Saharan dust layers at CG (Wagenbach et
31	al., 1996).
32	3) Regarding 2., we respond to the intriguing present situation at CG where we face i)
33	fundamental shortcomings in making quantitative use of the stable water isotope

34		thermometer (Bohleber et al. 2013) and ii) the already known co-variation between
35		trends in Ca2+ and delta O-18 (Wagenbach et al. 1996, Wagenbach and Geis, 1989)
36		as well as delta 0-18 and instrumental temperature (Bohleber et al. 2013). This
37		raises the question to what extent a relationship exists between temperature and
38		Ca2+ trends, and if this may serve as a potential substitute for quantitative
39		temperature reconstruction at CG.
40	4)	While we explore the suggested relation between Ca2+ trends and temperature, we
41		strongly emphasize that it is not our intention to introduce a new ice core
42		temperature proxy. We evaluate the Ca2+ trends solely regarding their <i>site-specific</i>
43		temperature connection. This is an analogue approach as pursued for NH4+ in the
44		Bolivian Andes (Kellerhals et al. 2010).
45	5)	We also emphasize that we by no means disregard the influence of snow deposition
46		and post-depositional effects. In fact, the main goal in using the semi-quantitative
47		snow deposition model (section 2.2) is to demonstrate that post-depositional
48		influence may not be disregarded when evaluating the temperature coupling to
49		Ca2+-trends.
50	In orde	er to eliminate the apparent ambiguities in the original version and in order to make
51	our lin	e of argument more clear we have made the following major changes to the
52	manus	cript. We feel that by means of these changes the most important issues raised by the
53	review	ers have been properly addressed and the clarity of the paper has been substantially
54	improv	red. Detailed responses to the referees' comments are given separately for referee #1
55	and #2	, respectively.
56	Chang	es to manuscript:
57	•	We have clarified the abstract and the conclusions according the above points. We
58		now present two additional tables and two additional figures as supporting
59		evidence in the appendix / as supplementary material.
60	•	Page 4 Lines 29ff.: We added a clear statement regarding the separate treatment of
61		Ca2+-spikes and long-term variability in this study.
62	•	We have split up the previous section 2.2. as follows:
63		• Page 3 Lines 16ff.: We combined with the original section 2.1 the
64		fundamental description of snow preservation at CG and its consequences
65		for interpreting the isotope and mineral dust proxies. This also includes the

66	basic reasoning for expecting a temperature-related imprint in the long-			
67	term Ca2+-variability.			
68	• Since we feel like it has diverted the attention from our main line of			
69	argument, we have moved the details of the semi-quantitative treatment of			
70	snow deposition to the supplementary material in the appendix.			
71	• Page 18, Line 4ff.: We now refer to the semi-quantitative analysis at a later			
72	point in the manuscript. The discussion of potential causes of the observed			
73	Ca2+-temperature co-variation is now presented within 5. Results and			
74	Discussion. We believe this makes it easier to follow for the reader, since the			
75	results have been presented at that point.			
76	• Page 18, Line 21ff: We have included a clear statement regarding the site-specific			
77	nature of the observed Ca2+-temperature connection.			
78				
79	Response to anonymous referee #2			
80	The paper presents an excellent dataset of stable water isotopes and other 'dust' proxies			
81	(i.e. insoluble particles and Ca2+) from two separate ice cores drilled at Colle Gnifetti in the			
82	Pennine Alps, reaching back in time as far as a thousand year, a remarkable achievement for			
83	a European alpine ice core. This study combines a very good quality of data retrieval with a			
84	robust strategy regarding the dating, and therefore deserves to be published in Climate of			
85	the Past. The data treatment and statistical approach is also adequate and robust and only			
86	minor changes should be made. I will illustrate now few of the weaknesses that the			
87	manuscript presents and some suggestions on how to strengthen these points before the			
88	final publication. Detailed comments follow.			
89	We thank the referee for the comments and encouragement to further strengthen the			
90	manuscript.			
91				
92	Firstly, the manuscript fails a bit in illustrating the reason why it is important to obtain a			
93	Ca2+-derived temperature profile and what advantages/disadvantages this would have			
94	compared to a conventional $\delta 180$ -derived temperature profile. As mentioned in the			
95	abstract, the high and potentially non-stationary isotope/temperature sensitivity limits the			
96	quantitative use of the stable isotope ($\delta 180$) variability and therefore a Ca2+-derived			
97	temperature profile could provide essential information for a better constrain of			
98	temperature variability in the deepest (oldest) section of the two ice cores. This point			

99 should be highlighted more considering, however, that: i) Ca2+ sensitivity to temperature 100 changes might be, and it is likely to be, non-stationary as well over the last 1000 yrs; ii) the 101 relationship between Ca2+ and temperature could very well derive from post-depositional 102 processes. This last point is particularly relevant (also considering that NH4 show a similar 103 temperature dependance) and the authors should elaborate more on why they think this is 104 not the case. For example, if there is any data available of density, DEP or occurrence of melt 105 layers, I suggest that the authors should use these data to back up some of their assumption 106 regarding the summer-signal preservation by consolidation and its relationship with the 107 seasonality of Ca2+.

108 We thank the referee for this comment, in particular for the suggestion to include

109 considering the density profile of the core. As discussed in the initial remarks, considering

110 the reviews we realized that a few issues need to be clarified, and see some of these points

arising here, too. In fact, we believe that post-depositional processes must be considered

112 when explaining the apparent coupling between temperature and long-term variability of

113 Ca2+. We have clarified and extended our discussion of this point. The comparison between

density and Ca2+ data clearly shows that dust-rich layers are coinciding with locally

115 enhanced density, that stem from fast snow consolidation. This "self-preserving"

116 characteristic of Ca2+ (and other dust-related species) against wind erosion is one of the

117 main differences with respect to the stable isotope signal. We have also added text to

118 discuss the fact that, while a non-stationary character of the Ca2+-temperature relationship

119 is certainly a possibility, we find no evidence for this within the instrumental period (in

120 contrast to the stable isotopes). Following the referees comment we have also elaborated

121 that, for this reason, fundamental shortcoming exists in quantitatively interpreting the

122 isotope-thermometer over long time scales at CG. Although we do not intend to introduce

123 Ca2+ as a new general temperature indicator, we see our findings as a strong indication of

124 the potential for using the long-term variability of Ca2+ as a *site-specific* temperature proxy.

125 We have clarified this view also in our conclusions.

126 **Changes to manuscript**:

Page 4, Lines 4ff.: Rewrote part of this section accordingly. Specifically regarding the
 motivation for expecting a temperature-related imprint in Ca2+.

Page 18, Lines 4ff.: Moved and rewrote part of the paragraph (originally in section
2.2), specifically mentioning the self-preserving character of Ca2+.

• Page 13, Lines 14ff: Included additional mentioning of the shortcomings of the

- 132 stable isotope thermometer at CG.
- Page 19, Lines 13ff.: Included a statement to clarify the lack of evidence for a non stationary Ca2+-temperature relationship.
- Page 18, Lines 21ff: Emphasized the site-specific role of the Ca2+-temperature
 association.
- 137

138Furthermore, the assumption that the Ca2+ signal is almost entirely expression of a dust

- 139 input from Saharan region is not enough justified in the text. The fact that the Ca2+ profile
- 140 might derive from both wet and dry deposition and both proximal and distal sources cannot
- 141 be ruled out from the data shown in the manuscript. Since the isotope/impurity co-
- 142 variation on the inter-annual scale is mainly related to changes in the amount of winter
- 143 precipitation contributing to annual mean values, I think is necessary to briefly consider
- 144 different scenarios concerning the (although marginal) role of dry deposition in the Colle
- 145 Gnifetti area and how these could change the Ca2+ signal in the different cases.
- 146 We would like to refer here to the initial comments and point out that at CG, mineral
- background aerosol levels are generally low (including summer) and the Ca2+ record is
- 148 dominated by inputs of Saharan dust, which has been demonstrated in previous studies
- 149 (Wagenbach et al. 1996). Thank you also for pointing out the role of dry deposition, which
- 150 we have so far not explicitly mentioned in the manuscript.
- **151 Changes to manuscript:** Page 4, Lines 13ff.: We have included a brief discussion of the
- 152 contribution made by dry deposition to the mineral dust content at CG.
- 153
- 154 While provenance studies (Sr and Nd isotopes for example) go beyond the scope of the
- 155 work, I think a more detailed discussion on the comparison of the insoluble dust profile vs
- the Ca2+ profile is necessary to utilize the calcium signal a proxy for Saharan dust input.
- 157 We agree with the referee that a provenance study based on isotopic trace element analysis
- 158 exceeds the scope of this study. At the same time the identification of Saharan dust input
- based on Ca2+ (and an alkalinity measure) has already been established in a previous study
- 160 (Wagenbach et al. 1996). Thus we did not intend to develop a new (and arguably more
- 161 precise) proxy for Saharan dust events at CG, but intended to use this already established
- tool. We have clarified this in the respective introductory section 2.
- 163 **Changes to manuscript:** Page 4, Lines 26ff.: Added a clarifying statement regarding the
- 164 tool to identify Saharan dust events.

166 Whether Saharan dust-Ca2+ data is a reliable proxy for palaeotemperature is yet again 167 another point that needs to be better illustrated in the text. I think the authors should 168 provide more justification regarding why the Ca2+ variability is mainly related to 169 temperature changes and not, for instance, to changes at the dust source (Saharan desert). 170 As outlined above it is not our intention to directly link the Saharan-dust component of the 171 Ca2+ data (spikes) to temperature, but rather investigate for this purpose the long-term 172 variability of Ca2+. We find the Ca2+ trends in surprisingly good correlation with 173 instrumental temperature throughout the full instrumental period, and go on to discuss 174 how snow preservation plays a decisive role in introducing this Ca2+-temperature coupling. 175 It seems likely that only large and systematic changes at the dust source would change the 176 long-term Ca2+ variability, or eventually override the coupling to temperature. On the other 177 hand, these changes (e.g. increased dust mobilization) would likely also influence the 178 Saharan dust spikes and their frequency of occurrence. However, the only instance where 179 we find an outstanding according feature is the increased dust occurrence in the medieval 180 period of our record. We thank the referee for this suggestion and now consider this issue 181 in our discussion. 182 Changes to manuscript: Page 19, Line 18: Added text to discuss the role of changes at the 183 dust source. 184 185 Detailed comments: 186 Page 1 Line 1-2: I would update this statement in view of the recent 7000-yrs long ice core 187 record from the Ortles (Gabrielli et al., 2017). 188 Changed accordingly to clarify. In contrast to Ortles, Colle Gnifetti is a non-temperate site. 189 190 Page 3 Line 11-12: "which prevents any link of the climatologic precipitation rate to the net 191 snow accumulation rate". I am not sure I understand here: Does this mean that the 192 seasonality in the proxies is not governed by accumulation rate? Or is rather the longer-193 time variability? In any case I suggest changing the word "prevents" with "limits". 194 What we intend to say is that due to the highly variably snow deposition at CG, it is not 195 possible to infer precipitation changes based on e.g. annual layer thickness (e.g. as done 196 with Greenland ice cores). We have clarified the wording accordingly. 197 Changes to manuscript: Page 3, Line 10-11: "limits linking the net snow accumulation rate

198	to the climatologic precipitation rate"
199	
200	Page 3 Line 17: I found the wording a bit confusing. What "chemical/isotopic conditions"
201	means? Do you mean chemical and isotopic signatures?
202	Yes we mean the signature of chemical and isotopic species measured in the CG ice cores.
203	We have clarified the wording accordingly.
204	Changes to manuscript: Page 3, Line 16: "chemical and isotopic signatures "
205	
206	Page 4 Line 1-2: "the isotope/impurity co-variation on the inter-annual scale reflects to a
207	large degree changes in the amount of winter precipitation contributing to annual mean
208	values" I think is important here to highlight why the authors think dry deposition is
209	playing a marginal role.
210	See our response above, we now include a short discussion of the role of dry deposition.
211	Changes to manuscript: Page 4, Line 14ff.: Added text regarding dry deposition.
212	
213	Page 4 line 10-11:"Therefore, the Ca2+ record of the CG ice cores is primarily related to
214	mineral dust and dominated by Saharan dust". It's hard to tell without provenance studies. I
215	suggest using "dominated by dust, most likely originating in the Saharan desert".
216	Thank you, we have reworded this statement and included the respective reference.
217	Changes to manuscript: Page 4, Line 13-14: Reworded the previous statement.
218	
219	Page 7 Line 3: "Deviations from a CPP of 50% indicate higher or lower contribution of large
220	and small particles respectively". You have to exclude local sources of dust then if you want
221	to use the threshold to distinguish Saharan dust layers. I would add a sentence justifying
222	this.
223	Thank you. We now point out the findings of Wagenbach and Geis (1988) in this context,
224	who showed that Saharan dust in fact differs in volume size distribution in comparison to
225	local and background sources.
226	Changes to manuscript: Page 5, Line 16: " The threshold was chosen such that it
227	corresponds to the expected median particle diameter of Saharan dust particles at CG,
228	which was shown to be distinguishable from background sources"

230	Page 8 Line 1: I would specify what "Ca signal" means. Is it Intensity in counts per second?
231	Or total counts? Please add this also to the relevant figures.
232	Thank you for pointing this out- in this case it is in fact intensity in counts per second,
233	although it is possible to achieve an according calibration of the LA-ICP-MS signal (Sneed et
234	al. 2015).
235	Changes to manuscript: Added text to captions of Figures 2 and 3, respectively.
236	
237	Page 9 Line 8: "Below 26 m WE the identification of annual layers became ambiguous and
238	was abandoned". Maybe I missed this information, but why then LA-ICPMS was not
239	performed on the KCI core? Please provide justification, if it is not provided somewhere
240	else.
241	There was actually a pilot study for LA-ICP-MS performed on KCI (Sneed et al. 2015),
242	however, not targeting yet the identification and counting of annual layers. Given the
243	sophisticated and time-consuming nature of LA-ICP-MS we have so far only analysed KCC in
244	a continuous manner. We take the comment as encouragement to further pursue the LA-
245	ICP-MS analysis, potentially revisiting KCI in the future. We have added text to provide this
246	information.
247	Changes to manuscript: Page 8, Lines 10ff.: Added text regarding LA-ICP-MS on KCI.
248	
249	Page 13 Line 12: "due to the strong effect of isotope diffusion at CG, inter-annual or even
250	seasonal isotope variability is effectively eliminated". What about Ca2+ diffusion? While
251	dust does not diffuse, the contribution of soluble particles to the Ca44 signal should be
252	briefly addressed too, together with their possible diffusion.
253	The effect of diffusion is certainly smaller for Ca2+ than for the stable water isotopes, as we
254	do not see any evidence of diffusion hampering the identification of the annual layers at the
255	high resolution afforded by LA-ICP-MS. However, we are now mentioning this effect, and in
256	particular also point out the contribution of soluble Ca to the LA-ICP-MS signal.
257	Changes to manuscript:
258	• Page 6, Line 13: "The 44Ca signal comprises contributions of soluble and insoluble
259	Ca"
260	• Page 9, Line 16-17: "The annual layer signal remains clearly identifiable for the
261	remaining part of the depth-range investigated here (e.g. apparently not affected by
262	diffusion of soluble Ca)"

264	Page 14 Line 31-32: "From a preliminary inspection of snow pit data recently obtained for
265	the KCI-KCC flow line, there is no clear indication of a systematic trend in mean $\delta 180$ levels
266	upstream of KCC, however." It might be worthy to consider adding a plot (at least in the
267	supplementary material) showing this.
268	As mentioned in the text the detailed investigation of the isotope-upstream effect is still
269	ongoing based on sophisticated 3D-flow modelling (PhD thesis Carlo Licciulli at Heidelberg
270	University). However, we have provided additional information regarding the preliminary
271	inspection.
272	Changes to manuscript: Page 13, Lines 24ff.: Added text accordingly.
273	
274	Page 15 Line 12: "higher sensitivity values for KCI than KCC, revealing 2.3 vs. 1.4 $\%$ /°C,
275	respectively".
276	This discrepancy seems surprisingly high even considering the difference in accumulation
277	rate that you correctly highlight. Could it be related also to the strong isotope diffusion at
278	CG?
279	The degree of isotope diffusion could certainly be another difference between KCI and KCC,
280	thank you for pointing this out. This is especially so in the firn section, and here (due to the
281	difference in accumulation rate) the age interval represented by the firn column differs for
282	the two cores. Although it is difficult at this stage to give a more quantitative evaluation
283	regarding its effect on isotope sensitivity, we now include mention isotope diffusion. We
284	will also consider this in a potential future investigation on the enhanced isotope sensitivity.
285	Changes to manuscript: Page 14, Line 13-14: Added text accordingly.
286	
287	Page 20 Line 10-17: This entire section seems a bit far-fetched. As the authors said, the
288	summer-bias signal at CG strongly advocate against a NAO imprint on the KCC and KCI
289	temperature reconstruction. I suggest adding few more considerations to justify this link or
290	remove the entire section.
291	After considering the comments of both referees in this direction, we decided to remove
292	this section from the discussion.

- Page 21 Line 1-20: I suggest to the authors to add a sentence outlining the feasibility of
- using Ca2+ records for temperature reconstruction in other alpine site, or generally in other
- low accumulation ice core site.
- 297 We are now generally trying to be more clear about the site-specific nature of the potential
- temperature significance of the Ca2+ long-term variability. However, it would be interesting
- to test if the Ca2+-temperature association observed at CG holds also at other alpine sites.
- **300 Changes to manuscript:** Page 20, Line 28: Included a respective statement in the
- 301 conclusions.
- 302
- 303 References
- 304 Gabrielli, P., Barbante, C., Bertagna, G., Bertó, M., Carturan, L., Dinale, R., & Seppi, R. (2017,
- April). 7000 year European climate record from the Ortles ice core. In EGU General
- 306 Assembly Conference Abstracts (Vol. 19, p. 9932).
- 307

Temperature and mineral dust variability recorded in two low accumulation Alpine ice cores over the last millennium

Pascal Bohleber^{1,2,3}, Tobias Erhardt^{4, 5}, Nicole Spaulding¹, Helene Hoffmann², Hubertus Fischer^{4, 5}, and Paul Mayewski¹

¹Climate Change Institute, University of Maine, Orono, Maine, USA

²Institute of Environmental Physics, Heidelberg University, Heidelberg, Germany

³Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences, Innsbruck, Austria

⁴Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland

⁵Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

Correspondence to: Pascal Bohleber (Pascal.Bohleber@iup.uni-heidelberg.de)

Abstract. Among ice core drilling sites in the European Alps, Colle Gnifetti (CG) is the only non-temperate glacier to offer climate records back to at least 1000 years. This unique long-term archive is the result of an exceptionally low net accumulation driven by wind erosion and rapid annual layer thinning. However, the full exploitation of the CG time series has been hampered by considerable dating uncertainties and the seasonal summer bias in snow preservation. Using a new core drilled in 2013 we

- 5 extend annual layer counting, for the first time at CG, over the last 1000 years and add additional constraints to the resulting age scale from radiocarbon dating. Based on this improved age scale, and using a multi-core approach with a neighboring ice core, we explore the time series of stable water isotopes and the mineral dust proxies Ca^{2+} and insoluble particles. Also in our latest ice core we face the already known limitation to the quantitative use of the stable isotope variability based on a high and potentially non-stationary isotope/temperature sensitivity at CG. Decadal trends in Ca^{2+} reveal substantial agreement
- 10 with instrumental temperature and are explored here as a potential site-specific supplement to the isotope-based temperature reconstruction. The observed coupling between temperature and Ca^{2+} -trends likely results from snow preservation effects and the advection of dust-rich air masses coinciding with warm temperatures. We find that if calibrated against instrumental data, the Ca^{2+} -based temperature reconstruction is in robust agreement with the latest proxy-based summer temperature reconstruction, including a "Little Ice Age" cold period as well as a medieval climate anomaly. Part of the medieval climate period around
- 15 1100–1200 AD clearly stands out through an increased occurrence of dust events, potentially resulting from a relative increase in meridional flow and/or dry conditions over the Mediterranean.

1 Introduction

Glaciers and ice caps of high mountain ranges can provide climate records of mid- and low latitudes complementary to polar ice cores. In comparison to their polar counterparts, mountain drilling sites are characterized by a comparatively small-scale

20

glacier geometry and their proximity to continental source areas. As a consequence, cold mountain glaciers are an especially worthwhile target for ice core studies focusing on Holocene climate, e.g. in view of the envisaged IPICS 2k array (Brook et al.,

2006) and the present underrepresentation of ice core records contributing to the PAGES 2k Network (Ahmed et al., 2013). In the European Alps, ice core studies have been performed at Col du Dôme, Mont Blanc (Preunkert et al., 2000), Fiescherhorn, Bernese Alps (Schwerzmann et al., 2006), Ortles, Eastern Alps (Gabrielli et al., 2016) as well as at Colle Gnifetti and Colle del Lys in the Monte Rosa region (e.g. Wagenbach et al., 2012, and references therein). Among these glaciers, Colle Gnifetti

- 5 (CG) in spite of its limited glacier depth stands out as the only non-temperate site where net snow accumulation is low enough to provide records over the last millennium and potentially beyond at a reasonable time resolution. The exceptionally low net accumulation at CG is a result of seasonal net snow loss by wind erosion: Since snow consolidation is most effective during the summer half year, winter precipitation is more likely to be removed from the surface (Wagenbach, 1992). This has far-reaching consequences with respect to the interpretation of the CG ice cores, hampering to-date the full exploitation of their
- 10 unique long climate time series. On the one hand, considerable uncertainty in the individual ice core chronologies becomes an obstacle already after a few hundred years. Difficulties in deploying annual layer counting as the main dating tool arise from snow scouring, rapid layer thinning associated with strongly non-linear time-depth relationships and the extremely low time resolution achieved in the bottom part of the glacier by conventional cm-resolution analyses. As a consequence, dating the deeper part of CG ice cores is commonly based on simple extrapolation combined with constraints from radiocarbon analysis
- 15 (e.g. Jenk et al., 2009). On the other hand, irregular and summer-biased snow deposition makes the annual or long-term levels of ice core proxy signals with a prominent seasonal cycle a primary function of the relative winter snow fraction preserved, as opposed to their common climatological meaning. In addition, net snow accumulation is characterized by substantial spatial and temporal variability, leading to considerable influence of upstream flow effects and depositional noise (Wagenbach, 1992). In contrast to the strong signals of anthropogenic aerosol increase, depositional noise especially challenges the detection of the
- 20 comparatively weak stable water isotope trends (δ^{18} O and δ D). Under these circumstances, the comparison of multiple cores drilled at the same site can be used to identify an atmospheric signal as shared variability among the cores (Bohleber et al., 2013; Wagenbach et al., 2012).

Here we present new results to tackle the two-fold challenge above with a new core drilled at Colle Gnifetti in 2013, integrating datasets from an additional ice core drilled in 2005 on the same flow line. In order to obtain a reliable long-term chronology for

- 25 the 2013 core, we utilize state-of-the-art continuous flow analysis for ice core impurity profiling and, to identify even highly thinned annual layers, laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) at sub-mm depth resolution (Della Lunga et al., 2017; More et al., 2017; Haines et al., 2016; Mayewski et al., 2014). We combine annual layer counting in the resulting impurity profiles with absolute age constraints from radiocarbon analysis, taking advantage of recent progress in applying this technique to mountain ice cores (Hoffmann et al., 2017b; Uglietti et al., 2016). Based on a refined long-term
- 30 chronology, the time series of stable water isotopes and mineral dust proxies (Ca^{2+} and insoluble particles) are investigated, with special emphasis on their relation to temperature.



Figure 1. The ice core array at Colle Gnifetti, at 4450 m asl in the Monte Rosa summit range. The drilling sites of the two cores KCI and KCC are located on approximately the same flow line (black line) towards the eastern flank (downwind of the main wind direction), hence providing their same upstream catchment area. Locations of previous drillings initiated by the Institute of Environmental Physics are also shown as small black dots for reference.

2 Glaciological settings of the CG drilling site

Details on the glaciological features of CG are described thoroughly in the literature; e.g. Haeberli et al. (1988); Lüthi and Funk (2000); Konrad et al. (2013) for geometry and glacier flow, Haeberli and Funk (1991); Hoelzle et al. (2011) for englacial temperature and Alean et al. (1983) for surface accumulation. Here, we present only a brief overview, mainly dedicated to

5 explaining the role of snow deposition in relation to recording atmospheric temperature and mineral dust variability in the CG ice cores.

With a horizontal scale of 400 m and a maximum ice thickness of around 140 m, the CG site forms a small firn saddle at around 4500 m asl between two summits of the Monte Rosa massif. The orientation of the convex, central saddle axis coincides with the main westerly wind direction, thereby making the downwind-situated ice cliff a perfect sink for drifting snow

- 10 (Figure 1). Hence, a substantial fraction of the annual fresh snow precipitation is removed at CG, which limits linking the net snow accumulation rate to the climatologic precipitation rate. The latter ranges from 0.15 m water equivalent (WE) per year in the north-facing flank to about 1.2 m WE per year in the southern one, where the higher abundance of ice layers and ice crusts significantly reduces the snow erosion rate (Alean et al., 1983). Within the CG north flank (comprising our CG ice core array) fresh snow consolidation is faster during the summer half year (additionally supported by refreezing surface melt). Accord-
- 15 ingly, the mean net snow accumulation is mainly made up by precipitation of the warm seasons, which entails a systematic over-representation of the summer half-year in chemical and isotopic signatures (Wagenbach, 1989). A changing amount of winter precipitation contributing to annual mean values may introduce a coupling on the inter-annual scale among seasonal varying signals, including δ^{18} O and most impurities (Wagenbach, 1992). Notably this also includes a potential link to temper-

ature, since warm summers feature increased vertical mixing and hence a higher atmospheric impurity load. In addition, faster fresh snow consolidation favored by higher temperatures may lead to an increased relative amount of impurity-rich summer snow deposition.

A long-term co-variation between δ^{18} O and Ca²⁺ suggesting a possible relationship between climate and dust deposition at CG

- 5 has already been noted but was left for future investigation (Wagenbach and Geis, 1989; Wagenbach et al., 1996). A later study specifically explored the link between the δ¹⁸O signal and air temperature changes in the presence of the snow preservation influence at CG. A dominant influence of atmospheric temperature on decadal isotope variability shared among the CG cores was found, although a high and potentially non-stationary isotope/temperature sensitivity hampered the quantitative use of the CG isotope variability (Bohleber et al., 2013). Considering the co-variation of the long-term variability of i) Ca²⁺ and δ¹⁸O,
- 10 and ii) δ^{18} O and temperature, suggests that atmospheric temperature variability could also be reflected in the Ca²⁺ trends. In view of the shortcomings in quantitatively using the isotope-thermometer at CG, identifying a temperature-related imprint in the Ca²⁺ variability could provide a valuable supplement in this respect.

At CG, mineral background aerosol levels are generally low, making the Ca^{2+} record is dominated by episodic inputs of dust, most likely originating in the Saharan desert (Wagenbach et al., 1996). While dry deposition may add to the average mineral

- 15 dust content (Haeberli et al., 1983), it appears less important in case of Saharan dust events (Schwikowski et al., 1995). In addition, only a marginal contribution to changes in the particle size distribution is expected from changes in the dry deposition (Ruth et al., 2003). As for most of the impurity species at CG, the seasonal contrast in Ca²⁺ concentration is primarily connected to the seasonal gradient in vertical atmospheric mixing, with an additional component from sporadic Saharan dust inputs (Preunkert et al., 2000; Preunkert and Wagenbach, 1998). Saharan dust deposition events are a frequent phenomenon
- 20 in the Alps with main occurrence in spring and summer (Prodi and Fea, 1979). A single deposition event typically lasts less than a few days (Sodemann et al., 2005; Schwikowski et al., 1995). The associated warm air temperature and the substantially lowered snow albedo both support surface snow consolidation and partly protect the dust layer from wind erosion (Haeberli et al., 1983). Intensive Saharan dust events of the summer half year, associated with directly northward transport of air masses, are most likely to become preserved at CG. Saharan dust layers in CG ice cores can be characterized by high concentrations
- of insoluble particles, SO_4^{2-} and Ca^{2+} coinciding with buffered low acidity, as well as to some extent by increased $\delta^{18}O$ and deuterium excess values (Wagenbach et al., 1996; Wagenbach and Geis, 1989). Accordingly, the combination of Ca^{2+} with an alkalinity measurement is a specific tool to identify Saharan dust influenced layers in CG ice cores (Wagenbach et al., 1996), which will be employed in the following.

The above considerations warrant a general distinction and separate evaluation of the following two features of the Ca^{2+} record

30 of the CG ice cores: i) The long-term average Ca²⁺ concentration, and its potential coupling with δ^{18} O and temperature via snow preservation. ii) Spikes in Ca²⁺ typically two orders of magnitude above background, which are dominated by Saharan dust input (Wagenbach et al., 1996). Regarding ii), changes in the dust peak occurrence rate can originate from changes in the meridional versus zonal circulation and/or in the desert dust source strength. Here detecting the frequency of dust peaks in the ice core matters, which is expected to be comparatively more robust against snow preservation influence.

Table 1. Basic glaciological parameters of the two CG ice cores

Core name	KCI	КСС
Position GPS (WGS84)	N 45.92972 E 7.87696	N 45.92893 E 7.87627
Year of drilling	2005	2013
Total depth [m abs]	61.84	71.81
Total depth [m WE]	48.44	53.77
Surface net accumulation [cm WE/yr]	14	22
Firn-ice-transition [m WE]	17	21

3 Ice core analysis

The two cores used in this study, denoted as KCI and KCC, were drilled in 2005 and 2013, respectively. Both cores were drilled roughly on the same flow line, making them the natural choice for our inter-core comparison, i.e. opposed to using previously deep cores drilled on another flow line (Figure 1). Table 1 summarizes basic glaciological parameters of the two cores. The depth sections used in this study were chosen to comprise roughly the last 1000 years, i.e., the upper 44 m WE (corresponding to 81% relative depth) and 35 m WE (73% relative depth) of KCC and KCI, respectively. Table 2 provides an overview of the carefully co-registered datasets used in this study. The various methods of analysis are discussed briefly in the following.

3.1 Impurity profiles from continuous flow analysis

 Continuous flow analysis (CFA) of the KCC core was performed with the setup at the Division for Climate and Environmental
 Physics, Physics Institute, at the University of Bern. Analyses performed on the meltwater flow included meltwater conductivity, insoluble particle concentration and size distribution as well as selected ion species (Ca²⁺, NH₄⁺, NO₃⁻, Na⁺, see Table 2). In addition, stable water isotopes were analyzed using a Picarro instrument coupled directly to the meltwater flow. The size distribution of insoluble particles recorded by the optical particle sensor was used to derive a profile of the "coarse particle percentage" (CPP). The CPP was calculated based on particle volume, and represents the percentage of particles exceeding a

- 15 threshold of 4.0 μ m. The threshold was chosen such that it corresponds to the expected median particle diameter of Saharan dust particles at CG, which was shown to be distinguishable from background sources (Wagenbach and Geis, 1989). Deviations from a CPP of 50% indicate higher or lower contribution of large and small particles respectively. The melt rate was adjusted to provide the necessary amount of water for all analyses resulting in an effective depth resolution ranging from 1.2 cm at the very top of the core to about 0.5 cm for all depth below approximately 25 m WE. Electrical conductivity measurements (ECM)
- 20 performed at the Institute of Environmental Physics, Heidelberg University were used primarily to obtain a qualitative record

5

of the acidity of the ice in connection to the detection of Saharan dust events. The KCI core was analyzed using the reduced CFA setup at the Institute of Environmental Physics, Heidelberg University.

Meltwater conductivity and insoluble particle concentration were measured by CFA at about 0.7 cm effective resolution. Con-

Table 2. Overview on ice core analyses and datasets used in this study

Core	Parameters	Sampling	Effective resolution [cm]
KCC	Meltwater Conductivity, NH_4^+ , NO_3^- , Na^+	Continuous Flow	> 0.5
	Insoluble Particles, Ca ²⁺	Continuous Flow	> 0.5
	Stable water isotopes (δ^{18} O and δ D)	Continuous Flow	> 0.5
	Electric Conductivity	ECM	> 0.5
	⁴⁴ Ca	Laser ablation ICP-MS	$120~\mu{\rm m}$
KCI	Meltwater Conductivity, Insoluble Particles	Continuous Flow	> 0.7
	Stable water isotopes (δ^{18} O or δ D)	Discrete Sampling	10 - 1.5

tinuous sub-sampling of the core for stable water isotope analyses was conducted at a depth resolution typically ranging between 5 and 10 cm. Due to the relatively high firn temperature at CG, isotope smoothing is much faster compared to polar sites with similar annual layer thickness. Hence re-sampling most of KCI even at 1.5 cm depth resolution did not significantly restore any high-frequency isotope variability (Bohleber et al., 2013).

5 3.2 Ultra-high resolution Ca-profile of the KCC core by laser ablation ICP-MS

Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) was conducted in the WM Keck Laser Ice Facility at the Climate Change Institute (University of Maine) and used to analyze ⁴⁴Ca at ultra-high depth resolution (better than 120 μ m). The more abundant ⁴⁰Ca is blocked by mass interference from ⁴⁰Ar used as carrier gas. Details regarding the method, sample preparation and calibration routine can be found in Sneed et al. (2015). Briefly, the components of this system

- 10 include a Thermo Element 2 ICP-MS, a New Wave UP-213 laser, and a cryo-cell chamber, designed to seal a 1 m ice core from the surrounding air while maintaining a uniform temperature of -15° C. In order to ensure a complete seal of the ablation chamber, porous firn parts could not be measured. From 29.5 m WE to bedrock, the KCC ice core was analyzed for ⁴⁴Ca along a single ablation track. The ⁴⁴Ca signal comprises contributions of soluble and insoluble Ca (Sneed et al., 2015). Crucial for further deployment for annual layer counting, the trend components in the LA-ICP-MS measured Ca signal have been shown
- 15 to be in good correspondence with the lower resolution CFA Ca signal, as shown in Figure 2 and previously by Sneed et al. (2015).

3.2.1 Radiocarbon analysis

The measurements for radiocarbon dating of the ice core have been conducted at the Institute of Environmental Physics (Heidelberg, Germany) under close collaboration with the accelerator mass spectrometer (AMS) facility at the Klaus-Tschira-Lab

20 in Mannheim, Germany. The microscopic particulate organic carbon fraction (POC) incorporated into the ice matrix was extracted, combusted and analyzed for ¹⁴C content. Calibration of the retrieved ¹⁴C ages was performed using OxCal version 2.4



Figure 2. Ca signals obtained from the KCC ice core at around 65% relative depth using LA-ICP-MS and CFA in direct comparison after careful alignment of the two depth scales. Top row: Raw (black) and filtered LA-ICP-MS Ca signal (blue). Bottom: CFA Ca (red) vs filtered LA-ICP-MS Ca signal (blue). Note i) additional peaks and high-frequency information revealed by LA-ICP-MS, ii) a general agreement of CFA and low frequency LA-ICP-MS components is consistently observed over core parts measured by LA-ICP-MS. LA-ICP-MS intensity is reported as counts per second.

(Ramsey, 2016) and by convention the 1-sigma error range is shown (Stuiver and Polach, 1977). For details on the sample preparation and measurement procedure see Hoffmann et al. (2017a, b); Hoffmann (2016). The average ice sample masses were for both cores in a range of ca. 300–500 g ice resulting in absolute POC masses below 10 μ gC. For the KCC core, a fraction of the ice core with a cross section of 17 cm² was reserved for the POC ¹⁴C analysis. Within the upper 44 m WE, a

5 total of six samples were analysed, typically comprising between 40–60 cm of core. For the KCI ice core more core material (one third) was available, resulting in depth intervals of 40 cm length used for radiocarbon dating. Within the upper 40 m WE of the KCI core five samples have been analyzed so far.

4 Ice core dating

Ice core chronologies were established by annual layer counting as the main dating tool in combination with additional age constraints from ¹⁴C for the lower core parts. For roughly the last 100 years, dated time horizons (1963 bomb-radioactivity, and the Saharan dust layers of 1977, 1947 and 1901, cf. Figure 4 below) are available to constrain the counting (Table A1 with the depths of the respective horizons for KCC and KCI is included in the appendix). The 1963 horizon was used to cross-check that the annual signal had been identified correctly (cf. sub-annual and multi-year signals). The dust events were independently used for verification and typically lie within one to two years of the counted age scale (four years at maximum

15 for the 1901 horizon). Regarding additional absolute age markers beyond 1901, the identification of volcanic eruptions solely based on basic ice chemistry profiles is not feasible at CG. This is due to the fact that the relatively weak signals of volcanic



Figure 3. Examples for annual layer counting in KCC impurity profiles for three different depth sections, labeled a), b) and c), and corresponding roughly to 100, 250 and 1000 years before 2013, respectively (cf. Table 3). In the upper core parts (firn sections) CFA measured impurities were used for counting, with special emphasis on NH_4^+ (a). Counted years are marked as full stars, uncertain years as white stars. The middle row (b) shows an example of overlap in counting between CFA and LA-ICP-MS Ca, showing (10 ± 3) and (11 ± 3) years, respectively. The LA-ICP-MS Ca raw signal is shown in black together with Gaussian smoothing (blue). Note that a minor depth offset (at most a few cm) may exist between the CFA and LA-ICP-MS datasets. Accordingly, no one-to-one match of the individual peaks is attempted. Counting within one of the deepest sections analysed for this study is shown in (c). Here, only the LA-ICP-MS Ca allows a reliable identification of almost sub-cm thin annual layers. LA-ICP-MS intensity is reported as counts per second.

sulphate or volcanic acidity are easily overlooked at CG since they are embedded into the relatively large variability of Saharan dust associated sulphate (mainly from gypsum) and (acidity consuming) carbonate. More promising in this respect is the investigation of relatively volatile trace elements (Kellerhals et al., 2010b), or the detection of tephra markers, which are beyond the scope of this work, however.

5 4.1 The KCI chronology

For KCI, insoluble particle concentration and meltwater conductivity were used for annual layer counting, extending down to about 26 m WE. Below 26 m WE the identification of annual layers became ambiguous and was abandoned. This depth corresponds (taking the uncertainty in layer counting into account) to (1492 ± 30) AD. A two-parameter model (based on a simple analytical expression for the decrease of the annual layer thickness with depth) was used to extrapolate a continuous age-depth relation to greater depth (Nye, 1963; Jenk et al., 2009). Note that high-resolution annual layer counting could only be performed in KCC (see below) since only small sections of KCI have been analyzed by LA-ICP-MS so far (Sneed et al., 2015).

4.2 The KCC chronology

15

10

All impurity species measured by CFA (Table 2) were used in combination for annual layer counting. Annual layers were defined as local maxima in at least two of the six impurity signals, with special emphasis on NH_4^+ featuring the largest seasonal amplitude. An example of counting annual layers in the CFA profiles in shown in Figure 3 a). In order to identify highly

Core	Depth [m WE]	Age [year AD]	Uncertainty [years]	Annual Layer Thickness [cm WE]
KCC	10	1971	1	18
	20	1912	4	11.5
	30	1762	12	2.7
	40	1000	72	1
KCI	10	1917	4	7.4
	20	1700	20	3.2
	30	1312	62	1.7
	35	939	77	1

Table 3. Ice core age, dating uncertainty and annual layer thickness for selected depths

thinned, sub-cm annual layers expected to dominate the deeper core sections, an independent counting was established using the LA-ICP-MS Ca profile starting at 29.5 m WE (corresponding to 1760 AD). At this depth, the average annual layer thickness was estimated from CFA-based counting as around 3 cm. The LA-ICP-MS Ca record was investigated at full resolution and as a smoothed version (using the leading components in singular spectrum analysis or Gaussian smoothing). In its upper section,

- 5 the LA-ICP-MS Ca profile is characterized by regular occurrence of several distinct peaks grouped together going along with an elevated baseline of Ca concentration (Figure 3 b)). The groups of peaks are separated by a comparatively stable signal of low Ca concentrations. The latter is interpreted as resulting from the varying degree of winter snow being included in the record otherwise dominated by summer snow. Accordingly, the grouped peaks correspond to sub-annual snow deposition events of elevated Ca concentration during the summer period, which is also observed in the most shallow parts of KCC (see Figure
- 10 A1 in the appendix). For the depth interval 29.5–32.5 m WE, counting separated groups of peaks (typically 3–5 peaks per annual layer) in the LA-ICP-MS Ca record results in good agreement with counting performed on the CFA profile (typically within ±1 year per 10 counted years, see Figure 3 b)). Below 32.5 m WE, average annual layer thickness becomes close to 1 cm and counting in the CFA profile becomes increasingly difficult (i.e. frequent "shoulder type" annual layers merged into a single impurity peak). The LA-ICP-MS Ca profile continues to show distinct groups of peaks that become increasingly closely
- 15 spaced and eventually merge into single broad peak events (Figure 3 c)). Accordingly, LA-ICPMS Ca was the dominant source of annual layer counting after around 32.5 m WE (1600 AD). The annual layer signal remains clearly identifiable for the remaining part of the depth-range investigated here (e.g. apparently not affected by diffusion of soluble Ca).

4.3 Age constraints from radiocarbon analysis

20

For KCC, results from ¹⁴C analysis are found to back the annual layer counted age scale. Five of six ¹⁴C dates agree with the counting within their 1-sigma range (Figure 4), corresponding to a root mean square deviation of 118 years (227 years including the outlier). The outlier ¹⁴C point contradicts a monotonic increase of age with depth and is thus disregarded. This is justified, because the ¹⁴C age of this sample matches with a very sensitive section of the ¹⁴C-calibration curve. Therefore

already a small, unknown blank contribution would be able to shift the calibrated age of this sample significantly. Within the 2 sigma error range it also hits the error range of the annual layer counting chronology in the present configuration. Its deviation is therefore not of consequence.

For the KCI ice core, the radiocarbon ages are found to agree with the extension of the existing age scale based on the two-

- 5 parameter model. It seems worth noting, however, that four out of five 14 C points lie systematically above the extrapolated age scale (albeit in agreement within their 1-sigma range). Only the sample at 28.4 m WE shows an age that is significantly older than expected. This can on one hand be due to the extremely small (also compared to the other KCI samples) sample size of only 2.2 µgC making this sample prone to even very small potential blank contributions. In this context also a potential influence of aged organic material (e.g. from Saharan dust) has to be regarded. At present, the age of this sample is therefore
- 10 regarded as an outlier. Additional radiocarbon measurements of this core section above and below the critical sample are planned to further refine the match, and to test if the systematic deviation of the ¹⁴C ages persists. The ¹⁴C ages of the KCC and KCI samples are summarized in the appendix, Table A2.

4.4 Dating uncertainty

Potential sources of uncertainty in annual layer counting stem from i) erroneously identifying or missing of existing annual

- 15 layers, ii) interpolating data gaps and iii) an incomplete stratigraphy missing years due to annual snowfall fully eroded from the surface. Regarding i), we estimated the likelihood of miscounting layers by marking "uncertain years" (Figure 3). In view of the high snow erosion at CG, uncertain layers were defined as additional peaks in close proximity to an annual layer (e.g. "shoulder type" peaks). To quantify counting uncertainty from uncertain layers, we followed the approach successfully employed for Greenland ice cores. This is to count uncertain layers as 0.5±0.5 years and to estimate the maximum counting error (MCE)
- from N uncertain layers as N x 0.5 years (Andersen et al., 2006; Rasmussen et al., 2006). With 144 uncertain layers detected within the upper 40 m WE of KCC, this corresponds to an uncertainty of \pm 72 years at 1000 AD. With respect to ii), the depth interval considered in this work was completely recovered without core loss. The ends of the CFA core sections were trimmed in case of irregular core breaks. This resulted typically in less than a centimeter of missing CFA data, thus not interfering with annual layer counting. The ECM profile was used as an alternative backup across these short CFA data gaps. Likewise, the
- 25 CFA data was used as an alternative indicator where the LA-ICP-MS profile was incomplete, which only concerned one major instance of missing LA-ICP-MS data between about 33.8–34.24 m WE. Contribution iii) constitutes a fundamental difference relative to Greenland conditions, since CG is not a closed system with respect to precipitation and loss of the annual snowfall in selected years can occur. The frequency of occurrence in these total snow loss events is, however, extremely hard to quantify. Counting annual layers in between the above mentioned (dust) hori-
- 30 zons within the last century, reveals an offset of typically only one to two years as compared to the known age of the horizons. Thus, the counting appears not to be systematically flawed by missing years. Hence we regard uncertainty i) as dominant and use the MCE as an uncertainty estimation of the KCC age scale. Notably, the uncertainty refers to the unconstrained counting approach used here and could be further refined in the future with new absolute dating horizons, especially for the per-1900 AD period.



Figure 4. Age-depth relations over the last 1000 years for KCC and KCI, shown in the top and bottom row respectively. Age is plotted on a logarithmic axis, together with the according estimates of maximum dating uncertainty (dashed lines) and 14 C age constraints (with 1-sigma range) for KCC and KCI. Also shown is the adjusted age scale of KCI based on the stable water isotope time series comparison (solid black line, within less than 15 years of the original dating and thus hardly distinguishable here, see text). Absolute dating horizons used roughly within the last 100 years (see text) are shown as black squares. Note that the KCI chronology is based on a simple extrapolation below 26 m WE and has large uncertainty beyond the last 1000 years (thus indicated as light gray line only).

The uncertainty of the KCI age scale was obtained in a consistent manner, using the MCE for the annual layer counted interval and extrapolating the upper and lower uncertainty limits with the two-parametric model. Figure 4 shows the resulting age-depth relation and uncertainty bands for KCC and KCI, together with ¹⁴C dates (shown with their 1–sigma uncertainty range) available for the respective depth interval. Table 3 gives a complementary summary of the age-depth relation, uncertainties and annual layer thickness for 10 m depth intervals. It is important to note that we are less confident about the age-depth relation of KCI compared to KCC, due to KCI featuring i) annual layering counting using only two bulk parameters and only down to

4.5 Inter-core time series comparison and age scale alignment

26 m WE and ii) the extrapolation by the two-parameter model.

10

5

To investigate potential offsets between the KCI chronology and the presumably more reliably dated KCC, we compared the stable water isotope time series of the two cores. This is motivated by the fact that the decadal isotope trends among the CG ice cores have been previously shown to agree over the last 250 years (Bohleber et al., 2013). Without substantial dating



Figure 5. Stable water isotope time series of KCC and KCI, shown in blue and red, respectively. The top row shows both records on their original time scale over the last 1000 years. The KCI age scale was adjusted using the algorithm of (Lisiecki and Lisiecki, 2002) to optimise the match with KCC.

offset between the cores, this inter-core agreement should hold also on longer time intervals. It is important to note that due to the strong effect of isotope diffusion at CG, inter-annual or even seasonal isotope variability is effectively eliminated. As a consequence, the records (except for the last 100 years in KCC) resolve only decadal-scale variability at best. Hence we did not apply any further smoothing to the time series. In order to avoid potential biases from increasing sampling resolution, both

5 time series were sub-sampled to nominal biennial resolution. Figure 5 shows the comparison of the respective time series on their original time scales for the last 1000 years.

The two original time series of KCC and KCI already feature striking similarities, although frequently separated by a lag between the two time series (e.g. note the distinct isotope minima around 1360 AD). The direction and magnitude of this lag varies with time, hampering an absolutely straightforward adjustment to match the two records. Aiming to adjust the KCI

- 10 record to KCC time series, we employed the powerful algorithm developed by Lisiecki and Lisiecki (2002) for correlating paleoclimate time series. In doing so, we left the last 150 years of the KCI age scale unchanged (since considered reliably dated) but did not prescribe any further user-defined tie-points to the algorithm. The result shows the original lag between the two time series eliminated (Figure 5). A maximum shift of around 15 years is needed to align the two records (e.g. around the 1360 AD isotope minimum), which is within the estimated dating uncertainty of KCI (Figure 4). As a result, the aligned
- 15 time series are significantly correlated (r = 0.47). This degree of correlation is within the typical range of correlating CG isotope time series on the decadal scale within the last 250 years (Bohleber et al., 2013). However, this is the first time that the correlation holds to this extent also for the comparatively old core sections of CG ice cores, e.g. we find a correlation coefficient of r = 0.50 when considering the interval 1500–1000 AD only. In the following, all KCI time series are considered on their aligned time scale.

5 Results and Discussion

The age scale of KCC provides the first chronology of the last millennium for a CG ice core that is fully based on annual layer counting. The new KCC age scale offers the to-date most accurate foundation to study the CG proxy time series over long time scales, e.g. regarding the recent investigation pursuing the link with historical evidence by More et al. (2017) who used

a slightly adjusted version of the age scale presented here (albeit not significantly different with respect to uncertainty). The novel technique of LA-ICP-MS was crucial for a reliable identification of cm and sub-cm thin layers in the deeper parts of the core. Thereby, this work adds to recent studies (e.g. Della Lunga et al., 2017; Haines et al., 2016; Mayewski et al., 2014) to demonstrate the potential of the high-resolution impurity records afforded by LA-ICP-MS for investigating highly thinned sections of polar and alpine ice cores. The combination of high-resolution annual layer counting and radiocarbon analysis
promises a break-through also for dating highly thinned deep parts of ice cores drilled at other sites.

5.1 Stable water isotope records

The covariation of the δ^{18} O time series between KCI and KCC strongly suggests a common atmospheric driver, i.e. temperature. At first glance Figure 5 shows an increasing trend over the last 100 years but also generally higher mean isotope levels prior to about 1900 AD. This is in line with earlier findings suggesting an "early instrumental period" warmer than instru-

- 15 mental data by about $+0.4^{\circ}$ C derived from the CG isotope signal (Bohleber et al., 2013). Here we find the generally higher average isotope levels to persist over much of the pre-industrial period: For instance, the mean δ^{18} O level in KCC between 1860-1000 AD is higher by about 0.75% than the 2000-1860 AD average. A quantitative use of the common isotope signal would therefore require addressing systematic so-called "upstream effects" and a reliable calibration of the isotope signal against instrumental temperature.
- 20 Upstream-effects concern the systematic variation in seasonality of the net accumulation upstream of the drilling site and have the potential to bias long-term core averages. Quantifying this effect requires accurate identification of the upstream catchment area (typically by sophisticated flow modeling) and evaluating the spatial variability in mean isotope levels. Dedicated efforts to evaluate the upstream-effect for the KCI-KCC flow line are currently underway (pers. comm. Carlo Licciulli and Josef Lier IUP Heidelberg). From a preliminary inspection of snow pit data recently obtained for the KCI-KCC flow line, there is no clear
- 25 indication of a systematic trend in mean δ¹⁸O levels upstream of KCI, however: Comprising roughly the years 2016–2014, three snow pits evaluated thus far show mean δ¹⁸O levels of -13.22, -11.94 and -15.04 ‰ at about 60, 195 and 300 m distance upstream of KCI, respectively (KCC is located roughly 110 m upstream of KCI, cf. Figure 1). In order to calibrate the stable water isotope signal, we used the instrumental temperature dataset compiled in an earlier study (Bohleber et al., 2013). This temperature dataset (referred to here as "CG modified temperature") was specifically adjusted
- 30 to the CG ice core conditions, taking into account the summer bias in precipitation and snow deposition. To calculate an iso-tope/temperature sensitivity, we considered both KCC and KCI individually as well as a stack of the two stable isotope records (calculated as their simple average at nominal annual resolution). Using 2000–1860 AD as calibration period (thus deliberately avoiding the "early instrumental period" prior to 1860) our results reproduce earlier findings of Bohleber et al. (2013). This



Figure 6. Comparison of KCC δ^{18} O and Ca²⁺ against the CG modified instrumental temperature (orange), shown in top and bottom, respectively, and covering the full instrumental period back to 1760 AD. Shown are anomalies relative to the respective 2000-1860 AD mean, and as decadal trends obtained from Gaussian smoothing.

specifically includes showing i) an overall agreement between the isotope and temperature record interrupted by characteristic decadal mismatch periods (Figure 6), ii) an increase in isotope/temperature correlation for multi-annual and decadal averages (e.g. r = 0.33, 0.47, 0.64 for discretely binned annual, 5 and 10 year averages, respectively in case of KCC) and iii) higher correlations obtained from the stack vs. the individual time series (e.g. r = 0.48, 0.67, 0.79 for annual, 5 and 10 year averages, respectively).

5

Regarding sensitivity values, we also find an increase with length in averaging period as well as substantially higher sensitivity values for KCI than KCC, revealing 2.3 vs. 1.4 $\%_{o}$ /°C, respectively, when using discretely binned 10-year averages (and 1.8 $\%_{o}$ /°C for the stacked record). Hence we obtain sensitivity values about threefold of what is expected, e.g. based on the isotope/temperature relationship of 0.65 $\%_{o}$ /°C reported by Rozanski et al. (1992) for European temporal trends in precipita-

- 10 tion. Since a more strict confinement towards sampling the high summer season can be expected for the lower accumulation KCI, the sensitivity difference among KCI and KCC points towards the seasonal bias in snow samping to be connected with enhancing sensitivity. It is important to note that the high isotope-sensitivity deserves a separate thorough investigation, ideally comprising regional climate-isotope modeling and taking into account post-depositional effects such as snow preservation, upstream-effects and isotope diffusion, which is outside the scope of this study. Until an adequate long-term calibration of
- 15 the CG isotope signal is achieved, however, we do not attempt a quantitative temperature reconstruction based on the isotope composite record so far.

Figure 7 shows the time series of δ^{18} O in comparison with Ca²⁺, deuterium excess and CPP. Due to the logarithmic distribution of the Ca²⁺ data, we generally use a log-scale to show the Ca²⁺ time series. For KCC, we find δ^{18} O and Ca²⁺ to be significantly correlated over the last 1000 years (at r = 0.49, cf. Figure 7). This suggests that the common decadal-scale



Figure 7. Records of KCC, displaying δ^{18} O together with Ca²⁺ (top row) and deuterium excess together with the coarse particle percentage (CPP, bottom row), shown in blue and orange, respectively. All time series are shown as anomalies relative to the respective 2000–1860 AD mean at biennial resolution. Note the high levels of deuterium excess and CPP around 1100–1200 AD.

signal driver behind the shared variability between the δ^{18} O time series of KCC and KCI also holds for Ca²⁺. Prior to about 1860 AD, the δ^{18} O time series constitutes nearly an upper envelope signal as compared to Ca²⁺. Short excursions to low Ca²⁺ concentrations missing a respective counterpart in δ^{18} O may have been smoothed out by isotope diffusion. Worth noting in this respect, the firn-ice transition in KCC coincides roughly with the last 100 years in the record (Tables 1 and 3).

5 5.2 Mineral dust proxy records

Following our general distinction (section 2), two signal components of the Ca^{2+} time series are investigated separately: i) the frequency of occurrence in Saharan dust deposition, and ii) the long-term average Ca^{2+} concentration. In order to investigate to what extent the time series agreement observed for $\delta^{18}O$ also holds in case of mineral dust related species, we use the insoluble particle signal as a surrogate for Ca^{2+} (since Ca^{2+} has not been measured for KCI). Figure 8 shows an overview of

- 10 the insoluble particle datasets of KCC and KCI. KCC data shows that the insoluble particle signal and Ca^{2+} concentrations are generally highly correlated (e.g. r = 0.9 within the last 1000 years). The KCC-KCI inter-core comparison of insoluble particle records reveals agreement of decadal scale features, as well as similarities regarding periods of low concentrations and higher peak abundance (e.g. 1800–1820 vs. 1780–1800, respectively). Differences in the magnitude of individual peak events as well as mean levels of particle concentrations can be explained in light of i) the KCI particle signal measured on
- 15 diluted sample meltwater, ii) potential calibration differences in the optical particle sensor and iii) inter-site snow deposition variability. Accordingly, it was not attempted to construct a composite record of insoluble particles from the two cores.



Figure 8. Inter-core comparison of the insoluble particle signal of KCC (blue) and KCI (orange). The top and middle row show the insoluble particle time series on a linear and logarithmic scale, respectively. The bottom row shows decadal trends of anomalies with respect to their 2000–1860 AD mean, highlighted by Gaussian smoothing. The KCI record is on the adjusted time scale after matching the stable water isotope records.

5.2.1 Detection of Saharan dust peak events and frequency of occurrence

Essential for the calculation of a robust occurrence rate of dust events are adequate means to distinguish desert dust from background and from deposition events of long-range transported anthropogenic pollutants. While the particle signal alone is not sufficient for differentiating these events, Saharan dust layers in CG ice cores can be reliably identified based on the

- 5 analyses of Ca²⁺, supplemented by alkalinity measurements and, in principle, particle size distribution (Wagenbach et al., 1996). The central criteria used in this study in order to identify Saharan dust are strongly elevated concentrations of Ca²⁺ coinciding with acidity values reduced to alkaline levels. Since no direct acidity measurements are available in our case from CFA, we rely on the ECM record for this purpose. At CG high dust levels are able to reduce the ECM signal to almost zero (rendering the ECM to be a qualitative dust indicator rather than an quantitative acidity gauge). Dust anomalies were identified
- 10 as "peaks over threshold". A robust spline smoothing was used to remove the general trend from the Ca^{2+} -data. Peak events then needed to exceed three times the median absolute deviations (Figure 9). We have also used the particle size distribution to investigate exemplarily a small number of dust events, finding that dust events show systematically higher CPP with respect to dust-free core sections.
- To detect the frequency of occurrence in dust peak events, we followed the statistical tool outlined in Chapter 6 of Mudelsee
 (2010). For a non-parametric occurrence rate estimate we used a moving Gaussian kernel (bandwidth 51 years) and accounted for boundary effects. For KCC, only the subset of peak events coinciding with a vanishing ECM signal was considered to be of Saharan dust origin. For KCI, we employed the same peak detection scheme to the insoluble particle signal. However, due



Figure 9. Results from detecting Saharan dust events in the ice core records and estimating their long-term frequency of occurrence. Figure part a) corresponds to KCC, part b) to KCI (see text). The bottom row shows the detected events (blue) and the frequency of occurrence kernel estimate with a 51-year bandwidth (bottom rows, in red) together with 90% confidence intervals.

to the lack of a full ECM profile, no subset corresponding to low acidity could be defined. Using a direct comparison with the insoluble particle signal of KCC, with and without ECM correction, we found that the respective uncorrected frequency of occurrence is expected to contain a minor bias towards higher peak abundances, but leaves the overall features unchanged. As the main robust features for KCC and KCI, the frequency of occurrence in dust peaks is systematically increased prior

- 5 to 1250 AD with respect to the rest of the record (Figure 9). Dust anomalies are found clustered in periods around 1100– 1200 (extending into the 1200s), around 1400–1450 and, for KCC only, between 1500–1800. This is in broad agreement with enhanced Saharan dust deposition reported by Thevenon et al. (2009) for periods around 1200–1300, 1430–1520, 1570–1690, 1780–1800, and after 1870. The latter periods were identified by Thevenon et al. (2009) based on sophisticated elemental analysis in a CG ice core, albeit at much coarser resolution and larger dating uncertainty, which hampers a more detailed
- 10 comparison. However, in our data we also recognize a large dust peak located between 1780 and 1800 which was suggested as a dating reference horizon by Thevenon et al. (2009).

5.2.2 The long-term Ca²⁺ variability in relation to temperature

Within the calibration period 2000–1860 AD, we find the overall increasing trend in instrumental temperature to be represented also in increasing levels of δ^{18} O and Ca²⁺ (Figure 6). The Ca²⁺ signal correlates significantly with the CG modified instru-

also in increasing levels of δ^{16} O and Ca²⁺ (Figure 6). The Ca²⁺ signal correlates significantly with the CG modified instrumental temperature at r = 0.41, 0.56, 0.71 using biennial, 5 and 10 year averages, respectively, within the calibration period. Nearly identical correlation values are obtained for the full instrumental period back to 1760 AD. Within the calibration period (Figure 6), we compared the decadal trends (highlighted by Gaussian smoothing) of the CG modified instrumental temperature with δ^{18} O and Ca²⁺, respectively. The comparison reveals that the Ca²⁺ signal performs similarly to δ^{18} O in explaining variance of the temperature data (both at around 25%, although only interpreted with caution due to the autocorrelation of the smoothed curves).

Potential drivers for a Ca²⁺-temperature coupling can be expected from i) the advection of air masses comprising a high Sa-

- 5 haran dust load generally being associated with warm temperatures (Wagenbach et al., 1996), ii) the deposition of dust leading to lowered snow albedo thus supporting surface snow consolidation (Haeberli et al., 1983) and iii) warm temperature favoring snow consolidation (e.g. Fauve et al., 2002; Haeberli et al., 1983). However, it generally remains difficult to quantify the influence of the above processes. It seems worth pointing out that, process iii) acts independently from the type of impurity / isotope species considered. Likewise, a similar process as i) may be envisaged in case of δ¹⁸O. However, process ii) mainly
- 10 concerns dust-related species such as Ca²⁺ providing an essential "self-preserving" character for these species with respect to snow deposition. Regarding this connection between dust content and presumed faster snow preservation, supporting evidence is provided by including the high-resolution density profile of KCC. Comparing profiles of Ca²⁺ and density reveals that layers with a high dust load generally coincide with layers of enhanced density (Figure A2 in the appendix). Regarding process iii), we made an attempt to semi-quantitatively explore the imprint of snow preservation on the long-term variability in Ca²⁺
- 15 (appendix A1). Previous studies already used simplified conceptual models to investigate the influence of snow deposition on seasonal ice core signals, and demonstrated the decisive role played by the amplitude of the seasonality (Wagenbach et al., 2012; Fisher and Koerner, 1988). We employed the model by Wagenbach et al. (2012), with parameters reflecting to CG snow preservation conditions. This revealed that the resulting bias from incomplete snow preservation on the average Ca²⁺ level is already in the same order of magnitude as the long-term Ca²⁺ trends found in previous studies (Wagenbach et al., 1996) and
- 20 also in the core investigated here. As a result, the influence of snow deposition appears non-negligible in explaining the apparent Ca²⁺-temperature co-variation. Notably this implies that, at best, the Ca²⁺-temperature coupling allows for using Ca²⁺ trends as a site-specific temperature proxy only. This is in analogue to the study by Kellerhals et al. (2010a) demonstrating temperature-related variability for NH_4^+ at a low-latitude site, albeit explained by a different mechanism than discussed for Ca²⁺ at CG here.

25 5.3 Temperature and mineral dust variability over the last millennium

Based on the above considerations and the agreement within the instrumental period (Figure 6), the potential of the Ca²⁺ signal to quantitatively record temperature variability is explored further over the full 1000 year period. For this purpose the biennial logarithmic Ca²⁺ is calibrated tentatively against instrumental temperature using linear regression within the time period 2006–1860 AD. The respective 90% confidence intervals are used to calculate a temperature reconstruction with uncertainty bands (0.7–1–1.8 °C/log Ca²⁺ [ppb]). Decadal trends are again highlighted by Gaussian smoothing in Figure 10. The result-

30 bands (0.7–1–1.8 °C/log Ca²⁺ [ppb]). Decadal trends are again highlighted by Gaussian smoothing in Figure 10. The resulting 1000-year record is shown with a tentative 200-year extension. Regarding its overall features and in view of remaining dating uncertainties, the record provides evidence of "Little Ice Age conditions" systematically cooler than the reference and calibration period, with an average of -0.3°C between 1800–1200 AD. A shorter warm interval of about +0.3°C is found in the late 1100s These features are especially noteworthy considering the above-average mean of the δ^{18} O values with respect



Figure 10. Comparison of decadal temperature trends as anomalies with respect to the mean of 2006–1860 AD. Shown are calibrated temperatures obtained from the KCC Ca^{2+} variability (blue lines, with uncertainty indicated as light blue bands). Also shown are instrumental temperature data (black) and the summer temperature reconstruction of Luterbacher et al. (2016) in red (uncertainty as gray bands). Note that the overall co-variation among the two reconstructions persists for at least another 200 years beyond 1000 AD (light gray shaded area). Black bars on the bottom indicate maximum dating uncertainty.

to 2006–1860 AD (Figure 7). The comparison with δ^{18} O suggests that the Ca²⁺-temperature coupling may be less affected by non-stationary sensitivity (and upstream) effects.

In this context we further explored our Ca^{2+} -based temperature-reconstruction attempt in comparison with other proxy reconstructions of European summer temperature. We show here results from using the mean European summer temperature

- 5 anomalies reconstructed by Luterbacher et al. (2016), considering their composite-plus-scaling method (CPS) adjusted to biennial resolution and our reference time period of 2006–1860 AD. The decadal trends (represented by Gaussian smoothing) of both reconstructions shown in Figure 10 are generally consistent in their overall features (and formally correlate at r = 0.4). These features comprise the recent warming trend and below average conditions during the "Little Ice Age" (LIA), and a warm episode in the late 1100s ("Medieval Climate Anomaly", MCA). The only major feature of disagreement occurs around the
- 10 already noted minimum around 1250–1230 AD. The overall low levels of impurities (especially NH_4^+) and $\delta^{18}O$ may point to increased deposition of winter snow during this time, or exceptionally cold summer conditions. It is worth noting that if, tentatively, extending the comparison for another 200 years beyond 1000 AD, the agreement between the two reconstructions continues to last until 800 AD, consistently showing a relatively warm interval lasting between about 1000–850 AD. The overall agreement is especially noteworthy in the light of: i) Only small offsets exist between the general features of the two
- 15 records, which may stem from the remaining dating uncertainty of KCC. The comparison is not intended as a dating validation, however. ii) The absence of evidence of a non-stationary sensitivity. The magnitude of the general features (LIA, MCA) and decadal scale temperature variability derived from our ice core record is consistent overall with the other proxy reconstruction. A potential systematic bias to the observed Ca²⁺-temperature coupling could arise from strong and long-term changes at the

dust source, e.g. increased dust mobilization, which could also affect the long-term Ca^{2+} variability. At the same time, increased dust mobilization would likely also influence the Saharan dust spikes and their frequency of occurrence. However, we only find one instance of long-term changes in dust occurrence rate. Being the most outstanding period in the dust event occurrence warrants taking a closer look at 1100–1200 AD. This outstanding period is characterized by i) an increased frequency of

- 5 Saharan dust events, ii) its above average levels, both in CPP and deuterium excess, starting to rise around 1100 AD, and iii) a delayed relative increase in δ^{18} O and Ca²⁺, e.g. first showing minimum concentrations between 1130–1170 AD followed by a shorter maxima around 1170–1200 AD. The CPP maximum constitutes the dominant feature of the entire record (notably, this result does not depend of the exact threshold chosen to calculate the CPP). The median of the CPP record within this 1100–1200 AD time period (0.61 ± 0.11 , reported with one median absolute deviation) indicates an increase in coarse particles by
- 10 about 12% relative to the median of the rest of the 1000-year time period (0.48 ± 0.05) . While an increase in dust mobilization cannot be ruled out, the connection of the distinct increase in coarse particles with enhanced dust event frequency rather suggests an increase in the direct transport of Saharan dust (as opposed to indirect advection with longer pathway and thus stronger decrease in coarse particles). Finding increased values of deuterium excess supports the view of a relative increase in direct Saharan dust advection, as increased deuterium excess could be expected from warm and dry air masses collecting moisture
- 15 over the Mediterranean (e.g. Pfahl and Sodemann, 2014). Overall these findings motivate getting an even more detailed picture of this outstanding period, e.g. using isotopic fingerprinting of the dust layers for a precise provenance investigation, which is left for future investigations.

6 Conclusions and Outlook

A combination of state-of-the-art methods in ice core analysis allowed us to date the latest CG ice core KCC with unprecedented confidence. The breakthrough in this respect was to extend annual layer counting, for the first time at CG, over more than the last 1000 years and finding the resulting age scale corroborated by radiocarbon analyses. The combination of high-resolution annual layer counting afforded by LA-ICP-MS with constraints from radiocarbon analyses could be employed with great success also at deep sections of other (mountain) ice cores. By means of the improved age scale it became possible, for the first time, to demonstrate that the inter-core agreement in decadal isotope variability among two cores on the same flow line extends

25 over the last 1000 years. The inter-core agreement suggests a common driver of the shared signal, also extending to the longterm variability in Ca^{2+} . We find substantial agreement among the decadal trends of Ca^{2+} and temperature at CG, over the entire instrumental period. Since snow preservation plays a key role for the observed coupling between Ca^{2+} and instrumental temperature, this makes Ca^{2+} trends at best a site-specific temperature proxy, although it remains to be tested to what degree the association with temperature also holds at other alpine drilling sites. In contrast to the stable isotope signal at CG, however,

30

we find no evidence of non-stationary temperature sensitivity for Ca^{2+} : Exploiting the Ca^{2+} -temperature agreement for a quantitative reconstruction i) proves to be consistent with other latest summer temperature reconstructions, and ii) reproduces overall features regarding the "Little Ice Age" and the "Medieval Climate Anomaly". Parameters less influenced by snow preservation (dust event occurrence rate and particle size distribution) reveal an exceptional medieval period around 1100–

1200 AD, suggesting a relative increase in meridional flow and dry conditions over the Mediterranean during that time. Future and ongoing investigations will target the application of our new dating approach to the bottom 10 m WE of KCC, and an improved quantitative understanding of the isotope-thermometer at CG. In this context a central question remains whether the isotope-based temperature signal can be reconciled quantitatively with the Ca^{2+} -based reconstruction.

- 5 Acknowledgements. We are grateful to numerous colleagues for their commitment regarding field work, ice core drilling and ice core analyses. In particular we would like to acknowledge the support of the Initiative for the Science of the Human Past at Harvard University and all its project members. Additional invaluable support in ice core processing was provided by the Alfred-Wegener-Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven (AWI). The Klaus-Tschira-Lab Mannheim is acknowledged for their support in radiocarbon analysis. We also would like to thank Johanna Kerch, Carlo Licciulli, Josef Lier and Lars Zipf from IUP Heidelberg for their support. We
- 10 thank Johannes Freitag (AWI Bremerhaven) for the high-resolution density data. Recovery and analysis of the 2013 CG ice core KCC were supported by the Arcadia Fund of London (AC3450) and the Helmholtz Climate Initiative REKLIM. Work on the 2005 CG ice core KCI has been funded by the European Union under contract ENV4-CT97-0639 (project ALPCLIM) and within the project ALP-IMP through grant EVK2-CT2002-00148. LA-ICP-MS ice core analyses were conducted in the Climate Change Institute's W. M. Keck Laser Ice Facility at the University of Maine supported from the W. M. Keck Foundation and the National Science Foundation (PLR-1042883, PLR-1203640).
- 15 Financial support was provided to P.B. by the Deutsche Forschungsgemeinschaft (BO 4246/1-1). We would like to especially thank and acknowledge our late colleague Dietmar Wagenbach (Heidelberg University) for his long-standing contributions to glaciological research at Colle Gnifetti, and in particular for sharing his unique expertise with us at the early stage of our project.

References

19, 131-147, 1983.

5

25

Ahmed, M., Anchukaitis, K. J., Asrat, A., Borgaonkar, H. P., Braida, M., Buckley, B. M., Büntgen, U., Chase, B. M., Christie, D. A., Cook,E. R., et al.: Continental-scale temperature variability during the past two millennia, Nature Geoscience, 6, 339, 2013.

Alean, J., Haeberli, W., and Schädler, B.: Snow accumulation, firn temperature and solar radiation in the area of the Colle Gnifetti core drilling site (Monte Rosa, Swiss Alps): distribution patterns and interrelationships, Zeitschrift für Gletscherkunde und Glazialgeologie,

Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M.-L., Steffensen, J. P., Dahl-Jensen, D., et al.: The Greenland ice core chronology 2005, 15–42ka. Part 1: Constructing the time scale, Quaternary Science Reviews, 25, 3246–3257, 2006.

10 Bohleber, P., Wagenbach, D., Schöner, W., and Böhm, R.: To what extent do water isotope records from low accumulation Alpine ice cores reproduce instrumental temperature series?, Tellus B, 65, 2013.

Brook, E. J., Wolff, E., Dahl-Jensen, D., Fischer, H., Steig, E. J., et al.: The future of ice coring: international partnerships in ice core sciences (IPICS), PAGES news, 14, 6–10, 2006.

Della Lunga, D., Müller, W., Rasmussen, S. O., Svensson, A., and Vallelonga, P.: Calibrated cryo-cell UV-LA-ICPMS elemental concentra-

- 15 tions from the NGRIP ice core reveal abrupt, sub-annual variability in dust across the GI-21.2 interstadial period, The Cryosphere, 11, 1297–1309, doi:10.5194/tc-11-1297-2017, http://www.the-cryosphere.net/11/1297/2017/, 2017.
 - Fauve, M., Rhyner, H., and Schneebeli, M.: Pistenpräparation und Pistenpflege: ein Handbuch für den Praktiker, Eidgen. Inst. für Schnee-u. Lawinenforschung, 2002.
- Fisher, D. A. and Koerner, R. M.: The effects of wind on $\delta({}^{18}O)$ and accumulation give an inferred record of seasonal δ amplitude from the Agassiz ice cap, Ellesmere Island, Canada, Annals of Glaciology, 10, 34–37, 1988.
- Gabrielli, P., Barbante, C., Bertagna, G., Bertó, M., Binder, D., Carton, A., Carturan, L., Cazorzi, F., Cozzi, G., Dalla Fontana, G., et al.: Age of the Mt. Ortles ice cores, the Tyrolean Iceman and glaciation of the highest summit of South Tyrol since the Northern Hemisphere Climatic Optimum, The Cryosphere, 10, 2779, 2016.

Haeberli, W., Schotterer, U., Wagenbach, D., Schwitter, H. H., and Bortenschlager, S.: Accumulation characteristics on a cold, high-Alpine firn saddle from a snow-pit study on Colle Gnifetti, Monte Rosa, Swiss Alps, Journal of Glaciology, 29, 260–271, 1983.

Haeberli, W., Schmid, W., and Wagenbach, D.: On the geometry, flow and age of firn and ice at the Colle Gnifetti core drilling site (Monte Rosa, Swiss Alps), Zeitschrift für Gletscherkunde und Glazialgeologie, 24, 1–19, 1988.

- 30 Haines, S. A., Mayewski, P. A., Kurbatov, A. V., Maasch, K. A., Sneed, S. B., Spaulding, N. E., Dixon, D. A., and Bohleber, P. D.: Ultra-high resolution snapshots of three multi-decadal periods in an Antarctic ice core, Journal of Glaciology, 62, 31–36, 2016.
 - Hoelzle, M., Darms, G., Lüthi, M. P., and Suter, S.: Evidence of accelerated englacial warming in the Monte Rosa area, Switzerland/Italy, The Cryosphere, 5, 231–243, 2011.

Hoffmann, H., Friedrich, R., Kromer, B., and Fahrni, S.: Status report: Implementation of gas measurements at the MAMS 14C AMS facility

in Mannheim, Germany, Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms,
 410, 184 – 187, doi:https://doi.org/10.1016/j.nimb.2017.08.018, 2017a.

Haeberli, W. and Funk, M.: Borehole temperatures at the Colle Gnifetti core-drilling site (MonteRosa, Swiss Alps), Journal of Glaciology, 37, 37–46, 1991.

Hoffmann, H., Preunkert, S., Legrand, M., Leinfelder, D., Bohleber, P., Friedrich, R., and Wagenbach, D.: A new sample preparation system for micro-14C dating of glacier ice with a first application to a high Alpine ice core from Colle Gnifetti (Switzerland), Radiocarbon, in press, 2017b.

Hoffmann, H. M.: Micro radiocarbon dating of the particulate organic carbon fraction in Alpine glacier ice: method refinement, critical

5 evaluation and dating applications, Ph.D. thesis, Heidelberg University, http://www.ub.uni-heidelberg.de/archiv/20712, 2016.

Jenk, T. M., Szidat, S., Bolius, D., Sigl, M., Gäggeler, H. W., and co-authors: A novel radiocarbon dating technique applied to an ice core from the Alps indicating late Pleistocene ages, Journal of Geophysical Research, 114, D14 305, 2009.

Kellerhals, T., Brütsch, S., Sigl, M., Knüsel, S., Gäggeler, H. W., and Schwikowski, M.: Ammonium concentration in ice cores: A new proxy for regional temperature reconstruction?, Journal of Geophysical Research: Atmospheres, 115, 2010a.

10 Kellerhals, T., Tobler, L., Brütsch, S., Sigl, M., Wacker, L., Gäggeler, H. W., and Schwikowski, M.: Thallium as a tracer for preindustrial volcanic eruptions in an ice core record from Illimani, Bolivia, Environmental Science & Technology, 44, 888–893, 2010b.

Konrad, H., Bohleber, P., Wagenbach, D., Vincent, C., and Eisen, O.: Determining the age distribution of Colle Gnifetti, Monte Rosa, Swiss Alps, by combining ice cores, ground-penetrating radar and a simple flow model, Journal of Glaciology, 59, 179–189, 2013.

- Lisiecki, L. E. and Lisiecki, P. A.: Application of dynamic programming to the correlation of paleoclimate records, Paleoceanography, 17, 2002.
 - Luterbacher, J., Werner, J., Smerdon, J. E., Fernández-Donado, L., González-Rouco, F., Barriopedro, D., Ljungqvist, F. C., Büntgen, U., Zorita, E., Wagner, S., et al.: European summer temperatures since Roman times, Environmental Research Letters, 11, 024 001, 2016.
 Lüthi, M. P. and Funk, M.: Dating ice cores from a high Alpine glacier with a flow model for cold firn, Annals of Glaciology, 31, 69–79, 2000.
- 20 Mayewski, P. A., Sneed, S. B., Birkel, S. D., Kurbatov, A. V., and Maasch, K. A.: Holocene warming marked by abrupt onset of longer summers and reduced storm frequency around Greenland, Journal of Quaternary Science, 29, 99–104, doi:10.1002/jqs.2684, http://doi. wiley.com/10.1002/jqs.2684, 2014.

More, A. F., Spaulding, N. E., Bohleber, P., Handley, M. J., Hoffmann, H., Korotkikh, E. V., Kurbatov, A. V., Loveluck, C. P., Sneed, S. B., McCormick, M., and Mayewski, P. A.: Next generation ice core technology reveals true minimum natural levels of lead (Pb) in the atmo-

25 sphere: insights from the Black Death, GeoHealth, pp. n/a–n/a, doi:10.1002/2017GH000064, http://dx.doi.org/10.1002/2017GH000064, 2017GH000064, 2017GH000064, 2017.

Mudelsee, M.: Climate Time Series Analysis, Springer, 2010.

Nye, J.: Correction factor for accumulation measured by the thickness of the annual layers in an ice sheet, Journal of Glaciology, 4, 785–788, 1963.

30 Pfahl, S. and Sodemann, H.: What controls deuterium excess in global precipitation?, Climate of the Past, 10, 771, 2014.

Preunkert, S. and Wagenbach, D.: An automatic recorder for air/firn transfer studies of chemical aerosol species at remote glacier sites, Atmospheric Environment, 32, 4021–4030, 1998.

Preunkert, S., Wagenbach, D., Legrand, L., and Vincent, C.: Col du Dôme (Mt Blanc Massif, French Alps) suitability for ice-core studies in relation with past atmospheric chemistry over Europe, Tellus, 52B, 993–1012, 2000.

35 Prodi, F. and Fea, G.: A case of transport and deposition of Saharan dust over the Italian peninsula and southern Europe, Journal of Geophysical Research: Oceans (1978–2012), 84, 6951–6960, 1979.

Ramsey, B.: C. OxCal Program v4. 2.4, Radiocarbon Accelerator Unit, Univ. Oxford, 2016.

Rasmussen, S. O., Andersen, K. K., Svensson, A., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M.-L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., et al.: A new Greenland ice core chronology for the last glacial termination, Journal of Geophysical Research: Atmospheres, 111, 2006.

Rozanski, K., Araguas-Araguas, L., and Gonfiantini, R.: Relation Between Long-Term Trends of Oxygen-18 Isotope Composition of Pre-

5

Ruth, U., Wagenbach, D., Steffensen, J. P., and Bigler, M.: Continuous record of microparticle concentration and size distribution in the central Greenland NGRIP ice core during the last glacial period, Journal of Geophysical Research: Atmospheres, 108, 2003.

Schwerzmann, A., Funk, M., Blatter, H., Lüthi, M., Schwikowski, M., and Palmer, A.: A method to reconstruct past accumulation rates in alpine firn regions: A study on Fiescherhorn, Swiss Alps, Journal of Geophysical Research: Earth Surface, 111, 2006.

- 10 Schwikowski, M., Seibert, P., Baltensperger, U., and Gaggeler, H.: A study of an outstanding Saharan dust event at the high-alpine site Jungfraujoch, Switzerland, Atmospheric Environment, 29, 1829–1842, 1995.
 - Sneed, S. B., Mayewski, P. A., Sayre, W. G., Handley, M. J., Kurbatov, A. V., Taylor, K. C., Bohleber, P., Wagenbach, D., Erhardt, T., and Spaulding, N. E.: New LA-ICP-MS cryocell and calibration technique for sub-millimeter analysis of ice cores, Journal of Glaciology, 61, 233, 2015.
- 15 Sodemann, H., Palmer, A., Schwierz, C., Schwikowski, M., and Wernli, H.: The transport history of two Saharan dust events archived in an Alpine ice core, Atmospheric Chemistry and Physics Discussions, 5, 7497–7545, 2005.

Stuiver, M. and Polach, H.: Discussion: Reporting of 14C data., Radiocarbon, 19, 355-63., 1977.

cipitation and Climate, Science, 258, 981-985, 1992.

Thevenon, F., Anselmetti, F. S., Bernasconi, S. M., and Schwikowski, M.: Mineral dust and elemental black carbon records from an Alpine ice core (Colle Gnifetti glacier) over the last millennium, Journal of Geophysical Research: Atmospheres, 114, 2009.

- 20 Uglietti, C., Zapf, A., Jenk, T. M., Sigl, M., Szidat, S., Salazar Quintero, G. A., and Schwikowski, M.: Radiocarbon dating of glacier ice: overview, optimisation, validation and potential, The Cryosphere, 10, 3091–3105, 2016.
 - Wagenbach, D.: Environmental Records in Alpine Glaciers, in: The Environmental Record in Glaciers and Ice Sheets, edited by Oeschger,H. and C. C. Langway, Jr., pp. 69–83, John Wiley and Sons, 1989.

Wagenbach, D.: Special problems of mid-latitude glacier ice-core research. In: Greenhouse Gases, Isotopes and Trace Elements in Glaciers

- 25 as Climatic Evidence of the Holocene, vol. Arbeitsheft Nr.14 of *Report of the ESF/EPC Workshop, Zürich 27-28 October 1992*, Versuchsanstalt für Wasserbau, Hydrologie und Glaziologie der Eidgenössischen Technischen Hochschule Zürich, 1992.
 - Wagenbach, D. and Geis, K.: The mineral dust record in a high altitude glacier (Colle Gnifetti, Swiss Alps), in: Paleoclimatology and Paleometeorology: Modern and Past Patterns of Global Atmospheric transport, edited by M, L. and M, S., pp. 543–564, Kluwer Academic Publishers, Dordrecht, 1989.
- 30 Wagenbach, D., Preunkert, S., Schäfer, J., Jung, W., and Tomadin, L.: Northward transport of Saharan dust recorded in a deep Alpine ice core, in: The impact of desert dust across the Mediterranean, pp. 291–300, Springer, 1996.

Wagenbach, D., Bohleber, P., and Preunkert, S.: Cold alpine ice bodies revisited: What may we learn from their isotope and impurity content?, Geografiska Annaler: Series A, Physical Geography, 94, 245–263, 2012.

Appendix A: Supplementary Material

A1 A semi-quantitative treatment of the snow deposition influence on average Ca²⁺-concentrations and a possible coupling to atmospheric temperature

We attempt to semi-quantitatively explore the imprint of snow preservation on the long-term variability in Ca²⁺, using the 5 conceptual model by Wagenbach et al. (2012) and considering δ^{18} O as a reference. The model assumes sinusoidal cycles for the precipitation-borne signal S(t) and the surface accumulation pattern A(t), and a phase-lag t_{ϕ} between S(t) and A(t). The deviation Δ of the mean signal recorded in the firm at CG (\bar{S}_{firm}) with respect to the overall mean signal (S_0) in the precipitation then becomes (corresponding to equation (5) in Wagenbach et al. (2012)):

 $S(t) = S_0(1 + s_r \sin\omega(t + t_\phi)) \qquad A(t) = A_0(1 + a_r \sin\omega t) \qquad \Delta = \bar{S}_{\rm firm} - S_0 = S_0 s_r a_r \cos\omega t_\phi \qquad (A1)$

- 10 with a_r and s_r denoting the relative signal amplitudes, t_{ϕ} the temporal phase shift between the two cycles and ω the cycle frequency equal to $2\pi/T$ (period T = 1 yr). We reproduced here the calculation for δ^{18} O by Wagenbach et al. (2012) with typical seasonality parameters at Colle Gnifetti (e.g. $a_r = 0.8$). To estimate Δ for Ca²⁺, we used values reported by Preunkert et al. (2000) for CG, i.e. taking $S_0 = 112$ ppb (the average of typical summer and winter concentrations) as well as $s_r = 0.64 \pm 0.26$. Notably, this value of s_r for Ca²⁺ is close to the absolute value of $s_r = 0.5$ obtained for δ^{18} O by Wagenbach et al. (2012).
- 15 Estimating the radiation-induced control of snow consolidation on the seasonal net accumulation cycle at CG is consistent with a phase shift of $t_{\phi} = 1.5$ months (broadly equal to the delay of the isotope/aerosol peak with respect to insolation (Wagenbach et al., 2012)). Figure A3 shows the deviation Δ for Ca²⁺ and δ^{18} O as a function of phase shift t_{ϕ} and a_r . The framework of the conceptual model allows us to semi-quantitatively explore a potential coupling of the long-term Ca²⁺

signal to atmospheric temperature via snow preservation effects. For instance, the effect of warmer atmospheric summer tem-

- 20 peratures on snow preservation is envisaged as increased summer snow deposition, corresponding to an according change in a_r and, an additional possibility, a change in phase-shift t_{ϕ} . In the model, a more efficient snow-preservation for dust-rich layers would correspond to a decrease in phase-shift t_{ϕ} . We follow Wagenbach et al. (2012) and highlight the sensitivity in the deviation corresponding to an arbitrarily chosen variability of a_r of $\pm 10\%$ and $t_{\phi} = 1.5 \pm 0.5$ months. This already shifts the mean Ca²⁺ level by about 15 ppb, which is in the same order of magnitude as the long-term trends of around ± 50 ppb found in pre-
- vious studies (Wagenbach et al., 1996) and also in the core investigated here. The potential variability in the Ca²⁺-seasonality due to the episodic input of Saharan dust can add a substantial contribution to the deviation (cf. dashed lines in Figure A3). However, the influence of a single Saharan dust event is rather short-term in comparison to the envisaged systematic shifts to a_r and t_{ϕ} imposed by atmospheric temperature change.

Horizon	Year AD	Depth KCC	Depth KCI
		[m WE]	[m WE]
Dust	1977	8.8	4.7
Tritium	1963	11.2	5.8
Dust	1947	15.3	7.7
Dust	1901/02	22.2	11.5

Table A1. Absolute age horizons used in datings of KCC and KCI

Core	Depth]	14 C cal	¹⁴ C cal sigma
	[m WE]	[yr b1950]	[yrs]
KCC	38.5	736	349
	39.2	899	695
	40.9	555	179
	41.3	1007	167
	42.6	1395	300
	43.1	1405	27
KCI	26	365	200
	28.4	1217	336
	30	354	214
	33.9	647	242
	36.2	850	276
	38.5	1178	179

Table A2. Radiocarbon ages used in datings of KCC and KCI



Figure A1. KCC δ^{18} O and Ca²⁺ data shown for the uppermost core section (the first 1.6 m were excavated by a snow trench for drilling). The vertical grey dashed lines indicate individual annual layers determined based on the NH₄⁺ profile. Black arrows show multiple sub-seasonal local maxima in the Ca²⁺ concentration, supporting the notion of "grouped peaks" revealed by the high resolution LA-ICP-MS Ca profile at greater depths.



Figure A2. Example view on comparing the KCC Ca^{2+} record with the density profile. Density was measured at high resolution by computer tomography at the Alfred Wegener Institute Bremerhaven (pers. comm. Johannes Freitag). The comparison illustrates the connection between dust content and presumed faster snow preservation: Layers of high dust load coincide with layers of enhanced density (with respect to ambient layers).



Figure A3. Sensitivity of the average Ca²⁺ signal to snow preservation effects, in comparison to δ^{18} O. The plot shows the deviation of the mean annual ice core signal from the respective precipitation mean as a function of phase shift t between the seasonal cycles of the signal and the accumulation rate. The thick solid black line shows the deviation for δ^{18} O (from Wagenbach et al. (2012)) and Ca²⁺, plotted on the y-axes on the left and right axis, respectively, and using typical seasonality parameters at Colle Gnifetti (see text). Indicated as thin lines are ranges corresponding to a variability of the relative seasonal amplitude a_r of 10% (arbitarily chosen), and using the estimated maximum uncertainty range of s_r for Ca²⁺ (solid and dashed lines, respectively).