

# Enhanced Mediterranean water cycle explains increased humidity during MIS 3 in North Africa - Response to reviewers

We thank the two anonymous Reviewers of our draft manuscript for their detailed and constructive reviews, and are extremely pleased that they find our work both interesting and worthy of publication. We fully concur that the interpretation of the data we present is complicated by structure of the dataset, and we are happy both reviewers agree with us that the data itself is so unique as to make a pressing case for publication and, that our analysis of it is fair, balanced and reasonable.

Below, we respond to the reviewers in order (first Reviewer 1, then Reviewer 2). The location code given for each comment and response represents the page the comment occurs in, followed by paragraph and lines (C#, #, #-#).

## **Anonymous Reviewer 1**

C2, 2, 3-4: *".....however, the clarity of the manuscript could be improved by changing the structure, especially the Discussion section."*

and C2, 2, 6-10: *".....the current structure of the manuscript makes it hard for the reader to follow the arguments..... and the factors influencing the stable isotopic composition of rainfall that need to be clarified / included."*

Response: obviously, we set out to make the draft manuscript as clear as possible, but we are happy to clarify further through editing and re-structuring as recommended by this reviewer. This will include re-ordering the Discussion so that information arrives in the most useful order, and further improving the clarity of the figures. It is important to us to make our work as accessible as possible! The *"few additional points"* are discussed below.

C2, 3, 4-9 continuing to C3, 1, 1-5: *"However, I disagree with some of the statements made in the paragraph starting in line 66..... A northward shift of the tropical monsoon belt to ~25oN would be sufficient for this."*

Response: The Reviewer's concern is essentially that we have exaggerated the lack of agreement between empirical evidence of wet conditions between 30-35°N and models, which generally do not get the monsoon so far north. They are also concerned we neglect the role of water recycling in the region 25-35°N. We are very happy to improve the discussion by including the water recycling argument, which we do overlook. On the other hand, we consider that the uncertainty in reconciling the empirical and physical lines of evidence is interesting and unresolved, and deserves to be highlighted in the way we do.

C2, 3, 5-7: *"I would also like to see a bit more detail about what the lake and vegetation records from the Sahara suggest for the actual time period covered by the speleothems."*

Response: Although beginning to be recognised as a humid period elsewhere in the Mediterranean basin (Langgut et al., 2018), MIS3 is not well expressed in the Sahara region. Consequently, there is limited pollen or lake constraints to develop our understanding from. Generally, the Libyan interior is considered arid or hyperarid throughout the last glaciation (Cancellieri E. et al., 2016). Recent re-evaluation of lake levels in southwest Egypt indicates a groundwater fed system was active around 41 ka (Nicoll, 2018), which is similar to dates for springline tufa systems at Kharga Oasis (Smith et

al., 2007). We are not aware of continental MIS3 pollen records from the region, but marine pollen from Tunisia indicates more arid conditions through the last glacial than during the Holocene (Brun, 1991). There is a triple peak in runoff from the Nile recorded in the marine sediment record, with maxima at ~60, ~55 and ~35ka, indicating higher rainfall within the upper Nile catchment (Revel et al., 2010). We will include a summary of this evidence in the introduction, to provide better context for our new data.

*C2, 3, 10-12: "A short discussion of the effects of different boundary conditions and how it could affect northern African moisture should be included."*

Response: We agree this would be useful, although keeping it short is a challenge for this complex system! Very briefly, the boundary conditions on northern African atmospheric moisture supply are 1) the sea surface temperature of the Atlantic and Mediterranean, 2) the surface water  $\delta^{18}\text{O}$  of the same ocean regions, 3) land surface temperature of Africa and to a lesser extent southern Europe, 4) insolation (especially with respect to ITCZ position) and 5) the zonal pressure gradient across northern Africa.

*C3, 1, 12-25: "I also think there should be a section in the introduction about the present day rainfall systems in the region..... [error in citation of Celle-Jeanton et al 2001].....How were they [the rainfall end members] defined?.....How were the averages in Figure 9 calculated?"*

Response: First – apologies for the error in the citation – the reference given in the References section is correct.

We agree that including more detail about the modern rainfall system in the Introduction would be helpful. We also agree that we can improve description and definition of the end members. The Bet Dagan and Tunisian datasets we use are shown in Figure 5a, and do indeed have different meteoric water lines and D-excess characteristics. The sub-categories of rainfall within the Sfax dataset occupy different positions on the same meteoric water line (Celle-Jeanton et al., 2001). We have checked, and confirm that the same is true for the Bet Dagan data.

The moisture sources used for the Bet Dagan site is taken from the regional meteorology (Black et al., 2010; Gat et al., 2003). Moisture sources for Tunisia are as defined by Celle-Jeanton et al (2001), and we refer readers to this primary reference. Averages are arithmetic means, as this better reflects behaviour over time within systems where water throughput in and out of the karst system is likely very rapid compared to speleothem growth – we do not expect to be able to resolve synoptic events in this record.

*C3, 1, 25-27: "What synoptic processes were involved in the formation of rain clouds – convection, advection? What circumstances lead to convective rainfall in the region?"*

Response: Convective systems, cyclones, upper-level troughs and static instabilities can all drive rainfall patterns in the Mediterranean basin and these modes are reviewed in (Dayan et al., 2015). Convection essentially reflects the relatively high SST of the Mediterranean during the winter, but rising air masses generally also need significant advection of moisture to drive significant rainfall. Upper level troughs reflect large-scale circulation (e.g. Red Sea Trough) or reflect lee effects downstream of mountains in the western Mediterranean, and promote rainfall in their regions of formation. The dominant cyclogenetic centre is in the Gulf of Genoa, and secondary centres are placed in south Italy, Crete and Cyprus. Cyclonic systems can also penetrate from the Atlantic, where the high SST of the winter Mediterranean tends to sustain and amplify them, in close analogy to convection forcing. The key static instability is the penetration of the tropical air mass into the

subtropical Mediterranean, forming a 'Saharan Cloud Band' at middle and upper atmospheric levels. These originate from within the ITCZ. As Libya is very sparsely instrumented, there is no literature we can find to specifically identify the synoptic processes involved in cloud formation precisely over our site. However, the Levant region is very well instrumented. Here, most rainfall falls under winter, low pressure conditions, and is convective (Peleg and Morin, 2012). The responsible low pressure systems can relate to transient, shallow lows over northern Israel, or less frequently more long-lasting Cyprus Lows or Red Sea Trough systems (Peleg and Morin, 2012).

C3, 1, 27-29 continuing to C4, 1, 1-2: *"Further adding to the point above, it seems that the discussion of factors influencing stable isotopic reconstruction of rainfall is focussed on only the effects of rainfall amounts, temperature and moisture source. These are important factors, but I think one important factor is missing and that is the effect of cloud formation processes on the stable isotopic composition of rainfall"*.

And C4, 1, 9-11: *"The cloud formation processes of these winter storms and convection will affect the isotopic composition of the resulting rainfall....."*

Response: This is indeed an aspect we neglect, and deserves some further discussion. The high level of agreement between the absolute values of the fluid inclusion data and modern precipitation isotope data make it likely that similar condensation processes are responsible for the MIS3 rainfall as are responsible today, making source effects the first order control on composition. Moreover, the modern precipitation and meteoric water lines derived from them already encompass the range of different condensation styles found in the modern Mediterranean. There are undoubtedly considerable further advances to be made from northern African speleothem fluid inclusion research, and we expect these nuances to be delineated by these future studies.

C4, 2, 3-7: *"I think the arguments made for the mixing of the different end members would be much more clear if the discussion of the stable isotopic composition and the d-excess would be combined including clear definitions of the values for the three depositional phases of the speleothem and the modern rainfall end members"*.

Response: If the Reviewer feels this will make our work more accessible, we will be very happy to follow their guidance.

C4, 3, 1-5 continuing to C5, 1, 1-7: *"There seems to be a discrepancy between some of the statements in the manuscript with regards to the Atlantic rainfall source..... I think this needs to be clarified..... First it is stated that increased convection during phases with a low precession parameter must be related to a northward shift of the ITCZ, then the convection is attributed to enhance internal convection."*

Response: We do not see the core discrepancy that troubles the Reviewer on this point. We can provide little positive evidence for Atlantic-sourced water in our record, and it is likely that other sites on the Mediterranean margin may show the same. Equally, it is undeniably true that to alter the Mediterranean freshwater budget the water must be external – and both the winter westerlies and the monsoon source much of that water from the Atlantic. We conclude that the key water responsible for Mediterranean freshening is likely to be arriving as runoff, not direct rainfall (which is not controversial – (Grant et al., 2016)). Primarily, we are aiming to warn colleagues working in the Mediterranean terrestrial sphere that their evidence of wet / dry changes may not relate directly to fresh / salty conditions in the Mediterranean Sea. Hopefully, this will become clearer with the restructuring this Reviewer recommends.

The suggestion that recycled water from the Sahara could be important is interesting. Water re-evaporated from rainfall in northern Africa should be isotopically light, reflecting this relatively depleted source. We do not see a population of depleted fluid inclusions that sit outside of the modern rainfall system that would suggest there is a substantial contribution from such a source.

C5, 4, 1-3: *"The last paragraph of the section starting in line 123 is not really about 'the central North African speleothem record'..... and should maybe be in its own chapter"*.

Response: agreed.

C5, 6, 1-3 (repeated in C5, 9, 1-2): *"The speleothems carbonate stable isotopic composition were published, so the sentence..... can be removed"*

Response: Agreed.

C5, 7,1-3: *"I think this section should include a clear definition of the three depositional phases, giving the range of fluid inclusion stable isotopes and d-excess. This would make the comparison much easier."*

Response: Agreed.

C5, 11, 1-2 continued to C6, 1, 1-8: *"Technical corrections"*.

Response: Agreed – these changes should be made.

C6 continued into C7: *"Figures"* (presentational and formatting considerations for figures 1, 4, 5, 6, 7 and 8).

Response: Agreed – these changes should be made.

## **Anonymous Reviewer 2**

C2, 2, 6-10: *"What I would like to see in the ms is an assessment of the extent to which the fluid inclusion isotope data are in isotopic equilibrium with the calcite..... While isotope equilibrium is not a given for many speleothems, at least some consistency is to be expected between  $d_{18O_{cc}}$  and  $d_{18O_{fi}}$ ."*

Response: We concur with the reviewer's sentiment that difficulties in the correlation of the time-series are "uncomfortable", and this is why we approach the data by analysing large groups of datapoints rather than using a time-series approach. We are extremely pleased that this reviewer also feels that collection of the fluid inclusion dataset is "technically sound" (C1, 2, 7), and that the general accuracy of the dataset is supported by our quantitative analysis of it compared to modern rainfall isotopes (C2, 1, 1-2). It can only be concluded that these data are a good representation of this isotopic system, even though they look unusual.

We suggest that the apparently poor correlation of time-series likely arises from aliasing of a complicated signal in the fluid inclusions, and emphasise that it is unlikely the data presented are sufficiently resolved to demonstrate the two datasets actually have different structure. The correlation of the datasets is therefore ambiguous, rather than disproven. Sadly, the at least order-of-magnitude increase in the size of the fluid inclusion dataset needed to resolve this point is not realistic: indeed, this Reviewer notes that this dataset is already *"comparatively large"* (C1, 2, 1). The most appropriate way forward in this situation is to minimise the interpretation of the temporal

structure of the fluid inclusion data we present, and this is what we have done. We are pleased that despite their discomfort, this Reviewer supports publication of this “*interesting study*” (C3, 3, 13-17).

A conventional equilibrium test is difficult to perform for this dataset, as each measurement comprises a mixture of inclusions with different compositions from each layer, and therefore an unknown position on the mixing line between these end members. Should we compute mean values (arithmetic or volume weighted); or extrapolate end members, and test for equilibrium of both? All these judgements require assumptions we are not in a position to make. Consequently, we are only able to test for equilibrium in the subset of samples where the end members are sufficiently close together for the analysis to be fully “duplicable” (the subset shown in Figure 6). Modern mean winter temperature in Dernah (the nearest city to Susah Cave) is 11.9°C, with maximum 17.7°C and minimum 7.1°C. These fluid inclusions are therefore certainly at least close to isotopic equilibrium with the carbonate hosting them.

Table R1

Duplicated Sample	Mean $\delta^{18}\text{O}_{\text{fi}}$	Mean $\delta^2\text{H}_{\text{fi}}$	Distance From Base (mm)	Age	$\delta^{18}\text{O}_{\text{cc}}$	Temperature (°C) <sup>1</sup>	Growth Phase
9	-4.35	-21.51	15-20	66394-66388	-3.78	13.3	I
11	-5.69	-19.95	56-61	66493-64393	-3.96	8.7	I
18	-5.92	-21.75	109-118	63487-63302	-3.98	7.9	I
23	-5.45	-25.62	260-267	59106-58490	-4.51	11.8	I
25	-5.38	-25.54	395-400	51687-51648	-5.21	15.0	II
33	-4.70	-26.72	556-561	37260-37221	-4.37	14.3	III
29	-5.08	-22.72	597-602	36957-36921	-4.6	13.7	III
35	-5.03	-21.37	652-657	36581-36284	-3.82	10.8	III
7	-4.96	-23.74	767-772	35688-35647	-4.32	13.1	III

<sup>1</sup> Calculated using the equation presented by (Sharp, 2017).

C2, 2, 17-19: “It would perhaps be useful if the authors discuss that a bit more in the context of the interpretation of their record”.

Response: Agreed.

C3, 1, 5-10: “*Bringing a third water source in, as is suggested in the ms, cannot really be supported by the data from my perspective..... One could perhaps argue that slight isotope changes within each of these moisture sources can cause similar isotope patterns?*”

Response:, The Reviewer agrees with our analysis that the data does show a mixing pattern of western and eastern Mediterranean sources (C3, 1, 4-5). So, we assume the ‘third source’ mentioned above is therefore the Atlantic external water we argue for, and find in relatively small amounts. We happily agree this is the most speculative part of our analysis. However, we also note

---

<sup>1</sup> Calculated using the equation presented by Sharp, Z.: Principles of stable isotope geochemistry, 2017.

that *finding no Atlantic moisture at all* in this dataset is a rather more startling interpretation than our suggestion that we find only a little. Atlantic-sourced moisture contributes to rainfall in central northern Africa today, and this mode of rainfall has previously been argued to be greater in past humid phases (Toucanne et al., 2015). We therefore find this point of speculation actually rather conservative in its nature.

C4, 2, 1-2: *"I'd like to know where your duplicable samples from Fig. 6 are located in the stalagmite (stratigraphically). All in one period, or distributed all over?"*

Response: See Table 1. All three Growth Phases are represented by at least one fully duplicated sample.

C4, 2, 2-4: *"Do you have a better correlation with the d18O values of the carbonate when you consider the duplicable dataset only?"*

Response: Beyond the differences between the three phases (see next response), it is difficult to judge whether there is true correlation between the reduced fluid inclusion dataset and the calcite isotope dataset, because the former is rather small. To make interpretations based on such a "correlation" would seem to us rather speculative. We are safer limiting the discussion of the time series.

C4, 2, 4-5: *"Further, I'd like to see if, based on the duplicable set only, one can still observe clear differences between the three wet intervals."*

Response: The sample from Phase II is more depleted both in  $\delta^{18}\text{O}_{\text{fi}}$  and  $\delta^{18}\text{O}_{\text{cc}}$  than any of the samples from Phases I and III, which show similar compositions. Our interpretation that the water driving this middle growth phase is different to the other two is therefore supported by this additional analysis.

C4 paragraphs 3, 4 and 5: *"Figures"*

Response: See response to Reviewer 1.

C4, 6, 1-2: Error in text.

Response: Correction will be made.

C4, 7, 1-5: *"It would be interesting to know..... could you have any sea spray effect?"*

Response: Given that we find no clear signal (or indeed, no variance beyond measurement uncertainty) in the Sr isotope record, these points cannot alter the interpretation and we are consequently unclear about their relevance (?).

C4, 8, 1-2 continued to C5, 1, 1-5: *"Towards the end of the discussion, d13CC plays an important role. These data, however, are not shown.....Shouldn't d18Ofi do the same as d2Hfi if your claim is correct?"*

Response:  $\delta^{13}\text{C}_{\text{cc}}$  is actually shown, within Figure 8b. We do make greater use of  $\delta^2\text{H}_{\text{fi}}$  towards the end of the discussion, because we are attempting to use the fluid inclusion data to better understand the carbonate isotope datasets, and hydrogen cannot be measured in carbonate and therefore provides valuable independent evidence of changes. As the fluid inclusion isotopes do correlate, it is indeed true that the  $\delta^{18}\text{O}_{\text{fi}}$  shows a similar pattern.

C5, 2, 1: *"Your statement in line 392 to 394 is not clear to me.*

Response: We have clarified that statement.

## **References**

Black, E., Brayshaw, D. J., and Rambeau, C. M. C.: Past, present and future precipitation in the Middle East: Insights from models and observations, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 368, 5173-5184, 10.1098/rsta.2010.0199, 2010.

Brun, A.: Reflections on the pluvial and arid periods of the Upper Pleistocene and of the Holocene in Tunisia, *Palaeoecology of Africa and the surrounding islands*. Vol. 22. Proc. symposium on African palynology, Rabat, 1989, 157-170, 1991.

Cancellieri E., Cremaschi M., Zerboni A., and S., d. L.: Climate, Environment, and Population Dynamics in Pleistocene Sahara, in: *Africa from MIS 6-2. Vertebrate Paleobiology and Paleoanthropology.*, edited by: Jones S., and B., S., Springer, Dordrecht, 2016.

Celle-Jeanton, H., Zouari, K., Travi, Y., and Daoud, A.: Caractérisation isotopique des pluies en Tunisie. Essai de typologie dans la région de Sfax, *Sciences de la Terre et des planètes*, 333, 625-631, 2001.

Dayan, U., Nissen, K., and Ulbrich, U.: Review Article: Atmospheric conditions inducing extreme precipitation over the eastern and western Mediterranean, *Nat. Hazards Earth Syst. Sci.*, 15, 2525-2544, 10.5194/nhess-15-2525-2015, 2015.

Gat, J. R., Klein, B., Kushnir, Y., Roether, W., Wernli, H., Yam, R., and Shemesh, A.: Isotope composition of air moisture over the Mediterranean Sea: An index of the air-sea interaction pattern, *Tellus, Series B: Chemical and Physical Meteorology*, 55, 953-965, 10.1034/j.1600-0889.2003.00081.x, 2003.

Grant, K. M., Grimm, R., Mikolajewicz, U., Marino, G., Ziegler, M., and Rohling, E. J.: The timing of Mediterranean sapropel deposition relative to insolation, sea-level and African monsoon changes, *Quaternary Science Reviews*, 140, 125-141, <http://dx.doi.org/10.1016/j.quascirev.2016.03.026>, 2016.

Langgut, D., Almogi-Labin, A., Bar-Matthews, M., Pickarski, N., and Weinstein-Evron, M.: Evidence for a humid interval at ~56–44 ka in the Levant and its potential link to modern humans dispersal out of Africa, *Journal of Human Evolution*, 124, 75-90, 10.1016/j.jhevol.2018.08.002, 2018.

Nicoll, K.: A revised chronology for Pleistocene paleolakes and Middle Stone Age – Middle Paleolithic cultural activity at Bîr Tîrfawi – Bîr Sahara in the Egyptian Sahara, *Quaternary International*, 463, 18-28, <https://doi.org/10.1016/j.quaint.2016.08.037>, 2018.

Peleg, N., and Morin, E.: Convective rain cells: Radar-derived spatiotemporal characteristics and synoptic patterns over the eastern Mediterranean, *J. Geophys. Res. D Atmos.*, 117, 10.1029/2011JD017353, 2012.

Revel, M., Ducassou, E., Grousset, F. E., Bernasconi, S. M., Migeon, S., Revillon, S., Mascle, J., Murat, A., Zaragosi, S., and Bosch, D.: 100,000 Years of African monsoon variability recorded in sediments of the Nile margin, *Quaternary Science Reviews*, 29, 1342-1362, 10.1016/j.quascirev.2010.02.006, 2010.

Rohling, E. J.: Environmental control on Mediterranean salinity and delta O-18, *Paleoceanography*, 14, 706-715, 1999.

Sharp, Z.: *Principles of stable isotope geochemistry*, 2017.

Smith, J. R., Hawkins, A. L., Asmerom, Y., Polyak, V., and Giegengack, R.: New age constraints on the Middle Stone Age occupations of Kharga Oasis, Western Desert, Egypt, *Journal of Human Evolution*, 52, 690-701, 2007.

Toucanne, S., Angue Minto'o, C. M., Fontanier, C., Bassetti, M.-A., Jorry, S. J., and Jouet, G.: Tracking rainfall in the northern Mediterranean borderlands during sapropel deposition, *Quaternary Science Reviews*, 129, 178-195, <http://dx.doi.org/10.1016/j.quascirev.2015.10.016>, 2015.



1 Enhanced Mediterranean water cycle  
2 explains increased humidity during MIS 3  
3 in North Africa  
4

5 Mike Rogerson<sup>1</sup>

6 Yuri Dublyansky<sup>2</sup>

7 Dirk L. Hoffmann<sup>3</sup>

8 Marc Luetscher<sup>2,4</sup>

9 Christoph Spötl<sup>2</sup>

10 Paul Töchterle<sup>2</sup>

11 1 School of Environmental Sciences, University of Hull, Cottingham Road, Hull, HU6 7RX, UK.

12 2 Institute of Geology, University of Innsbruck, Innrain 52, 6020 Innsbruck, Austria.

13 3 Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher  
14 Platz 6, 04103, Leipzig, Germany

15 4 Swiss Institute for Speleology and Karst Studies (ISSKA), Serre 68, CH-2300 La Chaux-de-Fonds

16

## 17 Abstract

18 *We report a new fluid inclusion dataset from Northeast Libyan speleothem SC-06-01, which is the*  
19 *largest speleothem fluid inclusion dataset for North Africa to date. The stalagmite was sampled in*  
20 *Susah cave, a low altitude coastal site, in Cyrenaica, on the northern slope of the Jebel Al-Akhdar.*  
21 *Speleothem fluid inclusions from latest Marine Isotope Stage (MIS) 4 and throughout MIS 3 (~67 to*  
22 *~30 ka BP) confirm the hypothesis that past humid periods in this region reflect westerly rainfall*  
23 *advected through the Atlantic storm track. However, most of this moisture was sourced from the*  
24 *Western Mediterranean, with little direct admixture of water evaporated from the Atlantic. Moreover,*  
25 *we identify a second moisture source likely associated with enhanced convective rainfall within the*  
26 *Eastern Mediterranean. The relative importance of the western and eastern moisture sources seems to*  
27 *differ between the humid phases recorded in SC-06-01. During humid phases forced by precession,*  
28 *fluid inclusions record compositions consistent with both sources, but the 52.5 – 50.5 ka interval*  
29 *forced by obliquity reveals only a western source. This is a key result, showing that although the*  
30 *amount of atmospheric moisture advections changes, the structure of the atmospheric circulation over*  
31 *the Mediterranean does not fundamentally change during orbital cycles. Consequently, an arid belt*  
32 *must have been retained between the Intertropical Convergence Zone and the mid-latitude winter*  
33 *storm corridor during MIS 3 pluvials.*

## 34 Introduction

35 Atmospheric latent heat is a major component of global and regional climate energy budgets and  
36 changes in its amount and distribution are key aspects of the climate system (Pascale et al., 2011).  
37 Equally, in mid- and low-latitude regions, changes in the water cycle have more impact on landscapes  
38 and ecosystems than changes in sensible heat (Black et al., 2010). Rainfall in semi-arid regions is thus  
39 one of the key climate parameters that understanding future impact on human societies depends upon  
40 (IPCC, 2014), making constraining of mid-latitude hydrology a globally significant research priority.  
41 These regions, however, have a particularly sparse record of palaeoclimate due to typically poor  
42 preservation of surface sedimentary archives (Swezey, 2001). North Africa is a region that fully

43 exhibits these limitations, and large areas present either no pre-Holocene record or else they present  
44 highly discontinuous deposits indicating major reorganisation of the hydroclimate, which are  
45 challenging to date (Armitage et al., 2007). North Africa also fully exhibits the progress  
46 palaeoclimatologists have made in understanding continental hydrological change from its impact on  
47 the marine system; our understanding of past North African hydroclimate is disproportionately drawn  
48 from records from the Mediterranean Sea (Rohling et al., 2015) and the eastern Central Atlantic  
49 (Goldsmith et al., 2017; deMenocal et al., 2000.; Adkins et al., 2006).

50

### 51 Past changes in North African hydroclimate

52 Marine-based evidence offers a coherent model in which changes in the spatial distribution of  
53 insolation alter atmospheric circulation on orbital timescales ( $10^4$  to  $10^5$  years) and force major  
54 reorganisations of rainfall in semi-arid regions such as the Sahel and southern Saharan regions  
55 (Rohling et al., 2015; Goldsmith et al., 2017). This result is at least partially confirmed in climate  
56 modelling experiments (Bosmans et al., 2015; Tuenter et al., 2003) and provides a conceptual  
57 framework in which fragmentary evidence of hydrological change on the adjacent continent can be  
58 understood (Rowan et al., 2000). There is 1) strong geochemical evidence that runoff from the  
59 African margin initiated the well-known “sapropel” thermohaline crises of the eastern Mediterranean  
60 (Osborne et al., 2010; Osborne et al., 2008) and, 2) convincing evidence that the southern margin of  
61 the Mediterranean was more variable than the northern in terms of the relative magnitude of  
62 precipitation changes and the distribution of flora, fauna and hominid populations (Drake et al., 2011).  
63 However, we emphasise the fact that this understanding is largely drawn from evidence from outside  
64 continental North Africa, and that this limits our knowledge about the nature and impact of  
65 hydrological changes in this region.

66 There is strong evidence for a more humid climate throughout the Sahara and Sahel regions during the  
67 early Holocene (Gasse and Campo, 1994; Gasse, 2002; Fontes and Gasse, 1991; Prentice and Jolly,  
68 2000; Jolly et al., 1998; Collins et al., 2017), and in older interglacial periods (Drake et al.,

69 2008;Armitage et al., 2007;Vaks et al., 2013). This evidence has been interpreted to indicate that  
70 humid conditions extended from the modern Sahel (~15°N) to the Mediterranean coast (30-35°N).  
71 However, this only partially agrees with model results, which do indicate orbitally forced migration of  
72 the monsoon belt but not across such a large spatial scale as suggested by the empirical data. Model  
73 experiments indicate that monsoonal rainfall occurring within the Intertropical Convergence Zone  
74 (ITCZ) likely extended no further north than ~23°N (Harrison et al., 2015). This well-recognised lack  
75 of agreement between rainfall fields in model experiments for the past and reconstructed  
76 hydrographies from the distribution of lakes and vegetation (via pollen) (Peyron et al., 2006) remains  
77 ~~ana~~ a major research problem. While some models also suggest that during times of high Northern  
78 Hemisphere insolation, enhanced westerlies advected Atlantic moisture into the basin (Brayshaw et  
79 al., 2009;Tuenter et al., 2003;Bosmans et al., 2015), high-resolution regional modelling indicates that  
80 this primarily affected the northern Mediterranean margin only (Brayshaw et al., 2009). This result is  
81 consistent with evidence of enhanced runoff at these times from the southern margin of Europe  
82 (Toucanne et al., 2015). On the African coast east of Algeria, the southern limit of enhanced  
83 precipitation arising from increased westerly activity within model experiments essentially lies at the  
84 coastline (~32°N), and does not appear to drive terrestrial hydrological changes. Overall, there is  
85 therefore a striking mismatch between the apparent humidity of Africa between 23 and 32°N in the  
86 empirical record (a zonally oriented belt ~1000 km in width) and the climate models. This region  
87 encompasses southern Tunisia, in which multiple lines of evidence for distinct and widespread  
88 periods of increased humidity provide a highly secure basis for enhanced rainfall during Northern  
89 Hemisphere insolation maxima (Ballais, 1991;PETIT-MAIRE et al., 1991), the Fezzan basin, in  
90 which compelling evidence for multiple lake highstands exists (Drake et al., 2011) and western Egypt,  
91 where large tufa deposits attest to higher past groundwater tables (Smith et al., 2004).

92 An emerging picture of MIS 3 as a humid period within the Mediterranean basin is developing  
93 (Langgut et al., 2018), and the current study focusses on this time period. However, MIS3 is not well  
94 expressed in the Sahara region. The Libyan interior is considered arid or hyperarid throughout the last  
95 glaciation (Cancellieri E. et al., 2016). Recent re-evaluation of lake levels in southwest Egypt

106 indicates a groundwater fed system was active around 41 ka (Nicoll, 2018), which is similar to dates  
107 for springline tufa systems at Kharga Oasis (Smith et al., 2007). We are not aware of continental  
108 MIS3 pollen records from the region, but marine pollen from Tunisia indicates more arid conditions  
109 through the last glacial than during the Holocene (Brun, 1991). There is a triple peak in runoff from  
110 the Nile recorded in the marine sediment record, with maxima at ~60, ~55 and ~35ka, indicating  
111 higher rainfall within the upper Nile catchment (Revel et al., 2010).

112 It is unlikely that significant further progress will be made in understanding the palaeoclimate of  
113 North Africa without new empirical evidence of regional hydrological changes from which  
114 atmospheric dynamics can be delineated.

#### 115 The central North African speleothem record

116 Speleothem palaeoclimatology has high potential for North Africa, but is only recently becoming  
117 established through key records developed for Morocco (Wassenburg et al., 2013; Ait Brahim et al.,  
118 2017; Wassenburg et al., 2016). Until recently, the only speleothem record published from central  
119 North Africa was a single continuous record from 20 to 6 ka BP from northern Tunisia (Grotte de la  
120 Mine). This record shows a large deglacial transition in both  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  (Genty et al., 2006), with  
121 oxygen isotopes indicating a 2-step change from a relatively isotopically heavy (-5‰) LGM (20-16 ka  
122 BP), through an intermediate (-6 to -7‰) deglacial period (16-11.5 ka BP) to a relatively isotopically  
light early Holocene. The  $\delta^{13}\text{C}$  record indicates cool periods exhibiting higher carbon isotope values,  
more clearly delineating the Bølling-Allerød / Younger Dryas oscillation than  $\delta^{18}\text{O}$ . This is assumed  
to reflect higher soil respiration during warm periods (Genty et al., 2006). A major change in the  
carbon isotopic composition occurred across the transition from the relatively arid glacial to the more  
humid Early Holocene, and indicates a significant reorganisation of the regional hydroclimate.  
However, it is difficult to interpret these data in isolation. A recently reported speleothem record (SC-  
06-01) indicates that conditions in northern Libya during Marine Isotope Stage 3 (MIS 3) were more  
humid than today, and shows isotopic evidence of a teleconnection between temperature in Greenland  
and rainfall at the southern Mediterranean margin (Hoffmann et al., 2016). The oxygen isotope record  
indicates that the water dripping into the cave during MIS 3 was isotopically too heavy for the

123 moisture to be sourced from within the monsoon system (Hoffmann et al., 2016). However, beyond  
124 ruling out a southern source  $\delta^{18}\text{O}_{\text{cc}}$  values alone are not sufficient to determine the origin of  
125 atmospheric vapour. Three distinct humid phases within MIS3 are reported from this speleothem: 65-  
126 61 ka, 52.5-50.5 ka and 37.5-33 ka. Phases I and III occur during times of low precession parameter,  
127 when summer insolation on the northern hemisphere is relatively increased. Phase II represents the  
128 first evidence for high obliquity being able to cause a pluvial period in the north African subtropics in  
129 the same manner as precession (Hoffmann et al., 2016). In SC06-01, all three growth phases are  
130 fractured into multiple short periods of growth, and show a marked temporal coherence with  
131 Greenland Dansgaard-Oeschger interstadials (Hoffmann et al., 2016). Here, we report fluid inclusion  
132 data from this speleothem and discuss how this helps resolve some of the issues discussed above.

### 133 Fluid Inclusions

134 Speleothem fluid inclusions are small volumes of water that were enclosed between or within calcite  
135 crystals as they grew, ranging in size from less than 1  $\mu\text{m}$  to hundreds of  $\mu\text{m}$  (Schwarcz et al., 1976).  
136 This water represents quantities of ancient drip-water that can be interrogated directly to ascertain the  
137 isotopic properties of the oxygen ( $\delta^{18}\text{O}_{\text{fi}}$ ) and hydrogen ( $\delta^2\text{H}_{\text{fi}}$ ) it comprises. This powerful approach  
138 circumvents some of the uncertainty inherent in the interpretation of the stable isotopic values  
139 preserved in the calcite comprising the speleothem itself ( $\delta^{18}\text{O}_{\text{cc}}$ ,  $\delta^{13}\text{C}_{\text{cc}}$ ). Fluid inclusion isotopes have  
140 been used to demonstrate changes in air temperatures (Wainer et al., 2011; Meckler et al.,  
141 2015; Arienzo et al., 2015) and in the origin of the moisture from which precipitation was sourced  
142 (McGarry et al., 2004; Van Breukelen et al., 2008). Fluid inclusions from speleothems in Oman have  
143 also been used to identify monsoon-sourced precipitation during interglacial phases (Fleitmann et al.,  
144 2003), providing a rationale for similar investigation of fluid inclusion isotope behaviour in North  
145 Africa.

146 In the case of fluid inclusions from northeastern Libyan speleothems, the boundary conditions for  
147 atmospheric moisture supply are 1) the sea surface temperature of the Atlantic and Mediterranean, 2)  
148 the surface water  $\delta^{18}\text{O}_{\text{sw}}$  of the same ocean regions, 3) land surface temperature of Africa and to a

149 lesser extent southern Europe, 4) insolation (especially with respect to ITCZ position) and 5) the zonal  
150 pressure gradient across northern Africa.

### 151 **Modern rainfall system**

152 Modern rainfall in central North Africa is dominated by relatively wet winters, and summers with  
153 little, if any, precipitation. Convective systems, cyclones, upper-level troughs and static instabilities  
154 can all drive rainfall patterns in the Mediterranean basin and these modes are reviewed in (Dayan et  
155 al., 2015). Convection essentially reflects the relatively high SST of the Mediterranean during the  
156 winter, but rising air masses generally also need significant advection of moisture to drive significant  
157 rainfall. Upper level troughs reflect large-scale circulation (e.g. Red Sea Trough) or reflect lee effects  
158 downstream of mountains in the western Mediterranean, and promote rainfall in their regions of  
159 formation. The dominant cyclogenetic centre is in the Gulf of Genoa, and secondary centres are placed  
160 in south Italy, Crete and Cyprus. Cyclonic systems can also penetrate from the Atlantic, where the  
161 high SST of the winter Mediterranean tends to sustain and amplify them, in close analogy to  
162 convection forcing. The key static instability is the penetration of the tropical air mass into the  
163 subtropical Mediterranean, forming a ‘Saharan Cloud Band’ at middle and upper atmospheric levels.  
164 These originate from within the ITCZ. Libya is very sparsely instrumented, so we assume that  
165 synoptic processes are similar to the Levant region. Here, most rainfall falls under winter, low  
166 pressure conditions, and is convective (Peleg and Morin, 2012). The responsible low pressure systems  
167 can relate to transient, shallow lows north of the area in which rainfall is occurring, or less frequently  
168 more long-lasting Cyprus Lows or Red Sea Trough systems (Peleg and Morin, 2012).

### 169 Material and Methods

170 SC-06-01 is a 93-cm long stalagmite from Susah Cave (Fig. 1, 32°53.419' N, 21°52.485' E), which  
171 lies on a steep slope ~200 m above sea level in the Al Akhdar massif in Cyrenaica, Libya (Fig. 1). The  
172 region is semi-arid today, with mean annual temperature ~20°C and receiving less than 200 mm  
173 precipitation per year, mostly in the winter (October to April). The Al Akhdar massif has thin soil  
174 cover and a Mediterranean “maquis” vegetation. Susah Cave is hydrologically inactive today, and all

175 formations are covered with dust. The chronology of the speleothem and the general features of its  
176 growth and  $\delta^{18}\text{O}_{\text{cc}}$  record are published elsewhere (Hoffmann et al., 2016), and this study focuses on  
177 fluid inclusion isotopes, their impact on the interpretation of  $\delta^{18}\text{O}_{\text{cc}}$  and to a lesser extent on  $\delta^{13}\text{C}_{\text{cc}}$  and  
178 Sr isotopes.

179 ~~Calcite isotopes were measured using a ThermoFisher Delta<sup>plus</sup>XL isotope ratio mass spectrometer~~  
180 ~~(IRMS) equipped with a Gasbench II interface at the University of Innsbruck, according to standard~~  
181 ~~methods (Spötl, 2011).~~ Fluid inclusions were examined in doubly-polished thick section (100  $\mu\text{m}$ )  
182 slides, using a Nikon Eclipse E400 POL microscope. The isotope composition of fluid inclusion water  
183 was measured at the University of Innsbruck using a Delta V Advantage IRMS coupled to a Thermal  
184 Combustion/Elemental Analyser and a ConFlow II interface (Thermo Fisher) using the line, crusher  
185 and cryo-focussing cell described in Dublyansky and Spötl (2009). Samples were cut with a diamond  
186 band saw along visible petrographic boundaries in the speleothem, and therefore represent specific  
187 growth increments. Samples were analysed at least in duplicate, with the standard sampling protocol  
188 used on the Innsbruck instrument (Dublyansky and Spötl, 2009). To exclude the possibility of post-  
189 depositional diagenetic alteration, petrographic thin sections were investigated using transmitted-light  
190 microscopy. Results are detailed in Supplemental Information 1.

191 Optical emission spectroscopy (OES) was used to measure a variety of elemental concentrations,  
192 including Sr, along the main growth axis of SC-06-01. The low spatial resolution of trace elemental  
193 analyses (every 10 mm) does not allow to investigate time series of elemental variation but was useful  
194 to assess Sr contents of the samples for Sr isotope measurements by thermal ionisation mass  
195 spectrometry (TIMS). The samples for TIMS analyses were drilled using a hand held micro drill with  
196 a tungsten carbide drill bit. Sample sizes range between 2 and 4 mg, thus we achieved a minimum Sr  
197 load of 100 ng on the Re filaments for TIMS. Chemical sample preparation and subsequent TIMS  
198 measurement were done following standard protocols (Charlier et al., 2006). No spike was added to  
199 the samples prior to chemical purification. The Sr isotope measurements were done on a Triton TIMS  
200 housed at the Bristol Isotope Group laboratory, University of Bristol.



201 Results

202 Fluid inclusions

203 Petrographic analysis of the thick sections indicates that the distribution of fluid inclusions is highly  
204 variable, with macroscopically opaque “milky” calcite typical of rapidly growing intervals containing  
205 sometimes very abundant inclusions and the discoloured, translucent calcite of the slowly growing  
206 intervals being almost inclusion-free (Fig. 2). In most samples, two distinct populations of inclusions  
207 were identified with numerous small intra-crystalline inclusions and larger, but less frequent, inter-  
208 crystalline inclusions. Consequently, the volume of water analysed per sample was very variable (Fig.  
209 3). Indeed, a significant proportion of individual fluid inclusion measurements had analyte volumes  
210 too small (<0.1  $\mu\text{L}$ ) to have confidence in the isotope results. A small number of analyses failed due  
211 to excessive water saturating the detector, and these have not been included in the datasets presented  
212 here. The major impact of the highly variable availability of inclusions in the speleothem is a  
213 significant bias in the analyses towards the most rapidly growing, and therefore probably humid, time  
214 periods. Three rapidly-growing phases are reported in SC-06-01, named Phase I (62-67ka), Phase II  
215 (53-50 ka) and Phase III (37-33 ka) (Hoffmann et al., 2016). Fluid inclusions for Phases I and II are  
216 isotopically similar (with  $\delta^{18}\text{O}_{\text{FI}}$  ranging from -7.5 ‰ to -3.8 ‰ and from -8.5 ‰ to -3.2‰  
217 respectively and  $\delta^2\text{H}_{\text{FI}}$  ranging from -26.7 ‰ to -18.6 ‰ and from -29.4 ‰ to -16.1 ‰ respectively).  
218 However, compositions for Phase II are different, particularly with respect to deuterium ( $\delta^{18}\text{O}_{\text{FI}}$   
219 ranging from -8.9 ‰ to -4.5 ‰ and  $\delta^2\text{H}_{\text{FI}}$  ranging from -38.3 ‰ to -25.1 ‰).

220 In most samples, achieving within-error replication ( $\delta^2\text{H} \pm 1.5\text{‰}$ ,  $\delta^{18}\text{O}: \pm 0.5\text{‰}$ ) of both  $\delta^{18}\text{O}_{\text{fi}}$  and  
221  $\delta^2\text{H}_{\text{fi}}$  was difficult. This must reflect more than one population of inclusions with different properties  
222 being present within at least some samples, and each replicate analysis represents some proportion of  
223 mixing between these populations. This suggests significant short-term variability in the composition  
224 of the water stored in the presumably rather small soil/epikarst zone overlying the cave.

225 Consequently, any given time interval risks being under-sampled with regard to variability at that  
226 time. Although there is some visual correspondence between the  $\delta^{18}\text{O}_{\text{fi}}$ ,  $\delta^2\text{H}_{\text{fi}}$  and  $\delta^{18}\text{O}_{\text{cc}}$  data series

227 (Fig. 4), it seems that the fluid inclusion time series risks aliasing changes seen in the calcite  
228 isotope time series. Consequently, the usefulness of interpretation that can be drawn from the episodic  
229 SC-01-06 fluid inclusion dataset when arranged as a time series is limited. We and we therefore  
230 largely limit our discussion to the properties of the population of waters as a full dataset. This  
231 approach minimises the impact the different populations can have on interpretation.

232 Figure 5 shows the SC-06-01 fluid inclusion dataset alongside Global Natural Network of Isotopes in  
233 Precipitation (GNIP) datasets from Tunis World Meteorological Office (WMO station 6071500), Sfax  
234 (6075000) and Bet Dagan (4017900) (locations in Fig. 1) and other published precipitation datasets.  
235 The Tunisian datasets fit within a trend typical of the Global Meteoric Water Line (GMWL) ( $\delta^2\text{H} =$   
236  $8\delta^{18}\text{O} + 10$ ). However, all this data lies along a single moisture evolution trend, and the Tunis and  
237 Sfax populations overlap. The data from Bet Dagan exhibits a trend which is extremely close to being  
238 parallel to the global trend dominating in Tunisia, but translated by +10 ‰ in  $\delta^2\text{H}$ , reflecting greater  
239 deuterium excess. This is typical of the Mediterranean Meteoric Water Line (MMWL) (Ayalon et al.,  
240 1998; Gat et al., 2003), and reflects internal recycling of water with consequent deuterium enrichment  
241 in the eastern Mediterranean and its bordering continental areas.

242 The values of  $\delta^2\text{H}_f$  and  $\delta^{18}\text{O}_f$  fit within the range of values for modern precipitation, giving  
243 confidence that these measurements do reflect past precipitation composition despite the influence of  
244 multiple inclusion populations. The lack of apparent scatter towards positive  $\delta^{18}\text{O}$  values both in the  
245 precipitation and fluid inclusion datasets further indicates that the data represent little-altered  
246 precipitation values, and that surface re-evaporation was minor at least during humid phases.  
247 However, the range of fluid inclusion values is inconsistent with either an exclusively Tunis-type or  
248 an exclusively Bet Dagan-type moisture source for precipitation in Cyrenaica during MIS 3. Even  
249 when all but the subset of fluid inclusion analyses who replicates are similar are excluded (Fig. 6), the  
250 population is split between the Tunisian and Israeli precipitation end-members.

251

252 **Strontium isotopes**

253 The  $^{87}\text{Sr}/^{86}\text{Sr}$  signal in the SC-06-01 record is rather invariable (Fig. 7), with all analyses indicating  
254 values within analytical error. Mean values vary between 0.708275 and 0.708524 and although there  
255 is an apparent trend from maxima at 34 and 64 ka BP with a minimum at 52 ka BP, which mimics the  
256 precession history, this is too weak to be significant relative to the error.

257 **Calcite carbon isotopes**

258 Both  $\delta^{13}\text{C}_{\text{cc}}$  and  $\delta^{18}\text{O}_{\text{cc}}$  show similar trends throughout the record (Fig. 8), indicating that depleted  
259 oxygen isotopes coincide with depleted carbon isotope values. This does not appear to arise from  
260 fractionation on the speleothem surface (Hoffmann et al., 2016), and so represents changes in soil  
261 bioproductivity acting in concert with changes in precipitation.

262 **Discussion**

263 **Moisture advection during Libyan humid phases**

264 The range of values of both individual and replicated fluid inclusion measurements can only be  
265 reconciled with multiple moisture sources. Most of the fluid inclusion data cluster between the  
266 weighted mean value for precipitation collected at Sfax with a mixed source from the Atlantic and  
267 western Mediterranean, (“Sfax Mixed”  $\delta^{18}\text{O}_{\text{ppt}} = -4.93 \text{ ‰}$ ,  $\delta^2\text{H}_{\text{ppt}} = -26 \text{ ‰}$ ; Fig. 9) and High

268 Precipitation events at Bet Dagan ( $\delta^{18}\text{O}_{\text{ppt}} = -6.33 \text{ ‰}$ ,  $\delta^2\text{H}_{\text{ppt}} = -21.46 \text{ ‰}$ ; Fig. 9). ~~This value is  
269 representative of many of the largest individual precipitation events at Sfax in the period 1992–1999  
270 associated with a Western Mediterranean moisture source (Celle Jeanton et al., 2001).~~ However, the  
271 fluid inclusion data cluster also extends to the end member reflecting pure western Mediterranean  
272 sources at Sfax ( $\delta^{18}\text{O}_{\text{ppt}} = -3.99 \text{ ‰}$ ,  $\delta^2\text{H}_{\text{ppt}} = -20.3 \text{ ‰}$ ; Fig. 9), indicating a third end member  
273 composition with higher  $\delta^{18}\text{O}_{\text{ppt}}$ . ~~Consequently, we consider that this data reflects a dynamic balance  
274 of moisture sources contributing to rainfall in Cyrenaica which resembles modern precipitation in  
275 Tunisia and Israel in roughly equal proportions.~~

276 The weighted mean value for Atlantic-sourced precipitation events in Sfax ( $\delta^{18}\text{O}_{\text{ppt}} = -6.7 \text{ ‰}$ ,  $\delta^2\text{H}_{\text{ppt}} =$   
277  $-37.7 \text{ ‰}$ ) is distant from any observed fluid inclusion value (Fig. 9). ~~Likewise, compositions similar to~~  
278 ~~the high amount Atlantic-sourced rainfall events in Sfax ( $\delta^{18}\text{O}_{\text{ppt}} = -8 \text{ ‰}$ ,  $\delta^2\text{H}_{\text{ppt}} = -46 \text{ ‰}$ ) are not~~  
279 ~~reflected in the fluid inclusion data in Figure 9 suggesting a relatively low admixture of water from~~  
280 ~~this source-9).~~ A simple 3-end-member unmixing of fluid inclusion isotope values using the  
281 quantitative approach of (Rogerson et al., 2011) indicates that Atlantic-sourced water supplied no  
282 more than 15 % of the mass for any given fluid inclusion analysis. However, the coherence of fluid  
283 inclusion isotope ratios with the weighted mean of “mixed” Atlantic and Mediterranean precipitation  
284 at Sfax suggests that this small Atlantic influence is nevertheless persistent, and this must reflect  
285 synoptic westerly storms (Celle-Jeanton et al., 2001).

286 The simplest interpretation of the Susah Cave fluid inclusion data is therefore that they reflect a  
287 dynamic balance of moisture sources contributing to rainfall in Cyrenaica which resembles modern  
288 precipitation in Tunisia and Israel in roughly equal proportions. An alternative way to explain the  
289 trend of some points towards enriched  $\delta^{18}\text{O}$  values on the GMWL would be the temperature-  
290 dependent fractionation that would be caused by a shift to summertime precipitation. We do not  
291 favour this explanation, as it requires a more fundamental reorganisation of regional atmospheric  
292 circulation than our suggestion that the winter storms observed today penetrated further east in the  
293 past.

294 ~~Within the data presented in Figure 9, the Phase II fluid inclusions are exceptional, because none~~  
295 ~~show compositions consistent with a Bet Dagan source. Indeed, all the measurements for this period~~  
296 ~~resemble GMWL compositions. This seems to reflect a fundamental difference between this period~~  
297 ~~and Phases I and III, where all precipitation is drawn from synoptic westerly storms in the winter.~~  
298 ~~Consequently, it would seem that during the Obliquity forced period of humidity the Israeli mode~~  
299 ~~precipitation did not occur in the manner that it did during both Precession forced periods of~~  
300 ~~humidity.~~

301 Although the isotopic composition of Mediterranean water will have been more enriched during MIS  
302 3 due to ice-volume effects and increased Mediterranean water residence time (Rohling and Bryden,  
303 1994), the similar mean values of the SC-06-01 fluid inclusion waters compared to modern  
304 precipitation indicates the meteoric waterline at this time was not displaced to more enriched isotope  
305 values. This could reflect balancing of source water effects by changes in kinetic fractionation during  
306 evaporation (Goldsmith et al., 2017), which is controlled by normalised relative humidity. This would  
307 imply that the Mediterranean air masses were less saturated with moisture than today during MIS 3,  
308 which is consistent with the high deuterium excess  $\delta^2\text{H}_{\text{excess}}$  values found in some fluid inclusion  
309 samples (Fig. 10), but is difficult to reconcile with the increased precipitation recorded in SC-06-01.

310 In addition, changes in cloud height and cloud formation processes could possibly alter the isotopic  
311 fractionation in the atmosphere. Alternatively, the source water effect may be countered by increased  
312 runoff from the margins of the Mediterranean supplying isotopically depleted water to evaporating  
313 surface water. Isotopic “residuals” consistent with this argument are identified throughout MIS 3 in  
314 the eastern Mediterranean marine core LC21 (Grant et al., 2016), and this is also consistent with  
315 higher rainfall in Cyrenaica. We therefore favour the latter explanation.

316 ~~We conclude~~Although we find that our results likely reflect patterns of atmospheric transport in MIS3  
317 comparable to today, it is possible that some moisture was drawn from re-evaporation of monsoon  
318 rain falling further south, with no modern analogue in the region (Aggarwal et al., 2016). This water  
319 would likely be extremely isotopically light, reflecting both monsoon-type compositions and further  
320 fractionation during secondary evaporation. Moreover, a shift to more southerly-sourced regions is  
321 inconsistent with Sr-isotope data from Susah Cave. Sr-isotopes are known to be sensitive to changes  
322 in transport of Saharan dust (Frumkin and Stein, 2004), but even considering the most slow-growing  
323 and most rapidly-growing parts of SC-06-01, no significant difference in  $^{87}\text{Sr}/^{86}\text{Sr}$  has been identified.  
324 Although at times of extreme rainfall in the region, Saharan / Sahellian dust production is suppressed,  
325 this is not true during MIS3 (Collins et al., 2013). It seems that despite changes in the intensity of  
326 moisture transport during the period 65-30 ka BP, there is no large-scale change in atmospheric dust  
327 transport direction. This further supports our conclusion from the fluid inclusions that the Eastern

328 Mediterranean rainfall operating during precession parameter minima reflects enhanced internal  
329 convection rather than transport of moisture from the east or south with an atmospheric circulation  
330 pattern that prevails today.

331 Different sources at different times?

332 Phase II fluid inclusions are exceptional, because none show compositions consistent with a Bet  
333 Dagan source. This is most clearly reflected in the  $\delta^2\text{H}_{\text{excess}}$  values (Fig. 10), which show consistently  
334 low values across Phase II comparing well to the Western water end-member (~10 ‰) and not the  
335 Eastern water end-member (~30 ‰). The lack of Eastern water during Phase II seems to reflect a  
336 fundamental difference between this period and Phases I and III, as during this time all precipitation  
337 was drawn from synoptic westerly storms in the winter. Consequently, it would seem that during the  
338 Obliquity-forced period of humidity, the Israeli-mode precipitation did not occur in the manner that it  
339 did during both Precession-forced periods of humidity. This difference in the origin of the moisture  
340 feeding rainfall may explain the difference in average  $\delta^{18}\text{O}_{\text{cc}}$  during these different phases (Hoffmann  
341 et al., 2016), and why during some periods in Susah Cave show strong correlation with North Atlantic  
342 temperature, whereas others do not (Hoffmann et al., 2016).

343 Palaeoclimatological significance

344 ~~most~~Most of the precipitation supplied to Cyrenaica during MIS 3 was sourced from within the  
345 Mediterranean basin, which exhibited a similar meteoric water cycle to that observed today, albeit  
346 with more freshwater influence. This is a critical observation, as ~~internally-cycled water cannot alter~~  
347 ~~the basin-scale hydrological balance and therefore is a minor influence on deep convection in the~~  
348 ~~Mediterranean Sea (Bethoux and Gentili, 1999). The~~the precipitation feeding runoff must be  
349 externally sourced if it is to materially change Mediterranean functioning, as is observed during  
350 sapropel events (Rohling et al., 2015). ~~As most of the precipitation identified in SC-06-01 is sourced~~  
351 ~~internally to the Mediterranean, only the small, Atlantic-sourced portion of this water can be assumed~~  
352 ~~to play a role in~~The internally-cycled water we report from Susah Cave cannot alter the basin-scale  
353 hydrological balance, and therefore is a minor influence on deep convection in the Mediterranean Sea  
354 (Bethoux and Gentili, 1999): put simply, this means evidence of increased rainfall in the coastal

355 Mediterranean ~~freshening. This conclusion is likely transferable to any site on the continental margins~~  
356 ~~of the Mediterranean, does not provide evidence for decreased net evaporation in the marine system.~~  
357 This observation is critical, as it decouples the processes of precipitation on the Mediterranean  
358 margins with sapropel formation, and consequent changes in ~~moentum~~momentum transfer to the  
359 North Atlantic (Rogerson et al., 2012). ~~Consequently, we recommend that great care is taken to~~  
360 ~~determine whether past precipitation peaks reflect significantly enhanced external water advection~~  
361 ~~before any continental record can be used as a basis for inferring Mediterranean freshening.~~

### 362 ~~Palaeoclimatological significance~~

363 ~~The consistency of MIS 3 and modern precipitation isotope values permits comparison of fluid~~  
364 ~~inclusion values and precipitation magnitude records at Sfax and Bet Dagan. Most of the water~~  
365 ~~reaching Susah Cave seems to have been derived from large magnitude rainfall sourced from the~~  
366 ~~Western or Eastern Mediterranean surface water. The primary difference between these end members~~  
367 ~~is the level of  $D_{\text{excess}}$ , with the Western water  $\sim 10\%$  and Eastern water  $\sim 30\%$ . This difference allows~~  
368 ~~the influence of these two sources to be compared between the three major humid phases (Hoffmann~~  
369 ~~et al., 2016) recorded in SC-06-01 (Fig. 10). These phases reflect changes in the distribution of~~  
370 ~~insolation as a consequence of changes in orbital tilt, with Phase I (65 to 61 ka BP) and Phase III~~  
371 ~~(37.5 to 33.5 ka BP) associated with reflecting Northern Hemisphere heating during precession~~  
372 ~~minima and Phase II (52.5 to 50.5 ka BP) which has been associated with a change in obliquity. In all~~  
373 ~~cases, the peak in rainfall recorded by the speleothem leads the orbital peak by  $\sim 3$  ka. Phases I and III~~  
374 ~~both show very elevated  $D_{\text{excess}}$ , whereas no such values were found in Phase II. This provides further~~  
375 ~~support to our conclusion that the Eastern Mediterranean source contributed significant moisture to~~  
376 ~~Cyrenaica during precession related humid events, but that it did not during the obliquity related~~  
377 ~~humid event. This difference in the origin of the moisture feeding rainfall may explain the difference~~  
378 ~~in average  $\delta^{18}\text{O}_{\text{ee}}$  during these different phases (Hoffmann et al., 2016).~~

379 ~~The varying balance between Eastern and Western precipitation is diagnostic of changing basin-scale~~  
380 ~~atmospheric structure during the past~~Despite the low level of Atlantic moisture contributing to rainfall  
381 in Libya in MIS 3. ~~Eastern sourced rainfall may occasionally relate to wintertime storms, as today~~

382 ~~(Gat et al., 2003), but essentially reflects convective rainfall with relatively small advection distances.~~  
383 ~~The significant enhancement of the magnitude and regional significance of this convective rainfall~~  
384 ~~observed at Susah Cave must reflect greater atmospheric convergence due to northward displacement~~  
385 ~~of the annual average position of the ITCZ (Tuenter et al., 2003). Contrary to this, the Western-~~  
386 ~~sourced moisture is transported ~1500 km eastwards to reach Cyrenaica, which must reflect the mid-~~  
387 ~~latitude storm track (Brayshaw et al., 2009). Consequently, although it does not seem that Atlantic~~  
388 ~~moisture is important to the climatology of Cyrenaica, the momentum derived from Atlantic winter~~  
389 ~~storms predicted by regional climate modelling (Brayshaw et al., 2009) and observed on the northern~~  
390 ~~Mediterranean margin (Toucanne et al., 2015) remains pivotal to supplying moisture to North Africa.~~  
391 Consequently, the North Atlantic heat budget provides an important control on northern African  
392 rainfall in the past. In contrast, this control cannot explain changes in the Eastern-sourced rainfall  
393 revealed by our analysis. Eastern-sourced rainfall may occasionally relate to wintertime storms, as  
394 today (Gat et al., 2003), but essentially reflects convective rainfall with relatively small advection  
395 distances. Within obliquity forced phases, advective transport of moisture alone drives humidity. In  
396 contrast, we conclude that during precession forced humid phases, the impact of advective transport  
397 of moisture from the Western to the Eastern Mediterranean basin occurs alongside strong convergence  
398 and convective rainfall within the eastern basin. The dilution of the advective signal by internal  
399 convective rainfall may be the reason why Dansgaard-Oeschger cycles in the North Atlantic are well  
400 reflected at Susah Cave during high precession (Hoffmann et al., 2016), whereas there is weaker  
401 correspondence of Cyrenaican rain and North Atlantic heat during low precession. It is likely this  
402 arises due to greater atmospheric convergence due to northward displacement of the annual average  
403 position of the ITCZ (Tuenter et al., 2003)  
404 ~~Further constraint on large scale atmospheric advection can be provided by Sr isotopes, which are~~  
405 ~~known to be sensitive to changes in transport of Saharan dust (Frumkin and Stein, 2004). Even~~  
406 ~~considering the most slow growing and most rapidly growing parts of SC-06-01, no significant~~  
407 ~~difference in  $^{87}\text{Sr}/^{86}\text{Sr}$  was identified. This is unexpected and significant, as climate driven changes in~~  
408  ~~$^{87}\text{Sr}/^{86}\text{Sr}$  have previously been reported from speleothems in the Mediterranean region (Frumkin and~~



409 Stein, 2004). Palaeoclimatologically, our analysis reveals that 1) during northern hemisphere  
410 insolation peaks reflecting Precession, coastal Libya experiences greater westerly advection of water  
411 due to an increase in Atlantic heat and greater convective rainfall due to migration of the ITCZ  
412 whereas 2) insolation peaks reflecting obliquity show increased Atlantic heat and westerlies, but no  
413 comparable change in the ITCZ position.

414 ~~It seems that despite changes in the intensity of moisture transport during the period 65–30 ka BP,~~  
415 ~~there is no large-scale change in atmospheric dust transport direction. This further supports our~~  
416 ~~conclusion from the fluid inclusions that the Eastern Mediterranean rainfall operating during~~  
417 ~~precession minima reflects enhanced internal convection rather than transport of moisture from the~~  
418 ~~east or south with an atmospheric circulation pattern that prevails today.~~

#### 419 *Implications for Susah Cave $\delta^{18}\text{O}_{cc}$*

420 Aside from those data with high deuterium excess, which reflect influence from the Eastern  
421 Mediterranean source, much of the variance in the fluid inclusion dataset is captured by a two end-  
422 member mixing system resembling modern rainfall in Tunisia. One end-member is the Western  
423 Mediterranean source of Celle-Jeanton et al. (~~2003.~~ (2001)), but the other is isotopically too heavy to be  
424 identified with the Atlantic source. Rather, it resembles the “Sfax Mixed” population defined by  
425 Celle-Jeanton et al. (~~2003.~~ (2001)), reflecting a mixed source of moisture from both the Western  
426 Mediterranean and Atlantic. Consequently, although quantitatively minor amounts of Atlantic water  
427 reached the site, changes in the moisture advection driven by westerly winds had a strong influence on  
428  $\delta^{18}\text{O}_{\text{dripwater}}$  trends in time. At Sfax today, this influence causes a prominent bimodal behaviour with  
429 two rainfall maxima with different  $\delta^{18}\text{O}_{\text{ppt}}$ , which eliminates a simple and quantitative rainfall amount  
430 control on precipitation, which can be observed at Tunis (WMO code 6071500,  
431 <https://nucleus.iaea.org/wiser/gnip.php>). Furthermore, addition of heavy rain events derived from the  
432 Eastern Mediterranean aliases the tendency towards depleted  $\delta^{18}\text{O}_{\text{dripwater}}$ , as this water is also more  
433 depleted than modern Western Mediterranean precipitation. In the Bet Dagan data, there is also a  
434 tendency to lower  $\delta^{18}\text{O}_{\text{ppt}}$  with higher precipitation amount, but the relationship between rainfall  
435 amount and rainfall isotope composition is not identical to Tunis. Ultimately, it seems likely that

436 rainfall amount changes at Susah Cave do cause depleted (enriched)  $\delta^{18}\text{O}_{\text{cc}}$  values to be associated  
437 with high (low) rainfall, but this is too complicated by independent changes increases (decreases) in  
438 westerly moisture advection and increases (decreases) in convergence. Qualitatively, all these  
439 parameters are expected symptoms of North African humid phases and so these trends remain a  
440 valuable expression of climatic variability. Quantitatively, more information is required to translate  
441 the trends into fully-functional palaeoclimatologies, and this analysis pivots on whether  $\delta^{18}\text{O}_{\text{cc}}$  trends  
442 reflect changes in water deficit / surplus in Cyrenaica.

443 Although it is likely the oxygen isotope fractionation during calcite precipitation occurred close to  
444 isotope equilibrium (Hoffmann et al., 2016), there is a good degree of correspondence between  
445 positive and negative phases in  $\delta^{18}\text{O}_{\text{cc}}$  and  $\delta^{13}\text{C}_{\text{cc}}$ , indicating a shared control. Indeed,  $\delta^{13}\text{C}_{\text{cc}}$  has a  
446 markedly higher amplitude variability than  $\delta^{18}\text{O}_{\text{cc}}$ . More isotopically depleted carbon may represent  
447 increased incorporation of respired soil carbon, increased dominance of C3 over C4 plants, and/or  
448 decreased degassing of aquifer water (Baker et al., 1997). Today, the Susah Cave location on Jebel  
449 Malh has very thin soil cover, colonised by shrubby maquis vegetation. Soil respiration and  
450 colonisation by C3 plants is limited by the strong water deficit of the region, and aquifer water  
451 outgassing is enhanced by long residence times due to low water infiltration. Increased water  
452 availability will progressively deplete the  $\delta^{13}\text{C}$  of dripwater by all three mechanisms described above.  
453 Consequently, all three of these processes promote correlation between  $\delta^{13}\text{C}_{\text{cc}}$  and precipitation  
454 amount. Within the  $\delta^{18}\text{O}_{\text{cc}}$  data series, peak growth rates occur both during relatively enriched and  
455 relatively depleted isotope stages. This is not the case for  $\delta^{13}\text{C}_{\text{cc}}$ , which more consistently shows  
456 depleted values during times of rapid growth (SC-06-01 growth phases shown in Fig. 11). We  
457 therefore consider it likely that  $\delta^{13}\text{C}_{\text{cc}}$  indeed more accurately records rainfall amount than  $\delta^{18}\text{O}_{\text{cc}}$  does.

## 458 Conclusions and Implications

459 A key feature of this combined dataset is the long-term sinusoidal trend in both the  $\delta^{18}\text{O}_{\text{cc}}$  and  $\delta^2\text{H}_{\text{fi}}$ ,  
460 reflecting the differing rainfall regimes dominant between Humid Phases I and III compared to Phase  
461 II. This is not developed in  $\delta^{13}\text{C}_{\text{cc}}$ , implying that the process forcing the long-term cycle in moisture

462 source is not impacting on carbon dynamics in the soil and epikarst. We therefore conclude that there  
463 is a mixed amount and source control on  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  in the SC-01-06 record, whereas  $\delta^{13}\text{C}$  is  
464 dominantly controlled by water availability.

465 The fluid inclusions from SC-06-01 show that rainfall compositions in the southeast Mediterranean  
466 region during MIS 3 were comparable to modern rainfall compositions recorded in regional GNIP  
467 datasets. However, the diversity of compositions is impossible to explain with a single rainfall source,  
468 rather indicating that moisture derived from the Atlantic, the Western Mediterranean and the Eastern  
469 Mediterranean basins have all contributed to MIS 3 precipitation in Libya. This requires both  
470 enhanced westerly advection of moisture to this region, reflecting the Atlantic storm track, and  
471 enhanced convective rainfall within the Eastern Mediterranean basin. There is some indication that  
472 these two mechanisms differ in terms of their response to orbital forcing, with precession parameter  
473 minima enhancing westerly advection and internal convection, whereas obliquity minima enhance  
474 westerly advection without significantly altering internal convection.

475 Crucially, this picture is most consistent with atmospheric circulation over the Mediterranean  
476 remaining essentially unchanged during precession cycles. This is consistent with regional climate  
477 model experiments showing major enhancement of winter westerly storm activity, but it not  
478 consistent with the extreme migration of the ITCZ, where the monsoon belt approaches the North  
479 African coast. The strong implication is that a significant arid belt is retained between the  
480 Mediterranean and the ITCZ, even when northernmost Africa is experiencing significantly enhanced  
481 rainfall.

482 It is likely that rainfall amount played a role in controlling the isotopic composition of the calcite in  
483 this speleothem ( $\delta^{18}\text{O}_{\text{cc}}$ ). However, the more depleted values reflecting higher rainfall are also  
484 consistent with different mixing between the end members identified by the fluid inclusion analysis.  
485 The structure of the  $\delta^{13}\text{C}_{\text{cc}}$  record provides an independent means of assessing changes in water  
486 surplus / deficit, as more depleted values will reflect lower aquifer residence times, enhanced soil  
487 respiration and changes in vegetation structure, all of which are limited by water availability in this

488 semi-arid environment. Combined analysis of the proxies provides a powerful new demonstration that  
489 the northeast Libyan climate was more humid during millennial-scale warm periods in the North  
490 Atlantic realm, but quantification will be dependent on generating unambiguous independent evidence  
491 for water availability in the soil and epikarst.

## 492 Acknowledgements

493 We thank the Royal Geographical Society for the pump-priming investment that began this work  
494 (Thesiger-Oman International Fellowship 2009), the Natural Environment Research Council for  
495 providing the funds that made the analytical work on this project possible (NE/J014133/1) and The  
496 Leverhulme Trust for funding activities within the associated International Network (IN-2012-113).

## 497 References

- 498 Arienzo, M.M. et al. 2015. Bahamian speleothem reveals temperature decrease associated with  
499 Heinrich stadials. *EPSL* 430, 377-386.
- 500 [Adkins, J., Demenocal, P., and Eshel, G.: The "African humid period" and the record of marine  
501 upwelling from excess 230Th in Ocean Drilling Program Hole 658C, \*Paleoceanography and  
502 Paleoclimatology\*, 21, 2006.](#)
- 503 [Ait-Brahim, Y., Cheng, H., Sifeddine, A., Wassenburg, J. A., Cruz, F. W., Khodri, M., Sha, L., Pérez-  
504 Zanón, N., Beraaouz, E. H., Apaéstegui, J., Guyot, J.-L., Jochum, K. P., and Bouchaou, L.: Speleothem  
505 records decadal to multidecadal hydroclimate variations in southwestern Morocco during the last  
506 millennium, \*Earth and Planetary Science Letters\*, 476, 1-10,  
507 <https://doi.org/10.1016/j.epsl.2017.07.045>, 2017.](#)
- 508 [Arienzo, M. M., Swart, P. K., Pourmand, A., Broad, K., Clement, A. C., Murphy, L. N., Vonhof, H. B.,  
509 and Kakuk, B.: Bahamian speleothem reveals temperature decrease associated with Heinrich  
510 stadials, \*Earth and Planetary Science Letters\*, 430, 377-386, 2015.](#)
- 511 [Armitage, S. J., Drake, N. A., Stokes, S., El-Hawat, A., Salem, M., White, K., Turner, P., and McLaren, S.  
512 J.: Multiple phases of north African humidity recorded in lacustrine sediments from the fazzan basin,  
513 \*Libyan sahara\*, \*Quaternary Geochronology\*, 2, 181-186, 2007.](#)
- 514 [Ayalon, A., Bar-Matthews, M., and Sass, E.: Rainfall-recharge relationships within a karstic terrain in  
515 the Eastern-Mediterranean semi-arid region, Israel:  \$\delta^{18}\text{O}\$  and  \$\delta\text{D}\$  characteristics, \*Journal of  
516 Hydrology\*, 207, 18-31, \[10.1016/S0022-1694\\(98\\)00119-X\]\(https://doi.org/10.1016/S0022-1694\(98\)00119-X\), 1998.](#)
- 517 [Baker, A., Ito, E., Smart, P. L., and McEwan, R. F.: Elevated and variable values of  \$^{13}\text{C}\$  in speleothems  
518 in a British cave system, \*Chemical Geology\*, 136, 263-270, 1997.](#)
- 519 [Ballais, J.-L.: Evolution holocène de la Tunisie saharienne et présaharienne, \*Méditerranée\*, 74, 31-38,  
520 1991.](#)
- 521 [Bethoux, J. P., and Gentili, B.: Functioning of the Mediterranean Sea: past and present changes  
522 related to freshwater input and climate changes, \*Journal of Marine Systems\*, 20, 33-47, 1999.](#)
- 523 [Black, E., Brayshaw, D. J., and Rambeau, C. M. C.: Past, present and future precipitation in the Middle  
524 East: Insights from models and observations, \*Philosophical Transactions of the Royal Society A:  
525 Mathematical, Physical and Engineering Sciences\*, 368, 5173-5184, \[10.1098/rsta.2010.0199\]\(https://doi.org/10.1098/rsta.2010.0199\), 2010.](#)

526 Bosmans, J. H. C., Drijfhout, S. S., Tuenter, E., Hilgen, F. J., Lourens, L. J., and Rohling, E. J.: Precession  
527 and obliquity forcing of the freshwater budget over the Mediterranean, *Quaternary Science*  
528 *Reviews*, 123, 16–30, 10.1016/j.quascirev.2015.06.008, 2015.

529 Brayshaw, D. J., Woollings, T., and Vellinga, M.: Tropical and Extratropical Responses of the North  
530 Atlantic Atmospheric Circulation to a Sustained Weakening of the MOC, *Journal of Climate*, 22, 3146–  
531 3155, 10.1175/2008jcli2594.1, 2009.

532 Celle Jeanton, H., Zouari, K., Travi, Y., and Daoud, A.: Caractérisation isotopique des pluies en  
533 Tunisie. Essai de typologie dans la région de Sfax, *Sciences de la Terre et des planètes*, 333, 625–631,  
534 2001.

535 Charlier, B., Ginibre, C., Morgan, D., Nowell, G., Pearson, D., Davidson, J., and Ottley, C.: Methods for  
536 the microsampling and high-precision analysis of strontium and rubidium isotopes at single crystal  
537 scale for petrological and geochronological applications, *Chemical Geology*, 232, 114–133, 2006.

538 Collins, J. A., Prange, M., Caley, T., Gimeno, L., Beckmann, B., Mulitza, S., Skonieczny, C., Roche, D.,  
539 and Schefuß, E.: Rapid termination of the African humid period triggered by northern high-latitude  
540 cooling, *Nature Communications*, 8, 1372, 2017.

541 deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., and Yarusinsky, M.: Abrupt  
542 onset and termination of the African Humid Period: rapid climate responses to gradual insolation  
543 forcing., *Quaternary Science Reviews*, 19, 347–361, 2000.

544 Drake, N. A., El Hawat, A. S., Turner, P., Armitage, S. J., Salem, M. J., White, K. H., and McLaren, S.:  
545 Palaeohydrology of the Fazzan Basin and surrounding regions: The last 7 million years,  
546 *Palaeogeography Palaeoclimatology Palaeoecology*, 263, 131–145, 10.1016/j.palaeo.2008.02.005,  
547 2008.

548 Drake, N. A., Blench, R. M., Armitage, S. J., Bristow, C. S., and White, K. H.: Ancient watercourses and  
549 biogeography of the Sahara explain the peopling of the desert, *Proc. Natl. Acad. Sci. U. S. A.*, 108,  
550 458–462, 10.1073/pnas.1012231108, 2011.

551 Dublyansky, Y. V., and Spötl, C.: Hydrogen and oxygen isotopes of water from inclusions in minerals:  
552 Design of a new crushing system and on-line continuous-flow isotope ratio mass spectrometric  
553 analysis, *Rapid Communications in Mass Spectrometry*, 23, 2605–2613, 10.1002/rcm.4155, 2009.

554 Fleitmann, D., Burns, S. J., Neff, U., Mangini, A., and Matter, A.: Changing moisture sources over the  
555 last 330,000 years in Northern Oman from fluid-inclusion evidence in speleothems, *Quaternary*  
556 *Research*, 60, 223–232, [http://dx.doi.org/10.1016/S0033-5894\(03\)00086-3](http://dx.doi.org/10.1016/S0033-5894(03)00086-3), 2003.

557 Fontes, J. C., and Gasse, F.: PALHYDAF (Palaeohydrology in Africa) program: objectives, methods,  
558 major results, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 84, 191–215, 10.1016/0031-  
559 0182(91)90044-R, 1991.

560 Frumkin, A., and Stein, M.: The Sahara–East Mediterranean dust and climate connection revealed by  
561 strontium and uranium isotopes in a Jerusalem speleothem, *Earth and Planetary Science Letters*,  
562 217, 451–464, 10.1016/S0012-821X(03)00589-2, 2004.

563 Gasse, F., and Campo, E. v.: Abrupt post-glacial climate events in west Asia and north Africa  
564 monsoon domains, *Earth and Planetary Science Letters*, 126, 435–456, 1994.

565 Gasse, F.: Diatom-inferred salinity and carbonate oxygen isotopes in Holocene waterbodies of the  
566 western Sahara and Sahel (Africa), *Quaternary Science Reviews*, 21, 737–767, 2002.

567 Gat, J. R., Klein, B., Kushnir, Y., Roether, W., Wernli, H., Yam, R., and Shemesh, A.: Isotope  
568 composition of air moisture over the Mediterranean Sea: An index of the air–sea interaction pattern,  
569 *Tellus, Series B: Chemical and Physical Meteorology*, 55, 953–965, 10.1034/j.1600-  
570 0889.2003.00081.x, 2003.

571 Genty, D., Blamart, D., Ghaleb, B., Plagnes, V., Causse, C., Bakalowicz, M., Zouari, K., Chkir, N.,  
572 Hellstrom, J., Wainer, K., and Bourges, F.: Timing and dynamics of the last deglaciation from  
573 European and North African  $\delta^{13}\text{C}$  stalagmite profiles – comparison with Chinese and South  
574 Hemisphere stalagmites, *Quaternary Science Reviews*, 25, 2118–2142, 2006.

575 Goldsmith, Y., Polissar, P. J., Ayalon, A., Bar-Matthews, M., deMenocal, P. B., and Broecker, W. S.:  
576 The modern and Last Glacial Maximum hydrological cycles of the Eastern Mediterranean and the

577 Levant from a water isotope perspective, *Earth and Planetary Science Letters*, 457, 302-312,  
578 <http://dx.doi.org/10.1016/j.epsl.2016.10.017>, 2017.

579 Grant, K. M., Grimm, R., Mikolajewicz, U., Marino, G., Ziegler, M., and Rohling, E. J.: The timing of  
580 Mediterranean sapropel deposition relative to insolation, sea level and African monsoon changes,  
581 *Quaternary Science Reviews*, 140, 125-141, <http://dx.doi.org/10.1016/j.quascirev.2016.03.026>,  
582 2016.

583 Harrison, S., Bartlein, P., Izumi, K., Li, G., Annan, J., Hargreaves, J., Braconnot, P., and Kageyama, M.:  
584 Evaluation of CMIP5 palaeo-simulations to improve climate projections, *Nature Climate Change*, 5,  
585 735-743, 2015.

586 Hoffmann, D. L., Rogerson, M., Spötl, C., Luetscher, M., Vance, D., Osborne, A. H., Fello, N. M., and  
587 Moseley, G. E.: Timing and causes of North African wet phases during MIS 3 and implications for  
588 Modern Human migration, *Nature Scientific Reports*, 6, 36367, 2016.

589 IPCC: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects.*  
590 *Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on*  
591 *Climate Change*, United Kingdom and New York, 688, 2014.

592 Jolly, D., Prentice, I. C., Bonnefille, R., Ballouche, A., Bengo, M., Brenac, P., Buchet, G., Burney, D.,  
593 Cazet, J. P., Cheddadi, R., Ederh, T., Elenga, H., Elmoutaki, S., Guiot, J., Laarif, F., Lamb, H., Lezine, A.  
594 M., Maley, J., Mbenza, M., Peyron, O., Reille, M., Reynaud-Farrera, I., Riollet, G., Ritchie, J. C., Roche,  
595 E., Scott, L., Ssemmanda, I., Straka, H., Umer, M., Van Campo, E., Vilimumbalo, S., Vincens, A., and  
596 Waller, M.: Biome reconstruction from pollen and plant macrofossil data for Africa and the Arabian  
597 peninsula at 0 and 6000 years, *Journal of Biogeography*, 25, 1007-1027, 1998.

598 McGarry, S., Bar-Matthews, M., Matthews, A., Vaks, A., Schilman, B., and Ayalon, A.: Constraints on  
599 hydrological and paleotemperature variations in the Eastern Mediterranean region in the last 140 ka  
600 given by the  $\delta D$  values of speleothem fluid inclusions, *Quaternary Science Reviews*, 23, 919-934,  
601 <http://dx.doi.org/10.1016/j.quascirev.2003.06.020>, 2004.

602 Meckler, A. N., Affolter, S., Dublyansky, Y. V., Krüger, Y., Vogel, N., Bernasconi, S. M., Frenz, M.,  
603 Kipfer, R., Leuenberger, M., Spötl, C., Carolin, S., Cobb, K. M., Moerman, J., Adkins, J. F., and  
604 Fleitmann, D.: Glacial-interglacial temperature change in the tropical West Pacific: A comparison of  
605 stalagmite-based paleo-thermometers, *Quaternary Science Reviews*, 127, 90-116,  
606 <http://dx.doi.org/10.1016/j.quascirev.2015.06.015>, 2015.

607 Osborne, A., Vance, D., Rohling, E., Barton, N., Rogerson, M., and Fello, N.: A humid corridor across  
608 the Sahara for the migration of early modern humans out of Africa 120,000 years ago, *Proc. Natl.*  
609 *Acad. Sci. U. S. A.*, doi\_10.1073\_pnas.0804472105, 2008.

610 Osborne, A. H., Marino, G., Vance, D., and Rohling, E. J.: Eastern Mediterranean surface water Nd  
611 during Eemian sapropel S5: monitoring northerly (mid-latitude) versus southerly (sub-tropical)  
612 freshwater contributions, *Quaternary Science Reviews*, 29, 2473-2483,  
613 <http://dx.doi.org/10.1016/j.quascirev.2010.05.015>, 2010.

614 Pascale, S., Gregory, J. M., Ambaum, M., and Tailleux, R.: Climate entropy budget of the HadCM3  
615 atmosphere-ocean general circulation model and of FAMOUS, its low-resolution version, *Climate*  
616 *Dynamics*, 36, 1189-1206, 10.1007/s00382-009-0718-1, 2011.

617 PETIT-MAIRE, N., BUROLLET, P. F., A-BALLAIS, J. L., A-FONTUGNE, M., A-ROSSO, J. C., A-LAZAAR, A.,  
618 and Gauthier-Villars, I.: Paléoclimats holocènes du Sahara septentrional. Dépôts lacustres et  
619 terrasses alluviales en bordure du Grand Erg Oriental à l'extrême Sud de la Tunisie, *Comptes rendus*  
620 *de l'Académie des sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de*  
621 *la Terre* 312, 1661-1666, 1991.

622 Peyron, O., Jolly, D., Braconnot, P., Bonnefille, R., Guiot, J., Wirmann, D., and Chalief, F.: Quantitative  
623 reconstructions of annual rainfall in Africa 6000 years ago: Model-data comparison, *Journal of*  
624 *Geophysical Research Atmospheres*, 111, D24110  
625 Artn d24110, 2006.

626 Prentice, I. C., and Jolly, D.: Mid-Holocene and glacial-maximum vegetation geography of the  
627 northern continents and Africa, *Journal of Biogeography*, 27, 507-519, 2000.

628 Rogerson, M., Schönfeld, J., and Leng, M.: Qualitative and quantitative approaches in  
629 palaeohydrography: A case study from core top parameters in the Gulf of Cadiz, *Marine Geology*,  
630 **280**, 150–167, 2011.

631 Rogerson, M., Rohling, E. J., Bigg, G. R., and Ramirez, J.: Palaeoceanography of the Atlantic-  
632 Mediterranean Exchange: Overview and first quantitative assessment of climatic forcing., *Reviews*  
633 *of Geophysics*, **50**, DOI: 8755-1209/8712/2011RG000376, 2012.

634 Rohling, E., Marino, G., and Grant, K.: Mediterranean climate and oceanography, and the periodic  
635 development of anoxic events (sapropels), *Earth Science Reviews*, **143**, 62–97, 2015.

636 Rohling, E. J., and Bryden, H. L.: Estimating past changes in the Eastern Mediterranean freshwater  
637 budget, using reconstructions of sea level and hydrography, *Proceedings Koninklijke Nederlandse*  
638 *Akademie van Wetenschappen, Serie B*, **97**, 201–217, 1994.

639 Rowan, J. S., Black, S., Macklin, M. G., Tabner, B. J., and Dore, J.: Quaternary environmental change  
640 in Cyrenaica evidenced by U-Th, ESR and OSL of coastal alluvial fan sequences., *Libyan Studies*, **31**,  
641 5–16, 2000.

642 Schwarcz, H. P., Harmon, R. S., Thompson, P., and Ford, D. C.: Stable isotope studies of fluid  
643 inclusions in speleothems and their paleoclimatic significance, *Geochimica et Cosmochimica Acta*,  
644 **40**, 657–665, [http://dx.doi.org/10.1016/0016-7037\(76\)90111-3](http://dx.doi.org/10.1016/0016-7037(76)90111-3), 1976.

645 Smith, J. R., Giegengack, R., Schwarcz, H. P., McDonald, M. M. A., Kleindienst, M. R., Hawkins, A. L.,  
646 and Churcher, C. S.: A reconstruction of quaternary pluvial environments and human occupations  
647 using stratigraphy and geochronology of fossil spring tufas, Kharga Oasis, Egypt, *Geoarchaeology an*  
648 *International Journal*, **19**, 407–439, 2004.

649 Spötl, C.: Long term performance of the Gasbench isotope ratio mass spectrometry system for the  
650 stable isotope analysis of carbonate microsamples, *Rapid Communications in Mass Spectrometry*,  
651 **25**, 1683–1685, [10.1002/rcm.5037](https://doi.org/10.1002/rcm.5037), 2011.

652 Swezey, C.: Eolian sediment responses to late Quaternary climate changes: temporal and spatial  
653 patterns in the Sahara, *Palaeogeography, Palaeoclimatology, Palaeoecology*, **167**, 119–155, 2001.

654 Toucanne, S., Angue Minto'o, C. M., Fontanier, C., Bassetti, M. A., Jorry, S. J., and Jouet, G.: Tracking  
655 rainfall in the northern Mediterranean borderlands during sapropel deposition, *Quaternary Science*  
656 *Reviews*, **129**, 178–195, <http://dx.doi.org/10.1016/j.quascirev.2015.10.016>, 2015.

657 Tuenter, E., Weber, S. L., Hilgen, F. J., and Lourens, L. J.: The response of the African summer  
658 monsoon to remote and local forcing due to precession and obliquity, *Global and Planetary Change*,  
659 **36**, 219–235, 2003.

660 Vaks, A., Woodhead, J., Bar-Matthews, M., Ayalon, A., Cliff, R., Zilberman, T., Matthews, A., and  
661 Frumkin, A.: Pliocene–Pleistocene climate of the northern margin of Saharan–Arabian Desert  
662 recorded in speleothems from the Negev Desert, Israel, *Earth and Planetary Science Letters*, **368**, 88–  
663 100, 2013.

664 Van Breukelen, M., Vonhof, H., Hellstrom, J., Wester, W., and Kroon, D.: Fossil dripwater in  
665 stalagmites reveals Holocene temperature and rainfall variation in Amazonia, *Earth and Planetary*  
666 *Science Letters*, **275**, 54–60, 2008.

667 Wainer, K., Genty, D., Blamart, D., Daëron, M., Bar-Matthews, M., Vonhof, H., Dublyansky, Y., Pons-  
668 Branchu, E., Thomas, L., van Calsteren, P., Quinif, Y., and Caillon, N.: Speleothem record of the last  
669 180 ka in Villars cave (SW France): Investigation of a large  $\delta^{18}\text{O}$  shift between MIS6 and MIS5,  
670 *Quaternary Science Reviews*, **30**, 130–146, <http://dx.doi.org/10.1016/j.quascirev.2010.07.004>, 2011.

671 Wassenburg, J. A., Immenhauser, A., Richter, D. K., Niedermayr, A., Riechelmann, S., Fietzke, J.,  
672 Scholz, D., Jochum, K. P., Fohlmeister, J., Schröder-Ritzrau, A., Sabaoui, A., Riechelmann, D. F. C.,  
673 Schneider, L., and Esper, J.: Moroccan speleothem and tree ring records suggest a variable positive  
674 state of the North Atlantic Oscillation during the Medieval Warm Period, *Earth and Planetary Science*  
675 *Letters*, **375**, 291–302, <http://dx.doi.org/10.1016/j.epsl.2013.05.048>, 2013.

676 Wassenburg, J. A., Dietrich, S., Fietzke, J., Fohlmeister, J., Jochum, K. P., Scholz, D., Richter, D. K.,  
677 Sabaoui, A., Spötl, C., Lohmann, G., Andreae, Meinrat O., and Immenhauser, A.: Reorganization of

678 [the North Atlantic Oscillation during early Holocene deglaciation, Nat. Geosci., 9, 602,](#)  
679 [10.1038/ngeo2767](#)

680 <https://www.nature.com/articles/ngeo2767#supplementary-information>, 2016.

681 [Adkins, J., Demenocal, P., and Eshel, G.: The “African humid period” and the record of marine](#)  
682 [upwelling from excess 230Th in Ocean Drilling Program Hole 658C, Paleoceanography and](#)  
683 [Paleoclimatology, 21, 2006.](#)

684 [Aggarwal, P. K., Romatschke, U., Araguas-Araguas, L., Belachew, D., Longstaffe, F. J., Berg, P.,](#)  
685 [Schumacher, C., and Funk, A.: Proportions of convective and stratiform precipitation revealed in](#)  
686 [water isotope ratios, Nat. Geosci., 9, 624, 2016.](#)

687 [Ait Brahim, Y., Cheng, H., Sifeddine, A., Wassenburg, J. A., Cruz, F. W., Khodri, M., Sha, L., Pérez-](#)  
688 [Zanón, N., Beraaouz, E. H., Apaéstegui, J., Guyot, J.-L., Jochum, K. P., and Bouchaou, L.: Speleothem](#)  
689 [records decadal to multidecadal hydroclimate variations in southwestern Morocco during the last](#)  
690 [millennium, Earth and Planetary Science Letters, 476, 1-10,](#)  
691 <https://doi.org/10.1016/j.epsl.2017.07.045>, 2017.

692 [Arienzo, M. M., Swart, P. K., Pourmand, A., Broad, K., Clement, A. C., Murphy, L. N., Vonhof, H. B.,](#)  
693 [and Kakuk, B.: Bahamian speleothem reveals temperature decrease associated with Heinrich](#)  
694 [stadials, Earth and Planetary Science Letters, 430, 377-386, 2015.](#)

695 [Armitage, S. J., Drake, N. A., Stokes, S., El-Hawat, A., Salem, M., White, K., Turner, P., and McLaren, S.](#)  
696 [J.: Multiple phases of north African humidity recorded in lacustrine sediments from the fazzan basin,](#)  
697 [Libyan sahara, Quaternary Geochronology, 2, 181-186, 2007.](#)

698 [Ayalon, A., Bar-Matthews, M., and Sass, E.: Rainfall-recharge relationships within a karstic terrain in](#)  
699 [the Eastern Mediterranean semi-arid region, Israel:  \$\delta^{18}O\$  and  \$\delta D\$  characteristics, Journal of](#)  
700 [Hydrology, 207, 18-31, 10.1016/S0022-1694\(98\)00119-X, 1998.](#)

701 [Baker, A., Ito, E., Smart, P. L., and McEwan, R. F.: Elevated and variable values of  \$^{13}C\$  in speleothems](#)  
702 [in a British cave system, Chemical Geology, 136, 263-270, 1997.](#)

703 [Ballais, J.-L.: Evolution holocène de la Tunisie saharienne et présaharienne, Méditerranée, 74, 31-38,](#)  
704 [1991.](#)

705 [Bethoux, J. P., and Gentili, B.: Functioning of the Mediterranean Sea: past and present changes](#)  
706 [related to freshwater input and climate changes, Journal of Marine Systems, 20, 33-47, 1999.](#)

707 [Black, E., Brayshaw, D. J., and Rambeau, C. M. C.: Past, present and future precipitation in the Middle](#)  
708 [East: Insights from models and observations, Philosophical Transactions of the Royal Society A:](#)  
709 [Mathematical, Physical and Engineering Sciences, 368, 5173-5184, 10.1098/rsta.2010.0199, 2010.](#)

710 [Bosmans, J. H. C., Drijfhout, S. S., Tuenter, E., Hilgen, F. J., Lourens, L. J., and Rohling, E. J.: Precession](#)  
711 [and obliquity forcing of the freshwater budget over the Mediterranean, Quaternary Science](#)  
712 [Reviews, 123, 16-30, 10.1016/j.quascirev.2015.06.008, 2015.](#)

713 [Brayshaw, D. J., Woollings, T., and Vellinga, M.: Tropical and Extratropical Responses of the North](#)  
714 [Atlantic Atmospheric Circulation to a Sustained Weakening of the MOC, Journal of Climate, 22, 3146-](#)  
715 [3155, 10.1175/2008jcli2594.1, 2009.](#)

716 [Brun, A.: Reflections on the pluvial and arid periods of the Upper Pleistocene and of the Holocene in](#)  
717 [Tunisia, Palaeoecology of Africa and the surrounding islands. Vol. 22. Proc. symposium on African](#)  
718 [palynology, Rabat, 1989, 157-170, 1991.](#)

719 [Cancellieri E., Cremaschi M., Zerboni A., and S., d. L.: Climate, Environment, and Population](#)  
720 [Dynamics in Pleistocene Sahara, in: Africa from MIS 6-2. Vertebrate Paleobiology and](#)  
721 [Paleoanthropology., edited by: Jones S., and B., S., Springer, Dordrecht, 2016.](#)

722 [Celle-Jeanton, H., Zouari, K., Travi, Y., and Daoud, A.: Caractérisation isotopique des pluies en](#)  
723 [Tunisie. Essai de typologie dans la région de Sfax, Sciences de la Terre et des planètes, 333, 625-631,](#)  
724 [2001.](#)

725 [Charlier, B., Ginibre, C., Morgan, D., Nowell, G., Pearson, D., Davidson, J., and Ottley, C.: Methods for](#)  
726 [the microsampling and high-precision analysis of strontium and rubidium isotopes at single crystal](#)  
727 [scale for petrological and geochronological applications, Chemical Geology, 232, 114-133, 2006.](#)



728 [Collins, J. A., Govin, A., Mulitza, S., Heslop, D., Zabel, M., Hartmann, J., Röhl, U., and Wefer, G.:](#)  
729 [Abrupt shifts of the Sahara–Sahel boundary during Heinrich stadials, \*Climate of the Past\*, 9, 1181-](#)  
730 [1191, 2013.](#)

731 [Collins, J. A., Prange, M., Caley, T., Gimeno, L., Beckmann, B., Mulitza, S., Skonieczny, C., Roche, D.,](#)  
732 [and Schefuß, E.: Rapid termination of the African humid period triggered by northern high-latitude](#)  
733 [cooling, \*Nature Communications\*, 8, 1372, 2017.](#)

734 [Dayan, U., Nissen, K., and Ulbrich, U.: Review Article: Atmospheric conditions inducing extreme](#)  
735 [precipitation over the eastern and western Mediterranean, \*Nat. Hazards Earth Syst. Sci.\*, 15, 2525-](#)  
736 [2544, 10.5194/nhess-15-2525-2015, 2015.](#)

737 [deMenocal, P., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., and Yarusinsky, M.: Abrupt](#)  
738 [onset and termination of the African Humid Period: rapid climate responses to gradual insolation](#)  
739 [forcing. , \*Quaternary Science Reviews\*, 19, 347- 361, 2000.](#)

740 [Drake, N. A., El-Hawat, A. S., Turner, P., Armitage, S. J., Salem, M. J., White, K. H., and McLaren, S.:](#)  
741 [Palaeohydrology of the Fazzan Basin and surrounding regions: The last 7 million years,](#)  
742 [Palaeogeography Palaeoclimatology Palaeoecology, 263, 131-145, 10.1016/j.palaeo.2008.02.005,](#)  
743 [2008.](#)

744 [Drake, N. A., Blench, R. M., Armitage, S. J., Bristow, C. S., and White, K. H.: Ancient watercourses and](#)  
745 [biogeography of the Sahara explain the peopling of the desert, \*Proc. Natl. Acad. Sci. U. S. A.\*, 108,](#)  
746 [458-462, 10.1073/pnas.1012231108, 2011.](#)

747 [Dublyansky, Y. V., and Spötl, C.: Hydrogen and oxygen isotopes of water from inclusions in minerals:](#)  
748 [Design of a new crushing system and on-line continuous-flow isotope ratio mass spectrometric](#)  
749 [analysis, \*Rapid Communications in Mass Spectrometry\*, 23, 2605-2613, 10.1002/rcm.4155, 2009.](#)

750 [Fleitmann, D., Burns, S. J., Neff, U., Mangini, A., and Matter, A.: Changing moisture sources over the](#)  
751 [last 330,000 years in Northern Oman from fluid-inclusion evidence in speleothems, \*Quaternary\*](#)  
752 [Research, 60, 223-232, \[http://dx.doi.org/10.1016/S0033-5894\\(03\\)00086-3\]\(http://dx.doi.org/10.1016/S0033-5894\(03\)00086-3\), 2003.](#)

753 [Fontes, J. C., and Gasse, F.: PALHYDAF \(Palaeohydrology in Africa\) program: objectives, methods,](#)  
754 [major results, \*Palaeogeography, Palaeoclimatology, Palaeoecology\*, 84, 191-215, 10.1016/0031-](#)  
755 [0182\(91\)90044-R, 1991.](#)

756 [Frumkin, A., and Stein, M.: The Sahara-East Mediterranean dust and climate connection revealed by](#)  
757 [strontium and uranium isotopes in a Jerusalem speleothem, \*Earth and Planetary Science Letters\*,](#)  
758 [217, 451-464, 10.1016/S0012-821X\(03\)00589-2, 2004.](#)

759 [Gasse, F., and Campo, E. v.: Abrupt post-glacial climate events in west Asia and north Africa](#)  
760 [monsoon domains, \*Earth and Planetary Science Letters\*, 126, 435-456, 1994.](#)

761 [Gasse, F.: Diatom-inferred salinity and carbonate oxygen isotopes in Holocene waterbodies of the](#)  
762 [western Sahara and Sahel \(Africa\), \*Quaternary Science Reviews\*, 21, 737-767, 2002.](#)

763 [Gat, J. R., Klein, B., Kushnir, Y., Roether, W., Wernli, H., Yam, R., and Shemesh, A.: Isotope](#)  
764 [composition of air moisture over the Mediterranean Sea: An index of the air-sea interaction pattern,](#)  
765 [Tellus, Series B: Chemical and Physical Meteorology, 55, 953-965, 10.1034/j.1600-](#)  
766 [0889.2003.00081.x, 2003.](#)

767 [Genty, D., Blamart, D., Ghaleb, B., Plagnes, V., Causse, C., Bakalowicz, M., Zouari, K., Chkir, N.,](#)  
768 [Hellstrom, J., Wainer, K., and Bourges, F.: Timing and dynamics of the last deglaciation from](#)  
769 [European and North African  \$\delta^{13}\text{C}\$  stalagmite profiles-comparison with Chinese and South](#)  
770 [Hemisphere stalagmites, \*Quaternary Science Reviews\*, 25, 2118-2142, 2006.](#)

771 [Goldsmith, Y., Polissar, P. J., Ayalon, A., Bar-Matthews, M., deMenocal, P. B., and Broecker, W. S.:](#)  
772 [The modern and Last Glacial Maximum hydrological cycles of the Eastern Mediterranean and the](#)  
773 [Levant from a water isotope perspective, \*Earth and Planetary Science Letters\*, 457, 302-312,](#)  
774 [http://dx.doi.org/10.1016/j.epsl.2016.10.017, 2017.](#)

775 [Grant, K. M., Grimm, R., Mikolajewicz, U., Marino, G., Ziegler, M., and Rohling, E. J.: The timing of](#)  
776 [Mediterranean sapropel deposition relative to insolation, sea-level and African monsoon changes,](#)  
777 [Quaternary Science Reviews, 140, 125-141, <http://dx.doi.org/10.1016/j.quascirev.2016.03.026>,](#)  
778 [2016.](#)

779 [Harrison, S., Bartlein, P., Izumi, K., Li, G., Annan, J., Hargreaves, J., Braconnot, P., and Kageyama, M.: Evaluation of CMIP5 palaeo-simulations to improve climate projections, \*Nature Climate Change\*, 5, 735-743, 2015.](#)

780

781

782 [Hoffmann, D. L., Rogerson, M., Spötl, C., Luetscher, M., Vance, D., Osborne, A. H., Fello, N. M., and Moseley, G. E.: Timing and causes of North African wet phases during MIS 3 and implications for Modern Human migration, \*Nature Scientific Reports\*, 6, 36367, 2016.](#)

783

784

785 [IPCC: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, United Kingdom and New York, 688, 2014.](#)

786

787

788 [Jolly, D., Prentice, I. C., Bonnefille, R., Ballouche, A., Bengo, M., Brenac, P., Buchet, G., Burney, D., Cazet, J. P., Cheddadi, R., Ector, T., Elenga, H., Elmoutaki, S., Guiot, J., Laarif, F., Lamb, H., Lezine, A. M., Maley, J., Mbenza, M., Peyron, O., Reille, M., Reynaud-Farrera, I., Riollet, G., Ritchie, J. C., Roche, E., Scott, L., Ssemmanda, I., Straka, H., Umer, M., Van Campo, E., Vilimumbalo, S., Vincens, A., and Waller, M.: Biome reconstruction from pollen and plant macrofossil data for Africa and the Arabian peninsula at 0 and 6000 years, \*Journal of Biogeography\*, 25, 1007-1027, 1998.](#)

789

790

791

792

793

794 [Langgut, D., Almogi-Labin, A., Bar-Matthews, M., Pickarski, N., and Weinstein-Evron, M.: Evidence for a humid interval at ~56–44 ka in the Levant and its potential link to modern humans dispersal out of Africa, \*Journal of Human Evolution\*, 124, 75-90, 10.1016/j.jhevol.2018.08.002, 2018.](#)

795

796

797 [McGarry, S., Bar-Matthews, M., Matthews, A., Vaks, A., Schilman, B., and Ayalon, A.: Constraints on hydrological and paleotemperature variations in the Eastern Mediterranean region in the last 140 ka given by the  \$\delta D\$  values of speleothem fluid inclusions, \*Quaternary Science Reviews\*, 23, 919-934, <http://dx.doi.org/10.1016/j.quascirev.2003.06.020>, 2004.](#)

798

799

800

801 [Meckler, A. N., Affolter, S., Dublyansky, Y. V., Krüger, Y., Vogel, N., Bernasconi, S. M., Frenz, M., Kipfer, R., Leuenberger, M., Spötl, C., Carolin, S., Cobb, K. M., Moerman, J., Adkins, J. F., and Fleitmann, D.: Glacial–interglacial temperature change in the tropical West Pacific: A comparison of stalagmite-based paleo-thermometers, \*Quaternary Science Reviews\*, 127, 90-116, <http://dx.doi.org/10.1016/j.quascirev.2015.06.015>, 2015.](#)

802

803

804

805

806 [Nicoll, K.: A revised chronology for Pleistocene paleolakes and Middle Stone Age – Middle Paleolithic cultural activity at Bîr Tîrfawi – Bîr Sahara in the Egyptian Sahara, \*Quaternary International\*, 463, 18-28, <https://doi.org/10.1016/j.quaint.2016.08.037>, 2018.](#)

807

808

809 [Osborne, A., Vance, D., Rohling, E., Barton, N., Rogerson, M., and Fello, N.: A humid corridor across the Sahara for the migration of early modern humans out of Africa 120,000 years ago, \*Proc. Natl. Acad. Sci. U. S. A.\*, doi 10.1073 pnas.0804472105, 2008.](#)

810

811

812 [Osborne, A. H., Marino, G., Vance, D., and Rohling, E. J.: Eastern Mediterranean surface water Nd during Eemian sapropel S5: monitoring northerly \(mid-latitude\) versus southerly \(sub-tropical\) freshwater contributions, \*Quaternary Science Reviews\*, 29, 2473-2483, <http://dx.doi.org/10.1016/j.quascirev.2010.05.015>, 2010.](#)

813

814

815

816 [Pascale, S., Gregory, J. M., Ambaum, M., and Tailleux, R.: Climate entropy budget of the HadCM3 atmosphere-ocean general circulation model and of FAMOUS, its low-resolution version, \*Climate Dynamics\*, 36, 1189-1206, 10.1007/s00382-009-0718-1, 2011.](#)

817

818

819 [Peleg, N., and Morin, E.: Convective rain cells: Radar-derived spatiotemporal characteristics and synoptic patterns over the eastern Mediterranean, \*J. Geophys. Res. D Atmos.\*, 117, 10.1029/2011JD017353, 2012.](#)

820

821

822 [PETIT-MAIRE, N., BURROLLET, P. F., A BALLAIS, J.-L., A FONTUGNE, M., A ROSSO, J.-C., A LAZAAR, A., and Gauthier-Villars, I.: Paléoclimats holocènes du Sahara septentrional. Dépôts lacustres et terrasses alluviales en bordure du Grand Erg Oriental à l'extrême-Sud de la Tunisie, \*Comptes rendus de l'Académie des sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre\* 312, 1661-1666, 1991.](#)

823

824

825

826

827 [Peyron, O., Jolly, D., Braconnot, P., Bonnefille, R., Guiot, J., Wirmann, D., and Chalief, F.: Quantitative reconstructions of annual rainfall in Africa 6000 years ago: Model-data comparison, \*Journal of Geophysical Research-Atmospheres\*, 111, D24110](#)

828

829

830 [Artn d24110, 2006.](#)

831 [Prentice, I. C., and Jolly, D.: Mid-Holocene and glacial-maximum vegetation geography of the](#)

832 [northern continents and Africa, \*Journal of Biogeography\*, 27, 507-519, 2000.](#)

833 [Revel, M., Ducassou, E., Grousset, F. E., Bernasconi, S. M., Migeon, S., Revillon, S., Mascle, J., Murat,](#)

834 [A., Zaragosi, S., and Bosch, D.: 100,000 Years of African monsoon variability recorded in sediments of](#)

835 [the Nile margin, \*Quaternary Science Reviews\*, 29, 1342-1362, 10.1016/j.quascirev.2010.02.006,](#)

836 [2010.](#)

837 [Rogerson, M., Schönfeld, J., and Leng, M.: Qualitative and quantitative approaches in](#)

838 [palaeohydrography: A case study from core-top parameters in the Gulf of Cadiz, \*Marine Geology\*,](#)

839 [280, 150-167, 2011.](#)

840 [Rogerson, M., Rohling, E. J., Bigg, G. R., and Ramirez, J.: Palaeoceanography of the Atlantic-](#)

841 [Mediterranean Exchange: Overview and first quantitative assessment of climatic forcing. , \*Reviews\*](#)

842 [of Geophysics, 50, DOI: 8755-1209/8712/2011RG000376, 2012.](#)

843 [Rohling, E., Marino, G., and Grant, K.: Mediterranean climate and oceanography, and the periodic](#)

844 [development of anoxic events \(sapropels\), \*Earth-Science Reviews\*, 143, 62-97, 2015.](#)

845 [Rohling, E. J., and Bryden, H. L.: Estimating past changes in the Eastern Mediterranean freshwater](#)

846 [budget, using reconstructions of sea level and hydrography, \*Proceedings Koninklijke Nederlandse\*](#)

847 [Akademie van Wetenschappen, Serie B](#), 97, 201-217, 1994.

848 [Rowan, J. S., Black, S., Macklin, M. G., Tabner, B. J., and Dore, J.: Quaternary environmental change](#)

849 [in Cyrenaica evidenced by U-Th, ESR and OSL of coastal alluvial fan sequences. , \*Libyan Studies\*, 31,](#)

850 [5-16, 2000.](#)

851 [Schwarcz, H. P., Harmon, R. S., Thompson, P., and Ford, D. C.: Stable isotope studies of fluid](#)

852 [inclusions in speleothems and their paleoclimatic significance, \*Geochimica et Cosmochimica Acta\*,](#)

853 [40, 657-665, \[http://dx.doi.org/10.1016/0016-7037\\(76\\)90111-3\]\(http://dx.doi.org/10.1016/0016-7037\(76\)90111-3\), 1976.](#)

854 [Smith, J. R., Giegengack, R., Schwarcz, H. P., McDonald, M. M. A., Kleindienst, M. R., Hawkins, A. L.,](#)

855 [and Churcher, C. S.: A reconstruction of quaternary pluvial environments and human occupations](#)

856 [using stratigraphy and geochronology of fossil-spring tufas, Kharga Oasis, Egypt, \*Geoarchaeology-an\*](#)

857 [International Journal](#), 19, 407-439, 2004.

858 [Smith, J. R., Hawkins, A. L., Asmerom, Y., Polyak, V., and Giegengack, R.: New age constraints on the](#)

859 [Middle Stone Age occupations of Kharga Oasis, Western Desert, Egypt, \*Journal of Human Evolution\*,](#)

860 [52, 690-701, 2007.](#)

861 [Swezey, C.: Eolian sediment responses to late Quaternary climate changes: temporal and spatial](#)

862 [patterns in the Sahara, \*Palaeogeography, Palaeoclimatology, Palaeoecology\*, 167, 119-155, 2001.](#)

863 [Toucanne, S., Angue Minto'o, C. M., Fontanier, C., Bassetti, M.-A., Jorry, S. J., and Jouet, G.: Tracking](#)

864 [rainfall in the northern Mediterranean borderlands during sapropel deposition, \*Quaternary Science\*](#)

865 [Reviews](#), 129, 178-195, <http://dx.doi.org/10.1016/j.quascirev.2015.10.016>, 2015.

866 [Tuenter, E., Weber, S. L., Hilgen, F. J., and Lourens, L. J.: The response of the African summer](#)

867 [monsoon to remote and local forcing due to precession and obliquity, \*Global and Planetary Change\*,](#)

868 [36, 219-235, 2003.](#)

869 [Vaks, A., Woodhead, J., Bar-Matthews, M., Ayalon, A., Cliff, R., Zilberman, T., Matthews, A., and](#)

870 [Frumkin, A.: Pliocene–Pleistocene climate of the northern margin of Saharan–Arabian Desert](#)

871 [recorded in speleothems from the Negev Desert, Israel, \*Earth and Planetary Science Letters\*, 368, 88-](#)

872 [100, 2013.](#)

873 [Van Breukelen, M., Vonhof, H., Hellstrom, J., Wester, W., and Kroon, D.: Fossil dripwater in](#)

874 [stalagmites reveals Holocene temperature and rainfall variation in Amazonia, \*Earth and Planetary\*](#)

875 [Science Letters](#), 275, 54-60, 2008.

876 [Wainer, K., Genty, D., Blamart, D., Daëron, M., Bar-Matthews, M., Vonhof, H., Dublyansky, Y., Pons-](#)

877 [Branchu, E., Thomas, L., van Calsteren, P., Quinif, Y., and Caillon, N.: Speleothem record of the last](#)

878 [180 ka in Villars cave \(SW France\): Investigation of a large  \$\delta^{18}O\$  shift between MIS6 and MIS5,](#)

879 [Quaternary Science Reviews](#), 30, 130-146, <http://dx.doi.org/10.1016/j.quascirev.2010.07.004>, 2011.

880 [Wassenburg, J. A., Immenhauser, A., Richter, D. K., Niedermayr, A., Riechelmann, S., Fietzke, J.,](#)  
881 [Scholz, D., Jochum, K. P., Fohlmeister, J., Schröder-Ritzrau, A., Sabaoui, A., Riechelmann, D. F. C.,](#)  
882 [Schneider, L., and Esper, J.: Moroccan speleothem and tree ring records suggest a variable positive](#)  
883 [state of the North Atlantic Oscillation during the Medieval Warm Period, Earth and Planetary Science](#)  
884 [Letters, 375, 291-302, <http://dx.doi.org/10.1016/j.epsl.2013.05.048>, 2013.](#)  
885 [Wassenburg, J. A., Dietrich, S., Fietzke, J., Fohlmeister, J., Jochum, K. P., Scholz, D., Richter, D. K.,](#)  
886 [Sabaoui, A., Spötl, C., Lohmann, G., Andreae, Meinrat O., and Immenhauser, A.: Reorganization of](#)  
887 [the North Atlantic Oscillation during early Holocene deglaciation, Nat. Geosci., 9, 602,](#)  
888 [10.1038/ngeo2767](#)  
889 <https://www.nature.com/articles/ngeo2767#supplementary-information>, 2016.

890  
891  
892  
893

#### 894 Figure Captions

895 Figure 1: Map showing the location of Susah Cave (filled circle) and GNIP sites used in the discussion  
896 (open circles). [Blue stars indicate sources of marine water evaporation discussed in the text. Grey](#)  
897 [arrows indicate recent average winter wind direction.](#)

898 Figure 2) Macroscopic structure of SC-06-01 speleothem, showing alternation of transparent and  
899 milky fabrics

900 Figure 3) Variability of water content ( $\mu\text{L}$ ) per unit mass of speleothem (g) in SC-06-01 fluid inclusion  
901 samples. Grey area shows working range of instrument.

902 Figure 4a) Fluid inclusion oxygen isotope values ( $\delta^{18}\text{O}_{\text{fi}}$ ; ~~blueblack~~ crosses) compared to calcite  
903 oxygen isotope values ( $\delta^{18}\text{O}_{\text{cc}}$ ; ~~blackblue~~ circles and line); 4b) Fluid inclusion hydrogen isotope values  
904 ( $\delta^2\text{H}_{\text{fi}}$ ; ~~blueblack~~ crosses) compared to  $\delta^{18}\text{O}_{\text{cc}}$  (~~blackblue~~ circles and line). Growth Phases I, II and III  
905 are shown as grey areas.

906 Figure 5a) Regional precipitation isotope data. Thick line represents Global Meteoric Water Line,  
907 dashed thick line represents Mediterranean Meteoric Water Line and thin lines representing  
908 expected range of deviation ( $\pm 10\text{‰}$   $\delta^2\text{H}_{\text{ppt}}$ ) below GMWL and above MMWL. Bet Dagan, Tunis and  
909 Sfax GNIP datasets ([http://www-naweb.iaea.org/napc/ih/IHS\\_resources\\_gnip.html](http://www-naweb.iaea.org/napc/ih/IHS_resources_gnip.html)). Sfax Atlantic  
910 and Mediterranean Rainfall are taken from Celle-Jeanton et al. ([20032001](#)). 5b-d) Summarised  
911 precipitation isotopes, and fluid inclusion measurements for SC-06-01 [for Phases I, II and II](#)  
912 [respectively](#).

913 Figure 6) Double-replicated fluid inclusion measurements from SC-06-01, and regional precipitation  
914 isotope trends.

915 Figure 7)  $^{87}\text{Sr}/^{86}\text{Sr}$  record for SC-06-01, compared to calcite  $\delta^{18}\text{O}_{\text{cc}}$  record (light grey line). Error bars  
916 are  $2\sigma$ . Growth Phases I, II and III are shown as grey areas.

917 Figure 8) Carbon isotope ( $\delta^{13}\text{C}_{\text{cc}}$ ) record for SC-06-01 compared to oxygen isotope record ( $\delta^{18}\text{O}_{\text{cc}}$ ;  
918 (Hoffmann et al., 2016)). Growth Phases I, II and III are shown as grey areas.

919 Figure 9) Fluid inclusion measurements relative to summarised precipitation data and the modern  
920 precipitation end members used in the discussion. Solid lines are the Meteoric Water Lines as in Fig.  
921 5a. Precipitation and fluid inclusion measurements are as shown in Figure 5b. “Mean Atlantic”, “Sfax  
922 Mixed”, “Sfax Med” and “High Precip Atlantic” indicate the mean of measurements in Celle-Jeanton  
923 et al. (~~2003~~, 2001) originating from Atlantic moisture, mixed source, Mediterranean moisture and  
924 High Precipitation measurements from an Atlantic moisture source (as described in Discussion)  
925 respectively. “Mean Bet Dagan” is the mean of GNIP measurements from this location, and “High  
926 Precip Bet Dagan” is the subset of high precipitation measurements as described in the Discussion.

927 Figure 10) Fluid inclusion deuterium excess ( $\delta^2\text{H}_{\text{excess-FI}}$ ) relative to calcite  $\delta^{18}\text{O}_{\text{cc}}$ . Note some fluid  
928 inclusions (70 to 60 ka BP and 40 to 30 ka BP) show high  $\text{D}_{\text{excess}}$  ( $\delta^2\text{H}_{\text{excess-FI}}$ ) indicative of an Eastern  
929 Mediterranean source. Growth Phases I, II and III are shown as grey areas.

930