



Spiky Fluctuations and Scaling in High-Resolution EPICA Ice Core Dust Fluxes

Shaun Lovejoy¹, Fabrice Lambert²

¹Physics Department, McGill University, 3600 University st., Montreal, Que. H3A 2T8, Canada

⁵²Department of Physical Geography, Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, Santiago, Chile

Abstract.

Atmospheric variability as a function of scale has been divided in various dynamical “regimes” with alternating 10 increasing and decreasing fluctuations: weather, macroweather, climate, macroclimate, megaclimate. Although a vast amount of data is available at small scales, the larger picture is not well constrained due to the scarcity and low resolution of long paleoclimatic time-series. Using statistical techniques originally developed for the study of turbulence, we analyse the 15 fluctuations of a centimetric resolution dust flux time-series from the EPICA Dome C ice-core in Antarctica that spans the past 800,000 years. The temporal resolution is 5 years over the last 400 kyr, and 25 years over the last 800 kyr, enabling the detailed statistical analysis and comparison of eight glaciation cycles, and the subdivision of each cycle into eight consecutive phases. The unique span and resolution of the dataset allows us to analyze the macroweather and climate scales 20 in detail, i.e. fluctuations with periodicities from 1 year to 100,000 years.

We find that the interglacial and glacial maximum phases of each cycle showed particularly large macroweather to 25 climate transition scale τ_c (around 2 kyr), whereas mid-glacial phases feature centennial transition scales (average of 300 yr). This suggests that interglacials and glacial maxima are exceptionally stable when compared with the rest of a glacial cycle. The Holocene (with $\tau_c \approx 7.9$ kyr) had a particularly large τ_c but it was not an outlier when compared with the phase 1 and 2 of other cycles. For each phase, we quantified the drift, intermittency, amplitude, and extremeness of the variability. Phases close to the interglacials (1, 2, 8) show low drift, moderate intermittency, and strong extremes, while the “glacial” middle phases 3-7 display strong drift, weak intermittency, and weaker extremes. Our results suggest that despite the large climatic changes occurring during glacial-interglacial transitions, glacial maxima, interglacials, and glacial inceptions were characterized by relatively stable atmospheric conditions, but punctuated by more frequent and severe droughts, than during the more unstable mid-glacial conditions. The low amplitude during phases 6-8 also suggests that the Patagonian ice sheet 30 was not yet fully developed before 30 kyr after glacial inception.



1 Introduction

Over the late Pleistocene, surface temperature variability is strongly modulated by insolation, both at orbital (Jouzel et al., 2007), and daily time scales. In between these two extremes, temperature variability has been shown to scale according 5 to power-law relationships, thus evidencing a continuum of variability at all frequencies (Huybers and Curry, 2006). However, although a vast amount of high-resolution data exists for modern conditions, our knowledge of climatic variability at glacial-interglacial time scales is usually limited by the lower resolution of paleoclimatic archive records, thus restricting high frequency analyses during older time sections. Previous analyses using marine and terrestrial temperature proxies from 10 both hemispheres suggest a generally stormier and more variable atmosphere during glacial times than during interglacials (Ditlevsen et al., 1996; Rehfeld et al., 2018).

One of the difficulties in characterizing climate variability is that ice core paleo-temperature reconstructions rapidly lose their resolutions as we move to the bottom of the ice column. Fig. 1 shows this visually for the EPICA Antarctic ice core (5787 measurements in all); the curve becomes noticeably smoother as we move back in time. In terms of data points, the most recent 100 kyr period has more than 3000 points (\approx 30 year resolution) whereas the most ancient 100 kyr period has 15 only 137 (\approx 730 year resolution). This implies that while the most recent glacial-interglacial cycle can be perceived with reasonable detail, it is hard to compare it quantitatively to previous cycles - or to deduce any general cycle characteristics.

Fluctuation analysis (Lovejoy and Schertzer, 2013; Nilsen et al., 2016), gives a relatively simple picture of atmospheric temperature variability (Fig. 2). The figure shows a series of regimes each with variability alternately increasing and decreasing with scale. From left to right we see weather scale variability, in which fluctuations tend to persist, building 20 up with scale - they are unstable - increasing up to the lifetime of planetary structures (about 10 days), followed by a macroweather regime with fluctuations tending to cancel each other out, decreasing with scale, displaying stable behaviour. In the last century, anthropogenically forced temperature changes dominate the natural (internal, macroweather) variability at about 10- 20 years. In pre-industrial periods the lower frequency climate regime starts somewhere between 100 and 1000 25 years (the macroweather-climate transition scale τ_c). Further to the right of Fig. 2, we can see the broad peak associated with the glacial cycles at about 50kyrs (half the 100 kyr period) and then at very low frequencies, the megaclimate regime again shows increasing variability with scale. In between the climate and megaclimate regimes, the fluctuations decrease with scale over a relatively short range from about 100 kyrs to 500 kyrs. However, the temperature fluctuations shown in Fig. 2 display average behavior, which can potentially hide large variations from epoch to epoch. In this paper, we use a uniquely long and high-resolution paleo dataset to analyze the macroweather and climate scales in detail.

30 We focus on the EPICA Dome C dust flux record, which has a 55 times higher resolution than the temperature record, including high resolution over even the oldest cycle (Lambert et al., 2012, Fig. 1). Antarctic dust fluxes are well correlated with temperature at orbital frequencies (Lambert et al., 2008; Ridgwell, 2003). But the fluxes are also affected by climatic conditions at the source and during transport (Lambert et al., 2008; Maher et al., 2010). The analysis of the dust record



presented here can therefore be thought of as a more “holistic” climatic parameter that includes not only temperature changes, but describes atmospheric variability as a whole (including wind strength and patterns, and the hydrological cycle).

2 Method

5 In order to proceed to a further quantitative analysis of the types of statistical variability, and of the macroweather-climate transition scale, we need to make some definitions. A commonly used way of quantifying fluctuations is the Fourier analysis. It quantifies the contribution of each frequency range to the total variance of the process. However, the interpretation of the spectrum is neither intuitive, nor straightforward (section 2.3). The highly non-Gaussian spikiness – for both dust flux and its logarithm (e.g. Fig. 3b, c), implies strong Fourier space spikes; indeed, (Lovejoy, 2018) found that the
10 probability distribution of spectral amplitudes can themselves be power laws. This has important implications for interpreting spectra, especially those estimated from single series (“periodograms”): if the spectral amplitudes are highly non-Gaussian, then we will typically see strong spectral spikes that are purely random in origin. This makes it very tempting to attribute quasi-oscillatory processes to what are in fact random spectral peaks. It therefore makes sense to consider the real (rather than Fourier) space variability (fluctuations). The problem here is that the spectrum is a second order statistical
15 moment (the spectrum in the Fourier transform of the autocorrelation function). While second order moments are sufficient for characterizing the variability of Gaussian processes, in the more general and usual case - especially with the highly variable dust fluxes - we need to quantify statistics of higher orders. In particular, the higher order statistics that characterize the extremes. Here, we will use two simple concepts to describe variability and intermittency (or spikiness) of the data.

The theoretical framework that we use in this paper is that of scaling, multifractals, the outcome of decades of
20 research attempting to understand turbulent intermittency. Intermittent – spiky transitions – characterized by different scaling exponents for different statistical moments - turns out to be the generic consequence of turbulent cascade processes. Although the cascades are multiplicative, the extreme probabilities generally turn out to be power laws (Mandelbrot, 1974; Schertzer and Lovejoy, 1987) - not log-normals (as was originally proposed by (Kolmogorov, 1962)). The analyses are based on scaling regimes and their statistical characteristics. Because scaling is a symmetry (in this case invariance of
25 exponents under dilations in time), the broad conclusions of our dust flux analyses – scaling regimes and their break points, stability/instability - are expected to be valid for the more usual climate parameters including the temperature. Although it is beyond our present scope, we will explore the scale by scale relationship between EPICA dust fluxes and temperatures in a future publication.

30 2.1 Haar Fluctuations

The basic tool we use to characterize variability in real space is the Haar fluctuation, which is simply the absolute difference of the mean over the first and second halves of an interval:



$$\Delta F(\Delta t) = \frac{2}{\Delta t} \int_{t-\Delta t/2}^t F(t') dt' - \frac{2}{\Delta t} \int_{t-\Delta t}^{t-\Delta t/2} F(t') dt' \quad (1)$$

We can characterize the fluctuations by their statistics. For example, by analyzing the whole dataset using intervals of various lengths, we can thus define the variability as a function of scale (i.e. interval length). If over a range of time scales Δt , there is no characteristic time, then this relationship is a power law, and the mean absolute fluctuation varies as:

$$5 \quad \langle |\Delta F(\Delta t)| \rangle \propto \Delta t^H \quad (2)$$

where “ $\langle \rangle$ ” indicates ensemble average, here an average over all the available disjoint intervals. A positive H implies that the average fluctuations increase with scale. This situation corresponds to unstable behavior identified with the climate regime. In contrast, when H is negative, variability converges towards a mean state with increasing scale. This is the situation found in the stable macroweather regime.

10 More generally, we can consider other statistical moments of the fluctuations, the “generalized structure functions”, $S_q(\Delta t)$:

$$S_q(\Delta t) = \langle |\Delta F(\Delta t)|^q \rangle \propto \Delta t^{\xi(q)} \quad (3)$$

If the fluctuations are from a Gaussian process, then their exponent function is linear: $\xi(q) = qH$. More generally however, $\xi(q)$ is concave and it is important to characterize this, since the nonlinearity in $\xi(q)$ is due to intermittency, i.e. 15 sudden, spiky transitions (for more details on Haar fluctuations and intermittency we refer to (Lovejoy and Schertzer, 2012)). We therefore decompose $\xi(q)$ into a linear and a nonlinear (convex) part $K(q)$, with $K(1)=0$:

$$\xi(q) = qH - K(q) \quad (4)$$

so that $K(q) = 0$ for quasi-Gaussian processes. Since the spectrum is a second order moment, the spectrum of a scaling process at frequency ω is a power law:

$$20 \quad E(\omega) \approx \omega^{-\beta} \quad (5)$$

where the spectral exponent $\beta = 1 + \xi(2) = 1 + 2H - K(2)$; $K(2)$ is sometimes termed the “intermittency correction”.

2.2 Intermittency

A simple way to quantify the intermittency is thus to compare, the mean and Root Mean Square (RMS) Haar 25 fluctuations:

$$S_1(\Delta t) = \langle |\Delta F(\Delta t)| \rangle \propto \Delta t^{\xi(1)} = \Delta t^H \quad (6)$$

$$S_2(\Delta t)^{1/2} = \left\langle (\Delta F(\Delta t))^2 \right\rangle^{1/2} \propto \Delta t^{\xi(2)/2} = \Delta t^{H-K(2)/2} \quad (7)$$



with ratio:

$$S_1(\Delta t)/S_2(\Delta t)^{1/2} = \langle |\Delta F(\Delta t)| \rangle / \left\langle (\Delta F(\Delta t))^2 \right\rangle^{1/2} \propto \Delta t^{K(2)/2} \quad (8)$$

where we estimate $S(\Delta t)$ using all available disjoint intervals of size Δt . These expressions are valid in a scaling regime. Since the number of disjoint intervals decreases as Δt increases, so does the sample size, hence the statistics are less reliable
 5 at large Δt .

For theoretical reasons (Lovejoy and Schertzer, 2013; Schertzer and Lovejoy, 1987), it turns out that the intermittency near the mean ($q=1$) is best quantified by the parameter $C_1 = K'(1)$. Since $K(1) = 0$, it turns out that for log-normal multifractals, (approximately relevant here) the ratio exponent $K(2)/2 \approx C_1$.

While the mean to RMS ratio is intuitive, a more accurate estimate of C_1 uses the intermittency function $G(\Delta t)$:

$$10 \quad G(\Delta t) = \left\langle \Delta F \right\rangle \left[\frac{\left\langle \Delta F^{1-\Delta q} \right\rangle}{\left\langle \Delta F^{1+\Delta q} \right\rangle} \right]^{1/(2\Delta q)} \propto \Delta t^{\xi(1)-\xi(1)} = \Delta t^{C_1}; \quad \Delta q \rightarrow 0 \quad (9)$$

whose exponent is C_1 . The intermittency exponent C_1 quantifies the *rate* at which the clustering near the mean builds up as a function of the range of scales over which the dynamical processes act; it only partially quantifies the spikiness. For this, we need other exponents, in particular the exponent q_D that characterizes the tails of the
 15 probability distributions. This is because scaling in space and/or time generically gives rise to power law probability distributions (Mandelbrot, 1974; Schertzer and Lovejoy, 1987). Specifically, the probability (Pr) of a random dust flux fluctuation ΔF exceeding a fixed threshold s is:

$$Pr(\Delta F > s) \approx s^{-q_D}; \quad s \gg 1 \quad (10)$$

Where the exponent q_D characterizes the extremes, for example, $q_D \approx 5$ has been estimated for wind or temperature
 20 (Lovejoy and Schertzer, 1986) and for paleotemperatures (Lovejoy and Schertzer, 2013) whereas $q_D = 3$ for precipitation (Lovejoy et al., 2012). A qualitative classification of probability distributions describes classical exponential tailed distributions (such as the Gaussian) as “thin tailed”, log normal (and log-Levy) distributions as “long-tailed”, and power law distributions as “fat tailed”. Whereas thin and long tailed distributions have convergence of all statistical moments, power distributions only have finite moments for orders $q < q_D$.

25

2.3 How Fluctuations help interpret spectra



Although spectra may be familiar, their physical interpretations are nontrivial, a fact that was underscored in (Lovejoy, 2015). In a scaling regime – a good approximation to the macroweather and climate regimes discussed here – the spectrum is a power law form (eq. 5) where the spectral exponent β characterizes the spectral *density*. Although β tells us how quickly the variance changes per frequency *interval*, its physical significance is neither intuitive nor obvious.

5 Integrating the spectrum over a frequency range is already easier to understand: it is the total variance of the process contributed by the range. Therefore we already see that $\beta-1$ (the exponent of the integrated spectrum) is more directly relevant than β . But even to understand this, we need to consider whether over a range of frequencies the process is dominated by either high or low frequencies. For this, we can compare the total variance contributed by neighboring octaves. For a power law spectrum, the variance ratio of one octave to its neighboring higher frequency octave is $2^{1-\beta}$.

10 From this, we see that $\beta > 1$ yields a ratio $2^{1-\beta} < 1$ implying low frequency dominance whereas when $\beta < 1$, we have $2^{1-\beta} > 1$ and high frequency dominance.

But what does low frequency or high frequency “dominance” mean physically? For this, it is easier to consider the situation in real space using fluctuations; the simplest relevant fluctuations being the Haar fluctuations ΔF discussed in section 2.2 that varies with time interval Δt as $\Delta F \approx \Delta t^H$. We saw that the exponents in real and spectral space were simply

15 related by $\beta = 1+2H - K(2)$ where $K(2) > 0$ due to the spikiness (intermittency). This formula leads to two important conclusions. First, if we ignore intermittency (putting $C_1 = 0$, hence $K(2) = 0$) and assume that the mean fluctuations scale with the same exponent as the RMS fluctuations, then $H = (\beta - 1)/2$ showing again that it is the sign of $\beta - 1$ that is fundamental: $\beta > 1$ implies $H > 0$ hence fluctuations grow with scale and the process “drifts” or “wanders”, it is unstable. Conversely $\beta < 1$ implies $H < 0$ hence fluctuations decrease with scale and the process “cancels”, “converges”, it is “stable”.

20 The second conclusion is that if intermittency is strong (here we typically have $C_1 \approx 0.1$, $K(2) \approx 0.2$), then the relationship between the second and first order statistical moments is a little more complex so that for example, with these values and a $\beta \approx 0.9$ we would have high frequencies dominating the variance ($\beta < 1$) but low frequencies dominating the mean ($H > 0$).

3 Results

3.1 Looking at the data

25 Unlike water isotopes that diffuse and lose their temporal resolution in the bottom section of an ice core at high pressures and densities, the relatively large dust particles diffuse much less and have been used to estimate the dust flux over every centimetre of the 3.2 km long EPICA core (298,203 measurements, (Lambert et al., 2012)). The temporal resolution of this series varies from 0.81 years to 11.1 yrs (the averages over the most recent and the most ancient 100 kyrs respectively). The worst temporal resolution of 25 years per centimeter occurs around 3050 m depth, with the result that at that resolution,

30 there are virtually no missing data points in the whole record (Fig. 1).



Dust measurements cannot be assigned to one particular atmospheric variable, like temperature for the water isotopes. The amount of dust deposited in East Antarctica will depend on the vegetation cover at the source region (mostly Patagonia for East Antarctic dust (Delmonte et al., 2008)), on the amount of dust available in the source region (can depend on the presence of glaciers), on the strength of the prevailing winds between South America and Antarctica, and the strength of the hydrological cycle (more precipitation will wash out more dust from the atmosphere (Lambert et al., 2008)). At low frequencies the dust variability will be driven by conditions at the source (presence of glaciers, vegetation cover), which is primarily driven by Southern Hemisphere temperature, explaining the high correlation between dust and temperature in ice cores. At high frequencies however, dust and temperature are decoupled and dust variability will be driven by changes in wind and the hydrological cycle. A single dust peak within a low background may therefore reflect a short-term atmospheric disturbance like drought over South America or low precipitation over the Southern Ocean. The analysis presented here focuses heavily on the occurrence of dust fluctuations, the physical interpretation of which will depend on the scale of the phenomenon.

Fig. 3a shows a succession of 10 factors of 2 “blowdowns” (upper left to lower right at 11 different resolutions). In order to avoid smoothing, the data was “zoomed” in depth rather than time, but the point is clear: the signal is very roughly scale invariant, at no stage is there any sign of obvious smoothing, and the quasi-periodic 100 kyr oscillations is the only obvious time scale (we quantify this below). In comparison with more common paleoclimate signals such as temperature proxies, the dust flux itself is already quite spiky. However, it also displays spiky transitions. In Fig. 3b we show the absolute change in dust flux and one can visually see the strong spikiness associated with strongly non Gaussian variability: the intermittency. At each resolution, the solid red line indicates the maximum spike expected if the process was Gaussian, and the upper dashed lines the expected level for a (Gaussian) spike with probability 10^{-6} . Again, without sophisticated analysis, we can see that the spikes are wildly non-Gaussian, frequently exceeding the 10^{-6} level even though each segment has only 290 points, with the spikiness being nearly independent of resolution.

Taking the logarithms of the dust flux is a common practice since it reduces the extremes and makes the signal closer to the temperature and other more familiar atmospheric parameters. We therefore show the corresponding spike plot for the log transformed data (fig. 3c). Although the extreme spikes are indeed less extreme (see also fig. 6a, b), we see that the transformation has not qualitatively changed the situation with spikes still regularly exceeding (log) Gaussian probability levels of 10^{-5} and occasionally 10^{-8} .

3.2 Spectra

Figure 4 shows various spectral analyses (for the corresponding fluctuation analyses, see fig. 5). There is a clear periodicity at about $(100 \text{ kyr})^{-1}$. In the double power law fit (line plot), the transition frequencies are a little lower: $\omega_0 = (160 \text{ kyr})^{-1}$ (flux) and $\omega_c = (145 \text{ kyr})^{-1}$ (log flux), although a Gaussian fit near the max gives a spike at $(94 \pm 9 \text{ kyr})^{-1}$. Note that it is actually a little bit “wide” (two peaks) hence it is not perfectly periodic, and the amplitude is only about a factor 4 above the



background. In comparison, the amplitude of the annual temperature frequency peak is several thousand times above the background (depending on the location) and is narrower (not shown).

Since this is a log-log plot, power laws appear as straight lines. We show in the figure the fits to the bi-scaling function

5
$$E(\omega) = \frac{a}{(\omega/\omega_0)^{\beta_h} + (\omega/\omega_0)^{\beta_l}}$$

that smoothly transitions between a spectrum with $E(\omega) \propto \omega^{-\beta_h}$ at $\omega > \omega_0$ and $E(\omega) \propto \omega^{-\beta_l}$ at $\omega < \omega_0$. The figure shows the regressions with $\beta_l = -2.5$, $\beta_h = 1.7$, and $a = 7.5$ $(\text{mg/m}^2/\text{yr})^2\text{yr}$, $\omega_0 \approx (145\text{kyrs})^{-1}$ for the fluxes, and $a = 0.375$ $(\text{mg/m}^2/\text{yr})^2\text{yr}$, $\omega_0 \approx (160\text{kyrs})^{-1}$ for the logarithms of fluxes. According to the figure, the high frequency climate regime scaling continues to about $(300 \text{ yrs})^{-1}$ before flattening to a very high frequency scaling ($\beta_m \approx 0.8$) “macroweather” regime 10 (Lovejoy and Schertzer, 2013). Note that this spectral transition scale is close but not identical to the transition scale estimated in real space with fluctuation analysis (around 250 years, Fig. 5). The scaling exponents $\beta_h = 1.7$ and $\beta_m = 0.8$ corresponding to the climate and macroweather regime respectively, may be compared with the values 2.1 and 0.4 for the EPICA paleotemperatures discussed in a future publication (compare however the red and black curves in Fig. 2). A review covering nearly 2 dozen β_h estimates for temperature proxies from both hemispheres was given in (Lovejoy and Schertzer, 15 2013) (see especially table 11.4; also (Ditlevsen et al., 1996; Huybers and Curry, 2006; Shao and Ditlevsen, 2016)). Although the dust and temperature exponents are not identical - implying scale-varying correlations – these results do support the use of dust as a proxy for atmospheric variability.

The plot graphically counterposes two views of the variability. Although we clearly see a spectral maximum at around $(100 \text{ kyrs})^{-1}$, the broad bispectral scaling model already accounts for 96% of the spectral energy (variance) leaving 20 only 4% for the (extra) contribution from the (near) $(100\text{kyrs})^{-1}$ Milankovitch frequency. If it is argued that the logarithm of the flux is more physically relevant (blue spectrum) the situation is barely changed. Alternatively, we may take a narrow spectral spike model that approximates the spectral spike near $(100 \text{ kyr})^{-1}$ as a Gaussian shaped profile. With this model, the spike is localised at $(94 \pm 9 \text{ kyrs})^{-1}$ and contributes a total of 31% of the total variance. However, not all of this is above what we would expect from a scaling background; the exact amount depends on how the background is defined. For example, 25 over the range from the 6th to the 11th highest frequencies in this discrete spectrum (from $(133 \text{ kyrs})^{-1}$ to $(72 \text{ kyrs})^{-1}$), in comparison to the background over this range, there is an enhancement of about 80% due to the strong peaks (the enhancement is about 100% for the 7th to the 12th frequencies). This means although the $(94 \pm 9 \text{ kyr})^{-1}$ peak represents 31% of the total variability over the range from $(800 \text{ kyrs})^{-1}$ to $(25 \text{ yrs})^{-1}$, it is only about 15% above the “background” (note that only 5% of the total variance is between $(25\text{yrs})^{-1}$ and $(1 \text{ kyr})^{-1}$). The overall conclusion is that the background represents 30 between 85% and 96% of the total variance.



3.3 Haar Fluctuation Analysis

Figure 5 shows that Haar fluctuations have simple interpretations in terms of the variability of the dust flux. For example, typical variations over a glacial-interglacial cycle (half cycle ≈ 50 kyr) are about $\pm 3 \text{ mg/m}^2/\text{yr}$ (dashed line). From the figure we see there is a short regime with $H < 0$ (up to about 250 yrs), a scaling regime with $H > 0$ (up to glacial-interglacial periods (≈ 50 kyr)) and finally a long time scale decrease in variability that is possibly (but not obviously) scaling. As expected, the regimes correspond to those indicated in Fig. 4 with the relation $\beta = 1 + \xi(2)$ where $E(\omega) \approx \omega^{-\beta}$ and represent macroweather, climate, and macroclimate, respectively.

Fig. 6a shows the fluctuation probabilities of the entire 800 kyr series at 25 year resolution. We see that the large fluctuations (the tail) part of the distribution is indeed quite linear on a log-log plot with exponents $q_D \approx 2.75$ and 2.98 in time and depth respectively (both fit to the extreme 0.1% of the distributions). To get an idea of how extreme these distributions are, consider the depth distribution with $q_D = 2.98$. With this exponent, dust flux fluctuations 10 times larger than typical fluctuations occur only $10^{2.98} \approx 1000$ times less frequently. In comparison, for a Gaussian, they would be $\approx 10^{23}$ times less likely; they would never be observed.

While the fluxes are positive definite and so cannot be Gaussians, the increments analyzed here could easily be approximately so. Nevertheless, a common way of trying to tame the spikes is by making a log transformation of fluxes. Fig. 4 already showed that this did not alter the spectrum very much; here it similarly has only a marginal effect. For example Fig. 6b shows that the extreme tails on the log dust flux distribution has $q_D = 3.60$ in time (25 yrs) and 4.59 in depth (at 1 cm resolution; this is close to the value $q_D \approx 5$ reported for both GRIP and Vostok paleotemperatures in (Lovejoy and Schertzer, 2013)). The log-transformed variable still displays huge extremes with the extreme log flux corresponding to a log-Gaussian probability of 10^{-30} and 10^{-50} (time, depth respectively). Whether or not taking logarithms yields a more climate relevant parameter, it does not significantly change the problem of intermittency or of the extremes.

These power law fluctuations are so large that according to the classical assumptions, they would be outliers. While Gaussians are mathematically convenient and can be justified when dealing with measurement errors, in atmospheric science thanks to the scaling, very few processes are Gaussian. This has important applications in tipping point analysis, where noise induced tipping points are generally studied using well behaved white or Gaussian noise (e.g. Dakos et al., 2012).

3.4 Phases

Scaling is a statistical symmetry. In our case, it means that *on average* the statistics at small, medium and large scales are the same in some way. The difficulty is that on a single realization – such as that available here, a single core from a single planet earth – the symmetry will necessarily be broken. For example, in the spectrum Fig. 4, in each of the proposed scaling regimes, scaling only predicts that the actual spectrum from this single core will vary about the indicated straight lines that represent the ensemble behaviour. Since this variability is strong, we made the potential scaling regimes more obvious by either averaging the spectrum over frequency bins (the red and blue spectra) – or by breaking the series into



shorter parts and averaging the spectra over all the parts, effectively treating each segment as a separate realization of a single process (green).

This already illustrates the general problem: in order to obtain robust statistics we need to average over numerous realizations – and since here we have a single series, the best we can do is to break the series into disjoint segments and 5 average the statistics over them. Yet at the same time, in order to see the wide-range scaling picture (which also helps to more accurately estimate the scaling properties/exponents), we need segments that are as long as possible. The compromise that we chose between numerous short segments and a small number of long ones was to break the series into 8 glacial-interglacial cycles, and each cycle into 8 successive phases. As a first approximation, we defined eight successive 100kyr periods (hereafter called “segments”, Fig. 7, top set), corresponding fairly closely to the main periodicity of the series. As we 10 discussed, the spectral peak is broad implying that the duration of each cycle is variable – the cycles are only “quasi-periodic”. It is therefore of interest to consider an additional somewhat flexible definition of cycles defining them as the period from one interglacial to the next (hereafter called “cycle”, Fig. 7, bottom set). The break points were taken at interglacial optima: 0.4, 128.5, 243.5, 336, 407.5, 490, 614, 700, 789 kyrs BP, i.e. 96.9 ± 18.7 kyrs per cycle. Using this latter 15 definition, the cycles were nondimensionalized so that nondimensional time was defined as the fraction of the cycle, effectively stretching or compressing the cycles by $\pm 19\%$.

With either of these definitions, we have 8 segments or cycles, each with 8 phases. Note that in our nomenclature, phase 1 and 8 are the youngest and oldest phases, respectively, and that time flows from phase 8 to phase 1. Fig. 8 shows the phase by phase information summarized by the average flux over each cycle including the dispersion of each cycle about the mean (for the segments in the top set, and the cycles in the bottom set). We see that the variability is highest in the middle of 20 a cycle and lowest at the ends.

Since the spectra in Fig. 4 showed that there were wide scale ranges that are scale invariant – power laws - we are interested in characterizing the scaling properties over the different phases. In Fig. 9 we compares the statistics averaged over cycles and the statistics averaged over phases. The figure shows that the phase to phase differences are much more important than the cycle to cycle differences. Particularly noticeable are the phase to phase differences in the average 25 fluctuations $\langle|\Delta F(\Delta t)|\rangle$ (lower left).

From the global statistics (e.g. Figs. 4, 5), it is clear that in each glacial-interglacial cycle there are two regimes, so that before characterizing the structure functions by their exponents (e.g. $H = \xi(1)$ for the mean fluctuations), we have to determine the macroweather-climate transition time scale τ_c whose average (from Fig. 4, 5) is 250-300 years.

One way of estimating the transition scale τ_c is to make a bilinear fit of $\log_{10} S_1(\Delta t)$ (i.e. Haar with $q = 1$, the mean 30 absolute fluctuation) with the mean slopes -0.05 (small Δt) and slope +0.25 (large Δt ; the values were chosen because they are roughly the H estimates from the average over all the cycles) (Fig. 10). Bilinear fits were made for each phase of each segment (blue) as well as for each phase of each cycle (black). For each phase there were thus 8 transition scales, which were used to calculate the mean and its standard deviation, (shown here as representative black arrows). From the figure we see



that at first (phases 8-3) the transition scale is relatively short (250–400 yr), but that it rapidly moves to longer (1 – 2 kyr) scales for the final phases 2 and 1. The average transition scale over all phases is around 300 years.

The figure shows that our results are robust since the results are not very different using dimensional and nondimensional time (segments and cycles). Comparing the blue and black curves, we see that in all cases the late phases have much larger τ_c than the early and middle phases. Also shown in Fig. 10 (dashed) is a plot of the break points estimated by a more subjective method that attempts to visually determine a break point on $\log S_1 - \log \Delta t$ plots. Again, we reach the same conclusion with quantitatively very similar results: a transition of millennia for phases 1 and 2, and a few centuries in the middle of the cycle. The cycle average value ($\tau_c \approx 300$ years) is therefore not representative of the latest phases where τ_c is many times larger (glacial maxima and interglacials). The Holocene has an even larger transition scale ($\tau_c = 7.9$ kyr), marked by an X in Fig. 10), but it lies just outside the standard deviation of the first nondimensional phases (red arrows in Fig. 10). Although the Holocene value of τ_c is the largest in phase 1, it corresponds to 1.55 standard deviations above the mean with a corresponding p value of 0.12, roughly the expected extreme of a sample of 8; it is therefore not a statistical outlier.

Alternatively, rather than fixing a phase and determining the variation of the mean fluctuation and intermittency function (Fig. 10), we can consider the variation of the Haar fluctuations at fixed time scales and see how they vary from phase to phase (Fig. 11). The figure shows the phase to phase variation of Haar fluctuations at 50, 100, 200, 400, 800, 1600, 3200, 7000 years scales (bottom to top). Over the macroweather regime (up to 400 to 800 years) the fluctuations tend to cancel so that the variability is nearly independent of time scale. In contrast, once we reach the scales in the climate regime (800 to 7000 years), the fluctuations increase noticeably as the time interval Δt is increased. For every time scale, there is a clear cyclicity (left to right), with fluctuation amplitudes largest in the middle phases. We note that the cycle to cycle variability is fairly large; about a factor of 2 (for clarity the error bars indicating this cycle to cycle spread were not shown).

Finally, we describe for each phase the drift tendency and the intermittency, as well as fluctuation amplitude and extremeness of the data. In Figure 12 we show the result on the nondimensional phases of the range 500 years $< \Delta t <$ 3000 years, (upper left and right; the range was chosen to be mostly with $\Delta t > \tau_c$, and it was fixed so as to avoid any uncertainty associated with the algorithm used to estimate τ_c). Recall that the fluctuation exponent $H > 0$ quantifies the rate at which the average fluctuations increase with time scale. Similarly, the exponent C_1 characterizes the rate at which the spikiness near the mean (the intermittency exponent) increases with scale. We see (upper left) that H is fairly high in the early phases with H reaching small value in the later phases (with H actually a little bit negative on average in phase 1 due to the large τ_c value). C_1 on the other hand (upper right) decreases a bit in the middle the phases. The error bars show that there is quite a lot of cycle to cycle variability.

If H quantifies the “drift” and C_1 the “spikiness”, then Fig. 12 shows that the early phases have high drift and medium spikiness, the middle phases have high drift and lower spikiness, while phases 1-2 have low drift but medium spikiness. To understand this better, consider the transition time scales in Fig. 10. The youngest 2 phases with the low drift and spikiness



are also the phase with the longest transition scales. This means that the rate at which the variability builds up is small and that it only builds up over a short range of scales (from τ_c to roughly $\Delta t = 50$ kyrs, the half cycle duration, this can be checked on Fig. 9 that shows the phase by phase structure functions and intermittency functions). Conversely, phases 3 and 4 with high drift and high intermittency also have a smaller τ_c so that both the fluctuations and spikiness build up faster (Fig. 5 11) and over a wider range of scales (Fig. 10).

Another useful characterisation of the phases is to directly consider the flux variability at a fixed reference scale, taken here as the 25 year resolution; quantifying the amplitude of the variability of each segment by its standard deviation A at 25yr time scale (Fig. 12, lower left). This is not the *difference* between neighbouring values or fluctuation (as in figure 11), it is rather the variability of the series itself at 25 year resolution. For each of the phases, we have 8 estimates (one from 10 each cycle); these are used to calculate the mean (black) and standard deviation shown by the error bars. We can see that the amplitude of the 25 yr scale fluctuations is about four times higher in the middle of the ice age (phase 4) than at interglacial (phase 1). The figure clearly shows the strong change of variability across the cycle.

Whereas C_1 characterizes the intermittency near the mean, we have seen that the probability exponent q_D characterizes the extreme spikiness. Fig. 12 lower right, compares q_D phase by phase. Recalling that small q_D implies more 15 extreme extremes, we see that the extremes are stronger in the beginning and end of the cycle, and somewhat less pronounced in the middle phases of the cycle (note the overall mean is 2.62 ± 0.42 , this can be compared to the value $q_D = 3.60$ for the overall log transformed data, fig. 6b). Notice that for phase 8, $q_D=2.03$ (the mean); this is close to the value $q_D = 2$ below which the extremes are so strong that the variance (and hence spectrum) does not converge. An extreme (low) 20 exponent q_D phase implies that most of the time the changes in flux are small, but occasionally, there are huge transitions. Conversely, a high (less extreme) q_D implies that there is a wider range of different flux changes so that most of the changes tend to be in a restricted range. We can now categorize the phase by phase spikiness as: extremes strong, and medium spikiness (phases 1, 2, 8), and extremes intermediate and low spikiness (phases 3-7). For the cycle to cycle estimates (not shown), the value $q_D = 2.75 \pm 0.41$, seems to be fairly representative of all the cycles, although there is a slight tendency for q_D to decrease for the older cycles implying that they may have been a bit more extreme than the recent ones.

25

4 Discussion

An attractive aspect of dust fluxes is that they are paleo indicators with unparalleled resolutions over huge ranges of temporal scales. However, they come with two difficulties. First, their physical interpretation is not clear: while they depend on temperature, wind, and precipitation, and so are holistic climate indicators, the precise climate significance of dust flux variability is hard to nail down. Second, their appearance as a sequence of strong spikes is unlike that of any of the familiar 30 proxies. Indeed, we argued that their highly spiky (intermittent) nature is outside the purview of conventional statistical frameworks including autoregressive, moving average or more generally of quasi-Gaussian or even quasi log-Gaussian processes.



Due to the dominance of the continuum, (spectral, background) variability, physical interpretations must be based on an understanding of climate variability as a function of scale. We will first consider overall analyses over the whole dust flux series, and then focus on the phases. The spectral analysis (Fig. 4) is the most familiar and it is qualitatively similar to previous results obtained with temperature data, although temperature spectra with anything approaching the resolution of Fig. 4 are only possible over the most recent glacial cycle. The most striking spectral feature is the peak over the background at 100 kyr periodicity. The broadness of this peak already indicates the irregularity of the eccentricity-forced Milankovitch cycles. More surprising is the (near) absence of obliquity frequencies at 41 kyr. Although there is definitely power in that frequency range, it is not significantly larger than the background continuum, suggesting low internal feedback to that forcing. Finally, our high-resolution data allows us to discern two different power-law regimes, one at low frequencies with an exponent $\beta = 1.7$, and one at high frequencies with exponent $\beta = 0.8$, with the transition between the two at around 300 years.

In section 2.3, we discussed some of the difficulties inherent in interpreting spectra and showed that the exponent of the integrated spectrum $\beta-1$ is more directly relevant than β . Applying this understanding to the dust exponents, we see that in macroweather, there is a weak high frequency dominance ($1-\beta \approx 0.2 > 0$) whereas the climate regime is dominated by low frequencies ($1-\beta \approx -0.7 < 0$). A plausible physical explanation is that over long periods of time (at large scales), the amount of dust in the SH atmosphere is driven by changes in glacier and vegetation coverage, which is itself forced by SH temperature change. There is therefore a very strong correlation between dust and temperature at climatic scales. At higher frequencies in the macroweather regime, temperature oscillations are too fast to overcome the inertia of ice sheet and vegetation responses. Instead, dust deposition in Antarctica will be more sensitive to temporary atmospheric disturbances in the winds and the hydrological cycle.

To interpret the analysis by phase of the dust record (Fig. 12) one must understand the significance of A and of the exponents H , C_1 , and q_D in the context of dust deposition. The H exponent and the amplitude A are linked to oscillations/fluctuations. We saw that the H -exponent signifies a tendency to “drift”, meaning that when $H < 0$, the dust oscillations will tend to cancel each other out and the record will cluster around a mean value. In contrast, $H > 0$ indicates that the dust fluxes will not cluster around a mean value, in essence, the process wanders and does not stay constant. The low H numbers during phases 1 and 2 (interglacial and glacial maximum) indicate a very constant, stable climatic state, with Patagonian dust production being either very low during interglacials (low glacier activity, large vegetation cover) or very high (Patagonian ice cap fully grown, large outwash plains on the Argentinean side). In contrast, the high H and amplitude A values during the mid-glacial may have been due to strong variability in glacier extent during that time, and therefore a very variable dust supply (see also Fig. 11 that shows how the amplitude of the fluctuations at different time scales varies with the phase). The glacial inception (phases 7 and 8) features low A but a high H exponent. This implies that the mean dust level was highly variable, but the dust supply was still low, thus not allowing for large amplitude fluctuations. The higher



amplitudes in phases 6 and 7 indicates that dust supply became abundant then, which suggests that the Patagonian ice cap had already grown to a substantial size about 30 kyr after glacial inception.

The exponents C_1 and q_D are associated with the intermittency, or spikiness of the data. While C_1 is a measure of the sparseness of the spikes whose amplitudes contribute most to the mean, q_D characterizes how extreme these extreme spike values are. The dust flux record is generally more intermittent in phases 8, 1, and 2 (glacial inception, interglacial, glacial maximum) than in the mid-glacial, with also more extreme spike values. Since the C_1 and q_D exponents are calculated from the derivative of the signal, a spike and a fast change in the system state (e.g. Dansgaard-Oeschger event in the NH) will both produce a similar signal. However, such fast changes in system state do not occur in the SH where the corresponding signal to NH DO events is more triangular and gradual in shape. We therefore interpret the C_1 and q_D exponents as purely indicative of spikes in the dust signal. A short large spike (<100 years) in dust deposition cannot be associated to ice sheet changes which have generally larger reaction times. Its origin is therefore due to vegetation and/or atmospheric changes. Large-scale natural fires could alter the landscape in a very short time, allowing for more dust uptake by the winds and a sudden rise in atmospheric dust. The recuperation of vegetation cover would be more gradual, though, resulting in a saw-tooth shape of the dust spike that we do not observe in the data. The most likely explanation for a dust spike is therefore a short-term disturbance in the atmosphere, involving either the winds or the hydrological cycle (or both at the same time). The obvious candidate for a perturbation that would lead to increased dust in the atmosphere is drought. We will therefore interpret short dust spikes as multiannual to multidecadal drought events in southern South America. With this interpretation, we can conclude that glacial maxima, interglacials, and glacial inceptions were characterized by more frequent and more severe drought events than during the mid-glacial. During glacial maxima, such extreme dust events could have contributed to Southern Hemisphere deglaciation by significantly lowering ice sheet albedo at the beginning of the termination (Ganopolski and Calov, 2011). In contrast, more frequent dust events could have contributed to glacial inception through negative radiative forcing of the atmosphere.

5 Conclusions

Until now, a systematic comparison of the different glacial-interglacial cycles has been hindered by a limitation of the most common paleoclimate indicators – the low resolution of temperature reconstructions from ice or marine sediment cores. Due to this intrinsic characteristic, the older cycles are poorly discerned; we gave the example of EPICA paleo temperatures whose resolution in the most recent cycle was 25 times higher than the resolution in the oldest one. In this paper, we therefore took advantage of a unique dust flux dataset with 1 cm resolution measuring 320,000 cm. The most recent four cycles were discerned at 5 year resolution throughout (20,000 points per cycle) and the entire record of eight glaciations could be resolved at 25 years, and this, without signs of over-sampling or smoothing.

Dust fluxes are challenging not only because of their high resolutions, but also because of their unusually high spikiness (intermittency) and their extreme transitions that occur over huge ranges of time scales. Standard statistical



methodologies are inappropriate for analyzing such data. They typically assume exponential decorrelations (e.g. autoregressive or moving average processes) that have variability confined to narrow ranges of scale. In addition, they assume that the variability is quasi Gaussian or at least that it can be reduced to quasi Gaussian through a simple transformations of variables (e.g. by taking logarithms). In this paper, using standard spectral and probability distribution analysis, we show that both the spectral and the probability tails were power laws, not exponential, requiring nonstandard approaches.
5

The high resolution of the data allowed us to not only quantitatively compare glacial-interglacial cycles with each other, but also to subdivide each cycle into 8 successive phases that could also be compared to one another. One of the key findings was that there was a great deal of statistical similarity between the different cycles and that within each cycle there
10 were systematic variations of the statistical properties with phase. These conclusions would not have been possible with the corresponding much lower resolution temperature data.

Our variability analysis using real space (Haar) fluctuations confirmed that the majority of the variability was in the macroweather and climate scaling regime “backgrounds” with an average transition scale τ_c of about 300 years. In the climate regime (time scales above τ_c), dust variability is more affected by long-term hemispheric-wide climate changes
15 affecting slow response subsystems like glaciers and vegetation. In contrast, dust variability in the macroweather regime (time scales below τ_c) would have been more influenced by short-term atmospheric perturbations.

Using various techniques, τ_c was found to be systematically larger in the youngest two phases than in the middle and oldest phases; about 2 kyr but with nearly a factor of 4 cycle to cycle spread and equal to 300 years (with a factor of 2 spread) for the six remaining phases. For the Holocene, τ_c was found to be 7.9 kyr, which makes it an exceptionally stable
20 interglacial, but not a statistical outlier compared to other interglacials. Similarly, the typical (RMS) variation in flux amplitude was smaller in the early phase increases by (on average) a factor of 4 from $\pm 0.13 \text{ mg/m}^2/\text{yr}$ to about $\pm 0.5 \text{ mg/m}^2/\text{yr}$ in the middle and later phases. The Holocene (with an amplitude of $\pm 0.08 \text{ mg/m}^2/\text{yr}$) was again particularly stable with respect to the phase 1 of other cycles, but it was not an outlier.

The task of statistically characterizing the cycles reduced primarily to the problem of characterizing the phases’
25 variability exponents H , C_1 , q_D and amplitude A We show that the atmosphere was relatively stable during glacial maxima and interglacials, but highly variable during glacial inception and mid-glacial. However, the low amplitude of dust variability during glacial inceptions indicates that the Patagonian ice sheet was not very active until ~30 kyr after glacial inception.

We interpret the intermittency indicators as suggesting a higher frequency of drought events and more severe
30 droughts during glacial inception, interglacials, and glacial maxima than during mid-glacial conditions. These short-term spikes in atmospheric dust could have helped trigger southern hemisphere deglaciation through albedo feedback of ice-sheet surfaces, or glacial inception through negative radiative forcing.




This paper is an early attempt to understand this unique very high resolution data set. In future work, we will extend our methodology to the EPICA paleo temperatures and to the scale by scale statistical relationship between the latter and the dust fluxes.

5 Acknowledgements

SL's contribution to this fundamental research was unfunded and there were no conflicts of interest. FL acknowledges support by CONICYT projects Fonadap 15110009 and Fondecyt 1171773, and the Millennium Nucleus Paleoclimate. MN Paleoclimate is a Millennium Nucleus supported by the Millennium Scientific Initiative of the Ministry of Economy, Development and Tourism (Chile). Data available here: <https://doi.pangaea.de/10.1594/PANGAEA.779311>



Figures



5 Figure 1: Temperature (blue) and dust flux (red) from the EPICA Dome C ice core (Jouzel et al., 2007; Lambert et al., 2012). The dust flux time series has 32,000 regularly spaced points (25 year resolution), the temperature series, has 5,752 points. The temperature data are irregularly spaced, and lose resolution as we go back into the past (number of temperature data points in successive ice ages: 3022, 1117, 521, 267, 199, 331, 134, 146). In both cases we can make out the glacial cycles but they are at best only quasi-periodic.




Figure 2: A composite showing root mean square (RMS) Haar fluctuations (ΔT in units of $^{\circ}\text{C}$) black, and RMS dust

5 fluctuations analysed in this paper (red, in units of $\text{mg}/\text{m}^2/\text{yr}$, (Lambert et al., 2012)). From left to right: thermistor temperatures at 0.0167s resolution (Lovejoy, 2018), hourly temperatures from Landers Wyoming (Lovejoy, 2015), daily temperatures from 75 °N (Lovejoy, 2015), EPICA Dome C temperatures (Jouzel et al., 2007), and two marine benthic stacks (Veizer et al., 1999; Zachos et al., 2001). The macroweather-climate transition is not in phase between the different records because the left ones (industrial side) are influenced by anthropogenic climate change, while the right data is pre-industrial natural variability. As elsewhere in this
 10 paper, the fluctuations were multiplied by the canonical calibration constant of 2 so that when the slopes are positive, the fluctuations are close to difference fluctuations. The various scaling regimes are indicated at the bottom. Adapted from (Lovejoy, 2017).




Figure 3a: Zooming out of the Holocene dust fluxes by octaves, by doubling the depth resolution from 1 cm (upper left) to

5 11m (lower right) resolution. Starting at the left and moving to the right and from top to bottom (see the ellipses on the first three in the sequence) we zoom out by factors of 2 in depth maintaining exactly 290 data points (effectively nondimensionalizing the depth; the small number of missing data points were not interpolated so that the final resolution is not exactly 2^{10} cm = 10.24m). The temporal resolution is not exactly doubled due to the squashing of the ice column, the total duration (in years) of each section is indicated in each plot, the average temporal resolution of plots are: 0.24, 0.48, 0.98, 2.02, 4.32, 10.1 24.5, 54.1, 184, 434, 2710 yr.

10 In order to fit all the curves on the same vertical scale, the dust fluxes were normalized by their mean over each segment. The means (in mg/m²/yr) are: 0.44, 0.38, 0.30, 0.36, 0.35, 0.33, 0.34, 0.39, 2.48, 2.18, 2.41 i.e. the first 8 plots have nearly the same vertical scales whereas the last three are about 6 times larger range. This means that all the plots except the last three are at nearly constant normalization.




Figure 3b: Same as Fig. 3a but for the absolute changes between neighbouring values in dust flux normalized by the
5 corresponding mean over the segment (290 points). The horizontal lines indicate the Gaussian probability levels for $p = 1/290$
(representing the mean extreme for a 290 point segment, full line), as well as $p = 10^{-6}$ (lower dashed) and $p = 10^{-9}$ (upper dashed).




Figure 3c: Same as Fig. 3a but for the absolute changes between neighbouring values in the logarithms of dust flux normalized by the corresponding mean over the segment (290 points). The horizontal lines indicate the Gaussian probability levels for $p = 1/290$ (representing the mean extreme for a 290 point segment, full line), as well as $p = 10^{-5}$ (lower dashed) and $p = 10^{-8}$ (upper dashed, not the same as in fig. 3b).




Figure 4: Log-log plot of the Fourier spectrum of the (25yr) $^{-1}$ resolution dust concentration in frequency units of kyrs $^{-1}$ (red) and the same but of the logarithms of the flux (blue). Also shown is the average spectrum of the 5 year resolution data over the last 400 kyrs (green). For the latter, the periodograms of each the four most recent 100 kyr cycles were averaged, but the full spectral resolution (5yrs) $^{-1}$ was retained. The beta parameters are the exponents of the theoretical spectrum (see main text, the negative of the logarithmic slope) for the macroclimate (-2.5), climate (1.7), and macroweather (0.8) regimes. The spectra were analyzed using FFT with standard Hanning windows.




Figure 5: The Haar fluctuation analysis of the entire 800 kyr dust flux data set (thin lines). The dashed black and solid pink (top pair) represent RMS fluctuations for dimensional and non-dimensional time, respectively. The solid black and blue curves are the same but for the mean absolute ($q = 1$) fluctuations. The curves with non-dimensional time lags have nominal (average) resolutions of 25 years and the fluctuation statistics are averaged over the 8 cycles. The thick black line shows the Haar fluctuations for the most recent 400 kyr at 5 year resolution. Note that the peak in the curves occurs as expected at $\Delta t \approx 50$ kyr i.e. at about a half cycle; and the horizontal dashed line shows that at this scale – corresponding to the largest difference in phases – the change in the mean absolute dust flux is about ± 3 mg/m²/yr. Also shown (dashed vertical line) is the (average) time scale $\tau_c \approx 250$ yrs at which the transition from macroweather to climate occurs. Several reference lines (with the slopes/exponents indicated) are shown showing approximate scaling behaviours.




Figure 6a: The probability distribution $\text{Pr}(\Delta F > s)$ of random changes in dust flux (ΔF) exceeding a fixed threshold s ; in time at 25 year resolution (brown, 32,000 points), and in depth at 1cm resolution (black, 251,075 points corresponding to the last 5 400 kyrs). The frequency scales at the right give the number (N) of jumps in each of the series that exceeds the threshold s . The straight lines indicate power law probability tails with exponents q_D indicated. Also shown (parabolas) are the Gaussians with the same mean and standard deviations. In time, the maximum change in flux corresponds to about 28 standard deviations (i.e. to a Gauss probability $\approx 10^{-91}$), in depth, to 51 standard deviations (i.e. to $p \approx 10^{-455}$). On the right, we provide axes giving the actual number of flux increments that exceed s , brown for the fluctuations in time, black for those in depth.




Figure 6b: Same as 6a except for the increments of the log of the dust flux (brown is in time, 25 year resolution, black is in depth, 1 cm resolution), the curves are the closest fitting (log) Gaussians.




Figure 7: Top set: successive segments of theoretical 100 kyr-long glacial cycles using usual (dimensional) time (present to past: bottom to top, the segment number is at the far right) with the 12.5 kyr phases indicated by vertical dashed lines. The short red lines indicate the interglacial dust minima. Each glacial-interglacial cycle is shifted by 25 units in the vertical for clarity. The red markers in the upper plot get mapped to the first dashed blue line in the lower plot.

Bottom set: successive cycles using nondimensional time (interglacial to interglacial) and then shifted by one phase to better line up with the usual time segments (the left most phase of the bottom line of the lower plot is zeroed). The average (nominal) resolution is 25 years. The interglacial dust minima were taken as 128.5, 243.5, 336, 407.5, 490, 614, 700, 789 kyr B.P. and the data start at 373 yrs B.P. Each cycle is shifted by 25 units in the vertical for clarity. The data older than 789 kyr were not used in these nondimensional cycles.




Figure 8: Top set: Averaging over the 8 cycles at 25 year resolution, we get the above picture: the mean is brown and the 5 one standard deviation cycle to cycle variability is shown by the red. The dashed vertical lines give a further division into 8 x 12.5kyr segments, the 8 “phases” of the cycle.

Bottom set: the same but for the nondimensional time. The relative position of the interglacial minimum at the first dashed line is indicated.




Figure 9: The top row shows the intermittency function $G(\Delta t)$ (whose slope on the log-log plot is C_1) and the bottom row,
 5 the mean absolute Haar fluctuation $S_1(\Delta t)$ (whose slope on the log-log plot is H), the left column shows the result for each phase
 after averaging over the 8 cycles with the numbers next to each line indicate the phase number; the right hand column shows the
 result for each cycle after averaging over the phases. Whereas each cycle is fairly similar to every other cycle (the right column),
 each phase is quite different (the left column). We see the most significant difference is the fluctuation amplitude as a function of
 phase (lower left).




Figure 10: The transition scale τ_c estimated in two ways for each of the 8 phases and from two definitions of the phases. The first method (solid lines) used a bilinear fit to the (logarithm) of the Haar $q=1$ structure function (i.e. mean absolute fluctuation) as a function of log time lag Δt . To obtain robust results, a small Δt region with the slope -0.05 and a large Δt slope +0.25 was imposed with the transition point (τ_c) determined by regression. This was done for each segment and cycle. For each phase there were thus 8 transition scales, which were used to calculate the mean of the logarithm of τ_c and its standard deviation. Results are shown for dimensional (segments, blue) and nondimensional time (cycles, black).

The second method used to estimate τ_c was graphical and relied on a somewhat subjective fitting of scaling regimes and transitions, but without imposing small and large Δt slopes (exponents H). The results are shown in dashed lines, they are quite similar although we can note some differences for the first phase (dimensional, blue) and the middle phases (nondimensional, black). There is also considerable cycle to cycle spread that was quantified by the standard deviations. In order to avoid clutter, typical spreads are shown by the double headed black arrows. Dashed horizontal lines show the ensemble mean transition scale (about 250 years) as well as ensemble mean for phases 1 and 2 (around 2 kyrs), which stands out compared to the rest of the phases. The red arrow shows one standard deviation for the nondimensional first phases, while the X marks the value of the Holocene τ_c (7.9 kyr) just outside the 1-sigma limit.




Figure 11: Using nondimensional time, the amplitude of the Haar fluctuations are averaged over all the cycles. The curves from bottom to top are for time scales of $\Delta t = 50, 100, 200, 400, 800, 1600, 3500, 7000$ years, alternating solid and dashed (for clarity, only some of the Δt 's are marked). The cycle to cycle variability (the dispersion around each line) is about a factor of 2 (it is not shown to avoid clutter).




Fig.12: The fluctuation and intermittency exponents H and C_1 (top row) are estimated over the range 500 – 3000 years, as

5 a function phase with the standard deviations from the cycle to cycle variability (all using nondimensional time). The upper left (H) plot shows low drift in phases 1 and 2 but become driftier in the middle and older phases. The intermittency (C_1 , upper right) is moderate at the beginning and end of the cycles, and a little weaker in the middle. The lower left shows the amplitude of the fluctuations at 25 years determined by the standard deviation of the dust flux (units: $\text{mg/m}^2/\text{yr}$). We see that the flux has low amplitude fluctuations at the beginning and end of the cycles and 3–4 times higher amplitude fluctuations in the middle. The lower 10 right shows the probability exponent q_D estimated from the 25 year resolution data for each phase; the extreme 5% of the flux changes were used to determine the exponent in each phase; the cycle to cycle spread is indicated by the error bars (overall average over the phases: $q_D = 2.62 \pm 0.42$).



References

- Dakos, V., Carpenter, S. R., Brock, W. A., Ellison, A. M., Guttal, V., Ives, A. R., Kéfi, S., Livina, V., Seekell, D. A., van Nes, E. H. and Scheffer, M.: Methods for detecting early warnings of critical transitions in time series illustrated using simulated ecological data, *PLoS One*, 7(7), doi:10.1371/journal.pone.0041010, 2012.
- 5 Delmonte, B., Andersson, P. S., Hansson, M., Schöberg, H., Petit, J. R., Basile-Doelsch, I. and Maggi, V.: Aeolian dust in East Antarctica (EPICA-Dome C and Vostok): Provenance during glacial ages over the last 800 kyr, *Geophys. Res. Lett.*, 35(7), 2–7, doi:10.1029/2008GL033382, 2008.
- Ditlevsen, P. D., Svensmark, H. and Johnsen, S.: Contrasting atmospheric and climate dynamics of the last-glacial and Holocene periods, *Nature*, 379(6568), 810–812, doi:10.1038/379810a0, 1996.
- 10 Ganopolski, A. and Calov, R.: The role of orbital forcing, carbon dioxide and regolith in 100 kyr glacial cycles, *Clim. Past*, 7(4), 1415–1425, doi:10.5194/cp-7-1415-2011, 2011.
- Huybers, P. and Curry, W.: Links between annual, Milankovitch and continuum temperature variability, *Nature*, 441(7091), 329–332, doi:10.1038/nature04745, 2006.
- Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S., Leuenberger, M., Louergue, L., Lüthi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J. P., Stenni, B., Stocker, T. F., Tison, J. L., Werner, M. and Wolff, E. W.: Orbital and millennial Antarctic climate variability over the past 800,000 years., *Science* (80-.), 317(5839), 793–796, doi:10.1126/science.1141038, 2007.
- Kolmogorov, A. N.: A refinement of previous hypotheses concerning the local structure of turbulence in a viscous incompressible fluid at high Reynolds number, *J. Fluid Mech.*, 13(01), 82, doi:10.1017/S0022112062000518, 1962.
- 20 Lambert, F., Delmonte, B., Petit, J., Bigler, M., Kaufmann, P., Hutterli, M., Stocker, T., Ruth, U., Steffensen, J. and Maggi, V.: Dust-climate couplings over the past 800,000 years from the EPICA Dome C ice core., *Nature*, 452(7187), 616–619, doi:10.1038/nature06763, 2008.
- Lambert, F., Bigler, M., Steffensen, J., Hutterli, M. and Fischer, H.: Centennial mineral dust variability in high-resolution ice 25 core data from Dome C, Antarctica, *Clim. Past*, 8(2), 609–623, doi:10.5194/cp-8-609-2012, 2012.
- Lovejoy, S.: A voyage through scales, a missing quadrillion and why the climate is not what you expect, *Clim. Dyn.*, 44(11–12), 3187–3210, doi:10.1007/s00382-014-2324-0, 2015.
- Lovejoy, S.: How scaling fluctuation analysis transforms our view of the climate, *Past Glob. Chang. Mag.*, 25(3), 136–137, doi:10.22498/pages.25.3.136, 2017.
- 30 Lovejoy, S.: The spectra, intermittency and extremes of weather, macroweather and climate, *Nat. Sci. Reports*, in press, 2018.
- Lovejoy, S. and Schertzer, D.: Scale invariance in climatological temperatures and the local spectral plateau, *Ann. Geophys.*, 4(B), 401–410, 1986.
- Lovejoy, S. and Schertzer, D.: Haar wavelets, fluctuations and structure functions: convenient choices for geophysics, 35 Nonlinear Process. Geophys., 19(5), 513–527, doi:10.5194/npg-19-513-2012, 2012.
- Lovejoy, S. and Schertzer, D.: The Weather and Climate: Emergent Laws and Multifractal Cascades, Cambridge University Press, Cambridge., 2013.
- Lovejoy, S., Pinel, J. and Schertzer, D.: The global space–time cascade structure of precipitation: Satellites, gridded gauges and reanalyses, *Adv. Water Resour.*, 45, 37–50, doi:10.1016/J.ADVWATRES.2012.03.024, 2012.
- 40 Maher, B. a., Prospero, J. M., Mackie, D., Gaiero, D., Hesse, P. P. and Balkanski, Y.: Global connections between aeolian dust, climate and ocean biogeochemistry at the present day and at the last glacial maximum, *Earth-Science Rev.*, 99(1–2), 61–97, doi:10.1016/j.earscirev.2009.12.001, 2010.
- Mandelbrot, B. B.: Intermittent turbulence in self-similar cascades: divergence of high moments and dimension of the carrier, *J. Fluid Mech.*, 62(02), 331, doi:10.1017/S0022112074000711, 1974.
- 45 Nilsen, T., Rypdal, K. and Fredriksen, H.-B.: Are there multiple scaling regimes in Holocene temperature records?, *Earth Syst. Dyn.*, 7(2), 419–439, doi:10.5194/esd-7-419-2016, 2016.
- Rehfeld, K., Münch, T., Ho, S. L. and Laepple, T.: Global patterns of declining temperature variability from the Last Glacial Maximum to the Holocene, *Nature*, 554(7692), 356–359, doi:10.1038/nature25454, 2018.
- Ridgwell, A. J.: Implications of the glacial CO₂ “iron hypothesis” for Quaternary climate change, *Geochemistry Geophys.*



Geosystems, 4(9), 1–10, doi:10.1029/2003GC000563, 2003.

Schertzer, D. and Lovejoy, S.: Physical modeling and analysis of rain and clouds by anisotropic scaling multiplicative processes, J. Geophys. Res., 92(D8), 9693, doi:10.1029/JD092iD08p09693, 1987.

Shao, Z. G. and Ditlevsen, P. D.: Contrasting scaling properties of interglacial and glacial climates, Nat. Commun., 7, 1–8,
5 doi:10.1038/ncomms10951, 2016.

Veizer, J., Ala, D., Azmy, K., Bruckschen, P., Buhl, D., Bruhn, F., Carden, G. A. F., Diener, A., Ebneth, S., Godderis, Y.,
Jasper, T., Korte, C., Pawellek, F., Podlaha, O. G. and Strauss, H.: $^{87}\text{Sr}/^{86}\text{Sr}$, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ evolution of Phanerozoic
seawater, Chem. Geol., 161(1–3), 59–88, doi:10.1016/S0009-2541(99)00081-9, 1999.

Zachos, J., Pagani, M., Sloan, L., Thomas, E. and Billups, K.: Trends, rhythms, and aberrations in global climate 65 Ma to
10 present., Science, 292(5517), 686–93, doi:10.1126/science.1059412, 2001.