

Point-by-point reply to Short Comment #1:

1. Could you later provide a much better map of the study area than the present Fig. 1?

In the final version of the article, we have added a precise map of the Beaune vineyard, with indication of altitudes and relief (Fig. 2)




Figure 1: The vineyard of the city of Beaune

Point-by-point reply to Referee Comment #1

1. Improved introduction and clarification of notation

See our comment in the Reply

We have improve the introduction with the addition at the very beginning of the introduction of the sentences “Since the Middle Ages the opening day of the grape harvest is each year, in a given territory, the outcome of a collective decision. Either in pre-industrial history, when authorities set each year an official ban after which it was permitted for everybody to pick up the grapes, and in recent history, the opening day of grape harvest has always induced an important amount of documentary data.”

2. Consistency of determining the GHD

See our comment in the Reply

We have added this table in the supplementary material (Table S 1), to illustrate our methodology.

Table S 1 : Comparison between dates of the last meeting of the Notre-Dame of Beaune church chapter before the grape harvest and the official ban harvest set by the city council (1554-1620)

	1554	1555	1557	1558	1569	1574	1583	1613	1617	1619	1620
Ban dates set by the city council of Beaune	3 Sep	19 Sep	7 Sep	6 Sep	9 Sep	13 Sep	5 Sep	21 Sep	27 Sep	20 Sep	25 Sep
Dates given by the protocols of the church of Notre-Dame	23 Aug	13 Sep	3 Sep	2 Sep	2 Sep	15 Sep	1 Sep	18 Sep	20 Sep	6 Sep	16 Sep
Estimated dates	26 Aug	16 Sep	6 Sep	5 Sep	5 Sep	18 Sep	8 Sep	26 Sep	28 Sep	14 Sep	24 Sep

3. Harmonisation of time series

See our comment in the Reply

To illustrate the special behaviour of the Beaune GHD time series in comparison with other European GHD times series, we have improved the Table 1 in the final version of the article. The new Table 1 contains a comparison with the Aubonne series, and a new column "Difference of days between Period 1 and Period 2", which did not appear in the discussion paper.

Table 2: Comparison of mean GHD time series in Beaune (this article), Aubonne (Angot 1885), Salins (Angot 1885), Switzerland (Wetter et al. 2013) and Czech Lands (Mózny et al. 2016).

	Period 1: 1599-1717	Period 2: 1718-1875	1354-1717	1718-2018	Difference of days between period 1 and period 2
Mean Beaune GHD	21 Sep	28 Sep	20 Sep	27 Sep	+7
Mean Swiss GHD	20 Oct	15 Oct			-5
Mean Czech Lands GHD	14 Oct	14 Oct			0
Mean Salins GHD	11 Oct	09 Oct			-2
Mean Aubonne GHD	21 Oct	18 Oct			-3

Furthermore, we have partially rephrased paragraph 2.6, explaining the reasons why we have chosen to homogenised the Beaune GHD raw data adding 7 days to all dates prior to 1718.

l. 189-192: “Otherwise it would have mean April-to-August GHD temperature based reconstructions with maxima and minima c. 1°C warmer before 1718 than afterwards on the decadal scale, even though temperatures measurements available in France since 1658 do not provide any evidences of such a warming during the period 1658-1718.”

l. 202-205: “In the course of the 18th century, agronomists all underlined the fact that more mature grapes favour the process of fermentation and can prevent the acidification of wines. The works of Jean-Antoine Chaptal (Chaptal 1807) who gave his name to the process of *chaptalisation*, i.e. addition of sugar in grape must, illustrate this evolution.”

4. Statistical modelling

See our comment in the Reply

Section 3.2 has been largely rephrased accordingly to the reviewer comment. Additionally we have reworked the terminology of Model A and Model B that was found too complicated. (especially as the two denote something different in the forward and backward way). Therefore we have chosen a new and more intuitive terminology:

F_{linear} , $F_{\text{transformed}}$, $F_{\text{transformed.EKF400}}$, $F_{\text{transformed.CCC400}}$, B_{linear} , $B_{\text{weighted.simulations}}$

Here F stands for forward and B for backward, with ".CCC400" and ".EKF400" we denote the application of the calibrated model to these data sets. The rest (linear, transformed, weighted simulations) should be self-explanatory. Furthermore, we now give the explicit regression equation with the fitted coefficients for all models.

Point-by-point reply to Referee Comment #2

1. “It is particularly interesting that the new reconstruction does not corroborate the putative heatwave in 1540 CE [...] ; an aspect that should be discussed in a more balanced and nuanced way than currently done.”

See our comment in the reply.

We have given more information in the final version of the article on this especially important issue.

We have rephrased Section 4.5 with the following input:

“Subsequently conditions in two outstanding years - 1540 and 1556 - are considered in more detail. First of all, it is puzzling that the exceptional heat and drought in 1540 ranks only 19th in the statistics of Beaune GHD. Possible reasons to deemphasise this event based on tree ring evidence were brought forward by Büntgen et al. (2015). However, their arguments are thought to be questionable (Pfister et al. 2015). An interpretation of this paradox is attempted using vine phenological evidence available from vineyards around Biel-Bienne, Zürich and Schaffhausen (Switzerland) in 1540 and 1556 in comparison with the Beaune evidence. Source references are provided in the Euro-Climhist data-platform (<https://echdb.unibe.ch/>). Elevation matters for the comparison: the three Swiss vine growing areas are located at altitudes of about 430 metres asl, i.e. about 130 to 200 metres higher than those of Beaune region vineyards. Vine cultivars (CV), i.e. varieties, need to be considered besides altitudes. In Biel-Bienne CV Chasselas used to be grown while in Zurich and Schaffhausen CV Räschling was cultivated which survives in a few vineyards until today. The maturity of the Chasselas and Räschling cultivars is 10 to 14 days earlier than that of Pinot noir grown in the Beaune area (Altwegg 2018). Table 6 lists the main phenological grapevine stages - ‘end of flowering’, ‘beginning of veraison’ (colour change and softening of berries) and harvest dates for the years 1540, 1556, 2003 and 2018. Gladstones (2011) refers to the widely observed fact that “the date of flowering, can usually predict quite closely the dates of veraison (colour change and softening of berries) and maturity to follow [...]. The later phenological intervals show little response to temperature and tend to be constant from year to year”. His assessment is confirmed by Chuine et al. (2004) and Jüstrich (2018). In 1540, however, the interval between véraison and harvest took 67 days in the Zürich area, i.e. 27 days longer than usual, while this interval was somewhat shorter in 1556. In both years the grapes were harvested some time after full ripening. For 1540 this assumption is confirmed by the sources. “Vine-growers in Schaffhausen (Switzerland) were “long waiting for rain to begin the harvest”, as chronicler Oswald Huber relates. He writes, however, they “finally tackled the work nevertheless, because the plants withered.” Likewise, vine-growers at the shores of Lake Constance and in the Upper Alsace interrupted the vintage after picking a few juicy grapes, because the remaining ones were quasi dried

out. The vintage was then resumed after a two-day spell of rain. At harvest time grapes in many vineyards had become raisins. They yielded a sweet sherry-like wine which made people more rapidly drunk than usual (Wetter and Pfister 2013). It needs to be stressed, that meteorological conditions were almost matchless in 1540. The evidence for this mega-drought taken from documentary data offers a broad spectrum of evidence on past weather and climate and their societal impacts. It is described in a sample of 312 first-hand documentary weather reports originating from continental Europe (Wetter and Pfister, 2013). The hot and dry period in 1540 lasted from April to the end of the year. The heat-wave peaked at the end of a 46 day long rainless period between 23th June and 7th August during which many forests and settlements in a large area from the Ardennes to Poland went up in flames (Pfister 2018). Maximum temperatures from late July may have exceeded 40°C (Orth et al. 2016). Vagrants and homeless people were hunted down for raising town and forest fires (Pfister 2017). The heatwave of 1540 undoubtedly struck the region of Burgundy. The church of Notre-Dame of Beaune organised 8 processions to call for rains from the beginning of May to the end of August (Labbé, Gaveau 2011). In the near city of Besançon (c. 100 km eastwards) a chronicler wrote that warm temperatures lasted from April to November and that the heatwave was hardly bearable by humans during summer (Wetter et al. 2014). It cannot be excluded, that even the date of *véraison* was also somewhat delayed due to drought stress. Research in grapevine biology established that under conditions of extreme heat and drought the development of grapes is slowed down or stopped (Keller 2016). When occurring before *véraison*, extreme hydric stress alters grape quality and the onset of ripening, possibly because it induces leaf defoliation and therefore carbon assimilation limitation (Basile et al., 2012; Girona et al., 2009; Ollat and Gaudillere, 1998). In fact, this phenomenon was observed during the temperature peak in the hot summer 1947 in Schaffhausen (Amtsblatt 1947). In 1556 grapes in Beaune were harvested on 16th August. In central France, the 1556 heat and drought began in mid-April, i.e. about 45 days later than in 1540, following a wet winter. The heat-wave to mid-June spurred vegetation growth. In the Loir-et-Cher region, a variety of red Pinot noir cultivar named Auvernat was already blossoming around the 25 of April (Nouel, 1878, 235). Not a drop of rain fell during this period. On 14th June, it was pouring down for three to four hours which visibly refreshed the vegetation. Before 10th July the first grapes were ripe. In July the ground became so hot that it burnt people's feet walking barefoot. Like in 1540 the heat and drought peaked in early August (Bourquelot, 1857, 30-31; Hiver 1867, 81 and 90; Nouel 1878, 235-237). Following the deliberation protocols of the chapter of Notre-Dame of Beaune, seven processions for obtaining rain were held in the city from 15th June to 15th August (Arch. Dep. Côte d'Or, G 2499).

The 1556 harvest of Chasselas grapes in Biel-Bienne was estimated to have occurred between 25th and 30th August i.e. at about the same time as that of Pinot noir in Beaune considering the chronicler's remark that the abundant harvest already ended on 10th September (Gregorian Style). Like in the Beaune area, the preceding winter 1556 had been very wet considering the daily weather observations by Wolfgang Haller in Zürich. Disregarding June, which was completely rainless, the

spring-summer period included 7 precipitation days in April, 3 in May and 5 in July (Haller in Pfister, Rohr <https://echdb.unibe.ch/>, 15th Oct 2018). A hot and almost rainless period began on 29th July and lasted to 11th September, i.e. more than a month later than in 1540.

The heat-wave in summer 2003 was reassessed by Pfister (2018). The water deficit was not sufficiently strong to "block" grape ripening through extremely limited photosynthesis. Obviously, ripening made its course quickly. Additionally, wine growers harvested pretty soon to maintain sufficient acidity in wines mostly (and in some cases avoid excessively high alcohol content in wines).

Since the early 21st century, higher temperatures combined with increase control of grape sanitary status (grey mould disease mostly) makes ripening duration (i.e. the lag during veraison and harvest) more winemaker dependent. The winemaker choices depend on both cultivar and style of wine. For instance, the number of days between veraison and harvest for CV Cabernet-Sauvignon has nearly been doubled in a famous Château in the appellation Margaux near Bordeaux (Van Leeuwen and Destrac-Irvine, 2017)

In sum, the decline of 13 days in the average date of GHD since 1988 went along with large increase in the number of extreme spring to summer seasons. These include involve situations such in 1540 in which both human and ecological systems behave non-linearly outside the normal range of biological and probability laws. Documentary data may be helpful to describe such conditions in the necessary detail. “

2. “[...] the article would benefit from an assessment of the persistence (i.e. autocorrelation structure) in Beaune’s new GHD-based April-July temperature reconstruction compared to the possibly inflated long-term persistence in tree ring width records”

See our comment in the Reply

We entirely agree with the proposition made by the reviewer. Therefore, we have added in Section 4.3 the following paragraph that develops some considerations on this issue:

“The autocorrelation structure of the Beaune GHD-based temperature reconstruction (black in figure S2; regression model F and B lead to equal results) is very similar to the tree-ring reconstruction from the Pyrenees (blue) but has clearly less autocorrelation than the tree-ring reconstruction from the Lötschental, Switzerland (red). The Lötschental time series experiences much more low frequency variability than the Beaune and Pyrenees records. However, it is hard to argue that one of them should be the correct one. All have uncertainties with regard to their low frequency behaviour. In the tree-ring series many studies discuss these issues that stem from age detrending, temporally inconsistent tree-age distributions, etc. On the other hand documentary data for GHD are also not free from issues concerning low frequency variability. In the case of vine there may be adaptation, breeding, gene (de)activation processes over the decades that may dampen low frequency variability. There may also be changes in taste altering harvest dates in both directions.

Another point worth mentioning in comparison with tree-ring reconstructions is that the Beaune temperature reconstruction rather underestimates interannual variability ($SD=0.58$ K, period 1659-2007) compared to the Paris observations ($SD=0.87$ K) whereas both tree-ring reconstructions overestimate it (both $SD=1.10$ K).” This paragraph is accompanied with a new figure provided as supplementary material (Figure S 2; see below).




Figure S 2 : Autocorrelation structure of Beaune GHD-based temperature reconstruction (in black), Löchtental tree-ring-based temperature reconstruction (in red) and Pyrénées tree ring-based temperature reconstruction (in blue)

3. “I kindly ask the authors to consider the spatially explicit European summer temperature reconstruction by Luterbacher et al. (2016) for comparison

See our comment in the reply: “In the article we have compared the Beaune record with the two closest proxy records, which actually are the basis for the Luterbacher reconstruction, too (i.e. the Alps and Pyrenees, east and south-west of Beaune respectively). Luterbacher et al. do not have any additional proxy information to north or west. Hence, the interpolation to a coarse grid, even with a sophisticated method, could hardly add significant information at our site.”

Accordingly we did not change anything on this point.

The longest homogeneous series of grape harvest dates, Beaune 1354-2018, and its significance for the understanding of past and present climate

Thomas Labbé,^{1,2} Christian Pfister,³ Stefan Brönnimann,^{3,4} Daniel Rousseau,⁵ Jörg Franke,^{3,4} Benjamin Bois^{6,7}

¹Leibniz Institute for the History and Culture of Eastern Europe (GWZO), University of Leipzig, Specks Hof, Reichsstraße 4-6, 04109 Leipzig, Germany

²Maison des Sciences de l'Homme de Dijon, USR 3516 CNRS, University of Burgundy, 6 Esplanade Erasme, BP 26611, 21066 Dijon cedex, France

³Oeschger Centre for Climate Change Research, University of Bern, Hochschulstrasse 4, 3012 Bern, Switzerland

⁴Institute of Geography, University of Bern, Hallerstrasse 12, 3012 Bern, Switzerland

⁵Conseil Supérieur de la Météorologie, 73 avenue de Paris, 94160 Saint-Mandé, France

⁶Biogéosciences UMR 6282 CNRS, University of Burgundy, 6 boulevard Gabriel, 21000 Dijon, France

⁷Institut Universitaire de la Vigne et du Vin, University of Burgundy, 1 rue Claude Ladrey, 21000 Dijon, France

Correspondence to: thomas.labbe@u-bourgogne.fr

Abstract. Records of grape harvest dates (GHD) are the oldest and the longest continuous phenological data in Europe. However, many available series including the well-known (Dijon) Burgundy series are error prone, because scholars so far uncritically drew the data from nineteenth century publications instead of going back to the archives. The GHD from the famous vine region of Beaune (Burgundy) were entirely drawn from the archives, critically cross checked with narrative evidence. In order to reconstruct temperature, the series was calibrated against the long Paris temperature series comprising the 360 years from 1659 to 2018. The 664-year-long Beaune series from 1354 to 2018 is also significantly correlated with tree-ring and documentary proxy evidence as well as with the Central European temperature series (from 1500). The series is clearly subdivided into two parts. From 1354 to 1987 grapes were on average picked from 28 September on, whereby during the last 31 year long period of rapid warming from 1988 to 2018 harvests began 13 days earlier. **Early harvest dates are shown to be accompanied by high pressure over western-central Europe and atmospheric blocking over Denmark.** The 33 extremely early harvests comprising the 5% percentile bracket of GHD are unevenly distributed over time. 21 of them occurred between 1393 and 1719, whereby this is the case for just five years between 1720 and 2002. Since the hot summer 2003, 8 out of 16 spring-summer periods were outstanding according to the statistic of the last 664 years, no less than 5 among them within the last 8 years. **In the Paris temperature measurements since 1659, April-to-July temperature reached**

the highest value ever in 2018. In sum, the 664-year-long Beaune GHD series demonstrates that outstanding hot and dry years in the past were outliers, whereby they became the norm since the transition to rapid warming in 1988.

35 1. Introduction

Since the Middle Ages the opening day of the grape harvest is each year, in a given territory, the outcome of a collective decision. Either in pre-industrial history, when authorities set an official ban after which it was permitted for everybody to pick up the grapes, and in recent history, the opening day of grape harvest has always produced an important amount of documentary data. Records of grape harvest dates (GHD) provide the longest continuous series of phenological data in Europe and have been repeatedly used for estimating spring-summer temperatures (Chuine et al. 2004; Guiot et al. 2005; Menzel 2005; Le Roy Ladurie et al. 2006; Krieger et al. 2011; Garcia de Cortázar-Atauri et al. 2010; Daux et al. 2012; Rousseau 2015). In the literature, the GHD series (1385-1905) from the town of Dijon situated in Burgundy (France, see Fig. 1) is the longest available series. Going back to the late 14th century, it constitutes the backbone of all the reconstructions of “Burgundian” GHD series. The most widely quoted article published by Chuine et al. (2004) in “Nature” provides a “Burgundian” GHD dataset (1370-2003) in which Dijon GHD constitute the reference series.

However, two biases affect the reliability of this dataset. First, scholars have until now uncritically drawn on the data from nineteenth century publications. The original data of the Dijon series have been recently revisited directly in the local archives (Labbé and Gaveau 2011) and this reassessment makes obvious that the formerly published “Dijon series” is thoroughly unreliable, due to compilation errors. Secondly, the available Burgundian GHD dataset is not homogeneous. Due to a lack of information concerning Dijon where the vineyard has disappeared from the 19th century because of the urbanisation, the series is complemented for the 18-20th centuries by a mix of data taken from different locations disseminated throughout the vineyard region of Burgundy.

An entirely unedited series based on manuscript material discovered in the archives of the town of Beaune situated 45 km south of Dijon has recently been collected for the period 1371-2010 (Labbé and Gaveau 2013a). Unlike Dijon, Beaune is still surrounded with vineyards situated on altitudes between 220 and 300 metres (Fig. 2). Since the end of the Middle Ages the territory of the city is dominated by the culture of grapes. The region produces labelled wines certified by the “Appellation d’Origine Contrôlée” system. In this paper, we have extended the series back to 1354 and updated it to 2018. It is validated using the long Paris temperature series that goes back to 1658 (Rousseau 2009, Rousseau 2013, updated to 2018) and used to assess April-to-July temperatures from 1354 to 2018. The phenological data are available on the “Euro-Climhist” data-platform: <http://www.euroclimhist.unibe.ch/en/> while reconstructed temperatures are included in the supplementaries.

The study is organized as follows. The first section reviews the former “Burgundian” GHD series highlighting its inadequacies. The second section presents the generation of the new Beaune series. Section three outlines the methodological steps used for reconstructing April-to-July temperatures from 1354 to 2018 and for checking the reliability of the data. In section four the Beaune series is presented and compared to other GHD series, other documentary data series as well as to tree-ring evidence. This section also provides the reconstruction of April-to-July temperatures from 1354 to 2018 and a detailed analysis of two extremely early years. In the last section we summarize the main conclusions that can be drawn from the study, in particular in view of comparing isolated outliers in the remote past with the increasing frequency of such events in the last three decades of rapid warming.

2. Data

2.1 The inadequacies of the “Dijon/Burgundian” GHD series

Etienne Noirot, land surveyor in Dijon, is one of the first scientists who showed some interest in long GHD time-series. In 1836, he gathered a Dijon GHD series from 1385 to his time, mainly to demonstrate that climate did not change significantly since five hundred years (Noirot 1836). In the course of the cold fluctuation leading to the glacier maximum around 1850 (Nussbaumer 2018) GDH’s became important for scientists as indicators of past climate (Labbé and Gaveau 2013b). In this context Jules Lavalle (1855) re-published the same dataset for the period 1366-1842. His series forms the backbone of all further publications using the Dijon evidence. In the early 1880’s Alfred Angot, Director of the Paris Meteorological Research Office, instructed the meteorological commissions of the departments to extract grape harvest dates from documentary sources and assembled a compilation of 606 GHD series from France, Switzerland and Germany. This publication was a milestone in the field. The Dijon series, continued up to 1879, constitutes the longest series of his data compilation (Angot 1885). The German geographer Eduard Brückner, son of a historian, attempted an analysis of GHD’s based on Angot’s data in his 1890 study on “Climatic Change since 1700” (Stehr and von Storch 2000). The French historian Emmanuel Le Roy Ladurie valued the Angot data for his synthesis on climate changes since AD 1000 (Le Roy Ladurie 1971).

Subsequently, three main sources of inhomogeneity have to be addressed concerning available Burgundian GHD.

2.2 Flawed data

First, the “Dijon series” is riddled with printing, typing and copying errors. The investigation in the original archives of the City Council in Dijon shows 132 differences with the Angot series for the period 1385 to 1879 (Labbé and Gaveau, 2011). The mismatches reach 5-10 days for 17 years, 10-20 days in 9 occasions, and more than 20 days in 1448, 1522, 1523, 1540,

1659, 1660 and 1842. A serious mismatch concerns the year 1540. According to the supplementary material on internet the GHD in this year is 4 October (day-of-year (DOY) 278) (Chuine et al., 2004) whereas the correct value found in the archive is 3 September (DOY 247). This flaw is the main reason why the outstanding extreme spring-summer temperature of this year (Wetter and Pfister 2013, Wetter et al. 2014) was overlooked in the “Nature” article by Chuine et al. (2004). In particular, Angot’s series is especially not reliable prior to 1420, in the first half of the sixteenth century, around 1840, and for several isolated years (Fig. 3).

2.3 Lack of homogeneity

The vineyards around Dijon were built over from the early nineteenth century. In 1906 the City Council did not set official ban date anymore. For the nineteenth and twentieth century the Dijon series was thus complemented with data from the southern part of the Burgundian vine growing area without, however, taking into account the resulting differences in mean grape ripening. It needs to be known that Dijon is situated at the northernmost point of the Burgundian wine region (Fig. 1) which involves a delay in the mean date of grape harvest compared to other locations along the latitudinal (North-South) orientation of the Burgundian wine area. Grapes in Dijon between 1600 and 1800 were picked on average 5 days after those in the town of Beaune (Fig. 4). In fact, the “Burgundian series” used by Chuine et al. (2004), indiscriminately combines the pre-1800 Dijon series with the 19th and 20th century evidence from the southern part of the Burgundian vine area.

2.4 Assembling a 664 year long GHD series for Beaune

To correct for the above mentioned inadequacies we constructed a new 664-year long almost homogenous and nearly uninterrupted series of GHD focussing the archival study on the Beaune vine growing area (available on the “Euro-Climhist” data-plateform, <https://echdb.unibe.ch/>). Thereby we used different kind of archival information.

Wage payment data (1354-1506): Late medieval accounts can be used as a source for day-specific information about the actual start and end of agricultural work (Brázdil and Kotyza 1999, Wetter and Pfister 2011, Pribyl et al. 2012, Wetter and Pfister 2013).

In the case of Beaune, data on daily wage payments made to day-labourers for picking grapes are available from 1354 to 1506. The oldest accounts were kept for a c.18 hectares domain owned by the dukes of Burgundy. The start dates of the harvest (1354-1426) in these estates have already been published (Guerreau 1995). Nevertheless, the most detailed and numerous series of accounts refers to the c.10 hectares domain of the church chapter of Notre-Dame in Beaune, whereby we could find almost without any lacunae unedited GHD from 1371 to 1506. The parcels documented in these accounts are continuously planted with vines since the late Middle Ages. Thus, these GHD are related to the maturity of grapes in domains such as *Corton Clos-du-Roi*, *Beaune-Sanvignes*, *Beaune-Tuvillains*, *Beaune-Bressandes* among others, which

produce today first class red wines designated as *Grand cru*, *Premier cru* or *Beaune-village*. The “Beaune” wine produced in these domains had already a reputation for quality since the 13th century (Dion 1959). The chapter of Notre-Dame sold for example barrels of wine to the merchants of the king of France and to other key persons. It is however not known which grape varieties were grown in these estates. Fine red and “clairret” (almost rosé) wines massively produced during the Middle Ages, were made with varieties of Pinot noir, whereby Gamay was rated second class. In 1395 the duke of Burgundy even prohibited the cultivation of Gamay vines around the cities of Dijon, Beaune and Chalon-sur-Saône. But in reality his direction never became effective (Dion 1959). In any case, the documentation on grape harvests never refers to the varieties cultivated in the domains. Nonetheless, these data are very reliable and even refine in which parcel the grapes were picked on a specific day and how many male and female labourers were at work.

Meetings of the Notre Dame Church chapter (1507-1699): In 1506 both aforementioned estates were leased out to tenants, so that precise information on their cultivation are not kept anymore in the accounting documentation. At the same time, the deliberations of the city council containing the setting of the vintage ban are fragmentary prior to 1700. The books of deliberations of the Notre-Dame Church chapter of Beaune offer an acceptable substitute. This documentation provides the date of the last meeting before the vacation of the chapter who had to organize the harvest. As church chapter meetings took regularly place twice a week, three days were added to the date of the last meeting to assess the harvest date. Since 1583, the books of deliberation indicate at what time the members of the chapter had to organise the food supply needed to feed the day labourers during the harvest. With regard to getting perishable food such as cheese and meat, it was assumed that the harvest began eight days later. *The accuracy of these estimations, though we must admit some uncertainties, is confirmed with the comparison of fragmentary preserved official ban dates set by the city of Beaune for some few years within this period, as summarized in table S1.*

Deliberations of the city council (1700 to 1965): From 1700 onwards the books of deliberation of the city council are continuously preserved and the opening date of the harvest can be easily drawn. However, setting the vintage ban is a social outcome and does not depend only of the observation of the full maturity of grapes. It resulted from a consensus between vine-growers and the local town administration. In practice, experts inspected the maturity of the grapes and proposed a ban date to the city council. The city council then took a decision, considering also the availability of day labourers coming from outside for harvesting as well as eventual military threats and plague outbreaks, as Garnier and co-authors (2011) have shown for Besançon (eastern France). In the case of Beaune, the main bias that potentially affects the date of setting the ban on the long run is nonetheless different. After the French Revolution actually, the re-organization of the territory into administrative districts called “arrondissements” induced a change in the decision-making process. The dates of the ban now resulted from a consensus among the mayors of all villages contained in the Beaune “arrondissements”. Though Beaune continues to play the major role in the decision process, the lifting of the harvest ban subsequently concerned a larger area

than before extending more to the south. Furthermore, the outbreak of the Phylloxera disease after 1870 induced a new organization of the vine sector in Burgundy in which local institutions further lost weight in the setting-up of the harvest on their territory. From the second half of the 20th century up to 2007 prefects promulgated a uniform harvest ban which did not any more consider local conditions. Moreover, to avoid dealing with requests for harvesting prior to the official date the prefectural ban tended to be artificially early, which affected the value for 2003 in the “Dijon series” (Chuine et al. 2004). Therefore, another kind of information was used after 1965.

Newspaper reports (1966 to 2018): From 1966 articles in the local newspaper “Le Bien Public” are used. Every year an article reports the date when the vine-growers start the harvest in the town vineyard. These dates are closer to the reality than the official bans set by prefectural decrees. Between 1980 and 2007 the Pearson correlation coefficient r between Beaune “newspaper” GHD and the mean April-to-July (AMJJ) temperatures of Paris is stronger ($r = -0.833$) than with the official administrative GHD ($r = -0.789$). In 2007, the official opening of GHD was set on 13th of August whereas wine-growers actually began the harvest in the territory of Beaune on the 1st of September according to the newspapers.

2.5 Interpolations

Despite an exhaustive investigation in the local archives, 61 dates for Beaune are still missing before 1645. To fill in these lacunae, when possible we used the evidence of the corrected Dijon series (Labbé and Gaveau 2011), taking into account the mean difference of days between the two raw series.

The dates from 1358 to 1364 have been taken from the Torino GHD series (Rotelli 1973), taking into account the mean differences of days between the two series.

Certain dates are affected by regional political and military biases and need to be interpolated as well with an extra-regional series, e.g. the corrected Swiss series (Wetter and Pfister 2013). The political situation has been particularly critical in the region of Beaune and Dijon in the context of the Thirty Years war (1618-1648). As in the nearby city of Besançon documentary records are sketchy in the second quarter of the 17th century because troop movements often prevented wine-growers to properly organize the harvest (Garnier et al. 2011). In Beaune, the archives do not provide any data from 1631 to 1638 which constitutes the longest undocumented period of the series. We interpolated these data from the Dijon series, but the dates for 1636 (4 September) and 1637 (3 September) still turned out to be artificially early in comparison with the series from the Swiss Mittelland, respectively 5 October and 1 October (Wetter and Pfister 2013). In 1636, the city council deliberations of Beaune inform us that the region was actually threatened by both enemy troops and by an outbreak of plague, which certainly disorganised the harvest process. In 1636 and in 1637, we have then interpolated the dates with the Swiss series.

2.6 Homogenisation

A comparison of Beaune GHD times series with the GHD of nearby regions of Switzerland (Wetter and Pfister 2013), Czech Lands (Mózný et al. 2016) and of Salins (Angot 1885), makes obvious that the Beaune GHD are artificially early before c. 1718 (Fig. 5). In the period 1599-1875 for which we can compare the three time series without lacunae, the stability of the mean GHD is stronger in Switzerland, in the Czech Lands, in Salins and in Aubonne (Jura, France), whereas in Beaune the average GHD occurred 7 days earlier before 1718 (21 September) than after this date (28 September) (Table 1).

In the perspective of reconstructing past spring-summer temperatures, this bias must be taken into account. Otherwise it would have mean April-to-August GHD temperature based reconstructions with maxima and minima c. 1°C warmer before 1718 than afterwards on the decadal scale, even though temperatures measurements available in France since 1658 do not provide any evidences of such a warming during the period 1658-1718. To homogenise the Beaune GHD series we have then added 7 days to raw data prior to 1718. The Pearson correlation with April-to-July mean temperature in Paris for the period 1658-2018 is stronger with the homogenised Beaune GHD time series ($r = 0.76$) than with the non-homogenised one ($r = 0.74$).

Anthropogenic changes in wine-making are the most likely explanation. This break is actually synchronous with a very important change in Burgundian practices of grape cultivation. The early 18th century is a turning point in wine history. The production and the commercialization of wines shifted to a new model distinguishing between ordinary and fine wines and focussing upon more coloured and longer keeping red wines (Dion 1959, Lachiver 1988). In Burgundy, the turnaround is described in a treatise written in 1728 by Claude Arnoux who referred to the emerging distinction between short- and long-keeping wines (Arnoux 1728). Unlike the production of the common “clairet” wines produced in premodern Burgundy, the manufacturing of stronger wines called for harvesting more mature grapes (Lachiver 1988). In the course of the 18th century, agronomists all underlined the fact that more mature grapes favour the process of fermentation and can prevent the acidification of wines. The works of Jean-Antoine Chaptal (Chaptal 1807) who gave his name to the process of *chaptalisation*, i.e. addition of sugar in grape must, illustrate this evolution. A similar retardation of mean harvest dates is also known from the region of Montpellier and Beziers in the same period (Blanchemanche 2009).

Our homogenisation approach is based on arguments of viticulture history. Statistical homogenisation might reveal further inhomogeneities. Based on simple visual tests (Craddock, 1979) we estimate a remaining error of 7 days or less.

3. Methodology

The new Beaune GHD series was first compared with other grape harvest series as well as with other climate-related proxy time series to test the robustness of the series. Then it was used to reconstruct April-to-July temperature back to 1354.

Furthermore, years with extremely early GHD in Beaune were analysed climatologically, with a special emphasis on atmospheric blocking.

3.1. Comparison with other time series

210 The quality of the improved series was first of all tested involving the long Swiss GHD series from 1444 to 2012 (Wetter and Pfister 2013) and the long series of the Czech Lands (from 1499 to 2015) (Možny et al. 2016). Additionally we have investigated the similarity between the grape harvest dates and tree-ring based temperature reconstructions. Two tree-ring based temperature reconstructions are chosen from the N-TREND data set (Wilson et al. 2016). This is a global data set, containing the best tree-ring based temperature reconstructions, selected by experts in the field. The spatially closest reconstructions in N-TREND are from the Spanish Pyrenees (Dorado Liñán et al. 2012) and from the Swiss Alps (Büntgen et al. 2006).

215 The first part of the Beaune series (1354 to 1431) was compared with estimated April-to-July temperatures in Norfolk (South-East England) obtained from the first-time dates of wages paid to grain harvest workers (Pribyl 2017). The second part of the series was compared with the detailed GHD obtained for the period 1420 to 1537 in Metz (France) (Litzenburger 220 2015). The last part was correlated with the estimated April-to-July temperatures in Central Europe from 1500 to 1759 that are based on Pfister Indices from Germany, the Czech Lands and Switzerland (Dobrovolný et al. 2010). Likewise, the Beaune GHD were compared with the series for Switzerland and the Czech Lands (Fig. 7).

3.2. Statistical models

225 We compared the GHD series not only with indirect climate-observations, but also directly with temperature. Both forward (i.e. reconstruction of GHD from temperatures, denoted F) and backward (i.e. reconstruction of temperatures from GHD, denoted B) reconstruction models were used. Correlating GHD month-by-month with Paris mean temperature from 1659 to 2018 (Rousseau 2009, updated to 2018) we found statistically significant correlations between GHD and temperature for all months from March to September. Enologically, using August and September is questionable, in line with the results of stepwise regression analysis attempted by Legrand (1979), Pfister (1984) and Guerreau (1995), all the more as in Beaune 230 harvest sometimes starts already in late August. Therefore we excluded August and September.

We modelled harvest dates in Beaune from Paris monthly temperature (i.e., the forward approach), starting in 1659, using two multiple linear regression models relating GHD to temperature in March, April, May, June and July, and fitted with ordinary least squares. Model F_{linear} is linear while model $F_{\text{transformed}}$ used transformed variables. Since the relation between grape growth and temperature is assumed to be non-linear, all temperatures were logit transformed in that model such that

235 curves flatten out for very low and very high temperatures and have the steepest slope at 18 °C (the same transformation was used for all calendar months):

$$T' = \frac{1}{1 + e^{\frac{-(T-18^{\circ}\text{C})}{3^{\circ}\text{C}}}} \quad (1)$$

Further, the harvest date was log-transformed such that a difference in harvest date of one day obtains more weight at the beginning of September than at the end of October:

240 $GHD' = \ln(GHD - 150)$ (2)

where \ln denotes the natural logarithm. Both models were calibrated in the period 1659-1850, while the period 1851-2018 was used for evaluation.

The calibrated model F_{linear} reads

$$GHD = 396.75 [\text{days}] - 0.624 T_{\text{Mar}} [^{\circ}\text{C}] - 1.259 T_{\text{Apr}} [^{\circ}\text{C}] - 2.200 T_{\text{May}} [^{\circ}\text{C}] - 2.773 T_{\text{Jun}} [^{\circ}\text{C}] - 1.655 T_{\text{Jul}} [^{\circ}\text{C}] \quad (3)$$

245

and the calibrated model $F_{\text{transformed}}$ reads

$$GHD' = 5.145 - 1.026 T_{\text{Mar}}' - 0.500 T_{\text{Apr}}' - 0.319 T_{\text{May}}' - 0.289 T_{\text{Jun}}' - 0.178 T_{\text{Jul}}' \quad (4)$$

For analysing atmospheric circulation conducive to early harvest dates, climate simulations can be used if they are able to reproduce harvest dates. We therefore applied model $F_{\text{transformed}}$ (without recalibration) to the global monthly ensemble climate reconstructions EKF400 back to 1603 (termed $F_{\text{transformed.EKF400}}$), which is based on assimilating instrumental data, documentary data and tree ring proxies into an ensemble of 30 global climate model simulations (Franke et al. 2017) as well as to the underlying model simulations CCC400 (Bhend et al. 2012), termed $F_{\text{transformed.CCC400}}$. The closest grid point to Beaune was extracted, and the data were debiased according to the calibration period mean value of each calendar month.

250

April-to-July mean temperatures were reconstructed from GHD (i.e., the backward approach) using two models, again calibrated in the period 1659-1850 and evaluated in 1851-2018. Model B_{linear} is a simple linear regression, yielding the relation:

255

$$T_{\text{AMJJ}} = 31.59 [^{\circ}\text{C}] - 0.061 [^{\circ}\text{C} / \text{day}] \cdot GHD [\text{days}] \quad (5)$$

Additionally we defined a model **B_{weighted.simulations}** starting from the simulated GHD in the CCC400 simulations (**F_{transformed.CCC400}**). We simulated all **prephylloxera** years (1603-1850; before the phylloxera outbreak that destroyed European cultivars and the subsequent reconstruction of the vineyard with American rootstocks) in the 30 member ensemble after debiasing, yielding 7440 years. We assume that any of these years may serve as an analog provided its GHD is close to the observed one. The reconstruction $x_{rec,j}$ for year j is then:

$$x_{rec,j} = \frac{\sum_i w_{i,j} \cdot x_i}{\sum_i w_{i,j}} \quad (6)$$

where i refers to the 7440 model years, x_i is the variable to be reconstructed (here Apr-Jul mean temperature, but the same equation could be used for other variables or other definitions of the warm season) in the model simulation for year i and $w_{i,j}$ is the weight of model year i , taken as the density of a normal distribution $N(GHD_j, r_{EKF}^2)$ at the location of the modelled GHD_i . GHD_j is the observed GHD and r_{EKF} is the standard deviation of EKF400 residuals (see above).

Note that this procedure is equivalent to a Bayesian approach, where the modelled Apr-Jul mean temperatures from all (equally likely) 7440 years serve as the prior for the distribution x_j . According to Bayes theorem

$$P(x_j | GHD_j) \propto f(GHD_j | x_j)P(x_j) \quad (7)$$

where $f(GHD_j | x_j)$ is the conditional density of GHD_j given x_j , which corresponds to the weights w_j . The expectation value is then equivalent to (6).

3.3. Atmospheric circulation blocking

For addressing anomalous atmospheric circulation causing early or late harvest dates, we analyse 500 hPa geopotential height (GPH) and atmospheric blocking. The blocks were defined based on 500 hPa GPH according to the algorithm of Tibaldi and Molteni (1990) (see also Tibaldi et al., 1994, Scherrer et al. 2006 for more details). For the period after 1850 we use version 2c of the „Twentieth Century Reanalysis“ (20CRv2c, Compo et al. 2011). It provides an ensemble of 56 realisations of an atmospheric reanalyses with 6-hourly time steps. For analysing blocking for early harvest dates prior to 1850, we used CCC400. The performance of blocking algorithms for 20CRv2c and CCC400 has been evaluated in Rohrer et al. (2018). For addressing the latest summer season (2018), we used the ERA5 reanalysis (Hersbach and Dee, 2017).

500 hPa GPH and blocking was reconstructed using equation (6), with x denoting Apr-Jul mean fields of 500 GPH and blocking, respectively. This gives atmospheric circulation statistics that are consistent with the corresponding harvest date.

4. Results and discussion

4.1 Presentation of the Beaune GHD series

285 The 664-year-long Beaune series is quite homogeneous showing nearly identical averages and standard deviations over the four sub-periods mentioned in section 2 prior to 1988 (Table 3). It is significantly correlated at $r = -0.76$ with April-to-July mean temperatures in Paris over the 360-year-long period 1659-2018 (Table 2).

290 The curve is clearly divided in two parts. Grapes were on average picked on 28 September from 1354 to 1987 comprising most of the Little Ice Age and the 20th century period of slow warming. In contrast, GHD's were 13 days earlier (15 September), during the last 31-year-long period of rapid warming from 1988 to 2018 (Table 3). The main phenological phases in the development of grapes (*Vitis vinifera*) – bud break, flowering, veraison (change colour and softening of the berries) went in step with the harvest dates (Jüstrich 2018).

295 Besides the climatic shift in 1988 several warm (positive) and cold (negative) fluctuations stand out: GHD were 6.5 days earlier between 1383 and 1435 than between 1354 and 1382. These fluctuations agree with those of glacier length. The Gorner glacier (Canton Valais, Switzerland) advanced since the 1340's culminated in 1385 on its first Little Ice Age maximum that corresponds to the position of the glacier in 1859. Then the glacier melted back to a low level which cannot exactly be established (Holzhauser 2010). Likewise the GHD curve mirrors the well-known 1520-1560 and 1720-1739 warm phases as well as the cold c.1600, c.1640 and 1820-1860 phases documented through the waxing and waning of Alpine glaciers (Nussbaumer and Zumbühl 2018). In 1520-1560 grapes were on average harvested four days earlier (24 Sep) than the mean value prior to 1988, and six days earlier (22 Sep) in the period 1720-1739. Between 1820 and 1860 on the contrary GHD occurred 4 days later (2 Oct.) than the mean value. This phase is strongly influenced by multiple volcanic eruptions from which the climate system only recovered slowly (Brönnimann et al. 2019).

4.2 Correlation with other proxy time series

305 Correlations between documentary-based proxy series and the Beaune GHD series turned out to be significant (Fig. 7). We focused on the period prior to 1850 and significant Pearson correlation was found between wheat harvest dates in Norfolk and grape harvest dates in Beaune despite the considerable distance between the two locations and the different nature of the proxy. The GHD available for Metz from 1420 to 1537 are well correlated to Beaune GHD as well ($r = 0.60$). On the other

hand the coefficient is surprisingly high between the GHD from the Czech Lands and the Beaune series despite the distance of 900 km between Beaune and the region northwest of Prague.

310 Tree-ring based temperature reconstructions were used to investigate the similarity with the grape harvest dates. Both have annual resolution and are influenced by the summer growing season. The closest tree-ring reconstructions are found at a distance of a few hundred kilometers from Beaune, similar to the long instrumental measurements from Paris. Nevertheless, seasonal average temperatures should be highly correlated over regions of several hundred kilometers.

315 Pearson correlation coefficients are expected to be negative because the warmer a growing season is, the earlier the harvest date and the thicker the tree ring or the denser the latewood. Correlation coefficient for both sites are high and clearly significant ($p < 0.05$), obviously not as high as with instrumental temperature because tree-ring proxies include additional noise from non-climatic influences. Correlations coefficient are robust and remain constant throughout all the tested sub-periods (Table 4).

4.3. Statistical model for harvest dates

320 GHD can be well reconstructed from temperature, using either model F_{linear} or $F_{\text{transformed}}$ (see Table 5, Fig. 8; scatterplots are shown in Fig. S1). Interestingly, both models produce a better correlation (>0.8) in the evaluation period (1851-2018) than in the calibration period (1659-1850), probably due to the strong and well reproduced trend of GHD in the evaluation period. Given the fact that the observed temperature refers to a location more than 300 km away and is based on early instruments with presumably substantial errors, a correlation coefficient of $r=0.8$ over the entire period is indeed surprising.

325 Correlations are still high when applying model $F_{\text{transformed}}$ to EKF400 (Table 5 top shows the range over the entire ensemble; see also the scatter plots in Fig. S1). This suggests that GHD can be modelled from climate simulation output, which is important for the analogue reconstruction approach.

The reconstruction of April-to-July mean temperature from GHD using models B_{linear} and $B_{\text{weighthed.simulations}}$ yields similar statistics as the forward approach (Table 5, Fig. 8, bottom). In the observations, April-to-July temperature reached the highest value ever in 2018. No year in the past was warmer in the observations (although there were two years with earlier harvest).

330 The autocorrelation structure of the Beaune GHD-based temperature reconstruction (black in figure S2; models B_{linear} and $B_{\text{weighthed.simulations}}$ lead to equal results) is very similar to the tree-ring reconstruction from the Pyrenees (blue) but has clearly less autocorrelation than the tree-ring reconstruction from the Löttschental, Switzerland (red). The Löttschental time series experiences much more low frequency variability than the Beaune and Pyrenees records. However, it is hard to argue that one of them should be the correct one. All have uncertainties with regard to their low frequency behaviour which could be due to age detrending, temporally inconsistent tree-age distributions, and other factors. On the other hand documentary data

for GHD are also not free from issues concerning low frequency variability. In the case of vine there may be adaptation, breeding, gene (de)activation processes over the decades that may dampen low frequency variability. There may also be changes in taste altering harvest dates in both directions.

Another point worth mentioning in comparison with tree-ring reconstructions is that the Beaune temperature reconstruction rather underestimates interannual variability ($SD=0.58$ K, period 1659-2007) compared to the Paris observations ($SD=0.87$ K) whereas both tree-ring reconstructions overestimate it (both $SD=1.10$ K).

4.4. Role of atmospheric blocking

What atmospheric conditions are conducive to early GHDs? Using the analogue approach we can analyse the April-to-July averaged 500 GPH and blocking statistics over the North Atlantic-European region for past years. Figure 9 shows GPH (anomalies in contours) and blocking (anomalies in colour, climatology in contours) fields that are consistent with the GHD of 1556; the second earliest on record after 2003. For comparison, we also show a composite of summer blocking for the 10 earliest GHDs in the period 1851 to 1980 from 20CRv2c reanalysis (we excluded the last decades due the strong anthropogenic warming effect) relative to the average over that period. Finally we also show blocking anomalies for summer 2018 relative to the average 2000-2018 from ERA5 reanalysis.

Early GHDs are related to high-pressure anomalies centred over Western or North Central Europe. High pressure situations are accompanied by increased radiation and high temperatures. With respect to blocking, anomalies are weak over the study area (central-western Europe). Rather, blocking during early GHD years occurred more frequently over Denmark (less frequently over Northern Scandinavia). In such situations the study area lies to the southwest of the block and receives dry and warm continental air masses. A similar blocking pattern is also found for the 10 earliest GHD years in the 20CR reanalysis. The year 2018 follows a similar pattern. Early harvest dates are thus related to blocking over Denmark. Late harvest dates (not shown) do not imprint significantly onto blocking.

4.5 Grape ripening in extremely hot and dry years

The 33 extremely warm events comprising the 5% percentile bracket of GHD are unevenly distributed over time (Fig. 10). 21 of them occurred between 1393 and 1719, i.e. one out of 15 years included a hot spring-summer period. In contrast, this is the case for just five years between 1720 and 2002, i.e. one out of 56. Under those circumstances, the memory of outstandingly warm years faded. No wonder that the hot summer 2003 came as a surprise. Since then 8 out of 16 spring-summer periods were outstanding according the statistic of the last 664 years, no less than 5 among them within the last 8 years. This implies that the extremes in the past have now become normal. The acceleration of extreme temperatures in the

last decade went along with an increased melting back or decay of Alpine glaciers which lost about 20% of their remaining volume (Swiss Glaciers 2017).

Subsequently conditions in two outstanding years - 1540 and 1556 - are considered in more detail. First of all, it is puzzling that the exceptional heat and drought in 1540 ranks only 19th in the statistics of Beaune GHD. Possible reasons to deemphasise this event based on tree ring evidence were brought forward by Büntgen et al. (2015). However, their arguments are thought to be questionable (Pfister et al. 2015). An interpretation of this paradox is attempted using vine phenological evidence available from vineyards around Biel-Bienne, Zürich and Schaffhausen (Switzerland) in 1540 and 1556 in comparison with the Beaune evidence. Source references are provided in the Euro-Climhist data-platform (<https://echdb.unibe.ch/>). Elevation matters for the comparison: the three Swiss vine growing areas are located at altitudes of about 430 metres asl, i.e. about 130 to 200 metres higher than those of Beaune region vineyards. Vine cultivars (CV), i.e. varieties, need to be considered besides altitudes. In Biel-Bienne CV Chasselas used to be grown while in Zurich and Schaffhausen CV Räuschling was cultivated which survives in a few vineyards until today. The maturity of the Chasselas and Räuschling cultivars is 10 to 14 days earlier than that of Pinot noir grown in the Beaune area (Altwegg 2018). Table 6 lists the main phenological grapevine stages - ‘end of flowering’, ‘beginning of veraison’ (colour change and softening of berries) and harvest dates for the years 1540, 1556, 2003 and 2018. Gladstones (2011) refers to the widely observed fact that “the date of flowering, can usually predict quite closely the dates of veraison (colour change and softening of berries) and maturity to follow [...]. The later phenological intervals show little response to temperature and tend to be constant from year to year”. His assessment is confirmed by Chuine et al. (2004) and Jüstrich (2018). In 1540, however, the interval between véraison and harvest took 67 days in the Zürich area, i.e. 27 days longer than usual, while this interval was somewhat shorter in 1556. In both years the grapes were harvested some time after full ripening. For 1540 this assumption is confirmed by the sources. “Vine-growers in Schaffhausen (Switzerland) were “long waiting for rain to begin the harvest”, as chronicler Oswald Huber relates. He writes, however, they “finally tackled the work nevertheless, because the plants withered.” Likewise, vine-growers at the shores of Lake Constance and in the Upper Alsace interrupted the vintage after picking a few juicy grapes, because the remaining ones were quasi dried out. The vintage was then resumed after a two-day spell of rain. At harvest time grapes in many vineyards had become raisins. They yielded a sweet sherry-like wine which made people more rapidly drunk than usual (Wetter and Pfister 2013). It needs to be stressed, that meteorological conditions were almost matchless in 1540. The evidence for this mega-drought taken from documentary data offers a broad spectrum of evidence on past weather and climate and their societal impacts. It is described in a sample of 312 first-hand documentary weather reports originating from continental Europe (Wetter and Pfister, 2013). The hot and dry period in 1540 lasted from April to the end of the year. The heat-wave peaked at the end of a 46 day long rainless period between 23th June and 7th August during which many forests and settlements in a large area from the Ardennes to Poland went up in flames (Pfister 2018).

Maximum temperatures from late July may have exceeded 40°C (Orth et al. 2016). Vagrants and homeless people were hunted down for raising town and forest fires (Pfister 2017). The heatwave of 1540 undoubtedly struck the region of Burgundy. The church of Notre-Dame of Beaune organised 8 processions to call for rains from the beginning of May to the end of August (Labbé, Gaveau 2011). In the near city of Besançon (c. 100 km eastwards) a chronicler wrote that warm temperatures lasted from April to November and that the heatwave was hardly bearable by humans during summer (Wetter et al. 2014). It cannot be excluded, that even the date of véraison was also somewhat delayed due to drought stress. Research in grapevine biology established that under conditions of extreme heat and drought the development of grapes is slowed down or stopped (Keller 2016). When occurring before véraison, extreme hydric stress alters grape quality and the onset of ripening, possibly because it induces leaf defoliation and therefore carbon assimilation limitation (Basile et al., 2012; Girona et al., 2009; Ollat and Gaudillere, 1998). In fact, this phenomenon was observed during the temperature peak in the hot summer 1947 in Schaffhausen (Amtsblatt 1947).

In 1556 grapes in Beaune were harvested on 16th August. In central France, the 1556 heat and drought began in mid-April, i.e. about 45 days later than in 1540, following a wet winter. The heat-wave to mid-June spurred vegetation growth. In the Loir-et-Cher region, a variety of red Pinot noir cultivar named Auvernat was already blossoming around the 25 of April (Nouel, 1878, 235). Not a drop of rain fell during this period. On 14th June, it was pouring down for three to four hours which visibly refreshed the vegetation. Before 10th July the first grapes were ripe. In July the ground became so hot that it burnt people's feet walking barefoot. Like in 1540 the heat and drought peaked in early August (Bourquelot, 1857, 30-31; Hiver 1867, 81 and 90; Nouel 1878, 235-237). Following the deliberation protocols of the chapter of Notre-Dame of Beaune, seven processions for obtaining rain were held in the city from 15th June to 15th August (Arch. Dep. Côte d'Or, G 2499).

The 1556 harvest of Chasselas grapes in Biel-Bienne was estimated to have occurred between 25th and 30th August i.e. at about the same time as that of Pinot noir in Beaune considering the chronicler's remark that the abundant harvest already ended on 10th September (Gregorian Style). Like in the Beaune area, the preceding winter 1556 had been very wet considering the daily weather observations by Wolfgang Haller in Zürich. Disregarding June, which was completely rainless, the spring-summer period included 7 precipitation days in April, 3 in May and 5 in July (Haller in Pfister, Rohr <https://echdb.unibe.ch/>, 15th Oct 2018). A hot and almost rainless period began on 29th July and lasted to 11th September, i.e. more than a month later than in 1540.

The heat-wave in summer 2003 was reassessed by Pfister (2018). The water deficit was not sufficiently strong to "block" grape ripening through extremely limited photosynthesis. Obviously, ripening made its course quickly. Additionally, wine growers harvested pretty soon to maintain sufficient acidity in wines mostly (and in some cases avoid excessively high alcohol content in wines).

Since the early 21st century, higher temperatures combined with increase control of grape sanitary status (grey mould disease mostly) makes ripening duration (i.e. the lag during veraison and harvest) more winemaker dependent. The winemaker choices depend on both cultivar and style of wine. For instance, the number of days between veraison and harvest for CV Cabernet-Sauvignon has nearly been doubled in a famous Château in the appellation Margaux near Bordeaux (Van Leeuwen and Destrac-Irvine, 2017)

In sum, the decline of 13 days in the average date of GHD since 1988 went along with large increase in the number of extreme spring to summer seasons. These include involve situations such in 1540 in which both human and ecological systems behave non-linearly outside the normal range of biological and probability laws. Documentary data may be helpful to describe such conditions in the necessary detail.

5. Conclusions

Time series of documentary proxy data such as GHD need to be critically evaluated by historians prior to statistical analysis. 19th century's publications need to be cautiously examined before being used and in any case first-hand documentary material should be preferred, which takes a lot of meticulous detail work in the archives.

The 664-year-long Beaune GHD series assembled from the archives of the city is significantly correlated with the long Paris temperature series from 1659 to 2018 and with documentary and tree-ring proxy data from 1354 to 1658. Statistical models describe Beaune GHD very well, with Pearson correlations around 0.8. The climate shift of 1988 divides the series in two different parts. Over the period of the Little Ice Age and the “warm twentieth century” up to 1987 grapes in the Beaune area were picked on 28 September on average. After the climate shift in 1988 the harvest date declined by 13 days to an average of 15th of September during the last 31-year-long period of rapid warming from 1988 to 2018. It is noteworthy that the 33 values below the 5% percentile are unevenly distributed over time. While 21 of them occurred between 1354 and 1719, only 4 of them were registered between 1720 and 1987. In contrast, 8 outstanding extremes occurred within the last 30 years, 3 between 2000 and 2010 and 5 between 2011 and 2018, which probably witnessed the warmest warm season temperatures since 1354.

Early harvest dates coincide with high pressure influence and increased blocking over Denmark. Conversely, the most outstanding heat and drought years were not necessarily the earliest in the ranking of harvest dates. It is concluded that grape development slowed down or even stopped during very long rainless periods and extreme maximum temperatures such as in 1540 and 1473 (Camenisch et al. in review). In sum, the long homogenized Beaune series visually demonstrates that warm extremes in the past were outliers, whereby they have become the norm in the present time.

Author contribution:

Thomas Labbé provided the Beaune GHD dataset by archival researches, and wrote section 1 and 2 as well as 4.1.together with Christian Pfister

Christian Pfister wrote section 1, and 4.1 together with Thomas Labbé, section 4.5. together with Benjamin Bois and wrote sections 4.2 and 5.

Stefan Brönnimann and Jörg Franke performed the statistical reconstructions and wrote sections 3.2, 3.3, 4.3, 4.4.

Daniel Rousseau provided the updated Paris mean temperature series (1659-2018) and wrote section 2.6.

Benjamin Bois provided information on vine phenology and helped writing section 4.5.

Competing interests:

The authors declare that they have no conflict of interest.

Acknowledgments:

Stefan Brönnimann and Jörg Franke were supported by the Swiss National Science Foundation (project RE-USE) and by the the European Research Council (AdG 787574 “PALAEO-RA”). Simulations were performed at the Swiss National Supercomputing Centre CSCS. The Twentieth Century Reanalysis Project datasets are supported by the U.S. Department of Energy (DOE) Office of Science Innovative and Novel Computational Impact on Theory and Experiment (INCITE) program, and Office of Biological and Environmental Research (BER), and by the National Oceanic and Atmospheric Administration Climate Program Office

The authors thank the team of the GEOBFC of the Maison des Sciences de l’Homme of Dijon (USR CNRS – Ub 3516) for the making of Figure 2.

References:

Alzwegg, A. Weinlandschaft Zürichsee, Geschichte und Gesichter, Th. Gut editions, Stäga 2018, 213 pp.

Amtsblatt: Amtsblatt des Kantons Schaffhausen 7/1947 : 4, 1947.

Angot, A.: Études sur les vendanges en France, Annales du bureau central météorologique de France, année 1883, B29-B120, 1885.

Arnoux, C.: Dissertation sur la situation de la Bourgogne, sur les vins qu’elle produit, sur la manière de cultiver les vignes, de faire le vin et de l’éprouver, Londres, Samuel Jallason, 1728.

- 485 Basile, B., Girona, J., Behboudian, M. H., Mata, M., Rosello, J., Ferré, M. and Marsal, J.: Responses of “Chardonnay” to deficit irrigation applied at different phenological stages: vine growth, must composition, and wine quality, *Irrig. Sci.*, 30(5), 397–406, doi:10.1007/s00271-012-0353-1, 2012.
- Bhend, J., Franke, J., Folini, D., Wild, M., Brönnimann, S.: An ensemble-based approach to climate reconstructions, *Climate Past*, 8, 963–976, 2012.
- 490 Blanchemanche, P.: Cruces historiques et vendanges en Languedoc méditerranéen oriental : la source, le signal et l’interprétation, *Archéologie du Midi Médiéval*, 27, 225-235, 2009 .
- Bourquelot, F.: Mémoires de Claude Haton, contenant le récit des événements accomplis de 1553 à 1582, principalement dans la Champagne et la Brie, vol. 1, Paris, Impr. Impériale, 1857.
- Brázdil, R., Kotyza, O.: History of Weather and Climate in the Czech Lands III, Daily Weather Records in the Czech Lands in the Sixteenth Century, Brno, Masaryk University, 1999.
- 495 Brázdil, R., Pfister, C., Wanner, H., Von Storch, H., Luterbacher, J.: Historical Climatology in Europe—The State of the Art, *Climatic Change*, 70, 363–430, 2005.
- Brönnimann, S., Franke, J., Nussbaumer, S. U., Zumbühl, H. J., Steiner, D., Trachsel, M., Hegerl, G. C., Schurer, A., Worni, M., Malik, A., Flückiger, J., Raible, C. C.: Last phase of the Little Ice Age forced by volcanic eruptions. *Nature Geoscience*, in press, 2019.
- 500 Büntgen, U., Frank, D. C., Nievergelt, D., Esper, J.: Summer temperature variations in the European Alps, AD 755–2004. *J Climate*, 19(21), 5606–5623, 2006.
- Büntgen, U., Tegel, W., Carrer, M., Krusic, P. J., Hayes, M., Esper, J.: Commentary to Wetter et al. (2014): Limited tree-ring evidence for a 1540 European ‘Megadrought’, *Clim. Chang.*, 131/2, 183-190, doi: 10.1007/s10584-015-1423-1, 2015.
- 505 Chaptal, J.A.: L’art de faire le vin, Paris, Impr. de Crapelet, 1807.
- Chuine, I., Yiou, P., Viovy, N., Seguin, B., Daux, V., and Le Roy Ladurie, E.: Grape ripening as a past climate indicator, *Nature*, 432, 89–290, 2004.
- Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Brönnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., Mauder, M., Mok, H. Y., Nordli, Ø., Ross, T. F., Trigo, R. M., Wang, X. L., Woodruff, S. D. and Worley, S. J.: The Twentieth Century Reanalysis Project, *Quarterly Journal of the Royal Meteorological Society*, 137(654), 1–28, doi:10.1002/qj.776, 2011.
- 510 Craddock, J. M.: Methods of comparing annual rainfall records for climatic purposes. *Weather*, 34, 332–346, 1979.

Daux, V., Garcia de Cortazar-Atauri, I., Yiou, P., Chuine, I., Garnier, E., Le Roy Ladurie, E., Mestre, O., and Tardaguila, J.:
 515 An openaccess database of grape harvest dates for climate research: data description and quality assessment, *Clim. Past*, 8, 1403–1418, doi:10.5194/cp-8-1403-2012, 2012.

Dion, R.: *Histoire de la vigne et du vin en France, des origines au XIX^e siècle*, Paris, chez l'auteur, 1959.

Dobrovolný, P., Moberg, A., Brázdil, R., Pfister, C., Glaser, R., Wilson, R., Van Engelen, A., Limanówka, D., Kiss, A.,
 520 Halíčková, M., Macková, J., Riemann, D., Luterbacher, J., Böhm, R.: Monthly, seasonal and annual temperature reconstructions for central Europe derived from documentary evidence and instrumental records since AD1500. *Clim Change* 101:69–107, 2010.

Dorado Liñán, I., Buentgen, U., Gonzalez-Rouco, F., Zorita, E., Montavez, J. P. , Gomez-Navarro, J. J., Brunet, M.,
 Heinrich, I., Helle, G. , Gutiérrez, E.: Estimating 750 years of temperature variations and uncertainties in the Pyrenees by
 tree ring reconstructions and climate simulations. *Climate of the Past* 8(3), 919–933, 2012.

525 Dufour, M. L.: Problème de la variation du climat. *Bulletin de la Société Vaudoise des Sciences naturelles*, 10, 359–556, 1870.

Egli, E. (ed): *Heinrich Bullingers Diarium (Annales Vitae) der Jahre 1504-1574, Zum 400. Geburtstag Bullingers am 18. Juli 1504*, Basel Basler Buch- und Antiquariatshandlung, vormals Adolf Geering, 1904.

Franke, J., Brönnimann S., Bhend J., and Brugnara Y.: A monthly global paleo-reanalysis of the atmosphere from 1600 to
 530 2005 for studying past climatic variations. *Scientific Data* 4, 170076, 2017.

Garcia de Cortázar-Atauri, I., Daux, V., Garnier, E., Yiou, P., Viovy, N., Seguin, B., Boursiquot, J.-M., Parker, A.-K., van
 Leeuwen, C. and Chuine, I.: Climate reconstructions from grape harvest dates: Methodology and uncertainties, *Holocene*
 20(4), 599-608, 2010.

Garnier, E., Daux, V., Yiou, P., and Garcia de Cortázar-Atauri, I.: Grapevine harvest dates in Besançon (France) between
 535 1525 and 1847: social outcomes or climatic evidence?, *Climatic Change*, 104, 703–727, doi:10.1007/s10584-010-9810-0, 2011.

Girona, J., Marsal, J., Mata, M., Del Campo, J. and Basile, B.: Phenological sensitivity of berry growth and composition of
 Tempranillo grapevines (*Vitis vinifera* L.) to water stress, *Aust. J. Grape Wine Res.*, 15(3), 268–277, doi:10.1111/j.1755-
 0238.2009.00059.x, 2009.

540 Gladstones, J.: *Wine, Terroir and Climate Change*, Kent Towns, Wakefield Press, 2011.

Guerreau, A.: *Climat et Vendanges: Révisions et compléments*, *Histoire et Mesure*, 10, 89–147, 1995.

Guiot, J., Nicault, A., Rathgeber, C., Edouard, J.-L., Guibal, F. and Till, C.: Last-millennium summer-temperature variations
 in western Europe based on proxy data, *Holocene* 15, 489-500, 2005.

Hersbach, H., Dee, D.: ERA5 reanalysis is in production, *ECMWF Newsletter* 147, 7, 2016.

- 545 Hiver, M. (ed): *Journal de Jehan Glaumeau*. Bourges, 1541-1562, Bourges, 1867.
- Holzhauser, H.: Zur Geschichte des Gornergletschers. Ein Puzzle aus historischen Dokumenten und fossilen Hölzern aus dem Gletschervorfeld. *Geographica Bernensia* G84, Bern, 253 pp, 2010.
- Jüstrich, H.: Klimawandel- Fakt oder Fiktion? *Schweizer Z. für Obst- und Weinbau*, 16/13, 1-7, 2018.
- Keller, M.: *The Science of Grapevines, Anatomy and Physiology* Second Edition, Academic Press, Elsevier Inc., Amsterdam 2016, 2016.
- 550 Krieger, M., Lohmann, G. and Laepple, T.: Seasonal climate impacts on the grape harvest date in Burgundy (France), *Clim. Past* 7, 425-435, 2011.
- Labbé, T., Gaveau, F.: Les dates de vendange à Dijon : établissement critique et révision archivistique d'une série ancienne, *Revue historique*, 657, 19-51, 2011.
- 555 Labbé, T., Gaveau, F.: Les dates de vendange à Beaune (1371-2010). Analyse et données d'une nouvelle série vendémiologique, *Revue historique*, 666, 333-367, 2013a.
- Labbé, T., Gaveau F.: Deux cents ans de réflexion sur les bans de vendange : un bilan historiographique de la question, *in* Ducos J. (ed), *Météores et climats d'hier. Décrire et percevoir le temps qu'il fait de l'Antiquité au XIX^e siècle*, Paris, Hermann, 179-203, 2013b.
- 560 Lachiver, M.: *Vins, vignes et vigneron*. Histoire du vignoble français, Paris, Fayard, 1988.
- Lavalle, J.: *Histoire et statistique de la vigne et des grands vins de la Côte d'Or*, Paris, Dusacq, 1855.
- Legrand, J.-P.: L'expression de la vigne au travers du climat depuis le Moyen-Age, *Revue Française d'oenologie*, 75, 23-52, 1979.
- Le Roy Ladurie, E.: *Times of Feast, Times of Famine: A History of Climate since the Year 1000*, Allen & Unwin, London (Original: *Histoire du climat depuis l'an mil*, Paris, Flammarion, 1971).
- 565 Le Roy Ladurie: *Histoire humaine et comparée du Climat. Canicules et Glaciers (XIII^e à XVIII^e siècles)*, Paris, Fayard, 2004.
- Le Roy Ladurie, E., Daux, V., and Luterbacher, J.: Le climat de Bourgogne et d'ailleurs, *Histoire, Économie et Sociétés* 3, 421-436, 2006.
- 570 Le Roy Ladurie, E. and Daux, V.: The climate in Burgundy and elsewhere, from the fourteenth to the twentieth century, *Interdisc. Sci. Rev.*, 33, 10-24, 2008.
- Litzenburger, L.: *Une ville face au climat. Metz à la fin du Moyen Âge (1400-1530)*, Nancy, Presses Univ. de Nancy, 2015.
- Meier, N., Rutishauser, T., Pfister, C., Wanner, H., Luterbacher, J.: Grape harvest dates as a proxy for Swiss April to August temperature reconstructions back to AD 1480, *Geophysical Research Letters* 34, L20705, doi:10.1029/2007GL031381, 2007.
- 575

Menzel, A.: A 500 year pheno-climatological view in the 2003 heatwave in Europe assessed by grape harvest dates, *Meteorologische Zeitschrift* 14, 75-77, 2005.

Možný, M., Brázdil, R., Dobrovolný, P., Trnka, M.: Cereal harvest dates in the Czech Republic between 1501–2008 as a proxy for March–June temperature reconstruction, *Climatic Change*, 110, 808-821, 2012.

580 Možný, M., Brázdil, R., Dobrovolný, P., Trnka, M.: April-August temperatures in the Czech Lands, 1499-2015, reconstructed from grape-harvest dates, *Climate of the Past*, 12, 1421-1434, doi: 10.5194/cp-12-1421-2016, 2016.

Noirot, E.: Recherches sur les époques de la maturité des raisins, *Le Cultivateur*, 18 août 1836, 1836.

Nouel, M.E. (ed.): Chronique de Michel Garault, chanoine de Troo (1543-1598), *Bulletin de la Société archéologique, scientifique et littéraire du Vendômois*, 17, 222-257, 1878.

585 Nussbaumer, S.U., Zumbühl, H.J.: Evidence from the Archives of Societies: Historical Sources in Glaciology, in White, S., Pfister, C., Mauelshagen, F. (ed), *The Palgrave Handbook of Climate History*, London, Palgrave Macmillan, 93-97, 2018.

Ollat, N. and Gaudillere, J. P.: The Effect of Limiting Leaf Area During Stage I of Berry Growth on Development and Composition of Berries of *Vitis vinifera* L. cv. Cabernet Sauvignon, *Am. J. Enol. Vitic.*, 49(3), 251–258, 1998.

590 Orth, R., Vogel, M., Luerbacher, J., Pfister, C., Seneviratne, S.I.: Did European Temperatures in 1540 Exceed Present-day Records?, *Environmental Research Letters*, 11, 114021, doi: 10.1088/1748-9326/11/11/14021, 2016.

Otto-Bliesner, B.L., E.C. Brady, J. Fasullo, A. Jahn, L. Landrum, S. Stevenson, N. Rosenbloom, A. Mai, G. Strand: Climate Variability and Change since 850 C.E.: An Ensemble Approach with the Community Earth System Model (CESM), *Bulletin of the American Meteorological Society*, 735-754, 2016.

595 Pfister, C.: *Das Klima der Schweiz von 1525 bis 1860 und seine Bedeutung in der Geschichte von Bevölkerung und Landwirtschaft*. 2 vols. Paul Haupt. Bern, 1984.

Pfister, C., Wetter, O., Brázdil, R., Dobrovolný, P., Glaser, R., Luterbacher, J., Seneviratne, S. I., Zorita, E., Alcoforado, M.-J., Barriendos, M., Bieber, U., Burmeister, K. H., Camenisch, C., Contino, A., Grünewald, U., Herget, J., Himmelsbach, I., Labbé, T., Limanówka, D., Litzemberger, L., Kiss, A., Kotyza, O., Nordli, Ø., Pribyl, K., Restö, D., Riemann, D., Rohr, C., Werner, S., Spring, J.-L., Söderberg J., Wagner, S., Werner, J. P.: Tree-rings and people – different views on the 1540 Megadrought. Reply to Büntgen et al. 2015, *Clim. Chang.*, 131/2, 191-198, doi: 10.1007/s10584-015-1429-8, 2015.

600 Pfister, C.: When Europe was burning. The multi-seasonal drought of 1540 and the arsonist paranoia, in Gerrit Jasper Schenk (ed.), *Disasters, Risks and Cultures. A Comparative and Transcultural Survey of Historical Disaster Experiences between Asia and Europe*, Springer, 155-185, 2017.

605 Pfister, C.: The “Black Swan” of 1540. Aspects of a European Megadrought, in Klaus Leggewie and Franz Mauelshagen (eds.), *Climatic Change and Cultural Transition in Europe*, Leiden, Brill, 156-196, 2018.

Pfister, C., Camenisch, C., Dobrovolný, C.: *Analysis and Interpretation: Temperature and Precipitation Indices*, in White S., Pfister C., Mauelshagen F. (eds), *The Palgrave Handbook of Climate History*, London, Palgrave, 115-129, 2018.

Pribyl, K., Cornes, R., Pfister, C.: Reconstructing Medieval April–July Mean Temperatures in East Anglia, 1256–1431, *Climatic Change*, 113, 393-412, 2012.

610 Pribyl, K., *Farming, Famine and Plague. The Impact of Climate in Late Medieval England*, Cham, Springer, 2017.

Rohrer, M., Brönnimann S., Martius O., Raible C. C., Wild M., Compo G. P.: Representation of cyclones, blocking anticyclones, and circulation types in multiple reanalyses and model simulations., *J. Climate*, 31, 3009–3031, 2018.

Rotelli, C., *Storia agraria del Piemonte fra il 1250 e il 1450*, Torino, 1973.

Rousseau, D.: Les températures mensuelles en région parisienne de 1676 à 2008, *La météorologie*, 67, 43-55, 2009.

615 Rousseau, D. : Les moyennes mensuelles de températures à Paris de 1658 à 1675, d’Ismael Boulliau à Louis Morin, *La météorologie*, 81, 11-22, 2013.

Rousseau, D.: Fluctuations des dates de vendanges bourguignonnes et fluctuations des températures d’avril à septembre de 1378 à 2010, XXVII^e Colloque de l’Association Internationale de Climatologie, 2-5 juillet 2014 – Dijon (France), 757-763, 2015.

620 Stehr, N., von Storch, H., Brückner, E., 2000: *The Sources and Consequences of Climate Change and Climate Variability in Historical Times*, Dordrecht, Kluwer Academic Press.

Scherrer, S. C., Croci-Maspoli, M., Schwierz, C., and Appenzeller, C.: Two-dimensional indices of atmospheric blocking and their statistical relationship with winter climate patterns in the Euro-Atlantic region. *Int. J. Climatol.*, 26, 233–249, 2006.

625 Swiss Glaciers: *The Swiss Glaciers Yearbook of the Cryospheric Commission of the Swiss Academy of Sciences (SCNAT)*, published since 1964 by the Laboratory of Hydraulics, Hydrology and Glaciology (VAW) of ETH Zürich, No. 136, <http://www.glamos.ch>, 2017.

Tibaldi, S., Molteni, F.: On the operational predictability of blocking. *Tellus*, 42A, 343–365, 1990.

Tibaldi, S., Tosi, E., Navarra, A., Pedulli, L.: Northern and Southern Hemisphere seasonal variability of blocking frequency and predictability, *Mon. Wea. Rev.*, 122, 1971–2003, 1994.

630 Van Leeuwen, C., Destrac-Irvine, A.: Modified grape composition under climate change conditions requires adaptations in the vineyard, *OENO One*, 51(2), 147, doi:[10.20870/oenone.2016.0.0.1647](https://doi.org/10.20870/oenone.2016.0.0.1647), 2017.

Wetter, O. and Pfister, C.: Spring-summer temperatures reconstructed for northern Switzerland and southwestern Germany from winter rye harvest dates, 1454–1970, *Clim. Past*, 7, 1307–1326, doi:10.5194/cp-7-1307-2011, 2011.

635 Wetter, O., Pfister, C.: An underestimated record breaking event – why summer 1540 was likely warmer than 2003. *Clim Past*, 9, 41–56, 2013.

640 Wetter, O., Pfister, C., Werner, J.P., Zorita, E., Wagner, S., Seneviratne, S.I., Herget, J., Grünewald, U., Luterbacher, J., Alcoforado, M.J., Barriendos, M., Bieber, U., Brázdil, R., Burmeister, K.H., Camenisch, C., Contino, A., Dobrovolný, P., Glaser, R., Himmelsbach, I., Kiss, A., Kotyza, O., Labbé, T., Limanówka, D., Lützenburger, L., Nordli, Ø., Pribyl, K., Retsö, D., Riemann, D., Rohr, C., Siegfried, W., Söderberg, J., Spring, J.L.: The year-long unprecedented European heat and drought of 1540 – a worst case, *Clim Chang.*, 125, 349-363, doi:10.1007/s10584-014-1184-2, 2014.

645 Wilson, R., Anchukaitis, K., Briffa, K., Büntgen, U., Cook, E., D'Arrigo, R., Davi, N., Esper, J., Frank, D., Gunnarson, B., Hegerl, G., Helema, S., Klesse, S., Krusic, P., Linderholm, H.W., Myglan, V., Osborn, T., Rydval, M., Schneider, L., Schurer, A., Wiles, G., Zhang, P., Zorita, E.: Last millennium Northern Hemisphere summer temperatures from tree rings: Part I: the long term context. *Quaternary Science Reviews* 134: 1-18, <http://dx.doi.org/10.1016/j.quascirev.2015.12.005>, 2016.




Figure 2: Geographical area of the study.






Figure 3: The vineyard of the city of Beaune



655

Figure 4: Mismatches between “Angot 1885” and “Labbé-Gaveau 2011” Dijon GHD series (1385-1905).

660



665 *Figure 5: Time interval between the GHD of Dijon and the GHD of Beaune (1600-1800).*






Figure 6: 11-years low- pass filtered GHD series of Beaune (homogenised and non-homogenised), Switzerland (Wetter et al. 2013) and Salins (Angot 1885)



675


Figure 7: Relation between April-to-July mean temperature (°C) in Paris and Beaune GHD. Green dots indicate the calibration period, blue dots evaluation period. The linear regression line is shown.



680

Figure 8: Comparison of times series of GHD from Beaune (this work), Czech Lands (Mózný et al. 2016), Switzerland (Wetter et al. 2013), and Metz (Litzenburger 2015), as well as temperature reconstructions for Norfolk (Prybil et al. 2012; right scale, inverted). The number indicate the correlation between the series prior to 1850.

685



690 Figure 9: (top) Harvest dates in observations and modelled from observed and reconstructed (EKF400) temperatures. (bottom) April-to-
July mean temperature from observations and reconstructed from models B_{linear} and $B_{weighted.simulations}$. Blue shading denotes the 95%
695 prediction interval for B_{linear} .

695




Figure 10: Anomalies in April-to-July 500 hPa GPH (purple contours in gpm) and blocking frequency (in % of time steps blocked, note the non-linear scale) in (left) model $B_{weighted}$ simulations for 1556 (relative to all years prior to 1850) as well as (middle) averaged for the 10 earliest harvest dates in 1851 to 1980 in 20CRv2c (relative to 1851-1980). The right figure shows anomalous 500 hPa GPH and blocking in April to July 2018 in ERA5 (relative to 2000-2018). Grey solid contours give the corresponding blocking climatologies (in % of time steps blocked). Red lines indicate where the 5% to 95% confidence interval for anomalous blocking is exceeded based on 1000 repetitions of a Monte Carlo sampling of the weights (for model B) or years in 20CRv2c, respectively.

700

705




Figure 11: Beaune GHD time series (1354-2018), with indication of the 5% earliest dates.

710

715

720

725

Table 3: Comparison of mean GHD time series in Beaune (this article), Aubonne (Angot 1885), Salins (Angot 1885), Switzerland (Wetter et al. 2013) and Czech Lands (Mózny et al. 2016).

	Period 1: 1599-1717	Period 2: 1718-1875	1354-1717	1718-2018	Difference of days between period 1 and period 2
Mean Beaune GHD	21 Sep	28 Sep	20 Sep	27 Sep	+7
Mean Swiss GHD	20 Oct	15 Oct			-5
Mean Czech Lands GHD	14 Oct	14 Oct			0
Mean Salins GHD	11 Oct	09 Oct			-2
Mean Aubonne GHD	21 Oct	18 Oct			-3

730

Table 4: Pearson correlation (r) between Beaune GHD time series and March to September mean Paris temperatures (1659-2018).

March (M)	Apr (A)	May (M)	Jun (J)	Jul (J)	Aug (A)	Sept (S)	AMJJAS	AMJJA	AMJJ
-0.28	-0.42	-0.51	-0.53	-0.43	-0.31	-0.22	-0.72	-0.74	-0.76

735

740

Table 5: Mean GHD and Standard deviation for various subperiods of the Beaune GHD homogenised time series.

	1354-1506	1507-1699	1700-1965	1966-1987	1988-2018
Mean GHD	28 Sep	27 Sep	28 Sep	28 Sep	15 Sep
Std. Dev.	9.88	10.60	8.91	7.20	9.93

745

750

Table 6: Pearson correlation coefficients (r) for various subperiods between wine harvest dates and tree-ring based temperature reconstructions. All correlations are statistically significant ($p < 0.05$).

	1354-2010	1354-1659	1659-2010	1711-1770	1771-1850
Pyrenees	-0.41	-0.36	-0.44	-0.48	-0.48
Lötschental	-0.32	-0.35	-0.34	-0.41	-0.56

755

760

765

Table 7: Reconstruction statistics for the forward (F) and backward (B) models. Obs. denotes observations. All models were calibrated against observations in the 1659-1850 period ($B_{\text{weighted.simulations}}$ is not calibrated but uses the calibrated model $F_{\text{transformed.CCC400}}$) and evaluated in the period 1851-2018. $F_{\text{transformed.EKF400}}$ denotes the application of the calibrated model $F_{\text{transformed}}$ to EKF400. „Calib.“ and „Eval.“ indicate the model statistics in the calibration and evaluation periods, respectively, sd. indicates its standard deviation (in days and °C, respectively) for the common period (1659-2004). Numbers for EKF400 refer to the ensemble mean of 30 reconstructions.

	Reconstructing GHD from Temperature				Reconstructing $T_{\text{Apr-Jul}}$ from GHD		
	Obs.	F_{linear}	$F_{\text{transformed}}$	$F_{\text{transformed.EKF400}}$	Obs.	B_{linear}	$B_{\text{weighted.simulations}}$
r_{cal}		0.726	0.731	0.651		0.704	0.702
r_{eval}		0.832	0.843	0.720		0.824	0.822
bias_{cal}		0.00	-0.20	0.28		0.00	-0.18
$\text{bias}_{\text{eval}}$		1.06	0.48	-1.46		0.02	-0.15
sd	9.37	6.91	7.10	6.14	0.85	0.57	0.60

770

775

780

785

Table 8: Phenological stages (flowering, veraison, ripening of the grapes and harvest dates) of the vine during extremely early years. S (Schaffhausen, white Räuschling cultivar); Z (Zürich, white Räuschling); Bi (Biel-Bienne, white Chasselas), M (Malans, red Pinot noir), B (Beaune, red Pinot noir)

Year	flower (end)	DOY	veraison	DOY	Ripe (grapes)	DOY	GHD	DOY	veraison to GHD (days)
1540	10 Jun (S)	162	5 Jul (S)	187	16 Jul (Z)	198	11 Sep (Z)	254	67 (S-Z)
1540							3 Sep (B)	247	
1556	10 Jun (Z)	162	4 Jul (S)	186	19 Jul (Z)	201	25-30 Aug (Bi)	238- 243	52 -57 (Bi-S)
1556							26 Aug (B)	239	
2003	30 May (B)	150	28 Jul 28 (B)	209			20 Aug (B)	232	23(B)
2018	31 May (B)	151	31 Jul 31	212			30 Aug (B)	242	30 (B)
2018	31 May (M)	151	1 Aug (M)	213			17 Sep (M)	261	48 (M)

790