

1 **Response of *Pinus sylvestris* var. *mongolica* to water change and drought** 2 **history reconstruction in the past 260 years, northeast China**

3 Liangjun Zhu^{1, 2}, Qichao Yao¹, David J. Cooper², Shijie Han^{3,4}, Xiaochun Wang^{1*}

4 ¹School of Forestry, Northeast Forestry University, Harbin 150040, China;

5 ²Department of Forest and Rangeland Stewardship, Colorado State University, Fort Collins, CO 80523, USA;

6 ³School of Life Science, Henan University, Kaifeng 475004, China

7 ⁴Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

8 Correspondence to: Xiaochun Wang (wangx@nefu.edu.cn)

9 **Abstract.** We present a 260-year annual PDSI reconstruction based on a tree-ring width chronology of
10 Scots pine (*Pinus sylvestris* var. *mongolica*) from four sample sites in the Central Daxing'an Mountains,
11 northeast China. The reconstruction equation explained 38.2 % of the variance of annual PDSI in the
12 calibration period from 1911 to 2010. Our reconstruction confirmed the local historical documents and
13 other nearby hydroclimate reconstructions. Drought in the 1920s-1930s was more severe in the
14 Daxing'an Mountains than the surrounding areas. A slight moisture increase was identified in the study
15 area, while a warm-dry pattern was found in the West-Central Mongolian Plateaus (mild drier) and their
16 transition zones: The West-Central Mongolian Plateaus (severe drier). Overall, the variation of drought
17 in the Daxing'an Mountains and its relationship with surrounding areas may be affected by the Pacific
18 or Atlantic oscillations (e.g., ENSO, PDO, AMO, NAO and SNAO), which can affect the Asian
19 monsoon, change the local temperature and precipitation, and lead to drought.

20 **Keywords** PDSI reconstruction; Central Daxing'an Mountains; Tree rings; *Pinus sylvestris* var.
21 *mongolica*; PDO; AMO; Drought

22

23 **1 Introduction**

24 Drought as one of the major natural disasters is being more frequently with climate change in the
25 world (Cook et al., 2010; Dai, 2011, 2013; Davi et al., 2006; Li et al., 2016). Severe droughts can
26 threaten agriculture and human social activities, and also have a devastating impact on human lives and
27 the survival of native and domestic plants and animals (Cook et al., 2010; Dong et al., 2013; Shen,
28 2008; Sun, 2007). Drought is one of the most severe and frequent natural disasters in China, especially
29 in semi-arid and arid regions (Bao et al., 2015; Chen et al., 2015; Cook et al., 2010; Dong et al., 2013;
30 Liang et al., 2006; Shen, 2008; Sun and Liu, 2013; Xu, 1998). For example, the drought in the 1920s
31 affected almost all of northern China, accompanied by severe economic and social losses (Dong et al.,
32 2013; Liang et al., 2006; Shen, 2008; Sun, 2007). Natural droughts are recorded in tree rings in the arid
33 or semi-arid regions (Bao et al., 2015; Chen et al., 2015; Liang et al., 2006; Sun and Liu, 2013; Wang
34 and Song, 2011). Recent studies indicate a trend of increasing drought frequency, persistence and
35 severity due to global warming in many regions of the world (Bao et al., 2015; Cook et al., 2010; Dai,
36 2011, 2013; Schrier et al., 2013). A rapid and pronounced warming accompanied by a decrease in
37 precipitation has occurred in China, especially in high latitude and high altitude regions (Bao et al.,
38 2015; Chen et al., 2015; Cook et al., 2010; Dai, 2013; Sun and Liu, 2013; Zhu et al., 2017), leading to
39 severe and prolonged drought in recent decades, such as from 1999 to 2002 (Bao et al., 2015; Liu et al.,
40 2009; Shen, 2008).

41 The Daxing'an Mountains in northeast China is a transition area from semi-humid climate in the
42 east to more arid conditions in the west. (Bao et al., 2015; Zhao et al., 2002). The Asian monsoon

43 system **directly affects** the occurrence, intensity and severity of droughts and floods (Bao et al., 2015;
44 Cook et al., 2010; Liang et al., 2006; Wang et al., 2013; Wang et al., 2005; Zhao et al., 2002) that has a
45 devastating effects on human society and economy as well as natural ecosystems (Sun, 2007; Xu,
46 1998). For example, the drought in 2009 affected 81 million people in northeast China and more than
47 720,000 hectares of farmland suffered from water shortages ([http://www.chinadaily.com.cn/cndy/2009-](http://www.chinadaily.com.cn/cndy/2009-08/13/content_8562996.htm)
48 [08/13/content_8562996.htm](http://www.chinadaily.com.cn/cndy/2009-08/13/content_8562996.htm)). In addition, drought also can increase the occurrence of large wildfires.
49 Drought in Daxing'an Mountains, especially in spring or early summer, often leads to high risk of forest
50 wildfires (Sun, 2007). The fire caused by drought in northern Daxing'an Mountains in May 1987 killed
51 over 200 people and burned ~17,000 km² (Sun, 2007; Yao et al., 2017).

52 To better characterize current drought conditions and project those of the future, an improved
53 understanding of past drought variability and potential forcing mechanisms is necessary. However, the
54 shorter meteorological records in Daxing'an Mountains only started in 1950s limited our understanding
55 of the long-term regime of past droughts. Tree rings can serve as an important high resolution proxy for
56 long-term drought reconstructions (Cook et al., 2010; Dai, 2011; Pederson et al., 2013), and several
57 hydroclimate reconstructions (Bao et al., 2015; Lv and Wang, 2014; Wang and Lv, 2012) have been
58 conducted in northern China. Cook et al. (2010) also reconstructed the June-July-August Dai-PDSI in
59 534 grid points (Monsoon Asia Drought Atlas, MADA) in monsoon Asia using 327 tree-ring width
60 chronologies. However, some disagreements occur between the MADA and the tree-ring-based local
61 drought reconstructions and instrumental drought data, especially in eastern Asia, which might be an
62 insufficient tree-ring data in eastern Asia used in MADA (Li et al., 2015; Liu et al., 2017). Additional
63 drought reconstructions in eastern Asia are needed to gain a more thorough understanding of the

64 variability in the Asian Monsoon. Many researchers use the Palmer drought severity index (PDSI),
65 calculated from a water balance equation, incorporating air temperature and precipitation, to estimate
66 drought periodicity and intensity (Bao et al., 2015; Cook et al., 2010; Dai, 2011; Sun and Liu, 2013).
67 Here, we present a 260-year reconstruction of annual PDSI using tree-ring chronologies from the
68 Daxing'an Mountains to identify the timing of droughts and their correlation with eastern Mongolian
69 Plateaus climate as well as their potential forcing mechanisms.

70

71 2 Materials and methods

72 2.1 Study area

73 The Daxing'an Mountains, in northeast Inner Mongolia and northwest Heilongjiang Province, form
74 an important natural geographic divide between the Pacific Ocean and the north-western semi-arid
75 inland (Fig. 1). It is known to be a transition zone from the semi-humid to semi-arid region or from a
76 monsoon to non-monsoon climates (Zhao et al., 2002). The summer monsoons from the south-east are
77 blocked by the mountains and cannot penetrating further to the northwest. The western region is more
78 arid, and the eastern region is wetter. Summer weather is clarified by periodic incursions of warm,
79 humid air masses from low-latitude oceans, while dry and cold air in winter persists air masses invade
80 from high latitudes.

81 This study was conducted in high-latitude forests in the Daxing'an Mountains, northeast China. The
82 forests are dominated by Dahurian larch (*Larix gmelinii* Rupr.) and Scots pine (*Pinus sylvestris* L. var.
83 *mongolica* Litv.). Soils are predominantly brown coniferous and dark-brown forest peat (Xu, 1998).
84 Meteorological data was collected from stations nearest our sample sites (Xiaoergou station; Table S1).

85 The annual mean temperature ranges from -2.6 to 2.0 °C. The coldest and hottest month is January (-
86 39.5 °C) and July (32.8 °C), respectively. Annual precipitation ranges from 289 to 1000 mm (average
87 500 mm) with high interannual variations. Rain from June to August accounts for 68% of total annual
88 precipitation (Fig. 2). The relative humidity is low except for the growing season. Severe drought
89 occurs frequently, especially in spring and early summer (Sun, 2007), and leads to high fire risk. This
90 region has the highest average annual burned area in China (Sun, 2007).

91 2.2 Tree-ring data

92 Tree-ring cores were sampled from four Scots pine-dominated sites that are rarely disturbed in the
93 central Daxing'an Mountains in May 2011 and 2012. Each sampled tree was selected to avoid the
94 influence of identifiable stand disturbances (including animal and human disturbance, windstorm, snow,
95 and fire damage) and any obvious abnormal growth. The distance between sample sites is more than
96 100 km (Fig. 1). A total of 120 cores were obtained from living old trees at breast height (ca. 1.3 m)
97 (Table 1) using a 5.15-mm-diameter increment borer (500 mm length, two screws, Haglöf Sweden,
98 Långsele, Sweden). All cores were dried, mounted, surfaced, and cross-dated following standard
99 techniques of dendrochronology (Cook and Kairiukstis, 1990; Fritts, 1976). Ring widths were measured
100 with a precision of 0.001 mm using a Velmex measuring system (Velmex, Inc., Bloomfield, NY, USA).

101 The quality of cross-dating and measurement was evaluated using the COFECHA program
102 (Holmes, 1983). Two cores with weak correlation to the master chronology were excluded from further
103 analysis. Successively, the age-related trends were removed by fitting a cubic smoothing spline with a
104 50% frequency response cut-off at 2/3 of the series length using the ARSTAN program (Cook and
105 Kairiukstis, 1990). Tree-ring index was calculated as the ratio of the observed value to the estimated

106 growth curves. Autocorrelation was removed by autoregressive modelling, and the site chronology was
107 calculated using a bi-weighted robust mean (Cook and Kairiukstis, 1990).

108 Four chronologies have high values of standard deviation, mean sensitivity, mean series correlation
109 and agreement within population. They reflect high inter-annual variation and a strong common signal
110 and are excellent proxies for regional climate. Since the four chronologies fit well (Table 2), we merged
111 all samples to develop a single robust regional chronology (Fig. S1). Running RBAR (mean correlation
112 between series) and EPS (expressed population signal) statistics were calculated using a 51-year interval
113 of the chronology with a 25-year overlap to assess confidence in the chronology. RBAR averages
114 variance among ring width series in a chronology, which estimates chronology signal strength (Cook
115 and Kairiukstis, 1990). EPS estimates the degree to which the chronology represents a hypothetical
116 chronology based on a finite number of trees that match a hypothetically perfect chronology; EPS
117 values greater than 0.85 are generally considered acceptable threshold for a reliable chronology (Wigley
118 et al., 1984). The regional chronology spanned the period from 1725 to 2010, and the reliable interval
119 (EPS > 0.85) was 1751-2010 corresponding to eight trees (Fig. S1).

120 2.3 Climate and statistical analyses

121 Climate data were obtained from the National Meteorological Information Center
122 (<http://data.cma.cn/>). The weather station nearest to the sample sites is Xiaoergou (Table S1 and Fig. 1),
123 about 70-91 km away. Large-scale climate data (e.g. El Niño-Southern Oscillation, ENSO, Atlantic
124 Multidecadal Oscillation, AMO; Pacific Decadal Oscillation, PDO; North Atlantic Oscillation, NAO)
125 and high-resolution gridded climate data (Table S1; e.g. gridded temperature, precipitation and drought
126 indices) were downloaded from the website: <http://climexp.knmi.nl/>. Pearson correlation analysis was

127 conducted to estimate climate-growth relationships. The gridded climate dataset is much longer and has
128 higher homogeneity and coherency than station data (Fig. 2), the gridded monthly total precipitation
129 (CRU GPCC; Schneider et al. (2015)) and mean temperature (CRU TS3.23; Jones and Harris (2013))
130 nearest to our sites were used for climate response analyses. In addition, the nearby gridded monthly
131 Palmer Drought Severity Index (PDSI) data from Dai (2011) (Dai-PDSI, hereafter) was used to assess
132 the effects of drought. Correlation analyses between the regional chronology and monthly climatic
133 records were calculated from the previous July to the current July.

134 A linear regression model was used to reconstruct the drought variation, and a traditional split-
135 period calibration and verification method was applied to examine the model fitness (Fritts, 1976).
136 Statistical parameters included the R^2 , Sign test (ST), reduction of error (RE), coefficient of efficiency
137 (CE), product means test (PMT) and root mean square error (RMSE) (Cook and Kairiukstis, 1990;
138 Fritts, 1976). Spatial correlation of the measured and reconstructed drought variables with regional
139 gridded CRU-PDSI (Schrier et al., 2013) were performed to examine the spatial representativeness of
140 our reconstruction using the KNMI climate explorer. Local historical drought data recorded in
141 “Meteorological disasters dictionary of China” (Shen, 2008; Sun, 2007) were used to verify our PDSI
142 reconstruction.

143 We also carried out the superposed epoch analysis (SEA) between the nearby forest fire events and
144 the drought series to further validate the accuracy of our reconstruction because seasonal or annual
145 droughts are usually a key factor of forest fire severity in the Daxing'an Mountains (Shen, 2008; Sun,
146 2007). Two regional forest fire chronologies (Mengkeshan and Pangu) reconstructed by tree-ring scars
147 in nearby forests were used (Yao et al., 2017). The SEA was carried out using the software package

148 FHAES V2.0.0 (<https://www.frames.gov/partner-sites/fhaes/download-fhaes/>). In addition, the
149 consistency between our reconstruction and other local drought related time series including the gridded
150 **Standardized** Precipitation-Evapotranspiration Index (SPEI), Monsoon Asia Drought Atlas (MADA)
151 from [Cook et al. \(2010\)](#) (Cook-PDSI, hereafter) and Self-calibrating PDSI from [Schrier et al. \(2013\)](#)
152 (scPDSI, hereafter). **We also compared our reconstruction with the** nearby tree-ring-based hydroclimatic
153 reconstructions (the December-March precipitation reconstruction of the A'li River (AR) in the
154 Daxing'an Mountains from [Lv and Wang \(2014\)](#), the April-August SPEI reconstruction of the Hulun
155 Buir steppe (HB) on the east edge of Mongolian Plateaus in the western Daxing'an Mountains ([Bao et](#)
156 [al., 2015](#)), and the tree-ring-based streamflow reconstruction of Selenge River (SR) from [Davi et al.](#)
157 [\(2006\)](#) in the Mongolian Plateaus, Mongolia) to assess the reliability of reconstruction by filtering and
158 moving correlations.

159 To identify **spatiotemporal** patterns of drought variation in Northeast Asia and their relationship
160 with our reconstructed drought series, we analyzed the correlations between our series and other four
161 hydroclimatic reconstruction series in the Daxing'an Mountains and the Mongolian Plateau (Fig. 1). To
162 **better visualize the** comparison all series **described above** were standardized using Z-scores and
163 smoothed with a 21-year moving averaged to highlight low-frequency drought signals.

164 To evaluate the extreme dry and wet years in the historical period, we defined extreme dry and wet
165 years with the annual PDSI value being lower or higher than average ± 1.5 **standard deviation**. We
166 assessed the multiyear dry/wet periods based on the intensity (average departure values from the long-
167 term mean) and magnitude (cumulative departure values from the long-term mean).

168 A spectral analysis **was** applied to identify the periodicity of dry/wet **variation** and possible effects
169 of large-scale climate using **the** Multi-taper method (MTM) (Mann and Lees, 1996). To further confirm
170 the linkage between large-scale climate and regional drought, we **analyzed** their relationship with
171 Pearson correlation analysis. Teleconnections between the reconstructed drought **series** and the global
172 sea surface temperature ($0.5^{\circ} \times 0.5^{\circ}$) were carried out to verify the potential drivers of large-scale
173 climate, **such as ENSO, PDO and AMO**, on local drought. **To explore the linkages between the**
174 **reconstructed Dai-PDSI extreme events and atmospheric circulation patterns in Asia, the NCEP climate**
175 **data (Kalnay et al., 1996) were used to create January-December composite anomaly maps of the SSTs,**
176 **the 200-hPa geopotential height and vector wind in the wettest 10 years and driest 10 years during the**
177 **period 1948-2010.**

178

179 **3 Results**

180 **3.1 Tree growth-climate relationships**

181 The radial growth of Scots pine was significantly ($p < 0.05$) positively correlated with precipitation
182 in all months except the previous November and current February (Fig. 3a) and temperature from the
183 previous November to current May (except for the current April) (Fig. 3a). **The highest Pearson's**
184 **correlation coefficients occurred in October precipitation ($R = 0.35, p < 0.05$) and the previous**
185 **December mean temperature ($R = 0.35, p < 0.05$).** Radial growth of Scots pine in the Daxing'an
186 Mountains was influenced by both precipitation and temperature **simultaneously**, but the effects of
187 precipitation were stronger, **revealing the** annual precipitation sensitivity of Scots pine during the last
188 century (Fig. 3a). Furthermore, we calculated the correlation between the tree-ring index and Dai-PDSI

(common period of 1901-2010), which takes into account temperature and precipitation (Dai, 2011). Significant ($p < 0.05$) positive correlations between tree rings and PDSI was found in all months from the previous July to the current July (Fig. 3b). The correlation between tree growth and annual (Jan-Dec) average PDSI showed the highest correlation coefficient ($R = 0.62$, $p < 0.0001$, $n = 110$) among all seasonal PDSI compositions. The results confirmed that water availability had a significant limiting influence on Scots pine growth in the last century (Fig. 3).

3.2 PDSI reconstruction

The linear model for PDSI reconstruction is:

$$D_t = 6.69 I_t - 7.13, (R = 0.62, N = 100, F = 60.52, p < 0.0001) \quad (1)$$

where D_t is the annual PDSI and I_t is the tree-ring index at year t . The split calibration-verification test showed that the explained variances were high during the two calibration periods (1911-1960 and 1961-2010). The statistics of R , R^2 , ST, PMT are all significant at $p < 0.05$, which indicated that the model was reliable (Table 3). The most rigorous tests, RE and CE, were also positive for both verification periods (Cook and Kairiukstis, 1990; Fritts, 1976) (Table 3). For the calibration period 1911-2010, the reconstruction explained 38.2% of the PDSI variation (37.6% after accounting for the loss of degrees of freedom). These results suggest that the linear model is robust for PDSI reconstruction.

The instrumental and reconstructed PDSI of Central Daxing'an Mountains have similar trends and are parallel to each other during the calibration period (Fig. 4). However, the reconstructed PDSI did not capture the magnitude of extreme dry or wet conditions. Spatial correlation analysis showed that the

instrumental and reconstructed PDSI had a strong and similar spatial correlation pattern with the Northeast Asia gridded Dai-PDSI (Fig. 5).

3.3 Historical PDSI variability

The reconstructed annual PDSI with an 11-year moving average exhibited a mean of 0.48 and a standard deviation (SD) of ± 1.15 during the past 260 years (Fig. 4b). Reconstruction of the annual PDSI displayed strong interannual to decadal scale variability throughout the period 1751-2010. During the last 260 years, there were 22 extreme dry years (accounting for 8.5%) and 15 extreme wet years (5.8%) (Table 4). Most extreme dry years occurred in the 19th (12 years, accounting for 48%) and 20th (9 years, accounting for 36%) centuries, and most extreme wet years occurred in the 20th century (9 years, accounting for 60%). Among the extreme years, 1784, 1853, 1818, 1862 and 1863 were the five driest years, and 1998, 1952, 1770, 1993 and 1766 were the five wettest years (Table 4). We also found that many extreme dry or wet years occurred in succession, for example 1862 and 1863.

Compared with the severe single-year droughts, multi-year droughts had a greater effect on tree growth, and we defined the dry and wet periods as those when the 11-year moving average PDSI was more than 0.5 SD from the mean for at least 2 consecutive years. Four dry periods, AD 1751-1752, 1812-1817, 1847-1866 and 1908-1927, and four wet periods 1757-1771, 1881-1902, 1952-1955 and 1989-2004 were identified (Table 5). The dry periods of 1847-1866 and 1906-1927 were the longest, spanning 20 years, while the longest wet period, from 1881-1902, lasted for 22 years (Table 5). The multiyear drought in 1847-1866 was the most serious due to the long duration and intensity, and the period 1906-1927 was the second most significant drought (Table 5). Wet periods in 1757-1771 and 1989-2004 were the most remarkable in terms of their intensity and duration (Table 5).

230 Spectral analysis revealed that the **historical PDSI variation** in the Daxing'an Mountains **showed**
231 **several** significant (95% or 99% confidence level) periodicities at 46.5-78.7 (99%), 12, 5-6 (99%), and
232 2-3 (99%) years, which corresponded to significant cycle peaks **presented** in Fig. 6.

233

234 **4 Discussion**

235 **4.1 Climate-growth relationship**

236 Scots pine is an drought-tolerant species and drought stress is thought to be the **main climate factor**
237 **limiting its radial growth** in semi-arid or semi-humid regions, such as in the **Mongolian** Plateaus and
238 western Daxing'an Mountains (Bao et al., 2015; Davi et al., 2006; Liu et al., 2009; Pederson et al.,
239 2013). Previous dendroclimatic studies from these regions suggest that radial growth of Scots pine is
240 sensitive to humidity, precipitation or drought (e.g. PDSI, SPEI), and most analyses have reconstructed
241 hydroclimatic history (Bao et al., 2015; Liu et al., 2009). In these areas, the radial growth of Scots pine
242 usually has a typical **climate** (drought) response pattern with positive tree growth response to increasing
243 precipitation and **a** negative response to increasing temperature (Bao et al., 2015; Davi et al., 2006; Liu
244 et al., 2009). This typical drought response pattern **is** usually found in other drought or wetness tree ring
245 reconstructions (Li et al., 2016; Liu et al., 2017). In this study, the correlation between tree-ring **index**
246 and monthly precipitation and temperature revealed that the radial growth of Scots pine was mainly
247 limited by water, which is consistent with the physiological characteristics of tree species living in
248 **semi-arid** regions. A significant positive relationship between the tree-ring index and PDSI in all
249 months supported moisture as the main limiting factor of Scots pine radial growth (Fig. 3b).

250 The drought response was also found in Dahurian larch (Wang and Lv, 2012), another important
251 conifer tree species in the study area. However, the typical drought response to temperature was not
252 obvious, and the radial growth of Scots pine was not significantly negatively correlated with growing
253 season (July-September) temperature (Fig. 3a). On the contrary, a significant positive response of radial
254 growth to the non-growing season temperature was found. It is possible that higher winter temperatures
255 could protect dormant buds from frost damage (Chen et al., 2012). The positive correlation with spring
256 temperature could be due to earlier and larger snow melting, which supplies the spring soil water, and
257 eventually stimulates tree growth (Hollesen et al., 2015; Zhu et al., 2017). This unusual drought
258 response pattern might be due to the relatively humid climate and the northern latitude of our study
259 sites, where the positive effect of temperature was greater than the negative effect resulting from
260 drought stress (Wang and Song, 2011). Similar drought response patterns were also found in tree-ring-
261 based drought reconstructions in the middle Qilian Mountains (Sun and Liu, 2013) and the Tianshan
262 Mountains of western China (Chen et al., 2015).

263 4.2 Comparison with regional records

264 We used the local historical document records to verify our PDSI reconstruction for the timing of
265 extreme dry years or periods. During the last 260 years, 60.1% (13/22) of extreme dry years were noted
266 in historical documents (Shen, 2008; Sun, 2007). Tree rings cannot fully record the continuous drought
267 events (years) resulting in a limited percentage or correspondence. For example, only 1861 was
268 recorded in our reconstruction during the extreme drought period 1860-1865. Thus, some severe
269 drought events affect radial tree growth in some but not all years (Fritts, 1976). In addition, the lag
270 response of radial growth to climate (drought) might have a great contribution to unrecorded extreme

drought events (Fritts, 1976). For example, the local historical documents recorded the dry years of 1817 and 1855 that showed narrower rings or as an extreme dry event in the following year. Two multiyear droughts recorded in tree-rings, 1847-1866 and 1908-1927, can both be identified in historical documents (Shen, 2008; Sun, 2007). Moreover, the SEA between our reconstructed drought series and forest wildfire history revealed that a significant drop of PDSI values occurred during the year of the forest fire in Mengkeshan and Pangu (Fig. S2), further validating the accuracy of our reconstruction.

Spatial correlation analysis indicated a strong pattern between our reconstruction and gridded scPDSI in Northeast Asia (Fig. 7), and our reconstruction also represented drought/wet variations in surrounding geographic regions. During the common periods, our reconstruction shared a similar dry/wet fluctuation with precipitation of the A'li River (Wang and Lv, 2012) and SPEI of Hulun Buir steppe (Bao et al., 2015) both in the low and high frequency (Fig. 7b-d). Significant ($p < 0.05$) correlations among them were found in low and high frequency and some common dry/wet periods, highlighted in Fig. 7, which confirmed that our drought reconstruction could account for the most dry/wet variations in the Daxing'an Mountains.

It's important to note that our drought reconstruction and the MADA by Cook et al. (2010) from the same PDSI grid showed a complete opposite trend ($R_L = -0.19$; $p < 0.01$) in low frequency (Fig. 7). Negative correlations between the MADA and the SPEI ($R_L = -0.31$; $p = 0.03$) and scPDSI ($R_L = -0.126$; $p = 0.236$), positive correlations between our drought reconstruction and the SPEI ($R_L = 0.95$; $p < 0.01$) and scPDSI ($R_L = 0.807$; $p < 0.01$) were also found, although it had a seasonal difference with our drought reconstruction. These imply that the MADA by Cook et al. (2010) might be inaccurate or even reverse in the timing of dry/wet variations in the Daxing'an Mountains. Similar divergence of tree-

ring-based drought reconstruction between the MADA and individual sampling sites was also found by Li et al. (2015) from Guancen Mountain and Liu et al. (2017) from central Inner Mongolia. The insufficient spatiotemporal distribution of tree-ring networks, especially in eastern China, used in MADA might be the main reason for this divergence/inaccuracy (Cook et al., 2010; Li et al., 2015; Liu et al., 2017). Therefore, our drought reconstruction is necessary to improve our understanding of the East Asian Monsoon climate variability.

On a larger spatial scale, the streamflow reconstruction of Selenge River in the West-Central Mongolian Plateaus from Davi et al. (2006) presented a significant positive correlation with our drought reconstruction in low frequency ($R_L = 0.29$; $p < 0.01$) during the full periods. Our reconstructed PDSI also displayed some common variation trends for dry/wet periods with the reconstructed streamflow variations from the Selenge River (Davi et al., 2006), especially at the decadal scale. These relationships suggest that there are common drivers affecting dry/wet variations of the Daxing'an Mountains and the West-Central Mongolian Plateau, although there might be some discordance. Among those differences, the most obvious one is the completely different dry/wet variation trends among the Daxing'an Mountains (wetter), the West-Central Mongolian Plateaus (mild drier) as well as their transition zones, the East Mongolian Plateaus (Hulun Buir steppe; drier) since the late 1970s (Fig. 8a). Similar results were also found by Dai (2013), which presented a different dry-wet pattern under global warming using observations and models. In the Tibetan Plateau, Li et al. (2016) found moisture increases related to rapid warming (warm-wet). Although the reason for this divergence should be further studied, it might be related to the different response to the phase shift (negative to positive) of the PDO in 1976 and 1977 (Ma, 2007; Wang et al., 2014). Ma (2007) found that the positive PDO phase usually corresponds to the

313 drought period with warming and less precipitation, while the negative PDO phase often matches the
314 wet period with low temperature and more precipitation. **Simultaneously**, the drought trend caused by
315 the persistent significant warming in semi-arid or semi-humid regions might be more serious than in
316 humid regions (Dai, 2013). In addition, a different record of severe drought that occurred over a large
317 geographic area in northern Asia during the period 1920s-1930s, has been reported by many other
318 studies in north China (Bao et al., 2015; Chen et al., 2015; Liang et al., 2006; Liu et al., 2009). As
319 indicated by the tree-ring series, the drought event during **the period** 1920s-1930s in the Daxing'an
320 Mountains **was** more severe than **on** the East Mongolian Plateaus (the Hulun Buir steppe), which was
321 consistent with the result by Dong et al. (2013). The drought, however, was not found in the West-
322 Central Mongolian plateaus (the Selenge River). On the **contrary**, it was very **wet at** that time (Fig. 8).
323 Different spatial patterns of severe drought over northeast Asian might be associated with the intensity
324 and scope of the strong ENSO during this period (Dong et al., 2013).

325 **4.3 Linkages to the Pacific and Atlantic Oceans**

326 Spectral analysis revealed that several significant cycles existed in our drought **series** (Fig. 6). **The**
327 significant high-frequency 2.0–5.8-year periodicities were within the 2–7 year cycles of ENSO (Li et
328 al., 2013), so the **drought** variations in the Daxing'an Mountains might be related to ENSO. Similarly,
329 **the local dry-wet changes related to ENSO has been confirmed by other** tree-ring-based hydroclimatic
330 reconstructions in northeast China (Bao et al., 2015; Lv and Wang, 2014; Wang and Lv, 2012),
331 northwest China (Chen et al., 2015; Sun and Liu, 2013) and the Mongolian Plateaus (Davi et al., 2006).
332 **A** strong connection appears between our reconstruction and annual SSTs over the Pacific Ocean,
333 especially nearby the equator, the north Pacific, as well as the east and west coasts of the Pacific Ocean

(Fig. S3). The significant positive correlation between the Niño 3 index and the dry-wet index in both low and high frequencies (Table 6, Fig. 8b). also confirmed the potential links between ENSO and the dry/wet variations in the Daxing'an Mountains. Although the mechanisms need to be further studied, the close relationship between the oscillatory changes of North Atlantic SST and the Asian monsoon have been demonstrated (Zuo et al., 2013). ENSO might indirectly influence dry-wet changes in the Daxing'an Mountains by affecting the local climate (Shuai et al., 2016). Wang et al. (2013) found that the ENSO could potentially drive or affect the Asian monsoon, which in turn affects temperature and precipitation to drive local drought variations, as a possible driving mechanism (Fig. 9). Significant positive correlations between the Niño 3 index and local climate (temperature and precipitation) further confirms our inference (Table 6).

The 12-year cycle indicated that dry-wet changes in the Daxing'an Mountains might be influenced by solar activity (Shindell et al., 1999). Several previous studies have demonstrated that solar activity can influence the local dry-wet variations (Chen et al., 2015; Hodel et al., 2001; Sun and Liu, 2013). In northeastern China, Hong et al. (2001) also found the signals of solar activity in a 6000-year record of drought and precipitation. Significant positive correlations between the Total Solar Irradiance (TSI; reconstruction from IPCC AR5) and the dry-wet index in the Daxing'an Mountains in low and high frequencies, and between the TSI and the local climate (temperature and precipitation) further confirmed a possible relationship between solar activity and local drought (Table 6, Fig. 8b). Wang et al. (2005) found a potential link between the Asian monsoon and solar changes. Dry-wet changes in the Daxing'an Mountains might be driven by the Asian monsoon which is influenced by solar activities (Fig. 9).

355 Cycles of 46.5 - 48.8 years might be related to the PDO since it coincided with the 50-70 year cycle
356 of PDO (Macdonald and Case, 2005). This was verified by the strong connection between our drought
357 reconstruction and annual SSTs over the Pacific Ocean (Fig. S3). The cycles/signals of PDO widely
358 exist in most tree-ring-based drought reconstructions (Bao et al., 2015; Chen et al., 2015; Sun and Liu,
359 2013; Wang and Lv, 2012), and many studies have confirmed that PDO can influence drought
360 conditions in China (Bao et al., 2015; Cook et al., 2010; Ma, 2007). The potential linkages between the
361 PDO and local drought in the Daxing'an Mountains is further confirmed by the significant positive
362 correlations between the PDO index (Mann and Lees, 1996) and the dry-wet index in low and high
363 frequencies (Table 6, Fig. 8b). The positive/warm phase of PDO usually corresponds to the dry period,
364 while the negative/cold phase corresponds to the wetting period (Ma, 2007). For example, the severe
365 drought in 1920s-1930s corresponds to the PDO negative phase. Significant positive correlations
366 between the PDO index and local climate (Fig. 9) suggest that the PDO might affect the dry-wet
367 changes in the Daxing'an Mountains by regulating the intensity or location of the Asian monsoon (Bao
368 et al., 2015; Cook et al., 2010; Ma, 2007). Similar results were found in a nearby tree-ring-based
369 drought reconstruction (Bao et al., 2015).

370 The 73-years drought cycle might be derived from oscillatory changes in the North Atlantic SST-
371 (Knudsen et al., 2011). Spatial correlations between our drought series and annual SSTs also show a
372 strong teleconnection across the Atlantic Ocean (Fig. S3), which further confirmed potential linkages
373 between the North Atlantic SSTs and dry-wet changes in the Daxing'an Mountains. Although our
374 research area is far from the Atlantic, some studies have confirmed that large-scale climate oscillations
375 in the Atlantic Ocean (such as the Atlantic Multidecadal Oscillation (AMO), North Atlantic Oscillation

(NAO) as well as Summer NAO (SNAO)) could affect local climate and tree growth in China (Bates, 2007; Linderholm et al., 2011; Linderholm et al., 2013; Sun et al., 2008; Wang et al., 2011). Most tree-ring drought reconstructions also found the signals of oscillatory changes correlated with the North Atlantic SSTs (e.g. AMO, NAO and SNAO), such as in the Daxing'an Mountains (Lv and Wang, 2014; Wang and Lv, 2012), eastern Mongolian Plateaus (Bao et al., 2015; Liu et al., 2009), West-Central Mongolia (Davi et al., 2006), and northwest China (Chen et al., 2015; Sun and Liu, 2013). Furthermore, we also identified a significant negative/positive correlation between the dry-wet change in the Daxing'an Mountains and the AMO, NAO and SNAO index both in low or high frequency (Table 6, Fig. 8c). The strong AMO signal (Wang et al., 2011) and teleconnections with SNAO (Linderholm et al., 2013) also have been found in tree-ring widths of Scots pine in northeast China and east central Siberia during the last 400 years. These studies all confirmed that oscillatory changes in the North Atlantic SST (e.g. AMO, NAO and SNAO) could drive dry-wet changes in the Daxing'an Mountains. Although its mechanism needs to be further studied, the close relationship between the oscillatory changes in the North Atlantic SST and the Asian monsoon has been demonstrated. Recent studies have shown that the AMO (Wang et al., 2013), NAO (Feng and Hu, 2008) and SNAO (Linderholm et al., 2011) all could drive or affect the Asian monsoon. In this study, although only the AMO index was significantly correlated with local climate (Table 6, Fig. 8c), it also confirmed that the oscillatory changes in the North Atlantic SST, especially the AMO, could drive wet-dry changes in the Daxing'an Mountains by influencing the Asia Monsoon (Bao et al., 2015; Chen et al., 2015; Cook et al., 2010; Li et al., 2015; Linderholm et al., 2011; Sun et al., 2008).

396 Previous studies have found that drought variation in northeast Asia may be associated with Asian
397 monsoon activity (Bao et al., 2015; Chen et al., 2015; Cook et al., 2010; Li et al., 2015; Linderholm et
398 al., 2011; Sun et al., 2008). In wet years, the strengthened southerlies and easterlies entered inland
399 China associated with a positive pattern over northeast Asia and some negative height-anomaly centers
400 in west Russia and south Asia as well as the Indian and north Pacific oceans, which strengthened the
401 westerly circulation (Fig. 10a, c). In dry years, however, strengthened southerlies and south-westerlies
402 entered northeast China associated with a positive pattern over east Asia and western Russia, and some
403 negative height-anomaly centers in southern Russia and south Asia as well as the Indian and south
404 Pacific oceans (Fig. 10a, c).

405 The composite of 200-hPa geopotential height of the most humid 10 years (positive anomaly) in the
406 central-north Daxing'an Mountains is opposite to that of the most arid 10 years (negative anomaly)
407 (Fig. 10c, d). Positive and negative SST anomalies were also found in the western and northern Pacific
408 Ocean during the wettest and driest years (Fig. 10e, f). In the wet years, abundant moisture is
409 transported from the Pacific Ocean through Mongolian Plateau to the Daxing'an Mountains *via* the
410 strong east Asian monsoon's southeasterly moisture flux joined with a strong Westerly circulation (Fig.
411 10a). This negative anomaly combined with positive SST in the western and northern Pacific Ocean
412 lead to an enhanced dry jet (south-westerlies) across/toward the Daxing'an Mountains (Fig. 10b, c, e).
413 Several studies have reported that the dry and wet variations in northeast Asia are strongly linked with
414 the Asian monsoon and SSTs in the Pacific and Atlantic oceans (Bao et al., 2015; Chen et al., 2015;
415 Cook et al., 2010; Li et al., 2015; Linderholm et al., 2011). In addition, the potential evaporation pattern

416 in the Daxing'an Mountains is extremely low in the wettest years, and it also supports the above
417 remote-connection assumptions (Fig. S4).

418 5 Conclusion

419 We developed a 260-year (1751 to 2010) tree-ring chronology of Scots pine (*Pinus sylvestris* L.
420 var. *mongolica* Litv.) from four sample sites in the central Daxing'an Mountains, northeast China.
421 Radial growth of Scots pine was mainly limited by water availability ($R = 0.62$, $p < 0.01$). A 260-year
422 dry-wet change history was reconstructed, and the reconstruction equation explained 38.2 % of the
423 PDSI variance for the period 1911-2010. Four dry and wet periods were found in the past 260 years,
424 respectively. The extreme dry years in our reconstruction series are consistent with the local historical
425 document records. Our reconstruction series revealed the dry-wet changes in the Daxing'an Mountains,
426 and also was the representative of the dry-wet changes in the West-Central Mongolian Plateaus,
427 especially at the decadal scale. Droughts during the 1920s-1930s in the Daxing'an Mountains were
428 more severe than in surrounding areas. In addition, the reconstruction series showed that the Daxing'an
429 Mountains is getting warm and wet since the late 1970s. This is not in line with the situation in the
430 Mongolian Plateaus, especially in the transition zones. Our reconstruction also suggests that the MADA
431 by Cook et al. (2010) may not be accurate in the Daxing'an Mountains likely due to the insufficient
432 spatiotemporal distribution of the tree-ring data in this area. Overall, drought variability in the central
433 Daxing'an Mountains and its relationship with the surrounding areas might be driven by climate
434 oscillations of the Pacific and Atlantic Oceans (e.g., ENSO, PDO, AMO, NAO and SNAO). These
435 large-scale climate oscillations affect the Asian monsoon and then lead to dry and wet changes in
436 Daxing'an Mountains.

438 **Acknowledgements** This research was supported by the Key Project of the China National Key
439 Research and Development Program (2016YFA0600800), the National Natural Science Foundation of
440 China (Nos. 41471168 and 31770490), the Program for Changjiang Scholars and Innovative Research
441 Team in University (IRT-15R09), and the Fundamental Research Funds for the Central Universities
442 (2572016AA32). We also thank Yongxian Lu and Lei Zhang of Northeast Forestry University for their
443 assistance in the field.

444

445 **References**

446 Bao, G., Liu, Y., Liu, N., and Linderholm, H. W.: Drought variability in eastern Mongolian Plateau and
447 its linkages to the large-scale climate forcing, *Climate Dynamics*, 44, 717-733, 2015.

448 Bates, G. T.: Influence of the Atlantic multidecadal oscillation on the winter climate of East China,
449 *Advances in Atmospheric Sciences*, 24, 126-135, 2007.

450 Chen, F., Yuan, Y., Wei, W., Yu, S., Zhang, T., Shang, H., Zhang, R., Qin, L., and Fan, Z.: Tree-ring
451 recorded hydroclimatic change in Tianshan mountains during the past 500 years, *Quaternary*
452 *International*, 358, 35-41, 2015.

453 Chen, F., Yuan, Y., Wei, W., Yu, S., Zhang, T.: Tree ring-based winter temperature reconstruction for
454 Changting, Fujian, subtropical region of Southeast China, since 1850: linkages to the Pacific
455 Ocean. *Theoretical and Applied Climatology*, 109, 141-151, 2012.

456 Cook, E. R., and Kairiukstis, L. A.: *Methods of dendrochronology: applications in the environmental*
457 *sciences*, Springer, 1990.

458 Cook, E. R., Anchukaitis, K. J., Buckley, B. M., D'Arrigo, R. D., Jacoby, G. C., and Wright, W. E.:
 459 Asian monsoon failure and megadrought during the last millennium, *Science*, 328, 486-489, 2010.
 460 Dai, A.: Characteristics and trends in various forms of the Palmer Drought Severity Index during 1900-
 461 2008, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
 462 Dai, A.: Increasing drought under global warming in observations and models, *Nature Climate Change*,
 463 3, 52-58, 2013.
 464 Davi, N., Jacoby, G., Curtis, A., and Baatarbileg, N.: Extension of Drought Records for Central Asia
 465 Using Tree Rings: West-Central Mongolia, *Journal of Climate*, 19, 288-299, 2006.
 466 Dong, A., Wang, J., and Li, Y.: The Disaster and Causes of Serious Drought in Seven Provinces of
 467 Northern China in 1920, *Journal of Arid Meteorology*, 31, 750-755, 2013. Feng, S., and Hu, Q.:
 468 How the North Atlantic Multidecadal Oscillation may have influenced the Indian summer
 469 monsoon during the past two millennia, *Geophysical Research Letters*, 35, 2008.
 470 Fritts, H. C.: *Tree rings and climate*, Elsevier, Amsterdam, Netherland, 1976.
 471 Hodell, D. A., Brenner, M., Curtis, J. H., and Guilderson, T.: Solar forcing of drought frequency in the
 472 Maya lowlands, *Science*, 292, 1367-1370, 2001.
 473 Hollesen, J., Buchwal, A., Rachlewicz, G., Hansen, B. U., Hansen, M. O., Stecher, O., and Bo, E.:
 474 Winter warming as an important co-driver for Betulanana growth in western Greenland during the
 475 past century, *Global Change Biology*, 21, 2410--2423, 2015.
 476 Holmes, R. L.: Computer-assisted quality control in tree-ring dating and measurement, *Tree-ring*
 477 *Bulletin*, 43, 69-78, 1983.
 478 Hong, Y., Wang, Z., Jiang, H., Lin, Q., Hong, B., Zhu, Y., Wang, Y., Xu, L., Leng, X., and Li, H.: A

479 6000-year record of changes in drought and precipitation in northeastern China based on a $\delta^{13}\text{C}$
480 time series from peat cellulose, *Earth and Planetary Science Letters*, 185, 111-119, 2001.

481 Jones, P., and Harris, I.: CRU TS3. 20: Climatic Research Unit (CRU) Time-Series (TS) Version 3.20 of
482 High Resolution Gridded Data of Month-by-Month Variation in Climate (January 1901-December
483 2011), NCAS British Atmospheric Data Centre, 2013.

484 Knudsen, M. F., Seidenkrantz, M.-S., Jacobsen, B. H., and Kuijpers, A.: Tracking the Atlantic
485 Multidecadal Oscillation through the last 8,000 years, *Nature Communications*, 2, 178, 2011.

486 Li, J., Xie, S., Cook, E. R., Morales, M. S., Christie, D. A., Johnson, N. C., Chen, F., D'Arrigo, R.,
487 Fowler, A. M., and Gou, X.: El Niño modulations over the past seven centuries, *Nature Climate*
488 *Change*, 3, 822-826, 2013.

489 Li, J., Shi, J., Zhang, D., Yang, B., Fang, K., and Yue, P.: Moisture increase in response to high-altitude
490 warming evidenced by tree-rings on the southeastern Tibetan Plateau, *Climate Dynamics*, 1-12,
491 2016.

492 Li, Q., Liu, Y., Song, H., Yang, Y., and Zhao, B.: Divergence of tree-ring-based drought reconstruction
493 between the individual sampling site and the Monsoon Asia Drought Atlas: an example from
494 Guancen Mountain, *Science Bulletin*, 1688-1697, 2015.

495 Liang, E., Liu, X., Yuan, Y., Qin, N., Fang, X., Lei, H., Zhu, H., Wang, L., and Shao, X.: The 1920S
496 drought recorded by tree rings and historical documents in the semi-arid and arid areas of northern
497 China, *Climatic Change*, 79, 403-432, 2006.

498 Linderholm, H. W., Ou, T., Jeong, J. H., Folland, C. K., Gong, D., Liu, H., Liu, Y., and Chen, D.:
499 Interannual teleconnections between the summer North Atlantic Oscillation and the East Asian

500 summer monsoon, Journal of Geophysical Research: Atmospheres, 116, 2011.

501 Linderholm, H. W., Seim, A., Ou, T., Jeong, J.-H., Liu, Y., Wang, X., Bao, G., and Folland, C.:
502 Exploring teleconnections between the summer NAO (SNAO) and climate in East Asia over the
503 last four centuries-A tree-ring perspective, Dendrochronologia, 31, 297-310, 2013.

504 Liu, Y., Bao, G., Song, H., Cai, Q., and Sun, J.: Precipitation reconstruction from Hailar pine (*Pinus*
505 *sylvestris* var. *mongolica*) tree rings in the Hailar region, Inner Mongolia, China back to 1865 AD,
506 Palaeogeography, Palaeoclimatology, Palaeoecology, 282, 81-87, 2009.

507 Liu, Y., Zhang, X., Song, H., Cai, Q., Li, Q., Zhao, B., Liu, H., and Mei, R.: Tree-ring-width-based
508 PDSI reconstruction for central Inner Mongolia, China over the past 333 years, Climate Dynamics,
509 48, 867-879, 2017.

510 Lv, S., and Wang, X. C.: Growth-climate response and winter precipitation reconstruction of *Pinus*
511 *sylvestris* var. *mongolicain* A'li River of Greater Khingan Range, Journal of Northeast Normal
512 University (Natural Science Edition), 46, 110-116, 2014.

513 Ma, Z.: The interdecadal trend and shift of dry/wet over the central part of North China and their
514 relationship to the Pacific Decadal Oscillation (PDO), Chinese Science Bulletin, 52, 2130-2139,
515 2007.

516 Macdonald, G. M., and Case, R. A.: Variations in the Pacific Decadal Oscillation over the past
517 millennium, Geophysical Research Letters, 32, 93-114, 2005.

518 Mann, M. E., and Lees, J. M.: Robust estimation of background noise and signal detection in climatic
519 time series, Climatic Change, 33, 409-445, 1996.

520 Mann, M. E., Zhang, Z., Rutherford, S., Bradley, R. S., Hughes, M. K., Shindell, D., Ammann, C.,

521 Faluvegi, G., and Ni, F.: Global signatures and dynamical origins of the Little Ice Age and
 522 medieval climate anomaly, *Science*, 326, 1256-1260, 2009.

523 Pederson, N., Leland, C., Nachin, B., Hessler, A., Bell, A., Martin-Benito, D., Saladyga, T., Suran, B.,
 524 Brown, P., and Davi, N. K.: Three centuries of shifting hydroclimatic regimes across the
 525 Mongolian Breadbasket, *Agricultural and Forest Meteorology*, 178, 10-20, 2013.

526 Schneider, U., Becker, A., Finger, P., Meyer-Christoffer, A., Rudolf, B., and Ziese, M.: GPCC full data
 527 reanalysis version 7.0 at 0.5: monthly land-surface precipitation from rain-gauges built on GTS-
 528 based and historic data, *FD_M_V6_050*, 10.5676/DWD_GPCC_FD_M_V7_050, 2015.

529 Schrier, G., Barichivich, J., Briffa, K. R., and Jones, P. D.: A scPDSI - based global data set of dry and
 530 wet spells for 1901–2009, *Journal of Geophysical Research: Atmospheres*, 118, 4025-4048,
 531 2013.

532 Shen, J.: Meteorological disasters dictionary of China: vol. Inner Mongolia, China Meteorological
 533 Press, Beijing (in Chinese), 2008.

534 Shindell, D., Rind, D., Balachandran, N., Lean, J., and Lonergan, P.: Solar cycle variability, ozone, and
 535 climate, *Science*, 284, 305-308, 1999.

536 Shuai, J., Zhang, Z., Tao, F., and Shi, P.: How ENSO affects maize yields in China: understanding the
 537 impact mechanisms using a process - based crop model, *International Journal of Climatology*, 36,
 538 424-438, 2016.

539 Sun, J., Wang, H., and Yuan, W.: Decadal variations of the relationship between the summer North
 540 Atlantic Oscillation and middle East Asian air temperature, *Journal of Geophysical Research:*
 541 *Atmospheres*, 113, **D15107**, 2008.

542 Sun, J., and Liu, Y.: Drought variations in the middle Qilian Mountains, northeast Tibetan Plateau, over
543 the last 450 years as reconstructed from tree rings, *Dendrochronologia*, 31, 279-285, 2013.

544 Sun, Y.: Meteorological disasters dictionary of China: vol. Heilongjiang, China Meteorological Press,
545 Beijing, China, 2007.

546 Trouet, V., Esper, J., Graham, N. E., Baker, A., Scourse, J. D., and Frank, D. C.: Persistent positive
547 North Atlantic Oscillation mode dominated the medieval climate anomaly, *Science*, 324, 78-80,
548 2009.

549 Wang, B., Liu, J., Kim, H.-J., Webster, P. J., Yim, S.-Y., and Xiang, B.: Northern Hemisphere summer
550 monsoon intensified by mega-El Niño/southern oscillation and Atlantic multidecadal oscillation,
551 *Proceedings of the National Academy of Sciences*, 110, 5347-5352, 2013.

552 Wang, X., Brown, P. M., Zhang, Y., and Song, L.: Imprint of the Atlantic multidecadal oscillation on
553 tree-ring widths in Northeastern Asia since 1568, *PLoS ONE*, 6, e22740, 2011.

554 Wang, X., and Song, L.: Climate-tree growth relationships of *Pinus sylvestris* var. *mongolica* in the
555 northern Daxing'an Mountains, China, *Chinese Journal of Plant Ecology*, 35, 294-302, 2011.

556 Wang, X., and Lv, S.: Tree-ring reconstructions of January-March streamflow in the upper Nenjiang
557 River since 1804, China, *Arid Land Geography*, 4, 35, 537-544, 2012.

558 Wang, X., Li, Z., and Ma, K.: Decreased sensitivity of tree growth to temperature in Southeast China
559 after the 1976/'77 regime shift in Pacific climate, *Sains Malaysiana*, 43, 9-19, 2014.

560 Wang, Y., Cheng, H., Edwards, R. L., He, Y., Kong, X., An, Z., Wu, J., Kelly, M. J., Dykoski, C. A., and
561 Li, X.: The Holocene Asian monsoon: links to solar changes and North Atlantic climate, *Science*,
562 308, 854-857, 2005.

563 Wigley, T. M., Briffa, K. R., and Jones, P. D.: On the average value of correlated time series, with
564 applications in dendroclimatology and hydrometeorology, *Journal of Climate and Applied*
565 *Meteorology*, 23, 201-213, 1984.

566 Xu, H.: *Daxing'an Mountains Forests in China*, Science Press, Beijing, China, 1998.

567 Yao, Q., Brown, P. M., Liu, S., Rocca, M. E., Trouet, V., Zheng, B., Chen, H., Li, Y., Liu, D., and Wang,
568 X.: Pacific - Atlantic Ocean influence on wildfires in northeast China (1774 to 2010), *Geophysical*
569 *Research Letters*, 44, 1025-1033, 2017.

570 Zhao, H., Zhao, X., Zhang, T., and Zhou, R.: Boundary line on agro-pasture zigzag zone in north China
571 and its problems on eco-environment, *Advances in Earth Science*, 17, 739-747, 2002.

572 Zhu, L., Li, Z., Zhang, Y., and Wang, X.: A 211-year growing season temperature reconstruction using
573 tree-ring width in Zhangguangcai Mountains, Northeast China: linkages to the Pacific and Atlantic
574 Oceans, *International Journal of Climatology*, 37, 3145-3153, 2017.

575 Zuo, J., Li, W., Sun, C., Xu, L., Ren, H.: Impact of the North Atlantic sea surface temperature tripole on
576 the East Asian summer monsoon, *Advances in Atmospheric Sciences*, 30, 1173-1186, 2013.

577

578

579 **Table 1** Site description and statistical characteristic of the *Pinus sylvestris* var. *mongolica*
 580 dendrochronologies in the Daxing'an Mountains.

Site	Latitude (N)	Longitude (E)	Elevation (m)	Number of trees	Time span	EPS ^a	RBAR ^b
Keyihe (KY)	50°39'44.8"	122°23'13.3"	550	36	1725-2010	0.93	0.57
Alihe (AL)	50°38'37.7"	124°28'28.7"	380	32	1742-2010	0.87	0.52
Ganhe (GH)	50°43'51.9"	123°05'56.9"	760	19	1793-2010	0.88	0.59
Jinhe (JH)	50°26'16.7"	121°59'46.7"	830	33	1769-2011	0.95	0.61
Region (RE)	-	-	-	-	1725-2010	0.97	0.51

581 ^a Expressed population signal statistic.

582 ^b RBAR = the mean correlation coefficient between all tree-ring series used in a chronology.

583 **Table 2** Five-chronology correlation matrix over the common period 1793-2010.

	AL	GH	JH	Region
KY	0.38**	0.46**	0.55**	0.81**
AL		0.33**	0.32**	0.68**
GH			0.32**	0.72**
JH				0.74**

584 ** Significance level ($p < 0.01$). The site codes are identical with those in Table 1.

585 **Table 3** Calibration and verification statistics of the PDSI reconstruction

Calibration	<i>R</i>	Verification	<i>R</i> ²	RE	CE	ST	PMT	RMSE
1911-2010	0.62**	-	-	0.38	-	(74, 26)**	8.04**	1.4
1961-2010	0.53**	1911-1960	0.47**	0.47	0.47	(39, 11)**	5.24**	1.34
1911-1960	0.69**	1961-2010	0.28**	0.28	0.25	(34, 16)*	6.23**	1.25

586 * = $p < 0.05$, ** = $p < 0.01$. RE-reduction of error, CE-coefficient of efficiency, ST-sign test, PMT-
587 product means test, RMSE-root mean square error.

588

589 **Table 4** Reconstructed extreme dry/wet years and annual PDSI of the Daxing'an Mountains.

Dry year (Rank)	PDSI	Dry year (Rank)	PDSI	Wet year (Rank)	PDSI
1784 (1)	-3.574	1909 (16)	-2.484	1998 (1)	2.521
1853 (2)	-3.315	1916 (17)	-2.479	1952 (2)	2.091
1818 (3)	-3.238	1854 (18)	-2.405	1770 (3)	2.020
1862 (4)	-3.006	1865 (19)	-2.314	1993 (4)	2.011
1863 (5)	-3.001	1861 (20)	-2.310	1766 (5)	1.790
1918 (6)	-2.991	1864 (21)	-2.283	1897 (6)	1.728
1919 (7)	-2.977	1856 (22)	-2.275	1996 (7)	1.663
1915 (8)	-2.882			1899 (8)	1.655
1917 (9)	-2.777			1755 (9)	1.576
1852 (10)	-2.733			1999 (10)	1.548
1851 (11)	-2.716			2000 (11)	1.488
1860 (12)	-2.695			1997 (12)	1.451
1967 (13)	-2.671			1994 (13)	1.422
1925 (14)	-2.660			1769 (14)	1.405
1911 (15)	-2.581			1764 (15)	1.279

590

591

592 **Table 5** The long-term droughts and pluvial in the Central Daxing'an Mountains during the last 260
593 years.

Year	Dry/Wet	Duration (year)	Magnitude (sum of PDSI)	Intensity (mean PDSI)
1751-1752	Dry	2	-2.36	-1.33
1757-1771	Wet	15	19.52	1.30
1812-1817	Dry	6	-3.73	-0.62
1847-1866	Dry	20	-32.70	-1.64
1881-1902	Wet	22	18.98	0.86
1906-1927	Dry	20	-31.79	-1.59
1952-1955	Wet	4	2.78	0.69
1989-2004	Wet	16	19.67	1.23

594

Table 6 Correlation coefficients between the large-scale climate index (AMO, PDO, NAO, SNAO, TSI, and Niño 3) and the local annual mean temperature, total precipitation, instrumental Dai-PDSI as well as the Z-score of dry-wet variation in the Daxing'an Mountains (DM_{Z-score})

	Temperature			Precipitation			Dai-PDSI			DM _{Z-score}		
	<i>R</i>	<i>p</i>	N	<i>R</i>	<i>p</i>	N	<i>R</i>	<i>p</i>	N	<i>R</i>	<i>p</i>	N
AMO	0.44**	0.00	106	0.30**	0.00	106	0.44**	0.00	96	0.35**	0.00	282
PDO	0.46**	0.00	106	0.39**	0.00	106	0.51**	0.00	96	0.34**	0.00	282
NAO	0.17	0.08	106	-0.04	0.71	106	-0.08	0.43	96	-0.21**	0.00	282
SNAO	0.22*	0.02	110	0.08	0.39	110	0.08	0.42	100	0.13*	0.05	246
TSI	0.23*	0.01	110	0.35**	0.00	110	0.34**	0.00	100	0.12*	0.04	286
Niño 3	0.34**	0.00	106	0.26**	0.01	106	0.28**	0.01	96	0.14**	0.02	282

Note: The AMO, PDO, NAO, SNAO, TSI and Niño 3 is the Atlantic Multidecadal Oscillation reconstruction from Mann et al. (2009), the Pacific Decadal Oscillation reconstruction from Mann et al. (2009), the Multi-decadal Winter North Atlantic Oscillation reconstruction from Trouet et al. (2009), the summer NAO based on the 20C reanalysis sea-level pressure reconstruction (SNAO), the Total Solar Irradiance reconstruction from IPCC AR5 and the Niño 3 reconstruction from Mann et al. (2009), respectively. All above data were downloaded from <http://climexp.knmi.nl/>. * $p < 0.05$, ** $p < 0.01$

604 **Figures captions**

605 **Fig. 1** Sampling sites and weather station distribution map. The red circles, start and black square are
606 the sampled sites, PDSI grid point and the weather station, respectively. The red box represents the
607 northern Daxing'an Mountains (this study). The blue and green box represents the east and west-
608 central Mongolian Plateau, respectively.

609 **Fig. 2** Monthly total precipitation (P) and mean temperature (T) at the Xiaoergou (a) meteorological
610 station (1957-2014) and grids (b) data (1901-2014); the annual total precipitation (c), and the
611 annual mean temperature (d) and PDSI (e). The dashed line indicates the linear fitting values.

612 **Fig. 3** Pearson correlation coefficients between tree-ring index of *Pinus sylvestris* var. *mongolica* and
613 the monthly total precipitation, mean temperature (a) and Dai-PDSI (b). Significant correlations (p
614 < 0.05) are indicated by above or below the 95% confidence line (dash line). The minus sign “-” in
615 abscissa represents the previous year, for example, “-7” represents the previous July.

616 **Fig. 4** The reconstruction PDSI series in the Daxing'an Mountains, northeast China. (a) Comparison of
617 the observed (pink line) and reconstructed (red line) annual PDSI during the calibration period
618 1911-2010; (b) The reconstruction series of annual PDSI, plotted annually from 1725 to 2010
619 (green line), along with a smoothed 11-year moving average (red bold line); Blue filled triangles
620 indicate a forest fire in nearby area reconstructed from tree-ring fire scars in Mengkeshan (higher)
621 and Pangu (lower), northern Daxing'an Mountains.

622 **Fig. 5** Spatial correlation fields between (a) the instrumental and (b) reconstructed annual Dai-PDSI for
623 the Daxing'an Mountains and the regional Dai-PDSI during the period 1911-2010
624 (<http://climexp.knmi.nl/>). The blue circle is the reconstructed PDSI grid.

625 **Fig. 6** Multi-taper method power spectrum of the reconstructed PDSI during the period 1751-2010. The
626 95% and 99% confidence level relative to red noise are shown and the numbers refer to the
627 significant period in years.

628 **Fig. 7** Comparisons of (a) the drought reconstruction derived from the Monsoon Asia Drought Atlas
629 (MADA, Cook et al. (2010)), (b) the winter precipitation reconstruction of the A'li River in
630 northeastern China (AR, Lv and Wang (2014)), (c) the mean annual PDSI reconstruction in the
631 Daxing'an Mountains (TS, in this study), (d) the April-August SPEI reconstruction of the Hulun
632 Buir steppe in eastern Mongolian Plateaus (HB, Bao et al. (2015)) and (e) the April-October
633 streamflow reconstruction of the Selenge River in northeastern Mongolia(SR, Davi et al. (2006)).
634 All above series were standardized using Z-scores (high frequency) and then smoothed with a 21-
635 year moving average (low frequency; red bold line). Blue and red shade areas represent a
636 consistent period of drought and wetness, respectively. Correlation coefficients between our
637 reconstruction series and other series in low (R_L) and high (R_H) frequency are shown on the
638 diagram. $** p < 0.01$

639 **Fig. 8** Comparisons of the drought reconstruction and other large-scale climate oscillations. (a) the dry-
640 wet changes in the Daxing'an Mountains (DM, the average of our reconstruction and the
641 precipitation reconstruction of the A'li River), the Mongolian Plateaus (MP, the streamflow
642 reconstruction of the Selenge River) as well as their transition zones (TZ, the SPEI reconstruction
643 of the Hulun Buir steppe); (b) the drought reconstruction in the Daxing'an Mountains (DM), the
644 Pacific Decadal Oscillation (PDO) and the Niño 3 index reconstruction from Mann et al. (2009)
645 (Nino3) as well as the Total Solar Irradiance reconstruction from IPCC AR5 (TSI); (c) the drought

reconstruction in the Daxing'an Mountains (DM), the Atlantic Multidecadal Oscillation reconstruction from Mann et al. (2009) (AMO), the Multi-decadal Winter North Atlantic Oscillation reconstruction by Trouet et al. (2009)(NAO) and the summer NAO based on the 20C reanalysis sea-level pressure reconstruction (SNAO). All above series were standardized using Z-scores and then smoothed with a 21-year moving averaged to highlight low-frequency drought signals. Significant correlation coefficients (** $p < 0.01$) are listed in the figure.

Fig. 9 Spatial correlations between the annual East Asian monsoon index and the local (a) temperature, (b) precipitation and (c) scPDSI from AD 1948 to 2010.

Fig. 10 Composite anomaly maps of the 200-hPa vector wind and geopotential height, and the SSTs (from January to December) for the 10 wettest (a, b and e) and 10 driest (c, d and f) years of the Dai-PDSI reconstruction during the period 1948-2010.






Fig. 1 Sampling sites and weather station distribution map. The red circles, star and black square are the sampled sites, PDSI grid point and the weather station, respectively. The red box represents the northern Daxing'an Mountains (this study). The blue and green box represents the east and west-central Mongolian Plateau, respectively.




664

665 **Fig. 2** Monthly total precipitation (P) and mean temperature (T) at the Xiaoergou (a) meteorological
 666 station (1957-2014) and grids (b) data (1901-2014); the annual total precipitation (c), and the annual mean
 667 temperature (d) and PDSI (e). The dashed line indicates the linear fitting values.



668

669 **Fig. 3** Pearson correlation coefficients between tree-ring index of *Pinus sylvestris* *var. mongolica* and
 670 the monthly total precipitation, mean temperature (a) and Dai-PDSI (b). Significant correlations ($p <$
 671 0.05) are indicated by above or below the 95% confidence line (dash line). The minus sign “-” in
 672 abscissa represents the previous year, for example, “-7” represents the previous July.



673

674 **Fig. 4** The reconstruction PDSI series in the Daxing'an Mountains, northeast China. (a) Comparison of
 675 the observed (pink line) and reconstructed (red line) annual PDSI during the calibration period 1911–
 676 2010; (b) The reconstruction series of annual PDSI, plotted annually from 1725 to 2010 (green line),
 677 along with a smoothed 11-year moving average (red bold line); Blue filled triangles indicate a forest fire
 678 in nearby area reconstructed from tree-ring fire scars in Mengkeshan (higher) and Pangu (lower),
 679 northern Daxing'an Mountains.






Fig. 5 Spatial correlation fields between (a) the **instrumental** and (b) reconstructed annual Dai-PDSI for the Daxing'an Mountains and the regional **Dai-PDSI** during the period 1911-2010 (<http://climexp.knmi.nl/>). The **blue circle** is the reconstructed PDSI grid.



684


685 **Fig. 6** Multi-taper method power spectrum of the reconstructed PDSI during the period 1751-2010. The
 686 95% and 99% confidence level relative to red noise are shown and the numbers refer to the significant
 687 period in years.



688

689 **Fig. 7** Comparisons of (a) the drought reconstruction derived from the Monsoon Asia Drought Atlas
 690 (MADA, Cook et al. (2010)), (b) the winter precipitation reconstruction of the A'li River in


691 northeastern China (AR, Lv and Wang (2014)), (c) the mean annual PDSI reconstruction in the
692 Daxing'an Mountains (TS, in this study), (d) the April-August SPEI reconstruction of the Hulun Buir
693 steppe in eastern Mongolian Plateaus (HB, Bao et al. (2015)) and (e) the April-October streamflow
694 reconstruction of the Selenge River in northeastern Mongolia(SR, Davi et al. (2006)). All above series
695 were standardized using Z-scores (high frequency) and then smoothed with a 21-year moving average
696 (low frequency; red bold line). **Blue and red shade areas represent a consistent period of drought and**
697 **wetness, respectively**. Correlation coefficients between our reconstruction series and other series in low
698 (R_L) and high (R_H) frequency are **shown on the diagram**. ** $p < 0.01$



699

700 **Fig. 8** Comparisons of the drought reconstruction and other large-scale climate oscillations. (a) the dry-
 701 wet changes in the Daxing'an Mountains (DM, the average of our reconstruction and the precipitation
 702 reconstruction of the A'li River), the Mongolian Plateaus (MP, the streamflow reconstruction of the
 703 Selenge River) as well as their transition zones (TZ, the SPEI reconstruction of the Hulun Buir steppe);
 704 (b) the drought reconstruction in the Daxing'an Mountains (DM), the Pacific Decadal Oscillation
 705 (PDO) and the Niño 3 index reconstruction from Mann et al. (2009) (Nino3) as well as the Total Solar
 706 Irradiance reconstruction from IPCC AR5 (TSI); (c) the drought reconstruction in the Daxing'an
 707 Mountains (DM), the Atlantic Multidecadal Oscillation reconstruction from Mann et al. (2009) (AMO),


708 the Multi-decadal Winter North Atlantic Oscillation reconstruction by Trouet et al. (2009)(NAO) and
709 the summer NAO based on the 20C reanalysis sea-level pressure reconstruction (SNAO). All above
710 series were standardized using Z-scores and then smoothed with a 21-year moving averaged to highlight
711 low-frequency drought signals. Significant correlation coefficients (** $p < 0.01$) are listed in the figure.



712

713 **Fig. 9** Spatial correlations between the annual East Asian monsoon index and the local (a) temperature,

714 (b) precipitation and (c) scPDSI from AD 1948 to 2010.



715

716 **Fig. 10** Composite anomaly maps of the 200-hPa vector wind and geopotential height, and the SSTs

717 (from January to December) for the 10 wettest (a, b and e) and 10 driest (c, d and f) years of the Dai-

718 PDSI reconstruction during the period 1948-2010.