



# Neoglacial Climate Anomalies and the Harappan Metamorphosis

Authors:

Liviu Giosan<sup>1\*</sup>, William D. Orsi<sup>2,3</sup>, Marco Coolen<sup>4</sup>, Cornelia Wuchter<sup>4</sup>,  
Ann G. Dunlea<sup>1</sup>, Kaustubh Thirumalai<sup>5</sup>, Samuel E. Munoz<sup>1</sup>, Peter D. Clift<sup>6</sup>,  
Jeffrey P. Donnelly<sup>1</sup>, Valier Galy<sup>7</sup>, Dorian Q. Fuller<sup>8</sup>

Affiliations:

<sup>1</sup>Geology & Geophysics, Woods Hole Oceanographic Institution, MA, USA

<sup>2</sup>Department of Earth and Environmental Sciences, Paleontology & Geobiology, Ludwig-Maximilians-Universität München, 80333 Munich, Germany

<sup>3</sup>GeoBio-CenterLMU, Ludwig-Maximilians-Universität München, 80333 Munich, Germany

<sup>4</sup>Curtin University, Perth, Australia

<sup>5</sup>Brown University, Providence, RI, USA

<sup>6</sup>Geology & Geophysics, Louisiana State University, USA

<sup>8</sup>Marine Chemistry & Geochemistry, Woods Hole Oceanographic Institution, MA, USA

<sup>7</sup>Institute of Archaeology, University College London, London, UK

\*Correspondence: [lgiosan@whoi.edu](mailto:lgiosan@whoi.edu)



27 Abstract:

28

29 Climate exerted constraints on the growth and decline of past human societies but our knowledge  
30 of temporal and spatial climatic patterns is often too restricted to address causal connections. At  
31 a global scale, the inter-hemispheric thermal balance provides an emergent framework for  
32 understanding regional Holocene climate variability. As the thermal balance adjusted to gradual  
33 changes in the seasonality of insolation, the Inter-Tropical Convergence Zone migrated  
34 southward accompanied by a weakening of the Indian summer monsoon. Superimposed on this  
35 trend, anomalies such as the Little Ice Age point to asymmetric changes in the extratropics of  
36 either hemisphere. Here we present a reconstruction of the Indian winter monsoon in the Arabian  
37 Sea for the last 6000 years based on paleobiological records in sediments from the continental  
38 margin of Pakistan at two levels of ecological complexity: sedimentary paleo-DNA reflecting  
39 water column environmental states and planktonic foraminifers sensitive to winter conditions.  
40 We show that strong winter monsoons between ca. 4,500 and 3,000 years ago occurred during an  
41 interval of weak interhemispheric temperature contrast, which we identify as the Early  
42 Neoglacial Anomaly (ENA), and were accompanied by changes in wind and precipitation  
43 patterns across the eastern Northern Hemisphere and Tropics. This coordinated climate  
44 reorganization may have helped trigger the metamorphosis of the urban Harappan civilization  
45 into a rural society through a push-pull migration from summer flood-deficient river valleys to  
46 the Himalayan piedmont plains with augmented winter rains. Finally, we speculate that time-  
47 transgressive landcover changes due to aridification of the Tropics may have led to a generalized  
48 instability of the global climate during ENA at the transition from the warmer Holocene  
49 Optimum to the cooler Neoglacial.

50

51



## 1. Introduction

The growth and decline of human societies can be affected by climate (e.g., Butzer, 2012; DeMenocal, 2001) but addressing causal connections is difficult, especially when no written records exist. Human agency sometimes confounds such connections by acting to mitigate climate pressures or, on the contrary, increasing the brittleness of social systems in face of climate variability (Rosen, 2007). Moreover, our knowledge of temporal and spatial climatic patterns remains too restricted to fully address social dynamics. Still, the coalescence of migration phenomena, profound cultural transformations and/or collapse of societies regardless of geographical and cultural boundaries during certain time periods characterized by climatic anomalies, events or regime shifts suggests that large scale climate variability may be involved (e.g., Donges et al., 2015 and references therein). At the global scale, the interhemispheric thermal balance provides an emergent framework for understanding such major Holocene climate events (Boos and Korty, 2016; Broecker and Putnam, 2013; McGee et al., 2014; Schneider et al., 2014). As this balance adjusted over the Holocene to gradual changes in the seasonality of insolation (Berger and Loutre, 1991), the Inter-Tropical Convergence Zone (ITCZ) migrated southward (e.g., Arbuszewski et al., 2013; Haug et al., 2001) accompanied by a weakening of the Indian summer monsoon (e.g., Fleitmann et al., 2003; Ponton et al., 2012). Superimposed on this trend, centennial- to millennial-scale anomalies point to asymmetric changes in the extratropics of either hemisphere (Boos and Korty, 2016; Broccoli et al., 2006; Chiang and Bitz, 2005; Chiang and Friedman, 2012; Schneider et al., 2014).

The most extensive but least understood among the early urban civilizations, the Harappan (Fig. 1), collapsed ca. 3900 years ago (e.g., Shaffer, 1992). At their peak, the Harappans spread over the alluvial plain of the Indus and its tributaries encroaching onto the Ghaggar-Hakra (G-H) interfluvium that separates the Indus and Ganges drainage basins (Fig. 1). In the late Harappan phase that was characterized by more regional artefact styles and trading networks, cities and settlements along the Indus and tributaries declined while the number of rural sites increased on the upper G-H interfluvium (Gangal et al., 2001; Kenoyer, 1998; Mughal, 1997; Possehl, 2002; Wright, 2010). The agricultural Harappan economy showed a large degree of versatility by adapting to water availability (e.g., Fuller, 2011; Giosan et al., 2012; Madella and Fuller, 2006; Petrie et al., 2017; Weber et al., 2010; Wright et al., 2008). Two precipitation sources, the summer monsoon and winter westerlies (Fig. 1), provide rainfall to the region (Bookhagen and Burbank, 2010; Petrie et al., 2017; Wright et al., 2008). Previous simple modeling exercises suggested that winter rain increased in Punjab over the late Holocene (Wright et al., 2008). During the hydrologic year, part of this precipitation, stored as snow and ice in surrounding mountain ranges, is redistributed as meltwater by the Indus and its Himalayan tributaries to the arid and semi-arid landscape of the alluvial plain (Karim and Veizer, 2002).



92 The climatic trigger for the urban Harappan collapse was probably the decline of the summer  
93 monsoon (e.g., Dixit et al., 2014; Kathayat et al., 2017; MacDonald, 2011; Singh et al., 1971;  
94 Staubwasser et al. 2003; Stein, 1931) that led to less extensive and more erratic floods making  
95 inundation agriculture less sustainable along the Indus and its tributaries (Giosan et al., 2012)  
96 and may have led to bio-socio-economic stress and disruptions (e.g., Meadow, 1991; Schug et  
97 al., 2013). Still, the remarkable longevity of the decentralized rural phase until ca. 3200 years  
98 ago in the face of persistent late Holocene aridity (Dixit et al., 2014; Fleitmann et al., 2003;  
99 Ponton et al., 2012; Prasad and Enzel, 2006) remains puzzling. Whether the Harappan  
100 metamorphosis was simply the result of habitat tracking toward regions where summer monsoon  
101 floods were still reliable or also reflected a significant increase in winter rain remains unknown  
102 (Giosan et al., 2012; Madella and Fuller, 2006; Petrie et al., 2017; Wright et al., 2008). To  
103 address this dilemma, we present a proxy record of the Indian winter monsoon in the Arabian  
104 Sea and show that its variability was an expression of large scale climate reorganization across  
105 the eastern Northern Hemisphere and Tropics affecting precipitation patterns across the  
106 Harappan territory. Aided by an analysis of Harappan archaeological site redistribution, we  
107 speculate that the Harappan relocation after the collapse of its urban phase may have conformed  
108 to a push-pull migration model.

109

110

## 111 2. Background

112

113 Under modern climatological conditions (Fig. 2), the summer monsoon delivers most of the  
114 precipitation to the former Harappan territory, but winter rains are also significant in quantity  
115 along the Himalayan piedmont (i.e., between 15 and 30% annually). Winter rain is brought in  
116 primarily by extra-tropical cyclones embedded in the Westerlies (Dimri et al. 2017) and are  
117 known locally as Western Disturbances (WD). These cyclones distribute winter rains to a zonal  
118 swath extending from the Mediterranean through Mesopotamia, the Iranian Plateau and  
119 Baluchistan, all across to the western Himalayas (Fig. 2). Stronger and more frequent WD rains  
120 in NW India are associated with southern shifts of the Westerly Jet in the upper troposphere  
121 (e.g., Dimri et al. 2017). Surface winter monsoon winds are generally directed towards the  
122 southwest but they blow preferentially toward the southeast in the northernmost Arabian Sea  
123 (Fig. 2). An enhanced eastward zonal component over the northern Arabian Sea is typical for  
124 more rainy winters (Dimri et al. 2017). Although limited in space and time, modern  
125 climatologies indicate a strong, physical linkage between winter sea-surface temperatures (SST)  
126 in the northern Arabian Sea and precipitation on the Himalayan piedmont, including the upper  
127 G-H interfluvium (see supplementary materials).

128

129 In contrast to the wet summer monsoon, winds of the winter monsoon flow from the continent  
130 toward the ocean and are generally dry. That explains in part why Holocene reconstructions of  
131 the winter monsoon are few and contradictory suggesting strong regional variabilities (Jia et al.,



2015; Kotlia et al., 2017; Li and Morrill, 2015; Sagawa et al., 2014; Wang et al., 2012; Yancheva et al., 2007). Holocene eolian deposits linked to the winter monsoon are also geographically-limited (Li and Morrill, 2015). However, in the Arabian Sea indirect wind proxies have been used based on changes in planktonic foraminifer assemblages and other mixing proxies to reconstruct distinct hydrographic states caused by seasonal winds (Boll et al., 2014; Curry et al., 1992; Luckge et al., 2001; Munz et al., 2015; Schiebel et al., 2004; Schulz et al., 2002). Winter monsoon winds blowing over the northeast Arabian Sea cool its surface waters via evaporation and weaken thermal stratification promoting convective mixing (Banse and McClain, 1986; Luis and Kawamura, 2004). Cooler SSTs and the injection of nutrients into the photic zone lead in turn to changes in the plankton community (Madhupratap et al., 1996; Luis and Kawamura, 2004; Schulz et al., 2002). To reconstruct the history of winter monsoon we thus employed complementary proxies for convective winter mixing, at two levels of ecological complexity: (a) sedimentary paleo-DNA to assess the water column plankton community structure, and (b) the relative abundance of *Globigerina falconensis*, a planktonic foraminifer sensitive to winter conditions (Munz et al., 2015; Schulz et al., 2002).

### 3. Methods

#### 3.1 Sediment Core

We sampled the upper 2.3 m comprising the Holocene interval in the 13-m-long piston core Indus 11C (Clift et al., 2014) retrieved during *R/V Pelagia* cruise 64PE300 in 2009 from the oxygen minimum zone (OMZ) in the northeastern Arabian Sea (23°07.30'N, 66°29.80'E; 566 m depth) (Fig. 1). The chronology for the Holocene section of the core was previously reported in Orsi et al. (2017) and is based on calibrated radiocarbon dates of five multi-specimen samples of planktonic foram *Orbulina universa* and one mixed planktonic foraminifer sample. Calibration was performed using Calib 7.129 with a reservoir age of  $565 \pm 35$  radiocarbon years following regional reservoir reconstructions by Staubwasser et al. (2002). Calibrated radiocarbon dates were used to derive a polynomial age model (see supplementary materials). The piston corer did not recover the last few hundred years of the Holocene record probably due to overpenetration. However, indistinct but continuous laminations downcore with no visual or X-radiograph discontinuities, together with the radiocarbon chronology indicate that the sedimentary record recovered is continuous.

#### 3.2. DNA Analyses

A total of five grams of wet weight sediment were extracted inside the ancient DNA-dedicated lab at Woods Hole Oceanographic Institution (WHOI), aseptically as described previously (Coolen et al., 2013) and transferred into 50 mL sterile tubes. The sediments were homogenized



for 40 sec at speed 6 using a Fastprep 96 homogenizer (MP Biomedicals, Santa Ana, CA) in the presence of beads and 15 ml of preheated (50 °C) sterile filtered extraction buffer (77 vol% 1M phosphate buffer pH 8, 15 vol% 200 proof ethanol, and 8 vol% of MoBio's lysis buffer solution C1 [MoBio, Carlsbad, CA]). The extraction was repeated with 10 ml of the same extraction buffer but without C1 lysis buffer (Orsi et al., 2017). After centrifugation, the supernatants were pooled and concentrated to a volume of 100 µl without loss of DNA using 50,000 NMWL Amicon® Ultra 15 mL centrifugal filters (Millipore) and contaminants were removed from the concentrated extract using the PowerClean® Pro DNA Clean-up Kit (MoBio). The exact same procedures were performed in triplicate without the addition of sediment as a control for contamination during extraction and purification of the sedimentary DNA.

DNA was quantified fluorometrically using Quant-iT PicoGreen dsDNA Reagent (Invitrogen), and ~20 nanograms of each extract was used as template for PCR amplification of preserved planktonic 18S rRNA genes. The short (~130 base pair) 18S rDNA-V9 region was amplified using the domain-specific primer combination 1380F (5'-CCC TGC CHT TTG TAC ACA C-3') and 1510R (5'CCT TCY GCA GGT TCA CCT AC-3')(REF). qPCR was performed using a SYBR®Green I nucleic acid stain (Invitrogen) and using a Realplex quantitative PCR system (Eppendorf, Hauppauge, NY). The annealing temperature was set to 66 °C and all reactions were stopped in the exponential phase after 35-42 cycles. 18S libraries were sequenced on an Illumina MiSeq sequencing using the facilities of the W.M. Keck Center for Comparative and Functional Genomics, University of Illinois at Urbana-Champaign, IL, USA sequenced 18S libraries that resulted in approximately 12 million DNA sequences.

rRNA gene sequences were processed in QIIME (Caporaso et al., 2010). Reads passing quality control (removal of any sequence containing an 'N', minimum read length 250 bp, minimum Phred score=20) were organized into OTUs sharing 95% sequence identity with UCLUST (Edgar et al., 2010) and assigned to taxonomic groups through BLASTn searches against the SILVA database (Pruesse et al., 2007). OTU tables were rarefied to the sample with the least number of sequences, and all OTUs containing less than one sequence were removed. OTUs that were detected in only one sample were also removed. Metagenomes were directly sequenced bi-directionally on an Illumina HiSeq, at the University of Delaware Sequencing and Genotyping Center (Delaware Biotechnology Institute). Contigs were assembled de novo as described in Orsi et al. (2017). To identify contigs containing chlorophyll biosynthesis proteins, open reading frames on the contig sequences were detected using FragGeneScan (Rho et al., 2010), and protein homologs were identified through BLASTp searches against the SEED database ([www.theseed.org](http://www.theseed.org)). Only hits to reference proteins with at least 60% amino acid similarity over an alignment length >50 amino acids were considered true homologs and used for downstream analysis. Assignment of ORFs to biochemical pathway classes were made based on the SEED metabolic pathway database and classification scheme. The relative abundance of reads mapping



to ORFs was normalized against values of a suite of 35 universally conserved single copy genes (Orsi et al., 2015), per metagenome sample.

### 3.3 Factor Analysis

Q-mode Factor Analysis (QFA) was employed to simplify the paleo-DNA dataset. Prior to the factor analysis the DNA database was reduced to 124 most abundant taxonomic units from a total of 1,462 units identified by considering only those present in two or more samples with a cumulative abundance higher than  $0.5 \pm 0.1\%$  (Table S1). The data was pretreated with a range-normalization and run through the QFA with a VARIMAX rotation (Pisias et al., 2013). QFA identified taxonomic groups that covary in our dataset and determined the minimum number of components (i.e., factors) needed to explain a given fraction of the variance of the data set (Fig. 3; see supplementary materials). Each VARIMAX-rotated factor indicates an association of taxonomic groups that covary (i.e., behave similarly amongst the samples). Taxonomic groups that covary strongly within a factor will have high factor scores for that factor. We primarily used dominant taxa with scores higher than 0.2 in a factor to interpret the plankton taxonomic groups in that factor. The importance of a factor in any given sample is recorded by the factor loading that we used to interpret the importance of that factor with depth/time downcore.

### 3.4 Foraminifera Counts

Samples for counting planktonic foraminifer *Globigerina falconensis* were wet-sieved over a 63- $\mu\text{m}$  screen. Typical planktonic foraminifer assemblages for the NE Arabian Sea were observed: *Globigerinoides ruber*, *Neoglobobulimina dutertrei*, *Globigerina falconensis*, *Orbulina universa*, *Globigerinoides sacculifer*, *Pulleniatina obliquiloculata*, *Globobulimina menardii*. Counts of *Globigerina falconensis* were conducted on the size fraction  $>150 \mu\text{m}$ . We report counts for the samples yielding  $>300$  foraminifer individuals (see supplementary materials).

### 3.5 Harappan Sites

Archaeological site distribution provides an important line of evidence for social changes in the Harappan domain (e.g., Possehl, 2000). We analyzed the redistribution of small ( $<20 \text{ ha}$ ), rural vs. large ( $>20 \text{ ha}$ ), possibly urban sites on the G-H interfluvium from the Early Harappan period, through the Mature and Late periods to the post-Harappan Grey Ware culture. Compared to settlements along the Indus and its tributaries affected by fluvial erosion (Giosan et al., 2012), the distribution of archaeological sites on G-H, where large laterally-incising Himalayan rivers were absent during the Holocene, is probably more complete and representative of their original distribution. To observe trends related to partial or complete drying of the G-H system (Clift et al., 2012; Giosan et al., 2012; Singh et al., 2017), we divided the settlements into upper and lower G-H sites located in the modern regions of Punjab and Haryana in India, respectively





Cholistan in Pakistan. For archaeological site locations and their radiocarbon and/or archaeological ages we follow Giosan et al. (2012) using data from the compilation by Gangal et al. (2001) with additions from regional gazetteers and surveys (Kumar, 2009; Mallah, 2010; Mughal, 1996 and 1997; Possehl, 1999; Wright et al., 2005).

#### 4. Results

Exceptional preservation of organic matter in the OMZ (Altabet et al., 1995; Schulz et al., 2002) allowed us to reconstruct the history of the plankton communities based on their fossil sedimentary DNA (see also Orsi et al., 2017). The factor analysis of the dominant DNA species identified three significant factors that together explain 48% of the variability in the dataset (see supplementary materials). Additional factors were excluded as they would have increased the variability explained by an insignificant amount for each (< 3%). We interpret these factors as corresponding to the SST regime, nutrient availability, and sea level state, respectively (Fig. 3). Factor 1 explains 20% of the variability and is largely dominated by radiolarians (*Polycystinea*) that prefer warmer sea surface conditions (e.g., Cortese and Ablemann, 2002; Kamikuri et al., 2008). High scores for jellyfish (*Cnidaria*) that thrive in warm, eutrophic waters (Purcell, 2005) also support interpreting Factor 1 as a proxy for a plankton community adapted to high sea surface temperatures. A general increase of the Factor 1 loadings since early Holocene is in accordance with the  $U^{K}_{37}$ -reconstructed warming of Orsi et al. (2017). During the Holocene, relatively colder conditions are evident in Factor 1 between ~4500 and 3000 years BP (Fig. 3) as previously detected in the higher resolution  $U^{K}_{37}$  record from a core located nearby on the Makran continental margin (Dooze-Rolinski et al., 2001).

Factor 2 explains 18% of the variability and is dominated by marine dinoflagellates indicative of high nutrient, bloom conditions (e.g., Worden et al., 2015), flagellates (*Cercozoa*) and fungi. Parasitic Alveolates (*Hematodinium* and *Syndiniales*) that typically appear during blooms (Worden et al., 2015) are also important. Increased representation of chlorophyll biosynthesis genes (Fig. 3) in sediment metagenomes (Orsi et al., 2017) indicate higher productivity (Worden et al., 2015) during the Factor 2 peak. All these associations suggest that Factor 2 is a nutrient-sensitive proxy with a peak that overlaps with the colder conditions between ~4500 and 3000 years BP. The inland retreat of Indus fluvial nutrient source as sea level rose (see below) probably explains the asymmetry in Factor 2 that exhibits higher scores in the early vs. late Holocene. Overall, Factors 1 and 2 suggests enhanced winter convective mixing between ~4500 and 3000 years BP that brought colder, nutrient-rich waters to the surface.

Factor 3 explains 10% variability and is dominated by a wide group of taxa. The main identified contributors to Factor 3 include the coastal diatom *Eucampia* (Werner, 1977), the fish-egg parasite dinoflagellate *Ichthyodinium*, also reported from coastal habitats (Shadrin, 2010), and





soil ciliates (*Colpodida*), which altogether suggest a nearshore environment with fluvial inputs. The plankton community described by Factor 3 was dominant in the first half of the Holocene and became scarce as the sea level rose (Camoin et al., 2004) and the Indus coast retreated inland (Fig. 3).

At a simpler ecological level, *Globigerina falconensis* is the dominant planktonic foraminifer in the NE Arabian Sea under strong winter wind mixing conditions (Munz et al., 2015; Schulz et al., 2002). Over the last six millennia, after the sea level approached the present level, and when the plankton community was consistently outside the influence of coastal and fluvial processes, *G. falconensis* shows a peak in relative abundance between ~4500 and 3000 years during the cold reversal previously identified by the paleo-DNA (Fig. 3). A similar peak in *G. falconensis* was detected in core SO42-74KL from the western Arabian Sea upwelling area (Schulz et al., 2002) suggesting that mixing occurred in the whole northern half of the Arabian Sea (Fig. 3).

## 5. Discussion with Conclusions

### 5.1 Winter Monsoon Variability in the Neoglacial

In concert with previous data from the northern Arabian Sea, our reconstructions suggest that convective mixing conditions indicative of a stronger winter monsoon occurred between ~4,500 and 3,000 years. Another cold yet variable period in the northern Arabian Sea (Dooze-Rolinski et al., 2001) occurred after ~1500 years ago under strong winter monsoon mixing (Boll et al., 2014; Munz et al., 2015) and is seen in *G. falconensis* record of Schulz et al. (2002) but is not captured completely in our top-incomplete record. In accordance with modern climatologies colder SSTs in the northern coastal Arabian Sea correspond to increased westerly extratropical cyclones bringing winter rains as far as Baluchistan and the western Himalayas (Fig. 3). Pollen records offshore the Makran coast where rivers from Baluchistan and ephemeral streams flood during winter (von Rad et al., 1999) indeed indicate enhanced winter monsoon precipitation during between ~4,500 and 3,000 years BP (Ivory and Lezine, 2009). Bulk chemistry of sediments from the same Makran core were used to infer enhanced winter-monsoon conditions between 3900 and 3000 years ago (Luckge et al., 2001).

During the Holocene, two periods of weak interhemispheric thermal gradient (Fig. 4) for areas poleward of 30°N and 30°S occurred on top of more gradual, monotonic changes driven by the seasonality of insolation (Marcott et al., 2013; Schneider et al., 2014). These intervals are coeval within the limitations of age models with the strong winter monsoon phases in the Arabian Sea (Fig. 4) and southward swings of the Intertropical Convergence Zone (ITCZ) in the western Atlantic Ocean (Haug et al., 2001). Occurring when Neoglacial conditions became pervasive across the Northern Hemisphere (Solomina et al., 2015), we identify the two late Holocene



intervals of low interhemispheric thermal gradient as the Early Neoglacial Anomaly (ENA) between ca. 4,500 and 3,000 years ago and the Late Neoglacial Anomaly (LNA) after ~1,500, respectively.

LNA includes well-known cold events such as the Little Ice Age (Mann et al., 2009) and the preceding cold during the European Migration Period (Büntgen et al., 2016), whereas ENA is more enigmatic. The high resolution Cariaco ITCZ record showing successive southward excursions suggests a series of Little Ice Age-like events (LIALE in short - a term proposed by Sirocko, 2015). Furthermore, during ENA, similar to synoptic conditions during the Little Ice Age, a dominantly negative phase of the North Atlantic Oscillation – NAO (Fig. 4; Olsen et al., 2012) occurred concurrent with moderate increases in storminess in the high-latitude North Atlantic region, as shown by sea-salt sodium in Greenland's GISP2 core (Fig. 4; O'Brien et al., 1995). During both ENA and LNA the tropical North Atlantic was remarkably quiescent in terms of hurricane activity (Fig. 4), which appears to be the direct result of the prevailing southward position of the ITCZ (Donnelly and Woodruff, 2007; van Hengstum et al., 2016).

At mid latitudes, a southward position for the Westerlies wind belt, as expected during negative NAO conditions, is supported at the western end of our domain of interest by well-defined increases in spring floods in the Southern Alps (Fig. 4) during both ENA and LNA (Wirth et al., 2013). A higher precipitation-evaporation state in the northern Levant (Fig. 4; Cheng et al., 2015) and positive balances from lake isotope records in the Eastern Mediterranean (Fig. 4; Roberts et al., 2011), including lakes in Iran, occur further along the southward Westerlies precipitation belt. The preferential southward track of the Westerlies during ENA and LNA is also in agreement with a stronger Siberian Anticyclone, the dominant mode of winter and spring climate in Eurasia, as interpreted from increases in the GISP2 non-sea-salt potassium (Fig. 4). At the Far East end, support comes from dust reconstructions in the Sea of Japan (Nagashima et al. 2013) and modeling (Kong et al., 2017), which suggest that the Westerlies stayed preferentially further south in the late Holocene. Like in modern climatologies, this suite of paleorecords supports our interpretation that stronger winter monsoon winds during ENA and LNA in the northernmost Arabian Sea, that ought to have driven more convective mixing at our core site, were accompanied by increased precipitation penetration along the Westerlies' path across the Iranian Plateau, Baluchistan and Makran to the western Himalayas.

In addition to its paleoclimatological value for the Harappan domain (see discussion below), a more fundamental question emerges from our analysis: what triggered ENA and LNA? The reduced influence of insolation on the ITCZ during the late Holocene (e.g., Haug et al., 2001; Schneider et al., 2014) could have provided favorable conditions for internal modes of climate variability, either tropical or polar, to become dominant (e.g., Wanner et al., 2008; Debret et al., 2009; Thirumalai et al., 2018). In order to explain intervals of tropical instabilities that did not extend over the entire Neoglacial various trigger mechanisms and/or coupling intensities between climate subsystems could be invoked. For example, the weaker orbital forcing increased



the susceptibility of climate to volcanic and/or solar irradiance, which have been proposed to explain decadal to centennial time events such as the Little Ice Age (e.g., Mann et al., 2009; McGregor et al., 2005). For the recently defined Late Antique Little Ice Age between 536 to about 660 AD, a cluster of volcanic eruptions sustained by ocean and sea-ice feedbacks and a solar minimum have been proposed as triggers (Buntgen et al., 2016). However, during ENA the solar irradiance was unusually stable without prominent minima (Stuiver and Braziunas, 1989; Steinhilber et al., 2012). The volcanic activity in the northern hemisphere was also not particularly higher during ENA than after (Zielenski et al., 1996) and it was matched by an equally active southern hemisphere volcanism (Castellano et al., 2005). As previously suggested for the Little Ice Age (Dull et al. 2010; Nevle and Bird, 2008), we speculate that mechanisms related to changes in landcover and possibly landuse could have instead been involved in triggering ENA.

Biogeophysical effects of aerosol, albedo and evapotranspiration due to landcover changes were previously shown to be able to modify the position of ITCZ and lead to significant large scale geographic alterations in hydrology (e.g., Chung and Soden, 2017; Dallmeyer et al., 2017; Devaraju et al. 2015; Kang et al., 2018; Sagoo and Storelvmo, 2017; Tierney et al., 2017). Similarly, tropical changes in albedo and concurrent changes in regional atmospheric dust emissions due to aridification in the Neoglacial could have affected the ITCZ. Anthropogenic early land use changes could have also led to large scale biogeophysical impacts (e.g., Smith et al., 2016). Such landcover- and landuse-driven changes were time-transgressive across Asia and Africa (e.g., Lezine et al., 2017; Jung et al., 2004; Prasad and Enzel, 2006; Shanahan et al., 2015; Tierney et al., 2017; Wang et al. 2010; Kaplan et al., 2011) and could have led to a generalized instability of the global climate as it passed from the warmer Holocene Optimum state to the cooler Neoglacial state. Therefore the instability seen during ENA may reflect threshold behaviour of the global climate system characterized by fluctuations or flickering (Dakos et al., 2008; Thomas, 2016) or a combination of different mechanisms affecting the coupling intensity between climate subsystems (Wirtz et al. 2010).

## 5.2 Climate Instability and the Harappan Metamorphosis

In contrast to other urban civilizations of the Bronze Age, such as Egypt and Mesopotamia, Harappans did not employ canal irrigation to cope with the vagaries of river floods despite probable knowledge about this agricultural technology through their western trade network (e.g., Ratnagar, 2004). Instead, they relied on a multiple cropping system that started to develop prior to their urban rise (Madella and Fuller, 2006; Petrie et al., 2017) and integrated the winter crop package imported from the Fertile Crescent (e.g., wheat, barley, peas, lentil) with local summer crops (e.g., millets, sesame, limited rice). A diverse array of cropping practices using inundation and/or dry agriculture that were probably supplemented by labor-intensive well irrigation was employed across the Indus domain, dependent on the regional characteristics of seasonal rains



412 and river floods (e.g., Weber 2003; Pokharia et al. 2014; Petrie and Bates, 2017; Petrie et al.,  
413 2017). The alluvial plains adjacent to the foothills of the Himalayas were probably the Harappan  
414 region most amenable to multiple crops using summer monsoon and WD rains directly or  
415 redistributed via the perennial and/or ephemeral streams of the G-H interfluvial. The  
416 orographically-controlled stability and availability of multiple water sources that could be used  
417 to mitigate climate risks probably made this area more attractive as the inundation agriculture  
418 faltered along the Indus and its tributaries when the summer monsoon became more erratic.  
419

420 Aridity intensified over most of the Indian subcontinent as the summer monsoon rains started to  
421 decline after 5,000 years ago (Ponton et al., 2012; Prasad et al., 2014). The closest and most  
422 detailed summer monsoon reconstruction to the Harappan domain shows a highly variable  
423 multicentennial trend to drier conditions between ca. 4,300 and 3,300 years ago (Fig. 5; Kathayat  
424 et al., 2017). Thresholds in evaporation-precipitation affecting lakes on the upper G-H interfluvial  
425 occurred during the same period (Fig. 5; Dixit et al., 2014). The flood regime controlled by this  
426 variable and declining summer monsoon became more erratic and/or spatially restricted (Giosan  
427 et al., 2012; Durcan et al., 2017) making inundation agriculture less dependable. Whether fast or  
428 over generations, the bulk of Harappan settlements relocated toward the Himalayan foothills on  
429 the plains of the upper G-H interfluvial (Possehl, 2002; Kenoyer, 1998; Wright, 2010; Madella  
430 and Fuller, 2006; Giosan et al., 2017). Abandoned by Himalayan rivers since the early Holocene  
431 (Giosan et al., 2012; Clift et al., 2012; Singh et al., 2017), this region between the Sutlej and  
432 Yamuna was watered by orographically-enhanced rain feeding an intricate small river network  
433 (e.g., Yashpal et al., 1980; van Dijk et al., 2016; Orengo and Petrie, 2017).  
434

435 During the aridification process the number of large, urban-sized settlements on the G-H  
436 interfluvial decreased and the number of small settlements drastically expanded (Fig. 5). The  
437 rivers on the G-H interfluvial merged downstream to feed flows along the Hakra into Cholistan, at  
438 least seasonally, until the latest Holocene (Giosan et al., 2012). Regardless if these settlements on  
439 the lower G-H interfluvial were temporary and mobile (Petrie et al., 2017) most of them were  
440 abandoned (Fig. 5) as the region aridified, suggesting that flows became less reliable in this  
441 region. However, the dense stream network on the upper G-H interfluvial must have played an  
442 important role in more uniformly watering that region, whether perennially or seasonally.  
443 Remarkably, Late Harappan settling did not extend toward the northwest along the entire  
444 Himalayan piedmont despite the fact that this region must have received orographically-  
445 enhanced rains too (Fig. 2). One possible reason is that interfluvies between Indus tributaries (i.e.,  
446 Sutlej, Beas, Ravi, Chenab, Jhelum) are not extensive. These Himalayan rivers are entrenched  
447 and collect flows inside their wide valleys rather than supporting extensive interfluvial stream  
448 networks (Giosan et al., 2012).  
449

450 Our winter monsoon reconstruction suggests that WD precipitation increased during the time of  
451 urban Harappan collapse (Fig. 5). As the summer monsoon flickered and declined in the same



time, the classical push-pull model (e.g., Dorigo and Tobler, 1983; Ravenstein, 1885; 1889) could help explain the Harappan migration. Push-pull factors induce people to migrate from negatively affected regions to more favorable locations. Inundation agriculture along the summer flood-deficient floodplains of the Indus and its tributaries became too risky, which pushed people out, in the same time as the upper G-H region became increasingly attractive due to augmented winter rain, which pulled migrants in. These winter rains would have supported traditional winter crops like wheat and barley, while drought tolerant millets could still be grown in rotation during the monsoon season. Although present throughout the Harappan period, a greater reliance on summer crops after the urban Mature Harappan collapse implies that intense efforts were made to adapt to hydroclimatic stress at the arid outer edge of the monsoonal rain belt (Giosan et al., 2012; Madella and Fuller, 2006; Petrie and Bates, 2017; Wright et al., 2008). The longevity of the Late Harappan settlements in this region may be due to a consistent availability of multiple year-round sources of water. Summer monsoon remained high enough locally due to orographic rainfall while winter precipitation increased during ENA and both these sources provided relief from labor-intensive alternatives such as well irrigation.

The metamorphosis of Indus civilization remains an episode of great interest. The degradation of cities and disintegration of supra-regional elements of the Indus cultural system such as its script need not be sudden to be defined as a collapse. However, recent contributions of geoarchaeological and settlement patterns studies, together with refinements in chronology, require higher levels of sophistication for addressing links between climatic shifts and cultural decline. While variation in coverage and imprecision in dating sites require further efforts (Petrie et al., 2017), it remains clear that there were shifts in the distribution of population and the range of site sizes, with decline in the size of the largest sites. The impacts of climatic shifts while remarkable from recent chronological correlations (e.g., Katahayat et al 2017) must now be assessed regionally through a nuanced appreciation of rainfall quantities as well as its seasonality (e.g., Madella and Fuller, 2006; MacDonald, 2011; Petrie et al., 2017; Wright et al., 2008). How precipitation was distributed seasonally would have affected the long-term stability and upstream sources of the stream and river network (Giosan et al 2012; Singh et al 2017). Our study suggests broad spatial and temporal patterns of variability for summer and winter precipitation across the Harappan domain but the role of seasonal gluts or shortage of rain on river discharge need also to be considered. For example, did the increase in winter rain during ENA lead to more snow accumulation in the Himalayas that affected the frequency and magnitude of floods along the Indus and its tributaries? Or did settlements in Kutch and Saurashtra, regions of relatively dense habitation during Late Harappan times, also benefit from increases in winter rains despite the fact that modern climatologies suggest scarce local precipitation?

Local reconstructions of seasonal hydroclimatic regimes would greatly enhance our ability to understand social and economic choices made by Harappans. Attempts made to reconstruct WD precipitation in the western Himalayas (e.g., Kotlia et al., 2017) are confounded by the dominant



492 summer monsoon (c.f., Kathayat et al., 2017). Developing local proxies based on summer vs.  
493 winter crop remains may provide a more fruitful route for disentangling the sources of water in  
494 the Harappan domain (e.g., Bates et al., 2017). The Indus civilization especially in the northern  
495 and eastern regions had a broad choice of crops of both seasons. Mixed cropping may have  
496 become increasingly important, including drought-tolerant, but less productive, summer millets  
497 that suited weakening monsoon and winter cereals, including drought-tolerant barley, that were  
498 aided by the heightened winter rains of Late Harappan era. Facilitated by this climatic  
499 reorganization during ENA, the eastward shift in settlements, while it may have undermined the  
500 pre-eminence of the largest urban centres like Harappan, can be seen as a strategic adjustment in  
501 subsistence to the summer monsoon decline. If and how ENA may have affected human  
502 habitation at the scale of the entire eastern Northern Hemisphere, and particularly in the Fertile  
503 Crescent and Iran that also depend on winter rains, remains to be assessed.

504

505

506 Acknowledgements

507

508 This work was supported by the NSF OCE Grant #0634731 and internal WHOI funds to LG,  
509 NSF MGG Grant #1357017 to MJLC, VG, and LG, and a C-DEBI grant #OCE-0939564 to  
510 WDO. We thank the editors and reviewers. Thanks go to Mary Carman for help with  
511 foraminifera and Pakistani and Indian colleagues who helped with acquiring and/or provided  
512 access to data including Kavita Gangal, Ronjoy Adhikari, Ali Tabrez, and Asif Inam.

513





## References

- Altabet, M.A., Francois, R., Murray, D.W. and Prell, W.L., 1995. Climate-related variations in denitrification in the Arabian Sea from sediment 15N/14N ratios. *Nature*, 373(6514), pp.506-509.
- Arbuszewski, J. A., deMenocal, P. B., Cleroux, C., Bradtmiller, L. & Mix, A. Meridional shifts of the Atlantic intertropical convergence zone since the Last Glacial Maximum. *Nature Geosci.* 6, 959–962 (2013).
- Banse, K. and McClain, C.R., 1986. Winter blooms of phytoplankton in the Arabian Sea as observed by the Coastal Zone Color Scanner. *Marine Ecology Progress Series*, pp.201-211.
- Bates, J., Singh, R.N. and Petrie, C.A., 2017. Exploring Indus crop processing: combining phytolith and macrobotanical analyses to consider the organisation of agriculture in northwest India c. 3200–1500 BC. *Vegetation History and Archaeobotany*, 26(1), pp.25-41.
- Berger, W. H., and M. F. Loutre (1991), Insolation values for the climate of the last 10 m. y. *Quat. Sci. Rev.*, 10, 297–317.
- Berkelhammer, M., Sinha, A., Stott, L., Cheng, H., Pausata, F.S.R., and Yoshimura, K., 2012, An abrupt shift in the Indian monsoon 4000 years ago, in Giosan, L., et al., eds., *Climates, landscapes, and civilizations: American Geophysical Union Geophysical Monograph* 198, p. 75–87, doi:10.1029/2012GM001207.
- Bhadra, B.K.; Gupta, A.K.; Sharma, J.R. Saraswati Nadi in Haryana and its linkage with the Vedic Saraswati River - Integrated study based on satellite images and ground Based information. *J. Geol. Soc. India* 2009, 73, 273–288.
- Böll, A., Lückge, A., Munz, P., Forke, S., Schulz, H., Ramaswamy, V., Rixen, T., Gaye, B. and Emeis, K.C., 2014. Late Holocene primary productivity and sea surface temperature variations in the northeastern Arabian Sea: Implications for winter monsoon variability. *Paleoceanography*, 29(8), pp.778-794.
- Bookhagen B, Burbank DW (2010) Towards a complete Himalayan hydrological budget: the spatiotemporal distribution of snow melt and rainfall and their impact on river discharge. *J Geophys Res-Earth* 115:1–25.
- Boos, W.R. and Korty, R.L., 2016. Regional energy budget control of the intertropical convergence zone and application to mid-Holocene rainfall. *Nature Geoscience*.
- Broccoli, A. J., Dahl, K. A. & Stouffer, R. J. Response of the ITCZ to Northern Hemisphere cooling. *Geophys. Res. Lett.* 33, L01702 (2006).
- Broecker, W.S. and Putnam, A.E., 2013. Hydrologic impacts of past shifts of Earth's thermal equator offer insight into those to be produced by fossil fuel CO<sub>2</sub>. *Proceedings of the National Academy of Sciences*, 110(42), pp.16710-16715.
- Büntgen, U., Myglan, V.S., Ljungqvist, F.C., McCormick, M., Di Cosmo, N., Sigl, M., Jungclaus, J., Wagner, S., Krusic, P.J., Esper, J. and Kaplan, J.O., 2016. Cooling and societal change during the Late Antique Little Ice Age from 536 to around 660 AD. *Nature Geoscience*, 9(3), pp.231-236.
- Butzer KW (2012) Collapse, environment, and society. *Proc Natl Acad Sci USA* 109(10):3632–3639.
- Camoin, G.F., Montaggioni, L.F., and Braithwaite, C.J.R., 2004, Late glacial to post glacial sea levels in the western Indian Ocean: *Marine Geology*, v. 206, p. 119–146, doi: 10.1016/j.margeo.2004.02.003.
- Caporaso, J. G. et al. QIIME allows analysis of high-throughput community sequencing data.





- 560 Nat Methods 7, 335-336, (2010).
- 561 Castellano, E., Becagli, S., Hansson, M., Hutterli, M., Petit, J.R., Rampino, M.R., Severi, M.,  
562 Steffensen, J.P., Traversi, R. and Udisti, R., 2005. Holocene volcanic history as recorded in  
563 the sulfate stratigraphy of the European Project for Ice Coring in Antarctica Dome C  
564 (EDC96) ice core. *Journal of Geophysical Research: Atmospheres*, 110(D6).
- 565 Cheng, H., Sinha, A., Verheyden, S., Nader, F.H., Li, X.L., Zhang, P.Z., Yin, J.J., Yi, L., Peng,  
566 Y.B., Rao, Z.G. and Ning, Y.F., 2015. The climate variability in northern Levant over the  
567 past 20,000 years. *Geophysical Research Letters*, 42(20), pp.8641-8650.
- 568 Chiang, J. C. H. & Bitz, C. M. Influence of high latitude ice cover on the marine Intertropical  
569 Convergence Zone. *Clim. Dynam.* 25, 477–496 (2005).
- 570 Chung, E.S. and Soden, B.J., 2017. Hemispheric climate shifts driven by anthropogenic aerosol-  
571 cloud interactions. *Nature Geoscience*, 10(8), p.566.
- 572 Clift PD, et al. (2012) U-Pb zircon dating evidence for a Pleistocene Sarasvati River and Capture  
573 of the Yamuna River. *Geology* 40:211–214.
- 574 Clift, P.D., Giosan, L., Henstock, T.J. and Tabrez, A.R., 2014. Sediment storage and reworking  
575 on the shelf and in the Canyon of the Indus River-Fan System since the last glacial  
576 maximum. *Basin Research*, 26(1), pp.183-202.
- 577 Cortese, G. and Abelmann, A., 2002. Radiolarian-based paleotemperatures during the last 160  
578 kyr at ODP Site 1089 (Southern Ocean, Atlantic Sector). *Palaeogeography*,  
579 *Palaeoclimatology, Palaeoecology*, 182(3), pp.259-286.
- 580 Curry, W.B., Ostermann, D.R., Guptha, M.V.S. and Ittekkot, V., 1992. Foraminiferal production  
581 and monsoonal upwelling in the Arabian Sea: evidence from sediment traps. *Geological*  
582 *Society, London, Special Publications*, 64(1), pp.93-106.
- 583 Dakos, V., Scheffer, M., van Nes, E.H., Brovkin, V., Petoukhov, V. and Held, H., 2008. Slowing  
584 down as an early warning signal for abrupt climate change. *Proceedings of the National*  
585 *Academy of Sciences*, 105(38), pp.14308-14312.
- 586 Dallmeyer, A., Claussen, M., Ni, J., Cao, X., Wang, Y., Fischer, N., Pfeiffer, M., Jin, L., Khon,  
587 V., Wagner, S. and Haberkorn, K., 2017. Biome changes in Asia since the mid-Holocene-an  
588 analysis of different transient Earth system model simulations. *Climate of the Past*, 13(2),  
589 p.107.
- 590 Debret, M., Sebag, D., Crosta, X., Massei, N., Petit, J.R., Chapron, E. and Bout-Roumazeilles,  
591 V., 2009. Evidence from wavelet analysis for a mid-Holocene transition in global climate  
592 forcing. *Quaternary Science Reviews*, 28(25), pp.2675-2688.
- 593 deMenocal PB (2001) Cultural responses to climate change during the late Holocene. *Science*  
594 292(5517):667–673.
- 595 Devaraju, N., Govindasamy Bala, and Angshuman Modak. "Effects of large-scale deforestation  
596 on precipitation in the monsoon regions: Remote versus local effects." *Proceedings of the*  
597 *National Academy of Sciences* 112.11 (2015): 3257-3262.
- 598 Dimri AP, Niyogi D, Barros AP, Ridley J, Mohanty UC, Yasunari T, Sikka DR (2015) Western  
599 disturbance: a review. *Rev Geophys.* 53, 225–246, doi:10.1002/2014RG000460.
- 600 Dimri, A. P. (2006), Surface and upper air fields during extreme winter precipitation over the  
601 western Himalayas, *Pure Appl. Geophys.*, 163(8), 1679–1698, doi:10.1007/s00024-006-  
602 0092-4.
- 603 Dixit Y, Hodell DA, Petrie CA (2014) Abrupt weakening of the summer monsoon in northwest  
604 India ~4100 yr ago. *Geology* 42(4):339–342.
- 605 Donges, J.F., Donner, R., Marwan, N., Breitenbach, S.F., Rehfeld, K. and Kurths, J., 2015. Non-



- linear regime shifts in Holocene Asian monsoon variability: potential impacts on cultural change and migratory patterns. *Climate of the Past*, 11(5), pp.709-741.
- Donnelly, J.P. and Woodruff, J.D., 2007. Intense hurricane activity over the past 5,000 years controlled by El Niño and the West African monsoon. *Nature*, 447(7143), p.465.
- Dooe-Rolinski, H., Rogalla, U., Scheeder, G., Lückge, A. and Rad, U., 2001. High-resolution temperature and evaporation changes during the late Holocene in the northeastern Arabian Sea. *Paleoceanography*, 16(4), pp.358-367.
- Dorigo, G., and W. Tobler (1983), Push-pull migration laws, *Ann. Assoc. Am. Geogr.*, 73(1), 1–17.
- Dull, R.A., Nevle, R.J., Woods, W.I., Bird, D.K., Avnery, S. and Denevan, W.M., 2010. The Columbian encounter and the Little Ice Age: Abrupt land use change, fire, and greenhouse forcing. *Annals of the Association of American Geographers*, 100(4), pp.755-771.
- Durcan, J.A., Thomas, D.S., Gupta, S., Pawar, V., Singh, R.N. and Petrie, C.A., 2017. Holocene landscape dynamics in the Ghaggar-Hakra palaeochannel region at the northern edge of the Thar Desert, northwest India. *Quaternary International*.
- Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26, 2460-2461, (2010).
- Fleitmann, D. et al. Holocene forcing of the Indian monsoon recorded in a stalagmite from southern Oman. *Science* 300, 1737–1739 (2003)
- Fuller DQ (2011) Finding plant domestication in the Indian subcontinent. *Curr Anthropol* 52:S347–S362.
- Gangal K, Vahia M, Adhikari R (2010) Spatio-temporal analysis of the Indus urbanization. *Current Sci India* 98:846–852.
- Giosan, L., Clift, P.D., Blusztajn, J., Tabrez, A., Constantinescu, S. and Filip, F., 2006, December. On the control of climate- and human-modulated fluvial sediment delivery on river delta development: the Indus. In *AGU Fall Meeting Abstracts*.
- Giosan, L., Clift, P.D., Macklin, M.G., Fuller, D.Q., Constantinescu, S., Durcan, J.A., Stevens, T., Duller, G.A.T., Tabrez, A., Adhikari, R., Gangal, K., Alizai, A., Filip, F., VanLaningham, S., Syvitski, J.P.M., 2012. Fluvial Landscapes of the Harappan Civilization. *Proc. Natl. Acad. Sci.* 109 (26), 1688–1694.
- Haug, G. H., Hughen, K. A., Sigman, D. M., Peterson, L. C. & Rohl, U. Southward migration of the Intertropical Convergence Zone through the Holocene. *Science* 293, 1304–1308 (2001).
- Hermann, C. F. (1997) “Harappan” Gujarat: the Archaeology-Chronology connection. *Paleorient* 22(2): 77-112
- Herzschuh U (2006) Palaeo-moisture evolution in monsoonal Central Asia during the last 50,000 years. *Quaternary Sci Rev* 25:163–178.
- Huffman, G.J., Bolvin, D.T., Nelkin, E.J., Wolff, D.B., Adler, R.F., Gu, G., Hong, Y., Bowman, K.P. and Stocker, E.F., 2007. The TRMM multisatellite precipitation analysis (TMPA): Quasi-global, multiyear, combined-sensor precipitation estimates at fine scales. *Journal of Hydrometeorology*, 8(1), pp.38-55.
- Ivory, S.J. and Lézine, A.M., 2009. Climate and environmental change at the end of the Holocene Humid Period: A pollen record off Pakistan. *Comptes Rendus Geoscience*, 341(8), pp.760-769.
- Jia, G., Bai, Y., Yang, X., Xie, L., Wei, G., Ouyang, T., Chu, G., Liu, Z. and Peng, P.A., 2015. Biogeochemical evidence of Holocene East Asian summer and winter monsoon variability from a tropical maar lake in southern China. *Quaternary Science Reviews*, 111, pp.51-61.



- 652 Jung, S.J.A., Davies, G.R., Ganssen, G.M. and Kroon, D., 2004. Stepwise Holocene aridification  
653 in NE Africa deduced from dust-borne radiogenic isotope records. *Earth and Planetary*  
654 *Science Letters*, 221(1), pp.27-37.
- 655 Kamikuri, S.I., Motoyama, I. and Nishimura, A., 2008. Radiolarian assemblages in surface  
656 sediments along longitude 175 E in the Pacific Ocean. *Marine Micropaleontology*, 69(2),  
657 pp.151-172.
- 658 Kang, S.M., Shin, Y. and Xie, S.P., 2018. Extratropical forcing and tropical rainfall distribution:  
659 energetics framework and ocean Ekman advection. *npj Climate and Atmospheric Science*,  
660 1(1), p.2.
- 661 Kaplan, J. O. et al. Holocene carbon emissions as a result of anthropogenic land cover change.  
662 *Holocene* 21, 775–791 (2011).
- 663 Karim A, Veizer J (2002) Water balance of the Indus river basin and moisture source in the  
664 Karakoram and western Himalayas: implications from hydrogen and oxygen isotopes river  
665 water. *J Geophys Res* 107:4362, doi: 10.1029/2000JD000253.
- 666 Kathayat, G., Cheng, H., Sinha, A., Yi, L., Li, X., Zhang, H., Li, H., Ning, Y. and Edwards, R.L.,  
667 2017. The Indian monsoon variability and civilization changes in the Indian subcontinent.  
668 *Science advances*, 3(12), p.e1701296.
- 669 Kenoyer JM (1998) *Ancient Cities of the Indus Valley Civilization* (Oxford University Press,  
670 Oxford).
- 671 Kong, W., Swenson, L.M. and Chiang, J.C., 2017. Seasonal transitions and the westerly jet in the  
672 Holocene East Asian summer monsoon. *Journal of Climate*, 30(9), pp.3343-3365.
- 673 Kotlia, B.S., Singh, A.K., Joshi, L.M. and Bisht, K., 2017. Precipitation variability over  
674 Northwest Himalaya from~ 4.0 to 1.9 ka BP with likely impact on civilization in the foreland  
675 areas. *Journal of Asian Earth Sciences*.
- 676 Kumar M(2009) *Linguistics, Archaeology and the Human Past*, Occasional Paper 7, eds Osada  
677 T, Uesugi A (Research Institute for Humanity and Nature, Nakanishi Printing Co. Ltd.,  
678 Kyoto), pp. 1–75.
- 679 Lézine, A.M., Ivory, S.J., Braconnot, P. and Marti, O., 2017. Timing of the southward retreat of  
680 the ITCZ at the end of the Holocene Humid Period in Southern Arabia: Data-model  
681 comparison. *Quaternary Science Reviews*, 164, pp.68-76.
- 682 Li, Y. and Morrill, C., 2015. A Holocene East Asian winter monsoon record at the southern edge  
683 of the Gobi Desert and its comparison with a transient simulation. *Climate dynamics*, 45(5-  
684 6), pp.1219-1234.
- 685 Lückge, A., Doose-Rolinski, H., Khan, A.A., Schulz, H. and Von Rad, U., 2001. Monsoonal  
686 variability in the northeastern Arabian Sea during the past 5000years: geochemical evidence  
687 from laminated sediments. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 167(3),  
688 pp.273-286.
- 689 Luis, A.J. and Kawamura, H., 2004. Air-sea interaction, coastal circulation and primary  
690 production in the eastern Arabian Sea: a review. *Journal of Oceanography*, 60(2), pp.205-  
691 218.
- 692 MacDonald, G., 2011. Potential influence of the Pacific Ocean on the Indian summer monsoon  
693 and Harappan decline. *Quaternary International*, 229(1-2), pp.140-148.
- 694 Madella M, Fuller DQ (2006) *Palaeoecology and the Harappan Civilisation of South Asia: a*  
695 *reconsideration*. *Quaternary Sci Rev* 25:1283–1301.
- 696 Madhupratap, M., Kumar, S.P., Bhattathiri, P.M.A., Kumar, M.D., Raghukumar, S., Nair,  
697 K.K.C. and Ramaiah, N., 1996. Mechanism of the biological response to winter cooling in



- 698 the northeastern Arabian Sea. *Nature*, 384(6609), p.549.
- 699 Mallah QH (2010) Current Studies on the Indus Civilization Rohn-Manohar Indus Project Series,  
700 eds Osada T, Uesugi A (Manohar Publishers, India), pp 27–76.
- 701 Mann, M. E., Zhang, Z., Rutherford, S., Bradley, R. S., Hughes, M. K., Shindell, D., Ammann,  
702 C., Faluvegi, G., and Ni, F.: Global signatures and dynamical origins of the Little Ice Age  
703 and Medieval Climate Anomaly, *Science*, 326, 1256–1260, 2009.
- 704 Marcott, S. A., Shakun, J. D., Clark, P. U. & Mix, A. C. A reconstruction of regional and global  
705 temperature for the past 11,300 years. *Science* 339, 1198–1201 (2013).
- 706 McGee, D., Donohoe, A., Marshall, J. and Ferreira, D., 2014. Changes in ITCZ location and  
707 cross-equatorial heat transport at the Last Glacial Maximum, Heinrich Stadial 1, and the mid-  
708 Holocene. *Earth and Planetary Science Letters*, 390, pp.69–79.
- 709 McGregor, H. V. et al. (2015), Robust global ocean cooling trend for the pre-industrial Common  
710 Era, *Nature Geosci.*, 8(9), 671–677, doi:10.1038/ngeo2510.
- 711 Meadow RH (1991) Harappa excavations 1986–1990: a multidisciplinary approach to Third  
712 Millennium urbanism. Madison: Prehistory Press. 275 pp.
- 713 Mughal MR (1996) Pakistan Archaeology Vol. 29, eds Iqbal F, Khan MA, Hassan M  
714 (Department of Archaeology and Museums, Pakistan; Karachi).
- 715 Mughal, M.R., 1997. Ancient Cholistan: archaeology and architecture. Ferozsons.
- 716 Muntazir Mehdi, Syed & Pant, Naresh & Saini, Hari & Mujtaba, S & Pande, Prabhas. (2016).  
717 Identification of Palaeochannel Configuration in the Saraswati River Basin in Parts of  
718 Haryana and Rajasthan, India, through Digital Remote Sensing and GIS. *Episodes*. 39.  
719 10.18814/epiugs/2016/v39i1/89234.
- 720 Munz, P.M., Siccha, M., Lückge, A., Böll, A., Kucera, M. and Schulz, H., 2015. Decadal-  
721 resolution record of winter monsoon intensity over the last two millennia from planktic  
722 foraminiferal assemblages in the northeastern Arabian Sea. *The Holocene*, 25(11), pp.1756-  
723 1771.
- 724 Nagashima, K., Tada, R. and Toyoda, S., 2013. Westerly jet-East Asian summer monsoon  
725 connection during the Holocene. *Geochemistry, Geophysics, Geosystems*, 14(12), pp.5041-  
726 5053.
- 727 Nevle, R.J. and Bird, D.K., 2008. Effects of syn-pandemic fire reduction and reforestation in the  
728 tropical Americas on atmospheric CO<sub>2</sub> during European conquest. *Palaeogeography,*  
729 *Palaeoclimatology, Palaeoecology*, 264(1-2), pp.25–38.
- 730 O'Brien, S.R., Mayewski, P.A., Meeker, L.D., Meese, D.A., Twickler, M.S. and Whitlow, S.I.,  
731 1995. Complexity of Holocene climate as reconstructed from a Greenland ice core. *Science*,  
732 270(5244), pp.1962–1964.
- 733 Olsen, J., Anderson, N.J. and Knudsen, M.F., 2012. Variability of the North Atlantic Oscillation  
734 over the past 5,200 years. *Nature Geoscience*, 5(11), pp.808–812.
- 735 Orsi, W. D. et al. Ecophysiology of uncultivated marine euryarchaea is linked to particulate  
736 organic matter. *ISME J* 9, 1747–1763, (2015).
- 737 Orsi, W.D., Coolen, M.J., Wuchter, C., He, L., More, K.D., Irigoien, X., Chust, G., Johnson, C.,  
738 Hemingway, J.D., Lee, M., Galy, V., Giosan, L., 2017. Climate oscillations reflected within  
739 the microbiome of Arabian Sea sediments. *Scientific Reports*, 7(1), p.6040.
- 740 Orengo, H.A. and Petrie, C.A., 2017. Large-scale, multi-temporal remote sensing of palaeo-river  
741 networks: a case study from northwest India and its implications for the Indus civilisation.  
742 *Remote Sensing*, 9(7), p.735.
- 743 Petrie, C.A. and Bates, J., 2017. 'Multi-cropping', Intercropping and Adaptation to Variable



- 744 Environments in Indus South Asia. *Journal of World Prehistory*, 30(2), pp.81-130.
- 745 Petrie, C.A., Singh, R.N., Bates, J., Dixit, Y., French, C.A., Hodell, D.A., Jones, P.J., Lancelotti,  
746 C., Lynam, F., Neogi, S. and Pandey, A.K., 2017. Adaptation to Variable Environments,  
747 Resilience to Climate Change: Investigating Land, Water and Settlement in Indus Northwest  
748 India. *Current Anthropology*, 58(1)
- 749 Pokharia, A. K., Kharakwal, J. S., & Srivastava, A. (2014). Archaeobotanical evidence of millets  
750 in the Indian subcontinent with some observations on their role in the Indus civilization.  
751 *Journal of Archaeological Science* 42: 442-455.
- 752 Ponton, C., Giosan, L., Eglinton, T.I., Fuller, D.Q., Johnson, J.E., Kumar, P., and Collett, T.S.,  
753 2012, Holocene aridification of India: Geophysical Research Letters, v. 39, L03704,  
754 doi:10.1029/2011GL050722.
- 755 Possehl GL (1999) *Indus Age. The Beginnings* (University of Pennsylvania Press, Philadelphia).
- 756 Possehl GL (2000) in *The drying up of the Sarasvati: environmental disruption in South Asian*  
757 *prehistory. Environmental Disaster and the Archaeology of Human Response*, eds Bawden  
758 G, Reyecraft M. (Maxwell Museum of Anthropology, University of New Mexico,  
759 Albuquerque, NM) Paper no. 7.
- 760 Possehl GL (2002) *The Indus Civilization: A Contemporary Perspective* (Altamira Press,  
761 Lanham, MD).
- 762 Prasad, S. and Enzel, Y., 2006. Holocene paleoclimates of India. *Quaternary Research*, 66(3),  
763 pp.442-453.
- 764 Prasad, S., A. Anoop, N. Riedel, S. Sarkar, P. Menzel, N. Basavaiah, R. Krishnan, D. Fuller, B.  
765 Plessen, B. Gaye, U. Rohl, H. Wilkes, D. Sachse, R. Sawant, M.G. Wiesner, M. Stebich,  
766 (2014), Prolonged monsoon droughts and links to Indo-Pacific warm pool: a Holocene record  
767 from Lonar Lake, central India *Earth Planet. Sci. Lett.*, 391, pp. 171–182
- 768 Pruesse, E. et al. SILVA: a comprehensive online resource for quality checked and aligned  
769 ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Research* 35, 7188-  
770 7196, (2007).
- 771 Purcell, J. E. (2005) Climate effects on formation of jellyfish and ctenophore blooms: a review.  
772 *Journal of the Marine Biological Association of the United Kingdom*. 85, 461-476.
- 773 Rao, R.R., Molinari, R.L. and Festa, J.F., 1989. Evolution of the climatological near-surface  
774 thermal structure of the tropical Indian Ocean: 1. Description of mean monthly mixed layer  
775 depth, and sea surface temperature, surface current, and surface meteorological fields.  
776 *Journal of Geophysical Research: Oceans*, 94(C8), pp.10801-10815.
- 777 Ratnagar, S., 2004. *Trading encounters: From the Euphrates to the Indus in the Bronze Age*.  
778 Oxford University Press.
- 779 Ravenstein, E. 1885. The laws of migration. *Journal of the Statistical Society* 48.167-235.
- 780 Ravenstein, E. 1889. The laws of migration: second paper. *Journal of the Royal Statistical*  
781 *Society* 52:241-305.
- 782 Reimer PJ, et al. (2013) Intcal13 and Marine13 Radiocarbon Age Calibration Curves 0-50,000  
783 Years Cal BP. *Radiocarbon* 55(4): 1869–1887.
- 784 Rho, M., Tang, H. & Ye, Y. FragGeneScan: predicting genes in short and error-prone reads.  
785 *Nucleic Acids Res* 38, e191, (2010).
- 786 Roberts N, Eastwood WJ, Kuzucuoğlu C, Fiorentino G, Caracuta V (2011) Climatic, vegetation  
787 and cultural change in the eastern Mediterranean during the mid-Holocene environmental  
788 transition. *Holocene* 21(1):147–162.
- 789 Rosen, A.M., 2007. *Civilizing climate: social responses to climate change in the ancient Near*





- 790 East. Rowman Altamira.
- 791 Sagoo, N. and Storelvmo, T., 2017. Testing the Sensitivity of Past Climates to the Indirect
- 792 Effects of Dust. *Geophysical Research Letters*.
- 793 Sarkar, S., Prasad, S., Wilkes, H., Riedel, N., Stebich, M., Basavaiah, N., and Sachse, D.:
- 794 Monsoon source shifts during the drying mid-Holocene: biomarker isotope based evidence
- 795 from the core “monsoon zone” (CMZ) of India, *Quaternary Sci. Rev.*, 123, 144–157, 2015.
- 796 Schug, G.R., Blevins, K.E., Cox, B., Gray, K. and Mushrif-Tripathy, V., 2013. Infection, disease,
- 797 and biosocial processes at the end of the Indus Civilization. *PLoS One*, 8(12), p.e84814.
- 798 Shadrin, A.M., Kholodova, M.V. and Pavlov, D.S., 2010, June. Geographic distribution and
- 799 molecular genetic identification of the parasite of the genus *Ichthyodinium* causing mass
- 800 mortality of fish eggs and larvae in coastal waters of Vietnam. In *Doklady Biological*
- 801 *Sciences* (Vol. 432, No. 1, pp. 220-223). SP MAIK Nauka/Interperiodica.
- 802 Shaffer, J.G., 1992, The Indus Valley, Baluchistan, and Helmand traditions: Neolithic through
- 803 Bronze Age, in Ehrich, R.W., ed., *Chronologies in Old World archaeology: Volumes 1 and 2*
- 804 (third edition): Chicago, Illinois, University of Chicago Press, p. I.441–I.464, II.425–II.446.
- 805 Schiebel, R., Zeltner, A., Treppke, U.F., Wanek, J.J., Bollmann, J., Rixen, T. and Hemleben, C.,
- 806 2004. Distribution of diatoms, coccolithophores and planktic foraminifers along a trophic
- 807 gradient during SW monsoon in the Arabian Sea. *Marine Micropaleontology*, 51(3), pp.345-
- 808 371.
- 809 Schneider T, Bischoff T, Haug GH (2014) Migrations and dynamics of the intertropical
- 810 convergence zone. *Nature* 513(7516):45–53.
- 811 Schulz, H., von Rad, U. and Ittekkot, V., 2002. Planktic foraminifera, particle flux and oceanic
- 812 productivity off Pakistan, NE Arabian Sea: modern analogues and application to the
- 813 palaeoclimatic record. *Geological Society, London, Special Publications*, 195(1), pp.499-
- 814 516.
- 815 Shanahan, T.M., McKay, N.P., Hughen, K.A., Overpeck, J.T., Otto-Bliesner, B., Heil, C.W.,
- 816 King, J., Scholz, C.A. and Peck, J., 2015. The time-transgressive termination of the African
- 817 Humid Period. *Nature Geoscience*, 8(2), pp.140-144.
- 818 Singh, G., 1971. The Indus Valley culture seen in the context of postglacial climatic and
- 819 ecological studies in north-west India. *Archaeology and Physical Anthropology of Oceania* 6,
- 820 177–189.
- 821 Singh, A., Thomsen, K.J., Sinha, R., Buylaert, J.P., Carter, A., Mark, D.F., Mason, P.J.,
- 822 Densmore, A.L., Murray, A.S., Jain, M. and Paul, D., 2017. Counter-intuitive influence of
- 823 Himalayan river morphodynamics on Indus Civilisation urban settlements. *Nature*
- 824 *communications*, 8(1), p.1617.
- 825 Sirocko, F., 2015, Winter climate and weather conditions during the Little-Ice-Age-like cooling
- 826 events of the Holocene: implications for the spread of Neolithisation? In Meller et al.
- 827 (Editors) “2200BC - A climatic breakdown as a cause for the collapse of the old world?”,
- 828 Tugengen des Landesmuseum für Vorgeschichte Halle, 12/II, 978-3-944507-29-3
- 829 Smith, M.C., Singarayer, J.S., Valdes, P.J., Kaplan, J.O. and Branch, N.P., 2016. The
- 830 biogeophysical climatic impacts of anthropogenic land use change during the Holocene.
- 831 *Climate of the Past*, 12(4), pp.923-941.
- 832 Solomina, O.N., Bradley, R.S., Hodgson, D.A., Ivy-Ochs, S., Jomelli, V., Mackintosh, A.N.,
- 833 Nesje, A., Owen, L.A., Wanner, H., Wiles, G.C. and Young, N.E., 2015. Holocene glacier
- 834 fluctuations. *Quaternary Science Reviews*, 111, pp.9-34.
- 835 Souza-Egipsy, V. et al. Prokaryotic community structure in algal photosynthetic biofilms from



- 836 extreme acidic streams in Rio Tinto (Huelva, Spain). *Int Microbiol* 11, 251-260 (2009).
- 837 Staubwasser, M., Sirocko, F., Grootes, P.M. and Erlenkeuser, H., 2002. South Asian monsoon  
838 climate change and radiocarbon in the Arabian Sea during early and middle Holocene.  
839 *Paleoceanography and Paleoclimatology*, 17(4).
- 840 Staubwasser M, Sirocko F, Grootes PM, Segl M (2003) Climate change at the 4.2 ka BP  
841 termination of the Indus valley civilization and Holocene south Asian monsoon variability.  
842 *Geophys. Res. Lett.* 30: 1425
- 843 Stein, M. A., 1931. An archaeological tour of Gedrosia. *Memoires of the Archaeological Survey*  
844 *of India* 43. New Delhi: Government of India Press.
- 845 Steinhilber et al., *Proc. Natl. Acad. Sci. U.S.A.* 109, 5967 (2012).
- 846 Stuiver, M., Braziunas, T.F., 1989. Atmospheric  $^{14}\text{C}$  and century-scale solar oscillations. *Nature*  
847 388, 405– 407.
- 848 Thomas, Z.A., 2016. Using natural archives to detect climate and environmental tipping points in  
849 the Earth system. *Quaternary Science Reviews*, 152, pp.60-71.
- 850 Thirumalai, K., Quinn, T.M., Okumura, Y., Richey, J.N., Partin, J.W., Poore, R.Z. and Moreno-  
851 Chamarro, E., 2018. Pronounced centennial-scale Atlantic Ocean climate variability  
852 correlated with Western Hemisphere hydroclimate. *Nature Communications*, 9(1), p.392.
- 853 Tierney, J.E., Pausata, F.S., deMenocal, P.B., 2017. Rainfall regimes of the Green Sahara.  
854 *Science advances*, 3(1), p.e1601503.
- 855 Uppala, S.M., Kållberg, P.W., Simmons, A.J., Andrae, U., Bechtold, V.D., Fiorino, M., Gibson,  
856 J.K., Haseler, J., Hernandez, A., Kelly, G.A. and Li, X., 2005. The ERA-40 re-analysis.  
857 *Quarterly Journal of the Royal Meteorological Society*, 131(612), pp.2961-3012.
- 858 van Dijk, W.M.; Densmore, A.L.; Singh, A.; Gupta, S.; Sinha, R.; Mason, P.J.; Joshi, S.K.;  
859 Nayak, N.; Kumar, M.; Shekhar, S.; et al. Linking the morphology of fluvial fan systems to  
860 aquifer stratigraphy in the Sutlej-Yamuna plain of northwest India. *J. Geophys. Res-Earth*  
861 2016, 121, 201–222.
- 862 Van Hengstum, P.J., Donnelly, J.P., Fall, P.L., Toomey, M.R., Albury, N.A. and Kakuk, B.,  
863 2016. The intertropical convergence zone modulates intense hurricane strikes on the western  
864 North Atlantic margin. *Scientific reports*, 6, p.21728.
- 865 von Rad, U., Schaaf, M., Michels, K.H., Schulz, H., Berger, W.H. and Sirocko, F., 1999. A  
866 5000-yr record of climate change in varved sediments from the oxygen minimum zone off  
867 Pakistan, Northeastern Arabian Sea. *Quaternary research*, 51(1), pp.39-53.
- 868 Wanner, H., Beer, J., Bütikofer, J., Crowley, T.J., Cubasch, U., Flückiger, J., Goosse, H.,  
869 Grosjean, M., Joos, F., Kaplan, J.O. and Küttel, M., 2008. Mid-to Late Holocene climate  
870 change: an overview. *Quaternary Science Reviews*, 27(19), pp.1791-1828.
- 871 Wang L, Li J, Lu H, Gu Z, Rioual P, Hao Q, Mackay AW, Jiang W, Cai B, Xu B, Han J, Chu G  
872 (2012) The East Asian winter monsoon over the last 15, 000 years: its links to high-latitudes  
873 and tropical climate systems and complex correlation to the summer monsoon. *Quat Sci Rev*  
874 32:131–142
- 875 Wang, Y., Liu, X. and Herzschuh, U., 2010. Asynchronous evolution of the Indian and East  
876 Asian Summer Monsoon indicated by Holocene moisture patterns in monsoonal central Asia.  
877 *Earth-Science Reviews*, 103(3), pp.135-153.
- 878 Weber, S.A., 2003. Archaeobotany at Harappa: indications for change. In: Weber, S.A., Belcher,  
879 W.R. (Eds.), *Indus Ethnobiology. New Perspectives from the Field*. Lexington Books,  
880 Lanham, pp. 175–198.
- 881 Weber, Steven A., Tim Barela, and Heather Lehman. 2010. Ecological continuity: an explanation





- 882 for agricultural diversity in the Indus Civilisation and beyond. *Man and Environment*  
883 35(1):62–75.
- 884 Werner, D. ed., 1977. *The biology of diatoms* (Vol. 13). Univ of California Press.
- 885 Wirth, S.B., Glur, L., Gilli, A. and Anselmetti, F.S., 2013. Holocene flood frequency across the  
886 Central Alps–solar forcing and evidence for variations in North Atlantic atmospheric  
887 circulation. *Quaternary Science Reviews*, 80, pp.112–128.
- 888 Wirtz, K.W., Lohmann, G., Bernhardt, K. and Lemmen, C., 2010. Mid-Holocene regional  
889 reorganization of climate variability: Analyses of proxy data in the frequency domain.  
890 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 298(3), pp.189–200.
- 891 Worden A. Z., Follows M. J., Giovannoni S. J., Wilken S., Zimmerman A. E., Keeling P. J.  
892 (2015) Rethinking the marine carbon cycle: Factoring in the multifarious lifestyles of  
893 microbes. *Science* 347, 735–745.
- 894 Wright RP (2010) *The Ancient Indus: Urbanism, Economy and Society* (Cambridge, Cambridge  
895 University Press).
- 896 Wright RP, Schuldenrein J, Mughal MR (2005) *South Asian Archaeology 2001*, eds C Jarrige  
897 and V Lefèvre (CNRS, Paris), pp 327–33
- 898 Wright RP, Bryson R, Schuldenrein J (2008) Water supply and history: Harappa and the Beas  
899 regional survey. *Antiquity* 82:37–48.
- 900 Yashpal, S.B.; Sood, R.K.; Agarwal, D.P. Remote sensing of the ‘Lost’ Sarasvati River. *P Indian*  
901 *AS-Earth* 1980, 89, 317–331. Zielinski, P. A. Mayewski, L. D. Meeker, S. Whitlow, M. S.  
902 Twickler, 1996 *Quat. Res.* 45, 109
- 903 Zielinski, G.A., Mayewski, P.A., Meeker, L.D., Whitlow, S. and Twickler, M.S., 1996. A  
904 110,000-yr record of explosive volcanism from the GISP2 (Greenland) ice core. *Quaternary*  
905 *Research*, 45(2), pp.109–118.
- 906



907 Figure Captions

908

909 Fig. 1. Physiography and precipitation sources for the Harappan domain. The extent of the Indus  
 910 basin and Ghaggar-Hakra (G-H) interfluve are shown with purple and brown masks,  
 911 respectively. Locations for the cores discussed in the text are shown.

912

913 Fig. 2. Modern seasonal climatology for South Asia. Average precipitation as well as wind  
 914 direction and intensity for the summer (June-July-August or JJA) and winter (December-  
 915 January-February or DJF) months are presented in the left and right panels, respectively. Note  
 916 the differences in scales between panels for both rainfall and winds. Data used come from the  
 917 ERA-40 reanalysis dataset (Uppala et al., 2005) for winds (averaged from 1958-2001) and the  
 918 TRMM dataset (Huffman et al., 2007) for rainfall (averaged from 1998-2014). The white box  
 919 encompasses the upper G-H interfluve.

920

921 Fig. 3. Holocene variability in plankton communities as reflected by their sedimentary DNA  
 922 factor loadings and winter mixing-sensitive % *G. falconensis* in core Indus 11C in the NE  
 923 Arabian Sea (see text). Relative chlorophyll biosynthesis proteins abundances are also shown.  
 924 Sea level points are from Camoin et al. (2004); SSTs are from Dooze-Rolinski et al. (2001); and  
 925 *G. falconensis* census from the NW Arabian Sea is from Schulz et al. (2002). Triangles show  
 926 radiocarbon dates. The Early Neoglacial Anomaly (ENA) is shaded.

927

928 Fig. 4. Northern Hemisphere hydroclimatic conditions since the middle Holocene. ENA interval  
 929 is shaded. From high to low: Greenland dust from non-sea-salt  $K^+$  showing the strength of the  
 930 Siberian Anticyclone (O'Brien et al., 1995); NAO proxy reconstruction (Olsen et al., 2012) and  
 931 negative NAO-indicative floods in S Alps (Wirth et al., 2013); grainsize-based hurricane  
 932 reconstruction in the N Atlantic (van Hengstum et al., 2016); interhemispheric temperature  
 933 anomaly (Marcot et al., 2013); ITCZ reconstruction at the Cariaco Basin (Haug et al., 2011);  
 934 winter monsoon paleo-DNA-based reconstruction for the NE Arabian Sea (this study);  
 935 speleothem  $\delta^{18}O$ -based precipitation reconstruction for northern Levant (Cheng et al., 2015); and  
 936 stacked lake isotope records as a proxy precipitation-evaporation regimes over Middle East and  
 937 Iran (Roberts et al., 2011).

938


939 Fig. 5. Monsoon hydroclimate changes since the middle Holocene and changes in settlement  
 940 distribution on the Ghaggar-Hakra interfluve. From high to low: variability in summer monsoon  
 941 calculated as 200-year window moving standard deviation of the detrended monsoon record of  
 942 Katahayat et al. (2017); speleothem  $\delta^{18}O$ -based summer monsoon reconstruction of Katahayat et  
 943 al. (2017); lacustrine gastropod  $\delta^{18}O$ -based summer monsoon reconstruction of Dixit et al.  
 944 (2014); changes in the number of settlements on the Ghaggar-Hakra interfluve as a function of  
 945 size and location; and winter monsoon paleo-DNA-based reconstruction for the NE Arabian Sea



946 (this study). ENA interval is shaded and durations for Early (E), Mature (M) and Late (L)  
947 Harappan phases are shown with dashed lines.




948 Fig. 1  
 949



950  
 951  
 952




953 Fig. 2.  
954






961 Fig. 3.  
 962



963  
 964



965 Fig. 4.  
 966




967





968 Fig. 5.  
 969



970