- Siberian tree-ring and stable isotope proxies as indicators of temperature and moisture
 changes after major stratospheric volcanic eruptions

-	
4	Olga V. Churakova (Sidorova) ^{1,2*} , Marina V. Fonti ² , Matthias Saurer ^{3,4} , Sébastien Guillet ¹ ,
5	Christophe Corona ⁵ , Patrick Fonti ³ , Vladimir S. Myglan ⁶ , Alexander V. Kirdyanov ^{2,7,8} ,
6	Oksana V. Naumova ⁶ , Dmitriy V. Ovchinnikov ⁷ , Alexander V. Shashkin ^{2,7} , Irina P. Panyush-
7	kina ⁹ , Ulf Büntgen ^{3,8} , Malcolm K. Hughes ⁹ , Eugene A. Vaganov ^{2,7,10} , Rolf T.W. Siegwolf ^{3,4} ,
8	Markus Stoffel ^{1,11,12}
9	
10	¹ Institute for Environmental Sciences, University of Geneva, CH-1205 Geneva, 66 Bvd Carl
11	Vogt, Switzerland
12	² Institute of Ecology and Geography, Siberian Federal University RU-660049 Krasnoyarsk,
13	Svobodniy pr 79/10, Russian Federation
14	³ Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse 111,
15	CH-8903 Birmensdorf, Switzerland
16	⁴ Paul Scherrer Institute, CH- 5232 Villigen - PSI, Switzerland
17	⁵ Université Blaise Pascal, Geolab, UMR 6042 CNRS, 4 rue Ledru, F-63057 Clermont-Fer-
18	rand, France
19	⁶ Institute of Humanities, Siberian Federal University RU-660049 Krasnoyarsk, Svobodniy pr
20	82, Russian Federation
21	⁷ V.N. Sukachev Institute of Forest SB RAS, Federal Research Center "Krasnoyarsk Science
22	Center SB RAS" RU-660036 Krasnoyarsk, Akademgorodok 50, bld. 28, Russian Federation
23	⁸ Department of Geography, University of Cambridge, Downing Place, Cambridge CB2 3EN
24	⁹ Laboratory of Tree-Ring Research, University of Arizona, 1215 E. Lowell St., Tucson,
25	85721, USA

26	¹⁰ Siberian Federal University, Rectorate, RU-660049 Krasnoyarsk, Svobodniy pr 79/10, Rus-
27	sian Federation
28	¹¹ dendrolab.ch, Department of Earth Sciences, University of Geneva, 13 rue des Maraîchers,
29	CH-1205 Geneva, Switzerland
30	¹² Department F.A. Forel for Environmental and Aquatic Sciences, University of Geneva, 66
31	Boulevard Carl-Vogt, CH-1205 Geneva, Switzerland
32	
33	Corresponding author: Olga V. Churakova (Sidorova)*
34	E-Mail: olga.churakova@hotmail.com
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	

48 Abstract

49 Stratospheric volcanic eruptions have far-reaching impacts on global climate and society.

- 50 Tree rings can provide valuable climatic information on these impacts across different spatial
- 51 and temporal scales. To detect temperature and hydro-climatic changes after strong strato-
- 52 spheric Common Era (CE) volcanic eruptions for the last 1500 years (CE 535 Unknown, CE
- 53 540 Unknown, CE 1257 Samalas, CE 1640 Parker, CE 1815 Tambora, and CE 1991
- 54 Pinatubo), we measured and analyzed tree-ring width (TRW), maximum latewood density
- 55 (MXD), cell wall thickness (CWT), and δ^{13} C and δ^{18} O in tree-ring cellulose chronologies of
- 56 climate-sensitive larch trees from three different Siberian regions (Northeastern Yakutia -
- 57 YAK, Eastern Taimyr TAY, and Russian Altai ALT).
- 58 All tree-ring proxies proved to encode a significant and specific climatic signal of the grow-
- 59 ing season. Our findings suggest that TRW, MXD, and CWT show strong negative summer
- air temperature anomalies in 536, 541-542, and 1258-1259 at all studied regions. Based on
- 61 δ^{13} C, 536 was extremely humid in YAK, 537-538 in TAY. No extreme hydro-climatic anom-
- 62 alies occurred in Siberia after the volcanic eruptions in 1640, 1815 and 1991, except for 1817
- 63 in ALT. The signal stored in δ^{18} O indicated significantly lower summer sumshine duration in
- 64 542, 1258-1259 in YAK, and 536 in ALT. Our results show that trees growing at YAK and
- 65 ALT mainly responded the first year after the eruptions, whereas at TAY, the growth re-
- 66 sponse occurred after two years.

67 The fact that differences exist in climate responses to volcanic eruptions – both in space and 68 time – underlines the added value of a multiple tree-ring proxy assessment. As such, the vari-69 ous indicators used clearly help to provide a more realistic picture of the impact of volcanic 70 eruption to past climate dynamics, which is fundamental for an improved understanding of 71 climate dynamics, but also for the validation of global climate models.

- 72 Key words: Dendrochronology, δ^{13} C and δ^{18} O in tree-ring cellulose, tree-ring width, maxi-
- 73 mum latewood density, cell wall thickness, temperature, precipitation, sunshine duration, va-
- 74 por pressure deficit

75

76 1. Introduction

77 Major stratospheric volcanic eruptions can modify the Earth's radiative balance and substan-

tially cool the troposphere. This is due to the massive injection of sulphate aerosols, which

reduce surface temperatures on timescales ranging from months to years (Robock, 2000).

80 Volcanic aerosols significantly absorb terrestrial radiation and scatter incoming solar radia-

tion, resulting in a cooling that has been estimated to about 0.5°C during the two years fol-

82 lowing the Mount Pinatubo eruption in June 1991 (Hansen et al., 1996).

83 Since trees – as living organisms – are impacted in their metabolism by environmental

84 changes, their responses to these changes are recorded in the biomass, as it is found in tree-

ring parameters (Schweingruber, 1996). The decoding of tree-ring archives is used to recon-

struct past climates. A summer cooling of the Northern Hemisphere ranging from 0.6°C to

87 1.3°C has been reported after the strongest known volcanic eruptions of the past 1500 years

88 (CE 1257 Samalas, 1815 Tambora and 1991 Pinatubo) based on temperature reconstructions

using tree-ring width (TRW) and maximum latewood density (MXD) records (Briffa et al.,

90 1998; Schneider et al., 2015; Stoffel et al., 2015; Wilson et al., 2016; Esper et al., 2017, 2018;

91 Guillet et al., 2017; Barinov et al., 2018).

92 Climate simulations show significant changes in the precipitation regime after large volcanic

93 eruptions. These include, among others, rainfall deficit in monsoon prone regions and in

94 Southern Europe (Joseph and Zeng, 2011), and wetter than normal conditions in Northern

Europe (Robock and Liu 1994; Gillet et al., 2004; Peng et al., 2009; Meronen et al., 2012;

96 Iles et al., 2013; Wegmann et al., 2014). However, despite recent advances in the field, the

97 impacts of stratospheric volcanic eruptions on hydro-climatic variability at regional scales re-

98 main largely unknown. Therefore, further knowledge about moisture anomalies is critically

99 needed, especially at high-latitude sites where tree growth is mainly limited by summer tem-

100 peratures.

As dust and aerosol particles of large volcanic eruptions affect primarily the radiation regime, three major drivers of plant growth (i.e. photosynthetic active radiation (PaR), temperature and vapor pressure deficit (VPD)) will be affected by volcanic activity. This reflects in low TRW as a result of reduced photosynthesis but even more so due to low temperature. As cell division is temperature dependent, its rate (tree-ring growth) will exponentially decrease with decreasing temperature below +3°C (Körner, 2015), outweighing the "low light / low-photosynthesis" effect by far.

Furthermore, over the last years some studies using mainly carbon isotopic signals (δ^{13} C) in tree rings showed eco-physiological responses of trees to volcanic eruptions at the mid- (Battipaglia et al., 2007) or high- (Gennaretti et al., 2017) latitudes. By contrast, a combination of both carbon (δ^{13} C) and oxygen (δ^{18} O) isotopes in tree rings has been employed only rarely to trace volcanic eruptions in high-latitude or high-altitude proxy records (Churakova (Si-

113 dorova) et al., 2014).

115

114 Application of TRW, MXD, and cell wall thickness (CWT) as well as δ^{13} C and δ^{18} O in tree

116 winter and early spring temperatures at high-latitude and high-altitude sites (Kirdyanov et al.,

cellulose chronologies is a promising tool to disentangle hydro-climatic variability as well as

117 2008; Sidorova et al., 2008, 2010, 2011; Churakova (Sidorova) et al., 2014; Castagneri et al.,

118 2017). In that sense, recent CWT measurements allowed generating high-resolution, seasonal

information of water and carbon limitations on growth during springs and summers (Pa-

nyushkina et al., 2003; Sidorova et al., 2011; Fonti et al., 2013; Bryukhanova et al., 2015).

121 Depending on site conditions, δ^{13} C variations reflect light (stand density) (Loader et al.,

122 2013), water availability (soil properties) and air humidity (proximity to open waters, i.e. riv-

123 ers, lakes, swamps and orography) as these parameters have been recognized to modulate sto-

124 matal conductance (g_l) controlling carbon isotopic discrimination.

125 Depending on the study site, a decrease in the carbon isotope ratio can be expected after stratospheric volcanic eruptions due to limited photosynthetic activity and higher stomatal con-126 127 ductance, which in turn would be the result of decreased temperatures, VPD, and a reduction 128 in light intensity. By contrast, volcanic eruptions have also been credited for an increase in photosynthesis as dust and aerosol particles cause an increased light scattering, compensating 129 for the light reduction (Gu et al., 2003). A significant increase in δ^{13} C values in tree-ring cel-130 lulose should be interpreted as an indicator of drought (stomatal closure) or high photosyn-131 thesis (Farguhar et al., 1982). In the past, very little attention has been paid to the elemental 132 133 and isotopic composition of tree rings for years during which they may have been subjected to the climatic influence of powerful, but remote, and often tropical, volcanic eruptions. 134 135 In this study, we aim to fill this gap by investigating the response of different components of 136 the Siberian climate system (i.e. temperature, precipitations, VPD, and sunshine duration) to stratospheric volcanic events of the last 1500 years. By doing so, we seek to extend our un-137 138 derstanding of the effects of volcanic eruptions on climate by combining multiple climate-139 sensitive variables measured in tree rings that were clustered around the time of the major 140 volcanic eruptions (Table 1). We focus our investigation on remote tree-ring sites in Siberia, two at high latitudes (northeastern Yakutia - YAK and eastern Taimyr - TAY), and one at 141 142 high altitude (Russian Altai - ALT), for which long tree-ring chronologies were developed previously with highly climate sensitive trees. We assemble a dataset from five tree-ring 143 proxies: TRW, MXD, CWT, δ^{13} C and δ^{18} O in larch tree-ring cellulose chronologies in order 144 145 to: (1) determine the major climatic drivers of the tree-ring proxies and to evaluate their indi-146 vidual and integrative response to climate change, and to (2) reconstruct the climatic impacts of volcanic eruptions over specific periods of the past (Table 1). 147

- 148
- 149

150 **2.** Material and methods

151 *2.1. Study sites*

152 The study sites are situated in Siberia (Russian Federation), far away from industrial centers

- 153 (and 1500–3400 km apart from each other), in the zone of continuous permafrost in north-
- 154 eastern Yakutia (YAK: 69°N, 148°E) and eastern Taimyr (TAY: 70°N, 103°E), and mountain
- permafrost in Altai (ALT: 50°N, 89°E) (Fig. 1a, Table 2). Tree-ring samples were collected
- during several field trips and included old relict wood and living larch trees: *Larix cajanderi*
- 157 Mayr (up to 1216 years) in YAK, Larix gmelinii Rupr. (max. 640 years) in TAY and Larix
- 158 sibirica Ldb. (max. 950 years) in ALT. TRW chronologies have been developed and pub-
- lished in the past (Fig. 1, Hughes et al., 1999; Sidorova and Naurzbaev, 2002; Sidorova, 2003
- 160 for YAK; Naurzbaev et al., 2002; Panyushkina et al., 2003 for TAY; Myglan et al., 2008 for

161 ALT).

- 162 Due to the remote location of our study sites, we used meteorological data from monitored
- 163 weather stations located at distances ranging from 50-200 km from the sampled sites. Tem-
- 164 perature data from these weather stations are significantly correlated (r>0.91; p < 0.05) with
- 165 gridded data (<u>http://climexp.knmi.nl</u>). However, poor correlation is found with precipitation
- 166 data (r<0.45; p<0.05), which most likely is the result of local topography (Churakova (Si-
- 167 dorova) et al., 2016).
- 168 Mean annual air temperature is lower at the high-latitude YAK and TAY sites than at the
- high-altitude ALT site (Table 2). Annual precipitation is low (153-269 mm yr⁻¹) at all sites.
- 170 The growing season calculated with the tree growth threshold of $+5^{\circ}$ C (Fritts, 1976;
- 171 Schweingruber, 1996) is very short (50-120 days) at all locations (Table 2). Sunshine dura-
- 172 tion is higher at YAK and TAY (ca. 18-20 h/day in summer) compared to ALT (ca. 18 h/day
- in summer) (Sidorova et al., 2005; Myglan et al., 2008; Sidorova et al., 2011; Churakova (Si-
- 174 dorova) et al., 2014).



Fig. 1. Location of the study sites (stars) and known volcanos from the tropics (black dots)
considered in this study (a). Annual tree-ring width index (light lines) and smoothed by 51year Hamming window (bold lines) from the northeastern Yakutia (YAK - blue, b) (Hughes
et al., 1999; Sidorova and Naurzbaev, 2002; Sidorova, 2003), eastern Taimyr (TAY - green,
c) (Naurzbaev et al., 2002), and Russian Altai (ALT - red, d) (Myglan et al., 2009). Photos
show the larch stnads at YAK, TAY (M.M. Naurzbaev) and ALT (V.S. Myglan) sites.

182

183 2.2. Selection of volcanic events and larch subsamples

Identification of the events used in this study was based on volcanic aerosols deposited in ice
core records (Zielinski 1994; Robock 2000), and more precisely on Toohey and Sigl (2017),
where the authors listed the top 20 eruptions over the past 2000 years, based on volcanic

187 stratospheric sulfur injection (VSSI). From that list, we selected those reconstructed VSSI

188	and events that are well recorded in tree-ring proxies and may thus have had a noticeable im-
189	pact on the forest ecosystems in high-latitude and high-altitude regions (Briffa et al., 1998;
190	D'Arrigo et al., 2001; Churakova (Sidorova) et al., 2014; Büntgen et al., 2016; Gennaretti et
191	al., 2017; Helama et al., 2018). Therefore, based on our previously published TRW and de-
192	veloped MXD, CWT, δ^{13} C and δ^{18} O in tree-ring cellulose chronologies, we selected the peri-
193	ods CE 520-560, 1242-1286, 1625-1660, 1790-1835, and 1950-2000 with strong volcanic
194	eruptions in CE 535, 540, 1257, 1640, 1815, and 1991, as they have had far-reaching climatic
195	effects (Table 1). The recent period 1950-2000 is used to calibrate the tree-ring proxy against
196	available climate data.
197	Tree-ring material was prepared from the 2000-year TRW chronologies available at each of
198	site from the previous studies (Fig. 1 b-d). According to the level of conservation of the mate-
199	rial, the largest possible number of samples was prepared for each of the proxies. Unlike
200	TRW, which could be measured on virtually all samples, some of the material was not availa-
201	ble with sufficient quality to allow for tree-ring anatomy and stable isotope analysis. We
202	therefore use a smaller sample size for CWT ($n=4$) and stable isotopes ($n=4$) than for TRW
203	(n=12) or MXD $(n=12)$. Nonetheless, replications are still comparable with those used in ref-
204	erence papers on stable isotopes and CWT (Loader et al., 1997; Panyushkina et al., 2003).
205	

Study period	Date of eruption	Volcano	Volcanic	Location,	References		
(CE)	Month/Day/Year	name	Explosivity	coordinates			
			Index (VEI)				
520-560	NA/NA/535	Unknown	?	Unknown	Stothers, 1984		
	NA/NA/540	Unknown	?	Unknown	Sigl et al., 2015; Toohey, Sigl 2017		
1242-1286	May-October/NA/ 1257	Samalas	7	Indonesia, 8.42°N, 116.47°E	Stothers, 2000; Lavigne et al., 2013; Sigl et al.,		
					2015		
1625-1660	December/26/1640	Parker	5	Philippines, 6°N, 124°E	Zielinski et al., 1994; 2000		
1790-1835	April/10/1815	Tambora	7	Indonesia, 8°S, 118°E	Zielinski et al., 1994; 2000		
1950 - 2000	June/15/1991	Pinatubo	6	Philippines, 15°N, 120°E	Zielinski et al., 1994; Sigl et al., 2015		

Table 1. List of stratospheric volcanic eruptions used in the study.

213 Table 2. Tree-ring sites in northeastern Yakutia (YAK), eastern Taimyr (TAY), and Altai (ALT) and weather stations used in the study. Monthly

214 air temperature (T, °C), precipitation (P, mm), sunshine duration (S, h/month) and vapor pressure deficit (VPD, kPa) data were downloaded from

215 t	he meteorological	database:	http://	/aisori	.meteo.r	u/ClimateR.
-------	-------------------	-----------	---------	---------	----------	-------------

Site	Tree	Location	Weather	Meteorological parameters			Length of	Thawing	Annual air	Annual	
	species		station					growing	permafrost	temperature	precipitation
				T (°C)	P (mm)	S (h/month)	VPD (kPa)	season	depth	(°C)	(mm)
				Periods	(IIIII)	(in month)	(KI ŭ)	(day)	(max, cm)		
ҮАК	Larix cajanderi Mayr.	69°N, 148°E	Chokurdach 62°N, 147°E, 61 m. a.s.l.	1950- 2000	1966- 2000	1961-2000	1950- 2000	50-70*	20-50*	-14.7	205
TAY	Larix gmelinii Rupr.	70°N, 103°E	Khatanga 71°N, 102°E, 33m. a.s.l.	1950- 2000	1966- 2000	1961-2000	1950- 2000	90**	40-60**	-13.2	269
ALT	<i>Larix</i> sibirica Ledeb.	50°N, 89°E	Mugur Aksy 50°N, 90°E 1850 m. a.s.l.	1963- 2000	1966- 2000			90-120***	80-100***	-2.7	153
			Kosh-Agach 50°N, 88°E 1758 m.a.s.l.			1961-2000	1950- 2000				

216 *Abaimov et al., 1996; Hughes et al., 1999; Churakova (Sidorova) et al., 2016

217 **Naurzbaev et al., 2002

218 ***Sidorova et al., 2011

219 2.3. Tree-ring width analysis

Ring width of 12 trees was re-measured for each selected period. Cross-dating was checked by 220 comparison with the existing full-length 2000-yr TRW chronologies (Fig. 1). The TRW series 221 222 were standardized using the ARSTAN program (Cook and Krusic, 2008) with negative exponential curve (k>0) or a linear regression (any slope) prior to bi-weight robust averaging (Cook and 223 Kairiukstis, 1990). Signal strength in the regional TRW chronologies was assessed with the Ex-224 225 pressed Population Signal (EPS) statistics as it measures how well the finite sample chronology 226 compares with a theoretical population chronology with an infinite number of trees (Wigley et 227 al., 1984). Mean inter-series correlation (RBAR) and EPS values of stable isotope chronologies 228 were calculated for the period 1950-2000, for which individual trees were analyzed separately. All series have RBAR ranges between 0.59 and 0.87, and the common signal exceeds the EPS 229 threshold of 0.85. Before 1950, we used pooled cellulose only. For all other tree-ring parameters 230 231 and studied periods, the EPS exceeds the threshold of 0.85, and RBAR values range from 0.63 to 232 0.94.

233

234 2.4. Image analysis of cell wall thickness (CWT)

Analysis of wood anatomy was performed for all studied periods with an AxioVision scanner 235 (Carl Zeiss, Germany). Micro-sections were prepared using a sliding microtome and stained with 236 237 methyl blue (Furst, 1979). Tracheids in each tree ring were measured along five radial files of cells (Munro et al., 1996; Vaganov et al., 2006) selected for their larger tangential cell diameter 238 239 (T). For each tracheid, CWT was computed separately. In a second step, tracheid anatomical parameters were averaged for each tree ring. Site chronologies are presented for the complete an-240 nual ring chronology without standardization due to the lack of low-frequency trend. CWT data 241 242 from ALT for the periods 1790-1835 and 1950-2000 were used from the past studies (Sidorova et al., 2011; Fonti et al., 2013) and for YAK for the period from 1600-1980 from Panyushkina et 243

- al. (2003). Unfortunately, the remaining sample material for the CE 536 ring at TAY was insuffi-
- cient to produce a clear signal. As a result, CWT is missing for CE 536 at TAY (Fig. 2).
- 246
- 247 2.5. Maximum latewood density (MXD)

Maximum latewood density chronologies from ALT were available continuously for the period 248 CE 600-2007 from Schneider et al. (2015) and Kirdyanov A.V. (personal communication), and 249 from YAK and TAY for the period CE 1790-2004 from Sidorova et al. (2010). For any other pe-250 251 riods, at least six cross-sections and for CE 520-560 four sections are used. The wood is subsampled with a double-bladed saw at 1.2 mm thickness with the angle to the fiber direction. The 252 253 samples were exposed to X-rays for 35-60 min (Schweingruber, 1996). MXD measurements were obtained at 0.01 mm resolution and brightness variations calibrated to g/cm³ (Lenz et al., 254 1976: Eschbach et al., 1995) using a Walesch X-ray densitometer 2003. All MXD series were 255 256 detrended in the ARSTAN program by calculating deviation from straight-line function (Fritts, 1976). Site MXD chronologies were developed for each volcanic period using the bi-weight ro-257 258 bust averaging.

259

260 2.6. Stable carbon ($\delta^{13}C$) and oxygen ($\delta^{18}O$) isotopes in tree-ring cellulose

During photosynthetic CO₂ assimilation ¹³CO₂ is discriminated against ¹²CO₂, leaving the newly 261 produced assimilates depleted in ¹³C. The carbon isotope discrimination ($^{13}\Delta$) is partitioned in 262 the diffusional component with a = 4.4% and the biochemical fractionation with b = 27%, for 263 C3 plants, during carboxylation via Rubisco. The ${}^{13}\Delta$ is directly proportional to the c_i/c_a ratio, 264 where c_i is the leaf intercellular, and c_a the ambient CO₂ concentration. This ratio reflects the bal-265 ance between stomatal conductance (g_l) and photosynthetic rate (A_N) . A decrease in g_l at a given 266 A_N results in a decrease of ${}^{13}\Delta$, as c_i/c_a decreases and vice versa. The same is true when A_N in-267 creases or decreases at a given g₁. Since CO₂ and H₂O gas exchange are strongly interlinked with 268 the C-isotope fractionation ${}^{13}\Delta$ is controlled by the same environmental variables i.e. PaR, CO₂, 269

270 VPD and temperature (Farquhar et al., 1982, 1989; Cernusak et al., 2013). The oxygen isotopic compositions of tree-ring cellulose record the δ^{18} O of the source water derived from precipita-271 tion, which itself is related to temperature variations at middle and high latitudes (Craig, 1961; 272 Dansgaard, 1964). It is modulated by evaporation at the soil surface and to a larger degree by 273 274 evaporative and diffusion processes in leaves; the process is largely controlled by the vapor pressure deficit (Dongmann et al., 1972, Farquhar and Loyd, 1993, Cernusak et al., 2016). A further 275 step of fractionation occurs as sugar molecules are transferred to the locations of growth (Roden 276 277 et al., 2000). During the formation of organic compounds the biosynthetic fractionation leads to a positive shift of the δ^{18} O values by 27‰ relative to the leaf water (Sternberg, 2009). The oxy-278 gen isotope variation in tree-ring cellulose therefore reflects a mixed climate information, often 279 dominated by a temperature, source water or sunshine duration modulated by the VPD influence. 280 281 The cross-sections of relict wood and cores from living trees used for the TRW, MXD and CWT measurements were then selected for the isotope analyses. We analyzed four subsamples for 282 each studied period according to the standards and criteria described in Loader et al. (2013). The 283 first 50 yrs. of each sample were excluded to limit juvenile effects (McCarroll and Loader, 284 2004). After splitting annual rings with a scalpel, the whole wood samples were enclosed in filter 285 286 bags. α-cellulose extraction was performed according to the method described by Boettger et al. (2007). For the analyses of ${}^{13}C/{}^{12}C$ and ${}^{18}O/{}^{16}O$ isotope ratios, 0.2-0.3 mg and 0.5-0.6 mg of cel-287 lulose were weighed for each annual ring, into tin and silver capsules, respectively. Carbon and 288 289 oxygen isotopic ratios in cellulose were determined with an isotope ratio mass spectrometer 290 (Delta-S, Finnigan MAT, Bremen, Germany) linked to two elemental analyzers (EA-1108, and EA-1110 Carlo Erba, Italy) via a variable open split interface (CONFLO-II, Finnigan MAT, Bre-291 292 men, Germany). The ¹³C/¹²C ratio was determined separately by combustion under oxygen excess at a reactor temperature of 1020°C. Samples for ¹⁸O/¹⁶O ratio measurements were pyrolyzed 293 to CO at 1080°C (Saurer et al., 1998). The instrument was operated in the continuous flow mode 294

for both, the C and O isotopes. The isotopic values were expressed in the delta notation multi-

296 plied by 1000 relative to the international standards (Eq. 1):

297
$$\delta$$
 sample = $R_{sample}/R_{standard}$ -1 (Eq. 1)

298 where R_{sample} is the molar fraction of ${}^{13}C/{}^{12}C$ or ${}^{18}O/{}^{16}O$ ratio of the sample and $R_{standard}$ the molar

fraction of the standards, Vienna Pee Dee Belemnite (VPDB) for carbon and Vienna Standard

300 Mean Ocean Water (VSMOW) for oxygen. The precision is $\sigma \pm 0.1\%$ for carbon and $\sigma \pm 0.2\%$

301 for oxygen. To remove the atmospheric δ^{13} C trend after CE 1800 from the carbon isotope values

in tree rings (i.e. Suess effect, due to fossil fuel combustion) we used atmospheric $\delta^{13}C$ data from

303 Francey et al. (1999), http://www.cmdl.noaa.gov./info/ftpdata.html). These corrected series were

304 used for all statistical analyses. The δ^{18} O cellulose series were not detrended.

305

306 *2.7. Climatic data*

Meteorological series were obtained from local weather stations close to the study sites and used for the computation of correlation functions between tree-ring proxies and monthly climatic parameters (Table 2, http://aisori.meteo.ru/ClimateR).

310

311 2.8. Statistical analysis

All chronologies for each period were normalized to z-scores (Fig. 2). To assess post-volcanic
climate variability, we used Superposed Epoch Analysis (SEA, Panofsky and Brier, 1958) with
the five proxy chronologies available at each of the three study sites. In this study, intervals of 15
years before and 20 years after a volcanic eruption were analyzed. SEA is applied to the six annually dated volcanic eruptions (Table 1).
To test the sensitivity of the studied tree-ring parameters to climate bootstrap correlation func-

tions were computed between proxy chronologies and monthly climate predictors using the

319 'bootRes' package of R software (R Core Team 2016) for the period 1950 (1966)-2000.

320 To estimate whether volcanic years can be considered as extreme and how anomalous they are

321 compared to non-volcanic years, we computed Probability Density Functions (PDFs, Stirzaker,

322 2003) for each study site and for each tree-ring parameter over a period of 219 years for which

323 measurements are available (Fig. S1). A year is considered (very) extreme if the value of a given

324 parameter is below the $(5^{\text{th}}) 10^{\text{th}}$ percentile of the PDF.

325

326 3. Results

327 *3.1. Anomalies in tree-ring proxy chronologies after stratospheric volcanic eruptions*

328 Normalized TRW chronologies show negative deviations the year following the eruptions at all

329 studied sites (Fig. 2). Regarding CWT, a strong decrease is observed in CE 537 at all study sites.

330 Only two layers of cells were formed in CE 537 (-1.8 σ) and 541 (-2.4 σ) for YAK as compared

to the 11-20 layers of cells formed on average during "normal" years. In addition, we also ob-

serve the formation of frost rings in ALT between CE 536 and 538, as well as in 1259. An abrupt

333 CWT decrease is recorded in TAY in 537 (-3.1 σ).

Furthermore, we found decreasing MXD values at ALT (-4.4 σ) in CE 537 and YAK (-2.8 σ) in

335 CE 536. However, for TAY, data show a less pronounced pattern of MXD variation (Fig. 2). In

this regard, the sharpest decrease was observed in the CWT chronologies from YAK in CE 540

 (-1.9σ) and $541(-2.4\sigma)$, whereas the response was smaller in TAY and ALT for the same years

338 (Fig. 2). The ALT δ^{18} O chronology recorded a drastic decrease in 536 CE with - 4.8 σ (Fig. 2,

Fig. S1). A δ^{18} O decrease for YAK was found after the CE 1257 Samalas eruption in CE 1258 (-

340 1.5 σ) and in 1259 (-2.9 σ), which is opposite to the increased δ^{18} O value found in CE 1259 at

341 ALT (Fig. 2; Fig. S1).

342 Regarding the carbon isotope ratio, negative anomalies are observed in ALT already in 1258 (-

 2.3σ). The CE 540 eruption was less clearly recorded in tree-ring proxies from TAY, compared

to YAK and ALT (Fig. 2). With respect to the CE 1257 Samalas eruption (Fig. 2), the year fol-

lowing the eruption was recorded as very extreme in the TRW, MXD, δ^{18} O, while less extreme

- 346 in CWT and δ^{13} C from YAK. ALT chronologies show a synchronous decrease for all proxies
- following two years after the eruption (Fig. 2, Fig. S1).



- 358 The impacts of the more recent CE 1640 Parker, 1815 Tambora, and 1991 Pinatubo eruptions
- are, by contrast, far less obvious. In CE 1642, decreasing values are observed in all tree-ring
- 360 proxies from the high-latitude sites YAK and TAY, whereas tree-ring proxies are not clearly
- affected at ALT (Fig. 2; Fig. S1).
- Hardly any strong anomalies are observed in CE 1816 in Siberia regardless of the site and the
- 363 tree-ring parameter analyzed. The ALT δ^{13} C value (-3.3 σ) in CE 1817 and YAK MXD (-
- 364 2.4σ) in 1816 can be seen as an exception to the rule here as they evidenced extreme values,
 365 respectively (Fig. S1).
- 366 Finally, the Pinatubo eruption is mainly captured by the MXD (-2.8 σ) and CWT (-2.2 σ) chro-
- 367 nologies from YAK in CE 1992. Simultaneous decreases of all tree-ring proxies from ALT
- are observed in 1993 (Fig. 2), which, however, cannot be classified as extreme (Fig. S1).
- 369 Overall, the SEA (Fig. 3) shows that volcanic eruptions centered around CE 535, 540, 1257,
- 370 1640, 1815, and 1991 have led to decreasing values for all tree-ring proxies following next
- 371 two years afterwards. A short-term response by two years after the eruptions is observed in
- 372 the TRW and CWT proxies for TAY, while for YAK and ALT, the CWT decrease lasts
- 373 longer (up to 5-6 years in ALT and YAK, respectively) (Fig. 3). The δ^{18} O isotope chronolo-
- 374 gies (z-score) show a distinct decrease the year after the eruptions. At ALT, however, the du-
- 375 ration of negative anomalies were shorter (5 years) than at the high-latitude TAY (12 years)
- and YAK (9 years) sites. At the YAK site, two negative years followed the events, intermitted
- 377 with one positive value, to remain negative during the following 7 years. The duration of neg-
- ative anomalies recorded in δ^{13} C values (z-score) lasts also longer at the high-latitude YAK
- 379 site 10 years after the eruptions and 13 years at TAY compared to 7 years at ALT (Fig. 3).
- 380 The largest decrease in MXD values (in terms of z-score) is found at the high-latitude YAK
- site. The SEA for TRW, MXD, δ^{13} C, and CWT from YAK as well as TRW and MXD from

proxies and study sites (Fig. 3). 383 $\delta^{18}O$ 384 YAK TAY ALT 0.5 0.5 0.5 z-score z-score z-score -0.5 -0.5 -0.5 -1.5 -1.5 -1.5 $\delta^{18}O$ ALT δ^{18} O YAK δ^{18} O TAY -2.5 -2.5 -2.5 4 -2.5 15 6 15 ò 5 10 00 20 385 Year before and after the eruption Year before and after the eruption Year before and after the eruption $\delta^{13}C$ 386 0.5 0.5 0.5 -0.5 z-Score z-score -0.5 z-score -0.5 -1.5 -1.5 -1.5 δ^{13} C YAK $\delta^{13}C$ ALT $\delta^{13}C$ TAY -2.5 4 -2.5 4 -2.5 4 15 10 15 20 10 15 20 5 20 387 388 Year before and a Year before and after the eruption Year before and after the eruption the eruption CWT 0.5 0.5 0.5 -0.5 z-score e.0-z-score e.0-z-score -1.5 -1.5 -1.5 CWT YAK CWT TAY CWT ALT -2.5 -2.5 -2.5 <u>6</u> 2 15 2 2 5 2 20 9 s, 2 20 9 ĥ 20 389 390 and after th hefore eruption ar before and after the eruption Year before and eruption TRW 0.5 0.5 0.5 d a f -0.5 z-score z-score -0.5 z-score -0.5 -1.5 -1.5 -1.5 TRW YAK TRW ALT TRW TAY -2.5 4 -2.5 L -2.5 10 15 Ϋ́ 20 10 15 12 9 20 2 20 5 391 392 Year before and after the eruption Year before and after the eruption Year before and after the eruption MXD 0.5 0.5 0.5 e.0-z-score -0.5 z-score e.o-z-score -1.5 -1.5 -1.5 MXD TAY MXD ALT MXD YAK -2.5 -2.5 -2.5 -2.5 -2.5 -2.5 -2.5 0 15 0 12 9 15 9 LC. c ŝ 20 9 ĥ 20 -10 ŵ 20 393 Year before and after the eruption Year before and after the eruption Year before and after the eruption

382 ALT show a more drastic decrease of values during the first year when compared to other

- **Fig. 3.** Superposed epoch analysis (SEA) of δ^{18} O, δ^{13} C, CWT, TRW, and MXD chronologies
- 395 for the Yakutia (YAK), Taimyr (TAY), and Altai (ALT) sites, summarizing negative anoma-
- lies 15 years before and 20 years after the volcanic eruptions in CE 535, 540, 1257, 1640,
- 1815, and 1991. Statistically negative anomalies are marked with a red star (*p < 0.05).
- 398

399 **3.2.** Tree-ring proxies versus meteorological series

- 400 *3.2.1. Monthly air temperatures and sunshine duration*
- 401 Bootstrapped functions calculated for the instrumental period (1950-2000) show significant
- 402 positive correlations (p < 0.05) between TRW and MXD chronologies and mean summer
- 403 (June-July) temperatures at all sites. Temperatures at the beginning (June) and the end of the
- 404 growing season (mid-August) influenced the MXD chronology in ALT (r = 0.57) and YAK (r
- 405 = 0.55), respectively (Fig. 4). July temperatures appear as a key factor for determining tree
- 406 growth as they significantly impact CWT, δ^{13} C, and δ^{18} O (with the exception of TAY for the
- 407 latter) chronologies (r=0.28-0.60) at YAK and ALT.
- 408 Correlation analysis between July temperature and July sunshine duration indicate significant
- 409 (p < 0.05) correlation for YAK (r=0.56) and ALT (r=0.34). July sunshine duration are strongly
- 410 and positively correlated with δ^{18} O in larch tree-ring cellulose chronologies from YAK
- 411 (r=0.73) and ALT (r=0.51) for the period 1961-2000 (available sunshine duration data set).



412 Fig. 4. Significant correlation coefficients between tree-ring parameters: TRW, MXD, CWT, 413 δ^{13} C and δ^{18} O versus weather station data: temperature (T, red), precipitation (P, blue), vapor 415 pressure deficit (VPD, green), and sunshine duration (S, yellow) from September of the previ-416 ous year to August of the current year for three study sites were calculated. Table 2 lists sta-417 tions and periods used in the analysis.

418

419 *3.2.2. Monthly precipitation*

420 The strongest July precipitation signal is observed at ALT (r=-0.54) and TAY (r=-0.51) with

421 δ^{13} C chronologies (p < 0.05). In addition, the ALT data shows a significant relationship

422 (p < 0.05) between March precipitation and TRW (r=0.37) and MXD (r=0.32), whereas April

423 precipitation correlates positively with CWT (r=0.34). At YAK, July precipitation showed

424 negative relationship with δ^{18} O in tree-ring cellulose (r=-0.34; p<0.05) only.

- 425 *3.2.3. Vapor pressure deficit (VPD)*
- 426 June VPD is significantly and positively correlated with the δ^{18} O chronology from ALT
- 427 (r=0.67 p < 0.05, respectively) for the period 1950-2000. The δ^{13} C in tree-ring cellulose from
- 428 YAK correlate with July VPD only (r=0.69 p < 0.05). We did not find significant influence of
- 429 VPD in TAY tree-ring and stable isotope parameters.
- 430
- 431 *3.2.4. Synthesis of the climate data analysis*
- 432 In summary, during the instrumental period of weather station observations (Table 2) summer
- 433 temperature impacts TRW, MXD and CWT at the high-latitude sites (YAK, TAY), while
- 434 summer precipitation affects stable carbon and oxygen isotopes (YAK, TAY, ALT), sunshine
- 435 duration (YAK, ALT), and vapor pressure deficit (YAK, ALT).
- 436

437 3.3. Response of Siberian larch trees to climatic changes after the major volcanic erup438 tions

- 439 Based on the statistical analysis above for the calibration period, we assumed that these rela-
- tionships would not change over time and will provide information about climatic changes
- 441 during the past volcanic periods (Fig. 5).

442





457 *3.3.1. Temperature proxies*

We found strong negative summer air temperature anomalies at all sites after the CE 535 and 458 459 1257 volcanic eruptions. The temperature decrease was found in the TRW and CWT datasets at all sites, and also in the MXD datasets at YAK and ALT (Fig. 5). For the volcanic erup-460 461 tions in later centuries, the evidence for a decrease in temperature was not as pronounced. Whereas no strong decline of summer temperature was found at ALT in CE 1642, we observe 462 463 a slight decrease in TRW, MXD and CWT values in 1643. By contrast, a cold summer was recorded by most tree-ring parameters at YAK, except for δ^{18} O. The absence of strong cool-464 ing is even more so striking during the years that followed the CE 1815 Tambora eruption. In 465 CE 1816, only the MXD from YAK shows colder than normal conditions (Fig. 5). CE 1992 466 was recorded as a cold year in MXD and CWT from YAK, but again not at the other regions 467 468 and by other proxies.

469

470 *3.3.2. Moisture proxies: precipitation and VPD*

471 Based on the climatological analysis with the local weather stations data (Table 2, Fig. 4) for all studied sites we considered δ^{13} C in tree-ring cellulose as a proxy for precipitation and va-472 por pressure deficit changes. Yet, CWT from ALT could be considered as a proxy with mixed 473 temperature and precipitation signal (Fig. 4). Accordingly, the δ^{13} C values point to humid 474 summers at YAK in 536, 1258, 1259, 1642, and 1643, at TAY in 536-538, and 1259, and at 475 476 ALT the years of 541, 542, 1258, 1259 and 1817. Compared to other proxies and sites, the years 536-538 were neither extremely humid nor dry at ALT (Fig. 5). No negative hydrologi-477 cal anomalies were recorded after the Tambora and Pinatubo eruptions at the high-latitude 478 sites (YAK, TAY). However, positive anomalies were recorded in δ^{13} C values, pointing to 479 dry conditions at TAY in CE 1817 (Fig. 2). A rather wet summer was reconstructed for the 480

- 481 high-altitude ALT site in CE 1817 compared to 1816 (Fig. 5). Overall, there were mostly hu482 mid anomalies after the eruptions at YAK.
- 483

484 *3.3.3. Sunshine duration proxies*

Instrumental measurements of sunshine duration (Table 2) at YAK and ALT during the recent period showed a significant link with δ^{18} O cellulose. The sunshine duration is decreased after various eruptions at YAK (538, 542, 1258, and 1259) and in 536 at ALT site.

488

489 4. Discussion

490 In this paper, we analyze climatic anomalies in years following selected large volcanic erup-

491 tions using long-term tree-ring multi-proxy chronologies for δ^{13} C and δ^{18} O, TRW, MXD,

492 CWT for the high-latitude (YAK, TAY) and high-altitude (ALT) sites. Since trees as living

493 organisms respond to various climatic impacts, the carbon assimilation and growth patterns

494 accordingly leave unique "finger prints" in the photosynthates, which is recorded in the wood

in the tree rings specifically and individually for each proxy.

496

497 *4.1. Evaluation of the applied proxies in Siberian tree-ring data*

This study clearly shows that each proxy has to be analyzed and interpreted specifically for its
validity at each studied site and evaluated for its suitability for the reconstruction of abrupt
climatic changes.

The TRW in temperature-limited environments is an indirect proxy for summer temperature reconstructions, as growth is a temperature-controlled process. Temperature clearly determines the duration of the growing season and the rate of cell division (Cuny et al., 2014). Accordingly, low temperature of growing season is recorded by narrow tree rings. The upper limit of temperature is specific to tree species and biome. In most cases, tree growth is limited

by drought rather than by high temperatures, since water shortage and VPD increase with in-506 creasing temperature. Still this does not make TRW a suitable proxy to determine the influ-507 508 ence of water availability and air humidity, especially at the temperature-limited sites. MXD chronologies obtained for the Eurasian subarctic record mainly a July-August tempera-509 510 ture signal (Vaganov et al., 1999; Sidorova et al., 2010; Büntgen et al., 2016) and add valua-511 ble information about climate conditions toward the end of the growth season. Similarly, 512 CWT is an anatomical parameter, which contains information on carbon sink limitation of the 513 cambium due to extreme cold conditions (Panyushkina et al., 2003; Fonti et al., 2013; Bryu-514 khanova et al., 2015). There is a strong signal of low cell number within a growing season, for example, strong decreasing CWT in CE 537 at YAK or the formation of frost rings at ALT in 515 516 (CE 536-538, and 1259) has been shown in our study. Low $\delta^{13}C$ values can be explained by a reduction in photosynthesis caused by volcanic dust 517 veils. For the distinction whether δ^{13} C is predominantly determined by A_N or g_l the combined 518 evaluation with δ^{18} O or TRW is needed. High δ^{18} O values indicate high VPD, which induces 519 520 a reduction in stomatal conductance, reducing the back diffusion of depleted water molecules 521 from the ambient air. This confirms a sunny CE 1993 at ALT with mild weather conditions according to observational data from the closest weather station (Table 2). Interestingly, we 522 also find less negative values for δ^{13} C in the same period. This shows that the two isotopes 523 524 correlate with each other and indicates the need for a combined evaluation of the C and O isotopes (Scheidegger et al., 2000) taking into account precautions as suggested by Roden and 525 Siegwolf (2012). 526

527

528 *4.2. Lag between volcanic events and response in tree rings*

529 Most discussed events suggest a lag between the eruption and the tree-ring response for one530 year or more (Fig. 3). This lag is explained by the tree's use of stored carbohydrates, which

are the substrate for needle and early wood production. These stored carbohydrates carry the
isotopic signal of previous years and depend on their remobilization, as such the signals may
be masked in freshly produced biomass. The delayed signal could also reflect the time needed
for the dust veil to be transported to the study regions.

535

536 4.3. Temperature and sunshine duration changes after stratospheric volcanic eruptions 537 Correlation functions show that MXD and CWT (with the exception of TAY in the latter 538 case), and to a lesser extent TRW chronologies, portray the strongest signals for summer 539 (June-August) temperatures. In addition, significant information about sunshine duration can be derived from the YAK and ALT δ^{18} O series. Thus, we hypothesize that extremely narrow 540 541 TRW and very negative anomalies observed in the MXD and CWT chronologies of YAK and to a lesser extent at ALT, along with low δ^{18} O values reflect cold and low sunshine duration 542 conditions in summer. Presumably, the temperatures were below the threshold values for 543 growth over much of the growing season (Körner, 2015). This hypothesis of a generalized re-544 545 gional cooling after both eruptions is further confirmed by the occurrence of frost rings at 546 ALT site in CE 538, 1259 (Myglan et al., 2008; Guillet et al., 2017), as well as in neighboring 547 Mongolia (D'Arrigo et al., 2001). The unusual cooling in CE 536-542 is also evidenced by a very small number of cells formed at YAK (Churakova (Sidorova) et al., 2014). Although 548 δ^{18} O is an indirect proxy for needle temperature, low δ^{18} O values in CE 538, 542, 1258, and 549 550 1259 for YAK and in CE 536 for ALT are a result of low irradiation, leading to low temperature and low VPD (high stomatal conductance), both likely a result from volcanic dust veils. 551 552 Similarly, in the aftermath of the Samalas eruption, the persistence of summer cooling is lim-553 ited to CE 1258 and 1259 at the three studied sites, which is in line with findings of Guillet et

al., (2017). Interestingly, a slight decrease in oxygen isotope chronologies, which can be related to low levels of summer sunshine duration (i.e. low leaf temperatures), allows for hypothesizing that cool conditions could have prevailed.

557 For all later high-magnitude CE eruptions, temperature-sensitive tree-ring proxies do not evi-558 dence a generalized decrease in summer temperatures. Paradoxically, the impacts of the Tam-559 bora eruption, known for its triggering of a widespread "year without summer" (Harrington, 560 1992), did only induce abnormal MXD at YAK in 1816, but no anomalies are observed at TAY and ALT, except for the positive deviation of δ^{13} C at TAY and the negative anomaly at 561 562 ALT in CE 1817 (Fig. 2, Fig. 5, Fig. S1). While these findings may seem surprising, they are in line with the TRW and MXD reconstructions of Briffa et al. (1998) or Guillet et al. (2017), 563 564 who found limited impacts of the CE 1815 Tambora eruption in Eastern Siberia and Alaska 565 using TRW and MXD data only. The inclusion of CWT chronologies by Barinov et al. (2018) 566 confirms the absence of a significant cooling signal after the second largest eruption of the 567 last millennium (CE 1815) in larch trees of the Altai-Sayan mountain region. 568 Finally, in CE 1992, our results evidence cold conditions at YAK, which is consistent with 569 weather observations showing that the below-average anomalies of summer temperatures (af-570 ter Pinatubo eruption) were indeed limited to Northeastern Siberia (Robock, 2000). As both isotopes indicate a reduction in stomatal conductance, we found that warm (in agreement with 571 572 MXD and CWT) and dry conditions were prevalent at ALT at this time. This isotopic constellation was confirmed by the positive relationships between VPD and δ^{18} O and δ^{13} C at ALT. 573 574 However, temperature and sunshine duration are not always highly coherent over time due to 575 the influence of other factors, like Arctic Oscillations as suggested for Fennoscandia regions 576 by Loader et al. (2013).

577

578 *4.4. Moisture changes*

Water availability is a key parameter for Siberian trees as they are growing under extremely 579 580 continental conditions with hot summers and cold winters, and even more so with very low 581 annual precipitation (Table 2). Permafrost plays a crucial role and can be considered as a buffer for additional water sources during hot summers (Sugimoto et al., 2002; Boike et al., 582 583 2013; Saurer et al., 2016). Yet, thawed permafrost water is not always available to roots due 584 to the surficial structure of the root plate or extremely cold water temperature (close to 0° C), 585 which can hardly be utilized by trees (Churakova (Sidorova) et al., 2016). Thus, Siberian trees 586 are highly susceptible to drought, induced by dry and warm air during July and therefore the 587 stable carbon isotopes can be sensitive indicators of such conditions. After volcanic eruptions, however, low light intensity due to dust veils induce low temperatures and reduced VPD, the 588 589 driver for evapotranspiration. Under such conditions drought stress is unlikely to occur. How-590 ever, the transition phases with changes from cool and moist to warm and dry conditions are 591 more critical when drought is more likely to occur.

In our study, higher δ^{13} C values in tree-ring cellulose indicate increasing drought conditions as a consequence of reduced precipitation for two years after the CE 1815 volcanic eruption at TAY site. No further extreme hydro-climatic anomalies occurred at Siberian sites in the aftermath of the Pinatubo eruption.

596

597 4.5. Synthetized interpretation from the multi-parameter tree-ring proxies

598 Our analysis demonstrates the added value of a tree-ring derived multi-proxy approach to bet-599 ter capture the climatic variability after large volcanic eruptions. Besides the well-documented 600 effects of temperature derived from TRW and MXD, CWT, stable carbon and oxygen iso-601 topes in tree-ring cellulose provide important and complementary information about moisture 602 and sunshine duration changes (an indirect proxy for leaf temperature effective for air-to-leaf 603 VPD) after stratospheric volcanic eruptions.

Our results reveal the complex behavior of the Siberian climatic system to the stratospheric 604 605 volcanic eruptions of the Common Era. The CE 535 and CE 1257 Samalas eruptions caused 606 substantial cooling – very likely induced by dust veils (Churakova (Sidorova) et al., 2014; 607 Guillet et al., 2017; Helama et al., 2018) – as well as humid conditions at both the high-lati-608 tude and high-altitude sites. Conversely, only local and limited climate responses were ob-609 served after the CE 1641 Parker, 1815 Tambora, and 1991 Pinatubo eruptions. Similar site-610 dependent impacts referred to the coldest summers of the last millennium in the Northern 611 Hemisphere based on TRW and MXD reconstructions (Schneider et al., 2015; Stoffel et al., 612 2015; Wilson et al., 2016; Guillet et al., 2017). This absence of widespread and intense cooling or missing drastic changes in hydrological regime over vast regions of Siberia may result 613 614 from the location and strength of the volcanic eruption, atmospheric transmissivity as well as from the modulation of radiative forcing effects by regional climate variability. These results 615 616 are consistent with other regional studies, which interpreted the spatial-temporal heterogeneity of tree responses to past volcanic events (Wiles et al., 2014; Esper et al., 2017; Barinov et 617 618 al., 2018) in terms of regional climates.

619

620 **5.** Conclusions

In this study, we demonstrate that the consequences of large volcanic eruptions on climate are rather complex between sites and among events. The different locations and magnitudes of eruptions, but also regional climate variability, may explain some of this heterogeneity. We show that each tree-ring and isotope proxy alone cannot provide the full information of the volcanic impact on climate, but that they, when combined, contribute to the formation of the full picture, which is critical for a comprehensive description of climate dynamics induced by volcanism and the inclusion of these phenomena in global climate models. The analyses with a larger number of samples in the investigations of Siberian and other Northern Hemispheric sites will indeed to provide higher certainty in terms of data interpretation of climatic dynamics of these boreal regions. However, the multi-proxy approach as applied in our study also provides a strong set of complementary information to the research field, as it allows the refinement of the interpretations and thus improves our understanding of the heterogeneity of climatic signals after CE stratospheric volcanic eruptions, as recorded in multiple tree-ring and stable isotope parameters.

635

636 Author contribution: TRW analysis was performed at V.N. Sukachev Institute of Forest SB RAS by O.V. Churakova (Sidorova), D.V. Ovchinnikov, V.S. Myglan and O.V. Naumova. 637 638 CWT analysis was carried out at the V. N. Sukachev Institute of Forest SB RAS, Krasnoyarsk, Russia by M.V. Fonti and at the University of Arizona by I.P. Panyushkina. Stable iso-639 640 tope analysis was conducted at the Paul Scherrer Institute (PSI), by O.V. Churakova (Sidorova), M. Saurer, and R. Siegwolf. MXD measurements were realized with a DENDRO 641 642 Walesh 2003 densitometer at WSL and at the V.N. Sukachev Institute of Forest SB RAS, 643 Krasnovarsk, Russia by O.V. Churakova (Sidorova) and A.V. Kirdyanov. Samples from YAK 644 and TAY were collected by M.M. Naurzbaev. All authors contributed significantly to the data 645 analysis and paper writing.

646

Acknowledgements: This work was supported by Marie Curie International Incoming Fellowship [EU_ISOTREC 235122], Re-Integration Marie Curie Fellowship [909122] and UFZ
scholarship [2006], RFBR [09-05-98015_r_sibir_a] granted to Olga V. Churakova (Sidorova); SNSF Matthias Saurer [200021_121838/1]; Era.Net RusPlus project granted to
Markus Stoffel [SNF IZRPZ0_164735] and RFBR [№ 16-55-76012 Era_a] granted to Eugene

A. Vaganov; project granted to Vladimir S. Myglan RNF, Russian Scientific Fond [№ 15-14-

33

- 653 30011]; Alexander V. Kirdyanov was supported by the Ministry of Education and Science of
- the Russian Federation [#5.3508.2017/4.6] and RSF [#14-14-00295]; Scientific School
- 655 [3297.2014.4] granted to Eugene A. Vaganov; and US National Science Foundation (NSF)
- 656 grants [#9413327, #970966, #0308525] to Malcolm K. Hughes and US CRDF grant # RC1-
- 657 279, to Malcolm K. Hughes and Eugene A. Vaganov. We thank Tatjana Boettger for her sup-
- port and access to the stable isotope facilities within UFZ Haale/Saale scholarship 2006; Anne
- 659 Verstege, Daniel Nievergelt for their help with sample preparation for the MXD and Paolo
- 660 Cherubini for providing lab access at the Swiss Federal Institute for Forest, Snow and Land-
- 661 scape Research (WSL).
- 662 We thank two anonymous reviewers and handling Editor Juerg Luterbacher for their construc-
- tive comments on this manuscript.

664 Figure legends

665

666	Fig. 1. Location of the study sites (stars) and known volcanos from the tropics (black dots)
667	considered in this study (a). Annual tree-ring width index (light lines) and smoothed by 51-
668	year Hamming window (bold lines) from the northeastern Yakutia (YAK - blue, b) (Hughes
669	et al., 1999; Sidorova and Naurzbaev 2002; Sidorova 2003), eastern Taimyr (TAY - green, c)
670	(Naurzbaev et al., 2002), and Russian Altai (ALT - red, d) (Myglan et al., 2009). Photos show
671	the larch stnads at YAK, TAY (M.M. Naurzbaev) and ALT (V.S. Myglan) sites.
672	
673	Fig. 2. Normalized (z-score) individual tree-ring index chronologies (TRW, black), maxi-
674	mum latewood density (MXD, purple), cell wall thickness (CWT, green), δ^{13} C (red) and
675	δ^{18} O (blue) in tree-ring cellulose chronologies from northeastern Yakutia (YAK), eastern Tai-
676	myr (TAY) and Altai (ALT) for the specific periods 520-560, 1242-1286, 1625-1660, 1790-
677	1835, 1950-2000 before and after the eruptions CE 535, 540, 1257, 1640, 1815 and 1991 are
678	presented. Vertical lines show year of the eruptions.
679	
680	Fig. 3. Superposed epoch analysis (SEA) of δ^{18} O, δ^{13} C, CWT, TRW, and MXD chronologies
681	for the Yakutia (YAK), Taimyr (TAY), and Altai (ALT) sites, summarizing negative anoma-
682	lies 15 years before and 20 years after the volcanic eruptions in CE 535, 540, 1257, 1640,
683	1815, and 1991. Statistically negative anomalies are marked with a red star (* $p < 0.05$).
684	
685	Fig. 4. Significant correlation coefficients between tree-ring parameters: TRW, MXD, CWT,
686	δ^{13} C and δ^{18} O versus weather station data: temperature (T, red), precipitation (P, blue), vapor

pressure deficit (VPD, green), and sunshine duration (S, yellow) from September of the previous year to August of the current year for three study sites were calculated. Table 2 lists stations and periods used in the analysis.

690

691 Fig. 5. Responses of larch trees from Yakutia (YAK), Taimyr (TAY) and Altai (ALT) to volcanic eruptions (Table 1). Squares, rhombs, circles, and triangles indicate the years following 692 693 each eruption that can be considered as very extreme (negative values < 5th percentile of the PDFs, intensive color), extreme (negative values >5th, <10th percentile of the PDFs, light 694 color) and non-extreme (>10th percentile of the PDFs, white color). July temperature changes 695 are presented with squares. Summer vapor pressure deficit (VPD) variability is shown with 696 circles. July precipitations are presented with rhombs, and July sunshine duration is shown as 697 698 triangles.

699

700 **Table 1.** List of stratospheric volcanic eruptions used in the study

701

702 Table 2. Tree-ring sites in northeastern Yakutia (YAK), eastern Taimyr (TAY), and Altai

703 (ALT) and weather stations used in the study. Monthly air temperature (T, °C), precipitation

704 (P, mm), sunshine duration (S, h/month) and vapor pressure deficit (VPD, kPa) data were

705 downloaded from the meteorological database: <u>http://aisori.meteo.ru/ClimateR</u>.

706

Fig. S1. Probability density function (Pdf) computed for each of the tree-ring parameter for northeastern Yakutia (YAK), eastern Taimyr (TAY) and Russian Altai (ALT). Tree-ring parameters (TRWi - **black**, MXD – **purple**, CWT – **green**, δ^{18} O - **blue** and δ^{13} C - **red**) in bold lines represent the probability density function. Dotted lines represent the anomalies (z-score)

- observed for the first and second years following the CE 535, 540, 1257, 1640, 1815 and 1991
- 712 volcanic eruptions for each tree-ring parameter.

713

714 References

- Abaimov, A.P., Bondarev, A.I., Yzrzanova, O.V., Shitova, S.A.: Polar forests of Krasnoyarsk
 region. Nauka Press, Novosibirsk. 208 p, 1997.
- Battipaglia, G., Cherubini, P., Saurer, M., Siegwolf, R.T.W., Strumia, S., Cotrufo, M.F.: Volcanic explosive eruptions of the Vesuvio decrease tree-ring growth but not photosynthetic rates in the surrounding forests. Global Change Biology. 13, 1-16, 2007.
- Barinov, V.V., Myglan, V.S., Taynik, A.V., Ojdupaa, O.Ch., Agatova, A.R., Churakova (Sidorova) O.V.: Extreme climatic events in Altai-Sayan region as indicator of major
 volcanic eruptions. Geophysical processes and biosphere. 17, 45-61, 2018. doi:
- 723 10.21455/GPB2018.3-3.
- Boettger T., Haupt, M., Knöller, K., Weise, S., Waterhouse, G.S. ... Schleser, G.H.: Wood
 cellulose preparation methods and mass spectrometric analyses of δ¹³C, δ¹⁸O, and
 non ex-changeable δ²H values in cellulose, sugar, and starch: An inter-laboratory
 comparison. Anal. Chem. 79, 4603–4612, doi:10.1021/ac0700023, 2007.
- 728 Boike, J., Kattenstroth, B., Abramova, K., Bornemann, N., Cherverova, A., Fedorova, I.,
- 729 Fröb, K., Grigoriev, M., Grüber, M., Kutzbach, L., Langer, M., Minke, M., Muster,
- 730 S., Piel, K., Pfeiffer, E.-M., Stoff, G., Westermann, S., Wischnewski, K., Wille, C.,
- 731 Hubberten, H.-W.: Baseline characteristics of climate, permafrost and land cover
- from a new permafrost observatory in the Lena Rive Delta, Siberia (1998-2011). Biogeosciences. 10, 2105-2128, 2013.
- Briffa, K.R., Jones, P.D., Schweingruber, F.H., Osborn, T.J.: Influence of volcanic eruptions
 on Northern Hemisphere summer temperature over the past 600 years. Nature. 393,
 450–455, 1998.
- 737 Bryukhanova, M.V., Fonti, P., Kirdyanov, A.V., Siegwolf, R., Saurer, M., Pochebyt, N.P.,
 738 Churakova (Sidorova), O.V., Prokushkin, A.S.: The response of δ¹³C, δ¹⁸O and cell

- anatomy of *Larix gmelinii* tree rings to differing soil active layer depths. Dendrochronologia. 34, 51-59, 2015.
- Büntgen, U., Myglan, V.S., Ljungqvist, F.C., McCormick, M., Di Cosmo, N., Sigl M.,Kirdyanov, A.V.: Cooling and societal change during the Late Antique Little Ice Age
 from 536 to around 660 AD. Nature Geoscience. 9, 231-236, 2016.
- Castagneri, D., Fonti, P., von Arx, G., Carrer, M.: How does climate influence xylem morphogenesis over the growing season? Insights from long-term intra-ring anatomy in *Picea abies*. Annals of Botany. 19, 1011-1020, doi:10.1093/aob/mcw274, 2017.
- 747 Cernusak, L., Ubierna, N., Winter, K., Holtum, J.A.M., Marshall, J.D., Farquhar, G.D.: Envi748 ronmental and physiological determinants of carbon isotope discrimination in terres749 trial Plants. Transley Review New Phytologist. 200, 950-965, 2013.
- Cernusak, L., Barbour, M., Arndt, S., Cheesman, A., English, N., Field, T., Helliker, B., Holloway-Phillips, M., Holtum, J., Kahmen, A., Mcnerney F, Munksgaard N, Simonin
 K, Song X, Stuart-Williams H, West J and Farquhar G.: Stable isotopes in leaf water
- of terrestrial plants. Plant, Cell & Environment. 39 (5), 1087-1102, 2016.
- 754 Churakova (Sidorova), O.V., Bryukhanova, M., Saurer, M., Boettger, T., Naurzbaev, M.,
- Myglan, V.S., Vaganov, E.A., Hughes, M.K., Siegwolf, R.T.W.: A cluster of stratospheric volcanic eruptions in the AD 530s recorded in Siberian tree rings. Global and
 Planetary Change. 122, 140-150, 2014.
- 758 Churakova (Sidorova), O.V., Shashkin, A.V., Siegwolf, R., Spahni, R., Launois, T., Saurer
- 759 M., Bryukhanova, M.V., Benkova, A.V., Kupzova, A.V., Vaganov, E.A., Peylin, P.,

760	Masson-Delmotte, V., Roden, J.: Application of eco-physiological models to the cli-
761	matic interpretation of δ^{13} C and δ^{18} O measured in Siberian larch tree-rings. Dendro-
762	chronologa, doi:10.1016/j.dendro.2015.12.008, 2016.
763	Cook, E., Briffa, K., Shiyatov, S., Mazepa, V.: Tree-ring standardization and growth trend es-
764	timation. In: Methods of dendrochronology: applications in the environmental sci-
765	ences, Eds: Cook, E.R., Kairiukstis, L.A. 104-123, 1990.
766	Cook, E.R., Krusic, P.J.: A Tree-Ring Standardization Program Based on De-trending and
767	Autoregressive Time Series Modeling, with Interactive Graphics (ARSTAN). (Ed.
768	by E.R., Cook and P.J., Krusic), 2008.
769	Craig, H.: Isotopic variations in meteoric waters. Science. 133, 1702-1703, 1961.
770	Crowley, T.J., Unterman, M.B.: Technical details concerning development of a 1200 yr.
771	proxy index for global volcanism. Earth Syst. Sci. Data. 5, 187-197, 2013.
772	Cuny, H.E., Rathgeber, C.B.K., Frank, D., Fonti, P., Fournier, M.: Kinetics of tracheid devel-
773	opment explain conifer tree-ring structure. New Phytologist. 203, 1231–1241, 2014.
774	D'Arrigo, R.D., Jacoby, G.C., Frank, D., Pederson, N.D., Cook, E., Buckly, B.M., Nachin, B.,
775	Mijidorj, R., Dugarjav, C.: 1738-years of Mongolian temperature variability inferred
776	from a tree-ring width chronology of Siberian pine. Geophisycal Research Letters.
777	Vol. 28 (3), 543-546, 2001.
778	Dansgaard, W.: Stable isotopes in precipitation. Tellus. 16, 436–468, 1964.
779	Dawson, T.E., Mambelli, S., Plamboeck, A.H., Templer, P.H., Tu, K.P.: Stable isotopes in
780	plant ecology Ann. Review of Ecology and Systematics. 33, 507-559, 2004.
781	Dongmann, G., Förstel, H., Wagener, K.: ¹⁸ O-rich oxygen from land photosynthesis. Nature
782	New Biol. 240, 127–128, 1972.

- Eschbach, W., Nogler, P., Schär, E., Schweingruber, F.H.: Technical advances in the radiodensitometrical determination of wood density. Dendrochronologia. 13, 155–168,
 2015.
- Esper, J., Büntgen, U., Hartl-Meier, C., Oppenheimer, C., Schneider, L.: Northern Hemisphere temperature anomalies during 1450s period of ambiguous volcanic forcing.
 Bull. Volcanology. 79, 41, 2017.
- 789 Esper, J., St. George, S., Anchukaitis, K., D'Arrigo, R., Ljungqvist, F., Luterbacher, J.,
- Schneider, L., Stoffel, M., Wilson, R., Büntgen, U.: Large-scale, millennial-length
 temperature reconstructions from tree-rings. Dendrochronologia. 50, 81–90, 2018.
- **792** Farquhar, G. D.: Eds. Stable Isotopes and Plant Carbon-Water Relations. Academic Press,
- 793San Diego. 47–70, 1982.
- Farquhar, G.D., Ehleringer, J.R., Hubick, K.T.: Annu. Rev. Plant Physiol. Plant Mol. Biol. 40,
 503 p, 1989.
- Farquhar, G.D., Lloyd, J.: Carbon and oxygen isotope effects in the exchange of carbon dioxide between terrestrial plants and the atmosphere. In: Ehleringer, J.R., Hall, A.E.,
- Farquhar, G.D. (Eds) Stable Isotopes and Plant Carbon-Water Relations. Academic
 Press, San Diego, 47–70, 1993.
- 800 Fonti, P., Bryukhanova, M.V., Myglan, V.S., Kirdyanov, A.V., Naumova, O.V., Vaganov,
- 801 E.A.: Temperature-induced responses of xylem structure of *Larix sibirica* (Pinaceae)
 802 from Russian Altay. American Journal of Botany. 100 (7), 1-12, 2013.
- Francey, R.J., Allison, C.E., Etheridge D.M., Trudinger, C.M., Langenfelds, R.L., Michel, E.,
 Steele, L.P.: A 1000-year high precision record of δ¹³C in atmospheric CO₂. Tellus.
 Ser. B (51), 170-193, 1999.
- 806 Fritts, H.C.: Tree-rings and climate. London. New York; San Francisco: Acad. Press. 567 p,
 807 1976.

- Furst, G.G.: Methods of Anatomical and Histochemical Research of Plant Tissue. Nauka,
 Moscow. 156 p, 1979.
- Gao, C., Robock, A., Ammann, C.: Volcanic forcing of climate over the past 1500 years: An
 improved ice core-based index for climate models. J. Geophys. Res. Atmos.
- 812 113:D23111. doi:10.1029/2008jd010239, 2008.
- 813 Gennaretti, F., Huard, D., Naulier, M., Savard, M., Bégin, C., Arseneault, D., Guiot, J.:
- 814 Bayesian multiproxy temperature reconstruction with black spruce ring widths and 815 stable isotopes from the northern Quebec taiga. Clym. Dyn. doi: 10.1007/s00382-
- 816 017-3565-5, 2017.
- Gillett, N.P., Weaver, A.J., Zwiers, F.W. Wehner, M.F.: Detection of volcanic influence on
 global precipitation. Geophysical Research Letters. 31 (12),
- doi:10.1029/2004GL020044 R, 2004.
- B20 Groisman, P.Ya.: Possible regional climate consequences of the Pinatubo eruption. Geophys.
 B21 Res. Lett., 19, 1603–1606, 1992.
- 822 Gu, L., Baldocchi, D.D., Wofsy, S.C., Munger, J.W., Michalsky, J.J., Urbanski, S.P., Boden,
- T.A.: Response of a deciduous forest to the Mount Pinatubo eruption: Enhanced photosynthesis, Science. 299 (5615), 2035–2038, 2003.
- 825 Guillet, S., Corona, C., Stoffel, M., Khodri M., Lavigne F., Ortega, P.,....Oppenheimer, C.:
- 826 Climate response to the 1257 Samalas eruption revealed by proxy records. Nature
 827 geoscience, doi:10.1038/ngeo2875, 2017.
- 828 Hansen, J., Sato, M., Ruedy, R., Lacis, A., Asamoah, K., Borenstein S.,Wilson, H.: A
- 829 Pinatubo climate modeling investigation. In The Mount Pinatubo Eruption: Effects
- 830 on the Atmosphere and Climate, NATO ASI Series Vol. I 42. G. Fiocco, D. Fua, and
- G. Visconti, Eds. Springer-Verlag, 233-272, 1996.

- Harrington, C.R.: The Year without a summer? World climate in 1816. Ottawa: Canadian
 Museum of Nature, ISBN 0660130637, 1992.
- 834 Helama, S., Arppe, L., Uusitalo, J., Holopainen, J., Mäkelä, H.M., Mäkinen, H., Mielikäinen,
- 835 K., Nöjd, P., Sutinen, R., Taavitsainen, J.-P., Timonen, M., Oinonen, M.: Volcanic
- 836 dust veils from sixth century tree-ring isotopes linked to reduced irradiance, primary
- production and human health. Scientific reports 8, 1339, doi:10.1038/s41598-018-

838 19760-w, 2018.

- Hughes, M.K., Vaganov, E.A., Shiyatov, S.G., Touchan, R. & Funkhouser, G.: Twentiethcentury summer warmth in northern Yakutia in a 600-year context. The Holocene.
 9(5), 603-608, 1999.
- 842 Iles, C.E., Hegerl, G.C.: The global precipitation response to volcanic eruptions in the CMIP5
 843 models. Environ. Res. Lett. 9, doi:10.1088/1748-9326/9/10/104012, 2014.
- Joseph, R., Zeng, N.: Seasonally modulated tropical drought induced by volcanic aerosol. J.
 Climate, 24, 2045–2060, 2011.
- Kirdyanov, A.V., Treydte, K.S., Nikolaev, A., Helle, G., Schleser, G.H.: Climate signals in
 tree-ring width, wood density and δ¹³C from larches in Eastern Siberia (Russia)
 Chemical Geology, 252, 31-41, 2008. doi:10.1016/j.chemgeo.2008.01.023
- Körner, Ch.: Paradigm shift in plant growth control. Curr. Opinion Plant Biol. 25, 107-114,
 2015.
- Lavigne, F., Degeai, J.-P., Komorowski, J.-C., Guillet, S., Robert, V., Lahitte, P., Oppenheimer, C., Stoffel, M., Vidal, C.M., Suro, I.P., Wassmer, P., Hajdas, I., Hadmoko,
- $\mathbf{352} \qquad \mathbf{1161}, \mathbf{C}_{10}, \mathbf{5101101}, \mathbf{101}, \mathbf{V10a1}, \mathbf{C}_{1011}, \mathbf{5010}, \mathbf{1.1}, \mathbf{W} \mathbf{assinc1}, \mathbf{1}, \mathbf{11ajuas}, \mathbf{1}, \mathbf{11au110K0}, \mathbf{1}$
- 853D.S., Belizal, E.: Source of the great A.D. 1257 mystery eruption unveiled, Samalas
- 854 volcano, Rinjani Volcanic Complex, Indonesia. Proc Natl Acad Sci 110, 16742–
- 855 16747, doi:10.1073/pnas.1307520110, 2013.

Lenz, O., Schär, E., Schweingruber F.H.: Methodische Probleme bei der radiographisch-densitometrischen Bestimmung der Dichte und der Jahrrinbreiten von Holz.

858 Holzforschung, 30, 114-123, 1976.

- Loader, N.J., Robertson, I., Barker, A.C., Switsur, V.R., Waterhouse, J.S.: Improved technique for the batch processing of small whole wood samples to alpha-cellulose.
 Chemical Geology. 136, 313-317, 1997.
- Loader, N.J., Young, G.H.F., Grudd, H., McCarroll.: Stable carbon isotopes from Torneträsk,
 norther Sweden provide a millennial length reconstruction of summer sunshine and
- its relationship to Arctic circulation. Quaternary Science Reviews. 62, 97-113, 2013.
- 865 McCarroll, D., Loader, N.J.: Stable isotopes in tree rings. Quaternary Science Review. 23,

866 771-801, 2004.

Meronen, H., Henriksson, S.V., Räisänen, P., Laaksonen, A.: Climate effects of northern
hemisphere volcanic eruptions in an Earth System Model. Atmospheric Research,

869 114-115: 107-118, 2012.

- Munro, M.A.R., Brown, P.M., Hughes, M.K., Garcia, E.M.R.: Image analysis of tracheid
 dimensions for dendrochronological use. Radiocarbon, Eds. by M.D. Dean, J.

872 Swetnam T), pp. 843-851. Tucson, Arizona, 1996.

- Myglan, V.S., Oidupaa, O. Ch., Kirdyanov, A.V., Vaganov, E.A.: 1929-year tree-ring chronology for Altai-Sayan region (Western Tuva). Journal of archeology, ethnography
 and anthropology of Eurasia. 4 (36), 25-31, 2008.
- 876 Naurzbaev, M.M., Vaganov, E.A., Sidorova, O.V., Schweingruber, F.H.: Summer tempera-
- 877 tures in eastern Taimyr inferred from a 2427-year late-Holocene tree-ring chronology
 878 and earlier floating series. The Holocene. 12(6), 727-736, 2002.

- Panofsky, H.A., Brier, G.W.: Some applications of statistics to meteorology. University Park,
 PA. Mineral industries extension services, college of mineral industries, Pennsylvania State University, 224 p, 1958.
- Panyushkina, I.P., Hughes, M.K., Vaganov, E.A., Munro, M.A.R.: Summer temperature in
 northern Yakutia since AD 1642 reconstructed from radial dimensions of larch tracheids. Canadian Journal of Forest Research. 33, 1-10, 2003.
- Peng, Y., Shen, C., Wang, W.-C., Xu, Y.: Response of summer precipitation over Eastern
 China to large volcanic eruptions. Journal of Climate. 23, 818-824, 2009.
- R Core Team.: R: A Language and Environment for Statistical Computing. Vienna, Austria,
 2016.
- Robock, A.: Volcanic eruptions and climate. Reviews of Geophysics. 38(2), 191-219, 2000.
- Robock, A., Liu, Y.: The volcanic signal in Goddard Institute for Space Studies three-imensional model simulations. J. Climate. 7, 44-55, 1994.
- Roden, J.S., Siegwolf, R: Is the dual isotope conceptual model fully operational? Tree Physiology. 32, 1179-1182, 2012.
- 894 Saurer, M., Robertson, I., Siegwolf, R., Leuenberger, M.: Oxygen isotope analysis of
- cellulose: an interlaboratory comparison. Analytical cemistry, 70, 2074-2080, 1998.
- Saurer, M., Kirdyanov, A. V., Prokushkin, A. S., Rinne, K. T., Siegwolf, R.T.W.: The impact
 of an inverse climate-isotope relationship in soil water on the oxygen-isotope

898 composition of *Larix gmelinii* in Siberia. New Phytologist. 209(3), 955-964, 2016.

- 899 Scheidegger, Y., Saurer, M., Bahn, M., Siegwolf, R.: Linking stable oxygen and carbon iso-
- 900 topes with stomatal conductance and photosynthetic capacity: a conceptual model.
- 901 Oecologia. 125, 350–357. doi: 10.1007/s004420000466, 2000.
- 902 Schneider, L., Smerdon, J.E., Büntgen, U., Wilson, R.J.S., Myglan, V.S., Kirdyanov, A.V.,
- 903 Esper, J.: Revising mid-latitude summer temperatures back to A.D. 600 based on a

- 904 wood density network. Geophys. Res. Lett. 42, GL063956,
- 905 doi:10.1002/2015gl063956, 2015.
- Schweingruber, F.H.: Tree rings and environment dendroecology. Paul Haupt Publ Bern,
 Stuttgart, Vienna 1996. pp. 609, 1996.
- Sidorova, O.V., Naurzbaev, M.M.: Response of *Larix cajanderi* to climatic changes at the
 upper timberline and in the Indigirka River valley. Lesovedenie (in Russian). 2, 7375, 2002.
- 911 Sidorova, O.V.: Long-term climatic changes and the larch radial growth on the northern
- 912 Middle Siberia and the Northeastern Yakutia in the Late Holocene. Abs. PHD Diss,
- 913 V.N. Sukachev Institute of Forest, Krasnoyarsk, 2003.
- 914 Sidorova, O.V., Naurzbaev, M.M., Vaganov, E.A.: Response of tree-ring chronologies growing
 915 on the Northern Eurasia to powerful volcanic eruptions. Problems of ecological moni916 toring and ecosystem modeling, XX, 60-72, 2005.
- 917 Sidorova, O.V., Saurer, M., Myglan, V.S., Eichler, A., Schwikowski, M., Kirdyanov, A.V.,
- 918 Bryukhanova, M.V., Gerasimova, O.V., Kalugin, I., Daryin, A., Siegwolf, R.: A
- 919 multi-proxy approach for revealing recent climatic changes in the Russian Altai. Cli920 mate Dynamics, 38 (1-2), 175–188, 2011.
- Sidorova, O.V., Siegwolf, R., Myglan, V.S., Loader, N.J., Helle, G., Saurer, M.: The application of tree-rings and stable isotopes for reconstructions of climate conditions in the
 Altai-Sayan Mountain region. Climatic Changes, doi: 10.1007/s10584-013-0805-5,
- **924** 2012.
- 925 Sidorova, O.V., Siegwolf, R., Saurer, M., Naurzbaev, M., Shashkin, A.V., Vaganov, E.A.:
- 926 Spatial patterns of climatic changes in the Eurasian north reflected in Siberian larch
- 927 tree-ring parameters and stable isotopes. Global Change Biology, doi:
- 928 10.1111/j.1365-2486.2009.02008.x, 16, 1003-1018, 2010.

- Sidorova, O.V., Siegwolf, R.T.W., Saurer, M., Naurzbaev, M.M., Vaganov, E.A.: Isotopic
 composition (δ¹³C, δ¹⁸O) in Siberian tree-ring chronology. Geophysical research
 Biogeosciences. 113, 1-13, 2008.
- Sigl, M., Winstrup, M., McConnell, J.R.: Timing and climate forcing of volcanic eruptions
 for the past 2500 years. Nature. 523, 543-549. doi:10.1038/nature14565, 2015.
- 934 Sprenger, M., Tetzlaff, D., Buttle, J. M., Laudon, H., Leistert, H., Mitchell, C., Snelgrove, J.,
- Weiler, M., Soulsby, C.: Measuring and modelling stable isotopes of mobile and
 bulk soil water, Vadose Zone Journal, https://doi.org/10.2136/vzj2017.08.0149, 20,
 2017.
- Sternberg, L.S.O.: Oxygen stable isotope ratios of tree-ring cellulose: The next phase of understanding. New Phytologist. 181 (3), 553-562, 2009.
- 940 Stirzaker, D.: Elementary Probability density functions. Cambridge. Sec. Ed. 538 p, 2003.
- 941 Stoffel, M., Khodri, M., Corona, C., Guillet, S., Poulain, V., Bekki, S., Guiot, J., Luckman,
- 942 B.H., Oppenheimer, C., Lebas, N., Beniston, M., Masson-Delmotte, V.: Estimates of
 943 volcanic-induced cooling in the Northern Hemisphere over the past 1500 years. Na-
- 944 ture Geoscience. 8, 784–788, 2015.
- Stothers, R.B.: Climatic and Demographic Consequences of the Massive Volcanic Eruption of
 1258. Climatic Change. 45, 361-374, 2000.
- 947 Stothers, R.B.: Mystery cloud of AD 536. Nature. 307, 344-345, doi:10.1038/307344a0,
 948 1984.
- 949 Sugimoto, A., Yanagisawa, N., Fujita, N., Maximov, T.C.: Importance of permafrost as a
 950 source of water for plants in east Siberian taiga. Ecological Research. 17 (4), 493951 503, 2002.

- Toohey, M., Sigl, M.: Volcanic stratospheric sulphur injections and aerosol optical depth
 from 500 BCE to 1900 CE. Earth System Science Data. doi:10.5194/essd-9-8092017, 2017.
- Vaganov, E.A., Hughes, M.K., Kirdyanov, A.K., Schweingruber, F.H., Silkin, P.P.: Influence
 of snowfall and melt timing on tree growth in subarctic Eurasia. Nature. 400, 149151, 1999.
- 958 Vaganov, E.A., Hughes, M.K., Shashkin, A.V.: Growth dynamics of conifer tree rings.
 959 Springer Verlag, Berlin., pp. 353, 2006.
- 960 Wegmann, M., Brönnimann, S., Bhend, J., Franke, J., Folini, D., Wild, M., Luterbacher, J.:
- 961 Volcanic influence on European summer precipitation through monsoons: Possible
 962 cause for "years without summer". AMS, doi.org/10.1175/JCLI-D-13-00524.1, 2014.
- Wigley, T.M.L., Briffa, K.R., Jones, P.D.: On the Average Value of Correlated Time Series,
 with Applications in Dendroclimatology and Hydrometeorology. Journal of Climate
 and Applied Meteorology. 23 (2), 201-213, doi:10.1175/15200450(1984)023.0201,
 1984.
- 967 Wiles, G.C., D'Arrigo, R.D., Barclay, D., Wilson, R.S., Jarvis, S.K., Vargo, L., Frank, D.:
- 968 Surface air temperature variability reconstructed with tree rings for the Gulf of
 969 Alaska over the past 1200 years. The Holocene. 6, 10.1177/0959683613516815,
 970 2014.
- Wilson, R.J.S., Anchukaitis, K., Briffa, K. et al.: Last millennium Northern Hemisphere summer temperatures from tree rings. Part I: the long-term context. Quaternary Science
 Review. 134, 1–18, 2016.
- 274 Zielinski, G.A., Mayewski, P.A., Meeker, L.D., Whitlow, S., Twickler, M.S., Morrison, M.,
 275 Meese, D.A., Gow A.J., Alley, R.B.: Record of volcanism since 7000 BC from the

- 976 GISP2 Greenland ice core implications for the volcano-climate system. Science. 264
- 977 (5161), 948-952, 1994.
- 978 Zielinski, G.A.: Use of paleo-records in determining variability within the volcanism-climate
- 979 system. Quaternary Science Reviews. 19, 417-438, 2000.