

The 4.2-ka event, ENSO, and coral-reef development

Lauren T. Toth¹ and Richard B. Aronson²

¹U.S. Geological Survey, St. Petersburg Coastal and Marine Science Center, St. Petersburg, FL, 33701, USA

²Department of Ocean Engineering and Marine Sciences, Florida Institute of Technology, Melbourne FL, 32901, USA

5 Correspondence to: Lauren T. Toth (ltoth@usgs.gov)

Variability of sea-surface temperature related to shifts in the mode of the El Niño–Southern Oscillation (ENSO) has been implicated as a possible forcing mechanism for the changes in global-scale, tropical and subtropical precipitation known as the 4.2-ka event. We review records of coral-reef development and paleoceanography from the tropical eastern Pacific (TEP) to evaluate the potential impact of the 4.2-ka event on coral reefs. Our goal is to identify the regional climatic and oceanographic drivers of a 2500-year shutdown of vertical reef accretion in the TEP after 4.2 ka. The 2500-year hiatus represents ~40% of the Holocene history of reefs in the TEP and appears to have been tied to increased variability of ENSO. When ENSO variability abated approximately 1.7–1.6 ka, coral populations recovered and vertical accretion of reef framework resumed apace. There is some evidence that the 4.2-ka event suppressed coral growth and reef accretion elsewhere in the Pacific Ocean as well. Although the ultimate causality behind the global 4.2-ka event remains elusive, correlations between shifts in ENSO variability and the impacts of the 4.2-ka event suggest that ENSO could have played a role in climatic changes at that time, at least in the tropical and subtropical Pacific. We outline a framework for testing hypotheses of where and under what conditions ENSO may be expected to have impacted coral-reef environments around 4.2 ka. Although most studies of the 4.2-ka event have focused on terrestrial environments, we suggest that understanding the event in marine systems may prove to be the key to deciphering its ultimate cause.

1 Introduction

The abrupt climatic shift at ~4200 years before present (BP; expressed as years before 1950), known as the 4.2-ka event, is now recognized by many scientists to be the most significant climatic perturbation of the middle to late Holocene (Walker et al., 2012; ka stands for thousands of years ago). On 25 June 2018, the International Union of Geological Sciences officially subdivided the Holocene epoch, with the 4.2-ka event marking the start of the late Holocene, or the Meghalayan age. Although the signature of the 4.2-ka event is not as ubiquitous as the north-Atlantic cooling episode known as the 8.2-ka event, which is the demarcation between the early and middle Holocene, it is now clear that the 4.2-ka event had global climatic, ecological, and cultural impacts (Mayewski et al., 2004; Weiss, 2016). In most locations, the 4.2-ka event was manifested as extreme and abrupt changes in terrestrial hydroclimate (Straubwasser et al., 2003; Marchant and Hooghiemstra, 2004; Walker et al. 2012; Xiao et al., 2018), which drove environmental changes (Marchant and Hooghiemstra, 2004; Booth et al., 2005) and cultural collapses (Weiss et al., 1993; Straubwasser et al., 2003; Weiss, 2016; Wu et al., 2016) on a global scale. The ultimate cause(s) of the event and its impact and connection with other regional- to global-scale climatic changes remain unknown (Walker et al., 2012).

Whereas shifts in the climate of high-latitude regions in general, and the northern Atlantic in particular, have been implicated in the majority of climate events during the late Quaternary (Dansgaard et al., 1993; Mayewski et al., 2004; Booth et al., 2005; Walker et al., 2012), there is little evidence for a high-latitude driver of the 4.2-ka event (Walker et al., 2012). The event is only weakly recorded in Greenland ice-core records (Johnsen et al., 2001; Marchant and Hooghiemstra, 2004; Mayewski et al., 2004; Rasmussen et al., 2006): major cooling in the northern Atlantic, and the associated onset of northern-hemisphere neoglaciation, preceded the 4.2-ka event by at least 1000 years (Svenden and Mangerud, 1997; Mayewski et al., 2004; Walker et al., 2012; Marcott et al., 2013). The lack of a clear, high-latitude forcing mechanism for the 4.2-ka event has led some researchers to suggest that there may have been a tropical driver of the climatic changes around this time (Marchant and Hooghiemstra, 2004).

The most significant source of climatic variability in the tropics is the El Niño–Southern Oscillation (ENSO). Although ENSO originates in the tropical Pacific, it causes global-scale thermal anomalies and changes in hydroclimate (McPhaden et al., 2006). The climatological manifestations of individual El Niño and La Niña events can vary, but many of the broadscale impacts of ENSO are similar to the global-scale changes inferred to have taken place during the 4.2-ka event. For example, some of the most striking changes associated with the 4.2-ka event were apparent shifts in the strength of the Asian Monsoon (Staubwasser et al., 2003; Wang et al., 2003). ENSO variability has been closely linked with the dynamics of the Asian Monsoon throughout the Holocene (Liu et al. 2000). Whereas El Niño events are typically associated with a weaker Monsoon and resultant drought conditions, the Monsoon is stronger during La Niña events (Liu et al., 2000; Wang et al., 2003). The similarity in the climatic changes associated with ENSO and the 4.2-ka event, coupled with the evidence for increasing ENSO variability around 4.2 ka (Conroy et al., 2008; Koutavas and Joanides, 2012; Carré et al., 2014), has led a number of researchers to suggest that ENSO may have played a salient role in the 4.2-ka event (e.g., Marchant and Hooghiemstra, 2004; Booth et al., 2005; Walker et al., 2012; Li et al., 2018). It is not yet clear, however, whether changing ENSO variability around 4.2 ka is likely to have been an ultimate driver of broad-scale climatic changes around this time or whether ENSO was a proximal response to other climatic forcing.

Coral reefs are excellent models for elucidating the causes and consequences of climatic excursions such as the 4.2-ka event. Reef-building (hermatypic) corals, which secrete aragonitic skeletons that build the geologic frameworks of the reefs, are highly sensitive to extremes in temperature and light. As a result, coral reefs are among the most vulnerable ecosystems to modern, anthropogenic climate change (e.g., Hoegh-Guldberg, 1999; Hughes et al., 2018). Photosynthetic dinoflagellates, called zooxanthellae, which live endosymbiotically within the cnidarian hosts, provide the animal component of the coral holobiont (the entire animal–plant–microbial symbiosis) with nearly all of its energetic needs (Muscatine and Porter, 1977). Persistently high temperature and light levels, which are now occurring with increasing frequency as a result of anthropogenically-driven climate change, disrupt the mutualistic relationship between the coral animal and the zooxanthellae, leading to coral bleaching and associated morbidity and mortality (Fig. 1; Brown, 1987; Glynn, 1991; Hoegh-Guldberg, 1999; Donner et al., 2005; Anthony, 2016; and many others). Cold-water conditions are also inimical to coral growth and reef development (Ginsburg and Shinn, 1994; Precht and Aronson, 2004; Lirman et al., 2011; Kemp et al., 2016; Toth et al., 2018). Corals are

vulnerable to a variety of other stresses and disturbances, including sedimentation, subaerial exposure, nutrient loading, hurricane damage, low-pH conditions, and low salinity (Kleypas et al., 1999; Aronson and Precht, 2001; Buddemeier et al., 2004; Goldberg, 2013), which can increase as a result of (or in tandem with) climatic shifts in one
75 direction or another. Reef frameworks and the coral skeletons that constitute them preserve well in the fossil and subfossil record, providing a historical record of the response of coral reefs to climate change (Jackson, 1992). Moreover, many of the perturbations listed above leave geochemical and taphonomic traces that can be used as proxies to reconstruct changes in environmental conditions (e.g., Cobb et al., 2013; Toth et al., 2012, 2015a, 2015b; Flannery et al. 2018).

80 Whereas the majority of the records of the 4.2-ka event have come from terrestrial environments, the contemporary impacts of ENSO are often felt most keenly in marine ecosystems. Understanding whether and how marine ecosystems responded to climatic changes around 4.2 ka is, therefore, critical to discovering the ultimate drivers of the 4.2-ka event. Here, we explore the hypothesis that ENSO played a role in the 4.2 ka event by reviewing paleoecological and paleoceanographic records from marine environments in the tropical Pacific. We
85 focus on the long-term collapse of coral-reef development in the tropical eastern Pacific (TEP), which appears to be related to changes in ENSO. We conclude that the relationship between ENSO and the 4.2 ka event warrants further study and outline a conceptual framework for future studies to investigate the linkages between these climatic phenomena using records from coral-reef environments in the tropical Pacific.

2 Climate and coral-reef development in the eastern tropical Pacific

90 2.1 Ecology of Coral Reefs in the Tropical Eastern Pacific

Holocene reefs of the TEP are, for the most part, small and poorly developed, as is generally the case for reefs on the eastern margins of ocean basins (Darwin, 1862; Cortés, 1993). Regional and local diversities of hard corals are low in the TEP. Several living reefs, however, overlie well-developed Holocene frameworks that preserve millennial-scale records of regional and larger-scale climatic variability, and the responses of coral assemblages to that
95 variability.

The shallowest habitats of contemporary reefs off the Pacific coast of Panamá, ranging from 1–5 m water depth, are characterized by fields of branching corals of the genus *Pocillopora* (Fig. 1A; Glynn and Macintyre, 1977; Cortés, 1997; Glynn and Ault, 2000). The stands of *Pocillopora* are dominated by what has been termed *P. damicornis*, but they include several other species in the genus (many or most of which are hybrids; see below).
100 Massive corals, primarily *Pavona* spp., *Porites* spp., and *Gardinerocoris planulata*, become more abundant with increasing depth below the zone of essentially monogeneric occupation by *Pocillopora* (Glynn, 1976; Glynn and Maté, 1997).

Richmond (1987, 1990) suggested that *P. damicornis* in the TEP had adopted a life-history strategy of exclusive, or nearly-exclusive, asexual reproduction. In this scenario, the populations were seeded by sexually
105 produced planula larvae from the central Pacific, which were transported by the North-Equatorial Countercurrent and by eastward equatorial flow during El Niño conditions (Glynn and Ault, 2000). Oceanographic-modeling

exercises, however, cast doubt on Richmond's proposed teleconnection between the central and eastern Pacific, at least so far as *Pocillopora* is concerned (Wood et al., 2016; Romero-Torres et al., 2018). Furthermore, genetic analysis by Combosch et al. (2008) demonstrated that the eastern-Pacific populations of *Pocillopora* reproduce sexually. The genetics showed a difference in reproductive strategy in the genus *Pocillopora* between the eastern Pacific and the rest of its range in the Indo-Pacific: *Pocillopora* reproduce sexually by broadcast-spawning in the eastern Pacific, whereas they brood internally-fertilized or parthenogenetic eggs in central and western Pacific, and in the Indian Ocean.

Broadcast-spawning is correlated with interspecific hybridization among the species of *Pocillopora* in the TEP. Hybridization in the TEP is in turn associated with a radically different functional ecology of (hybridized) *P. damicornis* in the eastern Pacific, including a far more robust skeletal morphology, ecological dominance, and the construction of reef-frameworks composed predominantly of *Pocillopora* branches. Within the eastern Pacific, limited gene flow among populations of (hybridized) *P. damicornis* (henceforth simply '*Pocillopora*') raises the possibility of adaptation to regional-scale oceanographic gradients of upwelling (Combosch and Vollmer, 2011).

2.2 Response to ENSO Events

Contemporary populations of corals on reefs of the TEP (see Fig. 6.1 in Toth et al., 2017 for a map of locations where coral reefs occur) highlight the responses of coral populations to sequential perturbations by ENSO. The 1982–83 El Niño was one of the strongest to affect the eastern Pacific in recent history (Glynn, 1988a; Glynn et al., 2001). (The impacts of this event were less severe elsewhere in the tropical Pacific, however.) Persistently elevated water temperatures for 14 months resulted in severe, regional-scale bleaching of corals (Fig. 1B) and other cnidarians bearing endosymbiotic algae (Wellington and Glynn, 2007). Coral mortality ranged from 50% in Costa Rica to 97% in the Galápagos Islands and correlated with the degree of thermal stress the reefs experienced in each location (Glynn, 1990). Mortality levels were somewhat lower off the Pacific coast of Panamá. In the Gulf of Panamá, where seasonal upwelling is strong, coral mortality was 85%. In the adjacent Gulf of Chiriquí, where seasonal upwelling is weak or absent, mortality was similar, averaging 76% (Glynn, 1990, 1991; Glynn and Colgan, 1992; Wellington and Glynn, 2007). The similarity in the level of mortality throughout Pacific Panamá reflects the fact that El Niño suppressed seasonal upwelling in the Gulf of Panama in 1982–83 so the level of warming was similar in both gulfs (Glynn et al., 2001).

Ecological observations prior to 1982 suggested that dense stands of *Pocillopora* had persisted throughout the TEP for decades at least. *Pocillopora* had in fact been the dominant benthic component in shallow reef habitats prior to the 1982–83 event, covering up to 90% of the available substratum at 3–5 m depth (Fig. 1A). *Pocillopora* proved especially vulnerable to bleaching-induced mortality and vast fields were killed, although mortality was variable on multiple spatial scales (Glynn, 1990; Macintyre and Glynn, 1990). A collateral effect of *Pocillopora* mortality was the decline of symbiotic crustaceans that defend the corals against attack by the predatory crown-of-thorns starfish, *Acanthaster planci*. At Uva Island, in the Gulf of Chiriquí, *Acanthaster* were able to crawl across freshly killed thickets of *Pocillopora* to attack and kill massive corals that had not bleached as severely and did not harbor the

crustaceans (Glynn, 1985, 1990). *Acanthaster* did not then and does not now occur in the Gulf of Panamá (Glynn, 2004).

145 Algal turfs growing on the dead-coral surfaces enhanced populations of regular echinoids by increasing their food supplies. Higher densities of grazing sea urchins greatly increased bioerosion and suppressed coral recruitment, resulting in a switch from net vertical reef accretion to net framework erosion (Glynn, 1988a, 1988b, 1990; Glynn, and Colgan 1992; Eakin, 1996; Reaka-Kudla et al., 1996). Centennial-scale increases in the frequency of severe ENSO events such as the one in 1982–83 could have been responsible for the poor development of reefs observed in many areas of the eastern Pacific (Glynn and Colgan, 1992; Glynn, 2000).

150 A second extreme ENSO event killed an estimated 16% of corals worldwide in 1997–98 (Wilkinson, 2000). The 1997–98 and 1982–83 events were of approximately equal magnitude and duration in the eastern Pacific. Both events may have been enhanced by global warming and they were the two most intense events in the preceding 50 yr (Hansen, et al. 1999; Karl et al., 2000; Enfield, 2001; Hughes et al., 2018).

155 Coral mortality in the eastern Pacific was lower in 1997–98 (Guzmán and Cortés, 2001; Vargas-Ángel et al., 2001). There was essentially no mortality in the Gulf of Panamá because, whereas seasonal upwelling was suppressed during the 1982–83 El Niño event, upwelling was unaltered in 1997–98 and cooled the water column (Riegl and Piller, 2003; Glynn et al., 2001). Muted levels of coral mortality elsewhere, including only 13% mortality in the Gulf of Chiriquí, were correlated with the presence of thermally resistant zooxanthellae in the corals (Glynn et al., 2001; Baker et al., 2004; see also D’Croz and Maté, 2004). Collateral effects following the 1982–83 event, particularly enhanced predation by *Acanthaster* (in the Gulf of Chiriquí) and increased bioerosion by sea urchins, were also far less severe after the 1997–98 ENSO, due to precipitous declines of both taxa (Eakin, 2001; Glynn et al., 2001; see also Fong and Glynn, 2001). The effects of the 2015–16 El Niño event were also minimal in Pacific Panamá. Upwelling again buffered the reefs in the Gulf of Panamá from elevated temperatures, and most reefs in the Gulf of Chiriquí experienced coral bleaching, but only minor bleaching-related mortality.

160 Coral populations have recovered to varying degrees from the El Niño events in 1982–83 and 1997–98, apparently having been seeded by populations that persisted within refugia in the eastern Pacific (Wood et al., 2016). Coral cover declined in the Gulf of Chiriquí at 3–7 m depth after 1983. By 2002, coral cover was 12% at Uva Island (down significantly from 35%) and 8% at Secas Island (down non-significantly from 11%; Wellington and Glynn, 2007). Coral cover at Saboga Island in the Gulf of Panamá recovered at 2–3 m depth from 0% in 1984 to the pre-ENSO level of 50% by 1992. At 3–5 m depth, some areas of the Saboga reef recovered to 50% within a few years, but large areas of coral rubble have persisted to this day (Fig. 1C), and this has also been the case on many other reefs. The areas of coral rubble consist primarily of taphonomically degraded *Pocillopora* branches, which are covered in algal turfs, encrusted by coralline algae, and colonized at low frequency by the early-successional coral *Psammocora stellata*.

175 In summary, elevated water temperatures and high irradiance (resulting from low cloud cover) associated with the strong El Niño events in 1982–83 and 1997–98 in Pacific Panamá caused widespread coral bleaching, which was associated with mass coral mortality in the earlier event (Glynn et al., 2001). La Niña is also problematic for corals in Pacific Panamá. Lowered sea level in the TEP during La Niña events causes more frequent coral mortality

180 associated with subaerial exposure (Eakin and Glynn, 1996; Toth et al., 2017). In addition, La Niña is associated with elevated rainfall and enhanced upwelling in Pacific Panamá. Elevated rainfall increases turbidity, and enhanced upwelling reduces water temperatures, decreases pH, and increases nutrient levels. All of these changes act to suppress coral growth (Glynn, 1976).

185 Food webs on eastern Pacific reefs are complex despite their low diversity (Glynn, 2004). Corallivores other than *Acanthaster*, including pufferfish (Tetraodontidae) and gastropods (Pediculariidae and Muricidae), have effects that vary on multiple scales. On the other hand, because it is situated close to the equator, Panamá has not experienced a hurricane since at least as early as 1871 (Neumann et al., 1999), removing hurricanes as a factor—but not storms and wave-action in the winter dry season.

2.3 The Question of Prior Occurrence

190 The TEP has been affected by more than 30 strong ENSO events during the past 500 yr and perhaps hundreds of events as strong as the 1982–83 and 1997–98 ENSOs during the Holocene (Colgan, 1990; Moy et al., 2002; Rein et al., 2005). Cores drilled from massive coral colonies (coral heads) throughout the Indo-Pacific display a trend toward more intense ENSO events after 1976 (Urban et al., 2000); however, although ENSO variability during the last century may be significantly higher than many times in the past, there is precedent in the fossil record for the intensity of recent events (Cobb et al., 2013). Whereas gross patterns of reef development have been examined in the eastern Pacific (Glynn and Macintyre, 1977; Macintyre et al., 1992; Cortés et al., 1994), biotic turnover, its climatic drivers, and its geological consequences remain understudied.

200 At least some effects of the 1982–83 ENSO event were unprecedented on a scale of centuries. First, colonies of the massive corals *Porites lobata* and *Pavona clavus* in the Galápagos were 350–425 years old at the time of their death during the event (Glynn, 1990). Second, Dunbar et al. (1994) examined a colony of *Pavona clavus* that was part of the reef community at Urvina Bay in the Galápagos. In 1954, Urvina Bay was tectonically uplifted more than 7 m, preserving in excellent detail the effects of a strong El Niño in 1941 (Colgan, 1990). The *Pavona* colony, which had survived the 1941 event, showed more than 350 yr of continuous growth before its death in the 1954 uplift. Clearly, some massive corals had lived through previous, strong ENSO events. On the other hand, partial colony mortality and regeneration of the massive species *G. planulata* at Uva Island in the Gulf of Chiriquí marked both the 1982–83 and 1997–98 ENSOs, so even the latter event was not without ecological consequences (Wellington and Glynn, 2007).

210 More germane are Colgan's (1990) observations on uplifted stands of *Pocillopora* at Urvina Bay. Much of the *Pocillopora* was in poor taphonomic condition (but excellently preserved in that poor condition; Colgan and Malmquist, 1987), undoubtedly the result of 13 yr of bioerosion by urchin grazing after the ENSO of 1941 and prior to the uplift. The uplifted reef contained clear evidence of bioerosive undercutting and collapse of the *Pocillopora* framework caused by sea urchins, *Eucidaris galapagensis*, quite similar to what was observed in living communities in the Galápagos after the 1982–83 event. Furthermore, *Eucidaris* tests were scattered in large numbers at the base of the eroded *Pocillopora* framework.

215 The question of prior occurrence is important because, as discussed above, repeated El Niño events could
explain why reefs are poorly developed in the eastern Pacific. Push-cores that we extracted from uncemented reef-
frameworks in both the Gulf of Panamá and the Gulf of Chiriquí appear to corroborate the pre-1982 ecological data,
suggesting continuous dominance of *Pocillopora* for at least the last 1500 years (Toth et al., 2012, 2017). The
ecological and paleobiological lines of evidence are not strictly comparable, however, because the temporal
precision of cores may not be great enough to detect decadal-scale events. Can we identify mass coral-kills and
220 excursions from the *Pocillopora*-dominated state in the subfossil record of eastern Pacific reefs? Colgan's (1990)
results and post-1983 observations throughout the region imply that under the right conditions it should be possible
to detect bioeroded horizons of *Pocillopora* and shifts in dominance between coral species.

Pocillopora-dominated reefs are not well-developed in the eastern Pacific due to the narrow shelves on which
they grow; exposure in at least some locations to cold and acidic, upwelling waters; and episodic bleaching during
225 ENSO events (Cortés et al., 1994; Glynn and Maté, 1997; Kleypas et al., 2006). Glynn and Colgan (1992; Glynn,
2000) argued that eastern Pacific reefs accrete poorly because the corals are periodically killed by severe El Niño
events and then bioeroded by echinoids. Our study of the geologic record, in contrast, points to a protracted phase
shift, during which the vertical accretion of Holocene reef-frameworks off the Pacific coast of Panamá was
suppressed for far longer than the return time of strong ENSO events.

230 Our push-cores from three fringing Panamanian reefs, which spanned a gradient of upwelling and water depths
(-0.8 to -4.1 m relative to mean sea level), and which we dated with radiocarbon and uranium-series techniques,
showed that vertical accretion essentially ceased from ~4100 to 1600 BP (Fig. 2; Toth et al., 2012). There is no
evidence that the reefs shifted to lateral accretion at this time, as the reefs at all coring sites were growing over a
paleodepth-range of ~1.6 m at the time they shut down (L. T. Toth, unpubl. data). The hiatus in growth lasted
235 approximately 2500 years, meaning the reefs were in a phase of negligible growth for as much as 40% of their
history (Toth et al., 2012). Detailed core logs are provided in Toth et al. (2012, 2013).

Active coral growth was represented in the cores as long intervals of well-preserved *Pocillopora* branch-
fragments. The hiatus was manifested as a thin layer of taphonomically degraded *Pocillopora* branch-fragments,
coralline algae, and *Psammocora stellata*, all of which characterize ENSO-generated rubble-fields on contemporary
240 reefs in Pacific Panamá. During the rest of their history, however, the reefs grew upward at rates comparable with
the vertical-accretion rates of Caribbean reefs (Toth et al., 2012). The 'poor' Holocene development of these reefs is,
therefore, a consequence of the hiatus. The tempo and causality of reef development in Panamá represent a vastly
scaled-up version of the earlier model of Glynn and Colgan (1992; 2000), which had suggested that poor reef
development was a consequence of decadal-scale disturbance by El Niño. Although ENSO suppresses populations
245 of *Pocillopora* on subdecadal to multidecadal scales in Panamá, slowed growth on those scales has not been the
cause of low bulk-accretion rates through the Holocene.

3 ENSO and Coral-Reef Collapse at 4.2 ka

There is now broad consensus across both climate models and paleoclimate reconstructions that the middle
Holocene was a period of relatively low ENSO variability (Sandweiss et al., 2001; Tudhope et al., 2001; Rein et al.,

250 2005; Koutavas et al., 2006; Zheng et al., 2008; Cobb et al., 2013; Carré et al., 2014; Liu et al., 2014; Emile-Geay et al., 2016; Leonard et al., 2016b; Thompson et al., 2017). In particular, the interval 5–3 ka stands out as a period when paleoclimate records from throughout the tropical Pacific show a significant reduction in ENSO activity (Emile-Geay et al., 2016). Although the exact timing of the middle-Holocene ENSO minimum varies among reconstructions, some studies have suggested that there was exceptionally low ENSO variability between 4.3 and 4.2 ka (Cobb et al., 2013; McGregor et al., 2013; Leonard et al., 2016b; Emile-Geay et al., 2016). Paradoxically, ~4.2 ka is also a point of inflection in most ENSO records, after which time ENSO variability increased, eventually leading to the establishment of the modern ENSO regime (Conroy et al., 2008; Koutavas and Joanides, 2012; Carré et al., 2014).

260 ENSO appears to be a primary control on modern coral growth and reef development in the TEP (Glynn and Colgan, 1992; Glynn et al., 2001; Toth et al., 2012, 2017). The dominant role of ENSO in modulating the ecology of contemporary reefs in the TEP led us to hypothesize that ENSO variability may also have shaped the dynamics of Panamanian reefs over millennial timescales. We tested the hypothesis that changes in ENSO activity around ~4.2 ka could have triggered reef shutdown in Pacific Panamá by evaluating geochemical proxy records from corals in our cores in the Gulf of Panamá (Fig. 2; Toth et al., 2015a, 2015b). For obvious reasons, the availability of coral samples from the time of the hiatus was limited; however, we were able to analyze the Sr/Ca and $\delta^{18}\text{O}$ of seven coral skeletons that our age model suggested grew at the beginning of the hiatus (~3.9–3.6 ka; Toth et al., 2015a). We also have a measurement of oceanic radiocarbon, a proxy for ocean circulation, from one coral that grew during this period (Toth et al., 2015b). Therefore, although we do not have any records at precisely 4.2 ka, our reconstructions give insights into the mean climatic and oceanographic states bounding 4.2 ka, which we evaluate in the context of records of ENSO variability from throughout the tropical Pacific presented in Fig. 3. Note that the schematic (Fig. 3) provides an overview of changes in ENSO, suggested by our work and the paleoclimate literature, that could have contributed to the shutdown of eastern Pacific reef development. It is not meant as a comprehensive review of the literature.

275 Our record of oceanic radiocarbon variability suggests that upwelling was enhanced beginning at 4.3 ka and continued to intensify over at least the next 500 years (Fig. 2A; Toth et al., 2015b). The hiatus in reef development was more protracted in the Gulf of Panamá, implying that upwelling may have made reefs more sensitive to further environmental perturbations (Toth et al., 2012). Paleoclimate reconstructions, which we based on Sr/Ca and $\delta^{18}\text{O}$ in the coral skeletons, suggest that these oceanographic changes were followed by a transition to a cooler, wetter climate beginning by ~3.9 ka (Fig. 2B, 2C; Toth et al., 2015a). A lake record from the Galápagos recorded a contemporaneous period of anomalous drying, which in that location points to enhanced La Niña activity (Fig. 3; Conroy et al., 2008). Together, the inferred conditions indicate that an enhanced La Niña-like climate regime was established just after 4.2 ka, which could have provided the initial trigger that shut down reef development in Pacific Panamá for the next 2500 years (Toth et al., 2012, 2015a).

285 Whereas the period from 5–3 ka may have been a time of low ENSO variability *on average* (Emile-Geay et al., 2016), the time just after 4.2 ka was likely characterized by high climatic volatility in the tropical Pacific (Fig. 3). At least one record, from Vanuatu in the western Pacific, indicates that a multi-decadal period of enhanced

ENSO variability followed just 500 years after the putative low in ENSO activity at ~4.2 ka (Corrège et al., 2000); however, we note that there is less evidence for elevated ENSO variability in a contemporaneous record from nearby New Caledonia (Emile-Geay et al., 2016). Our data from Pacific Panamá suggest that the period after 4.2 ka would have been followed by a transition to La Niña-like conditions by 3.9 ka (Toth et al., 2015a, 2015b), and another recent study suggests that La Niña events were also more frequent from 3.5–3 ka (Thompson et al., 2017). A record of El Niño-related flooding from Peru implies that the strongest El Niño events of the Holocene occurred 4–2 ka (Rein et al., 2005), and a series of records from the same region suggest a possible peak in the frequency of El Niño events around 3 ka (Sandweiss et al., 2001; Moy et al., 2002). We note, however, that recent reanalysis has called these conclusions into question (Emile-Geay and Tingley 2016). The beginning of the late Holocene was also characterized by a period of high variability in the position of the Intertropical Convergence Zone (ITCZ), supporting the idea that regional climatic variability was high at the time (Haug et al., 2001). Taken together, these studies support the conclusion that ENSO variability was enhanced after 4.2 ka (Conroy et al., 2008; Koutavas and Joanides, 2012; Carré et al., 2014).

Stronger and/or more frequent El Niño and La Niña events would have acted to suppress reef development after 4.2 ka (Toth et al., 2012). ENSO variability continued to increase after the end of the hiatus in reef development (Conroy et al., 2008; Cobb et al., 2013; Thompson et al., 2017); however, major changes in the mode of ENSO coincided with its termination (Fig. 3). First, the variability in the mean position of the ITCZ had declined by ~2.5 ka (Haug et al., 2001). Second, although ENSO variability and the absolute number of El Niño events may have increased beginning 2 ka (Sandweiss et al., 2001; McGregor and Gagan, 2004; Conroy et al., 2008; Thompson et al., 2017), El Niño events were likely stronger (Rein et al., 2005) and La Niña event may have been more frequent (Thompson et al., 2017) during the hiatus. Our records suggest that upwelling was more moderate after the hiatus (Fig. 2A), which would indicate that there were fewer or less extreme La Niña events. Furthermore, because the climate remained relatively cool after the hiatus (Fig. 2B), it is likely that the influence of El Niño events had also decreased. We hypothesize that the waning influence of La Niña and reduction in El Niño strength permitted accretion to resume between ~1.8–1.5 ka (Toth et al., 2012, 2015a).

We conclude that ENSO was likely the primary driver of the collapse of coral reefs in Pacific Panamá and, perhaps, other locations around the tropical Pacific at ~4.2 ka. The fact that the shutdown in reef accretion occurred in both the Gulf of Panamá and the Gulf of Chiriquí excludes upwelling and outbreaks of *Acanthaster* as drivers: upwelling is weak or absent in the Gulf of Chiriquí, and *Acanthaster* is absent from the Gulf of Panamá. Other possible factors, including changes in relative sea level, tectonics, and bioerosion, cannot explain the observed patterns either (Toth et al., 2012, 2015a). The potential role of changing ENSO variability in ecological changes elsewhere at this time is less clear, but there is some evidence to support the idea: increased ENSO-like variability in the climate of eastern China is temporally correlated with an inferred increase in coral bleaching around 4.2 ka (Li et al., 2018; Xu et al. 2018). Furthermore, even if enhanced ENSO variability triggered reef-shutdown in the TEP at 4.2 ka, that does not mean ENSO-related changes necessarily accounted for the next 2500 years of suppressed reef development; Fig. 3 implies a concatenation of conditions that might or might not have been dependent or semi-

independent. Some evidence does in fact point to climatic shifts around the time of the 4.2-ka event that persisted for millennia (Selvaraj et al. 2008).

325 There is still considerable disagreement about timing and magnitude of the putative shifts in ENSO
variability during the middle to late Holocene (Emile-Geay et al., 2016). One problem in linking ENSO as a driver
or response to the 4.2-ka event is the asymmetry in the ENSO signal between the eastern and central Pacific that can
be observed from both recent El Niño events (Fig. 4; Takahashi et al., 2011) and in the Holocene record (Carré et
al., 2014; Karamperidou et al., 2015). Indeed, there appear to be two distinct “flavors” of ENSO based on the
330 location of the strongest thermal anomalies—eastern Pacific and central Pacific events—and the dominance of these
ENSO modes may have varied over millennial timescales (Karamperidou et al., 2015). Another problem is the large
temporal uncertainties associated with many of the existing paleoclimate records of ENSO. Currently, it is not
possible to resolve clearly the centennial-scale shifts in ENSO variability around 4.2 ka. Reducing the temporal
uncertainties in paleo-ENSO records at 4.2 ka and the dominant flavor of ENSO at this time, by targeting ENSO
335 proxies from throughout the tropical Pacific, will be a critical step to determining whether shifts in ENSO variability
can be linked to the 4.2-ka event on a geographic scale and the subsequent 2.5 millennia of impacts on reefs
throughout the tropical Pacific.

4 Regional- to Global-Scale Impacts of ENSO and the 4.2 ka Event

340 ENSO is the most prominent interannual driver of global-scale climate variability. Although it originates in the
tropical Pacific, it impacts regional climates around the world (McPhaden et al., 2006). The global impacts of ENSO
are primarily a result of the influence of ENSO-based sea-surface temperature anomalies on other drivers of climate
variability such as the ITCZ, the Asian Monsoon, and the North Atlantic Oscillation (NAO). For example, the warm
(cool) tropical Pacific sea-surface temperatures associated with El Niño (La Niña) generally drive a southerly
(northerly) shift in the mean position of the ITCZ, which results in changes in precipitation from the tropical Pacific
345 to the neotropics, to Africa (Haug et al., 2001; Chiang et al., 2002; Sachs et al., 2009). Similarly, the Asian Monsoon
is generally suppressed during El Niño events and enhanced during La Niña events (Liu et al., 2000; Wang et al.,
2003). Although the relationship between ENSO and the NAO, which has a strong influence on European climates,
is less straightforward, the NAO is typically negative (positive) during El Niño (La Niña) events (Rodríguez-
Foncesca et al., 2016).

350 The global impacts of El Niño and La Niña events can vary significantly from event to event; however,
there are striking spatial correlations between the global signature of ENSO (summarized in Davey et al., 2013) and
that of the 4.2-ka event. In the majority of the records to date, the 4.2-ka event has been characterized by changes in
terrestrial hydroclimate (Weiss, 2016). Whereas the event was characterized by widespread aridification in southern
Europe, the Middle East, northern Africa, midcontinental North America, and parts of eastern and southwest Asia
355 (Weiss et al., 1993; Straubwasser et al., 2003; Marchant and Hooghiemstra, 2004; Booth et al., 2005; Xiao et al.,
2018), other locations, such as western South America, experienced marked increases in precipitation (Marchant and
Hooghiemstra, 2004). With the exception of Europe, which is weakly teleconnected with ENSO, the mean shifts in
precipitation that occurred during the 4.2-ka event are generally consistent with those observed during a typical El

Niño event (Davey et al., 2013). The changes in hydroclimate during typical El Niño events even replicate the zonal
360 gradient in precipitation observed across northern South America at ~4.2 ka (Marchant and Hooghiemstra, 2004;
McPhaden et al., 2006; Davey et al., 2013).

Like many records of ENSO variability from the tropical Pacific, our records from Pacific Panamá do not
display a sudden change in El Niño activity at 4.2 ka (Toth et al., 2015a; Emile-Geay et al., 2016); however, most
reconstructions of Holocene ENSO suggest that there was a transition to increasing ENSO variability around this
365 time (Conroy et al., 2008; Koutavas and Joanides, 2012; Cobb et al., 2013; Carré et al., 2014). Whereas the 4.2-ka
event was manifested as an abrupt, short-lived climatic event in many locations, the larger-scale climatic and
ecosystem responses to the event may have been more gradual and more protracted. Increased ENSO variability
could in this sense be a response to the abrupt climate changes associated with the 4.2 ka event. In Pacific Panamá,
the onset of reef collapse just after 4.2 ka was associated with a transition to a cooler, La Niña-like climate (Toth et
370 al., 2015; see also Cabarcos et al., 2014), which triggered a regime shift that lasted for millennia. Similarly, there is
evidence that the onset of aridification in the mid-continent United States was followed by glacial advances in
western Canada, from 4.2–3.8 ka, associated with a cooler, wetter climate (Menounos et al., 2008; Mayewksi et al.,
2004) that is characteristic of La Niña conditions in that region (Davey et al., 2013). Similarly, whereas a number of
records indicate a weaker East Asian Monsoon around 4.2 ka, indicative of an El Niño-like climate (Straubwasser et
375 al., 2003), there is also evidence that some regions of southern Asia became wetter, or more La Niña-like, after 4.2
ka (reviewed in Wu and Liu, 2004). Thus, whereas many of the abrupt changes in terrestrial hydroclimate during the
4.2-ka event are similar to those observed during El Niño events, the ecosystem changes following the 4.2-ka event
could also reflect responses to secondary changes in regional climatic systems.

We are not arguing that ENSO was necessarily an ultimate cause of the 4.2-ka event; however, the shift in
380 ENSO variability after 4.2 ka and the similarity between the global impacts of ENSO and the global footprint of the
4.2-ka event suggest a potential connection. The climatic forcing that produced the changes in ENSO variability
during the middle Holocene are still being debated, but it is generally thought that changes in ENSO during the
Holocene have been driven by gradual changes in the climate system associated with changing insolation and/or
feedbacks with annual climate cycles (McPhaden et al., 2006; McGregor et al., 2013; Emile-Geay et al., 2016;).
385 For example, several researchers (Koutavas et al., 2006; McGregor et al., 2013) have suggested that the weaker ENSO
variability observed in the central Pacific at ~4.3 ka was related to the more northerly ITCZ at this time, which
would have also enhanced eastern-Pacific upwelling, produced cooler sea temperatures in the TEP, and driven a
stronger zonal gradient in sea-surface temperature that suppressed ENSO (Clement et al., 2000). In the absence of a
clear high-latitude driver for the 4.2-ka event, these changes in the tropical ocean must be considered as potential
390 contributors to the 4.2-ka event (Marchant and Hooghiemstra, 2004).

5 Prospectus

Because the global teleconnections of El Niño and La Niña can vary significantly from event to event (Fig. 4;
McPhaden et al., 2006), ecosystems of the tropical and subtropical Pacific are likely to yield the best evidence for
evaluating the potential role of ENSO in the 4.2-ka event. Corals and coral-reef communities that were living under

395 chronically stressed conditions should have been particularly vulnerable to increased ENSO variability after 4.2 ka. If the hiatus was triggered by La Niña-like conditions (Fig. 4), the 4.2 ka event should be most strongly manifested in reef records from areas of upwelling and in seasonally cooler, subtropical environments (compared with less-seasonal, tropical environments). Nearshore habitats, which are more influenced by the thermal swings and high sedimentation rates associated with terrigenous input, should be more vulnerable compared with reefs exposed to clearer oceanic waters. Reefs off the Pacific coast of Panamá and Costa Rica, which are exposed to a strong terrigenous influence and (in some areas) strong upwelling, display the hiatus (Table 1; Cortés et al., 1994; Toth et al., 2012). Latitudinally marginal lagoonal habitats elsewhere in the Pacific should also manifest the hiatus (see Cruz et al., 2018). Preliminary evidence corroborates that hypothesis in the western Pacific, in turbid, lagoonal habitats south of the Great Barrier Reef, in southern China, and in Japan (Table 1; Lybolt et al., 2011; Hamanaka et al., 2012; Yamano et al., 2012; Xu et al., 2018); however, falls or stillstands in relative sea-level in the western Pacific after the middle Holocene likely also contributed to stalled vertical reef growth during the late Holocene in many locations (e.g., Dechnik et al., 2018), making it more difficult to discern the impacts of climatic variability and also potentially introducing variability into the start-times of hiatuses in reef accretion. Other possible causes of variations in the start-times listed in Table 1 include the artifactual absence of coral material of particular ages in particular samples and the uncertainty associated with the dating techniques. Whether La Niña-driven reef shutdown occurred broadly on subtropical reefs of the central Pacific remains a topic of active research, but there is some evidence that increased wave energy tied to ENSO suppressed vertical reef accretion in Hawaii around the time of the 4.2-ka event (Table 1; Grossman and Fletcher, 2004; Rooney et al., 2004).

By contrast, the impacts of increased El Niño intensity that followed the putative increase in La Niña activity (Figs. 3, 4), might have been attenuated in lagoonal and nearshore habitats, where turbidity associated with terrigenous runoff from increased precipitation (in many locations, though not in Panamá) would have decreased light penetration and thereby offered a measure of protection from bleaching (Perry et al., 2012; Cacciapaglia and van Woesik, 2015). Above some threshold amplitude, however, the ENSO swings would have overwhelmed the capacity of even turbid-water reefs to resist bleaching (e.g., Aronson et al., 2002). Furthermore, El Niño conditions are accompanied by increased precipitation in most places, with potentially negative impacts on corals.

Across the equatorial Pacific, contemporary corals and coral reefs at longitudes that have experienced higher temperatures over the previous century have proven more resistant to bleaching events. The eastern and central Pacific, including tropical as well as subtropical areas, have thus been more vulnerable for historical reasons than the western Pacific (Thompson and van Woesik, 2009). If the near-term geographic patterns of ENSO history are indicative of patterns earlier in the Holocene, then on the basis of El Niño-type impacts we would expect the 4.2-ka event to have manifested strongly in the eastern and central Pacific, and less so in the western Pacific. This hypothesis could be tested by coring Holocene reef-frameworks across the Pacific near the equator. The physiographic marginality of individual reefs, resulting from additional sources of environmental variability (such as upwelling) and their particular ecologies (including their species diversity), would have decreased their resilience and thereby increased the likelihood that El Niño disturbances would have left a signature in their fossil record.

Cobb et al. (2013) documented low ENSO variability at 4.2 ka at Kiritimati in the Line Islands of the central Pacific. That result potentially falsifies our hypothesis of an event at 4.2 ka being strongly manifested in reef frameworks of the central Pacific, as well as the idea that contemporary geographic patterns of ENSO variability reflect patterns that prevailed 4.2 ka. There is some evidence that the eastern Pacific mode of ENSO was dominant after ~4.5 ka (Carré et al. 2014), which may explain the muted response to the 4.2 ka event in the central Pacific. Parsing the spatially variable impacts of ENSO and other drivers, such as relative sea level, upwelling, climatic effects other than ENSO, and local physiography and hydrography, to explain patterns of reef development will be a significant challenge going forward (for the Great Barrier Reef: Dechnik et al., 2015; Leonard et al., 2016a, 2016b; Ryan et al., 2018; Webster et al., 2018; see Toth et al., 2016 for the eastern Pacific).

The preponderance of research to date on the causes and consequences of the 4.2-ka event has focused on terrestrial environments and cultural impacts. In this paper we have shown that there were significant perturbations to coral-reef environments in Pacific Panamá and throughout the tropical Pacific around 4.2 ka. Without a clear high-latitude driver for the 4.2-ka event, it stands to reason that the climatic changes around this time could have originated in the tropical oceans and may have been related to contemporaneous changes in ENSO variability.

Coral-reef archives could provide the key to discerning the role of ENSO in the 4.2-ka event.

A strong role for ENSO in the 4.2-ka event would have important implications for living coral populations and future reef accretion in the TEP. Contemporary climate change will likely increase ENSO variability in the eastern Pacific (Cai et al. 2018). A robust connection between ENSO variability and reef development could, therefore, portend another shutdown of eastern Pacific reefs.

Acknowledgements. This paper grew out of an international workshop, ‘The 4.2 ka BP Event,’ which was organized by G. Zanchetta, H. Weiss, and M. Bini and held in the Dipartimento di Scienze della Terra at the University of Pisa in January 2018. We thank the organizers for the opportunity to participate in the workshop. R. Bradley, M. Bush, E. Edinger, N. Graham, W. Precht, J. Richey, R. van Woesik, H. Weiss, and two anonymous reviewers provided valuable advice and discussion. Our research was supported by grant OCE-1535007 from the U.S. National Science Foundation. Toth’s research is also funded by the Coastal and Marine Hazards and Resources Program of the U.S. Geological Survey. This is contribution number 203 from the Institute for Global Ecology at the Florida Institute of Technology. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

References

- Anthony, K. R. N.: Coral reefs under climate change and ocean acidification: challenges and opportunities for management and policy, *Ann. Rev. Environ. Resour.*, 41, 59–81, 2016.
- Aronson, R. B., and Precht, W. F.: Evolutionary paleoecology of Caribbean coral reefs, in: *Evolutionary paleoecology: the ecological context of macroevolutionary change*, edited by: Allmon, W. D., and Bottjer, D. J., Columbia University Press, New York, 171–233, 2001.
- Aronson, R. B., Precht, W. F., Toscano, M. A., and Koltes, K. H.: The 1998 bleaching event and its aftermath on a

- coral reef in Belize, *Mar. Biol.*, 141, 435–447, 2002.
- Baker, A. C., Starger, C. J., McClanahan, T. R., and Glynn, P. W.: Corals' adaptive response to climate change, *Nature*, 430, 741, 2004.
- 470 Booth, R. K., Jackson, S. T., Forman, S. L., Kutzbach, J. E., Bettis Iii, E. A., Kreigs, J., and Wright, D. K.: A severe centennial-scale drought in midcontinental North America 4200 years ago and apparent global linkages, *The Holocene*, 15, 321–328, 2005.
- Brown, B. E.: Worldwide death of corals: natural cyclical events or man-made pollution, *Mar. Poll. Bull.*, 18, 9–13, 1987.
- 475 Buddemeier, R. W., Kleypas, J. A. and Aronson, R. B.: Coral reefs and global climate change: potential contributions of climate change to stresses on coral reef ecosystems, Pew Center on Global Climate Change, Arlington, Virginia, USA, 2004.
- Cabarcos, E., Flores, J. -A., and Sierro, F. J.: High-resolution productivity record and reconstruction of ENSO dynamics during the Holocene in the Eastern Equatorial Pacific using coccolithophores, *The Holocene*, 24, 176–187, 2014.
- 480 Cacciapaglia, C., and van Woesik, R.: Reef-coral refugia in a rapidly changing ocean, *Global Change Biol.*, 21, 2272–2282, 2015.
- Carré, M., Sachs, J. P., Purca, S. Schauer, A., Braconnot, P., Falcón, R. A., Julien, M., and Lavallée, D.: Holocene history of ENSO variance and asymmetry in the eastern tropical Pacific, *Science*, 345, 1045–1048, 2014.
- 485 Chaing, J. C. H., Kushnir, Y., and Giannini, A.: Deconstructing Atlantic Intertropical Convergence Zone variability: influence of the local cross-equatorial sea surface temperature gradient and remote forcing from the eastern equatorial Pacific, *J. Geophys. Res.*, 107, JD000307, 2002.
- Cai, W., Wang, J., Dewitte, B., Wu, L., Santoso, A., Takahashi, K., Yang, Y., Carréric, A., and McPhaden, M. J.: Increased variability of eastern Pacific El Niño under greenhouse warming, *Nature*, 564, 201–206, 2018.
- 490 Clement, A., Seager, R., Cane, M. A., and Zebiak, S. E.: An ocean dynamical thermostat, *J. Clim.*, 9, 2910–2196, 2000.
- Cobb, K. M., Charles, C. D., Cheng, H., and Edwards, R. L.: El Niño/Southern Oscillation and tropical Pacific climate during the last millennium, *Nature*, 424, 271–276, 2003.
- Cobb, K. M., Westphal, N., Sayani, H. R., Watson, J. T., Di Lorenzo, E., Cheng, H., Edwards, R. L., and Charles, C. D.: Highly variable El Niño–Southern Oscillation throughout the Holocene, *Science*, 339, 67–70, 2013.
- 495 Colgan, M. W.: El Niño and the history of eastern Pacific reef building, in: *Global ecological consequences of the 1982–83 El Niño–Southern Oscillation*, edited by: Glynn, P. W., Elsevier, Amsterdam, 183–232, 1990.

- Colgan, M. W., and Malmquist, D. L.: The Urvina Bay uplift: a dry trek through a Galápagos coral reef, *Oceanus*, 30, 61–66, 1987.
- 500 Combosch, D. J., Guzmán, H. M, Schuhmacher, H., and Vollmer, S. V.: Interspecific hybridization and restricted trans-Pacific flow in the tropical eastern Pacific *Pocillopora*, *Mol. Ecol.*, 17, 1304–1312, 2008.
- Combosch, D. J., and Vollmer, S. V.: Population genetics of an ecosystem-defining reef coral *Pocillopora damicornis* in the tropical eastern Pacific, *PLoS ONE*, 6, e21200, doi: 10.1371/journal.pone.0021200, 2012.
- Conroy, J. L., Overpeck, J. T., Cole, J. E., Shanahan, T. M., and Steinitz-Kannan, M.: Holocene changes in eastern tropical Pacific climate inferred from a Galápagos lake sediment record, *Quat. Sci. Rev.*, 27, 1166–1180, 2008.
- 505 Cortés, J.: Comparison between Caribbean and eastern Pacific coral reefs, *Revista de Biología Tropical*, 41, 19–21, 1993.
- Cortés, J.: Biology and geology of eastern Pacific coral reefs, *Coral Reefs*, 16, S39–S46, 1997.
- Cortés, J., Macintyre, I. G., and Glynn, P. W.: Holocene growth history of an eastern Pacific fringing reef, Punta Islotes, Costa Rica, *Coral Reefs*, 13, 65–73, 1994.
- 510 Corrège, T., Delcroix, T., Réey, J., Beck, W., Cabioch, G., and Le Cornec, F.: Evidence for stronger El Niño–Southern Oscillation (ENSO) events in a mid-Holocene massive coral, *Paleoceanography*, 15, 465–470, 2000.
- Cruz, I. G. S., Waters, L. G., Kikuchi, R. K. P., Leão, Z. M. A. N., and Turra, A.: Marginal coral reefs show high susceptibility to phase shift, *Mar. Pollut. Bull.*, 135, 551–561, 2018.
- Dansgaard, W., Johnsen, S.J., Clausen, H.B., Dahl-Jensen, D., Gundestrup, N.S., Hammer, C.U., Hvidberg, C.S., Steffensen, J.P., Sveinbjörnsdóttir, A.E., Jouzel, J. and Bond, G.: Evidence for general instability of past climate from a 250-kyr ice-core record, *Nature*, 364, 218, 1993.
- 515 Darwin, C.: The structure and distribution of coral reefs (reprinted). University of California Press, Berkeley, California, 214 pp, 1862.
- Davey, M. K., Brookshaw, A., and Ineson, S.: The probability of the impact of ENSO on precipitation and near-surface temperature, *Clim. Risk Manage.*, 1, 5–24, 2014.
- 520 D’Croz, L., and Maté, J. L.: Experimental responses to elevated water temperature in genotypes of the reef coral *Pocillopora damicornis* from upwelling and non-upwelling environments in Panama, *Coral Reefs*, 23, 473–483, 2004.
- Dechnik, B., Webster, J. M., Davies, P. J., Braga, J.-C., and Reimer, P. J.: Holocene “turn-on” and evolution of the southern Great Barrier Reef: revisiting reef cores from the Capricorn Bunker Group, *Mar. Geol.*, 363, 174–190, 2015.
- 525 Donner, S. D., Skirving, W. J., Little, C. M., Oppenheimer, M., and Hoegh-Guldberg, O.: Global assessment of coral bleaching and required rates of adaptation under climate change, *Global Change Biol.*, 11, 2251–2265, 2005.

- 530 Dunbar, R. B., Wellington, G. M., Colgan, M. W., and Glynn, P. W.: Eastern Pacific sea surface temperature since 1600 A.D.: the ^{18}O record of climate variability in Galápagos corals, *Paleoceanography*, 9, 291–315, 1994.
- Eakin, C. M.: Where have all the carbonates gone? A model comparison of calcium carbonate budgets before and after the 1982–1983 El Niño at Uva Island in the eastern Pacific, *Coral Reefs*, 15, 109–119, 1996.
- Eakin, C. M.: 2001. A tale of two ENSO events: carbonate budgets and the influence of two warming disturbances and intervening variability, Uva Island, Panama, *Bull. Mar. Sci.* 69, 171–186, 2001.
- 535 Eakin, C. M., and Glynn, P. W.: Low tidal exposures and reef mortalities in the eastern Pacific, *Coral Reefs*, 15, 120, 1996.
- Emile-Geay, J., and Tingley, M.: Inferring climate variability from nonlinear proxies: application to palaeo-ENSO studies, *Clim. Past*, 12, 31–50. 2016.
- 540 Emile-Geay, J., Cobb, K. M., Carré, M., Braconnot, P., Leloup, J., Zhou, Y., Harrison, S. P., Corrège, T., McGregor, H. V., Collins, M., Driscoll, R., Elliot, M., Schneider, B., and Tudhope, A.: Links between tropical Pacific seasonal, interannual and orbital variability during the Holocene, *Nat. Geosci.*, 9, 168–175, 2016.
- Enfield, D. B.: Evolution and historical perspective of the 1997–1998 El Niño–Southern Oscillation event, *Bull. Mar. Sci.*, 69, 7–25, 2001.
- 545 Flannery, J. A., Richey, J. N., Toth, L. T., Kuffner, I. B., and Poore, R. Z.: 2018. Quantifying uncertainty in coral Sr/Ca-based SST estimates from *Orbicella faveolata*: A basis for multi-colony SST reconstructions, *Paleoceanography and Paleoclimatology*. doi: 10.1029/2018PA003389.
- Fong, P., and Glynn, P. W.: Population abundance and size-structure of an eastern tropical Pacific reef coral after the 1997–98 ENSO: a simulation model predicts field measures, *Bull. Mar. Sci.*, 69, 187–202, 2001.
- 550 Ginsburg, R. N., and Shinn, E. A.: Preferential distribution of reefs in the Florida reef tract: the past is the key to the present, in: *Proceedings of the Colloquium on Global Aspects of Coral Reefs: Health, Hazards, and History*, compiled by: Ginsburg, R. N., Rosenstiel School of Marine and Atmospheric Science, University of Miami, Coral Gables, Florida, 21–26, 1994.
- Glynn, P. W.: Some physical and biological determinants of coral community structure in the eastern Pacific, *Ecol. Monogr.*, 46, 431–456, 1976.
- 555 Glynn, P. W.: El Niño-associated disturbance to coral reefs and post-disturbance mortality by *Acanthaster planci*, *Mar. Ecol. Prog. Ser.*, 26, 295–300, 1985.
- Glynn, P. W.: El Niño–Southern Oscillation 1982–1983: nearshore population, community, and ecosystem responses, *Ann. Rev. Ecol. Syst.*, 19, 309–345, 1988a.
- 560 Glynn, P. W.: El Niño warming, coral mortality and reef framework destruction by echinoid bioerosion in the eastern Pacific, *Galaxea* 7, 129–160, 1988b.

- Glynn, P. W.: Coral mortality and disturbances to coral reefs in the tropical eastern Pacific, in: Global ecological consequences of the 1982–83 El Niño–Southern Oscillation, edited by: Glynn, P. W., Elsevier, Amsterdam, 55–125, 1990.
- 565 Glynn, P. W.: Coral reef bleaching in the 1980s and possible connections with global warming, *Trend. Ecol. Evol.*, 6, 175–179, 1991.
- Glynn, P. W.: El Niño–Southern Oscillation mass mortalities of reef corals: a model of high temperature marine extinctions?, in: Carbonate platform systems: components and interactions, edited by: Insalaco, E., Skelton, P. W., and Palmer, T. J., Geological Society of London Special Publication 178, London, 117–133, 2000.
- 570 Glynn, P. W.: High complexity food webs in low-diversity eastern Pacific reef-coral communities, *Ecosystems*, 7, 358–367, 2004.
- Glynn, P. W., and Ault, J. S.: A biogeographic analysis and review of the far eastern Pacific coral reef region, *Coral Reefs*, 19, 1–23, 2000.
- Glynn, P. W., and Colgan, M. W.: Sporadic disturbances in fluctuating coral reef environments: El Niño and coral reef development in the eastern Pacific, *Am. Zool.*, 32, 707–718, 1992.
- 575 Glynn, P. W., Maté, J. L.: Field guide to the Pacific coral reefs of Panamá, *Proc. 8th Int. Coral Reef Symp.*, Panama, 1, 145–166, 1997.
- Glynn, P. W., Maté, J. L., Baker, A. C., and Calderón, M.: Coral bleaching and mortality in Panama and Ecuador during the 1997–1998 El Niño–Southern Oscillation event: Spatial/temporal patterns and comparisons with the 1982–1983 event, *Bull. Mar. Sci.*, 69, 79–109, 2001.
- 580 Glynn, P. W., and Macintyre, I. G.: Growth rate and age of coral reefs on the Pacific coast of Panama, *Proc. 3rd Int. Coral Reef Symp.*, Miami, 2, 251–259, 1977.
- Goldberg, W. M.: The biology of reefs and reef organisms. University of Chicago Press, Chicago, Illinois, USA, 2013.
- 585 Grossman, E. E., and Fletcher III, C. H.: Holocene reef development where wave energy reduces accommodation space, Kailua Bay, windward Oahu, Hawaii, U.S.A, *J. Sed. Res.*, 74, 49–63, 2004.
- Guzmán, H., and Cortés, J.: Changes in reef community structure after fifteen years of natural disturbances in the eastern Pacific (Costa Rica), *Bull. Mar. Sci.*, 69, 133–149, 2001.
- Hamanaka, N., Kan, H., Yokoyama, Y., Okamoto, T., Nakashima, Y., and Kawana, T.: Disturbances with hiatuses in high-latitude coral reef growth during the Holocene: correlation with millennial-scale global climate change, *Global Planet. Change*, 80, 21–35, 2012.
- 590 Hansen, J., Ruedy, R., Glascoe, J., and Sato, M.: GISS analysis of surface temperature change, *J. Geophys. Res.*, 107, 30997–31022, 1999.

- 595 Haug, G. H., Hughen, K. A., Sigman, D. M., Peterson, L. C., and Röhl, U.: Southward migration of the Intertropical Convergence Zone through the Holocene, *Science*, 293, 1304–1308, 2001.
- Hoegh-Guldberg, O.: 1999. Climate change, coral bleaching and the future of the world's coral reefs, *Mar. Freshw. Res.*, 50, 839–866, 1999.
- Hughes, T. P., Anderson, K. D., Connolly, S. R., Heron, S. F., Kerry, J. T., Lough, J. M., Baird, A. H., Baum, J. K., Berumen, M. L., Bridge, T. C. and Claar, D. C.: Spatial and temporal patterns of mass bleaching of corals in the
600 Anthropocene, *Science*, 359, 80–83, 2018.
- Jackson, J. B. C.: Pleistocene perspectives of coral reef community structure, *Am. Zool.*, 32, 719–731, 1992.
- Johnsen, S.J., Dahl-Jensen, D., Gundestrup, N., Steffensen, J.P., Clausen, H.B., Miller, H., Masson-Delmotte, V., Sveinbjörnsdottir, A.E. and White, J.: Oxygen isotope and palaeotemperature records from six Greenland ice-core stations: Camp Century, Dye-3, GRIP, GISP2, Renland and NorthGRIP, *J. Quat. Sci.*, 16, 299–307, 2001.
- 605 Karamperidou, C., Di Nezio, P. N., Timmermann, A., Jin, F.-F., and Cobb K. M.: The response of ENSO flavors to mid-Holocene climate: implications for proxy interpretation, *Paleoceanography*, 30, 527–547, 2015.
- Karl, T. R., Knight, R. W., and Baker, B.: The record breaking global temperatures of 1997 and 1998: evidence for an increase in the rate of global warming? *Geophys. Res. Lett.*, 27, 719–722, 2000.
- Kemp, D. W., Colella, M. A., Bartlett, L. A., Ruzicka, R. R., Porter, J. W., and Fitt, W. K.: Life after cold death: reef
610 coral and coral reef responses to the 2010 cold water anomaly in the Florida Keys, *Ecosphere*, 7, e01373, 2016.
- Kleypas, J. A., Buddemeier, R. W., Archer, D., Gattuso, J.-P., Langdon, C., and Opdyke, B. N.: Geochemical consequences of increased atmospheric carbon dioxide on coral reefs, *Science*, 284, 118–120, 1999.
- Kleypas, J. A., Feely, R. A., Fabry, V. J., Langdon, C., Sabine, C. L., and Robbins, L. L.: Impacts of ocean acidification on coral reefs and other marine calcifiers: a guide for future research, Report of a workshop held
615 18–20 April 2005, St. Petersburg, Florida. NSF, NOAA and USGS, www.isse.ucar.edu/florida/, 2006
- Koutavas, A., deMenocal, P. B., Olive, G. C., and Lynch-Stieglitz, J.: Mid-Holocene El Niño-Southern Oscillation (ENSO) attenuation revealed by individual foraminifera in eastern tropical Pacific sediments, *Geology*, 34, 993–996, 2006.
- Koutavas, A., and Joanides, S.: El Niño–Southern Oscillation extrema in the Holocene and Last Glacial Maximum,
620 *Paleoceanography*, 27, PA4208, 2012.
- Leonard, N. D., Zhao, J.-x., Welsh, K. J., Feng, Y.-x., Smithers, S. G., Pandolfi, J. M., and Clark, T. R.: Holocene sea level instability in the southern Great Barrier Reef, Australia: high-precision U–Th dating of fossil microatolls, *Coral Reefs*, 35, 625–639, 2016.

- Leonard, N. D., Welsh, K. J., Lough, J. M., Feng, Y. -x., Pandolfi, J. M., Clark, T. R., and Zhao, J. -x.: Evidence of reduced mid-Holocene ENSO variance on the Great Barrier Reef, Australia, *Paleoceanography*, 31, PA002967, 2016.
- Li, C.-H., Li, Y.-X., Zheng, Y.-F., Yu, S.-Y., Tang, L.-Y., Li, B.-B., and Cui, Q.-Y. 2018. A high-resolution pollen record from East China reveals large climate variability near the Northgrippian-Meghalayan boundary (around 4200 years ago) exerted societal influence. *Palaeogeogr. Palaeoclimatol., Palaeoecol.*, 512, 156–165, 2018.
- 630 Lirman, D., Schopmeyer, S., Manzello, D., Gramer, L. J., Precht, W. F., Muller-Karger, F., Banks, K., Barnes, B., Bartels, E., Bourque, A., Byrne, J., Donahue, S., Duquesnel, J., Fisher, L., Gilliam, D., Hendee, J., Johnson, M., Maxwell, K., McDevitt, E., Monty, J., Rueda, D., Ruzicka, R., and Thanner, S.: Severe 2010 cold-water event caused unprecedented mortality to corals of the Florida reef tract and reversed previous survivorship patterns, *PLoS ONE* 6, e23047, 2011.
- 635 Liu, Z., Kutzbach, J., and Wu, L.: Modeling climate shift of El Niño variability in the Holocene, *Geophys. Res. Lett.*, 27, 2265–2268, 2000.
- Liu, Z., Lu, Z., Wen, X., Otto-Bliesner, B. L., Timmermann, A., and Cobb, K. M.: Evolution and forcing mechanisms of El Niño over the past 21,000 years, *Nature*, 515, 550–553, 2014.
- Lybolt, M., Neil, D., Zhao, J., Feng, Y., Yu, K.-F., and Pandolfi, J.: Instability in a marginal coral reef: the shift from natural variability to a human-dominated seascape. *Front. Ecol. Environ.*, 9, 154–160, 2011.
- 640 Macintyre, I. G., and Glynn, P. W.: Upper limit of El Niño killoff, *Coral Reefs*, 9, 92, 1990.
- Macintyre, I. G., Glynn, P. W., and Cortés, J.: Holocene reef history in the eastern Pacific: mainland Costa Rica, Caño Island, Cocos Island, and Galápagos Islands, *Proc. 7th Int. Coral Reef Symp.*, Guam 2, 1174–1184, 1992.
- Macintyre, I. G.: Demise, regeneration, and survival of some western Atlantic reefs during the Holocene transgression, in: *Geological approaches to coral reef ecology*, edited by: Aronson, R. B., Springer, New York, NY, 181–200, 2007.
- 645 Marchant, R., and Hooghiemstra, H.: Rapid environmental change in African and South American tropics around 4000 years before present: a review, *Earth-Sci. Rev.*, 66, 217–260, 2004.
- Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional and global temperature for the past 11,300 years, *Science*, 339, 1198–1201, 2013.
- 650 Mayewski, P. A., Rohling, E. E., Stager, J. C., Karlén, W., Maasch, K. A., Meeker, L. D., Meyerson, E. A., Gasse, F., van Kreveld, S., Holmgren, K. and Lee-Thorp, J.: Holocene climate variability, *Quat. Res.*, 62, 243–255, 2004.
- McGregor, H. V., and Gagan, M. K.: Western Pacific coral $\delta^{18}\text{O}$ records of anomalous Holocene variability in the El Niño–Southern Oscillation, *Geophys. Res. Lett.*, 31, L11204, 2004.
- 655

- McGregor, H. V., Fischer, M. J., Gagan, M. K., Fink, D., Phipps, S. J., Wong, H., and Woodroffe, C. D.: A weak El Niño/Southern Oscillation with delayed seasonal growth around 4,300 years ago, *Nat. Geosci.* 6, 949–953, 2013.
- 660 McPhaden, M. J., Zebiak, S. E., and Glantz, M. H.: ENSO as an integrating concept in earth science, *Science*, 314, 1740–1745, 2006.
- Menounos, B., Clague, J. J., Osborn, G., Luckman, B. H., Lakeman, T. R., and Minkus, R.: Western Canadian glaciers advance in concert with climate change circa 4.2 ka, *Geophys. Res. Lett.*, 35, GL033172, 2008.
- Moy, C. M., Seltzer, G. O., Rodbell, D. T., and Anderson, D. M.: Variability of El Niño/Southern Oscillation activity at millennial timescales during the Holocene epoch, *Nature*, 420, 162–165, 2002.
- 665 Muscatine, L., and Porter, J. W.: Reef corals: mutualistic symbioses adapted to nutrient-poor environments, *Bioscience*, 27, 454–460, 1977.
- Neumann, C. J., Jarvinen, B. R., McAdie, C. J., and Hammer, G. R.: Tropical cyclones of the North Atlantic Ocean, 1871–1998, NOAA National Climatic Data Center, Asheville, North Carolina, 1999.
- Perry, C. T., Smithers, S. G., Gulliver, P., and Browne, N. K.: Evidence of very rapid reef accretion and reef growth under high turbidity and terrigenous sedimentation, *Geology*, 40, 719–722, 2012.
- 670 Precht, W. F., and Aronson, R. B.: Climate flickers and range shifts of reef corals, *Front. Ecol. Environ.*, 2, 307–314, 2004.
- Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., and Bigler, M.: A new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res.: Atmospheres*, 111(D6), 2006.
- 675 Reaka-Kudla, M. L., Feingold, J. S., and Glynn, P. W.: Experimental studies of rapid bioerosion of coral reefs in the Galápagos Islands, *Coral Reefs*, 15, 101–107, 1996.
- Rein, B., Lückge, A., Reinhardt, L., Sirocko, F., Wolf, A., and Dullo, W. -C.: El Niño variability off Peru during the last 20,000 years, *Paleoceanography*, 20, PA4003, 2005.
- 680 Richmond, R. H.: Energetic relationships and biogeographical differences among fecundity, growth and reproduction in the reef coral *Pocillopora damicornis*, *Bull. Mar. Sci.*, 41, 594–604, 1987.
- Richmond, R. H.: The effects of the El Niño / Southern Oscillation on the dispersal of corals and other marine organisms, in: *Global ecological consequences of the 1982–83 El Niño–Southern Oscillation*, edited by: Glynn, P. W., Elsevier, Amsterdam, 127–140, 1990.
- 685 Riegl, B., and Piller, W. E.: Possible refugia for reefs in times of environmental stress, *Int. J. Earth Sci.*, 92, 520–531, 2003.

- Rodríguez-Fonseca, B., Suárez-Moreno, R., Ayarzagüena, B., López-Parages, J., Gómara, I., Villamayor, J., Mohino, E., Losada, T. and Castaño-Tierno, A.: A review of ENSO influence on the North Atlantic, A non-stationary signal, *Atmosphere*, 7, atmos7070087, 2016.
- 690 Romero-Torres, M., Treml, E., Acosta, A., and Paz-García, D. A.: The eastern tropical Pacific coral population connectivity and the role of the Eastern Pacific Barrier, *Sci. Rep.*, 8, 9354, 2018.
- Rooney, J., Fletcher, C., Grossman, E., Engels, M., and Field, M.: El Niño influence on Holocene reef accretion in Hawai'i, *Pac. Sci.*, 58, 305–324, 2004.
- Ryan, E. J., Smithers, S. G., Lewis, S. E., Clark, T. R., and Zhao, J.-x.: The variable influences of sea level, 695 sedimentation and exposure on Holocene reef development over a cross-shelf transect, central Great Barrier Reef, *Diversity*, 10, 110, 2018.
- Sachs, J. P., Sachse, D., Smittenberg, R. H., Zhang, Z., Battisti, D. S., and Golubic, S.: Southward movement of the Pacific intertropical convergence zone AD 1400–1850, *Nat. Geosci.* 2, 519–525, 2009.
- Sandweiss, D. H., Maasch, K. A., Burger, R. L., Richardson III, J. B., Rollins, H. B., and Clement, A.: Variation in 700 Holocene El Niño frequencies: climate records and cultural consequences in ancient Peru, *Geology*, 29, 603–606, 2001.
- Selvaraj, K., Chen, C.-T. A., and Lou, J.-Y.: Holocene weak summer East Asian monsoon intervals in subtropical Taiwan and their global synchronicity, *Clim. Past Discuss.*, 4, 929–953 (2008).
- Staubwasser, M., Sirocko, F., Grootes, P. M., and Segl, M.: Climate change at the 4.2 ka BP termination of the Indus 705 valley civilization and Holocene south Asian monsoon variability, *Geophys. Res. Lett.*, 30, GL016822, 2003.
- Stuiver, M., Reimer, P. J., and Reimer, R.W.: CALIB 7.2 at <http://calib.org>, accessed 2018-7-25, 2018.
- Svendsen, J. I., and Mangerud, J.: Holocene glacial and climatic variations on Spitsbergen, Svalbard, *The Holocene*, 7, 45–57, 1997.
- Takahashi, K., Montecinos, A., Goubanova, K., and Dewitte, B.: ENSO regimes: reinterpreting the canonical and 710 Modoki El Niño, *Geophys. Res. Lett.*, 38, GL10704, 2011.
- Thompson, D. M., Conroy, J. L., Collins, A., Hlohowskyj, S. R., Overpeck, J. T., Riedinger-Whitmore, M., Cole, J. E., Bush, M. B., Whitney, H., Corley, T. L., and Kannan, M. S.: Tropical Pacific climate variability over the last 6000 years as recorded in Bainbridge Crater Lake, Galápagos, *Paleoceanography*, 32, PA003089, 2017.
- Toth, L. T., Aronson, R. B., Vollmer, S. V., Hobbs, J. W., Urrego, D. H., Cheng, H., Enochs, I. C., Combosch, D. J., 715 van Woesik, R., and Macintyre, I. G.: ENSO drove 2500-year collapse of eastern Pacific coral reefs, *Science*, 337, 81–84, 2012.
- Toth, L. T.: Holocene coral-reef development in the tropical eastern Pacific, Florida Inst. Tech. (Doctoral Dissertation), 402 p., 2013.

- 720 Toth, L. T., Aronson, R. B., Cobb, K. M., Cheng, H., Edwards, R. L., Grothe, P. R., and Sayani, H. R.: Climatic and biotic thresholds of coral-reef shutdown, *Nat. Clim. Change*, 41, 369–374, 2015a.
- Toth, L. T., Aronson, R. B., Cheng, H., and Edwards, R. L.: Holocene variability in the intensity of wind-gap upwelling in the tropical eastern Pacific, *Paleoceanography*, 30, 1113–1131, 2015b.
- Toth, L. T., Aronson, R. B., and Macintyre, I. G.: Holocene reef development in the eastern tropical Pacific, in: Coral Reefs of the Eastern Tropical Pacific, in: *Coral Reefs of the World 8*, edited by: Glynn, P. W., Manzello, D. P., and Enochs, I. C., Springer Science+Business Media, Dordrecht, 2017.
- 725 Toth, L. T., Kuffner, I. B., Stathakopoulos, A., and Shinn, E. A.: A 3000-year lag between the geological and ecological collapse of Florida’s coral reefs, *Global Change Biol.*, 2018. doi: 10.1111/gcb.14389.
- Urban, F. E., Cole, J. E., and Overpeck, J. T.: Influence of mean climate change on climate variability from a 155-year tropical Pacific coral record, *Nature*, 407, 989–993, 2000.
- 730 Vargas-Ángel, B., Zapata, F. A., Hernández, H., and Jiménez, J. M.: Coral and coral reef responses to the 1997–98 El Niño event on the Pacific coast of Colombia, *Bull. Mar. Sci.*, 69, 111–132, 2001.
- Walker, M. J., Berkelhammer, M., Björck, S., Cwynar, L. C., Fisher, D. A., Long, A. J., Lowe, J. J., Newnham, R. M., Rasmussen, S. O., and Weiss, H.: Formal subdivision of the Holocene Series/Epoch: a discussion paper by a working group of INTIMATE (Integration of ice-core, marine and terrestrial records) and the subcommission on Quaternary Stratigraphy (International Commission on Stratigraphy), *J. Quat. Sci.*, 27, 649–659, 2012.
- 735 Wang, B., Wu, R., and Li, T.: Atmosphere–warm ocean interaction and its impacts on Asian–Australian monsoon variation, *J. Clim.*, 16, 1195–1211, 2003.
- Webster, J. M., Braga, J. C., Humblet, M., Potts, D. C., Iryu, Y., Yokoyama, Y., Fujita, K., Bourillot, R., Esat, T. M., Fallon, S., Thompson, W. G., Thomas, A. L., Kan, H., McGregor, H. V., Hinestrosa, G., Obrochta, S. P., and Lougheed, B. C.: Response of the Great Battier Reef to sea-level and environmental changes over the past 30,000 years, *Nat. Geosci.*, 11, 426–432, 2018.
- 740 Weiss, H., Courty, M. A., Wetterstrom, W., Guichard, F., Senior, L., Meadow, R., and Curnow, A.: The genesis and collapse of third millennium north Mesopotamian civilization, *Science*, 261, 995–1004, 1993.
- Weiss, H.: Global megadrought, societal collapse and resilience at 4.2–3.9 ka BP across the Mediterranean and West Asia, *Clim. Chang. Cult. Evol. PAGES Mag.*, 24, 62, 2016.
- 745 Wellington, G. M., and Glynn, P. W.: Responses of coral reefs to El Niño–Southern Oscillation sea-warming events, in: *Geological approaches to coral reef ecology*, edited by: Aronson, R. B., Springer-Verlag, New York, 342–385, 2007.
- Wilkinson, C. (ed.): *Status of coral reefs of the world: 2000*, Australian Institute of Marine Science, Cape Ferguson and Dampier, 2000.
- 750

- Wood, S., Baums, I. B., Paris, C. B., Ridgwell, A., Kessler, W. S., and Hendy, E. J.: El Niño and coral larval dispersal across the eastern Pacific marine barrier, *Nat. Comm.*, 7, 12571, doi:10.1038/ncomms12571, 2016.
- Wu, Q., Zhao, Z., Liu, L., Granger, D. E., Wang, H., Cohen, D. J., Wu, X., Ye, M., Bar-Yosef, O., Lu, B., and Zhang, J.: Outburst flood at 1920 BCE supports historicity of China's Great Flood and the Xia dynasty, *Science*, 353, 579–582, 2016.
- 755
- Xiao, J., Zhang, S., Fan, J., Wen, R., Zhai, D., Tian, Z., Jiang, D.: The 4.2 ka BP event: multi-proxy records from a closed lake in the northern margin of the East Asian summer monsoon, *Clim. Past*, 13, 1417–1425.
- Xu, S., Yu, K., Tao, S., Wu, C. -C., Wang, Y., Jiang, W., Wang, S., and Shen, C. -C.: Evidence for the thermal bleaching of *Porites* corals from 4.0 ka B.P. in the northern South China Sea, *J. Geophys. Res.: Biogeosciences*, 123, 79–94, 2018.
- 760
- Yamano, H., Sugihara, K., Watanabe, T., Shimamura, M. and Hyeong, K.: Coral reefs at 34°N, Japan: Exploring the end of environmental gradients, *Geology*, 40, 835–838, 2012.
- Zheng, W., Braconnot, P., Guilyardi, E., Merkel, U., and Yu, Y.: ENSO at 6ka and 21ka from ocean-atmosphere coupled model simulations, *Clim. Dynam.*, 3, 745–762, 2008.

765 **Table 1: Summary of records of perturbations to coral-reef ecosystems around the time of the 4.2-ka event. Reported radiocarbon ages were calibrated with the marine radiocarbon curve and regional marine reservoir age corrections, using CALIB software (v.7.0.2; Stuiver et al., 2018).**

Type of impact	Location	Time period (ky BP)	Reference
Hiatus in reef development	Pacific Panamá		
	<i>Contadora</i>	4.3–1.5	Toth et al. 2012
	<i>Iguana</i>	4.2–1.6	Toth et al. 2012
	<i>Canales de Tierra</i>	4.0–1.8	Toth et al. 2012
	Costa Rica		
	<i>Golfo Dulce</i>	4.6–2.0	Cortés et. al 1994
	Great Barrier Reef		
	<i>Moreton Bay</i>	4.2–2.0	Lybolt et al. 2011
Shutdown of reef development	Japan		
	<i>Kodakara Is.</i>	4.4–4.0	Hamanaka et al. 2012
	Hawaii		
	<i>Kailua, O'ahu</i>	5.0–present	Rooney et al. 2004
	<i>Punalu'u, O'ahu</i>	4.9–present	Rooney et al. 2004
	<i>Hale o Lono, Moloka'i</i>	4.8–present	Rooney et al. 2004
	<i>Mana, Kaua'i</i>	4.5–0	Rooney et al. 2004
Coral mortality	China		
	<i>South China Sea</i>	4.2–3.8	Xu et al. 2018
Decline in coral $\delta^{13}\text{C}$	Pacific Panamá		
	<i>Contadora</i>	3.9–3.6	Toth et al. 2015
	China		
	<i>South China Sea</i>	3.8	Xu et al. 2018

770

775 **Figure 1: Photographs depicting the typical states of coral reefs in Pacific Panamá (A) before, (B) during, and (C) after an El Niño event. (A) shows an example of a well-developed coral reef dominated by branching *Pocillopora* corals. (B) shows a reef dominated by *Pocillopora* experiencing high levels of coral bleaching during the 2015–16 El Niño event. (C) shows a large area of *Pocillopora* rubble that has persisted since the 1982–83 El Niño event.**

780 **Figure 2. Coral-based reconstructions of oceanographic and climatic variability in Pacific Panamá from 5500 BP to present (after Toth et al., 2015a, 2015b). (A) provides a reconstruction of upwelling intensity in the Gulf of Panamá based on the location-specific radiocarbon-reservoir age-offset, ΔR (Toth et al., 2015b). (B & C) provide reconstructions of temperature and salinity variability for Pacific Panamá based on 200-yr running means ($\pm 95\%$ CIs) of Sr/Ca and $\delta^{18}\text{O}$ of corals sampled from a core collected from the reef at Contadora Island in the Gulf of Panamá. The individual data points used to generate the curves can be found in Toth (2013; Toth et al. 2015a). Modern values are means ($\pm 95\%$ CIs) measured from living corals collected from the same reef in 2011 (described in Toth et al. 2015a). The timing of the 4.2-ka event and the hiatus in the development of reefs in Pacific Panamá are indicated by the red and gray bars, respectively.**

790 **Figure 3: Schematic diagram of the reconstructed changes in ENSO variability in the tropical Pacific in relation to vertical reef-accretion in Pacific Panamá. Colored bars are based on a palaeoceanographic reconstruction from Pacific Panamá shown in Fig. 2; gray bars summarize the results of other paleoclimate studies from throughout the tropical Pacific. Superscripts reference the studies that support the climatic and oceanographic changes indicated in the bars: ¹Sandweiss et al., 2001; ²Rein et al., 2005; ³Zheng et al., 2008; ⁴Koutavas and Joanides, 2012; ⁵Cobb et al. 2013; ⁶Carré et al., 2014; ⁷Emile-Geay et al., 2016; ⁸Leonard et al., 2016b; ⁹Thompson et al., 2017; ¹⁰Toth et al. 2015a; ¹¹McGregor et al., 2013; ¹²Toth et al., 2015b; ¹³Conroy et al., 2008; ¹⁴Corrège et al. 2000; ¹⁵Haug et al. 2001; ¹⁶Moy et al., 2002.**

800 **Figure 4: Thermal anomalies associated with two of the strongest El Niño events (1997–98 and 2015–16) and La Niña events (1988–89 and 2010–11) in recent history. Black points on the maps indicate the locations of coral reefs. The imagery was created with data from NOAA’s Smith and Reynolds Extended, Reconstructed Sea-Surface Temperature, Level 4 Monthly Dataset (V5) using NASA’s PO.DAAC LAS V8.6.1 visualization tool (<https://podaac-tools.jpl.nasa.gov/las/>).**

805