



1

2

3

4 **Reassessing long-term drought risk and societal impacts in Shenyang,**
5 **Liaoning province, Northeast China (CE 1200 - 2015)**

6

7

8

9 LingYun Tang^{1*}, Neil Macdonald¹, Heather Sangster¹, Richard Chiverrell¹ and

10 Rachel Gaulton²

11

12 ¹Department of Geography, School of Environmental Planning, University of Liverpool,
13 Liverpool, L69, 3BX, U.K.

14 ² School of Natural and Environmental Sciences, Newcastle University, Newcastle upon
15 Tyne, NE1 7RU, UK

16

17

18

19 *Corresponding author E-mail: psltang@liverpool.ac.uk

20

21



22 **Reassessing long-term drought risk and societal impacts in Shenyang,**
23 **Liaoning province, Northeast China (CE 1200 - 2015)**

24

25 **Abstract**

26 The occurrence of two severe droughts in Northeastern China since CE 2000 has raised
27 attention in the risk presented by droughts. This paper presents a historic drought series for
28 Shenyang in the Liaoning province, NE China since CE 1200 to present, with a reconstructed
29 long precipitation series (1906-2015), augmented with historical documentary accounts.
30 Analysis of the instrumental series using a standardised precipitation index (SPI) and extending
31 it using historical records has produced a combined series spanning over eight centuries. The
32 combined long series was analysed for patterns in drought frequency, severity and typology.
33 Three droughts comparable to those since CE 2000 occur in the instrumental series during early
34 twentieth century (e.g. 1907, 1916-18 and 1920-21), and coeval archival sources reveal the
35 human impacts of these severe droughts. The archival sources demonstrate how reduced
36 vulnerability resulting from societal and cultural changes in the early twentieth century helped
37 prevent the loss of life experienced during comparable severe droughts at the end of the
38 nineteenth century (1887 and 1891). Incorporating a longer temporal perspective to drought
39 analysis shows that onset is often earlier than is documented explicitly within the archives, and
40 so combined SPI series for a region could provide an early warning of drought development
41 expressed as a water deficit in the previous year. Analysis of archival data provides a rich
42 historical description of impacts and societal responses to severe drought. The archives provide
43 a rich historical description of drought impacts and responses at the personal and community
44 level, whilst also detailing the different roles played by communities, state and international
45 organisations in responding to events.

46

47 **Keywords:** Drought; Reconstruction; Historical; Shenyang; Liaoning, China



48 **1 Introduction**

49 Drought is an world-wide problem, causing more deaths globally than any other natural disaster
50 (Delbiso et al., 2017); with over 485,000 deaths and more than 1.6 billion people adversely
51 affected during the last decade (2010-2019; EM-DAT, 2019). Drought is often a slow
52 developing pervasive environmental disaster that is hard to predict and manage, and a variety
53 of definitions for drought in operational use around the world, and there is no universal
54 definition of the term *drought* (Lloyd-Hughes, 2014), but many definitions focus on a
55 deficiency in precipitation over a period of time (Wilhite, 2000; Belal *et al.*, 2014). From a
56 macro perspective, drought is a long-term water deficit that develops slowly under long-term
57 natural conditions or human intervention, with a negative impact on nature and humans,
58 resulting in a shortage of water that causes adverse impacts on activities (e.g. food production)
59 or societal groups (e.g. farmers) (Dai, 2011a). Drought often begins following a prolonged
60 period of moisture deficiency (Lanen, 2006; Palmer, 1965) propagating through the
61 hydrological cycle, exhibiting differing spatial and temporal characteristics depending on a
62 variety of factors e.g. antecedent conditions and soil moisture (Heim, 2002; Todd et al., 2013).
63 Wilhite and Glantz (1985), classified droughts into four types: meteorological, hydrological,
64 agricultural, socioeconomic, with Mishra and Singh (2010) recommending the inclusion of a
65 fifth classification ‘ground water’ drought. Drought has been referred to as a ‘creeping
66 phenomenon’ (Mishra and Singh, 2010), and its impacts vary from region to region, with
67 drought effects exacerbated by other meteorological elements, such as temperature, wind, and
68 humidity (Brázdil et al., 2008). Palmer (1965, pp.1) notes that ‘drought means various things
69 to various people, depending on specific interest’. Droughts are complex so-called ‘natural’
70 hazards – the term ‘natural’ in natural hazards, although etymologically doubtful, because in a
71 sense all hazards are natural, maybe considered as ‘natural’ as sanctioned by a long-term use
72 in disaster research (Sangster et al., 2018), with droughts causing significant environmental,
73 social and economic impacts (Van Loon et al., 2016). Drought is an international phenomenon
74 with notable drought episodes throughout the twentieth and twenty-first centuries, e.g. 1930s
75 ‘Dust Bowl’ in the USA (Schubert et al., 2004); 1975-76 in Europe (Parry et al., 2012; Zaidman
76 et al., 2010); China 1994 & 2010-2011 (Zhang et al., 2019) and South Africa 2015-17 (Wolski,
77 2018). Over the last decade a number of studies have started to explored historical droughts
78 (Brázdil et al., 2009, 2018b) and the impacts experienced over decades to centuries on water
79 resources (Lennard et al., 2015); agriculture (Brázdil et al., 2018a); infrastructure (Harvey-
80 Fishenden et al., 2019); stream and river flows (Zaidman et al., 2010); groundwater
81 (Bloomfield and Marchant, 2013); with recent calls (e.g. Trnka et al., 2018) for more to be



82 done with existing data, particularly in understanding past socio-drought responses and
83 changes in vulnerability. Considerable work has been undertaken in recent decades in
84 developing robust and long flood and drought chronologies using combinations of archival
85 (Brázdil et al., 2018b; Yan et al., 2014; Zheng et al., 2006) and instrumental (Brázdil et al.,
86 2009) sources from around the globe, although much work to date has focused on Europe
87 (Wilhelm et al., 2018). The development of new online digitised sources has facilitated greater
88 historical analysis (Black and Law, 2004; Wang et al., 2018) with greater recognition from
89 regulatory authorities of the value of historical information (Kjeldsen et al., 2014).

90

91 China is one of the most natural disaster-prone countries in the world (Dai, 2011b; He et al.,
92 2011b; Lorbach et al., 2011), and droughts are a recurrent feature of the Chinese climate (He
93 et al., 2011a). Drought can be considered as the most disastrous natural hazard within China,
94 with over 465,000 deaths and more than 3.1 billion adversely affected from 1970-present and
95 12 million deaths since 1900 (EMDAT, 2019). Historically notable droughts in 1876-1878,
96 1928-1930 and 1958-62 resulted in widespread loss of life, poor harvest, leading to serious
97 social consequences of famine, robbery, unrest, and political instability (De Châtel, 2014;
98 Janku, 2018; Teklu et al., 1992; Yang et al., 2012). Between BC 206 and AD 1948, 1056 severe
99 droughts are recorded in Chinese history (Zhang, 2004, 2013); with 11 years of micro-drought,
100 13 years of light-drought, 14 years of mid-drought, 10 years of heavy-drought, and four years
101 of extreme-drought during the period 1949 to 2000 (Zhang et al., 2008b). Precipitation
102 recording in China has developed through time, with some of the most advanced recording
103 globally during the Qing Dynasty (CE 1644-1912), with both rainfall and snow depth recorded
104 in 1736 (Ge et al., 2005). The installation of better equipment through the 1920s and 1950s
105 saw many stations upgraded, with meteorological stations often retained; however, the
106 availability of metadata on early recorders is more limited. Past droughts have had a far-
107 reaching impact on society in China; a clear understanding current and future drought risk is
108 therefore critical. With population growth, economic development, urbanisation and climatic
109 change, drought is a global challenge, but felt acutely in China, producing threats to
110 environmental ecology, food security, impacting environmental ecology, urban and rural water
111 supply (Bohle et al., 1994; Homer-Dixon, 1994).

112

113 This paper examines the history of drought in the Shenyang region of Northeast China, the
114 spatial and temporal variability in droughts, the characteristics of droughts, and mechanisms
115 responsible and impacts on society. Our objectives are:



- 116 i. To develop and analyse a record of droughts and the documentary evidence for
117 associated impacts (CE 1200 AD - present), using a variety of sources including the
118 compendium of Chinese droughts produced by Zhang (2004, 2013);
119 ii. Identify and analyse contemporary droughts using the instrumented daily precipitation
120 series at Shenyang Meteorological Observatory (Station 54342: 1961-2015), and
121 augment this series with the longer monthly precipitation data for Shenyang (CE 1906-
122 1988);
123 iii. Generate a Standardised Precipitation Index (SPI-1, -6 and -12) for the augmented
124 precipitation series spanning the period CE 1906-2015; construct one of the longest
125 drought series (CE 1200 - present) combining the augmented instrumental series (ii)
126 with historical data (i), and then classify the different types of drought and event
127 severity; and,
128 iv. Analyse the patterns in drought frequency, severity and type for Shenyang, examining
129 the documented impacts and responses to drought to better understand how societal
130 vulnerability has changed through time.

131

132 **2 Study Area**

133 Shenyang (41.8°N 123.4°E) is the capital city of Liaoning Province in Northeast China (Figure
134 1), with a temperate continental monsoon climate, with temperature ranging from -17°C
135 (January) to 29°C (July), decreasing from southwest to northeast (plain to mountain) (Chen et
136 al., 2016); whilst average annual precipitation (500-1000 mm a⁻¹) increases from west to east
137 (Zhang et al., 2013). The Shenyang municipality is home to approximately 8M people in 2016.
138 The region has witnessed reductions (at 78% of stations) in annual precipitation over the period
139 1961-2008 (Liang et al., 2011). The Liaoning province is a primary grain producing region in
140 China; as such droughts and associated impacts on regional agricultural production are of
141 national importance, with previous studies detecting recent warming and reductions in
142 precipitation (Chen et al., 2016).

143

144 **3 Data and Methods**

145 **3.1 Data sources**

146 This study uses a variety of source materials including historical and instrumental datasets
147 detailed below.

148

149 **3.1.1 Documentary data**



150 The ‘A compendium of Chinese Meteorological Records of the Last 3,000 Years’ produced by
151 Zhang (2004) and updated in 2013, summarises 7835 historical sources from the earliest
152 existent materials in the Chinese language, the ‘Oracle Bones Collection’ (c.1600 BC) through
153 to more recent sources which describe meteorological incidences in China. The ‘Oracle Bones’
154 have a long history of being studied for meteorological information, with early studies
155 undertaken by Wittfogel (1940). There are also a small number of private diaries and court
156 memorial files of the Qing Dynasty, though the ‘History of Drought Archives in the Qing
157 Dynasty’ (Tan, 2012) provides a summary of the collection spanning from 1689 to 1911, with
158 more than one million pieces present in the Qing Dynasty palace archive. The China
159 Meteorological Disasters Ceremony (Liaoning volume) from Wen et al. (2005) provides
160 detailed accounts of drought alongside records of other disasters which may have been caused
161 by drought, such as famine and plague; a full list of source materials can be found in Table 1.
162 Over recent decades considerable effort has been placed into collating the archival materials
163 present across China detailing natural hazards, this wealth of information provides valuable
164 opportunities for further exploration; however, such volume limits the capacity for cross
165 checking and validation, with many sources not easily accessible. This has raised questions of
166 reliability and transparency, but as Bradley (2006) notes, the compendium produced by Zhang
167 (2004) clearly illustrates critical analysis, with careful checking for consistency and
168 discrepancies identified. Recent developments include a move to digitise these databases,
169 ensuring and maintaining high levels of archival practice, with the development of the
170 REACHES climate database (Wang et al., 2018).

171

172 In addition to the meteorological sources identified, information from sources detailing
173 agricultural activity provide valuable auxiliary reference materials, including the following
174 items: Shenyang local records (Meng, 1989; Shenyang Municipal People’s Government Local
175 Records Office (1994-2011), 2011); The year of flood and drought in Shenyang from 1276 to
176 1985 (Shenyang Municipal People’s Government Local Records Office, 1998). The following
177 datasets have been acquired from the Office of State Flood Control and Drought Relief (1999);
178 Farmland affected area from 1949 to 1990 in Liaoning Province Statistics on Drought Area of
179 Heavy Drought in Liaoning Province; Drought rating assessment in various regions of
180 Liaoning Province from 1949 to 1990; Drought Statistics in the Province from 1470 to 1949;
181 Comparison of Precipitation in Liaoning Province from 1949 to 1964 and from 1965 to 1990;
182 Comparison of grain yield per plant, drought frequency and drought reduction in various
183 regions of Liaoning Province; hydrological station data for Liaoning Province; Regular



184 frequency of continuous drought in dry season in Liaoning Province. Local newspapers have
185 also been accessed to corroborate records of droughts e.g. the Shengjing Times (reflecting the
186 old city name).

187

188 3.1.2 Instrumental data

189 Instrumental climate data are taken from two datasets, the first is long-term meteorological
190 data, including monthly precipitation (05/1905 to 12/1988) from the Research Data Archives
191 Computational & Information Systems Lab (NCAR, 1996), no records present for 1944-1946.
192 This dataset covers 60 relatively evenly distributed sites in China, with long records. The
193 second precipitation series was retrieved from the National Disaster Reduction Centre of China
194 (NDRCC), which provides daily data for air pressure (V10004), daily average temperature
195 (V12001), daily highest temperature (V12052), daily lowest temperature (V12053),
196 precipitation (V13201), average wind speed (V11002), sunshine hours (V14032), for the period
197 01/01/1961 to 31/05/2016. Analysis of these datasets permits varying temporal analysis of the
198 precipitation, with a long overlap period that can be used to compare the association of these
199 two data sets.

200

201 Previous studies have illustrated a strong relationship between droughts and ENSO anomalies
202 (Li et al., 2019; Zhang et al., 2018) for differing regions of China. However many of these
203 studies use relatively short series (1960-). The extended precipitation series (CE 1906-2015)
204 presented here provides a valuable opportunity to explore this relationship over a longer
205 timescale. The Niño 3.4 sea surface temperature index, defined as the area-averaged SST
206 anomalies over (5N–5S, 170–120W), compiled from PSD using the HadISST1 dataset for the
207 period CE 1870-2015 by Rayner et al., (2003) is used.

208

209 **3.2 Data processing**

210 3.2.1 Documentary data

211 The compendium provided by Zhang (2004, 2013) provides the framework for the early record
212 (pre-1911); however, great care was undertaken in assessing the historical record.

213

214 In the process of analysing documentary sources for Shenyang, it is necessary to pay particular
215 attention to historical changes to the name of Shenyang and the borders of the provinces. For
216 example, in the book “Zhong Guo Dong Bei Yu Dong Bei Ya Gu Dai Jiao Tong Shi” (Wang
217 and Pu, 2016), it is noted that during the Han Dynasty, ‘Liao Dong Jun’ was used for the



218 Shenyang area, whereas during the Dong Han Dynasty, the southern part of Shenyang
219 continued to belong to Liao Dong Jun, and the northern part belonged to Xuan Tu Jun (Zhao,
220 2006). In addition, the Gao Xian region is the recent Sujiatun area in Shenyang (Wang and Pu,
221 2016); Yan (2012) detailed the historical changes in the Shenyang (Table 2).

222

223 Historical records for all drought years are included where records exist, but historical records
224 for the following situations are excluded:

225 i. Information unclear - the disasters cause or event location is unclear. For example, in 1549,
226 the drought and locust disaster occurred in Xingcheng County of Liaoning Province
227 ("Ming Shi Zong Shi Lu", Vol. 353). In 1549 Xingcheng belonging to Liaoxi; however,
228 Shenyang belonged to Liaodong, therefore, this record is not in the target region and is
229 excluded.

230 ii. The record does not clearly state drought or that caused by drought. Although there are
231 many types of event that are associated/related to droughts, such as locusts, epidemic
232 disease or famine, where historical records do not directly state drought or attribute the
233 cause to drought they are excluded. For example, in October 1551, the Liaodong area did
234 not collect grain tax because of disasters ("Ming Shizong Record", Vol. 3, 7:8). The record
235 does not specifically state that a drought occurred though this is a common response to a
236 drought.

237

238 3.2.2 Instrumental data

239 Data quality assessment and management of both long and shorter series was required to ensure
240 homogenisation and data suitability. Total precipitation includes both liquid and equivalent
241 frozen precipitation. All meteorological variables are recorded as one-tenth of their specific
242 units (mm), but are converted to mm throughout. For both instrumental series, care and
243 attention was taken with the original data series quality, with the data descriptors recorded in
244 Table 3. At Shenyang meteorological station, missing data occurred eight times (representing
245 0.826% of the record), and rainfall was marked three times with 'R', reflecting monthly totals
246 identical to the previous month, raising concerns as to the validity of the data (01-02/1906,
247 12/1908-01/1909 and 12/1968-01/1969). There is a reduction of available meteorological data
248 during the years 1943-46 following WWII across much of eastern China, as such no suitable
249 local sites could be identified to infill this series; for other missing monthly data, the monthly
250 averages are included where single months are missing, as often other local stations also have



251 missing data. For the shorter instrumental daily precipitation series (source 2), data descriptors
252 are included in Table 3, including percentage of record impacted.

253

254 Analysis of the two series coeval years of record (1961-1988) was undertaken, a Q-Q plot was
255 undertaken to verify that both data sources are normally distributed (Figure 2a). Figure 2b
256 shows a good linear distribution (p-value of 0.028); however, differences between the series
257 exist. During the period 1961-1988, the average difference between the two datasets is 12.72
258 mm and the maximum is 313.2, which occurred in October 1974; further examination reveals
259 that all the differences occurring in the period 1961-1979, with the two datasets producing
260 identical values for all months from 1980 onwards, this replicability in the later records
261 provides confidence in extending Source 1 through to the present (2015). Analysis of the
262 dispersion and outliers for each month was also undertaken (Figure 2c), the months with
263 greatest discrepancy are March and April, possibly reflecting challenges of recording snow/ice
264 fall. Comparison of the monthly and seasonal precipitation patterns presented in Figure 3 for
265 Shenyang for the period 1906-2015 using the new augmented series illustrate that some of the
266 anomalous values from source 2 from the period 1961-1979 appear unrealistic, e.g. 04/1964,
267 285.9 mm, with an average normally of c. 50 mm. An analysis of the variability in the
268 precipitation is presented (Figure 4), with the lowest precipitation (the driest, 1913: 341.1 mm
269 a^{-1}) and highest (wettest) years noted (1923: 1064.9 mm a^{-1} ; Figure 4a); a seasonal analysis and
270 long term trend is also presented (Figure 4b-e) with a 30-year Savitzky-Golay filter presented
271 (Savitzky and Golay, 1964).

272

273 **3.3 Drought Identification**

274 Using the combined instrumental and archival source materials, a record of droughts will be
275 reconstructed for Shenyang, the droughts will be explored and examined from a number of
276 perspectives including: type of drought (classification), intensity/magnitude, frequency and
277 trends; together these characterise the drought structure.

278

279 **3.3.1 Standardized Precipitation Index (SPI)**

280 A number of drought indices have been developed Heim (2002). Meteorological drought
281 indicators can be divided into two categories focused on either the physical mechanisms of
282 drought or the statistical distribution of meteorological elements; the SPI belongs to the latter
283 group and is widely used (Lennard et al., 2015; Mckee et al., 1993). As the long precipitation
284 series includes only monthly data, the Standardised Precipitation Index (SPI) is used, this index



285 has a number of advantages when used over long timescales compared to other potential
286 drought indices. The SPI developed by Mckee et al., (1993), is a widely applied meteorological
287 drought index that quantifies precipitation deficits or excess across different climates at
288 multiple timescales, typically of 1–24 months, however the simplicity of the SPI (precipitation
289 as the only input) causes some limitations too, e.g. no consideration of evaporative demand
290 (Vicente-Serrano et al., 2014). SPI values are dimensionless units, with negative values
291 indicating drier than normal conditions and positive values wetter than normal conditions.
292 Drought onset is generally assumed to occur at SPI values exceeding ≤ 1 , however the National
293 Standards of People’s Republic of China (2017) classification uses ≤ 0.50 as indicative of
294 drought onset, with drought termination identified as when SPI returns to ≥ 0 (Table 4a). SPI
295 can be used to characterise drought duration, severity and timing of onset and termination
296 (together known as the drought structure), based on the classifications identified in Table 4a;
297 the SPI classification recommended in China (National Standards of People’s Republic of
298 China, 2017) differs slightly from that of the WMO (2012; Table 4 (a and c), though others
299 have also proposed regionally specific SPI versions based on Mckee et al. (1993) e.g. Moreira
300 et al. (2008) for Portugal. Drought duration is determined by the number of months between
301 drought onset (SPI ≤ 0.49) and termination (SPI ≥ 0), drought severity is categorised using the
302 SPI classification system with peak severity the minimum SPI value recorded during the
303 drought. Within this study SPI will be examined at 3 temporal scales SPI-1 (1 month), SPI-6
304 (6 months), and SPI-12 (12 months) (Figure 5a-c). The SPI was determined by fitting a
305 probability density function to selected accumulation periods using L-moments to estimate
306 parameters. A gamma probability density distribution was found to be the most appropriate fit,
307 using a Kolmogorov-Smirnov (K-S) test to compare empirical and theoretical fit, calculating
308 the cumulative probability. This was then converted into the standard normal distribution, with
309 transformation of the cumulative probability of the fitted distribution to standard normal
310 distribution to define the SPI value (Lloyd-Hughes and Saunders, 2002; Vicente-Serrano et al.,
311 2010). Other univariate distributions have been recommended where a gamma distribution is
312 not appropriate (Barker et al., 2016; Stagge et al., 2015).

313

314 3.3.2 Documentary analysis

315 Documentary data provides additional information beyond that offered by instrumental series,
316 providing valuable information detailing both societal impacts and responses to past events
317 (Pfister, 2010). At Shenyang, the first recorded drought occurs in 347 AD, but only three events
318 are recorded during the period CE 347-1200, therefore the records analysed within this paper



319 start post CE 1200, as the frequency of records increases. Previous studies (e.g. Brázdil et al.,
320 2009; Hanel et al., 2018; Todd et al., 2013) using historical archival sources have examined
321 qualitative records and used a variety of different indices or grades of drought. The use of
322 ordinal index systems for the classification of descriptive accounts in historical climatology is
323 common, with a range of classes used e.g. Nash et al., (2016) used a +2 to -2 classification in
324 examining wet/dry phases in Natal and Zululand in Southern Africa. In augmenting the
325 instrumental with the historical series, clear benefits can be achieved if the descriptive
326 classification is comparable to the SPI drought classification applied in China (Table 4a).
327 Therefore, five drought classes are used in considering the historical descriptions, allowing
328 alignment between the two data forms, typical types of descriptor for each of the five classes
329 are presented in Table 5.

330

331 Analysing the historical records unearthed different forms of drought which broadly reflect the
332 five drought classes identified by Mishra and Singh (2010); meteorological, hydrological and
333 agricultural are comparable, the difference being few accounts detail groundwater droughts
334 (incorporated into hydrology within this study), with the socio-economic class being split into
335 economic (impacts of clear cost) and social impact (impacts on people e.g. health). In splitting
336 the socio-economic class into economic and social impact the wealth of materials present in
337 the historical record examining these aspects can be examined in greater depth. Each of the
338 different classes of drought increases in impact severity (Table 5) in documenting each of these
339 an assessment of the interrelationship between different types of impact can be made, for
340 example, the point at which food relief may be initiated, or tax payments suspended (typically
341 class 2/3), others such as praying for rain/snow are associated with high classes (4/5), reflecting
342 personal, community and governmental responses (e.g. government control of food prices).

343

344 Annual drought values for the instrumental period (CE 1906-2015) are represented by the
345 minimum SPI-12 value within each calendar year; within the documentary accounts the most
346 severe class of drought is used to determine the classification.

347

348 3.3.3 Drought trend and frequency analysis

349 The combined long-term drought series for Shenyang (CE 1200-2015) permits an analysis of
350 the long term drought trends and patterns. Clearly over such a long timescale a number of
351 socio-political and cultural changes will have occurred (Bavel et al., 2019), which may
352 influence the extent or severity of a particular drought and the capacity a population has to



353 respond to a drought of any given magnitude or severity (Keenan and Krannich, 2010; Kreibich
354 et al., 2019; Mechler and Bouwer, 2015). Human interventions may mitigate and/or exacerbate
355 the impacts of drought downstream through hydrological system management and engineering
356 (He et al., 2017). The socio-political and cultural circumstances during each recorded drought
357 will represent an important underpinning in considering long-term drought trends and
358 variability and will be considered individually in each instance (see discussion by Brázdil et
359 al., 2020).

360

361 An analysis of the different types of drought will be undertaken, assessing long term variability,
362 severity and frequency, including examination of where droughts have been documented
363 during the instrumental period. The severity of droughts will be considered using the different
364 classes of drought, examining whether any notable differences in drought type emerge, which
365 may help determine underlying changes in vulnerability through time. The reliability of the
366 historical account classification process was assessed for the period 1906-2015 by statistical
367 analysis (Spearman - ordinal drought class) of the assigned drought class to annual minimum
368 SPI.

369

370 The principal challenge identified within this study is in attempting to assess droughts defined
371 between those characterised by the historical analysis which is subjective and that defined by
372 the indices (SPI), which assumes a distribution with predefined probabilities attributed to each
373 class (Guttman, 1998). Whilst an advantage in drought risk analysis, this makes it challenging
374 for comparison to a subjective classification.

375

376 **4 Results and Discussion**

377 **4.1 Temporal analysis of instrumental time series**

378 The augmented precipitation series illustrates the range of precipitation experienced at
379 Shenyang over the last 100 years, with a maximum annual rainfall of 1064.9mm (1923) and a
380 minimum of 341mm (1913). The mean of 704 mm is slightly higher than the median value (red
381 dashed line; Figure 4a). Of the 28 years annual rainfall below the quartile, 10 occur prior to
382 1960 and 18 after. Precipitation at Shenyang is concentrated in the summer months, with little
383 winter precipitation (Figure 3), typical of a continental climate. Documentary accounts often
384 discuss spring droughts in Shenyang, which hinders the development of crops at the start of the
385 growing season (Wang et al., 2019).

386



387 Seasonal analysis of precipitation (CE 1906-2015: Figure 4b & 4e) illustrates that precipitation
388 in winter and spring gradually increases with time, with a slight reduction of summer and
389 autumn precipitation, but are statistically insignificant (at 0.05 level; Figure 4c and d). The
390 most severe spring drought occurred in 2001, with only 33.7mm spring precipitation, this is
391 supported with widespread media coverage of the drought in Shenyang and more widely in
392 Liaoning. The worst summer drought occurred in 2014 (170.6mm), with precipitation less than
393 fifty percent of the norm, presenting the worst summer drought since 1961; in response the
394 Liaoning provincial government instigated a level III drought emergency response, this
395 included additional funding from central government (150M yuan) and provincial departments
396 (70M yuan) (Wang, 2014), with drought relief teams created to support community water
397 infrastructure projects (Sun, 2015).

398

399 The SPI generated from the long precipitation series is analysed at SPI-1, -6 and -12, with SPI-
400 1 suited to short-term (monthly) analysis, with SPI-6 appropriate for seasonal drought analysis
401 and SPI-12 for annual to multi-annular droughts. SPI-6, with scores of ≤ -2 (severe droughts)
402 occur 14 times during the 110 year record (Figure 5b and c). There are six severe drought years
403 before (1907, 1913, 1914, 1917, 1920, 1926) and eight (1961, 1963, 1965, 1989, 1997, 2000,
404 2014, 2015) after 1960, with several of these constituting multi-annular droughts. There are
405 seven droughts that exceed ≤ -2 in the SPI-12 series (Figure 5c).

406

407 **4.2 Drought classification and trends**

408 The reconstruction of historical droughts in Shenyang is divided into two parts. The first
409 obtains drought class information from the SPI for the period CE 1906-2015 from an
410 augmented instrumental series. The second uses historical documents and is defined based on
411 specific classification criteria shown in Table 3 producing a long drought reconstruction from
412 1200 AD to 2015, with documentary (coloured) and instrumental data (black) for Shenyang
413 (Figure 6b). Analysis of the period 1906-2015 demonstrates a non-statistically significant
414 correlation exists in the relationship between annual minimum SPI-12 and documentary
415 drought class for any given year, of the 107 years of record, 42 record both an SPI and
416 descriptive account of drought. The relative absence of class 1 events in the documentary
417 record suggests that no account is often made during 'normal' conditions, with absence of
418 record often likely reflecting no drought, therefore the analysis was repeated, years with no
419 description were attributed to class 1, as a result a statistically significant relationship is
420 identified (Spearman, $p < 0.05$). There is a clustering of events during the fifteenth and sixteenth



421 centuries, coeval with a previously identified reduced monsoon phase in Central China (Zhang
422 et al., 2008a) and the Spörer period (CE 1460-1550) of reduced solar activity, a relative
423 quiescent phase is then noted between CE 1600-1750 with few droughts recorded (Figure 6b).
424 A number of droughts occurring in the period CE 1750-1880 AD are documented; however,
425 the frequency and severity of droughts increases thereafter (Figure 6c). The first drought year
426 with an assessment of class 5 occurs in March 1883, with the Shenyang chronicles referring to
427 drought, a cholera epidemic, and more than 20,000 deaths in a week (Shenyang Municipal
428 People's Government Local Records Editing Office, 1989). This was followed by a second
429 event in 1891, with documentary sources detailing famine and over 20,000 estimated deaths
430 (Wen et al., 2005). Table 7 summarizes the frequency of droughts at Shenyang in each century,
431 with a small peak in Shenyang drought frequency from CE 1501-1600, drought frequency then
432 decreased until the nineteenth century (Figure 6c). The frequency of class 4-5 drought events
433 indicates an increase during the nineteenth century, but this is not evenly distributed with most
434 of those events occurring in the period 1906-1921 (1907, 1913-14, 1916-18 and 1920-21), with
435 only three severe droughts events after 1921 in Shenyang in 1968-9, 1999-2002 and 2014-15
436 (Figure 6a). The documentary accounts in the period 1906-2015, record a number of class 2-4
437 drought events, but few events are classified as either 1 or 5, although the presence of the early
438 droughts in the period 1906-21 are corroborated.

439

440 The types of drought recorded within the records are indicated in Figure 6b, these illustrate that
441 the majority of records document meteorological drought conditions followed by economic
442 impacts. The drought severity in the descriptive accounts places most of documented droughts
443 in class 2, 3 or 4 (Figure 6b) The absence of deaths being documented restricts the number of
444 class 5 socio-drought, although the drought of 1920-21 is documented as a class 5 hydrological
445 drought, the only documentary class 5 event in the twentieth century. It may be that such
446 information was not published, and/or that the droughts within the Liaoning province did not
447 lead to such impacts, as few events prior to the late nineteenth century approach class 5. In
448 focussing on the city of Shenyang, there is also a risk that the impacts differed within the city
449 to those experienced in rural communities within the province, thereby reducing the number of
450 agricultural drought documented. Future works should therefore focus at the provincial scale
451 to incorporate a wider diversity of impact.

452



453 **4.3 Societal vulnerability to droughts**

454 The transformation of responses in Shenyang from *pre-industrial (folk)*, to *industrial*
455 (*technological*) and subsequently *post-industrial* (Chester et al., 2012; White, 1974) during the
456 period of study presents challenges in assessing and comparing impacts. Recent droughts of
457 comparable meteorological severity, e.g. 2014 (SPI -2.8) to those of the early twentieth century,
458 namely 1906 (-2.6), 1917 (-2.8) or 1921 (-2.5) illustrate how the responses and resulting
459 impacts potentially differed. In analysing these events the consequences of the droughts
460 differed considerably, whilst these events do not record deaths among the population in
461 Shenyang and/or Liaoning province they are severe, with the 1920-21 drought described as
462 “Spring drought for several months, well and river dry up, land dry up, no harvest at all, winter
463 disaster victims everywhere, people live in hunger and cold move out from the mountain village,
464 village empty” [class 4 socio-drought but class 5-hydrological] (Office of State Flood Control
465 and Drought Relief, 1999, p.388), across China an estimated 500,000 died (Edwards, 1922).
466 Analysis of the international media at the time reporting on the event is shaped by the socio-
467 political circumstances, with The Times (London) recording 3M as being displaced (9 Nov.
468 1920 p.11); however, as Fuller (2011) importantly notes this is often viewed from an
469 international perspective, with local relief providers often failing to receive recognition. The
470 responses to the drought varied, but included those expected within an *industrial* framework,
471 with both national and international relief occurring, but also local support complimenting *pre-*
472 *industrial* responses, with the Shengjing Times (1920) reporting on the 1st July that “Chief
473 Zhang set up an alter begging for rain” (6080, p.4). However, as Li (2007) notes in north China,
474 population increases without apparent agricultural intensification or expansion during the late
475 nineteenth century may have contributed to an increased susceptibility to drought associate
476 harvest fluctuations. In comparison during the 2014 drought which resulted in a Level III
477 emergency response, itself a notable difference from 1920 as a plan was in place, a number of
478 responses were deployed to mitigate the impacts of the drought, these included: the provision
479 of central and provincial relief funds (see section 4.1); water transfer of 400M m³ from the
480 Hun River, securing domestic and agricultural provisions (Sun, 2015); and the provision of
481 relief service teams to support local infrastructure improvements e.g. drilling new wells and
482 supply or water to over 32,000 people suffering shortages (Wang, 2014). The impacts of the
483 drought were widely reported in the media, with notably commentary focused on the impacts
484 to water supplies and food production: “Food production in Liaoning... estimated to decline
485 by 5 billion kg this year” (China Daily, 2014). Whilst both events 1920-21 and 2014 were
486 severe droughts, the relief planning and coordinated effort coupled with improved



487 infrastructure and a more stable socio-political environment facilitated a more efficient
488 response.

489

490 **4.4 Contemporary droughts and generating mechanisms**

491 Analysis of contemporary droughts through coupled documentary sources and SPI provide
492 valuable insights into the importance of drought severity and duration on associated impacts.

493 The ‘severe drought’ as defined by the SPI of 1968 (SPI -2.13, duration 26 months) appears to
494 have a relatively limited impact in Liaoning province, with few accounts recording particularly
495 notable impacts beyond reduced agricultural output, whereas, interestingly, the drought of
496 08/1979-07/1983, whilst not a severe from the perspective of the SPI (-1.8) but of longer
497 duration (47 months) receives greater coverage within the documentary accounts, possibly
498 reflecting the duration and cumulative impact on agriculture. This is further supported as the
499 drought of 07/1999-04/2002 (SPI -2.3, duration 34 months) receives similar levels of
500 documentary coverage to that of 1979-83 and 07/2014-15 (SPI -2.8; 18 months but extends
501 beyond the end of the record) also receives more detailed descriptions.

502

503 Documentary accounts often identify that droughts begin in the spring months, but the SPI
504 results suggest that deficits often appear in the previous late summer (e.g. 1968-1969 and 1999-
505 2002 droughts), suggesting that the impacts of dry previous summer and/or autumn are not
506 particularly noted within the documentary accounts, and it is only when the impacts are felt
507 that the consequences are noted. Analysis of the seasonal precipitation to the seasonal ENSO3.4
508 series shows no significant correlations, but annual minimum SPI has a significant (95% level)
509 correlation with ENSO3.4 Summer ($p=0.0168$) and Autumn ($p=0.0228$) for the period 1906-
510 2015.

511

512 **5 Summary**

513 Our analysis capitalises on the long instrumental and documentary accounts available for
514 Shenyang and the Liaoning province in NE China, by constructing a homogenised precipitation
515 (SPI) series for CE 1906-2015, and a long documentary drought series CE 1200-2015.
516 Previously documented notable droughts in the early twentieth century (1907, 1916-18, 1920-
517 21) are compared to the droughts of the last two decades (1999-2002 and 2014-15), illustrating
518 that these have comparable drought structures, with duration potentially being more important
519 than the specific drought severity when considering the societal impacts. It illustrates that
520 recent severe droughts (1999-2002 and 2014-15), whilst notable, are not unusual within the



521 region, with several similar magnitude events in the early twentieth century. Societally the most
522 impactful droughts in the region occurred in the late nineteenth century (1883 and 1891), whilst
523 appearing of comparable structure to those that occurred later (e.g. 1920-21 and 2014-15),
524 social and cultural circumstances resulted in greater social disruption and vulnerability.
525 Reduced vulnerability to severe droughts is evident from the early twentieth century as greater
526 drought mitigation planning and central support are available (see responses to 1920-21 and
527 2014-15 drought, section 4.3). The relative low number (one) of documentary accounts
528 recording class 1 events reflects preferential recording of notable events, and remains
529 challenging in any documentary analysis reconstructing climate, as mundane conditions are
530 often overlooked and therefore unrecorded.

531

532 The calibration and augmentation of historical records with the instrumental series using the
533 SPI presents challenges. Whilst there appears to be good agreement of drought classes 2-4, the
534 probabilistic underpinning of the SPI inevitably ensures some high magnitude drought events
535 are present (class 5), however this is not necessarily reflected within the documentary sources
536 for all drought types. The impact of the probabilistic SPI structure potentially over recording
537 class 5 events is mitigated to some degree with the application of a long precipitation series,
538 where the potential of such events to be recorded increases. Analysis of the documentary
539 droughts in the late nineteenth century suggests that the duration is comparable to those of the
540 early twentieth century, with similar generating mechanisms, a dry winter and/or spring
541 followed by a hard drought in late summer, often spanning multiple years, however the impacts
542 on the communities differ. Our identification of a ‘build-up’ period prior the severest droughts
543 (and their associated impacts) is notable, which is further reinforced by the significant relationship
544 to summer and autumn ENSO3.4 and should be incorporated into future drought management
545 plans, enabling the effective preparation of drought plans.

546

547 **Data availability**

548 The precipitation series are available from 1. Carbon Dioxide Information Analysis
549 Center/Environmental Sciences Division/Oak Ridge National Laboratory/U. S. Department of
550 Energy (1996): Two Long-Term Instrumental Climatic Data Bases of the People's Republic of
551 China. Research Data Archive at the National Center for Atmospheric Research,
552 Computational and Information Systems Laboratory. <http://rda.ucar.edu/datasets/ds578.5/>.
553 Accessed† 10-12-2018. The second series (1961-2015) daily precipitation was supplied by



554 National Disaster Reduction Centre of China, data use and access permitted through their
555 involvement in project (NE/P015484/1).

556

557 **Acknowledgement**

558 NM, RG were funded through a NERC-GCRF grant (NE/P015484/1). LT was supported by
559 the China Scholarship Council studentship at the University of Liverpool, supervised by NM,
560 HS and RC. We would like to thank the National Disaster Reduction Centre of China and the
561 National Engineering Research Center for Information Technology in Agriculture for
562 assistance within the DRIER-China project and data provision.

563



564 **References**

- 565 Barker, L. J., Hannaford, J., Chiverton, A. and Svensson, C.: From meteorological to
566 hydrological drought using standardised indicators, *Hydrol. Earth Syst. Sci.*, 20, 2483–2505,
567 doi:10.5194/hess-20-2483-2016, 2016.
- 568 Bavel, B. J. P., Curtis, D. R., Hannaford, M. J., Moatsos, M., Roosen, J. and Soens, T.: Climate
569 and society in long-term perspective: Opportunities and pitfalls in the use of historical datasets,
570 *Wiley Interdiscip. Rev. Clim. Chang.*, Online first, doi:10.1002/wcc.611, 2019.
- 571 Belal, A. A., El-Ramady, H. R., Mohamed, E. S. and Saleh, A. M.: Drought risk assessment
572 using remote sensing and GIS techniques, *Arab. J. Geosci.*, 7(1), 35–53, doi:10.1007/s12517-
573 012-0707-2, 2014.
- 574 Black, A. R. and Law, F. M.: Development and utilization of a national web-based chronology
575 of hydrological events/Développement et utilisation sur internet d'une chronologie nationale
576 d'événements hydrologiques, *Hydrol. Sci. J.*, 49(2), 237–246,
577 doi:10.1623/hysj.49.2.237.34835, 2004.
- 578 Bloomfield, J. P. and Marchant, B. P.: Analysis of groundwater drought building on the
579 standardised precipitation index approach, *Hydrol. Earth Syst. Sci.*, 17(12), 4769–4787,
580 doi:10.5194/hess-17-4769-2013, 2013.
- 581 Bohle, H. G., Downing, T. E. and Michael, J.: Climate change and social vulnerability, *Glob.*
582 *Environ. Chang.*, 4(1), 37–48, doi:10.1016/0959-3780(94)90020-5, 1994.
- 583 Bradley, R. S.: Book Review: A compendium of Chinese meteorological records of the last
584 3,000 years, *The Holocene*, 16(4), 621–622, doi:10.1177/095968360601600415, 2006.
- 585 Brázdil, R., Trnka, M., Dobrovolný, P., Chromá, K., Hlavinka, P. and Žalud, Z.: Variability of
586 droughts in the Czech Republic, *Theor. Appl. Climatol.*, doi:10.1007/s00704-008-0065-x,
587 2008.
- 588 Brázdil, R., Trnka, M., Dobrovolný, P., Chromá, K., Hlavinka, P. and Zcaron;alud, Z.:
589 Variability of droughts in the Czech Republic, 1881–2006, *Theor. Appl. Climatol.*, 97(3–4),
590 297–315, doi:10.1007/s00704-008-0065-x, 2009.
- 591 Brázdil, R., Možný, M., Klír, T., Řezníčková, L., Trnka, M., Dobrovolný, P. and Kotyza, O.:
592 Climate variability and changes in the agricultural cycle in the Czech Lands from the sixteenth
593 century to the present, *Theor. Appl. Climatol.*, 1–21, doi:10.1007/s00704-018-2508-3, 2018a.
- 594 Brázdil, R., Kiss, A., Luterbacher, J., Nash, D. J. and Řezníčková, L.: Documentary data and
595 the study of past droughts: a global state of the art, *Clim. Past*, 14(12), 1915–1960,
596 doi:10.5194/cp-14-1915-2018, 2018b.
- 597 Brázdil, R., Kiss, A., Řezníčková, L. and Barriendos, M.: Droughts in Historical Times in
598 Europe, as Derived from Documentary Evidence, pp. 65–96, Springer, Cham., 2020.
- 599 De Châtel, F.: The Role of Drought and Climate Change in the Syrian Uprising: Untangling
600 the Triggers of the Revolution, *Middle East. Stud.*, 50(4), 521–535,
601 doi:10.1080/00263206.2013.850076, 2014.
- 602 Chen, T., Xia, G., Wilson, L. T., Chen, W. and Chi, D.: Trend and Cycle Analysis of Annual
603 and Seasonal Precipitation in Liaoning, China, *Adv. Meteorol.*, 2016, 1–15,
604 doi:10.1155/2016/5170563, 2016.



- 605 Chester, D. K., Duncan, A. M. and Sangster, H.: Human responses to eruptions of Etna (Sicily)
606 during the late-Pre-Industrial Era and their implications for present-day disaster planning, *J.*
607 *Volcanol. Geotherm. Res.*, 225–226, 65–80, doi:10.1016/j.jvolgeores.2012.02.017, 2012.
- 608 Dai, A.: Drought under global warming: A review, *Wiley Interdiscip. Rev. Clim. Chang.*, 2(1),
609 45–65, doi:10.1002/wcc.81, 2011a.
- 610 Dai, A.: Drought under global warming: A review, *Wiley Interdiscip. Rev. Clim. Chang.*, 2(1),
611 45–65, doi:10.1002/wcc.81, 2011b.
- 612 Delbiso, T. D., Rodriguez-Llanes, J. M., Donneau, A. F., Speybroeck, N. and Guha-Sapir, D.:
613 Drought, conflict and children’s undernutrition in Ethiopia 2000–2013: A meta-analysis, *Bull.*
614 *World Health Organ.*, 95(2), 94–102, doi:10.2471/BLT.16.172700, 2017.
- 615 Edwards, D. W.: The North China famine of 1920-1921, with special reference to the west
616 Chihli area. Being the report of the Peking united international famine relief committee.,
617 [Printed by the Commercial Press works Ltd.], Peking. [online] Available from:
618 [https://www.worldcat.org/title/north-china-famine-of-1920-1921-with-special-reference-to-](https://www.worldcat.org/title/north-china-famine-of-1920-1921-with-special-reference-to-the-west-chihli-area-being-the-report-of-the-peking-united-international-famine-relief-committee/oclc/5452960)
619 [the-west-chihli-area-being-the-report-of-the-peking-united-international-famine-relief-](https://www.worldcat.org/title/north-china-famine-of-1920-1921-with-special-reference-to-the-west-chihli-area-being-the-report-of-the-peking-united-international-famine-relief-committee/oclc/5452960)
620 [committee/oclc/5452960](https://www.worldcat.org/title/north-china-famine-of-1920-1921-with-special-reference-to-the-west-chihli-area-being-the-report-of-the-peking-united-international-famine-relief-committee/oclc/5452960), 1922.
- 621 EM-DAT: The Emergency Events Database, edited by D. Guha-Sapir, EM-DAT Emerg.
622 Events Database, (Brussels, Belgium.) [online] Available from: www.emdat.be (Accessed 6
623 August 2019), 2019.
- 624 Fuller, P.: ‘Barren Soil, Fertile Minds’: North China Famine and Visions of the ‘Callous
625 Chinese’ *Circa* 1920, *Int. Hist. Rev.*, 33(3), 453–472, doi:10.1080/07075332.2011.595236,
626 2011.
- 627 Ge, J.: *Chinese Immigration History*, Wu-Nan Book Inc., Shandong., 2005.
- 628 Ge, Q.-S., Zheng, J.-Y., Hao, Z.-X., Zhang, P.-Y. and Wang, W.-C.: Reconstruction of
629 Historical Climate in China: High-Resolution Precipitation Data from Qing Dynasty Archives,
630 *Bull. Am. Meteorol. Soc.*, 86(5), 671–680, doi:10.1175/bams-86-5-671, 2005.
- 631 Ge, Y., Dou, W., Gu, Z., Qian, X., Wang, J., Xu, W., Shi, P., Ming, X., Zhou, X. and Chen, Y.:
632 Assessment of social vulnerability to natural hazards in the Yangtze River Delta, China, *Stoch.*
633 *Environ. Res. Risk Assess.*, 27(8), 1899–1908, doi:10.1007/s00477-013-0725-y, 2013.
- 634 Guttman, N. B.: Comparing the palmer drought index and the standardized precipitation index,
635 *J. Am. Water Resour. Assoc.*, 34(1), 113–121, doi:10.1111/j.1752-1688.1998.tb05964.x, 1998.
- 636 Hanel, M., Rakovec, O., Markonis, Y., Máca, P., Samaniego, L., Kyselý, J. and Kumar, R.:
637 Revisiting the recent European droughts from a long-term perspective, *Sci. Rep.*, 8(1), 9499,
638 doi:10.1038/s41598-018-27464-4, 2018.
- 639 Harvey-Fishenden, A., Macdonald, N. and Bowen, J. P.: Dry weather fears of Britain’s early
640 ‘industrial’ canal network, *Reg. Environ. Chang.*, 1–13, doi:10.1007/s10113-019-01524-5,
641 2019.
- 642 He, B., Lü, A., Wu, J., Zhao, L. and Liu, M.: Drought hazard assessment and spatial
643 characteristics analysis in China, *J. Geogr. Sci.*, 21(2), 235–249, doi:10.1007/s11442-011-
644 0841-x, 2011a.
- 645 He, B., Lü, A., Wu, J., Zhao, L. and Liu, M.: Drought hazard assessment and spatial
646 characteristics analysis in China, *J. Geogr. Sci.*, 21(2), 235–249, doi:10.1007/s11442-011-



- 647 0841-x, 2011b.
- 648 He, X., Wada, Y., Wanders, N. and Sheffield, J.: Intensification of hydrological drought in
649 California by human water management, *Geophys. Res. Lett.*, 44(4), 1777–1785,
650 doi:10.1002/2016GL071665, 2017.
- 651 Heim, R. R.: Century Drought Indices Used in the United States, *Bull. Am. Meteorol. Soc.*,
652 (August), 1149–1165, 2002.
- 653 Homer-Dixon, T. F.: Environmental Scarcities and Violent Conflict: Evidence from cases, *Int.*
654 *Secur.*, 19(1)(1), 1–36, 1994.
- 655 Janku, A.: Drought and famine in northwest china: a late Victorian tragedy?, *J. Chinese Hist.*,
656 2(2), 373–391, doi:10.1017/jch.2018.4, 2018.
- 657 Keenan, S. P. and Krannich, R. S.: The Social Context of Perceived Drought Vulnerability1,
658 *Rural Sociol.*, 62(1), 69–88, doi:10.1111/j.1549-0831.1997.tb00645.x, 2010.
- 659 Kjeldsen, T. R., Macdonald, N., Lang, M., Mediero, L., Albuquerque, T., Bogdanowicz, E.,
660 Brazdil, R., Castellarin, A., David, V., Fleig, A., Gu'l, G. O., Kriauciuniene, J., Kohnova', S.,
661 Merz, B., Nicholson, O., Roald, L. A., Salinas, J. L., Sarauskiene, D., S'raj, M., Strupczewski,
662 W., Szolgay, J., Toumazis, A., Vanneuville, W., Veijalainen, N. and Wilson, D.: Documentary
663 evidence of past floods in Europe and their utility in flood frequency estimation, *J. Hydrol.*,
664 517, 963–973, doi:10.1016/j.jhydrol.2014.06.038, 2014.
- 665 Kreibich, H., Blauhut, V., Aerts, J. C. J. H., Bouwer, L. M., Van Lanen, H. A. J., Mejia, A.,
666 Mens, M. and Van Loon, A. F.: How to improve attribution of changes in drought and flood
667 impacts, *Hydrol. Sci. J.*, 64(1), 1–18, doi:10.1080/02626667.2018.1558367, 2019.
- 668 Lanen, H. A. J. Van: Drought propagation through the hydrological cycle, *Clim. Var. Chang.*
669 *Impacts*, 308, 122–127, 2006.
- 670 Lennard, A. T., Macdonald, N., Clark, S. and Hooke, J. M.: The application of a drought
671 reconstruction in water resource management, *Hydrol. Res.*, nh2015090, 2015.
- 672 Li, B. and Meng, Q.: *China Meteorological Disasters Ceremony (Liaoning volume)*, China
673 Meteorological Press., Beijing., 2005.
- 674 Li, L. M.: *Fighting famine in North China : state, market, and environmental decline, 1690s-*
675 *1990s*, Stanford University Press., 2007.
- 676 Li, Y., Strapasson, A. and Rojas, O.: Assessment of El Niño and La Niña impacts on China:
677 Enhancing the Early Warning System on Food and Agriculture, *Weather Clim. Extrem.*,
678 100208, doi:10.1016/J.WACE.2019.100208, 2019.
- 679 Liang, L., Li, L. and Liu, Q.: Precipitation variability in Northeast China from 1961 to 2008, *J.*
680 *Hydrol.*, 404(1–2), 67–76, doi:10.1016/j.jhydrol.2011.04.020, 2011.
- 681 Lloyd-Hughes, B.: The impracticality of a universal drought definition, *Theor Appl Clim.*, 117,
682 607–611, doi:10.1007/s00704-013-1025-7, 2014.
- 683 Lloyd-Hughes, B. and Saunders, M. A.: A drought climatology for Europe, *Int. J. Climatol.*,
684 22(13), 1571–1592, doi:10.1002/joc.846, 2002.
- 685 Van Loon, A. F., Stahl, K., Di Baldassarre, G., Clark, J., Rangelcroft, S., Wanders, N., Gleeson,
686 T., Van Dijk, A. I. J. M., Tallaksen, L. M., Hannaford, J., Uijlenhoet, R., Teuling, A. J., Hannah,
687 D. M., Sheffield, J., Svoboda, M., Verbeiren, B., Wagener, T. and Van Lanen, H. A. J.: Drought



- 688 in a human-modified world: Reframing drought definitions, understanding, and analysis
689 approaches, *Hydrol. Earth Syst. Sci.*, 20(9), 3631–3650, doi:10.5194/hess-20-3631-2016, 2016.
- 690 Loorbach, D., Kemp, R., Wilson, S., Bray, R., Cooper, P., Committee on Climate Change
691 Adaptation, Authorities, L., Sofoulis, Z., Schitter, G. P., Scenarios, U., Urban, E., Futures, U.
692 K., Literature, F., Directions, N., Whittle, R., Medd, W., Deeming, H., Kashefi, E., Mort, M.,
693 Twigger Ross, C., Walker, G., Waton, N., Wisner, B., Blaikie, P., Cannon, T., Davis, I., Engel,
694 K., Jokieli, D., Kraljevic, A., Geiger, M., Smith, K., Needs, S., Sector, E., Matters, W., Paton,
695 D., Committee on Climate Change- Adaptation Sub-Committee, Farmers, S., Agents, B.,
696 Officers, F., Contents, F., Tool-kit, T., Council, C., Stiglitz, J. E., Publication, R., Balmforth,
697 D., susDrain, Mccarry, T., Horth, D., Development, O., Knowledges, C. F., Barr, S., Woodley,
698 E., The Environment Food and Rural Affairs Committee, McBain, W., Wilkes, D., Retter, M.,
699 Act, W. M., Guidance, R. D., Natural, M., Risks, E., He, X., Keyaerts, N., Azevedo, I., Meeus,
700 L., Hancher, L., Glachant, J. M., Wedawatta, H., Ingirige, M., Proverbs, D., Ofwat, Stockholm
701 Resilience Centre, Cabinet Office, United Kingdom, CCWater, Rice, C., Kennedy, M.,
702 Manager, S. S., Breu, F., Guggenbichler, S., Wollmann, J., Metag, J., Fuchslin, T., Schafer, M.
703 S., Stevens, R., Ogunyoye, F., Ofwat, Lyness, N., West, S., Cook, J., Ofwat, Cobbing, P., Leach,
704 K., Bott, J., Flood, D., Risk, D., Hutchings, M., Bott, J., Project, F., March, R., et al.: At Risk:
705 natural hazards, people's vulnerability and disasters, *Challenges*, 33(July), 1–5,
706 doi:10.1080/10417946809371961, 2011.
- 707 Lu, Y. and Teng, Z.: *History of Chinese Population*, Shandong people's publishing house.,
708 2000.
- 709 Mckee, T. B., Doesken, N. J. and Kleist, J.: The relationship of drought frequency and duration
710 to time scales, *AMS 8th Conf. Appl. Climatol.*, (January), 179–184 [online] Available from:
711 [http://www.droughtmanagement.info/literature/AMS_Relationship_Drought_Frequency_Dur](http://www.droughtmanagement.info/literature/AMS_Relationship_Drought_Frequency_Duration_Time_Scales_1993.pdf)
712 [ation_Time_Scales_1993.pdf](http://www.droughtmanagement.info/literature/AMS_Relationship_Drought_Frequency_Duration_Time_Scales_1993.pdf) (Accessed 13 August 2018), 1993.
- 713 Mechler, R. and Bouwer, L. M.: Understanding trends and projections of disaster losses and
714 climate change: is vulnerability the missing link?, *Clim. Change*, 133(1), 23–35,
715 doi:10.1007/s10584-014-1141-0, 2015.
- 716 Meng, J.: *Shenyang Chronicles*, Integrated volume one., Shenyang Publishing House.,
717 Shenyang., 1989.
- 718 Mishra, A. K. and Singh, V. P.: A review of drought concepts, *J. Hydrol.*, 391(1–2), 202–216,
719 doi:10.1016/j.jhydrol.2010.07.012, 2010.
- 720 Moreira, E. E., Coelho, C. A., Paulo, A. A., Pereira, L. S. and Mexia, J. T.: SPI-based drought
721 category prediction using loglinear models, *J. Hydrol.*, 354(1–4), 116–130,
722 doi:10.1016/J.JHYDROL.2008.03.002, 2008.
- 723 Nash, D. J., Pribyl, K., Klein, J., Neukom, R., Endfield, G. H., Adamson, G. C. D. and Kniveton,
724 D. R.: Seasonal rainfall variability in southeast Africa during the nineteenth century
725 reconstructed from documentary sources, *Clim. Change*, 134(4), 605–619,
726 doi:10.1007/s10584-015-1550-8, 2016.
- 727 National Standards of People's Republic of China: Grades of meteorological drought, , GB/T
728 20481-201, 2017.
- 729 NCAR: Carbon Dioxide Information Analysis Center/Environmental Sciences Division/Oak
730 Ridge National Laboratory/U. S. Department of Energy National Laboratory/U. S. Department
731 of Energy, Two Long-Term Instrumental Climatic Data Bases of the People's Republic of ,



- 732 [online] Available from: <https://rda.ucar.edu/datasets/ds578.5/> (Accessed 6 August 2019),
733 1996.
- 734 Office of State Flood Control and Drought Relief, B. of H. and W. R. S. of L. P.: Liaoning
735 Flood and Drought Disaster, Shenyang., 1999.
- 736 Palmer, W. .: Meteorological Drought, Washington DC. [online] Available from:
737 <https://www.ncdc.noaa.gov/temp-and-precip/drought/docs/palmer.pdf> (Accessed 7 August
738 2018), 1965.
- 739 Parry, S., Hannaford, J., Lloyd-Hughes, B. and Prudhomme, C.: Multi-year droughts in Europe:
740 analysis of development and causes, *Hydrol. Res.*, 43(5), 689, doi:10.2166/nh.2012.024, 2012.
- 741 Pfister, C.: The vulnerability of past societies to climatic variation: a new focus for historical
742 climatology in the twenty-first century, *Clim. Change*, 100(1), 25–31, doi:10.1007/s10584-
743 010-9829-2, 2010.
- 744 Pu, Z.: Extraordinary drought in Liaoning Province during 2000, Liaoning Science and
745 Technology Publishing House, Shenyang., 2001.
- 746 Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P.,
747 Kent, E. C. and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night
748 marine air temperature since the late nineteenth century, *J. Geophys. Res.*, 108(D14), 4407,
749 doi:10.1029/2002JD002670, 2003.
- 750 Sangster, H., Jones, C. and Macdonald, N.: The co-evolution of historical source materials in
751 the geophysical, hydrological and meteorological sciences: Learning from the past and moving
752 forward, *Prog. Phys. Geogr.*, 42(1), 61–82, doi:10.1177/0309133317744738, 2018.
- 753 Savitzky, A. and Golay, M. J. E.: Smoothing and Differentiation of Data by Simplified Least
754 Squares Procedures., *Anal. Chem.*, 36(8), 1627–1639, doi:10.1021/ac60214a047, 1964.
- 755 Schubert, S. D., Suarez, M. J., Pegion, P. J., Koster, R. D. and Bacmeister, T.: On the Cause of
756 the 1930s Dust Bowl Published by : American Association for the Advancement of Science
757 Stable URL : <http://www.jstor.org/stable/3836515> ., *Science* (80-.), 303(5665), 1855–1859,
758 2004.
- 759 Shengjing Times: Drought, 1st July 1920, Issue 6080: p. 4, 1920.
- 760 Shenyang Municipal People’s Government Local Records Office: Shenyang chronicle (volume
761 eight), Shenyang., 1998.
- 762 Shenyang Municipal People’s Government Local Records Office (1994-2011): Shenyang
763 chronicles, Shenyang., 2011.
- 764 Stagge, J. H., Tallaksen, L. M., Gudmundsson, L., Van Loon, A. F. and Stahl, K.: Candidate
765 Distributions for Climatological Drought Indices (SPI and SPEI), *Int. J. Climatol.*, 35(13),
766 4027–4040, doi:10.1002/joc.4267, 2015.
- 767 Sun, Y.: 2014 China Drought and Drought Relief Operations, *China Flood Drought Manag.*,
768 25(1), 21–24, 2015.
- 769 Tan, X.: Historical materials of Drought Archives in the Qing Dynasty., China Book Publishing
770 House, Beijing., 2013.
- 771 Teklu, T., Braun, J. Von and Zaki, E.: IFPRI report Drought and famine relationships in Sudan :
772 Policy implications, , 14(2), 1–3, 1992.



- 773 Todd, B., Macdonald, N., Chiverrell, R. C., Caminade, C. and Hooke, J. M.: Severity, duration
774 and frequency of drought in SE England from 1697 to 2011, *Clim. Change*, 121(4), 673–687,
775 2013.
- 776 Trnka, M., Hayes, M., Jurečka, F., Bartošová, L., Anderson, M., Brázdil, R., Brown, J.,
777 Camarero, J., Cudlín, P., Dobrovolný, P., Eitzinger, J., Feng, S., Finnessey, T., Gregorič, G.,
778 Havlik, P., Hain, C., Holman, I., Johnson, D., Kersebaum, K., Ljungqvist, F., Luterbacher, J.,
779 Micale, F., Hartl-Meier, C., Možný, M., Nejedlik, P., Olesen, J., Ruiz-Ramos, M., Rötter, R.,
780 Senay, G., Vicente-Serrano, S., Svoboda, M., Susnik, A., Tadesse, T., Vizina, A., Wardlow,
781 B., Žalud, Z. and Büntgen, U.: Priority questions in multidisciplinary drought research, *Clim.*
782 *Res.*, 75(3), 241–260, doi:10.3354/cr01509, 2018.
- 783 Vicente-Serrano, S. M., Beguería, S., López-Moreno, J. I., Vicente-Serrano, S. M., Beguería,
784 S. and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming: The
785 Standardized Precipitation Evapotranspiration Index, *J. Clim.*, 23(7), 1696–1718,
786 doi:10.1175/2009JCLI2909.1, 2010.
- 787 Vicente-Serrano, S. M., Lopez-Moreno, J.-I., Beguería, S., Lorenzo-Lacruz, J., Sanchez-
788 Lorenzo, A., García-Ruiz, J. M., Azorin-Molina, C., Morán-Tejeda, E., Revuelto, J., Trigo, R.,
789 Coelho, F. and Espejo, F.: Evidence of increasing drought severity caused by temperature rise
790 in southern Europe, *Environ. Res. Lett.*, 9(4), 044001, doi:10.1088/1748-9326/9/4/044001,
791 2014.
- 792 Wang, F.: Spring Drought Report of Liaoning Province in 2001., Liaoning Science and
793 Technology Publishing House, Shenyang., 2002.
- 794 Wang, M. and Pu, W.: History of Ancient Traffic in Northeast China and Northeast Asia,
795 Liaoning People's Publishing House., 2016.
- 796 Wang, P. K., Lin, K. H. E., Liao, Y. C., Liao, H. M., Lin, Y. S., Hsu, C. T., Hsu, S. M., Wan,
797 C. W., Lee, S. Y., Fan, I. C., Tan, P. H. and Ting, T. T.: Construction of the reaches climate
798 database based on historical documents of China, *Sci. Data*, 5(1), 180288,
799 doi:10.1038/sdata.2018.288, 2018.
- 800 Wang, Y.: Causes of summer drought in Liaoning Province in 2014 and countermeasures',
801 *China Flood Drought Manag.*, 24(5), 9–11, 2014.
- 802 Wang, Y., Zhao, W., Zhang, Q. and Yao, Y.: Characteristics of drought vulnerability for maize
803 in the eastern part of Northwest China, *Sci. Rep.*, 9(1), 964, doi:10.1038/s41598-018-37362-4,
804 2019.
- 805 White, G. F.: Natural hazards, local, national, global, Oxford University Press., 1974.
- 806 Wilhelm, B., Ballesteros Cánovas, J. A., Macdonald, N., Toonen, W. H. J., Baker, V.,
807 Barriandos, M., Benito, G., Brauer, A., Corella, J. P., Denniston, R., Glaser, R., Ionita, M.,
808 Kahle, M., Liu, T., Luetscher, M., Macklin, M., Mudelsee, M., Munoz, S., Schulte, L., St.
809 George, S., Stoffel, M. and Wetter, O.: Interpreting historical, botanical, and geological
810 evidence to aid preparations for future floods, *Wiley Interdiscip. Rev. Water*, e1318,
811 doi:10.1002/wat2.1318, 2018.
- 812 Wilhite, D. A. and Glantz, M. H.: Understanding: the Drought Phenomenon: The Role of
813 Definitions, *Water Int.*, 10(3), 111–120, doi:10.1080/02508068508686328, 1985.
- 814 Wittfogel, K. A.: American Geographical Society Meteorological Records from the Divination
815 Inscriptions of, *Source Geogr. Rev.*, 30(1), 110–133 [online] Available from:



- 816 https://www.jstor.org/stable/210452?seq=1&cid=pdf-reference#references_tab_contents
817 (Accessed 6 August 2019), 1940.
- 818 Wolski, P.: How severe is Cape Town's "Day Zero" drought?, *Significance*, 15(2), 24–27,
819 doi:10.1111/j.1740-9713.2018.01127.x, 2018.
- 820 World Meteorological Organization (WMO): Standardized Precipitation Index User Guide, (M.
821 Svoboda, M. Hayes and D. Wood), WMO-No. 1090, Geneva. [online] Available from:
822 [http://www.droughtmanagement.info/literature/WMO_standardized_precipitation_index_user](http://www.droughtmanagement.info/literature/WMO_standardized_precipitation_index_user_guide_en_2012.pdf)
823 [_guide_en_2012.pdf](http://www.droughtmanagement.info/literature/WMO_standardized_precipitation_index_user_guide_en_2012.pdf) (Accessed 9 August 2019), 2012.
- 824 Yan, J. H., Liu, H. L., Hao, Z. X., Zhang, X. Z. and Zheng, J. Y.: Climate extremes revealed
825 by Chinese historical documents over the middle and lower reaches of the Yangtze river in
826 winter 1620, *Adv. Clim. Chang. Res.*, 5(3), 118–122, doi:10.1016/j.accre.2014.11.001, 2014.
- 827 Yan, W.: The historical memory of Shenyang place names all the way from ancient times,
828 *Chinese place names*, 9, 31–33, 2012.
- 829 Yang, J., Friedman, E., Guo, J. and Mosher, S.: Tombstone : the great Chinese famine, 1958-
830 1962, Farrar, Straus and Giroux. [online] Available from:
831 https://books.google.co.uk/books?id=nadqrYU10eMC&redir_esc=y (Accessed 19 July 2019),
832 2012.
- 833 Zaidman, M. ., Rees, H. . and Young, A. .: Spatio-temporal development of streamflow
834 droughts in north-west Europe, *Hydrol. Earth Syst. Sci.*, 6(4), 733–751, doi:10.5194/hess-6-
835 733-2002, 2010.
- 836 Zhang, D.: A compendium of Chinese meteorological records of the last 3,000 years, Jiangsu
837 Education Publishing House, Nanjing., 2004.
- 838 Zhang, D.: A Compendium of Chinese Meteorological Records of the Last 3000 Years (In
839 Chinese), Phoenix House Ltd., Jiangsu, China., 2013.
- 840 Zhang, J., Chen, H. and Zhang, Q.: Extreme drought in the recent two decades in northern
841 China resulting from Eurasian warming, *Clim. Dyn.*, 52(5–6), 2885–2902,
842 doi:10.1007/s00382-018-4312-2, 2019.
- 843 Zhang, L., Wu, P., Zhou, T., Xiao, C., Zhang, L., Wu, P., Zhou, T. and Xiao, C.: ENSO
844 Transition from La Niña to El Niño Drives Prolonged Spring–Summer Drought over North
845 China, *J. Clim.*, 31(9), 3509–3523, doi:10.1175/JCLI-D-17-0440.1, 2018.
- 846 Zhang, P., Cheng, H., Edwards, R. L., Chen, F., Wang, Y., Yang, X., Liu, J., Tan, M., Wang,
847 X., Liu, J., An, C., Dai, Z., Zhou, J., Zhang, D., Jia, J., Jin, L. and Johnson, K. R.: A test of
848 climate, sun, and culture relationships from an 1810-year Chinese cave record, *Science* (80-.),
849 322(5903), 940–942, doi:10.1126/science.1163965, 2008a.
- 850 Zhang, S.: China Historical Drought from 1949 to 2000, Nanjing, Hohai University Press.,
851 2008.
- 852 Zhang, S. F., Su, Y. S. and Song, D. D.: China historical drought 1949–2000, Nanjing., 2008b.
- 853 Zhao, H.: Xuan Tu Tai Shou Kao Lv., *Historiography*, 2, 89–93, 2006.
- 854 Zheng, J., Wang, W. C., Ge, Q., Man, Z. and Zhang, P.: Precipitation variability and extreme
855 events in eastern China during the past 1500 years, *Terr. Atmos. Ocean. Sci.*, 17(3), 579–592,
856 doi:10.3319/TAO.2006.17.3.579(A), 2006.



857 Zou, D.: Shenyang chronicles 1986-2005, Shenyang Publishing House., Shenyang., 2010.

858

859



860 Table 1: Historical source materials used in the drought reconstruction for Shenyang

Years	Location	Notes in material	Author/ Year	Source
23rd century BC - 1911 AD	China	The collection of various weather, climate, and atmospheric physical phenomena in history, including flood, drought, rain and snow, cold and warm weather, freezing, frost and other records. There are 7835 kinds of historical materials used in the data set, including local chronicles, historical biography, notes, inscriptions, private diaries, and court memorial files of the Qing Dynasty. Earlier weather phenomena can be converted into modern language descriptions through Oracle bones records.	(Zhang, 2004)	Meteorological Records of the Last 3,000 Years
308AD - 2000AD	Liaoning	The drought chapter of this book provides a description of the drought in Liaoning Province from 308 to 2000 AD. And from 352 to 2000 AD, there were descriptions of insect disasters, famine, epidemic diseases, and some unexplained disasters.	(Li and Meng, 2005)	China Meteorological Disasters Ceremony (Liaoning volume)
352AD - 1948AD	Liaoning	Based on historical data, drought descriptions and statistics were provided for the Liaoning area from 352 to 1948. For the 12 key cities in Liaoning Province (including Shenyang), the drought rating was listed by year. This drought level assessment was based on the reduction rate of grain yield. And a statistical table of light drought years and heavy drought years for several rivers in Liaoning area is provided.	(Office of State Flood Control and Drought Relief, 1999)	Liaoning Flood and Drought Disaster
1949 - 2000	China	It provides the annual and seasonal changes of agricultural drought, the change of disaster areas, the degree of drought risk, and the measures of drought prevention and mitigation against agriculture after 1949.	(Zhang, 2008)	China Historical Drought from 1949 to 2000
2000	Liaoning	This book provides the causes, characteristics and the degree of drought and the statistics of surface water resources in each region. The degree of drought in Liaoning Province in 2000 was respectively analysed by precipitation, river runoff, crop yield reduction and farmland drought rate, and comprehensive indicators.	(Pu, 2001)	Extraordinary drought in Liaoning Province during 2000
2001	Liaoning	Data and description of drought causes, precipitation distribution, and the multi-year comparison of the net flow of rivers are provided. The drought level is determined by the extent of agricultural disasters, meteorological factors, precipitation frequency, and water supply and demand balance.	(Wang, 2002)	Spring Drought Report of Liaoning Province in 2001
1986-2005	Shenyang	This multi-year Shenyang chronicle provided the major events that occurred in Shenyang from 1986 to 2005, including some meteorological disasters. The natural environment section records the climate, rainfall, and natural disasters during the period.	(Zou, 2010)	Shenyang chronicles 1986-2005, volume one
1994-2011	Shenyang	The annual Shenyang chronicle records the climatic conditions, meteorological disasters, and some water conservation measures of the year.	(Shenyang Municipal People's Government Local Records Office (1994-2011), 2011)	Shenyang chronicles 1994-2011 (separate volumes)
1276-1985	Shenyang	In integrate Shenyang chronicle, there are statistics on flood and drought in suburbs region, Xinmin region and Liaozhong region in Shenyang city from 1276-1985.	(Shenyang Municipal People's Government Local Records Office, 1998)	Shenyang chronicle, volume eight



1840-1987	Shenyang	The big events which happened in Shenyang from 1840 to 1987. In physical geography part, it described the seasonal climate and precipitation characters in Shenyang, and natural disasters.	(Meng, 1989)	Shenyang chronicles, Integrated volume one
1689 - 1911	China	This information comes from more than 1 million pieces of Qing dynasty memorial to the throne, including rain, floods, droughts, water conservancy projects.	(Tan, 2013)	Historical materials of drought archives in the Qing Dynasty

861



862 Table 2. Shenyang historical place names

Periods /start time	Region Name	References
Western Han Dynasty (140BC)	Gao Xian, Hou Cheng	(Yan, 2012)
Eastern Han Dynasty (108AD)	Liao Dong Jun, Xuan Tu Jun	(Lu and Teng, 2000; Yan, 2012)
Three Kingdoms Period (238AD)	Xuan Tu Ju, Gao Xian	(Yan, 2012)
Jin Dynasty (404AD)	Gai Mou Cheng	(Yan, 2012)
Tang Dynasty (670AD)	Gai Mou Zhou	(Yan, 2012)
Five Dynasties and Ten Kingdoms Period (916AD)	Shen Zhou	(Ge, 2005; Yan, 2012)
Yuan Dynasty(1296AD)	Shen Yang Lu	(Yan, 2012)
Northern Yuan Dynasty (1386AD)	Shen Yang Zhong Wei	(Yan, 2012) (Yan, 2012)
Ming Dynasty (1634AD)	Shengjing or Sheng Jing	(Yan, 2012)
Qing Dynasty (1657AD)	Feng Tian	(Yan, 2012)
Qing Dynasty (1664AD)	Cheng De Xian	(Yan, 2012)
Republic of China (1929AD)	Shen Yang Shi	(Yan, 2012)

863



864
 865
 866

Table 3. Data Information Description Table
 (Source 1: <https://rda.ucar.edu/datasets/ds578.5/docs/ndp039.des>; Source 2: NDRCC)

Source 1			Source 2			
Value	Meaning	Impacted record (%)	Value	Meaning	Treatment	Impacted record (%)
-9999	Error	0.83	32700	Microscale	Ignore	8.97
R	Total is identical to the previous or following month's total.	0.62	32744	Black	Ignore	0
H	Total is especially high for this station and is considered spurious	0	32766	Missing	Ignore	0
E	Original total was considered suspect too high for the station.	0	30xxx	Rain and snow	Keep	0.32
			31xxx	Snow	1/10	1.51
			32xxx	Fog frost	Ignore	9.25

867



868 Table 4. SPI drought classifications applied within different regions, a) China Grades of
 869 meteorological drought (National Standards of People’s Republic of China, 2017); b) the arbitrary
 870 drought intensity classes originally defined by Mckee et al. (1993); and, c) as used by the WMO
 871 (World Meteorological Organization (WMO), 2012).
 872

Grade / class	A		B		C	
	SPI value	Drought level	SPI value	Drought level	SPI value	Drought level
1	0.49 to -0.49	Normal	0 to -0.99	Mild drought	-0.99 to 0.99	Near normal
2	-0.5 to -0.99	Mild drought	-1.00 to -1.49	Moderate drought	-1.0 to -1.49	Moderately dry
3	-1.00 to -1.49	Medium drought	1.50 to -1.99	Severe drought	-1.5 < to ≤ -1.99	Severely dry
4	-1.50 to -1.99	Severe drought	≤ -2.00	Extreme drought	≤ -2.00	Extremely dry
5	≤ -2.00	Extreme drought				

873



874
 875

Table 5. Drought class and phenomenon comparison table

	Class 1: Normal	Class 2: Mild Drought	Class 3: Moderate Drought	Class 4: Severe Drought	Class 5: Extreme Drought
Meteorology	Less record or no record/ Hot weather	Less rain for several month / rain delay/ drought	No rain for several months / drought deviant, frequently or in a wide range	Heavy annual drought	Heavy drought lasting for several years
Agriculture	Soil a bit dry/ dust cover	Wheat a bit dry or slightly reduced/ soil very dry	Injury to crop field/ wheat seedling withered/ no seeding/ difficult farming/	No harvest	Long-term wide-range land dry and no harvest at all
Hydrologic		River or canal water level slightly reduced	Slight interruption of the river/ soil is not moist	Canal or land dry up	Long-term river dry up
Economic		Food price instability	Food price rise	Food price suddenly very expensive	Sell important items at a low price in exchange for food
Social Impact		Social complaints/ unrest	Displaced or loss of home/ famine/ lack of food/ people beg for food/ people living hard	Large number of displaced people/ heavy famine/ locusts as food/ death/ people snatch supplies	Corpses everywhere/ cannibalism/ selling children or women
Derived disaster		Locust disaster/ windy and haze/	Locust disaster affect traffic (people and horses)/ epidemic/ turbid red moon	Flying locust shading sky/ fire/ Plague epidemic/	Extensive epidemics

876


877



878 Table 6. The frequency of droughts in Shenyang since 1200 AD and associated drought class
 879 (see Table 5). The average drought reflects the average class achieved for each period.

Year	Average drought class	Number of droughts recorded	Class 1	Class 2	Class 3	Class 4	Class 5	Classes 1-3	Classes 4-5
1201-1300	2.5	4	0	2	2	0	0	2	0
1301-1400	2.3	3	0	2	1	0	0	2	0
1401-1500	2.6	14	0	7	6	1	0	7	1
1501-1600	2.6	17	0	9	5	3	0	9	3
1601-1700	2.5	6	0	3	3	0	0	3	0
1701-1800	2.1	7	0	6	1	0	0	6	0
1801-1900	3.1	12	0	9	3	0	2	9	2
1901-2000	2.4	74	23	16	21	9	5	60	14
2001-2015	2.9	14	2	4	3	3	2	9	5

880




881
882
883
884

Figure 1. The geographical location of Shenyang in Liaoning province and China




885
886
887



888


889 Figure 2. a) QQ plot of two precipitation (mm) data sources (p-value 0.028); b) monthly
890 precipitation comparison of two datasets (significance Analysis of Precipitation from 1961 to
891 1988); c) monthly precipitation distribution and outliers (a-l: January to December)



892

893

Figure 3. 1906-2015 Monthly and seasonal precipitation box chart




894

895 Figure 4. Annual and seasonal precipitation from 1906 to 2015, a) annual; b) spring; c) summer; d)
896 autumn; and, e) winter. A 30-year Savitzky-Golay filter is presented (bold line b-e).

897
898
899
900
901
902
903
904




905



906
907

Figure 5. Standard Precipitation Index from 1906 to 2015 a): SPI-1; b) SPI-6; and, c) SPI-12



908
 909
 910
 911
 912

Figure 6. Shenyang drought classification with instrumental (black) and documentary sources (colour/shaped) for a) 1906-present; b) augmented period (1200-2015); and, running 30-year mean drought frequency.