Changes in high-intensity precipitation on the Northern Apennines (Italy) as revealed by multidisciplinary data over the last 9000 years

5 Stefano Segadelli¹, Federico Grazzini^{2,6}, Veronica Rossi³, Margherita Aguzzi², Silvia Marvelli⁴, Marco Marchesini⁴, Alessandro Chelli⁵, Roberto Francese⁵, Maria Teresa De Nardo¹, Sandro Nanni² ¹Geological, Seismic and Soil Service, Emilia-Romagna Region Administration, Bologna, Italy.

² Regional Agency for Prevention, Environment and Energy of Emilia-Romagna, Hydro-Meteo-Climate Service (ARPAE-SIMC), Bologna, Italy.

10

³ Department of Biological, Geological, and Environmental Sciences - BiGeA, Alma Mater Studiorum University of Bologna, Bologna, Italy.

⁴Laboratory of Palynology and Archaeobotany - C.A.A. Giorgio Nicoli, San Giovanni in Persiceto (Bologna, Italy).

⁵ Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Italy.

⁶Ludwig-Maximilians-Universität, Meteorologisches Institut, München, Germany

15

Correspondence to: Federico Grazzini (fgrazzini@arpae.it)

Abstract. Several record-breaking precipitation events have stricken the mountainous area of Emilia-Romagna Region (Northern Apennines, Italy) over the last years. As a consequence, severe geomorphological processes, like debris

20 avalanches and debris flows, shallow landslides and overbank flooding affected the territory, causing severe damages to human-made structures. The unusual intensity of these phenomena prompted an investigation on their frequency in the past, beyond the instrumental time. Are these phenomena unprecedented in the region?

Peat bog and lake deposits can provide useful elements to reply to this question and to infer the frequency of extreme precipitation events occurred in the past,

- 25 Here we present the results of a dedicated field campaign performed in summer 2017 at Lake Moo in the Northern Apennines, a 0.15 km² peat bog located at an altitude of 1130 m a.s.l. During the extreme precipitation event of 13-14 September 2015, several debris flows, generated by small streams, have affected the Lake Moo plain. In such a small drainage basin (< 2km²), high-density flood can be triggered only by high-intensity precipitation events.
- The sedimentary succession (ca. 13 m-thick) was studied through the drilling of two cores and one trench. The sequence,
 characterised by clusters of coarse-grained alluvial deposits interbedded with organic-rich silty clays and peat layers, was analysed combining sedimentological, pollen, microanthracological, pedological data and radiocarbon dating (AMS ¹⁴C), in an innovative multidisciplinary approach for this area.

Original data acquired during the field campaign were also correlated with other specific paleoclimatic proxies available in the literature for the Northern Apennines area. We discover that the increase of extreme paleoflood, associated with

35 coarse-grained deposits similar to the ones observed recently, correlates well with the warm phases of the Holocene Thermal Maximum and with the ongoing warming trend observed which started at the beginning of the last century.

Keywords

extreme precipitation, Holocene flood activity, Northern Apennines, climate change, water cycle, global warming

40 **1. Introduction**

High-intensity precipitation (HIP) events, also known as torrential rainstorms for their capacity to generate flash flood on small streams, represents a significant component of the Mediterranean water cycle. More than half of the annual precipitation is often concentrated in a few major precipitation events (e.g., Frei and Schär 1998; Isotta et al., 2014). A better knowledge of their expected frequency (and maximum intensity) is crucial for planning adequate hydraulic defences and sustainable water resources management in the present and future climate.

50

45

Under the threat of the global warming, a growing number of studies have investigated the link between a temperature rise of air masses and intensity of extreme precipitation, highlighting spatial and seasonal differences across the globe (e.g., Lehmann et al., 2015; Papalexiou and Montanari, 2019). There is a consensus that HIP events are increasing with global warming, more than mean precipitation which paradoxically could instead decrease in some regions (Berg et al., 2013). Myhre et al. (2019) show that HIP precipitation over Europe almost doubles per degree of warming, due to the combined effect of increasing frequency (the major driver) and an increase in intensity.

At the local scale, different responses of weather patterns and limitations in moisture availability can alter the uniform expected raise of HIP due to the increase of saturation water vapour pressure (6-7% K⁻¹) (e.g., 55 Westra et al., 2014; Prein et al., 2017). For example, on the Swiss Alps Scherrer et al. (2016) found a significant increase in daily extreme precipitation indices since the beginning of the last century (i.e., between 1901-2014) while Brugnara and Maugeri (2019) documented regional differences in HIP trends across the whole Alpine area. These results suggest that instrumental data do not cover a suitable period to infer reliable local changes in HIP frequency.

60

The investigation of sedimentary archives, like peat bogs and lakes, allow testing the robustness of the hypothesised linkage between extreme precipitation and temperatures in the distant past, widening substantially the period where we can verify this relation.

In literature, there is ample documentation of the use of these sedimentary archives to reconstruct 65 chronologies of past flood events (Zavala et al., 2006; Zavala et al., 2011; Giguet-Covex et al., 2012; Gilli et al., 2013; Glur et al., 2013; Stoffel et al., 2013; Wirth, 2013; Wirth et al., 2013; Anselmetti et al., 2014; Longman et al., 2017; Schillereff et al., 2014; Stoffel et al., 2016; Swierczynski et al., 2017; Ahlborn et al., 2018; Wilhelm et al., 2012, 2018; Zavala and Pan, 2018), alongside others as tree rings (Ballesteros-Cánovas et al., 2015), speleothems (Regattieri et al., 2014; Zanchetta et al., 2011) and torrential fans and cones 70 (Schneuwly-Bollschweiler et al., 2013).

75

Interestingly, over the Alpine area a synchronisation between increasing flood frequency and Holocene cooling periods characterised by cold and wet summers, like the Little Ice Age-LIA, has been documented by a large number of studies (e.g., Glur et al., 2013; Henne et al., 2018). Other authors reported that under warm paleoclimate conditions floods in the Alps became rarer but stronger in intensity, highlighting the complexity of this issue (Giguet-Covex et al., 2012; Brönnimann et al., 2018a). Alpine data mostly derive from lacustrine sedimentary successions that are known to be influenced by a combination of factors (e.g.,

precipitation intensity, duration, seasonality and changes in atmospheric-circulation and land use) that might

overshadow the physically-based relation between temperature increase and extreme precipitation events (Utsumi et al., 2011; Brönnimann et al., 2018b).

80

85

90

In contrast, less attention was paid on the Holocene flood activity in the Northern Apennine area (N Italy in Fig. 1), which has a different precipitation climatology compared to the inner Alpine region. Indeed, the link between cold summers and high values of flood frequency appears more logical for the Alpine region, which shows a precipitation maximum in the Summer. In contrast, in the Northern Apennine area, precipitation maxima occur in Autumn with more than sixty per cent of the annual total precipitation concentrated in few days characterised by severe meteorological events (Isotta et al., 2014).

Focusing the analysis on the Emilia-Romagna Region (ER Region; Fig. 1), an increase in interannual variability of torrential rainfall has been observed over the last ten years with marked or even exceptional droughts, such as those occurring in 2012 and 2017 (Grazzini et al., 2012), followed by years of record-breaking rainfall (2014 and 2018). Between September 2014 and September 2015, the ER Region has been affected by three events of exceptional intensity with an estimated return period of several centuries (Grazzini et al., 2016).

To consolidate HIP trends on the N Apennine area, we analysed the Holocene sedimentary succession of a small-size peat bog (Lake Moo in Fig. 1) through the application of a multidisciplinary approach that integrates sedimentological and environmental data (e.g., pollen-derived paleovegetation patterns) with climatological observations and physical arguments.

95

100

105

110

The Lake Moo is located in the proximity of the main N Apennines crest that is very exposed to moist maritime airflow coming from the central Mediterranean Sea. The dimension of the drainage basin is small (total area $<2 \text{ km}^2$) and the anthropic impact is low as the site belongs to the Natura 2000 network (site code and name: IT4020008 - Monte Ragola, Lago Moo, Lago Bino). Moreover, the activation of debris flows (*sensu* Hungr et al., 2013) during the severe flash flood event occurred on 13^{th} - 14^{th} September 2015, with an estimated rainfall intensity of 112 mm/3h over the Lake Moo basin (Segadelli et al., 2020; Grazzini et al., 2016), offers the unique opportunity to use these recent deposits as the archetype of past events. According to literature (Milliman and Syvitski, 1992; Mulder and Syvitski, 1995; Mutti et al., 1996), in such a small drainage basin high-density floods can be triggered only by HIP events lasting enough time for the water to infiltrate and mobilise large quantities of debris. All these geomorphological features make the Lake Moo an ideal site to investigate past changes in HIP events in terms of sedimentary expression, frequency and forcing factors, and to address the following research questions:

- a) Could we consider the recent events unprecedented on a millennial timescale (i.e., the Holocene period)?
- b) Is the frequency of HIP events coupled with (paleo)temperature variations?

2. The study area

2.1. Geographic and climate context

115

The Lake Moo plain (44°37'29"N, 9°32'25"E), about 0.15 Km^2 wide, is located near the boundary between Emilia-Romagna and Liguria regions (Piacenza province, Italy; Fig. 1a) in the upper valley of the Nure stream at an altitude of 1130 m above sea level (a.s.l.). A high tree cover density characterises the present-day catchment area (i.e., the total woodland cover is 65.55%; Corticelli et al., 2011), with a high richness of plant species and an exceptional concentration of protected mountain taxa with a particular phytogeographical

interest. The vegetation landscape shows the widespread occurrence of *Fagus sylvatica*, locally interrupted by grazing areas and blueberry moorlands with the presence of rare *Juniperus nana* and *Sorbus chamaemespilus*. Reforestation of *Pinus nigra* is also documented (Fig. 1b).



125

Figure 1. (a) Location map of the Lake Moo plain. The area affected by the extreme precipitation event of September 13th and 14th 2015 is highlighted by the red rectangle. (b) View of the Lake Moo landscape from the southern side.

Present-day climate conditions primarily reflect the interaction of the prevalent atmospheric flow and the mountain range surrounding the Lake Moo plain. The proximity to the Ligurian Sea (about 40 km SW) makes the location particularly favourable to orographic precipitation enhancement in case of moisture-laden southerly-flow. The largest precipitation is usually observed in Spring and the Autumn, with a marked peak in early Autumn. In this period we observe a particular synergy of mid-latitude synoptic disturbances, becoming more frequent towards the cold seasons, and strong convective systems still developing over the warm Mediterranean Sea at the end of Summer and in Autumn (Grazzini et al., 2020); a situation very favourable the genesis of extreme precipitation events in the area.

135

2.2. Geological and geomorphological setting

- The study area mainly consists of extensively fractured serpentinites that represent the accreted fragments of the Ligure-Piemontese oceanic basin, separating the European and Adria plates during the Middle-Upper Jurassic (Marroni et al., 2010). The ultramafites are bordered by polygenic breccias, made of blocks of limestones or marly limestones embedded within a fine-grained matrix (Mt. Ragola Complex, Late Santonian Early Campanian; Elter et al., 1997).
- The geomorphological landscape includes flat areas and steep slopes located at different altitudes. On
 the former, marshy environments commonly occur recording the last filling phases of small lacustrine basins, some of which still exists as the Lake Moo and the Lake Bino (Figs. 1-2). However, the origin of the Lake Moo is still a matter of debate as some Authors point to a glacial origin (Elter et al., 1997; Marchetti and Fraccia, 1988; Carton and Panizza, 1988), while others have interpreted the basin as the expression of Holocene deep-seated gravitational slope deformations (Geological, Seismic and Soil Service of the Emilia-Romagna
 Region, 2012) mainly controlled by changes in the incision rates of the Nure stream (Elter et al., 1997). The complexity of past and present morphological processes are likely enhanced by the superposition of lithological units with strong mechanical contrast, such as ophiolites and the underlying a predominantly clayey unit (Elter et al., 1997).

155 **3. Data and methods**

160

165

3.1. Field investigation

Taking into account the criteria indicated by Gilli et al. (2013) and Schillereff et al. (2014), the Lake Moo shares several advantageous characteristics for the reconstruction of the Holocene flood activity:

- steep slopes (average inclination of 24°) composed of deposits highly susceptible to erosion (i.e., polygenic and monogenic breccias with a pelitic matrix the Mt. Ragola Complex, Elter et al., 1997; Fig. 2);
 - absence of lacustrine basins in the upstream part of the catchment;
 - small drainage basin area (1.94 km²);
- one dominant inflow into the lake;
- lack of regulated flow structures;
 - lack of natural pre-lake sediment storage zones.

The field campaign led to the acquisition of two sedimentary cores (S1 and S2), 14m and 6m-long respectively, and one trench (T1), undertaken to investigate the sedimentary succession capped by the high-density flood deposits formed during the recent HIP event (13th-14th September 2015). The location of the cores and the trench benefited from a detailed geomorphological map (Fig. 2) and a high-resolution reflection seismic survey (Fig. SUP1 in the supplementary material), both originally produced for this research. The latter provided useful information about the lake basin-floor morphology and the thickness of the infilling succession.



175 Figure 2. Geomorphological map showing the main geological units and the hydrological elements of the Lake Moo plain. The detailed mapping of flood deposits formed during the recent HIP event (13th-14th September 2015), the location of the geognostic investigations and the geophysical surveys tracks are also reported. Original data from field campaign overlaid to the Geological database of Emilia-Romagna region.

The two cores, S1 and S2, were undertaken using a continuous drilling system that guaranteed an undisturbed stratigraphy and a high recovery percentage (about 90%). A trench (6m long, 3m wide, 2m depth) was excavated between the two cores allowing a detailed analysis of the most recent deposits.

We focused our attention on the longest core (S1) that shows the complete record of the Lake Moo infilling succession, as it reaches the ophiolite bedrock. Therefore, all the laboratory analyses (grain size, radiocarbon and palynological analysis) were undertaken on core S1, while the other core (S2) and the trench were used in support to the stratigraphic reconstruction. Elevation and geographic coordinates of reference core S1 were acquired using Garmin eTrex 10 GPS receiver: 1120.2 m. a.s.l. and 44° 37' 25"N – 9° 32' 43"E, respectively.

190 *3.2. Facies analysis and chronology*

Facies characterisation of the Lake Moo sedimentary succession was mainly performed integrating the observable macroscopic physical characteristics (i.e., grain size, sedimentary structures, Munsell chart colour and types of bounding surfaces) with the grain-size data available for the reference core S1. The interpretation in terms of depositional environments also benefited from the precise position of the core(s) with respect to the spill point of the Lake Moo and the application of the facies tract concept. The latter strongly supported the interpretation of the coarse-grained intervals encountered throughout the studied succession. Specifically, a facies tract is defined as the association of genetic facies that can be observed within the flood deposits along with the down-slope motion transformation (Lowe, 1982; Mutti, 1992; Mutti et al., 1996). Thus, the application of the facies tract concept, if framed into the core(s) stratigraphy, may allow reconstructing how flood deposits changed through time at a fixed location (Figs. 3, 4).

200

195

180

The abundance of wood remains and peaty layers within the sedimentary succession under examination supported the reconstruction of a robust chronological framework. Twelve samples were selected from the reference core S1 and radiocarbon dated at the CEDAD Laboratory of the University of Salento (Italy). The data are collected within the Table TS1 (in supplementary material) and the conventional ¹⁴C ages were converted into calendar years using the OxCal software version 3.10 (Reimer et al., 2013).

205

An age-depth model was also constructed applying a linear interpolation. Coarse-grained intervals were not excluded from the processing because of the difficulty to evaluate the exact thickness of the deposits interpretable as instantaneous events. Indeed, the upper limit of coarse-grained intervals commonly appears transitional toward finer sediments and, in few cases, some intervals for a total thickness of 107 cm were not recovered

210 recovered.

3.3. Pollen and microanthracological analysis

In order to refine facies characterisation and highlight past vegetation changes at the study site, pollen analyses were carried out on 14 samples collected from the core S1. We focused the analyses on two key stratigraphic intervals, formed during periods of well-known different paleoclimate (and paleotemperature) conditions: i) Interval 1 (ca. 10.5-9 m core depth) covering the period between ca. 9600-7300 cal yr BP and, thus centred on the Holocene Thermal Maximum-HTM (Renssen et al., 2012) and ii) Interval 2 (ca. 5.5-4.5 m core depth) recording the final stages of HTM and the following cooling period between ca. 5500-3800 cal yr BP. In these two stratigraphic intervals, samples were collected from finegrained layers following a mean sampling resolution of ca. 30 cm.

220

A standard methodology already tested for pollen substrates was applied with some minor modifications (Lowe et al., 1996). The method includes a series of laboratory treatments: about 8-10 g were treated in 10% Na-pyrophosphate to deflocculate the sediment matrix; a Lycopodium spores tablet was added to calculate pollen concentration (expressed as pollen grains per gram = p/g) and the sediment residue was subsequently washed through 7 micron sieves. The sample was re-suspended in HC1 10% for remove calcareous material and subjected to Erdtman acetolysis; heavy liquid separation method was then introduced using Na-metatungstate hydrate of s.g. 2.0 and centrifugation at 2000 rpm for 20 minutes. Following this procedure, the retained fractions were treated with 40% HF for 24 h and then the sediment residue was washed previously in distilled water and after in ethanol with glycerol; the final residue was desiccated and mounted on slides by glycerol jelly and finally sealed with paraffin.

230

225

This method preserves the slides for many years after preparation and therefore it is suitable for pollen extractions from geological and paleo-archeological samples. Identification of the samples was performed at 400 magnification and only difficult pollen such as Triticum, Avena or Hordeum types were observed at 1000 magnification.

235 Determination of pollen grains was based on the Palinoteca of the "Centro Agricoltura Ambiente -CAA G. Nicoli" laboratory (Italy), atlases and a vast amount of specific morpho-palynological bibliography. Names of the families, genus and species of plants conform to the classifications of Italian Flora proposed by Pignatti (2017-2019) and European Flora (Tutin et al., 1964-1993). The pollen terminology was based on Berglund and Ralska-Jasiewiczowa (1986), Faegri and Iversen (1989) and Moore et al. (1991) with slight 240 modifications that tend to simplify plants nomenclature. The term "taxa" is used in a broad sense to indicate both the systematic categories and the pollen morphological types (Beug, 2004). For each sample, at least 500 pollen grains were counted, and the identified taxa have been expressed as percentages of the total pollen sum that includes only terrestrial pollen, no fern spores and no aquatic plants.

245 identified: Conifers (Pinus/Abies alba), Deciduous trees (this group includes quercetum taxa - Quercus, Carpinus betulus, Corylus avellana, Fraxinus, Ostrya carpinifolia, Tilia and Ulmus + other Deciduous trees), Meadow (this group mainly comprises Fabaceae and Asteroideae, Caryophyllaceae, Cichorioideae and Poaceae), Anthropic indicator (e.g. Cerealia, Chenopodium, Convolvulus arvensis, Plantago, Urtica) and Alia that includes all taxa excluded from previous groups. The group of hygro + aquatic plants was also 250 distinguished. It includes hygrophilous herbs (e.g., Cyperaceae), helophytes and hydrophytes (i.e., Lemna, Junchus, Nymphaea, Phragmites, Potamogeton and Sparganium), which are considered good proxy of humid conditions typical of wetlands.

On the basis of vegetational and ecological characteristics, the following main pollen groups were

The 14 samples prepared for pollen analysis were also investigated for the identification of microcharcoals. Microanthracological analysis has been used to track past changes in fire history, likely 255 connected to anthropogenic activities (Vescovi et al., 2010). Point count estimation of microscopic charcoal abundance was carried out, and charcoal fragments encountered during pollen counting were recorded in four, size classes based on the long axis length: 10–50, 50–125, 125–250, >250 µm (Whitlock and Millspaugh 1996; Clark 1982; Patterson et al., 1987; Whitlock and Larsen 2001, Fisinger et al., 2008). The former two classes were interpreted to be wind-blown transported, hence giving information about the regional fire events, 260 whereas the latter two were considered the result of local vegetation burning (Vittori Antisari et al., 2015).

Pollen and microcharcoal results are presented in section 5, to support facies stratigraphy and to discuss controlling factors on floods activation.

3.4. Temperature reconstruction and modern climatological dataset

265

Central to our study is the availability of a reliable (paleo)temperature dataset for the chronological period recorded by the Lake Moo sedimentary succession. In this respect, the Holocene paleoclimate reconstruction conducted by Samartin et al. (2017) on the nearby Lake Verdarolo represents a very important reference. The Lake Verdarolo site is located at 1390 m a.s.l., 270 m higher and 54 km SE from Lake Moo (Fig. 1), in a very similar climatic context. The Authors reconstructed the mean July air temperature using a 270 chironomid-based inference model, developed through a combination of data extracted from more than 200 lakes of Norway and Swiss Alps (Heiri et al., 2011). This vegetation-independent paleotemperature reconstruction is the first for the Northern Apennines and it agrees with that of the Gemini Lake (1350 m a.s.l. elevation and 48 km SE from Lake Moo) as well as several other records coming from central Italy (Samartin et al., 2017).

275 Modern temperature and precipitation time series (1961-2018) at Lago Verdarolo and Lake Moo were derived from the gridded high-resolution dataset of Emilia-Romagna (Eraclito4), described in Antolini et al. (2016). Trend estimation and Mann Kendall significance trend test were computed with the pyMannKendall package (Hussain et al., 2019).

4. Results 280

290

4.1. Holocene stratigraphy

4.1.1. Sedimentary facies

Fourteen lithofacies (LM1 to LM14) have been identified within the infilling succession of the Lake Moo basin. Detailed lithological description of each lithofacies is reported below.

285 LM1 to LM2: Clast-supported deposit with a thickness ranging between 5-15 cm. Fine boulders to coarse cobbles with abundant coarse to fine grained sand matrix, showing a low degree of sorting, occur. Polygenic clasts have low sphericity and a very angular shape. Wood fragments locally occur. The basal and top contacts are sharp; the basal one occasionally shows evidence of erosion;

LM3 to LM4: Clast-supported deposit with a thickness ranging between 5-15 cm. It is composed of fine cobbles to coarse pebbles with a very fine sand to silt matrix. Polygenic clasts have low sphericity

and a subangular shape. Scraps of wood are encountered. The basal and top contacts are sharp; the basal one occasionally shows evidence of erosion;

- LM5: Massive matrix-supported deposit composed of medium pebbles to granules embedded within
 a very fine sand to silt matrix. Polygenic clasts have low sphericity and a subangular shape. Scraps of wood are encountered. The basal and top contacts are sharp; the basal one occasionally shows evidence of erosion;
- LM6 to LM8: Generally clast-supported, poorly sorted deposit with a thickness ranging between 10-30 cm. Polygenic clasts, massive or crudely graded angular/subangular fine pebbles to very fine granules. Crudely horizontal laminae are locally recorded atop. Fragments of woods can be encountered at the base of the layers. The basal contact is sharp and occasionally shows evidence of erosion; the upper boundary is transitional;
- LM9 to LM10: Massive or crudely fining-upward graded very coarse to fine sands, showing a low degree of sorting. The total thickness ranges between 5-20 cm. Evidences of a planar lamination and scattered wood fragments occur close to the upper boundary, which is transitional. The basal contact is sharp;
- LM11: Fine sandy loam to clayey loam showing a dark grayish brown color (10YR 4/2 or 10YR 3/2).
 Polygenic fine to medium pebbles aligned with low sphericity and a very angular shape occur. Both basal and top contacts are sharp;
- LM12: Silty deposit of dark color (10YR 3/1) with a high content of decomposed organic matter.
 Occasional fine to medium pebbles aligned with low sphericity and very angular shape occasionally occur;

320

LM13: Loam to silty clay deposit showing a coarse angular blocky structure and dark olive gray color (5Y3/2), with yellowish-brown (10YR5/6) due to the presence of iron oxides. The deposit is deprived of calcium carbonate and a field pH 5.5 value is recorded;

- LM14: Massive clayey silts with a low amount of decomposed organic matter and a dark greenish gray color (5G 4\1). Scattered polygenic clasts (fine pebbles) with low sphericity and a very angular shape are recorded. Both basal and top contacts are sharp.



Figure 3. Sedimentological features and lithofacies characterisation of the reference core S1. Radiocarbon dates, palynological and grain-size samples as well as stratigraphic units (described in the section 4.2.) are also reported.

In terms of sedimentary processes, the coarse-grained LM1-LM10 lithofacies have been interpreted as flood deposits triggered by HIP events in the catchment area following plliman and Syvitski (1992), Mulder and Syvitski (1995) and Mutti et al. (1996). Particularly, the identified lithofacies were grouped according to their features (i.e., grain-size, color, sedimentary structures) and the facies tract concept (Lowe, 1982; Mutti, 1992; Mutti et al., 1996), distinguishing three main depositional setting along an idealised transect (Fig. 4): subaerial (LM1 to LM4), marginal (LM5) and subacqueous/lacustrine (LM6 to LM10).

The fine-grained facies (LM11-LM14 in Figs. 3, 4) are also gathered into two main groups. The first group includes LM11 and LM13 (paleosol) that are characterised by features indicative of subaerial conditions; the second group consists of LM12 (i.e., peaty deposits) and LM14, deposited along the marginal zone of the (paleo)lake under low-energy conditions and different degree of organic-matter enrichment.



340

345

Figure 4. Idealised genetic facies tract showing the fourteen lithofacies (LM1-LM14) identified within the reference core S1, whose stratigraphy is reported in Fig. 3.

4.1.2. Stratigraphic units

Above the ophiolite bedrock, five informal stratigraphic units have been distinguished within the S1 cored succession and described as follows, from the bottom to the top (Fig. 3). Each unit has also interpreted in terms of depositional environment.

- Unit 1 (12.25-11.30 m core depth), This unit, 95 cm-thick, is exclusively represented by the LM13 lithofacies
 that is, in turn, composed of two main horizons indicative of a mature paleosol. The lowermost horizon (Bw), recorded between 12.25-11.60 m core depth, corresponds to weathered deposits overlain by 30 cm-thick slickensides horizon (Bss). This unit is interpreted as a residual pedogenized colluvium marking the base of a structural depression produced by gravitational block sliding;
- Unit 2 (11.30-4.4 m core depth) This ca.7 m-thick unit is characterised by the presence of several, very coarse intervals (LM6-LM8) showing a thickness of about 10-30 cm and locally capped by sandy layers (LM9-LM10). Silty deposits with a variable organic-matter content, accompanied by a scarce (LM12) or moderate

(LM5) occurrence of pebbles/granules, separate the coarse-grained intervals. This unit, which is capped by a marked unconformity surface, is interpreted as a relative shallow lacustrine environment subjected to several flood events of remarkable intensity;

360

365

Unit 3 (4.4-2.3 m core depth) - This unit, ca. 2 m thick, shows sharp boundaries and the occurrence of two sub-units (3a and 3b). The lowermost sub-unit 3a, ranging between 4.4-3.2 m of depth, is exclusively represented by silty deposits with scattered pebbles/granules, several wood fragments and a mottled-like appearance corresponding to the LM5 lithofacies. These deposits are attributed to a marginal lacustrine environment subject to sporadic floods. Upwards, marginal lacustrine clayey deposits (LM14) capped the sequence (sub-unit 3b), documenting a period of apparent flood inactivity;

Unit 4 (2.3-0.9 m core depth) - This unit, 1.4 m-thick, is composed of alternating coarse (LM3-LM4) and finegrained deposits (LM12 and LM11) that record:

- the disappearance of truly lacustrine deposits replaced by subaerial ones through an erosional surface that marks the lower boundary of the Unit 4. The stratigraphic depth of this boundary is in accordance with the altitude of the spill point of Lake Moo (1116 m a.s.l.);
- the return of several, cm-thick flood deposits (LM3-LM4).

375

Unit 5 (uppermost 0.9 m) - This unit is characterised by the presence of several layers of very coarse sediments (LM1-LM2) separated by finer deposits (LM11). An erosional contact separates Unit 4 from Unit 5, which includes the flood deposits produced by the rainfall event occurred on September 13th and 14th 2015 (Figs. 3, 5).

380

As a whole, we interpret the lacustrine succession as an infill of a structural depression produced by gravitational block sliding that was induced by post-glacial fluvial incision (Elter et al., 1997).



Figure 5. Representative photograph of the trench (see Fig. 2 for location), showing the uppermost 50 cm-thick sedimentary succession. Though the exposure is quite small, the graded pebble-sand couplets belonging to the Unit 5 can be interpreted as a sheet-flood deposit.

4.2. Age-depth model

390

The age-depth model, based on eight out of twelve ¹⁴C dates available for the S1 core (Table TS1 in the supplementary material), indicates that the sedimentary succession above the bedrock encompasses almost all the Holocene period (about the last 10 kyrs; Fig. 6). Three ages (at ca. 9.60 m, 3.10 m and 2.40 m core depth) were excluded because resulting younger than 1950 AD, while one sample collected at 2.35 m core depth resulted not datable due to a very low organic matter content. The selected calibrated ages are stratigraphically coherent and mainly derived from wood fragments and peaty deposits. In order to reduce the potential bias due to long-lived plants and reworking processes, small twigs were selected (Oswald et al., 2005) 395 from fine-grained deposits and identified when possible (e.g., Pinus and Abies alba). Due to the small dimension of the basin and the ophiolites prevalence in the drainage area (Fig. 2), we consider negligible the hard-water effect on peats.

400

The resulting age-depth model suggests the occurrence of four main stratigraphic intervals in terms of sedimentation rates. The lowermost interval (about 9.5-7.3 cal kyr BP) denotes a mean accumulation rate of ca. 0.50 mm/yr that strongly increases up to ca. 2.38 mm/yr within the overlying section dated around 7.3-5.5 cal kyr B.P. An important change in the accumulation rate occurs around at 5 m of the core depth with a drop to ca. 0.3 mm/yr between the 5.5-4.0 cal kyr BP interval. Above, between ca. 4.5-3 m, the deposit shows an uncertain chronology due to the lack of reliable radiocarbon dates and the occurrence of an unconformity surface around 4.5 m, at the contact between Units 2-3 (Figs. 3, 6). During recent times (the last 146-14 cal yr BP), the accumulation rate strongly increases, reaching the remarkable value of ca. 12 mm/yr.





Figure 6. S1 core stratigraphy and the age-depth model obtained from radiocarbon dates. Red and orange lines represent the two-sigma probability envelope.

5. Discussion

As a basis for discussion we assemble, and temporally synchronise, the Lake Moo data with the most relevant paleoclimate proxy available from literature, focusing on the N Apennines paleoarchives (Figs. 1, 7). As a whole, the Holocene infilling succession of the lacustrine basin (i.e., Units 2-3) is composed of an 415 alternation of cm to dm-thick, coarse-grained flood deposits and silts characterised by a variable amount of organic matter (sub-sections 4.1. and 4.2.; Figs. 3, 4, 7). The coarse intervals are particularly well-developed and frequent within the portion of the stratigraphic Unit 2 chronologically correspondent to the HTM (Holocene Thermal Optimum), constrained between ca. 9-5 kyr cal BP in the nearby Verdarolo Lake (Samartin et al., 2017 in Fig. 7A; section 3.4.). Interestingly, stable oxygen isotope records from the nearby Apuan Alps 420 (Corchia caves; Fig. 1) document, within the same time interval, the wettest conditions interpreted as precipitation increases in the Northern Apennine (I and II humid periods in Fig. 7B) with a peak between ca. 8500 and 7500 cal yr BP (I humid period; Regattieri et al., 2014; Zanchetta et al., 2011). This peak is in phase with a clastic layer, dated back to 8.2 and 7.1 cal kyr BP, indicative of a period of enhanced cave flooding triggered by high-magnitude precipitation events (Zhornyak et al., 2011). The Authors attribute this maximum 425 fluvial activity to an increase of strong convective episodes, like the one that affected the Versilia region in 1996 (Cacciamani et al., 2000 for a description of the Versilia flood event).

M reals simulations on Northern Apennine, are showing higher precipitation in Autumn in response to the intense Summer heat of the HTM period, (e.g., Tinner et al., 2013). We speculate that the precipitation increase, reflected also in high frequency of flood activity, could have occurred during a phase of a progressive 430 reduction of the Hadley cell, paralleled by a reduction of the subtropical high-pressure after the insolation peak (Fig. 7C). This would lead to an enhanced meteorological activity over the Mediterranean area, due to the midlatitude weather systems, especially at the transition towards the cold season. Although in decline, the presence of African Monsoon (Skinner and Poulsen, 2016) could also have contributed to enrich sub-tropical air masses in the water vapour potentially extractable by mid-latitude synoptic disturbances. Accordingly, Krichak et al. 435 (2015) documented how this mechanism is relevant for Mediterranean extreme precipitations nowadays; moreover, the events occurring in the autumn seasons are characterised by a greater transport of water vapour

from the subtropical Atlantic even across North Africa.

440

With the end of the HTM, a drastic decrease in flood activity is documented at the Lake Moo, as the coarse intervals abruptly reduce in number and thickness within the uppermost portion of Unit 2 dated to ca. 5.5-3.8 kyr BP. This stratigraphic trend is reasonably interpretable as the expression of a decrease in frequency of HIP events over the study area under cooler and less humid climate conditions (Fig. 7). The limit between Unit 2 and Unit 3 corresponds to an unconformity that marks the passage to a marginal lacustrine succession of uncertain age, at least for the lowermost portion (sub-unit 3a). This facies interpretation is supported by palynological data, which show remarkable values of herbaceous hygrophytes (mainly Cyperaceae) and 445 aquatics typical of wetlands.

The upper portion of Unit 3 (i.e., sub-unit 3b), dated around 146-14 cal yr BP, is almost deprived of coarse materials documenting a period of apparent floods inactivity at the Lake Moo (Figs. 3, 7). This is followed by a reactivation of flood processes during recent times (i.e., coarse-grained flood deposits within

Units 4-5; Figs. 3, 7) with a minimum of five events to a maximum of twelve events every 100 years, as the confidence range associated with the calibrated age at 2.80 m core depth is rather wide (146-14 cal yr BP; Fig. 7). Despite the low degree of precision affecting radiocarbon ages younger than 200 years, this renewed increase in flood activity fits well with the instrumental record that points to a high frequency of HIP events during the last decades over the Northern Appennine (Libertino et al., 2019). These phenomena seem to be also responsible for the replacement of lake deposits by subaerial ones through an erosional contact (Unit 4 lower boundary).





Figure 7. Stratigraphy of core S1, main palynological features and microcharcoal content. Relative abundances of pollen groups, explained in text (sub-section 3.3), are reported along with the frequencies of hygrophilous herbs and aquatics. Asterisks point to samples containing coprophilous fungi and other spores like *Sporormiella*, *Dicrocoelium* and *Ascaris*. These data are compared with the most relevant paleoclimate reconstructions available from literature for the area of interest. Panel (A) Reconstructed mean July air temperature from Lago Verdarolo (from Samartin et al., 2017); Panel (B) δ^{18} O speleothem records and reconstructed precipitation trends (mean anomaly time series) from Corchia cave (slightly modified from Combourieu et al., 2013; Zanchetta et al., 2011; Zhornyak et al., 2011) and Panel (C) June (red) - December (blue) insolation values reported for 45° Nord (from Samartin et al., 2017). Main Holocene climate phases and events are also shown (Regattieri et a., 2014). Available radiocarbon ages are reported as calibrated years BP.

In order to compare the most recent stratigraphic units (Units 4-5) with quantitative climate data, the instrumental temperature values, available since the second half of the last century from the Eraclito ER dataset, has been added to the Verdarolo curve. The overlap with the latest part of Verdarolo curve confirms the good accuracy of the reconstruction technique in this region (Fig. 8). The recent temperature increase is striking if compared with the previous period, and the current July temperature is comparable with the maximum temperature reached at HTM over the study area. The actual trend of July temperature, estimated over the period 1961-2018 from monthly mean values, is +4.3°C in 100 yea the trend is highly statistically significant, with a p-value=0.0004 (see section 3.4.), and more than double of the maximum temperature gradient found in the Verdarolo curve, +2°C/100 years, at the end of the Little Ice Age 1850-1900. From Fig.

475 6 and 7 we observe a flood intensification corresponding to periods of rising temperatures. This is testified by the exceptionally high accumulation rate estimated on the recent period for the Units 4-5 (Fig. 6; section 4.3.), a rate of about 1 m/100 years calculated since ca. 1800 AD. The recent increased in flood activity is also supported by instrumental data from Fig. 9, where we also observe an emerging trend (not yet significant though) of precipitation intensity increase (both as daily maximum and as monthly cumulative values) in the 480 autumn months.



485

Figure 8. Comparison of current data and reconstructed July mean temperature at the Lake Verdarolo site (see Fig. 1 for location). The blue line is the reconstructed temperature and the shaded area is the sample-specific estimated standard error associated with the temperature reconstruction based on chironomid assemblage from Samartin et al. (2017). The orange line represents the July mean temperature (1961-2018) retrieved for the grid cell of Lago Verdarolo from Emilia-Romagna climate reanalysis ERACLITO (11 years running average). The shaded orange area is +/- one standard deviation. The full available period is reported on the left, while a zoom on the most recent period (since 1600 AD) is shown on the right.



Figure 9. Monthly mean, 2m temperature (a), accumulated precipitation (b), maximum daily precipitation (c), for 490 1961-1990 (blue) and 1991-2018 (green) computed over Lake Moo site grid cell of Eraclito reanalysis (grid cell 258, Ferriere Municipality). Confidence intervals at 95% significance (black vertical segments) are computed with a bootstrapping method (1000 iterations) as part of the Seaborn python library.

5.1. Holocene flood activity, paleotemperature changes and HIP events

495

A positive link emerges correlating Holocene (July) paleotemperatures, precipitation intensity reconstructions and flood history (Fig. 7) at the study site, as the stratigraphic units showing the highest frequency of coarse-grained flood deposits (i.e., Units 2, 4 and 5) invariably fall into periods of higher temperatures (i.e., HTM and the post-LIA time). This strongly supports the hypothesis that a greater warmth favours the occurrence of extreme precipitation events, probably more frequent at the end of the Summer/Autumn as already reported by other authors (e.g., Marcott et al., 2013; Giguet-Covex et al., 2012). 500 Although the process attribution and the geographical uniformity of the HTM wet phase are still matter of debate, the abundance of coarse-grained flood deposits within the Lake Moo record around 9.5-5.5 kyr BP suggests a precipitation increase also identified in other sites of the central and southern Mediterranean (Magny et al., 2012b).

However, local factors as changes in vegetation cover (i.e., tree cover percentage) due to human 505 disturbance and fires could have significantly contributed to enhancing slope erosion and then influenced the Lake Moo sedimentary record, partly weakening our reconstruction of the temperature-HIP-flood activity relationship at the Holocene scale. In order to overcome this issue, we explored and compared the pollen and microcharcoal content of two key intervals (I_1 and I_2 in Figs. 7, 10) that are characterised by a different flood record (high versus low flood activity) formed during consecutive periods of distinct climate conditions: the 510 HTM and the following cooling. The difference in mean temperature between I_1 and I_2 , obtained averaging the corresponding chronological intervals in the Verdarolo paleotemperature curve, is +1.3°C while the maximum difference reaches +3.1°C (Fig.8).

For what concerns the post-LIA interval, and in particular since the 1950 AD, no significant contribution on debris mobilisation can be attributed to vegetation landscape changes along the slopes as the 515 Lake Moo is a protected area characterised by dense forests (Corticelli et al., 2011; section 2.1.). Consistently, the most recent pollen sample (P14 yielding an age of 146-14 cal yr BP; Fig. 7) shows a tree cover of about 80%.

The main considerations arising from the in-depth analysis of I_1 and I_2 are the following:

 I_1 (ca. 9.5-7 kyr BP; 10.5-9 m core depth): the woody component (i.e., tree cover %) is invariably 520 _ higher than the herbaceous one and it shows the remarkable value of 80% on average. The Conifers, mainly represented by pines and silver fir, are dominant with respect to the deciduous trees (e.g., Corylus avellana and Quercus), reaching a maximum of 95% within the P4 sample dated between 9.3-9.1 cal kyr BP. Interestingly, Abies alba shows the highest percentages (ca. 15-55%) within the 525 chronological interval that corresponds to the warmest period of the HTM (Figs. 7, 10; Samartin et al., 2017). Moreover, A. alba reaches the peak of ca. 55% during the HTM wettest conditions (around 7.5-7.3 cal kyr B.P.; I₁ humid period in Fig. 7), as reconstructed by several Authors (Regatieri et al., 2014; Zanchetta et al., 2011; Zhornyak et al., 2011) from the Apuan Alps speleothems. This trend is consistent with the fact that A. alba is a warm-temperature tree with a preference for high moisture availability (Tinner et al., 2013), as clearly documented within the nearby Lake Greppo record (Vescovi et al., 2010, confirming higher rainfall accumulation already highlighted by lake-level dynamics from Jura Mountains and the central-northern Italy (Magny et al., 2004, 2009, 2012). Although no specific inferences can be made on past water levels of the Lake Moo, it is interesting to note that the low amount of hygrophytes and aquatics (Fig. 7) is fully consistent with highstand conditions. Anthropic indicators taxa (Fig. 7), as well as the Anthropic Index (AI; Fig. 10), show very low values throughout the interval, documenting a negligible human impact on the area in accordance with the extremely low content in microcharcoals (0.01-0.13 mm²/g; Fig. 10). Besides, the coarser microcharcoals (>125 micron) are not recorded suggesting no significant local fires. A slightly higher amount of anthropic indicators taxa occurs within the sample P6, dated between 8070-7880 ca yr BP; this is interpreted as an effect of categorisation difficulties about spontaneous taxa as *Artemisia vulgaris* type, *Chenopodium, Plantago, Urtica*, etc.

Palynological data, combined with the stratigraphic interpretation of the S1 core, support our hypothesis that warmer and wetter conditions typical of the HTM led to a high component of HIP and then high flood activity in the basin, independently by local factors. Actually, the presence of an extensive vegetation cover along the slopes implies the need of even more intense rainfall to trigger a debris mobilisation that leads to the deposition of thick, coarse-grained layers containing pebble of remarkable diameters (LM6, LM7 and LM8 in Fig. 10).

I₂ (ca. 5.5-3.8 kyr BP.; 5.5-4.5 m core depth): tree cover reduction (ranging between ca. 87.4-45.1 %; 550 Figs. 7, 9), in favour of grassland expansion, characterises this interval that sees a fall in A. alba percentages (commonly less than 10%) and a remarkable amount of meadow taxa (mainly *Poaceae*, *Cichorioideae* and *Asteroideae*; Fig. 7). The growth in hygrophytes and aquatics (up to ca. 18.2%) is consistent with a marginal lacustrine environment and then a relative lowering of the water level. Both ant ropic indicators taxa and AI increase, up to ca. 7.3% and 16.2% respectively (Figs. 7, 9), 555 suggesting a moderate degree of human impact on the vegetation cover (i.e., tree cover changes). This impact is reasonably due to pastoral activities more than agriculture ones (few grains of cereals, less than 4%, are locally encountered), as documented by the abundance of meadow (pasture) taxa and the sporadic occurrence of coprophilous fungi and other spores (Sporormiella, Dicrocoelium and Ascaris; Figs. 7, 9). By contrast, microcharcoals remain very low in amount, pointing to a low fire activity. 560 Although pollen-derived vegetation features (i.e., decrease in tree cover and a moderate degree of human impact) document local conditions more favorable for the debris flows activation with respect to those characterising I₁, this interval shows a low past flood activity with few and thin (<5 cm) sandy deposits (LM9 to LM10 in Fig. 10). Integrating the I_2 stratigraphic record with pollen data, such a low

565

humid period (Figs. 7, 9).

545

flood activity reasonably reflects low frequency (and intensity) of HIP events during a cooler and less

The paleoenvironmental-paleovegetation features characterising the Interval 2 persists upcore, in correspondence of the Unit 3 that records apparent flood inactivity (Figs. 3, 7). The uppermost sample P14 likely tracks a strong increment of *A. alba*.



Figure 10. Sedimentological and palynological features of two key stratigraphic intervals (Intervals 1 and 2) from the reference core S1. These features are compared with the most relevant paleoclimate data available from literature for the area of interest (for references, please see Figure 7). The Anthropic Index (AI) has been calculated as: (Anthropic indicators/tree percentage)*100 following Accorsi et al. (1999). Asterisks point to samples containing coprophilous fungi and other spores like *Sporormiella*, *Dicrocoelium* and *Ascaris*. Core S1 sample ages derive from the age-depth model reported in Figure 6. P samples correspond to the pollen samples; LM labels refer to the lithofacies explained in text (section 4.1.) and in Figures 3-4; dmax corresponds to the maximum diameter.

580 6. Conclusions

575

Placing present climate conditions into a geological perspective time, beyond the instrumental era, is essential for understanding changes in the hydrological cycle induced by anthropogenic climate warming, complementing projections of model simulations. Physical reasons and regional climate reconstructions are consistently pointing to an increase in precipitation intensity when water vapour is not limited. In the last decade, record-breaking rainfall events have frequently occurred around the world. This trend is emerging with variable strength in different areas. The short timescale of the instrumental series doesn't allow yet to draw firm reconstructions. Therefore, in the attempt to consolidate confidence about the extreme precipitation trend, we extended our an bis, passing from the instrumental to the geological time scale (i.e., the Holocene period). This choice implies a robust multidisciplinary approach, which includes, in addition to climate and meteorological data, proxy and competencies coming from geological, geomorphological, stratigraphic and palynological-microanthracological area. We acquired original data during a field campaign performed in Summer 2017.

The major outcomes of our work can be summarized as follows:

The matching of instrumental temperature data and paleoenvironmental data allow us to affirm that
 current summer temperature is comparable with the temperature recorded in the Northern Apennine during the
 HTM. The current temperature trend computed from the monthly values of July temperature at Lake Verdarolo
 site over the period 1961-2018 is +4.3°C/100y (highly statistically significant), and it is likely to be one the
 highest recorded in the Holocene;

- 600 2) The stratigraphic units showing a high frequency of coarse-grained flood deposits are Units 2, deposited in the HTM, and Units 4 and 5 which belongs to the post-LIA time. Both periods are characterized by higher temperatures. The human impact is almost absent in the HTM and very low in the post-LIA time, especially in the last part of the 20th century. This strongly supports the hypothesis that a greater warmth favours the occurrence of extreme precipitation events. In particular, the recent sedimentation rate computed 605 since c.a. 1800 AD (post-LIA) is at least 1 m/100yr. This high value of the sedimentation rate has to be correlated to the presence of numerous coarse-grained levels within Units 4 and 5, after a period of absence of floods deposits in unit 3b. This difference in flood layers must be linked to an increase of HIP over the Lake Moo basin, since we could not attribute it to any other changes in physiographic and vegetation dynamics (i.e., tree cover). During HTM and current time, the forested area coverage around Lake Moo persisted on high 610 values and this factor is considered disadvantageous for debris mobilisation. On the contrary, during about 6-4 kyr BP we did not find any increase of coarse-grained layers from flood events, even though the forested area declined significantly and the anthropic impact increased;
- 3) HIP increase in response to higher temperature is already detectable in observation series. We found
 evidence that this also occurred in the past, especially during HTM testified by higher deposition of coarsegrained levels. A comparison with the past helps to understand future projections on the area. However, we are aware that past evolution cannot be taken as a perfect analogy for the future due to the different forcing and consequent response of the climate system (D'Agostino et al., 2019). As temperature will continue to increase in the Mediterranean area, precipitation intensity would keep increasing over the Northern Apennines.
 620 We hypothesise that precipitation intensity increase will be more evident in months with cooler and moist air masses, like in Autumn and in Winter, when moisture availability is not limited. An increase in precipitation maxima in Autumn months is already emerging on the Northern-Apennine.

Finally, we would like to remark that Lake Moo basin represents an ideal study area to achieve a reconstruction between high-intensity precipitation and debris flow, for its position to the dominant atmospheric flow and the favourable and undisturbed geological, geomorphological and vegetation characteristics. We think this site deserves further investigations and, in association with other analogous sites along the Northern Apennine crest, could be used in the future to provide a multisite assessment on the dynamics of past extreme precipitation events in relation with temperature variations.

Data availability: Original data concerning pollen and radiometric data are available on the open data repository of ARPAE Emilia-Romagna (https://arpaeprv.datamb.it/dataset/lake-moo)

- Author contribution: SS and FG developed the idea and the research activity planning and execution. SN,
 MA, MTDN have taken care of the administrative aspects. All authors, except for SN and MTDN, have contributed to field activity. Granulometric analysis has been conducted by AC. Pollen analysis have been conducted by SM and MM, data elaboration and figures were performed by VR, SM and MM. SS and FG and VR prepared the manuscript with contributions from all co-authors. FG management and coordination.
- 640 **Competing interests**: The authors declare that they have no conflict of interest.

Acknowledgements

This research was funded by Regional Agency of Civil Protection of the Emilia-Romagna Region and Geological, Seismic and Soil Survey of Emilia-Romagna Region in the framework of cooperation agreement with ARPAE-SIMC. We are very grateful for all comments received in the public discussion and in particular, we thank reviewers Picotti and Tinner for their useful suggestions.

References

645

- 650 Accorsi, C.A., Bandini Mazzanti, M., Forlani, L., Mercuri, A.M., Trevisan Grandi, G.: An overview of Holocene forest pollen flora/vegetation of the Emilia Romagna region - Northern Italy in Archivio Geobotanico, 5 (1-2), 3-27, 1999.
 - Ahlborn, M., Armon, M., Ben Dor, Y., Neugebauer, I., Schwab, M.J., Tjallingii, R., Shoqeir, J.H., Morin, E., Enzel, Y., and Brauer, A.: Increased frequency of torrential rain storms during a regional late Holocene eastern Mediterranean drought, Quaternary Research, 89, 425-431, https://doi.org/10.1017/qua.2018.9, 2018.
- 655 Anselmetti, F., Wirth, S.B., Glur, L., and Gilli, A.: Holocene flood frequency as reconstructed by lake sediments from multiple archives: A record influenced by solar forcing and atmospheric circulation patterns, in: Late Pleistocene and Holocene climatic variability in the Carpathian-Balkan region, Abstracts Volume, edited by: Mindrescu, M., and Gradinaru, I., Special Issue, 24, 1-2, 2014.
- Antolini, G., Auteri, L., Pavan, V., Tomei, F., Tomozeiu, R. and Marletto, V.: A daily high-resolution gridded climatic
 data set for Emilia-Romagna, Italy, during 1961–2010. Int. J. Climatol., 36: 1970-1986, doi:10.1002/joc.4473, 2016.
 - Ballesteros-Cánovas, J.A., Stoffel, M., St. George, S., and Hirschboeck, K.: A review of flood records from tree rings, Progress in Physical Geography, 1-23, doi 10.1177/0309133315608758, 2015.
 - Berg, P., Moseley, C., and Haerter, J.O.: Strong increase in convective precipitation in response to higher temperatures. Nature Geosci. 6, 181–185, https://doi.org/10.1038/ngeo1731, 2013.
- 665 Berglund, B.E., and Ralska-Jasiewiczowa, M.: Pollen analysis and pollen diagrams, in Handbook of Holocene Palaeoecology and Palaeohydrology, edited by: B.E. Berglund, Chichester, 455 - 484, 1986.

Beug, H.J.: Leifaden der Pollenbestimmungen für Mitteleuropa und angrenzende Gebiete, Pfeil, München, 2004.

Brönnimann, S., Rohr, C., Stucki, P., Summermatter, S., Bandhauer, M., Barton, Y., Fischer, A., Froidevaux, P., Germann, U., Grosjean, M., Hupfer, F., Ingold, K., Isotta, F., Keiler, M., Martius, O., Messmer, M., Mülchi, R.,

- Panziera, L., Pfister, L., Raible, C.C., Reist, T., Rössler, O., Röthlisberger, V., Scherrer, S., Weingartner, R., Zappa,
 M., Zimmermann, M. Zischg, A.P.: 1868 L'alluvione che cambiò la Svizzera: Cause, conseguenze e insegnamenti per il futuro. Geographica Bernensia, G94, 52 S., doi:10.4480/GB2018.G94.02, 2018a.
- Brönnimann, S., Rajczak, J., Fischer, E. M., Raible, C. C., Rohrer, M., and Schär, C.: Changing seasonality of moderate and extreme precipitation events in the Alps. Natural Hazards and Earth System Sciences, 18 (7), 2047-2056.
 https://doi.org/10.5194/nhess-18-2047-2018, 2018b.
 - Brugnara, Y., and Maugeri, M.: Daily precipitation variability in the southern Alps since the late 19th century. Int J Climatol.; 39: 3492–3504. https://doi.org/10.1002/joc.6034, 2019
 - Cacciamani, C., Cesari, D., Grazzini, F, Paccagnella, T., and Pantone, M.: Numerical Simulation of Intense Precipitation Events South of the Alps: Sensitivity to Initial Conditions and Horizontal Resolution. Meteorol Atmos Phys 72, 147– 159. https://doi.org/10.1007/s007030050012, 2000.
 - Carton, A. and Panizza, M. (Eds.): Il paesaggio fisico dell'Alto Appennino Emiliano, Grafis Edizioni, Bologna, Italia, 1988.

- 685 Combourieu-Nebout, N., Peyron, O., Bout-Roumazeilles, V., Goring, S., Dormoy, I., Joannin, S., Sadori, L., Siani, G., and Magny, M.: Holocene vegetation and climate changes in the central Mediterranean inferred from a high-resolution marine pollen record (Adriatic Sea), Clim. Past, 9, 2023-2042. https://doi.org/10.5194/cp-9-2023-2013, 2013.
 - Corticelli, S., Garberi, M.C., Mariani, M.C., and Masi, S.: Uso del suolo 2008, http://geoportale.regione.emiliaromagna.it/it/download/dati-e-prodotti-cartografici-preconfezionati/pianificazione-e-catasto/uso-del-suolo-1/2008coperture-vettoriali-uso-del-suolo-edizione-2011, 2011.
 - D'Agostino, R., Bader, J., Bordoni, S., Ferreira, D., and Jungclaus, J.: Northern Hemisphere monsoon response to mid-Holocene orbital forcing and greenhouse gas-induced global warming. Geophysical Research Letters, 46, 1591–1601. https://doi.org/10.1029/2018GL081589, 2019.
- Elter, P., Ghiselli, F., Marroni, M., and Ottria, G. (Eds.): Note illustrative del Foglio 197 "Bobbio" della Carta Geologica
 d'Italia alla scala 1:50.000. Istituto Poligrafico e Zecca dello Stato, Roma, Italia, 1997.
- Faegri, K., and Iversen, J.: Textbook of Pollen analysis, K. Faegri, P.E. Kaland, K. Krzywinski (Eds.), Chichester, 1989.Finsinger, W., Tinner, W., and Hu, F.S.: Rapid and accurate estimates of microchacoal in pollen slide. Fiorentino, G., Magri D. (Eds.), Charcoals from the Past, BAR int Ser., pp. 121-124, 2008.
- Frei, C. and Schär, C.: A precipitation climatology of the Alps from high-resolution rain-gauge observations. Int. J.
 Climatol., 18: 873-900. doi:10.1002/(SICI)1097-0088(19980630)18:8<873::AID-JOC255>3.0.CO; 2-9, 1998.
 - Geological, Seismic and Soil Service of the Emilia-Romagna Region: Landslide characteristics in Emilia-Romagna, 2012 (data available at: https://ambiente.regione.emilia-romagna.it/en/geologia/temi/dissesto-idrogeologico/lecaratteristiche-dei-fenomeni-franosi-in-emilia-romagna, last access: 04 April 2020).
- Giguet-Covex, C., Arnaud, F., Dirk, E., Jérôme, P., Laurent, M., Pierre, F., Fernand, D., Pierre-Jérôme, R., Bruno, W.,
 and Jean-Jacques, D.: Frequency and intensity of high-altitude floods over the last 3.5ka in northwestern French Alps

(Lake Anterne), Quaternary Research, 77, 12-22, https://doi.org/10.1016/j.yqres.2011.11.003, 2012.

Gilli, A., Anselmetti, F.S., Glur, L., and Wirth S.B.: Lake Sediments as Archives of Recurrence Rates and Intensities of Past Flood Events, in: Dating Torrential Processes on Fans and Cones, Advances in Global Change Research, edited by: Schneuwly-Bollschweiler, M., Stoffel, M., and Rudolf-Miklau, F., Springer Science Business Media, Dordrecht,

710 Holland, 225-242, doi 10.1007/978-94-007-4336-6_15, 2013.

680

Clark, R.L.: Point count estimation of charcoal in pollen preparations and thin sections of sediments, pollen et spores, 24, 523-53, ISSN 0375-9636, 1982.

- Glur, L., Wirth, S.B., Büntgen, U., Gilli, A., Haug, G.H., Schär, C., Beer, J., and Anselmetti, F.S.: Frequent floods in the European Alps coincide with cooler periods of the past 2500 years, Nature Scientific Reports, 3, doi 10.1038/srep02770, 2013.
- Grazzini, F., Craig, J.G., Keil, C., Antolini, G., and Pavan V.: Extreme precipitation events over Northern Italy: Part (I)
- 715 a systematic classification with machine learning techniques, QJR Meteorol Soc. 2020;1–17. https://doi.org/10.1002/qj.3635, 2020.
 - Grazzini, F., Segadelli, S., and Fornasiero, A.: Precipitazioni estreme ed effetti al suolo sul reticolo minore: il caso del 14 Settembre 2015. Technical report ARPAE-SIMC/SGSS Emilia-Romagna, 27pp, Available in Italian https://www.arpae.it/dettaglio_notizia.asp?id=8017&idlivello=32, 2016.
- 720 Grazzini, F., Pratizzoli, W. and Tomei, F.: Drought and big snow, are extreme events the new normal? Ecoscienza, 1, Available in Italian https://www.arpae.it/dettaglio_documento.asp?id=3674&idlivello=1171, 2012.
 - Heiri, O., S.J., Brooks, H.J.B., Birks, and Lotter, A.F.: A 274-lake calibration data-set and interference model for chironomid-based summer air temperature reconstruction in Europe, Quaternary Science Reviews 30, 3445-3456, 2011.
- 725 Henne, P.D., Bigalke, M., Büntgen, U., Colombaroli, D., Conedera, M., Feller, U., Frank, D., Fuhrer, J., Grosjean, M., Heiri, O., Luterbacher, J., Mestrot, A., Rigling, A., Rössler, O., Rohr, C., Rutishauser, T., Schwikowski, M., Stampfli, A., Szidat, S., Theurillat, J.P., Weingartner, R., Wilcke, W., and Tinner, W.: An empirical perspective for understanding climate change impacts in Switzerland. Reg Environ Change 18, 205–221, https://doi.org/10.1007/s10113-017-1182-9, 2018.
- 730 Hussain, M., and Mahmud, I.: pyMannKendall: a python package for non parametric Mann Kendall family of trend tests, Journal of Open Source Software, 4, (39), 1556, https://doi.org/10.21105/joss.01556, 2019.
 - Hungr, O.,Leroueil, S., and Picarelli, L.: The Varnes classification of landslide types, an update. Landslides, 11, 167–194, https://doi.org/10.1007/s10346-013-0436-y, 2014.
 - Isotta, F., Frei, C., Weilguni, V., Percec Tadic, M., Lassègues, P., Rudolf, B., Pavan, V., Cacciamani, C., Antolini, G.,
- Ratto, S.M., Munari, M., Micheletti, S., Bonati, V., Lussana, C., Ronchi, C., Panettieri, E., Marigo, G., and Vartacnik,
 G.: The climate of daily precipitation in the Alps: development and analysis of a high-resolution grid dataset from pan-Alpine rain-gauge data, International Journal of Climatology, 34, 1657-1675, doi 10.1002/joc.3794, 2014.
- Krichak, S.O., Barkan, J., Breitgand, J.S., Gualdi, S., and Feldstein, S.: The role of the export of tropical moisture into midlatitudes for extreme precipitation events in the Mediterranean region. Theor Appl Climatol 121, 499–515.
 https://doi.org/10.1007/s00704-014-1244-6, 2015.
 - Lehmann, J., Coumou, D., and Frieler, K.: Increased record-breaking precipitation events under global warming, Climatic Change 132: 501, https://doi.org/10.1007/s10584-015-1434-y, 2015.
 - Longman, J., Ersek, V., Veres, D., and Salzmann, U.: Detritical events and hydroclimate variability in the Romanian Carpathians during the mid-to-late Holocene, Quaternary Science Reviews, 167, 78-95, 2017.
- Libertino, A., Ganora, D., and Claps, P.: Evidence for increasing rainfall extremes remains elusive at large spatial scales: The case of Italy. Geophysical Research Letters, 46, 7437–7446. https://doi.org/10.1029/2019GL083371, 2019.
 - Lowe, D.R.: Sediment gravity flows; Depositional models with special reference to the deposits of high-density turbidity currents, Journal of Sedimentary Petrology, 52, 279-297, 1982.

Lowe, J.J., Accorsi, C.A., Bandini Mazzanti, M., Bishop, V., Van der Kaars, S., Forlani, L., Mercuri, A.M., Rivalenti, C.,

750 Torri, P., Watson, C.: Pollen stratigraphy of sediment sequences from carter lakes Albano and Nemi (near Rome) and

from the central Adriatic, spanning the interval from oxygen isotope stage 2 to the present day, Mem. Ist. Ital. Idrobiol. 55, 71–98, 1996.

- Magny, M.: Holocene climate variability as reflected by mid-European lake-level fluctuations and its probable impact on prehistoric human settlements, Quaternary International, 113, 65-79, 2004.
- Magny, M., Galop, D., Bellintani, P., Desmet, M., Didier, J., Hass, J.N., Martinelli, N., Pedrotti, A., Scandolari, S., Stock,
 A., and Vannière, B.: Late-Holocene climatic variability south of the Alps as recorded by lake-level fluctuations at
 Lake Ledro, Trentino, Italy. Holocene, 19, 575-589, 2009.
 - Magny, M., S. Joannin, D. Galop, B. Vanniere, J. N. Haas, M. Bassetti, P. Bellintani, R. Scandolari, and M. Desmet. 2012. Holocene palaeohydrological changes in the northern Mediterranean borderlands as reflected by the lake-level record of Lake Ledro, northeastern Italy. Quaternary Research, 77, 382-396, 2012.
 - Magny, M., Peyron, O., Sadori, L., Ortu, E., Zanchetta, G., Vannière, B. and Tinner, W.: Contrasting patterns of precipitation seasonality during the Holocene in the south- and north-central Mediterranean. J. Quaternary Sci., 27: 290-296. doi:10.1002/jqs.1543, 2012b.
- Marchetti, G., and Fraccia, R.: Carta geomorfologica dell'alta Val Nure, Appennino piacentino. Scala 1:25.000, in: Il paesaggio fisico dell'Alto Appennino Emiliano. Regione Emilia-Romagna, Carton, A., and Panizza, M., (Eds.), Grafis Edizioni, Bologna, Italia, 1988.
 - Marcott, S., Shakun, J., Clark, P.U., and Mix, A.C.: A Reconstruction of Regional and Global Temperature for the Past 11,300 Years, Science, 339, 1198-1201, doi 10.1126/science.1228026, 2013.
- Marroni, M., Meneghini, F., and Pandolfi, L.: Anatomy of the Ligure-Piemontese subduction system: evidence from Late
 Cretaceous-middle Eocene convergent margin deposits in the Northern Apennines, Italy, International Geology
 - Review, iFirst article, 1-33, 2010. Milliman, J.D. and Syvitski, J.P.M.: Geomorphic/Tectonic Control of Sediment Discharge to the Ocean: The Importance

of Small Mountainous Rivers, The Journal of Geology, 100, 525-544, http://dx.doi.org/10.1086/629606, 1992.

Moore, P.D., Webb, J.A., and Collinson, M.E.: Pollen Analysis, Oxford, 1991.

760

790

775 Mulder, T. and Syvitski, J.P.M.: Turbidity Currents Generated at River Mouths during Exceptional Discharges to the World Oceans, Journal of Geology, 103, 285-299, http://dx.doi.org/10.1086/629747, 1995.

Mutti, E.: Turbidite Sandstones. San Donato Milanese, Agip-Istituto di Geologia, Università di Parma, 275 pp, 1982.

- Mutti, E., Davoli, G., Tinterri, R., and Zavala, C.: The importance of ancient fluvio-deltaic systems dominated by catastrophic flooding in tectonically active basins, Memorie di Scienze Geologiche, Università di Padova, 48, 233-291, 1996.
 - Myhre, G., Alterskjær, K., Stjern, C.W., Hodnebrog, Ø., Marelle, L., Samset, B.H., Sillmann, J., Schaller, N., Fischer, E., Schulz, M., and Stohl, A.: Frequency of extreme precipitation increases extensively with event rareness under global warming. Scientific Reports 9:16063, https://doi.org/10.1038/s41598-019-52277-4, 2019.
- Oswald, W.W., Anderson, P.M., Brown, T.A., Brubaker, L.B., Hu, F.S., Lozhkin, A.V., Tinner, W., and Petra
 Kaltenrieder, P.: Effects of sample mass and macrofossil type on radiocarbon dating of arctic and boreal lake sediments. The Holocene 15, 5, 758-767, 2005.
 - Papalexiou, S. M. and Montanari, A.: Global and regional increase of precipitation extremes under global warming. Water Resources Research, 55, 4901–4914. https://doi.org/10.1029/2018WR02406, 2019.

Patterson, W.A., Edwards, K.J., and Maguire, D.J.: Microscopic charcoal as a fossil indicator of fire, Quaternary Science Reviews, 6, 3-23, https://doi.org/10.1016/0277-3791(87)90012-6,1987.

Pignatti, S.: Flora d'Italia, Edagricole-New Business Media, Milano, 2017-2019.

Prein, A.F., Rasmussen, R.M., Ikeda, K., Liu, C., Clark, M.P., and Holland, G.J.: The future intensification of hourly precipitation extremes, Nature Climate Change, 7, 48-52, doi:10.1038/nclimate3168, 2017.

- Regattieri, E., Zanchetta, G., Drysdale, R.N., Isola. I., Hellstrom, J.C., and Dallai, L.: Lateglacial to Holocene trace
 element record (Ba, Mg, Sr) from Corchia Cave (Apuan Alps, central Italy): paleoenvironmental implications, Journal of Quaternary Science, 29, 381-392, https://doi.org/10.1002/jqs.2712, 2014.
 - Reimer, P.J., Bayliss, A., Warren Beck, J., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott,
- E.M., Southon, J.R., Staff, R.A., Turney, CSM., van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP, Radiocarbon, 55, 1869-1887, https://doi.org/10.2458/azu_js_rc.55.16947, 2013.
 - Renssen, H., Seppä, H., Crosta, X., Goosse, H., and Roche, D.M.: 2012. Global characterisation of the Holocene Thermal Maximum. Quaternary Science Reviews 48, 7-19, 2012.
- 805 Samartin, S., Heiri, O., Joos, F., Renssen, H., Franke, J., Brönnimann, S., and Tinner, W.: Warm Mediterranean mid-Holocene summers inferred from fossil midge assemblages, Nature Geoscience, 10, 207-212. doi: 10.1038/NGEO2891, 2017.

810

- Scherrer, S. C., Fischer, E.M., Posselt, R., Liniger, M.A., Croci-Maspoli, M., and Knutti, R.: Emerging trends in heavy precipitation and hot temperature extremes in Switzerland, J. Geophys. Res. Atmos., 121, doi:10.1002/2015JD024634, 2016.
- Schillereff, D., Chiverrell, R.C., Macdonald, N., and Hooke, J.M.: Flood stratigraphies in lake sediments: A review, Earth-Science Reviews, 135, 17-37, doi 10.1016/j.earscirev.2014.03.011, 2014.
 - Schneuwly-Bollschweiler, M., Stoffel, M., and Rudolf-Miklau, F. (Eds.): Dating Torrential Processes on Fans and Cones. Methods and Their Application for Hazard and Risk Assessment. Springer, doi 10.1007/978-94-007-4336-6, 2013.
- 815 Segadelli S., Grazzini F., Adorni M., De Nardo M.T., Fornasiero A., Chelli A., and Cantonati M.: Predicting Extreme-Precipitation Effects on the Geomorphology of Small Mountain Catchments: Towards an Improved Understanding of the Consequences for Freshwater Biodiversity and Ecosystems. Water 12: 79, https://doi.org/10.3390/w12010079, 2020.
- Skinner, C. B., and Poulsen, C.J.: The role of fall season tropical plumes in enhancing Saharan rainfall during the African
 Humid Period, Geophys. Res. Lett., 43, 349–358, doi:10.1002/2015GL066318, 2016.
 - Stoffel, M., Butler, D.R., and Corona, C.: Mass movements and tree rings: A guide to dendrogeomorphic field sampling and dating, Geomorphology, 200, 106-120, https://doi.org/10.1016/j.geomorph.2012.12.017, 2013.
 - Stoffel, M., Wyzga, B., and Marston, R.A.: Floods in mountain environments: A synthesis, Geomorphology, 272, 1-9, https://doi.org/10.1016/j.geomorph.2016.07.008, 2016.
- 825 Swierczynski, T., Ionita, M., and Pino, D.: Using archives of past floods to estimate future flood hazards, Eos 98, https://doi.org/10.1029/2017EO066221, 2017.
 - Tinner, W., Colombaroli D., Heiri, O., Henne, P., Steinacher, M., Untenecker, J., Vescovi, E., Allen, J., Carraro, G., Conedera, M., Joos, F., Lotter, A., Luterbacher, J., Samartin, S., and Valsecchi, V.: The past ecology of Abies alba provides new perspectives on future responses of silver fir forests to global warming. Ecological Monographs 83(4): 419-439, 2013.
 - Tutin, T.G., Heywood, V.H., Burges, N.A., and Valentine, D.H.: Flora Europaea, Cambridge University Press, 1964-1993.

Utsumi, N., Seto, S., Kanae, E., Maeda, E., and Oki T.: Does higher surface temperature intensify extreme precipitation?, Geophys. Res. Lett., 38, L16708, doi:10.1029/2011GL048426, 2011.

- 835 Vescovi, E., Ammann, B., Ravazzi, C., and Tinner, W.: A new Late-glacial and Holocene record of vegetation and fire history from Lago del Greppo, northern Apennines, Italy. Vegetation History and Archaeobotany, 19, 219–233, https://doi.org/10.1007/s00334-010-0243-5, 2010.
 - Vittori Antisari, L., Bianchini, G., Cremonini, S., Di Giuseppe D., Falsone, G., Marchesini, M. Marvelli, S., and Vianello G.: Multidisciplinary study of a Lateglacial-Holocene sedimentary sequence near Bologna (Italy): insights on natural
- 840 and anthropogenic impacts on the landscape dynamics. J Soils Sediments 16, 2 645-662, doi 10.1007/s11368-015-1266-4, 2015
 - Westra, S., Fowler, H.J., Evans J.P., Alexander, L.V., Berg, P., Johnson, F., Kendon, E.J., Lenderink, G., and Roberts, N.M.: Future changes to the intensity and frequency of short duration extreme rainfall, Reviews of Geophysics, 52, 522-555, doi:10.1002/2014RG000464, 2014.
- 845 Whitlock, C. and Larsen, C.: Charcoal as a fire proxy. In Smol J.P., Birks, H.J.B. and Last, W.M., (Eds.), Terrestrial, algal, and siliceous indicators, Dordrecht: Kluwer, 75-97, 2001.
 - Whitlock, C., and Millspaugh, S.H.: Testing the assumptions of fire-history studies: an examination of modern charcoal accumulation in Yellowstone National Park, USA, The Holocene, 6, 7-15, https://doi.org/10.1177/095968369600600102, 1996.
- 850 Wilhelm, B., Arnaud, F., Sabatier, P., Crouzet, C., Brisset, E., Chaumillon, E., Disnar, J.R., Guiter, F., Malet, E., Reyss, J.L., Tachikawa, K., Bard, E., and Delannoy, J.J.: 1400 years of extreme precipitation patterns over the Mediterranean French Alps and possible forcing mechanism, Quaternary Research, 78, 1-12, doi 10.1016/j.yqres.2012.03.003, 2012.
 - Wilhelm, B., Ballesteros-Cànovas, J.A., Macdonald, N., Toonen, W.H.J., Baker, V., Barriendos, M., Benito, G., Brauer,A., Corella, J.P., Denniston, R., Glaser, R., Ionita, M., Kahle, M., Liu, T., Luetscher, M., Macklin, M., Mudelsee, M.,
- 855 Muňoz, S., Schulte, L., George, S.St., Stoffel, M., and Wetter, O.: Interpreting historical, botanical, and geological evidence to aid preparations for future floods, WIREs Water, 1-22, https://doi.org/10.1002/wat2.1318, 2018.
 - Wirth, S.B.: The Holocene Flood History of the Central Alps Reconstructed from Lacustrine Sediments: Frequency, Intensity and Controlling Climate Factors. Ph.D. thesis, ETH Zürich Bibliography, Switzerland, 179pp, doi 10.3929/ethz-a-009775044, 2013.
- 860 Wirth, S.B., Glur, L., Gilli, A., and Anselmetti, F.S.: Holocene flood frequency across the Central Alps solar forcing and evidence for variations in North Atlantic atmospheric circulation, Quaternary Science Reviews, 80, 112-128, https://doi.org/10.1016/j.quascirev.2013.09.002, 2013.

865

- Zanchetta, G., Isola, I., Piccini, L., and Dini, A.: The Corchia Cave (Alpi Apuane) a 2 Ma long temporal window on the earth climate, Technical Periodicals of National Geological Survey of Italy ISPRA, Geological Field Trips, 3, 55pp, doi 10.3301/GFT.2011.02, 2011.
- Zavala, C. and Pan, S.: Hyperpycnal flows and hyperpycnites: Origin and distinctive characteristics. Lithologic Reservoirs, 30, 1-27, doi 10.3969/j.issn.1673-8926.2018.01.001, 2018.
- Zavala, C., Ponce, J.J., Arcuri, M., Drittanti, D., Freije, H., and Asensio, M.: Ancient Lacustrine Hyperpycnites: A Depositional Model from a Case Study in the Rayoso Formation (Cretaceous) of West-Central Argentina. Journal of Sedimentary Research, 76, 41-59, https://doi.org/10.2110/jsr.2006.12, 2006.
- Zavala, C., Arcuri, M., Di Meglio, M., Gamero Diaz, H., and Contreras, C.: A Genetic Facies Tract for the Analysis of Sustained Hyperpychal Flow Deposits. in: Sediment transfer from shelf to deep water - Revisiting the delivery system, edited by: Slatt, R.M. and Zavala, C., AAPG Studies in Geology, 61, 31-51, 2011.

Zhornyak, L.V., Zanchetta, G., Drysdale, R.N., Hellstrom, J.C., Isola, I., Regattieri, E., Piccini, L., Baneschi, I., and Couchoud, I.: Stratigraphic evidence for a "pluvial phase" between ca 8200-7100 ka from Renella cave (Central Italy), Quaternary Science Reviews, 30, 409-417 doi 10.1016/j.quascirev.2010.12.003, 2011.