

Changes in high-intensity precipitation on the Northern Apennines (Italy) as revealed by multidisciplinary data over the last 9000 years

5 Stefano Segadelli¹, Federico Grazzini^{2,6}, Veronica Rossi³, Margherita Aguzzi², Silvia Marvelli⁴,
Marco Marchesini⁴, Alessandro Chelli⁵, Roberto Francese⁵, Maria Teresa De Nardo¹, Sandro Nanni²

¹ Geological, Seismic and Soil Service, Emilia-Romagna Region Administration, Bologna, Italy.

² Regional Agency for Prevention, Environment and Energy of Emilia-Romagna, Hydro-Meteo-Climate Service (ARPAE-SIMC), Bologna, Italy.

10 ³ Department of Biological, Geological, and Environmental Sciences - BiGeA, Alma Mater Studiorum University of Bologna, Bologna, Italy.

⁴ Laboratory of Palynology and Archaeobotany - C.A.A. Giorgio Nicoli, San Giovanni in Persiceto (Bologna, Italy).

⁵ Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Italy.

⁶ Ludwig-Maximilians-Universität, Meteorologisches Institut, München, Germany

15

Correspondence to: Federico Grazzini (fgrazzini@arpae.it)

Abstract. Several record-breaking precipitation events have stricken the mountainous area of Emilia-Romagna Region (Northern Apennines, Italy) over the last 10 years. As a consequence, severe geomorphological processes such as debris
20 avalanches and debris flows, shallow landslides and overbank flooding have affected the territory, causing severe damage to human-made structures. The unusual intensity of these phenomena prompted an investigation into their frequency in the past, beyond instrumental time. In the quest for understanding, whether these phenomena are unprecedented in the region, peat bog and lake deposits were analysed to infer the frequency of extreme precipitation events that may have occurred in the past. We present the results of a dedicated field campaign performed in summer 2017 at Lake Moo in the
25 Northern Apennines, a 0.15 km² peat bog located at an altitude of 1130 m a.s.l. During the extreme precipitation event of 13-14 September 2015, several debris flows generated by small streams, have affected the Lake Moo plain. In such a small drainage basin (< 2km²), high-density floods can be triggered only by high-intensity precipitation events. The sedimentary succession (ca. 13 m-thick) was studied through the drilling of two cores and one trench. The sequence, characterised by clusters of coarse-grained alluvial deposits interbedded with organic-rich silty clays and peat layers, was
30 analysed combining sedimentological, pollen, microanthracological, pedological data and radiocarbon dating (AMS ¹⁴C), in an innovative multidisciplinary approach for this area. Original data acquired during the field campaign were also correlated with other specific paleoclimatic proxies available in the literature for the Northern Apennines area. We discover that the increase of extreme paleoflood, associated with coarse-grained deposits similar to the ones observed recently, correlates well with the warm phases of the Holocene Thermal Maximum and with the ongoing warming trend
35 observed which started at the beginning of the last century.

Keywords

extreme precipitation, Holocene flood activity, Northern Apennines, climate change, water cycle, global warming

1. Introduction

40 High-intensity precipitation (HIP) events, also known as torrential rainstorms for their capacity to generate flash floods in small streams, represents a significant component of the Mediterranean water cycle. More than half of the annual precipitation is concentrated in a few major precipitation events (e.g., Frei and Schär 1998; Isotta et al., 2014). A better knowledge of their expected frequency (and maximum intensity) is crucial for planning adequate hydraulic defences and sustainable water resources management in the present
45 and future climate.

Under the threat of global warming, a growing number of studies have investigated the link between a temperature rise in air masses and intensity of extreme precipitation, highlighting spatial and seasonal differences across the globe (e.g., Lehmann et al., 2015; Papalexiou and Montanari, 2019). There is a consensus that HIP events are increasing with global warming, while mean precipitation could decrease in some regions
50 (Berg et al., 2013). Myhre et al. (2019) show that HIP precipitation over Europe almost doubles per degree of warming due to the combined effect of increasing frequency (the major driver) and an increase in intensity.

At the local scale, different responses of weather patterns and limitations in moisture availability can alter the uniform expected raise of HIP due to the increase of saturation water vapour pressure ($6\text{-}7\% \text{ K}^{-1}$) (e.g., Westra et al., 2014; Prein et al., 2017). For example, in the Swiss Alps Scherrer et al. (2016) found a significant
55 increase in daily extreme precipitation indices since the beginning of the last century (i.e., between 1901-2014) while Brugnara and Maugeri (2019) documented regional differences in HIP trends across the whole Alpine area. These results suggest that instrumental data do not cover a suitable period to infer reliable local changes in HIP frequency.

The investigation of sedimentary archives, like peat bogs and lakes, allow verifying the hypothesised
60 linkage between extreme precipitation and temperatures in the distant past, widening substantially the period where we can verify this relationship.

In the literature there is ample documentation of the use of these sedimentary archives to infer chronologies of past flood events (Zavala et al., 2006; Zavala et al., 2011; Giguet-Covex et al., 2012; Gilli et al., 2013; Glur et al., 2013; Stoffel et al., 2013; Wirth, 2013; Wirth et al., 2013; Anselmetti et al., 2014;
65 Longman et al., 2017; Schillereff et al., 2014; Stoffel et al., 2016; Swierczynski et al., 2017; Ahlborn et al., 2018; Wilhelm et al., 2012, 2018; Zavala and Pan, 2018), alongside others such as tree rings (Ballesteros-Cánovas et al., 2015), speleothems (Regattieri et al., 2014; Zanchetta et al., 2011) and torrential fans and cones (Schneuwly-Bollschweiler et al., 2013).

Interestingly, over the Alpine area a synchronisation between increasing flood frequency and
70 Holocene cooling periods characterised by cold and wet summers, like the Little Ice Age-LIA, has been documented by several studies (e.g., Glur et al., 2013; Henne et al., 2018). Other authors reported that under warm paleoclimate conditions floods in the Alps became rarer but stronger in intensity, highlighting the complexity of this issue (Giguet-Covex et al., 2012; Brönnimann et al., 2018a). Alpine data mostly derive from lacustrine sedimentary successions that are known to be influenced by a combination of factors (e.g.,
75 precipitation intensity, duration, seasonality and changes in atmospheric-circulation and land use) that might overshadow the physically-based relationship between temperature increase and extreme precipitation events

(Utsumi et al., 2011; Brönnimann et al., 2018b).

In contrast, less attention was paid to the Holocene flood activity in the Northern Apennine area (N Italy in Fig. 1), which has a different precipitation climatology compared to the inner Alpine region. Indeed, the link between cold summers and high values of flood frequency appears more logical for the Alpine region, which shows a precipitation maximum in the Summer. In contrast, in the Northern Apennine area, precipitation maxima occur in Autumn with more than 60% of the annual total precipitation concentrated in few days characterised by severe meteorological events (Isotta et al., 2014).

Focusing the analysis on the Emilia-Romagna Region (ER Region; Fig. 1), an increase in interannual variability of torrential rainfall has been observed over the last ten years with marked or even exceptional droughts, such as those occurring in 2012 and 2017 (Grazzini et al., 2012), followed by years of record-breaking rainfall (2014 and 2018). Between September 2014 and September 2015, the ER Region has been affected by three events of exceptional intensity with an estimated return period of several centuries (Grazzini et al., 2016).

To consolidate HIP trends on the N Apennine area, we analysed the Holocene sedimentary succession of a small-size peat bog (Lake Moo in Fig. 1) through the application of a multidisciplinary approach that integrates sedimentological and environmental data (e.g., pollen-derived paleovegetation patterns) with climatological observations and physical arguments.

Lake Moo is in the proximity of the main N Apennines crest that is very exposed to moist maritime airflow coming from the central Mediterranean Sea. The dimension of the drainage basin is small (total area <2 km²) and the modern anthropogenic impact is low as the site belongs to the Natura 2000 network (site code and name: IT4020008 - Monte Ragola, Lago Moo, Lago Bino). Moreover, the activation of debris flows (*sensu* Hungr et al., 2013) during the severe flash flood event occurred on 13th-14th September 2015, with an estimated rainfall intensity of 112 mm/3h over the Lake Moo basin (Segadelli et al., 2020; Grazzini et al., 2016), offers the unique opportunity to use these recent deposits as the archetype of past events. According to literature (Milliman and Syvitski, 1992; Mulder and Syvitski, 1995; Mutti et al., 1996), in such a small drainage basin high-density floods can be triggered only by HIP events lasting enough time for the water to infiltrate and mobilise large quantities of debris. All these geomorphological features make Lake Moo an ideal site to investigate past changes in HIP events in terms of sedimentary expression, frequency and forcing factors. This research addressed the following questions:

- a) Are the recent events unprecedented on a millennial timescale (i.e., the Holocene period)?
- b) Is the frequency of HIP events coupled with (paleo)temperature variations?

2. The study area

2.1. Geographic and climate context

The Lake Moo plain (44°37'29"N, 9°32'25"E), about 0.15 km² wide, is located near the boundary between Emilia-Romagna and Liguria regions (Piacenza province, Italy; Fig. 1a) in the upper valley of the

115 Nure stream at an altitude of 1130 m above sea level (a.s.l.). A high tree cover density characterises the present-day catchment area (i.e., the total woodland cover is 65.55%; Corticelli et al., 2011), with a high richness of plant species and an exceptional concentration of protected mountain taxa with a particular phytogeographical interest. The vegetation landscape shows the widespread occurrence of *Fagus sylvatica*, locally interrupted by grazing areas and blueberry moorlands with the presence of rare *Juniperus nana* and *Sorbus chamaemespilus*. Reforestation of *Pinus nigra* is also documented (Fig. 1b).

120




Figure 1. (a) Location map of the Lake Moo plain. The area affected by the extreme precipitation event of September 13th and 14th 2015 is highlighted by the red rectangle. (b) View of the Lake Moo landscape from the southern side.

125

Present-day climate conditions primarily reflect the interaction of the prevalent atmospheric flow and the mountain range surrounding the Lake Moo plain. The proximity to the Ligurian Sea (about 40 km SW) makes the location particularly favourable to orographic precipitation enhancement in the case of moisture-laden southerly-flow. The largest precipitation is usually observed in Spring and the Autumn, with a marked peak in early Autumn. In this period we observe a particular synergy of mid-latitude synoptic disturbances, becoming more frequent towards the cold seasons, and strong convective systems still developing over the warm Mediterranean Sea at the end of Summer and in Autumn (Grazzini et al., 2020); a situation very favourable the genesis of extreme precipitation events in the area.

130

2.2. Geological and geomorphological setting

135 The study area mainly consists of extensively fractured serpentinites that represent the accreted fragments of the Ligure-Piemontese oceanic basin separating the European and Adria plates during the Middle-Upper Jurassic (Marroni et al., 2010). The ultramafites are bordered by polygenic breccias, made of blocks of limestones or marly limestones embedded within a fine-grained matrix (Mt. Ragola Complex, Late Santonian Early Campanian; Elter et al., 1997).

140 The geomorphological landscape includes flat areas and steep slopes located at different altitudes. On the former, marshy environments commonly occur recording the last filling phases of small lacustrine basins, some of which still exist as the Lake Moo and the Lake Bino (Figs. 1-2). However, the origin of Lake Moo is still a matter of debate as some authors point to a glacial origin (Elter et al., 1997; Marchetti and Fraccia, 1988; Carton and Panizza, 1988), while others have interpreted the basin as the expression of Holocene deep-seated
145 gravitational slope deformations (Geological, Seismic and Soil Service of the Emilia-Romagna Region, 2012) mainly controlled by changes in the incision rates of the Nure stream (Elter et al., 1997). The complexity of past and present morphological processes are likely enhanced by the superposition of lithological units with strong mechanical contrast, such as ophiolites and the underlying a predominantly clayey unit (Elter et al., 1997).

150


3. Data and methods

3.1. Field investigation

Taking into account the criteria indicated by Gilli et al. (2013) and Schillereff et al. (2014), Lake Moo shares several advantageous characteristics for the reconstruction of the Holocene flood activity:

- 155 - steep slopes (average inclination of 24°) composed of deposits highly susceptible to erosion (i.e., polygenic and monogenic breccias with a pelitic matrix – the Mt. Ragola Complex, Elter et al., 1997; Fig. 2);
- absence of lacustrine basins in the upstream part of the catchment;
- small drainage basin area (1.94 km²);
- 160 - one dominant inflow into the lake;
- lack of regulated flow structures;
- lack of natural pre-lake sediment storage zones.

The field campaign led to the acquisition of two sedimentary cores (S1 and S2), 14m and 6m-long
165 respectively, and one trench (T1), to investigate the sedimentary succession capped by the high-density flood deposits formed during the recent HIP event (13th-14th September 2015). The location of the cores and the trench benefited from a detailed geomorphological map (Fig. 2) and a high-resolution reflection seismic survey (Fig. SUP1 in the supplementary material), both originally produced for this research. The latter provided useful information about the lake basin-floor morphology and the thickness of the infilling succession.



170

Figure 2. Geomorphological map showing the main geological units and the hydrological elements of the Lake Moo plain. The detailed mapping of flood deposits formed during the recent HIP event (13th-14th September 2015), the location of the geognostic investigations and the geophysical surveys tracks are also reported. Original data from field campaign overlaid on the Geological database of Emilia-Romagna region.

175

The two cores, S1 and S2, were obtained using a continuous drilling system that guaranteed an undisturbed stratigraphy and a high recovery percentage (about 90%). A trench (6 m long, 3 m wide, 2 m depth) was excavated between the two coring sites allowing a detailed analysis of the most recent deposits.

180

We focused our attention on the longest core (S1) that shows the complete record of the Lake Moo infilling succession, as it reaches the ophiolite bedrock. Therefore, all the laboratory analyses (grain size, radiocarbon and palynological analysis) were undertaken on core S1, while the other core (S2) and the trench were used to support the stratigraphic reconstruction. Elevation and geographic coordinates of reference core S1 were acquired using Garmin eTrex 10 GPS receiver: 1120.2 m. a.s.l. and 44° 37' 25"N – 9° 32' 43"E, respectively.

3.2. *Facies analysis and chronology*

Facies characterisation of the Lake Moo sedimentary succession was mainly performed integrating the observable macroscopic physical characteristics (i.e., grain size, sedimentary structures, Munsell chart colour and types of bounding surfaces) with the grain-size data available for the reference core S1. The interpretation in terms of depositional environments also benefited from the precise position of the core(s) with respect to the spill point of Lake Moo and the application of the facies tract concept. The latter strongly supported the interpretation of the coarse-grained intervals encountered throughout the studied succession. Specifically, a facies tract is defined as the association of genetic facies that can be observed within the flood deposits along with the down-slope motion transformation (Lowe, 1982; Mutti, 1992; Mutti et al., 1996). Thus, the application of the facies tract concept, if framed into the core(s) stratigraphy, may allow the reconstruction of how flood deposits changed through time at a fixed location (Figs. 3, 4).

The abundance of wood remains and peaty layers within the sedimentary succession under examination supported the development of a robust chronological framework. Twelve samples were selected from the reference core S1 and radiocarbon dated at the CEDAD Laboratory of the University of Salento (Italy). The data are available in Table TS1 (in supplementary material) and the conventional ^{14}C ages were converted into calendar years using the OxCal software version 3.10 (Reimer et al., 2013).

An age-depth model was also constructed using linear interpolation. Coarse-grained intervals were not excluded from the processing because of the difficulty to evaluate the exact thickness of the deposits interpretable as instantaneous events. Indeed, the upper limit of coarse-grained intervals commonly appears transitional toward finer sediments and, in few cases, some intervals for a total thickness of 107 cm were not recovered.

3.3. *Pollen and microanthracological analysis*

In order to refine facies characterisation and highlight past vegetation changes at the study site, pollen analyses were carried out on 14 samples collected from the core S1. We focused the analyses on two key stratigraphic intervals, formed during periods of well-known different paleoclimate (and paleotemperature) conditions: i) Interval 1 (ca. 10.5-9 m core depth) covering the period between ca. 9600-7300 cal yr BP and, thus centred on the Holocene Thermal Maximum-HTM (Renssen et al., 2012) and ii) Interval 2 (ca. 5.5-4.5 m core depth) recording the final stages of HTM and the following cooling period between ca. 5500-3800 cal yr BP. In these two stratigraphic intervals, samples were collected from fine-grained layers following a mean sampling resolution of ca. 30 cm.

A standard methodology already tested for pollen substrates was applied with some minor modifications (Lowe et al., 1996). The method includes a series of laboratory treatments: about 8-10 g of sample was treated in 10% Na-pyrophosphate to deflocculate the sediment matrix; a Lycopodium tablet was added to calculate pollen concentration (expressed as pollen grains per gram = p/g) and the sediment residue was subsequently washed through a 7 micron sieve. The sample was re-suspended in HCl 10% to remove

calcareous material and subjected to Erdtman acetolysis; heavy liquid separation method was then introduced using Na-metatungstate hydrate of s.g. 2.0 and centrifugation at 2000 rpm for 20 minutes. Following this procedure, the retained fractions were treated with 40% HF for 24 h and then the sediment residue was washed
225 in distilled water and after in ethanol with glycerol; the final residue was desiccated and mounted on slides using glycerol jelly and finally sealed with paraffin.

Identification of the samples was performed at x400 magnification and only difficult pollen such as *Triticum*, *Avena* or *Hordeum* types were observed at x1000 magnification.

Determination of pollen grains was based on the Palinoteca of the “Centro Agricoltura Ambiente –
230 CAA G. Nicoli” laboratory (Italy), atlases and a vast amount of specific morpho-palynological literature. Names of the families, genus and species of plants conform to the classifications of Italian Flora proposed by Pignatti (2017-2019) and European Flora (Tutin et al., 1964-1993). The pollen terminology was based on Berglund and Ralska-Jasiewiczowa (1986), Faegri and Iversen (1989) and Moore et al. (1991) with slight modifications that tend to simplify plants nomenclature. The term “taxa” is used in a broad sense to indicate
235 both the systematic categories and the pollen morphological types (Beug, 2004). For each sample, at least 500 pollen grains were counted, and the identified taxa have been expressed as percentages of the total pollen sum that includes only terrestrial pollen, no fern spores and no aquatic plants.

On the basis of vegetational and ecological characteristics, the following main pollen groups were identified: Conifers (*Pinus/Abies alba*), Deciduous trees (this group includes quercetum taxa – *Quercus*,
240 *Carpinus betulus*, *Corylus avellana*, *Fraxinus*, *Ostrya carpinifolia*, *Tilia* and *Ulmus* + other Deciduous trees), meadow (this group mainly comprises *Fabaceae* and *Asteroideae*, *Caryophyllaceae*, *Cichorioideae* and *Poaceae*), anthropogenic indicators (e.g. *Cerealia*, *Chenopodium*, *Convolvulus arvensis*, *Plantago*, *Urtica*) and Alia that includes all taxa excluded from previous groups. The group of hygro + aquatic plants was also distinguished. It includes hygrophilous herbs (e.g., *Cyperaceae*), helophytes and hydrophytes (i.e., *Lemna*,
245 *Juncus*, *Nymphaea*, *Phragmites*, *Potamogeton* and *Sparganium*), which are considered good proxy of humid conditions typical of wetlands.

The 14 samples prepared for pollen analysis were also investigated for the identification of microcharcoals. Microanthracological analysis has been used to track past changes in fire history, likely connected to anthropogenic activities (Vescovi et al., 2010). Point count estimation of microscopic charcoal
250 abundance was carried out, and charcoal fragments encountered during pollen counting were recorded in four, size classes based on the long axis length: 10–50, 50–125, 125–250, >250 μm (Whitlock and Millsaugh 1996; Clark 1982; Patterson et al., 1987; Whitlock and Larsen 2001, Fisinger et al., 2008). The former two classes were interpreted to be wind-blown transported, hence giving information about regional fire events, whereas the latter two were considered the result of local vegetation burning (Vittori Antisari et al., 2015).

255 Pollen and microcharcoal results are presented in section 5, to support facies stratigraphy and to discuss controlling factors on floods activation.

3.4. Temperature reconstruction and modern climatological dataset

Central to our study is the availability of a reliable (paleo)temperature dataset for the chronological
260 period recorded by the Lake Moo sedimentary succession. In this respect, the Holocene paleoclimate
reconstruction conducted by Samartin et al. (2017) on the nearby Lake Verdarolo represents a very important
reference. The Lake Verdarolo site is located at 1390 m a.s.l., 270 m higher and 54 km SE from Lake Moo
(Fig. 1), in a very similar climatic context. The authors reconstructed the mean July air temperature using a
chironomid-based inference model, developed through a combination of data extracted from more than 200
265 lakes of Norway and Swiss Alps (Heiri et al., 2011). This vegetation-independent paleotemperature
reconstruction is the first for the Northern Apennines and it agrees with that of the Gemini Lake (1350 m a.s.l.
elevation and 48 km SE from Lake Moo) as well as several other records coming from central Italy (Samartin
et al., 2017).

Modern temperature and precipitation time series (1961-2018) at Lago Verdarolo and Lake Moo were
270 derived from the gridded high-resolution dataset of Emilia-Romagna (Eraclito4), described in Antolini et al.
(2016). Trend estimation and Mann Kendall significance trend test were computed with the pyMannKendall
package (Hussain et al., 2019).

4. Results

275 4.1. Holocene stratigraphy

4.1.1. Sedimentary facies

Fourteen lithofacies (LM1 to LM14) have been identified within the infilling succession of the Lake Moo
basin. Detailed lithological description of each lithofacies is reported below.

- LM1 to LM2: Clast-supported deposit with a thickness ranging between 5-15 cm. Fine boulders to
280 coarse cobbles with abundant coarse - to fine - grained sand matrix, showing a low degree of sorting,
occur. Polygenic clasts have low sphericity and a very angular shape. Wood fragments locally occur.
The basal and top contacts are sharp; the basal one occasionally shows evidence of erosion;
- LM3 to LM4: Clast-supported deposit with a thickness ranging between 5-15 cm. It is composed of
fine cobbles to coarse pebbles with a very fine sand to silt matrix. Polygenic clasts have low sphericity
285 and a subangular shape. Scraps of wood are encountered. The basal and top contacts are sharp; the
basal one occasionally shows evidence of erosion;
- LM5: Massive matrix-supported deposit composed of medium pebbles to granules embedded within
a very fine sand to silt matrix. Polygenic clasts have low sphericity and a subangular shape. Scraps of
290 wood are encountered. The basal and top contacts are sharp; the basal one occasionally shows evidence
of erosion;
- LM6 to LM8: Generally clast-supported, poorly sorted deposit with a thickness ranging between 10-
30 cm. Polygenic clasts, massive or crudely graded angular\subangular fine pebbles to very fine

- 295 granules. Crudely horizontal laminae are locally recorded atop. Fragments of woods can be encountered at the base of the layers. The basal contact is sharp and occasionally shows evidence of erosion; the upper boundary is transitional;
- 300 - LM9 to LM10: Massive or crudely fining-upward graded very coarse to fine sands, showing a low degree of sorting. The total thickness ranges between 5-20 cm. Evidences of a planar lamination and scattered wood fragments occur close to the upper boundary, which is transitional. The basal contact is sharp;
- 305 - LM11: Fine sandy loam to clayey loam showing a dark grayish brown color (10YR 4/2 or 10YR 3/2). Polygenic fine to medium pebbles aligned with low sphericity and a very angular shape occur. Both basal and top contacts are sharp;
- 310 - LM12: Silty deposit of dark color (10YR 3/1) with a high content of decomposed organic matter. Occasional fine to medium pebbles aligned with low sphericity and very angular shape occasionally occur;
- LM13: Loam to silty clay deposit showing a coarse angular blocky structure and dark olive gray color (5Y3/2), with yellowish-brown (10YR5/6) due to the presence of iron oxides. The deposit is deprived of calcium carbonate and a field pH 5.5 value is recorded;

- LM14: Massive clayey silts with a low amount of decomposed organic matter and a dark greenish gray color (5G 4\1). Scattered polygenic clasts (fine pebbles) with low sphericity and a very angular shape are recorded. Both basal and top contacts are sharp.




Figure 3. Sedimentological features and lithofacies characterisation of the reference core S1. Radiocarbon dates, palynological and grain-size samples as well as stratigraphic units (described in the section 4.2.) are also reported.

In terms of sedimentary processes, the coarse-grained LM1-LM10 lithofacies have been interpreted as flood deposits triggered by HIP events in the catchment area (Milliman and Syvitski, 1992; Mulder and Syvitski, 1995; and Mutti et al., 1996). Particularly, the identified lithofacies were grouped according to their features (i.e., grain-size, colour, sedimentary structures) and the facies tract concept (Lowe, 1982; Mutti, 1992; Mutti et al., 1996), distinguishing three main depositional settings along an idealised transect (Fig. 4): subaerial (LM1 to LM4), marginal (LM5) and subaqueous/lacustrine (LM6 to LM10).

The fine-grained facies (LM11-LM14 in Figs. 3, 4) also form two main groups. The first group includes LM11 and LM13 (paleosol) that are characterised by features indicative of subaerial conditions; the second group consists of LM12 (i.e., peaty deposits) and LM14, deposited along the marginal zone of the (paleo)lake under low-energy conditions and different degree of organic-matter enrichment.




Figure 4. Idealised genetic facies tract showing the fourteen lithofacies (LM1-LM14) identified within the reference core S1, whose stratigraphy is reported in Fig. 3.

4.1.2. Stratigraphic units

Above the ophiolite bedrock, five informal stratigraphic units have been distinguished within core S1 and are described as follows (Fig. 3). Each unit is interpreted in terms of depositional environment.

Unit 1 (12.25-11.30 m). This 95 cm-thick unit is exclusively represented by the LM13 lithofacies that is, in turn, composed of two main horizons indicative of a mature paleosol. The lowermost horizon (Bw), recorded between 12.25-11.60 m, corresponds to weathered deposits overlain by 30 cm-thick slickensides horizon (Bss). This unit is interpreted as a residual pedogenized colluvium marking the base of a structural depression produced by gravitational block sliding;

Unit 2 (11.30-4.4 m). This ca.7 m-thick unit is characterised by the presence of several, very coarse intervals (LM6-LM8) showing a thickness of about 10-30 cm and locally capped by sandy layers (LM9-LM10). Silty deposits with a variable organic-matter content, accompanied by a scarce (LM12) or moderate (LM5) occurrence of pebbles/granules, separate the coarse-grained intervals. This unit, which is capped by a marked

unconformity surface, is interpreted as a relative shallow lacustrine environment subjected to several flood events of remarkable intensity;

355 Unit 3 (4.4-2.3 m). This unit, ca. 2 m thick, shows sharp boundaries and the occurrence of two sub-units (3a and 3b). The lowermost sub-unit 3a, ranging between 4.4-3.2 m, is exclusively represented by silty deposits with scattered pebbles/granules, several wood fragments and a mottled-like appearance corresponding to the LM5 lithofacies. These deposits are attributed to a marginal lacustrine environment subject to sporadic floods. Upwards, marginal lacustrine clayey deposits (LM14) capped the sequence (sub-unit 3b), documenting a period of apparent flood inactivity;

360

Unit 4 (2.3-0.9 m). This 1.4 m-thick unit, is composed of alternating coarse (LM3-LM4) and fine-grained deposits (LM12 and LM11) that record:

- the disappearance of truly lacustrine deposits that are replaced by subaerial ones through an erosional surface that marks the lower boundary of the Unit 4. The stratigraphic depth of this boundary is in accordance with the altitude of the spill point of Lake Moo (1116 m a.s.l.);
- the return of several, cm-thick flood deposits (LM3-LM4).

365

Unit 5 (0.9-0 m). This unit is characterised by the presence of several layers of very coarse sediments (LM1-LM2) separated by finer deposits (LM11). An erosional contact separates Unit 4 from Unit 5, which includes the flood deposits produced by the rainfall event occurred on September 13th and 14th 2015 (Figs. 3, 5).

370

As a whole, we interpret the lacustrine succession as an infill of a structural depression produced by gravitational block sliding that was induced by post-glacial fluvial incision (Elter et al., 1997).




375

Figure 5. Representative photograph of the trench (see Fig. 2 for location), showing the uppermost 50 cm-thick sedimentary succession. Though the exposure is quite small, the graded pebble-sand couplets belonging to the Unit 5 can be interpreted as a sheet-flood deposit.

380 4.2. Age-depth model

The age-depth model, based on eight out of twelve ¹⁴C dates available for the S1 core (Table TS1 in the supplementary material), indicates that the sedimentary succession above the bedrock encompasses almost all of the Holocene period (about the last 10 kyrs; Fig. 6). Three ages (at 9.60 m, 3.10 m and 2.40 m core depth) were excluded as they show evidences of contamination (i.e., ages younger than 1950 AD), while one sample collected at 2.35 m was not datable due to a very low organic matter content. The selected calibrated ages are stratigraphically coherent and mainly derived from wood fragments and peaty deposits. In order to reduce the potential bias due to long-lived plants and reworking processes, small twigs were selected (Oswald et al., 2005) from fine-grained deposits and identified when possible (e.g., *Pinus* and *Abies alba*). Due to the small dimension of the basin and the ophiolites prevalence in the drainage area (Fig. 2), we consider negligible the hard-water effect on peats.

The resulting age-depth model suggests the occurrence of four main stratigraphic intervals in terms of sedimentation rates. The lowermost interval (about 9.5-7.3 cal kyr BP) denotes a mean accumulation rate of ca. 0.50 mm/yr that increases up to ca. 2.38 mm/yr within the overlying section dated around 7.3-5.5 cal kyr B.P. An change in the accumulation rate occurs around 5 m with a drop to ca. 0.3 mm/yr between the 5.5-4.0 cal kyr BP interval. Between ca. 4.5-3 m, the deposit shows an uncertain chronology due to the lack of reliable radiocarbon dates and the occurrence of an unconformity surface around 4.5 m, at the contact between Units 2-3 (Figs. 3, 6). During recent times (the last 146-14 cal yr BP), the accumulation rate increases, reaching ca. 12 mm/yr.



400 **Figure 6.** S1 core stratigraphy and the age-depth model obtained from radiocarbon dates. Red and orange lines represent the two-sigma probability envelope.

5. Discussion

As a basis for discussion we assemble, and temporally synchronize, the Lake Moo data with the most relevant paleoclimate proxy available from literature, focusing on the N Apennines paleoarchives (Figs. 1, 7). As a whole, the Holocene infilling succession of the lacustrine basin (i.e., Units 2-3) is composed of an alternation of a centimeter to a meter thick, coarse-grained flood deposits and silts characterised by a variable amount of organic matter (sub-sections 4.1. and 4.2.; Figs. 3, 4, 7). The coarse intervals are particularly well-developed and frequent within the portion of the stratigraphic Unit 2 which is chronologically correspondent to the HTM (Holocene Thermal Optimum), constrained between ca. 9-5 kyr cal BP in the nearby Verdarolo Lake (Samartin et al., 2017 in Fig. 7A; section 3.4.). Interestingly, stable oxygen isotope records from the nearby Apuan Alps (Corchia caves; Fig. 1) document that, within the same time interval, the wettest conditions were observed (I and II humid periods in Fig. 7B) with a peak between ca. 8500 and 7500 cal yr BP (I humid period; Regattieri et al., 2014; Zanchetta et al., 2011). This precipitation peak is in phase with a clastic layer, dated to 8.2 and 7.1 cal kyr BP, indicative of a period of enhanced cave flooding triggered by high-magnitude precipitation events (Zhornyak et al., 2011). The authors attribute this maximum fluvial activity to an increase in strong convective episodes, like the one that affected the Versilia region in 1996 (see Cacciamani et al., 2000 for a description of the Versilia flood event).

Climatic simulations (monthly temperature and precipitation) for the mid-Holocene climatic conditions (Tinner et al., 2013), show higher precipitation in Autumn over the Northern Apennines. We speculate that the precipitation increases, also reflected in high-frequency flood activity, could have occurred during a phase of a progressive reduction of the Hadley cell, paralleled by a reduction in the subtropical high-pressure after the insolation peak (Fig. 7C). This would lead to enhanced meteorological activity over the Mediterranean area, due to the mid-latitude weather systems, especially at the transition towards the cold season. Although in decline, the presence of the African Monsoon (Skinner and Poulsen, 2016) could also have contributed to enrich sub-tropical air masses in the water vapour potentially extractable by mid-latitude synoptic disturbances. Accordingly, Krichak et al. (2015) documented how this mechanism is relevant for modern extremes in precipitation in the Mediterranean; moreover, the events occurring in the autumn months are characterised by a greater transport of water vapour from the subtropical Atlantic even across North Africa.

With the end of the HTM, a drastic decrease in flood activity is documented at Lake Moo, as the coarse intervals abruptly reduce in number and thickness within the uppermost portion of Unit 2 (ca. 5.5-3.8 kyr BP). This stratigraphic trend is reasonably interpreted as the expression of a decrease in frequency of HIP events over the study area under cooler and less humid climate conditions (Fig. 7). The limit between Unit 2 and Unit 3 corresponds to an unconformity that marks the passage to a marginal lacustrine succession of uncertain age, at least for the lowermost portion (sub-unit 3a). This interpretation is supported by palynological data, which show high values of herbaceous hygrophytes (mainly Cyperaceae) and aquatic species.

The upper sub-unit 3b dated around 146-14 cal yr BP, is almost deprived of coarse materials documenting a period of apparent flood inactivity at Lake Moo (Figs. 3, 7). This is followed by a reactivation of flood processes from c.a. 1800 AD (i.e., coarse-grained flood deposits within Units 4-5; Figs. 3, 7) with a minimum of five events to a maximum of twelve events every 100 years, as the confidence range associated

with the calibrated age at 2.80 m is rather wide (146-14 cal yr BP; Fig. 7). Despite the low degree of precision affecting radiocarbon ages younger than 200 years, this renewed increase in flood activity fits well with the instrumental record that points to a significant increase of record-breaking (from 3h to 24h accumulations) precipitation events in the Northern Appennine (Libertino et al., 2019) starting from the second half of last century. These phenomena seem to be responsible for the replacement of lake deposits by subaerial ones through an erosional contact (Unit 4 lower boundary).




Figure 7. Stratigraphy of core S1, main palynological features and microcharcoal content. Relative abundances of pollen groups explained in the text (sub-section 3.3), are reported along with the frequencies of hygrophilous herbs and aquatics. Asterisks point to samples containing coprophilous fungi and other spores like *Sporormiella*, *Dicrocoelium* and *Ascaris*. These data are compared with the most relevant paleoclimate reconstructions available from literature for the area of interest. Panel (A) Reconstructed mean July air temperature from Lago Verdarolo (from Samartin et al., 2017); Panel (B) $\delta^{18}\text{O}$ speleothem records and reconstructed precipitation trends (mean anomaly time series) from Corchia cave (slightly modified from Combourieu et al., 2013; Zanchetta et al., 2011; Zhornyak et al., 2011) and Panel (C) June (red) - December (blue) insolation values reported for 45° Nord (from Samartin et al., 2017). Main Holocene climate phases and events are also shown (Regattieri et al., 2014). Available radiocarbon ages are reported as calibrated years BP.

To compare the most recent stratigraphic units (Units 4-5) with quantitative climate data, the instrumental temperature values, available since the second half of the last century from the Eraclito ER dataset, has been added to the Verdarolo curve. The overlap with the latest part of Verdarolo curve confirms the good accuracy of the reconstruction technique in this region (Fig. 8). The recent sharp temperature increase is striking if compared with the whole Holocene, and the current July temperature is comparable with the maximum temperature reached at HTM over the study area. The actual trend of July temperature, estimated over the period 1961-2018 from monthly mean values, is +4.3°C in 100 years. The trend is statistically significant, with a p-value=0.0004 (see section 3.4.), and more than double of the maximum temperature gradient found in the Verdarolo curve, +2°C/100 years, at the end of the Little Ice Age 1850-1900. From Fig. 6 and 7, we observe flood intensification corresponding to periods of rising temperatures. This is testified by

the exceptionally high accumulation rate estimated on the recent period for the Units 4-5 (Fig. 6; section 4.3.), a rate of about 1 m/100 years calculated since ca. 1800 AD. The recent increase in flood activity is also supported by instrumental data, where we also observe an emerging trend (not yet significant) of precipitation intensity increase (both as daily maximum and as monthly cumulative values) in the autumn months (Fig. 9).




Figure 8. Comparison of current data and reconstructed July mean temperature at the Lake Verdarolo site (see Fig. 1 for location). The blue line is the reconstructed temperature and the shaded area is the sample-specific estimated standard error associated with the temperature reconstruction based on chironomid assemblage from Samartin et al. (2017). The orange line represents the July mean temperature (1961-2018) retrieved for the grid cell of Lago Verdarolo from Emilia-Romagna climate reanalysis ERACLITO (11 years running average). The shaded orange area is +/- one standard deviation. The full available period is reported on the left, while a zoom on the most recent period (since 1600 AD) is shown on the right.




Figure 9. Monthly mean, 2m temperature (a), accumulated precipitation (b), maximum daily precipitation (c), for 1961-1990 (blue) and 1991-2018 (green) computed over Lake Moo site grid cell of Eraclito reanalysis (grid cell 258, Ferriere Municipality). Confidence intervals at 95% significance (black vertical segments) are computed with a bootstrapping method (1000 iterations) as part of the Seaborn python library.

5.1. Holocene flood activity, paleotemperature changes and HIP events

485 A positive link emerges correlating Holocene (July) paleotemperatures, precipitation intensity
reconstructions and flood history (Fig. 7) at the study site, as the stratigraphic units showing the highest
frequency of coarse-grained flood deposits (i.e., Units 2, 4 and 5) invariably fall into periods of higher
temperatures (i.e., HTM and the post-LIA time). This strongly supports the hypothesis that greater warmth
favours the occurrence of extreme precipitation events, probably more frequent at the end of the
490 summer/autumn as already reported by other authors (e.g., Marcott et al., 2013; Giguët-Covex et al., 2012).
Although the process attribution and the geographical uniformity of the HTM wet phase are still matter of
debate, the abundance of coarse-grained flood deposits within the Lake Moo record around 9.5-5.5 kyr BP
suggests a precipitation increase also identified in other sites of the central and southern Mediterranean (Magny
et al., 2012b).

495 However, local factors such as changes in vegetation cover (i.e., tree cover percentage) due to human
disturbance and fires could have significantly contributed to enhancing slope erosion and then influenced the
Lake Moo sedimentary record, partly weakening our reconstruction of the temperature-HIP-flood activity
relationship at the Holocene scale. In order to overcome this issue, we explored and compared the pollen and
microcharcoal content of two key intervals (I₁ and I₂ in Figs. 7, 10) that are characterised by a different flood
500 record (high *versus* low flood activity) formed during consecutive periods of distinct climate conditions: the
HTM and the following cooling. The difference in mean temperature between I₁ and I₂, obtained averaging the
corresponding chronological intervals in the Verdarolo paleotemperature curve, is +1.3°C while the maximum
difference reaches +3.1°C (Fig.8).

For what concerns the post-LIA interval, and in particular since the 1950 AD, no significant
505 contribution on debris mobilisation can be attributed to vegetation landscape changes along the slopes as the
Lake Moo is a protected area characterised by dense forests (Corticelli et al., 2011; section 2.1.). Consistently,
the most recent pollen sample (P14 yielding an age of 146-14 cal yr BP; Fig. 7) shows a tree cover of about
80%.

The main considerations arising from the in-depth analysis of I₁ and I₂ are the following:

510

- I₁ (ca. 9.5-7 kyr BP; 10.5-9 m core depth): the woody component (i.e., tree cover %) is invariably
higher than the herbaceous one and it shows a value of 80% on average. The Conifers, mainly
represented by pines and silver fir, are dominant with respect to the deciduous trees (e.g., *Corylus*
avellana and *Quercus*), reaching a maximum of 95% within the P4 sample dated between 9.3-9.1 cal
515 kyr BP. Interestingly, *Abies alba* shows the highest percentages (ca. 15-55%) within the chronological
interval that corresponds to the warmest period of the HTM (Figs. 7, 10; Samartin et al., 2017).
Moreover, *A. alba* reaches the peak of ca. 55% during the HTM wettest conditions (around 7.5-7.3 cal
kyr B.P.; I₁ humid period in Fig. 7), as reconstructed by several authors (Regatieri et al., 2014;
Zanchetta et al., 2011; Zhornyak et al., 2011) from the Apuan Alps speleothems. This trend is
520 consistent with the fact that *A. alba* is a warm-temperature tree with a preference for high moisture
availability (Tinner et al., 2013), as clearly documented within the nearby Lake Greppo record

(Vescovi et al., 2010, confirming higher rainfall accumulation already highlighted by lake-level dynamics from Jura Mountains and central-northern Italy (Magny et al., 2004, 2009, 2012). Although no specific inferences can be made on past water levels of the Lake Moo, it is interesting to note that the low amount of hygrophytes and aquatics (Fig. 7) is fully consistent with highstand conditions. Anthropogenic indicators taxa (Fig. 7), as well as the Anthropogenic Index (AI; Fig. 10), show very low values throughout the interval, documenting a negligible human impact on the area in accordance with the extremely low content in microcharcoals (0.01-0.13 mm²/g; Fig. 10). Besides, the coarser microcharcoals (>125 micron) are not recorded suggesting any significant local fires. A slightly higher amount of anthropogenic indicator taxa occurs within the sample P6, dated between 8070-7880 ca yr BP; this is interpreted as an effect of categorisation difficulties about spontaneous taxa as *Artemisia vulgaris* type, *Chenopodium*, *Plantago*, *Urtica*, etc.

Palynological data, combined with the stratigraphic interpretation of the S1 core, support our hypothesis that warmer and wetter conditions typical of the HTM led to a high component of HIP and then high flood activity in the basin, independently by local factors. The presence of an extensive vegetation cover along the slopes implies the need of even more intense rainfall to trigger a debris mobilisation that leads to the deposition of thick, coarse-grained layers containing pebbles of remarkable diameters (LM6, LM7 and LM8 in Fig. 10).

- I₂ (ca. 5.5-3.8 kyr BP.; 5.5-4.5 m core depth): tree cover reduction (ranging between ca. 87.4-45.1 %; Figs. 7, 9), in favour of grassland expansion, characterises this interval that sees a fall in *A. alba* percentages (commonly less than 10%) and a high amount of meadow taxa (mainly *Poaceae*, *Cichorioideae* and *Asterioideae*; Fig. 7). The growth in hygrophytes and aquatics (up to ca. 18.2%) is consistent with a marginal lacustrine environment and then a relative lowering of the water level. Both anthropogenic indicator taxa and AI increase, up to ca. 7.3% and 16.2% respectively (Figs. 7, 9), suggesting a moderate degree of human impact on the vegetation cover (i.e., tree cover changes). This impact is reasonably due to pastoral activities more than agriculture as few grains of cereals are locally encountered (less than 4%). Accordingly, the meadow (pasture) taxa are abundant and coprophilous fungi and others spores (*Sporormiella*, *Dicrocoelium* and *Ascaris*; Figs. 7, 9) are sporadically found. By contrast, microcharcoal remains very low in amount, pointing to a little fire activity.

Although pollen-derived vegetation features (i.e., decrease in tree cover and a moderate degree of human impact) document local conditions more favorable for the debris flows activation with respect to those characterising I₁, this interval shows low flood activity with few and thin (<5 cm) sandy deposits (LM9 to LM10 in Fig. 10). Integrating the I₂ stratigraphic record with pollen data, such a low flood activity reasonably reflects low frequency (and intensity) of HIP events during a cooler and less humid period (Figs. 7, 9).

The paleoenvironmental-paleovegetation features characterising the Interval 2 persists upcore, in correspondence of Unit 3 that records apparent flood inactivity (Figs. 3, 7). The uppermost sample P14 likely tracks a strong increment of *A. alba*.




Figure 10. Sedimentological and palynological features of two key stratigraphic intervals (Intervals 1 and 2) from the reference core S1. These features are compared with the most relevant paleoclimate data available from literature for the area of interest (for references, please see Figure 7). The Anthropogenic Index (AI) has been calculated as: (Anthropogenic indicator/tree percentage) * 100 following Accorsi et al. (1999). Asterisks point to samples containing coprophilous fungi and other spores like *Sporormiella*, *Dicrocoelium* and *Ascaris*. Core S1 sample ages derive from the age-depth model reported in Figure 6. P samples correspond to the pollen samples; LM labels refer to the lithofacies explained in text (section 4.1.) and in Figures 3-4; dmax corresponds to the maximum diameter.

6. Conclusions

Placing present climate conditions into a geological perspective, beyond the instrumental era, is essential for understanding changes in the hydrological cycle induced by anthropogenic climate warming, complementing projections of model simulations. Physical reasons and regional climate reconstructions are consistently pointing to an increase in precipitation intensity when water vapour is not limited. In the last decade, record-breaking rainfall events have frequently occurred around the world. This trend is emerging with variable strength in different areas. Therefore, in the attempt to consolidate confidence about the extreme precipitation trend, we extended our analysis, passing from the instrumental to the geological time scale (i.e., the Holocene period). This choice implies a robust multidisciplinary approach, which includes, in addition to climate and meteorological data, proxy and expertise coming from the geological, geomorphological, stratigraphic and palynological-microanthracological area.

The major outcomes of our work can be summarised as follows:

1) The matching of instrumental temperature data and paleoenvironmental data allow us to affirm that current summer temperature is comparable with the temperature recorded in the Northern Apennine during the HTM. The current temperature trend computed from the monthly values of July temperature at Lake Verdarolo site over the period 1961-2018 is $+4.3^{\circ}\text{C}/100\text{y}$ (highly statistically significant), and it is likely to be one the highest recorded in the Holocene;

2) The stratigraphic units showing a high frequency of coarse-grained flood deposits are Units 2, deposited in the HTM, and Units 4 and 5 which belongs to the post-LIA time. Both periods are characterised by higher temperatures. The human impact is almost absent in the HTM and very low in the post-LIA time, especially in the last part of the 20th century. This strongly supports the hypothesis that a greater warmth favours the occurrence of extreme precipitation events. In particular, the recent sedimentation rate computed since c.a. 1800 AD (post-LIA) is at least $1\text{ m}/100\text{yr}$. This high value of the sedimentation rate has to be correlated to the presence of numerous coarse-grained levels within Units 4 and 5, after a period of absence of floods deposits in unit 3b. This difference in flood layers must be linked to an increase of HIP over the Lake Moo basin since we could not attribute it to any other changes in physiographic and vegetation dynamics (i.e., tree cover). During HTM and current time, the forested area coverage around Lake Moo persisted on high values, and this factor is considered disadvantageous for debris mobilization. On the contrary, during about 6-4 kyr BP we did not find any increase of coarse-grained layers from flood events, even though the forested area declined significantly and the anthropogenic impact increased;

3) HIP increase in response to higher temperature is already detectable in observation series. We found evidence that this also occurred in the past, especially during HTM testified by higher deposition of coarse-grained levels. A comparison with the past helps to understand future projections on the area. However, we are aware that past evolution cannot be taken as a perfect analogy for the future due to the different forcing and consequent response of the climate system (D'Agostino et al., 2019). As temperature will continue to increase in the Mediterranean area, precipitation intensity would keep increasing over the Northern Apennines. We hypothesise that precipitation intensity increase will be more evident in months with cooler and moist air masses, like in Autumn and in Winter, when moisture availability is not limited. An increase in precipitation maxima in Autumn months is already emerging on the Northern-Apennine.

Lake Moo basin proves to be an ideal study area to achieve a reconstruction between high-intensity precipitation and debris flow, for its position to the dominant atmospheric flow and favourable geological, geomorphological and vegetation characteristics. Further investigations, in association with other analogous sites along the Northern Apennine crest, are planned to provide a multisite assessment on the dynamics of past extreme precipitation events.

625 **Data availability:** Original data concerning pollen and radiometric data are available on the open data repository of ARPAE Emilia-Romagna (<https://arpaepv.datamb.it/dataset/lake-moo>)

Author contribution: SS and FG developed the idea and the research activity planning and execution. SN, MA, MTDN have taken care of the administrative aspects. All authors, except for SN and MTDN, have
630 contributed to field activity. Granulometric analysis has been conducted by AC. Pollen analysis have been conducted by SM and MM, data elaboration and figures were performed by VR, SM and MM. SS and FG and VR prepared the manuscript with contributions from all co-authors. FG management and coordination.

Competing interests: The authors declare that they have no conflict of interest.

635

Acknowledgements

This research was funded by Regional Agency of Civil Protection of the Emilia-Romagna Region and Geological, Seismic and Soil Survey of Emilia-Romagna Region in the framework of the cooperation agreement with ARPAE-SIMC. We are very grateful for all comments received in the public discussion and
640 in particular, we thank reviewers Picotti and Tinner for their useful suggestions. Finally, we thank Francesca Staffilani (Geological, Seismic and Soil Service, Emilia-Romagna Region) for the pedological data interpretation.

References

- 645 Accorsi, C.A., Bandini Mazzanti, M., Forlani, L., Mercuri, A.M., Trevisan Grandi, G.: An overview of Holocene forest pollen flora/vegetation of the Emilia Romagna region - Northern Italy in *Archivio Geobotanico*, 5 (1-2), 3-27, 1999.
- Ahlborn, M., Armon, M., Ben Dor, Y., Neugebauer, I., Schwab, M.J., Tjallingii, R., Shoqeir, J.H., Morin, E., Enzel, Y., and Brauer, A.: Increased frequency of torrential rain storms during a regional late Holocene eastern Mediterranean drought, *Quaternary Research*, 89, 425-431, <https://doi.org/10.1017/qua.2018.9>, 2018.
- 650 Anselmetti, F., Wirth, S.B., Glur, L., and Gilli, A.: Holocene flood frequency as reconstructed by lake sediments from multiple archives: A record influenced by solar forcing and atmospheric circulation patterns, in: *Late Pleistocene and Holocene climatic variability in the Carpathian-Balkan region*, Abstracts Volume, edited by: Mindrescu, M., and Gradinaru, I., Special Issue, 24, 1-2, 2014.
- Antolini, G., Auteri, L., Pavan, V., Tomei, F., Tomozeiu, R. and Marletto, V.: A daily high-resolution gridded climatic
655 data set for Emilia-Romagna, Italy, during 1961–2010. *Int. J. Climatol.*, 36: 1970-1986, doi:10.1002/joc.4473, 2016.
- Ballesteros-Cánovas, J.A., Stoffel, M., St. George, S., and Hirschboeck, K.: A review of flood records from tree rings, *Progress in Physical Geography*, 1-23, doi 10.1177/0309133315608758, 2015.
- Berg, P., Moseley, C., and Haerter, J.O.: Strong increase in convective precipitation in response to higher temperatures. *Nature Geosci.* 6, 181–185, <https://doi.org/10.1038/ngeo1731>, 2013.
- 660 Berglund, B.E., and Ralska-Jasiewiczowa, M.: Pollen analysis and pollen diagrams, in *Handbook of Holocene Palaeoecology and Palaeohydrology*, edited by: B.E. Berglund, Chichester, 455 - 484, 1986.

- Beug, H.J.: Leifaden der Pollenbestimmungen für Mitteleuropa und angrenzende Gebiete, Pfeil, München, 2004.
- Brönnimann, S., Rohr, C., Stucki, P., Summermatter, S., Bandhauer, M., Barton, Y., Fischer, A., Froidevaux, P.,
665 Germann, U., Grosjean, M., Hupfer, F., Ingold, K., Isotta, F., Keiler, M., Martius, O., Messmer, M., Mülchi, R.,
Panziera, L., Pfister, L., Raible, C.C., Reist, T., Rössler, O., Röthlisberger, V., Scherrer, S., Weingartner, R., Zappa,
M., Zimmermann, M. Zischg, A.P.: 1868 – L’alluvione che cambiò la Svizzera: Cause, conseguenze e insegnamenti
per il futuro. *Geographica Bernensia*, G94, 52 S., doi:10.4480/GB2018.G94.02, 2018a.
- Brönnimann, S., Rajczak, J., Fischer, E. M., Raible, C. C., Rohrer, M., and Schär, C.: Changing seasonality of moderate
and extreme precipitation events in the Alps. *Natural Hazards and Earth System Sciences*, 18 (7), 2047-2056.
670 <https://doi.org/10.5194/nhess-18-2047-2018>, 2018b.
- Brugnara, Y., and Maugeri, M.: Daily precipitation variability in the southern Alps since the late 19th century. *Int J
Climatol.*; 39: 3492– 3504. <https://doi.org/10.1002/joc.6034>, 2019
- Cacciamani, C., Cesari, D., Grazzini, F., Paccagnella, T., and Pantone, M.: Numerical Simulation of Intense Precipitation
Events South of the Alps: Sensitivity to Initial Conditions and Horizontal Resolution. *Meteorol Atmos Phys* 72, 147–
675 159. <https://link.springer.com/article/10.1007/s007030050012>, 2000
- Carton, A. and Panizza, M. (Eds.): Il paesaggio fisico dell’Alto Appennino Emiliano, Grafis Edizioni, Bologna, Italia,
1988.
- Clark, R.L.: Point count estimation of charcoal in pollen preparations and thin sections of sediments, pollen et spores, 24,
523-53, ISSN 0375-9636, 1982.
- 680 Combourieu-Nebout, N., Peyron, O., Bout-Roumazeilles, V., Goring, S., Dormoy, I., Joannin, S., Sadori, L., Siani, G.,
and Magny, M.: Holocene vegetation and climate changes in the central Mediterranean inferred from a high-resolution
marine pollen record (Adriatic Sea), *Clim. Past*, 9, 2023-2042. <https://doi.org/10.5194/cp-9-2023-2013>, 2013.
- Corticelli, S., Garberi, M.C., Mariani, M.C., and Masi, S.: Uso del suolo 2008, [http://geoportale.regione.emilia-romagna.it/it/download/dati-e-prodotti-cartografici-preconfezionati/pianificazione-e-catasto/uso-del-suolo-1/2008-](http://geoportale.regione.emilia-romagna.it/it/download/dati-e-prodotti-cartografici-preconfezionati/pianificazione-e-catasto/uso-del-suolo-1/2008-coperture-vettoriali-uso-del-suolo-edizione-2011)
685 [coperture-vettoriali-uso-del-suolo-edizione-2011](http://geoportale.regione.emilia-romagna.it/it/download/dati-e-prodotti-cartografici-preconfezionati/pianificazione-e-catasto/uso-del-suolo-1/2008-coperture-vettoriali-uso-del-suolo-edizione-2011), 2011.
- D’Agostino, R., Bader, J., Bordoni, S., Ferreira, D., and Jungclaus, J.: Northern Hemisphere monsoon response to mid-
Holocene orbital forcing and greenhouse gas-induced global warming. *Geophysical Research Letters*, 46, 1591– 1601.
<https://doi.org/10.1029/2018GL081589>, 2019.
- Elter, P., Ghiselli, F., Marroni, M., and Ottria, G. (Eds.): Note illustrative del Foglio 197 “Bobbio” della Carta Geologica
690 d’Italia alla scala 1:50.000. Istituto Poligrafico e Zecca dello Stato, Roma, Italia, 1997.
- Faegri, K., and Iversen, J.: Textbook of Pollen analysis, K. Faegri, P.E. Kaland, K. Krzywinski (Eds.), Chichester, 1989.
- Finsinger, W., Tinner, W., and Hu, F.S.: Rapid and accurate estimates of microcharcoal in pollen slide. Fiorentino, G.,
Magri D. (Eds.), Charcoals from the Past, BAR int Ser., pp. 121-124, 2008.
- Frei, C. and Schär, C.: A precipitation climatology of the Alps from high-resolution rain-gauge observations. *Int. J.
695 Climatol.*, 18: 873-900. doi:10.1002/(SICI)1097-0088(19980630)18:8<873::AID-JOC255>3.0.CO; 2-9, 1998.
- Geological, Seismic and Soil Service of the Emilia-Romagna Region: Landslide characteristics in Emilia-Romagna, 2012
(data available at: [https://ambiente.regione.emilia-romagna.it/en/geologia/temi/dissesto-idrogeologico/le-](https://ambiente.regione.emilia-romagna.it/en/geologia/temi/dissesto-idrogeologico/le-caratteristiche-dei-fenomeni-franosi-in-emilia-romagna)
[caratteristiche-dei-fenomeni-franosi-in-emilia-romagna](https://ambiente.regione.emilia-romagna.it/en/geologia/temi/dissesto-idrogeologico/le-caratteristiche-dei-fenomeni-franosi-in-emilia-romagna), last access: 04 April 2020).
- Giguet-Covex, C., Arnaud, F., Dirk, E., Jérôme, P., Laurent, M., Pierre, F., Fernand, D., Pierre-Jérôme, R., Bruno, W.,
700 and Jean-Jacques, D.: Frequency and intensity of high-altitude floods over the last 3.5ka in northwestern French Alps
(Lake Anterne), *Quaternary Research*, 77, 12-22, <https://doi.org/10.1016/j.yqres.2011.11.003>, 2012.

- Gilli, A., Anselmetti, F.S., Glur, L., and Wirth S.B.: Lake Sediments as Archives of Recurrence Rates and Intensities of Past Flood Events, in: *Dating Torrential Processes on Fans and Cones, Advances in Global Change Research*, edited by: Schneuwly-Bollschweiler, M., Stoffel, M., and Rudolf-Miklau, F., Springer Science Business Media, Dordrecht, Holland, 225-242, doi 10.1007/978-94-007-4336-6_15, 2013.
- 705 Glur, L., Wirth, S.B., Büntgen, U., Gilli, A., Haug, G.H., Schär, C., Beer, J., and Anselmetti, F.S.: Frequent floods in the European Alps coincide with cooler periods of the past 2500 years, *Nature Scientific Reports*, 3, doi 10.1038/srep02770, 2013.
- Grazzini, F., Craig, J.G., Keil, C., Antolini, G., and Pavan V.: Extreme precipitation events over Northern Italy: Part (I) a systematic classification with machine learning techniques, *QJR Meteorol Soc.* 2020;1–17. <https://doi.org/10.1002/qj.3635>, 2020.
- 710 Grazzini, F., Segadelli, S., and Fornasiero, A.: Precipitazioni estreme ed effetti al suolo sul reticolo minore: il caso del 14 Settembre 2015. Technical report ARPAE-SIMC/SGSS Emilia-Romagna, 27pp, Available in Italian https://www.arpae.it/dettaglio_notizia.asp?id=8017&idlivello=32, 2016.
- 715 Grazzini, F., Pratizzoli, W. and Tomei, F.: Drought and big snow, are extreme events the new normal? *Ecoscienza*, 1, Available in Italian https://www.arpae.it/dettaglio_documento.asp?id=3674&idlivello=1171, 2012.
- Heiri, O., S.J., Brooks, H.J.B., Birks, and Lotter, A.F.: A 274-lake calibration data-set and interference model for chironomid-based summer air temperature reconstruction in Europe, *Quaternary Science Reviews* 30, 3445-3456, 2011.
- 720 Henne, P.D., Bigalke, M., Büntgen, U., Colombaroli, D., Conedera, M., Feller, U., Frank, D., Fuhrer, J., Grosjean, M., Heiri, O., Luterbacher, J., Mestrot, A., Rigling, A., Rössler, O., Rohr, C., Rutishauser, T., Schwikowski, M., Stampfli, A., Szidat, S., Theurillat, J.P., Weingartner, R., Wilcke, W., and Tinner, W.: An empirical perspective for understanding climate change impacts in Switzerland. *Reg Environ Change* 18, 205–221, <https://doi.org/10.1007/s10113-017-1182-9>, 2018.
- 725 Hussain, M., and Mahmud, I.: pyMannKendall: a python package for non parametric Mann Kendall family of trend tests, *Journal of Open Source Software*, 4, (39), 1556, <https://doi.org/10.21105/joss.01556>, 2019.
- Hungr, O., Leroueil, S., and Picarelli, L.: The Varnes classification of landslide types, an update. *Landslides*, 11, 167–194, <https://doi.org/10.1007/s10346-013-0436-y>, 2014.
- Isotta, F., Frei, C., Weilguni, V., Percec Tadic, M., Lassègues, P., Rudolf, B., Pavan, V., Cacciamani, C., Antolini, G., Ratto, S.M., Munari, M., Micheletti, S., Bonati, V., Lussana, C., Ronchi, C., Panettieri, E., Marigo, G., and Vartacnik, G.: The climate of daily precipitation in the Alps: development and analysis of a high-resolution grid dataset from pan-Alpine rain-gauge data, *International Journal of Climatology*, 34, 1657-1675, doi 10.1002/joc.3794, 2014.
- 730 Krichak, S.O., Barkan, J., Breitgand, J.S., Gualdi, S., and Feldstein, S.: The role of the export of tropical moisture into midlatitudes for extreme precipitation events in the Mediterranean region. *Theor Appl Climatol* 121, 499–515. <https://doi.org/10.1007/s00704-014-1244-6>, 2015.
- 735 Lehmann, J., Coumou, D., and Frieler, K.: Increased record-breaking precipitation events under global warming, *Climatic Change* 132: 501, <https://doi.org/10.1007/s10584-015-1434-y>, 2015.
- Longman, J., Ersek, V., Veres, D., and Salzmann, U.: Detrital events and hydroclimate variability in the Romanian Carpathians during the mid-to-late Holocene, *Quaternary Science Reviews*, 167, 78-95, 2017.
- 740 Libertino, A., Ganora, D., and Claps, P.: Evidence for increasing rainfall extremes remains elusive at large spatial scales: The case of Italy. *Geophysical Research Letters*, 46, 7437– 7446. <https://doi.org/10.1029/2019GL083371>, 2019.

- Lowe, D.R.: Sediment gravity flows; Depositional models with special reference to the deposits of high-density turbidity currents, *Journal of Sedimentary Petrology*, 52, 279-297, 1982.
- 745 Lowe, J.J., Accorsi, C.A., Bandini Mazzanti, M., Bishop, V., Van der Kaars, S., Forlani, L., Mercuri, A.M., Rivalenti, C., Torri, P., Watson, C.: Pollen stratigraphy of sediment sequences from carter lakes Albano and Nemi (near Rome) and from the central Adriatic, spanning the interval from oxygen isotope stage 2 to the present day, *Mem. Ist. Ital. Idrobiol.* 55, 71-98, 1996.
- Magny, M.: Holocene climate variability as reflected by mid-European lake-level fluctuations and its probable impact on prehistoric human settlements, *Quaternary International*, 113, 65-79, 2004.
- 750 Magny, M., Galop, D., Bellintani, P., Desmet, M., Didier, J., Hass, J.N., Martinelli, N., Pedrotti, A., Scandolari, S., Stock, A., and Vannièrè, B.: Late-Holocene climatic variability south of the Alps as recorded by lake-level fluctuations at Lake Ledro, Trentino, Italy. *Holocene*, 19, 575-589, 2009.
- Magny, M., S. Joannin, D. Galop, B. Vannièrè, J. N. Haas, M. Bassetti, P. Bellintani, R. Scandolari, and M. Desmet. 2012. Holocene palaeohydrological changes in the northern Mediterranean borderlands as reflected by the lake-level record of Lake Ledro, northeastern Italy. *Quaternary Research*, 77, 382-396, 2012.
- 755 Magny, M., Peyron, O., Sadori, L., Ortu, E., Zanchetta, G., Vannièrè, B. and Tinner, W.: Contrasting patterns of precipitation seasonality during the Holocene in the south- and north-central Mediterranean. *J. Quaternary Sci.*, 27: 290-296. doi:10.1002/jqs.1543, 2012b.
- Marchetti, G., and Fraccia, R.: Carta geomorfologica dell'alta Val Nure, Appennino piacentino. Scala 1:25.000, in: *Il paesaggio fisico dell'Alto Appennino Emiliano*. Regione Emilia-Romagna, Carton, A., and Panizza, M., (Eds.), Grafis Edizioni, Bologna, Italia, 1988.
- 760 Marcott, S., Shakun, J., Clark, P.U., and Mix, A.C.: A Reconstruction of Regional and Global Temperature for the Past 11,300 Years, *Science*, 339, 1198-1201, doi 10.1126/science.1228026, 2013.
- Marroni, M., Meneghini, F., and Pandolfi, L.: Anatomy of the Ligure-Piemontese subduction system: evidence from Late Cretaceous-middle Eocene convergent margin deposits in the Northern Apennines, Italy, *International Geology Review*, iFirst article, 1-33, 2010.
- 765 Milliman, J.D. and Syvitski, J.P.M.: Geomorphic/Tectonic Control of Sediment Discharge to the Ocean: The Importance of Small Mountainous Rivers, *The Journal of Geology*, 100, 525-544, <http://dx.doi.org/10.1086/629606>, 1992.
- Moore, P.D., Webb, J.A., and Collinson, M.E.: *Pollen Analysis*, Oxford, 1991.
- 770 Mulder, T. and Syvitski, J.P.M.: Turbidity Currents Generated at River Mouths during Exceptional Discharges to the World Oceans, *Journal of Geology*, 103, 285-299, <http://dx.doi.org/10.1086/629747>, 1995.
- Mutti, E.: *Turbidite Sandstones*. San Donato Milanese, Agip-Istituto di Geologia, Università di Parma, 275 pp, 1982.
- Mutti, E., Davoli, G., Tinterri, R., and Zavala, C.: The importance of ancient fluvio-deltaic systems dominated by catastrophic flooding in tectonically active basins, *Memorie di Scienze Geologiche*, Università di Padova, 48, 233-291, 1996.
- 775 Myhre, G., Alterskjær, K., Stjern, C.W., Hodnebrog, Ø., Marelle, L., Samset, B.H., Sillmann, J., Schaller, N., Fischer, E., Schulz, M., and Stohl, A.: Frequency of extreme precipitation increases extensively with event rareness under global warming. *Scientific Reports* 9:16063, <https://doi.org/10.1038/s41598-019-52277-4>, 2019.
- Oswald, W.W., Anderson, P.M., Brown, T.A., Brubaker, L.B., Hu, F.S., Lozhkin, A.V., Tinner, W., and Petra Kaltenrieder, P.: Effects of sample mass and macrofossil type on radiocarbon dating of arctic and boreal lake sediments. *The Holocene* 15, 5, 758-767, 2005.
- 780

- Papalexiou, S. M. and Montanari, A.: Global and regional increase of precipitation extremes under global warming. *Water Resources Research*, 55, 4901–4914. <https://doi.org/10.1029/2018WR02406>, 2019.
- 785 Patterson, W.A., Edwards, K.J., and Maguire, D.J.: Microscopic charcoal as a fossil indicator of fire, *Quaternary Science Reviews*, 6, 3-23, [https://doi.org/10.1016/0277-3791\(87\)90012-6](https://doi.org/10.1016/0277-3791(87)90012-6), 1987.
- Pignatti, S.: *Flora d'Italia, Edagricole-New Business Media, Milano*, 2017-2019.
- Prein, A.F., Rasmussen, R.M., Ikeda, K., Liu, C., Clark, M.P., and Holland, G.J.: The future intensification of hourly precipitation extremes, *Nature Climate Change*, 7, 48-52, doi:10.1038/nclimate3168, 2017.
- 790 Regattieri, E., Zanchetta, G., Drysdale, R.N., Isola, I., Hellstrom, J.C., and Dallai, L.: Lateglacial to Holocene trace element record (Ba, Mg, Sr) from Corchia Cave (Apuan Alps, central Italy): paleoenvironmental implications, *Journal of Quaternary Science*, 29, 381-392, <https://doi.org/10.1002/jqs.2712>, 2014.
- Reimer, P.J., Bayliss, A., Warren Beck, J., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, 795 E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP, *Radiocarbon*, 55, 1869-1887, https://doi.org/10.2458/azu_js_rc.55.16947, 2013.
- Renssen, H., Seppä, H., Crosta, X., Goosse, H., and Roche, D.M.: 2012. Global characterisation of the Holocene Thermal Maximum. *Quaternary Science Reviews* 48, 7-19, 2012.
- 800 Samartin, S., Heiri, O., Joos, F., Renssen, H., Franke, J., Brönnimann, S., and Tinner, W.: Warm Mediterranean mid-Holocene summers inferred from fossil midge assemblages, *Nature Geoscience*, 10, 207-212. doi: 10.1038/NGEO2891, 2017.
- Scherrer, S. C., Fischer, E.M., Posselt, R., Liniger, M.A., Croci-Maspoli, M., and Knutti, R.: Emerging trends in heavy precipitation and hot temperature extremes in Switzerland, *J. Geophys. Res. Atmos.*, 121, 805 doi:10.1002/2015JD024634, 2016.
- Schillereff, D., Chiverrell, R.C., Macdonald, N., and Hooke, J.M.: Flood stratigraphies in lake sediments: A review, *Earth-Science Reviews*, 135, 17-37, doi 10.1016/j.earscirev.2014.03.011, 2014.
- Schneuwly-Bollschweiler, M., Stoffel, M., and Rudolf-Miklau, F. (Eds.): *Dating Torrential Processes on Fans and Cones. Methods and Their Application for Hazard and Risk Assessment*. Springer, doi 10.1007/978-94-007-4336-6, 2013.
- 810 Segadelli S., Grazzini F., Adorni M., De Nardo M.T., Fornasiero A., Chelli A., and Cantonati M.: Predicting Extreme-Precipitation Effects on the Geomorphology of Small Mountain Catchments: Towards an Improved Understanding of the Consequences for Freshwater Biodiversity and Ecosystems. *Water* 12: 79, <https://doi.org/10.3390/w12010079>, 2020.
- Skinner, C. B., and Poulsen, C.J.: The role of fall season tropical plumes in enhancing Saharan rainfall during the African Humid Period, *Geophys. Res. Lett.*, 43, 349–358, doi:10.1002/2015GL066318, 2016.
- 815 Stoffel, M., Butler, D.R., and Corona, C.: Mass movements and tree rings: A guide to dendrogeomorphic field sampling and dating, *Geomorphology*, 200, 106-120, <https://doi.org/10.1016/j.geomorph.2012.12.017>, 2013.
- Stoffel, M., Wyzga, B., and Marston, R.A.: Floods in mountain environments: A synthesis, *Geomorphology*, 272, 1-9, <https://doi.org/10.1016/j.geomorph.2016.07.008>, 2016.
- 820 Swierczynski, T., Ionita, M., and Pino, D.: Using archives of past floods to estimate future flood hazards, *Eos* 98, <https://doi.org/10.1029/2017EO066221>, 2017.

- Tinner, W., Colombaroli D., Heiri, O., Henne, P., Steinacher, M., Untenecker, J., Vescovi, E., Allen, J., Carraro, G., Conedera, M., Joos, F., Lotter, A., Luterbacher, J., Samartin, S., and Valsecchi, V.: The past ecology of *Abies alba* provides new perspectives on future responses of silver fir forests to global warming. *Ecological Monographs* 83(4): 419-439, 2013.
- 825
- Tutin, T.G., Heywood, V.H., Burges, N.A., and Valentine, D.H.: *Flora Europaea*, Cambridge University Press, 1964-1993.
- Utsumi, N., Seto, S., Kanae, E., Maeda, E., and Oki T.: Does higher surface temperature intensify extreme precipitation?, *Geophys. Res. Lett.*, 38, L16708, doi:10.1029/2011GL048426, 2011.
- 830
- Vescovi, E., Ammann, B., Ravazzi, C., and Tinner, W.: A new Late-glacial and Holocene record of vegetation and fire history from Lago del Greppo, northern Apennines, Italy. *Vegetation History and Archaeobotany*, 19, 219–233, <https://doi.org/10.1007/s00334-010-0243-5>, 2010.
- Vittori Antisari, L., Bianchini, G., Cremonini, S., Di Giuseppe D., Falsone, G., Marchesini, M. Marvelli, S., and Vianello G.: Multidisciplinary study of a Lateglacial-Holocene sedimentary sequence near Bologna (Italy): insights on natural and anthropogenic impacts on the landscape dynamics. *J Soils Sediments* 16, 2 645-662, doi 10.1007/s11368-015-1266-4, 2015
- 835
- Westra, S., Fowler, H.J., Evans J.P., Alexander, L.V., Berg, P., Johnson, F., Kendon, E.J., Lenderink, G., and Roberts, N.M.: Future changes to the intensity and frequency of short duration extreme rainfall, *Reviews of Geophysics*, 52, 522-555, doi:10.1002/2014RG000464, 2014.
- 840
- Whitlock, C. and Larsen, C.: Charcoal as a fire proxy. In Smol J.P., Birks, H.J.B. and Last, W.M., (Eds.), *Terrestrial, algal, and siliceous indicators*, Dordrecht: Kluwer , 75-97, 2001.
- Whitlock, C., and Millspaugh, S.H.: Testing the assumptions of fire-history studies: an examination of modern charcoal accumulation in Yellowstone National Park, USA, *The Holocene*, 6, 7-15, <https://doi.org/10.1177/095968369600600102>, 1996.
- 845
- Wilhelm, B., Arnaud, F., Sabatier, P., Crouzet, C., Brisset, E., Chaumillon, E., Disnar, J.R., Guiter, F., Malet, E., Reyss, J.L., Tachikawa, K., Bard, E., and Delannoy, J.J.: 1400 years of extreme precipitation patterns over the Mediterranean French Alps and possible forcing mechanism, *Quaternary Research*, 78, 1-12, doi 10.1016/j.yqres.2012.03.003, 2012.
- Wilhelm, B., Ballesteros-Cànovas, J.A., Macdonald, N., Toonen, W.H.J., Baker, V., Barriandos, M., Benito, G., Brauer, A., Corella, J.P., Denniston, R., Glaser, R., Ionita, M., Kahle, M., Liu, T., Luetscher, M., Macklin, M., Mudelsee, M., Muñoz, S., Schulte, L., George, S.St., Stoffel, M., and Wetter, O.: Interpreting historical, botanical, and geological evidence to aid preparations for future floods, *WIREs Water*, 1-22, <https://doi.org/10.1002/wat2.1318>, 2018.
- 850
- Wirth, S.B.: *The Holocene Flood History of the Central Alps Reconstructed from Lacustrine Sediments: Frequency, Intensity and Controlling Climate Factors*. Ph.D. thesis, ETH Zürich Bibliography, Switzerland, 179pp, doi 10.3929/ethz-a-009775044, 2013.
- 855
- Wirth, S.B., Glur, L., Gilli, A., and Anselmetti, F.S.: Holocene flood frequency across the Central Alps – solar forcing and evidence for variations in North Atlantic atmospheric circulation, *Quaternary Science Reviews*, 80, 112-128, <https://doi.org/10.1016/j.quascirev.2013.09.002>, 2013.
- Zanchetta, G., Isola, I., Piccini, L., and Dini, A.: The Corchia Cave (Alpi Apuane) a 2 Ma long temporal window on the earth climate, *Technical Periodicals of National Geological Survey of Italy - ISPRA, Geological Field Trips*, 3, 55pp, doi 10.3301/GFT.2011.02, 2011.
- 860
- Zavala, C. and Pan, S.: Hyperpycnal flows and hyperpycnites: Origin and distinctive characteristics. *Lithologic Reservoirs*, 30, 1-27, doi 10.3969/j.issn.1673-8926.2018.01.001, 2018.

- 865 Zavala, C., Ponce, J.J., Arcuri, M., Drittanti, D., Freije, H., and Asensio, M.: Ancient Lacustrine Hyperpynites: A Depositional Model from a Case Study in the Rayoso Formation (Cretaceous) of West-Central Argentina. *Journal of Sedimentary Research*, 76, 41-59, <https://doi.org/10.2110/jsr.2006.12>, 2006.
- Zavala, C., Arcuri, M., Di Meglio, M., Gamero Diaz, H., and Contreras, C.: A Genetic Facies Tract for the Analysis of Sustained Hyperpynal Flow Deposits. in: *Sediment transfer from shelf to deep water - Revisiting the delivery system*, edited by: Slatt, R.M. and Zavala, C., *AAPG Studies in Geology*, 61, 31-51, 2011.
- 870 Zhornyak, L.V., Zanchetta, G., Drysdale, R.N., Hellstrom, J.C., Isola, I., Regattieri, E., Piccini, L., Baneschi, I., and Couchoud, I.: Stratigraphic evidence for a "pluvial phase" between ca 8200-7100 ka from Renella cave (Central Italy), *Quaternary Science Reviews*, 30, 409-417 doi 10.1016/j.quascirev.2010.12.003, 2011.