



## Changes in high intensity precipitation on the Northern Apennines (Italy) as revealed by multidisciplinary data over the last 9000 years

5 Stefano Segadelli<sup>1</sup>, Federico Grazzini<sup>2</sup>, Margherita Aguzzi<sup>2</sup>, Alessandro Chelli<sup>3</sup>, Veronica Rossi<sup>4</sup>,  
Maria T. De Nardo<sup>1</sup>, Roberto Francese<sup>3</sup>, Silvia Marvelli<sup>5</sup>, Marco Marchesini<sup>5</sup>, Sandro Nanni<sup>2</sup>

<sup>1</sup> Geological, Seismic and Soil Service, Emilia-Romagna Region Administration, Bologna, Italy.

<sup>2</sup> Regional Agency for Prevention, Environment and Energy of Emilia-Romagna, Hydro-Meteo-Climate Service (ARPAE-SIMC), Bologna, Italy.

10 <sup>3</sup> Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Italy.

<sup>4</sup> Department of Biological, Geological, and Environmental Sciences - BiGeA, Alma Mater Studiorum University of Bologna, Bologna, Italy.

<sup>5</sup> Laboratory of Palynology and Archaeobotany - C.A.A. Giorgio Nicoli, San Giovanni in Persiceto (Bologna, Italy).

15 *Correspondence to:* Federico Grazzini ([fgrazzini@arpae.it](mailto:fgrazzini@arpae.it))

**Abstract.** Several record-breaking precipitation events have stricken the mountainous area of Emilia-Romagna Region (northern Apennines, Italy) over the last years. As consequence, several geomorphological processes, like widespread debris flows along the slopes and hyperconcentrated flood in the stream channels, shallow landslides and overbank flooding affected the territory, causing serious damages to man-made structures. The intensity and wide spatial scale of these phenomena leads us to investigate their frequency in the past, beyond the instrumental time. A detailed study of these recent deposits compared with fossil peat bog and lake paleo deposits can provide useful insight to support a strong match between precipitation intensity and warm climatic phases in antecedent climatic periods, as expected by the increase air water vapour holding capacity at higher temperatures

20 Here we present the results of the field campaign performed in summer 2017 at Lake Moo a 0.15km<sup>2</sup> peat bog located at an altitude of 1130m a.s.l. The chosen area has been affected, during the flooding of the upper Trebbia and Nure valleys 13-14 September 2015, by several high-density flows generated by the stream that flow into the plain. Our main assumption is that, in such a small drainage basin (area <2 km<sup>2</sup>), with favourable geologic and geomorphic characteristics implying advantageous sediment transfer into lake, high density flood can be triggered only by high intensity precipitation events (HIP) lasting enough time for water to infiltrate and mobilize large quantities of debris.

30 The sedimentary succession (ca. 13 m-thick) was studied through the extraction of two cores and one trench. The facies/paleoenvironmental interpretation of the sedimentary succession, characterized by clusters of coarse-grained alluvial deposits interbedded with organic-rich silty clays and peaty layers, was achieved combining sedimentological and pollen data with pedological data and radiocarbon dating (AMS 14C).

35 Observed depositional cycles were put in relation with other specific paleoclimatic proxies available in literature for the North-Apennine area. This comparison illustrates that the increase of extreme paleo flood (associated with coarse-grained deposits similar to the ones observed recently) correlates well with warm phases with a maximum activity during the Holocene thermal maximum and from the small ice age to the present day.



40 *Keywords*

*extreme precipitation, Holocene flood history, northern Apennines, climate change, water cycle, global warming*

**1. Introduction**

45 The frequency of high intensity precipitation (HIP), also known as torrential rainstorm for their capacity to  
generate flash flood of small streams, is a key aspect of the water cycle of Mediterranean climate since a significant  
amount of annual precipitation is often concentrated in few major events (see for example Frei and Schär 1998 for the  
Alpine area). Therefore, the knowledge of the expected frequency (and their maximum intensity), in present and future  
climate, is a very important constraint for planning adequate hydraulic defences and water resources managing. In  
particular, on the northern Apennines, more than sixty percent of total precipitation of the year is concentrated in days  
50 with moderate to high-intensity precipitation (Isotta et al., 2014). Restricting the analysis to the Emilia-Romagna region  
(Northern-Italy; Fig. 1), over the last years there we have been observed an increase in interannual variability of torrential  
rainfall, with marked or even exceptional droughts such as those occurring in 2012 and 2017 (Grazzini et al., 2012),  
followed by years with record rainfall (2014 and 2018). Moreover, between September 2014 and September 2015, the  
region has been affected by three events of exceptional intensity, estimated to have return period of several centuries  
55 (Grazzini et al., 2016).

Under the threat of global warming, a growing number of studies are investigating the link between current  
temperature rise and precipitation intensity. A consistent increase in long-term trend of extreme daily precipitation has  
been already detected for the northern extratropics since 1980, although this is largely changing with the area of the globe  
considered (Lehmann et al. 2015, Papalexiou and Montanari 2019). On the Alpine domain Scherrer et al. (2016) found  
60 significant increases in daily extreme precipitation indices over Switzerland (1901-2014). Brugnara and Maugeri (2019),  
through the analysis of newly digitized data of the last 150 years across the whole Alpine area, found regional differences  
in the trend of the extremes suggesting that instrumental data are not covering a sufficient period of time to infer reliable  
changes in extreme events frequency. Although there is a consensus, based on physical arguments, that precipitation  
intensities are expected to increase with higher temperatures (see the review paper by Westra et al., 2014) limitations in  
65 moisture availability can prevent the potential for increased precipitation from being realized (Prein et al. 2017) explaining  
spatial and seasonal differences observed in the trends.

In this context, the stratigraphic record of current and fossil peat bogs and lakes can provide useful insights to  
verify/test the hypothesized strong linkage between precipitation intensity and warm phases during the present interglacial  
(i.e., Holocene). In literature there is ample documentation of the use of these sedimentary archives to reconstruct past  
70 flood events chronologies (Ahlborn et al., 2018; Anselmetti et al., 2014; Giguët-Covex et al., 2012; Gilli et al., 2013;  
Giraudi, 2014; Glur et al., 2013; Longman et al., 2017; Schillereff, 2014; Stoffel et al., 2016; Swierczynski et al., 2017;  
Wilhelm et al., 2018; Wirth, 2013; Wirth et al., 2013; Zavala and Pan, 2018; Zavala et al., 2006; Zavala et al., 2011),  
alongside others as tree rings (Ballesteros-Cánovas et al., 2015; Stoffel et al., 2013), speleothems (Regattieri et al., 2014;  
Zanchetta et al., 2011) and torrential fans and cones (Schneuwly-Bollschweiler et al., 2013).

75 Contrary to our hypothesis, some studies suggest a synchronization of high frequency of flood events with  
cooling periods. For example, Glur et al. (2013), presenting results from a multi-archive flood reconstruction based on  
sediments of ten Alpine lakes, found periods of high frequency of sediment deposition in concomitance with summer  
cool temperatures. However, those results need a careful interpretation since lake sediments depositions are influenced  
by combination of factors which involves precipitation intensity, duration, seasonality, changes in atmospheric-circulation



80 which might overshadow the physically based positive correlation between temperature increase and precipitation intensity (Utsumi et al., 2011; Brönnimann et al., 2018). Those caveats call for an even more multidisciplinary approach integrating paleoclimate (pollen), sedimentary archive information with climatological observations and physical arguments. In this study, we aim to fill this gap, extending and reinforcing observed trends with information from the past, beyond instrumental times, coherently derived from the same region during a specific field campaign.

85 The study area was chosen because, during the flash-flood of the upper Trebbia and Nure valleys 13-14 September 2015, it has been affected by several high-density floods generated by a small stream that flow into the plain. Our main assumption is that, in such a small drainage basin (area <2 km<sup>2</sup>), which has favourable geologic and geomorphic characteristics to achieve substantial sediment transfer into lake (Schillereff, 2014), high magnitude flood events can be reconducted only to HIP, necessary to mobilize large quantities of debris (Milliman and Syvitski, 1992; Mulder and  
90 Syvitski, 1995; Mutti et al., 1996). In the Trebbia-Nure case, a detailed analysis of precipitation (Grazzini et al., 2016) over Lake Moo site microbasin showed that the observed debris flow occurred with a peak intensity of 112mm/3h. This is a very high value compared to shallow landslide and debris flow thresholds found in literature for nearby areas, like the Apuane and Garfagnana regions (Giannecchini et al., 2012), which can be explained by the dense vegetation cover present now in the area. In addition, the absence of surrounding anthropic activities, the vicinity of the lake to the main Apennines  
95 crest, very exposed to maritime moist airflow coming from the central Mediterranean Sea, make this site particularly suitable for a detailed reconstruction of HIP events over the Holocene in terms of frequency, sedimentary expression and forcing factors.

Through the analysis of the data acquired in on the field campaign, in this paper we want to address the following research questions:


- 100 a) Could we consider these recent events unprecedented over longer time-scales?  
b) If yes, is the frequency of these events coupled with temperature variations?

The paper is organised as follows. The study area is presented in section 2 with a description of the basin, its morphology and vegetation. In sections 3 source of instrumental data and methods for collecting data during the fields campaign are described. In section 4 stratigraphic results of the field campaign are commented while in section 5 we develop a full  
105 multidisciplinary discussion. Finally, in section 6, conclusions are drawn.

## 2. The study area

### 2.1. Geographic and climatic context

110 The Lake Moo plain (44°37'29"N, 9°32'25"E) has a surface area of about 0.15Km<sup>2</sup>. It is located near the boundary between Emilia-Romagna and Liguria regions (Piacenza province, Italy), in the high valley of the Nure stream at an average altitude of 1130m a.s.l. (Fig. 1). Nowadays, the catchment area of the Lago Moo is characterized by a dense forest cover (total woodland cover of ca. 65.55%; Corticelli et al., 2011), with high vegetational richness and an exceptional concentration of protected mountain species with a high phyto geographical interest. The widespread presence of the *Fagus sylvatica* is locally interrupted by grazing land areas and blueberry moorland with the presence of the rare *Juniperus*  
115 *nana* and *Sorbus chamaemespilus*. Reforestation of *Pinus nigra* is also present (Table 1 and Fig. 1).



**Figure 1.** Location and photo of the Lake Moo site

120

| Tree type specie   | Cover area (km <sup>2</sup> ) | Woodland cover (%) | Forest cover classification in the catchment area of Lago Moo |
|--|-------------------------------|--------------------|---|
| <i>Fagus sylvatica</i> forest  | 1.17                          | 63.00              | Moderately dense forest                                       |
| Reforestation of <i>Pinus nigra</i> forest   | 0.02                          | 1.00               | Scrub   |
| Mixed wood forest ( <i>Pinus nigra</i> , <i>Fagus sylvatica</i> and <i>Carpinus</i> genus) | 0.03                          | 1.55               | Scrub   |
| Total  | 1.22                          | 65.55              | Moderately dense forest                                       |

**Table 1** - Actual vegetation cover of Lake Moo area deduced by Corticelli et al., 2011



The climate of the area is characterised by the altitude and the mountain range, with enhanced humidity given the short distance to the Tyrrhenian Sea (Nistor, 2016). The months with the largest amount of precipitation are in the transition seasons (Spring and Autumn), with a predominant peak in early autumn (see Fig. 9). HIP conditions are very favourable in Autumn months due to a particular synergy of higher frequency of synoptic disturbances and mesoscale convective systems which could still develop in high thermodynamic unstable environment (Grazzini et al., 2019).

## 2.2. Geological and geomorphological setting

The study area consists mainly of strongly serpentinized ultramafites extensively fractured and representing the original oceanic crust of the Ligure-Piemontese basin, developed in the Middle to Upper Jurassic, which separated the Europe plate from the Adria plate (Marroni et al., 2010). Due to the compositional heterogeneity of its vertical sequence, the ultramafic medium is made of lithological units, tectonically overlapped. The ultramafites are bordered by deposits which are predominantly characterized by polygenic breccias (Casanova complex, early Campanian age, Vescovi, 2002), made of blocks of limestones or marly limestones immersed in a fine-grained matrix or a mineral cement.

The morphology of the area includes flat areas and steep slopes at different altitude that some authors (Elter et al., 1997; Marchetti and Fraccia, 1988; Carton and Panizza, 1988) interpret as originated from the phase of last glacial retreat. On these morphological flat areas, marsh deposits have developed originated from the filling of small glacial lakes some of which still exist as the Lago Moo plain and Bino lake. Other authors such as the Geological, Seismic and Soil Service of the Emilia-Romagna Region - Inventory Map of landslides at 1:10000 scale of the Emilia-Romagna Region (available at: <http://ambiente.regione.emilia-romagna.it/geologia/temi/dyسة-hydrogeological/the-paper-inventory-of-landslides>, 2019) have interpreted the Moo lake plain as the product of gravitational processes that affect the entire slope area (available at: <http://ambiente.regione.emilia-romagna.it/geologia/temi/disruption-hydrogeological/the-paper-inventory-of-landslides>, 2019). From this point of view, the main factor controlling the morphological evolution of the area of Lake Moo, is the river incision produced by the Nure stream during the Holocene (Gunderson et al., 2014). This incision triggered the development of gravitational phenomena on the slopes. These complex morphological features of the area is also influenced by the presence of lithological elements with a strong mechanical contrast such as ophiolites superimposed on a predominantly clayey substrate (Elter et al., 1997).

## 3. Data and methods

### 3.1 Field investigation


The selection of the study site has been obtained following criteria indicated by Gilli et al. (2013) and Schillereff et al. (2014). In this respect, the Lake Moo site presents several advantageous characteristics to the archiving of paleo flood deposits:

- Steep relief (average inclination of 24°) with slopes composed of polygenic and monogenetic braccias in clay matrix, highly susceptible to erosion (Monte Ragola Complex, Elter et al., 1997) are present;
- No lakes upstream in catchment;
- Small drainage basin area (1.94 km<sup>2</sup>);
- One dominant inflow into the lake;
- Absence of regulated flow structures;
- Absence of natural pre-lake sediment storage zones.

The field campaign led to the acquisition of two sedimentary cores (S1 and S2) and one trench, investigating the sedimentary succession capped by the flood deposit formed during the last HIP event (14 September 2015). The location



of the cores and trench was planned on the basis of a high-resolution reflection seismic survey (Fig. SUP1 of the supplementary material) and a detailed geomorphological map, both of them originally produced for this purpose (Fig. 2). The former provided information on the lake basin floor morphology and the thickness of the infilling succession.



**Figure 2.** Location of the geognostic investigations, geophysical surveys tracks, and detailed geomorphological mapping of the flood deposit occurs between 13 and 14 September 2015 (upper Trebbia and Nure valleys, province of Piacenza, Italy). *See a larger version of this figure at the end of the manuscript*

The two cores S1 and S2, 14 and 6 meters long respectively, were performed through a continuous perforating system, which guaranteed an undisturbed core stratigraphy and a high recovery percentage (S1 90% and S2 91%). The trench, carried out in between the two cores, reached the depth of 2 meters, the length of 6 meters and the width of 6 meters. The subsurface succession was stratigraphically analysed, and the longest core (S1) was selected as reference and sampled for grain size, radiocarbon and palynological analysis.

### 3.2 Facies analysis and chronology

A facies characterization of the sedimentary succession has been performed on basis of the observable macroscopic physical characteristics (i.e., grain size, sedimentary structures, Munsell chart colour and types of bounding surfaces), and granulometric data, the latter available for the reference core S1.

Relatively to the different coarse-grained levels recognized, we follow the facies tract concept. A facies tract here means all facies observed within the same deposits relative of the same flood undergoing transformations along its down-slope motion (Lowe, 1982; Mutti, 1992; Mutti et al., 1996). The facies tract approach is important because allows to recognize the following important information like:

- 1) the position in the facies tract with respect to the whole fluvial-hyperpycnal system;
- 2) relative flood event magnitude;
- 3) expected related facies types in more proximal and distal areas.



190 The abundance of wood remains and peat levels in the sedimentary cores allowed  $^{14}\text{C}$  radiocarbon twelve samples that were analysed at the CEDAD Laboratory of the University of Salento (Province of Puglia, Italy). The data are collected in Table TS2 of the supplementary material. The  $^{14}\text{C}$  ages were converted into calendar years using the OxCal version 3.10 software (Reimer et al., 2013).

### 195 **3.3 Temperature reconstruction and modern climatological dataset**

Central to our analysis is the availability of a reliable temperature reconstruction for the chronological period explored through the coring. In this respect the chironomid analysis of the nearby Lake Verdarolo conducted by Samarti et al. (2017), represent a unique opportunity. Lake Verdarolo site is located at 1390m a.s.l., 270m higher and 54km south from Lake Moo (Fig. 1), in a very similar climatic context. They reconstructed the mean July air temperature using a chironomid-based inferred model developed from a combination of data of over 200 lakes from Norway and Swiss Alps (Heiri et al., 2011). This represent the first vegetation independent holocenic temperature reconstruction of the Northern Apennine.

200 Modern temperature and precipitation time series (1961-2018) of Lago Verdarolo and Lake Moo are derived from the gridded high-resolution dataset of Emilia-Romagna (Eraclito4), described in Antolini et al. (2016). Trend estimation and Mann Kendall significance trend test are computed with the pyMannKendall package (Hussain et al., 2019).

### 205 **3.4 Pollen analysis**

Palynological analyses were carried out on 14 samples collected from core S1 to refine facies characterization of fine-grained deposits and obtain vegetation-derived paleoclimate data (Fig. 3). After the radiocarbon dating, a further choice was made using only 11 samples falling into two stratigraphic intervals of our interest: the first one ( $I_1$ ), covering 10.77m to 9.33m depth and with temporal resolution 7300 - 9600 cal yr BP centred on the Holocene Thermal Maximum (HTM), the second one ( $I_2$ ) from 5.48m to 4.55 depth and corresponding to the exit of HTM (3900 - 5500 cal yr BP). In these two stratigraphic intervals, the mean sampling resolution is 28cm. The choice of these two periods ensure a significant thermal/climatic separation and at the same time the samples were taken from predominantly lacustrine fine layers where the stratigraphic series shows characteristics of continuity of sedimentation.

215 Identification of the pollen grains was performed at 1000 light microscope magnification and based on the atlases and a vast amount of specific morpho-palynological bibliography stored at the CAA Laboratory (S. Giovanni in Persiceto, Italy). Names of the families, genus and species of plants conform to the classifications of Italian Flora proposal by Pignatti (2017-2019) and European Flora (Tutin et al., 1964-1993). The pollen terminology is based on Berglund and Ralska-Jasiewiczowa (1986), Faegri and Iversen (1989) and Moore et al. (1991) with slight modifications that tend to simplify nomenclature of plants. The term “taxa” is used in a broad sense to indicate both the systematic categories and the pollen morphological types (Beug 2004). Identified pollen groups (between 300 and 400 pollen grains) have been expressed as percentages of the total (usually between 300 and 400 pollen grains). Pollen percentages are computed on the basis of the total pollen sum. All samples are characterized by fairly high concentration, ranging from 1,072 -30,659 p/g, and good conservation status. The discreet pollen biodiversity (60 pollen types: 22 woody, 32 herbaceous and 6 Monophyta) found, suggests that flood deposits formed in a rich vegetal environment, with high floristic biodiversity. Pollen groups are defined on the basis of common biological characteristics of the plants, useful to reconstruct vegetation dynamics due to climate fluctuations. In particular groups composition are described in Fig. 10.



230 **4. Results**

**4.1 Stratigraphical data and their geological context**


**4.1.1 Sedimentary facies**

The stratigraphic distribution in the core profiles and the synthetically description facies are shown in Fig. 3. Thirteen different facies types have been identified and named from LM1 to LM13:

235

- LM1 to LM2 Clast-supported deposit with thickness ranging from 5 to 15 cm. Extremely coarse (LM1) to medium pebbles (LM2) with abundant coarse to fine grained sandstone matrix, showing a low degree of sorting. Polygenic clasts have low sphericity and very angular shape. Occasional fragments of wood;
- LM3 Massive matrix-supported deposit composed of medium pebbles to granules with very fine sand to silt matrix. Polygenic clasts have low sphericity and subangular shape. Sharp basal and top contacts with evidence of occasionally erosion along the basal one. Presence of fragments of wood;
- LM4 to LM6 General clast-supported, poorly sorted deposit with thickness ranging from 10 to 30 cm. Polygenic clasts, massive or crudely graded angular/subangular medium pebbles to very fine granules. Crudely horizontal laminae at the top. Sharp basal and top contacts with evidence of occasionally erosion along the basal one. Occasional fragments of wood at the base of the layers;
- LM7 to LM9 Massive or crudely graded, poorly sorted very coarse to fine sands 5-20 cm thick. The top part of the detrital layer consists of the finest particles with planar lamination. Sharp basal and top contacts. Occasional fragments of wood at the top of the layers;
- LM10 Sandy loam to clayey loam. Presence of polygenic fine to medium pebbles aligned with low sphericity and very angular shape. Dark greyish brown colour (10YR 4/2 or 10YR 3/2). Sharp basal and top contacts;
- LM11 Peat layer and organic rich clay layers. Fine to medium pebbles aligned with low sphericity and very angular shape. Dark colour (10YR 3/1);
- LM12 Massive clayey silts with rare organic matter. Occasional polygenic clasts (fine pebbles) have low sphericity and very angular shape. Sharp basal and top boundaries. Dark greenish grey colour (5G 4/1);
- LM13 Loam to silty clay texture, strong coarse angular blocky, dark olive grey dry colour (5Y3/2) with yellowish brown redox concentrations (10YR5/6), no carbonate, field pH 5.5.

255



**Figure 3.** Sediment core description (see legend) and radiocarbon dates. *See a larger version of this figure at the end of the manuscript*


260

Relatively to the Lake Moo site, the different coarse-grained facies from LM1 to LM9 have been interpreted as the extreme flood events triggered only by high-intensity convective rainfall events in the catchment area that flow into the Lake Moo as hyperpycnal flow. Our main assumption is that, with favourable sediment transfer into lake and small catchment area (<2km<sup>2</sup>), high density flood can be triggered only by HIP due to erosion of material from the drainage system network (Milliman and Syvitski, 1992; Mulder and Syvitski, 1995; Mutti et al., 1996). The facies from LM1 to LM9 were grouped according to the genetic approach and therefore on the basis of facies tract concept as described in Fig. 4 (Lowe, 1982; Mutti, 1992; Mutti et al., 1996).

265



270 Fine-grained facies, from LM10 to LM13, are subdivided into two groups. In the first group belong the facies LM11 and LM12 expression of environment shore zone and subaqueous lacustrine deposits respectively. The second group includes the facies LM10 and LM13 expression of subaerial deposits.



**Figure 4.** Idealized genetic facies tract interpretation of clastic deposits associated with S1 core.

#### 275 4.1.2 Stratigraphic units

The sedimentary succession of the reference core S1 is subdivided into five informal units, described below from the bottom to the top (Fig. 3):

280 Unit 1: This unit, 95 cm-thick, is represented only by the LM13 facies. Two levels can be distinguished: from 11.30 m to 11.60 m core depth is present a slickensides horizon (Bss), while from 11.60 m to 12.00 m core depth is present a weathered horizon (Bw). This unit is interpreted as a mature paleo soil, likely developed on a colluvium forming the base of a structural depression;

285 Unit 2: This unit is 6.9 m thick and ranging from 11.30 m to 4.4 m core depth, and is characterized by the presence of several layers of coarse sediments (LM3) separated by deposits with a predominantly fine composition (LM11). The thickness of coarse layers, belonging to facies LM4-LM6 ranges from 10 cm to 30 cm. To the top of the succession and if age dating is reliable, the unit is closed by an unconformity surface recording a important time gap. The same surface is the base of a lacustrine transgression which make up the unit 3;

290 Unit 3: This unit is thick 2.1 m and ranging from 4.4 m to 2.3 m of depth. Sharp basal and top boundaries. The basal portion of this unit (from 4.4 m to 3.2 m of depth) present matrix-supported deposit with several fragments of wood which we attribute to the shore zone environment. Only this portion of core show a characteristic mottledling-like appearance. From the 3.2 m of depth follows a lacustrine deposit that closed the unit. This unit is characterized by a period of apparent inactivity of the fluvial system, until its reactivation documented in the uppermost part of the core (unit 4 and 5).

295 Unit 4: This unit is thick 1.4m and ranging from 2.3m to 0.9m of depth and registers the return of several coarse-grained levels which are in general less thick and less coarse than those present in unit 2. These levels are prevalent separated by marsh deposits. Sharp basal and top unit contacts.



300 Unit 5: This unit is thick 0.9m, ranging from 0.9m of depth to land surface and records the disappearance of lake deposits  
replaced by fluvial ones through an erosional contact. The flood deposits produced by the rainfall event of September 13<sup>th</sup>  
and 14<sup>th</sup> 2015 closes the sequence (Fig. 5).



305 **Figure 5.** Though the exposure is quite small, these graded pebble-sand couplets could be interpreted as a sheet-flood  
deposit (Unit 5).

As a whole, we interpret the local lacustrine succession (units 1, 2, 3 and 4) is like to the infill of a structural  
depression produced by gravitational block sliding that was induced by post-glacial fluvial incision (Gunderson et al.,  
310 2014).


#### 4.2 Age-depth model

The age model for the S1 core is based on eight <sup>14</sup>C dates (Fig. 6). Coarse levels are removed from the sediment  
record when constructing the age-depth model because represent instantaneous deposits. The 27 deposits interpreted as  
315 instantaneous events representing a total of 374.5 cm were removed, the remaining were used to build an event-free  
sedimentary record.

Four radiocarbon dates were rejected: one (LTL18275A bis code) because it was not possible to date the sample for poor  
presence of organic matter, the others (LTL18275A, LTL18575A and LTL18272A codes) because it was not possible to  
identify a calibrate age.

320


325



**Figure 6.** Age-depth model obtained from radiocarbon dates. Black and grey line represents the two-sigma probability envelope depth without coarse grained events. Red and orange line represents the two-sigma probability envelope depth with coarse grained events.

### 5. Discussion

Figure 7 shows the stratigraphic succession of the S1 core compared with the most relevant climate proxy available from literature for the area of interest. These includes: reconstructed mean July air temperature from Lago Verdarolo (from Samartin et al., 2017),  $\delta^{18}\text{O}$  speleothem records and reconstructed precipitation trends (mean anomaly time series) from Renella cave (data modified from Combourieu et al., 2013; Regattieri et al., 2014; Zanchetta et al., 2011; Zhomyak et al., 2011), distribution of the higher lake levels event reconstructed in Jura and Pre Alpine mountains (modified from Magny, 2004), and June (red) - December (blue) insolation values records reported for 45° Nord.



340

**Figure 7.** The stratigraphic succession of the S1 core compared with the most relevant climate proxy available from literature for the area of interest. *See a larger version of this figure at the end of the manuscript*

The lake Moo stratigraphic sequence (Fig. 3) is showing the presence of several coarse grained layers separated by deposit with a predominantly fine composition. These coarse levels are particularly developed within unit 2 (between 5 and 10.5m in depth) and correspond chronologically to range of the HTM of the Verdarolo curve and with the wet/high intensity phase reported at the nearby Renella and Corchia Cave (Fig. 7, Unit 2). Zhomyak et al. (2011) attributed this high fluvial activity, which reached levels of the caves flooded only during extreme events, to an increase of strong convective events like the one that affected the region in 1996. Other authors confirm, through model simulations, higher precipitation accumulations during HTM over the Mediterranean area, especially at end of summer/autumn seasons. This has been linked with a greater expansion of the African Monsoon which contributed to enrich of water vapour air masses extracted from north-Africa by mid-latitude synoptic disturbances (Skinner and Poulsen, 2016 and Tinner et al., 2013 in their Climatic simulations section, Fig. 4). With the end of the HTM a drastic decrease of the coarse deposit levels is observed which we are assuming attributed to a decrease in frequency of HIP. In particular, the transition between unit 2 and 3 is marked by an important unconformity contact. This unconformity is followed by deposits of shorezone environment and subsequently at the top of unit 3 by lacustrine deposits. The top of the unit 3 sequence has a calibrated age that is approximately 146-14 BP and is characterized by a period of apparent inactivity of the fluvial system, until its reactivation documented in the units 4 and 5 due to hypothesized new increase of HIP in recent times. The upper most part of the unit 4 has an age of about 149-11BP. Finally, the disappearance of lake deposits replaced by fluvial ones through an erosional contact is marked by unit 5.


From this first qualitative comparison we can affirm that there is a positive correlation (at the millennial scale) between temperatures and precipitation peak deduced from nearby speleothem records and the level of alpine lakes. This confirms findings from other authors (for example Marcott et al. 2013, Giguet-Covex et al., 2012), which are associating greater warmth (in the northern hemisphere) with periods of more intense precipitation.

365




Instrumental data, available since the second half of the last century from the Eraclito ER dataset, has been added to the Verdarolo curve to allow a comparison between geological time and recent instrumental data. The overlap with the latest part of Verdarolo reconstructed curve suggests a good accuracy of the reconstruction technique in this region (Fig. 8). The recent temperature increase is striking if compared with the previous period, and the current July temperature is comparable with the maximum temperature reached at HTM over this area. The actual trend of July temperature, estimated over the period 1961-2018, is  $+4.3^{\circ}\text{C}$  in 100 years. This value is highly statistically significant, with a p-value=0.0004 (see section 3.3 for a description of the method and significance). The recent trend is more than double of the maximum temperature gradient found in the Verdarolo curve,  $+2^{\circ}\text{C}/100\text{y}$ , found, for a short period of time, at end of the Little Ice Age 1850-1900. While the maximum positive temperature gradient found at the end of Younger Dryas age is about  $+1^{\circ}\text{C}/100\text{y}$ . It is therefore clear that summer (July) temperature is increasing very rapidly, with an unprecedented rate in the whole Holocene. Such a large difference in increase rate can hardly be explained by the lower frequency sampling of the Verdarolo reconstructed curve although uncertainty in comparison instrumental data with reconstructed temperature data has been discussed before (Marcott et al. 2013).

This rise of temperature is associated also with a comparable unprecedented intensification of fluvial deposition, testified in the core, by the deposition of units 4 and 5, which marks the transition (in a short time) from marshy to fluvial sheet-flood deposits (Fig. 5). In fact, the recent sedimentation rate (computed from the core S1) is  $1.27\text{m}/100\text{y}$  since 1850, that is 10 times higher than HTM ( $0.124\text{ m}/100\text{y}$  from 5455-9672 BP). In particular, whether the lower or upper end of the 2 sigma confidence level associated with the dating at 280cm depth (146-14 B.P.) is considered, the frequency of the coarse grained deposits (Unit 4 and 5 in figure kk), triggered hence by HIP, varies from a minimum of 5 events every 100 years to a maximum of 12 events every 100 years, respectively. This observation reinforces the trend of increasing precipitation intensity (both as daily maximum and as monthly cumulative values) that is already emerging in the autumn months from the instrumental data as shown in Fig. 9.



**Figure 8.** Comparison of current data and reconstructed July mean temperature at Lake Verdarolo site. The blue line is the reconstructed temperature and the shaded area is the sample-specific estimated standard error associated with the temperature reconstruction based on chironomid assemblage from Samartin et al. (2017). The orange line represent the July mean temperature (1961-2018) retrieved for the grid cell of Lago Verdarolo from Emilia-Romagna climate reanalysis ERACLITO (11 years running average). The shaded orange area is  $\pm$  one standard deviation. On the left the full available period while on the right a zoom on the most recent period from 1600 AD up to now.



400 **Figure 9.** Monthly mean, 2m temperature (a), accumulated precipitation (b), maximum daily precipitation (c), for 1961-  
1990 (blue) and 1991-2018 (green) computed over Lake Moo site grid cell of Eraclito reanalysis (grid cell 258, Ferriere  
Municipality). Confidence intervals at 95% significance (black vertical segments) are computed with a bootstrapping  
method (1000 iterations) as part of the Seaborn python library.

#### 405 5.1 Detailed comparison on two key intervals of the Holocene

In this section, we introduce the pollen data to fully exploit the multidisciplinary approach. As discussed in  
section 3.4, we focus on two specific intervals, I<sub>1</sub> and I<sub>2</sub>, synthesizing all available data for these periods in Fig. 10. The  
difference in mean temperature between I<sub>1</sub> and I<sub>2</sub>, obtained averaging the corresponding samples in the reconstructed  
temperature curve from Verdarolo site, is +1.3°C while the maximum difference reach +3.1°C.

410

The main considerations arising from this comparison are the following:


- I<sub>1</sub>) the arboreal component is always higher than the herbaceous one, with an average value of 80.1%. The  
presence of conifers with a prevalence of pines and silver fir is significant compared to deciduous hardwoods.  
415 Conifers reach a maximum of 95% in P4 sample. The peak occurs during a relative cooling, just before the 8.2  
event, in accordance with the bibliography (Allen et al., 2007; Kofler et al., 2005; Matero et al., 2017; Tinner  
and Lotter, 2001). Subsequently, samples P5 and P6 (HTM) show a decrease in the presence of conifers,  
respectively 64% and 45%, with a corresponding increase in deciduous species, from 1.5% in P4 to 20% in P5  
and P6. This occur in response to warmer conditions and pluvial phase, as indicated by Regatieri et al., 2014;  
420 Zhomyak et al., 2011; Zanchetta et al., 2011. It should also be noted the absence of species typical of humid  
environment (hygro+hydro in Fig.10), which suggests a high stationing of the lake-level. This confirms higher  
rainfall accumulation in the period, in accordance with lake-level evidence from Jura Mountains and the central-  
northern Italy Magny et al. (2004, 2009, 2012). In P7 sample, we observe a new increase in conifers (75%) and



425 a consequent reduction in deciduous trees (10%). These results, combined with the stratigraphic interpretation  
 of the S1 core, suggest that pluvial phase, in this period, had been characterized by a high component of HIP.  
 This can be deduced from the characteristics of the basin and the presence of an extensive vegetation cover  
 which implies the need of even more intense rainfall to able to trigger high flood activity (as recorded in this  
 interval) in a dense vegetated area, with forest coverage that ranges from 65 to 97%. The thickness of coarse  
 430 grained layers range from 30 and 13 cm with pebble-sands, crudely graded characterised by basal and top shap  
 contact. These coarse grained deposits are expressing of proximal facies (LM4 to LM6, in Fig. 10).

- I<sub>2</sub> instead is characterized by LM7 to LM8 distal facies from hyperpycnal flows, as a result of a low flood activity.  
 The thickness layers do not exceed 5 cm and they are medium-fine sands. From a bibliographic point of view,  
 this time interval is characterized by a cooling trend and dry conditions over the Mediterranean region occurred  
 435 approximately between 4.2 - 3.9 ka BP (Regatieri et al., 2014; Zhomyak et al., 2011; Zanchetta et al., 2011,  
 2016), recently confirmed by a review paper of Bini et al. (2019). Further supporting these evidences low pluvial  
 activity, pollen samples highlight a significant reduction of forest, whose coverage falls to a minimum of 30%  
 (in P10 sample), in favour of a growth of grassland area (with maximum extension of 31% in P10) and  
 hygro+hydro species (13.2%).

440 The increase of hygrophilous species, which double their presence, supports the hypothesis of reduction of the  
 lake level, due to colonization of swampy areas which were forming as the lake level drop. Although the  
 conditions in I<sub>2</sub> interval are favourable for the activation of hyperconcentrated flows (due to the lower vegetation  
 cover), the low flood activity observed suggests a low frequency of HIP events.



445



**Figure 10.** The stratigraphic intervals I<sub>1</sub> and I<sub>2</sub> from S1 core compared with the original pollinic data and the relevant climate proxy available from literature for the area of interest. *See a larger version of this figure at the end of the manuscript*

## 6. Conclusions

450 Placing present climate conditions into a geological perspective time, beyond instrumental record, is important for understanding changes in the hydrological cycle induced by anthropogenic climate warming, complementing projections of model climate simulations. Physical reasons and regional climate reconstructions are consistently pointing to an increase of precipitation intensity when water vapour is not limited (Prein et al., 2017). In the last decade record-breaking rainfall events have occurred frequently around the world, this trend is emerging with different strength in  
455 different area, and often, on the short timescale of the instrumental series, doesn't allow yet to draw firm conclusion. Therefore, in the attempt to consolidate confidence about extreme precipitation trend, we extended our analysis, passing from the instrumental time scale to the geological time. This choice implies a strong multidisciplinary approach which includes, in addition to climate and meteorological data, proxy and competences coming from geological, geomorphological, stratigraphical, paleobotanical area. We acquired original data during a field survey the months of  
460 June and July 2017.

Here we can summarize the principal results:

1) The matching of instrumental temperature data and the paleoenvironmental data allow us to affirm that recent temperature trends (over the period 1961-2018), and in particular those of the summer months (July), are in absolute terms the highest ever recorded in the Northern Apennines with an estimated of +4.3°C/100y (highly statistically significant),  
465 which is four times higher than the maximum recorded at the exit of last ice age (Younger Dryas).

2) The recent sedimentation rate (1.27m/100y since 1850) is 10 times higher than that observed during the HTM (0.124 m/100y from 5455-9672 BP) and therefore higher the previous warmer period. The observed sedimentation increase, and higher frequency of high flow discharge has to be linked to an increase of HIP over the Lake Moo basin since we could not attribute it to other changes in physiographic and vegetation factors. The afforestation area for example,  
470 during HTM and in current time persisted on high percentage, a factor which should be detrimental for the mobilization of debris.

3) HIP increase in response to higher temperature is already detectable in observation series, especially in seasons when moisture availability is not limited, like in Autumn (Brönnimann et al., 2018; Prein et al., 2017). We found evidences that this occurred also in the past, especially during HTM testified by higher deposition of large size sediment in to the  
475 lake. A comparison with the past help understanding future projections on the area, although we are aware that past evolution cannot be taken as an analogous for future due to the different forcing and consequent response of climate system (D'Agostino et al. 2019). As temperature will continue to increase on the Mediterranean area precipitation intensity would keep increasing over the Northern Apennine. We hypothesize that precipitation intensity increase will be evident in months with cooler and moist air masses, like in Autumn as it already emerging (see Fig.9), but gradually  
480 extending towards Winter. In summer, increasing conditions of moisture limitations will induce a decrease in frequency and intensity of precipitation (Dobrinski et al. 2016a and 2016b)

4) The concept of facies tract (Lowe, 1982; Mutti et al., 1996; Zavala and Pan, 2018) applied here, may represent an important approach for activity flood reconstruction dynamics and may be applied to other peat-bog deposits to further strengthen and regionalize the signal.

485



Lake Moo site, in the Northern Apennine, was chosen for its strategic geographical position with respect to the dominant atmospheric flow associated with heavy precipitation events. In addition to this, the site has the right geological, geomorphological, vegetation characteristics, basin size (< 2 km<sup>2</sup>), to achieve a strong and clear response between high intensity precipitation and debris flow triggering. For these reasons, Lake Moo basin represents an ideal study area which  
490 deserves further investigation on the relation between past climatic conditions and the dynamics of past flood.

**Data availability:** Original data concerning pollen and radiometric data are available on the open data repository of ARPAE Emilia-Romagna (<https://arpaepv.datamb.it/dataset/lake-moo>)

**Author contribution:** SS and FG developed the idea and the research activity planning and execution. SN, MA, MTDN  
495 have taken care of the administrative aspects. All authors, except for SN and MTDN, have contributed to field activity. Granulometric analysis has been conducted by AC. Pollen analysis have been conducted by VR, SM and MM. SS and FG prepared the manuscript with contributions from all co-authors. FG management and coordination.

**Competing interests:** The authors declare that they have no conflict of interest.

500

#### Acknowledgements

This research was funded by Regional Agency of Civil Protection of the Emilia-Romagna Region and Geological, Seismic and Soil Survey of Emilia-Romagna Region in the framework of cooperation agreement with ARPAE-SIMC.

#### References

- 505 Ahlborn, M., Armon, M., Ben Dor, Y., Neugebauer, I., Schwab, M.J., Tjallingii, R., Shoqeir, J.H., Morin, E., Enzel, Y., and Brauer, A.: Increased frequency of torrential rain storms during a regional late Holocene eastern Mediterranean drought, *Quaternary Research*, 89, 425-431, <https://doi.org/10.1017/qua.2018.9>, 2018.
- Allen, J.R.M., Long, A.J., Ottley, C.J., Pearson, D.G. and Huntley, B. 2007.: Holocene climate variability in northernmost  
510 Europe. *Quaternary Science Reviews*, 26, 1432-1453.
- Anselmetti, F., Wirth, S.B., Glur, L., and Gilli, A.: Holocene flood frequency as reconstructed by lake sediments from multiple archives: A record influenced by solar forcing and atmospheric circulation patterns, in: Late Pleistocene and Holocene climatic variability in the Carpathian-Balkan region, Abstracts Volume, edited by: Mindrescu, M., and Gradinaru, I., Special Issue, 24, 1-2, 2014.
- 515 Antolini, G., Auteri, L., Pavan, V., Tomei, F., Tomozeiu, R. and Marletto, V.: A daily high-resolution gridded climatic data set for Emilia-Romagna, Italy, during 1961–2010. *Int. J. Climatol.*, 36: 1970-1986. doi:10.1002/joc.4473, 2016.
- Antolini, G., Pavan, V., Tomozeiu, R., and Marletto, V. (Eds.): *Atlante idroclimatico dell'Emilia-Romagna 1961-2015*, Servizio Idro-Meteo-Clima, Arpa Emilia-Romagna, Italy, [https://www.arpae.it/dettaglio\\_generale.asp?id=3811&idlivello=1591](https://www.arpae.it/dettaglio_generale.asp?id=3811&idlivello=1591), 2017.
- 520 Ballesteros-Cánovas, J.A., Stoffel, M., St. George, S., and Hirschboeck, K.: A review of flood records from tree rings, *Progress in Physical Geography*, 1-23, doi 10.1177/0309133315608758, 2015.
- Berglund, B.E., and Ralska-Jasiewiczowa, M.: Pollen analysis and pollen diagrams, in *Handbook of Holocene Palaeoecology and Palaeohydrology*, a cura di B.E. Berglund, Chichester, 455-484, 1986.
- Beug, H.J.: *Leifaden der Pollenbestimmungen für Mitteleuropa und angrenzende Gebiete*, Pfeil, München, 2004.
- 525 Bini, M., Zanchetta, G., Perşoiu, A., Cartier, R., Català, A., Cacho, I., Dean, J.R., Di Rita, F., Drysdale, R.N., Finnè, M., Isola, I., Jalali, B., Lirer, F., Magri, D., Masi, A., Marks, L., Mercuri, A.M., Odile Peyron, O., Sadori, L., Sicre, M.A.,



- Welc, F., Zielhofer, C., and Brisset, E.: The 4.2 ka BP Event in the Mediterranean region: an overview, *Clim. Past*, 15, 555–577. <https://doi.org/10.5194/cp-15-555-2019>, 2019.
- 530 Braggio, G., Guido, M.A., and Montanari, C.: Palaeovegetational evidence in the upper Nure Valley (Ligurian-Emilian Apennines, Northern Italy), *Webbia*, 46, 173-185, 1991.
- Brönnimann, S., Rajczak, J., Fischer, E. M., Raible, C. C., Rohrer, M., and Schär, C.: Changing seasonality of moderate and extreme precipitation events in the Alps. *Natural Hazards and Earth System Sciences*, 18 (7), 2047-2056. <https://doi.org/10.5194/nhess-18-2047-2018>, 2018
- 535 Brugnara, Y., and Maugeri, M.: Daily precipitation variability in the southern Alps since the late 19th century. *Int J Climatol.*; 39: 3492– 3504. <https://doi.org/10.1002/joc.6034>, 2019
- Cacho, I., Grimalt, J.O., Canals, M., Sbañfi, L., Shackleton, N.J., Schönfeld, J., and Zahn, R.: Variability of the western Mediterranean Sea surface temperature during the last 25,000 years and its connection with the Northern Hemisphere climatic changes, *Paleoceanography*, 16, 40-52.
- 540 Carton, A. and Panizza, M. (Eds.): *Il paesaggio fisico dell'Alto Appennino Emiliano*, Grafis Edizioni, Bologna, Italia, 1988.
- Combourieu-Nebout, N., Peyron, O., Bout-Roumazeilles, V., Goring, S., Dormoy, I., Joannin, S., Sadori, L., Siani, G., and Magny, M.: Holocene vegetation and climate changes in the central Mediterranean inferred from a high-resolution marine pollen record (Adriatic Sea), *Clim. Past*, 9, 2023-2042. <https://doi.org/10.5194/cp-9-2023-2013>, 2013.
- 545 Corticelli, S., Garberi, M.C., Mariani, M.C., and Masi, S.: *Usa del suolo 2008*, <http://geoportale.regione.emilia-romagna.it/it/download/dati-e-prodotti-cartografici-preconfezionati/pianificazione-e-catasto/uso-del-suolo-1/2008-coperture-vettoriali-uso-del-suolo-edizione-2011>, 2011.
- D'Agostino, R., Bader, J., Bordoni, S., Ferreira, D., & Jungclaus, J.: Northern Hemisphere monsoon response to mid-Holocene orbital forcing and greenhouse gas-induced global warming. *Geophysical Research Letters*, 46, 1591– 1601. <https://doi.org/10.1029/2018GL081589>, 2019
- 550 Drobinski, P., Alonzo B., Bastin S., Da Silva N., and Muller C.J.: Scaling of precipitation extremes with temperature in the French Mediterranean region: what explains the hook shape? *J Geophys Res.* doi:10.1002/2015JD023497, 2016a.
- Drobinski, P., Silva, N.D., Panthou, G., Bastin, S., Muller, C., Ahrens, B., Borga, M., Conte, D., Fosser, G., Giorgi, F., Guttler, I., Kotroni, I., Li, L., Morin, E., Onol, B., Quintana-Segui, P., Romera, R., and Torma, C.Z.: Scaling precipitation extremes with temperature in the Mediterranean: past climate assessment and projection in anthropogenic scenarios, *Climate Dynamics*, 51, 1237-1257, doi:10.1007/s00382-016-3083-x, 2016b.
- 555 Elter, P., Ghiselli, F., Marroni, M., and Ottria, G. (Eds.): *Note illustrative del Foglio 198 "Bobbio" della Carta Geologica d'Italia alla scala 1:50.000*. Istituto Poligrafico e Zecca dello Stato, Roma, Italia, 1997.
- Faegri, K., and Iversen, J.: *Textbook of Pollen analysis*, 4<sup>o</sup> edizione a cura di K. Faegri, P.E. Kaland, K. Krzywinski, Chichester, 1989.
- 560 Frei, C. and Schär, C.: A precipitation climatology of the Alps from high-resolution rain-gauge observations. *Int. J. Climatol.*, 18: 873-900. doi:10.1002/(SICI)1097-0088(19980630)18:8<873::AID-JOC255>3.0.CO;2-9, 1998.
- Giannecchini, R., Galanti, Y., and D'Amato Avanzi, G.: Critical rainfall thresholds for triggering shallow landslides in the Serchio River Valley (Tuscany, Italy), *Nat. Hazards Earth Syst. Sci.*, 12, 829–842, <https://doi.org/10.5194/nhess-12-829-2012>, 2012.
- 565 Giannecchini, R., and D'Amato Avanzi, G.: Historical research as a tool in estimating hydrogeological hazard in a typical small alpine-like area: the example of the Versilia River basin (Apuan Alps, Italy), *Physics and Chemistry of the Earth*, 49, 32-43, doi 10.1016/j.pce.2011.12.005, 2012.



- Giguët-Covex, C., Arnaud, F., Dirk, E., Jérôme, P., Laurent, M., Pierre, F., Fernand, D., Pierre-Jérôme, R., Bruno, W., and Jean-Jacques, D.: Frequency and intensity of high-altitude floods over the last 3.5ka in northwestern French Alps (Lake Anterne), *Quaternary Research*, 77, 12-22, <https://doi.org/10.1016/j.yqres.2011.11.003>, 2012.
- 570 Gilli, A., Anselmetti, F.S., Glur, L., and Wirth S.B.: Lake Sediments as Archives of Recurrence Rates and Intensities of Past Flood Events, in: *Dating Torrential Processes on Fans and Cones, Advances in Global Change Research*, edited by: Schneuwly-Bollschweiler, M., Stoffel, M., and Rudolf-Miklau, F., Springer Science Business Media, Dordrecht, Holland, 225-242, doi 10.1007/978-94-007-4336-6\_15, 2013.
- 575 Giraudi, C.: Coarse sediments in Northern Apennine peat bogs and lakes: New data for the record of Holocene alluvial phases in peninsular Italy, *The Holocene*, 24, 932-943, <https://doi.org/10.1177/0959683614534738>, 2014.
- Glur, L., Wirth, S.B., Büntgen, U., Gilli, A., Haug, G.H., Schär, C., Beer, J., and Anselmetti, F.S.: Frequent floods in the European Alps coincide with cooler periods of the past 2500 years, *Nature Scientific Reports*, 3, doi 10.1038/srep02770, 2013.
- 580 Grazzini, F., Craig, J.G., Keil, C., Antolini, G., and Pavan V.: Extreme precipitation events over Northern Italy: Part (I) a systematic classification with machine learning techniques, *QJR Meteorol Soc.* 2019;1–17. <https://doi.org/10.1002/qj.3635>, 2019.
- Grazzini, F., Segadelli, S., and Fornasiero, A.: Precipitazioni estreme e effetti al suolo sul reticolo minore: il caso del 14 Settembre 2015. Technical report ARPAE-SIMC/SGSS Emilia-Romagna, 27pp, Available in Italian [https://www.arpae.it/dettaglio\\_notizia.asp?id=8017&idlivello=32](https://www.arpae.it/dettaglio_notizia.asp?id=8017&idlivello=32), 2016.
- 585 Grazzini, F., Praticcoli, W. and Tomei, F.: Drought and big snow, are extreme events the new normal? *Ecoscienza*, 1, Available in Italian [https://www.arpae.it/dettaglio\\_documento.asp?id=3674&idlivello=1171](https://www.arpae.it/dettaglio_documento.asp?id=3674&idlivello=1171), 2012.
- Gunderson, K.L., Pazzaglia, F.J., Picotti, V., Anastasio, D.A., Kodama, K.P., Rittenour, T., Frankel, K.F., Ponza, A., Berti, C., Negri, A., and Sabbatini, A.: Unraveling tectonic and climatic controls on synorogenic growth strata (Northern Apennines, Italy), *GSA Bulletin*, 126, 532-552, doi 10.1130/B30902.1, 2014.
- 590 Heiri, O., S.J., Brooks, H.J.B., Birks, and A.F.Lotter.: A 274-lake calibration data-set and interference model for chironomid-based summer air temperature reconstruction in Europe, *Quaternary Science Reviews* 30, 3445-3456, 2011.
- Hussain, M., and Mahmud, I.: pyMannKendall: a python package for non parametric Mann Kendall family of trend tests, *Journal of Open Source Software*, 4, (39), 1556, <https://doi.org/10.21105/joss.01556>, 2019.
- 595 Kofler, W., Krapf, V., Oberhuber, W., Bortenschlager, S.: Vegetation responses to the 8200 cal. BP cold event and to long-term climatic changes in the Eastern Alps: Possible influence of solar activity and North Atlantic freshwater pulses. *The Holocene*, 15, 779-788, 2005.
- Isotta, F., Frei, C., Weigluni, V., Percec Tadic, M., Lassègues, P., Rudolf, B., Pavan, V., Cacciamani, C., Antolini, G., Ratto, S.M., Munari, M., Micheletti, S., Bonati, V., Lussana, C., Ronchi, C., Panettieri, E., Marigo, G., and Vartacnik, G.: The climate of daily precipitation in the Alps: development and analysis of a high-resolution grid dataset from pan-Alpine rain-gauge data, *International Journal of Climatology*, 34, 1657-1675, doi 10.1002/joc.3794, 2014.
- 600 Lehmann, J., Coumou, D., and Frieler, K.: Increased record-breaking precipitation events under global warming, *Climatic Change* 132: 501. <https://doi.org/10.1007/s10584-015-1434-y>, 2015.
- 605 Longman, J., Ersek, V., Veres, D., and Salzmann, U.: Detrital events and hydroclimate variability in the Romanian Carpathians during the mid-to-late Holocene, *Quaternary Science Reviews*, 167, 78-95, 2017.
- Lowe, D.R.: Sediment gravity flows; Depositional models with special reference to the deposits of high-density turbidity currents, *Journal of Sedimentary Petrology*, 52, 279-297, 1982.



- 610 Magny, M.: Holocene climate variability as reflected by mid-European lake-level fluctuations and its probable impact on prehistoric human settlements, *Quaternary International*, 113, 65-79, 2004.
- Magny, M., Galop, D., Bellintani, P., Desmet, M., Didier, J., Hass, J.N., Martinelli, N., Pedrotti, A., Scandolari, S., Stock, A., and Vannièrè, B.: Late-Holocene climatic variability south of the Alps as recorded by lake-level fluctuations at Lake Ledro, Trentino, Italy. *Holocene*, 19, 575-589, 2009.
- 615 Magny, M., S. Joannin, D. Galop, B. Vannièrè, J. N. Haas, M. Bassetti, P. Bellintani, R. Scandolari, and M. Desmet. 2012. Holocene palaeohydrological changes in the northern Mediterranean borderlands as reflected by the lake-level record of Lake Ledro, northeastern Italy. *Quaternary Research*, 77, 382-396, 2012.
- Marchetti, G., and Fraccia, R.: Carta geomorfologica dell'alta Val Nure, Appennino piacentino. Scala 1:25.000, in: *Il paesaggio fisico dell'Alto Appennino Emiliano. Regione Emilia-Romagna*, edited by: Carton, A., and Panizza, M., Grafis Edizioni, Bologna, Italia, 1988.
- 620 Marcott, S., Shakun, J., Clark, P.U., and Mix, A.C.: A Reconstruction of Regional and Global Temperature for the Past 11,300 Years, *Science*, 339, 1198-1201, doi 10.1126/science.1228026, 2013.
- Marroni, M., Meneghini, F., and Pandolfi, L.: Anatomy of the Ligure-Piemontese subduction system: evidence from Late Cretaceous-middle Eocene convergent margin deposits in the Northern Apennines, Italy, *International Geology Review*, iFirst article, 1-33, 2010.
- 625 Matero, I.S.O., Gregoire, L.J., Ivanovic, R.F., Tindall, J.C., and Haywood, A.M.: The 8.2 ka cooling event caused by Laurentide ice saddle collapse. *Earth and Planetary Science Letters*, 473, 205-214, 2017.
- Milliman, J.D. and Syvitski, J.P.M.: Geomorphic/Tectonic Control of Sediment Discharge to the Ocean: The Importance of Small Mountainous Rivers, *The Journal of Geology*, 100, 525-544, <http://dx.doi.org/10.1086/629606>, 1992.
- Moore, P.D., and Webb, J.A., and Collinson, M.E.: *Pollen Analysis*, 2a edizione, Oxford, 1991.
- 630 Mulder, T. and Syvitski, J.P.M.: Turbidity Currents Generated at River Mouths during Exceptional Discharges to the World Oceans, *Journal of Geology*, 103, 285-299, <http://dx.doi.org/10.1086/629747>, 1995.
- Mutti, E.: *Turbidite Sandstones*. San Donato Milanese, Agip-Istituto di Geologia, Università di Parma, 275 pp, 1982.
- Mutti, E., Davoli, G., Tinterri, R., and Zavala, C.: The importance of ancient fluvio-deltaic systems dominated by catastrophic flooding in tectonically active basins, *Memorie di Scienze Geologiche*, Università di Padova, 48, 233-291, 1996.
- 635 Nesje, A., Dahl, O., Matthews, J.A., and Berrisdorf, M.S.: A 4500 years of river floods obtained from a sediment core in Lake Atnsjoen, eastern Norway, *Journal of Paleolimnology*, 25, 329-342, 2001.
- Nistor, M.M.: Spatial distribution of climate indices in the Emilia-Romagna region, *Meteorological Applications*, 23, 640 304-313, doi 10.1002/met.1555, 2016.
- Papalexioy, S. M. and Montanari, A.: Global and regional increase of precipitation extremes under global warming. *Water Resources Research*, 55, 4901– 4914. <https://doi.org/10.1029/2018WR02406>. 2019.
- Pavan, V., Antolini, G., Barbiero, R., Berni, N., Brunier, F., Cacciamani, C., Cagnati, A., Cazzuli, O., Cicogna, A., De Luigi, C., Di Carlo, E., Francioni, M., Maraldo, L., Marigo, G., Micheletti, S., Onorato, L., Panettieri, E., Pellegrini, U., Pelosini, R., Piccinini, D., Ratto, S., Ronchi, C., Rusca, L., Sofia, S., Stelluti, M., Tomozeiu, R. and Malaspina, T.T. : High resolution climate precipitation analysis for north-central Italy, 1961–2015. *Climate Dynamics*, 52 3435–3453, <https://doi.org/10.1007/s00382-018-4337-6>. 2019




- Peyron, O., Goring, S., Dormoy, I., Kotthoff, U., Pross, J., de Beaulieu, J.-L. and Magny, M.: Holocene seasonality changes in the central Mediterranean region reconstructed from the pollen sequences of Lake Accesa (Italy) and Tenaghi Philippon (Greece), *The Holocene*, 21, 131–146. <https://doi.org/10.1177/0959683610384162>, 2011.
- Pignatti, S.: *Flora d'Italia*, Edagricole-New Business Media, Milano, 2017-2019.
- Prein, A.F., Liu, C., Ikeda, K., Trier, S.B., Rasmussen, R.M., Holland, G.J., and Clark, M.P.: Increased rainfall volume from future convective storms in the US, *Nature Climate Change*, 7, 880-884, doi:10.1038/s41558-017-0007-7, 2017a.
- Prein, A.F., Rasmussen, R.M., Ikeda, K., Liu, C., Clark, M.P., and Holland, G.J.: The future intensification of hourly precipitation extremes, *Nature Climate Change*, 7, 48-52, doi:10.1038/nclimate3168, 2017b.
- Regattieri, E., Zanchetta, G., Drysdale, R.N., Isola, I., Hellstrom, J.C., and Dallai, L.: Lateglacial to Holocene trace element record (Ba, Mg, Sr) from Corchia Cave (Apuan Alps, central Italy): paleoenvironmental implications, *Journal of Quaternary Science*, 29, 381-392, <https://doi.org/10.1002/jqs.2712>, 2014.
- Reimer, P.J., Bayliss, A., Warren Beck, J., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hafliðason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP, *Radiocarbon*, 55, 1869-1887, [https://doi.org/10.2458/azu\\_js\\_rc.55.16947](https://doi.org/10.2458/azu_js_rc.55.16947), 2013.
- Sabatier, P., Wilhelm, B., Ficotola, G.F., Moiroux, F., Poulencard, J., Develle, A.L., Bichet, A., Chen, W., Pignol, C., Reyss, J.L., Gielly, L., Bajard, M., Perrette, Yves., Malet, E., Taberlet, P. and Arnaud, F.: 6-kyr record of flood frequency and intensity in the western Mediterranean Alps - Interplay of solar and temperature forcing, *Quaternary Science Reviews*, 170, 121-135, <https://doi.org/10.1016/j.quascirev.2017.06.019>, 2017.
- Samartin, S., Heiri, O., Joos, F., Renssen, H., Franke, J., Brönnimann, S., and Tinner, W.: Warm Mediterranean mid-Holocene summers inferred from fossil midge assemblages, *Nature Geoscience*, 10, 207-212. doi:10.1038/NGEO2891, 2017.
- Scherrer, S. C., Fischer, E.M., Posselt, R., Liniger, M.A., Croci-Maspoli, M., and Knutti, R.: Emerging trends in heavy precipitation and hot temperature extremes in Switzerland, *J. Geophys. Res. Atmos.*, 121, doi:10.1002/2015JD024634, 2016.
- Schillereff, D., Chiverrell, R.C., Macdonald, N., and Hooke, J.M.: Flood stratigraphies in lake sediments: A review, *Earth-Science Reviews*, 135, 17-37, doi 10.1016/j.earscirev.2014.03.011, 2014.
- Schneuwly-Bollschweiler, M., Stoffel, M., and Rudolf-Miklau, F. (Eds.): *Dating Torrential Processes on Fans and Cones. Methods and Their Application for Hazard and Risk Assessment*. Springer, doi 10.1007/978-94-007-4336-6, 2013.
- Skinner, C. B., and C. J. Poulsen: The role of fall season tropical plumes in enhancing Saharan rainfall during the African Humid Period, *Geophys. Res. Lett.*, 43, 349–358, doi:10.1002/2015GL066318, 2016.
- Stoffel, M., Butler, D.R., and Corona, C.: Mass movements and tree rings: A guide to dendrogeomorphic field sampling and dating, *Geomorphology*, 200, 106-120, <https://doi.org/10.1016/j.geomorph.2012.12.017>, 2013.
- Stoffel, M., Wyzga, B., and Marston, R.A.: Floods in mountain environments: A synthesis, *Geomorphology*, 272, 1-9, <https://doi.org/10.1016/j.geomorph.2016.07.008>, 2016.
- Swierczynski, T., Ionita, M., and Pino, D.: Using archives of past floods to estimate future flood hazards, *Eos* 98, <https://doi.org/10.1029/2017EO066221>, 2017.
- Tinner, W., and Lotter, A.F.: Central European vegetation response to abrupt climate change at 8.2 ka. *Geology*, 29, 551–554, 2001



- 690 Tinner, W., Colombaroli D., Heiri, O., Henne, P., Steinacher, M., Untenecker, J., Vescovi, E., Allen, J., Carraro, G.,  
Conedera, M., Joos, F., Lotter, A., Luterbacher, J., Samartin, S., and Valsecchi, V.: The past ecology of *Abies alba*  
provides new perspectives on future responses of silver fir forests to global warming. *Ecological Monographs* 83(4):  
419-439, 2013.
- Tutin, T.G., Heywood, V.H., Burges, N.A., Valentine, D.H.: *Flora Europaea*, Cambridge University Press, 1964.
- Utsumi, N., Seto, S., Kanae, E., Maeda, E., and Oki T.: Does higher surface temperature intensify extreme precipitation?,  
695 *Geophys. Res. Lett.*, 38, L16708, doi:10.1029/2011GL048426, 2011.
- Vescovi, P. (Ed.): Note illustrative della Carta Geologica d'Italia alla scala 1:50.000, Foglio n° 216 Borgo Val di Taro.  
Sheet n° 216 Borgo Val di Taro, Servizio Geologico della Regione Emilia-Romagna, Servizio, ISPRA, Rome,  
[http://www.isprambiente.gov.it/Media/carg/note\\_illustrative/216\\_Borgo\\_Val\\_di\\_Taro.pdf](http://www.isprambiente.gov.it/Media/carg/note_illustrative/216_Borgo_Val_di_Taro.pdf), 2002.
- 700 Walker, M.J.C., Berkelhammer, M., Björck, S., Cwynar, L.C., Fisher, D.A., Long, A.J., Lowe, J.J., Newnham, R.M.,  
Rasmussen, S.O., and Weiss, H.: Formal subdivision of the Holocene series/Epoch: A discussion paper by a working  
group of INTIMATE (Integration of ice-core marine and terrestrial records) and the subcommission on quaternary  
stratigraphy (International commission on stratigraphy), *Journal of Quaternary Science*, 27, 649–659, 2012.
- Westra, S., Fowler, H.J., Evans J.P., Alexander, L.V., Berg, P., Johnson, F., Kendon, E.J., Lenderink, G., and Roberts,  
N.M.: Future changes to the intensity and frequency of short duration extreme rainfall, *Reviews of Geophysics*, 52,  
705 522-555, doi:10.1002/2014RG000464, 2014.
- Wilhelm, B., Arnaud, F., Sabatier, P., Crouzet, C., Brisset, E., Chaumillon, E., Disnar, J.R., Guter, F., Malet, E., Reyss,  
J.L., Tachikawa, K., Bard, E., and Delannoy, J.J.: 1400 years of extreme precipitation patterns over the Mediterranean  
French Alps and possible forcing mechanism, *Quaternary Research*, 78, 1-12, doi 10.1016/j.yqres.2012.03.003, 2012.
- 710 Wilhelm, B., Ballesteros-Cánovas, J.A., Macdonald, N., Toonen, W.H.J., Baker, V., Barriendos, M., Benito, G., Brauer,  
A., Corella, J.P., Denniston, R., Glaser, R., Ionita, M., Kahle, M., Liu, T., Luetscher, M., Macklin, M., Mudelsee, M.,  
Muñoz, S., Schulte, L., George, S.St., Stoffel, M., and Wetter, O.: Interpreting historical, botanical, and geological  
evidence to aid preparations for future floods, *WIREs Water*, 1-22, <https://doi.org/10.1002/wat2.1318>, 2018.
- Wirth, S.B.: The Holocene Flood History of the Central Alps Reconstructed from Lacustrine Sediments: Frequency,  
Intensity and Controlling Climate Factors. Ph.D. thesis, ETH Zürich Bibliography, Switzerland, 179pp, doi  
715 10.3929/ethz-a-009775044, 2013.
- Wirth, S.B., Glur, L., Gilli, A., and Anselmetti, F.S.: Holocene flood frequency across the Central Alps – solar forcing  
and evidence for variations in North Atlantic atmospheric circulation, *Quaternary Science Reviews*, 80, 112-128,  
<https://doi.org/10.1016/j.quascirev.2013.09.002>, 2013.
- Zanchetta, G., Isola, I., Piccini, L., and Dini, A., 2011. The Corchia Cave (Alpi Apuane) a 2 Ma long temporal window  
720 on the earth climate, *Technical Periodicals of National Geological Survey of Italy - ISPRA*, *Geological Field Trips*,  
3, 55pp, doi 10.3301/GFT.2011.02, 2011.
- Zavala, C. and Pan, S.: Hyperpycnal flows and hyperpycnites: Origin and distinctive characteristics. *Lithologic  
Reservoirs*, 30, 1-27, doi 10.3969/j.issn.1673-8926.2018.01.001, 2018.
- 725 Zavala, C., Ponce, J.J., Arcuri, M., Dritanti, D., Freije, H., and Asensio, M.: Ancient Lacustrine Hyperpycnites: A  
Depositional Model from a Case Study in the Rayoso Formation (Cretaceous) of West-Central Argentina. *Journal of  
Sedimentary Research*, 76, 41-59, <https://doi.org/10.2110/jsr.2006.12>, 2006.
- Zavala, C., Arcuri, M., Di Meglio, M., Gamero Diaz, H., and Contreras, C.: A Genetic Facies Tract for the Analysis of  
Sustained Hyperpycnal Flow Deposits. in: *Sediment transfer from shelf to deep water - Revisiting the delivery system*,  
edited by: Slatt, R.M. and Zavala, C., *AAPG Studies in Geology*, 61, 31-51, 2011.



- 730 Zhomyak, L.V., Zanchetta, G., Drysdale, R.N., Hellstrom, J.C., Isola, I., Regattieri, E., Piccini, L., Baneschi, I., and Couchoud, I.: Stratigraphic evidence for a "pluvial phase" between ca 8200-7100 ka from Renella cave (Central Italy), *Quaternary Science Reviews*, 30, 409-417 doi 10.1016/j.quascirev.2010.12.003, 2011.



735 Figure 2




Figure 3



740





Figure 7



745 Figure 10