General comment on the revised manuscript:

We have revised the manuscript in response to the comments of 3 reviewers. We have also taken the opportunity to update the model database to add 4 new CMIP6/PMIP4 models to the study. This required modification of Table 1 and all Figures to include the new set of models but did not substantially alter any of the conclusions of the study.

In revising the paper in response to the reviewers, we have added a new Table 2 summarising the ensemble mean ENSO variability changes for all models, CMIP5 and CMIP6 models and added Supplementary Figures S1-S4 showing results for CMIP5 and CMIP6 models separately. We have also added Supplementary Figure S5 showing temperature changes relative to the tropical mean.

The Response to Reviewers and revised manuscript with tracked changes are provided below. 10

Response to Reviewer 1:

General response: We appreciate the positive comments of the reviewer and the kind words written about our text and the manner in which we've approached the manuscript. We have revised the text in response to the comments of the reviewer, as outlined below

15

5

Comment: Though the paper is well written, I think there are parts that need further clarification. Specifically, the authors should be clearer about the updated models compared to the previous models. At points, they suggest they will be comparing new simulations with previous generations models (I assume CMIP6/PMIP4 versus CMIP5/PMIP3?) but their results suggest that the models are more lumped together. The paper would be improved if the authors clarified when and how the newest generation models add to our understanding of changes in ENSO with respect to the simulations from previous

20 generations models.

Response: We agree that we have not discussed or quantified the differences between CMIP6 and CMIP5 model results as extensively as we should have. We have addressed this by including Supplementary Figures showing the two ensembles separately, as well as revising the text to highlight where the model ensembles agree on a consistent result.

25 Scientific Questions/Issues

Comment: Lines 55-58 - There is an even newer reference from White et al., 2020 that can be added to this set of references (https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2019GL085504)

Response: This reference was added.

Comment: Lines 74-84 - This paragraph could benefit from updated references. For example, Cobb et al. 2013 is now 30 superseded by Grothe et al. (in press at Geophysical Research Letters https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2019GL083906), which does show a strong reduction of ENSO variability during the mid-Holocene. Also, there's a paper by White et al. 2019 (https://agupubs.onlinelibrary.wiley.com/doi/full/10.1002/2017GL075433) that shows a long-term trend in ENSO strength through the Holocene, contradicting some of the mid-Holocene ENSO reduction studies.

Response: These references was added and the discussion updated. 35

Comment: Section 2.1 Models - I think it would be useful to add just a few more sentences here about the models as not all readers of this journal will have that background. This can be brief and may only include one or two sentences, and then refer the reader to the website for more information. For example, what are the main differences/ improvements in CMIP6, since this paper is really about using the new generation of coupled atmosphere-ocean climate models for both past and

40 future climates. Or maybe just more broadly, what is it about CMIP6/PMIP4 that allows for an updated view of looking at changes in ENSO? I see that the authors add little bits of this specifically sprinkled throughout section 2.2, so I think just a

more broad/general description to set up the context of this study would be beneficial.

Response: The discussion of models in Section 2.1 was expanded to describe the models in more detail, and we identified relevant changes in CMIP6/PMIP4 generation models that may provide new information about changes in ENSO.

45 Comment: In Sections 4-6, when talking about the model ensemble and trends, it is useful to mention the model agreement. The authors do this at times, for example, on lines 338- 343, but I think it would strengthen the observations if this were done more consistently throughout these sections.

Response: The extent of model agreement over changes in the mean state and ENSO teleconnections are already indicated in Figures 6, 7, 10, 11 and 12 with the inclusion of stippling to show where more than 2/3 of models agree on the sign of change. No additional discussion of this aspect of model agreement was added as this would be repetitive and would make

- 50 change. No additional discussion of this aspect of model agreement was added as this would be repetitive and would make the results sections too long. The extent of model agreement over changes in ENSO amplitude (Figures 8 and 9) is already discussed in Section 5, and this was updated with the inclusion of several new models as well as Table 2 showing the multimodel mean and model range. A paragraph was added to Section 5 (and Supplementary Figure S4) to address the agreement between CMIP5 and CMIP6 models over changes in ENSO amplitude.
- 55 Comment: Figures note what the stippled pattern indicates in the legend (as done in Figure 11 and 12). Response: The figure legends were updated to add this information.

Technical Issues

Comment: Line 223 should read "all available data are: : :" Response: This was corrected.

Comment: Line 407 and 410 should be Niño. Response: This was corrected.

60 Comment: Line 414 – remove parentheses around Collins et al. Response: This was corrected.

Comment: Line 484 – Merryfield 2006 reference is not in the parentheses with the other references. **Response:** This was corrected (and added to reference list).

Comment: Line 491 - Should read "This includes: :: " Response: This was corrected.

65 Response to Reviewer 2

General response: We thank the reviewer for their thoughtful and detailed comments. We have revised the manuscript in response to the reviewer comments, as outlined below.

Comment 1: I can understand the framing with respect to the Pliocene but no Pliocene model output is incorporated into the discussion. This may be because the PlioMIP2 simulations were not available when this manuscript was being prepared. If this output is available now I would include it otherwise the framing of the paper is distracting. With respect to the Pliocene

this output is available now I would include it otherwise the framing of the paper is distracting. With respect to the Pilocene there are a few key citations missing. In addition to the zonal temperature gradient, several studies have characterised the thermocline conditions and its coupling to the cold tongue (e.g. Steph et al, 2006; 2010; Ford et al. 2012; 2015).

Also, a recent study by White and Ravelo (GRL, 2020) shows reduced ENSO when the thermocline is deep during the early Pliocene. They suggest mechanistically a weak thermocline feedback dampened ENSO during the Pliocene because the thermocline was deep during the Pliocene. I wouldn't use the Ford and Ravelo 2019 as evidence for ENSO. Ford and Ravelo show that the variability in the western Pacific during the Pliocene was similar to the Holocene. The ENSO variability itself

is very weak in the western Pacific and the reconstructed variability largely reflects the seasonal cycle.

Response: The Introduction has been revised to shorten the discussion of the Pliocene as we do not analyse the PlioMIP simulations (which are the subject of other studies). We have not removed all discussion of the Pliocene as we feel that it outributes to the broad context of understanding ENSO responses to past and future climate. Specifically, the first paragraph of the Introduction is a survey of the geological time-scale history of ENSO which includes the Pliocene and now also the Last Interglacial. The discussion of the Pliocene in Section 7 is now significantly shortened and only mentioned in comparison with the mid-Holocene. Reference to White and Ravelo (2020) was added.

Comment 2: Some paleodata reconstructions that are missing from the paper introduction/ discussion: LGM: Leduc et al., 2009; Koutavas and Joanides (2012); Sadekov et al., 2013; Ford et al., 2015. Mid-Holocene: White et al., 2018. Last Millennium: Rustic et al., 2015. Response: Some of these references were already included and the others have been added.

Comment 3: Can you expand on Line 186-187? What do you mean there has been substantial progress toward it? Given the average residence time of a deep-water parcel is 1000 years how is 150 years close to equilibrium? In what respect? **Response:** The sentence has been deleted to avoid confusion.

- 90 Comment 4: Lines 403-410: This paragraph is confusing to me because "mean state" can suggest different things. I usually think about it as the zonal temperature gradient rather than the eastern Pacific meridional gradient. Can you re-write this to be specific about the gradient you're referring to? Response: This paragraph was expanded and rewritten to more clearly explain which aspects of the mean state were examined: annual cycle, equatorial Pacific zonal SST gradient and eastern Pacific meridional SST gradient.
- 95 Comment 5: Line 420: The paleo reconstructions indicate the cold tongue was productive during the Pliocene (or at least similar to today, Lyle et al., 2015) and the winds were similar (Hovan 1995, Proceedings ODP). Response: This paragraph was edited to reduce the focus on the Pliocene as noted in response to Comment 1, therefore additional references were not added.
- Comment 6: Line 480: This also included a deep thermocline which is mechanistically important for ENSO. This has also been suggested for the LGM: Ford et al. 2018. Response: The role of the deep thermocline in suppression of ENSO variability is now mentioned in this discussion and also in the discussion of the LGM in the Introduction, with added citation of Ford et al. 2018.

Comment 7: Figure 13 is not red green colour-blind friendly. I think for the discussion it would be useful to have two separate comparison between the CMIP5 and CMIP6 grade models. It doesn't sound like there has been much improvement in model performance between the CMIP3 synthesis done for the IPCC report and the models here. It would be nice to know how the models are mechanistically improving or where there are known model errors.

Response: Figure 13 has been redrawn with different symbols for each experiment to ensure it is accessible. We have also added a discussion of the difference between CMIP5 and CMIP6 models throughout the paper, including 4 new supplementary figures and a new Table 2 showing changes in ENSO amplitude for CMIP5 and CMIP6 models. There is

110 actually general agreement between the two generations of models, based on the available simulations. Detailed evaluation of CMIP6 model performance in simulating ENSO is beyond the scope of this study and will no doubt be the focus of many other papers.

Response to Reviewer 3

115 General response: We thank the reviewer for their thorough analysis of our research.

Comment: Overall opinion: This is an interesting and well structured study. The results are well elaborated and convincing. My major concern is the presentation of some SST Figures, as relative SST highlights in many cases the relation between SST change and precipitation/atmospheric circulation changes much better (see major point). A more detailed discussion of the presented results to the changes in the Walker Circulation would also allow a deeper insight into the cause of the

- 120 precipitation and ENSO amplitude changes. Response: We recognise that this work is predominantly descriptive which is unfortunately necessary given the large quantity of models and simulations being analysed here. We now include figures of changes in relative SST in addition to the existing figures of absolute SST change. We would prefer to leave evaluation of the changes in the Walker circulation for future studies as this is a substantial additional body of work that would greatly increase the size of the current manuscript.
- 125 Major comments:

Comment: Introduction: You don't say anything about ENSO in the Interglacial. Please add. **Response:** There was already a brief discussion of proxy and model studies of ENSO in the last interglacial provided in the Discussion. A summary of these studies is now added to the Introduction as well.

Comment: Fig. 1: As for the tropical circulation the relative SST reveals the relation to precipitation and atmospheric circulation much better, please show the relative SST bias in Fig. 1c-f) and give the area mean temperature in the header (Johnson and Xie 2010; Johnson and Kosaka 2016; Bayr et al. 2018; Izumo et al. 2019). Response: The relative SST bias (with area average tropical temperature removed) is very similar to the original Figure 1, so is not included. We have produced a new plot showing the relative SST change (Fig S5) – see response to next comment.

Comment: Fig. 6: As for Fig. 1 I would strongly suggest to show the relative SST change (and area mean SST change in the 135 header), as this indicates the change in Walker Circulation (Bayr et al. 2014, 2020), which would be helpful to understand the precipitation change and ENSO amplitude change. **Response:** We have produced a new plot showing the relative SST change (a new version of Figure 6) which is included as Supplementary Figure S5. The absolute SST change is retained as Figure 6.

- Comment: Further, the change of Walker Circulation under different global mean temperatures is partly driven by the overall (homogeneous) warming (weakening under warmer and strengthening under colder mean climate, (Held and Soden 2006; Vecchi et al. 2006; DiNezio et al. 2011) and partly by the inhomogeneous warming (depends on the change of the SST gradient, Bayr and Dommenget 2013; Bayr et al. 2014). The best would be a more detailed analysis of the Walker Circulation changes to understand the ENSO amplitude change and precipitation change. But maybe you already get a clearer picture, when looking at the relative SST change. Response: As outlined above, we now include the relative SST
- 145 change as a new Supplementary Figure S5, and discuss this in the Section 4. We would prefer to leave evaluation of the changes in the Walker circulation for future studies.

Comment: Fig. 8 & 9: can you please show the multi model ensemble mean for each subfigure and the spread around as box plot. **Response:** We now include the multi-model mean values in Figure 8 and 9 and also in the new Table 2. Table 2 also includes the maximum and minimum model ENSO change values for each experiment.

150 Comment: Fig. 13: Please give the correlation values for each scatter plot. Response: We now include the correlation coefficients for the scatterplot figures (and the text in the final paragraph of Section 6 was modified slightly to refer to these correlations).

Comment: Further, I suggest to also look on how the wind-SST feedback changes in the scenarios, as from my experience and the study of (Vijayeta and Dommenget 2018) the change in the wind-SST feedback explain a large part of ENSO

155 amplitude change. The change in wind-SST feedback is strongly influenced by the change in the Walker Circulation (Bayr et al. 2018, 2020). Response: We would rather defer the analysis of the wind-SST feedback for future research which is planned on the Bjerknes Index.

Minor comments:

Comment: 75: mid-Holocene - please give the years BP. Response: Done.

160 Comment: 120: "was replaced by the Central Pacific-type El Niño" When? At the beginning of 21st century? Please make clearer. Response: This is clarified: in recent decades.

Comment: 132: You should also cite here (Latif and Keenlyside 2009). Response: This was added.

Comment: Fig. 1 & 2: What is the stippling? It is not mentioned in the figure caption. **Response:** The stippling is now defined in the Figure caption.

165 Comment: 255: "The Intertropical Convergence Zone (ITCZ) is generally shifted to the north". But also the rising branch of the Walker Circulation is shifted to the west (Bayr et al. 2018, 2020). This weakens the atmospheric feedbacks and hampers simulated ENSO dynamics. This should be discussed somewhere in the paper. Response: As this section is focused on precipitation biases in the models, we do not directly examine the atmospheric circulation. However we added a brief mention of the displacement of the Walker circulation due to SST biases here.

170 Comment: 429: "must consist of several processes." An other explanation is the nonlinear behavior of ENSO amplitude and SST gradient/thermocline slope as shown in Fig. 6 in (Hu et al. 2013). Response: This is now noted.

Comment: 460: "increase of the negative feedback by the mean current thermal advection" Another possible factor can be an increase of the negative heat flux damping as found in (Prigent et al. 2020) for the Atlantic Nino reduction since the year 2000. **Response:** This is an interesting suggestion but investigation of this mechanism is beyond the scope of the current

175 study.

Comment: 498: you should also mention here the bias ENSO dynamics due to the error compensation of the underestimated wind-SST and heat flux-SST feedback (Bayr et al. 2019) and weaker oceanic response (Kim et al. 2014). Response: These reference were added.

Comparison of past and future simulations of ENSO in **CMIP5/PMIP3 and CMIP6/PMIP4 models**

Josephine R. Brown¹, Chris M. Brierley², Soon-Il An³, Maria-Vittoria Guarino⁴, Samantha Stevenson⁵, Charles J. R. Williams^{6,7}, Qiong Zhang⁸, Anni Zhao², Pascale Braconnot⁹, Esther C. Brady¹⁰, Deepak Chandan¹¹, Roberta D'Agostino¹², Chuncheng Guo¹³, Allegra N. LeGrande¹⁴, Gerrit Lohmann¹⁵, Polina 185 A. Morozova¹⁶, Rumi Ohgaito¹⁷, Ryouta O'ishi¹⁸, Bette L. Otto-Bliesner¹⁰, W. Richard Peltier¹¹, Xiaoxu Shi¹⁵, Louise Sime⁴, Evgeny M. Volodin¹⁹, Zhongshi Zhang¹³, and Weipeng Zheng²⁰

¹School of Earth Sciences, University of Melbourne, Parkville, VIC, 3010, Australia

190 ²Department of Geography, University College London, London, WCIE 6BT, UK ³Department of Atmospheric Sciences, Yonsei University, Seoul, Korea ⁴British Antarctic Survey, High Cross, Madingley Road, CB3 0ET, Cambridge, UK ⁵Bren School of Environmental Sciences and Management, University of California, Santa Barbara, CA, USA ⁶School of Geographical Sciences, University of Bristol, University Road, Bristol BS8 1SS, UK

⁷Department of Meteorology, University of Reading, Earley Gate, P.O. Box 243, Reading RG6 6BB, UK 195 ⁸Department of Physical Geography and Bolin Centre for Climate Research, Stockholm University, 10691, Stockholm, Sweden

⁹Laboratoire des Sciences du Climat et de l'Environnement – IPSL, unite mixte CEA-CNRS-UVSO, Gif-sur-Yvette, France. ¹⁰National Center for Atmospheric Research, 1850 Table Mesa Drive, Boulder, CO 80305, USA

- 200 ¹¹Department of Physics, University of Toronto, 60 St. George Street, Toronto, Ontario M5S1A7, Canada ¹²Max-Planck-Institut für Meteorologie, Bundesstrasse 53, 20146, Hamburg, Germanv ¹³NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, Bergen, Norway 14NASA Goddard Institute for Space Studies, 2880 Broadway, New York, NY 10025, USA ¹⁵Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research Bussestr. 24, 27570 Bremerhaven, Germany
- 205 ¹⁶Institute of Geography Russian Academy of Sciences, Staromonetny L. 29, Moscow 119017, Russia ¹⁷Japan Agency for Marine-Earth Science and Technology, 3173-25 Showamachi, Kanazawa-ward, Yokohama, 236-0001, Japan ¹⁸Atmosphere and Ocean Research Institute, University of Tokyo, Kashiwa, Japan

¹⁹Marchuk Institute of Numerical Mathematics of the Russian Academy of Sciences, Moscow, Russia

²⁰State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics. Institute of 210 Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

Correspondence to Josephine R. Brown (josephine.brown@unimelb.edu.au)

Abstract. El Niño-Southern Oscillation (ENSO) is the strongest mode of interannual climate variability in the current climate, influencing ecosystems, agriculture and weather systems across the globe, but future projections of ENSO frequency

- 215 and amplitude remain highly uncertain. A comparison of changes in ENSO in a range of past and future climate simulations can provide insights into the sensitivity of ENSO to changes in the mean state, including changes in the seasonality of incoming solar radiation, global average temperatures and spatial patterns of sea surface temperatures. As a comprehensive set of coupled model simulations are now available for both palaeoclimate time-slices (the Last Glacial Maximum, mid-Holocene and Last Interglacial) and idealised future warming scenarios (one percent per year CO₂ increase, abrupt four times
- 220 CO2 increase), this allows a detailed evaluation of ENSO changes in this wide range of climates. Such a comparison can assist in constraining uncertainty in future projections, providing insights into model agreement and the sensitivity of ENSO to a range of factors. The majority of models simulate a consistent weakening of ENSO activity in the Last Interglacial and mid-Holocene experiments, and there is an ensemble mean reduction of variability in the western equatorial Pacific in the Last Glacial Maximum experiments. Changes in global temperature produce a weaker precipitation response to ENSO in the
- 225 cold Last Glacial Maximum experiments, and an enhanced precipitation response to ENSO in the warm increased CO₂ experiments. No consistent relationship between changes in ENSO amplitude and annual cycle was identified across experiments.

1 Introduction

When the first El Niño-Southern Oscillation (ENSO) event occurred in Earth's history is unclear. However, in light of air-230 sea coupled feedbacks, the birth of ENSO must be strongly related to the emergence of the tropical eastern Pacific cold tongue and its zonal sea surface temperature (SST) contrast with the tropical western Pacific warm pool. It has been proposed that the gradual uplifting of the Central American continent since around 24 million years BP (before present) triggered the development of the Pacific cold tongue by reducing the surface water exchange between oceans (e.g. Chaisson and Ravelo, 2000). Palaeo proxies revealed that the Pliocene warm period (~4.5–3.0 million years BP) recorded a very weak

- 235 zonal SST contrast, sometimes referred to as a "permanent El Niño-like state" (Brierley et al., 2009; Fedorov et al., 2013; Fedorov et al., 2006; Ravelo et al., 2006; White and Ravelo, 2020). Despite the weak mean east-west SST gradient, some proxy records (Scroxton et al., 2011; Watanabe et al., 2011) and General Circulation Model (GCM) experiments (Burls and Fedorov, 2014; Haywood et al., 2007) suggest the existence of interannual ENSO variability during the mid-Pliocene, at least sporadically. During the last interglacial (approximately 116-129 thousand years BP, hereafter ka), orbital changes
- 240 appear to have influenced the strength of ENSO variability, with coral proxy records providing evidence for relatively weak ENSO amplitude (Hughen et al., 1999; Tudhope et al., 2001). Model studies have also simulated reduced ENSO variability during the last interglacial (An et al., 2017; Kukla et al., 2002; Salau et al., 2012).

Lee sheet dynamics have also been thought to influence the behaviour of ENSO (Liu et al., 2014). Large-scale ice sheets in the Northern Hemisphere expanded from 2.7 million years BP onwards (Jansen et al., 2000). Global climate subsequently

Deleted: Ma (
Deleted:)	

De	e	te	d:	M
----	---	----	----	---

Deleted:, prior to the complete blocking of Central American Seaway around 2.7 Ma BP (Bacon et al., 2015; O'Dea et al., 2016),.

Deleted: 1	
Deleted: Ma	

underwent a series of glacial-interglacial cycles, with the most recent glacial period reaching maximum levels of global cooling and lower sea levels around 21~18 ka, the so-called "Last Glacial Maximum" (LGM). The tropical climate state during the LGM was 1–3°C colder on average than the present day. Reconstructions of LGM ENSO activity are uncertain,

- 255 with some studies finding increased ENSO variability (Koutavas and Joanides, 2012; Sadekov et al., 2013) and others finding reduced ENSO variability (Leduc et al., 2009). A recent synthesis of evidence from planktonic foraminifera (Ford et al., 2015; Ford et al., 2018) supports reduced ENSO variability but increased seasonality and a deepened equatorial Pacific thermocline during the LGM. Model simulations using an isotope-enabled GCM (Zhu et al., 2017) further assist in reconciling the proxy records, as the model simulates a 30% weakening of ENSO during the LGM but an increased annual
- 260 cycle contributing to enhanced variability in the foraminifera records of Koutavas and Joanides (2012). Coupled climate models included in the 2nd and 3rd phases of the Paleoclimate Modelling Intercomparison Project (PMIP2 and PMIP3, respectively) simulate a wide range of ENSO changes for the LGM (Masson-Delmotte et al., 2014; Saint-Lu et al., 2015; Zheng et al., 2008).
- A number of proxy records provide evidence for <u>weakened</u> ENSO variability during the mid-Holocene (around 6 ka),
 although the timing of this weakening varies between records (e.g. Carré et al., 2014; Conroy et al., 2008; Donders et al., 2005; Koutavas and Joanides, 2012; Koutavas et al., 2006; McGregor and Gagan, 2004; McGregor et al., 2013; Rein et al., 2005; Riedinger et al., 2002; Tudhope et al., 2001; White et al., 2018). Cobb et al. (2013) argued that coral records from the central Pacific do not show a statistically significant reduction in mid-Holocene ENSO variability but a new ensemble of central Pacific records (Grothe et al., 2019) provides evidence for a significant reduction from 3-5 ka. Some studies have
- 270 suggested that disagreement between the magnitude of mid-Holocene ENSO reduction in different proxy records may be due to shifts in the spatial pattern of ENSO variability between the eastern and central Pacific (e.g. Carré et al., 2014; Karamperidou et al., 2015). A synthesis of Holocene ENSO proxy records (Emile-Geay et al., 2016) identifies a sustained reduction in ENSO variability from 3-5 ka, with a reduction of 64% in the central Pacific. During the earlier "mid-Holocene" period from 5.5-7.5 ka, reduced ENSO variance occurs in central, western and eastern Pacific (66%, 50% and 33% procession) with large uncertainty and the central Pacific (66%, 50% and 33%).

275 respectively) with larger uncertainty ranges (Emile-Geay et al., 2016).

Climate models generally simulate reduced mid-Holocene ENSO activity. For example, transient simulations for part or the whole Holocene period using an intermediate ocean-atmosphere coupled model of the tropical Pacific climate forced by the orbital forcing (Clement et al., 2000), a fully coupled general circulation model with the time-varying climate forcing including orbital, greenhouse gas, meltwater flux, and continental ice sheets (Liu et al., 2014), and a hybrid-type simulation

280 using a combination of the intermediate complexity of earth system model forced orbital forcing and intermediate coupled tropical pacific climate model with varying background state (An et al., 2018), all showed a significant reduction of ENSO intensity during mid-Holocene and its recovery to modern-day ENSO strength around the late Holocene. A study with another set of transient Holocene simulations with coupled climate models confirmed this result, but found that it was the result of chaotic processes (Braconnot et al., 2019). Deleted: (thousand years) BP

Deleted: a weakening of

Deleted: . On the other hand,

The mid-Holocene time-slice simulations of PMIP2 and PMIP3, all of which fixed climate forcing at 6,000 years ago (6,ka), also showed suppressed ENSO variability in most of the models compared to the pre-industrial perpetual simulations
(Braconnot et al., 2007; Chen et al., 2019b; Chiang et al., 2009; Zheng et al., 2008). The reduction of interannual variability in PMIP models was especially dominant over the equatorial central Pacific (An and Bong, 2018; Chen et al., 2019b). However, the reduction of ENSO intensity in the 6,ka run of PMIP3 compared to the 0,ka run (~ 5% reduction in the standard deviation of NINO3.4 index from 11 models), in which more state-of-the-art GCMs participated, was rather weaker than that in PMIP2 (~18% reduction in the standard deviation of NINO3.4 index from 6 models) (An and Choi, 2014;
Masson-Delmotte et al., 2014). A comprehensive model-data comparison (Emile-Geay et al., 2016) found that models underestimated the reduction in mid-Holocene ENSO variability compared with proxy records, and also simulated an inverse relationship between the amplitude of the seasonal cycle and ENSO variability which was not evident in proxy reconstructions.

Over the last millennium, ENSO has exhibited considerable natural variability (Cobb et al., 2003). Multi-proxy reconstructions of central tropical Pacific SST confirm that vigorous decadal to multi-decadal variability of ENSO occurred (Emile-Geay et al., 2013) while eastern Pacific ocean sediment records suggest a mid-millennium shift from damped to amplified ENSO variability (Rustic et al., 2015). However, as for all the paleoclimate intervals considered, the assessment of ENSO variability over the last millennium is rather uncertain due to the temporal and spatial sparseness of palaeo-ENSO proxy records (Cobb et al., 2003; Cobb et al., 2013; Khider et al., 2011). Last Millennium experiments from PMIP3 models
showed that ENSO behaviour may be strongly modulated on decadal- to centennial time scales over the last millennium, and that teleconnections between ENSO and tropical Pacific climate vary on these time scales (Brown et al., 2016; Lewis and LeGrande, 2015).

The instrumental records of ENSO for the 20th century clearly document the variety of ENSO behaviour, including both temporal and spatial complexity (Timmermann et al., 2018). ENSO complexity includes its seasonal phase locking (Neelin

- 310 et al., 2000; Stein et al., 2011), the interaction with other time-scale climate variability (Eisenman et al., 2005; Levine et al., 2016; Tang and Yu, 2008; Zhang and Gottschalck, 2002), El Niño-La Niña asymmetry in amplitude, duration, and transition (An and Jin, 2004; An and Kim, 2018, 2017; Im et al., 2015; Okumura et al., 2011), the diversity in the peak location (i.e., Central Pacific- and Eastern Pacific-type El Niño; (Capotondi et al., 2015), and the combination modes due to interaction between annual and interannual spectra (Stuecker et al., 2015; Timmermann et al., 2018). Interestingly, the dominant
- 315 tendency of Eastern Pacific-type El Niño occurrence during 20th century was replaced by the Central Pacific-type El Niño in recent decades (Ashok et al., 2007; Yeh et al., 2014; Yeh et al., 2009), and all extreme El Niño events (1982-83, 1997-98, and 2015-16) recorded by the modern instruments occurred around/after the late 20th century. The increased frequency of Central Pacific-type events in recent decades is unusual in the context of a palaeo record for the last 400 years (Freund et al., 2019). Such distinct changes in ENSO characteristics through 20th and 21st centuries may be related to low-frequency
- 320 modulation by natural variability or the global warming trend due to increasing greenhouse gas concentrations, or a
 - 9

combination of natural and anthropogenic factors (e.g. An et al., 2008; Cai et al., 2015a; Collins, 2000; Gergis and Fowler, 2009; Timmermann et al., 1999; Trenberth and Hoar, 1997; Yang et al., 2018; Yeh et al., 2014; Yeh and Kirtman, 2007).

Although it is still a topic of ongoing debate as to whether the future tropical Pacific climate state becomes "El Niño-like" or "La Niña-like" (referring only to the change in zonal SST gradient) in response to greenhouse warming (An et al., 2012;

- 325 Cane et al., 1997; Collins et al., 2010; Lian et al., 2018; Seager et al., 2019), recent multi-model studies of projected changes in ENSO under global warming suggested no significant change in terms of mean ENSO amplitude compared to the historical ENSO amplitude (An and Choi, 2015; An et al., 2008; Chen et al., 2017; Christensen et al., 2014; Latif and Keenlyside, 2009; Stevenson, 2012). However, studies have identified robust increases in the extreme hydrological changes associated with El Niño (Cai et al., 2014; Cai et al., 2015a) and La Niña (Cai et al., 2015c), and changes in ENSO-driven
- 330 precipitation variability (Power et al., 2013). Moreover, even if the global mean temperature is constrained to the limit of 1.5 °C above pre-industrial levels following the Paris Agreement, a doubling of the frequency of extreme El Niño events may occur (Wang et al., 2017).

In this paper, we assess ENSO change through time as simulated in the new generation of coupled atmosphere-ocean climate models for both past and future climates (see Section 2.2). We also compare these new simulations with previous generations

- 335 of climate models. Detailed comparison of the past climate simulations with proxy records is beyond the scope of the current study, and will be the focus of subsequent research. We consider the change in ENSO amplitude, and its dynamical relationship with the change in the mean climate state under past and future conditions spanning colder past climates (Last Glacial Maximum), past climates with an altered seasonal cycle (Last Interglacial and Mid-Holocene), and idealised warming projections (abrupt four times CO₂ and one percent per year CO₂) to provide a context for evaluating projections of
- 340 ENSO change. The methods, models and experiments are introduced in Section 2. Model evaluation is provided in Section 3. In Section 4, the mean state changes in each experiment relative to the pre-industrial control are described. ENSO amplitude changes are presented in Section 5 and changes to ENSO teleconnections are considered in Section 6. In Section 7, the proposed mechanism for the ENSO change through the time are briefly discussed, and conclusions are given in Section 8.

2 Methods

345 This research analyses a total of 140 simulations, across 7 different experiments and 32_r climate models. The description of individual simulations is therefore kept brief, and often only the ensemble mean response will be shown. The combined model ensemble will be described in Section 2.1, whilst an overview of the experimental designs are provided in Section 2.2. The common analysis procedure is outlined in Section 2.3.

Deleted: 28

350 2.1 Models

State-of-the-art coupled global climate models solve the physical equations of the atmosphere and ocean. They are some of the most sophisticated of numerical models, and have been constantly developed for several decades. Globally there are around 40 such models with varying degrees of independence (Knutti et al., 2013). Given the resources required to undertake a single GCM simulation, the international community has settled upon a series of coordinated experiments to facilitate

- 355 model to model "intercomparison". These are organised under the umbrella of the Coupled Model Intercomparison Project (CMIP). Here we evaluate and analyse simulations from both the previous phase 5 (CMIP5; Taylor et al., 2012), as well as early results from the current phase 6 (CMIP6; Eyring et al., 2016a). Simulations of past climate are included from the Paleoclimate Modelling Intercomparison Project (PMIP), which is part of CMIP. Some of the simulations were carried out as part of PMIP phase 3 (PMIP3; Braconnot et al., 2012) and other simulations are part of PMIP phase 4 (PMIP4; Kageyama
- 360 et al., 2018).

This study provides an opportunity to compare the simulation of ENSO in past and future climates in CMIP5 and CMIP6 generations of models. The simulation of ENSO may be improved in the CMIP6 ensemble relative to the CMIP5 models due to some improvements in the simulation of the mean state, such as a reduced "double ITCZ bias" in tropical Pacific precipitation (Tian and Dong, 2020) and a reduced "cold tongue bias" in equatorial Pacific SSTs (Grose et al., 2020), as

- 365 shown in Supplementary Figures S1 and S2. Some CMIP6 models also include a more sophisticated treatment of aerosols, higher spatial resolution and updated parameterisation of process such as convection compared with CMIP5 models. For inclusion in this study, a model must have both completed at least one palaeoclimate simulation and provided the required output fields for at least 30 years for both this simulation and the pre-industrial control (see section 2.3 for details). The resulting 32 models are listed in Table 1, combined they contain over 35,000 years of monthly ENSO information.
- 370
 -) Further information about the CMIP5 models is provided in Table 9.A.1 of Flato et al. (2014). The CMIP6 models used in this study are described in more detail in the Supplementary Information and also online on the PMIP4 website https://pmip4.lsce.ipsl.fr/doku.php/database:participants.

2.2 Simulations

This study uses simulations consisting of seven different experiments. Four of the experiments are part of the CMIP effort, and form part of the "DECK" set of core simulations (Eyring et al., 2016a): the Pre-industrial and Historical and two idealised future warming scenarios. The study also includes three past climate experiments from the PMIP database, the mid-Holocene and Last Glacial Maximum were included in both PMIP3/CMIP5 and PMIP4/CMIP6, whereas the Last Interglacial was only included in PMIP4/CMIP6.

The mean state and ENSO variability of the models are evaluated in simulations with prescribed historical forcings, known as *historical* simulations (Hoesly et al., 2018; Meinshausen et al., 2017; van Marle et al., 2017). The specification of the *historical* simulation differs slightly between CMIP5 and CMIP6, most notably by the CMIP6 simulations being extended

11

Deleted: 28

until 2015 C.E. This has minimal influence over the chosen climatological period of 1971-2000. The baseline simulation relative to which all climate changes are calculated is the pre_industrial control (*piControl*; Eyring et al., 2016a; Stouffer et al., 2004). The *piControl* simulations represent constant 1850 forcing conditions, and have all reached a quasi-stable equilibrium.

The two idealised warming scenarios (*abrupt4xCO2* and *lpctCO2*) in the CMIP DECK both involve increases in carbon dioxide concentrations. The *abrupt4xCO2* experiment imposes an instantaneous quadrupling of carbon dioxide, to which the coupled climate system is left to equilibrate. The experiment was devised to calculate the climate sensitivity (Gregory et al.,

390 2004). The *lpctCO2* experiment is forced with a carbon dioxide increase of one percent per year. This compound increase achieves a quadrupling of carbon dioxide after 140 years, but the climate system is still highly transient. This experiment can be used to calculate the transient climate response (Andrews et al., 2012).

The experimental design for the mid-Holocene (*midHolocene*) and Last Interglacial (*lig127k*) simulations is given by Otto-Bliesner et al. (2017). All *midHolocene* and *lig127k* simulations should have followed this protocol, such that the only significant differences to their corresponding DECK *piControl* simulation are the astronomical parameters and the atmospheric trace greenhouse gas concentrations. In short, astronomical parameters have been prescribed according to orbital constants from Berger and Loutre (1991) and atmospheric trace greenhouse gas concentrations are based on recent reconstructions from a number of sources (see section 2.2 in Otto-Bliesner et al. (2017) for details). Note that the different orbital configurations for *midHolocene* and *lig127k* result in different seasonal and latitudinal distribution of top-of-

400 atmosphere insolation compared to the DECK *piControl*. For other boundary conditions, these are either small and locally constrained (e.g. for ice sheets) or there is insufficient spatial coverage to give an informed global estimate (e.g. for vegetation). These other boundary conditions, including solar activity, palaeogeography, ice sheets, vegetation and aerosol emissions, were therefore kept as identical to each model's DECK *piControl* simulation. In cases where a boundary condition can either be prescribed or interactive, such as vegetation, the *midHolocene* and *lig127k* simulations followed the 405 set-up used in the *piControl* (Otto-Bliesner et al., 2017).

The Last Glacial Maximum (*lgm*) simulation is focused on representing the glacial climate of 21,000 years ago. During this period, carbon dioxide concentrations dropped by around 100 ppm and large ice-sheets covered the land masses in the Northern mid to high latitudes. The precise specification of the ice coverage and volume varies between the CMIP5 (Abe-Ouchi et al., 2015) and CMIP6 (Kageyama et al., 2017) specifications. Whilst this will impact the teleconnections (Jones et al., 2018), its impact on the tropical Pacific is unclear. The implementation of land-sea changes in regions such as Maritime

410 al., 2018), its impact on the tropical Pacific is unclear. The implementation of land-sea changes in regions such as Maritime Continent is also important (Di Nezio et al., 2016).

2.3 Indices and Analysis

This analysis uses a series of standard metrics and measures to describe the simulated ENSO response. These are achieved using the Climate Variability Diagnostics Package (Phillips et al., 2014), which is part of the ESMValTool (Eyring et al.,

12

Deleted: Whilst a full equilibrium of the coupled system will not have occurred within the specified 150 years, there has been substantial progress towards it. The

2016b). This software package has previously been used to explore variability in palaeoclimate simulations, although in the tropical Atlantic rather than Pacific (Brierley and Wainer, 2018). The model output variables required for the analysis are

- 420 monthly precipitation rate, monthly surface air temperature and monthly surface temperature. The <u>surface</u> temperature, also known as skin temperature, is utilised to provide SST on the atmospheric grid (Juckes et al., 2019). Prior to undertaking the ENSO analysis, the monthly fields of the palaeoclimate simulations are adjusted to represent the changes in the calendar (i.e. due to changes in the length of months or seasons over time, related to changes in the eccentricity of Earth's orbit and precession), using the PaleoCalAdjust tool (Bartlein and Shafer, 2019).
- The mean state of the present day tropical Pacific is determined using a climatology over the period 1971-2000 for both the *historical* simulations and observational datasets: the Global Precipitation Climatology Project (GPCP; Adler et al., 2003), the Hadley Centre Sea Ice and Sea Surface Temperature (HadISST; Rayner et al., 2003) and the 20th Century Reanalysis (C20; Compo et al., 2011)). For the transient *abrupt4xCO2* and *IpctCO2* simulations, the climatology is computed over the final thirty years. For the other quasi-equilibrium simulations, all available data <u>are</u>, considered to create the climatology.

interpolating onto a common 1° by 1° grid, before averaging across the ensemble members (Brierley and Wainer, 2018).

The state of the tropical Pacific is tracked by the SST anomalies in the NINO3.4 region (5°S-5°N, 120-170°W) (Trenberth, 1997). The anomalies are computed with respect to each simulation's own climatology, with a linear trend removed. All available years are used to assess and composite the NINO3.4 index (even in the transient simulations with a defined

435 climatological period). This choice maximises the number of ENSO events that can be assessed, although it does require the assumption that changes in the background climatology progress linearly. This assumption is less valid for the *abrupt4xCO*₂ experiment than the *lpctCO*₂ experiment, however the ENSO responses show expected coherence across the ensemble, implying the errors introduced are not significant.

The normalised NINO3.4 timeseries are used to composite all years greater than 1 standard deviation to represent El Niño year and all years less than -1 standard deviation to represent La Niña year (Deser et al., 2010). The standard behaviour of the Climate Variability Diagnostics Package is to use December values of the monthly NINO3.4 time series smoothed with a 3-point binomial filter to identify seasons to composite (Phillips et al., 2014). The process is modified here to allow for changes in the seasonal peak of ENSO activity, potentially associated with the orbital variations. Instead of using smoothed December NINO3.4 values to classify ENSO events, the 3-month smoothed NINO3.4 timeseries is calculated for every month, and the maximum anomaly identified for any month (with a year counted from June to May). This added flexibility

does not quite replicate the standard behaviour over the period 1960-2010, because it additionally identifies 1987 as an El Niño that peaked in August. Previous research has accepted this as a valid El Niño event (e.g. Ramanathan and Collins, 1991). Deleted: latter

(Deleted: is

3 Model Evaluation

The ability of the CMIP5/PMIP3 and CMIP6/PMIP4 models (hereafter, "CMIP" models) used in this study to simulate the present-day SST pattern is first evaluated in comparison with HadISST observations (Rayner et al., 2003) as shown in Figure 1. Consistent with other studies of coupled GCMs (Bellenger et al., 2014; Collins et al., 2010), the models are generally

- 455 biased toward overly cold SSTs in the central to western equatorial Pacific. Biases toward overly warm SSTs are present in the far eastern Pacific, again a common feature of GCMs generally resulting from a lack of sufficiently deep stratocumulus decks in eastern boundary regions (see review by Ceppi et al., 2017). The northern subtropics also appear to have similar biases, with colder-than-observed SSTs in the central Pacific and warm biases near the western coast of Mexico. The effect of comparison time period selection is apparent when the *historical* and *piControl* simulations are contrasted (Figure 1c,d
- 460 versus 1e,f); the *piControl* climate is roughly 1°C colder on average than the *historical*, leading to an apparent exacerbation of the cold-tongue bias and reduction in the warm bias in the eastern equatorial Pacific. <u>The tropical Pacific SST biases in the newer CMIP6 models are smaller on average than those in the CMIP5 models (see Supplementary Figure S1).</u>

SST biases contribute to errors in the representation of precipitation in the simulations. The Intertropical Convergence Zone (ITCZ) is generally shifted to the north (Figure 2c,d) leading to a dry bias in the equatorial Pacific, which is particularly

- 465 pronounced during DJF (Figure 2c). The equatorial cold SST bias also leads to the rising branch of the Walker Circulation being shifted to the west, weakening atmospheric feedbacks (Bayr et al., 2018). South of the equator, there is a wet bias in the location of the climatological South Pacific Convergence Zone (SPCZ) (Figure 2c,d) consistent with previously documented tendencies for CMIP-class models to produce a so-called "double ITCZ" (Adam et al., 2018; Zhang et al., 2015). Once again, there are substantial differences between precipitation fields in the *historical* and *piControl* simulations
- 470 (Figure 2c,d vs. 2e,f), with equatorial precipitation generally increased in *historical*. This is consistent with the expected intensification of the hydrological cycle under climate change (Held and Soden, 2006; Vecchi and Soden, 2007), for which some observational evidence exists during the 20th century (Durack et al., 2012). These differences are confounded somewhat by the slight differences in the composition of the ensembles for *piControl* and *historical* simulations (see Table 1). The CMIP6 models simulate smaller biases in tropical Pacific precipitation than CMIP5 models (see Supplementary Figure S2), consistent with the improved SST distribution.

The spatial pattern of ENSO sea surface temperature anomalies is illustrated using the ensemble mean difference between composite El Niño and La Niña events, shown in Figure 3. The magnitude of simulated *historical* and *piControl* events (Figure 3b,c) is quite close to the observed value (Figure 3a), with peak SST anomaly values of roughly 2.5°C. However, the "centre of action" for ENSO is shifted westward relative to observations; this is a known feature of coupled GCMs, and is related to the biases in mean SST (Bellenger et al., 2014). Because of this westward shift in the peak of El Niño and La Niña events, the magnitude of SST variability is overly weak in the far eastern Pacific (Figure 3b, c). There may be substantial variation between the individual model simulations, which is documented elsewhere (e.g. Bellenger et al., 2014). The ENSO

SST anomalies in CMIP6 models are stronger than those in CMIP5 models in both the western and eastern equatorial Pacific (see Supplementary Figure S3).

- We also evaluate the simulation of global temperature teleconnections with ENSO variability (Figure 4). The observed warming over northern South America, Australia, and much of Southeast Asia during DJF of the El Niño event peak (Figure 4a) is reproduced by the CMIP ensemble mean (Figure 4c), although the magnitudes of the temperature anomalies appear weaker than observed. The teleconnection to the Atlantic and Indian Oceans likewise appears reliable, with comparable magnitudes of surface warming appearing in the models relative to observations. The models appear to have the most
- 490 difficulty in representing teleconnections to the higher latitudes; the strong warming over northern North America during DJF (Figure 4a) is significantly underestimated in the models (Figure 4c,e), as is the cooling over northern Eurasia. The same general tendencies hold during JJA (Figure 4b,d,f); here, notable model-observation disagreements are apparent over the eastern half of North America, southern South America, and the southwestern Pacific. This latter feature may relate to model difficulties with representing SPCZ dynamics (Brown et al., 2013).
- 495 Model performance in simulating ENSO temperature teleconnections is reflected in the structure of ENSO precipitation teleconnection biases, shown in Figure 5. In DJF, when El Niño events typically peak, drying occurs in the western Pacific warm pool and over the Amazon in the reanalysis (Figure 5a); this drying persists in JJA but is reduced (Figure 5b). In both cases, the models underestimate the magnitude of South American precipitation teleconnections; additionally, the western Pacific drying is shifted westward due to the bias in mean SST (Figure 5c,d). Precipitation teleconnections to North America 500 are overly weak in the simulations during both DJF and JJA, as are the tropical Atlantic anomalies.

In summary, the spatial pattern of ENSO SST variability and the remote teleconnections of temperature and precipitation in response to ENSO are reasonably well simulated in the CMIP models and particularly in the ensemble mean. We therefore examine the changes in ENSO in these models under a range of past and future climate conditions.

4 Mean State Changes

- 505 Changes in the mean state of the tropical Pacific are evaluated for each experiment relative to the *piControl*. This provides the context for consideration of changes in ENSO amplitude and teleconnections in the subsequent Sections. Figures 6 and 7 summarise the seasonal response (DJF and JJA) of the model ensemble to different forcing during the three palaeoclimate experiments (*midHolocene*, *lgm*, and *lig127k*), and two idealised future warming scenarios (*lpctCO2*, and *abrupt4xCO2*) for surface temperature (Figure 6) and precipitation (Figure 7).
- 510 For the Last Interglacial (*lig127k*), ensemble changes in surface temperatures (Figure 6e, f) exhibit a strong seasonality that is consistent with *lig127k* minus *piControl* insolation anomalies (see Otto-Bliesner et al., 2017). More specifically, in JJA, regions located at tropical and subtropical latitudes show warming (of about 0.5°C to 2°C). Indeed, during boreal summer, positive insolation anomalies reach their maximum in the Northern Hemisphere (NH) and extend into the tropics and the

Southern Hemisphere (SH). In contrast, in DJF, negative insolation anomalies are large in SH and NH equatorward of 40°N,
and tropical and subtropical latitudes show cooling (mostly of about 1°C). Similar patterns, although much weaker and spatially constrained, are shown for the *midHolocene* simulation (Figure 6a and Figure 6b for DJF and JJA, respectively), with small anomalies in both seasons. A small cooling of ~0.5°C is shown over the equatorial regions of the eastern (central) Pacific during DJF (JJA). No warming, however, is shown in the *midHolocene*, in either season. Based on the sign and magnitude of the insolation anomalies for both the *midHolocene* and (more so) *lig127k*, the model ensemble mean state
changes for both of these simulations are consistent with previous modelling and proxy reconstruction studies (see e.g. Otto-

- Bliesner et al., 2013).¹ The *lgm* ensemble mean shows cooling of around 2–3°C in the tropical Pacific in both DJF and JJA (Figure 6c,d), consistent with previous modelling studies and proxy reconstructions (Ballantyne et al., 2005; MARGO Project Members et al., 2009;
- Masson-Delmotte et al., 2014; Otto-Bliesner et al., 2009). For the idealised future scenarios, the tropical Pacific warms (by 1-3°C in the *lpctCO2* case and more than 3°C in the *abrupt4xCO2* case), with largest warming in the equatorial region. In the case of the *abrupt4xCO2* simulations, the ensemble mean warming is largest in the eastern and central Pacific, particularly in DJF; whereas the *lpctCO2* ensemble shows enhanced warming extending across the equatorial Pacific. This enhanced equatorial warming is a recognised feature of anthropogenic climate warming (DiNezio et al., 2009; Liu et al., 2005; Xie et al., 2010) and has important implications for ENSO, leading to more frequent extreme El Niño events in a
- 530 warmer climate (Cai et al., 2014; Wang et al., 2017).

Regarding ensemble precipitation anomalies, in the *lig127k* simulations (Figure 7e,f), the weaker Australian monsoon (drier conditions over northern Australia in DJF) and the enhanced North America monsoon (wetter conditions over northern South America in JJA) are consistent with a northward shift of the mean seasonal position of the ITCZ (over the oceans) and the associated tropical rainfall belt (over the continents). As for surface temperatures, the *midHolocene* ensemble mean (Figure

- 535 7a,b) shows similar spatial patterns of mean state precipitation change to the *lig127k* case, but with smaller magnitude. The Australian monsoon (North American monsoon) is weaker (stronger) in the *midHolocene* ensemble mean relative to *piControl*, just less so than the *lig127k*. Drying in the *midHolocene* is larger in the western Pacific than for *lig127k*. Mechanisms and drivers for precipitation changes over the tropics in Last Interglacial climates are still unclear and represent an active area of research (see e.g. Scussolini et al., 2019; Otto-Bliesner et al. *in prep*.).
- 540 Precipitation changes in the *lgm* ensemble (Figure 7 c,d) show a drying over the Maritime Continent, Australia and Southeast Asia. Precipitation in the ITCZ over the tropical Pacific is also reduced, particularly in JJA, in response to cooler SSTs. Precipitation is increased in the western Pacific and on the northern edge of the SPCZ, indicating a northward displacement of the SPCZ as found in previous studies (Saint-Lu et al., 2015). In the *lpctCO2* and *abrupt4xCO2* simulations (Figure 7 g-j), precipitation increases in the equatorial Pacific where SST warming is greatest (e.g. Chadwick et al., 2013; Xie et al.,

¹ Further analysis and description of the results of these PMIP4 experiments can be found in other articles in this special issue – e.g. Brierley et al. for *midHolocene* and Otto-Bliesner et al. for *lig127k*.



545 2010), with some drying on the northern edge of the ITCZ, particularly in the eastern Pacific. Drying also occurs in the southeast Pacific, where warming is relatively small and trade winds are intensified, leading to drying of the eastern edge of the SPCZ in DJF (Brown et al., 2013; Widlansky et al., 2013).

Previous studies have noted that changes in tropical precipitation are strongly influenced by the spatial pattern of SST change (Chadwick et al., 2013; Xie et al., 2010). Such changes in SST gradients are also linked to changes in the Walker
550 Circulation (Bayr and Dommenget 2013; Bayr et al., 2014). We therefore also plot the relative SST change, with the mean over the tropical domain (25°N-25°S, 100°E-80°W) subtracted, in Supplementary Figure S5. Comparison of the precipitation changes in Figure 7 and the relative SST changes in Supplementary Figure S5 confirms that there is generally close agreement, with regions of relative warming experiencing increased precipitation and regions of relative cooling becoming drier in the ensemble mean.

555 5 ENSO Amplitude Changes

The amplitude of ENSO, as measured by the standard deviation of SST from the NINO3.4 region, is shown for each model in each experiment in Figure 8. The amplitudes of ENSO in the *piControl* simulation from each model are also shown for reference. The percentage change in ENSO amplitude in the experiments relative to *piControl* is shown in Figure 9 and the ensemble mean change and minimum and maximum model changes are given in Table 2.

- 560 In the *midHolocene* simulations, a large majority (26 out of 30) of models show a decrease in ENSO variability, with the only exceptions being CMIP5 models CSIRO-Mk3-6-0 and MRI-CGCM3 and CMIP6 models INM-CM4-8 and NorESM2-LM. Most of these changes are small, however, with few models showing more than a ~20% decrease (Figures 8a and 9a). This is consistent with previous model studies, which generally show a smaller reduction in *midHolocene* ENSO amplitude than implied by proxy records (e.g. Emile-Geay et al., 2016), as discussed in Section 1. A similarly consistent reduction in
- 565 ENSO amplitude is found for the *lig127k* simulations (Figure 8c and 9c), with <u>10 out of 12 models showing a reduction in</u> amplitude, typically of at least 20%.

In contrast, a much less consistent response is found for the *lgm*, *lpctCO2* and *abrupt4xCO2* simulations. In all of these simulations, the sign of change in amplitude of ENSO is approximately equally spread between increases and decreases across the set of models. In the *lgm* simulations (Figure 8b and 9b), some models (e.g. FGOALS-g2) show a large decrease

- 570 in variability of over 40% and others (e.g. IPSL-CM5A-LR) show large increases of up to ~40%. Likewise, in the *lpctCO2* simulations (Figure 8d and 9d), ENSO variability is again highly model-dependent, with the range including large decreases of over ~20% in some models (e.g. CCSM4) to large increases of up to ~40% in others (e.g. MPI-ESM-P). The same is true for the *abrupt4xCO2* simulations (Figure 8e and 9e), with the range including large decreases of over ~40% in some models (e.g. GISS-E2-R) to large increases of up to ~50% in others (e.g. CSIRO-Mk3-6-0). This is consistent with previous studies
- 575 showing little agreement on future projections of ENSO amplitude change (Collins et al., 2010; Collins et al., 2014).

17

Deleted:

Deleted:	4	
Deleted:	27	
Deleted:	INM CM4 8	

eleted: , INM-CM4-8

Deleted: 9

Deleted: s

Comparing the absolute magnitudes of ENSO amplitude in all simulations (Figure 8), the standard deviation ranges (i.e. between the models showing the smallest and largest standard deviations) are $\sim 0.7^{\circ}$ C in the *midHolocene*, $\sim 1.1^{\circ}$ C in the *lgm*,

- 585 ~0.2°C in the *lig127k*, ~0.7°C in the *lpctCO2* simulation and ~1°C in the *abrupt4xCO2* simulation. The cold climate simulation and the extreme future run therefore show the largest spread between models, suggesting a lack of model agreement, whereas the *mid-Holocene* and *lig127k* simulations as well as the gradual future run have a much smaller spread between models.
- Comparing the change in ENSO amplitude in all simulations (Figure 9), we find that the *midHolocene* and *lig127k*simulations have high inter-model agreement on the sign of the response, consistently showing lower ENSO variability relative to the *piControl* in both cases. The common factor between these simulations is the change in seasonality of insolation, which in both cases is increased in boreal summer, leading to a damped ENSO via a range of mechanisms discussed in Section 1 and also Section 7. In contrast, there is much less inter-model agreement in the cold climate simulation (i.e. the *lgm*) and the gradual and extreme future warming runs (i.e. the *lpctCO2* and *abrupt4xCO2* simulations, respectively).
- Comparison of ENSO amplitude changes in CMIP5 and CMIP6 model ensembles shows generally strong agreement between the two generations of models (see Table 2 and Supplementary Figure S4). The ensemble mean reduction in *midHolocene* ENSO strength is somewhat greater for CMIP6 models than CMIP5 models, while there is no *lig127k* experiment in CMIP5. Both sets of models simulate weaker ENSO amplitude on average in the *lgm* experiments, although there are only 3 CMIP6 models available for this experiment. Both CMIP5 and CMIP6 models simulate an ensemble mean increase in ENSO amplitude in the idealised future *lpctCO2* and *abrupt4xCO2* experiments, although with large inter-model

spread (see Figure 9 and Supplementary Figure S4).

6 ENSO Patterns and Teleconnections

The anomalous pattern of El Niño minus La Niña SST composite for each experiment relative to *piControl* (see Figure 3c) is shown in Figure 10. The *midHolocene* and *lig127k* patterns (Figure 10a, c) show negative SST anomalies in the central equatorial Pacific, indicating weakening of event amplitude, consistent with the average weakening of ENSO variability in these experiments (see Section 5 and Figure 9). There is a much larger weakening of SST variability in the *lig127k* than the *midHolocene* case. The *lgm* SST pattern (Figure 10b) shows negative anomalies in the central to western Pacific, indicating either an eastward shift of the ENSO pattern and/or weaker central Pacific variability. On the other hand, both the *lpctCO2* and *abrupt4xCO2* composites (Figure 10d, e) show positive Pacific SST anomalies associated with ENSO at the equator,

- and *abrupt xCO2* composites (Figure 10d, e) show positive Pacific SS1 anomalies associated with ENSO at the equator, with the largest values in the central Pacific, and high model agreement. This suggests an increased ENSO variability among the ensemble, particularly for the *abrupt4xCO2* simulations, despite the disagreement between model ENSO amplitude changes discussed in Section 5.
 - 18

The global temperature and precipitation teleconnections with ENSO for each experiment relative to piControl are shown in

- 615 Figures 11 and 12. As discussed above, both *lig127k* and *midHolocene* simulations show weaker ENSO SST variability relative to *piControl* (Figure 10a, c). The *lig127k* simulations have a much greater weakening of the ENSO SST and temperature patterns (Figure 10c and 11e, f) than any of the other simulations (although based on a small number of models), with cooler SSTs in the central Pacific. The *midHolocene* ENSO SST and temperature pattern (Figure 10a and 11a, b) is a weaker version of the *lig127k* response. The ENSO precipitation teleconnection in the *lig127k* simulations (Figure 12e, f)
- 620 consists of a weakening of the *piControl* ENSO precipitation pattern, with much drier conditions in the equatorial Pacific and over the SPCZ during El Niño events. The *midHolocene* ENSO precipitation pattern (Figure 12a, b) is again a weaker version of the *lig127k* ENSO precipitation response.

The *lgm* simulations show cooler SSTs in the western tropical Pacific in the ENSO composite (Figure 10b), consistent with a weakening of ENSO variability in this region. Cool anomalies over Australia and warm anomalies over North America are

625 also evident (Figure 11c, d). As expected, given the colder global mean temperatures (Figure 6), precipitation associated with ENSO is reduced in the tropical Pacific and the overall hydrological cycle is weaker (Figure 12 c, d). Remote responses include wetter conditions over Australia and drier conditions over North America, somewhat resembling a La Niña pattern, although with low model agreement.

In contrast to the lgm experiments, the lpctCO2 and abrupt4xCO2 simulations are much warmer than the other experiments

- 630 (Figure 6). The largest temperature anomalies are seen for the *abrupt4xCO2* simulations (Figure 11i, j), which also had increased amplitude of SST variability in the central Pacific (Figure 10e). The *abrupt4xCO2* experiment shows warmer temperatures globally during El Niño events, particularly over continents and high northern latitudes (except Greenland in DJF). It is possible that elements of the pattern in this experiment arise from the linear detrending failing to sufficiently remove the transient changes in mean state (Section 2.3). The *1pctCO2* simulations (Figure 11g, h) show a similar but much
- 635 weaker response. The precipitation response to ENSO (Figure 12) is enhanced in the *abrupt4xCO2* and *lpctCO2* simulations relative to *piControl*, with increases in precipitation in the equatorial Pacific and decreases in the subtropics, as ENSO influences tropical atmospheric circulation and therefore the hydrological cycle (Lu et al., 2008; Nguyen et al., 2013). This is consistent with previous studies showing intensified ENSO temperature and precipitation impacts in a warmer climate (e.g. Power et al., 2013; Power and Delage, 2018).
- 640 The relationship between changes in ENSO and changes in the annual cycle, zonal and meridional SST gradient are investigated for all models and experiments (Figure 13). The change in ENSO amplitude based on standard deviation of NINO3.4 (relative to *piControl*) is plotted against the change in annual cycle of NINO3.4 SST in Figure 13a, with a weak positive correlation between the two variables. The change in ENSO amplitude was found to have no significant correlation with the change in equatorial Pacific zonal SST gradient (defined as 5° S–5° N, 150° E–170° W minus 5° S–5° N, 120° W–
- 645 90° W), shown in Figure 13b. Finally, the relationship between <u>castern Pacific rainfall and meridional SST gradient</u> (Cai et al., 2014) is investigated (Figure 13c). The meridional SST gradient in the eastern Pacific is defined as the average SST over the off-equatorial region (5° N–10° N, 150° W–90° W) minus the average over the equatorial region (2.5° S–2.5° N, 150°

(Deleted: underlying
~(Deleted: s
\mathcal{A}	Deleted: aspects of the mean state and
(Deleted: no significant
~(Deleted: also
(Deleted: extreme El Nino and the mean state
(Deleted: mean state is represented as changes in
~(Deleted: -

W–90° W). The strength of <u>eastern Pacific</u> El Niño rainfall is represented as the changes in ENSO composite precipitation over the NINO3 region (5° S–5° N, 150° W–90° W) normalised by the NINO3.4 standard deviation used to identify the composited events. This normalisation aims to remove the impact of the changes in ENSO variability documented between the experiments (e.g. Power and Delage, 2018). The <u>significant negative correlation</u> is consistent with the analysis demonstrated by Cai et al. (2014) and Collins et al. (2019), but this analysis approach allows the relationship to be visualised across many more simulations and experiments. This relationship appears to be fundamental feature of ENSO behaviour, rather than just a response to greenhouse gas forcing.

7 Mechanisms and Discussion

- There is evidence that the mid-Holocene, a period of supressed ENSO variability, featured a stronger zonal gradient in the tropical Pacific mean SST than the 20th century, namely "La Niña-like conditions" (Barr et al., 2019; Gagan and Thompson, 2004; Koutavas et al., 2002; Luan et al., 2012; Shin et al., 2006). In contrast, proxy records suggest that the mean state of the tropical Pacific during the Pliocene warm period featured sustained El Niño-like conditions (Fedorov et al., 2006; Wara et al., 2005). It is likely that the weak zonal SST gradient in the Pliocene was less favourable for ENSO occurrence (Brierley, 2013; Manucharyan and Fedorov, 2014). These contradictory responses imply that the dynamical mechanisms determining
- 670 the relationship between the zonal gradient in mean SST and ENSO amplitude (e.g., Sadekov et al., 2013) must consist of several processes. The relationship between ENSO amplitude and SST gradient may also be nonlinear (Hu et al., 2013). This lack of consistent or linear relationship between zonal SST gradient and ENSO amplitude is supported by the results presented here, shown in Figure 13b.
- During the mid-Holocene, the reduced tropical insolation led to the cooling of the tropical Pacific, directly producing a La Niña-like condition. Under La Niña-like conditions, the air-sea coupling strength is reduced due to a suppressed convective instability, and thus ENSO variability is suppressed (Liu et al., 2000; Roberts et al., 2014). The stronger seasonality in insolation over the Northern Hemisphere associated with the precession cycle resulted in a stronger annual cycle, which could also act to reduce ENSO variability through the intensified annual-frequency entrainment (Liu, 2002; Pan et al., 2005). A similar but stronger precession effect due to the higher eccentricity during the last interglacial period was also found to
- 680 have a relatively weak ENSO amplitude in palaeo-proxy records (Hughen et al., 1999; Tudhope et al., 2001) and climate model simulations (An et al., 2017; Salau et al., 2012). However, the mid-Holocene simulations of PMIP2/3 mostly showed a reduction of both annual cycle and ENSO amplitude (An and Choi, 2014; Masson-Delmotte et al., 2014).

The reduced annual cycle over the tropical eastern Pacific is attributed to the relaxation of eastern Pacific upper-ocean stratification due to the annual downwelling Kelvin wave forced by western Pacific wind anomalies (Karamperidou et al.,

685 2015) or the deepening of ocean mixed layer depth associated with the northward shift of the ITCZ (An and Choi, 2014). Therefore, the mid-Holocene ENSO variability in PMIP2/3 may be deemed to the result of the counterbalance between the

20

Deleted: n

(Deleted: strong negatively
(Deleted: relationship
~~(Deleted: (
\mathbb{X}	Deleted: ,
$\langle \rangle$	Deleted:
Ì	Deleted: to that

Deleted: The mean state of the tropical Pacific during the Pliocene warm period featured sustained El Niño-like conditions (Fedorov et al., 2006; Wara et al., 2005). Such weak zonal gradient in mean equatorial Pacific SST and the deep eastern Pacific thermoeline are linked to strongly suppressed trade winds. The suppressed trade winds, or westerly so-called 'super-rotation flow', can be driven by the equatorward westerly eddy momentum flux originating from the enhanced poleward Rossby wave pumping due to the stronger tropical convection under the warm ocean surface (Arnold et al., 2012; Tziperman and Farrell, 2009). Whilst individual El Niño events occurred during the Pliocene warm period (Ford and Ravelo, 2019), it is likely that the weak zonal contrast in mean SST (i.e., El Niño-like mean condition) was less favourable for ENSO occurrence (Brierley, 2013; Manucharyan and Fedorov, 2014). ...

Deleted:	another
Deleted:	suppression period
Deleted:	annual cycle
Deleted:	between the Pliocene and
Deleted:	mid-Holocene
Deleted:	

reduction due to the weaker air-sea coupling and the intensification due to the reduced frequency entrainment (An and Choi,

715 2014). Other factors may include coupling of the circulation in the eastern Pacific with the North American monsoon (implying dynamical damping of upwelling in the eastern Pacific), and an increased southeast Asian monsoon which strengthens winds in the western Pacific. Alternatively, An et al. (2010) and An and Choi (2013) argue that the change in annual cycle amplitude is not a cause of change in ENSO amplitude, it is the changes in the mean climate state that modify both the ENSO and annual cycle amplitudes in the opposite way. The analysis presented here would appear to support this regument there is no consistent relationship between changes in the amplitude of the annual cycle and changes in the ENSO

720 argument there is no convariability (Figure, 13a).

ENSO variability can be suppressed or enhanced by remote forcing. For example, the enhanced Asian summer monsoon also leads to La Niña-like conditions via increasing strength of the tropical Pacific trade winds and the resultant enhanced equatorial upwelling (Liu et al., 2000). Sensitivity experiments with fully coupled climate models demonstrate that greening

- 725 of the Sahara during the mid-Holocene could reduce ENSO variability through affecting the Atlantic Niño (Zebiak, 1993) and Walker Circulation, finally decreasing upwelling and deepening of the thermocline in the eastern Pacific (Pausata et al., 2017). The freshwater perturbation experiments, so-called "water hosing experiment" that lead to a weakening of the Atlantic Ocean meridional overturning circulation, showed a reduced seasonal cycle and enhanced ENSO variability through the inter-basin atmospheric teleconnection (Braconnot et al., 2012; Masson-Delmotte et al., 2014; Timmermann et al., 2007).
- 730 More sophisticated feedback analysis revealed that the reduction of ENSO variability is due to either the increase of the negative feedback by the mean current thermal advection (An and Bong, 2018) or the reduction of the major positive feedback processes (thermocline, zonal advection and Ekman feedbacks) (Chen et al., 2019a; Tian et al., 2017). The negative feedback due to the thermal advection by the mean current was intensified by the stronger cross-equatorial winds associated with the northward migration of the ITCZ (e.g., An and Choi, 2014), and the positive dynamical feedback was suppressed
- 735 due to the strengthening of the mean Pacific subtropical cell (Chen et al., 2019a). Therefore, the linear stability of ENSO during the mid-Holocene was reduced through the dedicated balance among the various feedback processes, but the change in each feedback process is model-dependent. External processes were also proposed as a suppression mechanism for mid-Holocene ENSO. For example, the Pacific meridional mode became weaker during the mid-Holocene, thereby ENSO has relatively less chance to be triggered (Chiang et al., 2009); the weaker ocean stratification due to the warm water subduction
- 740 from the subtropical ocean decreases ENSO stability (Liu et al., 2000).

How ENSO activity will change in response to anthropogenic global warming still remains uncertain (Cai et al., 2015b; Christensen et al., 2014; Collins et al., 2010). Observations show that ENSO variability has increased under greenhouse warming in the recent past (Zhang et al., 2008), which is also shown in CMIP5 climate model simulations (Cai et al., 2018). During the transient period of global warming, the tropical SSTs warm much faster than the subsurface ocean and leads to a

745 shallower and stronger thermocline in the equatorial Pacific (An et al., 2008), which enhances the ocean-atmosphere coupling and amplifies the ENSO variability (Zhang et al., 2008). A gradually intensified ENSO from mid- to late Holocene

21

Deleted:

also appears in a long-transient simulation since the last 21,000 years (Liu et al., 2014) and of the last 6000 years (Braconnot et al., 2019). For the equilibrium response to global warming, the subsurface ocean will eventually warm up and reduce the

- 750 vertical temperature gradient and weaken the ENSO variability. For instance, during the Pliocene warm period, the most recent period in the past with carbon dioxide concentrations similar or higher than today, SST reconstructions from the tropical Pacific show a reduced zonal SST gradient during this period, implying sustained El Niño-like conditions (Dekens et al., 2007; Fedorov et al., 2006; Wara et al., 2005) as well as a deeper thermocline with weaker thermocline feedback (White and Ravelo, 2020).
- 755 The tropical climate mean state response to current global warming is still a topic of debate as to whether it will be El Niñolike or La Niña-like (Cane et al., 1997; Collins, 2005; Merryfield 2006; An et al., 2012; Collins et al., 2010; Lian et al., 2018). Moreover, the global warming-induced tropical Pacific SST pattern seems to be less effective in changing ENSO activity (An and Choi, 2015). The strong internal modulation of ENSO activity over decadal-to-centennial timescales also obscures the actual global warming impact on ENSO variability (e.g. Wittenberg, 2009). So far, the future ENSO activity 760 reflected in SST anomalies obtained from CMIP5 models is not distinguishable from the historical ENSO activity
- (Christensen et al., 2014).

The enhancement and increasingly frequent occurrence of ENSO-driven extreme atmospheric responses to future global warming are strongly supported by model studies. This includes extreme rainfall events and extreme equatorward swings of the SPCZ (Cai et al., 2014; Cai et al., 2015a; Cai et al., 2012) and extreme weather events through teleconnections (Cai et al.,

- 765 2015b; Yeh et al., 2018). The changes in these extremes are due to the nonlinearity of atmospheric response to ENSO SSTs, especially with a warmer ocean surface. The changing amplitude of the extremes with changing meridional SST gradient is a feature of past climates as well as future climates. This is demonstrated by Figure, 13c, which shows an increase in the ENSO composited precipitation with increased SST gradient. However, most current GCMs still inaccurately simulate many aspects of the historical ENSO such as the far westward extent of the Pacific cold tongue (Taschetto et al., 2014), ENSO-
- related precipitation anomalies (Dai and Arkin, 2017), ENSO feedbacks (Bayr et al., 2019; Bellenger et al., 2014; Kim et al., 2014; Kim and Jin, 2011; Lloyd et al., 2012), and ENSO asymmetry in amplitude, duration, and transition (e.g. Chen et al., 2017; Zhang and Sun, 2014). Therefore, the accuracy of the future projections of ENSO will be guaranteed only when most coupled GCMs can simulate the observed modern-day ENSO more skilfully than at present.

8 Summary and Conclusions

775 We have presented a summary of ENSO amplitude and teleconnections changes in the most recent previous generation (CMIP5) and the new generation (CMIP6) of coupled climate models for past and future climates. The analysed simulations include the last glacial maximum climate (*lgm*), past interglacial climates (*lig127k* and *midHolocene*), and future idealised projections (*abrupt4xCO2* and *lpctCO2*), using the pre-industrial climate (*piControl*) as the reference state.

22

Deleted:)

Deleted:

We first evaluated a 30-year climatology from *historical* simulation against HadISST observation from 1971-2000. Similarly to the previous generations of climate models, the CMIP5 and CMIP6 models have cold biases in SST in the central to western equatorial Pacific, as well as in the subtropical Pacific. Warm SST biases are present in the eastern equatorial

Pacific. The *piControl* climate is about 1°C colder on average than the *historical* climate, leading to an apparent exacerbation of the cold-tongue bias and reduction in the warm bias in the eastern equatorial Pacific. These biases in SST lead to a strong and northward shift of ITCZ, a dry bias along the equatorial Pacific, and appearance of a "double ITCZ" as in previous CMIP-class simulations. <u>SST and precipitation biases are generally slightly smaller in the new generation of CMIP6 models than in the CMIP5 models.</u> The simulated ENSO pattern well resembles the observations with a slight displacement to the west, similarly the ENSO temperature and precipitation teleconnections are well simulated compared with observations.

The mean state changes were examined relative to the *piControl*. In the *lig127k* simulations, strong seasonal insolation anomalies lead to tropical and subtropical SST cooling of 0.5-2°C in DJF and JJA. No large-scale warming is found in the *midHolocene* simulations, but a slight cooling of 0.5°C occurs in the eastern Pacific in DJF and in the central Pacific in JJA.

- In the *lgm* simulations, 2-3°C cooling is found in the tropical Pacific. For the future scenarios, the tropical Pacific warms by 1-3°C in the *lpctCO2* case and more than 3°C in the *abrupt4xCO2* case. During *lig127k* and *midHolocene* simulations, the ITCZ shifts northward, leads to a weakened Australian summer monsoon and enhanced North American summer monsoon. In the *lgm* simulations, the ITCZ is intensified over the tropical Pacific, with drier conditions over the Maritime Continent, Australia and Southeast Asia, while wetter conditions are found in the western Pacific and on the northern edge of the SPCZ. In the *lpctCO2* and *abrupt4xCO2* simulations, precipitation increases in the equatorial Pacific following the largest SST
- 800 warming, with some drying on the northern edge of the ITCZ and in southeast Pacific in DJF.

The majority of models show a decrease in ENSO variability in the *lig127k* and *midHolocene* simulations. The reduction of ENSO variability in *lig127k* ranges to more than 40%, while only one model shows more than a ~20% decrease in ENSO variability in *midHolocene*. This is consistent with previous model studies of mid-Holocene ENSO, which generally show a smaller reduction in amplitude than implied by proxy records.

805 The changes in ENSO variability in *lgm*, *lpctCO2* and *abrupt4xCO2* simulations are highly model dependent, with the sign of change approximately equally divided between increases and decreases of up to 40% to 50% across the set of models. This is also consistent with previous studies showing little agreement on ENSO amplitude change under LGM conditions (Masson-Delmotte et al., 2014) or in future projections (Collins et al., 2014).

The ensemble mean weakening of ENSO in the *lgm* simulations is characterised by a cooling in the central and western Pacific. Cooling over Australia and warming over North America are also evident. The changes in the ENSO temperature teleconnections show cooling in the central Pacific in *lig127k* and *midHolocene*, and significant global warming in the *abrupt4xCO2* simulation, with strong warming amplification over continents and high northern latitudes. The *lpctCO2* simulations show a similar warming pattern but with much weaker magnitude. Precipitation teleconnections follow the



change in ensemble mean ENSO amplitude, with a weakening of the climatological *piControl* ENSO precipitation response in *lgm*, *lig127k* and *midHolocene* cases. In the *lgm* simulations, a weaker hydrological cycle due to cooler temperatures may also play a role. The precipitation response to ENSO is significant in the *abrupt4xCO2* and *lpctCO2* simulations, with an amplification of the climatological ENSO teleconnection pattern. This is consistent with previous studies showing increased ENSO precipitation responses in a warmer climate.

This study has provided an overview of changes in the mean state and ENSO in a set of past and future climate simulations from PMIP3/CMIP5 and PMIP4/CMIP6 models. We have not provided a comprehensive analysis of any aspect of these changes, and the set of CMIP6 models included is necessarily incomplete. Future work will focus on deepening understanding of the complex mechanisms driving interactions between changes in the mean state and ENSO. A more detailed comparison with palaeo-ENSO proxy records will also be required to evaluate model simulations of ENSO in past climates (e.g. Lu et al., 2018). Improved understanding of changes in ENSO in past climates can contribute to model

825 evaluation, understanding of ENSO dynamics and constraining projections of future change.

Code and data availability

The CMIP5 and CMIP6 model data used in this study are available from the Earth System Grid Federation. All the scripts used for the analysis and to produce the figures in this study are provided on the GitHub repository: 830 <u>https://github.com/chrisbrierley/PMIP4-enso</u>. All the ENSO diagnostics are also available via the PMIP variability database at past2future.org.

Author contributions

J.R.B. coordinated the research. C.M.B. performed the simulation analysis. J.R.B, C.M.B., S.-I.A., M.-V.G.,S.S., C.J.R.W., Q.Z. and A.Z. authored the manuscript with edits from R.D. Other authors contributed simulations towards the analysis.

835 Competing interests

The authors declare no competing interests.

Acknowledgements

J.R.B acknowledges support from the ARC Centre of Excellence for Climate Extremes (CE170100023). C.M.B., CJ.R.W.,
P.B., R.D., G.L. and X.S. acknowledge the JPI-Belmont-funded PACMEDY programme (NE/P006752/1; ANR-15-JCLI-0003-01; BMBF 01LP1607A). S.-I.A. was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2018R1A5A1024958). C.M.B. was also funded in part by NERC (NE/S009736/1).
M.V.G. acknowledges support from NERC research grant NE/P013279/1. Q.Z. acknowledges support from the Swedish

Research Council VR projects 2013-06476 and 2017-04232. Ru.O. acknowledges support from the Integrated Research Program for Advancing Climate Models (TOUGOU programme) from the Ministry of Education, Culture, Sports, Science

- 845 and Technology (MEXT), Japan. The simulations using MIROC-ES2L were conducted on the Earth Simulator of JAMSTEC. <u>B.L.O.-B.</u>, and E.C.B. acknowledge the CESM project, supported primarily by the National Science Foundation (NSF). The CESM2 simulations are based upon work supported by the National Center for Atmospheric Research (NCAR), which is a major facility sponsored by the NSF under Cooperative Agreement No. 1852977. Computing and data storage resources, including the Cheyenne supercomputer (doi:10.5065/D6RX99HX), were provided by the Computational and
- 850 Information Systems Laboratory (CISL) at NCAR. P.A.M. was supported by the state assignment project 0148-2019-0009. E.M.V. was supported by RSF grant 20-17-00190. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling groups for producing and making available their model output, the Earth System Grid Federation (ESGF) for archiving the data and providing access, and the multiple funding agencies who support CMIP6 and ESGF. PMIP is endorsed by both
- 855 WCRP/WGCM and Future Earth/PAGES. This research arose out of a workshop hosted at University College London by the PMIP working group on Past2Future: insights from a constantly varying past.

References

Abe-Ouchi, A., Saito, F., Kageyama, M., Braconnot, P., Harrison, S. P., Lambeck, K., Otto-Bliesner, B. L., Peltier, W. R., Tarasov, L., Peterschmitt, J. Y., and Takahashi, K.: Ice-sheet configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments, Geosci Model Dev, 8, 3621-3637, 2015.

Adam, O., Schneider, T., and Brient, F.: Regional and seasonal variations of the double-ITCZ bias in CMIP5 models, Clim Dynam, 51, 101-117, 2018.

Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P. P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D.,
 Gruber, A., Susskind, J., Arkin, P., and Nelkin, E.: The version-2 global precipitation climatology project (GPCP) monthly
 precipitation analysis (1979-present), Journal of Hydrometeorology, 4, 1147-1167, 2003.

An, S.-I. and Choi, J.: Inverse relationship between the equatorial eastern Pacific annual-cycle and ENSO amplitudes in a coupled general circulation model, Clim Dynam, 40, 663-675, 2013.

An, S.-I., Ham, Y.-G., Kug, J.-S., Timmermann, A., Choi, J., and Kang, I.-S.: The Inverse Effect of Annual-Mean State and Annual-Cycle Changes on ENSO, J Climate, 23, 1095-1110, 2010.

An, S.-I., Kim, H.-J., Park, W., and Schneider, B.: Impact of ENSO on East Asian winter monsoon during interglacial periods: effect of orbital forcing, Clim Dynam, 49, 3209-3219, 2017.

An, S. I. and Bong, H.: Feedback process responsible for the suppression of ENSO activity during the mid-Holocene, Theor Appl Climatol, 132, 779-790, 2018.

875 An, S. I. and Choi, J.: Mid-Holocene tropical Pacific climate state, annual cycle, and ENSO in PMIP2 and PMIP3, Clim Dynam, 43, 957-970, 2014.

An, S. I. and Choi, J.: Why the twenty-first century tropical Pacific trend pattern cannot significantly influence ENSO amplitude?, Clim Dynam, 44, 133-146, 2015.

An, S. I., Im, S. H., and Jun, S. Y.: Changes in ENSO Activity During the Last 6,000 Years Modulated by Background Climate State, Geophys Res Lett, 45, 2467-2475, 2018.

An, S. I. and Jin, F. F.: Nonlinearity and asymmetry of ENSO, J Climate, 17, 2399-2412, 2004.

An, S. I. and Kim, J. W.: ENSO Transition Asymmetry: Internal and External Causes and Intermodel Diversity, Geophys Res Lett, 45, 5095-5104, 2018.

An, S. I. and Kim, J. W.: Role of nonlinear ocean dynamic response to wind on the asymmetrical transition of El Nino and La Nina, Geophys Res Lett, 44, 393-400, 2017.

An, S. I., Kim, J. W., Im, S. H., Kim, B. M., and Park, J. H.: Recent and future sea surface temperature trends in tropical pacific warm pool and cold tongue regions, Clim Dynam, 39, 1373-1383, 2012.

An, S. I., Kug, J. S., Ham, Y. G., and Kang, I. S.: Successive modulation of ENSO to the future greenhouse warming, J Climate, 21, 3-21, 2008.

890 Andrews, T., Gregory, J. M., Webb, M. J., and Taylor, K. E.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere-ocean climate models, Geophys Res Lett, 39, 2012.

Ashok, K., Behera, S. K., Rao, S. A., Weng, H. Y., and Yamagata, T.: El Nino Modoki and its possible teleconnection, J Geophys Res-Oceans, 112, 2007.

Ballantyne, A., Lavine, M., Crowley, T., Liu, J., and Baker, P.: Meta-analysis of tropical surface temperatures during the Last Glacial Maximum, Geophys Res Lett, 32, 2005.

Barr, C., Tibby, J., Leng, M. J., Tyler, J. J., Henderson, A. C. G., Overpeck, J. T., Simpson, G. L., Cole, J. E., Phipps, S. J., Marshall, J. C., McGregor, G. B., Hua, Q., and McRobie, F. H.: Holocene El Niño–Southern Oscillation variability reflected in subtropical Australian precipitation, Scientific Reports, 9, 1627, 2019.

Bartlein, P. J. and Shafer, S. L.: Paleo calendar-effect adjustments in time-slice and transient climate-model simulations 900 (PaleoCalAdjust v1.0): impact and strategies for data analysis, Geosci Model Dev, 12, 3889-3913, 2019.

Bayr, T., Wengel, C., Latif, M., Dommenget, D., Lubbecke, J., and Park, W.: Error compensation of ENSO atmospheric feedbacks in climate models and its influence on simulated ENSO dynamics, Clim Dynam, 53, 155-172, 2019.

Bayr, T., Latif, M., Dommenget, D., Wengel, C., Harlass, J. and Park, W.: Mean-state dependence of ENSO atmospheric feedbacks in climate models, Clim Dynam, 50, 3171-3194, 2018.

905 Bayr, T., Dommenget, D., Martin, T., and Power, S.B.: The eastward shift of the Walker Circulation in response to global warming and its relationship to ENSO variability, Clim Dynam, 43, 2747-2763, 2014.

Bayr, T., and Dommenget, D.: The tropospheric land-sea warming contrast as the driver of tropical sea level pressure changes, J Climate, 26, 1387-1402, 2013.

Bellenger, H., Guilyardi, E., Leloup, J., Lengaigne, M., and Vialard, J.: ENSO representation in climate models: from 910 CMIP3 to CMIP5, Clim Dynam, 42, 1999-2018, 2014.

Berger, A. and Loutre, M. F.: Insolation Values for the Climate of the Last 10000000 Years, Quaternary Sci Rev, 10, 297-317, 1991.

Braconnot, P., Crétat, J., Marti, O., Balkanski, Y., Caubel, A., Cozic, A., Foujols, M.-A., and Sanogo, S.: Impact of Multiscale Variability on Last 6,000 Years Indian and West African Monsoon Rain, Geophys Res Lett, n/a, 2019.

915 Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-Ouchi, A., Otto-Bliesner, B., and Zhao, Y.: Evaluation of climate models using palaeoclimatic data, Nat Clim Change, 2, 417-424, 2012.

Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J. Y., Abe-Ouchi, A., Crucifix, M., Driesschaert, E., Fichefet, T., Hewitt, C. D., Kageyama, M., Kitoh, A., Loutre, M. F., Marti, O., Merkel, U., Ramstein, G., Valdes, P., Weber, L., Yu, Y., and Zhao, Y.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum -Part 2: feedbacks with emphasis on the location of the ITCZ and mid- and high latitudes heat budget, Clim Past, 3, 279-296,

Brierley, C.: ENSO behavior before the Pleistocene, PAGES news, 21, 70-71, 2013.

26

Deleted: Arnold, N. P., Tziperman, E., and Farrell, B.: Abrupt transition to strong superrotation driven by equatorial wave resonance in an idealized GCM, J Atmos Sci, 69, 626-640, 2012.¶

Deleted: Bacon, C. D., Silvestro, D., Jaramillo, C., Smith, B. T., Chakrabarty, P., and Antonelli, A.: Biological evidence supports an early and complex emergence of the Isthmus of Panama, P Natl Acad Sci USA, 112, 6110-6115, 2015. 930 Brierley, C. and Wainer, I.: Inter-annual variability in the tropical Atlantic from the Last Glacial Maximum into future climate projections simulated by CMIP5/PMIP3, Clim Past, 14, 1377-1390, 2018.

Brierley, C. M., Fedorov, A. V., Liu, Z. H., Herbert, T. D., Lawrence, K. T., and LaRiviere, J. P.: Greatly Expanded Tropical Warm Pool and Weakened Hadley Circulation in the Early Pliocene, Science, 323, 1714-1718, 2009.

Brown, J. R., Hope, P., Gergis, J., and Henley, B. J.: ENSO teleconnections with Australian rainfall in coupled model simulations of the last millennium, Clim Dynam, 47, 79-93, 2016.

Brown, J. R., Moise, A. F., and Colman, R. A.: The South Pacific Convergence Zone in CMIP5 simulations of historical and future climate, Clim Dynam, 41, 2179-2197, 2013.

Burls, N. J. and Fedorov, A. V.: Simulating Pliocene warmth and a permanent El Nino-like state: The role of cloud albedo, Paleoceanography, 29, 893-910, 2014.

940 Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso, A., McPhaden, M. J., Wu, L., England, M. H., Wang, G., Guilyardi, E., and Jin, F.-F.: Increasing frequency of extreme El Niño events due to greenhouse warming, Nat Clim Change, 4, 111, 2014.

Cai, W., Santoso, A., Wang, G., Yeh, S.-W., An, S.-I., Cobb, K. M., Collins, M., Guilyardi, E., Jin, F.-F., Kug, J.-S., Lengaigne, M., McPhaden, M. J., Takahashi, K., Timmermann, A., Vecchi, G., Watanabe, M., and Wu, L.: ENSO and greenhouse warming, Nat Clim Change, 5, 849, 2015a.

Cai, W. J., Lengaigne, M., Borlace, S., Collins, M., Cowan, T., McPhaden, M. J., Timmermann, A., Power, S., Brown, J., Menkes, C., Ngari, A., Vincent, E. M., and Widlansky, M. J.: More extreme swings of the South Pacific convergence zone due to greenhouse warming, Nature, 488, 365-369, 2012.

Cai, W. J., Santoso, A., Wang, G. J., Yeh, S. W., An, S. I., Cobb, K. M., Collins, M., Guilyardi, E., Jin, F. F., Kug, J. S.,
Lengaigne, M., McPhaden, M. J., Takahashi, K., Timmermann, A., Vecchi, G., Watanabe, M., and Wu, L. X.: ENSO and
greenhouse warming, Nat Clim Change, 5, 849-859, 2015b.

Cai, W. J., Wang, G. J., Dewitte, B., Wu, L. X., Santoso, A., Takahashi, K., Yang, Y., Carreric, A., and McPhaden, M. J.: Increased variability of eastern Pacific El Nino under greenhouse warming, Nature, 564, 201-206, 2018.

Deleted: +

Cai, W. J., Wang, G. J., Santoso, A., McPhaden, M. J., Wu, L. X., Jin, F. F., Timmermann, A., Collins, M., Vecchi, G.,
 Lengaigne, M., England, M. H., Dommenget, D., Takahashi, K., and Guilyardi, E.: Increased frequency of extreme La Nina
 events under greenhouse warming, Nat Clim Change, 5, 132-137, 2015c.

Cane, M. A., Clement, A. C., Kaplan, A., Kushnir, Y., Pozdnyakov, D., Seager, R., Zebiak, S. E., and Murtugudde, R.: Twentieth-century sea surface temperature trends, Science, 275, 957-960, 1997.

Capotondi, A., Wittenberg, A. T., Newman, M., Di Lorenzo, E., Yu, J. Y., Braconnot, P., Cole, J., Dewitte, B., Giese, B.,
Guilyardi, E., Jin, F. F., Karnauskas, K., Kirtman, B., Lee, T., Schneider, N., Xue, Y., and Yeh, S. W.: Understanding ENSO Diversity, B Am Meteorol Soc, 96, 921-938, 2015.

Carré, M., Sachs, J. P., Purea, S., Schauer, A. J., Braconnot, P., Falcom, R. A., Julien, M. and Lavallee, D.: Holocene history of ENSO variance and asymmetry in the eastern tropical Pacific, Science, 345, 1045-1048, 2014.

Ceppi, P., Brient, F., Zelinka, M. D., and Hartmann, D. L.: Cloud feedback mechanisms and their representation in global climate models, Wires Clim Change, 8, 2017.

Chadwick, R., Boutle, I., and Martin, G.: Spatial patterns of precipitation change in CMIP5: Why the rich do not get richer in the tropics, J Climate, 26, 3803-3822, 2013.

Chaisson, W. P. and Ravelo, A. C.: Pliocene development of the east-west hydrographic gradient in the equatorial Pacific, Paleoceanography, 15, 497-505, 2000.

970 Chen, L., Li, T., Yu, Y. Q., and Behera, S. K.: A possible explanation for the divergent projection of ENSO amplitude change under global warming, Clim Dynam, 49, 3799-3811, 2017.

Chen, L., Wang, L., Li, T., and Liu, J.: Drivers of reduced ENSO variability in mid-Holocene in a coupled model, Clim Dynam, 52, 5999-6014, 2019a.

975 Chen, L., Zheng, W. P., and Braconnot, P.: Towards understanding the suppressed ENSO activity during mid-Holocene in PMIP2 and PMIP3 simulations, Clim Dynam, 53, 1095-1110, 2019b.

Chiang, J. C. H., Fang, Y., and Chang, P.: Pacific Climate Change and ENSO Activity in the Mid-Holocene, J Climate, 22, 923-939, 2009.

- Christensen, J. H., Kanikicharla, K. K., Aldrian, E., An, S. I., Cavalcanti, I. F. A., de Castro, M., Dong, W. J., Goswami, P.,
 P80 Hall, A., Kanyanga, J. K., Kitoh, A., Kossin, J., Lau, N. C., Renwick, J., Stephenson, D. B., Xie, S. P., Zhou, T. J., Abraham, L., Ambrizzi, T., Anderson, B., Arakawa, O., Arritt, R., Baldwin, M., Barlow, M., Barriopedro, D., Biasutti, M., Biner, S.,
 Bromwich, D., Brown, J., Cai, W. J., Carvalho, L. V., Chang, P., Chen, X. L., Choi, J., Christensen, O. B., Deser, C.,
 Emanuel, K., Endo, H., Enfield, D. B., Evan, A., Giannini, A., Gillett, N., Hariharasubramanian, A., Huang, P., Jones, J.,
 Karumuri, A., Katzlev, J., Kiellstrom, E., Knight, J., Knutson, T., Kulkarni, A., Kundeti, K. R., Lau, W. K., Lenderink, G.,
- 985 Lennard, C., Leung, L. Y. R., Lin, R. P., Losada, T., Mackellar, N. C., Magana, V., Marshall, G., Mearns, L., Meehl, G., Menendez, C., Murakami, H., Nath, M. J., Neelin, J. D., van Oldenborgh, G. J., Olesen, M., Polcher, J., Qian, Y., Ray, S., Reich, K. D., de Fonseca, B. R., Ruti, P., Screen, J., Sedlacek, J., Solman, S., Stendel, M., Stevenson, S., Takayabu, I., Turner, J., Ummenhofer, C., Walsh, K., Wang, B., Wang, C. Z., Watterson, I., Widlansky, M., Wittenberg, A., Woollings, T., Yeh, S. W., Zhang, C. D., Zhang, L. X., Zheng, X. T., and Zou, L. W.: Climate Phenomena and their Relevance for Future 990 Regional Climate Change, Climate Change 2013: The Physical Science Basis, 2014. 1217-1308, 2014.

Clement, A. C., Seager, R., and Cane, M. A.: Suppression of El Nino during the mid-Holocene by changes in the Earth's orbit, Paleoceanography, 15, 731-737, 2000.

Cobb, K. M., Charles, C. D., Cheng, H., and Edwards, R. L.: El Nino/Southern Oscillation and tropical Pacific climate during the last millennium, Nature, 424, 271-276, 2003.

- 995 Cobb, K. M., Westphal, N., Sayani, H. R., Watson, J. T., Di Lorenzo, E., Cheng, H., Edwards, R. L., and Charles, C. D.: Highly Variable El Nino-Southern Oscillation Throughout the Holocene, Science, 339, 67-70, 2013. Collins, M.: El Nino- or La Nina-like climate change?, Clim Dynam, 24, 89-104, 2005. Collins, M.: Understanding uncertainties in the response of ENSO to greenhouse warming, Geophys Res Lett, 27, 3509-3512, 2000.
- 1000 Collins, M., An, S. I., Cai, W. J., Ganachaud, A., Guilyardi, E., Jin, F. F., Jochum, M., Lengaigne, M., Power, S., Timmermann, A., Vecchi, G., and Wittenberg, A.: The impact of global warming on the tropical Pacific ocean and El Nino, Nat Geosci, 3, 391-397, 2010.

Collins, M., Knutti, R., Arblaster, J., Dufresne, J. L., Fichefet, T., Friedlingstein, P., Gao, X. J., Gutowski, W. J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A. J., Wehner, M., Allen, M. R., Andrews, T., Beyerle, U., Bitz, C. M.,

- 1005 Bony, S., Booth, B. B., Brooks, H. E., Brovkin, V., Browne, O., Brutel-Vuilmet, C., Cane, M., Chadwick, R., Cook, E., Cook, K. H., Eby, M., Fasullo, J., Fischer, E. M., Forest, C. E., Forster, P., Good, P., Goosse, H., Gregory, J. M., Hegerl, G. C., Hezel, P. J., Hodges, K. I., Holland, M. M., Huber, M., Huybrechts, P., Joshi, M., Kharin, V., Kushnir, Y., Lawrence, D. M., Lee, R. W., Liddicoat, S., Lucas, C., Lucht, W., Marotzke, J., Massonnet, F., Matthews, H. D., Meinshausen, M., Morice, C., Otto, A., Patricola, C. M., Philippon-Berthier, G., Prabhat, Rahmstorf, S., Riley, W. J., Rogelj, J., Saenko, O., Seager, R.,
- 1010 Sedlacek, J., Shaffrey, L. C., Shindell, D., Sillmann, J., Slater, A., Stevens, B., Stott, P. A., Webb, R., Zappa, G., and Zickfeld, K.: Long-term Climate Change: Projections, Commitments and Irreversibility, Climate Change 2013: The Physical Science Basis, 2014. 1029-1136, 2014.

Collins, M., Sutherland, M., Bouwer, L., Cheong, S.-M., Frölicher, T., Jacot Des Combes, H., Koll Roxy, M., Losada, I.,
 McInnes, K., Ratter, B., Rivera-Arriaga, E., Susanto, R. D., Swingedouw, D., and Tibig, L.: Extremes, Abrupt Changes and
 Managing Risk, IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, 2019. 2019.

- Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Bronnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., Maugeri, M., Mok, H. Y., Nordli, O., Ross, T. F., Trigo, R. M., Wang, X. L., Woodruff, S. D., and Worley, S. J.: The Twentieth Century Reanalysis Project, Q J Roy Meteor Soc, 137, 1-28, 2011.
- 1020 Conroy, J. L., Overpeck, J. T., Cole, J. E., Shanahan, T. M., and Steinitz-Kannan, M.: Holocene changes in eastern tropical Pacific climate inferred from a Galápagos lake sediment record, Quaternary Sci Rev, 27, 1166-1180, 2008.

Dai, N. and Arkin, P. A.: Twentieth century ENSO-related precipitation mean states in twentieth century reanalysis, reconstructed precipitation and CMIP5 models, Clim Dynam, 48, 3061-3083, 2017.

Dekens, P. S., Ravelo, A. C., and McCarthy, M. D.: Warm upwelling regions in the Pliocene warm period, Paleoceanography, 22, 2007.

Deser, C., Alexander, M. A., Xie, S. P., and Phillips, A. S.: Sea Surface Temperature Variability: Patterns and Mechanisms, Annu Rev Mar Sci, 2, 115-143, 2010.

 Di Nezio, P. N., Timmermann, A., Tierney, J. E., Jin, F. F., Otto-Bliesner, B., Rosenbloom, N., Mapes, B., Neale, R., Ivanovic, R. F., and Montenegro, A.: The climate response of the Indo-Pacific warm pool to glacial sea level, Paleoceanography, 31, 866-894, 2016.

DiNezio, P. N., Clement, A. C., Vecchi, G. A., Soden, B. J., and Kirtman, B. P.: Climate Response of the Equatorial Pacific to Global Warming, J Climate, 22, 4873-4892, 2009.

Donders, T. H., Wagner, F., Dilcher, D. L., and Visscher, H.: Mid- to late-Holocene El Nino-Southern Oscillation dynamics reflected in the subtropical terrestrial realm, P Natl Acad Sci USA, 102, 10904-10908, 2005.

1035 Durack, P. J., Wijffels, S. E., and Matear, R. J.: Ocean Salinities Reveal Strong Global Water Cycle Intensification During 1950 to 2000, Science, 336, 455-458, 2012.

Eisenman, I., Yu, L. S., and Tziperman, E.: Westerly wind bursts: ENSO's tail rather than the dog?, J Climate, 18, 5224-5238, 2005.

Emile-Geay, J., Cobb, K. M., Carre, M., Braconnot, P., Leloup, J., Zhou, Y., Harrison, S. P., Correge, T., McGregor, H. V.,
 Collins, M., Driscoll, R., Elliot, M., Schneider, B., and Tudhope, A.: Links between tropical Pacific seasonal, interannual and orbital variability during the Holocene, Nat Geosci, 9, 168-173, 2016.

Emile-Geay, J., Cobb, K. M., Mann, M. E., and Wittenberg, A. T.: Estimating Central Equatorial Pacific SST Variability over the Past Millennium. Part II: Reconstructions and Implications, J Climate, 26, 2329-2352, 2013.

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled
 Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, Geosci. Model Dev., 9, 1937-1958,
 2016a.

Eyring, V., Righi, M., Lauer, A., Evaldsson, M., Wenzel, S., Jones, C., Anav, A., Andrews, O., Cionni, I., Davin, E. L., Deser, C., Ehbrecht, C., Friedlingstein, P., Gleckler, P., Gottschaldt, K. D., Hagemann, S., Juckes, M., Kindermann, S., Krasting, J., Kunert, D., Levine, R., Loew, A., Mäkelä, J., Martin, G., Mason, E., Phillips, A. S., Read, S., Rio, C., Roehrig,

1050 R., Senftleben, D., Sterl, A., van Ulft, L. H., Walton, J., Wang, S., and Williams, K. D.: ESMValTool (v1.0) – a community diagnostic and performance metrics tool for routine evaluation of Earth system models in CMIP, Geosci. Model Dev., 9, 1747-1802, 2016b.

Fedorov, A. V., Brierley, C. M., Lawrence, K. T., Liu, Z., Dekens, P. S., and Ravelo, A. C.: Patterns and mechanisms of early Pliocene warmth, Nature, 496, 43-49, 2013.

1055 Fedorov, A. V., Dekens, P. S., McCarthy, M., Ravelo, A. C., deMenocal, P. B., Barreiro, M., Pacanowski, R. C., and Philander, S. G.: The Pliocene paradox (mechanisms for a permanent El Nino), Science, 312, 1485-1489, 2006.

Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., Rummukainen, M., AchutaRao, K., Anav, A., Andrews, T., Baehr, J., Bindoff, N. L., Bodas-Salcedo, A., Catto, J., Chambers, D., Chang, P., Dai, A. G., Deser, C., Doblas-

- 1060 Reyes, F., Durack, P. J., Eby, M., de Elia, R., Fichefet, T., Forster, P., Frame, D., Fyfe, J., Gbobaniyi, E., Gillett, N., Gonzalez-Rouco, J. F., Goodess, C., Griffies, S., Hall, A., Harrison, S., Hense, A., Hunke, E., Ilyina, T., Ivanova, D., Johnson, G., Kageyama, M., Kharin, V., Klein, S. A., Knight, J., Knutti, R., Landerer, F., Lee, T., Li, H. M., Mahowald, N., Mears, C., Meehl, G., Morice, C., Msadek, R., Myhre, G., Neelin, J. D., Painter, J., Pavlova, T., Perlwitz, J., Peterschmitt, J. Y., Raisanen, J., Rauser, F., Reid, J., Rodwell, M., Santer, B., Scaife, A. A., Schulz, J., Scinocca, J., Sexton, D., Shindell, D.,
- 1065 Shiogama, H., Sillmann, J., Simmons, A., Sperber, K., Stephenson, D., Stevens, B., Stott, P., Sutton, R., Thorne, P. W., van Oldenborgh, G. J., Vecchi, G., Webb, M., Williams, K., Woollings, T., Xie, S. P., and Zhang, J. L.: Evaluation of Climate Models, Climate Change 2013: The Physical Science Basis, 2014. 741-866, 2014.

29

Deleted: +

- Ford, H. L., McChesney, C. L., Hertzberg, J. E., and McManus, J. F.: A deep eastern equatorial Pacific thermocline during
 the Last Glacial Maximum, Geophys Res Lett, 45, 11,806-11,816, 2018.
- Ford, H. L., Ravelo, A. C. and Polissar, P. J.: Reduced El Nino-Southern Oscillation during the Last Glacial Maximum, Science, 347, 255-258, 2015.
- Freund, M. B., Henley, B. J., Karoly, D. J., McGregor, H. V., Abram, N. J., and Dommenget, D.: Higher frequency of Central Pacific El Nino events in recent decades relative to past centuries, Nat Geosci, 12, 450-455, 2019.
- 1075 Gagan, M. K. and Thompson, L. G.: Evolution of the Indo-Pacific warm pool and Hadley-Walker circulation since the last deglaciation. In: The Hadley circulation: present, past and future, Springer, 2004. Gergis, J. L. and Fowler, A. M.: A history of ENSO events since AD 1525: implications for future climate change, Climatic Change, 92, 343-387, 2009.
- Gregory, J. M., Ingram, W. J., Palmer, M. A., Jones, G. S., Stott, P. A., Thorpe, R. B., Lowe, J. A., Johns, T. C., and Williams, K. D.: A new method for diagnosing radiative forcing and climate sensitivity, Geophys Res Lett, 31, 2004.
- Grose, M.R., Narsey, S., Delage, F.P., Dowdy, A.J., Bador, M., Boschat, G., Chung, C., Kajtar, J.B., Rauniyar, S., Freund, M.B. and Lyu, K., Rashid, H., Zhang, X., Wales, S., Trenham, C., Holbrook, N.J., Cowan, T., Alexander, L., Arblaster, J.M. and Power, S.:Insights from CMIP6 for Australia's future climate, Earth's Future, 8, e2019EF001469, 2020.
- Grothe, P. R., Cobb, K. M., Liguori, G., Di Lorenzo, E., Capotondi, A., Lu, Y., Cheng, H., Edwards, R.L., Southon, J. R.,
 Santos, G. M., Deocampo, D. M., Lynch-Stieglitz, J., Chen, T., Sayani, H. R., Thompson, D. M., Conroy, J. L., Moore, A.L.,
 Townsend, K., Hagos, M., O'Connor, G., and Toth, L.T.: Enhanced El Niño–Southern oscillation variability in recent
 decades, Geophys Res Lett, 47, e2019GL083906, 2019.

Haywood, A. M., Valdes, P. J., and Peck, V. L.: A permanent El Nino-like state during the Pliocene?, Paleoceanography, 22, 2007.

- 1090 Held, I. M. and Soden, B. J.: Robust responses of the hydrological cycle to global warming, J Climate, 19, 5686-5699, 2006. Hoesly, R. M., Smith, S. J., Feng, L. Y., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Davidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z. F., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750-2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), Geosci Model Dev, 11, 369-408, 2018.
- 095 <u>Hu, Z., Kumar, A., Ren, H., Wang, H., L'Heureux, M., and Jin, F.: Weakened interannual variability in the tropical Pacific Ocean since 2000, J Climate, 26, 2601–2613, 2013.</u>

Hughen, K. A., Schrag, D. P., Jacobsen, S. B., and Hantoro, W.: El Nino during the last interglacial period recorded by a fossil coral from Indonesia, Geophys Res Lett, 26, 3129-3132, 1999.

Im, S. H., An, S. I., Kim, S. T., and Jin, F. F.: Feedback processes responsible for El Nino-La Nina amplitude asymmetry, 100 Geophys Res Lett, 42, 5556-5563, 2015.

Jansen, E., Fronval, T., Rack, F., and Channell, J. E. T.: Pliocene-Pleistocene ice rafting history and cyclicity in the Nordic Seas during the last 3.5 Myr, Paleoceanography, 15, 709-721, 2000.

Jones, T. R., Roberts, W. H. G., Steig, E. J., Cuffey, K. M., Markle, B. R., and White, J. W. C.: Southern Hemisphere climate variability forced by Northern Hemisphere ice-sheet topography, Nature, 554, 351-<u>355</u>, 2018.

105 Juckes, M., Taylor, K. E., Durack, P., Lawrence, B., Mizielinski, M., Pamment, A., Peterschmitt, J.-Y., Rixen, M., and Sénésis, S.: The CMIP6 Data Request (version 01.00. 31), Geosci. Model Dev. Discuss, 2019, 1-35, 2019.

Kageyama, M., Albani, S., Braconnot, P., Harrison, S. P., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Marti, O., Peltier, W. R., Peterschmitt, J. Y., Roche, D. M., Tarasov, L., Zhang, X., Brady, E. C., Haywood, A. M., LeGrande, A. N., Lunt, D. J., Mahowald, N. M., Mikolajewicz, U., Nisancioglu, K. H., Otto-Bliesner, B. L., Renssen, H., Tomas, R. A., Zhang, Q., Abe-

110 Ouchi, A., Bartlein, P. J., Cao, J., Li, Q., Lohmann, G., Ohgaito, R., Shi, X. X., Volodin, E., Yoshida, K., Zhang, X., and Zheng, W. P.: The PMIP4 contribution to CMIP6-Part 4: Scientific objectives and experimental design of the PMIP4-CMIP6 Last Glacial Maximum experiments and PMIP4 sensitivity experiments, Geosci Model Dev, 10, 4035-4055, 2017. Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J. H., Otto-Bliesner, B. L., Peterschmitt, J. Y.,

Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J. H., Otto-Bilesner, B. L., Peterschmitt, J. Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft,

30

Deleted: Ford, H. L. and Ravelo, A. C.: Estimates of Pliocene Tropical Pacific Temperature Sensitivity to Radiative Greenhouse Gas Forcing, Paleoceanogr Paleocl, 34, 2-15, 2019.Ford, H. L.,

Deleted: +

Deleted: +

120 P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q., and Zhou, T. J.: The PMIP4 contribution to CMIP6-Part 1: Overview and over-arching analysis plan, Geosci Model Dev, 11, 1033-1057, 2018.

Karamperidou, C., Di Nezio, P. N., Timmermann, A., Jin, F. F., and Cobb, K. M.: The response of ENSO flavors to mid-Holocene climate: Implications for proxy interpretation, Paleoceanography, 30, 527-547, 2015.

125 Khider, D., Stott, L. D., Emile-Geay, J., Thunell, R., and Hammond, D. E.: Assessing El Nino Southern Oscillation variability during the past millennium, Paleoceanography, 26, 2011.

Kim, S. T., Cai, W., Jin, F.-F., Yu, J.-Y.: ENSO stability in coupled climate models and its association with mean state, Clim Dynam, 42, 3313–3321, 2014.

Kim, S. T. and Jin, F.-F.: An ENSO stability analysis. Part II: results from the twentieth and twenty-first century simulations of the CMIP3 models, Clim Dynam, 36, 1609-1627, 2011.

Knutti, R., Masson, D., and Gettelman, A.: Climate model genealogy: Generation CMIP5 and how we got there, Geophys Res Lett, 40, 1194-1199, 2013.

Koutavas, A. and Joanides, S.: El Nino-Southern Oscillation extrema in the Holocene and Last Glacial Maximum, Paleoceanography, 27, PA4208, 2012.

135 Koutavas, A., Demenocal, P. B., Olive, G. C., and Lynch-Stieglitz, J.: Mid-Holocene El Nino-Southern Oscillation (ENSO) attenuation revealed by individual foraminifera in eastern tropical Pacific sediments, Geology, 34, 993-996, 2006.

Koutavas, A., Lynch-Stieglitz, J., Marchitto, T. M., and Sachs, J. P.: El Nino-like pattern in ice age tropical Pacific sea surface temperature, Science, 297, 226-230, 2002.

Kukla, G. J., Clement, A. C., Cane, M. A., Gavin, J. E., and Zebiak, S. E.: Last interglacial and early glacial ENSO, Quaternary Research, 58, 27-31, 2002.

Latif, M., and Keenlyside, N. S.: El Niño/Southern Oscillation response to global warming, Proceedings of the National Academy of Sciences, 106, 20578-20583, 2009.

Leduc, G., Vidal, L., Cartapanis, O., and Baird, E.: Modes of eastern equatorial Pacific thermocline variability: implications for ENSO dynamics over the last glacial period, Paleoceanography, 24, PA3202, 2009.

145 Levine, A., Jin, F. F., and McPhaden, M. J.: Extreme Noise-Extreme El Nino: How State-Dependent Noise Forcing Creates El Nino-La Nina Asymmetry, J Climate, 29, 5483-5499, 2016.

Lewis, S. C. and LeGrande, A. N.: Stability of ENSO and its tropical Pacific teleconnections over the Last Millennium, Clim Past, 11, 1347-1360, 2015.

Lian, T., Chen, D. K., Ying, J., Huang, P., and Tang, Y. M.: Tropical Pacific trends under global warming: El Nino-like or La Nina-like?, Natl Sci Rev, 5, 810-812, 2018.

Liu, Z.: A Simple Model Study of ENSO Suppression by External Periodic Forcing, J Climate, 15, 1088-1098, 2002.

Liu, Z., Kutzbach, J., and Wu, L.: Modeling climate shift of El Nino variability in the Holocene, Geophys Res Lett, 27, 2265-2268, 2000.

Liu, Z., Vavrus, S., He, F., Wen, N., and Zhong, Y.: Rethinking Tropical Ocean Response to Global Warming: The Enhanced Equatorial Warming, J Climate, 18, 4684-4700, 2005.

Liu, Z. Y., Lu, Z. Y., Wen, X. Y., Otto-Bliesner, B. L., Timmermann, A., and Cobb, K. M.: Evolution and forcing mechanisms of El Nino over the past 21,000 years, Nature, 515, 550-553, 2014.

Lloyd, J., Guilyardi, E., and Weller, H.: The role of atmosphere feedbacks during ENSO in the CMIP3 models. Part III: the shortwave flux feedback, J Climate, 25, 4275-4293, 2012.

1160 Lu, J., Chen, G., and Frierson, D. M. W.: Response of the Zonal Mean Atmospheric Circulation to El Nino versus Global Warming, J Climate, 21, 5835-5851, 2008.

Lu, Z. Y., Liu, Z. Y., Zhu, J., and Cobb, K. M.: A Review of Paleo El Nino-Southern Oscillation, Atmosphere-Basel, 9, 2018. Luan, Y., Braconnot, P., Yu, Y., Zheng, W., and Marti, O.: Early and mid-Holocene climate in the tropical Pacific: seasonal cycle and interannual variability induced by insolation changes, Clim Past, 8, 1093-1108, 2012.

- 1165 Manucharyan, G. E. and Fedorov, A. V.: Robust ENSO across a Wide Range of Climates, J Climate, 27, 5836-5850, 2014. MARGO Project Members, Waelbroeck, C., Paul, A., Kucera, M., Rosell-Melé, A., Weinelt, M., Schneider, R., Mix, A., Abelmann, A., and Armand, L.: Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum, Nat Geosci, 2, 127, 2009.
- Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolski, A., Rouco, J. F. G., Jansen, E., Lambeck, K.,
 Luterbacher, J., Naish, T., Osborn, T., Otto-Bliesner, B., Quinn, T., Ramesh, R., Rojas, M., Shao, X. M., Timmermann, A.,
 Anchukaitis, K., Arblaster, J., Bartlein, P. J., Benito, G., Clark, P., Comiso, J. C., Crowley, T., De Deckker, P., de Vernal, A.,
 Delmonte, B., DiNezio, P., Dokken, T., Dowsett, H. J., Edwards, R. L., Fischer, H., Fleitmann, D., Foster, G., Frohlich, C.,
 Govin, A., Hall, A., Hargreaves, J., Haywood, A., Hollis, C., Horton, B., Kageyama, M., Knutti, R., Kopp, R., Krinner, G.,
 Landais, A., Li, C., Lunt, D., Mahowald, N., McGregor, S., Meehl, G., Mitrovica, J. X., Moberg, A., Mudelsee, M., Muhs, D.
- 1175 R., Mulitza, S., Muller, S., Overland, J., Parrenin, F., Pearson, P., Robock, A., Rohling, E., Salzmann, U., Savarino, J., Sedlacek, J., Shakun, J., Shindell, D., Smerdon, J., Solomina, O., Tarasov, P., Vinther, B., Waelbroeck, C., Wolf-Gladrow, D., Yokoyama, Y., Yoshimori, M., Zachos, J., and Zwartz, D.: Information from Paleoclimate Archives, Climate Change 2013: The Physical Science Basis, 2014. 383-464, 2014.

McGregor, H. V. and Gagan, M. K.: Western Pacific coral delta O-18 records of anomalous Holocene variability in the El 180 Nino-Southern Oscillation, Geophys Res Lett, 31, 2004.

Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O'Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, R.: Historical greenhouse gas concentrations for climate modelling (CMIP6), Geosci Model Dev, 10, 2057-2116, 2017.

185 Merryfield, W.J.: Changes to ENSO under CO₂ doubling in a multimodel ensemble, J Climate, 19, 4009-4027, 2006. Neelin, J. D., Jin, F. F., and Syu, H. H.: Variations in ENSO phase locking, J Climate, 13, 2570-2590, 2000.

Teenin, 5. D., 5in, 1. 1., and 5yu, 11. 11. Variations in Excos phase locking, 5 Chinate, 15, 2576-2576, 2000.

Nguyen, H., Evans, A., Lucas, C., Smith, I., and Timbal, B.: The Hadley Circulation in Reanalyses: Climatology, Variability, and Change, J Climate, 26, 3357-3376, 2013.

- Okumura, Y. M., Ohba, M., Deser, C., and Ueda, H.: A Proposed Mechanism for the Asymmetric Duration of El Nino and La Nina, J Climate, 24, 3822-3829, 2011.
- Otto-Bliesner, B. L., Braconnot, P., Harrison, S. P., Lunt, D. J., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Capron, E., Carlson, A. E., Dutton, A., Fischer, H., Goelzer, H., Govin, A., Haywood, A., Joos, F., LeGrande, A. N., Lipscomb, W. H., Lohmann, G., Mahowald, N., Nehrbass-Ahles, C., Pausata, F. S. R., Peterschmitt, J. Y., Phipps, S. J., Renssen, H., and Zhang, Q.: The PMIP4 contribution to CMIP6-Part 2: Two interglacials, scientific objective and experimental design for
 Holocene and Last Interglacial simulations, Geosci Model Dev, 10, 3979-4003, 2017.
- Otto-Bliesner, B. L., Rosenbloom, N., Stone, E. J., McKay, N. P., Lunt, D. J., Brady, E. C., and Overpeck, J. T.: How warm was the last interglacial? New model–data comparisons, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences, 371, 20130097, 2013.
- Otto-Bliesner, B. L., Schneider, R., Brady, E., Kucera, M., Abe-Ouchi, A., Bard, E., Braconnot, P., Crucifix, M., Hewitt, C.,
 and Kageyama, M.: A comparison of PMIP2 model simulations and the MARGO proxy reconstruction for tropical sea surface temperatures at last glacial maximum, Clim Dynam, 32, 799-815, 2009.

Pan, A., Liu, Q., and Liu, Z.: Periodic forcing and ENSO suppression in the Cane-Zebiak model, Journal of oceanography, 61, 109-113, 2005.

Pausata, F. S., Zhang, Q., Muschitiello, F., Lu, Z., Chafik, L., Niedermeyer, E. M., Stager, J. C., Cobb, K. M., and Liu, Z.:
 Greening of the Sahara suppressed ENSO activity during the mid-Holocene, Nat Commun, 8, 16020, 2017.

Phillips, A. S., Deser, C., and Fasullo, J.: Evaluating Modes of Variability in Climate Models, Eos, Transactions American Geophysical Union, 95, 453-455, 2014.

Power, S., Delage, F., Chung, C., Kociuba, G., and Keay, K.: Robust twenty-first-century projections of El Nino and related precipitation variability, Nature, 502, 541-545, 2013.

Deleted: O'Dea, A., Lessios, H. A., Coates, A. G., Eytan, R. I., Restrepo-Moreno, S. A., Cione, A. L., Collins, L. S., de Queiroz, A., Farris, D. W., Norris, R. D., Stallard, R. F., Woodburne, M. O., Aguilera, O., Aubry, M. P., Berggren, W. A., Budd, A. F., Cozzuol, M. A., Coppard, S. E., Duque-Caro, H., Finnegan, S., Gasparini, G. M., Grossman, E. L., Johnson, K. G., Keigwin, L. D., Knowlton, N., Leigh, E. G., Leonard-Pingel, J. S., Marko, P. B., Pyenson, N. D., Rachello-Dolmen, P. G., Stolezon, E., Stolezon, L., Todd, J. A., Vermeij, G. J., and Jackson, J. B. C.: Formation of the Isthmus of Panama, Sci Adv. 2. 2016.

1220 Power, S. B. and Delage, F. P.: El Niño-southern oscillation and associated climatic conditions around the world during the latter half of the twenty-first century, J Climate, 31, 6189-6207, 2018.

Ramanathan, V. and Collins, W.: Thermodynamic Regulation of Ocean Warming by Cirrus Clouds Deduced from Observations of the 1987 El-Nino, Nature, 351, 27-32, 1991.

Ravelo, A. C., Dekens, P. S., and McCarthy, M.: Evidence for El Niño-like conditions during the Pliocene, GSA Today, 16, 4-11, 2006.

Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, J Geophys Res-Atmos, 108, 2003.

Rein, B., Luckge, A., Reinhardt, L., Sirocko, F., Wolf, A., and Dullo, W. C.: El Nino variability off Peru during the last 2000 years, Paleoceanography, 20, 2005.

Riedinger, M. A., Steinitz-Kannan, M., Last, W. M., and Brenner, M.: A similar to 6100 C-14 yr record of El Nino activity from the Galapagos Islands, J Paleolimnol, 27, 1-7, 2002.

Roberts, W. H., Battisti, D. S., and Tudhope, A. W.: ENSO in the Mid-Holocene according to CSM and HadCM3, J Climate, 27, 1223-1242, 2014.

235 Rustic, G. R., Koutavas, A., Marchitto, T. M., and Linsley, B. K.: Dynamical excitation of the tropical Pacific Ocean and ENSO variability by Little Ice Age cooling, Science, 350, 1537-1541, 2015.

Sadekov, A. Y., Ganeshram, R., Pichevin, L., Berdin, R., McClymont, E., Elderfield, H., and Tudhope, A. W.: Palaeoclimate reconstructions reveal a strong link between El Nino-Southern Oscillation and Tropical Pacific mean state, Nat Commun, 4, 2013.

1240 Saint-Lu, M., Braconnot, P., Leloup, J., Lengaigne, M., and Marti, O.: Changes in the ENSO/SPCZ relationship from past to future climates, Earth Planet Sc Lett, 412, 18-24, 2015.

Salau, O., Schneider, B., Park, W., Khon, V., and Latif, M.: Modeling the ENSO impact of orbitally induced mean state climate changes, Journal of Geophysical Research: Oceans, 117, 2012.

Sandweiss, D. H., Richardson, J. B., Reitz, E. J., Rollins, H. B., and Maasch, K. A.: Geoarchaeological evidence from Peru 1245 for a 5000 years BP onset of El Nino, Science, 273, 1531-1533, 1996.

Scroxton, N., Bonham, S. G., Rickaby, R. E. M., Lawrence, S. H. F., Hermoso, M., and Haywood, A. M.: Persistent El Nino-Southern Oscillation variation during the Pliocene Epoch, Paleoceanography, 26, 2011.

Scussolini, P., Bakker, P., Guo, C. C., Stepanek, C., Zhang, Q., Braconnot, P., Cao, J., Guarino, M. V., Coumou, D., Prange, M., Ward, P. J., Renssen, H., Kageyama, M., Otto-Bliesner, B., and Aerts, J. C. J. H.: Agreement between reconstructed and modeled boreal precipitation of the Last Interglacial, Sci Adv, 5, 2019.

Seager, R., Cane, M., Henderson, N., Lee, D. E., Abernathey, R., and Zhang, H. H.: Strengthening tropical Pacific zonal sea surface temperature gradient consistent with rising greenhouse gases, Nat Clim Change, 9, 517-522, 2019.

Shin, S.-I., Sardeshmukh, P. D., Webb, R. S., Oglesby, R. J., and Barsugli, J. J.: Understanding the Mid-Holocene Climate, J Climate, 19, 2801-2817, 2006.

255 Stein, K., Timmermann, A., and Schneider, N.: Phase Synchronization of the El Nino-Southern Oscillation with the Annual Cycle, Phys Rev Lett, 107, 2011.

Stevenson, S. L.: Significant changes to ENSO strength and impacts in the twenty-first century: Results from CMIP5, Geophys Res Lett, 39, 2012.

Stouffer, R. J., Weaver, A. J., and Eby, M.: A method for obtaining pre-twentieth century initial conditions for use in climate change studies, Clim Dynam, 23, 327-339, 2004.

Stuecker, M. F., Jin, F. F., and Timmermann, A.: El Nino-Southern Oscillation frequency cascade, P Natl Acad Sci USA, 112, 13490-13495, 2015.

Tang, Y. and Yu, B.: MJO and its relationship to ENSO, J Geophys Res-Atmos, 113, 2008.

Deleted: +

1265 Taschetto, A. S., Gupta, A. S., Jourdain, N. C., Santoso, A., Ummenhofer, C. C., and England, M. H.: Cold tongue and warm pool ENSO events in CMIP5: mean state and future projections, J Climate, 27, 2861-2885, 2014.

Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, B Am Meteorol Soc, 93, 485-498, 2012.

 Tian, B., and Dong, X.:The double-ITCZ Bias in CMIP3, CMIP5 and CMIP 6 models based on annual mean precipitation,

 270
 Geophys Res Lett, 47, e2020GL087232, 2020.

Tian, Z., Li, T., Jiang, D., and Chen, L.: Causes of ENSO weakening during the Mid-Holocene, J Climate, 30, 7049-7070, 2017.

Timmermann, A., An, S.-I., Kug, J.-S., Jin, F.-F., Cai, W., Capotondi, A., Cobb, K. M., Lengaigne, M., McPhaden, M. J., Stuecker, M. F., Stein, K., Wittenberg, A. T., Yun, K.-S., Bayr, T., Chen, H.-C., Chikamoto, Y., Dewitte, B., Dommenget,

- 1275 D., Grothe, P., Guilyardi, E., Ham, Y.-G., Hayashi, M., Ineson, S., Kang, D., Kim, S., Kim, W., Lee, J.-Y., Li, T., Luo, J.-J., McGregor, S., Planton, Y., Power, S., Rashid, H., Ren, H.-L., Santoso, A., Takahashi, K., Todd, A., Wang, G., Wang, G., Xie, R., Yang, W.-H., Yeh, S.-W., Yoon, J., Zeller, E., and Zhang, X.: El Niño–Southern Oscillation complexity, Nature, 559, 535-545, 2018.
- Timmermann, A., Oberhuber, J., Bacher, A., Esch, M., Latif, M., and Roeckner, E.: Increased El Nino frequency in a climate model forced by future greenhouse warming, Nature, 398, 694-697, 1999.

Timmermann, A., Okumura, Y., An, S. I., Clement, A., Dong, B., Guilyardi, E., Hu, A., Jungclaus, J. H., Renold, M., Stocker, T. F., Stouffer, R. J., Sutton, R., Xie, S. P., and Yin, J.: The influence of a weakening of the Atlantic meridional overturning circulation on ENSO, J Climate, 20, 4899-4919, 2007.

Trenberth, K. E.: The definition of El Nino, B Am Meteorol Soc, 78, 2771-2777, 1997.

1285 Trenberth, K. E. and Hoar, T. J.: El Nino and climate change, Geophys Res Lett, 24, 3057-3060, 1997.

Tudhope, A. W., Chilcott, C. P., McCulloch, M. T., Cook, E. R., Chappell, J., Ellam, R. M., Lea, D. W., Lough, J. M., and Shimmield, G. B.: Variability in the El Nino - Southern oscillation through a glacial-interglacial cycle, Science, 291, 1511-1517, 2001.

yan Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A. L., Field, R. D., Arneth, A., Forrest, M., Hantson, S.,
 Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), Geosci. Model Dev., 10, 3329-3357, 2017.

Vecchi, G. A. and Soden, B. J.: Global warming and the weakening of the tropical circulation, J Climate, 20, 4316-4340, 2007.

1295 Wang, G. J., Cai, W. J., Gan, B. L., Wu, L. X., Santoso, A., Lin, X. P., Chen, Z. H., and McPhaden, M. J.: Continued increase of extreme El Nino frequency long after 1.5 degrees C warming stabilization, Nat Clim Change, 7, 568-572, 2017.

Wara, M. W., Ravelo, A. C., and Delaney, M. L.: Permanent El Niño-like conditions during the Pliocene warm period, Science, 309, 758-761, 2005.

Watanabe, T., Suzuki, A., Minobe, S., Kawashima, T., Kameo, K., Minoshima, K., Aguilar, Y. M., Wani, R., Kawahata, H.,
Sowa, K., Nagai, T., and Kase, T.: Permanent El Nino during the Pliocene warm period not supported by coral evidence, Nature, 471, 209-211, 2011.

White, S. M., and Ravelo, A. C.: Dampened El Niño in the early Pliocene warm period, Geophys Res Lett, 47, e2019GL085504, 2020.

White, S. M., Ravelo, A. C., and Polissar, P. J.: Dampened El Niño in the early and mid-Holocene due to insolation-forced
 warming/deepening of the thermocline, Geophys Res Lett, 45, 316–326, 2018.

Widlansky, M. J., Timmermann, A., Stein, K., McGregor, S., Schneider, N., England, M. H., Lengaigne, M., and Cai, W. J.: Changes in South Pacific rainfall bands in a warming climate, Nat Clim Change, 3, 417-423, 2013.

Wittenberg, A. T.: Are historical records sufficient to constrain ENSO simulations?, Geophys Res Lett, 36, 2009.

Deleted: Tziperman, E. and Farrell, B.: Pliocene equatorial temperature: Lessons from atmospheric superrotation, Paleoceanography, 24, 2009.

Xie, S.-P., Deser, C., Vecchi, G. A., Ma, J., Teng, H., and Wittenberg, A. T.: Global warming pattern formation: Sea surface temperature and rainfall, J Climate, 23, 966-986, 2010.

Yang, S., Li, Z., Yu, J.-Y., Hu, X., Dong, W., and He, S.: El Niño–Southern oscillation and its impact in the changing climate, Natl Sci Rev, 5, 840-857, 2018.

Yeh, S.-W., Kug, J.-S., and An, S.-I.: Recent progress on two types of El Niño: Observations, dynamics, and future changes, Asia-Pacific Journal of Atmospheric Sciences, 50, 69-81, 2014.

Yeh, S. W., Cai, W. J., Min, S. K., McPhaden, M. J., Dommenget, D., Dewitte, B., Collins, M., Ashok, K., An, S. I., Yim, B.
 Y., and Kug, J. S.: ENSO Atmospheric Teleconnections and Their Response to Greenhouse Gas Forcing, Reviews of
 Geophysics, 56, 185-206, 2018.

Yeh, S. W. and Kirtman, B. P.: ENSO amplitude changes due to climate change projections in different coupled models, J Climate, 20, 203-217, 2007.

Yeh, S. W., Kug, J. S., Dewitte, B., Kwon, M. H., Kirtman, B. P., and Jin, F. F.: El Nino in a changing climate, Nature, 461, 511-U570, 2009.

225 Zebiak, S. E.: Air-sea interaction in the equatorial Atlantic region, J Climate, 6, 1567-1586, 1993.

Zhang, C. D. and Gottschalck, J.: SST anomalies of ENSO and the Madden-Julian oscillation in the equatorial Pacific, J Climate, 15, 2429-2445, 2002.

Zhang, Q., Guan, Y., and Yang, H. J.: ENSO amplitude change in observation and coupled models, Adv Atmos Sci, 25, 361-366, 2008.

230 Zhang, T. and Sun, D. Z.: ENSO Asymmetry in CMIP5 Models, J Climate, 27, 4070-4093, 2014.

Zhang, X. X., Liu, H. L., and Zhang, M. H.: Double ITCZ in Coupled Ocean-Atmosphere Models: From CMIP3 to CMIP5, Geophys Res Lett, 42, 8651-8659, 2015.

Zheng, W., Braconnot, P., Guilyardi, E., Merkel, U., and Yu, Y.: ENSO at 6ka and 21ka from ocean-atmosphere coupled model simulations, Clim Dynam, 30, 745-762, 2008.

1335 Zhu, J., Liu, Z., Brady, E., Otto-Bliesner, B., Zhang, J., Noone, D., Tomas, R., Nusbaumer, J., Wong, T., Jahn, A., and Tabor, C.: Reduced ENSO variability at the LGM revealed by an isotope-enabled earth system model, Geophys Res Lett, 44, 6984– 6992, 2017.

Model			historical		lam	1:~1271	InstCO2	· ·
	gen.	Control	nisioricai	Holocene	ıgm	llg12/K	IpciCO2	CO2
AWI-ESM-1-1-LR	CMIP6	100	-	100	100	100	-	-
BCC-CSM1-1	CMIP5	500	163	100	-	-	140	150
CCSM4	CMIP5	1051	156	301	101	-	156	151
CESM2	CMIP6	1200	165	700	-	700	150	150
CNRM-CM5	CMIP5	850	156	200	200	-	140	150
CNRM-CM6-1	<u>CMIP6</u>	<u>500</u>	<u>165</u>	Ξ	± l	<u>301</u>	<u>150</u>	<u>150</u>
COSMOS-ASO	CMIP5	400	-	-	600	-	-	-
CSIRO-Mk3-6-0	CMIP5	500	156	100	-	-	140	150
CSIRO-Mk3L-1-2	CMIP5	1000	150	500	-	-	140	-
EC-EARTH-3-LR	<u>CMIP6</u>	<u>201</u>	Ξ	<u>201</u>	± l	Ξ.	z.	Ξ
FGOALS-f3-L	CMIP6	561	165	500	-	500	160	160
FGOALS-g2	CMIP5	700	115	680	100	-	244	258
FGOALS-g3	CMIP6	700	-	500	-	500	-	-
FGOALS-s2	CMIP5	501	-	100	-	-	140	150
GISS-E2-1-G	CMIP6	851	165	100	-	100	51	151
GISS-E2-R	CMIP5	500	156	100	100	-	151	151
HadGEM2-CC	CMIP5	240	145	35	-	-	-	-
HadGEM2-ES	CMIP5	336	145	101	-	-	140	151
HadGEM3-GC31-	CMIP6	100	-	100	_	200	_	_
LL	civili u	100		100		200		
INM-CM4-8	CMIP6	531	165	200	200	100	150	150
IPSL-CM5A-LR	CMIP5	1000	156	500	200	-	140	260
IPSL-CM6A-LR	CMIP6	1200	165	550	-	550	150	900
MIROC-ES2L	CMIP6	500	165	100	100	100	150	150
MIROC-ESM	CMIP5	630	156	100	100	-	140	150
MPI-ESM-P	CMIP5	1156	156	100	100	-	140	150
MPI-ESM1-2-LR	CMIP6	<u>1000</u>	1	<u>500</u>	=	=	<u>-</u>	<u>165</u>
MRI-CGCM3	CMIP5	500	156	100	100	-	140	150

Table 1: List of models included in study and length of simulations based on number of years of data available for NINO3.41340in the CVDP archive. Additional information about CMIP6/PMIP4 models (indicated in bold) is provided in the
Supplementary Material.

MRI-ESM2-0	CMIP6	701	165	200	-	-	151	151
NESM3	CMIP6	100	165	100	-	100	150	150
NorESM1-F	CMIP6	200	-	200	-	200	-	-
NorESM2-LM	CMIP6	<u>391</u>	<u>65</u>	<u>100</u>	=	=	Ξ.	<u>380</u>
UofT-CCSM-4	CMIP6	100	-	100	100	-	-	-

Table 2: Multi-model mean change in ENSO amplitude (%) based on NINO3.4 standard deviation for each experiment
 relative to the pre-industrial control for all models, CMIP5/PMIP3 models and CMIP6/PMIP4 models. Note that CMIP5/PMIP3 does not include *lig127k* experiment. Values in brackets for All models are the minimum and maximum model values.

Experiment	All models ENSO change %	<u>CMIP5</u> ENSO change %	<u>CMIP5</u> number models	<u>CMIP6</u> ENSO change %	<u>CMIP6</u> <u>number models</u>
midHolocene	<u>-8.4 (-36.8 to 18.5)</u>	<u>-6.9</u>	<u>14</u>	<u>-9.7</u>	<u>16</u>
<u>lgm</u>	<u>-4.8 (-46.9 to 32.5)</u>	<u>-4.7</u>	<u>9</u>	<u>-4.8</u>	<u>3</u>
<u>lig127k</u>	<u>-20.6 (-54.2 to 10.3)</u>	=	Ξ.	-20.6	<u>12</u>
<u>lpctCO2</u>	<u>5.6 (-33.2 to 34.1)</u>	<u>1.4</u>	<u>13</u>	<u>11.6</u>	<u>9</u>
abrupt4xCO2	<u>2.2 (-46.3 to 57.7)</u>	<u>3.0</u>	<u>12</u>	<u>1.4</u>	<u>11</u>



Figure 1: Ability of the ensemble to simulate present-day sea surface temperature (SST) patterns: (a) DJF and (b) JJA SST climatology from HadISST observational dataset (Rayner et al., 2003) between 1971-2000, (c) DJF and (d) JJA model ensemble mean SST in historical simulations minus HadISST observations between 1971-2000, and (e) DJF and (f) JJA model ensemble mean SST in pre-industrial control simulations minus HadISST observations. Units are °C. <u>Stippling</u> indicates that more than 2/3 of the ensemble members agree on the sign of the anomaly.



Figure 2: Ability of the ensemble to simulate present-day precipitation patterns: (a) DJF and (b) JJA SST climatology from GPCP observational dataset (Adler et al., 2003) between 1979-1999, (c) DJF and (d) JJA model ensemble mean precipitation in historical simulations minus GPCP observations between 1979-1999, and (e) DJF and (f) JJA model ensemble mean precipitation in pre-industrial control simulations minus GPCP observations. Units are mm/day. Stippling indicates that more than 2/3 of the ensemble members agree on the sign of the anomaly.



370

Figure 3: Evaluation of the ENSO SST anomaly pattern. Composite El Niño minus La Niña sea surface temperature anomaly from (a) HadISST observational dataset between 1871-2012, (b) model ensemble average from the historical simulations between 1850-2005 (for CMIP5 models) or 1850-2015 (for CMIP6 models), and (c) model ensemble average from the pre-industrial control simulations. Units are °C.



1375 Figure 4: Evaluation of the ENSO temperature teleconnections. Composite El Niño minus La Niña surface temperature anomaly from (a) DJF and (b) JJA from the C20 Reanalysis (Compo et al., 2011) between 1871-2010, (c) DJF and (d) JJA model ensemble mean from the historical simulations between 1850-2005 (for CMIP5 models) or 1850-2015 (for CMIP6 models), and (e) DJF and (f) JJA model ensemble mean ensemble from the pre-industrial control simulations. Units are °C.



Figure 5: Evaluation of the ENSO precipitation teleconnections. Composite El Niño minus La Niña precipitation anomaly from (a) DJF and (b) JJA from the C20 Reanalysis (Compo et al., 2011) between 1871-2010, (c) DJF and (d) JJA model ensemble mean from the historical simulations between 1850-2005(CMIP5)/2015 (CMIP6), and (e) DJF and (f) JJA model ensemble mean ensemble from the pre-industrial control simulations. Units are mm/day.



Figure 6: Ensemble mean seasonal changes in sea surface temperature in experiment minus pre-industrial control simulations: (a) DJF *midHolocene*, (b) JJA *midHolocene*, (c) DJF *lgm*, (d) JJA *lgm*, (e) DJF *lig127k*, (f) JJA *lig127k*, (g) DJF *lpctCO2*, (h) JJA *lpctCO2*, (i) DJF *abrupt4xCO2*, (j) JJA *abrupt4xCO2*. The ensemble mean temperature pattern in the pre-industrial control simulations is shown as black contours. Units are °C. <u>Stippling indicates that more than 2/3 of the ensemble members agree on the sign of the change.</u>



Figure 7: Ensemble mean seasonal changes in precipitation in experiment minus pre-industrial control simulations: (a) DJF midHolocene, (b) JJA midHolocene, (c) DJF lgm, (d) JJA lgm, (e) DJF lig127k, (f) JJA lig127k, (g) DJF lpctCO2, (h) JJA lpctCO2, (i) DJF abrupt4xCO2, (j) JJA abrupt4xCO2. The ensemble mean temperature pattern in the pre-industrial control simulations is shown as black contours. Units are mm/day. Stippling indicates that more than 2/3 of the ensemble members agree on the sign of the change.



Figure 8: Amplitude of ENSO measured from standard deviation of NINO3.4 index (°C) in *piControl* simulations (grey bars) and (a) *midHolocene* (dark green bars), (b) *lgm* (blue bars), (c) *lig127k* (light green bars), (d) *lpctCO2* (dark red bars) and (e) *abrupt4xCO2* (light red bars) simulations. Model names are given below plots. <u>Multi-Model (MM) Mean is also shown.</u>



Figure 9: Change in amplitude of ENSO measured from standard deviation of NINO3.4 index relative to *piControl* amplitude (%) in (a) *midHolocene*, (b) *lgm*, (c) *lig127k*, (d) *1pctCO2* and (e) *abrupt4xCO2*. Model names are given below plots. Multi-Model (MM) Mean is also shown.



1410 Figure 10: The changes in the SST pattern associated with ENSO in each experiment compared with *piControl*. The ensemble mean difference between the composites of each models El Niño minus La Niña (defined as ±1 standard deviation) in (a) *midHolocene*, (b) *lgm*, (c) *lig127k*, (d) *1pctCO2* and (e) *abrupt4xCO2* experiments minus the same pattern for the *piControl* simulations The ensemble mean ENSO SST pattern in the *piControl* simulations are shown as black contours. Stippling indicates that more than 2/3 of the ensemble members agree on the sign of the change.





compared with the pre-industrial control. The ensemble mean difference between the composites of each models El Niño minus La Niña (defined as ±1 standard deviation) in (a, b) midHolocene, (c, d) lgm, (e, f) lig127k, (g, h) lpctCO2 and (i, j) 420 abrupt4xCO2 experiments minus the same pattern for the piControl simulations The ensemble mean ENSO pattern in the piControl simulations are shown as black contours. Stippling indicates that more than 2/3 of the ensemble members agree on the sign of the change.



Figure 12: The changes in the seasonal precipitation teleconnection pattern associated with ENSO in each experiment compared with the pre-industrial control. The ensemble mean difference between the composites of each models El Niño minus La Niña (defined as ±1 standard deviation) in (a, b) *midHolocene*, (c, d) *lgm*, (e, f) *lig127k*, (g, h) *1pctCO2* and (i, j) *abrupt4xCO2* experiments minus the same pattern for the *piControl* simulations The ensemble mean ENSO pattern in the *piControl* simulations are shown as black contours. Stippling indicates that more than 2/3 of the ensemble members agree on the sign of the change.



Figure 13: Relationships across all experiments and all models: (a) ENSO amplitude change (%) versus change in the annual cycle (%), (b) ENSO amplitude change (°C) versus zonal SST gradient change (°C), and (c) NINO3 precipitation for ENSO composite versus meridional SST gradient change (°C). All changes are relative to *piControl*, see text for details. Experiments are identified as follows: blue squares = *lgm*, dark green circles = *midHolocene*, light green diamonds = *lig127k*, dark red stars = *lpctCO2*, light red triangles = *abrupt4xCO2* (after bar charts in Figure 8). <u>Correlation coefficients are shown</u>

435 dark red stars for each plot.