

Response letter to

Editor Decision: Publish subject to minor revisions (review by editor) (07 May 2019)

Thank you for your responses to reviewers. As you saw, the two reviews are positive and conclude on minor revisions. I have carefully read your responses to the reviewers and your propositions to modify the text of your manuscript.

Your paper is very interesting and I appreciate your detailed responses. Nevertheless, your manuscript needs some additional amendments corresponding to some of the comments of the two reviewers and few questions for which I would like your responses.

1- Following the comments of reviewers 2, I think that Bergwinds have to be explained to readers in the text that may be interested on this paper but not familiarized on the studied area.

We insert: "offshore winds descending from the interior plateau" (Page 5, Line 6-7)

2- I have some problems with definitions of endmembers as reviewer 1. Probably it will be easier to understand if when describing the endmembers in the corresponding paragraph you separate the different endmembers and their composition by subtitles each one for an endmember and then the correspondence with the previous one. For my part I am a bit lost in this part.

Each paragraph concerns 1 endmember, which we mention at the beginning of the paragraph (Page 9, Line 14, Page 10, Lines 8, 16, 32. We also introduce these names in the preceding paragraph (Page 9, Line 1-13).

In fact I really think as the reviewer 1 that a short table in the main text will be helpful even if we have some details in the supplement with the curves of all taxa presented separately (do the curves of all taxa be in the supplementary figures?).

OK, we add Table 2 (Page 10)

Table 2. Interpretation of the endmembers

Endmember	Main pollen taxa
E-Heathland	<i>Podocarpus</i> , <i>Celtis</i> , <i>Olea</i>
E-Mountain-Forest	Cyperaceae, Ericaceae, <i>Phaeoceros</i> , Restionaceae, <i>Stoebe</i> type, <i>Anthoceros</i> , <i>Typha</i> <i>Lycopodium</i> , Restionaceae
E-Shrubland	Poaceae, Asteroideae, <i>Buxus</i> , Amaranthaceae, <i>Euphorbia</i> , Meliaceae-Sapotaceae, <i>Acacia</i> , <i>Riccia</i> type, <i>Tribulus</i> , Acanthaceae pp, Asteraceae Vernoniae, <i>Hypoestes-Dicliptera</i> type, <i>Gazania</i> type, <i>Dombeya</i>
E-Woodland	<i>Alchornea</i> , <i>Spirostachys africana</i> , <i>Pteridium</i> type, Polypodiaceae, <i>Myrsine africana</i> , <i>Cassia</i> type, Rhizophoraceae, Aizoaceae, Combretaceae pp, <i>Manilkara</i> , <i>Burkea africana</i> , <i>Brachystegia</i> , <i>Dodonaea viscosa</i> , <i>Pseudolachnostylis</i> , <i>Hymenocardia</i> , <i>Aloe</i> , Rhamnaceae pp, <i>Protea</i> , <i>Parinari</i>

All curves mentioned in the text are plotted in the supplementary figures. I consider putting the supplementary figures into the main text, but that might be overkill. What do you think?

In fact you write the term endmembers in page before define it as the reviewer 1 said. what you add in the matériel and method sections is very short. what do you mean with "characteristic combination?"

Something that can be interpreted. We rephrase as follows:

“We regard the pollen percentages as a series of pollen assemblage mixtures, whereby each modelled endmember may be interpreted as the representation of one or more biomes.” (Page 6, Line 30 - Page 7, Line 1)

You add a citation of table 1 and table 2 supplementary documents but they are not in the documents given with the responses, I did not find them.
Please explain me what happened and /or produce these tables.

Supplementary Tables 1 and 2 have not changed. I upload them again. The supplementary figures are trailing the manuscript.

3- Concerning your corrected discussion I have some remarks:

a. In the figures it will be better to indicate glacial and interglacial numbers, sometimes you mark the glacial numbers sometimes the interglacial ones sometimes the terminations. I did not understand the choice (numbering glacial and interglacial or terminations???) in the different figures even in the supplementary ones. Explain me.

The choice for Terminations is that they clearly separate the climate cycles. We denote interglacial stages in the figures showing curves of vegetation variability associated with interglacials, such as E-woodland and E-shrubland. We use Terminations in the figures denoting vegetation variability associated with glacials, such as E-Heathland and A-Mountain-Forest. We add even MIS numbers at Figure 5.

b. End of p. 11 (new text) you say that mountain forest is replaced by woodlands in 11c, 9e, 7e, 7c, 5e and 1. First will you mark the in the right order. I do not see that in 7e and 7c. I see only a slight regression of mountain taxa and a slight increase in woodland in these two events.

Yes, ‘replaced’ is not the right word. We change the sentence into: “Woodlands would have expanded at the cost of mountain forest during 11c, 9e, 5e and 1, and to a lesser extend during 7e and 7c, “ (Page 12, Line 7-10)

c. p. 12 first sentence you explain “the record.....during those parts of glacial stages”. I do not understand, you were talking about interglacials???? And then the term “those part of the glacial stages” refers to what? I am lost here. Why did you talk about glacial in the description of interglacials?

Actually, the biomes and not the glacials or interglacials are the main themes of those paragraphs. We rearranged the paragraph about mountain forest as follows (Page 12, Line 12-16, Page 13, Line 1-2):

The glacial stages showed the expansion of either mountain forest or heathland. The record indicates extension of mountain forests in SE Africa during those parts of the glacial stages with low temperatures and atmospheric $p\text{CO}_2$ exceeding ~ 220 ppmv (Figure 5). If low temperatures were the only driver of the extension of mountain forests, further spread into the lowlands during the coldest glacial phases should be expected. Instead, when $p\text{CO}_2$ dropped below ~ 220 ppmv during those colder glacial periods, mountain forest declined, in particular during MIS 18, 16, 14, 8, 6, and 2. A picture emerges of cool glacial stages in SE Africa in which tree cover broke down when atmospheric $p\text{CO}_2$ became too low. Additionally, mountain forests were important during the Interglacials 19c, 17c, 15e, 15a, 13a, and 7e, in which $p\text{CO}_2$ and Antarctic temperatures were subdued.

You talk about Podocarpus and you cited the fig. 5 why? Podocarpus is not on the figure 5 only in additional figures.

We changed Podocarpus forest into mountain forest (see above).

d. Line 4 p.12 you explain that you have mountain forest in the coldest glacial just before the interglacials marked by the replacement between the two types of vegetation so MIS 18, 16, 14, 8, 6 and 2. First, here fig. 5 has to be cited please and it will be helpful to write at least the MIS glacial numbers on the figure).

Figure 5 (Page 1, Line 15) is cited and we add MIS glacial stages to the figure.

Have you an explanation for the difference observed in the 12 and 10? They are not so different from the 14.

No, not really.

e. I need also some links between the C4 vegetation, the C4 sedges and the endmembers in the text. In the beginning of the discussion, I understood that the C4 vegetation is represented in you record mainly by sedges. Is it true?

Yes

Then it mainly corresponds to old EM2 and new E heathland? Is it true?

Yes, see discussion section about endmember interpretation

Please add this in some sentences of the discussion to be clear.

We add sedges percentages (Cyperaceae %) to Figure 5. We also insert “being an important constituent of the ericaceous fynbos-like vegetation” after C4 sedges (Page 13, Line 12-13) and “(see also the correlation between SST and EM2 in Dupont et al. 2011)” after correlating with the SST record (Page 13, Line 34).

Please clarify these points in the discussion and organize the discussion in a different way to be more clear for the reader (first the interglacials and secondly the glacials) and cite the supplementary figures if necessary.

I do not think that these last remarks and comments take a long time to be addressed. So I am waiting after your new text and then will be able to post my final decision on the publication of this very interesting manuscript

With my best regards
Nathalie Combourieu-Nebout

Thank you very much. I hope the changes and responses are satisfactory.

With kind regards and greetings,
Lydie Dupont

Effects of atmospheric CO₂ variability of the past 800 ka on the biomes of Southeast Africa

Lydie M. Dupont, MARUM - Center for Marine Environmental Sciences, University of Bremen

Thibaut Caley - EPOC, UMR 5805, CNRS, University of Bordeaux, Pessac, France

Isla S. Castañeda - University of Massachusetts Amherst, Department of Geosciences, Amherst, MA 01003 USA

Abstract

Very little is known about the impact of atmospheric carbon dioxide pressure ($p\text{CO}_2$) on the shaping of biomes. The development of $p\text{CO}_2$ throughout the Brunhes Chron may be considered a natural experiment to elucidate relationships between vegetation and $p\text{CO}_2$. While the glacial periods show low to very low values (~220 to ~190 ppmv, respectively), the $p\text{CO}_2$ levels of the interglacial periods vary from intermediate to relatively high (~250 to more than 270, respectively). To study the influence of $p\text{CO}_2$ on the Pleistocene development of SE African vegetation, we used the pollen record of a marine core (MD96-2048) retrieved from Maputo Bay south of the Limpopo River mouth in combination with stable isotopes and geochemical proxies. Applying endmember analysis, four pollen assemblages could be distinguished representing different biomes: heathland, mountain forest, shrubland and woodland. We find that the vegetation of the Limpopo River catchment and the coastal region of southern Mozambique is not only influenced by hydroclimate but also by temperature and atmospheric $p\text{CO}_2$. Our results suggest that the extension of mountain forest occurred during those parts of the glacials when $p\text{CO}_2$ and temperatures were moderate and that only during the colder periods when atmospheric $p\text{CO}_2$ was low (less than 220 ppmv) open ericaceous vegetation including C4 sedges extended. The main development of woodlands in the area took place after the Mid-Brunhes Event (~430 ka) when interglacial $p\text{CO}_2$ levels regularly rose over 270 ppmv.

Keywords

Palynology, Pleistocene, Limpopo River catchment, atmospheric carbon dioxide, Mid-Brunhes Event

Short summary

Multiproxy study of marine sediments off the Limpopo River mouth spanning the Late Pleistocene reveals the impact of atmospheric carbon dioxide on the development of the vegetation of Southeast Africa and indicates changes in the interglacial vegetation before and after the Mid-Brunhes Event.

Highlight

A multiproxy study of marine sediments off the Limpopo River mouth spanning the Late Pleistocene reveals the impact of atmospheric carbon dioxide on the development of the vegetation of Southeast Africa and indicates changes in the interglacial vegetation before and after the Mid-Brunhes Event. The unique record of detailed vegetation change in SE Africa over the entire Brunhes Chron demonstrates the expansion of glacial vegetation in southern Africa when atmospheric CO₂ concentration was low and the development of miombo woodland in SE Africa during successive interglacials after the Mid-Brunhes Transition.

1 Introduction

2 Understanding the role of atmospheric carbon dioxide pressure ($p\text{CO}_2$) is paramount for the
3 interpretation of the of the paleo-vegetation record. The effects of low $p\text{CO}_2$ on glacial vegetation
4 have been discussed in a number of studies [Ehleringer et al. 1997, Jolly & Haxeltine 1997, Cowling &
5 Sykes 1999, Prentice & Harrison 2009, Prentice et al. 2017] predicting that glacial increases in C4
6 vegetation favored by low atmospheric CO_2 would have opened the landscape and lowered the tree
7 line. Comparing records of this glacial C4-rich vegetation with modern analogues could have led to
8 estimating more severe aridity than actually occurred during the Last Glacial Maximum. These
9 studies [opt cit.], however, mostly cover the last glacial-interglacial transition and do not examine
10 periods with intermediate $p\text{CO}_2$ such as during the Early Glacial (MIS 5a-d) or interglacials prior to
11 430 thousand years ago (430 ka). Comparing the vegetation record of subsequent climate cycles
12 showing different CO_2 levels might help to better understand the effects of $p\text{CO}_2$ on the vegetation.

13 During the Brunhes Chron (past 780 ka) the length of the glacial cycles became much longer lasting
14 roughly 100 ka due to a strong non-linear response of the ice sheets to solar forcing [Mudelsee &
15 Stattegger 1997]. Model experiments of Ganopolski & Calov [2011] indicate that low atmospheric
16 CO_2 concentrations are a prerequisite for the long duration of the glacial cycles of the past 800 ka.
17 Then, roughly mid-way through the Brunhes Chron, the amplitude of the climate cycles shifted with a
18 change in the maximum CO_2 concentration during interglacials.

19 This so-called Mid-Brunhes Event (MBE) [Jansen et al. 1986] - also called Mid-Brunhes Transition -
20 occurred about 430 ka ago and marks the transition between interglacials characterized by rather
21 low atmospheric CO_2 around 240 ppm (parts of Marine Isotope Stages (MIS) 19, 17, 15, 13) to
22 interglacials in which CO_2 levels reached 270 ppmv or more (parts of MIS 11, 9, 7, 5, 1) [Lüthi et al.
23 2008, Bereiter et al. 2015]. The climate transition of the MBE has been extensively studied using
24 Earth System Models of Intermediate Complexity. Yin & Berger [2010] stress the importance of
25 forcing by austral summer insolation and Yin & Berger [2012] argue that the model vegetation (tree-
26 fraction) was forced by precession through precipitation at low latitudes. Both papers show the
27 necessity to include the change in atmospheric CO_2 in the explanation of the MBE [Yin & Berger,
28 2010, 2012]. Yin [2013], however, concludes that it is not necessary to invoke a sudden event around
29 430 ka to explain the increased interglacial CO_2 ; the differences between interglacials before and
30 after the MBE can be explained by individual responses in Southern Ocean ventilation and deep-sea
31 temperature to various combinations of the astronomical parameters. On the other hand, statistical
32 analysis suggests a dominant role of the carbon cycle, which changed over the MBE [Barth et al.
33 2018]. Paillard [2017] developed a conceptual model of orbital forcing of the carbon cycle in which
34 sea-level fluctuations and the effects on carbon burial are decisive during shifts in the climate
35 system. Further modelling by Bouttes et al. [2018] showed qualitative agreement with the paleodata
36 of pre- and post-MBE interglacials but largely underestimated the amplitude of the changes.
37 Moreover, the simulated vegetation seems to counteract the effects of the oceanic response
38 [Bouttes et al. 2018]. Thus the vegetation, in particular at low latitudes, may play a crucial but poorly
39 understood role in the climate system.

1 Comparing records of pre- and post-MBE interglacials could offer insight in the interglacial climate at
2 different levels of CO₂ [Foley et al. 1994, Swan et al. 2010]. We define interglacials after PAGES
3 [2016] listing MIS 19c, 17c, 15a, 15e, 13a as pre-MBE and MIS 11c, 9e, 7e, 7a-c, 5e, 1 as post-MBE.
4 Currently, only a handful of vegetation records covering the entire Brunhes Chron have sufficient
5 temporal resolution to enable comparisons between interglacials before and after the Mid-Brunhes
6 transition. These records are from the eastern Mediterranean, the Colombian Andes [PAGES 2016].




Figure 1. Upper panel: map of southern Africa with the main phytocoria after White (1983). Lower panel: site location of MD96-2048; main vegetation formations; main rivers; 100 m, 200 m, 500 m, and 1000m contours; 200 m, 500 m, and 1000m bathymetric contours; Agulhas (AC) and counter currents (CC) forming a coastal Maputo Bight Lee Eddy. Zambezian vegetation woodland and savannah north of ~25°30'S, Tongaland-Pondoland coastal forests south of ~25°30'S, Zanzibar-Inhambane coastal forests east of 33–34°E. West of the escarpment with Afromontane forest rises the interior plateau covered with Highveld grasslands rises.

West and East Africa [Dupont et al. 1989, Miller & Goslin 2014, Castañeda et al. 2016, Johnson et al. 2016, Ivory et al. 2018, Owen et al. 2018]. The Andean pollen record is strongly influenced by the immigration of oak from North America during MIS 12 [Torres et al. 2013]. For the eastern Mediterranean a decline in plant diversity is observed at Tenaghi Phillipon (Greece) where the modern Mediterranean oak forests gradual emerged in the interglacials after MIS 16 but before the MBE [Tzedakis et al. 2006, 2009]. The West African record of Lake Bosumtwi in Ghana allows identification of six forest assemblages since 540 ka related to the interglacials of MIS 13, 11, 9, 7, 5e, and 1. The forests assemblage of MIS 13, however, does not show a strong contrast with those of the interglacials after the MBE [Miller & Goslin 2014]. The marine pollen record of ODP Site 658 off Cape Blanc tracks the latitudinal position of the open grass-rich vegetation zones at the boundary between Sahara and Sahel suggesting shifting vegetation zones between glacials and interglacials [Dupont & Hooghiemstra 1989, Dupont et al. 1989]. The drier interglacials occurred after MIS 9, which indicates a transition after the MBE to more arid conditions. Additionally, stable carbon isotope records from Chinese loess sections indicate interglacial-glacial variability in the C3-C4 proportions of the vegetation [Lyu et al. 2018, Sun et al. 2019]. However, the latter records do not show a prominent vegetation shift over the MBE.

For East Africa two terrestrial records and a marine one are available. From Lake Malawi, Johnson et al. [2016] infer wetter conditions and increased woodland vegetation between 800 and 400 ka based on the stable carbon isotopic composition of plant wax shifting from less to more strongly depleted values. Also from Lake Malawi, Ivory et al. [2018] published a pollen record of the past 600 ka revealing a number of phases of miombo woodland and mountain forest alternating with savannah vegetation (dry woodland and wooded grassland). Recently, a new record from Lake Magadi (Kenya) has been published indicating a change from wetter conditions to more aridity after 500 ka contrasting the Lake Malawi results [Owen et al. 2018]. In Lake Magadi, the representation of *Podocarpus* decreased over the MBE, while open grassy vegetation and salinity of the lake increased [Owen et al. 2018]. Neither the Lake Malawi nor the Lake Magadi records show dominant interglacial-glacial variability.

The marine record retrieved south of the Limpopo River mouth (Core MD96-2048) allows inferences about vegetation and climate in the catchment area of the Limpopo River draining large areas of South Africa, Botswana, Zimbabwe and Mozambique. Based on sediment chemistry, Caley et al. [2018] reported the effects of increased summer insolation in increased fluvial discharge and variability associated with eccentricity, which modulates precession amplitudes. Superimposed on the orbital-scale precipitation variability, a long-term trend from 1000 to 600 ka towards increased aridity in southeastern Africa was found [Caley et al. 2018]. The plant leaf wax carbon isotopic (hereafter $\delta^{13}\text{C}_{\text{wax}}$) record of the same core was originally interpreted as reflecting a trend toward increasingly drier glacials and wetter interglacials over the past 800 ka [Castañeda et al. 2016]. Additionally, the average chain lengths of the plant leaf waxes exhibit a stepwise decrease at 430 ka suggesting a change from more shrub vegetation before the MBE to a larger contribution of trees during the post-MBE interglacials [Castañeda et al. 2016]. Thus, a pollen record of MD96-2048 has the potential to register the changes in interglacial vegetation cover over the MBE. We might expect a change of Southern Hemisphere vegetation being less ambiguous than the changes found on the Northern Hemisphere (see above), because modelling indicates that the effects of the MBE were more pronounced on the Southern Hemisphere [Yin & Berger 2010]. Until now, the palynology of only the last 350 ka has been published [Dupont et al. 2011] and, therefore, here we extend the pollen record of MD96-2048 to cover the past 800 ka in sufficient resolution. As described below, our new palynology results have led to the re-interpretation of the MD96-2048 $\delta^{13}\text{C}_{\text{wax}}$ record [Castañeda et al. 2016].

Previous work on Core MD96-2048

The sediments of MD96-2048 were retrieved in the middle of the Maputo Bight (Figure 1) from the southern Limpopo cone forming a depot center that has been build up at least since the Late Miocene [Martin 1981]. The site collects terrestrial material including pollen and spores mostly from the rivers that discharge into the Maputo Bight of which the Limpopo River is the biggest draining large areas of northern South Africa and southern Mozambique. Apart from the offshore winds descending from the interior plateau, so-called Bergwinds, the predominant wind direction is landward [Tyson & Preston-Whyte 2000] and aeolian input of terrestrial material is probably minor. Thus, pollen source areas would cover the region north of the Maputo Bight in southern Mozambique and the region west of Maputo of the Lebombo hills and the Drakensberg Escarpment [Dupont et al. 2011].

A wide variety of measurements have been performed on MD96-2048 sediments. Caley et al. [2011, 2018] recorded stable oxygen isotopes of benthic foraminifers (*Planulina wuellerstorfi*) providing a stable oxygen stratigraphy and age model aligned to the global stack LR04 [Lisiecki & Raymo 2005] for the past 2200 ka. Trace element (Mg/Ca ratios) of the planktic foraminifer *Globigernoides ruber sensu stricto* and foraminifer assemblages were combined to produce a robust sea surface temperature (SST) record [Caley et al., 2018]. High resolution (0.5 cm) XRF-scanning has been performed over the total core length, of which the iron-calcium ratios, $\ln(\text{Fe}/\text{Ca})$, were used to estimate fluvial terrestrial input variability [Caley et al., 2018]. At millennial resolution, higher plant leaf wax (*n*-alkane) concentrations and ratios and compound specific stable carbon isotopes ($\delta^{13}\text{C}_{\text{wax}}$) provided a record of vegetation changes in terms of open versus closed canopy and C4 versus C3 plants of the past 800 ka [Castañeda et al. 2016]. A very low resolution leaf wax deuterium isotopic record was generated [Caley et al. 2018], and in conjunction with other high-resolution proxies including $\ln(\text{Fe}/\text{Ca})$, was used to reconstruct rainfall and Limpopo River runoff during the past 2.0 Ma.

Present-day climate and vegetation

Modern climate is seasonal with the rainy season in summer (November to March). Yearly precipitation ranges from 600 mm in the lowlands to 1400 mm in the mountains, whereby rains are more frequent along the coast under the influence of SSTs [Jury et al. 1993, Reason & Mulenga 1999]. Annual average temperatures range from 24 to 16°C but in the highlands clear winter nights may be frosty.

The modern vegetation of this area belongs to the forest, Highveld grassland, and savannah biomes and also includes azonal vegetation (Figure 1) [Dupont et al. 2011 and references therein]. The natural potential vegetation of the coastal belt is forest, although at present it is almost gone; north of the Limpopo River mouth rain forests belong to the Inhambane phytogeographical mosaic and south of the Limpopo River the forest belongs to the Tongaland-Pondoland regional mosaic [White 1983]. The vegetation of the northern part of the Tongaland-Pondoland region is the Northern Coastal Forest [Mucina & Rutherford 2006]. Semi-deciduous forest is found in the Lebombo hills [Kersberg 1996]. Afromontane forest and Highveld grasslands grow along the escarpment and on the mountains. The savannahs of the Zambebian phytogeographical region including e.g. the miombo dry forest occur further inland [White 1983]. Azonal vegetation consists of freshwater swamps, alluvial, and seashore vegetation and mangroves [Mucina & Rutherford 2006].

Material and Methods

Pollen analysis of the 37.59 m long core MD96-2048 (26°10'S 34°01'E, 660m water depth) was extended with 65 samples down core to 12 m (790 ka). Average sampling distance for the Brunhes part was 7 cm reaching an average temporal resolution of 4 ka according to the age model based on

the stable oxygen isotope stratigraphy of benthic foraminifers [Caley et al. 2011, 2018]. Two older windows have been sampled; 20 samples between 15 and 26 m (943-1537 ka) and 19 samples between 30 and 36 m (1785-2143 ka) were taken with an average resolution of 31 and 20 ka, respectively.

Pollen preparation has been described in [Dupont et al. 2011]. In summary, samples were decalcified with HCl (~10%), treated with HF (~40%) for two days, ultrasonically sieved over an 8- μ m screen and, if necessary, decanted. The samples were spiked with two *Lycopodium* spore tablets (either of batch #938934 or batch #177745). Residues were mounted in glycerol and pollen and spores examined at 400x or 1000x magnification. Percentages are expressed based on the total of pollen and spores ranging from over 400 to 60 – only in six samples this sum amounts to less than 100. Confidence intervals (95%) were calculated after Maher [1972, 1981]. Pollen have been identified using the reference collection of African pollen grains of the Department of Palynology and Climate Dynamics




Figure 2. Indicators of C4 vegetation and terrestrial input. From top to bottom: elemental Fe/Ca ratios [Caley et al. 2018]; less negative values indicate relatively wetter conditions, $\delta^{13}\text{C}_{\text{C}_{31}}$ of the *n*-alkane C_{31} [Castañeda et al. 2016, Caley et al. 2018]; less negative values of around -24‰ indicate more C4 inputs while more negative values of around -28‰ indicate more C3 inputs, Cyperaceae (sedges) pollen percentages [Dupont et al., 2011, this study], Poaceae (grass) pollen percentages [Dupont et al., 2011, this study]. Shaded areas denote 95% confidence intervals after [Maher 1972]. Stars denote corresponding maxima in Cyperaceae pollen percentages and the stable carbon isotopes indicating C4 vegetation. VPDB: Vienna Pee Dee Belemnite. Note the scale break.

of the University of Göttingen, the African Pollen Database collection, and literature [e.g. Bonnefille & Rioulet 1980, Scott 1982, Köhler & Brückner 1982, 1989, Schüler & Hemp 2016].

We assigned pollen taxa to groups such as riparian, woodland, forest, etc. (Supplementary Table 1) using information given by Scott [1982], White [1983], Beentje [1994], Kersberg [1996], Coates-Palgrave [2002], Vincens et al. [2007]. Additionally, we carried out a multivariate analysis in the form of an endmember model unmixing procedure [Weltje, 1997], the statistics of which are specifically designed for the treatment of percentage data. We regard the pollen percentages as a series of pollen assemblage mixtures, whereby each modelled endmember may be interpreted as ~~a~~the

~~representation of one or more biomes~~ characteristic combination. This linear mixing model can be compared to a ternary diagram but allowing for more than three axes. We use a version of the unmixer algorithm programmed in MATLAB by Dave Heslop in 2008. Taxa occurring in 6 or more samples (listed in Supplementary Table 2) were used in the endmember modelling (148 of 231 taxa in 220 samples). We selected a model with four components explaining more than 95% of the variance ($r^2 = 0.953$). Iteration was stopped at 1000x resulting in a convexity at termination of -1.6881. Significance level at 99% for taxa to score on the assemblages was 0.018.

To study the correlations between different parameters, we used a linear regression model (least square regression) on linearly interpolated values (5 ka steps) from 0 to 790 ka. Correlation coefficients are given in Table 1. For interpolation and testing the correlation, we used the package PAST [Hammer et al. 2001].

Result and Discussion

Terrestrial input and provenance of the C4 plant wax

Pollen percentages of Cyperaceae (sedges) and Poaceae (grasses) are plotted in Figure 2 together with the $\delta^{13}\text{C}_{\text{wax}}$ of the C_{31} n -alkane and XRF-scanning data, $\ln(\text{Fe}/\text{Ca})$, the natural logarithm of elemental ratios of iron over calcium. Comparing the records of Cyperaceae and $\delta^{13}\text{C}_{\text{wax}}$ reveals that high relative amounts of C4 plant material co-varied with increased representation of sedges. They also co-varied with higher terrestrial input indicated by $\ln(\text{Fe}/\text{Ca})$, and increased precipitation as suggested by deuterium of the C_{31} n -alkane [Caley et al., 2018]. We substantiated the correlations for the Brunhes Chron between pollen percentages, leaf waxes and elemental ratios in Table 1. Leaf wax data are after Castañeda et al. [2016] including Average Chain Length (ACL) of the C_{27} - C_{33} n -alkanes, the ratio of $\text{C}_{31}/(\text{C}_{31}+\text{C}_{29})$ and $\delta^{13}\text{C}_{\text{wax}}$. XRF $\ln(\text{Fe}/\text{Ca})$ ratios are from Caley et al. [2018]. Significant correlation is found between the leaf wax parameters and Cyperaceae data but not between $\delta^{13}\text{C}_{\text{wax}}$ (indicative of C4 inputs) and Poaceae pollen percentages - although a correlation exists between Cyperaceae and Poaceae percentages. While the sedge pollen percentages fluctuate between 10 and 50% (mostly > 20%), the percentages of grass pollen are always lower than 20%. Such low grass pollen values have not been found adjacent C4 grass dominated biomes (mainly savannahs) on the western side of the continent [Dupont 2011]. It is, therefore, likely that in sediments of MD96-2048 the C4 component of the plant wax originated from C4 sedges rather than from C4 grasses.

South Africa has 68 species of Cyperaceae (sedges) of which 28 use the C4 pathway (among the 10 *Cyperus* species 8 are C4) predominantly growing in the northern part of the country [Stock et al. 2004]. They are an important constituent of tropical swamps and riversides [Chapman et al. 2001].

Table 1. Correlation coefficients calculated with PAST [Hammer et al 2001]. Significant correlations are underlined (95%) or bold and underlined (99%). Average Chain Length (ACL), ratio of concentrations of $\text{C}_{31}/(\text{C}_{31} + \text{C}_{29})$, and stable carbon isotope composition of the C_{31} n -alkane ($\delta^{13}\text{C}_{\text{wax}}$) after Castañeda et al. [2016]. Cyperaceae and Poaceae pollen percentages (of total pollen and spores) after Dupont et al. [2011] and this study. $\ln(\text{Fe}/\text{Ca})$ data after Caley et al. [2018].

		Ratio	$\delta^{13}\text{C}_{\text{wax}}$	Cyperaceae	Poaceae	XRF
r^2	ACL	$\text{C}_{31}/(\text{C}_{29}+\text{C}_{31})$	(‰)	(%)	(%)	$\ln(\text{Fe}/\text{Ca})$
ACL	1					
Ratio $\text{C}_{31}/(\text{C}_{31}+\text{C}_{29})$	<u>0.635</u>	1				
$\delta^{13}\text{C}_{\text{wax}}$	<u>0.079</u>	<u>0.180</u>	1			
Cyperaceae (%)	0.027	<u>0.140</u>	<u>0.142</u>	1		
Poaceae (%)	0.003	0.016	0.003	<u>0.165</u>	1	
XRF $\ln(\text{Fe}/\text{Ca})$	0.016	<u>0.032</u>	<u>0.227</u>	<u>0.110</u>	0.011	1

1 An inventory of six modern wetlands between 500 and 1900m in KwaZulu Natal shows that C4
2 grasses dominate the dry surroundings of the wetlands at all altitudes [Kotze & O'Connor 2000]. In
3 the wet parts of the wetlands, however, C4 sedges may make up to 60% of the vegetation cover at
4 550 m. At higher altitudes the coverage of C4 sedges declines [Kotze & O'Connor 2000].

5 Cyperaceae pollen concentration (Figure 3) and percentages correlate with $\ln(\text{Fe}/\text{Ca})$ and with
6 $\delta^{13}\text{C}_{\text{wax}}$ (Table 1, Figure 2). The ratios of terrestrial iron over marine calcium can be interpreted as a
7 measure for terrestrial input, which in this part of the ocean is mainly fluvial. Correlation between
8 increased fluvial discharge and increased C4 vegetation as well as increased Cyperaceae pollen has
9 been reported from sediments off the Zambezi [Schefuß et al. 2011, Dupont & Kuhlmann 2017].
10 Moreover, a fingerprint of C4 sedges was found in Lake Tanganyika [Ivory & Russel 2016]. As a
11 consequence, material (leaf waxes and pollen) from the riverine vegetation is probably better
12 represented than that from dry and upland vegetation. These results corroborate the
13 reinterpretation of the $\delta^{13}\text{C}_{\text{wax}}$ record, in which the increased representation of C4 plants (*n*-alkanes
14 enriched in ^{13}C) is instead attributed to stronger transport of material from the upper Limpopo
15 catchment and the extension of swamps containing C4 sedges under more humid conditions [Caley
16 et al., 2018]. Previously Castañeda et al. [2016] had interpreted increased C4 inputs as reflecting
17 increased aridity.

18 Relatively low values of Cyperaceae pollen and Fe/Ca ratios are found for most interglacials of the
19 Brunhes Chron (Figures 2 and 3), which could be interpreted as an effect of sea-level high-stands.
20 However, Caley et al. [2018] demonstrated that the fluvial discharge is not related to sea-level
21 changes. From the bathymetry of Maputo Bay, strong influence of sea-level is also not expected
22 because the shelf is not broad and the locality of Core MD96-2048 is relatively remote on the
23 Limpopo cone in the center of the clockwise flowing Maputo Bight Lee Eddy. The eddy transports
24 terrestrial material northeastwards before it is taken southwards (Figure 1) [Martin 1981] and likely
25 has not changed direction during glacial times. Thus, fluvial discharge was probably low during
26 interglacials (among other periods), which might be the combined result of more evapotranspiration
27 and less precipitation. Despite drier conditions, the representation of woodland and dry forest is
28 relatively high during the interglacial periods (Figure 3, see also next section).

29 Endmembers representing vegetation on land

30 Palynological results have been published for the past 350 ka [Dupont et al. 2011] providing a
31 detailed vegetation record for the past three climate cycles. Pollen and spore assemblages could be
32 characterized initially by three endmembers via endmember modelling (EM1, EM2, EM3). The
33 assemblage of EM1 was dominated by *Podocarpus* (yellow wood) pollen percentages being more
34 abundant during the non-interglacial parts of MIS 5, 7, and 9. EM2 was characterized by pollen
35 percentages of Cyperaceae (sedges), Ericaceae (heather) and other plants of open vegetation and
36 abundant during full glacial stages. EM3 constituted of pollen from woodland, forest, and coastal
37 vegetation and was interpreted to represent a mix of several vegetation complexes.

We repeated the endmember modelling for the extended record covering the entire Brunhes Chron and the two early Pleistocene windows. With The analysis of the extended dataset gave compatible results with the previous analysis (Dupont et al. 2011). The main difference is that the longer sequence allowed to distinguish two assemblages of interglacial vegetation. In terms of analysis, the cumulative increase of explanatory power lessened after four (instead of three) endmembers and. Therefore, a model with four endmembers was chosen. We used the scores of the different pollen taxa on the endmember assemblages for our interpretation of the endmembers (list of taxa and scores in Supplementary Tables 1 and 2). This interpretation is summarized in Table 2. We compare the new results with the previous analysis. To distinguish between the previous and current analysis (which show strong similarities), we have given new names to the endmember assemblages reflecting our interpretation: **E-heathland, E-Mountain-Forest, E-Shrubland, E-Woodland**. A selection of pollen percentage curves are plotted together with each endmember's fractional abundance in Supplementary Figures 1-4.




Figure 3. Summary of results of MD96-2048. Bottom to top: Pollen and spore count used to calculate percentages; Fractional abundance of endmembers E-Shrubland, E-Woodland, E-Mountain-Forest, and E-Heathland; Pollen summary diagram (woodland and forest taxa are listed in Supplementary Table 1; Cyperaceae pollen concentration per ml (shading denoted 95% confidence intervals after Maher [1981]; ln(Fe/Ca); global stack of stable oxygen isotopes of benthic foraminifers, LR04 [Liesiecki & Raymo 2005]; $\delta^{18}\text{O}_{\text{benthic}}$ of Core MD96-2048 [Caley et al. 2018]; Interglacial peaks after PAGES [2016]. VPDB: Vienna Pee Dee Belemnite.

E-Heathland. Of the ~~new~~ four endmember assemblages (Figure 3), one endmember had a counterpart in EM2 [Dupont et al. 2011] of the previous analysis. Not only composition but also the fractional abundances, which were high during glacial stages, are very much alike. We name this endmember 'E-Heathland', which is dominated by Cyperaceae (sedges) pollen percentages followed

by Ericaceae (heather) pollen and hornwort (Anthocerotaceae) spores (Table 2). Also *Lycopodium* (clubmoss) spore, Restionaceae and *Stoebe*-type pollen percentages score highest on this endmember. The E-Heathland assemblage represents a Fynbos-like open vegetation growing during full glacials. Other pollen records from SE Africa also indicate an open ericaceous vegetation with sedges and Restionaceae during glacial times [Scott 1999, Dupont & Kuhlmann 2017]. The record of MD96-2048 testifies that this type of open glacial vegetation regularly occurred since at least two million years.

E-Mountain-Forest. Like the endmember EM1 [Dupont et al. 2011] of the previous analysis, one endmember is dominated by *Podocarpus* (yellow wood) pollen percentages (Table 2). The assemblage is enriched by pollen of *Celtis* (hackberries) and *Olea* (olive trees) accompanied by undifferentiated fern spores. The interpretation as an assemblage representing mountain forest is rather straightforward and we name the assemblage ‘E-Mountain-Forest’. The fractional abundance of the E-Mountain-Forest is also high in glacials of the Brunhes Chron but not during the extreme glacial stages, when temperatures and pCO_2 are particularly low (Figure 3). It is low in the early - Pleistocene parts of the record.

E-Shrubland. The remaining endmember assemblages have no direct counterpart in the previous analysis, although summed together the pattern of fractional abundance is similar to that of EM3 [Dupont et al. 2011]. One endmember groups together 44 pollen taxa, mostly from coastal and dune vegetation, which we name ‘E-Shrubland’. It includes pollen of Asteraceae and Poaceae (grasses). The latter are not very specific as grass pollen values score almost as high on other endmember assemblages (E-Heathland and E-Woodland). Several taxa scoring on this endmember are known from coastal or halophytic settings such as *Gazania*-type, Amaranthaceae, *Tribulus*, Acanthaceae and *Euphorbia*-type. Arboreal taxa in this assemblage are *Dombeya*, *Acacia*, Meliaceae/Sapotaceae (Table 2). The most typical taxa are the *Buxus* species. We distinguished three types of *Buxus* pollen: *B. macowani* type, *B. hildebrandtii* type and *B. cf. madagascariensis* [Köhler & Brückner 1982, 1989]. *B. madagascariensis* grows on Madagascar and its pollen is only found sporadically, while the other two species inhabit bushland and forest on coastal dunes of the East African main-land. *B. hildebrandtii* nowadays is found in Somalia and Ethiopia and *B. macowani* is native in South Africa. The record of MD96-2048 indicates that these *Buxus* species were more common in the early Pleistocene than during the Brunhes Chron (Figure 3).

Table 2. Interpretation of the endmembers

Endmember	Main pollen taxa
E-Heathland	<i>Podocarpus</i> , <i>Celtis</i> , <i>Olea</i>
E-Mountain-Forest	Cyperaceae, Ericaceae, <i>Phaeoceros</i> , Restionaceae, <i>Stoebe</i> type, <i>Anthoceros</i> , <i>Typha</i> <i>Lycopodium</i> , Restionaceae
E-Shrubland	Poaceae, Asteroideae, <i>Buxus</i> , Amaranthaceae, <i>Euphorbia</i> , Meliaceae-Sapotaceae, <i>Acacia</i> , <i>Riccia</i> type, <i>Tribulus</i> , Acanthaceae pp, Asteraceae Vernoniae, <i>Hypoestes-Dicliptera</i> type, <i>Gazania</i> type, <i>Dombeya</i>
E-Woodland	<i>Alchornea</i> , <i>Spirostachys africana</i> , <i>Pteridium</i> type, Polypodiaceae, <i>Myrsine africana</i> , <i>Cassia</i> type, Rhizophoraceae, Aizoaceae, Combretaceae pp, <i>Manilkara</i> , <i>Burkea africana</i> , <i>Brachystegia</i> , <i>Dodonaea viscosa</i> , <i>Pseudolachnostylis</i> , <i>Hymenocardia</i> , <i>Aloe</i> , Rhamnaceae pp, <i>Protea</i> , <i>Parinari</i>

E-Woodland. The last endmember, which we name ‘E-Woodland’, groups together 39 pollen taxa from forest and woodland species with maximum values of less than 5 or 2% of the total of pollen and spores. To this assemblage belong *Pseudolachnostylis*, *Dodonaea viscosa* and *Manilkara*, which are woodland trees. *Protea* (sugarbush) and *Myrsine africana* (Cape myrtle) grow more upland and *Alchornea* is a pioneer forest tree often growing along rivers. Others include wide-range woodland taxa such as Combretaceae species. The occurrence of pollen of *Brachystegia* (miombo tree), *Burkea*

africana, *Spirostachys africana* and *Hymenocardia* in this assemblage is indicative of Miombo dry forest and woodland. The assemblage additionally includes Rhizophoraceae pollen from the coastal mangrove forest (Table 2). The fractional abundance of the E-Woodland assemblage is low during the early Pleistocene, increased during the interglacials prior to the MBE and had maximum values during Interglacials 9e, 5e, and 1 (Figures 3, 4). These interglacials also exhibited maximum percentages of arboreal pollen excluding *Podocarpus*.

In summary, the endmember analysis indicates a very stable open ericaceous vegetation with partially wet elements such as sedges and Restionaceae characterizing the landscape of full glacials (when global temperatures and $p\text{CO}_2$ were lowest). During the less extreme parts of the glacials, mountain *Podocarpus* forest was extensive as in most mountains of Africa [Dupont 2011, Ivory et al. 2018]. On the other hand, interglacials were characterized by coastal shrubs. In the course of the Brunhes, the woody component, which was relatively weak before the MBE, became more and more important reflecting the same long-term trend found in the leaf wax records [Castañeda et al. 2016]. It is likely that the Miombo dry forest and woodland migrated into the region in the successive interglacials of the Brunhes Chron. Particularly during Interglacials 9e and 1 the area might have been more forested than during the older interglacials of the Brunhes Chron.

Long-term trends in vegetation and climate of East Africa

The region of the Limpopo River becoming more and more wooded in the course of successive interglacials [Castañeda et al. 2016] somewhat paralleled the conditions around Lake Malawi [Johnson et al. 2016]. However, around Lake Malawi, forested phases of either mountain forest, seasonal forest, or Miombo woodland alternating with savannahs occurred during both glacial and interglacial stages [Ivory et al. 2018]. Also in contrast to the Lake Malawi record, the MD96-2048 Poaceae pollen percentages fluctuated little and remained relatively low (less than 20%) indicating that savannahs were of less importance in the Limpopo catchment area and the coastal region of southern Mozambique.

The trend to increased woodland in SE Africa after the MBE, noted at both Lake Malawi and in the Limpopo River catchment [Johnson et al. 2016, Caley et al. 2018, this study] contrasts with the trend around Lake Magadi at the equator. At Lake Magadi a trend to less forest around marks the Mid-Brunhes transition [Owen et al. 2018]. Antiphase behavior of SE African climate with that of West and East Africa emphasizes the importance of the average position of the tropical rainbelt shifting southwards during globally cold periods as has been inferred from Holocene to Last Glacial records of Lake Malawi [Johnson et al. 2002, Scholz et al. 2011]. Our results confirm this relationship existed over the entire Brunhes Chron.

The Lake Malawi pollen record as well as that of the equatorial Lake Magadi in Kenya [Owen et al. 2018] do not show much of a glacial-interglacial rhythm and are dominated by the precession variability in tropical rainfall [cf. Clement et al. 2004]. Obviously, in the tropical climate of the Southern Hemisphere north of $\sim 15^\circ\text{S}$ the hydrological regime had more effect on the vegetation than changes in temperature, while further south the impact of glacial-interglacial variability on the vegetation increased.

Effects of atmospheric $p\text{CO}_2$

While the hydroclimate of the region shows precession variability [Caley et al. 2018], the vegetation shows a glacial-interglacial rhythm (Supplementary Information) indicating that besides hydrology, temperature and/or atmospheric CO_2 levels were important drivers of the vegetation development. Combining the results of the pollen assemblages with stable carbon isotopes and elemental information indicates that during interglacials the region of SE Africa (northern South Africa,

Zimbabwe, southern Mozambique) was less humid. This is in accordance with other paleoclimate estimates for the region [see reviews by [Simon et al. 2015](#), [Singarayer & Burrough 2015](#)].

The interglacial woodlands (represented by E-Woodland, Figure 4) would probably have grown under warmer and drier conditions than the glacial mountain forest (represented by E-Mountain-Forest). The increase in maximum $p\text{CO}_2$ levels during the post-MBE interglacials might have favored tree growth as higher $p\text{CO}_2$ levels would have allowed decreased stomatal conductivity and thus relieved drought stress [[Jolly & Haxeltine 1997](#)]. ~~Woodlands would have expanded at the cost of mountain forest during 11c, 9e, 5e and 1, and to a lesser extend during 7e and 7c. During Interglacials 11c, 9e, 7c, 7e, 5e, and 1, when temperatures and $p\text{CO}_2$ are were high, the mountain forest is replaced by woodland~~ (Figure 4). It might be only after interglacial $p\text{CO}_2$ levels rose over ~ 270 ppmv that Miombo woodland could fully establish in the area during the warm and relatively dry post-MBE interglacials.

~~The glacial stages showed the expansion of either mountain forest or heathland. The record indicates extension of mountain forests in SE Africa indicates some extension of moist *Podocarpus* forest during those parts of the glacial stages with low temperatures and atmospheric $p\text{CO}_2$ exceeding ~ 220 ppmv (Figure 5). This holds also for the Interglacials 19c, 17c, 15e, 15a, 13a, and 7e, in which $p\text{CO}_2$ and Antarctic temperatures were subdued, too. However, if low temperatures were the only driver of the extension of mountain forests, further spread into the lowlands during the coldest glacial~~




Figure 4: Comparing pollen assemblages E-Mountain-Forest, E-Woodland and E-Shrubland with atmospheric CO_2 [[Bereiter et al. 2015](#), [PAGES 2016](#)]. On top Interglacial peaks of the past 800 ka [[PAGES 2016](#)] and stable oxygen isotopes of benthic foraminifera ($\delta^{18}\text{O}_{\text{benthic}}$) of MD96-2048 [[Caley et al. 2011](#)]. CO_2 -levels of 220 and 270 ppmv are indicated with green-red shading. Grey shading highlight periods with maximum atmospheric CO_2 and maximum values of the sum of E-Woodland and E-Mountain-Forest. VPDB: Vienna Pee Dee Belemnite.

phases should be expected. Instead, when $p\text{CO}_2$ dropped below ~ 220 ppmv during those colder glacial periods, mountain forest declined, in particular during MIS 18, 16, 14, 8, 6, and 2. A picture emerges of cool glacial stages in SE Africa in which tree cover broke down when atmospheric $p\text{CO}_2$

became too low. Additionally, mountain forests were important during the Interglacials 19c, 17c, 15e, 15a, 13a, and 7e, in which $p\text{CO}_2$ and Antarctic temperatures were subdued.

With an inverse modelling technique, Wu et al. [2007] estimated the climate inputs for the vegetation model BIOME4 using as information the biome scores of pollen records from equatorial East African Mountains. Wu et al. found that lowering of the tree line under glacial conditions (1-3°C lower temperatures, less precipitation, 200 ppmv $p\text{CO}_2$) depended hardly on temperature but primarily on increased aridity and somewhat on lower $p\text{CO}_2$, whereby lower $p\text{CO}_2$ amplified the effects of water limitation. However, Izumi & Lézine [2016] found contrasting results using pollen records of mountain sites on both sides of the Congo basin. At any rate, the lack of trees in the Southeast African Mountains during glacial extremes is unlikely the result of drought, because our record indicates that climate conditions in SE Africa were less dry during glacials than during interglacials (the post-MBE interglacials in particular). Instead, C4 sedges being an important constituent of the ericaceous fynbos-like vegetation increased during glacials when atmospheric $p\text{CO}_2$ and temperatures were low (Figure 5). However, low temperatures are not particularly favorable for C4 sedges as indicated by the altitudinal distribution of C4 sedges in modern wetlands of KwaZulu Natal [Kotze & O'Connor 2000]. We presume, therefore, that the extension of C4 sedges during the more humid phases of the glacials is the result of low atmospheric CO_2 concentrations rather than of low temperatures.

Pollen records of ericaceous vegetation suggest an extensive open vegetation existing in the East African Mountains [e.g. Coetzee 1967, Bonnefille & Riollet 1988, Marchant et al. 1997, Debussk 1998, Bonnefille & Chalié 2000] and in SE Africa and Madagascar [e.g. Botha et al. 1992, Scott 1999, Gasse and Van Campo 2001, Scott & Tackeray 1987] during the last glacial. In our study, ericaceous fynbos-like vegetation (E-Heathland) was found for those parts of the glacials having lower (less than ~220 ppmv) atmospheric $p\text{CO}_2$ (Figure 5). Exceptions were found for MIS 12 and 14 when the difference of $p\text{CO}_2$ with that of the preceding stage was small [Bereiter et al. 2015]. Dupont et al. [2011] argued that increase of C4 vegetation as the result of low $p\text{CO}_2$ was unlikely because no extension of grasses was recorded. However, this argument is flawed if sedges dominantly constituted the C4 vegetation in the area. We also note that in many parts of South Africa, no substantial increase of C4 grasses occurred but that many sites suggest an expansion of C3 grasses during the Last Glacial Maximum [Scott 2002].

As climate was wetter during most of the glacials in this part of the world, the question arises about the climatic implication of the ericaceous fynbos-like vegetation (represented by E-Heathland, Figure 5) extending during full glacials over the mountains of South Africa - and correlating with the SST record (see also the correlation between SST and EM2 in Dupont et al. 2011). The correlation with SST, however, is problematic. Singarayer & Burrough [2015] argued that the control of the Indian Ocean SSTs on the precipitation of South Africa shifted from a positive correlation during the interglacial to a negative correlation during the Last Glacial Maximum. They invoked the effects of the exposure of the Sunda Shelf (Indonesia) and Sahul Shelf (Australia) on the Walker circulation causing a wetter region over the western Indian Ocean but also weaker easterly winds to transport moisture inland. To question the link between SST and precipitation in SE Africa even further, Caley et al. [2018] found that the precession signature in the river discharge proxy [$\ln(\text{Fe}/\text{Ca})$, see also Supplementary Information] was absent in the SST record made on the same material from the same core. SE Africa would have been more humid during glacials when the temperature difference between land and sea increased.

The increase in C4 vegetation during relative cool and humid climate would be in conflict with the idea that C4 plants are more competitive in hot and dry climates [Ehleringer et al. 1997, Sage 2004]. However, this idea is mainly based on the ecology of grasses and the development of savannahs, while the C4 vegetation expansion in SE Africa during cool and humid phases seems to be driven by sedges. A survey of the distribution of C4 sedges in South Africa revealed that those Cyperaceae do not have the same temperature constraints as C4 grass species [Stock et al. 2004]. More important, South African C4 sedges appear to have evolved under wetland conditions rather than under aridity. C4 *Cyperus* species even occur in the wettest parts of lower altitude wetlands in KwaZulu-Natal [Kotze & O'Conner 2000].




Figure 5: Comparing Cyperaceae pollen percentages and fractional abundances of the glacial pollen assemblages E-Mountain-Forest and E-Heathland with atmospheric CO₂ [Bereiter et al. 2015, PAGES 2016] and sea surface temperatures of the southeastern Indian Ocean (SST PC1 scores of MD96-20468) [Caley et al. 2018]. On top Terminations of the past 9 glaciations, glacial marine isotope stages (MIS) and stable oxygen isotopes of benthic foraminifera ($\delta^{18}\text{O}_{\text{benthic}}$) of MD96-2048 [Caley et al. 2011]. CO₂-levels of 220 and 270 ppmv are indicated with blue-green-red shading. Grey shading highlight periods with minimum atmospheric CO₂, minimum values of E-Mountain-Forest and Cyperaceae pollen, and maximum values of E-Heathland. VPDB: Vienna Pee Dee Belemnite.

Conclusions

Palynology in combination with sediment chemistry and carbon isotope analysis of leaf waxes carried out on the marine sediments of MD96-2048 retrieved from the Limpopo River cone in the Maputo Bay (SE Africa) allowed a detailed reconstruction of the biome developments over the Brunhes Chron and comparison with earlier Pleistocene vegetation of SE Africa.

Using endmember modelling, we could distinguish four pollen assemblages: E-Heathland, E-Mountain-Forest, E-Shrubland, E-Woodland. The open sedge-rich and ericaceous vegetation

represented by E-Heathland occurred during those parts of the glacials with lower temperatures and atmospheric $p\text{CO}_2$. *Podocarpus*-rich mountain forest represented by E-Mountain-Forest extended during the less extrem parts of the glacials. E-Shrubland represents a shrublike vegetation with coastal elements and *Buxus* species and mainly occurred during the earlier Pleistocene (before 1 Ma). E-Woodland represents interglacial woodlands, Miombo woodland in particular, becoming more and more important in the successive interglacial stages of the Brunhes Chron and dominated the post-MBE interglacials.

Our results indicate the influence of atmospheric $p\text{CO}_2$ fluctuations on the shaping of the biomes in SE Africa during the Brunhes Chron. We argue that (1) the precessional rhythms of river discharge compared to the interglacial-glacial biome variability indicates that hydroclimate cannot be the only driver of vegetation change. The other options of forcing mechanisms on interglacial-glacial time-scales are temperature and $p\text{CO}_2$. (2) Because of the correlation between Cyperaceae pollen percentages and $\delta^{13}\text{C}_{\text{wax}}$ and the lack of correlation between Poaceae percentages and $\delta^{13}\text{C}_{\text{wax}}$ in combination with the relatively low grass pollen percentages, we deduce that the C4 plant imprint mainly derives from the sedges. (3) The expansion of C4 sedges during the colder periods of the glacials is unlikely to result from lower temperatures. Thus, during the colder phases of the glacials, low atmospheric $p\text{CO}_2$ might have favored the expansion of C4 sedges. (4) The confinement of mountain forest to the glacial periods with moderate temperatures and moderate $p\text{CO}_2$, and the lack of extension into the lowlands of mountain forest during the colder periods, suggests that low $p\text{CO}_2$ became restrictive to the forest. Mountain forests could thrive during glacials as long as $p\text{CO}_2$ levels exceeded ~ 220 ppmv. (5) Based on the elemental composition as a proxy for river discharge [$\ln(\text{Fe}/\text{Ca})$], we estimate the post-MBE interglacials as the drier intervals of the sequence. Nevertheless woodland extended during those periods, which we attribute to increased temperatures and $p\text{CO}_2$. Atmospheric $p\text{CO}_2$ levels over 250 ppmv might have been a prerequisite for the establishment of the Miombo woodlands in SE Africa, which extended during the post MBE interglacials.

The vegetation record of the Limpopo catchement area shows a greater impact of glacial-interglacial variability, mainly driven by CO_2 fluctuations, and less influence of hydroclimate compared to the more equatorial records of Lake Malawi and Lake Magadi. The long-term trend of increased woodiness in the course of the Brunhes Chron paralleled that of Lake Malawi but contrasted Lake Magadi suggesting a long-term southward shift in the average position of the tropical rainbelt.

Acknowledgements

This project was funded through DFG -Research Center / Cluster of Excellence „The Ocean in the Earth System“. T.C. is supported by CNRS-INSU. Funding from LEFE IMAGO CNRS INSU project SeaSalt is acknowledged. David Heslop is thanked for providing the Endmember model into a Matlab application.

Data availability

Pollen counts are available at <https://doi.pangaea.de/10.1594/PANGAEA.897922>.

Previously published data can be retrieved at <https://doi.pangaea.de/10.1594/PANGAEA.895364>;
<https://doi.pangaea.de/10.1594/PANGAEA.895361>;
<https://doi.pangaea.de/10.1594/PANGAEA.895362>;
<https://doi.pangaea.de/10.1594/PANGAEA.863919>;
<https://doi.pangaea.de/10.1594/PANGAEA.895357>.

Author contributions

LMD carried out the palynological analysis, conceived and wrote the manuscript, TC carried out the sedimentology and stratigraphy and contributed to the discussion, ISC conducted the stable isotope analysis on plant waxes and contributed to the discussion.

References

- Barth, A.M., Clark, P.U., Bill, N.S., He, F. & Pisias, N.G., 2018. Climate evolution across the Mid-Brunhes Transition. *Climate of the Past*, 14: 2071-2087.
- Beentje, H., 1994. Kenya trees, shrubs and lianas. National Museum of Kenya, Nairobi: 722p.
- Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T.F., Fischer, H., Kipfstuhl, S. & Chappelaz, J., 2015. Revision of the EPICA Dome C CO₂ record from 800 to 600kyr before present. *Geophysical Research Letters*, 42: 542-549.
- Bereiter, B., Shackleton, S., Baggenstos, D., Kawamura, K. & Severinghaus, J., 2018. Mean global ocean temperatures during the last glacial transition. *Nature*, 553: 39-44.
- Bonnefille, R. & Chalié, F., 2000. Pollen-inferred precipitation time-series from equatorial mountains, Africa, the last 40 kyr BP. *Global and Planetary Change*, 26: 25-50.
- Bonnefille, R. & Riollet, G., 1980. Pollens des savanes d'Afrique orientale. Éditions du Centre National de la Recherche Scientifique, Paris: 140p., 113pl.
- Bonnefille, R. & Riollet, G., 1988. The Kashiru pollen sequence (Burundi). Palaeoclimatic implications for the last 40,000 yr. B.P. in tropical Africa. *Quaternary Research*, 30: 19-35.
- Botha, G.A., Scott, L., Vogel, J.C. & Von Brunn, V., 1992. Palaeosols and palaeoenvironments during the Late Pleistocene Hypothermal in northern Natal. *South African Journal of Science*, 88: 508-512.
- Bouttes, N., Swingedouw, D., Roch, D.E., Sanchez-Goni, M.F. & Crosta, X., 2018. Response of the carbon cycle in an intermediate complexity model to the different climate configurations of the last nine interglacials. *Climate of the Past*, 14: 239-253.
- Caley, T., Extier, T., Collins, J.A., Schefuß, E., Dupont, L., Malaizé, B., Rossignol, L., Souron, A., McClymont, E.L., Jimenez-Espejo, F.J., García-Comas, C., Eynaud, F., Martinez, P., Roche, D.M., Jorry, S.J., Charlier, K., Wary, M., Gouvers, P.-Y., Billy, I. & Giraudeau, J., 2018. A two-million-year-long hydroclimatic context for hominin evolution in southeastern Africa. *Nature*, 560: 76-79.
- Caley, T., Kim, J.-H., Malaizé, B., Giraudeau, J., Laepple, T., Caillon, N., Charlier, K., Rebaubier, H., Rossignol, L., Castañeda, I.S., Schouten, S. & Sinninge Damsté, J.S., 2011. High-latitude obliquity as a dominant forcing in the Agulhas current system. *Climate of the Past*, 7: 1285-1296.
- Castañeda, I.S., Caley, T., Dupont, L., Kim, J.-H., Malaizé, B. & Schouten, S., 2016. Middle to Late Pleistocene vegetation and climate change insubtropical southern East Africa. *Earth and Planetary Science Letters*, 450: 306-316.
- Chapman, L.J., Balirwa, J., Bugenyi, F.W.B., Chapman, C. & Crisman, T.L., 2001. Wetlands of East Africa: biodiversity, exploitation, and policy perspectives. Gopal, B., Junk, W.J. & Davis, J.A. (Eds.) *Biodiversity in Wetlands: Assessment, Funktion and Conservation*, Volume 2. Backhuys Publishers, Leiden: 101-131.
- Clement, A.C., Hall, A., & Broccoli, A.J., 2004. The importance of precessional signals in the tropical climate. *Climate Dynamics*, 22: 327-341.
- Coates Palgrave, K., 2002. *Trees of Southern Africa*. 3rd edition, revised and updated, Struik, Cape Town: 1212p.
- Coetzee, J.A., 1967. Pollen analytical studies in east and southern Africa. *Palaeoecology of Africa*, 3: 146 p.
- Cowling, S.A. & Sykes, M., 1999. Physiological significance of low atmospheric CO₂ for plant-climate interactions. *Quaternary Research*, 52: 237-242.
- DeBusk, G.H., 1998. A 37,500-year pollen record from Lake Malawi and implications for the biogeography of afro-montane forests. *Journal of Biogeography*, 25: 479-500.
- Dupont, L.M., 2011. Orbital scale vegetation change in Africa. *Quaternary Science Reviews*, 30: 3589-3602.
- Dupont, L.M. & Hooghiemstra, H., 1989. The Saharan-Sahelian boundary during the Brunhes chron. *Acta Botanica Neerlandica*, 38: 405-415.
- Dupont, L.M. & Kuhlmann, H., 2017. Glacial-interglacial vegetation change in the Zambezi catchment. *Quaternary Science Reviews*, 155: 127-135.
- Dupont, L.M., Beug, H.-J., Stalling, H. & Tiedemann, R., 1989. First palynological results from ODP Site 658 at 21°N west off Africa: pollen as climate indicators. In: Ruddiman, W.F., Sarnthein, M. et al. *Proceedings ODP Scientific Results*, 108, College Station TX (Ocean Drilling Program): 93-111.
- Dupont, L.M., Caley, T., Kim, J.-H., Castañeda, I., Malaizé, B. & Giraudeau, J., 2011. Glacial-interglacial vegetation dynamics in south eastern Africa coupled to sea surface temperature variations in the Western Indian Ocean. *Climate of the Past*, 7: 1209-1224.
- Ehleringer, J.R., Cerling, T.E. & Helliker, B.R., 1997. C₄ photosynthesis, atmospheric CO₂, and climate. *Oecologia*, 112: 285-299.


- 1 Foley, J., Kutzbach, J.E., Coe, M.T. & Levis, S., 1994. Feedbacks between climate and boreal forests during the Holocene
2 epoch. *Nature*, 371: 52-54.
- 3 Ganopolski, A. & Calov, R., 2011. The role of orbital forcing, carbon dioxide and regolith in 100 kyr glacial cycles. *Climate of*
4 *the Past*, 7: 1415-1425.
- 5 Gasse, F. & Van Campo, E., 2001. Late Quaternary environmental changes from a pollen and diatom record in the southern
6 tropics (Lake Tritrivakely, Madagascar). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 167: 287-308.
- 7 Hammer, Ø., Harper, D.A.T. & Ryan, P.D., 2001. PAST: Paleontological Statistics Software Package for Education and Data
8 Analysis. *Palaeontologia Electronica*, 4(1): 1-9.
- 9 Ivory, S.J. & Russell, J., 2016. Climate, herbivory, and fire controls on tropical African forest for the last 60ka. *Quaternary*
10 *Science Reviews*, 148: 101-114.
- 11 Ivory, S.J., Lézine, A-M., Vincens, A. & Cohen, A.S., 2018. Waxing and waning of forests: Late Quaternary biogeography of
12 southeast Africa. *Global Change Biology*, 2018: 1-13.
- 13 Izumi, K. & Lézine, A-M., 2016. Pollen-based biome reconstructions over the past 18,000 years and atmospheric CO₂
14 impacts on vegetation in equatorial mountains of Africa. *Quaternary Science Reviews*, 152: 93-103.
- 15 Jansen, J.H.F., Kuijpers, A. & Troelstra, S.R., 1986. A mid-Brunhes climatic event: long-term changes in global atmosphere
16 and ocean circulation. *Science*, 232: 619-622.
- 17 Johnson, T.C., Brown, E.T., McManus, J., Barry, S., Barker, P. & Gasse, F., 2002. A high-resolution paleoclimate record
18 spanning the past 25,000 years in southern East Africa. *Science*, 296: 113-132.
- 19 Johnson, T.C., Werne, J.P., Brown, E.T., Abbott, A., Berke, M., Steinman, B.A., Halbur, J., Conteras, S., Grosshuesch, S.,
20 Deino, A., Lyons, R.P., Scholz, C.A., Schouten, S. & Sinninghe Damsté, J.S., 2016. A progressively wetter climate in
21 southern East Africa over the past 1.3 million years. *Nature*, 537: 220-224.
- 22 Jolly, D. & Haxeltine, A., 1997. Effect of low glacial atmospheric CO₂ on tropical African montane vegetation. *Science*, 276:
23 786-788.
- 24 Jury, M.R., Valentine, H.R. & Lutjeharms, J.R., 1993. Influence of the Agulhas Current on summer rainfall along the
25 southeast coast of South Africa. *Journal of Applied Meteorology*, 32: 1282-1287.
- 26 Kersberg, H., 1996. Beiheft zu Afrika-Kartenwerk Serie S: Südafrika (Moçambique, Swaziland, Republik Südafrika). Bl. 7:
27 Vegetationsgeographie. Gebrüder Bornträger, Berlin: 182p.
- 28 Köhler, E. & Brückner, P., 1982. De Pollenmorphologie der afrikanischen Buxus- und Notobuxus-Arten (Buxaceae) und ihre
29 systemstische Bedeutung. *Grana*, 21: 71-82.
- 30 Köhler, E. & Brückner, P., 1989. The genus Buxus (Buxaceae): aspects of its differentiation in space and time. *Plant*
31 *Systematics and Evolution*, 162: 267-283.
- 32 Kotze, D. & O'Connor, T.G., 2000. Vegetation variation within and among palustrine wetlands along an altitudinal gradient
33 in KwaZulu-Natal, South Africa. *Plant Ecology*, 146: 77-96.
- 34 Lisiecki, L.E. & Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records.
35 *Paleoceanography*, 20, PA1003: 1-17.
- 36 Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura,
37 K. & Stocker, T.F., 2008. High-resolution carbon dioxide concentration record 650,000–800,000 years before present.
38 *Nature*, 453: 379-382.
- 39 Lyu, A., Lu, H., Zeng, L., Zhang, H., Zhang, E., & Yi, S., 2018. Vegetation variation of loess deposits in the southeastern Inner
40 Mongolia, NE China over the past ~1.08 million years. *Journal of Asian Earth Sciences*, 155: 174-179.
- 41 Maher, L.J., 1972. Nomograms for computing 0.95 confidence limits of pollen data. *Review of Palaeobotany and Palynology*,
42 13: 85-93.
- 43 Maher, L.J., 1981. Statistics for microfossil concentration measurements employing samples spiked with marker grains.
44 *Review of Palaeobotany and Palynology*, 32: 153-191.
- 45 Marchant, R., Taylor, D. & Hamilton, A., 1997. Late Pleistocene and Holocene history at Mubwindi Swamp, Southwest
46 Uganda. *Quaternary Research*, 47: 316-328.
- 47 Martin, A.K., 1981. The influence of the Agulhas Current on the physiographic development of the northernmost Natal
48 Valley (S.W. Indian Ocean). *Marine Geology*, 39: 259-276.
- 49 Miller, S.M. & Gosling, W.D., 2014. Quaternary forest associations in lowland tropical West Africa. *Quaternary Science*
50 *Reviews*, 84: 7-25.
- 51 Mucina, L. & Rutherford, M.C., 2006. The vegetation of South Africa, Lesotho and Swaziland. *Strelitzia*, 19. South African
52 National Biodiversity Institute, Pretoria: 807p.
- 53 Mudelsee, M. & Stattegger, K., 1997. Exploring the structure of the mid-Pleistocene revolution with advanced methods of
54 time-series analysis. *Geologische Rundschau*, 86: 499-511.
- 55 Owen, R.B., Muiruri, V.M., Lowenstein, T.K., Renaut, R.W., Rabideaux, N., Luo, S., Deino, A.L., Sier, M.J., Dupont-Nivet, G.,
56 McNulty, E.P., Leet, K., Cohen, A., Campisano, C., Deocampo, D., Shen, C-C., Billingsley, A. & Mbuthia, A., 2018.
57 Progressive aridification in East Africa over the last half million years and implications for human evolution. *PNAS*, 115,
58 44: 11174-11179.
- 59 Past Interglacials Working Group of Pages, 2016. Interglacials of the last 800,000 years. *Review of Geophysics*, 54: 162-219.

- 1 Paillard, D., 2017. The Plio-Pleistocene climatic evolution as a consequence of orbital forcing on the carbon cycle. *Climate of*
- 2 *the Past*, 13: 1259-1267.
- 3 Prentice, I.C. & Harrison, S.P., 2009. Ecosystem effects of CO₂ concentration: evidence from past climates. *Climate of the*
- 4 *Past*, 5: 297-307.
- 5 Prentice, I.C., Cleator, S.F., Huang, Y.H., Harrison, S.P. & Roulstone, I., 2017. Reconstructing ice-age palaeoclimates:
- 6 Quantifying low-CO₂ effects on plants. *Global and Planetary Change*, 149: 166-176.
- 7 Reason, C.J.C. & Mulenga, H., 1999. Relationships between South African rainfall and SST anomalies in the Southwest Indian
- 8 Ocean. *International Journal of Climatology*, 19: 1651-1673.
- 9 Sage, J.P., 2004. The evolution of C4 Photosynthesis. *New Phytologist*, 161: 341-370.
- 10 Schefuß, E., Kuhlmann, H., Mollenhauer, G., Prange, M. & Pätzold, J., 2011. Forcing of wet phases in southeast Africa over
- 11 the past 17,000 years. *Nature*, 480: 509-512.
- 12 Scholz, C.A., Cohen, A.S., Johnson, T.C., King, J., Talbat, M.R. & Brown, E.T., 2011. Scientific drilling in the Great Rift Valley:
- 13 The 2005 Lake Malawi Scientific Drilling Project - An overview of the past 145,000 years of climate variability in
- 14 Southern Hemisphere East Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 303: 3-19.
- 15 Schüller, L. & Hemp, A., 2016. Atlas of pollen and spores and their parent taxa of Mt Kilimanjaro and tropical East Africa.
- 16 *Quaternary International*, 425: 301-386.
- 17 Scott, L., 1982. Late Quaternary fossil pollen grains from the Transvaal, South Africa. *Review of Palaeobotany and*
- 18 *Palynology*, 36: 241-278.
- 19 Scott, L., 1999. Vegetation history and climate in the Savanna biome South Africa since 190,000 ka: a comparison of pollen
- 20 data from the Tswaing Crater (the Pretoria Saltpan) and Wonderkrater. *Quaternary International*, 57-58: 215-223.
- 21 Scott, L. & Thackeray, J.F., 1987. Multivariate analysis of late Pleistocene and Holocene pollen spectra from Wonderkrater,
- 22 Transvaal, South Africa *South African Journal of Science*, 83: 93-98.
- 23 Simon, M.H., Ziegler, M., Bosmans, J., Barker, S., Reason, C.J.C. & Hall, I.R., 2015. Eastern South African hydroclimate over
- 24 the past 270,000 years. *Scientific Reports*, 5, 18153: 1-10.
- 25 Singarayer, J.S. & Burrough, S.L., 2015. Interhemispheric dynamics of the African rainbelt during the late Quaternary.
- 26 *Quaternary Science Reviews*, 124: 48-67.
- 27 Stock, W.D., Chuba, D.K. & Verboom, G.A., 2004. Distribution of South African C3 and C4 species of Cyperaceae in relation
- 28 to climate and phylogeny. *Austral Ecology*, 29: 313-319.
- 29 Sun, Y., Yin, Q., Crucifix, M., Clemens, S.C., Araya-Melo, P., Liu, W., Qiang, X., Liu, Q., Zhao, H., Liang, L., Chen, H., Li, Y.,
- 30 Zhang, L., Dong, G., Li, M., Zhou, W., Berger, A. & An, Z., 2019. Diverse manifestations of the mid-Pleistocene climate
- 31 transition. *Nature Communications*, 10:351. doi:10.1038/s41467-018-08257-9
- 32 Swann, A.L., Fung, I.Y., Levis, S., Bonan, G.B. & Doney, S.C., 2010. Changes in Arctic vegetation amplify high-latitude
- 33 warming through the greenhouse effect. *PNAS*, 107: 1295-1300.
- 34 Torres, V., Hooghiemstra, H., Lourens, L. & Tzedakis, P.C., 2013. Astronomical tuning of long pollen records reveals the
- 35 dynamic history of montane biomes and lake levels in the tropical high Andes during the Quaternary. *Quaternary*
- 36 *Science Reviews*, 63: 59-72.
- 37 Tyson, P.D. & Preston-Whyte, R.A., 2000. *The weather and climate of Southern Africa*. Oxford University Press, Cape Town:
- 38 396p.
- 39 Tzedakis, P.C., Hooghiemstra, H. & Pälike, H., 2006. The last 1.35 million years at Tenaghi Philippon: revised
- 40 chronostratigraphy and long-term vegetation trends. *Quaternary Science Reviews*, 25: 3416-3430.
- 41 Tzedakis, P.C., Raynaud, D., Mcmanus, J.F., Berger, A., Brovkin, V. & Kiefer, T., 2009. Interglacial diversity. *Nature*
- 42 *Geoscience*, 2: 753-755.
- 43 Vincens, A., Lézine, A-M., Buchet, G., Lewden, D., Le Thomas, A. & Contributors., 2007. African pollen database inventory of
- 44 tree and shrub pollen types. *Review of Palaeobotany and Palynology*, 145: 135-141.
- 45 Weltje, G.J., 1997. End-member modeling of compositional data: numerical-statistical algorithms for solving the explicit
- 46 mixing problem. *Mathematical Geology*, 29: 503-549.
- 47 White, F., 1983. *The vegetation of Africa*. Natural Resources Research, 20. UNESCO: 356p., 3maps.
- 48 Wu, H., Guiot, J., Brewer, S. & Guo, Z., 2007. Climatic changes in Eurasia and Africa at the last glacial maximum and mid-
- 49 Holocene: reconstruction from pollen data using inverse vegetation modelling. *Climate Dynamics*, 29: 211-229.
- 50 Yin, Q.Z., 2013. Insolation-induced mid-Brunhes transition in Southern Ocean ventilation and deep-ocean temperature.
- 51 *Nature*, 494: 222-225.
- 52 Yin, Q.Z. & Berger, A., 2010. Insolation and CO₂ contribution to the interglacial climate before and after the Mid-Brunhes
- 53 Event. *Nature Geoscience*, 3: 243-246.
- 54 Yin, Q.Z. & Berger, A., 2012. Individual contribution of insolation and CO₂ to the interglacial climates of the past 800,000
- 55 years. *Climate Dynamics*, 38: 709-724.

Supplement

REDFIT frequency analysis

We conducted a frequency analysis on the data of the $\ln(\text{Fe}/\text{Ca})$ ratios, the E-Heathland fractional abundance scores, and the Cyperaceae pollen concentration covering the Brunhes Chron using the algorithm of REDFIT [Schulz & Mudelsee 2002] from the statistical package PAST version 3.14 (1999-2006) [Hammer et al. 2001]. The E-Heathland and Cyperaceae curves each consisted of 181 data points between 0 and 790 ka. REDFIT was run with 2 times oversampling, a Blackman-Harris window, and 2 overlapping averaging segments resulting in a bandwidth of 0.004291; false alarm level was 99.17. The $\ln(\text{Fe}/\text{Ca})$ curve contained 2307 data points between 1 and 790 ka. REDFIT was run with 2 times oversampling, a Blackman-Harris window, and 3 overlapping averaging segments resulting in a bandwidth of 0.005726; false alarm level was 99.91. The figure shows the power of $\ln(\text{Fe}/\text{Ca})$ ratios (left), the power of the E-Heathland values (middle), and the power of the Cyperaceae pollen concentration (right) against frequency running from 0 - 0.08 cycles per ka. Denoted are the bandwidth for each spectrum and a parametric approximation of the level above the null hypothesis of a red noise model using χ^2 -test at 90% (dashed lines). Grey bars indicate the orbital periodicities of 100, 41, 23, and 19 ka). Note the maximum in spectral density at 23 ka (precession) in the power spectrum of $\ln(\text{Fe}/\text{Ca})$ and the lack of spectral density at the precession bands (23 and 19 ka) in the power spectrum of the E-Heathland values. The Cyperaceae pollen concentration, which is both influenced by the expansion of Cyperaceae (sedges) and by the transport of pollen by river discharge, shows significant power at both the 100 and 19 ka.



Supplementary References


- Hammer, Ø., Harper, D.A.T. & Ryan, P.D., 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*, 4(1): 1-9.
- Schulz, M. & Mudelsee, M., 2002. REDFIT: estimating red-noise spectra directly from unevenly spaced paleoclimatic time series. *Computers & Geosciences*, 28: 421-426.

Excel-Tables


Supplementary Table 1: Family, Pollen taxon, group, growth form, endmember assemblage of maximum score.

Supplementary Table 2: Family, Pollen taxon, group, growth form, score per endmember (E-Mountain-Forest, E-Heathland, E-Woodland, E-Shrubland), significance of taxon scores (r^2) for two endmember analyses [this study and Dupont et al. 2011].


Supplementary Figures 1-4




Supplementary Figure 1. Endmember assemblage scores and selected pollen taxa for E-Heathland (filled curves) against age in ka. On top Terminations and even-numbered marine isotope stages (MIS) of the Brunhes Chron are indicated. Stable oxygen isotopes of benthic foraminifers of Core MD96-2048 (violet line) and of global stack LR04 (black line, [Lisiecki & Raymo 2005](#)).



Supplementary Figure 2. Endmember assemblage scores and selected pollen taxa for E-Mountain-Forest (filled curves) against age in ka. On top Terminations and even-numbered marine isotope stages (MIS) of the Brunhes Chron are indicated. Stable oxygen isotopes of benthic foraminifers of Core MD96-2048 (violet line) and of global stack LR04 (black line, [Lisiecki & Raymo 2005](#)).



Supplementary Figure 3. Endmember assemblage scores and selected pollen taxa for E-Shrubland (filled curves) against age in ka. On top Terminations and even-numbered marine isotope stages (MIS) of the Brunhes Chron are indicated. Stable oxygen isotopes of benthic foraminifers of Core MD96-2048 (violet line) and of global stack LR04 (black line, [Lisiecki & Raymo 2005](#)).



Supplementary Figure 4. Endmember assemblage scores and selected pollen taxa for E-Woodland (filled curves) against age in ka. On top Terminations and even-numbered marine isotope stages (MIS) of the Brunhes Chron are indicated. Stable oxygen isotopes of benthic foraminifers of Core MD96-2048 (violet line) and of global stack LR04 (black line, [Lisiecki & Raymo 2005](#)).