Multiproxy evidence of the Neoglacial expansion of Atlantic Water to eastern Svalbard

3

4

Joanna Pawłowska^{1*}, Magdalena Łącka¹, Małgorzata Kucharska¹, Jan Pawlowski^{1,2}, Marek
Zajączkowski¹

7

8 ¹Institute of Oceanology Polish Academy of Sciences, Sopot, 81-712, Poland

9 ²Department of Genetics and Evolution, University of Geneva, Geneva, CH 1211, Switzerland

10 Correspondence to: Joanna Pawłowska (pawlowska@iopan.pl)

11

12

Abstract. The main goal of this study is to reconstruct the paleoceanographic development of 13 14 Storfjorden during the Neoglacial (~ 4 cal ka BP). Storfjorden is one of the most important 15 brine factories in the European Arctic and is responsible for deep water production. Moreover, 16 it is a climate-sensitive area influenced by two contrasting water masses: warm and saline Atlantic Water (AW) and cold and fresh Arctic Water (ArW). Herein, a multiproxy approach 17 was applied to provide evidence for existing interactions between the inflow of AW and sea-18 19 ice coverage, which are the major drivers of environmental changes in Storfjorden. The sedimentary and microfossil records indicate that a major reorganization of oceanographic 20 conditions in Storfjorden occurred at ~ 2.7 cal ka BP. The cold conditions and the less-21 pronounced presence of AW in Storfjorden during the early phase of the Neoglacial were the 22 prerequisite conditions for the formation of extensive sea-ice cover. The period after ~ 2.7 cal 23 ka BP was characterized by alternating short-term cooling and warming intervals. Warming 24 25 was associated with pulsed inflows of AW and sea-ice melting that stimulated phytoplankton blooms and organic matter supply to the bottom. The cold phases were characterized by heavy 26 and densely packed sea ice resulting in decreased productivity. The ancient environmental 27 DNA (aDNA) records of foraminifera and diatoms support the occurrence of the major pulses 28 29 of AW (~2.3 and ~1.7 cal ka BP) and the variations in sea-ice cover. The episodes of enhanced AW inflow were marked by an increase in the percentage of DNA sequences of 30 31 monothalamous foraminifera associated with the presence of fresh phytodetritus. Cold and 32 less productive intervals were marked by an increased proportion of monothalamous taxa known only from environmental sequencing. The diatom aDNA record indicates that primary
 production was continuous during the Neoglacial, regardless of the sea-ice conditions.
 However, the colder periods were characterized by the presence of diatom taxa associated
 with sea ice, whereas the present-day diatom assemblage is dominated by open-water taxa.

5

6 **1. Introduction**

7 The northward flow of Atlantic Water (AW) is one of the major contributors of heat to the Arctic Ocean (Polyakov et al., 2017). Recent oceanographic data indicate a warming trend 8 9 due to an increased inflow of AW towards the Arctic Ocean (Rudels et al., 2015, Polyakov et 10 al., 2017). AW has been present along the western margin of Svalbard for at least the last 11 12,000 years (e.g., Werner et al., 2011; Rasmussen et al., 2014). One of the major intrusions of AW occurred during the early Holocene (10.8 - 6.8 cal ka BP). A distinct cooling and 12 13 freshening of the west Spitsbergen shelf bottom water masses occurred during the mid-late Holocene (6.8 – 1 cal ka BP) and was accompanied by glacier readvances in Svalbard, leading 14 15 to the present-day conditions (Ślubowska-Woldengen et al., 2007; Telesiński et al., 2018). The paleoceanographic conditions in the Svalbard margins correlate closely to the sea surface 16 17 temperature (SST) variations in the Nordic Seas and confirm that the Svalbard area is highly sensitive to fluctuations in the inflow of AW (Slubowska-Woldengen et al., 2007; Werner et 18 al., 2013). Conversely, until the 1990s eastern Svalbard was recognized as an area exclusively 19 influenced by the East Spitsbergen Current (ESC), which carries cold, less saline Arctic Water 20 (ArW) from the Barents Sea (e.g., Quadfasel et al., 1988; Piechura et al., 1996). However, 21 recent studies have revealed that the oceanography of the area is much more complicated 22 (e.g., Skogseth et al., 2007; Geyer et al., 2010). Oceanographic data obtained from 23 conductivity-temperature sensors attached to Delphinapterus leucas show a substantial 24 contribution of AW to Storfjorden, East Spitsbergen (Lydersen et al., 2002). Recently, Hansen 25 26 et al. (2011) suggested the presence of AW in Storfjorden during the early Holocene warming (11 - 6.8 cal ka BP), which was further confirmed by the foraminiferal and sedimentary 27 records of Łącka et al. (2015). 28

The latter part of the Holocene, the so-called Neoglacial cooling (~ 4 cal ka BP), in the European Arctic is characterized by a declined summer insolation at northern latitudes (Berger, 1978) that correlates to a decline in summer SST (e.g., Andersen et al., 2004; Risebrobakken et al., 2010; Rasmussen et al., 2014; Ivanova et al., 2019). The cooling of the surface waters and the limited AW inflow towards the Nordic Seas led to the formation of an extended sea-ice cover in West Spitsbergen margin (Müller et al., 2012). In addition, the

southwestern and eastern shelf of Spitsbergen experienced a strengthening of the East 1 2 Spitsbergen Current and/or Jan Mayen Current leading to an intensification of ArW inflow and the formation of extensive sea-ice cover (e.g., Sarnthein et al., 2003; Berben et al., 2014). 3 Therefore, the Neoglacial is usually considered a generally cold period (e.g., Consolaro et al., 4 2018). However, the records from Storfjorden and the Barents Sea suggest that the Neoglacial 5 6 was a period of variable oceanographic conditions with strong temperature and salinity 7 gradients (Martrat et al., 2003; Risebrobakken et al. 2010; Sarnthein et al., 2003; Łącka et al., 2015; 2019). In addition, there is evidence of episodic intensifications of the warm AW 8 9 inflow towards western Svalbard at that time (e.g.; Rasmussen et al., 2012).

According to Nilsen et al. (2008), the critical parameter controlling the fjord-shelf 10 11 exchange is the density difference between the fjord water masses and the AW. The local winter ice production and the formation of brine-enriched waters determine the density of 12 13 local water masses, which is a key factor that enables AW to penetrate into fjords during the spring and summer. Moreover, the production of brine-enriched waters and the associated 14 15 deep-water overflow are key contributors to large-scale ocean circulation (Killworth, 1983). In this respect, Storfjorden is especially important because it is one of the few areas where 16 17 brine-enriched waters have been frequently observed (Haarpainter et al., 2001). In recent decades, reduced brine formation has occurred during the periods with the most intensive AW 18 advection to Storfjorden and less sea-ice formation in the Barents Sea, while intense brine 19 20 formation re-establishes during periods of recurrent cooling (Årthun et al., 2011;Rasmussen 21 and Thomsen, 2014).

The aim of this study is to reconstruct the paleoceanographic development of 22 Storfjorden during the Neoglacial at multicentennial resolution. We assumed that the periodic 23 intensification of the AW inflow to the West Spitsbergen shelf during the Neoglacial resulted 24 in the appearance of AW also in eastern Spitsbergen, similar to the conditions in the early 25 26 Holocene (e.g., Łącka et al., 2015), affecting the density and extent of sea-ice cover in the area. A multiproxy approach comprising sedimentary, microfossil and molecular records was 27 28 applied to provide evidence for the interactions between the inflow of AW and sea-ice coverage in Storfjorden. The ancient environmental DNA (aDNA) analysis targeted diatoms 29 30 and nonfossilized monothalamous foraminifera. Both these of groups are hardly preserved in fossil records from the Svalbard fjords (Pawłowska et al., 2014) and shelf areas (Zimmermann 31 et al., 2019 and references therein). Recent studies have demonstrated that analyses of genetic 32 material obtained directly from environmental samples (so-called environmental DNA) are an 33 34 efficient method for performing biodiversity surveys across time and space (Thomsen and

Willerslev, 2015). The content of environmental DNA samples may be analyzed by DNA 1 metabarcoding, which consists of high-throughput sequencing of taxonomically informative 2 DNA fragments called metabarcodes. The identification of short, species-specific DNA 3 fragments (so called "barcodes") allows us to obtain species-level assignments of modern and 4 ancient DNA sequences (Herbert et al., 2003). The further demonstration that DNA can be 5 preserved in the environment across geological timescales opened new avenues for 6 7 palaeoclimatic and palaeoceanographic studies. Recent studies have demonstrated the preservation of DNA in marine sediments for tens to hundreds thousands of years. An aDNA 8 9 approach was successfully applied to trace the Holocene history of dinoflagellates, haptophytes (e.g., Coolen et al., 2009, 2013; Boere et al., 2009) and foraminifera in deep sea 10 11 (Lejzerowicz et al., 2013) and coastal areas (Pawłowska et al., 2014; 2016). The study of Pawłowska et al. (2016) was the first attempt to utilize foraminiferal aDNA as a 12 13 paleoenivronmental proxy. This study supported the existence of extremely diverse foraminiferal assemblages. The richness of the foraminiferal community revealed by the 14 15 molecular record was much higher than that in the fossil record (Pawłowska et al., 2014), mainly due to the detection of nonfossilized monothalamous taxa. The molecular data 16 17 correlated well with environmental changes and revealed even small changes that were not clearly indicated by other proxy records. The combination of aDNA studies with the analysis 18 of microfossils and sedimentary proxies provides a powerful means to reconstruct past 19 20 environments more comprehensively.

21

22 **2. Study area**

Storfjorden is located in southeastern Svalbard between the islands of Spitsbergen,
Edgeøya and Barentsøya (Fig. 1). Storfjorden is ~190 km long and its main basin is ~190 m
deep. Two narrow and shallow passages (Heleysundet and Freemansundet) connect northern
Storfjorden to the Barents Sea. To the south, a 120-m-deep sill separates the main basin from
the Storfjordrenna. Storfjordrenna is 245 m long, with a depth varying from 150 m to 420 m.

The water masses in Storfjorden are composed primarily of exogenous Atlantic and Arctic waters as well as mixed waters that have formed locally. The Atlantic Water is transported northwards by North Atlantic Current (NAC), which branches off in the Barents Sea into the West Spitsbergen Current and North Cape Current (Rudels et al., 2015; Loeng et al. 1991) (Fig. 1a). In addition to AW, cold and less saline Arctic Water (ArW) is transported to the Barents Sea via East Spitsbergen Current and Bear Island Current (Hopkins, 1991) (Fig. 1a). Arctic water (ArW) from the Arctic Ocean and the Barents Sea enters Storfjorden via two

passages to the northeast and continues along the inner shelf of Svalbard as a Coastal Current 1 (Fig. 1b). AW is characterized by temperatures > 3 °C and salinity > 34.95, while the 2 temperature and salinity of ArW are < 0 °C and 34.3-34.8, respectively. The presence of 3 locally formed water masses is a result of the interactions between AW, ArW and melt water. 4 Skogseth et al. (2005) listed six local water masses: melt water (MW), polar front water (PW), 5 East Spitsbergen water (ESW), brine-enriched shelf water (BSW), Storfjorden surface water 6 7 (SSW), and modified Atlantic water (MAW). BSW is formed due to the release of a large amounts of brines during polynya events and the intensive formation of sea ice (Haarpainter 8 9 et al., 2001; Skogseth et al., 2004, 2005) and is characterized by salinities exceeding 34.8 and 10 temperatures below -1.5 °C (Skogseth et al., 2005).

11 The sedimentary environment in Storfjorden is classified as a low-energy, high-12 accumulation environment, which is characteristic of inner fjords. The area is sheltered from 13 along-shelf bottom currents and is affected by high terrigenous inputs; therefore deposition 14 prevails over sediment removal by bottom currents (Winklemann and Knies, 2005). The 15 primary productivity is high and strongly depends on the sea-ice formation as well as the 16 seasonal duration of the marginal ice zone (Winkelman and Knies, 2005).

17

18 **3. Materials and methods**

19 3.1 Marine sediment core

The 55-cm-long sediment core ST_1.5 was taken with a gravity corer in Storfjorden retrieved with the R/V *Oceania* in August 2014. The sampling station was located at 76° 53,181' N and 19° 27,559' E at a depth of 153 m (Fig. 1). The salinity and temperature of the water column at the coring station was measured with a Mini CTD Sensordata SD 204 at intervals of 1 s (Fig. 2). The core was stored at 4°C and shipped to the Institute of Oceanology PAS for further analyses.

In the laboratory, the core was extruded and cut into 1-cm slices. During cutting, sterile subsamples for ancient DNA (aDNA) analyses were taken at 4 cm intervals. To avoid external and/or cross-contamination the thin layers of sediment that were in contact with under- or overlying sediments were removed using a sterile spatula. Samples for aDNA analyses were kept frozen at -20°C. Samples for other proxy analyses were taken every 2 cm.

31

32 **3.2 Chronology**

The chronology of the marine sediment core is based on high-precision accelerator mass spectrometry (AMS) ¹⁴C dating performed on five bivalve shells retrieved from the

sediment layers at 2.5, 5.5, 14.5, 43.5, and 52.5 cm core depth and on the foraminifera 1 Nonionellina labradorica from the 46.5 cm core depth. The bivalve shells were identified to 2 the highest possible taxonomic level and processed on the 1.5 SDH-Pelletron Model 3 "Compact Carbon AMS" in the Poznań Radiocarbon Laboratory, Poznań, Poland. Dating of 4 5 foraminiferal tests was performed at the National Ocean Sciences AMS (NOSAMS) laboratory in the Woods Hole Oceanographic Institution, Woods Hole, MA, USA. The dates 6 7 were converted into calibrated ages using the calibration program CALIB Rev. 7.1.0 Beta (Stuiver and Reimer, 1993) and the Marine13 calibration dataset (Reimer et al., 2013). A 8 reservoir age correction (ΔR) of 105 ± 24 was applied (Mangerud et al., 2006). The calibrated 9 results are reported in units of thousand calibrated years BP (cal ka BP) (Table 1). 10

11

12 **3.3 Grain size analysis**

13 The samples for grain size analyses were freeze-dried and milled. The measurements were performed using a Mastersizer 2000 particle laser analyzer coupled to a Hydro MU 14 15 device (Malvern, UK). The samples were treated with ultrasound to avoid aggregation. The raw data were analyzed using GRADISTAT v.8.0 software (Blott and Pye, 2001). The mean 16 17 0-63- μ m grain size [ϕ] was calculated via the logarithmic method of moments. The sediment fraction $>500 \ \mu m$ was used to reconstruct an ice rafted debris (IRD) record. The grains were 18 counted under a stereomicroscope and the amount of IRD is reported as the concentration 19 (i.e., the number of grains per gram of dry sediment) [grains g^{-1}] and the flux [grains $cm^{-2} y^{-1}$]. 20 21

22 **3.4 Benthic foraminifera assemblages**

Prior to the analysis of testate benthic foraminifera, samples were wet sieved through a 23 meshes with 500-µm and 100-µm openings and dried at 60°C. Samples with large quantities 24 of tests were divided using a microsplitter. At least 300 specimens of benthic foraminifera 25 26 were isolated from each sample and collected on micropaleontological slides. Benthic foraminifera specimens were counted and identified to the lowest possible taxonomic level. 27 The quantity of foraminifera is presented as the concentration (i.e., the number of individuals 28 per gram of dry sediment) [ind. g^{-1}] and the flux [ind. $cm^{-2} y^{-1}$]. For a species were 29 grouped according to their ecological tolerances. Four groups of indicators were 30 distinguished: AW/frontal zone indicators, ArW indicators, bottom current indicators and 31 glaciomarine species (Majewski et al., 2009). The morphologically similar species Islandiella 32 norcrossi and Islandiella helenae are reported as Islandiella spp. 33

1 **3.5 Stable isotope analysis**

Carbon and oxygen stable isotope analyses were performed on *Cibicidoides lobatulus*tests selected from 27 sediment layers. Ca. 10 to 12 specimens were collected from each
sample and subjected to ultrasonic cleaning. The measurements were performed on a Finnigan
MAT 253 mass spectrometer coupled to a Kiel IV carbonate preparation device at the
University of Florida. The resulting values are expressed in standard δ notation relative to
Vienna Pee Dee Belemnite (VPDB).

8

9 **3.6 Ancient DNA analysis**

The total DNA was extracted from approximately 10 g of sediment using a Power 10 Max Soil DNA extraction kit (MoBio). The foraminiferal SSU rDNA fragments containing 11 the 37f hypervariable region were PCR amplified using primers tagged with unique sequences 12 13 of five nucleotides appended to their 5' ends (denoted by Xs), namely, the foraminiferaspecific forward primer s14F1 (5'-XXXXXCGGACACACTGAGGATTGACAG-3') and the 14 15 reverse primer s15 (5'-XXXXXCCTATCACATAATCATGAAAG-3'). The diatom DNA fragment located in the V4 region was amplified with the forward DIV4for (5'-16 17 XXXXXXXGCGGTAATTCCAGCTCCAATAG-3') and reverse DIV4rev3 (5'-XXXXXXXXCTCTGACAATGGAATACGAATA-3') primers tagged with a unique 18 combination of eight nucleotides (denoted by Xs) attached at each primer's 5'-end. The 19 20 amplicons were purified using the High Pure PCR Cleanup Micro Kit (Roche) and quantified using a Qubit 2.0 fluorometer. Samples were pooled in equimolar quantities, and the sequence 21 library was prepared using a TruSeq library-preparation kit (Illumina). The samples were then 22 loaded into a MiSeq instrument for a paired-end run of 2*150 cycles (foraminifera) and 2*250 23 cycles (diatoms). The processing of the HTS sequence data was performed according to 24 procedures described by Leizerowicz et al. (2013) and Pawłowska et al. (2014). The post-25 26 sequencing data processing was performed with the use of the SLIM web app (Dufresne et al., 2019) and included demultiplexing the libraries, joining the paired-end reads, chimera 27 removal, operational taxonomic units (OTUs) clustering, and taxonomic assignment. 28 Sequences were clustered into OTUs using the Swarm module (Mahe et al. 2014), and each 29 OTU was assigned to the highest possible taxonomic level using vsearch (Rognes et al., 2016) 30 against a local database and then reassigned using BLAST (Altschul et al., 1990). The results 31 are presented in OTU-to-sample tables and transformed in terms of the number of sequences, 32 number of OTUs and percentage (%) of sequences. 33

- 1 **4. Results**
- 2

3 4.1 Chronology

In total, six radiocarbon dates were obtained, all of which were recorded in 4 5 chronological order. The uppermost layer contained modern, post-bomb carbon indicating a post-1960 age (Table 1). Samples from the 2.5 cm and 5.5 cm core depths were not calibrated 6 7 because they revealed ages invalid for the selected calibration curve. The age model was, therefore, based on the four remaining dates using a linear interpolation. The age of the 8 bottom of the core was estimated to be approximately ~ 7.9 cal ka BP (Fig. 3). However, the 9 extremely low temporal resolution between ~ 7.9 cal ka BP and ~ 4 cal ka BP precluded 10 making any general conclusion about that interval. Therefore, this study focuses only on the 11 last ~ 4 cal ka BP (the Neoglacial). 12

13 4.2 Sediment grainsize

The sediment was classified as medium to coarse silt throughout the core. The 14 sediment accumulation rate (SAR) prior to ~ 2.7 cal ka BP was 0.002 cm y^{-1} . The 15 approximately 10-fold increase in SAR was noted at ~ 2.7 cal ka BP, where it increased to 16 0.023 cm y⁻¹. During the last 1.5 cal ka BP, SAR decreased to 0.01 cm y⁻¹ (Fig. 4). The 17 amount of IRD was the highest prior to ~ 2.7 cal ka BP, reaching up to 83 grains g^{-1} . After ~ 18 2.7 cal ka BP, the amount of IRD was relatively stable and did not exceed 18 grains g⁻¹. The 19 IRD flux decreased slightly over time to 0.37 grains g^{-1} cm⁻¹, except for one peak reaching 0.8 20 grains g^{-1} cm⁻¹ at ~ 2.6 cal ka BP (Fig. 4). 21

The mean grain size of the 0-63- μ m fraction had its highest value (5.8 ϕ) at ~ 2.7 cal ka BP (Fig. 4). After ~ 2.4 cal ka BP a slight but continuous reduction in the mean 0-63- μ m grain size was noted. The minimum grain size (6.23 ϕ) was recorded at the top of the core (Fig. 4).

26

27 **4.3 Stable isotopes**

The δ^{18} O values prior to ~ 2.7 cal ka BP changed slightly between 3.55‰ and 3.69‰ vs. VPDB. Between ~ 2.7 and ~ 1.5 cal ka BP, δ^{18} O showed the strongest variation, with values ranging from 3.28‰ to 3.77‰ vs. VPDB. After ~ 1.5 cal ka BP, δ^{18} O became slightly lighter (3.43‰ - 3.64‰ vs. VPDB), except for one peak noted in the uppermost layer of the core, where δ^{18} O reached 3.87‰ vs. VPDB (Fig. 4).

In the period prior to ~ 2.7 cal ka BP, δ^{13} C displayed relatively light values ranging from 0.92‰ to 1.12‰ vs. VPDB. Slightly heavier δ^{13} C (up to 1.46‰ vs. VPDB) was observed between ~ 2.7 and ~ 1.5 cal ka BP. The gradual decrease was recorded from ~ 1.5
cal ka BP to the present, reaching 0.81‰ vs. VPDB at the top of the core (Fig. 4).

3

4 **4.4 Benthic foraminifera assemblages**

5 A total of 8647 fossil foraminifera specimens belonging to 47 species were identified (Supplement 1; Supplementary Fig. 1). The number of foraminifera individuals in a sample 6 varied from 156 to 2610 ind. g^{-1} , and the lowest abundances were observed prior to ~ 2.7 cal 7 ka BP (Fig. 4). A short-term decrease in foraminifera abundance was observed between 2.2 8 and 1.7 cal ka BP, with values reaching as low as 304 ind. g^{-1} . The abundance maxima were 9 noted at 2.3, 1.5, and 0.6 ka BP, with values reaching 2524, 2584, and 2610 ind. g^{-1} , 10 respectively. The foraminiferal flux was low and relatively stable throughout the core, with 11 values that did not exceed 1 ind cm^{-2} y⁻¹, except for two peaks at 2.3 and 1.5 ka BP, when the 12 flux reached 2.2 ind cm^{-2} y⁻¹ for both peaks (Fig. 4). 13

The most abundant species was *Cassidulina reniforme*, with densities reaching up to 900 ind g⁻¹. The other species that constituted the majority of the foraminiferal assemblage were *Bucella frigida*, *Cibicidoides lobatulus*, *Elphidium excavatum*, *Islandiella* spp, *Melonis barleeanum*, and *Nonionellina labradorica*. The abundances of the dominant species followed a general trend, with maxima at ~ 2.3 cal ka BP and after ~ 1.7 cal ka BP and minima prior to ~ 2.7 cal ka BP and between 2.3 and 1.7 cal ka BP. (Fig. 5).

The foraminiferal assemblage prior to ~ 2.7 cal ka BP was dominated by indicators of 20 AW inflow and/or frontal zones and glaciomarine taxa (Fig. 5). The most abundant species 21 were Nonionellina labradorica and Melonis barleeanum, as well as Cassidulina reniforme 22 and Elphidium excavatum, which together accounted for up to 60% of the foraminiferal 23 abundance (Fig. 5). After ~ 2.7 cal ka BP, there were AW/frontal zone indicator peaks 24 recorded at 2.4 and 1.8 cal ka BP, where the percentages increased to 33% and 28% of the 25 total abundance, respectively. The period between ~ 2.4 cal ka BP and ~ 1.8 cal ka BP was 26 characterized by an increase in the percentage of sea-ice indicators (B. frigida and Islandiella 27 spp), which accounted for up to 25% of the total foraminiferal abundance. Additionally, a 28 short-term peak in the glaciomarine taxa, reaching up to 49% of the foraminiferal assemblage, 29 was recorded between 2.5 and 2.1 cal ka BP. Also, the overall maximum of glaciomarine 30 species abundance was recorded between 1.7 and 0.5 cal ka BP, ranging from 25% to 43% of 31 foraminiferal assemblage (Fig. 5). A decrease in the relative abundance of glaciomarine 32 species was observed after ~ 0.5 cal ka BP and was followed by an increase in the AW/frontal 33

zone indicators and a single peak in the percentage of bottom current indicators, which
 reached 42% and 19%, respectively (Fig. 5).

3

4 4.5 Foraminiferal aDNA sequences

5 A total of 1,499,889 foraminiferal DNA sequences were clustered into 263 OTUs, and 6 20 remained unassigned. The remaining OTUs were assigned to Globigerinida (5 OTUs), 7 Robertinida (1 OTU), Rotaliida (49 OTUs), Textulariida (18 OTUs), Monothalamea (163 8 OTUs), and Miliolida (7 OTUs). The majority of sequences belonged to Monothalamea 9 (60%) and Rotaliida (31%) (Supplement 2; Supplementary Fig. 2). Herein, we focus on 10 Monothalamea, which is the dominant component of the foraminiferal aDNA record.

11 The most important components of the monothalamous assemblage were Micrometula sp., Cylindrogullmia sp., Hippocrepinella hirudinea, Ovammina sp., Nemogullmia sp., 12 13 Tinogullmia sp., Cedhagenia saltatus, undetermined allogromiids belonging to clades A and Y (herein called "allogromiids"), and sequences belonging to taxa known exclusively from 14 15 environmental sequencing (herein called "environmental clades"). Herein, the term "allogromiid" refers to monothalamous foraminifera with organic or predominantly organic 16 17 test walls (Gooday, 2002). Morphological and molecular evidence indicate that 'allogromiids' are not a coherent taxonomic group but are scattered between several monothalamous clades 18 (Pawlowski et al., 2002). "Clade" refers to phylogenetic clades defined by molecular data. 19 20 The clade is traditionally defined as a group of organisms that includes a common ancestor and all the descendants. The sequences belonging to allogromiids were present throughout the 21 core, accounting for 16-31.7% of all the foraminiferal sequences. The exceptions were the 22 intervals from ~ 4.0 to 2.4 cal ka BP, and ~ 1.7 cal ka BP, when the contribution of 23 allogromiid sequences decreased to less than 10% (Fig. 6). The majority of the allogromiids 24 belonged to clade Y, which made up to 100% of the allogromiid sequences. Only at 1.6-1.7 25 26 cal ka BP and 2.4-2.6 cal ka BP, most of allogromiid sequences belonged to clade A. 27 Additionally, allogromiids belonging to clade I were noted at ~ 2.4 cal ka BP, where they 28 made up 0.88% of allogromiid sequences (Fig. 7).

The periods prior to ~ 2.4 cal ka BP and ~ 1.7 cal ka BP were marked by the disappearance of sequences belonging to *C. saltatus*, *Nemogullmia* sp., and the environmental clades, followed by an increase in the percentages of sequences belonging to *Micrometula* sp., *Ovammina* sp., *Tinogullmia* sp., *Shepheardella* sp. and *Cylindrogullmia* sp. (Fig. 6).

- 33
- 34

1 4.6 Diatom aDNA sequences

A total of 824,697 diatom DNA sequences were clustered into 221 OTUs (Supplement 2 3; Supplementary Figure 3). The most abundantly sequenced diatom taxa were Thalassiosira 3 spp, which made up 61.1% of diatom sequences. Other abundantly sequenced taxa were 4 5 Chaetoceros sp. and T. antarctica, which made up 8.5% and 11.5% of sequences, respectively. The sequences of *Thalassiosira* sp were most abundant between ~ 2.2 cal ka BP 6 7 and ~ 1.9 cal ka BP, accounting for up to 85% of all diatom sequences. The lowest percentage (14%) of *Thalassiosira* sp. was recorded at ~ 0.4 cal ka BP. Sequences assigned to T. 8 9 antarctica were recorded throughout the core and their percentages were the highest at ~ 3.3 and ~ 2.6 cal ka BP, reaching up to 13% and 19%, respectively (Fig. 8). Sequences of T. 10 hispida were also noted throughout the core and constituted 4.7% of diatom sequences in the 11 uppermost layer. In the remaining samples, T. hispida sequences did not exceed 1%. The 12 13 percentage of sequences of Chaetoceros sp. decreased downcore, from 76% at the surface to less than 1% at the bottom of the core (Fig. 8). Navicula sp. constituted an important part of 14 15 the diatom assemblage at ~3.3 cal ka BP and ~1.9 cal ka BP, accounting for up to 25.5% and 10% of all diatom sequences, respectively. In the remaining samples, the abundance of 16 17 Navicula sp. did not exceed 5% (Fig. 8).

18

19 **5. Discussion**

The ST_1.5 age model is based on the linear interpolation between the four $AMS^{14}C$ dates; thus, the age control of the core should be treated with caution. However, the timing of major environmental changes revealed by the ST_1.5 multiproxy record is in agreement with other records from the region (e.g., Sarnthein et al., 2003; Risebrobakken et al. 2010; Berben et al. 2017). Moreover, the major pulses of AW that were recorded ~ 2.3 and 1.7 cal ka BP correlated well with winter and summer SST maxima recorded in the 23258-2 core (Sarnthein et al., 2003).

27

28 5.1 The period from 4 cal ka BP to 2.7 cal ka BP

Prior to ~ 2.7 cal ka BP, the ST_1.5 sedimentary record displayed relatively higher IRD delivery and a relatively lower 0-63- μ m sediment fraction than in the following period (Fig. 4). These results are in agreement with the record from Storfjordrenna (Łącka et al., 2015), where peaks in IRD were noted during the Neoglacial and were attributed to increased iceberg rafting due to fluctuations in the glacial fronts (e.g. Forwick et al., 2010).The coarser 0-63 μm fraction may suggest the winnowing of fine grained sediment, however,
 foraminiferal fauna showed no clear response to sediment removal.

The foraminiferal flux and abundance prior to 2.7 cal ka BP reached their lowest 3 values (Fig. 4). Previous studies reported a decrease in te concentration of benthic 4 foraminifera in Storfjorden at that time, which was attributed to the presence of extensive ice 5 cover (Rasmussen and Thomsen, 2015; Knies et al. 2017). The dominant components of the 6 7 ST_1.5 foraminiferal assemblage were C. reniforme. and M. barleeanum (Fig. 5). The presence of C. reniforme and M. barleeanum is associated with cooled and salty AW (e.g., 8 9 Hald and Steinsund, 1996; Jernas et al., 2013). Moreover, these species are also associated with the presence of phytodetritus, which may be related to the delivery of fresh organic 10 matter observed in frontal zones and/or near the sea-ice edge (Jennings et al., 2004). The 11 presence of sea-ice may be indicated also by the relatively light foraminiferal δ^{13} C (Fig. 4), as 12 13 well as the highest percentage of the sea-ice species Thalassiosira antarctica (cf Ikävalko, 2004; Fig. 8). However, the low sampling resolution during that period precluded us from 14 15 making a general conclusion, and the latter assumptions should be confirmed by further studies. 16

17

5.2 The period from 2.7 cal ka BP to 0.5 cal ka BP. Episodes of AW inflow at ~ 2.3 and 1.7 cal ka BP.

After ~ 2.7 cal ka BP, the increase in SAR was followed by a decrease in the mean 20 grain size of the 0-63-µm fraction and in the IRD delivery (Fig. 4). The 10-fold increase in 21 SAR most likely resulted from the intensive supply of turbid meltwater from advancing 22 glaciers and the consequent intensive sedimentation. Moreover, the accumulation of fine 23 sediment may also be enhanced by the slowdown of the bottom currents, indicated by the 24 finer 0-63-µm sediment fraction (Fig. 4). On the other hand, a decrease in IRD delivery may 25 26 suggest that the central Storfjorden was not impacted by iceberg rafting at that time. In 27 contrast, Rasmussen and Thomsen (2015) suggested glacial advance, followed by intensive 28 ice rafting and meltwater delivery to Storfjorden at that time. According to Knies et al. (2017), the inner Storfjorden was covered by densely packed sea ice between ~ 2.8 and 0.5 cal 29 ka BP. Therefore, the decreasing IRD in the ST_1.5 core may result from the presence of a 30 sea-ice cover that reduced iceberg rafting while the majority of coarse-grained material settled 31 in the proximity to the glacial fronts. Similar conclusions have been stated by Forwick and 32 Vorren (2009) and Forwick et al. (2010), who assumed that the enhanced formation of sea ice 33 34 along the West Spitsbergen coast trapped icebergs inside the Isfjorden system.

The foraminiferal fauna in central Storfjorden revealed more than a 10-fold increase in 1 flux and abundance followed by short-term fluctuations after ~ 2.7 cal ka BP (Fig. 4). The 2 latter may suggest favorable conditions for foraminiferal growth. The major peaks in the total 3 foraminiferal abundance (Fig. 4) followed by the peaks in the percentage of AW foraminiferal 4 indicators (Fig. 5) were noted ~ 2.3 cal ka BP and ~ 1.7 cal ka BP. These peaks were 5 associated with the occurrence of sequences of T. hispida (Fig. 8), a diatom species 6 7 characteristic of subpolar and temperate regions (Katsuki et al., 2009). The timing of the changes described above is in accordance with the findings of Sarntheim et al. (2003), who 8 9 reported two intervals of the remarkably warmer sea surface on the western continental margin of the Barents Sea at ~ 2.2 and ~ 1.6 cal ka BP, which was attributed to short-term 10 11 pulses of warm AW advection. Other records also indicated AW inflow to the western and northern Barents Sea as well as to the western Spitsbergen continental margin during mid-late 12 13 Holocene (e.g., Risebrobakken et al., 2010; Berben et al., 2014; 2017; Müller et al., 2012), followed by change in environmental conditions at $\sim 2.5/2.7$ cal ka BP (Risebrobakken et al. 14 15 2010; Berben et al., 2017) Our foraminiferal and diatom aDNA records confirm the presence of AW intrusions that may have caused an episodic breakup of sea-ice cover and permitted 16 primary production and the development of benthic biota, including foraminifera. 17

The pulses of AW inflow at 2.3 cal ka BP and 1.7 cal ka BP were marked by the 18 maxima of the foraminiferal flux (Fig. 4) and by peaks in the abundance of species associated 19 20 with highly productive environments, such as *M. barleeanum* and *N. labradorica* (Fig. 5). Moreover, the presence of diatom aDNA sequences throughout the core (Fig. 8) may suggest 21 continuous primary production. However, the responses of the benthic foraminifera 22 assemblage to the pulses of AW at ~ 2.3 cal ka BP and ~ 1.7 cal ka BP are slightly different. 23 The dominant components of foraminiferal assemblage at ~ 2.3 cal ka BP were M. 24 barleeanum and E. excavatum, while at ~ 1.7 cal ka BP, N. labradorica and C. reniforme 25 26 were dominant (Fig. 5). The major difference in environmental conditions between these two "AW episodes" was noticeably coarser 0-63 µm sediment fraction noted at ~ 2.3 cal ka BP, 27 28 what may indicate more intensive winnowing of fine sediment grains,, which would have created favorable conditions for the development of opportunistic species, such as E. 29 *excavatum*. In contrast, the interval between 2.3 and 1.7 cal ka BP featured variable δ^{13} C and 30 δ^{18} O followed by a decrease in the foraminiferal flux and abundance (Fig. 4). The 31 foraminiferal assemblage at this time was dominated by glaciomarine and sea-ice taxa (Fig. 32 5), which indicate more severe environmental conditions with extensive ice cover and 33 34 suppressed productivity.

The alternate cooling and warming periods described above were also reflected in the 1 aDNA record of monothalamous foraminifera. During the periods with more severe 2 environmental conditions (i.e., time intervals of 2.2–1.9 cal ka BP and 1.3–0.4 cal ka BP), the 3 monothalamous foraminifera was dominated by allogromiids belonging to clade Y, 4 Nemogullmia sp., C. saltatus and monothalamids belonging to so called "environmental 5 clades" (Fig. 6). A considerable portion of the allogromiid sequences in the ST 1.5 core 6 7 belong to clade Y (Fig. 7), which is primarily composed of taxa known only from environmental sequencing that have previously been noted in modern sediments in the 8 9 Spitsbergen fjords (Pawłowska et al., unpubl.). Clade Y has also been abundantly sequenced in the coastal areas off Scotland, characterized by high levels of environmental disturbances 10 11 (Pawlowski et al., 2014); this might suggest its high tolerance to environmental stress. C. saltatus was found by Gooday et al. (2011) in the Black Sea and its occurrence in areas with 12 13 high levels of pollution suggests that it is an opportunistic species with a high tolerance for environmental disturbances. In addition, so called "environmental clades" are composed of 14 15 monothalamous taxa known exclusively from environmental sequencing (Lecroq et al., 2011).. The abovementioned taxa nearly disappeared during the episodes of enhanced AW 16 17 inflow at ~ 2.4 cal ka BP and ~ 1.7 cal ka BP, and the monothalamous assemblage was dominated at that time by Micrometula sp., Ovammina sp., Shepheardella sp., Tinogullmia 18 sp., Cylindrogullmia sp., and allogromiids belonging to clade A (Fig. 6; Fig. 7). All these taxa 19 have recently been observed in the fjords of Svalbard and Novaya Zemlya (e.g. Gooday et al., 20 2005; Majewski et al., 2005; Sabbattini et al., 2007; Pawłowska et al., 2014; Korsun & Hald, 21 22 1998; Korsun et al., 1995). Cylindrogullmia and Micrometula are dependent on the presence of fresh phytodetritus (Alve, 2010). Ovammina sp. feeds on diatoms and other forms of 23 microalgae (Goldstein & Alve, 2011). Similarly, the presence of *Tinogullmia* is largely 24 controlled by the presence of organic material on the seafloor. High concentrations of 25 26 Tinogullmia have been found in coastal (Cornelius & Gooday, 2004) and deep-sea regions (Gooday, 1993) within phytodetrital aggregates. 27

The taxa that dominated the monothalamous assemblage during warm intervals seem to be responsive to the delivery of organic matter and may flourish during phytoplankton blooms associated with the settling of organic matter (e.g., Alve, 2010; Sabbattini et al., 2012, 2013). The pulses of AW inflow may be associated with phytoplankton blooms stimulated by sea-ice melting and with the organic matter supply to the bottom (cf. Łącka et al., 2019). The continuous aDNA record of the sea-ice diatom *T. antarctica* (Fig. 8) suggests the presence of at least seasonal ice cover in the study area. On the other hand, the episodes of AW inflow

were associated with the occurrence of the open-water taxa T. hispida (Fig. 8). The 1 occurrence of sequences of both these taxa suggests the formation of ice cover during winter-2 spring, followed by ice-free summers. A similar scenario was proposed by Berben et al. 3 (2017), who suggested increased AW to the eastern Svalbard and partial summer sea ice 4 occurrence after 2.7 cal ka BP. According to the record of Łącka et al. (2019) from 5 Storfjordrenna, the sea-ice melting induced the production of brines that may launch 6 convective mixing and nutrient resupply from the bottom, which stimulated primary 7 8 production.

9 Conversely, the colder phases of the Neoglacial were characterized by heavy and 10 densely packed sea ice resulting in limited productivity (Knies et al., 2017). The presence of 11 *T. anatrctica* sequences and the disappearance of *T. hispida* (Fig. 8) may suggest that primary 12 production was associated with sea-ice. Furthermore, the monothalamous assemblage was 13 less diverse and was dominated by more opportunistic taxa, which may indicate a reduced 14 supply of organic matter to the bottom.

15

16 **5.3 The period after 0.5 cal ka BP.**

17 Modern-like conditions were established in Storfjorden at ~ 0.5 cal ka BP (Knies et al., 2017). The ST_1.5 record displayed a decrease in SAR compared to the preceding period, a 18 decreasing 0-63 µm fraction and low IRD delivery (Fig. 4), which may indicate reduced 19 glacial impact. Moreover, the peak of heavy δ^{18} O recorded on the core top (Fig. 4) may 20 suggest the presence of AW or slightly increased salinity. Similarly, Berben et al. (2014) 21 recorded δ^{18} O values that suggested a minor increase in salinity, while for a fauna 22 showed slightly lower salinities in the western Barents Sea at that time. The latter is in 23 accordance with records from the Fram Strait (e.g. Werner et al., 2013) and the western 24 Spitsbergen shelf (Cabedo-Sanz and Belt, 2016), which suggest episodes of freshening of the 25 26 surface water masses associated with alternating sea ice increases and ice-free conditions in 27 the late Holocene. Additionally, the records of Rasmussen and Thomsen (2014) and Knies et 28 al., (2017) from Storfjorden indicated seasonally variable sea-ice cover. Moreover, the 29 majority of diatom aDNA sequences found in the ST_1.5 record after ~ 0.5 cal ka BP belonged to Chaetoceros sp. (Fig. 8), a taxa that is observed in surface waters and is almost 30 entirely absent under sea ice (Różańska et al., 2008). High abundances of Chaetoceros are 31 often associated with highly productive surface waters (Cremer, 1999). Rigual-Hernández et 32 al. (2017) also noted increased abundance of *Chaetoceros* sp. and enhanced algal productivity 33 34 in Storfjorden after 2.0 cal ka BP, what was associated to the vicinity of the Arctic Front.

However, the aDNA record of the monothalamous foraminifera at ~ 0.4 cal ka BP displayed relatively high percentages of taxa that dominated during the colder intervals of the Neoglacial (Fig. 6). This may be related to the recovery from the Little Ice Age, and consequently, from the temporarily deteriorated environmental conditions (D'Andrea et al., 2012). However, due to the low resolution during the LIA, a detailed interpretation is not possible. Therefore, further studies are required to confirm the latter conclusion.

7 8

5.4 Paleoceanographic implications

9 Our record revealed a two-phase Neoglacial, with a major shift in environmental 10 conditions at ~ 2.7 cal ka BP. According to the ST_1.5 proxy records, the Neoglacial in Storfjorden was not a constantly cold period, but comprised alternating short-term cooling and 11 12 warming periods, associated with variability in sea-ice coverage and productivity. The Neoglacial cooling was documented in various proxy reconstructions from the Nordic Seas 13 14 (e.g., Jennings et al., 2002; Moros et al., 2004; Consolaro et al., 2018). However, there is growing evidence of shifts in environmental conditions in the Nordic Seas region in the 15 Neoglacial, whose timings are in accordance with our record.. Alkenone record from the 16 17 Norwegian Sea revealed a significant drop in sea surface temperature at 2.7 cal ka BP (Calvo et al., 2002). Risebrobakken et al. (2010) recorded a change in oceanographic conditions in 18 19 the SW Barents Sea ca. 2.5 cal ka BP. The episodes of reduced surface and subsurface salinity were recorded after 2.5 cal ka BP, what was attributed to the expansion of coastal waters and 20 21 the occurrence of more sea-ice (Risebrobakken et al., 2010). Berben et al. (2017) recorded a shift ~2.7 cal ka BP, from the marginal ice zone to Arctic frontal conditions in the eastern 22 Barents Sea. They observed continuous cooling trend from ~ 5.9 cal ka BP to 2.7 cal ka BP, 23 with increased seasonal sea ice with less open water conditions, lower temperatures and 24 25 decreased AW influence. Whereas, after 2.7 cal ka BP, the influence of AW was variable, but generally generally increasing. The period was characterized by low insolation, associated 26 27 with surface cooling and enhanced formation of sea ice/reduced sea ice melt (Berben et al., 2017). 28

Moreover, our evidence of the presence of AW in Storfjorden during the Neoglacial supported previous suggestions that AW inflow during the late Holocene was strong enough to reach also the eastern coasts of Svalbard (e.g., Łącka et al., 2015). Episodic increases of the AW during the late Holocene were also observed in the northern Barents Sea (Duplessy et al., 2001; Lubinski et al., 2001), the eastern Barents Sea (Berben et al., 2014) and the Svalbard margin (Jernas et al., 2013; Werner et al., 2013). Sarnthein et al. (2003) postulated pulses of
AW inflow to the western Barents Sea shelf at 2.2 and 1.6 cal ka BP. According to Perner et
al. (2015), the Neoglacial delivery of chilled AW to the Nordic Seas culminated between 2.3
and 1.4 cal ka BP. These results are in accordance with the timing of major AW inflows
revealed by our record.

6

7 **6.** Conclusions

The ST_1.5 multiproxy record revealed that the environmental variability in Storfjorden 8 9 during the Neoglacial was controlled primarily by the interplay between AW and ArW and 10 sea-ice cover variability. The molecular record supports and complements sedimentary and 11 microfossil records, which indicate that major changes in the environmental conditions in Storfjorden occurred at ~ 2.7 cal ka BP. The general cooling in the early phase of the 12 13 Neoglacial initiated conditions for the formation of extensive sea-ice cover. The latter part of the Neoglacial (after ~ 2.7 cal ka BP) was characterized by alternating short-term cooling and 14 15 warming periods. Warming was associated with pulsed inflows of AW and sea-ice melting, 16 which may stimulate phytoplankton blooms and organic matter supply to the bottom. The 17 cold phases were characterized by heavy and densely packed sea ice resulting in limited productivity. 18

Moreover, the aDNA diatom record supports the conclusion that primary production took place continuously during the Neoglacial, regardless of the sea-ice conditions. The early phase of the Neoglacial was characterized by the presence of diatom taxa associated with sea ice, whereas the present-day diatom assemblage was dominated by *Chaetoceros* spp, a taxa characteristic of open water.

The aDNA record of monothalamous foraminifera is in agreement with the microfossil record and revealed the timing of the major pulses of AW at 2.3 and 1.7 cal ka BP. The AW inflow was marked by an increase in the percentage of sequences of monothalamous taxa associated with the presence of fresh phytodetritus. The monothalamous assemblage during cold intervals was less diverse and was dominated by monothalamous foraminifera known only from environmental sequencing.

30

31 Author contributions

MZ and Jan P designed the study. Joanna P, MŁ and MZ collected the sediment core. MŁ and
MK performed the sedimentological and micropaleontological analyses. Joanna P performed
the molecular analyses and prepared the manuscript with contributions from all co-authors.

2 Acknowledgements

The study was supported by the National Science Centre grants no. 2015/19/D/ST10/00244 and 2016/21/B/ST10/02308, and Swiss National Science Foundation grant no. 31003A_179125. The Authors would like to thank anonymous Reviewers for constructive comments that helped to improve the manuscript.

7

8 **References**

- 9 Alve, E.: Benthic foraminiferal responses to absence of fresh phytodetritus: A two year
- experiment, Mar. Micropaleontol., 76, 67-76, https://doi.org/10.1016/j.marmicro.2010.05.003,
 2010.
- 12 Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J.: Basic local alignment search
- tool. J. Mol. Biol. 215, 403-410, https://doi.org/10.1016/S0022-2836(05)80360-2, 1990.
- 14 Andersen, C., Koç, N., Moros, M.: A highly unstable Holocene climate in the subpolar North
- 15 Atlantic: evidence from diatoms, Quat. Sci. Rev., 23, 2155-2166,
 16 https://doi.org/10.1016/j.quascirev.2004.08.004, 2004.
- 17 Årthun, M., Ingvaldsen, R.B., Smedsrud, L.H., Schrum, C.: Dense water formation and
- 18 circulation in the Barents Sea, Deep Sea Res. Part I: Oceanogr. Res. Pap., 58, 801-817,
- 19 https://doi.org/10.1016/j.dsr.2011.06.001, 2011.
- 20 Berben, S.M.P., Husum, K., Cabedo-Sanz, P., Belt, T.S.: Holocene sub-centennial evolution
- of Atlantic water inflow and sea ice distribution in the western Barents Sea, Clim. Past. 10,
- 22 181-198, doi:10.5194/cp-10-181-2014, 2014.
- 23 Berben, S.M.P., Husum, K., Navarro-Rodriguez, A., Belt, T., Aagard-Sørensen, S.: Semi-
- 24 quantitative reconstruction of early to late Holocene spring and summer sea ice conditions in
- 25 the northern Barents Sea, J. Quat Sci., 32, 587-603, doi: 10.1002/jqs.2953, 2017.
- 26 Berger, A.L.: Long-term variations of daily insolation and quaternary climatic changes, J.
- Atmos. Sci., 35, 2362-2367, https://doi.org/10.1175/15200469(1978)035<2362:LTVODI>2.0.CO;2, 1978.
- 29 Blott, S.J., Pye, K.: GRADISTAT: a grain size distribution and statistics package for the
- 30 analysis of unconsolidated sediments, Earth Surf. Process. Landf., 26, 1237 1248,
- 31 https://doi.org/10.1002/esp.261, 2001.
- 32 Boere, A.C., Abbas, B., Rijpstra, W.I.C., Versteegh, G.J., Volkman, J.K., Sinninghe Damsté,
- 33 J.S., Coolen, M.J.L.: Late-Holocene succession of dinoflagellates in an Antarctic fjord using a

- multi-proxy approach: paleoenvironmental genomics, lipid biomarkers and palynomorphs, 1
- Geobiol., 7, 265-281, https://doi.org/10.1111/j.1472-4669.2009.00202.x, 2009. 2
- Cabedo-Sanz, P., Belt, S.T.: Seasonal sea ice variability in eastern Fram Strait over the last 3
- 2000 years, Arktos, 2, 22, https://doi.org/10.1007/s41063-016-0023-2, 2016. 4
- 5 Consolaro, C., Rasmussen, T.L., Panieri, G.: Palaeocoeanographic and environmental changes
- in the eastern Fram Strait during the last 14,000 years based on benthic and planktonic 6
- 7 foraminifera, Marine Micropaleontology, 139, 84-101,
- https://doi.org/10.1016/j.marmicro.2017.11.001, 2018. 8
- Coolen, M.J.L., Saenz, J.P., Giosan, L., Trowbridge, N.Y., Dimitrov, P., Dimitrov, D., 9
- Eglinton, T.I.: DNA and lipid molecular stratigraphic records of haptophyte succession in the 10 Planet. 11 Black Sea during the Holocene, Earth Sci. Lett., 284, 610-621, https://doi.org/10.1016/j.epsl.2009.05.029, 2009.
- 12
- Coolen, M.J.L., Orsi, W.D., Balkema, C., Quince., C., Harris, K., Sylva, S.P., Filipova-13
- Marinova, M., Giosan, L.: Evolution of the plankton paleome in the Black Sea from the 14
- 15 Deglacial to Anthropocene, PNAS, 10, 8609-8614, https://doi.org/10.1073/pnas.1219283110, 2013. 16
- Calvo, E., Grimalt, J., Jansen, E.: High resolution U₃₇^K sea surface temperature reconstruction 17 in the Norwegian Sea during the Holocene. Quat. Sci. Rev., 21, 1385-1394, 18
- https://doi.org/10.1016/S0277-3791(01)00096-8, 2002. 19
- Cornelius, N., Gooday, A.J.: 'Live' (stained) deep-sea benthic foraminiferans in the western 20
- Weddell Sea: trends in abundance, diversity and taxonomic composition along a depth 21 transect, Deep Sea Res. II, 51, 1571-1602, https://doi.org/10.1016/j.dsr2.2004.06.024, 2004. 22
- Cremer, H.: Distribution patterns of diatom surface sediment assemblages in the Laptev Sea 23
- Mar. Micropaleontol., 38, 39-67, https://doi.org/10.1016/S0377-24 (Arctic Ocean), 8398(99)00037-7, 1999. 25
- 26 D'Andrea, W.J., Vaillencourt, D.A., Balascio, N.L., Werner, A., Roof, S.R., Retelle, M.,
- 27 Bradley, R.S.: Mid Little Ice Age and unpredecented recent warmth in an 1800 year lake
- sediment record from Svalbard, Geology, 40, 1007-1010, https://doi.org/10.1130/G33365.1, 28
- 2012. 29
- Dufresne, Y., Lejzerowicz, F., Apotheloz Perret-Gentil, L., Pawlowski, J., Cordier, T.: SLIM: 30
- a flexible web application for the reproducible processing of environmental DNA 31
- metabarcoding data. BMC Bioinformatics, 20, 88, https://doi.org/10.1186/s12859-019-2663-32
- 2, 2019. 33

- 1 Duplessy, J.C., Ivanova, E., Murdmaa, I., Paterne, M., Labeyrie, L.: Holocene
- 2 paleoceanography of the northern Barents Sea and variations of the northward heat transport
- 3 by the Atlantic Ocean. Boreas, 30, 2-16, https://doi.org/10.1111/j.1502-3885.2001.tb00984.x,
- 4 2001.
- 5 Forwick, M., Vorren, T.O.,: Late Weichselian and Holocene sedimentary environments and
- 6 ice rafting in Isfjorden, Spitsbergen, Palaeogeogr. Palaeoclimatol. Palaeoecol. 280, 258-274,
- 7 https://doi.org/10.1016/j.palaeo.2009.06.026, 2009.
- 8 Forwick, M., Vorren, T.O., Hald, M., Korsun, S., Roh, Y., Vogt, C., Yoo, K.-C.: Spatial and
- 9 temporal influence of glaciers and rivers on the sedimentary environment in Sassenfjorden
- 10 and Tempelfjorden, Spitsbergen. In: Geological Society, London, Special Publications, vol.
- 11 344: 163-193, https://doi.org/10.1144/SP344.13, 2010.
- 12 Geyer, F., Fer, I., Smedsrud, L. H.: Structure and forcing of the overflow at the Storfjorden
- sill and its connection to the Arctic coastal polynya in Storfjorden, Ocean Sci., 6, 401-411,
- 14 https://doi.org/10.5194/os-6-401-2010, 2010.
- 15 Goldstein, S.T., Alve, E.: Experimental assembly of foraminiferal communities from coastal
- propagule banks, Mar. Ecol. Prog. Ser. 437, 1-11, https://doi.org/10.3354/meps09296,
 2011.
- 18 Gooday, A.J.: Deep-sea benthic foraminiferal species which exploit phytodetritus:
- 19 Characteristic features and controls on distribution, Mar. Micropaleontol., 22, 187-205,
- 20 https://doi.org/10.1016/0377-8398(93)90043-W, 1993.
- 21 Gooday A.J.: Organic-walled allogromiids: aspects of their occurrence, diversity and ecology
- 22 in marine habitats. J Foramin. Res., 32, 384-399, https://doi.org/10.2113/0320384, 2002.
- 23 Gooday, A.J., Bowser, S.S., Cedhagen, T., Cornelius, N., Hald, M., Korsun, S., Pawłowski,
- 24 J.: Monothalamous foraminiferans and gromiids (Protista) from western Svalbard: A
- 25 preliminary survey, Mar. Biol. Res., 1, 290 312,
- 26 https://doi.org/10.1080/17451000510019150, 2005.
- 27 Gooday, A.J., Anikeeva, O.V., Pawlowski, J.: New genera and species of monothalamous
- 28 Foraminifera from Bataclava and Kazach'ya Bays (Crimean Peninsula, Black Sea), Mar.
- 29 Biodiv., 41, 481-494, https://doi.org/10.1007/s12526-010-0075-7, 2011.
- 30 Grootes, P.M., and M. Stuiver. 1997. Oxygen 18/16 variability in Greenland snow and ice
- 31 with 10⁻³ to 10⁻⁵-year time resolution. J. Geophys. Res., 102, 26455-26470,
 32 https://doi.org/10.1029/97JC00880, 1997.

- Haarpainter, J., Gascard, J.C., Haugan, P.M.: Ice production and brine formation in
 Storfjorden, Svalbard, J. Geophys. Res. 106, 14001–14013,
 https://doi.org/10.1029/1999JC000133, 2001.
- 4 Hald, M. Steinsund, P.I.: Benthic foraminifera and carbonate dissolution in the
 5 surface sediments of the Barents and Kara Seas, Berichte zur Polarforschung, 212,
 6 285–307, 1996.
- 7 Hansen, J., Hanken, N.-M., Nielsen, J.K., Nielsen, J.K., Thomsen, E.: Late Pleistocene and
- 8 Holocene distribution of *Mytilus edulis* in the Barents Sea region and its paleoclimatic
- 9 implications, J. Biogeogr, 38, 1197-1212, https://doi.org/10.1111/j.1365-2699.2010.02473.x,
 10 2011.
- 11 Herbert, P.D., Cywińska, A., Ball, S.L., deWaard, J.R.: Biological identifications through
- 12 DNA barcodes. Proc. Biol. Sci., 270, 313-321, doi:10.1098/rspb.2002.2218, 2003.
- 13 Hopkins, T.S.: The GIN Sea: A synthesis of its physicaloceanography and literature review,
- 14 1972–1985. Earth Sci. Rev., 30, 175–318, https://doi.org/10.1016/0012-8252(91)90001-V,
 15 1991.
- 16 Ikävalko, J.: Checklist of unicellular and invertebrate organisms within and closely associated
- 17 with sea ice in the Arctic regions. MERI Report Series of the Finnish Institute of Marine
- 18 Research, 52, Helsinki, Finland, Finnish Institute of Marine Research, 2004.
- 19 Ivanova, E., Murdmaa, I., de Vernal, A., Risebrobakken, B., Peyve, A., Brice, C., Seitkalieva,
- 20 Pisarev,: Postglacial paleoceanography and paleoenvironments in the northwestern Barents
- 21 Sea, Quat. Res., 1-20, https://doi.org/10.1017/qua.2019.18, 2019.
- 22 Jennings, A.E., Weiner, N.J., Helgadottir, G., Andrews, J.T.: Modern foraminiferal faunas of
- 23 the southwestern to northern Iceland Shelf; oceanographic and environmental controls, J.
- 24 Foramin. Res., 34, 180-207, https://doi.org/10.2113/34.3.180, 2004.
- 25 Jernas, P., Klitgaard Kristensen, D., Husum, K., Wilson, L., Koç, N.: Palaeoenvironmental
- changes of the last two millennia on the western and northern Svalbard shelf, Boreas, 42, 236-
- 27 255, https://doi.org/10.1111/j.1502-3885.2012.00293.x, 2013.
- 28 Katsuki, K., Takahashi, K., Onodera, J., Jordan, R.W., Suto, I.: Living diatoms in the vicinity
- of the North Pole, summer 2004, Micropaleontol. 55, 137-170, 2009.
- Killworth, P.D.: Deep convection in the World Ocean, Rev. Geophys., 21, 1-26,
 doi:10.1029/RG021i001p00001, 1983.
- 32 Knies, J., Pathirana, I., Cabedo-Sanz, P., Banica, A., Fabian, K., Rasmussen, T.L., Forwick,
- 33 M., Belt, S.: Sea-ice dynamics in an Arctic coastal polynya during the past 6500 years,
- 34 Arktos, 3, 1, https://doi.org/10.1007/s41063-016-0027-y, 2017.

- 1 Korsun, S., Hald, M.: Modern benthic Foraminifera off Novaya Zemlya tidewater glaciers,
- 2 Arctic and Alpine Research, 30, 61-77, https://doi.org/10.1080/00040851.1998.12002876,
- **3** 1998.
- 4 Korsun, S., Pogodina, I.A., Forman, S.L., Lubinski, D.J.: Recent foraminifera in glaciomarine
- 5 sediments from three arctic fjords of Novaja Zemlja and Svalbard, Polar Res., 14, 15 31,
- 6 https://doi.org/10.1111/j.1751-8369.1995.tb00707.x, 1995.
- 7 Lejzerowicz, F., Esling, P., Majewski, W., Szczuciński, W., Decelle, J., Obadia, C., Martinez
- 8 Arbizu, P., Pawlowski, J.: Ancient DNA complements microfossil record in deep-sea
- 9 subsurface sediments. Biol. Lett., 9, 20130283, https://doi.org/10.1098/rsbl.2013.0283, 2013.
- 10 Lecroq B., Lejzerowicz F., Bachar D., Christen R., Esling P., Baerlocher L., Østerås M.,
- 11 Frinelli L., Pawlowski J.: Ultra-deep sequencing of foraminiferal microbarcodes unveils
- 12 hidden richness in deep-sea sediments, PNAS, 108:13177-82,
 13 https://doi.org/10.1073/pnas.1018426108, 2011.
- Loeng, H.: Features of the physical oceanographic conditions of the Barents Sea. Polar
 Res.,10, 5-18, https://doi.org./10.3402/polar.v10i1.6723, 1991.
- 16 Lubinski, D.J., Polyak, L., Forman, S.L.: Freshwater and Atlantic water inflows to the deep
- 17 northern Barents and Kara seas since ca 13¹⁴C ka: foraminifera and stable isotopes, Quat. Sci.
- 18 Rev., 20, 1851-1879, https://doi.org/10.1016/S0277-3791(01)00016-6, 2001.
- 19 Lydersen, C., Nøst, O., Lovell, P., McConell, B., Gammelsrød, T., Hunter, C., Fedak, M.,
- 20 Kovacs, K.: Salinity and temperature structure of a freezing Arctic fjord monitored by white
- 21 whales (Delphinapterus leucas), Geophys. Res. Lett., 29, 2119,
 22 https://doi.org/10.1029/2002GL015462, 2002.
- 23 Łacka, M., Zajączkowski, M., Forwick, M., Szczuciński, W.: Late Weichselian and Holocene
- 24 palaeoceanography of Storfjordrenna, southern Svalbard, Clim. Past, 11, 587-603,
- 25 https://doi.org/10.5194/cp-11-587-2015, 2015.
- 26 Łącka, M., Cao, M., Rosell-Melé, A., Pawłowska, J., Kucharska, M., Forwick, M.,
- 27 Zajączkowski, M.: Postglacial paleoceanography of the western Barents Sea: Implications for
- alkenone-based sea surface temperatures and primary productivity, Quat. Sci. Rev., 224,
- 29 105937, https://doi.org/10.1016/j.quascirev.2019.105973, 2019.
- 30 Mahé, F., Rognes T., Quince C., de Vargas, C., Dunthorn, M.: Swarm: robust and fast
- clustering method for amplicon-based studies, Peer J, 2, e593, doi: 10.7717/peerj.593, 2014.
- 32 Majewski, W., Pawłowski, J., Zajączkowski, M.: Monothalamous foraminifera from West
- 33 Spitsbergen fjords: a brief overview, Polish Polar Res., 26(4), 269-285, 2005.

- Majewski, W., Szczuciński, W., Zajączkowski, M.: Interactions of Arctic and Atlantic water masses and associated environmental changes during the last millennium, Hornsund (SW
- 3 Svalbard). Boreas, 38, 529-544, https://doi.org/10.1111/j.1502-3885.2009.00091.x, 2009.
- 4 Mangerud, J., Bondevik, S., Gulliksen, S., Hufthammer, A.K., Høseter, T.: Marine ¹⁴C
- 5 reservoir ages for 19th century whales and mollusks from the North Atlantic, Quat. Sci. Rev.,
- 6 25, 3228-3245, https://doi.org/10.1016/j.quascirev.2006.03.010, 2006.
- Martrat, B., Grimalt, J.O., Villanueva, J., van Kreveld, S., Sarntheim, M.: Climatic
 dependence of the organic matter contributions in the north eastern Norwegian Sea over the
 last 15,000 years, Org. Geochem., 34, 1057-1070, https://doi.org/10.1016/S01466380(03)00084-6, 2003.
- Müller, J., Werner, K., Stein, R., Fahl, K., Moros, M., Jansen, E.: Holocene cooling
 culminates in sea ice oscillations in Fram Strait, Quat. Sci. Rev., 47, 1–14,
 https://doi.org/10.1016/j.quascirev.2012.04.024, 2012.
- Moros, M., Emeis, K., Risebrobakken, B., Snowball, I., Kuijpers, A., McManus, J., Jansen,
 E.: Sea surface temperatures and ice rafting in the Holocene North Atlantic: climate
 influences on northern Europe and Greenland, Quat. Sci. Rev., 23, 2113-2126,
 https://doi.org/10.1016/j.quascirev.2004.08.003, 2004.
- 18 Nilsen, F., Cottier, F., Skogseth, R., Mattson, S.: Fjord-shelf exchanges controlled by ice and
- 19 brine production: The interannual variation of Atlantic Water in Isfjorden, Svalbard, Cont.
- 20 Shelf Res., 28, 1838-1853, https://doi.org/10.1016/j.csr.2008.04.015, 2008.
- 21 Pawłowska, J., Lejzerowicz, F., Esling, P., Szczuciński, W., Zajączkowski, M., Pawlowski, J.:
- Ancient DNA sheds new light on the Svalbard foraminiferal fossil record from the last millennium, Geobiology, 12, 277-288, https://doi.org/10.1111/gbi.12087, 2014.
- Pawłowska, J., Zajączkowski, M., Łącka, M., Lejzerowicz, F., Esling, P., Pawlowski, J.: 24 Palaeoceanographic changes in Hornsund Fjord (Spitsbergen, Svalbard) over the last 25 26 millennium: new insights from ancient DNA, Clim. Past, 12, 1459-1472, https://doi.org/10.5194/cp-12-1459-2016, 2016. 27
- 28 Pawlowski J., Holzmann M., Berney C., Fahrni J., Cedhagen T., Bowser S.S.: Phylogeny of
- 29 alogromiid Foraminifera inferred from SSU rRNA gene sequences. J. Foramin. Res., 32, 334-
- 30 343, https://doi.org/10.2113/0320334, 2002.
- 31 Pawlowski, J., Esling, P., Lejzerowicz, F., Cedhagen, T., Wildings, T.A.: Environmental
- 32 monitoring through protest next-generation sequencing metabarcoding: assessing the impact

- 1 of fish farming on benthic foraminifera communities, Mol. Ecol. Res., 14, 1129-1140, doi:
- 2 10.1111/1755-0998.12261, 2014.
- 3 Perner, K., Moros, M., Lloyd, J.M., Jansen, E., Stein, R.: Mid to late Holocene strengthening
- 4 of the East Greenland Current linked to warm subsurface Atlantic water, Quat. Sci. Rev., 129,
- 5 296-307, https://doi.org/10.1016/j.quascirev.2015.10.007, 2015.
- 6 Piechura, J.: Dense bottom waters in Storfjord and Storfjordrenna, Oceanologia, 38, 285-292,
 7 1996.
- 8 Polyakov, I. V., Pnyushkov, A.V., Alkire, M.B., Ashik, I.M., Baumann, T.M., Carmack, E.C.,
- 9 Goszczko, I., Guthrie, J., Ivanov, V.V., Kanzow, T.T., Greater role for Atlantic inflows on
- sea-ice loss in the Eurasian Basin of the Arctic Ocean, Science, eaai8204,
 https://doi.org/10.1126/science.aai8204, 2017.
- 12 Quadfasel, D., Rudels, B., Kurz, K.: Outflow of dense water from a Svalbard fjord into the
- 13 Fram Strait, Deep Sea Res., 35, 1143-1150, https://doi.org/10.1016/0198-0149(88)90006-4,
- 14 1988.
- 15 Rasmussen, T.L., Forwick, M., Mackensen, A.: Reconstruction of inflow of Atlantic Water to
- Isfjorden, Svalbard during the Holocene: Correlation to climate and seasonality, Mar.
 Micropaleontol., 94-95, 80-90, https://doi.org/10.1016/j.marmicro.2012.06.008, 2012.
- 18 Rasmussen, T. L., Thomsen, E.: Brine formation in relation to climate changes and ice retreat
- 19 during the last 15,000 years in Storfjorden, Svalbard, 76-78°N, Paleoceanography, 29, 911–
- 20 929, https://doi.org/10.1002/2014PA002643, 2014.
- 21 Rasmussen, T.L., Thomsen, E., Skirbekk, K., Ślubowska-Woldengen, M., Klitgaard
- 22 Kristensen, D., Koç, N.: Spatial and temporal distribution of Holocene temperature maxima in
- 23 the northern Nordic seas: interplay of Atlantic-, Arctic- and polar water masses, Quat. Sci.
- 24 Rev., 92, 280-291, https://doi.org/10.1016/j.quascirev.2013.10.034, 2014.
- 25 Rasmussen, T.L., Thomsen, E.: Palaeoceanographic development in Storfjorden, Svalbard,
- 26 during the deglaciation and Holocene: evidence from benthic foraminiferal records. Boreas,
- 27 44, 24–44, https://doi.org/10.1111/bor.12098, 2015.
- 28 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., van der
- 29 Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0-50,000 Years Cal
- 30 BP. Radiocarbon, 55(4), 1869-1887, https://doi.org/10.2458/azu_js_rc.55.16947, 2013.
- 31 Rigual-Hernández, A.S., Colmenero-Hidalgo, E., Martrat, B., Bárcena, M.A., de Vernal, A.,
- 32 Flores, J.A., Grimalt, J.O., Henry, M., Lucchi, R.G.: Svalabrd ice-sheet decay after the Last
- 33 Glacial Maximum: New insights micropalaeontological and organic biomarker

- 1 paleoceanographical reconstructions, Palaeogeogr., Palaeoclimatol., Palaeoecol., 465, 225-
- 2 236, http://dx.doi.org/10.1016/j.palaeo.2016.10.034, 2017.
- 3 Risebrobakken, B., Moros, M., Ivanova, E.V., Chistyakova, N., Rosenberg, R.: Climate and
- 4 oceanographic variability in the SW Barents Sea during the Holocene, The Holocene, 20,
- 5 609-612, https://doi.org/10.1177/0959683609356586, 2010.
- 6 Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F.: VSEARCH: a versatile open source
- 7 tool for metagenomics. Peer J, 4, e2584, https://doi.org/10.7717/peerj.2584, 2016.
- 8 Różańska, M., Poulin, M., Gosselin, M.: Protist entrapment in newly formed sea ice in the
- 9 Coastal Arctic Ocean, J. Mar. Sys., 74, 887-901,
 10 https://doi.org/10.1016/j.jmarsys.2007.11.009, 2008.
- 11 Rudels, B., Korhonen, M., Schauer, U., Pisarev, S., Rabe, B., Wisotzki, A.: Circulation and
- 12 transformation of Atlantic water in the Eurasian Basin and the contribution of the Fram Strait
- 13 inflow branch to the Arctic Ocean heat budget, Prog. Oceanogr., 132, 128-152,
- 14 https://doi.org/10.1016/j.pocean.2014.04.003, 2015.
- 15 Sarnthein, M., Van Kreveld, S., Erlenkeuser, H., Grootes, P.M., Kucera, M., Pflaumann, U.,
- 16 Schulz, M.: Centennial-to-millennial-scale periodicities of Holocene climate and sediment
- 17 injections off the western Barents shelf, 75°N, Boreas, 32, 447-461,
 18 https://doi.org/10.1111/j.1502-3885.2003.tb01227.x, 2003.
- 19 Sabbattini, A., Morigi, C., Negri, A., Gooday, A.J.: Distribution and biodiversity of stained
- 20 Monothalamous foraminifera from Tempelfjord, Svalbard, J. Foramin. Res., 37, 93-106,
- 21 https://doi.org/ 10.2113/gsjfr.37.2.93, 2007.
- 22 Sabbattini, A., Bonatto, S., Bianchelli, S., Pusceddu, A., Danovaro, R., Negri A.:
- 23 Foraminiferal assemblages and trophic state in coastal sediments of the Adriatic Sea, J. Mar.
- 24 Syst., 105, 163-174, https://doi.org/10.1016/j.jmarsys.2012.07.009, 2012.
- 25 Sabbattini, A., Nardelli M.P., Morigi C., Negri, A.: Contribution of soft-shelled
- 26 monothalamous taxa to foraminiferal assemblages in the Adriatic Sea, Acta Protozool., 52,
- 27 181-192, https://doi.org/10.4467/16890027AP.13.0016.1113, 2013.
- 28 Skogseth, R., Haugan, P.M., Haarpaintner, J.: Ice and brine production in Storfjorden from
- 29 four winters of satellite and in situ observations and modeling, J. Geophys. Res., 109,
- 30 C10008, https://doi.org/10.1029/2004JC002384, 2004.
- 31 Skogseth, R., Haughan, P.M., Jakobsson, M.: Watermass transformations in Storfjorden,
- 32 Cont. Shelf Res., 25, 667-695, https://doi.org/10.1016/j.csr.2004.10.005, 2005.

- Skogseth, R., Sandvik, A. D., Asplin, L.: Wind and tidal forcing on the meso-scale
 circulation in Storfjorden, Svalbard, Cont. Shelf Res., 27, 208-227,
 https://doi.org/10.1016/j.csr.2006.10.001, 2007.
- 4 Stuiver, M., Reimer, P.J.: Extended ¹⁴C database and revised CALIB 3.0 ¹⁴C age calibration
- 5 program, Radiocarbon, 35, 215-230, 1993.
- 6 Ślubowska-Woldengen, M., Rasmussen, T.L., Koç, N., Klitgaard-Kristensen, D., Nilsen, F.,
- 7 Solheim, A.: Advection of Atlantic Water to the western and northern Svalbard shelf since
- 8 17,500 cal yr BP. Quat. Sci. Rev., 26 463-478,
 9 https://doi.org/10.1016/j.quascirev.2006.09.009, 2007.
- Taberlet, P., Bonin, A., Zinger, L., Coissac, E.: Enviroenmntal DNA: For biodiversity
 research and monitoring, Oxford University Press,
 https://doi.org/10.1093/oso/9780198767220.001.0001, 2018.
- 13 Telesiński, M.M., Przytarska, J.E., Sternal, B., Forwick, M., Szczuciński, W., Łacka, M.,
- 14 Zajączkowski, M.: Palaeoceanographic evolution of the SW Svalbard shelf over the last
- 15 14 000 years, Boreas, 47, 410-422, https://doi.org/10.1111/bor.12282, 2018.
- Thomsen, P.F., Willerslev, E.: Environmental DNA An emerging tool in conservation for
 monitoring past and present biodiversity, Biol. Conserv., 183, 4-18,
 https://doi.org/10.1016/j.biocon.2014.11.019, 2015.
- Werner, K., Spielhagen, R.F., Bauch, D., Hass, H., Kandiano, E.S., Zamelczyk, K.: Atlantic
 Water advection to the eastern Fram Strait multiproxy evidence for late Holocene
 variability. Palaeogeogr. Palaeoclimatol. Palaeoecol., 308, 264-276,
 https://doi.org/10.1016/j.palaeo.2011.05.030, 2011.
- 23 Werner, K., Spielhagen, R.F., Bauch, D., Hass, H.C., Kandiano, E.: Atlantic Water advection
- versus sea-ice advances in the eastern Fram Strait during the last 9 ka: Multiproxy evidence
- 25 for a two-phase Holocene, Paleoceanogr., 28, 283-295, doi:10.1002/palo.20028, 2013.
- Winkelmann, D., Knies, J.: Recent distribution and accumulation of organic carbon on the continental margin west off Spitsbergen. Geochem. Geophys. Geosyst., 6, Q09012,
- 28 https://doi.org/10.1029/2005GC000916, 2005.
- 29 Zimmermann, H.H., Stoof-Leichsering, K.R., Kruse, S., Müller, J., Stein, R., Tiedemann, R.,
- 30 Herzschuh, U.: Changes in the composition of marine and sea-ice diatoms derived from
- sedimentary ancient DNA of the eastern Fram Strait over the past 30,000 years. Ocean Sci.
- 32 Discuss., https://doi.org/10.5194/os-2019-113, 2019.
- 33

34 Figures captions

- Figure 1: The modern oceanography of the study area (A) and the location of the studied core
 ST_1.5 (B) and the other cores discussed in this paper (A,B). Abbreviations of the main
- 3 surface currents: WSC West Spitsbergen Current, NAC North Atlantic Current, NCaC –
- 4 North Cape Current, ESC East Spitsbergen Current, BIC Bear Island Current, CC –
- 5 Coastal Current.

Figure 2: Temperature and salinity profile from the core location. Temperature is marked
with a dashed line, and salinity is marked with a black line. Abbreviations: AW – Atlantic
Water, TAW – Transformed Atlantic Water, BSW – Brine-enriched Shelf Water.

9 Figure 3: Age-depth model of the ST_1.5 core. The gray silhouettes show the probability

11 dates used for the age model. The dotted line shows the age-depth model derived from linear

distribution of the calendar dates that were obtained by the calibration of the individual ¹⁴C

12 interpolation between the dates.

13 Figure 4: Sedimentological and micropaleontological data plotted versus age. The sediment

14 accumulation rate (SAR), mean grain size of the 0-63- μ m fraction, ice-rafted debris (IRD)

- 15 flux and number of grains per gram of sediment, oxygen (δ^{18} O) and carbon (δ^{13} C) stable
- isotopes in benthic foraminiferal tests, and the flux and abundance of foraminifera arepresented.
- Figure 5: The absolute abundance (expressed as the number of individuals per gram of drysediment) and the percentage of the dominant benthic foraminifera.
- 20 Figure 6: The dominant components of the monothalamous assemblages. The abundance is
- 21 expressed as the percentage of the monothalamous sequences and the most abundantly

sequenced taxa are presented. The trend (2-point average) is indicated with a dashed line.

Figure 7: The percentage share of certain clades in the allogromiid sequences.

Figure 8: The percentage of sequences of dominant diatom taxa vs. time. The trend (2-point

25 average) is indicated with the dashed line.

26

10

- 27 Table captions
- **Table 1:** Raw and calibrated AMS¹⁴C dates used in the age model.



- 1
- 2 Figure 1: The modern oceanography of the study area (A) and the location of the studied core ST_1.5 (B) and
- 3 the other cores discussed in this paper (A,B). Abbreviations of main surface currents: WSC West Spitsbergen
- 4 Current, NAC North Atlantic Current, NCaC North Cape Current, ESC East Spitsbergen Current, BIC –
- 5 Bear Island Current, CC Coastal Current.
- 6



Figure 2: Temperature and salinity profile from the sampling station. Temperature is marked with a dashed line, and salinity is marked with a black line. Abbreviations: AW – Atlantic Water, TAW – Transformed Atlantic Water, BSW – Brine-enriched Shelf Water.







10 age-depth model derived from a linear interpolation between the dates.



Figure 4: Sedimentological and micropaleontological data plotted versus age. The sediment accumulation rate (SAR), mean grain size of the 0-63- μ m fraction, ice-rafted debris (IRD) flux and number of grains per gram of sediment, oxygen (δ^{18} O) and carbon (δ^{13} C) stable isotopes in benthic foraminiferal tests, and the flux and abundance of foraminifera are presented.



Figure 5: The absolute abundance (expressed as the number of individuals per gram of dry sediment) and the
 percentage of the dominant benthic foraminifera.





[% sedneuces]

Figure 6: The dominant components of the monothalamous assemblages. The abundance is expressed as the
 percentage of the monothalamous sequences and the most abundantly sequenced taxa are presented. The trend
 (2-point average) is indicated with a dashed line.



2 Figure 7: The percentage share of certain clades in the allogromiid sequences.



Figure 8: The percentage of sequences of dominant diatom taxa vs. time. The trend (2-point average) is
indicated with a dashed line.

Core	Material	Raw AMS ¹⁴ C	Cal. a	Cal. a BP
depth			$BP\pm 2\sigma$	used in age
[cm]				model
2.5	Nuculana pernula (B)	$107.38\pm0.33\ pMC$	-	-
5.5	Yoldiella lenticula (B)	$290\pm30 \text{ BP}$	-	-
14.5	Turitella erosa (B)	$2020\pm30\;BP$	1356-1555	1500
43.5	Yoldiella solituda (B)	$3010\pm50~BP$	2484-2787	2700
46.5	Nonionellina labradorica (${f F}$)	4490 ± 40 BP	4400-4701	4500
52.5	Yoldiella lenticula (B)	$7545\pm35~\mathrm{BP}$	7803-7989	7890

Table 1: Raw and calibrated AMS¹⁴C dates used in the age model. B stands for bivlave shells, while F stands for
 benthic foraminifera tests.