### **Responses to Reviewers Comments**

We would like to take the opportunity to thank the three reviewers for highlighting some very important generalisations in our interpretations of changing diatoms, guilds and beta-diversity. We have taken on board all their comments, but this has necessitated some substantial changes to the manuscript.

- We have moved away from discussing beta-diversity and resources in general because (i) beta-diversity has so many different meaning to different disciplines, and (ii) the reviewers were right to point out that talking about resources in general in a palaeo paper was too vague.
- 2. We have restructured the discussion somewhat, where we now (i) interpret the major changes to species in an environmental context; (ii) discuss what changes to the diatom guilds might mean, before (iii) finally using this information to inform changes in compositional turnover (which is one form of beta-diversity).
- 3. In the original version of the manuscript we had undertaken PCA of the genera, but barely talked about these in the results. We have decided to remove the PCA of the genera from the revised version.
- 4. However, we have included TOC as an extra proxy (as all reviewers made a comment on only looking at diatoms), and this has been very useful in interpreting carbon sequestration in the lake during the neoglacial.

## Other general changes:

- Aifeng Zhou is now a co-corresponding author along with Anson Mackay
- Keywords are more reflective of the revised manuscript

### **Reviewer 1:**

Reviewer 1 has two points of general criticism. First that we should develop the assumption that temporal changes in diatoms are linked to lake-catchment resource availability, which in itself is related to climate variability. Second, that we show no other palaeolimnological data such as TOC.

The first set of comments are really interesting, and have highlighted us to new literature by Rimet et al. 2019, and Passy & Larson 2019, including the role that turbulence may play in influencing alpine diatom communities. We have taken on board other factors that may influence diatom communities in an alpine setting throughout the text. Rather than providing a new suggested paragraph on temporal beta diversity (linked to Korhonen et al. 2010), we have provided greater clarification on the form of beta-diversity used in this study. In fact, as now stated in the text, we have chosen to discuss our data in the form of compositional turnover, to avoid the confusion that the term beta-diversity can now have, given all its different meanings (*sens*u Anderson et al. 2011).

### Specific comments have all been addressed:

Page 2, line 56: for consistency with the rest of the manuscript, use Medieval Climatic Anomaly (not Optimum)

• Done

Page 3, line 76: delete "however"

• Done

Page 3, lines 82-83: rephrase such as:: : : but provide their habitats to many iconic species that are also vulnerable

• Done, P3, Line 73-74

Page 3, line 86: : : : where long-term historical and/or instrumental records are

• Done

Page 4, line 102: add a reference about the driver of the NH cooling

• Done. Marcott et al. 2013, P4, Line 107

Page 4, line 108: correct spelling of climatic

• Done

Page 4, line 115: spell in full Asian Summer Monsoon (ASM) as it here mentioned for the first time in the text

• Done

Page 5, lines 127-128: from where this big assumption comes from? Is diatom diversity always linked to resources? Also need to be more specific about the resources: is it just nutrients, light? What about the other factor: habitat availability, grazing by invertebrates?

• Response: we have deleted the sentence related to this assumption, as the reviewer is right, it is a very big assumption, and we have not demonstrated its validity.

Page 5, line 144: (in legend for Fig. 1): change central Asia by central China.

• Done

Are subalpine meadow (as written in the text, line 141) and tundra the same type of vegetation?

• Response: No they are different; we have deleted "tundra" from the legend of Figure 1

Page 6, line 163: From what it is written I assume that the core was sliced at 1-cm intervals: please specify and also give the actual number of samples analysed

• Done

Page 7, line 167: it's divinylbenzene (DVB), not bivinyl

• Done

Page 7, line 168: the concentration unit should be in nb of spheres per volume, ml or cm3 but not cm2, right?

• Correct – this should have been spheres/cm<sup>3</sup>

Page 7, line 171: add the registered symbol <sup>®</sup> after Naphrax, Zeiss Axiostar Plus

• Done

Page 7, line 184: siltation is also an important factor that promote motile benthic diatoms over attached forms (e.g. Battegazzore et al. 2004; Dickman et al. 2005).

• Response: We have added in reference to siltation, using the Battegazzore reference: P8, Line 196

Page 7, line 192- 193: a log-linear contrast PCA. I'm not familiar with this technique, please explain in one sentence how different it is from "conventional" PCA and what are the advantages

• Response: Into P8, Lines 205-206, we have added in the clause, which also necessitates a new reference: "(appropriate for closed relative abundance data (Lotter and Birks, 1993)) was undertaken..."

Page 8, line 207: give a reference for the C2 program

• Done: added in Juggins 2014)

Page 10, line 228: Puncticulata is a redundant generic name. Use Lindavia (see Nakov et al. 2015).

• Done

Page 10, line 250: you forgot a few words: : : :while high profile and motile diatoms are persistently

• Done

Page 11, line 256 (legend of Fig. 3): rephrase such as: Only diatoms with abundance greater than 3% are shown

• Done

Page 12, Figure 3: It would be useful to group the diatom species according to their ecological guild.

• Done

In Fig 3, Correct spelling of Staurosira construens, change Puncticulata to Lindavia

• Done

Page 14, Fig.5: correct spelling for "centered"

• Done

Page 15, line 303: : : : dominated by species belonging to the Fragilariaceae

• Done

Page 15, line 304: replace "growing well" by "dominating the assemblages"

• Done

Page 15, line 306: on the abundance of fragilarioids:

• Done

Page 15, line 307: add "(Russia)" after "Eastern Sayan mountains"

• Done

Page 15, lines 310-319: This explanation about the decline of Stauroforma is not entirely convincing. First the catchment of the lake appears treeless as it is located above the treeline (as shown on Fig. 1). In that case, how the mentioned shift from a deciduous-conifer forest to a steppe forest would have impacted the soils of the lake catchment if no forest was there in the first place?

• Agreed: we have removed this sentence

Then you need to be more specific than just saying "related to the provision of resources linked to catchment changes around the lake". Could the reduced input of DOC may be linked to deeper, longer permafrost instead of shift in catchment vegetation? (as the authors have suggested themselves on lines 423-424 but for the upper part of the record).

• Response: we have tried to be more nuanced here, and have made the observation that highest Stauroforma and TOC occur together, so its presence may be linked to the lake being dystrophic at the time. See new text in Section 4.1, P 15, Lines 329-338

Page 16, line 329: use replaced instead of replacing

• Done

Page 16, line 341: Why are you comparing your results with winter temperature? Summer temperature would be more relevant for benthic diatoms. Also note that summer temperature may have been relatively high in Northern China during the "Roman Warm Period" as suggested by the pollen reconstruction by Stebich et al. (2015).

• Response: we had not seen the Stebich et al. 2015 article, so as suggested we have replotted their mean temperature of the warmest month in our multi-archive Figure 6, and included relevant interpretation in the text. (e.g. see P15, Lines 334-335)

Page 17, legend of Fig. 6. Replace "Chinese" by "central China".

• Done

Fig 6: The number for each curves are wrong, should be 6a, 6b, 6c, etc: : : instead of 7a, 7b, 7c:

• Done

Page 17, line 364: correct spelling of climatic

• Done

Pages 18-19: For consistency and to make it easy to found the features that are discussed on the diagrams, please give all the ages in cal yrs BP, not just in CE calendar.

• Done

Page 19, line 419: replace "important" by abundant Page 20,

• Done

Page 20, line 427: give a reference about the ecology of Denticula subtilis

• Done: we have actually re-focused the ecology here, as other studies have found it to be linked to elevated conductivity, e.g. Antoniades et al. (2005): see P22, Lines 463-467

Page 20, line 446: is a bit confusing: was aridity declining or on the contrary increasing, as the curve for reconstructed precipitation (Fig 6d) would suggest?

• Done – sorry, yes aridity was increasing. We have corrected the text

Page 20, line 448: delete "of the" after around the time

• Done

Page 20, lines: 448-450: what about the 2-sample shift from Stauroforma to Lindavia+humidophila?

• Response: We have added in extra discussion at the end related to this period, P 23 Lines 488-492.

Page 21, line 465: rephrase such as: Increased summer precipitation during the MCA as inferred from nearby records (here add references!) resulted in increased diatom fluxes

• Done

General comment on adding in more data.

• Response: unfortunately we do not have any geochemical data, but we do have measurements for total organic carbon, and these have now been added to methods, results, and implications in discussion.

### **References used:**

Antoniades, D., Douglas, M.S. and Smol, J.P., 2005. Benthic diatom autecology and inference model development from the Canadian high arctic Archipelago 1. *Journal of Phycology*, *41*(1), pp.30-45.

Anderson, M.J., Crist, T.O., Chase, J.M., Vellend, M., Inouye, B.D., Freestone, A.L., Sanders, N.J., Cornell, H.V., Comita, L.S., Davies, K.F. and Harrison, S.P., 2011. Navigating the multiple meanings of  $\beta$  diversity: a roadmap for the practicing ecologist. *Ecology letters*, 14(1), pp.19-28.

Juggins S (2014) C2 version 1.7.7: software for ecological and palaeoecological data. University of Newcastle

Lotter, A.F., Birks, H.J.B., 1993. The impact of the Laacher See tephra on terrestrial and aquatic ecosystems in the Black Forest, southern Germany. Journal of Quaternary Science 8, 263-276

We would like to take the opportunity to thank the three reviewers for highlighting some very important generalisations in our interpretations of changing diatoms, guilds and beta-diversity. We have taken on board all their comments, but this has necessitated some substantial changes to the manuscript.

- 1. We have moved away from discussing beta-diversity and resources in general because (i) beta-diversity has so many different meaning to different disciplines, and (ii) the reviewers were right to point out that talking about resources in general in a palaeo paper was too vague.
- 2. We have restructured the discussion somewhat, where we now (i) interpret the major changes to species in an environmental context; (ii) discuss what changes to the diatom guilds might mean, before (iii) finally using this information to inform changes in compositional turnover (which is one form of beta-diversity).
- 3. In the original version of the manuscript we had undertaken PCA of the genera, but barely talked about these in the results. We have decided to remove the PCA of the genera from the revised version.
- 4. However, we have included TOC as an extra proxy (as all reviewers made a comment on only looking at diatoms), and this has been very useful in interpreting carbon sequestration in the lake during the neoglacial.

### Other general changes:

- Aifeng Zhou is now a co-corresponding author along with Anson Mackay
- Keywords are more reflective of the revised manuscript

### **Reviewer 2 comments**

Reviewer 2 found some of the interpretation "a bit vague" and lacked a "multiproxy approach". These two broad comments are similar to Reviewer 1, and we have addressed both. The former comment has been tackled by being more explicit what we mean when we talk about diatom responses to changing resources – see comments to R1 above, and changes made throughout the text, highlighted under track changes. And the latter comment has been addressed now by adding in TOC data (see Fig 5c).

### Specific comments:

Line 70: Describe what is beta-diversity.

- Response: As identified by R1, we have added in further clarification on beta diversity (eg P3, lines 81-84).
- Because the term now means different things to different groups of scientists, we refer to the data more specifically as compositional turnover (P3, Lines 88-90)

Line 77: What do you mean by elevation-dependent warming?

• Response: We have provided greater explanation: , "i.e. the amplification of warming at higher altitudes" P3, Lines 69-70

Line 97: Don't really like the term global warmth...is there another way to say this?

• Done: have changed to "globally warmer temperatures".

Line 99: "less monotonic" – is that the right term? Thought monotonic was used to describe sequences...?

 Done – we have simplified this sentence to read "The extent of cooling varied regionally, being most pronounced in the extra-tropical northern hemisphere (Marcott et al. 2013)." (P4, Lines 105-106)

Line 110: Define LIA

• Done

Line 115: Define ASM

• Done

Line 124: What do you mean by ecological guilds?

• Response: This is defined in the methods in Section 2.2, P8-9, Lines 188-198

132: On what account is it a biodiversity hotspot? Describe

• Response: We have added in more detail: "...as a refuge for both Tertiary plants ( Zhang et al. 2017) and the vulnerable Giant Panda (Fan et al. 2014). (P5, Lines 134-136)

Line 141: Describe what you mean by Larix forest - what species? what climate/soil conditions do they require? Etc

• Response: we have added in detail – P6, Lines 144-146:

Line 142: More detail on the lake itself would be good. Has the lake ever been studied before? Any more bathymetry of the lake? What is its riparian/shoreline vegetation? Is it in a small, steep catchment? Does it have macrophytes in its shallow waters? Is it dimictic/polymictic etc? Or is knowledge extremely limited here?

Response: we actually have very little information on the lake, due to its remoteness. We have included what extra info that we have. We observed no shoreline macrophytes (it was winter when the lake was visited, and frozen over), but we measured surface water pH at 6.84, using a YSI ProDSS multiparameter water quality meter. (P6, Lines 146-150)

Line 155: Which country is Beta Analytic?

• Done: USA

Line 224: Don't quite follow the sentence. Do you mean 120 of the species were rare?

• Done: we have rephrased

Line 316: State specifically how and why the change in catchment vegetation would have altered delivery of allochthonous material? Increased or decreased?

• Response: We agree with the statement that more detail needs to be given as to whether allochthonous material increased or decreased with shift in vegetation. But as highlighted in R1 above, the relevance for this for YHC site is not clear – we have deleted this statement therefore

Line 321: Not a clear sentence. Did limiting resources become more limited? Or more resources became more limited?

• Response: We have deleted this sentence

Line 323: "deterministic processes become more important" - more important than what? What do you mean by deterministic processes?

• Response: Here we have clarified that deterministic processes become more important than stochastic ones. Deterministic processes here are ones related to niche selection. See P17, Lines 374-379

Line 330: What "resources"? Be specific.

• Response: we have rewritten and reorganized the start of the Discussion

Line 342: Again, be more specific about what "resources" you mean and why and how this increased the prevalence of high profile diatoms.

• Response: we have rewritten and reorganized the start of the Discussion

Line 387: Is this a worthy comparison if the soils and altitudinal differences are dissimilar? If so, why? This could be a good opportunity to highlight the importance of lake and catchment characteristics is shaping the response of lake ecology to large-scale spatial and temporal drivers of change.

• Response: we have deleted comparison to this other site, as it was a bit random, and given catchments differences, may well not have been a robust comparison to make

Line 395-6: Don't understand what you mean by "several interacting, time-transgressive forcings", please be clear and describe what you mean.

Response: we have added information: "..., including reduced solar activity during the late 17<sup>th</sup> century (Shindell et al. 2001), increased volcanic activity during the early 19<sup>th</sup> century (Brönnimann et al. 2019) and reduced Gulf Stream flow (Lund et al. 2006)." P20, Lines 430-433

Line 411: What is K+ a proxy for? Please describe for readers who may not know

• Response: we have added in extra text + reference - , "...given that potassium is likely sourced from central Asian dust via long-range transport to the Greenland ice sheet (Meeker and Mayewski 2002)" P21, Lines 450-452

Line 414: What do you mean by phenological records?

• Response: we have added "beginning of tree flowering" P20, Line 415

Line 421: What are these conditions? Be specific.

• We have deleted this sentence.

Line 422: How does cold conditions and extended ice cover decrease these planktonic diatoms? Be specific. Changes to turnover events? Nutrient supply?

• We have deleted this sentence

Line 428: What do you mean by "resources" here? Nutrients? Be specific.

• Response: we have actually re-focused the ecology of D. subtilis here, as other studies have found it to be linked to elevated conductivity, e.g. Antoniades et al. 2005): see P22, Line463-465.

Line 434: Again what do you mean by "resources" becoming more available? Be specific.

• Response: we have rephrased as "...increased competition among species (Larson et al. 2016)" P22, Lines 473-474

Line 436: Why is this important? Describe.

• We have deleted this sentence

Line 448: "time of the"..?

• Corrected

Line 453: What is "their"? Be clear

• We have deleted this sentence

### **References used:**

Antoniades, D., Douglas, M.S. and Smol, J.P., 2005. Benthic diatom autecology and inference model development from the Canadian high arctic Archipelago 1. *Journal of Phycology*, *41*(1), pp.30-45.

Brönnimann, S., Franke, J., Nussbaumer, S.U., Zumbühl, H. J., Steiner, D., Trachsel, M., Hegerl, G. C., Schurer, A., Worni, M., Malik, A., Flückiger, J.: Last phase of the Little Ice Age forced by volcanic eruptions. *Nat. Geosci. 12*, 650-656, 2019.

Meeker, L.D. and Mayewski, P.A., 2002. A 1400-year high-resolution record of atmospheric circulation over the North Atlantic and Asia. *The Holocene*, *12*(3), pp.257-266

Stebich, M., Rehfeld, K., Schlütz, F., Tarasov, P., Liu, J., Mingram, J.: Holocene vegetation and climate dynamics of NE China based on the pollen record from Sihailongwan Maar Lake. Quat. Sci. Rev., 124, 275-289, 2015.

We would like to take the opportunity to thank the three reviewers for highlighting some very important generalisations in our interpretations of changing diatoms, guilds and beta-diversity. We have taken on board all their comments, but this has necessitated some substantial changes to the manuscript.

- 1. We have moved away from discussing beta-diversity and resources in general because (i) beta-diversity has so many different meaning to different disciplines, and (ii) the reviewers were right to point out that talking about resources in general in a palaeo paper was too vague.
- 2. We have restructured the discussion somewhat, where we now (i) interpret the major changes to species in an environmental context; (ii) discuss what changes to the diatom guilds might mean, before (iii) finally using this information to inform changes in compositional turnover (which is one form of beta-diversity).
- 3. In the original version of the manuscript we had undertaken PCA of the genera, but barely talked about these in the results. We have decided to remove the PCA of the genera from the revised version.
- 4. However, we have included TOC as an extra proxy (as all reviewers made a comment on only looking at diatoms), and this has been very useful in interpreting carbon sequestration in the lake during the neoglacial.

### Other general changes:

- Aifeng Zhou is now a co-corresponding author along with Anson Mackay
- Keywords are more reflective of the revised manuscript

### **Reviewer 3 comments**

Reviewer 3 wanted more information on the lake and it's environmental context (see main comments to reviewer 2 above; what little information we do have, we have now included), and greater interpretation of the diatom changes themselves rather than relying on other papers. Given the responses to Reviewers 1 and 2 above, our interpretation of diatom changes is much tighter throughout the manuscript.

### **Specific comments:**

Line 48: "Here we take..."

• Response: This is still grammatically correct, so no need to change

Line 49: "Multidecadal variability" on what data is this based?

• Response - this phrase has been deleted

Line 49: The dating seems to have a hard water effect (which has not been clearly addressed) and only five AMS dates. Please clarify how you have come up with this multidecadal (55 years) variability

• Response: the reservoir effect in bulk organic dating is common in the lake sediments (Zhou et al., 2015). The Yuhuangchi lake is located in a high altitude area, where organic matter mainly comes from terrigenous organic matter input and the lake's authigenic algae. Terrigenous organic matter often contains the old soil carbon input through to the lake. We also found that has obvious reservoir effect in 10cm dating result. A common way to judge

radiocarbon reservoir effect is by using a linear or quadratic regression, and interpret the extrapolated surface of age as a radiocarbon reservoir effect (Hou 2012 QSR).

• Extra text and reference to Hou et al. 2012 have been provided in Section 2.1, P7, Lines 164-165

Line 55: "Important": : : what does this mean here? Important compared to what?

• We have removed this word

Line 61: Consider to add "productivity" to keywords

• Done

Line 77. Is there a word missing after "sensitive" e.g. "area" or "environment"?

• No - the word sensitive here refers to the regions themselves

Lines 82-83. "..but their habitats to many: ::" I don't really understand this sentence? Is this needed?

 Sentence has been corrected to "they provide habitats to many..." which is clearer; P2, Line 73-74

Lines 140 and 145: Is the elevation of the lake 3370m asl or 3365m asl?

• Good spot – the elevation has been corrected in the Figure legend to 3370m asl

Line 152. What kind of piston corer did you use? Please specify

• We actually used a homemade piston drill, similar to UWITEC's piston corer, which a plastic core tube with a piston inside and can be hammered into the sediment. So I think we can leave text as is

Line 153. ": : :from the central region of Lake Yuhuang Chi". Why from the central area, was it the deepest part of the lake? or the main sedimentation area? or had it the longest sediment sequence? Please clarify

• Response: Yes, this was the deepest part of lake and also with flat lake floor

Line 154. Why did you use bulk organic sediment for dating? Where there no terrestrial plant macrofossil in the sediment?

• Response: Yes, that is correct; no macrofossils were found in the sediments. We have made this clear in the text – Lines 159-160

Line 158. Its rather unclear how you came up with the 1340 year reservoir age effect. Please clarify this in sufficient detail and add the number of dates to line 156

• Response – both things done – see P6, lines 160-161

Lines 163-164. What does this sentence mean? How did you came up with a resolution of 55 years considering the uncertainties of the datings? What is the frequency of the diatom samples, 2mm, 5 mm, 1 cm: : :?

• Response: We have added in that diatom analyse were performed on alternate 1cm-thick samples (P7, line 174). The 55 year resolution is got from the age model but seeing as we are still in the methods section we have taken this clause out.

Line 165. Please add a reference(s) after ": : :standard procedures"

• Done: we have added in the reference to Battarbee et al. 2001

Lines 169-170. ": : : such that suitable concentrations could be calculated". What does this mean, please clarify

• Response: we have taken this phrase out; it is not necessary

Line 207. Please add a reference after "C2 Data Analysis Version 1.7.2

• Response: we have added Juggins (2014)

Line 214-> this is a bit odd paragraph as it contains only Table 1 and no written results?

• Response: good point; we have added in text to describe the lithology and introduce Table 1 (P9, Lines 226-230)

In Table 1 all +- ages are 30 years, is this correct?

• Yes

Lines 239-240. This sentence could be combined with the previous one as the information is almost the same

• Response: it is not clear what is being meant here so we have chosen to leave sets of info separate

Lines 256-257. The first two sentences of the Figure caption could be e.g "Diatoms with a relative abundance >3% in more than one sample are shown"

• Response: the figure legend has been edited

Line 263. Please add "Relative abundance (%) below the diatom stratigraphy figure

• Response: rather than doing this, we have added in (%) to the Figure legend, making clear what relative abundance refers to

Line 272. Please add "Relative abundance (%) below Figure 4

• Response: rather than doing this, we have added in (%) to the Figure legend, making clear what relative abundance refers to

Line 282. The beta-diversity value of 1.033 SD units is rather low and indicates relative subtle changes in the core (compare e.g. to values in Smol et al. (2005, PNAS))

• Response: I don't think that we need to respond to this just here, as it is a comment. But note from comments above re. beta-diversity, we have moved away from this concept to one that characterizes compositional change

Line 303. Fragilariaceae are also very common in high latitude lakes: : :you may add a reference after this statement

• Response: this is made explicit in the statement: "Fragilarioids are often opportunistic, dominating assemblages in alpine and arctic lakes with a short growing season and long periods of ice cover (Lotter and Bigler 2000)" (P14, Lines 322-324)

Line 329. H. Schmassmannii is not really replacing S. exiguiformis as its relative abundance varies between ca. 10- 15% during this period and S. exiguiformis decreases from ca. 50 to 40%...this does not really seem as a replacement, or? This is a too strong interpretation, please rephrase

• Response: this paragraph has been reinterpreted according to other reviewers comments as well; we no long have one species replacing another as such, but that changes in species are likely linked to changes in lake dystrophy (see P14, Lines 329-338)

Lines 337-338. "..resources stabilised or even increased slightly: : :" How can this be seen in the results? Which species or index is confirming or suggesting this intepretation? At least not the relative abundances of Figs 3 & 4

 Response: we have changed this from "resources stabilized" to "compositional turnover stabilised" - this is now data led rather than an interpretation of resources, which we acknowledge above may have been over-interpreting what these scores can tell us (17, Line 389)

Line 343. High profile G2-diatoms dominate the whole zones 1 & 2 with an relative occurrence around 60% (see Fig. 4)...so there are no real changes here unlike is suggested

• Response: we agree, and the text here has been modified substantially (P16, lines 348-371)

Lines 337-344. This whole paragraph consists of rather vague speculation without much of supporting data from the study. I would suggest you to re-evaluate your results and rewrite this accordingly

• Response: we agree, and the Discussion has been rewritten, moving away from focus on interpretation of resources, which was not clear from the data

Lines 348-353. Please change the numbers of sub-figures to be 6b, 6c, 6d etc

• Response: done

Line 358. Please add "Winter" before "temperature"

• Winter-temperature anomaly graph has now been removed

Lines 368-369. This is not clear in Figure 6c

• Winter-temperature anomaly graph has now been removed

Line 379. "Increasing diatom flux: ::"the relative abundance data does not really support this: : :could there be some problems in counting the diatom flux as the dating results are not very convincing and the sedimentation rate has a large impact on the flux values? Line 380-381. "Driven mainly by increasing P. bodanica": : :that is not clear when looking at Figure 3:

: P. bodanica seems to have a relative abundance around 20% before and after this period. Please check and re-think

• Response: we have removed the clause "including planktonic species", as reviewer is right, relative abundance data do not show major changes in the planktic guild at this time. Flux rates are of course tied to sedimentation rates, so we're not clear of the point being made. We think the dating is rather robust, with all errors and uncertainty given in the manuscript.

Line 382. Beta-diversity does not increase between 1500-800 cal yrs BP. Please re-phrase

• Response: this has been amended

Line 398. How come beta-diversity is almost zero although in Fig. 3 a relatively diverse diatom population still remains?

• Response: this is because beta-diversity means the change in species composition, not numbers of species per se. But as highlighted above, we have shifted our interpretation away from beta-diversity to compositional turnover which describes the data more accurately.

Lines 421-422. ": : :planktonic diatoms show a distinct decline during the LIA". Where can this be seen? In Fig 3. the P:L ratio does not decrease nor does the abundance of P. bodanica and also the planktonic guild is high in Fig 4? Please clarify

• Response: This statement has been removed from our re-write of the discussion

Lines 425-426. I doubt that the appearance of H. Schmassmannii is due to low water temperature? Do you have any data when this species blooms? Or what its temperature optima is? This species can be found also in high latitude lakes

• Response: the paper by Buczkó et al. 2015 reviews a number of different papers which state that this species is thermophilic, but also mentions at least one at the end which suggests that the species has been found during the younger Dryas at Krakenes, Norway. On balance, it is likely that this species does decline during cooler temperatures – so we have left this statement as it is, but we have augmented by bringing in possibility of other factors such as siltation also being important (P21, Line 463-464)

Line 429. ": : : the lake becoming more shallow due to increased aridity": : : The planktonic species P. bodanica still occurs with a relative abundance of ca. 20%, which could indicate deeper water as it needs turbulence to sustain in the water column. Could one explanation be that the water became clearer due to the frozen ground and less inwash of e.g. DOC/humic substances?

• Response: we suggest the the lake become more shallow due to the appearance of Denticula subtilis which usually occurs in shallow lakes with high conductivity. (P21, Lines 463-465)

Lines 456-472. This whole paragraph is rather general based on other publications instead of your own results. I think this whole paragraph should be re-written so that it reflects the results of this study

• Response: we have made greater reference to our own data in the conclusions

Line 548. Please add the reference for the program C2 here:

• Response: this has been done – Juggins 2014

Lines 659-664. Please switch the alphabetical order of these two references

• Response: done

Line 448. ": : : around the time of the did: : :" please re-phrase

• Response: done

### **References used**

Battarbee, R.W., Jones, V.J., Flower, R.J., Cameron, N.G., Bennion, H., Carvalho, L., Juggins, S., 2001. Diatoms. In: Smol, J.P., Birks, H.J.B., Last, W.M. (Eds.), Tracking Environmental Change Using Lake Sediments. Kluwer Academic Publishers, Dordrecht, pp. 155e201.

Hou, J., D'Andrea, W.J. and Liu, Z., 2012. The influence of 14C reservoir age on interpretation of paleolimnological records from the Tibetan Plateau. *Quaternary Science Reviews*, *48*, pp.67-79.

Juggins S (2014) C2 version 1.7.7: software for ecological and palaeoecological data. University of Newcastle

Zhou, A.-F.\*, He, Y.-X., Wu, D., Zhang, X.-N., Zhang, C., Liu, Z.-H., Yu, J.-Q., 2015. Changes in the Radiocarbon Reservoir Age in Lake Xingyun, Southwestern China during the Holocene. PLoS ONE 10(3): e0121532.

1 Neoglacial trends in diatom dynamics from a small alpine lake in the Qinling 2 Mountains of central China. 3 4 Bo Cheng<sup>1</sup>, Jennifer Adams<sup>2</sup>, Jianhui Chen<sup>3</sup>, Aifeng Zhou<sup>3<sup>±</sup></sup>, Qing Zhang<sup>3</sup>, Anson Mackay<sup>4\*</sup> 5 6 <sup>1</sup>Bo Cheng, 7 College of Urban and Environmental Science, Central China Normal University, 8 Wuhan 430079 China 9 chengbo@mail.ccnu.edu.cn 10 11 <sup>2</sup>Jennifer Adams 12 Department of Earth Sciences, University of Toronto, Toronto, ON, Canada 13 j.adams@utoronto.ca 14 15 <sup>3</sup>JianHui Chen 16 Key Laboratory of West China's Environmental System (Ministry of Education), College of 17 Earth and Environmental Sciences, Lanzhou University, Lanzhou 730000 China 18 jhchen@lzu.edu.cn 19 20 <sup>3</sup>Aifeng Zhou<sup>\*</sup>, 21 Key Laboratory of Western China's Environmental Systems (Ministry of Education), College 22 of Earth and Environmental Sciences, Lanzhou University, Lanzhou 730000, China 23 zhouaf@lzu.edu.cn 24 **Co-corresponding Author** 25 26 <sup>3</sup>Qing Zhang, 27 Key Laboratory of Western China's Environmental Systems (Ministry of Education), College 28 of Earth and Environmental Sciences, Lanzhou University, Lanzhou 730000, China 29 Zhangging16@lzu.edu.cn 30 31 <sup>4</sup>Anson Mackay\* 32 ECRC, Department of Geography, UCL, London WC1E 6BT UK

- 33 <u>a.mackay@ucl.ac.uk</u>
- 34 <u>Co-c</u>orresponding Author

35

36 37

38

# 41 Abstract:

42

43	During the latter stages of the Holocene, and prior to anthropogenic global warming, the
44	Earth underwent a period of cooling called the neoglacial. The neoglacial is associated with
45	declining summer insolation and changes to Earth surface albedo. Although impacts varied
46	globally, in China the neoglacial was generally associated with a cooler climate and an
47	attenuated Asian summer monsoon. Few studies in central China, however, have explored
48	the impact of neoglacial cooling on freshwater diversity, especially in alpine regions. Here
49	we take a palaeolimnological approach to characterise multidecadal variability in diatom
50	community composition, ecological guilds, and compositional turnover over the past 3,500
51	years from the alpine Lake Yuhuang Chi on Mount Taibai in the Qinling Mountains. Diatoms
52	in the high profile guild dominate much of the record from 3,500 - 615 cal yrs BP, which
53	suggests that few nutrients in the lake were limiting overall, and disturbance and herbivory
54	were likely low. After 615 cal yrs BP, low profile and planktic guild diatoms increase,
55	suggesting greater turbulence in the lake, alongside a decline in available nutrients. Diatom
56	turnover highlights periods in the lake history when deterministic processes structured
57	diatom communities. For example, an abrupt decline in turnover is coincident with the shift
58	from high to low profile diatoms at 615 cal yrs BP, and is likely due to the onset of the Little
59	Ice Age in the region. We suggest that Lake Yuhuang Chi become more shallow during
60	peak regional aridity, which led to the short-lived community restructuring observed in the
61	record.
62	
63	Keywords:

64 Diatoms, ecological guilds, compositional turnover, productivity, Little Ice Age

# **1. Introduction**

68	Alpine ecosystems are some of the most sensitive to changing climate, due in part to
69	elevation-dependent warming, i.e. the amplification of warming at higher altitudes (Yan and
70	Liu 2014; Pepin et al. 2015). Understanding how high-altitude ecosystems respond to
71	changing climate is a matter of urgency, because not only do these regions act as 'water
72	towers' supplying water to huge populations downstream (Messerli et al. 2004; Buyteart et
73	al., 2017), they provide habitats to many iconic species that are classified by the IUCN as
74	being vulnerable (Fan et al. 2014). Alpine freshwaters have multiple ecosystem functions
75	(Messerli et al. 2004; Buytaert et al., 2017) and provide many ecosystem services such as
76	freshwater regulation and habitat provision (Grêt-Regamey et al. 2011). Their
77	multifunctionality depends on local species communities, and how species vary through
78	space and time (beta-diversity) driven by ecosystem properties, environmental gradients,
79	and species interactions (Korhonen et al. 2010).
80	
81	Beta-diversity links biodiversity at regional and local scales, and may take the form of non-
82	directional variation, or directional turnover (Anderson et al. 2011). Long-term records of
83	beta-diversity commonly focus on the amount of compositional turnover over time, which
84	provide <u>s</u> important information on ecosystem functioning (Birks 2007). For example,
85	estimating species turnover assumes that species are lost and gained over time in response
86	to resource availability, competition, historical events and environmental factors (Korhonen
87	et al. 2010), over both recent (Smol et al. 2005) and long timescales (Leprieur et al. 2011.
88	However, because the concept of beta-diversity has so many meanings to different
89	disciplines, we take the approach of Felde et al. (2019) and just focus on directional
90	compositional turnover, appropriate for palaeolimnological datasets from a single lake.
I	

92 Natural archives are an important resource for reconstructing past environments where long-93 term historical and/or instrumental records are either scarce or absent. In central China, 94 speleothems provide exceptional, high resolution records of monsoon intensity, allowing 95 periods of multiannual and multidecadal drought to be determined (Wang et al. 2005). Yet 96 there are relatively few studies (Liu et al. 2017) which have explored multidecadal records of 97 biodiversity change over similar timescales, leaving a fundamental gap in understanding as 98 to how biodiversity in freshwater ecosystems, especially at higher altitudes, responded to 99 periods of climate variability. Reconstructing the impacts of past climate on freshwater 100 ecosystems is fundamental to understanding how freshwater biodiversity may respond to 101 future climate, especially during periods of rapid change. Here we focus on the neoglacial, 102 which spans at least the past c. 3,500 years.

103

104 The neoglacial, characterised by increasingly cooler temperatures, follows on from globally 105 warmer temperatures of the early- to mid-Holocene. The extent of cooling varied regionally. 106 being most pronounced in the extra-tropical northern hemisphere (Marcott et al. 2013). The 107 most important driver of northern hemisphere cooling was declining summer insolation (ibid.) 108 in conjunction with changes in albedo on the Earth's surface, linked to feedbacks from 109 vegetation and snow / ice. In China, the neoglacial resulted in the persistent decline in 110 monsoon intensity in southern China (Wang et al. 2005) and rapid decline in precipitation in 111 northern China (Chen et al. 2015a) leading to increased aridity and major shifts in vegetation 112 communities (Zhou et al. 2010). Superimposed on the insolation-driven neoglacial were 113 notable periods of sub\_Milankovitch, centennial-scale climatic events (e.g. Mayewski et al. 114 2004; Mann et al. 2009; Wanner et al. 2014), including the 2,800 yr BP event (Hall et al. 115 2004), the Medieval Climatic Anomaly (MCA) (c. 1000-1300 AD) and the Little Ice Age (LIA) 116 (c. 1300 – 1850 AD). The latter two events are well expressed throughout China; Medieval

temperatures were generally warmer than the following centuries spanning the LIA (Cook et
al. 2013; Chen et al. 2015b). However, while the LIA generally resulted in periods of aridity
(e.g. Wang et al. 2005; Tan et al. 2011; Chen et al. 2015a), in depth research highlights a
more heterogenous response across China (e.g. Cook et al. 2010), with some central and
southern regions becoming wetter due to interplays between the Westerly jet stream and the
Asian summer monsoon (ASM) (Tan et al. 2018).

123

Freshwater ecosystems in the Qinling Mountains of central China provide natural capital and ecosystem services for local and regional populations, and understanding the impact of monsoon variability on ecosystem functioning has the potential to add insight into how freshwater biodiversity may respond to future climate change, and predicted increases in mean annual precipitation (Guo et al. 2017). In this study, we reconstruct neoglacial trends in diatom community composition, their ecological guilds, and compositional turnover at a multidecadal resolution over the past 3500 years.

131

### 132 **1.1 Study region**

133

134 The Qinling Mountains are widely recognised for their conservation importance and as a 135 biodiversity hotspot; they are a refuge for many Tertiary plants (Zhang et al. 2017) and 136 vulnerable species such as the Giant Panda (Fan et al. 2014). The region is climatically very 137 sensitive, as it separates the northern subtropical zone of China from the country's central 138 warm-temperate zone (Figure 1). Mount Taibai (34°N, 108°E; 3767 m), is the highest 139 mountain in the range, with a timberline at c. 3,370 m, and treeline at c. 3,600 m (Liu et al. 140 2002). The mountain is classified as a glacial heritage site because Quaternary glaciations 141 are well preserved, especially the last glaciation (Yang et al. 2018). On Mount Taibai there 142 are several clusters of cirque lakes, and our study site, Lake Yuhuang Chi (YHC), is found in

- 143 one of these clusters. It is a cirque and moraine lake at 3370 masl, <u>placing it in the Larix</u>
- 144 <u>forest subalpine meadow ecotone. Specifically, Larix chinensis Beissn, grows in podzolic</u>
- soils between 2,000 m to 3,500 m, and is associated with cold, arid climate. Regional annual
- 146 average temperature is below 8°C and annual precipitation is c. 840-960 mm. Lake Yuhuang
- 147 <u>Chi has</u> a maximum depth of 21.5m and an area of c. 23,600 m<sup>2</sup>. <u>No shoreline macrophytes</u>
- 148 were observed during coring, but it was winter when the lake was frozen. Using a YSI
- 149 ProDSS multiparameter water quality meter, surface water pH was measured at 6.84 pH
- 150 <u>units.</u>
- 151
- **Figure 1:** Regional position of Lake Yuhuang Chi in the Qinling Mountains of central <u>China</u>.
- 153 The photograph of the frozen lake to the right shows the small\_catchment\_with few trees.



- 154
- 155
- 156 **2. Methods:**
- 157

# 158 2.1 Coring, Age model, Total Organic Carbon

- A <u>135 cm</u> sediment core (YHC15A) was collected <u>in 2015 CE</u> using a 6cm diameter piston
- 160 corer from the central region of Lake Yuhuang Chi. The highly humified lake sediments
- 161 <u>contained no sizable macrofossils for radiocarbon analyses.</u> Radiocarbon dating was <u>instead</u>
- 162 carried out on <u>five</u> bulk organic sediment <u>samples</u> using accelerator mass spectrometry
- 163 (AMS) at Beta Analytic, USA. There is a radiocarbon reservoir effect evident in the data

164	(likely from old soil carbon input from the catchment), so we used a quadratic extrapolation
165	to determined reservoir ages (Hou et al. 2012). All the radiocarbon dates were quadratic
166	fitted_( $^{14}CAge = 0.0693depth^2 + 17.31depth + 1340$ , R <sup>2</sup> = 0.9994), so we determined the top
167	(0cm) with a 1340 year reservoir age effect. An age-depth model was developed with
168	smooth fit using CLAM 2.2 (Blaauw, 2010) in <i>R</i> , using Intcal13 (Reimer et al., 2013)
169	calibration curve. Total organic carbon (TOC) provides an estimate of the amount of organic
170	carbon that escapes remineralisation before being incorporated into lake sediments. TOC
171	was measured on contiguous 1cm samples using an elemental analyser (Flash EA 1112).
172	

### 173 **2.2 Diatoms**

174 Diatom analysis was performed on alternate <u>1cm-thick</u> sediment samples. Approximately 175 0.1g of wet sediment from each sample was prepared using standard procedures outlined in 176 Battarbee et al. 2001. Organic matter was removed by heating each sample in 30% H<sub>2</sub>O<sub>2</sub>, 177 before 10% HCl was added to remove carbonates and any excess H<sub>2</sub>O<sub>2</sub>. Diatom 178 concentrations were calculated through the addition of divinylbenzene (DVB) microspheres 179 (concentration 8.02 x 10<sup>5</sup> spheres/cm<sup>3</sup>) to diatom suspensions, and diatom fluxes calculated 180 using sediment accumulation rates. Diatom suspensions were diluted and then pipetted onto 181 coverslips to dry before being fixed onto microscope slides with Naphrax<sup>®</sup>. Using a Zeiss 182 Axiostar Plus light microscope, diatoms were counted at x1000 magnification under an oil-183 immersion objective and phase contrast. A minimum of 300 diatom valves (min 331, max 184 591) were counted for each of the 67 samples. Diatoms were identified using a variety of 185 flora including Krammer and Lange-Bertalot, 1986, 1988, 1991a, 1991b; Williams and 186 Round, 1987; Lange-Bertalot, 2001.

187

188 Diatom species were categorised according to ecological guilds commonly associated with
189 the abundance of available resources (e.g. light, nutrients) and disturbance (e.g. grazing and

190 turbulence) (after Passy 2007; Rimet and Bouchez 2012). The low profile guild includes 191 diatoms which attach themselves to substrates in erect, prostrate, and adnate forms, are 192 very slow moving (Passy 2007), and are generally adapted to low nutrient conditions. High 193 profile guild diatoms are those of tall stature (e.g. they are filamentous, or chain-forming, or 194 found in mucilage tubes), and are generally adapted to high nutrients and low levels of 195 disturbance (Passy 2007). Motile diatoms are relatively fast\_moving species, tolerant of high 196 nutrients (Passy 2007) and siltation processes (Battegazzore et al. 2004). A new planktic 197 guild was determined by Rimet and Bouchez (2012) which includes centric species able to 198 resist sedimentation in lake ecosystems.

199

### 200 **2.3 Multivariate analyses**

201 The magnitude of diatom turnover was initially estimated using detrended correspondence 202 analysis (DCA), with square root transformation of the species data to stabilise variance and 203 rare species downweighted. The axis 1 gradient length was 1.44 standard deviation units, so 204 diatom abundances were reanalysed using principal components analysis (PCA). A log-205 linear contrast PCA was undertaken (appropriate for closed, relative abundance data (Lotter 206 and Birks 1993)), with symmetric scaling of ordination scores so that scaling of both samples 207 and species were optimised. Species compositional turnover was estimated using detrended 208 canonical correspondence analyses (DCCA), with the diatom data constrained using dates 209 from the calibrated age model (Smol et al. 2005). We used DCCA to estimate compositional 210 turnover because sample scores are scaled to be standard units through the process of 211 detrending by segments and non-linear rescaling (Birks 2007). Sample scores can therefore 212 be interpreted as the amount of species turnover through time, making them ecologically 213 useful and ideal for estimating compositional turnover. Ordinations were undertaken using 214 Canoco5 (Šmilauer and Lepš 2014). Breakpoint analysis, a form of segmented regression 215 analysis, was used to determine major points of change in diatom guilds and turnover, using

216	the segmented package in R v. 3.5.1 (Muggeo 2008). Stringent p-values were adopted
217	(p<0.001) when determining any major changes observed. All stratigraphical profiles shown
218	were constructed using C2 Data Analysis Version 1.7.7 (Juggins 2014), and zones
219	determined using stratigraphical constrained cluster analysis by incremental sum of squares
220	(CONISS) and broken stick analysis using the rioja package in R v. 3.5.1 (Juggins 2017).
221 222 223	3. Results
224	
225	3.1 Core Description and Age Model
226	The litholoay of the 135 cm-long YHC15A core consisted entirely of arey-brown, highly

humified gyttja. Radiocarbon dates of TOC are given in Table 1, and an age-depth model
 determined using CLAM 2.2 (Blaauw, 2010), shown in Figure 2. Sediment rates are lower
 during the early part of the record, before c. 1,400 cal yrs BP (0.28mm/yr) than sediments
 deposited during the past 1,400 years (0.48mm/yr).

Table 1: AMS-<sup>14</sup>C radiocarbon dates from Lake Yuhuang Chi (core YHC15A)

Lab No.	Depth (cm)	Material	δ <sup>13</sup> C ( <u>‰</u> VPDB)	<sup>14</sup> C date ±error (yr BP)	<sup>14</sup> C date minus 1340 reservoir age (yr BP)	Weighted calibrated age (No error) ( <u>cal yrs</u> BP)
Beta- 425231	10	Bulk organic	-24.6	1530±30	190±30	168
Beta- 425232	30	Bulk organic	-24.7	1920±30	580±30	595
Beta- 425233	50	Bulk organic	-24.9	2370±30	1030±30	949
Beta- 417757	70	Bulk organic	-24.8	2870±30	1530±30	1423
Beta- 425234	110	Bulk organic	-24.8	4140±30	2800±30	2868
Beta- 417758	130	Bulk organic	-24.9	4730±30	3390±30	3584

**Figure 2:** The age model determined on 5 radiocarbon dates of organic bulk sediments from

- core YHC15A. The age-depth model was developed with smooth fitting, using CLAM 2.2
   (Blaauw, 2010).
- 238



239

240

### 241 **3.2 Diatoms**

242 A total of 170 species of diatom were identified from Lake Yuhuang Chi, with the majority 243 (120 species) being only of low occurrence (< 1% in one or more samples). For much of the 244 stratigraphy, diatoms were dominated by fragilarioids and naviculoids up to c. 930 cal yrs 245 BP, [1020 CE] after which they decline, to be replaced by monoraphid and Gomphonema-246 type taxa alongside the centric *Lindavia*. Stratigraphically constrained cluster analysis by 247 incremental sum of squares analyses (CONISS) on diatom relative abundance data reveals 248 three zones. Zone 1 (c. 3550 – 2300 cal yrs BP), Zone 2 (c. 2300 – 615 cal yrs BP), and 249 Zone 3 (c. 615 cal yrs BP – present) (Fig 3,\_4). Zone 1 is dominated by diatoms in the high 250 profile guild (Fig 4), notably fragilarioids Stauroforma exiguiformis and Staurosirella pinnata. 251 Diatoms in the motile guild are well represented by the naviculoid Humidophila 252 schmassmannii, together with Diadesmis gallica, Mayamaea atomus and Mayamaea 253 fossalis. The decline in S. exiguiformis at the top of the zone is accompanied by an increase 254 in Pseudostaurosira brevistriata, and decline in motile diatoms e.g. M. atomus. In Zone 1, 255 there is a gradual decline in <u>compositional turnover</u> and PCA1 samples scores. Zone 2 is 256 marked by a notable increase in the planktic Lindavia bodanica and increasing P. 257 brevistriata and Pseudostaurosira pseudoconstruens. Diversity in zone 2 exhibits a rather 258 stable, flora, dominated by *P. brevistriata*, *P. pseudoconstruens* and *L. bodanica*, while 259 Gomphonema olivaceoides and Karayevia suchlandtii appear in the record for the first time 260 at c. 1400 and 1070 cal yrs BP, respectively. Motile diatoms become persistently lower than 261 the mean at this time during zone 2, while low profile diatom abundances increase to 262 fluctuate about the average (Fig 4). Zone 3 occurs just before a major change in diatom 263 turnover (Fig 3). Several species decline from the record altogether including S. exiguiformis 264 and *H. schmassmannii*, while other species reach peak abundance for the whole profile, 265 including *L*. bodanica and G. olivaceoides, and diatoms which occupy low profile guild status 266 in general (Fig 4). Denticula subtilis appears in the record for the first time at c. 400 cal yrs 267 BP (1550 CE). During zone 3, low profile and planktic diatoms increase to their highest 268 values for the whole record, while high profile and motile diatoms are persistently lower than 269 the mean. Diatom fluxes range from 0.07 – 7.02 (mean 1.85) valves x10<sup>6</sup> cm<sup>-2</sup> yr<sup>-1</sup>. When 270 centred around the mean, fluxes are highest in zone 2, between c. 1500 - 800 cal yrs BP 271 (450 – 1150 CE), but decline at c. 800 cal yrs BP (1150 CE), to lowest values from c. 600 cal 272 yrs BP (1350 CE) to the present (Fig 3). <u>%TOC values are low, and range from 2.6 - 4.3%</u> 273 (mean 3.45%). TOC values were highest in zone 1, declined in zone 2, and reached lowest 274 values, coincident with lowest values for compositional turnover, c. 1645 CE. 275

Figure 3: <u>Only diatoms with abundance</u> greater than 3% in more than one sample are

shown. Diatom species are shown as relative abundances (%), and grouped according to
 ecological guild. Also shown are PCA axis 1 sample scores, diatom compositional turnover

values, planktonic-benthic (P/B) ratio data, and mean-centered diatom fluxes. Zones were delimited using CONISS – see text for details.

281



Figure 4: All diatoms were classified into one of four guilds (after Passy 2007, and Rimet and Bouchez 2012): low profile (guild 1), high profile (guild 2), motile (guild 3) and planktic (guild 4). Guilds are presented as relative abundances (%) to the left, and deviations around the mean to the right.



PCA highlights a very strong first axis gradient which accounts for over 45% of variation in the diatom data. Trends in PCA-1 are most clearly seen in Fig 5, as deviations around the mean. Breakpoint analysis indicates major (p<0.001) change in PCA axis 1 scores (Table 2), close to the transition when PCA values switch from being higher than the mean, to being lower than the mean, and low values persist for the rest of the record. Diatom compositional turnover (estimated from DCCA; 1.033 SD units) shows a similar pattern to PCA-1, with breakpoints identified at c. 515 cal yr BP ± 40 years, (c. 1435 CE) and 335 cal yr BP ± 33 years (c. 1615 CE) (Table 2; Fig 5). 

- 309

Figure 5: Ordination and biodiversity trends shown as deviations around the mean.



Table 2: Significant breakpoints in diatom trend data; p < 0.001). 

	Breakpoint 1 <u>(cal yrs BP)</u>	p value	Breakpoint 2 <u>(cal yrs BP)</u>	p value
Species PCA	1850 ±200	p<0.001	none	
Compositional turnover	515 ± 97	p<0.001	335 BP ± 33	p<0.001
<u>Guild 2 – High profile</u>	<u>2910 ± 127)</u>	<u>p&lt;0.001</u>	<u>1565 BP ± 175</u>	<u>p&lt;0.001</u>
Guild 3 – Motile	<u>2880 ± 69</u>	p<0.001	<u>1960 BP ± 128</u>	p<0.001

- **4. Discussion:**
- 319

### 320 4.1 Diatom assemblage change

321 Sedimentary diatom assemblages in Lake Yuhuang Chi are dominated by species in the 322 Fragilariaceae (Fig 3) from 3,500 - 615 cal yrs BP. Fragilarioids are often opportunistic, 323 dominating assemblages in alpine and arctic lakes with a short growing season and long 324 periods of ice cover (Lotter and Bigler 2000). For example, July air temperature and ice 325 cover duration have both been shown to have significant influence on the abundance of 326 fragilarioids in the European Alps (Schmidt et al. 2004), while in a sub-alpine lake in the 327 Eastern Sayan mountains (Russia), insolation and northern hemisphere air temperatures 328 played a strong role on modulating fragilarioid responses through the Holocene (Mackay et 329 al. 2012). The early part of our record is dominated by Stauroforma exiguiformis, a species 330 common in dystrophic lakes (Flower et al. 1996). Its high abundance may be related to 331 relatively high carbon sequestration during the very early part of the record in Zone 1 (Fig 332 5c). The decline in S. exiguiformis is concomitant with the increase in Humidophila 333 schmassmannii and may be indicative of Lake Yuhuang Chi becoming less dystrophic 334 (Buczkó et al. 2015), perhaps linked to progressively cooler (Stebich et al. 2015) and more 335 arid (Chen et al. 2015a) climate (Fig 6) leading to increasingly frozen soils, reducing the 336 supply of allochthonous carbon. The decline in H. schmassmannii after c. 2800 cal yrs BP 337 further tracks the switch to a progressively cooler and more arid climate (Wang et al. 2005; 338 Chen et al. 2015b). 339 340 Growth of the planktic diatom L. bodanica in oligotrophic lakes is related to increased mixing 341 depth (Saros and Anderson 2015), because it can tolerate relatively low light levels and take 342 advantage of increased nutrient availability (Malik and Saros 2016). As the neoglacial 343 progressed therefore, mixing of the lake may have increased gradually, allowing this taxon

344	to eventually dominate for most of the past 615 years, even though diatom fluxes (Fig 3) and
345	%TOC (Fig 5) were relatively low, indicative of overall low diatom productivity.
346	
347	<u>4.2 Diatom traits</u>
348	Even though we do not have quantitative estimates of nutrients in this remote lake, we can
349	start to make inferences about nutrient availability from the traits exhibited by the diatom
350	communities themselves. For example, fragilarioids identified at Lake Yuhuang Chi can be
351	classified as high profile diatoms (i.e. of tall stature), able to compete effectively for
352	resources such as nutrients and light. However, their tall stature also makes them
353	susceptible to disturbance (Passy 2007) including turbulence in alpine lakes with short
354	water-residency times (Rimet et al. 2019). Their dominance at Lake Yuhuang Chi from 3500
355	- 615 cal yrs BP suggests that nutrients and light were not limiting (Passy and Larson 2019)
356	for much of the sequence. Herbivory was likely low as well (ibid.), because high profile
357	diatoms are not well adapted to high grazing pressures, which are generally low in alpine
358	lakes (Rimet et al. 2019).
359	
360	The decline in high profile guild values with concomitant increase in low profile and planktic
361	guilds since 615 cal yrs BP suggests that environmental conditions in Lake Yuhuang Chi
362	changed. Diatoms belonging to the low profile guild attach themselves to substrates in erect,
363	prostrate and adnate forms (Passy 2007), which means that although they are able to
364	tolerate relatively high disturbance, they also need to withstand an increase in the number of
365	resources that becomes limiting to them, i.e. they are well adapted to low nutrient
366	concentrations (Berthon et al. 2011). The planktic guild was proposed by Rimet et al. 2012
367	alongside the high, low and motile guilds determined by Passy (2007), as diatoms able to
368	resist sedimentation through the water column, and in our lake, this guild is dominated by L.
369	bodanica. That L. bodanica can tolerate low light conditions, and through turbulence of the
I	

370	water column obtain the nutrients that it needs to grow, again indicative of resources such as
371	nutrients becoming more limiting since the middle of the 14 <sup>th</sup> century.
372	
373	4.3 Diatom compositional turnover
374	In aquatic environments, when disturbance increases, or the number of limiting resources
375	increases, such as the decline in nutrient and / or light availability, deterministic processes
376	become more important than stochastic ones in structuring aquatic communities (Chase
377	2010). Increasing deterministic processes leads to a decline in compositional turnover
378	because disturbance or stress can act as environmental filters (Larson et al. 2016), through
379	niche selection. The decline in turnover between 3,100 – 1,900 cal yrs BP is gradual, and
380	reflects the slow, long term change in low profile guild diatoms increasing, and motile guild
381	diatoms declining. An increase in low profile diatoms hints at increasing disturbance and / or
382	declining nutrients (Passy 2007), while a decline in motile diatoms may be indictive of a
383	decline in siltation processes within the lake (Battegazzore et al. 2004). Over this time period
384	our data also show a marked decline in TOC sequestration to the bottom sediments (Fig.
385	5c). We suggest therefore the diatom and TOC data both point to a reduction in dissolved
386	organic carbon in Lake Yuhuang Chi, concurrent with cooling regional temperatures and
387	<u>increased aridity (</u> Chen et al. 2015a) <u>.</u>
388	
389	There are also periods when compositional turnover stabilised or even increased slightly
390	during the neoglacial, e.g. between c. $1.900 - 1,400$ cal yrs BP (Fig 6a). This period
391	coincides with distinctly warmer Arctic and European temperatures (PAGES 2k Consortium
392	2013), commonly referred to as the 'Roman Warm Period'. Precipitation in central China is
392	2013), commonly referred to as the 'Roman Warm Period'. Precipitation in central China is

- 393 closely tied to the intensity of the Asian summer monsoon (ASM) (Chen et al. 2015a), and at
- this time the summer monsoon was rather stable (Fig. 6d). Diatoms were dominated by high

- profile species, indicative that nutrients were not in limited supply, perhaps being washed in
- 396 from the catchment with summer monsoon rains.

Figure 6: <u>Compositional turnover</u> plotted alongside internal and external climate forcings:
mean temperature stack records for low latitude temperature anomalies (Fig <u>6b</u>; Marcott et al. 2013); <u>NE China pollen-inferred mean</u> temperature <u>warmest month</u> (July) (Fig <u>6c</u>; <u>Stebich</u> et al. 20<u>15</u>); trends in pollen-inferred mean annual precipitation (Fig <u>6d</u>; Chen et al. 2015a);
K+ concentrations in the GISP ice core (Fig <u>6e</u>; Mayewski et al. 2004); sun spot numbers
(Fig <u>6f</u>; Solanki et al. 2004); June solar insolation at 30 N (W m<sup>-2</sup>) (Fig <u>6g</u>; Berger and Loutre 1991)





409 Between 1400 – 615 cal yrs BP (550 - 1335 CE), turnover although lower than average 410 (Fig 6a), did not decline (Fig 3), indicative of increased competition between species (Larson 411 et al. 2016). This period coincides with the Medieval Climatic Anomaly (MCA), sometimes 412 referred to the Medieval Warm Period. Sub-decadal isotopic records from a stalagmite from 413 Buddha cave in the Qinling Mountains indicate a period of warm, wet climate between c. 985 414 - 475 cal yrs BP (965 - 1475 CE) (Paulsen et al. 2003), while phenological records, 415 including the beginning of tree flowering along the Yellow and Yangtze rivers, show that 416 winter half-year temperatures were high between 1380 – 640 cal yrs BP (570 – 1310 CE) 417 (Ge et al. 2003). Increased presence of *Quercus* and *Betula* pollen on Taibai Mountain also 418 suggest warm, wet conditions (Li et al. 2005; Wang et al. 2016) with temperatures perhaps 419 being as much as 2 °C warmer than mean annual temperatures observed today (Li et al. 420 2005). These local climate indicators tie in well with regional monsoon patterns; monsoon 421 strength was higher in central (Paulsen et al. 2003; Chen et al. 2015b; Wang et al. 2016) 422 and north east China (Chen et al. 2015a,b) than north west China (Chen et al. 2015b), while 423 globally, low latitude temperatures increased (Fig 6d). 424

## 425 **4.4** Abrupt ecological change during centennial-scale cold events

426 Against a backdrop of low northern hemisphere summer insolation (Fig 6g), amplified by 427 centennial-scale oceanic variability (Renssen et al., 2006), late Holocene cold events were 428 caused by several "overlapping" factors (such as volcanic eruptions and solar minima) 429 (Wanner et al., 2014). The most recent wide-scale cold event is the period commonly known 430 as the Little Ice Age approximately <u>1300</u> – 1850 CE, caused by several interacting, time-431 transgressive forcings, including reduced solar activity during the late 17th century (Shindell 432 et al. 2001), increased volcanic activity during the early 19th century (Brönnimann et al. 433 2019) and overall reduced Gulf Stream flow (Lund et al. 2006). It is the cooling event that we 434 focus on in this study, because cluster analyses of diatom assemblages delineate the

boundary between zones 2 and 3 at <u>615 cal yrs BP (1335 CE)</u> and <u>there are two significant</u>
<u>breakpoints in compositional turnover including when turnover is at its</u> lowest in the whole
record <u>at c. 335 – 330 cal yrs BP (1615-1620 CE) (Table 2)</u>.

438

439 Describing the Little Ice Age as a period characterised by cooler climate and glacier 440 readvance is rather simplistic, but one that has proven quite resilient, even as its 441 complexities are better understood (e.g. Matthews and Briffa 2005). As more regions are 442 investigated, impacts extend to changes in aridity as well as temperature. For example, 443 Chen et al. (2015b) demonstrated that by and large, regions north of 34° latitude (where our 444 study site is located) were generally drier than regions further south, with the extent of aridity 445 being affected by ocean-atmospheric interactions, such as ENSO, and its teleconnections to 446 SE Asia. The LIA is especially characterised by a strengthened Siberian High (SH), a semi-447 permanent anticyclone centred over Eurasia which strengthens intensively every winter. A 448 strong Siberian High results in a strong East Asian Winter Monsoon (EAWM) (Zhang et al. 449 1997). K<sup>+</sup> concentrations in the GISP ice core clearly show that the Siberian High was 450 especially strong between c. 550 - 150 cal yrs BP (1400 - 1800 CE) (Fig 6f), given that 451 potassium is likely sourced from central Asian dust via long-range transport to the Greenland 452 ice sheet (Meeker and Mayewski 2002). Concurrent with increased aridity, global low 453 latitude temperature records show rapid cooling at this time (Fig 6b; Marcott et al. 2013), 454 which in China led to very low winter anomalies (Ge et al. 2003), and lowest summer pollen-455 inferred temperatures in NE China (Fig 6c). 456 457 Very low diatom fluxes characterise the past 800 years at Lake Yuhuang Chi (Fig 3), 458 indicative of reduced diatom productivity, coincident with prevailing colder climate. Lowest 459 compositional turnover at c. 335 - 330 cal yrs BP (1615-1620 CE), linked to the

disappearance of *S. exiguiformis*, may be due to enhanced frozen soils, leading to reduced

461 carbon transport to the lake, while the disappearance of H. schamassmannii may be 462 because it cannot tolerate such low water temperatures (Buczkó et al. 2015), alongside 463 other factors such as a decline in siltation (Battegazzore et al. 2004). Denticula subtilis is a 464 very motile diatom, and in the Canadian High Arctic is characteristic of shallow lakes with 465 elevated conductivity (Antoniades et al. 2005). It may therefore be reflective of the lake 466 becoming more shallow due to increased aridity. Elsewhere, precipitation minima were 467 reconstructed from nearby by Gonghai lake (Chen et al. 2015a), and at neighbouring Lake 468 Sanging Chi, high Larix and Ephedra pollen frequencies are interpreted as being indicative 469 of cold, dry conditions (Wang et al. 2016). That minimum turnover and highest values for low 470 profile diatoms are observed at this time, indicates nutrient limitation and / or significant 471 disturbance has led to specialised diatoms to occupy niches where competition was very 472 low (Larson et al. 2016). Following harshest conditions for diatom growth in Lake Yuhuang 473 Chi during the early 17th century turnover increases once more, indicative of increased 474 competition among species (Larson et al. 2016).

475

476 While cold and arid climate during the LIA had a major impact on diatom diversity in Lake 477 Yuhuang Chi, impacts from previous centennial-scale cold events such as the 2,800 yr BP 478 event, are less conclusive. Like the LIA, the event dated at c. 2,800 yr BP is concurrent with 479 a deep, abrupt reduction in solar activity (Fig 6f), which led to a decline in surface water 480 temperatures in the North Atlantic (Andersson et al., 2003), weaker meridional overturning 481 circulation (Hall et al., 2004) and sea-ice expansion (Renssen et al. 2006). But although 482 these events led to a rapid weakening in ASM intensity in southern China (Wang et al. 483 2005), reconstructed precipitation from Gonghai Lake in northern China suggests that aridity 484 was already increasing from c. 3,100 cal yrs BP (Fig 6d) (Chen et al. 2015b). There is a 485 small increase in GISP2 K<sup>+</sup> concentrations, which suggests that the Siberian High during the 486 time of the 2,800 yr BP event did not reach the strengths observed during the LIA (Fig 6e),

487	but still led to regional summer time cooling (Fig 6c, Stebich et al. 2015). At Lake Yuhuang
488	Chi, these changes in climate may have caused both the small, temporary decline in both
489	compositional turnover and total diatom flux, but also a significant decline in high profile
490	diatoms, concomitant with an increase in L. bodanica. This may suggest increased water
491	column instability and turbulence in the lake (Saros and Anderson 2015).
492	
493	5. Conclusions
494	
495	Diatom turnover in the Qinling mountains of central China demonstrates strong directional
496	change. The decline in turnover over the past 3,500 years mirrors declining low latitude June
497	insolation, which drives overall low latitude cooling (Marcott et al. 2013). This suggests a
498	strong link between orbitally-forced climate change and increasingly deterministic processes
499	affecting aquatic ecosystems in this high altitude region of central China. Over the last 1300
500	years, impacts related to the Medieval Climatic Anomaly and the Little Ice Age are also
501	expressed in the palaeolimnological records from Lake Yuhuang Chi. Increased diatom
502	fluxes (Fig. 3) are coincident with increased regional summer precipitation during the MCA
503	(Chen et al. 2015a). Colder, more arid conditions during the Little Ice Age, linked to a very
504	strong Siberian High, led to lowest diatom turnover and carbon sequestration for the past
505	3,500 years, possibly due to the lake becoming more shallow at this time. Our study
506	highlights the value of interpreting diatom changes through compositional turnover, and
507	ecological trait analyses of their growth forms.
508	
509	
510	
511	
512	

# 513 **6.** Author contribution

514

515 BC, JC & QZ designed the study. BC undertook the diatom analyses, and AZ the

516 radiocarbon dating. JA provided assistance with statistical analyses, and AWM prepared the

517 manuscript with contributions from all authors.

518

# 519 **7.** Acknowledgements:

- 520
- 521 Funding: The work was supported by a China Scholarship Council award to Dr Bo Cheng,
- 522 and by the National Natural Science Foundation of China (Grants No. 41771208; No.
- 523 41790421). The authors declare that they have no conflict of interest.

525	
526	8. References
527	
528	Andersson, C., Risebrobakken, B., Jansen, E., and Dahl, S. O.: Late Holocene surface
529	ocean conditions of the Norwegian Sea (Vøring Plateau), Paleoceanography, 18, 1044,
530	doi:10.1029/2001PA000654, 2003.
531	
532	Antoniades, D., Douglas, M.S. and Smol, J.P.: Benthic diatom autecology and inference model
533	development from the Canadian high arctic Archipelago 1. J Phycol., 41, 30-45, 2005.
534	
535	Battarbee, R.W., Jones, V.J., Flower, R.J., Cameron, N.G., Bennion, H., Carvalho, L.
536	and Juggins, S.: 2001. Diatoms. In: Smol, J.P., Birks, H.J.B., Last, W.M. (Eds.), Tracking
537	Environmental Change Using Lake Sediments. Kluwer Academic Publishers, Dordrecht,
538	<u>pp. 155e201.</u>
539	
540	Battegazzore, M.; Morisi, A.; Gallino, B.; Fenoglio, S.: Environmental quality evaluation of
541	Alpine springs in NW Italy using benthic diatoms. Diatom Res., 19, 149-165 2004
542	
543	Berthon, V., Bouchez, A., and Rimet, F.: Using diatom life-forms and ecological guilds to
544	assess organic pollution and trophic level in rivers: a case study of rivers in south-
545	eastern France. Hydrobiologia, 673, 259–271, 2011.
546	
547	Birks, H. J. B.: Estimating the amount of compositional change in late-Quaternary pollen-
548	stratigraphical data, Veg. Hist. Archaeobot., 16, 197-202, 2007.
549	
550	Blaauw, M.: Methods and code for 'classical' age-modelling of radiocarbon
551	sequences, Quat. Geochron., 5, 512-518, 2010.

552	
553	Brönnimann, S., Franke, J., Nussbaumer, S.U., Zumbühl, H. J., Steiner, D., Trachsel, M., Hegerl,
554	G. C., Schurer, A., Worni, M., Malik, A., Flückiger, J.: Last phase of the Little Ice Age forced by
555	volcanic eruptions. Nat. Geosci. 12, 650-656, 2019.
556	
557	
558	Buytaert, W., Moulds, S., Acosta, L., De Bievre, B., Olmos, C., Villacis, M., Tovar, C.,
559	and Verbist, K. M. J.: Glacial melt content of water use in the tropical Andes, Environ.
560	Res. Lett., 12, 114014, 2017.
561	
562	Buczkó, K., Wojtal, A. Z., Beszteri, B., and Magyari, E. K.: Morphology and distribution of
563	Navicula schmassmannii and its transfer to genus Humidophila, Studia Botanica
564	Hungarica, 46, 25-41, 2015.
565	
566	Chase, J. M.: Drought mediates the importance of stochastic community assembly, P.
567	Natl. Acad. Sci. USA, 104, 17430-17434, 2010.
568	
569	Chen, F., Xu, Q., Chen, J., Birks, H. J. B., Liu, J., Zhang, S., Jin, L., An, C., Telford, R. J.,
570	Cao, X., and Wang, Z.: East Asian summer monsoon precipitation variability since the
571	last deglaciation, Sci. Rep. UK, 5, 11186, 2015a.
572	
573	Chen, J., Chen, F., Feng, S., Huang, W., Liu, J., and Zhou, A.: Hydroclimatic changes in
574	China and surroundings during the Medieval Climate Anomaly and Little Ice Age: spatial
575	patterns and possible mechanisms, Quaternary Sci. Rev., 107, 98-111, 2015b.
576	

577	Cook, E. R., Krusic, P. J., Anchukaitis, K. J., Buckley, B. M., Nakatsuka, T., Sano, M.,
578	and PAGES Asia2k Members: Tree-ring reconstructed summer temperature anomalies
579	for temperate East Asia since 800 C.E., Clim. Dynam., 41, 2957-2972, 2013.
580	
581	Cook, E. R., Anchukaitis, K. J., Buckley, B. M., D'Arrigo, R. D., Jacoby, G. C., and
582	Wright, W. E.: Asian monsoon failure and megadrought during the last
583	millennium, Science, 328, 486-489, 2010.
584	
585	Fan, J. T., Li, J. S., Xia, R., Hu, L. L., Wu, X. P., and Guo, L.: Assessing the impact of
586	climate change on the habitat distribution of the giant panda in the Qinling Mountains of
587	China, Ecol. Model., 274, 12–20, 2014.
588	
589	Felde, V. A., Flantua, S. G., Jenks, C.R., Benito, B. M., De Beaulieu, J.L., Kuneš, P.,
590	Magri, D., Nalepka, D., Risebrobakken, B., ter Braak, C. J. and Allen, J.R.:
591	Compositional turnover and variation in Eemian pollen sequences in Europe. Veg. Hist.
592	<u>Archaeobot,1-9. 2019.</u>
593	
594	Flower, R. J., Jones, V. J., and Round, F. E.: The distribution and classification of the
595	problematic Fragilaria (virescens V.) exigua Grun./Fragilaria exiguiformis (Grun.) Lange-
596	Bertalot: A new species or a new genus?, Diatom Res., 11, 41-57. 1996.
597	
598	Ge, Q., Zheng, J., Fang, X., Man, Z., Zhang, Z., Zhang, P., and Wang, W.: Winter half-
599	year temperature reconstruction for the middle and lower reaches of the Yellow River
600	and Yangtze River, China, during the past 2000 years, Holocene, 13, 933-940, 2003.
601	

602	Grêt-Regamey, A., Brunner, S. H., and Kienast, F.: Mountain ecosystem services: who
603	cares?, Mt. Res. Dev., 32, S23-S34, 2012.
604	
605	Guo, J., Huang, G., Wang, X., Li, Y., and Lin, Q.: Investigating future precipitation
606	changes over China through a high-resolution regional climate model ensemble, Earth's
607	Future, 5, 285-303, 2017.
608	
609	Hall, I. R., Bianchi, G. G., and Evans, J. R.: Centennial to millennial scale Holocene
610	climate-deep water linkage in the North Atlantic, Quaternary Sci. Rev., 23, 1529-1536,
611	2004.
612	
613	Hou, J., D'Andrea, W.J. and Liu, Z.: The influence of 14C reservoir age on interpretation
614	of paleolimnological records from the Tibetan Plateau. Quaternary Sci. Rev., 48, 67-79,
615	<u>2012.</u>
616	
617	Juggins, S.: C2 version 1.7.7: software for ecological and palaeoecological data.
618	University of Newcastle. 2014
619	
620	Juggins, S.: rioja: Analysis of Quaternary Science Data, R package version
621	3.5.1.http://cran.r-project.org/package=rioja, 2017.
622	
623	Korhonen, J. J., Soininen, J., Hillebrand, H.: 2010. A quantitative analysis of temporal
624	turnover in aquatic species assemblages across ecosystems. Ecology, 91, 508-517.
625	
626	Krammer, K., and Lange-Bertalot, H.: Bacillariophyceae: Naviculaceae. In: Ettl, H.,
627	Gerloff, J., Heynig, H., Mollenhauer, D. (Eds.), Süsswasserflora von Mitteleuropa. Band

2, Teil 1. Gustav Fischer, Stuttgart, 1986.

- 630 Krammer, K., and Lange-Bertalot, H.: Bacillariophyceae: Bacillariaceae, Epithemiaceae, 631 Surirellaceae. In: Ettl, H., Gerloff, J., Heynig, H., Mollenhauer, D. (Eds.), Süsswasserflora 632 von Mitteleuropa. Band 2, Teil 2. Gustav Fischer, Stuttgart, 1988. 633 634 Krammer, K., and Lange-Bertalot, H.: Bacillariophyceae: Achnanthaceae, Kritische Erga" 635 nzungen zu Navicula (Lineolatae) und Gomphonema Gesamtliterarurverzeichnis. In: Ettl, 636 H., Gerloff, J., Heynig, H., Mollenhauer, D. (Eds.), Süsswasserflora von Mitteleuropa. 637 Band 2, Teil 4. Gustav Fischer, Stuttgart, 1991a. 638 639 Krammer, K., and Lange-Bertalot, H.: Bacillariophyceae: Centrales, Fragilariaceae, 640 Eunotiaceae. In: Ettl, H., Gerloff, J., Heynig, H., Mollenhauer, D. (Eds.), Süsswasserflora 641 von Mitteleuropa. Band 2, Teil 3. Gustav Fischer, Stuttgart, 1991b. 642 643 Lange-Bertalot, H.: Diatoms of Europe, Volume 2: Navicula sensu stricto. 10 genera 644 separated from Navicula sensu lato Frustulia. In: Lange- Bertalot, H. (Ed.) Diatoms of 645 Europe: diatoms of the European inland waters and comparable habitats. A.R.G. 646 Gantner Verlag K.G., Ruggell, Germany, 2001. 647 648 Larson, C. A., Adumatioge, L., and Passy, S. I.: The number of limiting resources in the 649 environment controls the temporal diversity patterns in the algal benthos, Microbial 650 Ecol., 72, 64-69, 2016.
- 651

652	Leprieur, F., Tedesco, P. A., Hugueny, B., Beauchard, O., Dürr, H. H., Brosse, S., and
653	Oberdorff, T.: Partitioning global patterns of freshwater fish beta diversity reveals
654	contrasting signatures of past climate changes, Ecol. Lett., 14, 325-334, 2011.
655	
656	Li, X., Dodson, J., Zhou, J., Wang, S., and Sun, Q.: Vegetation and climate variations at
657	Taibai, Qinling Mountains in central China for the last 3500 cal BP, Journal Integr. Plant
658	Biol., 47, 905-916, 2005.
659	
660	Liu, H., Tang, Z., Dai, J., Tang, Y., and Cui, H.: Larch timberline and its development in
661	North China. Mt. Res. Dev., 22, 359-367, 2002.
662	
663	Liu, J., Rühland, K. M., Chen, J., Xu, Y., Chen, S., Chen, Q., Huang, W., Xu, Q., Chen,
664	F., and Smol, J. P.: Aerosol-weakened summer monsoons decrease lake fertilization on
665	the Chinese Loess Plateau, Nat. Clim. Change, 7, 190-195, 2017.
666	
667	Lotter, A.F., and Birks, H.J.B.: The impact of the Laacher See tephra on terrestrial and aquatic
668	ecosystems in the Black Forest, southern Germany. J. Quat. Sci., 8, 263-276, 1993.
669	
670	Lotter, A. F., and Bigler, C.: Do diatoms in the Swiss Alps reflect the length of ice cover,
671	Aquat. Sci., 62, 125-141, 2000.
672	
673	Lund, D. C., Lynch-Stieglitz, J., Curry, W. B.: Gulf Stream density structure and transport during
674	the past millennium. Nature, 444, 601- 604, 2006
675	
676	Mackay, A. W., Bezrukova, E. V., Leng, M. J., Meaney, M., Nunes, A., Piotrowska, N.,
677	Self, A., Shchetnikov, A., Shilland, E., Tarasov, P., Wang, L., and White, D.: Aquatic

678	ecosystem responses to Holocene climate change and biome development in boreal,
679	central Asia, Quaternary Sci. Rev., 41, 119-131, 2012.
680	
681	Malik, H. I., and Saros, J. E.: Effects of temperature, light and nutrients on five Cyclotella
682	sensu lato taxa assessed with in situ experiments in arctic lakes. J. Plankton Res., 38,
683	431-442, 2016.
684	
685	Mann, M. E., Zhang, Z., Rutherford, S., Bradley, R. S., Hughes, M. K., Shindell, D.,
686	Ammann, C., and Faluvegi, G., Ni, F.: Global signatures and dynamical origins of the
687	Little Ice Age and Medieval Climate Anomaly, Science, 326, 1256-1260, 2009.
688	
689	Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional
690	and global temperature for the past 11,300 years, Science, 339, 1198-1201, 2013.
691	
692	Matthews, J. A., and Briffa, K. R.: The 'Little Ice Age': re-evaluation of an evolving
693	concept, Geogr. Ann. A., 87, 7-36, 2005.
694	
695	Mayewski, P. A., Rohling, E. E., Stager, J. C. et al.: Holocene climate variability,
696	Quaternary Res., 62, 243–255, 2004.
697	
698	Meeker, L. D., and Mayewski, P. A.: A 1400-year high-resolution record of atmospheric
699	circulation over the North Atlantic and Asia. The Holocene 12, 257-266, 2002.
700	
701	Messerli, B., Viviroli, D., and Weingartner, R.: Mountains of the world: vulnerable water
702	towers for the 21st century, Ambio, 29-34, 2004.
703	

704	Muggeo, V. M. R.: Segmented: An R Package to Fit Regression Models with Broken-
705	Line Relationships, R News, 8, 20-25, 2008.
706	
707	PAGES 2k Consortium: Continental-scale temperature variability during the
708	past two millennia, Nat. Geosci. 6, 339e346, 2013.
709	
710	Passy, S. I.: Diatom ecological guilds display distinct and predictable behaviour along
711	nutrient and disturbance gradients in running waters, Aquat. Bot., 86, 171–178, 2007.
712	
713	Passy, S.I., Larson, C.A.: Niche dimensionality and herbivory control stream algal
714	biomass via shifts in guild composition, richness, and evenness. Ecology e02831, 2019.
715	
716	Paulsen, D. E., Li, H. C., and Ku, T. L.: Climate variability in central China over the last
717	1270 years revealed by high-resolution stalagmite records, Quaternary Sci. Rev., 22,
718	691-701, 2003.
719	
720	Pepin, N., Bradley, R. S., Diaz, H. F., Baraër, M., Caceres, E. B., Forsythe, N., Fowler,
721	H., Greenwood, G., Hashmi, M. Z., Liu, X. D., and Miller, J. R.: Elevation-dependent
722	warming in mountain regions of the world, Nat. Clim. Change, 5, 424-430, 2015.
723	
724	Reimer, P. J., Bard, E., Bayliss, A. et al.: IntCal13 and Marine13 radiocarbon age
725	calibration curves 0–50,000 years cal BP. Radiocarbon, 55, 1869–1887, 2013.
726	
727	Renssen, H., Goosse, H., and Muscheler, R.: Coupled climate model simulation of
728	Holocene cooling events: oceanic feedback amplifies solar forcing, Clim. Past, 2, 79-90,
729	2006.

730	
731	Rimet, F., and Bouchez, A.: Life-forms, cell-sizes and ecological guilds of diatoms in
732	European rivers, Knowl. Manag. Aquat. Ecosyst. 406, 01, 2012.
733	Rimet, F., Feret, L., Bouchez, A., Dorioz, JM., Dambrine E.:. Factors influencing the
734	heterogeneity of benthic diatom communities along the shoreline of natural alpine lakes.
735	<u>Hydrobiologia, 839, 103-118, 2019</u>
736	
737	Saros, J. E., and Anderson, N. J. The ecology of the planktonic diatom Cyclotella and its
738	implications for global environmental change studies, Biol. Rev., 90, 522-541, 2015.
739	
740	Schmidt, R., Kamenik, C., Lange-Bertalot, H., and Klee, R.: Fragilaria and Staurosira
741	(Bacillariophyceaea) from sediment surfaces of 40 lakes in the Austrian Alps in relation
742	to environmental variables, and their potential for palaeoclimatology. J. Limnol., 63, 171-
743	189, 2004.
744	
745	Šmilauer, P., and Lepš, J.: Multivariate analysis of ecological data using CANOCO 5.
746	Cambridge University Press. 2014.
747	
748	Solanki, S. K., Usoskin, I. G., Kromer, B., Schüssler, M., and Beer, J: An unusually active
749	sun during recent decades compared to the previous 11,000 years, Nature, 431, 1084-
750	1087, 2004.
751	
752	Smol, J. P., Wolfe, A. P., Birks, H. J. B., et al.: Climate-driven regime shifts in the
753	biological communities of arctic lakes, P. Natl. Acad. Sci. USA, 102, 4397-4402, 2005.
754	

755	Stebich, M., Rehfeld, K., Schlütz, F., Tarasov, P., Liu, J., Mingram, J.: Holocene vegetation
756	and climate dynamics of NE China based on the pollen record from Sihailongwan Maar
757	Lake. Quat. Sci. Rev., 124, 275-289, 2015.
758	
759	Tan, L., Cai, Y., An, Z., Yi, L., Zhang, H., and Qin, S.: Climate patterns in north central
760	China during the last 1800 yr and their possible driving force, Clim. Past, 7, 685-692,
761	2011.
762	
763	Tan, L., Cai, Y., Cheng, H., Edwards, E. R., Gao, Y., Xu, H., Zhang, H., and An, Z.:
764	Centennial- to decadal-scale monsoon precipitation variations in the upper Hanjiang
765	River region, China over the past 6650 years, Earth Planet. Sci. Lett., 482, 580-590,
766	2018.
767	
768	Wang, Y., Cheng, H., Edwards, R. L., He, Y., Kong, X., An, Z., Wu, J., Kelly, M. J.,
769	Dykoski, C. A., and Li, X.: The Holocene Asian monsoon: links to solar changes and
770	North Atlantic climate, Science, 308, 854-857, 2005.
771	
772	Wang, H., Song, Y. Cheng, Y., Luo, Y., Gao, Y., Deng, L., and Liu, H.: Mineral
773	magnetism and other characteristics of sediments from a sub-alpine lake (3080 m asl) in
774	central east China and their implications on environmental changes for the last 5770
775	years, Earth Planet. Sci. Lett., 452, 44-59, 2016.
776	
777	Wanner, H., Mercolli, L., Grosjean, M., and Ritz, S.P.: Holocene climate variability and
778	change: a database review, J. Geol. Soc. Lond., 172, 254-263, 2014.
779	

780	Williams, D. M., and Round, F. E.: Revision of the genus Fragilaria. Diatom Res., 2, 267-
781	288, 1987.
782	
783	Yan, L., and Liu, X.: Has climatic warming over the Tibetan Plateau paused or continued
784	in recent years? J. Earth, Ocean Atmos. Sci., 1, 13-28, 2014.
785	
786	Yang, W., Yan, Y., Zhang, Y., Guo, W., and Zha, F.: Geoheritages in the Qinling
787	Orogenic Belt of China: Features and comparative analyses, Geol. J., 53, 98-413, 2018.
788	
789	Zhang, Y., Sperber, K. R., and Boyle, J. S.: Climatology and interannual variation of the
790	east Asian winter monsoon: Results from the 1979–95 NCEP/NCAR reanalysis, Mon.
791	Wea. Rev., 125, 2605–2619, 1997.
792	
793	Zhang, Y. B., Wang, Y. Z., Phillips, N., Ma, K. P., Li, J. S., and Wang, W.: Integrated
794	maps of biodiversity in the Qinling Mountains of China for expanding protected
795	areas. Biol. Conserv., 210, 64-71, 2017.
796	
797	Zhou, A., Sun, H., Chen, F., Zhao, Y., An, C., Dong, G., Wang, Z., and Chen, J.: High-
798	resolution climate change in mid-late Holocene on Tianchi Lake, Liupan Mountain in the
799	Loess Plateau in central China and its significance. Chinese Sci. Bull., 55, 2118-2121,
800	2010.
801	

## 803 Figure Legends

804

Figure 1: Regional position of Lake Yuhuang Chi in the Qinling Mountains of central Asia.

806 The lake is situated 33<u>70</u>m asl, and was formed by glacial activity. The photograph of the

807 frozen lake to the right shows the small catchment and tundra vegetation.

808

809 Figure 2: The age model determined on 5 radiocarbon dates of organic bulk sediments from

810 core YHC15A. The age-depth model was developed with smooth fitting using CLAM 2.2

811 (Blaauw, 2010).

812

813 Figure 3: Diatoms shown greater than 3% in more than one sample. Diatom species are

given as relative abundances. Also shown are PCA axes 1 scores for species and genera,

815 <u>compositional turnover</u> values, planktonic-benthic (P/B) ratio data, and mean-cente<u>re</u>d

816 diatom fluxes. Zones were delimited using CONISS – see text for details.

817

Figure 4: All diatoms were classified into one of four guilds (after Passy 2007, and Rimet and
Bouchez 2012): low profile (guild 1), high profile (guild 2), motile (guild 3) and planktic (guild
4). Guilds are presented as relative abundances (%) to the left, and deviations around the
mean to the right.

822

Figure 5: Ordination, <u>compositional turnover</u>, and <u>TOC</u> trends shown as deviations around the mean.

825

Figure 6: Compositional turnover (deviations around the mean) plotted alongside internal

827 and external climate forcings: mean temperature stack records for low latitude temperature

- 828 anomalies (Fig 6b; Marcott et al. 2013); NE China pollen-inferred mean temperature
- 829 warmest month (July) (Fig 6c; Stebich et al. 2015); trends in pollen-inferred mean annual
- 830 precipitation (Fig 6d; Chen et al. 2015a); K+ concentrations in the GISP ice core (Fig 6e;
- 831 Mayewski et al. 2004); sun spot numbers (Fig 6f; Solanki et al. 2004); June solar insolation
- at 30 N (W m<sup>-2</sup>) (Fig 6g; Berger and Loutre 1991)
- 833
- 834
- 835
- 836