

Dear Professor Fischer,

Please find below the revised versions of the manuscript and the supplement of cp-2019-94. The differences with the first versions are highlighted, with removed text in ~~strikeout red~~ and added text in underlined blue.

Following the comments of the reviewers, here are the main differences with the first version:

- We have added details to describe the CFA method, notably why gaps are present in the final data.
- We have clarified the supplement section explaining why we need to deconvolve the original NEEM record, and we have performed a sensitivity analysis to assess the impact of the deconvolution on the conclusions of the article.
- We have removed the calcium parameterization from the layering artifacts model as it lacked robustness. However, we have kept the discussion on the potential link between chemistry and the presence of layering artifacts (Section 3.3.1). We have clearly stated that this is a preliminary results, that needs to be confirmed with dedicated high-resolution measurements. However, we do not oppose removing this section from the article as well, if requested to improve the clarity of the article.
- We have added details on the algorithm to remove layering artifacts from the CFA, and explained why we do not clean the data purely manually.
- We have detailed the description of the GAD extraction method.
- We have added a more detailed discussion on why the DO21 period is a better discriminant for gas age distributions than the DO6-9 period.

The removal of the calcium parameterization in the layering model should facilitate the overall understanding of the paper. We are aware that different parts of the manuscript are not independent and that these overlaps don't facilitate the reading, but we did not find a simpler overall organization of the manuscript.

Best Regards,
Kévin Fourteau on behalf of all co-authors.

Estimation of gas record alteration in very low accumulation ice cores

Kévin Fourteau¹, Patricia Martinerie¹, Xavier Faïn¹, Alexey A. Ekaykin², Jérôme Chappellaz¹, and Vladimir Lipenkov²

¹Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, F-38000 Grenoble, France

²Climate and Environmental Research Laboratory, Arctic and Antarctic Research Institute, St. Petersburg, 199397, Russia

Correspondence: kevin.fourteau@univ-grenoble-alpes.fr or patricia.martinerie@univ-grenoble-alpes.fr

Abstract. We measured the methane mixing ratios of enclosed air in five ice core sections drilled on the East Antarctic ~~plateau~~Plateau. Our work aims to study two effects that ~~affect~~alter the recorded gas concentrations in ice cores: layered gas trapping artifacts and firn smoothing. Layered gas trapping artifacts are due to the heterogeneous nature of polar firn, where some strata might close early and trap abnormally old gases that appear as spurious values during measurements. The smoothing is due to the combined effects of diffusive mixing in the firn and the progressive closure of bubbles at the bottom of the firn. Consequently, the gases trapped in a given ice layer span a distribution of ages. ~~Concentration measurements thus only measure the average value in the ice layer~~This means that the gas concentration in an ice layer is the average value over a certain period of time, which removes the fast variability from the record. ~~We~~we focus on the study of East Antarctic ~~plateau~~Plateau ice cores, as these low accumulation ice cores are particularly affected by both layering and smoothing. ~~Our results suggest that the presence of layering artifacts in deep ice cores is linked with the chemical content of the ice.~~ We use high-resolution methane data to ~~parametrize~~test a simple model reproducing the layered gas trapping artifacts for different accumulation conditions typical of the East Antarctic ~~plateau~~Plateau. We also use the high-resolution methane measurements to estimate the gas age distributions of the enclosed air in the five newly measured ice core sections. It appears that for accumulations below ~~2cm.ice.yr⁻¹ (ice equivalent)~~2cm ice equivalent yr⁻¹ the gas records experience nearly the same degree of smoothing. We therefore propose to use a single gas age distribution to represent the firn smoothing observed in the glacial ice cores of the East Antarctic ~~plateau~~Plateau. Finally, we used the layered gas trapping model and the estimation of glacial firn smoothing to ~~estimate~~quantify their potential impacts on a hypothetical million-and-a-half years old ice core from the East Antarctic ~~plateau~~Plateau. Our results indicate that layering artifacts are no longer individually resolved in the case of very thinned ice near the bedrock. They nonetheless contribute to slight biases of the measured signal (less than 10ppbv and 0.5ppmv in the case of methane and carbon dioxide, respectively). However, these biases are small compared to the dampening experienced by the record due to firn smoothing.

Copyright statement. TEXT

1 Introduction

The East Antarctic ~~plateau~~ Plateau is characterized by low temperatures and low accumulation rates. ~~The low precipitations create~~ This creates the conditions for the presence of very old ice near the domes ~~of the~~ region (Raymond, 1983; Martín and Gudmundsson, 2012). Thanks to this particularity, the ~~ice core~~ retrieved at Dome C, within the EPICA project, has been dated
5 ~~back~~ to 800,000 years in the past (Bazin et al., 2013; Veres et al., 2013). The analysis of the ice, and of the bubbles within, has made ~~possible~~ the reconstruction of the Earth's past temperatures and atmospheric concentrations in major greenhouse gases over the last eight glacial cycles (Lüthi et al., 2008; Loulergue et al., 2008). In turn, this knowledge helps us to better understand the Earth climate system and its past and future evolutions (Shakun et al., 2012). Furthermore, there is currently an active search for ice ~~older than~~ one-and-a-half million years within the European Beyond EPICA-Oldest Ice project. Most
10 of the potential drilling sites for such old ice are located on the East Antarctic ~~plateau~~ Plateau (Fischer et al., 2013; Van Lief-feringe and Pattyn, 2013; Passalacqua et al., 2018). Within the next decade, we might therefore expect the retrieval of new deep ice cores at low accumulation drilling sites of East Antarctica, with ages reaching back to one-and-a-half million years or more.

However, the gas records in low accumulation ice cores ~~should not be directly interpreted as images~~ cannot be interpreted
15 as perfect records of the atmospheric history. Indeed, due to the ~~very~~ process of gas trapping in the ice, two distinct effects ~~might~~ create discrepancies between the actual atmosphere's history and its imprint in the ice. The first one is due to the heterogeneous structure of the firn when transforming into airtight ice. The overall structure of the firn column is characterized by a progressive increase in density with depth, associated with the constriction of the interstitial pore network (Stauffer et al., 1985; Arnaud et al., 2000; Salamatín et al., 2009). At high enough densities, pores in the firn pinch
20 and isolate the interstitial air from the atmosphere (Stauffer et al., 1985). However, firn is a highly stratified medium (Freitag et al., 2004; Fujita et al., 2009; Hörhold et al., 2012; Gregory et al., 2014) and some ~~of the strata~~ especially dense strata (respectively less dense strata) might experience early (respectively late) pore closure when compared to the rest of the firn (~~Etheridge et al., 1992; Martinerie et al., 1992~~) (Etheridge et al., 1992; Martinerie et al., 1992; Fourteau et al., 2019). Consequently, abnormal strata might enclose older (respectively younger) air than their immediate surroundings, creating
25 age irregularities in the gas record (Mitchell et al., 2015; Rhodes et al., 2016). In turn, the irregularities appear as spurious values in the measured gas record (Rhodes et al., 2016; Fourteau et al., 2017). These anomalies have been referred to as layered gas trapping artifacts and do not reflect actual atmospheric variations. The second effect that creates differences between the atmosphere and its imprint in the ice, is due to the combination of diffusive air mixing in the firn (Schwander, 1989) and the progressive closure of pores in a firn stratum (Schwander et al., 1993; Mitchell et al., 2015). Consequently, the air enclosed

in a given ice layer does not originate from a single point in time, but is rather characterized by a continuous age distribution covering tens to hundreds of years (Schwander et al., 1988; Schwander et al., 1993; Rommelaere et al., 1997). Therefore, the concentration measured in an ice stratum is an average of atmospheric concentrations over a period of time. This effect removes the fast variability from the record, and has therefore been referred to as the smoothing effect (Spahni et al., 2003; Joos and Spahni, 2008; Köhler et al., 2011; Ahn et al., 2014; Fourteau et al., 2017). The degree of alteration between the ~~atmosphere and its recording~~ actual atmospheric history and the signal recorded in ice cores is strongly dependent on the local ~~precipitation~~ accumulation rate, with low accumulation ice cores being particularly affected both in terms of ~~gas-layered-layered trapping~~ artifacts (Rhodes et al., 2016; Fourteau et al., 2017) and in terms of smoothing (Spahni et al., 2003; Joos and Spahni, 2008; Köhler et al., 2011; Ahn et al., 2014; Fourteau et al., 2017). For layered gas trapping, Fourteau et al. (2017) report layering artifacts reaching up to 50ppbv in the Vostok methane record during the Dansgaard-Oeschger event 17 period. For smoothing, Spahni et al. (2003) report a gas age distribution in the EPICA Dome C ice core dampening atmospheric variability faster than a few hundred years. Similar degrees of smoothing have also been reported by Köhler et al. (2015) and Fourteau et al. (2017) for the ice cores of Dome C and Vostok, respectively.

15 In order to properly evaluate the composition of past atmospheres based on the gas records in polar ice cores, it is thus necessary to characterize the specificities of those two effects. ~~For the smoothing, the gas~~ Gas age distributions can be calculated ~~, for the purpose of estimating firn smoothing, and in~~ the case of modern ice cores, this may be accomplished by using gas trapping models ~~parameterized by firn pumping parametrized by firn air~~ and pore closure data (Buizert et al., 2012; Witrant et al., 2012) (Buizert et al., 2012). However, to estimate the gas age distribution in bubbles it is necessary to use a ~~closed porosity profile in firn. Such closed porosity profiles~~ depth-profile of the progressive closure of pores in the firn, quantifying the transformation of open pores into closed bubbles. Due to the re-opening of closed pores at the surface of the firn samples used for porosity measurements, such profiles of pore closure are associated with large ~~uncertainties (Schaller et al., 2017)~~ measurement uncertainties (Schaller et al., 2017; Fourteau et al., 2017). Moreover, a specific problem arises in the case of glacial ice cores from East Antarctica. They were formed under very low ~~precipitation temperatures and accumulation~~ rates, below ~~2cm.ice.yr⁻¹ (ice equivalent, Veres et al., 2013; Bazin et al., 2013);~~ 2cm ice equivalent yr⁻¹ (cm ice yr⁻¹, Veres et al., 2013; Bazin et al., 2013) ~~and temperatures that have no known equivalent nowadays. Thus,~~ that are not encountered on the present-day Antarctic and Greenland ice sheets. Thus, it is not possible to sufficiently constrain gas trapping models cannot be constrained and used to estimate, prohibiting robust estimation of the gas age distributions that were responsible for smoothing during glacial periods. Concerning layered Layered gas trapping, it on the other hand, originates from firn heterogeneities and is ~~therefore~~ a stochastic process. Moreover,



current gas trapping models do not fully represent the centimeter scale variability of the firm. The proper modelling of layered gas trapping is thus limited by the lack of knowledge of glacial firm heterogeneities and the difficulty to model gas trapping in a layered medium. Fourteau et al. (2017) proposed an empirical model to reproduce the artifacts observed in the Vostok ice core during the Dansgaard-Oeschger event 17. However, this model has only been applied to ~~this-particular-a single~~ event and might not be directly applicable for different periods and conditions.

For this work, we analyzed methane concentrations in five ice core sections from the East Antarctic ~~plateau~~Plateau. These sections cover accumulation and temperature conditions representative of East Antarctica for both the glacial and inter-glacial periods. The new data are used to ~~parameterize a revised version of the gas-layered-test~~ a layered gas trapping model proposed by Fourteau et al. (2017). ~~Our goal is to render this-~~ with the goal of rendering the model applicable to a range of accumulation rates characteristic of glacial and inter-glacial conditions on the East Antarctic ~~plateau~~Plateau. Then, the gas age distributions for each of the five records are estimated by comparing the low accumulation records with much higher accumulation records (Fourteau et al., 2017). Indeed, gas trapping occurs faster at high-accumulation sites, which are thus less affected by both smoothing and layering artifacts and can therefore be used to produce atmospheric scenarios with low levels of alteration. Finally, we simulate gas trapping in a hypothetical million-and-a-half years old and thinned ice core, including ~~gas~~-layering artifacts. We then simulate the process of methane measurements using a continuous flow analysis system, and of carbon dioxide measurements using a discrete method. The synthetic signals are finally compared to the original atmospheric reference to quantify the deterioration of atmospheric information due to firm smoothing and layered gas trapping.

2 Methods

2.1 Choice and description of the studied ice cores

The newly measured ice core sections originate from the East Antarctic sites of Vostok, Dome C, and Lock-In. The three sites are displayed on a map in Figure 1. The five measured sections correspond to time periods where high resolution methane measurements from high-accumulation ice cores are also available.

Modern Lock-In ice core:

The first studied ice core section is the upper part of the Lock-In ice core. The site of Lock-In is located 136km away from Dome C, towards the coast (3209m above sea level, coordinates 74°08.310' S, 126°09.510' E). The local accumulation is ~~of~~




Figure 1. Left: Map of the East Antarctic plateau Antarctica with the sites of Lock-In, Dome C, and Vostok, WAIS Divide, and Fletcher Promontory shown (made with the Quantartica package). Right: Map of Greenland with the site of NEEM shown (using Greenland Ice Mapping Project data: Howat et al., 2014).

$3.9\text{cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ $3.9\text{cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ (Yeung et al., 2019). About 80m of ice was analyzed for methane, ranging from 116m (near the firn-ice transition) to 200m depth. The last firn air pumping during Firn air sampling during the drilling operation was conducted at down to 108.3m depth. The gas record ranges from 500 about 400 to 3000yrBP (Before Present, with present defined as 1950 CE in this article).

5

Modern Dome C ice core:

A shallow ice core from Dome C was analyzed for depths ranging from 108 to 178m. Similarly to Lock-In, this section corresponds to ages ranging from 500 about 400 to 3000yrBP, but is characterized by a lower accumulation rate of $2.7\text{cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ $2.7\text{cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ (Gautier et al., 2016).

10

Holocene Dome C ice core:

Ice from the second drilling of the EPICA Dome C ice core (referred to as EDC99 hereafter) was measured from 312 to 338m depth. According to the AICC2012 chronology, the The gas ages range from 7950 to 9000yrBP 7800 to 8900yrBP, with an average accumulation of $3.1\text{cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ $3.1\text{cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ (Bazin et al., 2013; Veres et al., 2013). This period includes the 8.2ka cold event (Thomas et al., 2007), notably characterized by a sharp decrease in global methane concentrations (Spahni et al., 2003; Ahn et al., 2014).

15

Table 1. Summary of the different ice core sections studied in the article with their associated atmospheric references.

Ice Core Section	Coordinates	Accumulation (cm ie yr ⁻¹)	Atmospheric Reference
Lock-In Modern	74°08'S 126°10'E	3.9 ^a	WAIS Divide discrete ^b
Dome C Modern	75°06'S 123°20'E	2.7 ^c	WAIS Divide discrete ^b
Dome C 8.2ka	Same as above	3.1 ± 0.15 ^d	Fletcher-EDC composite ^e
Dome C DO6-9	Same as above	1.5 ± 0.15 ^d	WAIS Divide CFA ^f
Vostok D021	78°28'S 106°50'E	1.5 ± 0.05 ^d	Deconvoluted NEEM CFA ^g

a: Yeung et al. (2019) ; b: Mitchell et al. (2013) ; c: Gautier et al. (2016) ; d: Bazin et al. (2013); Veres et al. (2013) , uncertainties given as standard deviation over the section; e: EDC data from Loulergue et al. (2008) used for the lower part of the composite; f: Rhodes et al. (2015) ; g: Chappellaz et al. (2013)

DO6-9 Dome C ice core:

A section from the first drilling of the EPICA Dome C ice core (referred to as EDC96 hereafter) was analyzed for depths ranging from 690 to 780m. The AICC2012 chronology indicates gas ages covering the period 33000 to 41000yrBP, and an average accumulation rate of ~~1.5cm. ie. yr⁻¹~~ 1.5cm ie yr⁻¹ (Bazin et al., 2013; Veres et al., 2013). The CH₄ excursion associated with the
5 Dansgaard-Oeschger (DO) events 6 to 9 are included in this gas record (~~Huber et al., 2006; Chappellaz et al., 2013~~) (Huber et al., 2006; Chappellaz et al., 2013).

DO21 Vostok ice core:

The last analyzed section originates from the Vostok 4G ice core, for depths between 1249 and 1290m. The expected gas ages span from 84000 to 86500yrBP, with an average accumulation of ~~1.5cm. ie. yr⁻¹~~ 1.5cm ie yr⁻¹ (Bazin et al., 2013; Veres
10 et al., 2013). This section was chosen as it includes the record of the CH₄ excursion associated with the DO21 event, the fastest methane increase of the last glacial period (Chappellaz et al., 2013).

2.2 High-resolution methane measurements

The five ice core sections were analyzed for methane concentrations using a Continuous Flow Analysis (CFA) system, including a laser spectrometer based on optical-feedback cavity enhanced absorption spectroscopy (OF-CEAS; Morville et al., 2005),
15 at the Institut des Géosciences de l'Environnement (IGE), Grenoble, France. The ~~IGE's CFA apparatus and data processing procedure are described in greater details in Fourteau et al. (2017)~~ laser spectrometer was calibrated to the NOAA2004 scale (Dlugokencky et al., 2005) using three synthetic air standards of known concentrations. The five ice core sections were melted at an average rate of ~~3.6cm. ie. yr⁻¹, which allows to access~~ 3.6cm min⁻¹, which resolves centimeter scale variations in the methane record. ~~However~~ (Fourteau et al., 2017). Yet, the measured concentrations are affected by the preferential dissolution

of methane compared to nitrogen and oxygen in the meltwater (Chappellaz et al., 2013; Rhodes et al., 2013) (Chappellaz et al., 2013; Rhodes et al., 2013).

It is therefore necessary to apply a correction factor to account for ~~this~~ the solubility effect. However, this factor is a priori not known and potentially ~~ice-core dependent~~ differs between ice core measurement campaigns depending on factors such as the air content of the ice, or the precise CFA set-up. The methodology for correcting for methane dissolution ~~will be~~ is addressed in

5 Section ~~3.1 below~~ 3.1 below. Using a mixture of de-ionized water and standard gases, the analytical noise of the CFA system has been determined to be about 10ppbv peak-to-peak (Fourteau et al., 2017).

The obtained records present numerous gaps, ranging from a few centimeters to several meters. Several reasons explain the presence of such gaps. First, the space between consecutive melting sticks let modern air enter the CFA system, resulting in a contamination and abnormally high methane concentrations. Moreover, the presence of cracks and fractures in the ice might also let modern air enter the measured ice stick itself, also resulting in abnormally high concentrations. The moments of potential air intrusions were recorded during the measurement campaigns, and the data were screened to remove the resulting contaminations, creating gaps in the record (Fourteau et al., 2017). Finally, some of the ice was simply not available for this study, resulting in further gaps in the records. This notably explains the gap visible around the 1260 m depth in the Vostok record (Figure 6).

15 3 Results and Discussion

3.1 Correction for methane dissolution in the meltwater

In order to correct the measured mixing ratios for the preferential dissolution of methane in water, a correction factor is applied to the data to raise them to absolute values (Chappellaz et al., 2013; Rhodes et al., 2013). One way to estimate this correction factor is to compare the dissolution-affected CFA data with records that are already on an absolute scale.

20 The discrete methane measurements from the high accumulation WAIS Divide (WD) ice core published by Mitchell et al. (2013) cover the period from 500 to 3000yrBP. We therefore use the WD data to estimate the correction factors for the modern sections of Lock-In and Dome C.

Loulergue et al. (2008) performed discrete methane measurements in the Dome C EPICA ice core, that include the 8.2ka, DO6 to 9, and DO21 events. We use these discrete methane data to correct our CFA data for the dissolution, for the Holocene and DO6-9 Dome C ice core sections and for the DO21 event section in Vostok. The comparison between an already calibrated methane dataset and our corrected CFA measurements is given for the DO6 to 9 period in Figure S1 of the Supplement.

Note that in the case where our CFA data and the ~~absolute scale~~ measurements do not originate from the same ice core, the

match between the datasets are performed on sections with low methane variability to reduce the influence of firm smoothing. The signals after correction for the dissolution of methane are displayed in light blue in the upper panels of the Figures 2 to 6. The gas age chronologies indicated in the figures ~~will be constructed~~ are described in Section 3.4.1 of this article. The black superimposed curves in Figures 4 to 6 correspond to the part of the signal cleaned for layering artifacts, as described in Section 3.3.3. The correction factors ~~found for the solubility effect range between 1.12 and 1.14. They~~ are close to 1.13 the value of 1.125 reported by Fourteau et al. (2017) for the Vostok ice core with the same IGE CFA system, but larger than the 1.079 correction factor reported by Rhodes et al. (2013) for the NEEM ice core with a different CFA system. This difference could be due to the larger air content in NEEM as well as the different CFA systems, including the melthead geometry.

3.2 Atmospheric references

In Sections 3.3.2 and 3.4.1 below, weakly smoothed methane ~~gas~~ records will be used as atmospheric references to study ~~gas trapping in the layering artifacts and smoothing of~~ low accumulation ice cores. ~~As it is important that these gas records contain enough fast variability to be used as atmospheric references, we use~~ To derive the atmospheric references we used methane gas records from ~~much~~ higher-accumulation ice cores ~~(Fourteau et al., 2017)~~ where higher frequency variations are preserved relative to the low-accumulation ice cores (Fourteau et al., 2017). When possible, we use high-accumulation records from ice cores drilled in Antarctica. Otherwise, in order to compare the high and low-accumulation records we need to estimate the methane inter-hemispheric gradient.

For the modern Lock-In and Dome C sections, we ~~use~~ used the discrete measurements from the WD ice core, also used for correcting the methane preferential dissolution (Mitchell et al., 2013). For the Holocene section of EDC99, we used CFA data from an ice core drilled at Fletcher ~~promontory~~ Promontory as the atmospheric reference (Robert Mulvaney and Xavier Faïn, personal communication), ~~completed with~~ discrete data from Louergue et al. (2008) for the oldest part of the period (depths below 330m in Figure 4). Using the low-accumulation EDC99 record for the lowest part does not deteriorate the quality of the composite for our application, as this part does not include fast atmospheric variations to be smoothed, and using a low-accumulation record is therefore appropriate in this specific case. The dating of the Fletcher-EDC99 composite was made consistent with the WAIS Divide chronology by manually selecting tie points (Buizert et al., 2015). For the DO6-9 events section in the EDC96 ice core, we ~~use~~ used high-resolution CFA data from the WD ice core (Rhodes et al., 2015). The WD gas record dating has been made consistent with the AICC2012 and GICC chronologies following Buizert et al. (2015). Finally, the atmospheric reference used for the Vostok DO21 period is based on the NEEM CFA ~~record~~ data published by Chappellaz et al. (2013). Chappellaz et al. (2013) propose two CFA records obtained with two different spectrometers. For this study, we used

the average values of the two instruments. As the Vostok and NEEM sites are located in different hemispheres, it is necessary to take into account the inter-hemispheric methane gradient between the two sites. We to make the two records as consistent as possible. Using the Vostok and NEEM records, we evaluated this inter-hemispheric gradient to be 30ppbv for the DO21 period. This value is in line with the work of Dällenbach et al. (2000), and has been corroborated using the EDML methane record (EPICA Community Members et al., 2006). Yet, it remains possible that the inter-hemispheric gradient is was not constant during this period of global methane rise. The use of a high resolution and weakly smoothed Antarctic record as the atmospheric reference would resolve this inter-hemispheric gradient problem, but such a record is not available. Using a spline version of the The original data of Chappellaz et al. (2013), we observe that it is not possible to find a gas age distribution able to reproduce the smoothing over the all section (Figure S1 present various gaps, that were filled by computing a splined version of the original NEEM CFA record. The spline was chosen not to induce smoothing in the NEEM record and to overlap with the original dataset in parts where data already existed. It is shown together with the original Chappellaz et al. (2013) data in Figure S2 of the Supplement). This suggests.

Yet, we observed that the NEEM record cannot be directly used as an atmospheric reference for Vostok, due to the smoothing of the fast event at the onset input for the DO21 period. As explained in Section S2 of the Supplement, the first feature of the DO21 event (around the 1260m depth in Figure 6) is partially smoothed in the NEEM record. To retrieve the full variation amplitude of this fast event, we used a "deconvolution" technique the deconvolution technique described in Witrant and Martinerie (2013) and Yeung et al. (2019). To be applicable this method needs as input the gas age distribution responsible for the smoothing of the NEEM record. For this, we chose the age distribution estimated with a gas trapping model at the modern site of Siple Dome which has an accumulation 10.8cm ie yr^{-1} (Witrant et al., 2012), similar to the ones in Witrant and Martinerie (2013) and Yeung et al. (2019) accumulation of NEEM during the DO21 period (around 11.3cm ie yr^{-1} Rasmussen et al., 2013). In order to test the sensitivity of our results to the deconvolution step, we performed several deconvolutions with age distributions of modern sites ongoing accumulation values above and below the range of potential NEEM accumulations during the DO21 period. The results are presented in Section S2 of the Supplement and indicate that the deconvolution of NEEM is well constrained, and only weakly depends on the specific choice of the age distribution. The effect of the deconvolution is mainly to increase the amplitude of the fast event feature at the onset of the DO21, which in turn increases the consistency between the NEEM and Vostok methane records. Note that the deconvolution method requires gas age distributions for the NEEM ice core as input. The age distributions enclosed in NEEM vary over the DO21 event due to the large variation in accumulation at this site during the period. The gas age distributions in the NEEM ice core are estimated with gas trapping models forced with various accumulation conditions (Boeck et al., 2012, personal communication Patricia Martinerie). The The drilling sites of the

atmospheric references are displayed as orange dashed lines in the lower panels of Figure 2 to 6. in Figure 1.




Figure 2. Measured and modeled modern Lock-In methane profile. Upper panel: High resolution methane measurements in light blue. Atmospheric reference in orange. Lower panel: Modeled layering artifacts. The black curve is the expected signal without artifacts. The blue and yellow spikes respectively are randomly distributed early and late closure artifacts, respectively. The blue and yellow shaded areas respectively correspond to the range of early and late artifacts with density anomalies up to two standard deviations. The dashed orange curve is the record used as the atmospheric reference.




Figure 3. Same as Figure 2 for the modern Dome C section.




Figure 4. Same as Figure 2 for the Holocene EPICA Dome C section. In addition the superimposed black signal in the upper panel shows the data cleaned of layering artifacts




Figure 5. Same as Figure 4 for the DO6-9 EPICA Dome C section.

3.3 Layered trapping artifacts

It has been observed in the high-accumulation firm of DE08 that not all layers close at the same depth (Etheridge et al., 1992), and that the air in summer layers is generally older than in winter layers. Building on this idea, Mitchell et al. (2015) and Rhodes et al. (2016) proposed that the layering observed at the firn-ice transition induces gas stratigraphic irregularities and




Figure 6. Same as Figure 4 for the DO21 Vostok section.

age inversions in the record. Rhodes et al. (2016) and Fourteau et al. (2017) used such stratigraphic heterogeneities to explain the observation of sudden variations in high resolution methane records. These abrupt methane variations do not correspond to actual atmospheric variations and have been called layered trapping artifacts, or more simply layering artifacts. The mechanism producing them is that layers closing especially early (or late) contain air older (or younger) than their immediate surroundings.

5 During periods of variations of atmospheric composition, the methane mixing ratios in these layers are thus different than in their surroundings, and appear as anomalous values (Rhodes et al., 2016).

In accordance with Rhodes et al. (2016) and Fourteau et al. (2017), we observe abrupt centimeter-scale variations in the methane records, records during periods when the atmospheric CH_4 concentration was varying, that we interpret as layering

artifacts. We visually identified such layering artifacts as spikes with widths of a few centimeters and whose concentrations

10 are larger than the analytical noise of $\sim 10\text{ppbv}$, during periods of variations in atmospheric concentrations. These layering

artifacts. Despite the effort to clean the record of modern air contaminations, it is possible that some contamination spikes remain and would then be wrongly interpreted as layering artifacts. However, the presence of negatively orientated spikes that cannot be attributed to contamination, and the overall occurrence of the spikes near periods of fast atmospheric methane variations confirm that these centimeter-scale abrupt variations are mostly due to the mechanism of layered gas trapping. They

15 are notably well marked during the onset of DO events 8 and 21, corresponding to 750 and 1252m depth in the EDC96 and Vostok ice cores, respectively, in Figure 5 and 6. A zoom of the EDC96 record is available in Figure S2-S4 of the Supplement to display the layering artifact examples of individual layering artifacts. They appear as reduced methane concentration layers

near the onset of the DO events, ~~and cannot therefore be explained by contamination issues~~. The layered trapping artifacts tend to disappear in the absence of marked atmospheric variations. This explains the absence of important layering artifacts in the Lock-In and Dome C modern records (Figures 2 and 3).

The usage of high-resolution measurements allow us to easily distinguish layering artifacts as abrupt spikes exceeding the analytical noise in the record. However, with a lower-resolution technique, such as discrete measurements, it is possible to inadvertently measure a layering artifact without realizing that it is not representative of its surrounding concentrations, which would result in an anomalous point in the record. As a specific example, Fourteau et al. (2017) pointed out that one of the data points of the DO8 event in the EPICA Dome C methane record published by Louergue et al. (2008) might have been sampled in an early closure layer, and thus represents a spurious value in their record. By comparing our new high-resolution record with the one of Louergue et al. (2008), we confirm that this data point does not correspond to atmospheric variability, and that it was sampled in a zone with a high number of early closure artifacts (see Figure S3-S5 of the Supplement). In the case where such an abrupt variation is observed with discrete measurements, an additional sample should be measured in the vicinity of the first sample in order to confirm that the variability is not due to layered gas trapping.

Important late closure artifacts are unusual in methane gas records. The new measurements confirm this observation. Late closure artifacts, that should appear as positive methane anomalies at the onset of the DO events, are almost absent in our measurements. This can be explained by the fact that the surrounding firn layers are already sealed and prevent ~~long-distance gas transport~~ gas transport to greater depths, which in turn prevents the late closure layers ~~to enclose~~ from enclosing young air (Fourteau et al., 2017).

3.3.1 Impact of chemistry on layered gas trapping

The EDC96 methane data suggest a lack of layering artifacts affecting the DO6 event (around ~~the~~ 700m depth in Figure 5). We counted only 14 layering artifacts that clearly stand out of the analytical noise on the 20m long section between 695 and 715m. For a comparison, we counted 15 artifacts on the 4m long section from 718 to 722m (DO7 event). These sections are not directly comparable as they did not undergo the same atmospheric methane variations during bubble enclosure, but it nonetheless suggests that the upper section of the DO6-9 record shows a lower number of layering artifacts per meter. Previous studies have highlighted the role of chemical impurities for the presence of dense strata in polar firn. Hörhold et al. (2012) and Fujita et al. (2016) argue that the presence of ions can soften the firn and facilitate the densification of some strata. To evaluate the potential impact of ions on layered gas trapping artifacts in the EDC96 methane record, we compared the high resolution methane data with total calcium data measured in the ice phase (Lambert et al., 2012). We chose to focus on calcium

since high resolution data are readily available and [since](#) Hörhold et al. (2012) observed a correlation between calcium ~~content~~ ~~and the density of firn layers~~ and [density variability in deep firn](#). However, as pointed out by Hörhold et al. (2012) and Fujita et al. (2016), ~~calcium should not be seen as it does not entail that calcium is~~ the ion responsible for the ~~preferential densification of firn, but rather as a proxy for the general ionic content~~ [establishment of deep firn stratification. Indeed, calcium is correlated](#)

5 [with other ion species that could be the cause for the preferential densification of some firn strata \(Fujita et al 2016\)](#). A comparison of the ~~two calcium and methane~~ datasets is shown in Figure 7. The depth difference between the EDC96 ice core, in which the methane measurements were performed, and the EDC99 ice core, in which the calcium measurements were performed, was taken into account using [18 individual](#) volcanic markers (Parrenin et al., 2012). The calcium data clearly show a transition between two parts: ice below the depth [718m](#) is characterized by relatively high calcium values and a high variability, while ice

10 above 718m is characterized by low calcium values and a low variability. The transition between these two parts corresponds to the high methane plateau of the DO7 event. It also matches with the transition described above between the two regions of large and small numbers of layering artifacts. It therefore appears that the number of significant artifacts (standing out of the analytical noise) ~~is~~ [may be](#) correlated to the variability of calcium concentrations. ~~It is consistent with the mechanisms proposed in the literature: calcium variability is associated with a general ionic concentration variability that creates particularly dense strata~~

15 ~~in the firn (Hörhold et al., 2012; Freitag et al., 2013; Fujita et al., 2016). These strata then close in advance and trap young air that appears as layering artifacts (Rhodes et al., 2016; Fourteau et al., 2017). We then evaluated the calcium levels for the other studied ice core sections. Using the data of Lambert et al. (2012), it appears that the 8.2ka and modern periods of Dome C are characterized by low calcium levels and variability. Similar low calcium levels and variability at Lock-In have been observed by chemistry measurements (data not presented here). Unfortunately, high resolution chemistry data are not available~~

20 ~~for Vostok. Assuming that the calcium variability is similar at Dome C and Vostok, the data from Lambert et al. (2012) indicate that the DO21 period displays low calcium level and variability, while the DO17 period studied by Fourteau et al. (2017) shows high calcium level and variability, similar to~~ [While this preliminary result appears to support the idea of ion-induced layering artifacts, further work is still required on this question. One possible perspective would be to obtain high resolution methane and chemistry records from the DO7 to 9 period same ice core section, in order to study the potential concordance of layering](#)

25 [artifacts with ion rich strata.](#)




Figure 7. Upper panel: calcium concentrations measured in the EPICA Dome C ice core section covering the DOs 6 to 9 (Lambert et al., 2012). Lower panel: methane mixing ratios measured in this study.

3.3.2 Modeling layered trapping artifacts

The ability to model layered gas trapping helps ~~quantifying and predicting~~ quantify and predict its impact on gas records. For this purpose, we revised the simple layered gas trapping model proposed by Fourteau et al. (2017) for the Vostok DO17 section. Our aim is to parametrize the model in order to reproduce the layering artifacts observed in the new five measured sections, as well as in the previously published DO17 section of the Vostok core. This layered trapping model is based on the physical assumption that the density anomaly of a layer (its density difference ~~with the bulk behavior~~ relative to the meter-scale density profile, once centimeter-scale variability is removed) can directly be converted into a trapping depth anomaly (the difference in closing depth ~~with the bulk-average~~ behavior). The ~~computation of~~ depth anomalies is given by:

$$\Delta z = \frac{\Delta \rho}{\partial_z \rho} \quad (1)$$

where Δz is the depth anomaly of a given layer, $\Delta \rho$ its density anomaly, and $\partial_z \rho$ the derivative of the ~~bulk-meter-scale~~ firm density with depth in the trapping zone. For the rest of the article $\partial_z \rho$ will be referred to as the densification rate. The methane concentration in an early trapping layer is then simply the concentration ~~trapped by the bulk firm~~ that would normally be trapped at a depth Δz below. The Fourteau et al. (2017) model ~~is~~ based on the computation of the age anomalies of abnormal layers (the time difference between the closure of the layers and the ~~bulk-average~~ behavior). However, this age-based formulation is sensitive to the dating of the ice core, and can lead to abnormally strong artifacts ~~where-if~~ the chronology is poorly constrained.

On the other hand, the depth-based formulation proposed here (Equation 1) is not affected by dating uncertainties.

The model requires a densification rate for the firn-ice transition zone. Observations of density profiles in Dome C and Vostok proposed by Bréant et al. (2017) reveal densification rates around $2.2\text{kg}\cdot\text{m}^{-4}$ $2.2\text{kg}\text{m}^{-3}\text{m}^{-1}$ for both sites. This value also applies to the Lock-In firn as inferred from high resolution density measurements (Fourteau et al., 2019). Therefore, the densification rate was set at $2.2\text{kg}\cdot\text{m}^{-4}$ $2.2\text{kg}\text{m}^{-3}\text{m}^{-1}$ for all sites studied here. Finally, the model requires a typical density anomaly in the closing part of the firn. Based on the linear relationships of density standard deviation with temperature and accumulation proposed by Hörhold et al. (2011), Fourteau et al. (2017) estimated the typical density anomaly to be $5 \pm 2\text{kg}\cdot\text{m}^{-3}$ $5 \pm 2\text{kg}\text{m}^{-3}$ for the Vostok firn during the glacial DO17 event. Moreover, Hörhold et al. (2011) report a density standard deviation of $4.6\text{kg}\cdot\text{m}^{-3}$ $4.6\text{kg}\text{m}^{-3}$ around the firn-ice transition of the Dome C site. As a first guess, we set the density anomalies $\Delta\rho$ to obey a zero-centered Gaussian distribution of standard deviation $5\text{kg}\cdot\text{m}^{-3}$ $5\text{kg}\text{m}^{-3}$. Similarly to Fourteau et al. (2017), the depth anomalies of late closure layers are ~~reduce~~ reduced by 75% in order to replicate their low impact in the measured records.

~~While this parametrization produces adequate results for most of the gas records studied here, it results in a clear underestimation of the layering artifacts for the DO events 7 and 8 in the EDC96 ice core (Figure S4 of the Supplement). Our understanding is that the layering model fails to take into account the impact of high calcium variability during this period. To introduce the influence of calcium in the model, we simply increase the typical density anomaly during period of high calcium variability, by applying an enhancing factor of 1.5. This enhancing factor was chosen to produce visually correct results in the case of the high calcium variability period of the DO7 to 9 (depths below 718m in the EDC96 record). For consistency, we also applied this enhancing factor for the DO17 period in Vostok, as this period displays a calcium variability similar to the DO 7 and 9 period.~~

The modeled artifacts are displayed below the newly measured signals in Figures 2 to 6, ~~and as well as~~ the modeled artifacts for the DO17 in Vostok with the new calcium parametrization are displayed event in Vostok in Figure 8. The model produces layering artifacts in the expected parts of the records (mostly right after the onsets of DO events) and with the expected ~~signs and amplitudes~~ sign and amplitude. This is visible on Figures 2 to 6, where the layering artifacts of the record fall within the expected envelopes of the model results. However, it is hard to reproduce the precise distribution of layering artifacts. This is due to the fact that the position of layering artifacts is random, and that the observed layering artifacts in an ice core record are one single outcome among many possibilities. ~~The simple parametrization we derived thus appears to provide~~ Nonetheless, the simple model presented here provides reasonable results for accumulation conditions ranging from 3.9 to

~~1.3cm.ie.yr⁻¹~~1.3cm ie yr⁻¹, encompassing most of the climatic conditions of the East Antarctic ~~plateau~~Plateau for both the glacial and interglacial periods.




Figure 8. Same as Figure 2 for the DO17 Vostok section. The methane data of the upper panel are from Fourteau et al. (2017) and the atmospheric reference ~~of the lower panel~~ is from Rhodes et al. (2015).

3.3.3 Removing layered trapping artifacts

Since layered gas trapping artifacts are not due to chronologically ordered atmospheric variations, a proper interpretation of gas records requires them to be ~~cleaned~~removed from the datasets. As pointed out by Fourteau et al. (2017), in some sections of the records these artifacts only exhibit negative or positive methane anomalies. They cannot therefore be removed by applying a running average without introducing bias in the signal.

~~To~~ As the layering artifacts can be visually distinguished, it would be possible to remove them manually from the data.
~~However, such procedure would be cumbersome due to the large amount of CFA data points. Hence, to~~ clean the new three methane signals presenting layering artifacts (EDC96, EDC99 and Vostok) we use ~~the cleaning procedure~~ a recursive cleaning procedure similar to the algorithm described by Fourteau et al. (2017). ~~Briefly, this~~ The cleaning algorithm starts by estimating a smooth signal that should represent the measured signal free of layering artifacts and analytical noise. For this purpose, a running median is first computed to remove the layering artifacts while minimizing bias. Then, the signal is smoothed with a binned average, and interpolated back to high resolution using an interpolating spline. Then, the analytical noise is estimated ~~and the data clipped to remove artifacts larger than the analytical noise~~ using the Normalized Median Absolute Deviation of

the data (NMAD, Rousseeuw and Hubert, 2011; Fourteau et al., 2017). The data are clipped above 2.5 times the value of the NMAD, in order to trim a part of the artifacts. The algorithm is ~~looped~~ until the signal is determined to be free of layering artifacts. The signal is considered to be free of artifacts when the NMAD (estimation of noise without the layering artifacts) and the standard deviation (estimation of noise with the layering artifacts) are similar (Fourteau et al., 2017). This recursive method produces a progressive removal of the layering artifacts.

This procedure was successfully applied to the Holocene section of EDC99, that only exhibits a few ~~artifacts~~ clear artifacts near the 323m depth. The resulting cleaned signal is displayed in the upper panel of Figure 4 as a black curve. ~~However, direct~~ Note that the cleaning algorithm also removes a small part of analytical noise. However, this small removal of the analytical noise is negligible and does not influence our conclusions.

Direct application of this algorithm to the EDC96 DO6-9, and Vostok DO21 records leads to an ineffective removal of layering artifacts in periods of fast methane rise. After investigation, it appears that the main issue the algorithm is facing is the determination of an artifacts free signal by the ~~rolling~~ median. In some ~~section~~ sections of the ice cores, the signal displays high ~~concentrations~~ numbers of artifacts with methane anomalies of 50ppbv or more, as seen for instance in the onset of the DO8 in Figure 5. In these conditions, the ~~rolling~~ median is influenced by the presence of the artifacts and yields a biased signal. To circumvent this problem, we manually provided a set of values that correspond to the artifact free part of the signal. As it is easy to visually distinguish the layering artifacts from the rest of the signal, the manual selection of these points is fairly unambiguous and straightforward. The rest of algorithm then proceeds as normal, which allows us to use the automated determination of analytical noise and cleaning of the data. This also ensures a better consistency of cleaning between the different CFA datasets. The cleaned EDC99 Holocene, EDC96 DO6-9 and Vostok DO21 signals are displayed in the upper panels of Figures 5 and 6 in black.

3.4 Smoothing in East Antarctic ice cores

The five ice core sections measured in this article all originate from the East Antarctic ~~plateau~~ Plateau. The low temperatures and aridity of this region result in a slow densification of the firn, and a slow bubble closure. Therefore, the gas enclosed in a given ice layer tends to have a broad age distribution. This leads to an ~~important~~ significant smoothing of atmospheric fast variability observed in East Antarctic ice cores (Spahni et al., 2003; Joos and Spahni, 2008; Köhler et al., 2011; Ahn et al., 2014; Fourteau et al., 2017).

3.4.1 Estimating gas age distributions

Since smoothing in an ice core record is a direct consequence of the gas age distribution (GAD), the knowledge of GADs for various temperature and accumulation conditions is necessary to predict the impact of smoothing on gas signals. We thus apply the GAD extraction method proposed by Fourteau et al. (2017) to the five new high resolution records.

- 5 This method, designed to estimate the GAD of low accumulation records, is based on the comparison with a weakly smoothed record derived from a high accumulation ice core, used as an input atmospheric reference (see Section 3.2). ~~An optimal log-normal age distribution is derived to best match the measured signal with the smoothing 3.2).~~ The idea of the method is to find a GAD that is able to smooth the atmospheric reference into the low-accumulation record. We thus searched for the GAD that minimizes the Root-Mean-Square Deviation between the CFA measurements and the smoothed version of the at-
- 10 mospheric reference ~~by the GAD. Using a log-normal law allows.~~ In order to have a well-defined problem in ~~the mathematical sense.~~ A mathematical sense, the GAD of the low-accumulation ice core is assumed to be a log-normal function. Such a log-normal distribution is fully defined by two ~~independents parameters. The GAD extraction problem is thus independent~~ parameters. Finding the best GAD to smooth the atmospheric reference is then reduced to the recovery of a pair of optimal parameters. Nonetheless, log-normal ~~laws distributions~~ exhibit a large range of shapes that can adequately represent ~~the~~
- 15 ~~age distributions (Köhler et al., 2011; Fourteau et al., 2017).~~ During the extraction procedure, a new gas age chronologies are produced ~~age distributions (Köhler et al., 2011; Fourteau et al., 2017).~~ In order for the GAD extraction to perform well, it is also necessary to have a well defined gas age chronology for the low-accumulation ~~records. They are constructed record, so that its methane variability is well aligned with the atmospheric reference. This chronology is built recursively during the GAD extraction procedure by manually selecting tie-points, in order~~ to be consistent with the already-existing ~~chronologies of the~~
- 20 ~~high-accumulation chronology of the~~ atmospheric reference (Fourteau et al., 2017). The GAD extraction procedure is designed to match the CFA record and the convoluted atmospheric reference only where CFA data are available. We thus do not have to extrapolate the CFA data within the gaps of the record. However, the absence of gaps in the records would have reduced the uncertainty of the estimated GADs, as the additional data would have provided more constraints on the GAD estimation.
- The gas age distributions obtained for the five new ice core sections are displayed as solid lines in Figure 9, and an uncertainty
- 25 analysis is available in Section S5 of the Supplement. The parameters defining the different log-normal distributions are listed in Table 2. The standard deviation in this table should not be directly interpreted as the broadness of the age distribution and therefore its degree of smoothing. Indeed, some distributions exhibit strong asymmetries that tend to increase the standard deviation values. For instance, the Vostok DO21 age distribution has a standard deviation value much larger than the other distributions, but does not induce a much larger smoothing.




Figure 9. Gas age distributions in various ice cores. The distributions are centered around their median values. The left panel displays Holocene GADs and the right panel glacial period GADs. The DO21 Vostok, DO6-9 EDC96, Holocene EDC99 and modern Lock-In and Dome C age distributions shown as solid lines were determined in this study. The dashed Vostok DO17 distribution was also determined by comparison with a high-accumulation record (Fourteau et al., 2017). The dashed Vostok firn pumping distribution was obtained with a gas trapping model, constrained with firn air pumping data (Witrant et al., 2012).

Table 2. Parameters defining the log-normal gas age distributions derived by comparison with weakly smoothed records. The location and scale parameters correspond to the μ and σ parameters used to define the log-normal laws in Equation 1 of Köhler et al. (2011).

Site and Period	Location	Scale	Mean (yr)	Std Dev (yr)
Lock-In Modern	4.063	1.333	141	313
Dome C late-Modern	4.156	1.611	234	823
Dome C 8.2ka	4.618	1.100	185	285
Dome C DO6-9	4.765	1.231	250	472
Vostok DO21	4.339	1.988	552	3950

3.4.2 Smoothing and accumulation

The degree of smoothing in a gas record is strongly linked to the accumulation rate under which the gases are trapped, with low accumulation sites exhibiting a stronger degree of smoothing (Spahni et al., 2003; Joos and Spahni, 2008; Köhler et al., 2011). First, a lower accumulation is generally associated with a lower temperature. Low-temperature ice deforms less easily and ~~slowing~~ the densification process ~~is therefore slower and such that~~ bubble closure spans a larger time period. Moreover, a low accumulation is also associated with a slow mechanical load increase of the ice material, also resulting in a slower densification and a longer bubble closure period.

Two Our results suggest that, during the glacial period, East Antarctica ice cores are affected by the same level of smoothing, that can be represented with the same gas age distribution. We produced two new glacial period gas age distributions were derived in this work (EDC96 DO6-9 and Vostok DO21). This is, that are to be added to the previously published GAD obtained for the Vostok site during the DO17 event using the same GAD extraction method ($1.3\text{ cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ accumulation rate; Fourteau et al., 2017).

5 These three age distributions of East Antarctic sites under glacial conditions are displayed in the right panel of Figure 9. The smoothing they induce is represented for the DO6 to 9 and DO21 events in Figure 10. It appears that the two Vostok distributions lead to a similar smoothing. On the other hand the GAD obtained for EDC96 for the DO6-9 events is significantly broader than the Vostok ones and results in a stronger smoothing, especially visible on at the onset of the DO21 event. This is surprising as the Vostok DO21 and Dome C DO6-9 records have the same accumulation rate, and should therefore present

10 similar age distributions.

However, the uncertainty analysis in Section S5 of the Supplement reveals that the age distribution of the EDC96 record is poorly constrained. This indicates that the smoothing of DO6-9 period is weakly sensitive to the choice of GAD, and therefore that a large range of GADs results in adequate smoothing for the EDC96 record. On the other hand, the DO21 period is very sensitive to the choice of GAD, and much fewer age distributions are able to reproduce the smoothing of the Vostok DO21

15 record. Our understanding is that because of its shape and its fast atmospheric rate of change (Chappellaz et al., 2013), the first feature of DO21 is very sensitive to the choice of GAD, despite a gap in the record, and is therefore a good discriminant between age distributions. On the other hand, the step-like features of the DO6-9 record are less sensitive to the choice of GAD, which leads to a less-well constrained extraction of age distributions. Consequently, the distributions obtained for glacial Vostok (DO17 and 21 events) also appear to be are also suited for the smoothing of DO6 to 9 events in EDC96, as seen in the

20 lower panel of Figure 10. Moreover, the uncertainty analysis reveals that the distribution extracted from the EDC96 record is poorly constrained, with a large range of GADs resulting in adequate smoothing (Figure S11 in Section S5 of the supplement). It appears that the This suggests that the smoothing of the three glacial records exhibits similar degree of smoothing, and therefore enclose gases with the same gas age distribution. We are similar, and that the three ice cores therefore present similar gas age distributions. We thus propose to use a common gas age distribution to represent the smoothing in ice cores with accumula-

25 tions below $2\text{ cm}\cdot\text{ie}\cdot\text{yr}^{-1}$ $2\text{ cm}\cdot\text{ie}\cdot\text{yr}^{-1}$. The distribution proposed by Fourteau et al. (2017) is a good candidate for this common distribution, as it reproduces well the smoothing in the EDC96 DO6-9 record and the Vostok DO17 and 21 records. Moreover, its shape is a compromise between the two other glacial GADs proposed in this article (right panel of Figure-9). Finally, the similarity of the smoothing encountered during the DO17 and DO21 events suggests the absence of significant impurity effect

on the GADs.




Figure 10. Comparison of smoothing resulting from for the three different glacial age distributions, and the modern Vostok GAD constrained with firm air pumping data. Atmospheric references (dashed lines) have been smoothed by the four different GADs and are displayed overlying the measurements (light blue). The upper panel shows the DO21 event as measured in Vostok, and the lower panel shows the DO-6 to 9 as measured in EDC96.

Comparing Our results also indicate that the glacial and inter-glacial smoothings of East Antarctic ice cores are relatively similar. Indeed, comparing the typical GAD of the glacial period with the ones obtained for the Holocene and modern periods indicates that the later-latter lead to a slightly smaller degree of smoothing. This is illustrated with the smoothing of the 8.2ka event displayed in Figure 11. It is consistent with the expectation that the smoothing during the low-accumulation glacial period should be more important-greater than during the higher-accumulation Holocene period. However, the doubling of accumulation between the DO17 event at Vostok and the 8.2ka event in EDC99 does not result in a drastic decrease of the smoothing. It is also noteworthy that for the Dome C site, the transition from 8.2ka conditions (3.1cm.ice.yr^{-1} 3.1cm ice yr^{-1} accumulation rate) to modern conditions (2.7cm.ice.yr^{-1} 2.7cm ice yr^{-1} accumulation rate) does not significantly change the

degree of smoothing. It thus appears that for accumulations below ~~3.9cm. ie. yr⁻¹~~ 3.9cm ie yr⁻¹ the degree of smoothing only slowly increases with the ~~diminution of the precipitation~~ decrease of the accumulation rate.




Figure 11. Comparison of the smoothing induced by three different Holocene and a modern age distributions. The atmospheric reference for the 8.2ka event (dashed line) has been smoothed by the four different GADs and displayed over the EDC99 measurements (light blue).

Finally, the expected smoothing using the firn pumping GAD estimation for modern Vostok is also displayed in Figure 10 in black (Witrant et al., 2012). This GAD was obtained with a method independent of the GAD extraction technique of this article. The smoothing estimated for modern Vostok is stronger than the one measured in the glacial Vostok ice core. This can be seen ~~on~~ in the upper panel of Figure 10, where the smoothing induced by the Vostok firn pumping distribution results in a signal with ~~less variability~~ a smaller magnitude than the CFA measurements. This confirms the observation of Fourteau et al. (2017) that the firn pumping GAD of modern Vostok leads to a stronger smoothing than the one observed during the DO17 event. This stronger smoothing in a modern ice core is in contradiction with the Holocene and modern GADs reported in this article for Dome C and Lock-In. This discrepancy could result from the large uncertainties associated with the firn-model-based GAD, which notably arise from the poor constraints on the closed porosity profiles used (Schaller et al., 2017). Unfortunately, high-resolution methane measurements of the late Holocene period in the Vostok ice core are not available to apply the GAD extraction method.

3.4.3 Effect of smoothing on record synchronization

Compared to the GADs of the Holocene and modern periods, the GADs of the glacial period ~~exhibits~~ exhibit a stronger degree of skewness (see Figure 9). This asymmetry of the distributions means that the smoothing not only removes variability, but also induces phase shifts during fast variations (Fourteau et al., 2017). This is especially visible ~~on~~ in the variation recorded at depth 1260m in the Vostok DO21 ice core section, displayed in the upper panel of Figure 10. The peak of the event is recorded a couple of meters before its position in the absence of smoothing. In terms of ages, this corresponds to an error of about 85yr

in the Vostok gas chronology. Studies relying on the synchronization of atmospheric variability between ice cores should be aware of this potential bias (Bazin et al., 2013; Veres et al., 2013). ~~The best synchronization should either~~ To produce consistent chronologies, the synchronization should be performed on signals with a similar degree of smoothing, ~~or (either by convolving the higher-accumulation record or deconvolving the lower-accumulation one).~~ Otherwise, the phase shifts should be taken into account in the age uncertainty estimates.

4 Loss of climatic information in a deep and thinned ice core from East Antarctica

East Antarctica is a region of particular interest for the drilling of deep ice cores. Indeed, thanks to low accumulation rates, it likely contains the oldest stratigraphically-undisturbed ice on earth. There is currently a search for very old ice, with ages potentially dating back one-and-a-half million years (Fischer et al., 2013; Passalacqua et al., 2018). When retrieved, such an ice core will be characterized by the gas trapping of low-accumulation East Antarctic ~~plateau~~ Plateau ice cores. Moreover, the old ice will be located close to the bedrock and therefore will be thinned by strain.

In this section, we estimate the potential differences between atmospheric signals and their measurements in a theoretical one-and-a-half million years old ice core, for methane and carbon dioxide. ~~For this purpose~~ Here, we consider that the gas trapping occurs under an accumulation rate of ~~2cm.1e.yr⁻¹~~ 2cm 1e yr⁻¹. We suppose that the enclosed gases follow the age distribution obtained for Vostok during the DO17 period, established to be representative of East Antarctic glacial smoothing in Section 3.4.1. The presence of layered trapping artifacts is simulated using the model parametrized in Section 3.3.2, ~~with the assumption of high calcium variability.~~ Finally, we assume that the thinning of the ice results in a resolution of ~~10,000yr.m⁻¹~~ 10,000yr m⁻¹ (Passalacqua et al., 2018). This implies that a given ice layer at the bottom of the core is 200 times thinner than at the firn-ice transition. This has to be taken into account as analytical techniques have limited ~~spatial-depth~~ spatial-depth resolutions below which variations are no longer resolved. In particular, layering artifacts will no longer be visible as individual events in the record. Note that in this section we do not take into the diffusive mixing of gases within the ice matrix, that would further smooth the trace gas records (Bereiter et al., 2014).

4.1 Methane alterations

As a first case, we study the alterations of a methane record measured using a CFA system analogous to the one currently used at IGE. For the sake of simplicity, we assume that the atmospheric history recorded in this theoretical one-and-a-half million year-old ice core is similar to the DO15 to 17 events of the last glacial period. Hence, we simply use the WD CFA methane measurements as the atmospheric reference (Rhodes et al., 2015). We first determine the initially enclosed signal

using the layering artifacts and smoothing models, and apply a thinning factor of 200~~times thinning factor~~. The resulting signal is displayed in ~~blue in the upper panel of~~ light blue in Figure 12. Then, we simulate the process of CH₄ measurements by applying the smoothing induced by the CFA system. For this, we use the CFA impulse response derived by Fourteau et al. (2017). The end result is displayed in ~~blue in the lower panel of~~ red in Figure 12. As this is a synthetic signal, we plotted the

5 data on a relative depth scale, whose zero was arbitrarily set in the record. The layering artifacts are no longer visible as they have been smoothed out by the CFA measurement. However, they affect the measured signal due to their asymmetry. This is illustrated by the difference between the ~~blue-red~~ and green curves in ~~the lower panel of~~ Figure 12, that ~~respectively~~ represent the CFA measurements with and without the presence of layering artifacts. However, even during periods of strong methane variations, the bias specifically due to layering artifacts does not exceed 10ppbv. This is of the order of the analytical noise

10 usually observed with the CFA system (Fourteau et al., 2017). It therefore appears that layering artifacts only negligibly affect the measured signal in the case of thinned ice cores. However, the potential impact of layering artifacts is sensitive to the parameters of the layering model. Notably, increasing the number of anomalous layers increases the amplitude of the biases.




Figure 12. Impacts of layered trapping and smoothing on a synthetic million-and-a-half years old methane record. Upper panel: The signal recorded in the ice taking into account without smoothing (in black) and layered trapping or artifacts (is displayed in blue) dashed orange. The atmospheric reference recorded signal with layering is displayed shown in orange light blue, and the recorded signal with smoothing only in black. The depth scale takes synthetic CFA measurements taking into account the thinning of artifacts are displayed in red, while the ice. Lower panel: CFA synthetic measurements with (in blue) and without layering artifacts (are displayed in green). The atmospheric reference synthetic depth scale is displayed orange and expressed relative to an arbitrary point of the initial recorded signal without layering artifacts record. For clarity the depth scale is also displayed expressed in black centimeters.

The internal smoothing of the CFA system also tends to deteriorate information recorded in the ice core by attenuating fast variability. It is therefore important to determine to what extent CFA smoothing adds to the already present firn smoothing.

15 To study this point, we compare the frequency responses of the CFA system and firn smoothing. These frequency responses represent the attenuation experienced by a sine signal, as a function of the sine period. To achieve this, we first expressed

the CFA impulse response (its smoothing function) on a gas age scale, taking into account the thinning of the ice. Then, the frequency responses were deduced using fast Fourier transforms. The gas age distribution and the CFA impulse response are displayed in the left panel of Figure 13 with the frequency responses in the right panel. For sine periods longer than 100yr the firm smoothing is larger than the CFA smoothing. Yet, for a sine period of 500yr the firm induces a smoothing of 50% while the CFA system smooths by about 20%. The CFA smoothing is therefore noticeable and adds to the already present firm smoothing. This is illustrated in the lower-panel of Figure 12, where the difference between the black and green curves is due to the presence of CFA smoothing. The influence of this analytical smoothing would be even stronger in the case of a narrower gas age distribution, for instance during interglacial periods. This is problematic as it deteriorates climatic information still present in the gas record. It thus appears important to improve current CFA gas systems in order to minimize their internal smoothing, in order to take full advantage of the gas records entrapped in deep ice cores.




Figure 13. Left panel: gas age distribution in the synthetic million-and-a-half years old ice core in blue, and CFA impulse response on a gas age scale in orange. Right panel: attenuation of a sine signal by the firm and CFA smoothing in blue and orange respectively. The horizontal dashed line indicates the 50% attenuation.

4.2 Carbon dioxide alterations

So far in this article only methane records have been considered. However, smoothing and layering artifacts also potentially affect other gaseous records, such as carbon dioxide records. Since CO_2 is one of the principal and most studied greenhouse gases, we propose to quantify the potential impacts of gas trapping on the record enclosed in a million-and-a-half years old ice core. Similar to the methane case of the previous section, we used the CO_2 measurements of the high accumulation WD ice core as the atmospheric reference (Marcott et al., 2014). The chosen data points correspond to the last deglaciation, as they constitute a fast atmospheric increase of CO_2 . As for CH_4 , the synthetic ice core signal is determined with the smoothing and layering model and ice layers are thinned 200 times. Finally, we emulate the process of discrete measurements. For that, we assume that the final ice core is continuously discretized into 4cm thick pieces, and that the measured value is the average

value of the concentrations enclosed in the ice piece.

The results are displayed in Figure 14. The ice core signal is displayed in blue in the left panel of Figure 14, with the artifact free record superimposed in black. It can be seen that during enclosure, the layering heterogeneities can produce artifacts reaching up to 10ppmv. The measurements obtained by a discrete method are displayed on the right side of Figure 14 with and without the presence of layering artifacts in blue and black respectively. These results show that the impact of layering artifacts on the final measurements does not exceed 0.5ppmv, even during abrupt CO₂ rises. However, as for methane the impact of the layering artifacts is sensitive to the amount of abnormal layers per meter.

We also compared the rate of change of carbon dioxide in the atmospheric reference and in the measured record. For some of the fast variations of atmospheric CO₂, a reduced rate of change is deduced from the ice core record. A specific example (the deeper part of the records in Figure 14) is shown in Figure 15, where the rates of change of CO₂ in the atmosphere, recorded in the ice, and deduced from discrete measurements are compared. For this event, the atmospheric rate of change peaks at about 0.055ppmv.yr^{-1} , while the corresponding imprint in the ice has only a rate of change of 0.035ppmv.yr^{-1} due to firn smoothing. The process of discrete measurement further diminishes the measured rate of change to about 0.02ppmv.yr^{-1} , a bit more than a third of the atmospheric value.




Figure 14. Impacts of layered trapping and smoothing on a synthetic million-and-a-half years old carbon dioxide record. Left panel: recorded signal in the ice taking into account smoothing (in black) and layered trapping artifacts (in blue). The atmospheric reference signal without smoothing or artifacts is shown in orange. Right panel: measured signal by a discrete system shown as blue points. The atmospheric reference signal without smoothing or artifacts (in orange) and the measured signal without layering artifacts (black points) are also displayed. The depth scale is expressed relative to an arbitrary point in the record, and in centimeters.




Figure 15. Impact of gas trapping processes and measurements on the CO₂ rate of change. The rate of change of the atmosphere is displayed in orange, while the rate change recorded in the ice is in black, and the measured one is in blue. The depth scale is the same relative depth scale as in Figure 14.

5 Conclusions

This work evaluated two gas trapping effects that affect the recorded atmospheric trace gas history in polar ice cores. The first one is the layered gas trapping, that produces stratigraphic heterogeneities appearing as spurious values in the measured record (Rhodes et al., 2016). The second one is the smoothing effect that removes a part of the variability in the gas records (Spahni et al., 2003; Joos and Spahni, 2008; Köhler et al., 2011). Our work focuses on the arid region of East Antarctica, as this is where the oldest ice cores are retrieved. Five new sections from East Antarctic ice cores have been analyzed at high resolution for methane concentration.

The characterization of layering artifacts in these new measurements is consistent with the physical mechanisms already proposed in the literature (Etheridge et al., 1992; Rhodes et al., 2016). In accordance with previous studies (Hörhold et al., 2012; Fujita et al., 2016), the data suggest that chemistry could promote the presence of especially dense layers closing in advance and creating the layering artifacts. We ~~parameterized~~ parametrized a simple model to reproduce the observed distribution and characteristics of layering artifacts, and were able to find a common set of parameters to reproduce the ~~measured-observed~~ distribution of artifacts in all studied East Antarctic ice cores.

In order to constrain the smoothing effect, we estimated the gas age distributions in each of the five ice core sections. For this

purpose, we use the method proposed by Fourteau et al. (2017) based on the comparison with higher accumulation methane records. It appears that during the last glacial period, the different methane records in ice cores formed under accumulation rates lower than 2 cm.ice.yr^{-1} all display a similar degree of smoothing. We therefore propose to use a common gas age distribution for glacial East Antarctic sites with accumulations below 2 cm.ice.yr^{-1} . The comparison of the glacial and interglacial GADs suggests that the smoothing is higher at Dome C during the glacial period than during the interglacial period. Yet, despite a doubling of accumulation during the interglacial period, the resulting smoothing is only slightly lower. This means that more climatic information than previously thought is preserved in glacial East Antarctica ice core gas records, which are thus suited to study multi-centennial atmospheric variability. Similarly to Fourteau et al. (2017), we observed that the smoothing during the glacial in the Vostok ice core is weaker than the one predicted for modern Vostok by the age distribution estimated by gas trapping models (Witrant et al., 2012).

Finally, we applied our methodology to the theoretical case of a million-and-a-half years old ice core drilled on the East Antarctic plateau. Our results suggest that due to thinning, the layering artifacts will no longer be resolved during measurements. However, it appears that their potential influence on the end result measurements is rather low, with impacts below 10ppbv and 0.5ppmv for methane and carbon dioxide respectively. In the case of methane variations during DO events, most of the record alterations originate from the firn smoothing. Yet, the limited spatial resolution of the CFA system induces a smoothing that is not entirely negligible compared to the one of the firn. For carbon dioxide, firn smoothing appears to significantly diminish the recorded rates of change of abrupt CO_2 increases, compared to their atmospheric values. The estimations of CO_2 rates of change are further altered by the process of discrete measurement, and measured values can be three time-times lower than the actual atmospheric rate of change.

Code availability. The programs used for data processing and modeling were developed using python3 and available packages. They will be provided upon request to the corresponding authors.

Data availability. The high resolution methane datasets will be made available on the World Data Center for Paleoclimatology.

Author contributions. This scientific project was designed by JC, XF, KF, and PM. JC and PM participated in the Lock-In drilling and XF participated in the shallow Dome C drilling. AAE and VL made available and pre-processed the Vostok ice core section. The high resolution

methane measurements were carried out by XF and KF. The codes for data processing and modeling were developed by KF and PM. All authors contributed to the interpretation of the data. The manuscript was written by KF with the help of all co-authors.

Competing interests. The authors declare having no competing interests

Disclaimer. TEXT

- 5 *Acknowledgements.* This work is a contribution to EPICA, a joint European Science Foundation/European Commission scientific program funded by the European Union and national contributions from Belgium, Denmark, France, Germany, Italy, the Netherlands, Norway, Sweden, Switzerland, and the United Kingdom. The Vostok ice core was made accessible in the framework of the Laboratoire International Associé (LIA) Vostok. The Lock-In drilling was supported by the IPEV project No 1153 and the European Community's Seventh Framework Programme under grant agreement No291062 (ERC ICE&LASERS). We are grateful to the Lock-In field personnel: David Colin,
- 10 Phillipe Dordhain, and Phillipe Possenti who performed the drilling, as well as Patrice Godon for setting up the logistics. The shallow Dome C drilling was performed by Phillipe Possenti and supported by the IPEV project No 902 and the French ANR program RPD COCLICO (ANR-10-RPDOC-002-01). We thank Robert Mulvaney and the British Antarctic Survey for the Fletcher ~~promontory~~ Promontory ice core drilling, and the permission to use the Fletcher methane data. We thank Grégoire Aufresne for his help processing the ice cores. We thank Grégory Teste for his help during the ice core processing and the CFA methane measurements. We thank Jochen Schmitt and Hubertus
- 15 Fischer for their constructive comments. This work was supported by the French INSU/CNRS LEFE projects NEVE-CLIMAT and HEP-IGANE. This work was also supported by the French ANR programs RPD COCLICO (ANR-10-RPDOC-002-01). We are thankful to the two anonymous referees for reviewing this work, as well as to Hubertus Fischer for editing it. This is EPICA publication no. XX and Beyond EPICA - Oldest Ice publication No. YY.

References

- Ahn, J., Brook, E. J., and Buizert, C.: Response of atmospheric CO₂ to the abrupt cooling event 8200 years ago, *Geophys. Res. Lett.*, 41, 604–609, <https://doi.org/10.1002/2013gl058177>, 2014.
- Arnaud, L., Barnola, J.-M., and Duval, P.: Physical modeling of the densification of snow/firn and ice in the upper part of polar ice sheets, *Phys. Ice Core Rec.*, pp. 285–305, <http://hdl.handle.net/2115/32472>, 2000.
- Bazin, L., Landais, A., Lemieux-Dudon, B., Toyé Mahamadou Kele, H., Veres, D., Parrenin, F., Martinerie, P., Ritz, C., Capron, E., Lipenkov, V. Y., Loutre, M.-F., Raynaud, D., Vinther, B., Svensson, A., Rasmussen, S. O., Severi, M., Blunier, T., Leuenberger, M., Fischer, H., Masson-Delmotte, V., Chappellaz, J., and Wolff, E. W.: An optimized multi-proxy, multi-site Antarctic ice and gas orbital chronology (AICC2012): 120-800 ka, *Clim. Past*, 9, 1715–1731, <https://doi.org/10.5194/cp-9-1715-2013>, 2013.
- 10 Bereiter, B., Fischer, H., Schwander, J., and Stocker, T. F.: Diffusive equilibration of N₂, O₂ and CO₂ mixing ratios in a 1.5-million-years-old ice core, *Cryosphere*, 8, 245–256, <https://doi.org/10.5194/tc-8-245-2014>, 2014.
- Bock, J., Martinerie, P., Witrant, E., and Chappellaz, J.: Atmospheric impacts and ice core imprints of a methane pulse from clathrates, *Earth Planet. Sci. Lett.*, 349-350, 98 – 108, <https://doi.org/https://doi.org/10.1016/j.epsl.2012.06.052>, 2012.
- Bréant, C., Martinerie, P., Orsi, A., Arnaud, L., and Landais, A.: Modelling firn thickness evolution during the last deglaciation: constraints on sensitivity to temperature and impurities, *Clim. Past*, 13, 833–853, <https://doi.org/10.5194/cp-13-833-2017>, 2017.
- 15 Buizert, C., Martinerie, P., Petrenko, V. V., Severinghaus, J. P., Trudinger, C. M., Witrant, E., Rosen, J. L., Orsi, A. J., Rubino, M., Etheridge, D. M., Steele, L. P., Hogan, C., Laube, J. C., Sturges, W. T., Levchenko, V. A., Smith, A. M., Levin, I., Conway, T. J., Dlugokencky, E. J., Lang, P. M., Kawamura, K., Jenk, T. M., White, J. W. C., Sowers, T., Schwander, J., and Blunier, T.: Gas transport in firn: multiple-tracer characterisation and model intercomparison for NEEM, Northern Greenland, *Atmos. Chem. Phys.*, 12, 4259–4277, <https://doi.org/10.5194/acp-12-4259-2012>, 2012.
- 20 Buizert, C., Cuffey, K. M., Severinghaus, J. P., Baggenstos, D., Fudge, T. J., Steig, E. J., Markle, B. R., Winstrup, M., Rhodes, R. H., Brook, E. J., Sowers, T. A., Clow, G. D., Cheng, H., Edwards, R. L., Sigl, M., McConnell, J. R., and Taylor, K. C.: The WAIS Divide deep ice core WD2014 chronology-Part 1: methane synchronization (68-31 ka BP) and the gas age-ice age difference, *Clim. Past*, 11, 153–173, <https://doi.org/10.5194/cp-11-153-2015>, 2015.
- 25 Chappellaz, J., Stowasser, C., Blunier, T., Baslev-Clausen, D., Brook, E. J., Dallmayr, R., Faïn, X., Lee, J. E., Mitchell, L. E., Pascual, O., Romanini, D., Rosen, J., and Schüpbach, S.: High-resolution glacial and deglacial record of atmospheric methane by continuous-flow and laser spectrometer analysis along the NEEM ice core, *Clim. Past*, 9, 2579–2593, <https://doi.org/10.5194/cp-9-2579-2013>, 2013.
- Dlugokencky, E. J., Myers, R. C., Lang, P. M., Masarie, K. A., Crotwell, A. M., Thoning, K. W., Hall, B. D., Elkins, J. W., and Steele, L. P.: Conversion of NOAA atmospheric dry air CH₄ mole fractions to a gravimetrically prepared standard scale, *J. Geophys. Res. Atmos.*, 110, <https://doi.org/doi.org/10.1029/2005jd006035>, 2005.
- 30

- Dällenbach, A., Blunier, T., Flückiger, J., Stauffer, B., Chappellaz, J., and Raynaud, D.: Changes in the atmospheric CH₄ gradient between Greenland and Antarctica during the Last Glacial and the transition to the Holocene, *Geophys. Res. Lett.*, 27, 1005–1008, <https://doi.org/10.1029/1999GL010873>, 2000.
- EPICA Community Members, Barbante, C., Barnola, J.-M., Becagli, S., Beer, J., Bigler, M., Boutron, C., Blunier, T., Castellano, E., Cattani, O., et al.: One-to-one coupling of glacial climate variability in Greenland and Antarctica, *Nature*, 444, 195, <https://doi.org/10.1038/nature05301>, 2006.
- Etheridge, D. M., Pearman, G. I., and Fraser, P. J.: Changes in tropospheric methane between 1841 and 1978 from a high accumulation-rate Antarctic ice core, *Tellus B Chem. Phys. Meteorol.*, 44, 282–294, <https://doi.org/10.3402/tellusb.v44i4.15456>, 1992.
- Fischer, H., Severinghaus, J., Brook, E., Wolff, E. W., Albert, M., Alemany, O., Arthern, R., Bentley, C., Blankenship, D., Chappellaz, J., Creyts, T., Dahl-Jensen, D., Dinn, M., Frezzotti, M., Fujita, S., Gallee, H., Hindmarsh, R., Hudspeth, D., Jugie, G., Kawamura, K., Lipenkov, V. Y., Miller, H., Mulvaney, R., Parrenin, F., Pattyn, F., Ritz, C., Schwander, J., Steinhage, D., van Ommen, T., and Wilhelms, F.: Where to find 1.5 million yr old ice for the IPICS "Oldest-Ice" ice core, *Clim. Past*, 9, 2489–2505, <https://doi.org/10.5194/cp-9-2489-2013>, 2013.
- Fourteau, K., Faïn, X., Martinerie, P., Landais, A., Ekaykin, A. A., Lipenkov, V. Y., and Chappellaz, J.: Analytical constraints on layered gas trapping and smoothing of atmospheric variability in ice under low-accumulation conditions, *Clim. Past*, 13, 1815–1830, <https://doi.org/10.5194/cp-13-1815-2017>, 2017.
- Fourteau, K., Martinerie, P., Faïn, X., Schaller, C. F., Tuckwell, R. J., Löwe, H., Arnaud, L., Magand, O., Thomas, E. R., Freitag, J., Mulvaney, R., Schneebeli, M., and Lipenkov, V. Y.: Multi-tracer study of gas trapping in an East Antarctic ice core, *Cryosphere Discuss.*, 2019, 1–34, <https://doi.org/10.5194/tc-2019-89>, 2019.
- Freitag, J., Wilhelms, F., and Kipfstuhl, S.: Microstructure-dependent densification of polar firn derived from X-ray microtomography, *J. Glaciol.*, 50, 243–250, <https://doi.org/10.3189/172756504781830123>, 2004.
- Freitag, J., Kipfstuhl, S., Laepple, T., and Wilhelms, F.: Impurity-controlled densification: a new model for stratified polar firn, *J. Glaciol.*, 59, 1163–1169, <https://doi.org/10.3189/2013JoG13J042>, 2013.
- Fujita, S., Okuyama, J., Hori, A., and Hondoh, T.: Metamorphism of stratified firn at Dome Fuji, Antarctica: A mechanism for local insolation modulation of gas transport conditions during bubble close off, *J. Geophys. Res. Earth Surf.*, 114, F03023, <https://doi.org/10.1029/2008JF001143>, 2009.
- Fujita, S., Goto-Azuma, K., Hirabayashi, M., Hori, A., Iizuka, Y., Motizuki, Y., Motoyama, H., and Takahashi, K.: Densification of layered firn in the ice sheet at Dome Fuji, Antarctica, *J. Glaciol.*, 62, 103–123, <https://doi.org/10.1017/jog.2016.16>, 2016.
- Gautier, E., Savarino, J., Erbland, J., Lanciki, A., and Possenti, P.: Variability of sulfate signal in ice core records based on five replicate cores, *Clim. Past*, 12, 103–113, <https://doi.org/10.5194/cp-12-103-2016>, 2016.
- Gregory, S. A., Albert, M. R., and Baker, I.: Impact of physical properties and accumulation rate on pore close-off in layered firn, *Cryosphere*, 8, 91–105, <https://doi.org/10.5194/tc-8-91-2014>, 2014.

- Hörhold, M. W., Kipfstuhl, S., Wilhelms, F., Freitag, J., and Frenzel, A.: The densification of layered polar firn, *J. Geophys. Res. Earth Surf.*, 116, <https://doi.org/10.1029/2009jf001630>, 2011.
- Hörhold, M. W., Laepple, T., Freitag, J., Bigler, M., Fischer, H., and Kipfstuhl, S.: On the impact of impurities on the densification of polar firn, *Earth Planet. Sci. Lett.*, 325, 93–99, <https://doi.org/10.1016/j.epsl.2011.12.022>, 2012.
- 5 Howat, I. M., Negrete, A., and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *Cryosphere*, 8, 1509–1518, <https://doi.org/10.5194/tc-8-1509-2014>, 2014.
- Huber, C., Leuenberger, M., Spahni, R., Flückiger, J., Schwander, J., Stocker, T. F., Johnsen, S., Landais, A., and Jouzel, J.: Isotope calibrated Greenland temperature record over Marine Isotope Stage 3 and its relation to CH₄, *Earth Planet. Sci. Lett.*, 243, 504–519, 2006.
- Joos, F. and Spahni, R.: Rates of change in natural and anthropogenic radiative forcing over the past 20,000 years, *Proc. Natl. Acad. Sci. USA*, 105, 1425–1430, <https://doi.org/10.1073/pnas.0707386105>, 2008.
- 10 Köhler, P., Knorr, G., Buiron, D., Lourantou, A., and Chappellaz, J.: Abrupt rise in atmospheric CO₂ at the onset of the Bølling/Allerød: in-situ ice core data versus true atmospheric signal, *Clim. Past*, 7, 473–486, <https://doi.org/10.5194/cp-7-473-2011>, 2011.
- Köhler, P., Völker, C., Knorr, G., and Bard, E.: High latitude impacts on deglacial CO₂: southern ocean westerly winds and northern hemisphere permafrost thawing, *Nova. Act. Lc.*, 121, 135–140, <https://doi.org/10013/epic.45322.d002>, 2015.
- 15 Lambert, F., Bigler, M., Steffensen, J. P., Hutterli, M., and Fischer, H.: Centennial mineral dust variability in high-resolution ice core data from Dome C, Antarctica, *Clim. Past*, 8, 609–623, <https://doi.org/10.5194/cp-8-609-2012>, 2012.
- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J.-M., Raynaud, D., Stocker, T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years, *Nature*, 453, 383–386, <https://doi.org/10.1038/nature06950>, 2008.
- 20 Lüthi, D., Floch, M. L., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000– 800,000 years before present, *Nature*, 453, 379–382, <https://doi.org/10.1038/nature06949>, 2008.
- Marcott, S. A., Bauska, T. K., Buizert, C., Steig, E. J., Rosen, J. L., Cuffey, K. M., Fudge, T. J., Severinghaus, J. P., Ahn, J., Kalk, M. L., McConnell, J. R., Sowers, T., Taylor, K. C., White, J. W. C., and Brook, E. J.: Centennial-scale changes in the global carbon cycle during the last deglaciation, *Nature*, 514, 616–619, <https://doi.org/10.1038/nature13799>, 2014.
- 25 Martín, C. and Gudmundsson, G. H.: Effects of nonlinear rheology, temperature and anisotropy on the relationship between age and depth at ice divides, *Cryosphere*, 6, 1221–1229, <https://doi.org/10.5194/tc-6-1221-2012>, 2012.
- Martinerie, P., Raynaud, D., Etheridge, D. M., Barnola, J.-M., and Mazaudier, D.: Physical and climatic parameters which influence the air content in polar ice, *Earth Planet. Sc. Lett.*, 112, 1–13, [https://doi.org/10.1016/0012-821x\(92\)90002-d](https://doi.org/10.1016/0012-821x(92)90002-d), 1992.
- 30 Mitchell, L., Brook, E., Lee, J. E., Buizert, C., and Sowers, T.: Constraints on the Late Holocene Anthropogenic Contribution to the Atmospheric Methane Budget, *Science*, 342, 964–966, <https://doi.org/10.1126/science.1238920>, 2013.

- Mitchell, L. E., Buizert, C., Brook, E. J., Breton, D. J., Fegyveresi, J., Baggenstos, D., Orsi, A., Severinghaus, J., Alley, R. B., Albert, M., Rhodes, R. H., McConnell, J. R., Sigl, M., Maselli, O., Gregory, S., and Ahn, J.: Observing and modeling the influence of layering on bubble trapping in polar firn, *J. Geophys. Res. Atmos.*, 120, 2558–2574, <https://doi.org/10.1002/2014jd022766>, 2015.
- Morville, J., Kassi, S., Chenevier, M., and Romanini, D.: Fast, low-noise, mode-by-mode, cavity-enhanced absorption spectroscopy by diode-laser self-locking, *Appl. Phys. B-Lasers O.*, 80, 1027–1038, <https://doi.org/10.1007/s00340-005-1828-z>, 2005.
- Parrenin, F., Petit, J.-R., Masson-Delmotte, V., Wolff, E., Basile-Doelsch, I., Jouzel, J., Lipenkov, V., Rasmussen, S. O., Schwander, J., Severi, M., Udisti, R., Veres, D., and Vinther, B. M.: Volcanic synchronisation between the EPICA Dome C and Vostok ice cores (Antarctica) 0–145 kyr BP, *Clim. Past*, 8, 1031–1045, <https://doi.org/10.5194/cp-8-1031-2012>, 2012.
- Passalacqua, O., Cavitte, M., Gagliardini, O., Gillet-Chaulet, F., Parrenin, F., Ritz, C., and Young, D.: Brief communication: Candidate sites of 1.5 Myr old ice 37 km southwest of the Dome C summit, East Antarctica, *Cryosphere*, 12, 2167–2174, <https://doi.org/10.5194/tc-12-2167-2018>, 2018.
- Rasmussen, S. O., Abbott, P. M., Blunier, T., Bourne, A. J., Brook, E., Buchardt, S. L., Buizert, C., Chappellaz, J., Clausen, H. B., Cook, E., Dahl-Jensen, D., Davies, S. M., Guillevic, M., Kipfstuhl, S., Laepple, T., Seierstad, I. K., Severinghaus, J. P., Steffensen, J. P., Stowasser, C., Svensson, A., Vallenga, P., Vinther, B. M., Wilhelms, F., and Winstrup, M.: A first chronology for the North Greenland Eemian Ice Drilling (NEEM) ice core, *Clim. Past*, 9, 2713–2730, <https://doi.org/10.5194/cp-9-2713-2013>, 2013.
- Raymond, C. F.: Deformation in the Vicinity of Ice Divides, *J. Glaciol.*, 29, 357–373, <https://doi.org/10.3189/S0022143000030288>, 1983.
- Rhodes, R. H., Faïn, X., Stowasser, C., Blunier, T., Chappellaz, J., McConnell, J. R., Romanini, D., Mitchell, L. E., and Brook, E. J.: Continuous methane measurements from a late Holocene Greenland ice core: atmospheric and in-situ signals, *Earth Planet. Sc. Lett.*, 368, 9–19, <https://doi.org/10.1016/j.epsl.2013.02.034>, 2013.
- Rhodes, R. H., Brook, E. J., Chiang, J. C. H., Blunier, T., Maselli, O. J., McConnell, J. R., Romanini, D., and Severinghaus, J. P.: Enhanced tropical methane production in response to iceberg discharge in the North Atlantic, *Science*, 348, 1016–1019, <https://doi.org/10.1126/science.1262005>, 2015.
- Rhodes, R. H., Faïn, X., Brook, E. J., McConnell, J. R., Maselli, O. J., Sigl, M., Edwards, J., Buizert, C., Blunier, T., Chappellaz, J., and Freitag, J.: Local artifacts in ice core methane records caused by layered bubble trapping and in situ production: a multi-site investigation, *Clim. Past*, 12, 1061–1077, <https://doi.org/10.5194/cp-12-1061-2016>, 2016.
- Rommelaere, V., Arnaud, L., and Barnola, J.-M.: Reconstructing recent atmospheric trace gas concentrations from polar firn and bubbly ice data by inverse methods, *J. Geophys. Res. Atmos.*, 102, 30 069–30 083, <https://doi.org/10.1029/97jd02653>, 1997.
- Rousseeuw, P. J. and Hubert, M.: Robust statistics for outlier detection, *Wires Data Min. Knowl. Discov.*, 1, 73–79, <https://doi.org/10.1002/widm.2>, 2011.
- Salamatin, A. N., Lipenkov, V. Y., Barnola, J. M., Hori, A., Duval, P., and Hondoh, T.: Snow/firn densification in polar ice sheets, *Low Temperature Science*, 68, 195–222, 2009.

- Schaller, C. F., Freitag, J., and Eisen, O.: Critical porosity of gas enclosure in polar firn independent of climate, *Clim. Past*, 13, 1685–1693, <https://doi.org/10.5194/cp-13-1685-2017>, 2017.
- Schwander, J.: The transformation of snow to ice and the occlusion of gases, in: *The environmental record in glaciers and ice sheets*, edited by Oeschger, H. and Langway, C. C. J., pp. 53–67, John Wiley, New York, 1989.
- 5 Schwander, J., Stauffer, B., and Sigg, A.: Air mixing in firn and the age of the air at pore close-off, *Ann. Glaciol.*, 10, 141–145, <https://doi.org/10.1017/S0260305500004328>, 1988.
- Schwander, J., Barnola, J.-M., Andri , C., Leuenberger, M., Ludin, A., Raynaud, D., and Stauffer, B.: The age of the air in the firn and the ice at Summit, Greenland, *J. Geophys. Res. Atmos.*, 98, 2831–2838, <https://doi.org/10.1029/92jd02383>, 1993.
- Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto-Bliesner, B., Schmittner, A., and Bard, E.: Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation, *Nature*, 484, 49–54, <https://doi.org/10.1038/nature10915>, 2012.
- 10 Spahni, R., Schwander, J., Fl ckiger, J., Stauffer, B., Chappellaz, J., and Raynaud, D.: The attenuation of fast atmospheric CH₄ variations recorded in polar ice cores, *Geophys. Res. Lett.*, 30, <https://doi.org/10.1029/2003gl017093>, 2003.
- Stauffer, B., Schwander, J., and Oeschger, H.: Enclosure of air during metamorphosis of dry firn to ice, *Ann. Glaciol.*, 6, 108–112, <https://doi.org/10.3189/1985AoG6-1-108-112>, 1985.
- 15 Thomas, E. R., Wolff, E. W., Mulvaney, R., Steffensen, J. P., Johnsen, S. J., Arrowsmith, C., White, J. W., Vaughn, B., and Popp, T.: The 8.2 ka event from Greenland ice cores, *Quat. Sci. Rev.*, 26, 70–81, <https://doi.org/10.1016/j.quascirev.2006.07.017>, 2007.
- Van Liefferinge, B. and Pattyn, F.: Using ice-flow models to evaluate potential sites of million year-old ice in Antarctica, *Clim. Past*, 9, 2335–2345, <https://doi.org/10.5194/cp-9-2335-2013>, 2013.
- 20 Veres, D., Bazin, L., Landais, A., Toy  Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi, M., Svensson, A., Vinther, B., and Wolff, E. W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Clim. Past*, 9, 1733–1748, <https://doi.org/10.5194/cp-9-1733-2013>, 2013.
- Witrant, E. and Martinerie, P.: Input Estimation from Sparse Measurements in LPV Systems and Isotopic Ratios in Polar Firns, *IFAC Proc. Vol.*, 46, 659 – 664, <https://doi.org/10.3182/20130204-3-FR-2033.00201>, 5th IFAC Symposium on System Structure and Control, 2013.
- 25 Witrant, E., Martinerie, P., Hogan, C., Laube, J. C., Kawamura, K., Capron, E., Montzka, S. A., Dlugokencky, E. J., Etheridge, D., Blunier, T., and Sturges, W. T.: A new multi-gas constrained model of trace gas non-homogeneous transport in firn: evaluation and behaviour at eleven polar sites, *Atmos. Chem. Phys.*, 12, 11465–11483, <https://doi.org/10.5194/acp-12-11465-2012>, 2012.
- Yeung, L. Y., Murray, L. T., Martinerie, P., Witrant, E., Hu, H., Banerjee, A., Orsi, A., and Chappellaz, J.: Isotopic constraint on the twentieth-century increase in tropospheric ozone, *Nature*, 570, 224–227, <https://doi.org/10.1038/s41586-019-1277-1>, 2019.
- 30

Supplementary material for:

Predicting gas record alteration in very low accumulation ice cores

Kévin Fourteau¹, Patricia Martinerie¹, Xavier Faïn¹, Alexey A. Ekaykin², Jérôme Chappellaz¹, and Vladimir Lipenkov²

¹Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, F-38000 Grenoble, France

²Climate and Environmental Research Laboratory, Arctic and Antarctic Research Institute, St. Petersburg, 199397, Russia

Correspondence to: kevin.fourteau@univ-grenoble-alpes.fr or patricia.martinerie@univ-grenoble-alpes.fr

S1 ~~NEEM atmospheric reference without deconvolution~~ Correction for the dissolution of methane

In the main article, we used as atmospheric reference for the DO21 period the methane CFA data of Chappellaz et al. (2013) after a deconvolution. The CFA records measured at IGE need to be corrected for the preferential dissolution of methane compared to oxygen and nitrogen in water (Chappellaz et al., 2013; Rhodes et al., 2013). This was done as using the data without deconvolution leads to an inconsistency between the atmospheric reference and the Vostok record. Indeed, as illustrated by applying correction factors chosen to match the CFA concentrations with already calibrated data. Such a match is displayed for the DO6 to 9 section in the EDC96 ice core in Figure S1. The discrete measurements used for the correction were obtained by a discrete melt-refreeze method (Louergue et al., 2008).




Figure S1. Illustration of a corrected CFA record (in gray) by matching an already-calibrated record (orange dots, Louergue et al., 2008).

S2 Deconvolving the NEEM CFA record

10 In the main article we use high-resolution methane records as input atmospheric scenarios to study the smoothing of low-accumulation records. However, high-accumulation records can only be used in such a way if the fast variability is well preserved and not affected by smoothing. In the case of NEEM, this appears not to be the case. It can be seen in Figure S2 that it is not possible

to find ~~a single gas age distribution that leads to adequate smoothing in the Vostok~~ an age distribution to transform the NEEM record into the Vostok DO21 record. In particular, it is not possible to reproduce the Vostok record during the first feature of the DO21 event (1260m depth in the record. ~~As it is unlikely that the gas age distribution of the enclosed gases changed in the Vostok record between the two methane increases of the DO21, we attribute this discrepancy to the fact the raw NEEM data cannot be used as atmospheric reference without deconvolution~~), as the methane peak is not high enough in the NEEM record. It means that in this case, NEEM cannot be directly used as the input atmospheric scenario. This suggests that the NEEM scenario needs to be deconvoluted first, to retrieve the full atmospheric variability.

To do this, one needs to assume an age distribution for the NEEM methane record. During the DO21 period, the NEEM accumulation has been estimated to be between 10.4 and 13 cm ie yr⁻¹, with an average value of 11.3 cm ie yr⁻¹ (Rasmussen et al., 2013).

We therefore chose to use the age distribution estimated at the modern site of Siple Dome (10.8 cm ie yr⁻¹, Witrant et al., 2012). In order to test the sensitivity of the results to the choice of GAD, we also performed the deconvolution with the age distribution of South Pole (8.0 cm ie yr⁻¹) and Berkner Island (14.1 cm ie yr⁻¹, Witrant et al., 2012). The three deconvolutions produce the atmospheric scenarios displayed as solid lines in Figure S3. When these three scenarios are smoothed by the glacial age distribution chosen in Section 3.4.2 of the main article, they result in similar ~~low accumulation~~ records, displayed as dashed lines in Figure S3. We also performed the gas age distribution extraction procedure of Section 3.4.1 of the main article, and found similar gas age distributions (within the uncertainty range) using the three deconvolutions as atmospheric references. It indicates that our results weakly depend on the particular choice of GAD for the deconvolution of the NEEM record.




Figure S2. Illustration of the ~~impossibility to find~~ failure in finding a gas age distribution that smooths the atmospheric scenario (in dashed black/orange, splined NEEM data without deconvolution) in order to match the Vostok CFA measurements (in light blue). The smoothed version in green underestimates the fast event measured in the ice core around the 1260m depth. The black line represents the original Chappellaz et al. (2013) data, that partially overlaps the spline.




Figure S3. [The NEEM deconvoluted scenarios and their convolution into Vostok records.](#) For each color, the solid line represents the deconvoluted atmospheric reference and the dashed line represents the atmospheric reference smoothed by an East Antarctica glacial age distribution. Orange, blue and green curves respectively represent the cases with a 8.0, 10.8, and 14.1cm ie yr⁻¹ accumulation for the NEEM gas age distribution during the DO21 period. The original NEEM data are also shown in black.

S2-S3 [Zoom over layering artifacts](#)

A closer look at the layering artifacts of the EDC96 DO6-9 section is shown in Figure S4. This highlights the structure of the layering artifacts, and their widths of a couple of centimeters. They are similar to the artifacts reported in Figure 1 [if of](#) Fourteau et al. (2017).




Figure S4. [Zoom over the EDC96 DO6-9 methane record.](#) The blue data are the raw CFA measurements. In black are the data cleaned for layering artifacts [with the recursive cleaning procedure presented in Section 3.3.3 of the main article.](#)

5 **S3-S4** [Comparison of discrete and continuous methane data over the DO8](#)

In this section we compare the discrete (Louergue et al., 2008) and continuous (this study) methane measurements over the DO8 event in the EPICA Dome C record. Fourteau et al. (2017) pointed out that one the of the data [point-points](#) of Louergue

et al. (2008) might ~~corresponds~~ correspond to an early closure artifact and ~~is therefore not climatically relevant variability~~ may not be climatically relevant. This data point is highlighted with a blue circle in Figure S5. The high-resolution measurements confirm that this data point corresponds to an early closure artifact, the sample selected by Louergue et al. (2008) being in a zone with a lot layering artifacts with similar methane ~~concentration~~ concentrations.




Figure S5. In orange: discrete methane measurements of the DO8 event in the EPICA Dome C record (Louergue et al., 2008). In black: continuous methane measurements of the same event in the EPICA Dome C record (this study). In blue we highlight a data point of the discrete record likely corresponding to an early closure artifact.

5 **S4 Layering artifacts model, without impurity effect**

In the main text, a parametrization was included to account for the enhancement of layering due to the presence of high calcium concentrations and variabilities. Here we present the result of the layering model for the DO6 to 9 period in the EDC96 ice core, without taking into account the enhancement due to chemistry. The output of the model is displayed in Figure ?? . As seen on the figure, the model does not predict enough layering artifacts on the onset of the DO7 and 8. This highlights the necessity

10 to take into account the effect of impurities on layering.

Results of the layering model. This figure is similar to Figure 5 of the main article, but does not include the effect of impurity driven enhancement of layering.

S5 Uncertainty on Gas Age Distributions

The extraction method yields an optimal GAD that minimizes the ~~RMSD~~Root-Mean-Square Deviation (RMSD) between the CFA measurements and the smoothed atmospheric reference. To quantify the uncertainties associated with the calculated GADs, we performed a dedicated analysis. For each of the five ice core sections, we studied the impact of choosing slightly modified location and scale parameters.

Practically, we modify the scale and location parameters until the resulting RMSD is 15% larger than its optimal value. The ~~location~~locations and scales were modified following straight lines in the {location, scale} space, starting from the optimal values (Figure S6). We then obtained a set of 10 GADs meant to represent the limit of acceptable distributions. These GADs and their corresponding smoothed methane records in the ice are displayed in Figures S7 to S11. In the right panel of these figures, we also represented the set of location and scale values that increase the RMSD by 15%. The regions enclosed by the those values can be viewed as the uncertainty ranges of the location and scale parameters.

To better appreciate visualize the uncertainty associates with the EDC96 and Vostok glacial GADs, we plotted their uncertainty ranges on the common in Figure S12. It confirms that the EDC96 age distribution is less well constrained than Vostok GAD.




Figure S6. Method used to modify the location and scale parameters for the uncertainty assessment. Starting from the optimal values (blue square), the location and scale parameters are moved away following straight lines (dashed lines). If the new RMSD differs by less than 15% with the optimal value (green dots) the location and scale are further modified, until the 15% are reached (red dots). Then the parameters are modified following a new line. In total 10 different lines are tested.




Figure S7. GAD for the DO21 event in Vostok. Left panel: Optimal gas age distribution as a bold solid line. Age distributions resulting in 15% RMSD increases are shown as dashed lines. Middle panel: ~~Methane~~-smoothed record using the optimal GAD as a bold solid line, and ~~methane~~-smoothed records using the 15% higher RMSD distributions are displayed as dashed lines. CFA measurements are displayed in gray. Right panel: Location and scale parameters defining the gas age distributions. The blue square marks the optimal parameters and the red dots represent age distributions resulting in a 15% increase of RMSD.




Figure S8. Same as Figure ~~S3~~S7 for DO6-8 events in EDC96.




Figure S9. Same as Figure S3-S7 for 8.2ka event in EDC99.




Figure S10. Same as Figure S3-S7 for the modern section of Lock-In.




Figure S11. Same as Figure S3-S7 for the modern section of Dome C.




Figure S12. Uncertainty range of the location and scale parameters for the EDC96 DO6-8 and Vostok DO21 GADs. The EDC96 uncertainty range is represented as red dots, and the optimal EDC96 parameters is marked as a blue square. The Vostok DO21 uncertainty range is represented as orange dots, and the optimal Vostok DO21 parameters is marked as a black square.

References

- Chappellaz, J., Stowasser, C., Blunier, T., Baslev-Clausen, D., Brook, E. J., Dallmayr, R., Faïn, X., Lee, J. E., Mitchell, L. E., Pascual, O., Romanini, D., Rosen, J., and Schüpbach, S.: High-resolution glacial and deglacial record of atmospheric methane by continuous-flow and laser spectrometer analysis along the NEEM ice core, *Clim. Past*, 9, 2579–2593, doi:10.5194/cp-9-2579-2013, 2013.
- 5 Fourteau, K., Faïn, X., Martinerie, P., Landais, A., Ekaykin, A. A., Lipenkov, V. Y., and Chappellaz, J.: Analytical constraints on layered gas trapping and smoothing of atmospheric variability in ice under low-accumulation conditions, *Clim. Past*, 13, 1815–1830, doi:10.5194/cp-13-1815-2017, 2017.
- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J.-M., Raynaud, D., Stocker, T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years, *Nature*, 453, 383–386, doi:10.1038/nature06950, 2008.
- 10 Rasmussen, S. O., Abbott, P. M., Blunier, T., Bourne, A. J., Brook, E., Buchardt, S. L., Buizert, C., Chappellaz, J., Clausen, H. B., Cook, E., Dahl-Jensen, D., Davies, S. M., Guillevic, M., Kipfstuhl, S., Laepple, T., Seierstad, I. K., Severinghaus, J. P., Steffensen, J. P., Stowasser, C., Svensson, A., Vallelonga, P., Vinther, B. M., Wilhelms, F., and Winstrup, M.: A first chronology for the North Greenland Eemian Ice Drilling (NEEM) ice core, *Clim. Past*, 9, 2713–2730, doi:10.5194/cp-9-2713-2013, 2013.
- 15 Rhodes, R. H., Faïn, X., Stowasser, C., Blunier, T., Chappellaz, J., McConnell, J. R., Romanini, D., Mitchell, L. E., and Brook, E. J.: Continuous methane measurements from a late Holocene Greenland ice core: atmospheric and in-situ signals, *Earth Planet. Sc. Lett.*, 368, 9–19, doi:10.1016/j.epsl.2013.02.034, 2013.
- Witrant, E., Martinerie, P., Hogan, C., Laube, J. C., Kawamura, K., Capron, E., Montzka, S. A., Dlugokencky, E. J., Etheridge, D., Blunier, T., and Sturges, W. T.: A new multi-gas constrained model of trace gas non-homogeneous transport in firn: evaluation and behaviour at
20 eleven polar sites, *Atmos. Chem. Phys.*, 12, 11465–11483, doi:10.5194/acp-12-11465-2012, 2012.