

Cryptotephra from the Icelandic Veidivötn 1477 CE eruption in a Greenland ice core: confirming the dating of 1450s CE volcanic events and assessing the eruption's climatic impact

5

Peter M. Abbott¹, Gill Plunkett², Christophe Corona³, Nathan J. Chellman⁴, Joseph R. McConnell⁴, John R. Pilcher², Markus Stoffel^{5,6,7}, Michael Sigl¹

10 ¹Climate and Environmental Physics and Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

²Archaeology and Palaeoecology, School of Natural and Built Environment, Queen's University Belfast, Belfast BT7 1NN, UK

³Geolab, Université Clermont Auvergne, CNRS, F-63000, Clermont-Ferrand, France

⁴Desert Research Institute, Nevada System of Higher Education, Reno, Nevada 89512, USA

15 ⁵Climatic Change Impacts and Risks in the Anthropocene (C-CIA), Institute for Environmental Sciences, University of Geneva, 1205 Geneva, Switzerland

⁶Department of Earth Sciences, University of Geneva, 1205 Geneva, Switzerland

⁷Department F.-A. Forel for Environmental and Aquatic Sciences, University of Geneva, 1205 Geneva, Switzerland

Correspondence to: Peter M. Abbott (peter.abbott@climate.unibe.ch)

20 **Abstract.** Volcanic eruptions are a key source of climatic variability and reconstructing their past impact can improve our understanding of the operation of the climate system and increase the accuracy of future climate projections. Two annually resolved and independently dated palaeoarchives – tree rings and polar ice cores – can be used in tandem to assess the timing, strength and climatic impact of volcanic eruptions over the past ~2,500 years. The quantification of post-volcanic climate responses, however, has at times been hampered by differences between simulated and observed temperature
25 responses that raised questions regarding the robustness of the chronologies of both archives. While many chronological mismatches have been resolved, the precise timing and climatic impact of two major sulphate emitting volcanic eruptions during the 1450s CE, including the largest atmospheric sulphate loading event in the last 700 years, has not been constrained. Here we explore this issue through a combination of tephrochronological evidence and high-resolution ice-core chemistry
30 measurements from a Greenland ice core, the TUNU2013 record.

30

We identify tephra from the historically dated 1477 CE eruption of the Icelandic Veidivötn-Bárðarbunga volcanic system in direct association with a notable sulphate peak in TUNU2013 attributed to this event, confirming that this peak can be used as a reliable and precise time-marker. Using seasonal cycles in several chemical elements and 1477 CE as a fixed chronological point shows that ages of 1453 CE and 1458 CE can be attributed, with a high precision, to the start of two
35 other notable sulphate peaks. This confirms the accuracy of a recent Greenland ice-core chronology over the mid- to late 15th

century and corroborates the findings of recent volcanic reconstructions from Greenland and Antarctica. Overall, this implies that large-scale Northern Hemisphere climatic cooling affecting tree-ring growth in 1453 CE was caused by a Northern Hemisphere volcanic eruption in 1452 or early 1453 CE and then a Southern Hemisphere eruption, previously assumed to have triggered the cooling, occurred later in 1457 or 1458 CE.

40

The direct attribution of the 1477 CE sulphate peak to the eruption of *Veidivötn*, one of the most explosive from Iceland in the last 1,200 years, also provides the opportunity to assess the eruption's climatic impact. A tree-ring based reconstruction of Northern Hemisphere summer temperatures shows a cooling in the aftermath of the eruption of $-0.35\text{ }^{\circ}\text{C}$ relative to a 1961-1990 CE reference period and $-0.1\text{ }^{\circ}\text{C}$ relative to the 30-year period around the event, and a relatively weak and spatially incoherent climatic response in comparison to the less explosive but longer-lasting Icelandic *Eldgjá* 939 CE and *Laki* 1783 CE eruptions. In addition, the *Veidivötn* 1477 CE eruption occurred around the inception of the Little Ice Age and could be used as a chronostratigraphic marker to constrain the phasing and spatial variability of climate changes over this transition if it can be traced into more regional palaeoclimatic archives.

45

1. Introduction

50

1.1 Assessing the timing, strength and climatic impact of volcanic eruptions

Volcanic eruptions are a key source of natural climatic variability, with the sulphate aerosols they emit into the stratosphere shielding the Earth from solar radiation, causing short-term cooling from a local to global scale (Robock, 2000; Sigl et al., 2015). Reconstructing climatic variability caused by past volcanic forcing improves understanding of responses, feedbacks and internal variability in the climate system that can help increase the accuracy of future climate predictions (Schurer et al., 2013; Bethke et al., 2017). Two key proxies provide evidence for these reconstructions: tree rings, which record terrestrial climatic changes that could be related to volcanic eruptions, and polar ice cores, which record the atmospheric burden of sulphate aerosols produced by past volcanic eruptions. Both archives have independent annually resolved chronologies that can be used to determine the timing of past volcanic events and their climatic impact. However, the quantification of post-volcanic climate responses have been hampered by differences in simulated and reconstructed temperature changes and potential mismatches in timing between the archives, which raised questions about the robustness of both tree-ring and ice-core chronologies (e.g. Baillie, 2008; Anchukaitis et al., 2012; Esper et al., 2013; Büntgen et al., 2014). Age uncertainties can arise from cumulative age uncertainties in the annual layer interpretation of ice cores, the brevity of the growth phase of tree rings during the Northern Hemisphere summer and the time lag between sulphur emissions at a volcanic source and subsequent deposition of sulphuric aerosols on polar ice-sheets, which can range between a few weeks up to a year depending on location and other eruption source parameters (e.g. Marshall et al., 2019). While the majority of the potential chronological mismatches have been resolved using new ice-core chronologies and long-term hemispheric-wide tree-ring based climate reconstructions (Sigl et al., 2015; Stoffel et al., 2015; Wilson et al., 2016), several enigmas remain. The most

60

65

notable of these unsolved questions is the apparent discrepancy between the timing of two major sulphur-emitting volcanic eruptions, including the largest sulphate loading event in the last 700 years, and large-scale climatic cooling observed during the 1450s CE (Esper et al., 2017). This time period also aligns with the inception of the ‘Little Ice Age’ (LIA; Grove, 2012) in the Northern Hemisphere (e.g. Owens et al., 2017; Büntgen et al., 2020).

The large sulphate loading eruption during the 1450s CE has most commonly been attributed to the formation of the submarine Kuwae caldera offshore from Vanuatu in the South Pacific. This attribution is based on radiocarbon dating of charcoal that places the event, first described in oral histories of local islands (see Ballard, 2020), in the mid-15th century (Monzier et al., 1994; Robin et al., 1994) and petrological evidence that it was a sulphur-rich eruption (Witter and Self, 2007). This attribution has become commonly accepted within the literature despite a lack of confirmatory evidence. In recent studies, however, this source has been challenged or not attributed to the event (e.g. Németh et al., 2007; Esper et al., 2017; Toohey and Sigl, 2017; Hartman et al., 2019).

80 **1.2 Timing of volcanic eruptions in the 1450s CE**

Through the study of bristlecone pines in the western USA, LaMarche and Hirschboeck (1984) identified a series of frost rings, including a widespread frost ring in 1453 CE, that they attributed to large volcanic eruptions. Filion et al. (1986) and Delwaide et al. (1991) also reported light rings related to 1453 CE in more than 25% of subfossil black spruce trees sampled in subarctic Quebec. Briffa et al. (1998) subsequently identified a distinct cooling event in 1453 CE, recorded at multiple sites within their circum-northern hemispheric network of tree-ring chronologies from the northern boreal forest, while Salzer and Hughes (2007) identified evidence for anomalous tree-ring growth in the early and late 1450s CE in western USA bristlecone pines. More recently, warm season temperature reconstructions from Northern Hemisphere tree-ring networks have revealed spatially coherent and exceptional cooling in 1453 CE, with cooling ranging spatially between -0.4 and -6.9°C , marking the onset of a 15-year cold period (Stoffel et al., 2015; Esper et al., 2017; Guillet et al., 2017).

90 Various glaciochemical studies on Antarctic ice cores have provided evidence for a large volcanic event around 1450 CE (e.g. Zanolini et al., 1985; Legrand and Kirchner, 1990; Moore et al., 1991). Delmas et al. (1992) suggested that this so-called “1450 event” was derived from a Southern Hemisphere eruption, based on a limited coeval signal in Greenland (Hammer et al., 1980), and associated it with the climate cooling in the tree-ring records to attribute it an age of 1452 CE. 95 Langway et al. (1995) subsequently identified a bipolar volcanic event to which they attributed an age of 1459 CE and Zielinski (1995) attributed this to the large mid-15th century volcanic event. Later, in their compilation of 33 ice cores from Greenland and Antarctica, Gao et al. (2006) suggested that the 1459 CE signal was from an eruption local to Greenland and concluded that the major Southern Hemisphere volcanic eruption occurred in late 1452 CE or early 1453 CE. However, this study overlooked that many of the ice-core chronologies from Antarctica were fixed to the tree-ring minima of 1453 CE 100 following Briffa et al. (1998). Within the recent high-resolution ice-core based independent volcanic reconstructions of

Plummer et al. (2012) and Sigl et al. (2013, 2014, 2015) from both Greenland and Antarctica, two distinct sulphate peaks can be observed in the 1450s CE (Fig. 1). To provide a consistent terminology these unknown volcanic eruptions have been labelled after the year in which the increase of volcanic sulphate is first detectable in the records (i.e. 1453 CE and 1458 CE). The actual eruption date is either in the same year or the previous year, due to the potential time delay of volcanic sulphate deposition on the ice-core sites (e.g. Marshall et al., 2019). The bipolar expression of these sulphate signals indicates that a comparatively smaller volcanic eruption, with a sulphate burden similar to that following Krakatau 1883 CE, occurred in the low latitudes in 1452 or early 1453 CE and was followed by a massive low latitude event in 1458 CE (Fig. 1; Sigl et al. 2013). Sulphur isotope analysis of both peaks in ice cores from Greenland and of the 1458 CE peak in Antarctica shows that both of these eruptions injected sulphur gases into the stratosphere and thus had the potential to impact global climate (Baroni et al., 2008; Cole-Dai et al., 2013; Gautier et al., 2019).

Sigl et al. (2013) argued that the first eruption in 1453 CE caused the major climatic changes recorded in the tree-ring archives, that 1458 CE is the correct age for the large mid-1450s CE sulphate peak recorded at both poles and that this event contributed towards the decadal persistency of the cooling, but triggered less distinct climatic changes that were only recorded in some tree-ring records (e.g. Salzer and Hughes, 2007). Subsequently, based on the new ages from two ice cores from Antarctica and observations from their tree-ring network, Esper et al. (2017) proposed two scenarios for the larger of the two 1450s CE volcanic events: either the original age of 1452 CE proposed by Delmas et al. (1992) for ice cores in Antarctica was correct and the cooling observed in the tree-ring records in 1453 CE was caused by this event or it occurred in 1458 CE and did not cause large-scale cooling in the Northern Hemisphere extratropics. While omitting a discussion of the second major event in the 1450s CE, Esper et al. (2017) suggested that the first scenario is more likely and contended that the dendroclimatological evidence Sigl et al. (2013) used to support their dating to 1458 CE provides insufficient support for the new age for the larger sulphate loading eruption. Here we further explore this debate and the scenarios proposed by Sigl et al. (2013) and Esper et al. (2017) by testing the accuracy of the recent ice-core chronologies using a tephra horizon from a historically dated Icelandic eruption.

1.3 Determining the source of volcanic eruptions in ice cores

While comparisons of the relative peak sulphate concentrations between ice cores from Greenland and Antarctica can be used to assess if volcanic events occurred in the Southern or Northern Hemisphere extratropics or in the tropics (Sigl et al., 2015; see Sect. 5.1), sulphate records alone cannot pinpoint the exact source of an eruption. Particles of volcanic ash (glass tephra shards) may be preserved in ice cores following eruptions in association with sulphate peaks and their chemical composition is a fingerprint of their volcanic source. Through geochemical analysis of individual glass shards and comparisons to characterisations of proximal deposits it is often possible to determine the volcanic source of these ice-core tephra horizons (Abbott and Davies, 2012). The fine-grained nature, low concentration of particles and lack of visible expression (i.e. cryptotephra) for many of these deposits makes their identification and subsequent geochemical analysis

challenging. However, a range of methods for the preparation and analysis of ice-core tephra samples has been developed
135 and successfully applied to ice cores from Greenland and Antarctica (e.g. Dunbar et al., 2003; Davies et al., 2010; Dunbar
and Kurbatov, 2011; Coulter et al., 2012; Bourne et al., 2015, 2016; Iverson et al., 2017; Cook et al., 2018; Narcisi et al.,
2019).

Identifying tephra in ice cores in direct association with sulphate peaks allows the volcanic source of those emissions to be
140 determined with a high certainty; however, it has been shown that tephra shards are not found in direct association with all
sulphate peaks (e.g. Abbott et al., 2012; Coulter et al., 2012; Bourne et al., 2015). Despite this, any tephra-based source
attributions are useful, especially for sulphate peaks assumed to link to known historical eruptions that provide keystone time
markers for constructing, testing and synchronising ice-core chronologies (e.g. Öræfajökull 1362 CE – Palais et al., 1991;
Laki 1783 CE – Fiacco et al., 1994; Katmai 1912 CE – Coulter et al., 2012). In addition, if such events occur in close
145 association with other sulphate peaks of interest the dating control for all the events will be improved as the relative age
differences between closely-spaced peaks are more precise than absolute age estimates: so-called differential dating
(Andersen et al., 2006). This approach has been used to resolve the timing, strength and climatic impacts of past volcanic
events; for example, the identification of tephra in a Greenland ice core from the ‘Millennium Eruption’ of Changbaishan
(China/North Korea) (Sun et al., 2014) has helped to resolve these factors for that event and the earlier Icelandic Eldgjá
150 eruption (Oppenheimer et al., 2017, 2018).

Here we explore whether tephra from the historically dated 1477 CE eruption of Veiðivötn-Bárðarbunga, Iceland, to which
an obvious sulphate peak is commonly correlated and used as age-marker in Greenland ice-core chronologies (Fig. 1; Sigl et
al., 2013, 2015), can be identified in a Greenland ice core. Direct attribution of this peak to the historical event will test and
155 help improve ice-core dating around this marker and contribute towards the debate regarding the strength and climatic
impact of volcanic eruptions during the 1450s CE. In addition, attribution of the sulphate peak to the 1477 CE Icelandic
eruption would provide the opportunity to explore the climatic impact of sulphate emitted during this eruption. The 1477 CE
eruption was a notable event as it is considered to be one of the most explosive Icelandic eruptions during historical times,
with a volcano explosivity index (VEI; Newhall and Self, 1982) of 6.

160

2. Veiðivötn 1477 CE

Iceland is the primary source of tephra horizons identified in the Greenland ice cores (e.g. Abbott and Davies, 2012; Bourne
et al., 2015). As such, due to the high activity of many Icelandic volcanoes during the historical period (i.e. since the
settlement of Iceland in the late 9th century CE), when many eruptions can be precisely dated to the year, month, day and
165 even hour using documentary records, Icelandic tephra horizons can provide key markers to test ice-core chronologies for
the last ~1,100 years (Larsen et al., 2014). The Veiðivötn-Bárðarbunga volcanic system was one of the most active during
the historical period, based on the frequency of eruptions, with notable eruptions recorded in 1797 CE, 1717 CE, 1500 CE,

1477 CE, ~1410 CE and ~1159 CE (Larsen et al., 1998, 2014). Of these events the large-volume phreatoplinian eruption in 1477 CE was the most explosive and widespread, erupting over 10 km³ of basaltic tephra from a 60–65 km long fissure along the Veiðivötn lake basin and ash from the early stages of the eruption, which lasted for at least 12 hours, was distributed over about half of Iceland to the E, NE and N of the fissure (Larsen, 2005; Larsen et al., 2014; Fig. 2b). The pattern of dispersal of highly fragmented lithic material from this highly explosive eruption suggests an eruption column during the main phase of >18 km height (Larsen, 2005) and the release of sulphuric gases from the sulphur-rich basaltic magma into the stratosphere. Tephra from the 1477 CE eruption has been identified at many proximal and medial sites in Iceland as a visible horizon, including in soil sections, lake sediment cores and in offshore marine cores (Fig. 2b). Two distal cryptotephra occurrences have been identified thus far (Fig. 2a), one in An Loch Mór, western Ireland (MOR-T1; Chambers et al., 2004) and another in Lake Getvaltjärnen, Sweden (Davies et al., 2007). These correlations are underpinned by major element analysis of individual glass shards, but are supported by stratigraphic and geochronological evidence due to geochemical similarities between the basaltic products of the Veiðivötn volcanic system (see Section 5.3).

Early descriptions of deposits from this eruption proximal to Veiðivötn did not ascribe a historical date to the event as it was thought not to have been mentioned in written sources; however, it was assigned an age of 1480 CE ± 11 years based on soil accumulation rates between dated tephra layers (Larsen, 1984). Subsequently, on the basis of geochemical evidence and field mapping, a correlation was found between the Veiðivötn deposits and tephra layer “a”, a widespread historically dated deposit previously thought to have originated from the Kverkfjöll volcanic system (Thórarinnsson, 1958; Benjamínsson, 1981; Larsen et al., 2013). Thórarinnsson (1958) associated tephra layer “a” with an observed tephra-fall in the Eyjafjörður area, North Iceland (Fig. 1) that was subsequently reported in a votive letter written on 11th March 1477 CE. Based on this a historical date of early 1477 CE, most likely February, can be ascribed to the Veiðivötn-Bárðarbunga eruption and it is henceforth referred to as V1477 (G. Larsen, pers. comm., 2020).

190

3. Materials and Methods

3.1 TUNU2013 ice-core chemistry and chronology

The 213 m-long TUNU2013 ice core was retrieved from northeast Greenland during summer 2013 (78.04°N, 33.88°W; Fig. 2a) and covers the period between 280 CE and 2013 CE. Ice-core sections were measured at the Desert Research Institute (DRI) using a continuous melter analysis system that includes two Element2 (Thermo Scientific) high resolution inductively coupled plasma mass spectrometers operating in parallel to provide directly co-registered measurements of a broad range of ~35 elements, including sulphur, calcium and sodium (McConnell et al., 2002; Sigl et al., 2015; Maselli et al., 2017). There are several sources of sulphur in ice core records, therefore the sea salt component was removed to isolate the non-sea-salt (nssS) concentrations (Sigl et al., 2013). In addition to the elemental measurements, co-registered measurements of insoluble particle concentrations, in the size ranges of 2.6–4.5 and 4.5–9.5 µm diameter, were made using an inline Abakus® laser-

200

based particle detector (Ruth et al., 2003). The nssS and fine particle concentrations were used to guide the tephra sampling towards depths with coeval peaks in both parameters. While the particle measurements are not a direct proxy or solely controlled by the presence of tephra shards, this approach has been used successfully to identify tephra horizons in both Greenland and Antarctic ice cores (e.g. Jensen et al., 2014; Dunbar et al., 2017; McConnell et al., 2017, 2020; Plunkett et al., 2020; Smith et al., 2020).

The data from TUNU2013 were synchronised to the annually counted and seasonally resolved NS1-2011 chronology of Sigl et al. (2015) using a series of volcanic markers as tie-points, including the sulphate peak thought to relate to V1477. The period covering 1450–1500 CE could be identified between ~75.79–82.20 m and clear peaks in sulphate thought to have started in 1453 CE, 1458 CE and 1477 CE were observed (Fig. 3a). During this time period the measurement resolution is ~1 cm and annual layers are ~12 cm thick therefore the data has a sub-annual, nominally monthly, resolution with a seasonal precision (i.e. ~2 months) due to variations in snowfall throughout the year. Annual layers were also counted in TUNU2013 between 1445–1485 CE using concentration data for elements that show clear seasonal cycles, namely calcium and sodium whose variations are driven by continental dust and oceanic sea spray deposition, respectively, and the nssS/Na ratio was used to define the annual layer boundary (Sigl et al., 2013; Maselli et al., 2017; Fig. 5). This annual layer count permits the full assessment of the ages of the two 1450s CE sulphate peaks in TUNU2013 using differential dating if the V1477 tephra is identified.

3.2 Cryptotephra analysis

The 1477 CE sulphate peak in TUNU2013 was accompanied by a coeval peak in particle concentration, thus archived ice from the interval between 78.515–78.655 m was sampled directly for tephra analysis to determine if it related to the V1477 eruption (sample code QUB-1965; Fig. 3b). Direct ice samples (QUB-1964, -1963 and -1962) also were taken for tephra analysis across a later peak in particle concentrations, which according to the NS1-2011 is centred around late 1478 CE and early 1479 CE, and precedes a minor sulphate peak in mid to late 1479 CE (Fig. 3b).

The direct ice samples were transferred to individual Nalgene bottles, melted at room temperature and the meltwater centrifuged to concentrate any particulate material present. The concentrated meltwater was evaporated onto a frosted 28 × 48 mm microprobe slide and covered with Buehler EpoxyCure 2 resin. The presence of glass tephra shards was assessed at 100-400x magnification using a polarising light microscope and a significant concentration of shards were identified in QUB-1965. To section the glass shards and create a flat surface for geochemical analysis the epoxy mount was ground using 12 µm alumina powder then polished using 6, 3 and 1 µm diamond paste. The surface of the resin and exposed shards was then carbon coated prior to the concentrations of 11 major and minor elements within individual glass shards being analysed using a JEOL FEGSEM 6500F at Queen's University Belfast (QUB), which integrates wavelength and energy dispersive spectrometry analysis (WDS and EDS respectively; Coulter et al., 2010). Operating conditions followed Coulter et al. (2010)

with the addition of Cl and P measured using EDS and WDS respectively. Further details on the operating conditions are
235 provided in Supplementary Table S3. Glass from the rhyolitic Lipari obsidian and the basaltic Laki 1783 CE tephra were
analysed as secondary standards during the analytical period and compared to recommended values from Kuehn et al. (2011)
and a sample of V1477 tephra from Suðurárbotnar, North Iceland (G. Larsen, pers. comm., 2020) was analysed as a QUB
Veidivötn internal standard. The raw sample and secondary standard analyses are provided in the supplementary information
(Tables S1 and S2).

240

Geochemical analyses from several occurrences of the V1477 tephra were collated for data analysis (Fig. 2; Table S4). To
facilitate graphical and statistical comparisons between all datasets the geochemical data were normalised to an anhydrous
basis (i.e. 100% total oxides) using eight elements consistently analysed in all studies. As such, the minor elements MnO,
P₂O₅ and Cl were not included and their concentrations were deducted from the analytical totals prior to normalisation (see
245 Table S4). Statistical comparisons were made between tephra occurrences using the statistical distance (D^2) test described in
Perkins et al. (1995, 1998) and the similarity coefficient (SC) function of Borchardt et al. (1972) (Tables S5 and S6).

3.3 Tree-ring based reconstructions of cooling induced by Icelandic eruptions

3.3.1 Impact of volcanic eruptions on Northern Hemisphere summer temperatures

In addition to the ice-core and tephra analysis, we quantified the cooling induced by the V1477 eruption, using the NVOLC
250 v2 dataset of Guillet et al. (2020), and compared it to other major Icelandic eruptions of the Common Era. The NVOLC v2
dataset includes 13 tree-ring width (TRW) and 12 maximum latewood density (MXD) chronologies spread across the
Northern Hemisphere (NH; for further details see Stoffel et al. (2015) and Guillet et al. (2017)). All chronologies in NVOLC
v2 have either been calibrated against regional instrumental climate data and transferred into temperature units or have been
interpreted by the original authors as a temperature proxy. Here, a principal component regression (PCR) approach was
255 chosen to transfer TRW and MXD data into NH summer (or JJA for June-July-August) temperature units, expressed as
anomalies with respect to the 1961–1990 reference period. A reduced space signal of the proxy records was then extracted
using principal component analysis (PCA) resulting in a set of principal component (PC) scores and PC loading patterns.
The first n PCs with eigenvalues >1 were retained as predictors to develop a multiple linear regression model. A multiple
cross validation using random calibration sets (bootstrapping) was applied to the PCR to estimate the skill of the
260 reconstruction and confidence intervals around the reconstructed anomalies. Because each chronology had different lengths,
an iterative nesting method was used to develop the temperature reconstruction. This procedure entails the sub-setting of the
original dataset into complete data matrices without missing values, so-called nests.

In NVOLC v2, 23 nests have been adjusted to the common period of all series. The earliest spans the period 500–551 CE
265 and is composed of six chronologies. The most replicated nest (1230–1972 CE) included 25 chronologies. The nested PCR

was computed schematically following a 3-step procedure. In each nest, firstly, the number of predictor variables was reduced using a principal component analysis; secondly, the PCs with eigenvalues >1 were retained as independent variables within Ordinary Least Square (OLS) multiple regression models while a mean NH JJA temperature series, obtained from the Berkeley Earth Surface Temperature (BEST) dataset over the period 1805–1972 CE, was used as a dependent variable. The robustness of each model was tested based on a traditional split calibration/verification procedure bootstrapped 1,000 times and the final reconstruction of each nest was computed as the median of the 1,000 realizations, given with their 95% confidence interval. The skill of each reconstruction has been evaluated based on the coefficient of determination (r^2 for the calibration and R^2 for the verification periods), reduction of error (RE) and coefficient of efficiency (CoE) statistics. The final reconstruction (500–2000 CE) was achieved by splicing all the nested time series after the mean and variance of each nested reconstruction segment had been adjusted to the best replicated nest (1230–1972 CE). Finally, reconstructed temperature anomalies are presented as deviations from the 30-yr mean climatology around the year of the volcanic eruption. For example, for V1477, a background was calculated by averaging the windows 1462–1476 CE and 1478–1493 CE then an anomaly was created by subtracting this background from the 1477 CE reconstructed temperature.

3.3.2 Regional variability of Northern Hemisphere summer cooling

In order to characterize regional-scale climate changes and to reveal spatial anomaly patterns in the aftermath of Icelandic eruptions, we additionally used a network of tree-ring proxy data to develop a climate field reconstruction of extratropical NH JJA temperatures spanning the last 1,500 years. In this approach, we grouped the 25 chronologies from NVOLC v2 in 11 regional clusters using a minimum coefficient correlation (r) threshold above 0.3 ($p < 0.05$) over the common period of the chronologies. Based on the later threshold, we included only one chronology in the Quebec, Central Europe, Siberia–Taymir, Siberia–Yakutia, and China–Qilian Mountains clusters whereas five composed the Western Europe region. In the six clusters that included multiple chronologies, we derived reconstructions from bootstrapped nested PCRs. In the remaining five clusters, we used a bootstrap OLS linear regression to quantify interannual variations of summer temperatures. Based on CoE statistics >0.1 , 3,486 grid points have been reconstructed back to 500 CE. The smallest (Indigirka) and largest (Siberia–Poral Ural) clusters included 53 and 918 grid points, respectively.

4. Results

4.1 Cryptotephra analysis

4.1.1 1477 CE sulphate peak

A closer look at the sub-annual resolution data around the sulphate peak attributed to 1477 CE shows particle concentrations peak in late winter 1477 CE before sulphate concentrations peak approximately 2 months later during the spring and elevated concentrations are broadly restricted to 1477 CE (Fig. 3b). Volcanic particles, including glass tephra shards, are always

transported and deposited rapidly (within days to weeks) following an eruption due to their relatively high density. However, the longer residence time of sulphate aerosols means that these concentrations can be elevated for an extended period following a volcanic eruption, up to several years after a large eruption injecting aerosols high into the stratosphere (Robock, 2000). The relative time delay between peak particle and sulphate deposition can reflect the proximity of the eruptive source, with greater delays observed for more distal eruptions (e.g. Koffman et al., 2013, 2017; Smith et al., 2020). The sharp and narrow peak in sulphate concentrations observed in 1477 CE (in contrast to the one starting in 1458 CE; Fig. 1) and the short delay following the particle concentration peak is more typical of a Northern Hemisphere eruption local to Greenland and the timing of their deposition early in 1477 CE, i.e. during winter, is consistent with the historical date for the V1477 eruption.

305

Within the TUNU2013 ice sample covering 78.515-78.655 m depth (QUB-1965), ~126 tephra glass shards with a greenish-brown platy to blocky morphology, no indication of physical alternation, occasional microlites and long axis diameters up to 30 μm were observed (Fig. 3c). The major element composition of 16 of these shards was analysed and they form a single homogenous population with a tholeiitic basaltic composition (Fig. 4ai). The most likely high-latitude Northern Hemisphere volcanic source of material of this composition is Iceland and comparisons to the geochemistry of material proximal to several Icelandic volcanic systems shows it was most likely sourced from the Veiðivötn-Bárðarbunga system (Fig. 4aai and iii).

Comparisons were made to the collation of geochemical analyses for proximal and distal occurrences of tephra produced during the V1477 event. Graphical comparisons show that for most element oxides, e.g. Al_2O_3 , TiO_2 , K_2O and SiO_2 , there are close similarities between the concentrations observed in the TUNU2013 shards and the proximal and distal V1477 occurrences (Fig. 4bi and ii). However, for elements such as Na_2O , FeO and MgO , some offsets, most notably of up to 1 wt% for MgO , can be observed between the main populations (Fig. 4biii). Some individual analyses from the V1477 occurrences, which could be dismissed as outliers, do however have MgO concentrations similar to those from the TUNU2013 78.655 cm shards (Fig. 4biii). The geochemical differences are reflected in statistical comparisons between the TUNU2013 tephra and the collation of V1477 data (Tables S5 and S6). Statistical distance tests show that in comparison to ~36 % of the V1477 occurrences the composition of TUNU2013 78.655 m is statistically different at the 99 % confidence interval and the D^2 values for several other comparisons are just below the critical value (Table S5). A matrix of similarity coefficient comparisons between the average compositions of the tephra characterisations shows that all of the comparisons of prior V1477 occurrences have high values above typical thresholds for accepting that characterisations of Icelandic basaltic eruptions are identical, with all comparisons >0.95 and the vast majority >0.97 (Table S6). In contrast, comparisons between TUNU2013 78.655 m and the V1477 occurrences range between 0.917 and 0.957, values typically thought to indicate that the tephtras originate from the same volcanic source but not the same eruption (Table S6; Begét et al., 1992; Abbott et al., 2018).

330

Geochemical offsets can result from tephra shards with the same composition being analysed under contrasting operating conditions on different microprobes and/or at different times (Kuehn et al., 2011). However, a comparison of secondary standard analyses from the analytical periods for the TUNU2013 shards and those from the Skaftártunga occurrence of V1477 (Streeter and Dugmore, 2014) strongly indicates that the geochemical offsets are not the result of analytical issues but are real differences in composition between the sample datasets (Fig. S1). Furthermore, the V1477 tephra analysed as a QUB Veidivötn internal standard during the same analytical period as the TUNU2013 shards did not display the same MgO offset, but MgO values consistent with the main V1477 population (Fig. S2).

Overall, despite the geochemical offset, the evidence supporting the correlation of the TUNU2013 78.655 m tephra layer to the V1477 event is strong. Firstly, the geochemical differences between TUNU2013 78.655 m and the V1477 occurrences are only slight and for most elements the concentrations are highly similar. Possible explanations for the differences in composition will be discussed in Sect. 5.4. Secondly, the ice-core chemistry and the shard geochemistry indicate that TUNU2013 78.655 m was sourced from a Northern Hemisphere volcanic region close to Greenland and has a composition matching products of the Veidivötn-Bárðarbunga volcanic system. Thirdly, while it is acknowledged that there can be uncertainties in ice-core chronologies, chronological uncertainties in this age range would be of the order of years rather than the tens to hundreds of years required for TUNU2013 78.655 m to be the product of one of the other known Veidivötn-Bárðarbunga eruptions (e.g. V1717, V1500, ~V1410), which have a similar geochemical composition. In addition, of these eruptions V1477 had the largest magnitude and thus the greatest potential for widespread deposition of its fallout. Lastly, as the Icelandic tephrostratigraphy is well documented over the last 1,000 years (e.g. Thordarson and Larsen, 2007; Larsen and Eiríksson, 2008), it is unlikely that TUNU2013 78.655 m was deposited following a previously unknown eruption of the Veidivötn-Bárðarbunga volcanic system of a sufficient magnitude to deposit tephra over Greenland.

4.1.2 1479 CE sulphate peak

A limited number of tephra shards were found in the ice samples from the later microparticle peak at the start of 1479 CE, with up to three brown shards identified in both QUB-1963 and QUB-1962. A tephra layer with a rhyolitic composition was identified previously around 1479 CE by Fiacco et al. (1993) in the GISP2 ice core and attributed to an eruption from Mount St Helens, USA, dated using dendrochronology (Yamaguchi, 1983), although the geochemical match with this source is ambiguous (Abbott and Davies, 2012). The relationship between microparticle and sulphate concentrations for the 1479 CE event in TUNU2013 matches the observations of Fiacco et al. (1993), which also showed a peak in microparticles preceding a sulphate peak by about half a year (Fig. 3b). The tephra shards from QUB-1963 and QUB-1962 have not yet been geochemically analysed due to their small size (<20 µm) and low concentration. Therefore, it was not possible to determine if they correlate to the rhyolitic 1479 CE tephra of Fiacco et al. (1993), however, a geochemical correlation is unlikely as the physical characteristics of the TUNU2013 shards are indicative of a mafic composition. Based on the physical characteristics

it is possible that these shards represent remobilised material from V1477, however, without geochemical analysis we cannot determine if this is the case or if the tephra derived from another eruption.

365 **4.2 Differential dating of 1450s CE volcanic events**

Through the identification of the V1477 tephra layer in TUNU2013 in direct association with the large sulphate peak at ~78.56 m, it is demonstrated that the prior assumption that it represents aerosol deposition from the 1477 CE eruption of Veidivötn is correct. As such, the assertion that 1477 CE can be used as the age for this ice-core depth with 0 years uncertainty is appropriate. Using this age as a fix-point for the annual layer counts for TUNU2013 between 1445 CE and 370 1485 CE it is shown that the ages of 1453 CE and 1458 CE for the start of the two 1450s CE sulphate peaks are in agreement with those derived from the NS1-2011 chronology (Fig. 5). These ages were based on annual layers counts from NEEM-2011-S1 using an age of 1477 CE for a notable sulphate peak in that sequence (Sigl et al., 2015). The close correspondence between the nssS records from TUNU2013 and NEEM-2011-S1 and the strong annual signals in TUNU2013 provide further evidence to support those ages for the events (Fig. S3).

375 **4.3 Tree-ring based reconstruction of summer temperature cooling induced by Icelandic eruptions**

Within the NVOLC v2 reconstruction, each nest passed all verification tests, R^2 and RE, for the calibration period, as well as r^2 and CoE for the verification period. For the 1420-1650 CE period (Fig. 6a), the unfiltered reconstruction indicates a cooling in the aftermath of V1477 of -0.35°C relative to the 1961–1990 CE reference period, the 356th coldest NH summer since 500 CE (i.e. rank 356). By comparison, stronger cooling is observed following the Icelandic Eldgjá 939 CE (-0.5°C in 380 940 CE, rank 205) and Laki 1783 CE (-1.2°C , rank 9) eruptions. More extreme negative anomalies are observed following other major explosive volcanic eruptions, including -1.6°C in 1601 CE (rank 1, Huaynaputina), -1.5°C in 1816 CE (rank 3, Tambora), -1.4°C in 536 CE (rank 4, unidentified volcano) and -1.3°C in 1453 CE (rank 5, unidentified volcano) (Fig. 6a). After filtering with a 30-yr running mean centered around the year of the calculated anomaly, the NH summer cooling prevailing in 1477 CE was -0.1°C (Fig. 6b), the 524th coldest NH summer since 500 CE. Our findings are corroborated 385 through comparisons with other recent, large-scale reconstructions, Sch2015 (Schneider et al., 2015) and NTREND2015 (Wilson et al., 2016), which rank 1477 CE as the 538th (-0.04°C) and 182th (-0.23°C) coolest summers, respectively (Fig. 6b). The Sch2015, NTREND2015 and NVOLC v2 reconstructions all show more significant cooling in the aftermath of the Eldgjá and Laki eruptions, with cooling ranging between -0.4 to -0.6°C and -0.7 to -1°C , respectively (Fig. 6b). Grouping the 25 chronologies into 11 clusters enables an estimation of the regional variability of summer cooling induced by the 390 Icelandic eruptions (Fig. 6c). Based on this approach, the JJA gridded temperature reconstructions for 940 CE and 1783 CE (Fig. 6c) indicate that volcanic forcing induced widespread cooling over the NH. Extremely cold conditions prevailed in Scandinavia, Western Europe and in Coastal Alaska in 940 CE, and on the Yamal Peninsula and in Alaska in 1783 CE. Less consistent patterns are observed in the reconstructed temperatures following the V1477 eruption, with cold temperatures in

Scandinavia, Western Europe and Central Asia, contrasting with above-average anomalies over North America, the Polar
395 Ural and on the Yamal Peninsula (Fig. 6c).

5. Discussion

5.1 Confirming the timing and hemisphere of mid-15th century volcanic eruptions

The identification of the V1477 tephra layer in TUNU2013 has demonstrated the veracity of the age of 1477 CE commonly
400 used for a keystone sulphate peak identified in many ice cores from Greenland. This provides evidence to support the
accuracy of the NS1-2011 Greenland ice-core chronology around this period and corroborates the ages proposed by Sigl et
al. (2013) for the mid-1450s CE sulphate peaks in the Greenland ice-core records deposited following large volcanic
eruptions in 1452 or early 1453 CE and 1458 CE. The ages, temporal separation and duration of sulphate deposition for these
events are consistent with those for two volcanic eruptions recorded in the Antarctic WDC core, whose ages were derived
405 using an independent annual-counted timescale and not bipolar synchronisation of the sulphate records (Sigl et al., 2013; Fig.
1). As highlighted by Sigl et al. (2013), the sulphate deposition in Greenland relating to the later event is greater than for the
earlier event, and is also consistent with the record from Antarctica. Further independent confirmation of the matching of the
events could be achieved if the temporal evolution of sulphur isotope anomalies over the peak starting in 1458 CE is
consistent between ice cores from both polar ice sheets, as was demonstrated recently for the Samalas 1257 CE eruption
410 (Burke et al., 2019).

While factors other than latitude may influence the stratospheric distribution of sulphate between both hemispheres, aerosol
modelling suggests that a likely range of latitudinal positions for eruption sources can be approximated based on the
asymmetry of sulphate deposition between Greenland and Antarctica (Toohey et al., 2016, 2019; Marshall et al., 2019).
415 Cumulative sulphate deposition for the 1453 CE event was higher in Greenland (25 kg km^{-2}) than in Antarctica (5 kg km^{-2}),
deduced from a network of polar ice cores (Sigl et al., 2014; Toohey and Sigl, 2017). This suggests that 83% of the
atmospheric sulphate burden remained in the Northern Hemisphere, which is more likely to occur after an eruption from a
source located in that hemisphere. Conversely, for the 1458 CE event, sulphate deposition was higher in Antarctica (64 kg
 km^{-2}) than in Greenland (35 kg km^{-2}) suggesting that 64% of the atmospheric sulphate burden remained in the Southern
420 Hemisphere and we can attribute the eruption to a source in that hemisphere.

As such, these findings demonstrate that the scenario proposed, but not favoured, by Esper et al. (2017), is correct, i.e. a
strongly asymmetric Southern Hemisphere sulphate loading event occurred in 1457 or 1458 CE without causing large-scale
Northern Hemisphere cooling. In addition, the findings back up the argument of Sigl et al. (2013) that the first eruption in
425 1452 or early 1453 CE occurred in the Northern Hemisphere and caused the distinct climatic cooling observed in the tree-

ring records. The close succession of the two large sulphur-rich volcanic eruptions was most likely the cause of the longevity of the 15-year cold period following 1453 CE.

5.2 Testing ice-core chronologies and determining volcanic source using tephra horizons

This study has shown the power of using historically dated tephra horizons to test the accuracy of ice-core chronologies and assumptions regarding the source of sulphate aerosols deposited over ice sheets, which has critical implications for assessing the climatic impact of eruptions. The example provided here can be added to others from Greenland, with tephra from Öræfajökull 1362 CE, Laki 1783 CE and Katmai 1912 CE having all been identified in ice cores from Greenland in association with chemical indicators previously attributed to those eruptions (Palais et al., 1991; Fiacco et al., 1994; Coulter et al., 2012). Determining the source of notable peaks in sulphate or other volcanic indicators (e.g. acidity and electrical conductivity) using tephra is necessary, when possible, as associations to specific events can become embedded in the literature with limited supporting evidence, even if only originally proposed as a possibility. For example, associations between chemical indicators and the Hekla 1104 CE and Vesuvius 79 CE eruptions, that previously have been used as historical tie points with no uncertainty while building chronologies (Hammer et al., 1980; Zielinski et al., 1994; Vinther et al., 2006), have been found to be inaccurate following re-evaluation of the Greenland ice-core chronologies (Sigl et al., 2015). The volcanic signal previously associated with Hekla 1104 CE has been re-dated to 1108 CE and Guillet et al. (2020) have recently proposed it can be linked to an eruption of Mount Asama, Japan; however, at present this is not supported by tephra evidence. The volcanic signal previously associated with Vesuvius 79 CE has been redated to 87/88 CE by Sigl et al. (2015). The re-evaluation of the Greenland ice-core chronology was aided by documented historical reports (Sigl et al., 2015) and cosmogenic radionuclides in tree rings and ice cores as another form of isochronous marker horizon, most likely produced by solar particle events (Miyake et al., 2012, 2015; Mekhaldi et al., 2015; Büntgen et al., 2018). This highlights that tephra is just one of the tools that can be used to evaluate and improve ice-core chronologies.

Testing associations between volcanic indicators in ice cores and specific events using tephra could be useful for determining the source of the two 1450s CE eruptions, whose sources are currently unknown. The 1458 CE event often has been attributed to the formation of the submarine Kuwae caldera, however, the evidence is inconclusive and circumstantial so the source is still open for debate. Hartman et al. (2019) characterised tephra from an Antarctic ice core found in association with the 1458CE event and concluded that it was distinct to eruptive products from Kuwae; however, despite some geochemical similarities to the products of a local Antarctic volcano, a definitive source was not proposed. Based on the size of the particles and the expression of the sulphate in different ice cores, Hartman et al. (2019) suggest that it resulted from a tropospheric aerosol cloud; however, this is inconsistent with sulphur isotope evidence from Antarctica for stratospheric injection of the sulphate cloud (Baroni et al., 2008; Gautier et al., 2019). A local tropospheric source is also inconsistent with the widespread deposition of volcanic sulphate that prevailed for 2–3 years as observed in a large network of 16 ice cores all over Antarctica (Sigl et al., 2014; Toohey and Sigl, 2017). Further studies of sulphur isotopes coupled

with attempts to identify tephra associated with both 1450s CE events and more work on proximal records, could aid in
460 conclusively identifying their sources.

5.3 Use of V1477 as a chronostratigraphic marker for the inception of the LIA

The mid-15th century is widely considered as the inception of the LIA (e.g. Owens et al., 2017; Büntgen et al., 2020), that potentially was intensified by the large 1450s CE volcanic events (Miller et al., 2012; Schurer et al., 2014; Slawinska and Robock, 2018). As such, if the V1477 tephra layer could be traced into more palaeoclimatic archives it could act as a useful
465 chronostratigraphic marker for constraining the chronology of different proxy records and assessing the phasing and spatial variability of climatic changes associated with the LIA onset. The identification of the V1477 tephra layer in a Greenland ice core corroborates the findings of past studies of terrestrial sequences from Ireland and Sweden (Chambers et al., 2004; Davies et al., 2007) that following the eruption its products were dispersed over a wide geographical area. Therefore, it has significant potential to act as a chronostratigraphic tie-line linking records from Greenland, the North Atlantic and Northwest
470 Europe. As one of the most explosive Icelandic eruptions of the last 1,200 years, it is notable that V1477 has not already been identified at more sites and established as a key horizon in the tephrochronological framework for northern Europe (Lawson et al., 2012). The generally lower explosivity of Icelandic basaltic eruptions reduces the likelihood of the injection of material high into the atmosphere (Lawson et al., 2012), and the potential transport distance of the tephra is then dependent upon tropospheric conditions at the time of the eruption. Differences between glass shards produced during
475 basaltic and rhyolitic eruptions, may further constrain the long-range transport of basaltic glass tephra shards with the greater density and generally larger grain-size making the shards more prone to rapid fallout (Watson et al., 2017). However, it seems unlikely that these factors can explain the lack of more widespread V1477 occurrences in NW Europe given the prior distal identifications and the known high explosivity and plume height of the eruption. It has been proposed that basaltic tephra shards are not identified in some terrestrial sequences as they can degrade in acidic peatland environments (Pollard et al., 2003) or are overlooked in lake sediments due to the difficulties in separating and identifying basaltic shards (Lawson et al., 2012). In addition, non-identification could be due to (1) spatial heterogeneity in distal ash deposition (Davies et al., 2010), (2) latitudinal differences in wind strength, since although rarely identified in continental European sequences basaltic shards have been identified in a Russian Arctic site located a similar distance from Iceland but further north (Vakhrameeva et al., 2020), and/or (3) this specific time interval not being investigated. Therefore, we propose that the potential use of
485 V1477 as a chronostratigraphic marker could be increased through focused cryptotephra investigations of specific time windows and, for lake sediments, incorporating magnetic separation processing (e.g. Mackie et al., 2002) to maximise the isolation of basaltic material.

A key consideration for any future studies aiming to trace V1477 is the strong similarity between the geochemical
490 compositions of V1477 and other notable Veidivötn sourced tephra layers such as V1717, ~V1410 and V877, the Landnám tephra (Cage et al., 2011; Harning et al., 2018). To date, only the V1477 and Landnám tephra have been identified

definitively in European records, although the potential presence of tephra shards from other Veidivötn eruptions (e.g. ~V1159, V1717, V1766 and V1797) has been discussed (Plunkett and Pilcher, 2018; Vakhrameeva et al., 2020). Therefore, in palaeoarchives with a lower temporal resolution than ice cores independent chronological evidence would be required to support any correlations to specific Veidivötn eruptions, such as the radiocarbon age Davies et al. (2007) used to support the correlation of the Lake Getvaltjärnen deposit to V1477.

5.4 Geochemical variability of V1477 deposits

A notable observation from this work was a slight geochemical offset between tephra shards from TUNU2013 78.655 m and the V1477 occurrences (Fig. 4b). However, as outlined in Sect. 4.1.1 it does not hinder the attribution of TUNU2013 78.655 m to V1477. The offsets are most notable for the MgO concentrations which, except for a few outlier analyses from other deposits, are ~0.75-1 wt% higher in the TUNU2013 shards (Fig. 4biii). MgO concentrations around 7-8 wt% are not atypical for Veidivötn products (Fig. 4aai), but are more characteristic of Veidivötn eruptions that occurred earlier in the Holocene than 1477 CE (Óladóttir et al., 2011; Caracciolo et al., 2020).

The difference in MgO concentrations could indicate that the TUNU2013 shards have a more primitive, i.e. less evolved, geochemical composition than shards from the other V1477 occurrences. A general trend of increasing MgO values with decreasing FeO concentrations can be observed in the V1477 data (Fig. 4biii) and is most obvious in the characterisation of V1477 from Skaftártunga (Streeter and Dugmore, 2014). Within the Skaftártunga characterisation, alongside shards with strong similarities to the other V1477 analyses, more geochemically evolved shards with lower MgO concentrations and higher FeO values than most of the other V1477 analyses can be observed as well as two shard analyses with MgO and FeO values comparable to the TUNU2013 78.655 m data (Fig. 4biii). This overall trend could be indicative of geochemical evolution during the eruption, however, such clear fractionation trends are not observed for other element pairs. The bilobate form of proximal V1477 deposits on isopach maps and differences in the size and morphology of tephra shards between the lobes indicates that differential dispersal occurred with plumes to the ENE and NNW of the fissure (Benjamínsson, 1981). It has been suggested that the two lobes resulted from either a shift in wind direction from SW to SE during the eruption (Benjamínsson, 1981) or bifurcation of the volcanic plume due to a crosswind (Ernst et al., 1994). A change in wind direction during the course of the eruption could have caused products from different temporal phases to be transported along the two dispersal axes and account for the slight differences between geochemical characterisations from specific sites. Geochemical variability between coarser and finer grain-size fractions deposited at more proximal and more distal sites respectively could also be a factor; however, significant differences in grain-size are unlikely for the more distal deposits and when compared they still display the offset in MgO.

525 As prior studies have not identified the geochemical differences highlighted in this study the offsets could have resulted from analytical uncertainty; however, this is unlikely based on the secondary and internal standard analyses made when the TUNU2013 shards were analysed (Figs. S1 and S2). Further investigation of V1477 occurrences at proximal sites and in other Greenland ice cores could determine if the geochemical offset is consistent for other deposits along the same dispersal axis or can be related to grain-size differences and the potential for offsets should be considered within any attempts to trace this horizon.

5.5 Climatic impact of V1477 relative to other Icelandic eruptions

The definitive attribution of the 1477 CE sulphate peak in Greenland to the Icelandic V1477 eruption also provided an opportunity to assess the climatic impact of this explosive eruption in comparison to the effects of other historical eruptions. Overall, despite having a VEI of 6, the NH cooling following V1477 was relatively weak and spatially incoherent in comparison to the less explosive Icelandic Laki and Eldgjá eruptions (VEI 4-5), and more explosive lower-latitude eruptions, e.g. Tambora and Huaynaputina (Fig. 6; Sect. 4.3). The greater impact of the more effusive Icelandic eruptions relative to the V1477 eruption can be attributed to the huge sulphate load emitted during the Laki and Eldgjá eruptions over prolonged periods of time, ~8 months and >2 years, respectively, in comparison to the short-lived, days to weeks, duration of the V1477 eruption (Thordarson et al., 2001; Thordarson and Larsen, 2007). The short residence time of sulphate in the atmosphere following the V1477 eruption is highlighted by the sharp and narrow nature of the 1477 CE sulphate peak in the Greenland ice cores (Fig. 1).

6. Conclusions

545 Using sulphate and particle concentration records as a guide, volcanic ash from the Icelandic eruption of Veidivötn in 1477 CE has been identified as a cryptotephra in the TUNU2013 Greenland ice core. This identification verifies the widely assumed source of a notable sulphate peak, present in many Greenland ice cores, and confirms the accuracy of the NS1-2011 chronology around this depth. In addition, using annual layer counting the ages of 1452 CE and 1458 CE attributed to two volcanic events in the 1450s CE are confirmed with a high accuracy. While being considered one of the most explosive Icelandic eruptions of the last 1,200 years, a tree-ring based reconstruction of summer temperatures demonstrates that V1477 caused weak and spatially incoherent summer cooling in comparison to other recent Icelandic eruptions that, while less explosive, emitted a higher sulphate load. The timing of the eruption around the inception of the LIA in the NH means V1477 could act as a chronostratigraphic marker for assessing climatic changes associated with this onset. However, potential geochemical variability between V1477 deposits and geochemical similarities with the products of other eruptions from Veidivötn must be considered when correlations are being assessed.

Data availability

The data presented and utilised in this study are available in the supplementary material.

Author contributions

560 All authors contributed towards obtaining and analysing the data utilised in the study. MSi and GP conceived the research. JRM, NJC and MSi contributed ice-core glaciochemical records. GP and JRP performed tephra geochemical analyses. MSt and CC produced the North Hemisphere tree-ring based summer temperature reconstruction. PA wrote the manuscript with MSt and CC contributing Sections 3.3 and 4.3 and Figure 6. All authors contributed towards improving and editing the manuscript.

565 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

Thanks to Carla Grimaldi for compiling some of the V1477 geochemical data. Thanks to Guðrún Larsen for a detailed description of the history of the dating of V1477 and Esther Ruth Guðmundsdóttir for providing geochemical data from the
570 V1477 occurrence in Lake Lögurinn. Thanks to Richard Streeter and Chris Hayward for providing secondary standard data related to the V1477 occurrence from Skaftártunga. We thank Karen Fontijn and an anonymous reviewer for their comments that have helped to improve the manuscript. Thanks also to Matthew Toohey, the guest editor for this manuscript, for his helpful comments. This research benefitted from the participation of the authors in the Volcanic Impacts on Climate and Society (VICS) working group of the Past Global Changes (PAGES) project.

575 Financial support

PA and MSi received funding from the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 820047). MSi received funding from the Geoscience Department and the NFR TOPPFORK project "VIKINGS" at the University of Oslo. The collection and analysis of the TUNU2013 core was supported by the U.S. National Science Foundation (grant #1204176) to JRM, with additional support (grant #1925417) to
580 JRM and NJC for the volcanic interpretation. MSt and CC received funding from the Swiss National Sciences Foundation through the SNSF Sinergia CALDERA project (grant agreement no. CRSII5_183571).

References

- Abbott, P.M. and Davies, S.M.: Volcanism and the Greenland ice-cores: the tephra record. *Earth-Sci. Rev.*, 115(3), 173-191, doi: 10.1016/j.earscirev.2012.09.001, 2012.
- 585
- Abbott, P.M., Davies, S.M., Steffensen, J.P., Pearce, N.J.G., Bigler, M., Johnsen, S.J., Seierstad, I.K., Svensson, A. and Wastegård, S.: A detailed framework of Marine Isotope Stage 4 and 5 volcanic events in two Greenland ice-cores. *Quaternary Sci. Rev.*, 36, 59-77, doi: 10.1016/j.quascirev.2011.05.001, 2012.
- 590
- Abbott, P.M., Griggs, A.J., Bourne, A.J., Chapman, M.R. and Davies, S.M.: Tracing marine cryptotephra in the North Atlantic during the last glacial period: Improving the North Atlantic marine tephrostratigraphic framework. *Quaternary Sci. Rev.*, 189, 169-186, doi: 10.1016/j.quascirev.2018.03.023, 2018.
- 595
- Anchukaitis, K. J., Breitenmoser, P., Briffa, K. R., Buchwal, A., Büntgen, U., Cook, E. R., D'Arrigo, R. D., Esper, J., Evans, M. N., Frank, D., Grudd, H., Gunnarson, B. E., Hughes, M. K., Kirilyanov, A. V., Körner, C., Krusic, P. J., Luckman, B., Melvin, T. M., Salzer, M. W., Shashkin, A. V., Timmermann, C., Vaganov, E. A. and Wilson, R. J. S.: Tree rings and volcanic cooling. *Nat. Geosci.*, 5, 836-837, doi: 10.1038/ngeo1645, 2012.
- 600
- Andersen, K.K., Svensson, A., Johnsen, S.J., Rasmussen, S.O., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M.-L., Steffensen, J.P., Dahl-Jensen, D., Vinther, B.M. and Clausen, H.B.: The Greenland Ice Core Chronology 2005, 15-42 ka. Part 1: constructing the time scale. *Quaternary Sci. Rev.*, 25(23-24), 3246-3257, doi: 10.1016/j.quascirev.2006.08.002, 2006.
- 605
- Baillie, M.G.L.: Proposed re-dating of the European ice core chronology by seven years prior to the 7th century AD. *Geophys. Res. Lett.*, 35, L15813, doi: 10.1029/2008GL034755, 2008.
- Ballard, C.: The Lizard in the Volcano: Narratives of the Kuwae Eruption. *Contemp. Pac.*, 32(1), 98-123, doi: 10.1353/cp.2020.0005, 2020.
- 610
- Baroni, M., Savarino, J., Cole-Dai, J. H., Rai, V. K. and Thiemens, M. H.: Anomalous sulfur isotope compositions of volcanic sulfate over the last millennium in Antarctic ice cores. *J. Geophys. Res.*, 113(D20), D20112, doi: 10.1029/2008JD010185, 2008.
- 615
- Begét, J., Mason, O. and Anderson, P.: Age, Extent and Climatic Significance of the c. 3400 BP Aniakhchak Tephra, Western Alaska, USA. *Holocene*, 2(1), 51-56, doi: 10.1177/095968369200200106, 1992.
- Benjamínsson, J.: Tephra layer "a", in: *Tephra Studies*, edited by Self, S. and Sparks, R.S.J., Springer, Dordrecht, The Netherlands, 331-335, 1981.
- 620
- Bergþórsdóttir, H.B.: A 3000 year high resolution multi-proxy record of environmental change from lake Gripdeild, eastern Iceland. MS. University of Iceland, Iceland, 85 pp., 2014.
- 625
- Bethke, I., Outten, S., Otterå, O. H., Hawkins, E., Wagner, S., Sigl, M. and Thorne, P.: Potential volcanic impacts on future climate variability. *Nat. Clim. Change*, 7, 799-805, doi: 10.1038/nclimate3394, 2017.
- Borchardt, G.A., Aruscavage, P.J. and Millard, H.T.: Correlation of the Bishop Ash, a Pleistocene marker bed, using instrumental neutron activation analysis. *J. Sediment. Res.*, 42(2), 301-306, doi: 10.1306/74D72527-2B21-11D7-8648000102C1865D, 1972.

- 630 Bourne, A.J., Abbott, P.M., Albert, P.G., Cook, E., Pearce, N.J.G., Ponomareva, V., Svensson, A. and Davies, S.M.: Underestimated risks of recurrent long-range dispersal from northern Pacific Arc volcanoes. *Sci. Rep.*, 6, 29837, doi: 10.1038/srep29837, 2016.
- Bourne, A.J., Cook, E., Abbott, P.M., Seierstad, I.K., Steffensen, J.P., Svensson, A., Fischer, H., Schüpbach, S. and Davies, S.M.: A tephra lattice for Greenland and a reconstruction of volcanic events spanning 25-45 ka b2k. *Quaternary Sci. Rev.*, 118, 122-141, doi: 10.1016/j.quascirev.2014.07.017, 2015.
- 635 Briffa, K.R., Jones, P.D., Schweingruber, F.H. and Osborn, T.J.: Influence of volcanic eruptions on Northern Hemisphere summer temperature over the past 600 years. *Nature*, 393, 450-455, doi: 10.1038/30943, 1998.
- 640 Büntgen, U., Arseneault, D., Boucher, É., Churakova, O. V., Gennaretti, F., Crivellaro, A., Hughes, M. K., Kirilyanov, A. V., Klippel, L., Krusic, P. J., Linderholm, H. W., Ljungqvist, F. C., Ludescher, J., McCormick, M., Mygland, V. S., Nicolussi, K., Piermattei, A., Oppenheimer, C., Reinig, F., Sigl, M., Vaganov, E. A., and Esper, J.: Prominent role of volcanism in Common Era climate variability and human history, *Dendrochronologia*, 64, 125757, doi: 10.1016/j.dendro.2020.125757, 2020.
- 645 Büntgen, U., Wacker, L., Diego Galván, J., Arnold, S., Arseneault, D., Baillie, M., Beer, J., Bernabei, M., Bleicher, N., Boswijk, G., Bräuning, A., Carrer, M., Charpentier Ljungqvist, F., Cherubini, P., Christl, M., Christie, D.A., Clark, P.W., Cook, E.R., D'Arrigo, R., Davi, N., Eggertsson, Ó., Esper, J., Fowler, A.M., Gedalof, Z., Gennaretti, F., Gießinger, J., Grissino-Mayer, H., Grudd, H., Gunnarson, B.E., Hantemirov, R., Herzig, F., Hessler, A., Heussner, K.-U., Jull, A.J.T., Kukarskih, V., Kirilyanov, A., Kolář, T., Krusic, P.J., Kyncl, T., Lara, A., LeQuesne, C., Linderholm, H.W., Loader, N.J., Luckman, B., Miyake, F., Mygland, V.S., Nicolussi, K., Oppenheimer, C., Palmer, J., Panyushkina, I., Pederson, N., Rybníček, M., Schweingruber, F.H., Seim, A., Sigl, M., Churakova (Sidorova), O., Speer, J.H., Synal, H.-A., Tegel, W., Treydte, K., Villalba, R., Wiles, G., Wilson, R., Winship, L.J., Wunder, J., Yang, B. and Young, G.H.F.: Tree rings reveal globally coherent signature of cosmogenic radiocarbon events in 774 and 993 CE. *Nat. Commun.*, 9, 3605, doi: 10.1038/s41467-018-06036-0, 2018.
- 650 Büntgen, U., Wacker, L., Nicolussi, K., Sigl, M., Gütler, D., Tegel, W., Krusic, P.J. and Esper, J.: Extraterrestrial confirmation of tree-ring dating. *Nat. Clim. Change*, 4, 404-405, doi: 10.1038/nclimate2240, 2014.
- 660 Burke, A., Moore, K.A., Sigl, M., Nita, D.C., McConnell, J.R. and Adkins, J.F.: Stratospheric eruptions from tropical and extra-tropical volcanoes constrained using high-resolution sulfur isotopes in ice cores. *Earth Planet. Sci. Lett.*, 521, 113-119, doi: 10.1016/j.epsl.2019.06.006, 2019.
- 665 Cage, A.G., Davies, S.M., Wastegård, S. and Austin, W.E.N.: Identification of the Icelandic Landnám tephra (AD 871 ± 2) in Scottish fjordic sediment. *Quatern. Int.*, 246(1-2), 168-176, doi: 10.1016/j.quaint.2011.08.016, 2011.
- 670 Caracciolo, A., Bali, E., Guðfinnsson, G.H., Kahl, M., Halldórsson, S.A., Hartley, M.E. and Gunnarsson, H.: Temporal evolution of magma and crystal mush storage conditions in the Bárðarbunga-Veiðivötn volcanic system, Iceland. *Lithos*, 352-353, 105234, doi: 10.1016/j.lithos.2019.105234, 2020.
- Chambers, F.M., Daniell, J.R.G., Hunt, J.B., Molloy, K. and O'Connell, M.: Tephrostratigraphy of An Loch Mór, Inis Oírr, western Ireland: implications for Holocene tephrochronology in the northeastern Atlantic region. *Holocene*, 14(5), 703-720, doi: 10.1191/0959683604hl749rp, 2004.
- 675 Cole-Dai, J., Ferris, D.G., Lanciki, A.L., Savarino, J., Thiemens, M.H. and McConnell, J.R.: Two likely stratospheric volcanic eruptions in the 1450s C.E. found in a bipolar, subannually dated 800 year ice core record. *J. Geophys. Res.*, 118, 7459-7466, doi: 10.1002/jgrd.50587, 2013.

- 680 Cook, E., Portnyagin, M., Ponomareva, V., Bazanova, L., Svensson, A. and Garbe-Schönberg, D.: First identification of cryptotephra from the Kamchatka Peninsula in a Greenland ice core: Implications of a widespread marker deposit that links Greenland to the Pacific northwest. *Quaternary Sci. Rev.*, 181, 200-206, doi: 10.1016/j.quascirev.2017.11.036, 2018.
- 685 Coulter, S.E., Pilcher, J.R., Hall, V.A., Plunkett, G. and Davies, S.M.: Testing the reliability of the JEOL FEGSEM 6500F electron microprobe for quantitative major element analysis of glass shards from rhyolitic tephra. *Boreas*, 39, 163-169, doi: 10.1111/j.1502-3885.2009.00113.x, 2010.
- 690 Coulter, S.E., Pilcher, J.R., Plunkett, G., Baillie, M., Hall, V.A., Steffensen, J.P., Vinther, B.M., Clausen, H.B. and Johnsen, S.J.: Holocene tephtras highlight complexity of volcanic signals in Greenland ice cores. *J. Geophys. Res.*, 117, D21303, doi: 10.1029/2012JD017698, 2012.
- Davies, S.M., Wastegård, S., Abbott, P.M., Barbante, C., Bigler, M., Johnsen, S.J., Rasmussen, T.L., Steffensen, J.P. and Svensson, A.: Tracing volcanic events in the NGRIP ice-core and synchronising North Atlantic marine records during the last glacial period. *Earth Planet. Sci. Lett.*, 294(1-2), 69-79, doi: 10.1016/j.epsl.2010.03.004, 2010.
- 695 Davies, S.M., Elmquist, M., Bergman, J., Wohlfarth, B. and Hammarlund, D.: Cryptotephra sedimentation processes within two lacustrine sequences from west central Sweden. *Holocene*, 17(3), 1-13, doi: 10.1177/0959683607076443, 2007.
- Davies, S.M., Larsen, G., Wastegård, S., Turney, C.S.M., Hall, V.A., Coyle, L. and Thordarson, T.: Widespread dispersal of Icelandic tephra: how does the Eyjafjöll eruption of 2010 compare to past Icelandic events?. *J. Quaternary Sci.*, 25(5), 605-611, doi: 10.1002/jqs.1421, 2010.
- 700 Delmas, R.J., Kirchner, S., Palais, J.M. and Petit, J.R.: 1000 years of explosive volcanism recorded at the South Pole. *Tellus B*, 44(4), 335-350, doi: 10.1034/j.1600-0889.1992.00011.x, 1992.
- 705 Delwaide, A., Fillion, L. and Payette, S.: Spatiotemporal distribution of light rings in subarctic black spruce, Quebec. *Can. J. Forest Res.*, 21, 1828-1832, doi: 10.1139/x91-252, 1991.
- Dunbar, N.W., Iverson, N.A., Van Eaton, A.R., Sigl, M., Alloway, B.V., Kurbatov, A.V., Mastin, L.G., McConnell, J.R. and Wilson, C.J.N.: New Zealand supereruption provides time marker for the Last Glacial Maximum in Antarctica. *Sci. Rep.*, 7, 12238, doi: 10.1038/s41598-017-11758-0, 2017.
- 710 Dunbar, N.W. and Kurbatov, A.V.: Tephrochronology of the Siple Dome ice core, West Antarctica: correlations and sources. *Quaternary Sci. Rev.*, 30(13-14), 1602-1614, doi: 10.1016/j.quascirev.2011.03.015, 2011.
- 715 Dunbar, N.W., Zielinski, G.A. and Voisins, D.T.: Tephra layers in the Siple Dome and Taylor Dome ice cores, Antarctica: Sources and correlations. *J. Geophys. Res.*, 108(B8), 2374, doi: 10.1029/2002JB002056, 2003.
- Ernst, G.G.J., Davies, J.P. and Sparks, R.S.J.: Bifurcation of volcanic plumes in a crosswind. *Bull. Volcanol.*, 56, 159-169, doi: 10.1007/BF00279601, 1994.
- 720 Esper, J., Büntgen, U., Hartl-Meier, C., Oppenheimer, C. and Schneider, L.: Northern Hemisphere temperature anomalies during the 1450s period of ambiguous volcanic forcing. *Bull. Volcanol.*, 79, 41, doi: 10.1007/s00445-017-1125-9, 2017.
- Esper, J., Büntgen, U., Luterbacher, J. and Krusic, P. J.: Testing the hypothesis of post-volcanic missing rings in temperature sensitive dendrochronological data. *Dendrochronologia*, 31, 216-222, doi: 10.1016/j.dendro.2012.11.002, 2013.
- 725 Fiacco, R.J., Palais, J.M., Germani, M.S., Zielinski, G.A. and Mayewski, P.A.: Characteristics and Possible Source of a 1479 A.D. Volcanic Ash Layer in a Greenland Ice Core. *Quaternary Res.*, 39(3), 267-273, doi: 10.1006/qres.1993.1033, 1993.

- 730 Fiacco, R.J., Thordarson, T., Germani, M.S., Self, S., Palais, J.M., Whitlow, S. and Grootes, P.M.: Atmospheric Aerosol Loading and Transport Due to the 1783-84 Laki Eruption in Iceland, Interpreted from Ash Particles and Acidity in the GISP2 Ice Core. *Quaternary Res.*, 42(3), 231-240, doi: 10.1006/qres.1994.1074, 1994.
- Filion, L., Payette, S., Gauthier, L. and Boutin, Y.: Light rings in subarctic conifers as a dendrochronological tool.
735 *Quaternary Res.*, 26(2), 272-279, doi: 10.1016/0033-5894(86)90111-0, 1986.
- Gao, C., Robock, A., Self, S., Witter, J.B., Steffensen, J.P., Clausen, H.B., Siggaard-Andersen, M.-L., Johnsen, S.J., Mayewski, P.A. and Ammann, C.: The 1452 or 1453 A.D. Kuwae eruption signal derived from multiple ice core records: Greatest volcanic sulfate event of the past 700 years. *J. Geophys. Res.*, 111(D12), D12107, doi: 10.1029/2005JD006710,
740 2006.
- Gautier, E., Savarino, J., Hoek, J., Erbland, J., Caillon, N., Hattori, S., Yoshida, N., Albalat, E., Albarede, F. and Farquhar, J.: 2600-years of stratospheric volcanism through sulfate isotopes. *Nat. Commun.*, 10, 466, doi: 10.1038/s41467-019-08357-0, 2019.
745
- Grove, J.M.: *The Little Ice Age*. Routledge, London, United Kingdom, 2012.
- Gudmundsdóttir, E.R., Eiríksson, J. and Larsen, G.: Identification and definition of primary and reworked tephra in Late Glacial and Holocene marine shelf sediments off North Iceland. *J. Quaternary Sci.*, 26(6), 589-602, doi: 10.1002/jqs.1474,
750 2011.
- Gudmundsdóttir, E.R., Larsen, G., Björck, S., Ingólfsson, Ó. and Striberger, J.: A new high-resolution Holocene tephra stratigraphy in eastern Iceland: Improving the Icelandic and North Atlantic tephrochronology. *Quaternary Sci. Rev.*, 150, 234-249, doi: 10.1016/j.quascirev.2016.08.011, 2016.
755
- Guillet, S., Corona, C., Stoffel, M., Khodri, M., Lavigne, F., Ortega, P., Eckert, N., Sielenou, P.D., Daux, V., Churakova (Sidorova), O.V., Davi, N., Edouard, J.-L., Zhang, Y., Luckman, B.H., Myglan, V.S., Guiot, J., Beniston, M., Masson-Delmotte, V. and Oppenheimer, C.: Climate response to the Samalas volcanic eruption in 1257 revealed by proxy records. *Nat. Geosci.*, 10, 123-128, doi: 10.1038/ngeo2875, 2017.
760
- Guillet, S., Corona, C., Ludlow, F., Oppenheimer, C. and Stoffel, M.: Climatic and societal impacts of a “forgotten” cluster of volcanic eruptions in 1108-1110 CE. *Sci. Rep.*, 10, 6715, doi: 10.1038/s41598-020-63339-3, 2020.
- Hammer, C.U., Clausen, H.B. and Dansgaard, W.: Greenland ice sheet evidence of post-glacial volcanism and its climatic impact. *Nature*, 288, 230-235, doi: 10.1038/288230a0, 1980.
765
- Harning, D.J., Thordarson, T., Geirsdóttir, Á., Zalzal, K. and Miller, G.H.: Provenance, stratigraphy and chronology of Holocene tephra from Vestfirðir, Iceland. *Quat. Geochronol.*, 46, 59-76, doi: 10.1016/j.quageo.2018.03.007, 2018.
- 770 Hartman, L.H., Kurbatov, A.V., Winski, D.A., Cruz-Uribe, A.M., Davies, S.M., Dunbar, N.W., Iverson, N.A., Aydin, M., Fegyveresi, J.M., Ferris, D.G., Fudge, T.J., Osterberg, E.C., Hargreaves, G.M. and Yates, M.G.: Volcanic glass properties from 1459 C.E. volcanic event in South Pole ice core dismiss Kuwae caldera as a potential source. *Sci. Rep.*, 9, 14437, doi: 10.1038/s41598-019-50939-x, 2019.
- 775 Iverson, N.A., Kalteyer, D., Dunbar, N.W., Kurbatov, A. and Yates, M.: Advancements and best practices for analysis and correlation of tephra and cryptotephra in ice. *Quat. Geochronol.*, 40, 45-55, doi: 10.1016/j.quageo.2016.09.008, 2017.

- 780 Jensen, B.J.L., Pyne-O'Donnell, S., Plunkett, G., Froese, D.G., Hughes, P.D.M., Sigl, M., McConnell, J.R., Amesbury, M.J., Blackwell, P.G., van den Bogaard, C., Buck, C.E., Charman, D.J., Clague, J.J., Hall, V.A., Koch, J., Mackay, H., Mallon, G., McColl, L. and Pilcher, J.R.: Transatlantic distribution of the Alaskan White River Ash. *Geology*, 42(10), 875-878, doi: 10.1130/G35945.1, 2014.
- 785 Koffman, B.G., Dowd, E.G., Osterberg, E.C., Ferris, D.G., Hartman, L.H., Wheatley, S.D., Kurbatov, A.V., Wong, G.J., Markle, B.R., Dunbar, N.W., Kreutz, K.J. and Yates, M.: Rapid transport of ash and sulfate from the 2011 Puyehue-Cordón Caulle (Chile) eruption to West Antarctica. *J. Geophys. Res. Atmos.*, 122, 8908-8920, doi: 10.1002/2017JD026893, 2017.
- 790 Koffman, B.G., Kreutz, K.J., Kurbatov, A.V. and Dunbar, N.W.: Impact of known local and tropical volcanic eruptions of the past millennium on the WAIS Divide microparticle record. *Geophys. Res. Lett.*, 40, 4712-4716, doi: 10.1002/grl.50822, 2013.
- Kuehn, S.C., Froese, D.G. and Shane, P.A.R.: The INTAV intercomparison of electron-beam microanalysis of glass by tephrochronology laboratories: Results and recommendations. *Quatern. Int.*, 246, 19-47, doi: 10.1016/j.quaint.2011.08.022, 2011.
- 795 LaMarche Jr, V.C. and Hirschboeck, K.K.: Frost rings in trees as records of major volcanic eruptions. *Nature*, 307, 121-126, doi: 10.1038/307121a0, 1984.
- 800 Langway Jr, C.C., Osada, K., Clausen, H.B., Hammer, C.U. and Shoji, H.: A 10-century comparison of prominent bipolar volcanic events in ice cores. *J. Geophys. Res.*, 110(D8), 16241-16247, doi: 10.1029/95JD01175, 1995.
- Larsen, D.J., Miller, G.H., Geirsdóttir, Á. and Thordarson, T.: A 3000-year varved record of glacier activity and climate change from the proglacial lake Hvítárvatn, Iceland. *Quaternary Sci. Rev.*, 30(19-20), 2715-2731, doi: 10.1016/j.quascirev.2011.05.026, 2011.
- 805 Larsen, G.: Recent volcanic history of the Veiðivötn fissure swarm, southern Iceland – an approach to volcanic risk assessment. *J. Volcanol. Geoth. Res.*, 22(1-2), 33-58, doi: 10.1016/0377-0273(84)90034-9, 1984.
- Larsen, G.: Explosive Volcanism in Iceland: Three Examples of Hydromagmatic Basaltic Eruptions on long Volcanic Fissures within the past 1200 Years. *Geophysical Research Abstracts*, 7, 10158, doi: 10.1016/j.epsl.2008, 2005
- 810 Larsen, G. and Eiríksson, J.: Late Quaternary terrestrial tephrochronology of Iceland – frequency of explosive eruptions, type and volume of tephra deposits. *J. Quaternary Sci.*, 23(2), 109-120, doi: 10.1002/jqs.1129, 2008.
- 815 Larsen, G., Eiríksson, J., Knudsen, K.L. and Heinemeier, J.: Correlation of late Holocene terrestrial and marine tephra markers, north Iceland: implications for reservoir age changes. *Polar Res.*, 21(2), 283-290, doi: 10.3402/polar.v21i2.6489, 2002.
- 820 Larsen, G., Eiríksson, J. and Gudmundsdóttir, E.R.: Last millennium dispersal of air-fall tephra and ocean-rafted pumice towards the north Icelandic shelf and the Nordic seas, in: *Marine Tephrochronology*, edited by: Austin, W.E.N., Abbott, P.M., Davies, S.M., Pearce, N.J.G. and Wastegård, S., Geological Society, London, 113-211, doi: 10.1144/SP398.4, 2014.
- Larsen, G., Gudmundsson, M.T. and Björnsson, H.: Eight centuries of periodic volcanism at the centre of the Iceland hotspot revealed by glacier tephrostratigraphy. *Geology*, 26(10), 943-946, doi: 10.1130/0091-7613(1998)026<0943:ECOPVA>2.3.CO;2, 1998.
- 825


- Larsen, G., Gudmundsson, M.T., Einarsson, P. and Thordarson, T.: Bárðarbunga, in: Náttúruvá á Íslandi - Eldgos og jarðskjálftar, edited by Sólves, S., Sigmundsson, F. and Bessason, B., Viðlagatrygging Íslands/Háskólaútgáfan, Reykjavík, 253-261, 2013.
- 830 Lawson, I.T., Gathorne-Hardy, F.J., Church, M.J., Newton, A.J., Edwards, K.J., Dugmore, A.J. and Einarsson, Á.: Environmental impacts of the Norse settlement: palaeoenvironmental data from Mývatnssveit, northern Iceland. *Boreas*, 36(1), 1-19, doi: 10.1111/j.1502-3885.2007.tb01176.x, 2007.
- 835 Lawson, I.T., Swindles, G.T., Plunkett, G. and Greenberg, D.: The spatial distribution of Holocene cryptotephra in north-west Europe since 7 ka: implications for understanding ash fall events from Icelandic eruptions. *Quaternary Sci. Rev.*, 41, 57-66, doi: 10.1016/j.quascirev.2012.02.018, 2012.
- 840 Legrand, M.R. and Kirchner, S.: Origins and variations of nitrate in south polar precipitation. *J. Geophys. Res.*, 95(D4), 3493-3507, doi: 10.1029/JD095iD04p03493, 1990.
- Le Maitre, R.W., Bateman, P., Dudek, A., Keller, J., Lameyre, J., Le Bas, M.J., Sabine, P.A., Schmid, R., Sorensen, H., Streckeisen, A. and Woolley, A.R.: A classification of igneous rocks and glossary of terms. Blackwell, London, 1989.
- 845 MacDonald, G.A. and Katsura, T.: Chemical composition of Hawaiian lavas. *J. Petrol.*, 5(1), 82-133, doi: 10.1093/petrology/5.1.82, 1964.
- Mackie, E.A.V., Davies, S.M., Turney, C.S.M., Dobbyn, K., Lowe, J.J. and Hill, P.G.: The use of magnetic separation techniques to detect basaltic microtephra in last glacial-interglacial transition (LGIT; 15-10 ka cal. BP) sediment sequences in Scotland. *Scot. J. Geol.*, 38, 21-30, doi: 10.1144/sjg38010021, 2002.
- 850 Marshall, L., Johnson, J.S., Mann, G.W., Lee, L., Dhomse, S.S., Regayre, L., Yoshioka, M., Carslaw, K.S., and Schmidt, A.: Exploring How Eruptive Source Parameters Affect Volcanic Radiative Forcing Using Statistical Emulation. *J. Geophys. Res.*, 124(2), 964-985, doi: 10.1029/2018JD028675, 2019.
- 855 Maselli, O. J., Chellman, N. J., Grieman, M., Layman, L., McConnell, J. R., Pasteris, D., Rhodes, R. H., Saltzman, E. S. and Sigl, M.: Sea ice and pollution-modulated changes in Greenland ice core methanesulfonate and bromine. *Clim. Past*, 13, 39-59, doi: 10.5194/cp-13-39-2017, 2017.
- 860 McConnell, J. R., Burke, A., Dunbar, N. W., Köhler, P., Thomas, J. L., Arienzo, M. M., Chellman, N. J., Maselli, O. J., Sigl, M., Adkins, J. F., Baggenstos, D., Burkhardt, J. F., Brook, E. J., Buizert, C., Cole-Dai, J., Fudge, T. J., Knorr, G., Graf, H. F., Grieman, M. M., Iverson, N., McGwire, K. C., Mulvaney, R., Paris, G., Rhodes, R. H., Saltzman, E. S., Severinghaus, J. P., Steffensen, J. P., Taylor, K. C. and Winckler, G.: Synchronous volcanic eruptions and abrupt climate change similar to 17.7 ka plausibly linked by stratospheric ozone depletion. *P. Natl. Acad. Sci. USA*, 114(38), 10035-10040, doi: 10.1073/pnas.1705595114, 2017.
- 865 McConnell, J.R., Lamorey, G.W., Lambert, S.W. and Taylor, K.C.: Continuous Ice-Core Chemical Analyses Using Inductively Coupled Plasma Mass Spectrometry. *Environ. Sci. Technol.*, 36(1), 7-11, doi: 10.1021/es011088z, 2002.
- 870 McConnell, J. R., Sigl, M., Plunkett, G., Burke, A., Kim, W.M., Raible, C. C., Wilson, A. I., Manning, J. G., Ludlow, F. M., Chellman, N. J., Innes, H. M., Yang, Z., Larsen, J. F., Schaefer, J. R., Kipfthuhl, S., Mojtavavi, S., Wilhelms, F., Opel, T., Meyer, H., and Steffensen, J. P.: Extreme climate after massive eruption of Alaska's Okmok volcano in 43 BCE and effects on the late Roman Republic and Ptolemaic Kingdom, *P. Natl. Acad. Sci. USA*, 202002722, doi: 10.1073/pnas.2002722117, 2020.
- 875

- Mekhaldi, F., Muscheler, R., Adolphi, F., Aldahan, A., Beer, J., McConnell, J.R., Possnert, G., Sigl, M., Svensson, A., Synal, H.-A., Welten, K.C. and Woodruff, T.E.: Multiradionuclide evidence for the solar origin of the cosmic-ray events of AD 774/5 and 993/4. *Nat. Commun.*, 6, 8611, doi: 10.1038/ncomms9611, 2015.
- 880 Miller, G. H., Geirsdóttir, A., Zhong, Y. F., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson, C., Björnsson, H. and Thordarson, T.: Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks. *Geophys. Res. Lett.*, 39(2), L02708, doi: 10.1029/2011GL050168, 2012.
- 885 Miyake, F., Nagaya, K., Masuda, K. and Nakamura, T.: A signature of cosmic-ray increase in AD 774-775 from tree rings in Japan. *Nature*, 486, 240-242, doi: 10.1038/nature11123, 2012.
- Miyake, F., Suzuki, A., Masuda, K., Horiuchi, K., Motoyama, H., Matsuzaki, H., Motizuki, Y., Takahashi, K. and Nakai, Y.: Cosmic ray event of AD 774-775 shown in quasi-annual ¹⁰Be data from the Antarctic Dome Fuji ice core. *Geophys. Res. Lett.*, 42(1), 84-89, doi: 10.1002/2014GL062218, 2015.
- 890 Monzier, M., Robin, C. and Eissen, J.-P.: Kuwae (≈ 1425 A.D.): the forgotten caldera. *J. Volcanol. Geoth. Res.*, 59(3), 207-218, doi: /10.1016/0377-0273(94)90091-4, 207-218.
- 895 Moore, J.C., Narita, H. and Maeno, N.: A continuous 770-year record of volcanic activity from east Antarctica. *J. Geophys. Res.*, 96(D9), 17353-17359, doi: 10.1029/91JD01283, 1991.
- Narcisi, B., Petit, J.R., Delmonte, B., Batanova, V. and Savarino, J.: Multiple sources for tephra from AD 1259 volcanic signal on Antarctic ice cores. *Quaternary Sci. Rev.*, 210, 164-174, doi: 10.1016/j.quascirev.2019.03.005, 2019.
- 900 Németh, K., Cronin, S.J. and White, J.D.L.: Kuwae Caldera and Climate Confusion. *The Open Geology Journal*, 1, 7-11, doi: 10.2174/1874262900701010007, 2007.
- Newhall, C.G. and Self, S.: The volcanic explosivity index (VEI) an estimate of explosive magnitude for historical volcanism. *J. Geophys. Res.*, 87(C2), 1231-1238.
- 905 Óladóttir, B.A., Larsen, G. and Sigmarsson, O.: Holocene volcanic activity at Grímsvötn, Bárðarbunga and Kverkfjöll subglacial centres beneath Vatnajökull, Iceland. *Bull. Volcanol.*, 73, 1187-1208, doi: 10.1007/s00445-011-0461-4, 2011.
- 910 Óladóttir, B.A., Thordarson, T., Geirsdóttir, Á., Jóhannsdóttir, G.E. and Mangerud, J.: The Saksunarvatn Ash and the G10ka series tephra. Review and current state of knowledge. *Quat. Geochronol.*, 56, 101041, doi: 10.1016/j.quageo.2019.101041, 2020.
- Oppenheimer, C., Orchard, A., Stoffel, M., Newfield, T.P., Guillet, S., Corona, C., Sigl, M., Di Cosmo, N. and Büntgen, U.: The Eldgjá eruption: timing, long-range impacts and influence on the Christianisation of Iceland. *Climatic Change*, 147, 369-381, doi: 10.1007/s10584-018-2171-9, 2018.
- 915 Oppenheimer, C., Wacker, L., Xu, J., Diego Galván, J., Stoffel, M., Guillet, S., Corona, C., Sigl, M., Di Cosmo, N., Hajdas, I., Pan, B., Breuker, R., Schneider, L., Esper, J., Fei, J., Hammond, J.O.S. and Büntgen, U.: Multi-proxy dating the 'Millennium Eruption' of Changbaishan to late 946 CE. *Quaternary Sci. Rev.*, 158, 164-171, doi: 10.1016/j.quascirev.2016.12.024, 2017.
- 920 Owens, M. J., Lockwood, M., Hawkins, E., Usoskin, I., Jones, G. S., Barnard, L., Schurer, A. and Fasullo, J.: The Maunder minimum and the Little Ice Age: an update from recent reconstructions and climate simulations, *J. Sp. Weather Sp. Clim.*, 7,


- 925 A33, doi:10.1051/swsc/2017034, 2017.
- Palais, J.M., Taylor, K., Mayewski, P.A. and Grootes, P.: Volcanic ash from the 1362 AD Öræfajökull eruption (Iceland) in the Greenland Ice Sheet. *Geophys. Res. Lett.*, 18(7), 1241-1244, doi: 10.1029/91GL01557, 1991.
- 930 Perkins, M.E., Brown, F.H., Nash, W.P., Williams, S.K. and McIntosh, W.: Sequence, age and source of silicic fallout tuffs in middle to late Miocene basins of the northern Basin and Range province. *Geol. Soc. Am. Bull.*, 110(3), 344-360, doi: 10.1130/00167606(1998)110<0344:SAASOS>2.3.CO;2, 1998.
- Perkins, M.E., Nash, W.P., Brown, F.H. and Fleck, R.J.: Fallout tuffs of Trapper Creek, Idaho – A record of Miocene explosive volcanism in the Snake River Plain volcanic province. *Geol. Soc. Am. Bull.*, 107(12), 1484-1506, doi: 10.1130/0016-7606(1995)107<1484:FTOTCI>2.3.CO;2, 1995.
- 935 Plummer, C.T., Curran, M.A.J., van Ommen, T.D., Rasmussen, S.O., Moy, A.D., Vance, T.R., Clausen, H.B., Vinther, B.M. and Mayewski, P.A.: An independently dated 2000-yr volcanic record from Law Dome, East Antarctica, including a new perspective on the dating of the 1450s CE eruption of Kuwae, Vanuatu. *Clim. Past.*, 8, 1929-1940, doi: 10.5194/cp-8-1929-2012, 2012.
- 940 Plunkett, G. and Pilcher, J.R.: Defining the potential source region of volcanic ash in northwest Europe during the Mid- to Late Holocene. *Earth-Sci. Rev.*, 179, 20-37, doi: 10.1016/j.earscirev.2018.02.006, 2018.
- 945 Plunkett, G., Sigl, M., Pilcher, J. R., McConnell, J. R., Chellman, N., Steffensen, J. P. and Büntgen, U.: Smoking guns and volcanic ash: the importance of sparse tephra in Greenland ice cores, *Polar Res.*, 39, doi: 10.33265/polar.v39.3511, 2020.
- Pollard, A.M., Blockley, S.P.E. and Ward, K.R.: Chemical alteration of tephra in the depositional environment: theoretical stability modelling. *J. Quaternary Sci.*, 18(5), 385-394, doi: 10.1002/jqs.760, 2003.
- 950 Robin, C., Monzier, M. and Eissen, J.-P.: Formation of the mid-fifteenth century Kuwae caldera (Vanuatu) by an initial hydroclastic and subsequent ignimbrite eruption. *B. Volcanol.*, 56, 170-183, doi: 10.1007/BF00279602, 1994.
- 955 Robock, A.: Volcanic eruptions and climate. *Rev. Geophys.*, 38(2), 191-219, doi: 10.1029/1998RG000054, 2000.
- Ruth, U., Wagenbach, D., Steffensen, J.P. and Bigler, M.: Continuous records of microparticle concentration and size distribution in the central Greenland NGRIP ice core during the last glacial period. *J. Geophys. Res.*, 108(D3), 4098, doi: 10.1029/2002JD002376, 2003.
- 960 Salzer, M.W. and Hughes, M.K.: Bristlecone pine tree rings and volcanic eruptions over the last 5000 yr. *Quaternary Res.*, 67(1), 57-68, doi: 10.1016/j.yqres.2006.07.004, 2007.
- 965 Schneider, L., Smerdon, J.E., Büntgen, U., Wilson, R.J.S., Myglan, V.S., Kirilyanov, A.V. and Esper, J.: Revising midlatitude summer temperatures back to A.D. 600 based on a wood density network. *Geophys. Res. Lett.*, 42(11), 4556-4562, doi: 10.1002/2015GL063956, 2015.
- Schurer, A.P., Hegerl, G.C., Mann, M. E., Tett, S.F.B. and Phipps, S.J.: Separating Forced from Chaotic Climate Variability over the Past Millennium. *J. Climate*, 26, 6954-6973, doi: 10.1175/JCLI-D-12-00826.1, 2013.
- 970 Schurer, A.P., Tett, S.F.B. and Hegerl, G.C.: Small influence of solar variability on climate over the past millennium. *Nat. Geosci.* 7, 104-108, doi: 10.1038/ngeo2040, 2014.

- 975 Sigl, M., Fudge, T.J., Winstrup, M., Cole-Dai, J., Ferris, D., McConnell, J.R., Taylor, K.C., Welten, K.C., Woodruff, T.E., Adolphi, F., Bisiaux, M., Brook, E.J., Buizert, C., Caffee, M.W., Dunbar, N.W., Edwards, R., Geng, L., Iverson, N., Koffman, B., Layman, L., Maselli, O.J., McGwire, K., Muscheler, R., Nishiizumi, K., Pasteris, D.R., Rhodes, R.H. and Sowers, T.A.: The WAIS Divide deep ice core WD2014 chronology – Part 2: Annual-layer counting (0-31 ka BP). *Clim. Past.* 12, 769-786, doi: 10.5194/cp-12-769-2016, 2016.
- 980 Sigl, M., McConnell, J.R., Layman, L., Maselli, O., McGwire, K., Pasteris, D., Dahl-Jensen, D., Steffensen, J.P., Vinther, B., Edwards, R., Mulvaney, R. and Kipfstuhl, S.: A new bipolar ice core record of volcanism from WAIS Divide and NEEM and implications for climate forcing of the last 2000 years. *J. Geophys. Res.*, 118(3), 1151-1169, doi: 10.1029/2012JD018603, 2013.
- 985 Sigl, M., McConnell, J.R., Toohey, M., Curran, M., Das, S.B., Edwards, R., Isaksson, E., Kawamura, K., Kipfstuhl, S., Krüger, K., Layman, L., Maselli, O., Motizuki, Y., Motoyama, H., Pateris, D.R. and Severi, M.: Insights from Antarctica on volcanic forcing during the Common Era. *Nat. Clim. Change*, 4, 693-697, doi: 10.1038/nclimate2293, 2014.
- 990 Sigl, M., Winstrup, M., McConnell, J.R., Welten, K.C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O.J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D.R., Pilcher, J.R., Salzer, M., Schüpbach, S., Steffensen, J.P., Vinther, B.M. and Woodruff, T.E.: Timing and climate forcing of volcanic eruptions for the past 2,500 years. *Nature*, 523, 543-549, doi: 10.1038/nature14565, 2015.
- 995 Slawinska, J. and Robock, A.: Impact of Volcanic Eruptions on Decadal to Centennial Fluctuations of Arctic Sea Ice Extent during the Last Millennium and on Initiation of the Little Ice Age. *J. Climate*, 31, 2145-2167, doi: 10.1175/JCLI-D-16-0498.1, 2018.
- 1000 Smith, V.C., Costa, A., Aguirre-Díaz, G., Pedrazzi, D., Scifo, A., Plunkett, G., Poret, M., Tournigand, P.-Y., Miles, D., Dee, M.W., McConnell, J.R., Sunyé-Puchol, Dávila Harris, P., Sigl, M., Pilcher, J.R., Chellman, N. and Gutiérrez, E.: The magnitude and impact of the 431 CE Tierra Blanca Joven eruption of Ilopango, El Salvador. *P. Natl. Acad. Sci. USA*, ____ doi: [10.1073/pnas.2003008117](https://doi.org/10.1073/pnas.2003008117), 2020.
- 1005 Stoffel, M., Khodri, M., Corona, C., Guillet, S., Poulain, V., Bekki, S., Guiot, J., Luckman, B.H., Oppenheimer, C., Lebas, N., Beniston, M. and Masson-Delmotte, V.: Estimates of volcanic-induced cooling in the Northern Hemisphere over the past 1,500 years. *Nat. Geosci.*, 8, 784-788, doi: 10.1038/ngeo2526, 2015.
- Streeter, R. and Dugmore, A.J.: Late-Holocene land surface change in a coupled social-ecological system, southern Iceland: a cross-scale tephrochronology approach. *Quaternary Sci. Rev.*, 86, 99-114, doi: 10.1016/j.quascirev.2013.12.016, 2014.
- 1010 Sun, C., Plunkett, G., Liu, J., Zhao, H., Sigl, M., McConnell, J.R., Pilcher, J.R., Vinther, B., Steffensen, J.P. and Hall, V.: Ash from Changbaishan Millennium eruption recorded in Greenland ice: Implications for determining the eruption's timing and impact. *Geophys. Res. Lett.*, 41(2), 694-701, doi: 10.1002/2013GL058642, 2014.
- 1015 Thórarinnsson, S.: The Öræfajökull Eruption of 1362. *Acta Nat. Isl.*, 2(2), 1-99, 1958.
- Thordarson, T. and Larsen, G.: Volcanism in Iceland in historical time: Volcano types, eruption styles and eruptive history. *J. Geodyn.*, 43(1), 118-152, doi: 10.1016/j.jog.2006.09.005, 2007.
- 1020 Thordarson, T., Miller, D.J., Larsen, G., Self, S. and Sigurdsson, H.: New estimates of sulfur degassing and atmospheric mass-loading by the 934 AD Eldgjá eruption, Iceland. *J. Volcanol. Geoth. Res.*, 108(1-4), 33-54, doi: 10.1016/S0377-0273(00)00277-8, 2001.


- 1025 Toohey, M., Krüger, K., Schmidt, H., Timmreck, C., Sigl, M., Stoffel, M. and Wilson, R.: Disproportionately strong climate forcing from extratropical explosive volcanic eruptions. *Nat. Geosci.*, 12(2), 100–107, doi:10.1038/s41561-018-0286-2, 2019.
- Toohey, M., Krüger, K., Sigl, M., Stordal, F. and Svensen, H.: Climatic and societal impacts of a volcanic double event at the dawn of the Middle Ages. *Clim. Change*, 136(3–4), 401–412, doi:10.1007/s10584-016-1648-7, 2016.
- 1030 Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500 BCE to 1900 CE. *Earth Syst. Sci. Data*, 9, 809-831, doi: 10.5194/essd-9-809-2017, 2017.
- 1035 Vakhrameeva, P., Portnyagin, M., Ponomareva, V., Abbott, P.M., Repkina, T., Novikova, A., Koutsodendris, A. and Pross, J.: Identification of Icelandic tephra from the last two millennia in the White Sea region (Vodoprovodnoe peat bog, northwestern Russia). *J. Quaternary Sci.*, 35(4), 493-504, doi: 10.1002/jqs.3190, 2020.
- 1040 Vinther, B.M., Clausen, H.B., Johnsen, S.J., Rasmussen, S.O., Andersen, K.K., Buchardt, S.L., Dahl-Jensen, D., Seierstad, I.K., Siggaard-Andersen, M.-L., Steffensen, J.P., Svensson, A., Olsen, J. and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene. *J. Geophys. Res.*, 111, D13102, doi: 10.1029/2005JD006921, 2006.
- Watson, E.J., Swindles, G.T., Savov, I.P., Lawson, I.T., Connor, C.B. and Wilson, J.A.: Estimating the frequency of volcanic ash clouds over northern Europe. *Earth Planet. Sci. Lett.*, 460, 41-49, doi: 10.1016/j.epsl.2016.11.054, 2017.
- 1045 Wilson, R., Anchukaitis, K., Briffa, K. R., Büntgen, U., Cook, E., D'Arrigo, R., Davi, N., Esper, J., Frank, D., Gunnarson, B., Hegerl, G., Helama, S., Klesse, S., Krusic, P. J., Linderholm, H. W., Myglan, V., Osborn, T. J., Rydval, M., Schneider, L., Schurer, A., Wiles, G., Zhang, P. and Zorita, E.: Last millennium northern hemisphere summer temperatures from tree rings: Part I: The long term context. *Quaternary Sci. Rev.*, 134, 1-18, doi: 10.1016/j.quascirev.2015.12.005, 2016.
- 1050 Witter, J.B. and Self, S.: The Kuwae (Vanuatu) eruption of AD 1452: potential magnitude and impact and volatile release. *B. Volcanol.*, 69, 301-318, doi: 10.1007/s00445-006-0075-4, 2007.
- Yamaguchi, D.K.: New tree-ring dates for recent eruptions of Mount St. Helens. *Quaternary Res.*, 20(2), 246-250, doi: 10.1016/0033-5894(83)90080-7, 1983.
- 1055 Zanolini, F., Delmas, R.J. and Legrand, M.: Sulphuric and nitric acid concentrations and spikes along a 200 m deep ice core at D 57 (Terre Adélie, Antarctica). *Ann. Glaciol.*, 7, 70-75, doi: 10.3189/S0260305500005930, 1985.
- 1060 Zielinski, G.A., Mayewski, P.A., Meeker, L.D., Whitlow, S., Twickler, M.S., Morrison, M., Meese, D.A., Gow, A.J. and Alley, R.B.: Record of Volcanism Since 7000 B.C. from the GISP2 Greenland Ice Core and Implications for the Volcano-Climatic System. *Science*, 264, 948-952, doi: 10.1126/science.264.5161.948, 1994.




1065 **Figure 1: Bipolar comparison of sub-annual resolution ice-core non-sea-salt sulphur concentrations between 1440–1490 CE.** Greenland data from the NEEM-2011-S1 ice core (Sigl et al., 2013), the TUNU2013 ice core (Sigl et al., 2015) and the North Greenland Ice Core Project (NGRIP) ice core (Plummer et al., 2012) all plotted using the annually counted NS-2011 chronology (Sigl et al., 2015). Antarctic data is from the annually counted West Antarctic Ice Sheet Divide ice core (WDC) on the WD2014 age scale (Sigl et al., 2016).



1070 **Figure 2: (a) Location map of the TUNU2013 ice-core drilling site and sites outside Iceland containing tephra attributed to the**
 1075 **Veidivötn 1477 CE eruption. (b) Location map of Icelandic sites containing tephra from the Veidivötn 1477 CE eruption, from**
which geochemical characterisations were used in this study, the location of the historical report of the eruption and the 60–65 km
long fissure active during the 1477 CE eruption. The green shaded area covers the 0.5 cm isopach of the Veidivötn 1477 CE
eruption. Adapted from Larsen et al. (2014).




1080 **Figure 3:** (a) Annual resolution sulphur and particle concentration data from TUNU2013 between 1450–1500 CE. Data plotted using the NS1-2011 chronology. (b) Sub-annual resolution sulphur and particle concentration data from TUNU2013 between 1475–1480 CE. Data plotted using the NS1-2011 chronology. Red bars denote depth intervals of tephra sampling. Tephra samples: (i) QUB-1965 = 78.515–78.655 m (ii) QUB-1964 = 78.435–78.515 m (iii) QUB-1963 = 78.395–78.435 m (iv) QUB-1962 = 78.315–78.395 m. (c) Example of a brown, platy glass tephra shard identified in the TUNU2013 78.655 m deposit.




● TUNU2013 78.655 m (QUB-1965)

Icelandic volcanic systems:



Veidivötn 1477 CE occurrences:




1085 **Figure 4: Geochemical characterisation of glass tephra shards from the TUNU2013 78.655 m deposit and comparisons to Icelandic proximal material and analyses of Veiðivötn 1477 CE occurrences. (a) (i) Inset of total alkali versus silica plot showing the tholeiitic basalt composition of the TUNU2013 78.655 m deposit. Chemical classification and nomenclature after Le Maitre et al. (1989). Division line between alkaline and sub-alkaline material from MacDonald and Katsura (1964). (ii and iii) Geochemical characterisation of the TUNU2013 78.655 m deposit compared to geochemical envelopes for several Icelandic tholeiitic volcanic systems. Geochemical envelopes derived from unnormalised whole rock and glass shard analyses from deposits proximal to the Icelandic volcanic systems, adapted from Óladóttir et al. (2020) and references within. Unnormalised TUNU2013 78.655 m data utilised to permit a direct comparison with the published geochemical envelopes. (b) (i-iii) Comparisons of individual analyses from the geochemical characterisation of TUNU2013 78.655 m to characterisations of several Icelandic and non-Icelandic occurrences of the Veiðivötn 1477 CE eruption. Geochemical data for V1477 occurrences from Svartárvatn (Larsen et al., 2002), Helluvaðstjörn (Lawson et al., 2007), Skaftártunga (Streeter and Dugmore, 2014), Hvítárvatn (Larsen et al., 2011), Lake Lögurinn (Gudmundsdóttir et al., 2016), Gripdeild (Bergþórsdóttir, 2014), a compilation of six sites around the Vatnajökull ice cap (Óladóttir et al., 2011), MD99-2275 (Larsen et al., 2002; Gudmundsdóttir et al., 2011), An Loch Mór (Chambers et al., 2004) and Lake Getvaltjärnen (Davies et al., 2007). All data provided in supplementary information and plotted on a normalised anhydrous basis.**

1090


1095

1100

1105



1110 **Figure 5: High-resolution nssS and annual layer dating of the TUNU2013 ice core between 1445–1485 CE using a multiparameter approach. Concentration records are shown for nssS, Ca, Na and nssS/Na with grey dashed lines indicating the assigned years. Red shaded bar indicates the location of the V1477 tephra layer in the TUNU2013 record. Green shaded bars indicate the two 1450s CE sulphate peaks in TUNU2013.**



1115 **Figure 6: (a) Northern hemispheric (40–90°N) summer (JJA) temperature reconstruction depicting the cooling induced by selected**
15th to 17th century volcanic eruptions. The red line depicts 30-yr smoothed temperatures whereas the blue line indicates annual
temperature anomalies. The blue shaded area gives the 95% confidence interval. (b) Cooling induced by the most prominent
Icelandic eruptions of the last 1500 years in 939 CE (Eldgjá), 1477 CE (Veidivötn) and 1783 CE (Laki) in the NVOLC v2 (Guillet
et al., 2020), Sch2015 (Schneider et al., 2015) and NTREND2015 (Wilson et al., 2016) as well as in the Scandinavian cluster of
NVOLC v2; (c) Spatial patterns of Northern Hemisphere temperatures the year following the three major Icelandic eruptions of
 1120 **the last 1,500 years.**