



## Cosmogenic isotope measurements from recently deglaciated bedrock as a new tool to decipher changes in Greenland Ice Sheet size

Nicolás E. Young<sup>1</sup>, Alia J. Lesnek<sup>2</sup>, Josh K. Cuzzone<sup>3</sup>, Jason P. Briner<sup>4</sup>, Jessica A. Badgeley<sup>5</sup>, Alexandra Balter-Kennedy<sup>1</sup>, Brandon L. Graham<sup>4</sup>, Allison Cluett<sup>4</sup>, Jennifer L. Lamp<sup>1</sup>, Roseanne Schwartz<sup>1</sup>, Thibaut Tuna<sup>6</sup>, Edouard Bard<sup>6</sup>, Marc W. Caffee<sup>7,8</sup>, Susan R.H. Zimmerman<sup>9</sup>, Joerg M. Schaefer<sup>1</sup>

<sup>1</sup>Lamont-Doherty Earth Observatory, Columbia University, Palisades NY, 10964, USA

<sup>2</sup>Department of Earth Sciences, University of New Hampshire, Durham NH, 03824, USA

<sup>3</sup>Department of Earth System Science, University of California Irvine, Irvine CA, 92697, USA

<sup>4</sup>Department of Geology, University at Buffalo, Buffalo NY, 14260, USA

10 <sup>5</sup>Department of Earth and Space Sciences, University of Washington, Seattle WA, 98195, USA

<sup>6</sup>CEREGE, Aix-Marseille University, CNRS, IRD, INRAE, Collège de France, Technopôle de l'Arbois, Aix-en-Provence, France

<sup>7</sup>Department of Physics and Astronomy, PRIME Lab, Purdue University, West Lafayette IN, 47907, USA

<sup>8</sup>Department of Earth, Atmospheric, and Planetary Sciences, Purdue University, West Lafayette IN, 47907, USA

15 <sup>9</sup>Center for Accelerator Mass Spectrometry, Lawrence Livermore National Laboratory, Livermore CA, 94550, USA

*Correspondence to:* Nicolás E. Young (nicolasy@ldeo.columbia.edu)

**Abstract.** During the middle to late Holocene (8.2 ka BP to present), the Greenland Ice Sheet (GrIS) was smaller than its current configuration. Determining the exact dimensions of the Holocene ice-sheet minimum and the duration that the ice margin rested inboard of its current position remains challenging. Contemporary retreat of the GrIS from its historical maximum extent in southwestern Greenland is exposing a landscape that holds clues regarding the configuration and timing of past ice-sheet minima. To quantify the duration of the time the GrIS margin was near its modern extent we develop a new technique on Greenland that utilizes *in situ* cosmogenic <sup>10</sup>Be-<sup>14</sup>C-<sup>26</sup>Al in bedrock samples that have become ice free only in the last few decades by the retreating ice-sheet margin at Kangiata Nunaata Sermia (n=12 sites; KNS), southwest Greenland. To maximize the utility of this approach, we refine the deglaciation history of the region with stand-alone <sup>10</sup>Be measurements (n=49) and traditional <sup>14</sup>C ages from sedimentary deposits contained in proglacial-threshold lakes. We combine our reconstructed ice-margin history in the KNS region with additional geologic records from southwestern Greenland and recent model simulations of GrIS change, to constrain the timing of the GrIS minimum in southwest Greenland, the magnitude of Holocene inland GrIS retreat, and explore the regional climate history influencing Holocene ice-sheet behavior. Our <sup>10</sup>Be-<sup>14</sup>C-<sup>26</sup>Al measurements reveal that 1) KNS retreated behind its modern margin just before 10 ka, but likely stabilized near the present GrIS margin for several thousand years before retreating farther inland, and 2) pre-Holocene <sup>10</sup>Be detected in several of our sample sites is most easily explained by several thousand years of surface exposure during the Last Interglaciation. Moreover, our new results indicate that the minimum extent of the GrIS likely occurred after ~5 ka, and the GrIS margin may have approached its eventual historical maximum extent as early as ~2 ka. Recent simulations of GrIS change are able to match the geologic record of ice-sheet change in regions dominated by surface mass balance, but produce a poorer model-data fit in



35 areas influenced by oceanic and dynamic processes. Simulations that achieve the best model-data fit suggest that inland retreat  
of the ice margin driven by early to middle Holocene warmth may have been mitigated by increased precipitation. Triple  $^{10}\text{Be}$ -  
 $^{14}\text{C}$ - $^{26}\text{Al}$  measurements in recently deglaciated bedrock provide a new tool to help decipher the duration of smaller-than-present  
ice over multiple timescales. Modern retreat of the GrIS margin in southwest Greenland is revealing a bedrock landscape that  
was also exposed during the migration of the GrIS margin towards its Holocene minimum extent, but has yet to tap into a  
40 landscape that remained ice covered throughout the entire Holocene.

## 1 Introduction

The Greenland Ice Sheet (GrIS) has expanded and contracted repeatedly throughout the Quaternary. During  
45 glaciations the GrIS margin extends onto the continental shelf whereas during interglaciations, the dimensions of the GrIS are  
often similar to or smaller than today (de Vernal and Hillaire-Marcel, 2008; Hatfield et al., 2016; Knutz et al., 2019). Direct  
evidence of former GrIS maxima is found in offshore sedimentary deposits (e.g. Ó Cofaigh et al., 2013; Knutz et al., 2019),  
and the pattern of retreat from the most recent ice-sheet maximum can be reconstructed in detail through a combination of  
well-dated marine and terrestrial sedimentary archives (Bennike and Bjork, 2002; Funder et al., 2011; Kelley et al., 2013;  
50 Hogan et al., 2016; Jennings et al., 2017; Young et al., 2020a). Reconstructing the size and timing of ice-sheet minima, however,  
is extremely challenging because terrestrial evidence relating to ice-sheet minima has been overrun and destroyed by  
subsequent glacier re-expansion or resides in a largely inaccessible environment beneath modern glacier footprints. In place  
of direct terrestrial evidence, sediment-based proxy records contained in off-shore depocenters have been used to infer the  
dimensions and timing of paleo GrIS minima (Colville et al., 2011; Reyes et al., 2014; Bierman et al., 2016; Hatfield et al.,  
55 2016). These sediment-based approaches are not able to provide direct constraints on the magnitude or timing of GrIS minima,  
but they have the advantage of generally providing continuous records of inferred ice-sheet change.

Cosmogenic isotope measurements from recently deglaciated bedrock surfaces or those still residing under ice  
provide key constraints on the timing and magnitude of glacier and ice-sheet minima (e.g. Goehring et al., 2010; Schaefer et  
al., 2016; Pendleton et al., 2019). These bedrock surfaces serve as fixed benchmark locations and nuclide accumulation can  
60 only occur under extremely thin ice (e.g. *in situ*  $^{14}\text{C}$ ) or, more commonly, in the absence of ice cover when that surface is  
exposed to the atmosphere (i.e. a direct ice-margin constraint). The primary caveat of this method, however, is that measured  
nuclide inventories have non-unique solutions and only provide a measure of integrated surface exposure and burial. Moreover,  
drilling through extant glaciers and ice sheets to bedrock is logistically challenging, expensive, and can only be done after  
lengthy site consideration (e.g. Spector et al., 2018). Nonetheless, ground-breaking measurements of cosmogenic *in situ*  $^{10}\text{Be}$   
65 and  $^{26}\text{Al}$  in bedrock beneath the GISP2 borehole revealed that the GrIS likely disappeared on several occasions during the  
Pleistocene (Schaefer et al., 2016).



70 Contemporary retreat of the GrIS margin from its historical maximum extent is exposing a fresh bedrock landscape, and inventories of cosmogenic nuclides in this newly exposed bedrock can provide clues to past ice-sheet minima without having to drill through ice. Abundant geological evidence reveals that sometime during the middle Holocene, the GrIS was slightly smaller than today (e.g. Weidick et al., 1990; Long et al., 2011; Lecavalier et al., 2014; Larsen et al., 2015; Young and Briner, 2015; Lesnek et al., 2020). The mid-Holocene minimum was forced by regional temperatures that were likely as warm or warmer than today, and elucidating the behavior of the GrIS during this interval can provide key insights into GrIS behavior in a warming world. Bedrock emerging today from beneath the GrIS margin was potentially ice free during the middle Holocene, and cosmogenic nuclides in these surfaces can constrain the magnitude and duration of inland GrIS retreat.

75 Here, we present *in situ* cosmogenic  $^{10}\text{Be}$ - $^{14}\text{C}$ - $^{26}\text{Al}$  measurements from recently exposed bedrock surfaces ( $n=12$  sites) in the Kangiata Nunaata Sermia (KNS) forefield, southwestern Greenland (Fig. 1; Fig. 2). To aid interpretation of our  $^{10}\text{Be}$ - $^{14}\text{C}$ - $^{26}\text{Al}$  measurements, we refine the early Holocene deglaciation history of the landscape immediately outboard the historical GrIS maximum extent, and constrain when the GrIS retreated inboard of its present position through a combination of standalone  $^{10}\text{Be}$  measurements ( $n=49$ ) and traditional  $^{14}\text{C}$ -dated sediment sequences from proglacial-threshold lakes. We  
80 combine our new results with previously published records of deglaciation in southwestern Greenland to estimate when the GrIS was behind its present position and reached its minimum extent. We compare geologic records of ice-sheet change to recent model simulations of Holocene GrIS change to further assess the timing and magnitude of mid-Holocene GrIS retreat.

## 2 Settings and Methods

### 85 2.1 Overview

The study region is characterized by mountainous terrain dissected by a dense fjord network in which KNS resides (Fig. 1). Bedrock in the region consists primarily of Archean gneiss (Henriksen et al., 2000). Decades of research have resulted in a robust record of regional deglaciation. Minimum-limiting radiocarbon ages and  $^{10}\text{Be}$  ages reveal that initial coastal  
90 deglaciation occurred  $\sim 11.3$ - $10.7$  ka BP and the inner fjord region was ice-free by  $\sim 10.5$ - $10.0$  ka BP (Fig. 1; Tables S1 and S2). Punctuating early Holocene deglaciation was deposition of an extensive moraine system during a period locally referred to as the Kapisigdlit stade. Although no direct moraine ages exist, deposition of the Kapisigdlit stade moraines likely occurred sometime between  $\sim 10.4$  ka BP and 10.0 ka BP, based on the timing of deglaciation from regional radiocarbon and  $^{10}\text{Be}$  constraints, and a single maximum limiting radiocarbon age of  $10.17 \pm 0.34$  cal ka BP from re-worked marine sediments  
95 (Weidick et al., 2012; Larsen et al., 2014; Table S1). Following early Holocene deglaciation, retreat of the GrIS continued inboard of its current margin before re-advancing during the late Holocene. In the study area, the GrIS reached its historical maximum extent during the early to mid 18<sup>th</sup> century (Weidick et al, 2012), which is marked by a prominent moraine and trim-line (Weidick et al., 2012; Fig. 1; Fig. 2). In some locations the GrIS still resides at or near its historical maximum extent (Kelley et al., 2012), whereas in the KNS forefield, GrIS retreat from the historical maximum is slightly more pronounced and  
100 has exposed fresh bedrock surfaces.



## 2.2 Field Methods

Fieldwork was completed in 2017 and was primarily concentrated in the KNS forefield and north of KNS at Qamanaarsuup Sermia, (Fig. 1; Fig. 2; Fig. 3). In addition, we collected samples for  $^{10}\text{Be}$  dating near Narsap Sermia, located  
105 ~55 km north of KNS, and near Kangârsuup Sermia located ~35-40 km south of KNS (Lesnek et al., 2020; Fig. 1). Moraine  
crests were mapped prior to fieldwork and updated in the field. This mapping follows previous efforts (Weidick, 1974; Weidick  
et al., 2012; Pearce et al., 2018), with the exception that we distinguish between early to middle Holocene moraines and  
moraines marking the GrIS historical maximum extent (Fig. 2; Fig. 3). Across the broader KNS region, the distinction is  
obvious. Moraines and trim-lines attributed to the historical maximum extent of the GrIS are close to the modern ice margin  
110 and are fresh in appearance due to a lack of vegetation and lichen cover. Early Holocene moraines are typically located well  
outboard the historical moraines, and have extensive lichen cover. The key exceptions to this spatial relationship are regions  
where ice is topographically confined near small outlet glaciers (Fig. 1; Fig. 3). In these locations, the historical maximum and  
early Holocene moraines are closely stacked, yet are still easily distinguishable based on their morphologies and degree of  
lichen cover (Fig. 3; Fig. 4).

115 Samples for cosmogenic-nuclide analysis were collected using a Hilti brand AG500-A18 angle grinder-circular saw  
with diamond bit blades, and a hammer and chisel. Sample locations and elevations were collected with a handheld GPS device  
with a vertical uncertainty of  $\pm 5\text{m}$ , and topographic shielding was measured using a handheld clinometer. GPS units were  
calibrated to a known elevation each day, either sea level or the stated elevation of a lake derived from topographic maps.

Sediment cores from two proglacial-threshold lakes at Qamanaarsuup Sermia were collected using a Universal  
120 Percussion Corer and a Nesje-style percussion-piston coring device (Fig. 3). Goose Feather Lake (informal name) is located  
~2 km from the GrIS margin and currently receives GrIS meltwater. We collected two piston cores at a water depth of 12.6  
m (17GOOF-A3 and 17GOOF-A4; 64.45328 °N, 49.44373 °W). Marshall Lake (informal name) is located ~1 km from the  
GrIS margin and does not presently receive meltwater from the GrIS. We collected two cores from Marshall Lake with the  
Universal Percussion Corer system at a water depth of 5.95 m (17MAR-A2 and 17MAR-C1; 64.46361 °N 49.44373 °W).

125

## 2.3 $^{10}\text{Be}$ and $^{26}\text{Al}$ geochemistry and AMS measurements

We completed 61  $^{10}\text{Be}$  and 12  $^{26}\text{Al}$  measurements. Fifty-four of the  $^{10}\text{Be}$  samples and all of the  $^{26}\text{Al}$  samples were  
processed at the Lamont-Doherty Earth Observatory (LDEO) cosmogenic dating laboratory (Table S3, S4, S5). The remaining  
130 7  $^{10}\text{Be}$  samples were processed at the University at Buffalo Cosmogenic Isotope Laboratory (Tables S3 and S4). In both  
laboratories, quartz separation and Be and Al isolation followed well established protocols (Schaefer et al., 2009). We  
quantified the amount of native  $^{27}\text{Al}$  in each quartz aliquot and then added varying amounts of  $^{27}\text{Al}$  carrier to ensure that ~1400-  
1750 mg  $^{27}\text{Al}$  was achieved (Table S5). Total  $^{27}\text{Al}$  was quantified after sample digestion using inductively coupled plasma





optical emission spectrometry analysis of replicate aliquots. AMS analysis for  $^{10}\text{Be}$  samples was split between the Purdue Rare  
135 Isotope Measurement (PRIME) Laboratory (n=36) and the center for Accelerator Mass Spectrometry at Lawrence Livermore  
National Laboratory (LLNL-CAMS; n=25); all  $^{26}\text{Al}$  samples were measured at PRIME.

All  $^{10}\text{Be}$  samples were measured relative to the 07KNSTD standard with  $^{10}\text{Be}/^9\text{Be}$  ratio of  $2.85 \times 10^{-12}$  (Nishiizumi et  
al., 2007) and  $^{26}\text{Al}$  samples were measured relative to the KNSTD standard with the value of  $1.82 \times 10^{-12}$  (Nishiizumi, 2004).  
For  $^{10}\text{Be}$  samples measured at PRIME, the  $1\sigma$  analytical error ranged from 1.9% to 3.9%, with an average of  $2.7 \pm 0.5\%$  (n=36;  
140 Table S3), and the  $1\sigma$  analytical error for  $^{26}\text{Al}$  measurements ranged from 1.9% to 5.0% with an average of  $3.8 \pm 0.9\%$  (n=12;  
Table S5). For  $^{10}\text{Be}$  samples measured at LLNL-CAMS,  $1\sigma$  analytical error ranged from 1.6% to 4.2%, with an average of  
 $2.3 \pm 0.8\%$  (n=25; Table S3). Process blank corrections for all  $^{10}\text{Be}$  and  $^{26}\text{Al}$  samples were applied by taking the batch-specific  
blank value (expressed as number of atoms) and subtracting this value from the sample atom count (Table S4 and S5). In  
addition, we propagate through a 1.5% uncertainty in the carrier concentration when calculating  $^{10}\text{Be}$  concentrations. We  
145 assume half-lives of 1.387 Ma and 0.705 Ma for  $^{10}\text{Be}$  and  $^{26}\text{Al}$  (Chmeleff et al., 2010; Nishiizumi, 2004).

## 2.4 *In situ* $^{14}\text{C}$ measurements

We completed 12 *in situ*  $^{14}\text{C}$  extractions at the LDEO cosmogenic dating laboratory following well-established LDEO  
150 extraction procedures (Lamp et al, 2019; Table S6). All measured fraction modern values are converted to  $^{14}\text{C}$  concentrations  
following Hippe and Lifton (2014). The LDEO *in situ*  $^{14}\text{C}$  extraction laboratory has historically converted samples to graphite  
prior to measurement by AMS and LLNL-CAMS; however, we have recently transitioned to gas-source measurements on the  
AixMICADAS instrument at CEREGE, which can directly measure  $\sim 10\text{-}100 \mu\text{g C}$  and largely removes the need for the  
addition of a carrier gas added to typical *in situ*  $^{14}\text{C}$  samples (Bard et al., 2015; Tuna et al., 2018). Here, two samples underwent  
155 traditional graphitization and were measured at LLNL-CAMS (Table S6), whereas the remaining 10 samples were measured  
on the CEREGE AixMICADAS instrument (Table S6). Both sets of samples underwent the same *in situ*  $^{14}\text{C}$  extraction and  
 $^{14}\text{C}$  sample gas clean-up procedures, with only the samples measured at LLNL-CAMS undergoing an additional graphitization  
procedure (Lamp et al., 2019). Because we use two different measurement approaches, and our extraction efforts span the  
transition between sample graphitization and gas-source measurements, we briefly discuss our data reduction methods for both  
160 sets of measurements (Table S6).

Samples 17GRO-14 and 17GRO-74 were measured at LLNL-CAMS with  $1\sigma$  analytical uncertainties of 2.2% and  
3.0% (Table S6).  $^{14}\text{C}$  concentrations were blank-corrected using a long-term mean blank value of  $116,894 \pm 37,307$   $^{14}\text{C}$  atoms  
with the uncertainty in the blank correction propagated in quadrature (n=27; updated from Lamp et al., 2019). In addition, we  
propagate an additional 3.6% uncertainty in  $^{14}\text{C}$  concentrations based on the long-term scatter in internal graphite-based  
165 CRONUS-A standard measurements ( $698,109 \pm 25,380$  atoms/g; n=13; updated from Lamp et al., 2019); stated  $^{14}\text{C}$   
concentrations for samples measured at LLNL-CAMS have total uncertainties of 7.7% and 10.4%, respectively (Table S6).



170 Samples measured at CEREGE have  $1\sigma$  analytical uncertainties of range between 1.0% to 2.8% with a mean of  $1.4 \pm 0.5\%$  ( $n=10$ ; Table S6). For our gas-source measurements presented here, and for future lab measurements, we re-  
characterized our extraction and measurement procedure with a new set of process blank and CRONUS-A standard  
175 measurements (Table S6). We completed 6 process blank gas-source measurements at CEREGE with values ranging from  
 $\sim 73,000$ - $175,000$   $^{14}\text{C}$  atoms (Table S6). One blank measurement is anomalously high ( $174,813 \pm 3,582$  atoms; Table S6) and  
we suspect this blank was contaminated by atmosphere during collection in a breakseal. The remaining blank values have a  
mean of  $85,768 \pm 12,070$   $^{14}\text{C}$  atoms ( $n=5$ ) and we tentatively suggest that removing the graphitization procedure may also  
175 of  $81,094 \pm 6,972$  ( $n=4$ ) and  $85,768 \pm 12,070$   $^{14}\text{C}$  atoms ( $n=5$ ) to correct sample  $^{14}\text{C}$  concentrations and uncertainties in the  
blank corrections are propagated in quadrature (Table S6). In addition, we made 5 CRONUS-A standard measurements at  
CEREGE. Our CRONUS-A measurements are remarkably consistent with a mean value of  $662,132 \pm 9,849$  atoms/g ( $n=5$ ;  
1.5% uncertainty) and comparable to our graphite-based value of  $698,109 \pm 25,380$  atoms/g (updated from Lamp et al., 2019).  
Nonetheless, despite the promising consistency of our gas-source CRONUS-A measurements, we conservatively propagate  
180 through an additional 3.6% uncertainty in our sample concentrations based on the scatter in long-term LDEO CRONUS-A  
measurements. Total  $^{14}\text{C}$  concentration uncertainties for samples measured at CEREGE range from 4.3% to 5.2% with a mean  
uncertainty of  $4.6 \pm 0.2\%$  (Table S6).

## 2.5 $^{10}\text{Be}$ and *in situ* $^{14}\text{C}$ age calculations

185  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  surface exposure ages are calculated using the Baffin Bay  $^{10}\text{Be}$  production-rate calibration dataset  
(Young et al., 2013a) and the West Greenland *in situ*  $^{14}\text{C}$  production-rate calibration dataset (Young et al., 2014). All ages are  
presented using time-variant ‘Lm’ scaling (Lal, 1991; Stone, 2000), which accounts for changes in the magnetic field although  
these changes are minimal at this high latitude ( $\sim 64^\circ\text{N}$ ). The  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  calibration datasets are both derived from  
190 sites on western Greenland with early Holocene exposure histories, and the *in situ*  $^{14}\text{C}$  calibration measurements are derived  
from the same geologic samples as one of the  $^{10}\text{Be}$  calibration sites (Young et al., 2014). This combination of calibration  
datasets ensures that the production rates and the  $^{14}\text{C}/^{10}\text{Be}$  production ratio are regionally constrained. All ages are calculated  
in *MATLAB* using code from version 3 of the exposure age calculator found at <https://hess.ess.washington.edu/>, which  
implements an updated treatment of muon-based nuclide production (Balco et al., 2008; Balco, 2017). We do not correct  
195 nuclide concentrations for snow-cover or subaerial surface erosion; samples are almost exclusively from windswept locations  
and many surfaces still retain primary glacial features. In addition, we make no correction for the potential effects of isostatic  
rebound on nuclide production because both the production-rate calibration sites and sites of unknown age have experienced  
similar exposure and uplift histories (i.e. the correction is “built in”; Young et al., 2020a, 2020b). Individual  $^{10}\text{Be}$  and *in situ*  
 $^{14}\text{C}$  ages are presented and discussed with  $1\sigma$  analytical uncertainties, and moraine ages exclude the  $^{10}\text{Be}$  production-rate



200 uncertainty when we are comparing other  $^{10}\text{Be}$ -dated features. When moraine ages are compared to independent records of  
climate variability or ice-sheet change, the production-rate uncertainty is propagated through in quadrature. To allow for direct  
comparison to traditional radiocarbon constraints in the region, all  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  surface exposure ages are presented in  
thousands of years BP (CE 1950); exposure ages relative to the year of sample collection can be found in the supplemental  
information (CE 2017; Table S3 and S6).

205

## 2.6 Traditional $^{14}\text{C}$ ages from proglacial threshold lakes

Five radiocarbon ages from aquatic macrofossils were obtained from Marshall Lake (Fig. 3; Fig. 4; Table S7), and  
one radiocarbon age from an aquatic macrofossil was obtained from lake Kap01 (Fig. 1; Table S7). In addition, we discuss  
210 two previously reported radiocarbon ages from Goose Feather Lake, located adjacent to Marshall Lake (Lesnek et al., 2020;  
Table S7). Aquatic macrofossils were isolated from surrounding sediment using deionized water washes through sieves.  
Samples were freeze-dried and sent to the National Ocean Sciences Accelerator Mass Spectrometry Facility (NOSAMS) at  
Woods Hole Oceanographic Institution for age determinations. We targeted aquatic macrofossils for dating because terrestrial  
macrofossils may persist on the relatively low-energy Arctic landscape for hundreds of years before washing into a lake basin;  
215 dating terrestrial macrofossils could skew our interpretations. Hard-water effects on the  $^{14}\text{C}$  ages, which could make age  
determinations erroneously old, are unlikely in our study area because lake catchments are dominated by Archean gneiss, and  
the study lakes are all well above local marine limits. All new radiocarbon ages are calibrated using *CALIB 8.2* and the  
INTCAL20 dataset, and previously reported radiocarbon ages are re-calibrated in the same manner using the INTCAL20 and  
MARINE20 datasets (Stuiver et al., 2020; Reimer et al., 2020; Heaton et al., 2020; Table S1; Table S7).

220

## 2.7 Ice-sheet model simulations of southwestern GrIS change

We utilize recent paleo-simulations of southwestern GrIS change using the high-resolution Ice Sheet and Sea-level  
System Model (ISSM; Larour et al., 2012; Cuzzone et al., 2018, 2019; Briner et al., 2020). The model set-up has been  
225 previously described in Cuzzone et al (2019) and Briner et al (2020), but here we briefly describe model attributes. The model  
domain extends from the present-day coastline to the GrIS divide. The northern and southern boundaries of the domain are far  
to the north and south of our study area. The model resolution relies on anisotropic mesh adaptation to produce an unstructured  
mesh that varies based on bedrock topography; bedrock topography is from BedMachine v3 (Morlighem et al., 2017). For the  
southwestern GrIS, high horizontal model mesh resolution is necessary in areas of complex bed topography to prevent artificial  
230 ice-margin variability resulting from interaction with bedrock artifacts that occur at coarser resolution (Cuzzone et al., 2019).  
Thus, the model mesh varies from 20 km in areas where gradients in the bedrock topography are smooth, to 2 km in areas  
where bedrock relief is high. In the KNS region the mesh varies from 2 to 8 km.



Ice-sheet simulations are forced with paleoclimate reconstructions from Badgeley et al (2020) who used paleoclimate data assimilation to merge information from paleoclimate proxies and global climate models. The temperature reconstruction  
235 relies on oxygen-isotope records from eight ice cores, the precipitation reconstruction uses accumulation records from five ice cores, and both are guided by spatial relationships derived from the transient climate-model simulation TraCE21ka (Liu et al., 2009; He et al., 2013.) The climate reconstruction is in good agreement with published paleoclimate reconstructions (Badgeley et al., 2020 and references therein). The climate reconstruction yields 9 combinations of temperature and precipitation that are used as transient climate boundary conditions to force 9 ice model simulations.

240

### 3 Results

Adjacent to Qamanaarsuup Sermia, 27 <sup>10</sup>Be ages from moraine boulders and boulders perched on bedrock range from 20.34 ± 0.45 ka BP to 8.91 ± 0.20 ka BP (Fig. 3; Fig. 5; Table S3). Sediments in Goose Feather Lake are composed of a lower  
245 gray silt unit, overlain by organic sediments, which is in turn overlain by gray silt (Lesnek et al., 2020). A single radiocarbon age from bulk sediments at the basal sediment contact is 8,280 ± 90 cal yr BP, and a radiocarbon age from aquatic macrofossils at the upper contact between organic and minerogenic sediments is 820 ± 80 cal yr BP. Sediments in Marshall Lake display the same silt-organic-silt stratigraphy as sediments in Goose Feather Lake (Fig. 6). A radiocarbon age from aquatic macrofossils from the basal sediment contact is 8,720 ± 350 cal yr BP and a radiocarbon age from aquatic macrofossils at the  
250 uppermost contact is 520 ± 20 cal yr BP (Fig. 6). Three additional radiocarbon ages from aquatic macrofossils between the lower- and upper-most contacts are 7,250 ± 70 cal yr BP, 3,650 ± 50 cal yr BP, and 940 ± 20 cal yr BP and are in stratigraphic order (Fig. 6).

In the KNS region, 27 <sup>10</sup>Be ages from moraine boulders, erratics perched on bedrock, and abraded bedrock surfaces, range from 23.93 ± 0.53 ka BP to 5.38 ± 0.23 ka BP (Table S3), and 12 *in situ* <sup>14</sup>C ages from bedrock range from 10.11 ± 0.89  
255 ka BP to 5.62 ± 0.84 ka BP (Table S6). In addition, 12 <sup>26</sup>Al-<sup>10</sup>Be ratios range from 7.35 ± 0.33 to 6.01 ± 0.25 (all bedrock). A single radiocarbon age from aquatic macrofossils at the basal contact between silt and organic sediments in lake Kap01 is 9450 ± 440 cal yr BP (Fig. 1; Table S7).

North of Narsap Sermia near Caribou Lake (Fig. 1), three <sup>10</sup>Be ages from boulders perched on bedrock are 9.07 ± 0.32 ka BP, 8.66 ± 0.31 ka BP, and 8.66 ± 0.31 ka BP (Fig. 1; Table S3). South of KNS at Deception Lake, two <sup>10</sup>Be ages from  
260 boulders perched on bedrock are 10.66 ± 0.34 ka BP and 9.52 ± 0.32 ka BP, and near One-way lake, two <sup>10</sup>Be ages from boulders perched on bedrock are 16.43 ± 0.49 ka BP and 7.72 ± 0.26 ka BP, respectively (Fig. 1; Table S3).

## 4 Deglaciation chronologies

### 4.1 Qamanaarsuup Sermia

265



Twenty-seven  $^{10}\text{Be}$  ages from the Qamanaarsuup Sermia region range  $20.34 \pm 0.45$  ka BP to  $8.91 \pm 0.20$  ka BP (Fig. 3; Fig. 5; Table S3); however,  $^{10}\text{Be}$  ages are from three distinct morphostratigraphic units. Here, the Kapisigdlit stade is marked by numerous closely spaced moraine crests located immediately outboard of the historical moraines. Three  $^{10}\text{Be}$  ages from boulders resting on bedrock located outside of the entire Kapisigdlit moraine suite are  $10.39 \pm 0.32$  ka BP,  $10.34 \pm 0.31$  ka BP, and  $10.12 \pm 0.29$  ka BP, and have a mean age of  $10.29 \pm 0.14$  ka BP, which serves as a maximum-limiting age for the Kapisigdlit stade moraines in the region. Three  $^{10}\text{Be}$  ages from boulders resting on bedrock immediately inboard of all Kapisigdlit stade moraines, but outboard the historical moraine, are  $9.35 \pm 0.27$  ka BP,  $9.31 \pm 0.30$  ka BP, and  $9.21 \pm 0.26$  ka BP and provide a minimum-limiting age of  $9.29 \pm 0.07$  ka BP (Fig. 3). Of the 21  $^{10}\text{Be}$  ages from Kapisigdlit stade moraine boulders, five  $^{10}\text{Be}$  ages are likely influenced by  $^{10}\text{Be}$  inheritance as they are older than the maximum-limiting  $^{10}\text{Be}$  ages and similar in age to deglacial constraints found  $\sim 140$  km west at the modern coastline ( $11.67 \pm 0.32$  ka BP and  $11.01 \pm 0.41$  ka BP; Fig. 1; Fig. 3; Table S3), or date to when the GrIS margin was likely situated  $>140$  km to the west somewhere on the continental shelf ( $20.34 \pm 0.45$  ka BP,  $16.72 \pm 0.40$  ka BP, and  $14.63 \pm 0.36$  ka BP; Fig. 1; Fig. 3; Table S3). The remaining 16  $^{10}\text{Be}$  ages from the Kapisigdlit moraine set show no trend with distance from the ice margin. These  $^{10}\text{Be}$  ages overlap at  $1\sigma$  uncertainties with either each other, the minimum-limiting  $^{10}\text{Be}$  ages, or the maximum-limiting  $^{10}\text{Be}$  ages, suggesting that deposition of this suite of moraine crests happened within dating resolution (i.e. we cannot resolve the ages of different moraine crests). Combined, the 16  $^{10}\text{Be}$  ages from moraine boulders, excluding outliers, have a mean age of  $9.57 \pm 0.33$  ka BP, which is morphostratigraphically consistent with bracketing maximum- and minimum-limiting  $^{10}\text{Be}$  ages of  $10.29 \pm 0.14$  ka BP and  $9.29 \pm 0.07$  ka BP, respectively (Fig. 3). Including the uncertainty in the  $^{10}\text{Be}$  production-rate calibration,  $^{10}\text{Be}$  ages from the Qamanaarsuup Sermia region reveal that the GrIS margin approached its modern extent at  $10.29 \pm 0.23$  ka BP, deposited the Kapisigdlit stade moraines at  $9.57 \pm 0.38$  ka BP, and retreated behind the position of the historical maximum at  $9.29 \pm 0.18$  ka BP.

Alternating silt-organic-silt sediment packages found in Goose Feather and Marshall lakes are typical of those found in proglacial-threshold lakes in southwestern Greenland (Fig. 6; Briner et al., 2010; Larsen et al., 2015; Young and Briner, 2015; Lesnek et al., 2020). Silt deposition occurs when the GrIS margin resides within the lake catchment, but does not override the lake, feeding silt-laden meltwater into the lake. Organic sedimentation occurs when the GrIS margin is not within the lake catchment and meltwater is diverted elsewhere. Despite Goose Feather and Marshall lakes residing adjacent to each other on the landscape, their radiocarbon ages suggest slightly different ice-margin histories. Marshall Lake has a small and highly localized drainage catchment and does not receive GrIS meltwater at present. In contrast, because Goose Feather Lake currently receives GrIS meltwater, its drainage catchment extends somewhere beneath the modern GrIS. Goose Feather Lake is fed by meltwater sourced from an outlet glacier resting in an overdeepening, and once ice thins below the valley edge, GrIS meltwater is diverted elsewhere, likely indicating that the Goose Feather drainage divide resides near the modern ice margin (e.g. Young and Briner, 2015; Lesnek et al., 2020). Indeed, sub-ice topography in the Qamanaarsuup Sermia region reveals that the topographic threshold that dictates whether meltwater is diverted to Goose Feather Lake or elsewhere is located within  $\sim 1$  km of the modern ice margin (Fig. 6). Despite the GrIS margin retreating behind the position of the historical maximum position



300 at  $9.29 \pm 0.18$  ka BP, silt deposition in Goose Feather Lake until  $8,280 \pm 90$  cal yr BP indicates that the ice margin remained within  $\sim 1$  km of its present position between  $\sim 9.3$  and  $8.3$  ka (Fig. 6).

#### 4.2 Kangiata Nunaata Sermia

305 Fifteen  $^{10}\text{Be}$  ages in the KNS region constrain the timing of deposition of the Kapisigdlit stade moraines and the timing of when the GrIS retreated behind the eventual historical maximum limit (Fig. 2; Fig. 7; Fig. 8; Fig. 9). West of the KNS terminus, three  $^{10}\text{Be}$  ages from erratic boulders perched on bedrock located immediately outboard of the Kapisigdlit stade moraine are  $11.44 \pm 0.34$  ka BP,  $11.15 \pm 0.34$  ka BP, and  $10.23 \pm 0.42$  ka BP. Seven  $^{10}\text{Be}$  ages from moraine boulders range from  $11.22 \pm 0.35$  ka BP to  $5.38 \pm 0.23$  ka BP, and two  $^{10}\text{Be}$  ages from erratic boulders perched on bedrock located immediately  
310 inside the moraine are  $10.27 \pm 0.29$  ka BP and  $10.25 \pm 0.31$  ka BP (Fig. 7). Deglaciation of the outer coast at Nuuk occurred  $\sim 11.3$ - $10.7$  ka BP, suggesting that the single  $^{10}\text{Be}$  age of  $10.23 \pm 0.42$  ka BP from just outboard of the Kapisigdlit stade moraine is likely the closest maximum-limiting age on moraine deposition, and  $^{10}\text{Be}$  ages of  $11.44 \pm 0.34$  ka BP and  $11.15 \pm 0.34$  ka BP are likely influenced by a slight amount of inheritance (Fig. 1; Fig. 7). This maximum-limiting  $^{10}\text{Be}$  age is consistent with a maximum-limiting radiocarbon age of  $10,170 \pm 340$  cal yr BP from a bivalve re-worked into Kapisigdlit stade till located  
315 down-fjord (Weidick et al., 2012; Larsen et al., 2014; Fig. 1; Table S1). The  $10.23$  ka BP age is also consistent with the  $^{10}\text{Be}$  age of boulders outboard of the Kapisigdlit stade moraines in the Qamanaarsuup Sermia study area of  $\sim 10.29$  ka BP. There is significant scatter in our moraine boulder  $^{10}\text{Be}$  ages; the oldest  $^{10}\text{Be}$  age of  $11.22 \pm 0.35$  ka BP is likely influenced by isotopic inheritance whereas the younger outliers of  $6.73 \pm 0.18$  ka BP and  $5.38 \pm 0.23$  ka BP reflect post-depositional boulder exhumation (Fig. 7). The remaining  $^{10}\text{Be}$  ages from the Kapisigdlit stade moraine have a mean age of  $10.24 \pm 0.31$  ka BP  
320 ( $n=4$ ), which is supported by our minimum-limiting  $^{10}\text{Be}$  ages of  $10.27 \pm 0.29$  ka BP and  $10.25 \pm 0.31$  ka BP from immediately inside the moraine (Fig. 7). Indeed,  $^{10}\text{Be}$  ages from erratic boulders perched on bedrock located immediately inside moraines across southwestern Greenland typically provide constraints that are near-identical to tightly clustered  $^{10}\text{Be}$  ages from moraine boulders (e.g. Young et al., 2011a; Young et al., 2013b; Lesnek and Briner, 2018). Including the production uncertainty, we directly date the Kapisigdlit stade moraine to  $10.24 \pm 0.36$  ka BP, and all available supporting  $^{10}\text{Be}$  and  $^{14}\text{C}$  ages further  
325 constrain moraine deposition to  $\sim 10.4$ - $10.2$  ka BP.

On the east side of KNS, three  $^{10}\text{Be}$  ages from immediately outboard the historical maximum are  $12.86 \pm 0.57$  ka BP,  $10.28 \pm 0.24$  ka BP, and  $10.00 \pm 0.24$  ka BP (Fig. 9). The  $^{10}\text{Be}$  age of  $12.86 \pm 0.57$  ka BP is, again, almost certainly influenced by inheritance as this  $^{10}\text{Be}$  age pre-dates the timing of deglaciation at the outer coastline. The remaining  $^{10}\text{Be}$  ages of  $10.28 \pm 0.24$  ka BP and  $10.00 \pm 0.24$  ka BP are consistent with the  $^{10}\text{Be}$  ages from inside the Kapisigdlit stade moraine (and outboard  
330 of the historical maximum limit) on the west side of the fjord. Our new and previously published age constraints reveal that deposition of the Kapisigdlit stade moraine in the KNS forefield occurred ca.  $10.4$ - $10.2$  ka, followed by retreat of the GrIS within the historical maximum limit shortly thereafter (Fig. 2). Any possible moraine correlative with the  $\sim 9.6$  ka moraine found at Qamanaarsuup Sermia would have been overrun by the historical advance of KNS.





335 Lastly, a basal minimum-limiting radiocarbon age of  $9450 \pm 440$  cal yr BP from Kap01 is, within uncertainties,  
identical to a previously reported basal radiocarbon age of  $9850 \pm 290$  cal yr BP from the same lake (Table S1; Table S7;  
Larsen et al., 2014). We note that our new radiocarbon age is from aquatic macrofossils whereas the previously published  
radiocarbon age is from bulk sediments. Despite the risk of bulk sediments yielding radiocarbon ages that are too old, basal  
radiocarbon ages from Kap01 suggest that the offset between macrofossil- and bulk sediment-based radiocarbon ages is  
minimal during the initial onset of organic sedimentation following landscape deglaciation.

340

#### 4.3 Auxiliary sites

At our site near Narsap Sermia, located  $\sim 55$  km north of KNS, three  $^{10}\text{Be}$  ages from boulders perched on bedrock  
located outboard of the GrIS historical maximum and inboard of the Kapisigdlit stade limit have a mean age of  $8.80 \pm 0.24$   
345 ka BP ( $8.80 \pm 0.29$  ka BP including the production rate uncertainty; Table S3) and is consistent with a minimum-limiting basal  
radiocarbon age of  $7460 \pm 110$  cal yr BP (Lesnek et al., 2020; Fig. 1; Table S1). Near Kangâsarsuup Sermia, located  $\sim 35$  km  
south of KNS, two  $^{10}\text{Be}$  ages from boulders perched on bedrock outboard of the historical limit and inboard of the Kapisigdlit  
stade limit are  $10.66 \pm 0.34$  ka BP and  $9.52 \pm 0.32$  ka BP. An existing  $^{10}\text{Be}$  and traditional  $^{14}\text{C}$  age of  $9.95 \pm 0.19$  ka BP ( $n=2$ )  
and  $8790 \pm 190$  cal yr BP from locations slightly more distal from the ice sheet suggest that our age of  $10.66 \pm 0.34$  ka BP is  
350 perhaps influenced by a small amount of isotopic inheritance (Larsen et al., 2014; Fig. 1; Table S2; Table S3). The remaining  
age of  $9.52 \pm 0.32$  ka BP is consistent with a minimum-limiting basal radiocarbon age of  $9210 \pm 190$  cal yr BP from Deception  
Lake (Fig. 1; Lesnek et al., 2020). Lastly, south of Kangaasarsuup Sermia near One-way Lake, two  $^{10}\text{Be}$  ages from boulders  
perched on bedrock outboard of the historical limit are  $16.43 \pm 0.49$  ka BP, and  $7.72 \pm 0.26$  ka BP (Fig. 1; Table S3). The older  
of these two  $^{10}\text{Be}$  ages is influenced by isotopic inheritance leaving a single  $^{10}\text{Be}$  age of  $7.72 \pm 0.26$  ka BP as the best estimate  
355 for the timing of local deglaciation.

#### 5 $^{10}\text{Be}$ - $^{14}\text{C}$ - $^{26}\text{Al}$ measurements from the KNS forefield

Prior to interpreting triple  $^{10}\text{Be}$ - $^{14}\text{C}$ - $^{26}\text{Al}$  measurements in abraded bedrock located between the historical maximum  
360 extent and the current ice margin, we use our new chronology of early Holocene ice-margin change and historical observations  
to quantify the maximum duration of Holocene exposure our bedrock samples sites could have experienced. First, we use  $^{10}\text{Be}$   
ages from immediately outboard of the historical limit on the northeastern side of KNS and  $^{10}\text{Be}$  ages from just inboard of the  
Kapisigdlit stade moraine on the southwestern side of KNS to define the potential onset of Holocene exposure at our  $^{10}\text{Be}$ - $^{14}\text{C}$ -  
 $^{26}\text{Al}$  bedrock sites.  $^{10}\text{Be}$  ages from outboard of the historical limit overlap at  $1\sigma$  uncertainties ( $n=4$ ; excluding one outlier), and  
365 we calculate a mean age of  $10.20 \pm 0.14$  ka BP ( $10.27 \pm 0.23$  ka BP with production-rate uncertainty) as the earliest onset of  
exposure at our inboard bedrock sites (Fig. 2). This age represents the timing of deglaciation immediately outboard of the  
historical limit and, assuming the continued retreat of the GrIS margin, the initial timing of exposure for the inboard bedrock



370 sites (e.g. Young et al., 2016). Next, we capitalize on historical observations in the KNS region that constrain ice-margin  
change beginning in the 18<sup>th</sup> century (Weidick et al., 2012; Lea et al., 2014 and references therein). Based on scattered first-  
person observations, the advance towards the eventual historical maximum extent likely began by CE 1723-1729 and  
culminated ca. CE 1750, with initial ice-margin thinning taking place ca. CE 1750-1800 (Weidick et al., 2012). Broadly  
supporting this record of ice-margin migration is an early photograph by Danish geologist Hinrich Rink dated to sometime in  
the CE 1850s. The photograph depicts the front of KNS as seen from the south and clearly delineates an existing historical  
maximum trim-line indicating the local GrIS historical maximum was achieved and initial thinning from this maximum began  
375 prior to CE 1850 (pg. 25, Weidick et al., 2012). Additional first-person descriptions indicate that the KNS ice margin was more  
extended than today between ca. CE 1850 and at least CE 1948, punctuated by the CE 1920 stade, which marks a significant  
readvance of the ice margin (Fig. 2). Aerial photographs reveal that the ice margin was only a few 10s of m east of our bedrock  
sites on the western side of KNS at CE 1968, suggesting site deglaciation shortly beforehand (Weidick et al., 2012). Our  
eastern ice-marginal bedrock sites were likely ice-free ca. CE 2000 based on satellite imagery.

380 The majority of our bedrock sites are directly adjacent to the ice margin and we assume that pre-imagery historical  
observations of ice-margin change apply to both sampling regions because any differences in ice-covered and ice-free intervals  
between the two sampling sites are likely negligible for our purposes. The available historical constraints indicate that our  
bedrock sites became ice-covered ca. CE 1725 (historical maximum advance phase) and our sites on the western side of KNS  
likely became ice free ca. CE 1968; ice marginal sites on the east side of KNS became ice-free ca. CE 2000. These observations  
385 indicate that the western bedrock sites experienced 243 years of historical ice cover, whereas the eastern sites experienced 275  
years of ice cover. With the earliest possible onset of exposure occurring at  $10.20 \pm 0.23$  ka BP as constrained by our <sup>10</sup>Be  
ages, we assume that the maximum duration of Holocene surface exposure at all of our sites is 10.0 kyr. We do note, however,  
that three of our bedrock sampling sites on the northeastern side of KNS are at a higher elevation and lie closer to the historical  
maximum limit than the sites adjacent to the modern ice margin (Fig. 2; Fig. 9). These high-elevation sites almost certainly  
390 experienced shorter historical ice cover than our lower elevation ice-marginal sites, likely on the order of ~150-200 years. But  
considering the uncertainties in our chronology and analytical detection limits, we assume they have the same maximum  
Holocene exposure duration of 10 kyr. Lastly, samples could inherit cosmogenic nuclides from an earlier exposure (i.e.  
inheritance), and in the strictest sense, the most recent period of exposure for our bedrock sites equates to only the last few  
decades. Because of the well-constrained maximum possible exposure duration provided by geologic constraints and historical  
395 observations, here isotopic inheritance refers to exposure ages older than  $10.20 \pm 0.23$  ka BP (i.e. pre-Holocene exposure).

### 5.1 Apparent *in situ* <sup>10</sup>Be and <sup>14</sup>C surface exposure ages

400 Apparent <sup>10</sup>Be ages from abraded bedrock surfaces on the northeastern side of KNS, listed from just inboard of the  
historical maximum limit towards the modern ice margin, are:  $17.46 \pm 0.47$ ,  $17.15 \pm 0.44$ ,  $12.95 \pm 0.31$ ,  $9.96 \pm 0.24$ ,  $19.83 \pm$   
 $0.40$ ,  $23.93 \pm 0.53$ ,  $15.16 \pm 0.40$ , and  $6.83 \pm 0.16$  ka BP (Fig. 2; Fig. 9; Fig. 10; Fig. 11; Table S3). On the southwestern side



of KNS adjacent to the modern ice margin,  $^{10}\text{Be}$  ages from abraded bedrock surfaces are  $6.94 \pm 0.16$ ,  $6.75 \pm 0.16$ ,  $6.56 \pm 0.27$ , and  $6.51 \pm 0.14$  ka BP, all roughly equal distance from the present ice margin (Fig. 2; Fig. 12; Table S3). Along our northeastern transect, 6 of the 8 apparent  $^{10}\text{Be}$  ages exceed the maximum allowable Holocene exposure duration ( $10.20 \pm 0.23$  ka). Apparent  $^{10}\text{Be}$  ages greater than this indicate the presence of inherited  $^{10}\text{Be}$  accumulated from a period of pre-Holocene exposure and insufficient sub-glacial erosion during the last glacial cycle to reset the cosmogenic clock. Of the two remaining  $^{10}\text{Be}$  ages not influenced by isotopic inheritance, the  $^{10}\text{Be}$  age of  $9.96 \pm 0.24$  ka BP is statistically identical to the maximum allowable duration of Holocene exposure for the landscape located between the historical moraine and the modern ice margin. In addition, a  $^{10}\text{Be}$  age of  $6.83 \pm 0.16$  ka directly adjacent to the modern margin is suggestive of less exposure and more burial (or more erosion; see Section 5.2) at this site and is also statistically identical to apparent  $^{10}\text{Be}$  ages from the southwestern side of KNS, indicating similar Holocene exposure histories.

Although the maximum amount of allowable Holocene exposure is well-constrained, we pair our  $^{10}\text{Be}$  measurements with *in situ*  $^{14}\text{C}$  measurements to 1) further assess the magnitude of isotopic inheritance in our bedrock samples, and 2) constrain post-10 ka fluctuations of the GrIS margin. Whereas long-lived nuclides such as  $^{10}\text{Be}$  must be removed from the landscape via sufficient subglacial erosion, the relatively short half-life of *in situ*  $^{14}\text{C}$  ( $t_{1/2} = 5700$  years) allows for previously accumulated *in situ*  $^{14}\text{C}$  to decay to undetectable levels after  $\sim 30$  ka of simple burial of a surface by ice; with the aid of subglacial erosion, *in situ*  $^{14}\text{C}$  can reach undetectable levels more quickly. In contrast to our apparent  $^{10}\text{Be}$  ages, which display varying degrees of Holocene exposure and isotopic inheritance, *in situ*  $^{14}\text{C}$  measurements are consistent with Holocene-only exposure histories. On the southwestern side of KNS, four apparent *in situ*  $^{14}\text{C}$  ages range from  $6.59 \pm 0.47$  ka BP to  $5.62 \pm 0.84$  ka BP and paired  $^{10}\text{Be}$ - $^{14}\text{C}$  measurements yield concordant exposure ages (Fig. 12). Within our northeastern bedrock transect, the two highest elevation samples near the historical maximum limit with  $^{10}\text{Be}$  ages  $17.46 \pm 0.47$  ka BP and  $17.15 \pm 0.44$  ka BP have significantly younger *in situ*  $^{14}\text{C}$  ages of  $10.11 \pm 0.89$  ka and  $9.46 \pm 0.78$  ka BP, respectively (Fig. 2; Fig. 9). The next sample along this transect has a  $^{10}\text{Be}$  age of  $12.95 \pm 0.31$  ka BP and an *in situ*  $^{14}\text{C}$  age of  $9.70 \pm 0.80$  ka BP, followed by a sample with concordant  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages of  $9.96 \pm 0.24$  ka BP and  $9.80 \pm 0.98$  ka, respectively (Fig. 2; Fig. 9). Within our cluster of samples closest to the ice margin along the northeastern transect, the one apparent  $^{10}\text{Be}$  age of  $6.83 \pm 0.16$  ka BP is matched by a statistically identical *in situ*  $^{14}\text{C}$  age of  $6.99 \pm 0.50$  ka BP, indicating that this  $^{10}\text{Be}$  age is likely not influenced by isotopic inheritance (Fig. 2; Fig. 9). The remaining three samples in this cluster with  $^{10}\text{Be}$  ages of  $23.93 \pm 0.53$ ,  $19.83 \pm 0.40$ , and  $15.16 \pm 0.40$  ka BP have significantly younger *in situ*  $^{14}\text{C}$  ages of  $6.68 \pm 0.48$ ,  $6.29 \pm 0.73$ , and  $6.49 \pm 0.46$  ka BP, respectively (Fig. 2; Fig. 9).

There are two modes of apparent *in situ*  $^{14}\text{C}$  exposure ages: a cluster of *in situ*  $^{14}\text{C}$  ages at  $\sim 10$  ka BP, and a second cluster at  $\sim 6$ -7 ka BP (Fig. 2; Fig. 13). Perhaps more importantly, however, is that these two modes correlate with two distinct morphostratigraphic surfaces. *In situ*  $^{14}\text{C}$  ages between  $\sim 6$ -7 ka BP are all from sites located directly adjacent to the modern ice margin. On the southwestern side of KNS, *in situ*  $^{14}\text{C}$  ages of  $\sim 6$ -7 ka BP are matched by concordant  $^{10}\text{Be}$  ages (Fig. 2; Fig. 13). On the northeastern side of KNS, *in situ*  $^{14}\text{C}$  ages of  $\sim 6$ -7 ka BP are found closest to the modern ice margin. One of these sites has concordant  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages, while at the remaining sites, *in situ*  $^{14}\text{C}$  ages of 6-7 ka BP are



440 significantly younger than their paired  $^{10}\text{Be}$  ages (Fig. 2; Fig. 13). *In situ*  $^{14}\text{C}$  ages of  $\sim 10$  ka BP only exist at our high-  
elevation sites more proximal to the historical maximum limit. Moreover, *in situ*  $^{14}\text{C}$  ages that range between  $10.11 \pm 0.89$  ka  
BP and  $9.46 \pm 0.78$  ka BP at our high-elevation sites are statically indistinguishable from the maximum duration of Holocene  
exposure these sites could have experienced ( $\sim 10$  kyr; Fig. 2; Fig. 13). Thus, the *in situ*  $^{14}\text{C}$  ages of  $\sim 10$  ka BP, including the  
445 paired  $^{10}\text{Be}$ - $^{14}\text{C}$  ages of  $\sim 10$  ka BP, indicate that following deglaciation of the landscape just outboard the historical moraine  
at  $10.20 \pm 0.23$  ka BP, the GrIS margin continued to retreat inland and expose our high-elevation bedrock sites immediately  
thereafter. Moreover, subglacial erosion during the brief period of historical ice cover was negligible at these high-elevation  
sites because any significant subglacial erosion would result in apparent *in situ*  $^{14}\text{C}$  ages that are younger than the maximum  
Holocene exposure history these sites could have experienced. Our younger  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages of 6-7 ka BP, on the  
445 other hand, reflect some combination of less Holocene exposure and/or more subglacial erosion than the high-elevation  
sites.

## 5.2 $^{14}\text{C}/^{10}\text{Be}$ ratios, transient burial, and subglacial erosion

450 Combining two or more nuclides with different half-lives quantifies the integrated amount of surface exposure and  
burial (e.g. Bierman et al., 1999; Goehring et al., 2010). Following a period of surface exposure, burial of a surface by  
overriding ice will cease nuclide production and lead to the faster decay of the short-lived nuclide relative to the nuclide with  
a longer (or stable) half-life. Typically, the longer-lived nuclide is used to constrain the total amount of surface exposure,  
whereas the nuclide with a shorter half-life functions as the burial chronometer. Because the production ratio of two nuclides  
455 of a constantly exposed surface is known, a measured sample ratio below the constant-production value represents the duration  
of surface burial. Over the Holocene timescale considered here,  $^{10}\text{Be}$  functions as an essentially stable nuclide because of its  
long half-life ( $t_{1/2}=1.387$  Ma), whereas  $^{14}\text{C}$ , with its short half-life ( $t_{1/2}=5700$  yrs), acts as the burial chronometer (e.g. Goehring  
et al., 2010).

460 We only consider measured  $^{14}\text{C}/^{10}\text{Be}$  ratios in samples that do not have inherited  $^{10}\text{Be}$ ; inherited  $^{10}\text{Be}$  results in  
physically unobtainable ratios. Using the average precision of our measured  $^{14}\text{C}/^{10}\text{Be}$  ratios (5.7%), we estimate a minimum  
burial detection limit resolvable to 625 years at  $1\sigma$  (Table S8). All six samples without  $^{10}\text{Be}$  inheritance have  $^{14}\text{C}/^{10}\text{Be}$  ratios  
that are indistinguishable from constant exposure (Fig. 13; Table S8). One of these pairings with concordant  $^{14}\text{C}$ - $^{10}\text{Be}$  ages of  
 $\sim 10$  ka BP on the northeastern side of KNS indicates that the high-elevation bedrock sites likely became re-occupied by the  
GrIS only once as the ice sheet advanced towards the historical maximum extent,  $<625$  years ago (Fig. 2; Fig. 13). The  
465 remaining samples with  $^{14}\text{C}$ - $^{10}\text{Be}$  ages of  $\sim 6$ -7 ka BP, also reveal  $<625$  years of ice burial (Fig. 13). Our measured  $^{14}\text{C}/^{10}\text{Be}$   
ratios are consistent with the observation of  $\sim 245$ -275 years of recent historical ice cover as being the only period of ice cover  
these sites experienced after early Holocene deglaciation. Our measured ratios, however, cannot rule out brief pre-18<sup>th</sup> century  
advances of the GrIS that may have covered our ice-marginal bedrock sites. Because our ice marginal sites reside so close to



470 the modern ice margin, it is likely that any brief advance of KNS prior to the 18<sup>th</sup> century would have covered our sampled bedrock locations.

475 Our measured  $^{14}\text{C}$ - $^{10}\text{Be}$  ratios do not indicate any significant amounts of burial at our ice-marginal sites, but concordant  $^{14}\text{C}$ - $^{10}\text{Be}$  ages of  $\sim 6$ -7 ka BP suggest an exposure history and/or degree of subglacial erosion fundamentally different than the high-elevation 10 ka BP landscape (Fig. 2; Fig. 13). The simplest interpretation is that the concordant  $^{14}\text{C}$ - $^{10}\text{Be}$  ages and constant production ratios reflect one period of surface exposure over the last  $\sim 6$ -7 ka BP, prior to the period of  
480 historical ice cover. In this scenario, deglaciation of the high-elevation bedrock sites occurred at  $10.20 \pm 0.23$  ka BP, but instead of continued inland retreat of the GrIS margin, the ice margin stabilized for several thousand years, and then retreated inboard of today's margin around 6-7 ka BP. However, another possibility is that these sites also deglaciated at  $10.20 \pm 0.23$  ka BP or later, but later experienced significant subglacial erosion during the period of historical ice cover. Subglacial erosion through the production-depth profile in bedrock would result in younger apparent exposure ages (e.g. Goehring et al., 2010; Young et al., 2016) while maintaining a  $^{14}\text{C}/^{10}\text{Be}$  ratio consistent with constant exposure as long as erosion was limited to the first few  
485 10s of cm where spallation dominates nuclide production.

To explore the possibility that concordant  $^{10}\text{Be}$  and  $^{14}\text{C}$  ages of  $\sim 6$ -7 ka BP are a product of significant subglacial erosion, we cast apparent  $^{10}\text{Be}$  and  $^{14}\text{C}$  ages as a function of total erosional depth into bedrock during the period of historical ice cover assuming varying lengths of total surface exposure (Fig. 14; Table S9). Because of the significantly larger  
490 uncertainties in the *in situ*  $^{14}\text{C}$  measurements, we only use the  $^{14}\text{C}$ -based erosion depths as a check on the  $^{10}\text{Be}$ -based erosion depths. The more precise  $^{10}\text{Be}$  measurements result in more precise estimates of erosional depth, and because we made paired  $^{10}\text{Be}$ - $^{14}\text{C}$  measurements (versus single nuclide measurements in separate geological samples at different locations), our  $^{10}\text{Be}$  measurements provide more robust constraints of simulated erosional depth. As an estimate of the maximum amount of total erosion our sample sites could have experienced, we assume that our ice marginal sites experienced 10 kyr of total exposure,  
495 which is constrained by the timing of deglaciation from just outboard of the historical moraines and the estimated duration of historical ice cover (section 5). We then project the  $^{10}\text{Be}$  and  $^{14}\text{C}$  production-depth profiles in bedrock at each site and determine the depth below the theoretical 10 kyr surface to which our measured  $^{10}\text{Be}$  and  $^{14}\text{C}$  concentrations equate (Fig. 14; Table S9). Assuming a total exposure duration of 10 kyr,  $^{10}\text{Be}$ - and  $^{14}\text{C}$ -based total erosional depths are  $24.8 \pm 1.8$  cm and  $20.1 \pm 4.1$  cm, respectively, during the period of historical ice cover (Fig. 14; Table S9). Using the known duration of historical ice cover  
495 (section 5), these erosional depths translate to abrasion rates of  $1.00 \pm 0.11$  mm/yr ( $^{10}\text{Be}$ ) and  $0.81 \pm 0.19$  mm/yr ( $^{14}\text{C}$ ). Using total exposure durations of 9 kyr, 8 kyr, and 7.5 kyr yields  $^{10}\text{Be}$ -based erosion depths of  $18.3 \pm 1.8$  cm,  $11.1 \pm 1.8$  cm, and  $7.2 \pm 1.8$  cm, respectively, which equate to abrasion rates of  $0.74 \pm 0.10$  mm/yr,  $0.45 \pm 0.08$  mm/yr, and  $0.29 \pm 0.08$  mm/yr (Fig. 14; Table S9).

500 Calculated abrasion rates are all well above the canonical polar subglacial abrasion rate of 0.01 mm/yr (Hallet et al., 1996), but notably, similar to abrasion rates inferred at the Jakobshavn Isbræ forefield constrained by similar methodology ( $0.72 \pm 0.26$  mm/yr; Young et al., 2016). The inferred abrasion rates at KNS are consistent with the abrasion rates at Jakobshavn Isbræ, but there are key differences between the two landscapes which aid in further limiting the plausible abrasion rates in the



505 KNS forefield. Dozens of  $^{10}\text{Be}$  ages from the Jakobshavn Isbræ region, and most notably from bedrock inside and outside the  
Jakobshavn Isbrae historical maximum limit, reveal a landscape entirely devoid of isotopic inheritance suggestive of a highly  
erosive environment (Young et al., 2013b; Young et al., 2016). In contrast, new  $^{10}\text{Be}$  measurements presented here influenced  
by isotopic inheritance, including from bedrock, coupled with previous  $^{10}\text{Be}$  measurements from the region (e.g. Larsen et al.,  
2014), at least qualitatively point to a less erosive GrIS in the KNS region. We think it is unlikely that subglacial abrasion rates  
at KNS can match or exceed those from the Jakobshavn Isbræ forefield and therefore we favor an interpretation that couples  
less site exposure over significant amounts of subglacial abrasion. Moreover, the *in situ*  $^{14}\text{C}$  ages (and one  $^{10}\text{Be}$  age) from our  
510 high-elevation bedrock sites on the northeastern side of KNS are indistinguishable from the timing of early Holocene  
deglaciation and indicate that at least at these high-elevation sites, subglacial abrasion during historical ice cover was  
negligible.

The most straightforward process to generate statistically identical  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages across all of the ice-  
marginal bedrock sites is for these sites to have undergone relatively little to no subglacial abrasion following the period of  
515 early to middle Holocene exposure. We also doubt that significant subglacial abrasion in the KNS forefield during the period  
of historical ice cover would result in such strikingly uniform apparent exposure ages (i.e. uniform abrasion rates). On the  
other hand, it is unrealistic to assume that our ice-marginal sites did not experience some degree of abrasion because this is  
not a cold-based ice environment and striations were routinely observed at sampling locations; these features were likely  
formed during the period of historical ice cover. To estimate the likely timing of deglaciation at our ice marginal sites that  
520 have concordant  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages of 6-7 ka BP, we present a range of deglaciation estimates. To constrain the latest  
possible age of deglaciation, we use all of the bedrock  $^{10}\text{Be}$  ages on the southwestern side of KNS (n=4), combined with the  
apparent *in situ*  $^{14}\text{C}$  ages on the northeastern side of KNS that have paired  $^{10}\text{Be}$  measurements influenced by inheritance  
(n=3), and a single  $^{10}\text{Be}$  age that is not influenced by isotopic inheritance. The mean age of these samples is  $6.63 \pm 0.21$  ka  
BP, which does not include the last ~275 years of historical ice cover when no, or minimal (i.e.  $^{14}\text{C}$ ) isotope production  
525 would have occurred, and assumes zero erosion during historical ice cover. Accounting for the period of historical ice cover,  
the timing of middle Holocene deglaciation from our ice-marginal bedrock sites is  $6.91 \pm 0.21$  ka BP. As an upper bound on  
the timing of deglaciation, we rely on our modeled scenario of 7.5 kyr of exposure resulting in an abrasion rate of  $0.29 \pm$   
 $0.08$  mm/yr. As with our measured apparent  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages, the modeled 7.5 kyr scenario does not include the  
~275 years of historical ice cover. Including historical ice cover results in a deglaciation age of 7.78 ka BP. Combined, our  
530 favored interpretation is that the ice-marginal bedrock sites likely first became ice-free sometime between ~7.8 and ~6.9 ka  
BP and experienced abrasion during the period of historical ice cover at no more than ~0.3 mm/yr. We suggest that this  
estimate sufficiently accounts for some degree of subglacial abrasion during the period of historical ice cover as suggested  
by field observations, while also acknowledging that tightly clustered apparent  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages are suggestive of  
bedrock surfaces that have undergone minimal modification following initial mid-Holocene exposure.

535

### 5.3 $^{26}\text{Al}/^{10}\text{Be}$ ratios





As with  $^{14}\text{C}/^{10}\text{Be}$  ratios,  $^{26}\text{Al}/^{10}\text{Be}$  ratios can be used to measure integrated surface exposure and burial, albeit over much longer timescales. The  $^{26}\text{Al}$ - $^{10}\text{Be}$  pairing utilizes preferential decay of  $^{26}\text{Al}$  ( $t_{1/2}=0.705$  Ma) relative  $^{10}\text{Be}$  ( $t_{1/2}=1.387$  Ma) to quantify exposure and burial over glacial-interglacial timescales or longer (e.g. Bierman et al., 1999; Fabel et al., 2002; 540 Gjermundsen et al., 2015). The canonical  $^{26}\text{Al}/^{10}\text{Be}$  production ratio is considered 6.75 (Balco and Rovey, 2008; Balco et al., 2008), but measurements from western Greenland constrain the  $^{26}\text{Al}/^{10}\text{Be}$  production ratio to  $7.3 \pm 0.3$  suggesting the production ratio scales with latitude and elevation (Corbett et al., 2017). Modelling suggests that the  $^{26}\text{Al}/^{10}\text{Be}$  production ratio at sea level/high latitude is  $\sim 7.0$ - $7.1$  (Argento et al., 2013).

Our  $^{26}\text{Al}/^{10}\text{Be}$  ratios range from  $7.39 \pm 0.33$  to  $6.01 \pm 0.25$  in the KNS forefield ( $n=12$ ; Table S5). Because of the 545 extremely close proximity of all of our bedrock samples, these surfaces must have the same exposure and burial histories over glacial-interglacial timescales; the value of  $6.01 \pm 0.25$  is anomalously low relative to our remaining measurements. After removing the lowest ratio, remaining ratios range from  $7.39 \pm 0.33$  to  $6.71 \pm 0.24$  with a mean of  $7.05 \pm 0.24$  ( $n=11$ ; Table S5). Further limiting this dataset to samples that have no detectable inherited  $^{10}\text{Be}$ , results in a mean value of  $7.17 \pm 0.20$  ( $n=6$ ; Table S5). With the exception of our lowest measured ratio ( $6.01 \pm 0.25$ ), each of our  $^{26}\text{Al}/^{10}\text{Be}$  ratios overlaps with the constant 550 production values at  $2\sigma$  uncertainty regardless of which constant production is used. Yet, our  $^{26}\text{Al}/^{10}\text{Be}$  ratios are systematically greater than 6.75, and suggest that the true production ratio is  $>6.75$  and more consistent recent modeled and empirical estimates (Table S5; Argento et al., 2013; Corbett et al., 2017).

The burial detection limit for the paired  $^{26}\text{Al}$ - $^{10}\text{Be}$  method is typically on the order of  $\sim 1$  glacial cycle, although this is dependent on the uncertainty in the measured  $^{26}\text{Al}/^{10}\text{Be}$  ratio. All of our measured  $^{26}\text{Al}/^{10}\text{Be}$  ratios suggest constant exposure, 555 including samples with inherited  $^{10}\text{Be}$  (apparent  $^{10}\text{Be}$  ages  $>10$  ka BP), yet the known ice-margin history requires that constant exposure could have only occurred over the last  $\sim 10$  kyr (Fig. 15; excluding the brief period of historical ice cover that is undetectable with the  $^{26}\text{Al}$ - $^{10}\text{Be}$  chronometer). The most likely source of the excess  $^{10}\text{Be}$  in samples with apparent  $^{10}\text{Be}$  ages  $>10$  ka BP is surface exposure during Marine Isotope Stage 5e (MIS;  $\sim 129$ - $116$  ka BP; Stirling et al., 1998). Brief exposure of our bedrock surfaces during MIS 5e, surface burial between MIS 5d and  $\sim 10$  ka BP, followed by re-exposure for the last 10 560 kyr is the most straightforward scenario to have measured  $^{26}\text{Al}/^{10}\text{Be}$  ratios consistent with constant exposure while also containing a slight amount of inheritance. Moreover, this scenario is consistent with the broad outline of GrIS change over the last glacial cycle. Greenland ice core data and offshore sediment records reveal that the GrIS was smaller than today during MIS 5e (Colville et al., 2011; NEEM, 2013), which suggests that bedrock currently emerging from beneath the GrIS was likely exposed for some period of time during MIS 5e.

At the same time, we cannot rule out that small amounts of inherited  $^{10}\text{Be}$  coupled with  $^{26}\text{Al}/^{10}\text{Be}$  ratios consistent with constant exposure, is a result of exposure during MIS 3. Terrestrial evidence of a restricted GrIS during MIS 3 is limited so far to select sites in northern Greenland (e.g. Larsen et al., 2018). Marine-based sediment records from off southwestern Greenland, however, point to significant MIS 3 recession of the GrIS, although the MIS 3 ice margin may have still been located off the modern coastline and thus our sample sites would remain ice covered (Seidenkrantz et al., 2019). In addition, 570 any significant exposure of our KNS bedrock sites during MIS 3, would require significant amounts of burial in order to have



any accumulated *in situ*  $^{14}\text{C}$  decay away to undetectable levels as there is no evidence of inherited *in situ*  $^{14}\text{C}$  in our samples. Moreover, eustatic sea-level curves indicate that sea level was at least 30-40 m below present during MIS 3 (Siddall et al., 2003; Grant et al., 2014). Most of this eustatic sea-level signature is driven by changes in the Laurentide Ice Sheet, but it is difficult to imagine a complete decoupling of Laurentide and Greenland ice sheet behavior where the Laurentide remains relatively large while the GrIS is smaller than today during MIS 3. It is certainly possible that our sites were exposed during the MIS 3, but the lack of any additional terrestrial evidence in southwestern Greenland for a MIS 3 ice-sheet configuration similar to or more restricted than today, makes us favor the more straightforward explanation of MIS 5e exposure. While perhaps unsurprising that our  $^{26}\text{Al}$ - $^{10}\text{Be}$  measurements in the KNS forefield suggest surface exposure during a previous interglacial, these measurements nonetheless confirm the GrIS margin in the KNS region was likely at or behind the present margin during MIS 5e.

## 6 Holocene evolution of the southwestern Greenland Ice Sheet

### 6.1 Early Holocene moraine deposition

Direct  $^{10}\text{Be}$  ages from moraine boulders constrain Kapisigdlit stade moraine deposition to  $10.24 \pm 0.31$  ka BP, which is supported by statistically identical bracketing  $^{10}\text{Be}$  ages from erratic boulders located outboard and inboard of the Kapisigdlit stade moraine (Fig. 2; Fig. 7). North of KNS at Qamanaarsuup Sermia,  $^{10}\text{Be}$  ages from moraine boulders indicate that deposition of so-called Kapisigdlit moraine occurred at  $9.57 \pm 0.33$  ka BP, which is supported by bracketing  $^{10}\text{Be}$  ages of  $10.29 \pm 0.14$  ka and  $9.29 \pm 0.07$  ka (Fig. 3). These two moraines share identical maximum-limiting ages, but the direct moraine and minimum-limiting  $^{10}\text{Be}$  ages between the two sites are statistically distinguishable. Maximum-limiting  $^{10}\text{Be}$  ages from erratic boulders located immediately outboard of a moraine can provide a close constraint on the age of moraine deposition if moraine deposition occurs via a brief stillstand of the ice margin (e.g. Young et al., 2013b). If moraine deposition occurs after a re-advance of the ice margin, however, maximum limiting  $^{10}\text{Be}$  ages need not be close limiting ages. In contrast, minimum limiting  $^{10}\text{Be}$  ages from immediately inside a moraine should provide close minimum-limiting constraints regardless if moraine deposition occurred through a stillstand of the ice margin during retreat or after a readvance. The distribution of  $^{10}\text{Be}$  ages presented here suggests that deposition of the Kapisigdlit stade moraine in the immediate KNS region occurred via a stillstand of the ice-margin or a brief re-advance that occurred within the resolution of our chronometer. At Qamanaarsuup Sermia, however, there is a several hundred-year gap between maximum-limiting  $^{10}\text{Be}$  ages and the moraine-based  $^{10}\text{Be}$  ages, and a total of  $\sim 1$  kyr between maximum- and minimum-limiting  $^{10}\text{Be}$  ages suggesting the Qamanaarsuup Sermia moraine suite was deposited after a re-advance of the GrIS.

Including the uncertainty in the  $^{10}\text{Be}$  production rate, the Kapisigdlit stade moraine in the KNS forefield was deposited at  $10.24 \pm 0.36$  ka BP, consistent with a maximum-limiting radiocarbon age of  $10,170 \pm 340$  cal yr BP (Weidick et al., 2012; Larsen et al., 2014), and moraine deposition at Qamanaarsuup Sermia occurred at  $9.57 \pm 0.38$  ka BP. Moraine deposition in the KNS forefield at  $10.24 \pm 0.36$  ka BP is, within resolution, synchronous with widespread moraine deposition in Baffin Bay



605 ca. 10.4-10.3 ka BP. Indeed,  $^{10}\text{Be}$  ages across Baffin Bay reveal that sectors of the GrIS, Laurentide Ice Sheet, and independent  
alpine glaciers on Greenland and Baffin Island all deposited moraines ca. 10.4-10.3 ka BP, likely in response to freshwater-  
induced regional cooling (Young et al., 2020a). Contemporaneous moraine deposition indicates that the KNS sector of the  
GrIS also likely responded to regional cooling ca. 10.4-10.3 ka. Moraine deposition  $9.57 \pm 0.38$  ka BP at Qamanaarsuup  
Sermia, however, does not fit a well-established regional pattern of moraine deposition. Similar to moraine deposition ca. 10.4-  
610 10.3 ka BP, widespread moraine deposition across Baffin Bay occurred ca. 9.3-9.0 ka BP and is thought to be driven by the  
9.3 ka cooling event displayed in Greenland ice cores (Young et al., 2020a). The Qamanaarsuup Sermia moraine age ( $9.57 \pm$   
 $0.38$  ka BP) is, within uncertainties, synchronous with the 9.3 ka cooling event, but  $^{10}\text{Be}$ -dated moraines in Baffin Bay  
consistently date to at or slightly after the 9.3 ka cooling event (Young et al., 2011b; Young et al., 2020a; Crump et al., 2020).  
North of KNS,  $^{10}\text{Be}$  ages constrain deposition of a moraine to  $9.7 \pm 0.7$  ka, consistent with moraine deposition at Qamanaarsuup  
615 Sermia despite somewhat larger uncertainties (Lesnek and Briner, 2018). In addition to widespread moraine deposition in  
Baffin Bay ca. 10.4-10.3 and 9.3-9.0 ka BP, these emerging  $^{10}\text{Be}$  ages tentatively suggest an additional mode of moraine  
deposition ca. 9.7 ka, perhaps in a response to freshwater related cooling (Lesnek and Briner, 2018). Regardless of our ability  
to correlate the Qamanaarsuup Sermia moraine dated to  $9.57 \pm 0.38$  ka BP to moraines beyond the KNS region, our  $^{10}\text{Be}$  ages  
reveal that early Holocene moraines across the broader KNS region are not equivalent features. Our results reveal at least two  
620 periods of moraine deposition occurred at  $10.24 \pm 0.36$  ka BP and  $9.57 \pm 0.38$  ka BP. Lastly, we note that there is no moraine  
associated with the 8.2 ka abrupt cooling event in the KNS and Qamanaarsuup Sermia regions. Whereas widespread moraine  
deposition in Baffin Bay occurred in response to the 8.2 ka event (Young et al., 2020a), the GrIS retreated inboard of the  
eventual historical ice limit well prior to 8.2 ka at KNS and Qamanaarsuup Sermia. Any 8.2 ka event related moraine that may  
have existed on the landscape was overrun and destroyed by the historical advance of the GrIS.

## 625 **6.2 Retreat of the GrIS behind the modern margin during the early Holocene**

We next consider the timing and significance of when the GrIS margin crossed the historical maximum/modern ice  
margin threshold during early Holocene deglaciation. At KNS,  $^{10}\text{Be}$  ages from just outboard of the historical maximum limit  
630 indicate that following deposition of the Kapisigdlit stade moraine at  $10.24 \pm 0.36$  ka BP, the GrIS crossed the historical  
maximum limit soon thereafter at  $10.20 \pm 0.23$  ka (Fig. 2). At Qamanaarsuup Sermia,  $^{10}\text{Be}$  ages from just outboard of the  
historical maximum reveal that this portion of the GrIS margin crossed the historical maximum limit at  $9.29 \pm 0.18$  ka BP,  $\sim 1$   
kyr later than at KNS (Fig. 3). Near Narsap Sermia (Fig. 1),  $^{10}\text{Be}$  ages from outboard of the historical maximum limit indicate  
that the area deglaciated at  $\sim 8.8$  ka BP (Fig. 1), and south of KNS,  $^{10}\text{Be}$  ages suggest that the landscape just outboard of the  
635 historical maximum deglaciated between  $\sim 10.0$  and 9.5 ka BP. Farther south, a single  $^{10}\text{Be}$  age suggests that the GrIS margin  
did not retreat behind its modern margin until  $\sim 7.7$  ka BP (Fig. 1). Lastly, at Saqqap Sermia, basal radiocarbon ages suggest  
that deglaciation of the landscape immediately outboard the modern margin occurred as late as  $\sim 5.4$ - $4.6$  ka BP, at least 5 kyr  
after deglaciation of the landscape outboard of the modern ice margin in the KNS region (Fig. 1; Levy et al., 2017).



At face value, there is >5 kyr spread in the timing of deglaciation immediately outboard of the historical maximum  
640 limit in the broader KNS region, perhaps suggestive of substantial differences in ice-margin behavior in the early to middle  
Holocene (Fig. 1). When considering all of the available ice-margin constraints, deglaciation occurred earliest in the immediate  
KNS region, with deglaciation occurring later in sectors beyond KNS. The timing of deglaciation of the landscape immediately  
outboard of the historical maximum, however, is dictated by expected differences in the rate of ice-margin retreat and  
differences in the magnitude of late Holocene re-advance of the ice margin. Older  $^{10}\text{Be}$  ages from outboard of the historical  
645 limit at KNS relative to adjacent ice margins could simply reflect the earlier deglaciation of KNS compared to neighboring ice  
margins. In this scenario, marine-based dynamic processes would likely drive early and rapid deglaciation of KNS, whereas  
deglaciation would lag behind in adjacent land-based ice margins where marine-based dynamical processes exert less control  
on ice-margin behavior. Alternatively, the pattern of early deglaciation at KNS with later deglaciation in adjacent margins can  
be entirely explained by the magnitude of the late Holocene re-advance of the ice margin. For example, if the late Holocene  
650 readvance of KNS was of greater magnitude than that of adjacent margins, then the KNS terminus would overrun and rest  
upon a landscape that deglaciated earlier, and thus  $^{10}\text{Be}$  ages from outboard of the historical moraine would be older. The  
greater the magnitude of ice-margin readvance, the older the  $^{10}\text{Be}$  ages just outboard of the historical moraine will be.

Additional chronological constraints that track the history of the GrIS margin prior to, and immediately following  
when the ice margin retreated behind the eventual historical maximum limit, places the apparent asynchrony of deglaciation  
655 across the KNS region within a broader context. At Qamanaarsuup Sermia,  $^{10}\text{Be}$  ages just outboard of the historical moraine  
are  $9.29 \pm 0.18$  ka, however, radiocarbon-dated lake sediments suggest that ice remained near the historical maximum extent  
until  $\sim 8.3$  ka BP (Fig. 3; Fig. 6). Including  $^{10}\text{Be}$  ages from just beyond the early Holocene moraines, all available ice-margin  
constraints at Qamanaarsuup Sermia indicate the position of the GrIS margin in this region underwent minimal changes  
between  $\sim 10.3$  ka BP and  $\sim 8.3$  ka BP and at least  $\sim 2$  kyr of ice-margin history is represented in a relatively restricted lateral  
660 zone on the landscape. Had the late Holocene readvance of the Qamanaarsuup Sermia margin been slightly more extensive,  
the ice margin would have overrun the early Holocene moraines currently residing immediately outboard of the historical  
maximum. The resulting historical maximum limit would abut a landscape that deglaciated just prior to 10 ka BP, similar to  
the relationship between the historical maximum ice limit and  $^{10}\text{Be}$  ages outboard of this limit observed at KNS. Or, had the  
late Holocene readvance been slightly less extensive, the historical moraine would abut a landscape that likely deglaciated  
665  $\sim 8.3$  ka BP, similar to the timing of deglaciation at other locations in a broader KNS region (Fig. 1). In a similar manner,  $^{10}\text{Be}$   
ages at KNS indicate that the ice margin retreated behind the historical maximum limit at  $10.20 \pm 0.23$  ka, but ice did not  
continue to retreat inland and instead remained near the current margin until  $\sim 7.8$ - $6.9$  ka BP based on concordant  $^{10}\text{Be}$  and *in*  
*situ*  $^{14}\text{C}$  ages from recently exposed bedrock (section 5.2). Near Narsap Sermia (Fig. 1),  $^{10}\text{Be}$  ages from outboard of the  
historical maximum limit, paired with a basal radiocarbon age from Caribou Lake, indicate that the area deglaciated at  $\sim 8.8$  ka  
670 BP, but ice remained in the lake catchment and likely near the modern margin until  $\sim 7.5$  ka BP (Fig. 1). Additional  $^{10}\text{Be}$  ages  
south of KNS near Deception Lake suggest that the landscape just outboard the historical maximum deglaciated between  $\sim 10.0$   
and  $9.5$  ka BP, but the ice margin was likely near its present limit until  $\sim 9.2$  ka BP based on basal radiocarbon ages.



675 Additional locations with paired  $^{10}\text{Be}$  ages and proglacial-threshold lake records along much of the western GrIS  
margin reveal that after the GrIS margin retreated behind the position of the historical maximum extent, the ice margin  
remained near this position for several hundred to thousands of years before retreating farther inland (Fig. 16). Notably, paired  
 $^{10}\text{Be}$  ages and radiocarbon constraints north of Jakobshavn Isbræ suggest that the ice margin was near its current position  
between  $\sim 10$  ka BP and 5.2 ka BP (Newspaper Lake; Cronauer et al., 2015; Fig. 1; Fig. 16). Closer to Jakobshavn Isbræ, paired  
 $^{10}\text{Be}$  ages and radiocarbon constraints suggest the ice margin was in a configuration similar to today between 7.8 ka BP and  
5.5 ka BP (Lake Lo; Håkansson et al., 2014; Fig. 1; Fig. 16). In the Kangerlussuaq region, radiocarbon ages from proglacial-  
680 threshold lakes reveal that the ice margin remained in a configuration similar to today following initial deglaciation (Fig. 1;  
Fig. 16; Young and Briner, 2015; Lesnek et al., 2020). Unique to the Kangerlussuaq region is proglacial lake Tasersuaq, where  
radiocarbon-dated sediments, combined with maps of sub-ice topography suggest the GrIS margin never retreated out the  
Tasersuaq catchment, which extends only  $\sim 1.9$  km behind the modern margin, during the Holocene (Lesnek et al., 2020).  
Additional proglacial-threshold lake records from separate drainage basins in the Kangerlussuaq region suggest that the ice  
685 margin retreated  $\sim 3.7$  km and  $\sim 26$  km inland during the Holocene (Lesnek et al., 2020), and broadly support minimal ice-  
margin recession during the middle Holocene.

We suggest that the large range in ages constraining the timing of deglaciation outboard of the historical moraine  
within the KNS region and along the broader western GrIS margin, coupled with ice-margin constraints from proglacial-  
threshold lakes and cosmogenic isotope measurements from recently exposed bedrock, broadly define a window between  $\sim 10$ -  
690 5 ka BP when the GrIS margin was near its current margin. The southwestern GrIS margin reached its late Holocene maximum  
extent during historical times, but records from proglacial-threshold lakes indicate that the ice margin had advanced back to  
near the modern margin by  $\sim 2$  ka BP (Fig. 16). Within a relatively narrow  $\sim 3$  kyr window between  $\sim 5$ -2 ka BP, the  
southwestern GrIS margin retreated inland, achieved its minimum extent, and readvanced towards the historical maximum  
extent, which likely precludes significant inland retreat of the southwestern GrIS margin during the Holocene.

### 695 **6.3 Geologic data-model comparison of ice-margin change in southwest Greenland**

Geologic reconstructions of ice-sheet change offer an ideal target for ice-sheet modeling efforts aimed at  
reconstructing the geometry and ice volume of past ice sheets through model tuning (Simpson et al., 2009; Lecavalier et al.,  
700 2014). These reconstructions, however, also serve as an ideal testbed for modeling efforts aimed at evaluating the sensitivity  
of past ice-margin migration to climatic and oceanic influences (Briner et al., 2020). Paleo ice-sheet modeling efforts typically  
rely on coarse resolution meshes and simplification of ice-flow approximations to achieve computational efficiency. While  
this approach has enabled more simple models to assess the sensitivity of ice-margin response to model parameters, crude  
climate forcings are typically used, making it difficult to properly assess ice-margin sensitivity to climate. Here, we further  
705 explore the deglaciation history of the KNS region by comparing the geologic record of ice-sheet change to recently completed  
model simulations of southwestern GrIS evolution through the Holocene (Cuzzone et al., 2019; Briner et al., 2020). These



710 Holocene ice model simulations use the highest horizontal mesh resolution across our field area to date, and use a range of state-of-the-science gridded climate reconstructions as the input climate (Badgeley et al., 2020; Briner et al., 2020), thus presenting an opportunity for new insights regarding glacier history and ice-sheet modeling. Our goals in exploring the model simulations are to (a) assess the magnitude of recession inboard of the present margin, (b) compare rates of retreat and timing of ice-margin change in both the model and in the observations, and (c) explore avenues for model improvement in a known problem area for ice-sheet modeling.

715 Across the broader southwestern Greenland region, simulations are able to reproduce the observed pattern of ice-margin migration through the Holocene – early to middle Holocene retreat, inland recession, and late Holocene readvance (Fig. 17). In particular, north of the KNS region between Sisimiut and Kangerlussuaq, the simulated pattern and timing of ice-margin migration generally reproduces the geologic record of ice-margin change (Briner et al., 2020). In the KNS region, these same simulations generally depict an ice-margin that retreats eastwards, achieves a minimum extent, and then readvances in the late Holocene, thus sharing the broad outline of GrIS change provided by the available geological observations (Fig. 17). However, these simulations consistently depict ice that is too extensive compared to the observed record of ice-margin change in the KNS region (Fig. 17). Similar to the modeled pattern of ice-margin change, simulated ice masses in southwestern Greenland decrease through the early Holocene and achieve a minimum value between ~7.6 ka BP and 6.3 ka BP (Fig. 17).

720 The better model-data fit in the Kangerlussuaq region compared to the KNS region is likely the result of distinctly different ice-margin environments. The Kangerlussuaq region hosts what is primarily a land-terminating sector of the GrIS where ice-sheet behavior is dominated by surface mass balance (Cuzzone et al., 2019; Downs et al., 2020). Thus, ice-sheet behavior in the Kangerlussuaq region is almost entirely dictated by the climate forcings used in the model runs (Briner et al., 2020). The KNS fjord system, however, hosted a marine-terminating sector of the GrIS influenced by dynamical processes, but the model setup does not include calving or submarine melting of floating ice. Therefore, there is no dynamic mechanism by which to rapidly remove ice from the KNS fjord system in the model. Because the ice sheet model is built on a high-resolution mesh, down to 2 km in the KNS region, it is able to resolve bed features such as the existing KNS topography and fjord bathymetry, allowing for better representation of mass transport and stress balance than in lower resolution models. By resolving these features, however, ice is funneled efficiently to the ice margin, allowing the ice to persist in the KNS fjord system in the absence of calving, which is likely responsible for the simulated ice-margin being too extensive compared to the geologic record.

735 Evaluating the fit between our ice-sheet model results and the geochronological data in the KNS region must be treated cautiously as climatic and dynamic processes may not be the only influences on model-data mismatches. For example, the simulated GrIS retreat through the KNS region is sensitive to how well bed topography is resolved (Cuzzone et al., 2019). In simulations using a lower resolution mesh (i.e. 10 km or greater), fjord bathymetry in the KNS region is not well resolved, and as the ice surface lowers in response to Holocene warming, it intersects bedrock bumps (i.e. pinning points) that would otherwise be resolved as fjords using a higher resolution mesh. In this scenario, ice-margin migration influenced by bedrock sticking points due to a lower resolution mesh might give a false impression of good model fit to the data, but for reasons





solely related to model resolution. Regardless, simulated ice in the KNS fjord system that is too extensive suggests that marine forcings not included in our model likely played an important role in ice-margin retreat. While implementation and treatment of calving in ice sheet models continues to improve (e.g. Benn and Åström, 2018), our results highlight that inclusion of calving is necessary towards fully understanding ice-margin sensitivity to climate change in the KNS region. Yet, it is difficult to properly simulate calving in paleoclimate ice-sheet model setups that use coarse resolution grids, because fjord systems in Greenland are typically <5km and high-resolution grids (1 km) are necessary to capture grounding line migration (Seroussi et al., 2018).

The simulated ice margin positions in the Kangerlussuaq and KNS regions relative to the geological constraints offer insights into model-data fit across the entire model domain and the possible climatic conditions influencing ice-sheet behavior through the Holocene. Among the individual model runs of Cuzzone et al (2019) and Briner et al (2020), some simulated ice masses generally achieve a minimum between ~7.6 ka BP and 7.1 ka BP (simulations 1-6; Fig. 17), but there are three notable exceptions where the ice-mass minimum occurs later at ~6.3 ka BP (simulations 7-9; Fig. 17). These simulations that result in a later ice-mass minimum also depict a minimum mass that is less extreme with significantly less regrowth of the ice sheet following the minimum (Fig. 17). At the same time, these simulations with a later ice-mass minimum depict a more subdued, although still discernible, response to the 8.2 ka cooling event relative to other simulations despite the well-documented response of the southwestern GrIS to 8.2 ka cooling (Fig. 16; Fig. 17; Young et al., 2011b, 2013b, 2020a). Nonetheless, relative sea-level records from southwestern Greenland fall below modern sea level at ~4-3 ka BP before later rising towards modern sea-level (Long et al., 2011; Lecavalier et al., 2014), which is interpreted to reflect the re-loading of the crust during late Holocene re-growth of the southwestern GrIS initiating in the last few kyr. Given all of the geologic ice margin constraints from across southwestern Greenland considered here, the minimum extent of the GrIS likely occurred sometime after ~5 ka BP. Thus, simulations that produce an ice-mass minimum between ~7.6 and 7.1 ka BP appear to be inconsistent with the geological record. Simulations that result in an ice-mass minimum at ~6.3 ka BP are more compatible with the geologic record, especially when considering these model runs simulate a subtle and broad post-6.3 ka BP plateau followed by slight late Holocene regrowth (Fig. 17).

Simulations 7 through 9 rely on a temperature history that has a muted early Holocene warming compared to other runs, followed by peak Holocene mean annual temperature anomalies above the 1850-2000 mean from 7 ka to 4 ka BP (Badgley et al., 2020; Briner et al., 2020; Fig. 17). Run 7 relies on a precipitation history that has increased precipitation during 8 ka to 4 ka BP relative to the CE 1850-2000 mean, run 8 uses precipitation anomalies that are similar to the CE 1850-2000 mean, and run 9 uses precipitation anomalies that are lower than the CE 1850-2000 mean. Indeed, in the SMB-dominated Kangerlussuaq region, runs 7 and 8 appear to provide the best model-data fit of ice margin position, while also having an ice-mass minimum most consistent with the geologic record (Fig. 17). The higher precipitation scenario used in run 7 is broadly supported by proxy evidence of enhanced wintertime snowfall in southwestern Greenland and inferred precipitation using an ice-sheet flowline model (Thomas et al., 2018; Downs et al., 2020). Despite early Holocene warming, limited evidence suggests that this warmth, and its effect on ice-sheet mass balance, may be offset to some degree by increased precipitation. The timing



775 of the minimum inland extent of the ice margin may occur at slightly different times across southwestern Greenland and  
therefore mass does not necessarily equate to the most retracted ice margin. However, across all model runs using different  
climatologies, the simulated ice-mass minimum generally equates to a modeled ice margin near or slightly inboard of its current  
position (Fig. 17).

Our results highlight the potential of proxy-informed gridded climate reconstructions and points to where continued  
improvements to climate reconstructions used in paleo ice sheet modeling efforts are having the biggest impact; each different  
climate history applied here results in a slightly different simulated ice-sheet history. Simulated ice-sheet histories provide a  
better fit to geologic constraints in the SMB-dominated domain in the Kangerlussuaq region, with a relatively poorer model-  
data fit in the KNS region. Although some model-data mismatch occurs across our domain when considering lateral ice-margin  
position, rates of GrIS mass loss inferred for our domain (i.e. Briner et al., 2020) remain robust as differences in simulated ice-  
margin migration (i.e. slightly too fast near Kangerlussuaq and too slow in the KNS compared to geologic constraints; Fig. 17)  
likely offset each other to some degree when considered in the context of total ice-mass loss. Ultimately, our results suggest  
that oceanic and dynamic processes that are not included in our modeling effort likely play a key role in dictating the ice-  
margin retreat pattern in the KNS region.

## 790 7 Conclusions

New  $^{10}\text{Be}$  ages from the KNS region, southwestern Greenland, constrain deposition of two separate segments of the  
Kapisigdlit stade moraines to  $10.24 \pm 0.36$  ka BP and  $9.57 \pm 0.38$  ka BP, indicating that these moraines are not equivalent  
features. The older moraine is synchronous with widespread moraine deposition in Baffin Bay at this time, whereas the younger  
moraine is consistent with an additional  $^{10}\text{Be}$ -dated moraine in southwestern Greenland, tentatively defining a new mode of  
GrIS moraine deposition at  $\sim 9.7$  ka BP. Following early Holocene moraine deposition in the KNS forefield, the GrIS margin  
retreated inboard of the eventual historical maximum extent/modern ice margin. The timing of deglaciation of the landscape  
immediately outboard of the historical maximum extent in the KNS region and along much of the southwestern GrIS margin  
varies by several thousand years. Yet, additional chronological constraints provided by proglacial-threshold lakes and  
cosmogenic nuclide measurements from recently exposed bedrock surfaces constrain a several-thousand-year interval of time  
when the GrIS margin was within, but near its modern position, and the minimum GrIS extent occurring sometime after  $\sim 5$  ka  
BP. The variability in  $^{10}\text{Be}$  ages just outboard of the historical maximum limit is likely the result of slight variations in ice-  
sheet retreat and the magnitude of the late Holocene readvance of the GrIS, rather than major differences in ice-margin history  
or regional climate variability. The southwestern GrIS margin may have advanced back to the eventual historical maximum  
extent as early as  $\sim 2$  ka BP, leaving an  $\sim 3$  kyr window for the ice margin to achieve its minimum inland position.

Triple  $^{10}\text{Be}$ - $^{14}\text{C}$ - $^{26}\text{Al}$  measurements in recently exposed bedrock fronting the modern GrIS help constrain the  
minimum inland extent of the GrIS margin over multiple timescales. Our paired  $^{26}\text{Al}$ - $^{10}\text{Be}$  and  $^{14}\text{C}$ - $^{10}\text{Be}$  measurements are  
unable to detect any surface burial and are consistent with constant exposure of our sampled bedrock sites.  $^{14}\text{C}$ - $^{10}\text{Be}$



810 measurements constrain the magnitude of pre-Holocene isotopic inheritance at our bedrock sites and reveal that the period of  
18<sup>th</sup>-20<sup>th</sup> century ice cover was the only extended period that these bedrock sites became re-occupied by the GrIS following  
middle Holocene deglaciation. <sup>26</sup>Al-<sup>10</sup>Be measurements also reveal constant exposure of these sites and suggest that the slight  
amount of inherited <sup>10</sup>Be present in a subset of our bedrock samples is due to a period of surface exposure prior to the LGM,  
likely during MIS 5e. *In situ* <sup>14</sup>C inventories indicate that bedrock presently emerging from beneath the GrIS in the KNS  
815 forefield was exposed during the middle Holocene. Contemporary retreat of GrIS has yet to access a landscape that remained  
ice covered throughout the Holocene.

820 Geologic reconstructions in southwestern Greenland constrain the Holocene behavior of the GrIS across a land  
terminating, surface mass balance dominated region, and an area where the ice margin retreated rapidly as the ice front  
responded to dynamical changes imposed through marine influences (i.e. calving and/or submarine melting of floating ice). As  
paleo ice sheet models continue to improve both in the representation of processes controlling ice-margin migration and use  
825 of high-resolution model meshes, these geologic reconstructions provide robust validation targets upon which to benchmark  
and improve models simulating the past behavior of ice sheets. Continued use of model-data comparisons as ice-sheet modeling  
efforts, paleoclimate datasets, and geologic reconstructions become more refined will improve our understanding of past ice-  
sheet sensitivity to climatic and dynamic forcing mechanisms.

## 825 **Data Availability**

All analytical information for new cosmogenic-nuclide and traditional radiocarbon measurements is listed in the tables in the  
Supplement. Analytical information is also available in the ICE-D:Greenland online database (<http://greenland.ice-d.org/>,  
830 Balco, 2020).

## **Acknowledgements**

We thank the 109<sup>th</sup> Airlift Wing of the New York Air National Guard for transport to and from Greenland, Air Greenland for  
helicopter support, and CH2MHill Polar Field Services for additional logistical support. We also thank J. Hanley and J. Frisch  
835 for help processing <sup>10</sup>Be and <sup>26</sup>Al samples. Funding was provided by the National Science Foundation Arctic Natural Sciences  
and Arctic System Sciences programs: Awards #1417675 and #1503959 to NEY and JMS, and Awards #1417783 and  
#1504267 to JPB.

## **Author contributions**

840 Project conceived by NEY, JPB and JMS. NEY, AJL, and JKC wrote the first draft of manuscript with input from JPB, JAB,  
ABK, and JMS. All authors commented on and edited the manuscript. NEY, AJL, JPB, JAB, BLG, and AC completed the  
fieldwork. JKC led the ice-sheet modelling component and JAB provided the regional input climatologies. RS, NEY, and JMS



completed the  $^{10}\text{Be}$  and  $^{26}\text{Al}$  extraction, and NEY completed the *in situ*  $^{14}\text{C}$  extraction. MWC and SRHZ conducted final  $^{10}\text{Be}$  and  $^{26}\text{Al}$  measurements. TT and EB completed final gas-source  $^{14}\text{C}$  measurements. NEY, ABK, and JLL completed the final  
845 *in situ*  $^{14}\text{C}$  data reduction and interpretations. AJL and JPB isolated organic remains for traditional  $^{14}\text{C}$  dating.

### Competing interests

The authors declare that they have no conflict of interest.

850

### References

- Argento, D.C., Reedy, R.C., and Stone, J.O.: Modeling the earth's cosmic radiation, Nucl. Instrum. Methods Phys. Res. Sect. B., 294, 464-469, doi:10.1016/j.nimb.2012.05.022, 2013.
- 855 Badgeley, J.A., Steig, E.J., Hakim, G.J., and Fudge, T.J.: Greenland temperature and precipitation over the last 20,000 years using data assimilation, Clim. Past., 16, 1325-1326, https://doi.org/10.5194/cp-16-1325-2020, 2020.
- Balco, G.: Production rate calculations for cosmic-ray-muon-produced  $^{10}\text{Be}$  and  $^{26}\text{Al}$  benchmarked against geological calibration data, Quat. Geochronol., 39, 150-173, doi:10.1016/j.quageo.2017.02.001, 2017
- Balco, G.: ICE-D:Greenland, available at: <http://greenland.ice-d.org>, last access: 27 August 2020.
- 860 Balco, G. and Rovey II, C.W.: An isochron method for cosmogenic-nuclide dating of buried soils and sediments, Am. J. Sci., 308, 1083-1114, doi:10.2475/10.2008.02, 2008
- Balco, G., Stone, J., Lifton, N., and Dunai, T.J.: A complete and easily accessible means of calculating surface exposure ages and erosion rates from  $^{10}\text{Be}$  and  $^{26}\text{Al}$  measurements, Quat. Geochronol., 3, 174-195, doi:10.1016/j.quageo.2007.12.001, 2008
- 865 Bard, E., Tuna, T., Fagault, Y., Bonvalot, L., Wacker, L., Fahrni, S., and Synal, H.-A.: AixMICADAS, the accelerator mass spectrometer dedicated to  $^{14}\text{C}$  recently installed in Aix-en-Provence, France, Nucl. Instrum. Methods Phys. Res. Sect. B, 361, 80-86, doi:10.1016/j.nimb.2015.01.075, 2015.
- Benn, D.I. and Åström, J.A.: Calving glaciers and ice shelves, Advances in Physics: X, 3, 1513819, doi:10.1080/23746149.2018.1513819, 2018.
- 870 Bennike, O. and Björck, S.: Chronology of the last recession of the Greenland ice sheet, J. Quat. Sci., 17, 211-219, doi:10.1002/jqs.670, 2002.
- Bierman, P.R., Marsella, K.A., Patterson, C., Thompson Davis, P., and Caffee, M.: Mid-Pleistocene cosmogenic minimum-age limits for pre-Wisconsin glacial surfaces in southwestern Minnesota and southern Baffin Island: a multiple nuclide approach, Geomorphology, 27, 25-39, doi:10.1016/S0169-555X(98)00088-9, 1999.
- 875 Bierman, P.R., Shakun, J.D., Corbett, L.B., Zimmerman, S.R., and Rood, D.H.: A persistent and dynamic East Greenland Ice Sheet over the past 7.5 million years, Nature, 540, 256-260, doi:10.1038/nature20147, 2016.



- Briner, J.P., Stewart, H.A.M., Young, N.E., Phillips, W., and Losee, S.: Using proglacial-threshold lakes to constrain fluctuations of the Jakobshavn Isbræ ice margin, western Greenland, during the Holocene, *Quat. Sci. Rev.*, 29, 3861-3874, doi:10.1016/j.quascirev.2010.09.055, 2010.
- 880 Briner, J.P., Cuzzone, J.K., Badgley, J.A., Young, N.E., Steig, E.J., Morlighem, M., Schlegel, N.-J., Hakim, G., Schaefer, J., Johnson, J.V., Lesnek, A.L., Thomas, E.K., Allan, E., Bennike, O., Cluett, A.A., Csatho, B., de Vernal, A., Downs, J., Larour, E., and Nowicki, S.: Rate of mass loss from the Greenland Ice Sheet will exceed Holocene values this century, *Nature*, doi:10.1038/s41586-020-2742-6, 2020.
- Chmeleff, J., von Blanckenburg, F., Kossert, K., and Jakob, D.: Determination of the  $^{10}\text{Be}$  half-life by multicollector icp-ms and liquid scintillation counting, *Nucl. Instrum. Methods Phys. Res. Sect. B*, 268, 192-199, doi:10.1016/j.nimb.2009.09.012, 2010.
- 885 Colville, E.J., Carlson, A.E., Beard, B.L., Hatfield, R.G., Stoner, J.S., Reyes, A.V., and Ullman, D.J.: Sr-Nd-Pb isotope evidence for ice-sheet presence on southern Greenland during the Last Interglacial, *Science*, 333, 620-623, doi:10.1126/science.1204673, 2011.
- 890 Corbett, L.B., Bierman, P.R., Rood, D.H., Caffee, M.W., Lifton, N.A., and Woodruff, T.E.: Cosmogenic  $^{26}\text{Al}/^{10}\text{Be}$  surface production ratio in Greenland, *Geophys. Res. Lett.*, 44, 1350-1359, doi:10.1002/2016GL071276, 2017.
- Cronauer, S.L., Briner, J.P., Kelley, S.E., Zimmerman, S.R.H., and Morlighem, M.:  $^{10}\text{Be}$  dating reveals early-middle Holocene age of the Drygalski Moraines in central West Greenland, *Quat. Sci. Rev.*, 147, 59-68, doi:10.1016/j.quascirev.2015.08.034, 2015.
- 895 Crump, S.E., Young, N.E., Miller, G.H., Pendleton, S.L., Tulenko, J.P., Anderson, R.S., and Briner, J.P.: Glacier expansion on Baffin Island during early Holocene cold reversals, *Quat. Sci. Rev.*, 241, 106419, doi:10.1016/j.quascirev.2020.106419, 2020.
- Cuzzone, J.K., Morlighem, M., Larour, E., Schlegel, N., and Seroussi, H.: Implementation of higher-order vertical finite elements in ISSM v4.13 for improved ice sheet flow modeling over paleoclimate timescales, *Geosci. Model Dev.*, 11, 1683-1694, doi:10.5194/gmd-11-1683-2018, 2018.
- 900 Cuzzone, J.K., Schlegel, N.-J., Morlighem, M., Larour, E., Briner, J.P., Seroussi, H., and Caron, L.: The impact of model resolution on the simulated Holocene retreat of the southwestern Greenland ice sheet using Ice Sheet System Model (ISSM), *Cryosphere*, 13, 879-893, https://doi.org/10.5194/tc-13-879-2019, 2019.
- de Vernal, A. and Hillaire-Marcel, C.: Natural variability of Greenland climate, vegetation, and ice volume during the past million years, *Science*, 320, 1622-1625, doi:10.1126/science.1153929, 2018.
- 905 Downs, J., Johnson, J., Briner, J., Young, N., Lesnek, A., and Cuzzone J.: Western Greenland ice sheet retreat history reveals elevated precipitation during the Holocene thermal maximum, *Cryosphere*, 14, 1121-1137, doi.org/10.5194/tc-14-1121-2020, 2020.
- Fabel, D., Stroeven, A.P., Harbor, J., Kleman, J., Elmore, D., Fink, D.: Landscape preservation under Fennoscandian ice sheets determined from in situ produced  $^{10}\text{Be}$  and  $^{26}\text{Al}$ , *Earth. Planet. Sci. Lett.*, 201, 397-406, doi:10.1016/S0012-
- 910



- 821X(02)00714-8, 2002.
- Funder, S., Kjeldsen, K.K., Kjær, K., and Ó Cofaigh, C.: The Greenland Ice Sheet during the past 300,000 years: a review. In: Ehlers, J., Gibbard, P.L., Hughes, P.D. (Eds), *Developments in Quaternary Sciences*. Elsevier, Amsterdam, The Netherlands, 699-713, 2011.
- 915 Gjermundsen, E.F., Briner, J.P., Akçar, N., Foros, J., Kubik, P.W., Salvigsen, O., Hormes, A.: Minimal erosion of Arctic alpine topography during late Quaternary glaciation, *Nat. Geosci.*, 8, 789-792, doi: 10.1038/ngeo2524, 2015.
- Goehring, B.M., Schaefer, J.M., Schluechter, C., Lifton, N.A., Finkel, R.C., Jull, A.J.T., Akçar, N., and Alley, R.B.: The Rhone Glacier was smaller than today for most of the Holocene, *Geology*, 39, 679-682, doi:10.1130/G32145.1, 2011.
- Grant, K.M., Rohling, E.J., Bronk Ramsey, C., Cheng, H., Edwards, R.L., Florindo, F., Heslop, D., Marra, F., Roberts, A.P., 920 Tamisiea, M.E., and Williams, F.: Sea-level variability over five glacial cycles, *Nat. Comm.*, 5, 5076, doi: 10.1038/ncomms6076, 2014.
- Håkansson, L., Briner, J.P., Andresen, C.S., Thomas, E.K., and Bennike, O.: Slow retreat of a land-based sector of the West Greenland Ice Sheet during the Holocene Thermal Maximum: evidence from threshold lakes at Paakitsoq, *Quat. Sci. Rev.*, 98, 74-83, doi:10.1016/j.quascirev.2014.05.016, 2014.
- 925 Hallet, B., Hunter, L. and Bogen, J.: Rates of erosion and sediment evacuation by glaciers: A review of field data and their implications, *Global. Planet. Change.*, 12, 213-235, doi:10.1016/0921-8181(95)00021-6, 1996.
- Hatfield, R.G., Reyes, A.V., Stoner, J.S., Carlson, A.E., Beard, B.L., Winsor, K., and Welke, B.: Interglacial responses of the southern Greenland ice sheet over the last 430,000 years determined using particle-size specific magnetic and isotopic tracers, *Earth. Planet. Sci. Lett.*, 454, 225-236, doi:10.1016/j.epsl.2016.09.014, 2016.
- 930 He, F., Shakun, J.D., Clark, P.U., Carlson, A.E., Liu, Z., Otto-Bliesner, B.L., and Kutzbach, J.E.: Northern Hemisphere forcing of Southern Hemisphere climate during the last deglaciation, *Nature*, 494, 81-85, doi:10.1038/nature11822, 2013.
- Heaton, T.J., Köhler, P., Butzin, M., Bard, E., Reimer, R.W., Austin, W.E.N., Bronk Ramsey, C., Grootes, P.M., Hughen, K.A., Kromer, B., Reimer, P.J., Adkins, J., Burke, A., Cook, M.S., Olsen, J., Skinner, L.C.: Marine20-The marine radiocarbon age calibration curve (0-55,000 cal BP), *Radiocarbon*, 1-42, doi:10.1017/RDC.2020.68, 2020.
- 935 Henriksen, N., Higgins, A. K., Kalsbeek, F., and Pulvertaft, T. C. R.: Crystalline rocks older than 1600 Ma: The Greenland Precambrian shield. In N. Henriksen, A. K. Higgins, F. Kalsbeek, & T. C. R. Pulvertaft (Eds.), *Greenland from Archaean to Quaternary. Descriptive text to the Geological map of Greenland, 1:2,500,000 (Vol. 185, pp. 12-24) (2nd ed.)*, Geological Survey of Denmark and Greenland Bulletin 18. Copenhagen, Geological Survey of Denmark and Greenland, 2000.
- 940 Hippe, K. and Lifton, N.A.: Calculating isotope ratios and nuclide concentrations for in situ cosmogenic <sup>14</sup>C analyses, *Radiocarbon*, 56, 1167-1174, [https://doi.org/ 10.2458/56.17917](https://doi.org/10.2458/56.17917), 2014.
- Hogan, K.A., Ó Cofaigh, C., Jennings, A.E., Dowdeswell, J.A., and Hiemstra, J.F.: Deglaciation of a major paleo-ice stream in Disko Trough, West Greenland, *Quat. Sci. Rev.*, 147, 5-26, doi:10.1016/j.quascirev.2016.10.018, 2016.
- Jennings, A.E., Andrews, J.T., Ó Cofaigh, C., St. Onge, G., Sheldon, C., Belt, S.T., Cabedo-Sanz, P., and Hillaire-Marcel, C.:





- 945 Ocean forcing of Ice Sheet retreat in central west Greenland from LGM to the early Holocene, *Earth. Planet. Sci. Lett.* 472, 1-13, doi:10.1016/j.epsl.2017.05.007, 2017.
- Kelley, S.E., Briner, J.P., Young, N.E., Babonis, G.S., and Csatho, B.: Maximum late Holocene extent of the western Greenland Ice Sheet during the 20<sup>th</sup> century, *Quat. Sci. Rev.*, 56, 89-98, doi:10.1016/j.quascirev.2012.09.016, 2012.
- Kelley, S.E., Briner, J.P., and Young, N.E.: Rapid ice retreat in Disko Bugt supported by <sup>10</sup>Be dating of the last recession of the western Greenland Ice Sheet, *Quat. Sci. Rev.*, 82, 13-22, doi:10.1016/j.quascirev.2013.09.018, 2013.
- 950 Knutz, P.C., Newton, A.M.W., Hopper, J.R., Huuse, M., Gregersen, U., Sheldon, E., and Dybkjær, K.: Eleven phases of Greenland Ice Sheet shelf-edge advance over the past 2.7 million years, *Nat. Geosci.*, 12, 361-368, doi:10.1038/s41561-019-0340-8, 2019.
- Kobashi, T., Menviel, L., Jeltsch-Thömmes, A., Vinther, B.M., Box, J.E., Muscheler, R., Nakaegawa, T., Pfister, P.L., Döring, M., Leuenberger, M., Wanner, H., and Ohmura, A.: Volcanic influence on centennial to millennial Holocene Greenland temperature change, *Sci. Rep.*, 7, 1441, doi:10.1038/s41598-017-0451-7, 2017.
- 955 Lal, D.: Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models, *Earth Planet. Sci. Lett.*, 104, 424-429, doi:10.1016/0012-821X(91)90220-C, 1991.
- Lamp, J.L., Young, N.E., Koffman, T., Schimmelpfennig, I., Tuna, T., Bard, E., and Schaefer, J.M.: Update on the cosmogenic *in situ* <sup>14</sup>C laboratory at the Lamont-Doherty Earth Observatory, *Nucl. Instrum. Methods Phys. Res. Sect. B*, 456, 157-162, doi:10.1016/j.nimb.2019.05.064, 2019.
- 960 Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *J. Geophys. Res. Earth Surf.*, 117, F01022, doi:10.1029/2011JF002140, 2012.
- 965 Larsen, N.K., Funder, S., Kjær, K.H., Kjeldsen, K.K., Knudsen, M.F., and Linge, H.: Rapid early Holocene ice retreat in West Greenland, *Quat. Sci. Rev.*, 92, 310-323, doi:10.1016/j.quascirev.2013.05.027, 2014.
- Larsen, N.K., Kjær, K.H., Lecavalier, B., Bjørk, A.A., Colding, S., Huybrechts, P., Jakobsen, K.E., Kjeldsen, K.K., Knudsen, K.L., Odgaard, B.V., Olsen, J.: The response of the southern Greenland ice sheet to the Holocene thermal maximum, *Geology*, 43, 291-294, doi:10.1130/G36476.1, 2015.
- 970 Larsen, N.K., Levy, L.B., Carlson, A.E., Buizert, C., Olsen, J., Strunk, A., Bjørk, A.A., Skov, D.S.: Instability of the northeast Greenland ice stream over the last 45,000 years, *Nat. Comm.*, 9, 1872, doi:10.1038/s41467-018-04312-7, 2018.
- Lea, J.M., Mair, D.W.F., Nick, F.M., Rea, B.R., Weidick, A., Kjær, K.H., Morlighem, M., van As, D., Schofield, J.E.: Terminus-driven retreat of a major southwest Greenland tidewater glacier during the early 19<sup>th</sup> century: insights from glacier reconstructions and numerical modelling, *J. Glaciol.*, 60, 333-344, doi: 10.3189/2014JoG13J163, 2014.
- 975 Lecavalier, B.S., Milne, G.A., Simpson, M.J.R., Wake, L., Huybrechts, P., Tarasov, L., Kjeldsen, K.K., Funder, S., Long, A.J., Woodroffe, S., Dyke, A.S., and Larsen, N.K.: A model of Greenland ice sheet deglaciation constrained by observations of relative sea level and ice extent, *Quat. Sci. Rev.*, 102, 54-84, doi:10.1016/j.quascirev.2014.07.018, 2014.



- 980 Lesnek, A.J. and Briner, J.P.: Response of a land-terminating sector of the western Greenland Ice Sheet to early Holocene climate change: Evidence from  $^{10}\text{Be}$  dating in the Søndre Isortoq region, *Quat. Sci. Rev.*, 180, 145-156, doi:10.1016/j.quascirev.2017.11.028, 2018.
- Lesnek, A.J., Briner, J.P., Young, N.E., and Cuzzone, J.K.: Maximum southwest Greenland Ice Sheet recession in the early Holocene, *Geophys. Res. Lett.*, 47, e2019GL083164, doi:10.1029/2019GL083164, 2020.
- 985 Levy, L.B., Larsen, N.K., Davidson, T.A., Strunk, A., Olsen, J., and Jeppesen, E.: Contrasting evidence of Holocene ice margin retreat, south-western Greenland, *J. Quat. Sci.* 32, 604-616, doi:10.1002/jqs.2957, 2017.
- Liu, Z., Otto-Bliesner, B.L., He, F., Brady, E.C., Tomas, R., Clark, P.U., Carlson, A.E., Lynch-Stieglitz, J., Curry, W., Brook, E., Erickson, D., Jacob, R., Kutzbach, J., and Cheng, J.: Transient Simulation of Last Deglaciation with a New Mechanism for Bølling-Allerød Warming, *Science*, 325, 310–314, doi:10.1126/science.1171041, 2009.
- 990 Long, A.J., Woodroffe, S.A., Roberts, D.H., and Dawson, S.: Isolation basins, sea-level changes and the Holocene history of the Greenland Ice Sheet, *Quat. Sci. Rev.*, 30, 3748-3768, doi:10.1016/j.quascirev.2011.10.013, 2011.
- Morlighem, M., Williams, C.N., Rignot, E., An, L., Arndt, J.E., Bamber, J., Catania, G., Chauché, N., Dowdeswell, J.A., Dorschel, B., Fenty, I., Hogan, K. Howat, I., Hubbard, A., Jakobsson, M., Jordan, T.M., Kjeldsen, K.K., Millan, R., Mayer, L., Mouginot, J., Noël, B.P.Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M.J., Slabon, P., Straneo, F., van den Broeke, M.R., Weinrebe, W., Wood, M., and Zinglensen, K.B.: BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multibeam echo sounding combined with mass conservation, *Geophys. Res. Lett.* 44, 11,051-11,061, doi:10.1002/2017GL074954, 2017.
- 995 NEEM Community Members: Eemian interglacial reconstructed from a Greenland folded ice core, *Nature*, 493, 489–494, doi:10.1038/nature11789, 2013.
- Nishiizumi, K.: Preparation of  $^{26}\text{Al}$  AMS standards, *Nucl. Instrum. Meth. Phys. Res. Sect. B.*, 223–224, 388–392, doi:10.1016/j.nimb.2004.04.075, 2004.
- 000 Nishiizumi, K., Imamura, M., Caffee, M., Southon, J., Finkel, R., and McAninch, J.: Absolute calibration of  $^{10}\text{Be}$  AMS standards, *Nucl. Instrum. Meth. Phys. Res. Sect. B.* 258, 403–413, doi:10.1016/j.nimb.2007.01.297, 2007.
- Ó Cofaigh, C., Dowdeswell, J.A., Jennings, A.E., Hogan, K.A., Kilfeather, A., Hiemstra, J.F., Noormets, R., Evans, J., McCarthy, D.J., Andrews, J.T., and Moros, M.: An extensive and dynamic ice sheet on the West Greenland shelf during the last glacial cycle, *Geology*, 41, 219-222, doi:10.1130/G33759.1, 2013.
- 005 Pearce, D.M., Mair, D.W.F., Rea, B.R., Lean, J.M., Schofield, E., Kamenos, N., and Schoenrock, K.: The glacial geomorphology of upper Godthåbsfjord (Nuup Kangerlua) in southwest Greenland, *J. Maps.* 14, 45-55, doi:10.1080/17445647.2017.1422447, 2018.
- Pendleton, S.L., Miller, G.H., Lifton, N., Lehman, S.J., Southon, J., Crump, S.E., and Anderson, R.S.: Rapidly receding Arctic Canada glaciers revealing landscapes continuously ice-covered for more than 40,000 years, *Nat. Commun.*, 10, 445, doi:10.1038/s41467-019-08307-w, 2019.
- 010 Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Cheng, H., Edwards, R.L., Friedrich,



- 015 M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W.,  
Muscheler, R., Palmer, J.G., Pearson, C., van der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R.,  
Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich,  
R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S.: INTCAL20  
Northern Hemisphere radiocarbon age calibration curve (0-55 CAL kBP), *Radiocarbon*, 1-33,  
doi:10.1017/RDC.2020.41, 2020.
- 020 Reyes, A.V., Carlson, A.E., Beard, B.L., Hatfield, R.G., Stoner, J.S., Winsor, K., Welke, B., and Ullman, D.J.: South Greenland  
ice-sheet collapse during Marine Isotope Stage 11, *Nature*, 510, 525-528, doi:10.1038/nature13456, 2014.
- Schaefer, J.M., Denton, G.H., Kaplan, M., Putnam, A., Finkel, R.C., Barrell, D.J.A., Andersen, B.G., Schwartz, R.,  
Mackintosh, A., Chinn, T., and Schlüchter, C.: High-frequency Holocene glaciers in New Zealand differ from the  
Northern signature, *Science*, 324, 622-625, doi:10.1126/science.1169312, 2009.
- 025 Schaefer, J.M., Finkel, R.C., Balco, G., Alley, R.B., Caffee, M.W., Briner, J.P., Young, N.E., Gow, A.J., and Schwartz, R.:  
Greenland was nearly ice-free for extended periods during the Pleistocene, *Nature*, 540, 252-255,  
doi:10.1038/nature20146, 2016.
- Seidenkrantz, M.-S., Kuijpers, A., Olsen, J., Pearce, C., Lindblom, S., Ploug, J., Przbyło, P., and Snowball, I.: Southwest  
Greenland shelf glaciation during MIS 4 more extensive than during the Last Glacial Maximum, *Scientific Reports*,  
9, 15617, doi:10.1038/s41598-019-51983-3, 2019.
- 030 Seroussi, H. and Morlighem, M.: Representation of basal melting at the grounding line in ice flow models, *Cryosphere*, 12,  
3085-3096. doi.org/10.5194/tc-12-3085-2018, 2018.
- Siddall, M., Rohling, E.J., Almagi-Labin, A., Hemleben, Ch., Meischner, D., Schmelzer, I., Smeed, D.A.: Sea-level  
fluctuations during the last glacial cycle, *Nature*, 423, 853-858, doi: 10.1038/nature01690.
- 035 Simpson, M.J.R., Milne, G.A., Huybrechts, P., and Long, A.J.: Calibrating a glaciological model of the Greenland ice sheet  
from the Last Glacial Maximum to present-day using field observations of relative sea level and ice extent. *Quat. Sci.  
Rev.*, 28, 1631-1657, doi:10.1016/j.quascirev.2009.03.004, 2009.
- Spector, P., Stone, J., Pollard, D., Hillebrand, T., Lewis, C., and Gombiner, J.: West Antarctic sites for subglacial drilling to  
test for past ice-sheet collapse, *Cryosphere*, 12, 2741-2757, https://doi.org/10.5194/tc-12-2741-2018, 2018.
- 040 Stirling, C.H., Esat, T.M., Lambeck, K., and McCulloch, M.T.: Timing and duration of the Last Interglacial: Evidence for a  
restricted interval of widespread coral reef growth, *Earth Planet. Sci. Lett.*, 160, 745-762, doi:10.1016/S0012-  
821X(98)00125-3, 1998.
- Stone, J.O.: Air pressure and cosmogenic isotope production, *J. Geophys. Res.*, 105, 23,753-23,759,  
doi:10.1029/2000JB900181, 2000.
- Stuiver, M., Reimer, P.J., and Reimer, R.W.: CALIB 8.2 [WWW program] at http://calib.org, accessed 2020-8-13, 2020.
- 045 Thomas, E.K., Castañeda, I.S., McKay, N.P., Briner, J.P., Salacup, J.M., Nguyen, K.Q., and Schweinsberg, A.D.: A wetter




- Arctic coincident with hemispheric warming 8,000 years ago, *Geophys. Res. Lett.*, 45, 10,637-10, 647, doi:10.1029/2018GL079517, 2018.
- Tuna, T., Fagault, Y., Bonvalot, L., Capano, M., Bard, W.: Development of small CO<sub>2</sub> gas measurements with AixMICADAS, *Nucl. Instrum. Methods Phys. Res. Sect. B*, 437, 93–97, doi:10.1016/j.nimb.2018.09.012, 2018.
- 050 Weidick, A.: Quaternary Map of Greenland, 1:500 000, Frederikshåbs Isblink – Søndre Strømfjord. Copenhagen, Geological Survey of Greenland, 1974.
- Weidick, A., Oerter, H., Reeh, N., Thomsen, H.H., and Thorning, L.: The recession of the Inland Ice margin during the Holocene climatic optimum in the Jakobshavn Isfjord area of West Greenland, *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 82, 389-399, doi:10.1016/S0031-0182(12)80010-1, 1990.
- 055 Weidick, A., Bennike, O., Citterio, M., and Nørgaard-Pedersen, N.: Neoglacial and historical glacier changes around Kangarsuneq fjord in southern West Greenland, *Geol. Surv. Den. Green. Bull.*, 27, 68 pp, 2012.
- Young, N.E., and Briner, J.P.: Holocene evolution of the western Greenland Ice Sheet: Assessing geophysical ice-sheet models with geological reconstructions of ice-margin change, *Quat. Sci. Rev.*, 114, 1-17, doi:10.1016/j.quascirev.2015.01.018, 2015.
- 060 Young, N.E., Briner, J.P., Stewart, H.A.M., Axford, Y., Csatho, B., Rood, D.H., and Finkel, R.C.: Response of Jakobshavn Isbræ, Greenland, to Holocene climate change, *Geology*, 39, 131-134, doi:10.1130/G313991, 2011a.
- Young, N.E., Briner, J.P., Axford, Y., Csatho, B., Babonis, G.S., Rood, D.H., and Finkel, R.C.: Response of a marine-terminating Greenland outlet glacier to abrupt cooling 8200 and 9300 years ago, *Geophys. Res. Lett.*, 38, L24701, doi:10.1029/2011GL049639, 2011b.
- 065 Young, N.E., Schaefer, J.M., Briner, J.P., and Goehring, B.M.: A <sup>10</sup>Be production rate calibration for the Arctic, *J. Quat. Sci.*, 28, 515-526, doi:10.1002/jqs.2642, 2013a.
- Young, N.E., Briner, J.P., Rood, D.H., Finkel, R.C., Corbett, L.B., and Bierman, P.R.: Age of the Fjord Stade moraines in the Disko Bugt region, western Greenland, and the 9.3 and 8.2 ka cooling events, *Quat. Sci. Rev.*, 60, 76-90, doi:10.1016/j.quascirev.2012.09.028, 2013b.
- 070 Young, N.E., Schaefer, J.M., Goehring, B., Lifton, N., Schimmelpfennig, I., and Briner, J.P.: West Greenland and global *in situ* <sup>14</sup>C production-rate calibrations, *J. Quat. Sci.*, 29, 401-406, doi:10.1002/jqs.2717, 2014.
- Young, N.E., Briner, J.P., Maurer, J., and Schaefer, J.M.: <sup>10</sup>Be measurements in bedrock constrain erosion beneath the Greenland Ice Sheet margin, *Geophys. Res. Lett.*, 43, doi:10.1002/2016GL070258, 2016.
- Young, N.E., Briner, J.P., Miller, G.H., Lesnek, A.J., Crump, S.E., Thomas, E.K. Pendleton, S.L., Cuzzone, J., Lamp, J., Zimmerman, S., Caffee, M., and Schaefer, J.M.: Deglaciation of the Greenland and Laurentide ice sheets interrupted by glacier advance during abrupt coolings, *Quat. Sci. Rev.*, 229, 106091, doi:10.1016/j.quascirev.2019.10609, 2020a.
- 075 Young, N.E., Briner, J.P., Miller, G.H., Lesnek, A.J., Crump, S.E., Thomas, E.K. Pendleton, S.L., Cuzzone, J., Lamp, J.,





080

Zimmerman, S., Caffee, M., and Schaefer, J.M.: Reply to Carlson (2020) comment on “Deglaciation of the Greenland and Laurentide ice sheets interrupted by glacier advance during abrupt coolings”, *Quat. Sci. Rev.*, 240, 106329, doi:10.1016/j.quascirev.2020.106329, 2020b.




085

Figure 1: (a) Southwestern Greenland with the locations of proglacial-threshold lakes discussed in the text. Map data: Google, Landsat, U.S. Geological Survey. jaks = Jakobshavn Isbræ; new = Newspaper Lake (Cronauer et al., 2015); lo = Lake Lo (Håkansson et al., 2014); sov = South Oval Lake, il = Ice Boom Lake, mer = Merganzer Lake, rav = Raven Lake, goo = Goose Lake, loo = Loon Lake, et = Eqaqut taserussuat (Briner et al., 2010); kt = Kuussuup Tasia, tv = Tininnillik Valley (Kelley et al., 2012); fh = Four Hare Lake (Lesnek et al., 2020); luc = Lake Lucy, spo = Sports Lake (Young and Briner, 2015); bl = Baby Loon Lake, ts = Tasersuaq (Lesnek et al., 2020); 09379, fl = Frederikshåb Isblink (Larsen et al., 2015). (b) Nuuk (N) region with existing radiocarbon ages (red dots) and <sup>10</sup>Be ages (yellow dots) in ka BP (CE 1950) that constrain the timing of regional deglaciation. The Kapisigdlit stade moraines are shown in pink and historical moraines are in blue (modified from Pearce et al., 2018). For figure clarity we only show the mean deglaciation age at each location without uncertainties; see Table S1 (radiocarbon) and Table S2 (<sup>10</sup>Be) for details. Also shown are the locations of proglacial-threshold lakes discussed in the text (orange dots) and location of marine bivalves reworked into the historical maximum in Fig. 17. is1, is9, is12, (Levy et al., 2017); car = Caribou Lake, gf = Goose Feather Lake (Lesnek et al., 2020); mr = Marshall Lake (this study); dec = Deception Lake, ow = One-way Lake (Lesnek et al., 2020); Kan01 (Larsen et al., 2015); Kap01 (Larsen et al., 2014 and this study; b = bulk sediments; m = macrofossils). Glaciers: kns = Kangiata Nunaata Sermia; kas = Kangaasarsuup Sermia; aks = Akullersuup Sermia; qms = Qamanaarsuup Sermia; nrs = Narsap Sermia; sqs = Saqqap Sermia.

090

095




100

105


110





115

**Figure 3:** (a) Qamanaarsuup Sermia region depicting the Kapisigdlit stage moraines (pink) and retreat since from the historical maximum extent (brown shading); moraines were mapped in the field.  $^{10}\text{Be}$  ages (ka BP  $\pm$  1 S.D.) are from three morphostratigraphic groups: 1) erratic boulders perched on bedrock outboard the Kapisigdlit stage moraines (black symbols and text), 2) Kapisigdlit stage moraine boulders (pink symbols and text), and 3) erratic boulders perched on bedrock inside the Kapisigdlit stage moraines (blue symbols and text). Also shown are the sediment coring locations in Goose Feather Lake (17GOOF) and Marshall Lake (17MAR). Dashed blue line marks the route of meltwater from the GrIS to the Goose Feather Lake inflow (I). The outflow (O) for Goose Feather Lake routes meltwater back towards the GrIS. Map data: Google, Maxar Technologies. (b) Normal density estimates for the Kapisigdlit stage moraines from panel (a). Age in bold includes the production-rate uncertainty.



120

125


**Figure 4:** (a) View to the northeast in the Qamanaarsuup Sermia region (Fig. 3). In the foreground is a Kapisigdlit stade moraine crest resting directly adjacent to the historical maximum extent of this sector of the GrIS (dashed line). The GrIS is in the background and, here, there has been minimal retreat from the historical maximum extent. (b) View to the southwest in the Qamanaarsuup Sermia region showing Marshall Lake and Goose Feather Lake. Note the color contrast between the two lakes. Marshall Lake currently does not receive meltwater from the GrIS. Goose Feather Lake is currently a proglacial lake that receives silt-laden GrIS meltwater; the lake catchment currently extends beneath the modern GrIS footprint (Fig. 3a; Fig. 6).



130

**Figure 5: Representative boulder samples from the Qamanaarsuup Sermia region (Fig. 3). Samples 17GRO-16, -32, and -39 are moraine boulder samples. 17GRO-25 is an erratic boulder perched on bedrock inside the Kapisigdlit stade moraine and located only a few meters outboard of the historical maximum moraine, which can be seen in the background. Samples 17GRO-46 and -47 are erratic boulders resting on bedrock located outboard of the Kapisigdlit stade moraine.**






135

140

**Figure 6:** (a) Sediment cores and calibrated radiocarbon ages ( $\pm 2\sigma$ ) from Marshall (MAR) and Goose Feather (GOOF) lakes. Note the distinct color differences between silt and organic sediments (see Fig. 3 and Fig. 4). Details for radiocarbon ages can be found in Table S7. (b) Sub-ice topography in the Qamanaarsuup Sermia region generated using the BedMachine v3 DEM (Mortlighem et al., 2017) compared to our chronology of ice-margin change developed from  $^{10}\text{Be}$  ages and radiocarbon-dated lake sediments (panel A). Shown is the average  $^{10}\text{Be}$  age from each feature on the landscape (including production rate uncertainty; see Fig. 3) and basal radiocarbon ages from panel (a). Deglaciation of the landscape just outboard of the Kapisigdlit stade moraines occurred at  $10.29 \pm 0.23$  ka BP followed by moraine deposition at  $9.57 \pm 0.38$  ka BP. Ice retreated behind the modern margin at  $9.29 \pm 0.18$  ka BP, but the ice margin remained within the drainage catchment of Goose Feather Lake until  $8.28 \pm 0.09$  cal ka BP before retreating farther inland. The dashed line delimits the topographic threshold (T) under the modern GrIS that the ice margin must cross in order for Goose Feather Lake to receive silt-laden meltwater. Inflow of meltwater ceases when the GrIS margin retreats behind this topographic threshold, which rests ~1 km behind the modern margin.



145

Figure 7: Kapisigdlit stade moraine located just west of KNS (Fig. 2).  $^{10}\text{Be}$  ages (ka BP  $\pm$  1 S.D.) are from three morphostratigraphic groups: 1) outboard of the moraine (black text and symbols), 2) moraine boulders (pink text and symbols), and 3) inside of the moraine (blue text and symbols).  $^{10}\text{Be}$  ages that are considered outliers are in italics. Map data: Google, Maxar Technologies.

150




Figure 8: Representative boulder samples related to the Kapisigdlit stade moraine west of KNS. 17GRO-64 and -66 are moraine boulders, whereas 17GRO-67 and -70 are erratic boulders resting on bedrock located immediately inboard and outboard the Kapisigdlit stade moraine.





155



160


Figure 9: (a) Oblique aerial view to the northwest depicting recently deglaciated bedrock that rests between the historical maximum extent and the modern ice margin on the northeastern side of KNS. (b)  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ages (ka BP ± 1 S.D.), along with measured  $^{26}\text{Al}/^{10}\text{Be}$  ratios at each bedrock sample site.  $^{10}\text{Be}$  ages influenced by inheritance are in italics. The historical maximum extent of KNS is marked in blue. To orient the reader, L and B mark the same feature in each panel. Panel (b) map data: Google, Maxar Technologies.





165


Figure 10: Sampled bedrock surfaces located between the historical maximum extent of the GrIS and the modern ice margin on the northeastern side of KNS.




170

Figure 11: Sampled bedrock surfaces adjacent to the modern ice margin on the northeastern side of KNS.






175 Figure 12:  $^{10}\text{Be}$  ages and *in situ*  $^{14}\text{C}$  ages (ka BP  $\pm$  1 S.D.), and measured  $^{26}\text{Al}/^{10}\text{Be}$  ratios at each bedrock sample site on the southwestern side of KNS. Also shown are representative bedrock sample sites 17GRO-72 and -74. Map data: Google, Maxar Technologies.




180 Figure 13. (a) <sup>10</sup>Be-<sup>14</sup>C apparent exposure ages from the northeastern and southwestern sides of KNS plotted against sample  
 elevation. Apparent <sup>10</sup>Be ages that are older than ~10.3 ka BP are influenced by isotopic inheritance, yet their corresponding *in*  
*situ* <sup>14</sup>C ages are younger and consistent with Holocene-only exposure histories. *In situ* <sup>14</sup>C ages versus sample elevation are  
 consistent with ice-margin thinning. (b) Paired <sup>14</sup>C-<sup>10</sup>Be diagram. The x-axis is the measured <sup>10</sup>Be concentration normalized by the  
 185 regionally constrained <sup>14</sup>C/<sup>10</sup>Be spallation production ratio of 3.12 (Young et al., 2014). The simple exposure region (black lines) is  
 defined by the continuous and steady-state erosion lines. Samples that are influenced by isotopic inheritance are excluded because  
 they yield artificially low and meaningless ratios. Remaining samples are all consistent with constant exposure; we estimate a  
 minimum burial detection limit of 625 years.



190 Figure 14: Modeled  $^{10}\text{Be}$  (black line) and *in situ*  $^{14}\text{C}$  (green line) production with depth in bedrock using a typical rock density of  $2.65\text{ g cm}^{-3}$  for gneiss. A normalized  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  concentration of 1 equals a prescribed exposure duration of 10 kyr, 9 kyr, 8 kyr, or 7.5 kyr. Symbols mark the depth below the prescribed deglacial surface along the production-depth profile that the GRIS would have to erode to during the period of historical ice cover that results in our measured nuclide concentrations. The  $^{10}\text{Be}$  production-depth profile is not fit to the data points and any  $^{10}\text{Be}$  measurement will fall somewhere along the depth profile. Bars on the left ( $^{10}\text{Be}$ ) and right ( $^{14}\text{C}$ ) sides of the diagram are the average erosional depths ( $\pm 1$  S.D.) of each exposure-erosion scenario (see Table S9). For figure clarity, we only show where our individual  $^{10}\text{Be}$  measurements fall along the production-depth profile; the average erosional depth based on the *in situ*  $^{14}\text{C}$  measurements are on the right side of the figure (Table S9). Production by spallation and muons are calculated independently; muons according to Balco et al (2017; Model 1A).  $Z$  is the mass depth below the surface expressed in  $\text{g cm}^{-2}$  and is the product of depth (cm) and material density ( $\text{g cm}^{-3}$ ).  $P_i$  is the production rate of  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  ( $\text{atoms g}^{-1}\text{ yr}^{-1}$ ) by spallation or muons at depth  $z$ .  $P_i(0)$  is the surface production rate via spallation or muons.  $\Lambda$  is the effective attenuation length ( $\text{g cm}^{-2}$ ). Included here are only  $^{10}\text{Be}$  and *in situ*  $^{14}\text{C}$  concentrations for samples with concordant apparent exposure ages of  $\sim 6\text{--}7$  ka BP (17GRO-71, -72, -73, -74, and -75; Tables S6 and S9).

195


200



205


**Figure 15:** Paired  $^{26}\text{Al}$ - $^{10}\text{Be}$  diagram with all measurements from the KNS forefield ( $n=12$ ). The x-axis is the measured  $^{10}\text{Be}$  concentration normalized by the site-specific production rate (years); the y-axis is the measured  $^{26}\text{Al}/^{10}\text{Be}$  ratio normalized to the production ratio. We use a production ratio of 7.07 (Argento et al., 2013). The black lines define the simple exposure region. All of our measured ratios, with the exception of 17GRO-13, indicate constant exposure of the sample site (values listed in Table S5). Because of the close proximity of these sites, they must have the same glacial-interglacial exposure history and thus we consider 17GRO-13 a likely outlier.

210



215 Figure 16. (a) Cosmogenic nuclide and  $^{14}\text{C}$ -dated proglacial-threshold lake records constraining the behavior of the southwestern  
 220 GrIS ice margin. Site abbreviations are from Fig. 1. Using sedimentological boundaries in proglacial-threshold lakes (organic  
 sediments versus silt), the orange bars mark periods of time at each site that the GrIS was less extensive than today (organic  
 225 sediments), and blue bars marks periods of time when the ice margin was in a position similar to today (silt). Orange dots are  
 traditional radiocarbon ages from the contact between basal silt and the overlying organic sediments that constrain when the  
 southwestern GrIS margin retreated out of each lake's catchment in the early to middle Holocene. Blue bullseyes are  $^{10}\text{Be}$  ages from  
 230 immediately outboard of the historical maximum extent in each proglacial-threshold lake drainage catchment, and purple bullseyes  
 are additional  $^{10}\text{Be}$  ages from within  $\sim 1$  km of the historical maximum extent. Black bullseyes define our estimated timing of inland  
 retreat of KNS based on *in situ*  $^{14}\text{C}$  measurements (section 5.2) Orange diamonds are traditional  $^{14}\text{C}$  ages from marine bivalves  
 reworked into the historical limit at Ujarassuit Paavat (up; Fig. 1), which mark times when the ice margin was behind the historical  
 maximum extent. Combined,  $^{10}\text{Be}$  from immediately outboard the historical maximum limit,  $^{10}\text{Be}$  ages from within  $\sim 1$  km of the  
 historical maximum limit, and basal  $^{14}\text{C}$  ages from threshold lakes, define a window when the ice margin was in a position similar  
 to today. Considering all constraints, the southwestern GrIS margin likely achieved its minimum extent after  $\sim 5$  ka, and was  
 approaching its modern position as early as  $\sim 2$  ka BP (blue shading). (b) Mean-annual temperatures at the Greenland Ice Sheet  
 Project 2 site (GISP2) reconstructed using gas-phase  $\delta\text{Ar-N}_2$  measurements ( $\pm 2\sigma$ ; Kobashi et al., 2017). (c) Temperature anomalies  
 over southwestern Greenland used in recent ISSM modeling runs (Badgeley et al., 2020; Briner et al., in press). (d) Corresponding  
 modeled ice mass for the southwestern GrIS for nine simulations using different climate reconstructions (Briner et al., in press);  
 note the reversed y-axis.





235

240

Figure 17: (a) Area-averaged mean-annual temperature using three different reconstructions across the ISSM model domain (Badgeley et al., 2020; Briner et al., in press). (b) Same as (a), but for mean-annual precipitation (Badgeley et al., 2020; Briner et al., in press). (c) Simulated margin position in the Sisimiut-Kangerlussuaq region through the Holocene compared to independent observations of ice-margin position based on dated moraines (black dots;  $1\sigma$  age uncertainty; Young et al., 2020a) and the present-day ice margin (orange dot; modified from Briner et al., in press). Letters next to each simulation mark the temperature and precipitation reconstructions from panels (a) and (b). (d) Simulated margin position in the KNS region through the Holocene compared to independent observations of ice-margin position based on deglaciation ages (see Fig. 1 for age constraints and up-fjord transect). (e) Simulated ice mass through the Holocene across the model domain (Briner et al., in press); note the y-axis is reversed. (f). Simulated lateral ice-margin extent at the time of minimum ice mass in each model run. The year (BP) of the ice-mass minimum in each model run is listed in the legend.