

1 **Deoxygenation dynamics on the western Nile deep-sea fan during**  
2 **sapropel S1 from seasonal to millennial time-scales**

3 Cécile L. Blanchet<sup>1</sup>, Rik Tjallingii<sup>1</sup>, Anja M. Schleicher<sup>2</sup>, Stefan Schouten<sup>3,4</sup>, Martin Frank<sup>5</sup>, Achim  
4 Brauer<sup>1</sup>

5 <sup>1</sup> GFZ German Research Centre for Geosciences Potsdam, Section of Climate and Landscape Dynamics,  
6 Telegrafenberg, 14743 Potsdam, Germany

7 <sup>2</sup> GFZ German Research Centre for Geosciences Potsdam, Section of Inorganic and Isotope Geochemistry,  
8 Telegrafenberg, 14743 Potsdam, Germany

9 <sup>3</sup> Royal NIOZ, Department of Marine Microbiology and Biogeochemistry, Landsdiep 4, 1797 SZ 't Horntje, Texel,  
10 The Netherlands

11 <sup>4</sup> University of Utrecht, Department of Earth Sciences, Utrecht, the Netherlands.

12 <sup>5</sup> GEOMAR Helmholtz Centre for Ocean Research Kiel, Research Unit Paleoceanography, Wischhofstrasse 1-3,  
13 D-24148 Kiel, Germany

14 *Correspondence to:* Cécile L. Blanchet ([blanchet@gfz-potsdam.de](mailto:blanchet@gfz-potsdam.de))

15

## 16 **Abstract**

17 Ocean deoxygenation is a rising threat to marine ecosystems and food resources under present climate warming  
18 conditions. Organic-rich sapropel layers deposited in the Mediterranean Sea provide a natural laboratory to study  
19 the processes that have controlled changes in seawater oxygen levels in the recent geological past. Our study is  
20 based on three sediment cores spanning the last 10,000 years and located on a bathymetric transect offshore the  
21 western distributaries of the Nile delta. These cores are partly to continuously laminated in the sections recording  
22 sapropel S1, which is indicative of bottom-water anoxia above the western Nile deep-sea fan. We used a  
23 combination of microfacies analyses and inorganic and organic geochemical measurements to reconstruct changes  
24 in oxygenation conditions at seasonal to millennial time-scales. Millimetre-thick laminations are composed of  
25 detrital, biogenic and chemogenic sublayers reflecting seasonal successions of sedimentation. Dark layers reflect  
26 the deposition of summer floods and two types of light layers correspond to autumn plankton blooms and authigenic  
27 carbonates formed in the water-column during spring-early summer, respectively. The isotopic signature of the  
28 authigenic carbonates suggests permanent anoxic to euxinic bottom waters resulting in high levels of anaerobic  
29 remineralisation of organic matter and highlights their potential to reconstruct seawater chemistry at times when  
30 benthic fauna was absent. Ratios of major elements combined to biomarkers of terrestrial and marine organic matter  
31 and redox-sensitive compounds allow to track changes in terrigenous input, primary productivity and past  
32 deoxygenation dynamics on millennial time-scales. Rapid fluctuations of oxygenation conditions in the upper 700  
33 m water depth occurred above the Nile deep-sea fan between 10 and 6.5 ka BP while deeper cores recorded more  
34 stable anoxic conditions. Synchronous changes in terrigenous input, primary productivity and past oxygenation  
35 dynamics after 6.5 ka BP show that runoff-driven eutrophication played a central role in rapid oxygenation changes  
36 in the south-eastern Levantine Basin. These findings are further supported by other regional records and reveal  
37 time-transgressive changes in oxygenation state driven by rapid changes in primary productivity during a period of  
38 long-term deep-water stagnation.

## 39 **Copyright statement**

40 To be provided by Copernicus.

## 41 **1 Introduction**

42 The present-day Mediterranean Sea is well-oxygenated due to a vigorous thermohaline circulation of intermediate-  
43 and deep-water initiated in the eastern part of the basin (Pinardi et al., 2015; Roether et al., 1996). However,  
44 recurring episodes of water-column stratification and severe oxygen depletion have been recorded in Mediterranean

45 sediments (Cramp and O’Sullivan, 1999; Rossignol-Strick, 1985). These episodes marked by sapropel deposits  
46 were caused by drastic modifications of the sea-surface hydrological balance (Rohling, 1994). Orbital-forced  
47 enhancement of monsoonal precipitation over North Africa led to high freshwater discharge into the Eastern  
48 Mediterranean (mostly through the Nile River) and generated large-scale perturbations of the Mediterranean  
49 thermohaline circulation (Rohling et al., 2015). Seawater freshening is also a well-known perturbator in the North  
50 Atlantic that has led to slowing down or even complete shutting down of Atlantic Meridional Overturning  
51 Circulation during the last glacial period (Ganopolski and Rahmstorf, 2001). In the Mediterranean Sea, the input  
52 of relatively low-saline Atlantic seawater also exerted a large control on deep-water ventilation (Rogerson et al.,  
53 2012). The inflow of Atlantic-derived seawater at the Gibraltar Strait as early as 16-17 ka BP during deglacial sea-  
54 level rise increased the buoyancy of surface waters and led to long-lasting deep-water stagnation in the eastern  
55 Mediterranean (Rogerson et al., 2010), which is seen as a pre-condition for the development of basin-scale anoxia  
56 during sapropel S1 (11-6 ka BP) (Cornuault et al., 2018; Grimm et al., 2015).

57 A specificity of the eastern Mediterranean lies in its current nutrient-limitation and the consequently low primary  
58 productivity (Pujo-Pay et al., 2011). Nutrient-poor surface waters are advected from the western to the eastern  
59 Mediterranean and riverine nutrients (e.g. discharged by the Nile River) are quickly utilized or extracted by  
60 intermediate water circulation flowing westward (Fig. 1a). Prior to the construction of the Aswan Dam in 1965, the  
61 surface waters in the Levantine Basin were seasonally fertilised by riverine nutrient input during the summer Nile  
62 floods (Halim et al., 1967). These so-called “Nile Blooms” of phytoplankton sustained traditional fisheries of  
63 sardines and prawns (Nixon, 2003). After 1965, the Nile River discharge was reduced by 90% and thereby largely  
64 annihilated the main nutrient source to the eastern Mediterranean, initiating the present-day ultra-oligotrophic  
65 conditions. Since the 1980s however, the transition from a flood-sustained to a fertiliser-sustained agriculture in  
66 the Nile valley has led to a larger release of nutrients by the Nile River, which boosted primary productivity and  
67 renewed the fish stocks (Nixon, 2003). Anthropogenic fertilisation processes in coastal waters might have potential  
68 adverse effects as well, such as water deoxygenation or harmful algal blooms (Nixon, 2003). Eutrophication and  
69 severe anoxia due to anthropogenic fertilisation have been indeed observed in semi-enclosed settings such as the  
70 Baltic Sea (Jilbert and Slomp, 2013). In this context of increasing nutrient loading, it is crucial to estimate the  
71 changes in intermediate and deep-water ventilation caused by climate change. Modelling experiments predict a  
72 temperature and salinity rise for the Mediterranean over the next decades, which overall will result in a weakening  
73 or a change in configuration of the thermohaline circulation in the Mediterranean (e.g., Adloff et al., 2015).  
74 Sensitivity simulations showed that reduced deep ventilation leading to basin-wide hypoxia might occur in the  
75 Mediterranean over long time scales (>1000 yr), but is unlikely over shorter time scales (ca. 100 yr) (Powley et al.,


76 2016). However, the predicted changes in O<sub>2</sub> distributions do not account for additional factors that may affect O<sub>2</sub>  
77 consumption rates, in particular, rising inputs of anthropogenic nutrients or oxygen loss due to rising sea  
78 temperature (i.e., decreased oxygen solubility in warmer waters) (Keeling et al., 2009; Powley et al., 2016).  
79 Sapropels provide a natural laboratory to investigate the complex interplay between oceanographic, hydrological  
80 and biogeochemical processes driving Mediterranean deoxygenation (Rohling et al., 2015; Rossignol-Strick et al.,  
81 1982). Here we present a set of unique sediment cores retrieved along a bathymetric section near the main western  
82 Nile tributary, the Rosetta channel, to investigate the relationships between terrigenous inputs, primary  
83 productivity, and the development of sub- to anoxic conditions in intermediate water masses. The location of our  
84 study site close to the mouth of the Nile River allows to reconstruct changes in fluvial nutrient supply and associated  
85 fertilisation of the marine ecosystem.

## 86 **2 Regional context**

87 Surface waters originating from the North Atlantic flow into the Levantine Basin (Pinardi et al., 2015), which  
88 gradually become more saline due to evaporation as they flow eastwards in the Mediterranean Sea (Fig. 1a).  
89 Elevated salinity of surface waters in the eastern Levantine Basin (ca. 39 PSU) (MEDAR Group, 2002) in  
90 combination with cooler temperatures in winter lead to high surface water density and formation of the Levantine  
91 Intermediate Water (LIW), which flows westwards at water depths between 200 and 600 m (Cornuault et al., 2018;  
92 Pinardi et al., 2015). Deeper water masses flow below the LIW and form by incorporation of cold surface waters  
93 from the Adriatic Sea and sometimes dense waters from the Aegean Sea to the LIW (Cornuault et al., 2018;  
94 Robinson et al., 1992; Roether et al., 2007). According to recent results, deep convection is also occurring in the  
95 northern Levantine Basin, resulting in the formation of Levantine Deep Water (LDW) that ventilates the whole  
96 Levantine Basin (Kubin et al., 2019). At present, the LDW bathing the core sites is well oxygenated due to high  
97 rates of LDW formation and low primary productivity (Powley et al., 2016).

98 The main source of freshwater to the Levantine Basin is the Nile River runoff through the active Rosetta channel  
99 located in the western part of the Nile Delta (Fig. 1a, b). Although the present Nile flow is drastically reduced  
100 compared to historical times (Halim et al., 1967), runoff still leads to marked salinity gradients (halocline) in the  
101 upper 200 m water-depth (w-d), in particular during the summer monsoon (Fig. 1a). The plume of Nile-derived  
102 freshwater is deflected to the East by the long-shore surface current called “Levantine Jet” (Fig. 1b) (Pinardi et al.,  
103 2015).

104



105  
 106 **Figure 1. Regional context and location of the cores.** (a) Sea-surface salinity during the summer months (June to September)  
 107 (MedAtlas, MEDAR Group, 2002) plotted with the Ocean Data View software (Schlitzer, 2018). White star: location of the Nile  
 108 River freshwater plume. Black arrows: surface currents ((Pinardi et al., 2015; Rohling et al., 2015). (b) Zoom on the western Nile  
 109 deep-sea fan (DSF), with locations of cores P33, P73 and P99 (this study), as well as neighbouring cores MS27PT and MD04-2726  
 110 (Revel et al., 2015), MD04-2722 (Tachikawa et al., 2015), GeoB7702-3 (Castañeda et al., 2010), PS009PC (Hennekam et al., 2014),  
 111 MD9509 (Matthews et al., 2017) (black diamonds) and the Amon mud volcano (black triangle). White contours: 500-m and 1000-  
 112 m isobaths and grey contours: 1500-m and 2000-m isobaths. (c) Bathymetric transect across the western Nile DSF with sediment  
 113 lithological features (black: laminations and stripes: faint laminations). Map and bathymetric transect in (a) and (b) plotted using  
 114 the R freeware (R Core Team, 2013), the marmap package (Pante and Simon-Bouhet, 2013) and gridded bathymetric data from  
 115 GebcO (Gebco team, 2014).

### 116 3 Material and methods

#### 117 3.1 Sediment cores

118 **Table 1. Sediment cores used for the present study, with names used throughout the paper, water-depths and coordinates.**

Core Name	Shortened	Water-depth (m)	Latitude (dec. deg.)	Longitude (dec. deg.)
P362/2-33	P33	738	31.700	29.768
P362/2-73	P73	569	31.987	30.156
P362/2-99	P99	396	31.953	30.160

119  
 120 For our study, we use three gravity cores that were retrieved during the oceanographic campaign P362/2 “West  
 121 Nile Delta” onboard the R.V. Poseidon in August 2008 (Feseker et al., 2010) (Table 1). These sediment cores are  
 122 located along a bathymetric section on the western Nile deep-sea fan (DSF), in the vicinity of the North Alex and  
 123 Giza mud volcanoes (Feseker et al., 2010) (Fig. 1b,c). Cores P362/2-73 (P73) and P362/2-99 (P99) are located

124 close to each other and ca. 75 km eastward of core P362/2-33 (P33), on the opposite side of the submarine Rosetta  
 125 canyon (Fig. 1b). Core P33 was recovered at 738 m water-depth, is 5.6 m-long and contains well-preserved mm-  
 126 scale laminations in the lower 5 m of the core (see Blanchet et al., 2013, 2014, for detailed descriptions) (Fig. 2a).  
 127 Core P73 was collected at 569 m depth, is 5.4 m-long and contains an alternation of finely laminated (similar to  
 128 P33) and bioturbated intervals (Fig. 2b). Core P99 is located at 396 m depth, is 5.3 m-long and contains a few  
 129 laminated intervals within otherwise bioturbated sediments (Fig. 2c). All cores are composed of clay-rich dark-  
 130 brown hemipelagic mud, which shows significantly lighter colours in the upper 1m of the cores. In cores P33 and  
 131 P73, several harder carbonate-rich layers were identified (Fig. 2a, b). A quantitative assessment of the number of  
 132 benthic foraminifera per gram of sediment was realised in the lower part of core P33 (between 75 and 558 cm).

133

134 **Table 2: Radiocarbon ages, tie-points and modelled ages for cores P33, P73 and P99. Radiocarbon ages for core P33 from Blanchet**  
 135 **et al. (2013), with new modelled ages with uncertainties for the entire core (using Bacon V2.3, Blaauw and Christen, 2011). Six tie**  
 136 **points were defined on the Ti/Ca record of P33 (T1 to T6, see sect. 3.2 and Fig. 2). New radiocarbon measurements, tie-points and**  
 137 **modelled ages with uncertainties for cores P73 and P99. In the column “Material”, “Pk foraminifera” and “Bk foraminifera” refer**  
 138 **to planktonic and benthic foraminifera, respectively.**

Sample name	Depth (cm)	Material	14C age $\pm 1\sigma$	C cont. (mg)	DeltaR $\pm 1\sigma$	Calib curve	cal. age BP $\pm 2\sigma$	modelled median age (Bacon)
<b>P362/2-33 (P33)</b>								
	1.5 $\pm$ 1.5	Steamboat cinders					75 $\pm$ 75	161 $\pm$ 210
KIA 38572	30 $\pm$ 0.5	Pk foraminifera	3825 $\pm$ 30	1.3	0	Marine13	3775 $\pm$ 100	3840 $\pm$ 180
KIA 38573	50 $\pm$ 0.5	Pk foraminifera	5695 $\pm$ 30	1.3	0	Marine13	6110 $\pm$ 100	6140 $\pm$ 150
KIA 38574	75 $\pm$ 0.5	Pk foraminifera	6855 $\pm$ 35	1.6	150 $\pm$ 30	Marine13	7235 $\pm$ 105	7260 $\pm$ 180
KIA 37800	100 $\pm$ 0.5	Pk foraminifera	7340 $\pm$ 45	1.2	150 $\pm$ 30	Marine13	7655 $\pm$ 110	7825 $\pm$ 150
KIA 38575	152 $\pm$ 2.5	Pk foraminifera	7920 $\pm$ 45	1.3	150 $\pm$ 30	Marine13	8240 $\pm$ 120	8165 $\pm$ 130
KIA 37801	200 $\pm$ 0.5	Pk foraminifera	8115 $\pm$ 55	1.1	150 $\pm$ 30	Marine13	8430 $\pm$ 125	8405 $\pm$ 120
KIA 38576	252 $\pm$ 2.5	Pk foraminifera	8360 $\pm$ 40	1.1	150 $\pm$ 30	Marine13	8725 $\pm$ 180	8610 $\pm$ 115
KIA 37802	300 $\pm$ 0.5	Pk foraminifera	8375 $\pm$ 60	0.9	150 $\pm$ 30	Marine13	8750 $\pm$ 200	8765 $\pm$ 110
KIA 38577	327 $\pm$ 3	Pk foraminifera	8495 $\pm$ 35	1.3	150 $\pm$ 30	Marine13	8925 $\pm$ 160	8850 $\pm$ 110
KIA 37803	399 $\pm$ 1	Pk foraminifera	8440 $\pm$ 45	1.4	150 $\pm$ 30	Marine13	8845 $\pm$ 175	9050 $\pm$ 105
KIA 38578	463 $\pm$ 2	Pk foraminifera	8680 $\pm$ 40	1	150 $\pm$ 30	Marine13	9165 $\pm$ 150	9280 $\pm$ 100
KIA 37804	499 $\pm$ 1	Pk foraminifera	8805 $\pm$ 45	1.5	150 $\pm$ 30	Marine13	9330 $\pm$ 145	9410 $\pm$ 100
KIA 38579	547 $\pm$ 3	Pk foraminifera	9010 $\pm$ 35	1.2	150 $\pm$ 30	Marine13	9515 $\pm$ 100	9595 $\pm$ 125
KIA 37805	557 $\pm$ 1	Pk foraminifera	8780 $\pm$ 50	0.7	150 $\pm$ 30	Marine13	9300 $\pm$ 160	9635 $\pm$ 135

T1	8							1000±440
T2	17.5							2250±520
T3	31.5							4030±150
T4	48							5965±140
T5	80.5							7385±110
<b>P362/2-73 (P73)</b>								
Poz-113950	15 ±1	Pk foraminifera + pteropod	505±30	0.9	0	Marine13	130 ±90	80 ±180
T1	18.5						1000±440	265 ±335
T2	47						2250±520	1665±535
Poz-112951	74 ±1	Pk foraminifera	2925±35	0.7	0	Marine13	2710±85	2720±170
T3	85.5						4030±150	3785±500
T5	172						7385±110	7140± 27
Poz-112952	189 ±1	Pk foraminifera	6860±50	>1	150±30	Marine13	7240±130	737 ±170
Poz-113951	473 ±1	Pk + Bk foraminifera	13910±180	0.04	Too little carbon retrieved			
T6	268.5						7700±115	7810±195
<b>P362/2-99 (P99)</b>								
Poz-112947	11 ± 1	Pk foraminifera	415 ±30	0.8	0	Post-bomb curve NHZ2	80 ±15	40 ±135
T1	31.5						1000±440	1150±615
T2	54.5						2250±520	2590±800
T3	73.5						4030±150	4255±920
Poz-112948	79 ±1	Pk foraminifera	5990±50	>1	0	Marine13	6400±120	5250±960
T4	91						5965±140	6155±620
T5	112						7385±110	7170±285
Poz-112949	322 ±2	Pk foraminifera	7890±60	0.5	150±30	Marine13	8210±150	8200±190
Poz-112950	518 ±1	Pteropod	8390±60	>1	150±30	Marine13	8770±200	9070±275

139

### 140 3.2 Age determination and transformation depth-to-age

141 The chronology of core P33 has been constructed based on a set of 14 radiocarbon samples (Blanchet et al., 2013).

142 We prepared another 8 samples for radiocarbon dating using planktonic foraminifera and pteropod shells, which

143 provide tie-points for cores P73 and P99 (Table 2). Only one sample in core P73 (Poz-113951) did not generate

144 enough carbon for an accurate dating. The new radiocarbon measurements were performed at the Poznan  
145 radiocarbon laboratory (Poland).

146 More detailed stratigraphic constrains were obtained from correlation of the titanium over calcium (Ti/Ca) records  
147 of cores P73 and P99 with core P33 (Fig. 2). It was shown that changes in sedimentation rates are coherent on the  
148 western Nile DSF (Hennekam et al., 2015) and Ti/Ca records show similar patterns (Fig. S1). Six tie-points (marked  
149 T1-T6 on Fig. 2) mark changes in Ti/Ca records identified in the upper parts of the cores and were used to further  
150 synchronise the records of P73 and P99 with core P33 (Table 2).

151 The age-depth modelling was performed using Bacon version 2.3 (Blaauw and Christen, 2011), which enables to  
152 discriminate between sections of the core with contrasting accumulation rates and to provide these as priors.  
153 Convergence and mixing of the Markov chain Monte Carlo iterations used to build the age model by Bacon was  
154 tested and the number of iterations was adjusted to obtain a Gelman and Rubin Reduction factor below 1.05  
155 (Blaauw and Christen, 2011; Brooks and Gelman, 1998). The new age-depth model of core P33 based on  
156 radiocarbon ages is similar to that presented in Blanchet et al. (2013), but age uncertainties are now available for  
157 the whole core (Table 2 and Fig. 2a). Radiocarbon ages and the six tie-points were used for age modelling for cores  
158 P73 and P99 using the Bacon program following a similar procedure to that used for core P33 (Fig. 2b, c) (Blaauw  
159 and Christen, 2011). For the lowest part of core P73, for which there is no tie-point, we evaluated the ages and  
160 uncertainties using the range of sedimentation rates observed in the other two cores for ages older than 7500 years  
161 and derived the median sedimentation rate applying the maximum and minimum sedimentation rates observed as  
162 uncertainty range (Fig. 2b). Age uncertainties range from 100 to 1000 yrs in core P33 (larger in the last 2000 yrs),  
163 from 100 to 700 yrs in P73 (larger in the last 6000 yrs) and 10 to 1000 kyr in P99 (larger in the last 7000 yrs) (Fig.  
164 2).

### 165 **3.3 Microfacies description and scanning electron microscopy**


166 A detailed examination of the microfacies was performed on the laminated part of core P33 (i.e., between 100 and  
167 559 cm) and on a few samples of cores P73 and P99. Sediment blocks of 10 cm-long, 2 cm-wide and 1 cm-thick  
168 were cut out of the fresh sediment with 2-cm overlaps to enable continuous microfacies analysis. Preparation of  
169 thin-sections from soft and wet sediment blocks followed a standard procedure minimizing process-induced  
170 disturbances of sediment micro-structures and included shock-freezing with liquid nitrogen, freeze-drying for 48  
171 h, and epoxy resin impregnation under vacuum (Brauer and Casanova, 2001).

172 Detailed microfacies analysis was performed on the overlapping series of large-scale petrographic thin sections.  
173 Microscopic analysis included the investigation of sediment using a petrographic microscope with non-polarized



174 and (cross)-polarized light, at 5x–40x magnifications (Carl Zeiss Axioplan) (Fig. 3a). Backscatter scanning electron  
 175 microscope (SEM) images and energy dispersive spectroscopy (EDS) elemental analyses were acquired at GFZ  
 176 Potsdam using a Phenom™ XL (Fig. 3b). SEM and EDS analyses were performed on dried powder samples mounted  
 177 on a graphite coated sample holder.

178



179

180 **Figure 2. Age models. (a) core P33, (b) core P73 and (c) core P99. For all cores, from left to right: core depth (m), half-core surface**  
 181 **images (for core P99, realised after sampling with empty sections shown as crosses), section number (roman numbers), schematic**  
 182 **lithological log (white: bioturbated, dashed: faint laminations, hatched: laminations, orange: hard layers labelled HL), linear**  
 183 **sedimentation rates (cm/ka, step curves), age-depth relationships as computed by Bacon V2.3 (Blauw and Christen, 2011) with**  
 184 **radiocarbon ages (blue violin distributions) and tie points (orange dots), Log(Ti/Ca) records with tie-points T1-T6 (dashed lines)**  
 185 **(Tab. 2).**

### 186 3.4 Mineralogy

187 The mineralogical composition was determined by X-ray diffractometry (XRD) on random powder samples of core  
 188 P33 (Fig. 3c). Therefore, the rock chips were powdered to a grain size of < 63 μm, and loaded from the back side  
 189 of the sample holders. XRD analyses were performed with a PANalytical Empyrean x-ray diffractometer operating  
 190 with Bragg-Brentano geometry at 40 mA and 40 kV with CuKα radiation and PIXel3D detector at a step size of

191 0.013 °2θ from 4.6 to 85 °2θ and 60 sec per step. The Mineralogy was determined qualitatively with the EVA  
192 software (version 11.0.0.3) by Bruker.

### 193 **3.5 X-Ray Fluorescence scanning**

194 The bulk sediment compositions for cores P33 and P99 were measured using an Avaatech™ X-Ray fluorescence  
195 (XRF) core-scanner at the Institute of Geosciences of the University of Kiel (Germany) (Table S1 and S2). Non-  
196 destructive XRF core-scanning measurements were performed every 1 cm using a Rh X-Ray source at 10 kV and  
197 0.65 mA to acquire the elements S, Cl, K, Ca and Ti. Core P73 was measured using an ITRAX XRF core-scanner  
198 at GFZ Potsdam (Germany) (Table S3). These measurements were conducted every 1 cm with a Cr- X-Ray source  
199 operated at 30 kV and 60 mA to cover the same elements as acquired for core P33 and P99. Element intensity  
200 records obtained by XRF core-scanning are susceptible to down-core variations of physical properties, sample  
201 geometry and non-linear matrix effects (Tjallingii et al., 2007). These effects can be minimized by log-ratio  
202 transformation of element intensities, which also provide the most easily interpretable signals of relative changes  
203 in chemical composition (Weltje et al., 2015; Weltje and Tjallingii, 2008).

204 Two epoxy embedded samples of core P33 that are representative for the fine (P33-5 77-87 cm) and thick (P33-2  
205 59-69 cm) laminations have been selected for μ-XRF element mapping (Fig. 4). Measurements are conducted every  
206 50 μm at 50 kV, 600 μA and 50 ms using a Bruker M4 Tornado, which is equipped with a Rh X-ray source in  
207 combination with poly-capillary X-ray optics generating an irradiation spot of 20 μm. Mapping results of the  
208 element K, Ca, and Ti representing solid-state chemical components are presented as normalized element intensities  
209 after initial spectrum deconvolution (Fig. 4). However, elements that occur predominantly in pore fluids (e.g. Cl  
210 and S) are not well preserved in epoxy embedded samples.

### 211 **3.6 Stable oxygen and carbon isotopes**

212 Measurements of stable oxygen and carbon isotopes on core P33 were realized both at GEOMAR and GFZ (Table  
213 S4). Carbonate-rich layers were sampled individually and all samples were subsequently freeze-dried and carefully  
214 ground. In order to dissolve only the carbonates in the powdered samples, we used orthophosphoric acid at 70 °C  
215 in a Kiel II (GEOMAR) or Kiel IV (GFZ-Potsdam) carbonate device. The CO<sub>2</sub> released was analysed using a  
216 Thermo Scientific MAT 252 (GEOMAR) or MAT 253 (GFZ-Potsdam) isotope ratio mass spectrometer (IRMS).  
217 The oxygen and carbon isotopic composition were calibrated against international limestone standard (NBS 19)  
218 and are reported relative to the Vienna Pee Dee Belemnite (VPBD) in the delta notation. Analytical precision of 0.06

219 ‰ for both the  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  measurements were determined from repeated analyses of internal limestone  
220 standards.

### 221 **3.7 Biomarkers**

222 All biomarkers (tetraether lipids, *n*-alkanes and alkenones) were measured at the organic geochemistry laboratory  
223 of NIOZ (Texel, Netherlands) on cores P33 and P99 (Table S5 and S6). The lipids were extracted from 63 sediment  
224 samples with a DIONEX Accelerated Solvent Extractor 200 at the NIOZ using a solvent mixture of 9:1 (v/v)  
225 dichloromethane (DCM)/methanol (MeOH). After addition of internal standards  $\text{C}_{22}$  anti-isoalkane (*n*-alkanes), 10-  
226 nonadecanone (alkenones) and  $\text{C}_{46}$  glycerol trialkyl glycerol tetraether, the total lipid extract was separated into  
227 apolar, ketone and polar fractions using pipette column chromatography loaded with aluminium oxide and the  
228 solvent mixtures 9:1 (v/v) hexane/DCM, 1:1 hexane/DCM and 1:1 DCM/MeOH as eluents, respectively. The  
229 apolar fraction was then separated into saturated hydrocarbon (long-chain odd *n*-alkanes and the  $\text{C}_{22}$  anti-iso  
230 standard) and aromatic fractions using pipette columns loaded with  $\text{Ag}^+$ -impregnated silica and hexane and  
231 ethylacetate as eluents, respectively.


232 Molecular identification of the alkenones and *n*-alkanes was performed on a Thermo Finnigan Trace Gas  
233 Chromatograph (GC) Ultra coupled to a Thermo Finnigan DSQ mass spectrometer (MS). The alkenones and the  
234 *n*-alkanes were quantified with gas chromatography coupled to flame ionization detection. For GC analysis we  
235 used an Agilent HP 6890 GCs with CP Sil-5 columns (50 m length for the alkenones and 25 m for the *n*-alkanes)  
236 and helium as the carrier gas. Within the saturated hydrocarbon fraction, we also identified lycopane  
237 (2,6,10,14,19,23,27,31-octamethyldotriacontane) and several pentacyclic triterpenes (molecular mass  $M=440$ ,  
238 thereafter referred to as “triterpenoids”) such as  $\beta$ -amyrin Methyl Ether (ME) (Olean-12-ene-3 $\beta$ -ol), Germanicol  
239 ME (Olean-18-ene-3 $\beta$ -ol) and Taraxerol ME (Taraxer-14-en-3 $\beta$ -ol) (Fig. S2).

240 The polar fraction of 61 samples, containing the glycerol dialkyl glycerol tetraethers (GDGTs), was dissolved in a  
241 mixture of 99:1 (v/v) hexane/propanol and filtered through 0.45 mm PTFE filters. GDGTs were analysed (in  
242 triplicate) by high performance liquid chromatography (HPLC)/MS in single ion monitoring mode on an Agilent  
243 1100 series LC/MSD SL (Hopmans et al., 2004; Schouten et al., 2007).


## 245 4.1 Sedimentary patterns on the western Nile deep-sea fan

246 The deeper core P33 exhibits a near-continuous 500-cm finally laminated interval between ca. 100 cm core depth  
 247 and the bottom of the core, with high Ti/Ca ratios and high sedimentation rates (between 100 and 350 cm/ka) (Fig.  
 248 2a). A 40-cm interruption with poorly defined laminations and traces of bioturbation occurs in the upper part of the  
 249 laminated interval between 140 and 100 cm core depth. The intermediate depth core P73 shows a series of laminated  
 250 intervals between 200 cm and the bottom of the core, which are accompanied by high Ti/Ca ratios and high  
 251 sedimentation rates (between 150 and 250 cm/ka) (Fig. 2b). Several interruptions with poorly-laminated to non-  
 252 laminated (bioturbated) intervals are identified at 480 cm, 370 cm, 240-290 cm and 215 cm. In contrast to the two  
 253 deeper cores, the shallower core P99 is generally non-laminated with the exception of faint to well-defined  
 254 laminations at 510-520 cm, 490 cm, 470 cm, 405 cm, 340 cm, and 180-200 cm core depth (Fig. 2c). However, both  
 255 the Ti/Ca records and sedimentation rates exhibit changes similar to the two deeper cores with higher values  
 256 between 200 cm core depth and the bottom of the core (sedimentation rates of ca. 200 cm/ka).


(a) Partly-polarised microscope image



(b) SEM image &amp; EDS element analyses



(c) XRD mineralogy




257

258 Figure 3. Microscopic, elemental and mineralogical characterisation of sublayers from core P33. (a) Micro-facies and mineralogy of  
 259 a typical sublayer succession in cores P33. Left to right: part-polarized microscope image and schematic log of sublayers

260 DL1-LL2-DL2-LL1. (b) SEM images of powdered dark (DL1+DL2) and light sublayers (LL2) with EDS elemental identification.  
261 (c) mineralogical assemblage of powdered DL, LL and HL samples using XRD scans (from 4 to 85 °2θ). In sample DL (P33-3-22.5),  
262 the increase of the intensity baseline towards higher angles is due to the presence of an Fe-rich phases, probably pyrite.


#### 263 4.2 Structure and composition of the laminations

264 The lamination structure is regular throughout core P33 and similar to the laminations in core P73 and P99 (Fig.  
265 S4). Well-defined laminations are composed of an alternation of dark and light bands forming sublayers, whereby  
266 dark sublayers (DL) are mostly thicker (0.2 and 15 mm) than light sublayers (0.1 to 0.4 mm) (Fig. 3a). The  
267 mineralogy of DL consists of clay minerals (kaolinite, illite and smectite) as well as silt-size detrital minerals  
268 (quartz, plagioclase, pyroxene, iron-titanium oxides) (Fig. 3c). Subdivisions of DL into potassium (K)-rich  
269 (labelled DL1) and titanium (Ti)-rich (labelled DL2) parts are observed in thicker laminations. These can be  
270 distinguished by XRF elemental mapping as well as by their texture, Ti-rich sublayers containing coarser particles  
271 (ca. 20-30 μm) as K-rich sublayers (Fig. 3a, 4b).



272  
273 Figure 4. Laminations in core P33. (a) Fine laminations and (b) thick laminations from sections located at ca. 180 and 360 cm core  
274 depth, respectively. From left to right: core depth (cm), cross-polarized scan of the thin section, elemental mapping showing relative  
275 amounts of calcium (Ca, yellow) versus titanium (Ti, turquoise), elemental mapping showing relative amounts of calcium (Ca,  
276 yellow), potassium (K, pink) and titanium (Ti, blue), schematic lithological log with color-coded sublayer type following Fig. 3a (LL1:  
277 pink, DL1, grey, LL2: yellow and DL2: black), down-section profile of  $\log(\text{Ti}/\text{Ca})$  (blue) and  $\log(\text{Ti}/\text{K})$  (red).

278 Light sublayers (LL) contain predominantly carbonate and can be subdivided into two main types (Fig. 3a). The  
 279 light-sublayer type 1 (LL1) are diffuse sublayers of silt-sized (ca. 10-20  $\mu\text{m}$ ) and unsorted particles containing  
 280 biogenic carbonates (coccoliths and foraminifera), organic matter and quartz grains (Fig. 3a). Foraminifera shells  
 281 generally contain small grains of iron sulphides, which may have formed on organic residues (Fig. S5). Some of  
 282 these sublayers present a slight cementation. The light sublayers type 2 (LL2) contain well-sorted fine (ca. 1-10  
 283  $\mu\text{m}$ ) and needle-shaped calcite minerals, mixed with detrital silicate grains, including clay minerals (Fig. 3b). The  
 284 detrital assemblage in LL2 is similar to that of dark layers but contains a larger amount of calcite (Fig. 3c). LL2 are  
 285 abiotic, have a wavy internal structure and contain concentrated lenses of carbonate grains (Fig. 3a). Some of these  
 286 sublayers show a sharp lower boundary and either an upward decrease or increase in grain concentration. In the  
 287 lower part of core P33, three prominent LL2 are cemented and consolidated (thereafter denoted “hard layers” HL)  
 288 and are generally thicker than the softer LL2 (Fig. 2a). Four similar HL are also found in core P73 (Fig. 2b). The  
 289 presence of organic matter is observed either within LL1 or at the base of DL1.



290  
 291 **Figure 5. Isotopic signature of LL2 and HL from core P33. Scatter plot of  $\delta^{18}\text{O}$  vs  $\delta^{13}\text{C}$  values of three LL2 (yellow) and three**  
 292 **cemented HL (orange), as compared to isotopic signatures of methane-derived carbonates from the Nile DSF (grey triangles) (Aloisi**  
 293 **et al., 2000; Bayon et al., 2013); P33 top-core and S1 planktonic foraminifera (light and dark blue diamonds, resp.); present and S1**  
 294 **epibenthic foraminifera (light blue circle and blue triangles; Grimm et al., 2015); P33 bulk organic matter  $\delta^{13}\text{C}$  (brown rectangle);**  
 295 **Black Sea  $\delta^{13}\text{C}$  of dissolved inorganic carbon (DIC) (green triangle) (Fry et al., 1991);  $\delta^{18}\text{O}$  value of biogenic and inorganic benthic**  
 296 **and planktonic calcite formed in equilibrium with seawater (blue triangles) (Table S4, see calculation details in Supplementary**  
 297 **Information).**

298 The sublayers are deposited in a remarkably constant succession over the length of the laminated interval (Fig. 4).  
 299 This succession is particularly clear in the thicker laminations and follows the pattern from bottom to top: LL1,  
 300 DL1, LL2 and DL2 (Fig. 4b). Sublayers LL1 and DL2 are always present in the succession, whereas DL1 is very  
 301 thin or not detectable in ca. 10% of the successions and LL2 is missing in ca. 30% of the successions. Sublayers  
 302 are on average thicker in the lower part of the of the laminated interval (between 260 and 570 cm) and thinner in  
 303 the upper part. At our core sites, we did not identify any remains of diatoms. Thirteen anomalously thick layers  
 304 (mm- to cm-scale) were identified throughout the record and labelled event layers (EL). These well-mixed layers  
 305 containing matrix-supported grains and showing a clear bimodal distribution of silt- (ca. 15-20  $\mu\text{m}$ ) and mud-sized  
 306 particles. In general, EL have sharp but non-erosive boundaries and either no clear internal structure or show a  
 307 fining upward.


308

309 **Table 3: Overview of the proxies used and their main interpretation**

Proxy	Abbreviation	Interpretation	References
<b>Oxygenation</b>			
Laminations		Absence of bioturbation due to the lack of oxygen in bottom waters, which prevents burrowing animals to live in the sediments, changes in seasonality	(Schimmelmann et al., 2016)
Log(S/Cl) ratio	S/Cl	Relative increase of sedimentary sulphur with respect to pore-water concentrations indicating accumulations due to the formation of iron sulphides, which form when deoxygenation increases.	(Liu et al., 2012; Revel et al., 2015)
Lycopane		Low oxygen levels	(Sinninghe Damsté et al., 2003)
Benthic foraminifera	BF	Existence/absence of oxygen in the bottom waters	(Abu-Zied et al., 2008; Le Houedec et al., 2020)
<b>Primary productivity</b>			
Alkenones		Abundance of the haptophyte algae coccolithophorid <i>Emiliana Huxley</i> in the surface waters	(Marlowe et al., 1984; Volkman, 2000)
Crenarchaeol		Abundance of thaumarcheota living in the upper part of the water column	(Besseling et al., 2019)
<b>Sedimentation and river runoff</b>			
Log(Ti/Ca) ratio	Ti/Ca	Relative variation of terrigenous versus marine sediments input	(Billi and el Badri Ali, 2010; Blanchet et al., 2013; Garzanti et al., 2015)
Log(Ti/K) ratio	Ti/K	Relative proportion of coarser Ti-rich minerals to finer K-rich clay minerals	(Billi and el Badri Ali, 2010)
Sedimentation rate	SR	Speed of sediment accumulation (in cm/ka)	

Branched GDGTs	brGDGTs	Soil organic matter (with potential minor contributions from riverine and marine organic matter) brought by river runoff	(De Jonge et al., 2014; Sinninghe Damsté, 2016; Xiao et al., 2016)
Triterpenoids and long-chain n-alkanes	Triterpenoids and n-alkanes	Terrestrial higher plants biomarkers brought by river runoff	(Blanchet et al., 2014; Bray and Evans, 1961; Diefendorf et al., 2012; Jacob et al., 2005)
<b>Water-mass physical &amp; chemical properties</b>			
Oxygen isotopes	$\delta^{18}\text{O}$	Changes in temperature, salinity and isotopic composition (often related to the origin) of the water	
Carbon isotopes	$\delta^{13}\text{C}$	Changes in organic matter consumption and biogeochemical processes in the water column	

310



311



312 **Figure 6. Proxies of paleo-oxygenation in cores (a) P99 (turquoise), (b) P73 (blue) and (c) P33 (dark blue) on age scale (0-10 ka BP).**  
313 **For each core, from top to bottom: Log-ratio of S/Cl (continuous lines), mass accumulation rates (MAR) of lycopane in P99 and P33**  
314 **(dashed lines and circles), lamination pattern (darker boxes: laminated, lighter boxes: faint laminations). For P33: bottom curve**  
315 **shows the number of benthic foraminifera per g of sediment (white crosses and cubic spline as thick blue curve). Open boxes in P73**  
316 **and P33: interval of gradual reoxygenation. Grey bars under each marker and around laminated intervals show the age uncertainty.**

317 Stable  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  isotope measurements were performed on a set of 6 LL2 and HL sublayers ranging in  
318 thickness from 0.1 to 0.8 mm (Fig. 5). Cemented HL have a strongly depleted  $\delta^{13}\text{C}$  (-10 to -15 ‰) and an enriched  
319  $\delta^{18}\text{O}$  (0.5 to 2.5 ‰) signature compared to the softer LL2 (with  $\delta^{13}\text{C}$  between -11 and -5 ‰ and  $\delta^{18}\text{O}$  between 0.5  
320 and -2 ‰). These carbonates show a wide range of stable oxygen and carbon isotope values and apparently fall on  
321 a mixing line between a depleted  $\delta^{13}\text{C}$ /heavy  $\delta^{18}\text{O}$  and an enriched  $\delta^{13}\text{C}$ /light  $\delta^{18}\text{O}$  end-member, which do not  
322 correspond to either biogenic carbonates or regional methanogenic carbonates (Fig. 5).


### 323 **4.3 Spatial and temporal variations**

324 To build on the results from sedimentary facies at mm-scale, we explored the variations of oxygenation, primary  
325 productivity and terrigenous input during the past 10 kyr over the Nile DSF. We used a wide range of proxies,  
326 which are listed in Table 3 together with their interpretation keys.

327 In order to reconstruct past changes in oxygenation conditions, we used four independent tracers: the presence of  
328 laminations and lycopane, the benthic foraminifera assemblage and the sulphur to chlorine (S/Cl) ratio (Table 3  
329 and Fig. 6). In all cores, laminated intervals coincide with high sedimentation rates ( $> 100$  cm/ka, Fig. 6).

330 In the shallowest core P99, several short-term faintly-laminated intervals are observed between 9 and 8.3 ka BP  
331 and between 7.6 and 7.4 ka BP, which contain well-preserved laminations and a peak value of lycopane mass  
332 accumulation rates (MAR) (up to  $0.013 \mu\text{g}/\text{cm}^2.\text{a}$ ) at 7.5 ka BP (Fig. 6a). High S/Cl values occur between 9 and  
333 8.5 ka BP, but are interrupted by a strong drop around 8.7 ka BP. Values decrease after 8.5 ka BP, but remain  
334 slightly elevated between 8.5 and 7.2 ka BP (Fig. 6a). In intermediate depth core P73, laminations occur as a nearly  
335 continuous interval between 9.2 and 7.4 ka BP with an interruption between 8.3 and at 7.6 ka BP (Fig. 6b). High  
336 S/Cl values occur between 9.6 and 7.5 ka BP, with a strong drop in the non-laminated interval around 7.9 ka BP.  
337 The S/Cl record shows a gradual decrease between 7.3 and 6 ka BP, after which the record remained low and only  
338 varied slightly (Fig. 6b). In the deepest core P33, laminations occur continuously between 9.6 and 8.1 ka BP,  
339 followed by an interval of faint laminations until 7.3 ka BP (Fig. 6c). Lycopane MAR follow the lamination pattern  
340 with high values in the laminated interval between 9.6 and 8.1 ka BP (between  $0.1$  and  $0.35 \mu\text{g}/\text{cm}^2.\text{a}$ ), when MAR  
341 abruptly drop to ca.  $0.01 \mu\text{g}/\text{cm}^2.\text{a}$  and then gradually decrease until ca. 6 ka BP to less than  $0.001 \mu\text{g}/\text{cm}^2.\text{a}$  (Fig.  
342 6c). The S/Cl record shows generally higher values between 10 and 7.5 ka BP with lower values between 9.5 and

343 9.0 and around 8.8, 8.3, and 8 ka BP (Fig. 6c). A steady decrease is observed between 7.5 and 6.5 ka BP followed  
 344 by low values in the younger part of the core. The laminated interval between 9.6 and 8.1 ka BP is devoid of benthic  
 345 foraminifera (except at ca. 9.5 and 8.55 ka BP). From then on, a few occurrences are observed with a peak around  
 346 7.9 ka BP but their presence remains very volatile until ca. 7.4 ka BP, after which their number grows gradually  
 347 (Fig. 6c).  
 348




349  
 350 **Figure 7. Proxies of primary productivity in cores (a) P99 (turquoise), (b) P33 (blue), (c) GeoB7702-3 (black) and MS27PT (grey) on**  
 351 **age scale (0-10 ka BP). MAR of alkenones (crosses and continuous lines) and MAR of crenarchaeol for cores P33 and P99 (triangles**  
 352 **and dashed lines), as well as for cores GeoB7702-3 (black diamonds) (Castañeda et al., 2010) and MS27PT (grey triangles) (Ménot**  
 353 **et al., 2020). Lamination pattern (similar colour coding as in Fig. 6). Grey bars show the age uncertainty.**

354

355 Changes in primary productivity off the Nile mouth was traced using MAR of C<sub>37</sub>-alkenones and crenarchaeols in  
356 cores P99 and P33 (Table 3 and Fig. 7). Alkenones and crenarchaeol MAR are of similar order of magnitude in  
357 both cores (up to 0.3  $\mu\text{g}\cdot\text{cm}^{-2}\cdot\text{a}^{-1}$  and 3.5  $\mu\text{g}\cdot\text{cm}^{-2}\cdot\text{a}^{-1}$ , respectively). The overall patterns of changes in biomarkers  
358 are similar in both cores, with highest fluxes being measured between 10 and 7 ka BP. Several short pulses of  
359 alkenone and crenarchaeol deposition are observed in shallowest core P99 at 9.1, 8.8, 8.5, 8 and 7.5 ka BP, although  
360 the alkenone MAR do not show large peaks between 9 and 8.5 ka BP, while crenarchaeol MAR are high (Fig. 7a).  
361 There is a clear correspondence between the occurrence of peaks in MAR of organic compounds and the presence  
362 of laminations (Fig. 7a). In deeper core P33, the sampling resolution is higher and the signal is spikier (Fig. 7b).  
363 The MAR of both crenarchaeol and alkenones are higher in the laminated interval between 9.6 and 8.2 ka BP and  
364 in the faintly laminated interval between 8.2 and 6.5 ka BP (Fig. 7b). In nearby cores MS27PT (1390 m) and  
365 GeoB7702-3 (562 m) (Fig. 7c), crenarchaeol MAR are also higher between 10 and 8 ka BP, but MAR are highest  
366 between 10 and 8.5 ka BP in core MS27PT while a marked peak occurred in core GeoB7702-3 at ca. 8.1 ka BP  
367 (Castañeda et al., 2010; Ménot et al., 2020).

368 The changes in river runoff and sediment input to the Nile DSF were reconstructed using major elemental contents,  
369 MAR of branched GDGTs (brGDGTs), odd long-chain *n*-alkanes and triterpenoids and sedimentation rates (Table  
370 3, Fig. 8 and Fig. S3). The Ti/Ca and Ti/K records of all three cores ratios show very similar patterns with relatively  
371 high amounts of Ti in the early Holocene (ca. 10-7.5 ka BP) and the late Holocene (1-0 ka BP). Especially in well-  
372 laminated cores P73 and P33, high-frequency fluctuations in the Ti/Ca records during the early Holocene are related  
373 to the presence of Ca-rich laminations (Fig. 4). Although the magnitude is not similar in all cores, a rapid decrease  
374 in Ti/Ca records is observed around 7.2 ka BP, followed by a more gradual decrease until ca. 3-4 ka BP (Fig. 8).  
375 The decrease in Ti/Ca at 7.2 ka BP is accompanied by a simultaneous decrease in Ti/K (Fig. 8). A stepwise increase  
376 in Ti/Ca and Ti/K is observed in all cores between 3-3.5 and 1 ka BP. Sedimentation rates (SR) were calculated  
377 linearly between dated points (Table 2) (Fig. 8). They are of the same order of magnitude and vary synchronously  
378 in the three cores with highest SR between ca. 10 and 7 ka BP (>100 cm/ka). An abrupt decrease is recorded in  
379 cores P99 and P73 around 7 ka BP (Fig. 8a, b) whereas SR decreased more gradually from 9 to 7 in core P33 (Fig.  
380 8c). After 7 ka, the SR remained below 20 cm/ka in all three cores. The MAR of *n*-alkanes (and triterpenoids, Fig.  
381 S3) are a factor 10 higher in core P33 than in core P99 (up to 7 and 0.6  $\mu\text{g}\cdot\text{cm}^{-2}\cdot\text{a}^{-1}$ , resp.) but brGDGTs MAR are  
382 similar in both cores (up to 4  $\mu\text{g}\cdot\text{cm}^{-2}\cdot\text{a}^{-1}$  and 2  $\mu\text{g}\cdot\text{cm}^{-2}\cdot\text{a}^{-1}$ , resp.) (Fig. 8a, c). Higher fluxes of terrestrial organic  
383 matter are observed in shallowest core P99 between 9.5 and 7.5 ka BP with pulses at 9.1, 8.8, 8.5, 8.2 and 7.5 ka  
384 BP in laminated intervals (Fig. 8a). In deeper core P33, the MAR of terrestrial biomarkers are higher in the

385 laminated interval between 9.6 and 8.2 ka BP and lower in the faintly laminated interval between 8.2 and 6.5 ka  
386 BP (Fig. 8c).  
387




388

389 Figure 8. Proxies of terrigenous input in cores (a) P99 (turquoise), (b) P73 (blue) and (c) P33 (dark blue) on age scale (0-10 ka BP).  
 390 For all cores: Log(Ti/Ca) (continuous lines) and Ti/K (dashed lines), sedimentation rates and lamination pattern (similar colour  
 391 coding as in Fig. 6). For cores P99 and P33 (a, c): MAR of odd long-chain *n*-alkanes (circles and continuous lines) and branched  
 392 GDGTs (squares and dashed lines). Grey bars show the age uncertainty.

## 393 5. Discussion

### 394 5.1 Seasonal dynamics over the Nile deep-sea fan as derived from microfacies analyses

395 As shown in Fig. 5, the sequence of dark and light sublayers (DL2, LL1, DL1, LL2) is consistent throughout the  
 396 laminated interval, even though DL1 and LL2 can be very thin and sometimes absent. We propose that the different  
 397 sublayers recorded distinct depositional regimes throughout the year and represent an annual cycle.



398

399 Figure 9. (a) Seasonal succession of sublayers and (b) schematic seasonal cycle on the Nile DSF. (a) From left to right: part-polarized  
 400 microscope image, sublayer identification, determined season for each sublayer type and number of annual cycles in the sample. (b)  
 401 sketches of an annual cycle as derived from microfacies analyses from summer (top) to late spring (bottom).

402 The coarser grained DL2 are proposed to represent sediments deposited during summer floods of the Nile River  
 403 (Fig. 9). The presence of silt-sized particles of smectite, plagioclases and iron-titanium oxides in the DL2 matches  
 404 the grain size of particulate matter transported during peak Nile discharge (Billi and el Badri Ali, 2010), as well as

405 its mineralogical composition, typical of basaltic rocks from the Ethiopian volcanic plateau (Garzanti et al., 2015,  
406 2018) (Fig. 3c). According to present-day and historical observations (Halim et al., 1967), summer floods lead to  
407 the formation of freshwater plumes above a saline seawater mass (Fig. 1a). During sapropel S1, this low-salinity  
408 subsurface water mass extended much further offshore in the Levantine Basin (Bale et al., 2019; Tang and Stott,  
409 1993; Vadsaria et al., 2019). The important sediment loading of the freshwater plume that entered the relatively  
410 saline Levantine Basin probably played an important role in driving sedimentation on the Nile DSF. Observations  
411 of mixing processes between freshwater plumes containing a high sediment load and the underlying saline seawater  
412 shows the formation of salt and sediment fingers below the plume that lead to a fast and efficient settling of  
413 sediments (Jazi and Wells, 2016; Parsons et al., 2001). On the Nile DSF, the deposition of flood particles is likely  
414 driven by the formation of sediment-laden salt fingers evolving into dense (hyperpycnal) sediment plumes (Fig 9b)  
415 (Ducassou et al., 2008). The thick DL2 observed throughout the record match the depositional expression of  
416 hyperpycnal plumes and associated Nile summer floods. In addition to this regular pattern, exceptional flood events  
417 have been identified in several sediment archives over the western deep-sea fan based on the occurrence of mud  
418 clasts, and correspond in our record to the thick EL (Mologni et al., 2020).

419 The detrital DL2 are overlain by coarse-grained light sublayers (LL1) that are enriched in organic matter and  
420 biogenic carbonates, mostly foraminifera (often *G. ruber*) and coccoliths (Fig. S5). High abundance of planktonic  
421 foraminifera and in particular of *G. ruber* during sapropel S1 was reported in a sediment core from the SE Levantine  
422 Basin also under the influence of the Nile outflow (Mojtahid et al., 2015). Although generally associated to  
423 oligotrophic conditions in the Mediterranean Sea (Pujol and Grazzini, 1995), it was found that *G. ruber* thrived in  
424 low salinity and nutrient rich waters of the Nile plume. This high planktonic productivity was related to sustained  
425 reproduction of foraminifera during the months following the Nile flood, when salinity increased while food  
426 resources remained high (Mojtahid et al., 2015). We propose here that this process was similar to the so-called  
427 “Nile blooms”, which occurred in September-October prior to the construction of the Aswan Dam in 1965 (Nixon,  
428 2003). Fertilisation of the surface waters off the Rosetta mouth by large nutrients input during the summer Nile  
429 floods triggered an algal bloom, upon which zooplankton grazed (although no foraminifera was reported by Halim  
430 et al., 1967). We did not identify any diatom frustules associated to the Nile blooms, probably due to the poor  
431 preservation of opal in the Levantine Basin. However, the combination of foraminifera, coccoliths and organic  
432 matter in LL1 (Fig. S5) is proposed to represent post-flood layers following a fertilisation process similar to that  
433 occurring during the Nile blooms (Fig 9b). Efficient transport of nutrients from the Nile freshwater plume to the  
434 underlying saline Mediterranean water might have also been boosted by the process of diapycnal mixing described  
435 earlier (Oschlies et al., 2003). High levels of primary productivity probably triggered a gradual eutrophication of

436 the underlying water masses due to bacterial oxidation of sinking organic matter. Occasionally, these layers contain  
437 quartz grains that might be related to autumn-early winter dust storms (Goudie and Middleton, 2001).

438 The following sublayer DL1 is generally a fine-grained homogenous dark layer, which is enriched in potassium  
439 and reflects autumn to spring deposition. Analysis of the present suspension load of the Nile River shows that the  
440 suspended load of the main Nile is enriched in clay particles during low Nile discharge periods (i.e., from late  
441 autumn to early summer) (Billi and el Badri Ali, 2010; Garzanti et al., 2006). At present, the core site is ventilated  
442 by formation of intermediate and deep-water masses as it lies at the transition between LIW and LDW flows (Kubin  
443 et al., 2019). During S1 deposition, it was suggested that a sluggish hypoxic water mass occupied the depth range  
444 500-1800 m, originating from the Aegean Sea and labelled “Sapropel Intermediate Water” (SIW) (Zirks et al.,  
445 2019). The persistence of a yearly cycle of fluvial input during S1, with a marked low flow regime in winter as  
446 observed by our data fits previous observations by Tang and Scott (1993) and suggests that surface waters in the  
447 Aegean Sea might have been dense enough to sink and circulate at intermediate depth. However, the ventilation by  
448 SIW was probably very sluggish because we observe no reoxygenation of bottom waters during the winter or spring  
449 at our core site (see §5.2).

450 Sublayers DL1 are often overlain by a final sublayer of homogenous fine calcite grains (LL2) that is sometimes  
451 cemented (HL). The mineralogical assemblage in these layers is similar to the detrital-rich layers but shows an  
452 enrichment in calcite (Fig. 3c). SEM pictures suggest that microcrystalline calcite might have formed as a coating  
453 on the detrital grains, which could have served as nucleus (Fig. 3b). The internal draping structure of these layers  
454 and their regular occurrence in the lamination sequence always intercalated between DL1 and DL2 without  
455 sedimentary disturbance suggest that these calcite grains formed seasonally close to the seafloor, either in seawater  
456 or in highly porous sediments at the water-sediment interface (Fig. 9a). Furthermore, LL2 are also identified both  
457 in cores P73 and P99 with sublayer structures similar to those of core P33 (Fig. S4). This points to a regionally  
458 ubiquitous formation of authigenic carbonates in the water column, rather than local diagenetic processes occurring  
459 in the sediments. Based on their isotopic signature, we hypothesize that these layers formed during the late spring-  
460 early summer season (Fig. 9b and discussion in §5.2.2). The formation process of cemented HL is not clearly  
461 determined as these layers show internal disturbances and might have formed either at the seafloor (Alloué, 1990)  
462 or deeper in the sediments (Kasten et al., 2004).

## 463 5.2. Water-column stratification and chemistry


### 464 5.2.1. Bottom-water chemistry

465 The occurrence of bottom water-derived authigenic calcite sublayers (LL2) in core P33 provides a unique  
466 opportunity to investigate seawater chemistry using their oxygen and carbon isotope signatures (Fig. 5, 10). The  
467  $\delta^{13}\text{C}$  signatures of soft LL2 calcite sublayers range from -11 to -5 ‰ and are therefore distinct from  $\delta^{13}\text{C}$  values of  
468 present epibenthic foraminifera (ca. 1.2 ‰; Grimm et al., 2015) or from values estimated during S1 deposition (ca.  
469 0.3‰; Grimm et al., 2015) (Fig. 5). The markedly depleted  $\delta^{13}\text{C}$  of LL2 are comprised between the  $\delta^{13}\text{C}$  of bulk  
470 organic matter in core P33 (-17 to -20 ‰) and bottom water dissolved inorganic carbon (DIC) in the euxinic Black  
471 Sea (-6.3 ‰, Fry et al., 1991) (Fig. 5, 10b). For the two older LL2 in core P33, the bathymetric  $\delta^{13}\text{C}$  gradients from  
472 surface to bottom waters at the core site are similar to that of the present-day Black Sea: from 0 ‰ at the surface  
473 (as indicated by *G. ruber*  $\delta^{13}\text{C}$ ) to ca. -7‰ in bottom waters offshore the Nile mouth (Fig. 10b), and from 0.8 in  
474 surface to -6.3 ‰ in bottom waters in the Black Sea (Fry et al., 1991). Based on these preliminary results, we  
475 propose that LL2 formed in euxinic (sulfidic) bottom waters dominated by respiratory processes (i.e., degradation  
476 of organic matter) but further analyses are required to confirm this hypothesis (e.g., isotopic signatures of  
477 molybdenum and iron; Matthews et al., 2017).

478 The  $\delta^{13}\text{C}$  of HL and of the youngest LL2 is too low to result from anaerobic oxidation of organic matter only and  
479 suggests that part of the bottom-water DIC might have incorporated methane diffusing upwards from the sediments  
480 to bottom waters. As soft LL2 are assumed to form from bottom waters, the presence of methane implies a shoaling  
481 of the sulphate-methane transition zone (SMTZ) in the water column, which has been identified in the deeper parts  
482 of the Black Sea basin (Michaelis et al., 2002; Riedinger et al., 2010; Schouten et al., 2001). The persistence of  
483 methane in bottom waters of the SE Levantine Basin during sapropel S1 has also been proposed to account for the  
484 formation of Mg-rich calcite crusts around methane seeps and vents on deeper parts of the Nile DSF (Aloisi et al.,  
485 2002; Bayon et al., 2013). Anoxic and alkaline bottom waters at that time likely prevented the oxidation of methane  
486 into  $\text{CO}_2$  and led to the precipitation of carbonate crusts with a very depleted  $\delta^{13}\text{C}$  signature (-25 to -45 ‰), typical  
487 of hydrocarbon sources and clearly distinct from that of authigenic carbonate layers found in core P33 (Fig. 5).  
488 However, the incorporation of methane generated at the SMTZ from fermentation of sedimentary organic matter  
489 would provide a likely source of depleted DIC. It is therefore possible that methane-rich conditions developed  
490 temporarily in bottom waters and led to the formation of the youngest LL2 in core P33. The cemented HL in core  
491 P33 have a homogenous  $\delta^{13}\text{C}$  value of ca. -15 ‰ and might either have formed at the seafloor from bottom water



492 similarly to the youngest LL2, or from pore-waters deeper in the sediments during diagenetic precipitation of calcite  
 493 around the SMTZ. Such processes have previously been attributed to non-steady state diagenesis and fixation of  
 494 the SMTZ at a certain depth due to changes in sediment supply (Kasten et al., 2004).



495  
 496 **Figure 10.** Changes in water-column stratification and chemistry as provided by isotopic composition of authigenic calcite in  
 497 individual LL2 and HL of core P33. (a)  $\delta^{18}\text{O}$  of planktonic foraminifera (blue curve and diamonds), LL2 (yellow dots) and HL  
 498 (orange squares), as compared to biogenic and inorganic calcite formed in equilibrium with seawater providing information on the  
 499 water-column stratification as indicated on the right-hand side. (b)  $\delta^{13}\text{C}$  of planktonic foraminifera (green curve and diamonds),  
 500 LL2 (yellow dots) and HL (orange squares) and bulk organic matter (brown triangles) showing the changes in water-mass  
 501 oxygenation state. (c) Thickness of LL2 and HL, picture of the core surface, depth scale and indications of age.

502

### 503 5.2.2. Water-column stratification and precipitation of authigenic calcite

504 The  $\delta^{18}\text{O}$  of LL2 and HL ranges from -2 to 2.5 ‰, which is comprised between the  $\delta^{18}\text{O}$  signature of calcite formed  
505 in equilibrium with surface waters (ca. -1 to -2 ‰) and bottom waters (ca. 2 ‰) (Fig. 5, 10a). The heavier  $\delta^{18}\text{O}$  of  
506 HL and the youngest LL2 is similar to that of recent benthic foraminifera and suggests that they formed in bottom  
507 waters having similar physical and chemical characteristics as today with a comparable water-column stratification  
508 (Fig. 10a). By contrast, the  $\delta^{18}\text{O}$  values of older LL2 are similar to those of *G. ruber* formed in Blue Nile-  
509 dominated/low-salinity water during the summer, suggesting homogenous temperature and salinity gradients  
510 throughout the water-column (Fig. 10a). This observation has several important implications.

511 The  $\delta^{18}\text{O}$  of older LL2 around -2 ‰ implies that either LL2 formed in waters that remained dominated by the Nile  
512 freshwater input all year long, or that the formation of calcite in LL2 is related to the beginning of the flood season.  
513 According to isotopic data from various foraminifera species analysed in a sediment core south of Crete, low  
514 salinities typical of large Nile runoff was only a pervasive feature in the Levantine Basin (Tang and Stott, 1993).  
515 We therefore propose that the formation of the LL2 likely occurred during the onset of the Nile floods at the end  
516 of the spring/beginning of the summer, providing cations to the bicarbonate-rich waters that developed as a result  
517 of anaerobic oxidation of organic matter off the Nile mouth (as derived from  $\delta^{13}\text{C}$  signature of LL2, § 5.2.1) (Fig.  
518 9b). At the end of the dry (winter) period, the Nile River water is indeed enriched in cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ) (Dekov  
519 et al., 1997). The supply of cations to bottom waters enriched in  $\text{CO}_3^{2-}$  ions might have led to the supersaturation  
520 of seawater with respect to calcite and its precipitation on sediment particles at the seafloor.


521 The large range of  $\delta^{18}\text{O}$  values in LL2 suggests important reorganisation of water masses and water-column  
522 stratification during sapropel S1 offshore the Nile River mouth. Low  $\delta^{18}\text{O}$  during the oldest part of the sapropel  
523 indicates that temperature and salinity gradients were homogenous throughout the water column, with water-  
524 masses down to 700 m having a Nile freshwater fingerprint (Fig. 10a). As discussed earlier, the formation of  
525 authigenic calcite in core P33 likely occurred at the beginning of the summer flood season and the measured  $\delta^{18}\text{O}$   
526 therefore suggests a deep penetration of the freshwater plume on the Nile DSF. The convective mixing of fresh-  
527 and seawater driven by the formation of salt fingers below the flood plume described previously would provide a  
528 potential mechanism to draw low-salinity water to bottom parts of the Nile DSF (§ 5.1). It was shown that this  
529 process of double-diffusive (diapycnal) mixing enhances vertical fluxes of salt, nutrients and oxygen in the ocean  
530 to intermediate depths (Oschlies et al., 2003; Schmitt et al., 2005). We therefore hypothesize that during the  
531 deposition of S1, the freshwater released during Nile floods was efficiently transported downwards on the Nile

532 DSF, reaching water depth up to 700 m. The younger LL2 has a  $\delta^{18}\text{O}$  value of 0.6 ‰, between the  $\delta^{18}\text{O}$  of HL and  
533 benthic foraminifera (ca. 1.5 ‰) and the  $\delta^{18}\text{O}$  of present-day *G. ruber* (ca. 0 ‰). This indicates either a smaller  
534 inflow of freshwater during the Nile floods or a reduced vertical mixing.

### 535 **5.3. Spatial and temporal variability of deoxygenation in the SE Levantine Basin during sapropel S1**

536 Based on the observations of seasonal sublayer deposits and in particular the occurrence of authigenic calcite, we  
537 derived that bottom waters offshore the Nile mouth were anoxic almost all-year round during sapropel S1  
538 deposition and potentially reached euxinic and methane-rich states in spring. This indicates that even though the  
539 Nile flow decreased during the winter-spring months, potentially permitting to form the SIW in the Aegean Sea  
540 (Zirks et al., 2019), the circulation of this intermediate water mass was too sluggish to efficiently ventilate the  
541 basin. During the large summer Nile floods, efficient vertical mixing was probably able to draw low-salinity  
542 nutrient-rich and sediment-laden water masses down to 700 m w-d.

543 By comparing available markers of deoxygenation in the three sediment cores (in particular the S/Cl records and  
544 occurrence of laminations), we can determine the changes in seawater chemistry at centennial- to millennial-scale  
545 along a bathymetric transect (Fig. 11d). We have added two deeper cores MD04-2726 and MS27PT (retrieved at  
546 1060 and 1390 m w-d, resp.) for which S/Cl ratios were published previously (Ménot et al., 2020; Revel et al.,  
547 2015). The S/Cl ratio traces the diagenetic formation of sulphur minerals such as pyrite ( $\text{FeS}_2$ ), greigite ( $\text{Fe}_3\text{S}_4$ ) or  
548 pyrrhotite ( $\text{Fe}_7\text{S}_8$ ) from the pore-water sulphides (Liu et al., 2012; Matthews et al., 2017; Revel et al., 2015). This  
549 process is favoured by oxygen depletion, which leads to the accumulation of dissolved sulphides in the pore-waters  
550 and precipitation of solid iron sulphides. These iron sulphides are generally resistant to post-depositional  
551 oxygenation and therefore permit the identification of past hypoxic intervals that might have be partly reoxidized  
552 (Larrasoña et al., 2006).



553

554 **Figure 11. Regional compilation of changes in terrigenous, primary productivity and oxygenation conditions during the Holocene.**  
 555 (a) Standardized (z-score) records of log (Ti/Ca) of cores P99 (light blue), P73 (blue), P33 (dark blue), MD04-2726 (dark grey),  
 556 MS27PT (grey) (Revel et al., 2015). Z-scoring ( $(x - \text{mean}(Ex)) / \text{stdev}$ ) allows to compare trends in a similar proxy from different archives  
 557 by scaling the y-range and aligning the means on 0. (b) MAR of crenarchaeol in cores P99 (light blue) and P33 (dark blue), planktonic  
 558 foraminifera in core PS009PC (purple) (Mojtahid et al., 2015). (c) Abundance of benthic foraminifera in core P33 (cubic spline, dark  
 559 blue, inverse y-scale) and core PS009PC (purple). (d) Log(S/Cl) records and lamination patterns in cores from the Nile DSF  
 560 (shallower to deeper core site from top to bottom): P99 (light blue), P73 (blue), P33 (dark blue), MD04-2726 (dark grey), MS27PT  
 561 (grey) (Revel et al., 2015). (e) V/Al records in cores PS009PC (purple) (Hennekam et al., 2014) and MD9509 (brown) (Matthews et  
 562 al., 2017). Grey vertical bars: sapropel S1a and S1b; lighter grey bars: periods of slight reoxygenation; white bar: S1 interruption  
 563 (at ca. 8 ka BP). Map in (b) showing the location of the cores along the path of the Levantine Jet.

564 In order to determine potential forcing factors, this bathymetric and temporal record of paleo-oxygenation will be  
565 compared with regional reconstructions of terrigenous input and primary productivity (Fig. 11a,b). The dynamics  
566 of terrigenous input and primary productivity are coherent over the western Nile DSF as shown by the good  
567 alignment of Ti/Ca ratios measured in all cores (Fig. 11a) and the crenarchaeol fluxes from P33, P99, MS27PT  
568 (Ménot et al., 2020; Revel et al., 2015) and GeoB7702-3 (Castañeda et al., 2010) (Fig. 7, 11b). Biomarker records  
569 can be affected by post-depositional oxidation, especially by the downward diffusion of oxygen into the reduced  
570 sapropelic sediments (i.e., the so-called burn-down process, Rutten and de Lange, 2003). However, comparable  
571 trends between biomarker records and other markers of primary productivity from regional archives that are not  
572 affected by burn-down processes (i.e., flux of planktonic foraminifera in core PS009PC, Mojtahid et al., 2015)  
573 suggest that productivity played a larger role than post-depositional oxidation in modulating the biomarker fluxes  
574 in our records (Fig. 11b).

### 575 **5.3.1. Waxing and waning of a hypoxic water-mass during the Holocene**

576 Prior to 10 ka BP, most sediment records from the eastern Mediterranean recorded the existence of oxic conditions  
577 as reflected by the presence of epibenthic foraminifera at water depths ranging from 500 to 2000 m (Cornuault et  
578 al., 2018; Le Houedec et al., 2020; Schmiedl et al., 2010) as well as low sulphur contents and redox sensitive  
579 element ratios (RSER) (Azrieli-Tal et al., 2014; Hennekam et al., 2014; Matthews et al., 2017; Tachikawa et al.,  
580 2015). At deep site MS27PT offshore the Nile River mouth, suboxic conditions occurred between 12 and 10 ka BP  
581 and were accompanied by higher primary productivity (Ménot et al., 2020). A peak of productivity was also  
582 identified in some records from the SE Levantine Basin prior to the onset of S1 deposition at depth similar to  
583 MS27PT (>1000 m) (Zirks et al., 2021; Zwiép et al., 2018) but not at shallower depths (Castañeda et al., 2010;  
584 Mojtahid et al., 2015).

585 All cores from the Nile DSF show a pronounced decrease in oxygenation between 10 and 6.5 ka BP, marked by  
586 higher S/Cl records and the quasi-absence of benthic foraminifera, followed by relatively stable and oxic conditions  
587 in the younger parts of the cores (Fig. 11d). In core P33, we propose that the bottom waters were sulfidic to methanic  
588 during S1 deposition, based on  $\delta^{13}\text{C}$  measurements on LL2 (see § 5.2.1). Such conditions have been detected only  
589 in deeper parts of the Levantine Basin during S1 (>2000 m; Azrieli-Tal et al., 2014) and shallower sites located  
590 along the path of the Levantine Jet were found to be anoxic but not euxinic between 10 and 7 ka BP (Hennekam et  
591 al., 2014; Matthews et al., 2017; Zirks et al., 2021). Recent results hypothesize that high input of iron through  
592 enhanced Nile runoff during S1 likely prevented the bottom waters on the coast of Israel (>1000 m w-d) to become  
593 sulfidic and instead remained anoxic (Zirks et al., 2021). Our core sites on the Nile DSF are shallower and under

594 direct influence of fluvial nutrient release compared to the more distal sites of Zirks et al. (2021). Higher primary  
595 productivity and stronger bacterial respiration of sinking organic matter might have been sufficient to balance the  
596 large input of iron- and manganese-rich reactive phases and shift the bottom waters in a euxinic state. In addition,  
597 according to our seasonal reconstructions, euxinia might have developed in the bottom waters during a short  
598 interval in the spring (§5.2.1 and Fig. 9c), which might not be recorded by sampling at centennial-scale resolution.  
599 A complete reoxygenation is achieved at all depths by 6.5 ka BP (and by 6 ka at 1060 m w-d, Revel et al., 2015).  
600 At the scale of the Levantine Basin, the recovery of benthic ecosystems appears to have followed a time-  
601 transgressive pattern, with shallower sites depicting oxic conditions as early as 8 ka BP (Schmiedl et al., 2010).  
602 Such a pattern is also evidenced on the Nile DSF with deoxygenation conditions being maintained until ca. 6.5-6  
603 ka BP at depths >1000 m while shallower sites already started recovering around 7.5 ka BP (Fig. 11d).

### 604 **5.3.2. Dynamic fluctuations in time and space**

605 Centennial fluctuations in oxygenation conditions are observed on the Nile DSF between 570 and 1400 m w-d  
606 during sapropel 1a (10-8 ka BP) (Fig. 11d). Multiple intervals of faint laminations occur in core P99 every 200-  
607 300 yrs (i.e., at 9, 8.8, 8.6, 8.3 ka BP), suggesting a dynamic vertical structure of the upper boundary of the hypoxic  
608 water-mass. In particular, a pronounced shoaling to 400 m w-d is observed around 8.5 ka (Fig. 11d). Cores at  
609 intermediate depth also show variations in the oxygenation state with lower S/Cl ratios around 9.2 and 8.3 ka BP  
610 recorded in both cores P33 and P73 (although more prominent in P33) and at 8.7 ka BP recorded in core P33 (Fig.  
611 6 and 11c). Millennial-scale variations in bottom-water oxygenation were also recorded at similar depths by nearby  
612 cores PS009PC and 9509 as evidenced by fluctuations in the RSER (Fig. 11e) and in the benthic fauna (Hennekam  
613 et al., 2014; Le Houedec et al., 2020; Matthews et al., 2017). Below 1000 m w-d on the western Nile DSF, anoxic  
614 conditions have likely been more stable during S1a, as suggested by continuously laminated sapropels and  
615 relatively constant S/Cl ratios in deeper cores MD04-2726 and MS27PT (Fig. 6, 11d). Due to complex diagenetic  
616 processes involving sulphur cycling, the S/Cl ratio may not have recorded rapid changes in the oxygenation state  
617 as shown by RSER (Matthews et al., 2017; Tachikawa et al., 2015). However, the presence of continuous  
618 laminations throughout the sapropel sequence strongly suggests stable anoxic conditions at depths >1000 m. During  
619 S1a, wide-spread anoxia is recorded between 500 and 3000 m w-d in the LB (De Lange et al., 2008; Zirks et al.,  
620 2021).

621 A pronounced but short-lasting interval of reoxidation at ca. 8 ka BP is observed on the western Nile DSF in the  
622 upper 740 m, marked by a lack of laminations at 400 and 570 m w-d, faint laminations and recolonization of the  
623 seafloor by benthic foraminifera at 740 m w-d and reduced S/Cl ratios at all depths (Fig. 7, 11d). The presence of

624 continuous laminations in the deeper cores indicates that conditions remained anoxic during this time interval at  
625 depths >1000 m. The reoxygenation event at 8 ka BP is a widespread marker in the eastern Mediterranean generally  
626 referred to as sapropel S1 interruption, which has been related to a drastic decrease in Nile runoff around the 8.2  
627 ka event (Blanchet et al., 2013; Rohling et al., 2015). In nearby cores located above 900 m w-d at the coast of Israel,  
628 the timing of this reoxygenation event was similar (Fig. 11e) (Hennekam et al., 2014; Matthews et al., 2017) but  
629 these more distal cores reached a fully oxic state as indicated by the presence of epibenthic fauna (Fig. 11c) (Le  
630 Houedec et al., 2020; Schmiedl et al., 2010). Most sites in the Levantine Basin from depths ranging from 500 to  
631 3000 m depict suboxic to oxic conditions during S1 interruption (e.g., Azrieli-Tal et al., 2014; Gallego-Torres et  
632 al., 2010; Kuhnt et al., 2008; Mercone et al., 2001; Tachikawa et al., 2015). Low values of bottom-water  $\delta^{13}\text{C}$   
633 during the S1 interruption, however, indicate stagnating water masses (Schmiedl et al., 2010).  
634 Hypoxic conditions developed again on the western NDSF around 7.8-7.5 ka BP (sapropel S1b) (Fig. 11d). If  
635 bottom waters were suboxic at 740 m as indicated by the presence of a few benthic foraminifera and faint  
636 laminations in core P33, laminated intervals and high lycopane fluxes in cores P73 and P99 suggest that anoxic  
637 conditions existed intermittently between 600 and 400 m w-d during S1b (Fig. 6a, 11d). At deeper sites, conditions  
638 remained anoxic until ca. 6.7 ka BP when laminations stopped in core MS27PT (Fig. 11d). This suggests the  
639 existence of split anoxia, during which both shallower and deeper water masses were deprived of oxygen due to  
640 the respiration of rapidly-sinking organic detritus, while intermediate water masses remained suboxic (Bianchi et  
641 al., 2006; Rush et al., 2019). In most cores from the Levantine Basin, suboxic to anoxic conditions were re-  
642 established by 7.8 ka BP but generally lasted only a few hundred years and were rather unstable (Fig. 11c, e)  
643 (Schmiedl et al., 2010). Suboxic conditions and a gradual recolonization of the seafloor by benthic foraminifera  
644 indicating a gradual reoxygenation between 7.5 and 6.5 ka BP were observed at several sites along the Levantine  
645 Jet pathway at depths <1000 m (Fig. 11c) (Le Houedec et al., 2020; Schmiedl et al., 2010).

### 646 **5.3.3. The role of runoff-driven eutrophication in the SE Levantine Basin**

647 Recent modelling experiments indicated that the formation of a basin-wide oxygen depletion in the deep  
648 Mediterranean Sea during sapropel S1 required a multi-millennial deep-water stagnation (Grimm et al., 2015). The  
649 rising sea-level during the deglaciation allowed the inflow of fresher Atlantic-derived water through the Gibraltar  
650 Strait, thereby increasing the buoyancy of surface waters and decreasing deep-water ventilation in the eastern  
651 Mediterranean (Cornuault et al., 2018). This initial stagnation may have been reinforced by larger inputs of  
652 freshwater from the Nile river, which enhanced water-column stratification from 14 ka BP onwards (Castañeda et  
653 al., 2016; Cornuault et al., 2018; Vadsaria et al., 2019).

654 The marked onset of anoxia in the eastern Mediterranean Sea at ca. 10 ka BP is striking as it is recorded uniformly  
655 at various depths in the basin (De Lange et al., 2008; Schmiedl et al., 2010). Detailed faunal and isotopic studies  
656 showed that deep-water stagnation gradually developed from 17 ka BP onwards, with a distinct drop in  $\delta^{13}\text{C}$  around  
657 11 ka BP (Cornuault et al., 2018; Schmiedl et al., 2010). It was suggested that conditions remained oligotrophic  
658 until 10 ka BP, allowing benthic fauna to dwell even in stagnating water masses, after which benthic ecosystems  
659 passed a threshold that led to the development of anoxic conditions (Abu-Zied et al., 2008; Kuhnt et al., 2008). In  
660 deeper parts of the LB, the development of anoxia might primarily result from long-term preconditioning of water  
661 masses as modelled by Grimm et al. (2015). However, this modelling experiment does not produce anoxic  
662 conditions in water depths above 1800 m, which suggests that other forcing factors might drive deoxygenation  
663 dynamics at intermediate depths. As seen in section 5.3.2, considerable spatial and temporal variability in  
664 oxygenation was observed in the SE Levantine Basin. We propose here that these changes were largely driven by  
665 modulation of runoff-derived nutrient loadings. Around 10 ka BP onwards, a drastic increase in primary  
666 productivity has been identified in the SE Levantine Basin (Fig. 11b) (Mojtahid et al., 2015), which coincides with  
667 high precipitation and runoff in the Nile River basin (Castañeda et al., 2016; Weldeab et al., 2014). It is therefore  
668 postulated that the large nutrient loading delivered by summer floods of the Nile River led to a switch from an  
669 oligotrophic to a meso- or eutrophic state in the SE Levantine Basin, with high levels of primary productivity in  
670 surface waters. As an extended version of the Nile blooms forming offshore the Nile mouth during historical times  
671 (Halim et al., 1967), peak Nile runoff during the mega-summer monsoon of the African Humid Period probably  
672 fertilised large parts of the SE Levantine Basin during sapropel S1 deposition (Hennekam et al., 2015; Schmiedl et  
673 al., 2010). Lower oxygenation indices in cores located closer to the Nile River as shown in our study further support  
674 the pivotal impact of Nile River-induced eutrophication on the rapid spread of hypoxia in the SE Levantine Basin  
675 (Schmiedl et al., 2010; Zirks et al., 2019). Moreover, it was recently proposed that rapid changes in oxygenation  
676 could occur within 200-500 years for water masses located between 500 and 1800 m w-d due to oxygen utilization  
677 by bacterial organic matter degradation (Zirks et al., 2019). We therefore propose that most of the changes in the  
678 strength and extent of hypoxia offshore the Nile River were related to runoff-induced fertilisation and subsequent  
679 eutrophication of the water-column. However, these centennial- to millennial-scale variations were superimposed  
680 on the multi-millennial development of deep-water stagnation and suggest complex interactions and feedback  
681 processes between deep-water circulation, stagnation/stratification and eutrophication for driving deoxygenation  
682 dynamics. The rapid deoxygenation of the SE Levantine Basin around 10 ka BP following widespread  
683 eutrophication of surface waters, after thousands of years of deep-water stagnation, also calls for modelling



684 experiments to include temporal variations in nutrient loading and river runoff, which are important variables in  
685 the present Nile coastal system (Nixon, 2003; Powley et al., 2016).

## 686 **6 Conclusions**

687 By combining microfacies analyses with downcore geochemical measurements partly at seasonal resolution, our  
688 study provides a first estimation of changes in oxygenation conditions of the bottom waters on the western Nile  
689 deep-sea fan in the SE Levantine Basin. The regular seasonal alternation of detrital, biogenic and chemical  
690 sublayers in the laminated sequence deposited during sapropel S1 in a set of cores from different water depths on  
691 the Nile deep-sea fan are ascribed to seasonal changes (Fig. 9b). Strong summer Nile floods during S1 led to the  
692 deposition of thick (up to a few mm) silt-sized detrital sublayers that dominate total layer thickness. The large  
693 floods likely triggered surface blooms of phyto- and zooplankton in autumn similar to historical “Nile blooms”  
694 (Halim et al., 1967). The subsequent deposition of clay-rich detrital sublayers was associated with the low discharge  
695 regime of the Nile during winter. The reduction of Nile runoff might have sufficiently decreased the surface water  
696 salinity in the Levantine Basin, thereby allowing the formation of a sluggish intermediate water mass in the Aegean  
697 Sea (Zirks et al., 2019). The occurrence of inorganic carbonate sublayers in several laminated cores from the  
698 western Nile deep-sea fan suggests that bottom waters reached a supersaturation state for calcite. The depleted  $\delta^{13}\text{C}$   
699 signature of these sublayers points to the existence of anoxic to euxinic (and sometimes methane-rich) bottom  
700 waters accompanied by a high level of anaerobic remineralisation of organic matter leading to high alkalinity. The  
701 most likely process initiating the deposition of these layers was the onset of the Nile floods triggered by increased  
702 precipitation, which supplied sufficient amounts of cations to the seawater. First measurements of individual sub-  
703 millimetre layers underpin their potential to reconstruct seawater chemistry at times when no benthic fauna existed.  
704 On millennial time-scales, we show that variations in oxygenation dynamics at intermediate depth in the SE  
705 Levantine Basin followed changes in primary productivity driven by nutrient fertilisation during high Nile runoff.  
706 Deoxygenation above the Nile deep-sea fan shoaled to water depths as shallow as 400 m and varied on centennial-  
707 to millennial-scales in the upper 750 m of the water column. In contrast, the records of cores located below 1000  
708 m water-depth reflect more stable anoxic conditions between 10 and 6.5 ka BP. The development and fluctuations  
709 of anoxic conditions during sapropel S1 are coherent along the path of the Levantine Jet suggesting a common  
710 control. The posited multi-millennial development of deep-water stagnation was likely a prerequisite to trigger  
711 anoxia in the deeper parts of the eastern Mediterranean Basin (Cornuault et al., 2018; Grimm et al., 2015), but  
712 changes in primary productivity in surface waters probably drove the rapid changes in oxygenation state of  
713 intermediate water masses in the SE Levantine Basin through eutrophication processes. Indeed, tight temporal links

714 between regional productivity records, oxygenation and runoff dynamics, as well as the evidence of stronger  
715 deoxygenation close to the Nile mouth point to a pivotal role of runoff-driven nutrient supply and fertilisation.  
716 Furthermore, the rapid switch towards anoxic conditions around 10 ka BP suggests the crossing of thresholds or  
717 tipping points, which still remain elusive. Such processes could be further explored using transient modelling  
718 experiments incorporating variations in nutrient loading and Nile runoff at various time-scales for understanding  
719 feedback processes and system sensitivity.

#### 720 **Data availability**

721 All data presented in this paper will be available at the PANGEA database upon publication of the paper and for  
722 the reviewers

723 Table S1: Elemental contents measured by XRF for core P362/2-33 and shown in Fig. 2, 6, 8 and 9.

724 Table S2: Elemental contents measured by XRF for core P362/2-99 and shown in Fig. 2, 6, 8 and 9.

725 Table S3: Elemental contents measured by XRF for core P362/2-73 and shown in Fig. 2, 6, 8 and 9.

726 Table S4: Oxygen and carbon isotope data for core P362/2-33 shown in Fig. 4.

727 Table S5: Biomarker data for core P362/2-33 shown in Fig. 6, 7, 8 and 9.

728 Table S6: Biomarker data for core P362/2-99 shown in Fig. 6, 7, 8 and 9.

#### 729 **Supplement Link**

730 For the present submission, supplementary figures are provided in a separate file.

#### 731 **Author contribution**

732 CLB designed the study, measured and analysed the data. RT measured and analysed the XRF elemental contents.  
733 AES assisted with XRD measurements. SS, MF and AB provided guidance, lab space and logistic support for  
734 biomarker measurements, sedimentology and micro-facies analyses, respectively. CLB wrote the manuscript, to  
735 which all co-authors contributed.

#### 736 **Competing interests**

737 The authors declare that they have no conflicts of interest.

## 738 **Acknowledgements**

739 Measurements for this paper were carried out during a DFG own-position grant to C.B. at GEOMAR (Kiel,  
740 Germany) and NIOZ (Texel, Netherlands) (BL2111/1-1, 2010-2013) and during a reintegration grant at GFZ  
741 (Potsdam, Germany) (2018-2020). We are grateful to our colleagues from section 4.3 at GFZ for their support in  
742 term of sample preparation (B. Brademann), measurements (Stable isotopes lab, B. Plessen and S. Pinkerneil) and  
743 discussions (esp. J. Mingram). We also acknowledge G. Ménot, M. Mojtahid, R. Hennekam, G. Bayon and  
744 A. Matthews who kindly provided and discussed their data. The paper was written during the lockdown due to  
745 COVID-19 (March to June 2020) with background support from Daniel Hope and Max Richter.

746

## 747 **References**

748 Abu-Zied, R. H., Rohling, E. J., Jorissen, F. J., Fontanier, C., Casford, J. S. L. and Cooke, S.: Benthic foraminiferal  
749 response to changes in bottom-water oxygenation and organic carbon flux in the eastern Mediterranean during  
750 LGM to Recent times, *Mar. Micropaleontol.*, 67(1–2), 46–68, <https://doi.org/10.1016/j.marmicro.2007.08.006>,  
751 2008.

752 Adloff, F., Somot, S., Sevault, F., Jordà, G., Aznar, R., Déqué, M., Herrmann, M., Marcos, M., Dubois, C., Padorno,  
753 E., Alvarez-Fanjul, E. and Gomis, D.: Mediterranean Sea response to climate change in an ensemble of twenty first  
754 century scenarios, *Clim. Dyn.*, 45(9), 2775–2802, <https://doi.org/10.1007/s00382-015-2507-3>, 2015.

755 Allouç, J.: Quaternary crusts on slopes of the Mediterranean Sea: A tentative explanation for their genesis, *Mar.*  
756 *Geol.*, 94(3), 205–238, [https://doi.org/10.1016/0025-3227\(90\)90070-Z](https://doi.org/10.1016/0025-3227(90)90070-Z), 1990.

757 Aloisi, G., Pierre, C., Rouchy, J.-M., Foucher, J.-P. and Woodside, J.: Methane-related authigenic carbonates of  
758 eastern Mediterranean Sea mud volcanoes and their possible relation to gas hydrate destabilisation, *Earth Planet.*  
759 *Sci. Lett.*, 184(1), 321–338, [https://doi.org/10.1016/S0012-821X\(00\)00322-8](https://doi.org/10.1016/S0012-821X(00)00322-8), 2000.

760 Aloisi, G., Bouloubassi, I., Heijs, S. K., Pancost, R. D., Pierre, C., Sinninghe Damsté, J. S., Gottschal, J. C., Forney,  
761 L. J. and Rouchy, J.-M.: CH<sub>4</sub>-consuming microorganisms and the formation of carbonate crusts at cold seeps, *Earth*  
762 *Planet. Sci. Lett.*, 203(1), 195–203, [https://doi.org/10.1016/S0012-821X\(02\)00878-6](https://doi.org/10.1016/S0012-821X(02)00878-6), 2002.

763 Azrieli-Tal, I., Matthews, A., Bar-Matthews, M., Almogi-Labin, A., Vance, D., Archer, C. and Teutsch, N.:  
764 Evidence from molybdenum and iron isotopes and molybdenum–uranium covariation for sulphidic bottom waters  
765 during Eastern Mediterranean sapropel S1 formation, *Earth Planet. Sci. Lett.*, 393, 231–242,  
766 <https://doi.org/10.1016/j.epsl.2014.02.054>, 2014.

767 Bale, N. J., Hennekam, R., Hopmans, E. C., Dorhout, D., Reichart, G.-J., van der Meer, M., Villareal, T. A.,  
768 Sinninghe Damsté, J. S. and Schouten, S.: Biomarker evidence for nitrogen-fixing cyanobacterial blooms in a  
769 brackish surface layer in the Nile River plume during sapropel deposition, *Geology*,  
770 <https://doi.org/10.1130/G46682.1>, 2019.

- 771 Bayon, G., Dupré, S., Ponzevera, E., Etoubleau, J., Chéron, S., Pierre, C., Mascle, J., Boetius, A. and de Lange, G.  
772 J.: Formation of carbonate chimneys in the Mediterranean Sea linked to deep-water oxygen depletion, *Nat. Geosci.*,  
773 6(9), 755–760, <https://doi.org/10.1038/ngeo1888>, 2013.
- 774 Besseling, M. A., Hopmans, E. C., Koenen, M., van der Meer, M. T. J., Vreugdenhil, S., Schouten, S., Sinninghe  
775 Damsté, J. S. and Villanueva, L.: Depth-related differences in archaeal populations impact the isoprenoid tetraether  
776 lipid composition of the Mediterranean Sea water column, *Org. Geochem.*, 135, 16–31,  
777 <https://doi.org/10.1016/j.orggeochem.2019.06.008>, 2019.
- 778 Bianchi, D., Zavatarelli, M., Pinardi, N., Capozzi, R., Capotondi, L., Corselli, C. and Masina, S.: Simulations of  
779 ecosystem response during the sapropel S1 deposition event, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 235(1–3),  
780 265–287, <https://doi.org/10.1016/j.palaeo.2005.09.032>, 2006.
- 781 Billi, P. and el Badri Ali, O.: Sediment transport of the Blue Nile at Khartoum, *Quat. Int.*, 226(1–2), 12–22,  
782 <https://doi.org/10.1016/j.quaint.2009.11.041>, 2010.
- 783 Blaauw, M. and Christen, J. A.: Flexible paleoclimate age-depth models using an autoregressive gamma process,  
784 *Bayesian Anal.*, 6(3), 457–474, <https://doi.org/10.1214/11-BA618>, 2011.
- 785 Blanchet, C. L., Tjallingii, R., Frank, M., Lorenzen, J., Reitz, A., Brown, K., Feseker, T. and Brückmann, W.:  
786 High- and low-latitude forcing of the Nile River regime during the Holocene inferred from laminated sediments of  
787 the Nile deep-sea fan, *Earth Planet. Sci. Lett.*, 364, 98–110, <https://doi.org/10.1016/j.epsl.2013.01.009>, 2013.
- 788 Blanchet, C. L., Frank, M. and Schouten, S.: Asynchronous Changes in Vegetation, Runoff and Erosion in the Nile  
789 River Watershed during the Holocene, *PLOS ONE*, 9(12), e115958, <https://doi.org/10.1371/journal.pone.0115958>,  
790 2014.
- 791 Brauer, A. and Casanova, J.: Chronology and depositional processes of the laminated sediment record from Lac  
792 d’Annecy, French Alps, *J. Paleolimnol.*, 25(2), 163–177, <https://doi.org/10.1023/A:1008136029735>, 2001.
- 793 Bray, E. E. and Evans, E. D.: Distribution of n-paraffins as a clue to recognition of source beds, *Geochim.*  
794 *Cosmochim. Acta*, 22(1), 2–15, [https://doi.org/10.1016/0016-7037\(61\)90069-2](https://doi.org/10.1016/0016-7037(61)90069-2), 1961.
- 795 Brooks, S. P. and Gelman, A.: General Methods for Monitoring Convergence of Iterative Simulations, *J. Comput.*  
796 *Graph. Stat.*, 7(4), 434–455, <https://doi.org/10.1080/10618600.1998.10474787>, 1998.
- 797 Castañeda, I., Schefuß, E., Pätzold, J., Sinninghe Damsté, J. S., Weldeab, S. and Schouten, S.: Millennial-scale sea  
798 surface temperature changes in the eastern Mediterranean (Nile River Delta region) over the last 27,000 years,  
799 *Paleoceanography*, 25(1), <https://doi.org/10.1029/2009PA001740>, 2010.
- 800 Castañeda, I. S., Schouten, S., Pätzold, J., Lucassen, F., Kasemann, S., Kuhlmann, H. and Schefuß, E.:  
801 Hydroclimate variability in the Nile River Basin during the past 28,000 years, *Earth Planet. Sci. Lett.*, 438, 47–56,  
802 <https://doi.org/10.1016/j.epsl.2015.12.014>, 2016.

- 803 Cornuault, M., Tachikawa, K., Vidal, L., Guihou, A., Siani, G., Deschamps, P., Bassinot, F. and Revel, M.:  
804 Circulation Changes in the Eastern Mediterranean Sea Over the Past 23,000 Years Inferred From Authigenic Nd  
805 Isotopic Ratios, *Paleoceanogr. Paleoclimatology*, 33(3), 264–280, <https://doi.org/10.1002/2017PA003227>, 2018.
- 806 Cramp, A. and O’Sullivan, G.: Neogene sapropels in the Mediterranean: a review, *Mar. Geol.*, 153(1), 11–28,  
807 [https://doi.org/10.1016/S0025-3227\(98\)00092-9](https://doi.org/10.1016/S0025-3227(98)00092-9), 1999.
- 808 De Jonge, C., Hopmans, E. C., Zell, C. I., Kim, J.-H., Schouten, S. and Sinninghe Damsté, J. S.: Occurrence and  
809 abundance of 6-methyl branched glycerol dialkyl glycerol tetraethers in soils: Implications for palaeoclimate  
810 reconstruction, *Geochim. Cosmochim. Acta*, 141, 97–112, <https://doi.org/10.1016/j.gca.2014.06.013>, 2014.
- 811 De Lange, G. J., Thomson, J., Reitz, A., Slomp, C. P., Speranza Principato, M., Erba, E. and Corselli, C.:  
812 Synchronous basin-wide formation and redox-controlled preservation of a Mediterranean sapropel, *Nat. Geosci.*,  
813 1(9), 606–610, <https://doi.org/10.1038/ngeo283>, 2008.
- 814 Dekov, V. M., Komy, Z., Araújo, F., Van Put, A. and Van Grieken, R.: Chemical composition of sediments,  
815 suspended matter, river water and ground water of the Nile (Aswan-Sohag traverse), *Sci. Total Environ.*, 201(3),  
816 195–210, [https://doi.org/10.1016/S0048-9697\(97\)84057-0](https://doi.org/10.1016/S0048-9697(97)84057-0), 1997.
- 817 Diefendorf, A. F., Freeman, K. H. and Wing, S. L.: Distribution and carbon isotope patterns of diterpenoids and  
818 triterpenoids in modern temperate C3 trees and their geochemical significance, *Geochim. Cosmochim. Acta*, 85,  
819 342–356, <https://doi.org/10.1016/j.gca.2012.02.016>, 2012.
- 820 Ducassou, E., Mulder, T., Migeon, S., Gonthier, E., Murat, A., Revel, M., Capotondi, L., Bernasconi, S. M., Mascle,  
821 J. and Zaragosi, S.: Nile floods recorded in deep Mediterranean sediments, *Quat. Res.*, 70(03), 382–391,  
822 <https://doi.org/10.1016/j.yqres.2008.02.011>, 2008.
- 823 Feseker, T., Brown, K. R., Blanchet, C., Scholz, F., Nuzzo, M., Reitz, A., Schmidt, M. and Hensen, C.: Active mud  
824 volcanoes on the upper slope of the western Nile deep-sea fan—first results from the P362/2 cruise of R/V  
825 Poseidon, *Geo-Mar. Lett.*, 30(3–4), 169–186, <https://doi.org/10.1007/s00367-010-0192-0>, 2010.
- 826 Fry, B., Jannasch, H. W., Molyneaux, S. J., Wirsén, C. O., Muramoto, J. A. and King, S.: Stable isotope studies of  
827 the carbon, nitrogen and sulfur cycles in the Black Sea and the Cariaco Trench, *Deep Sea Res. Part Oceanogr. Res.*  
828 *Pap.*, 38, S1003–S1019, [https://doi.org/10.1016/S0198-0149\(10\)80021-4](https://doi.org/10.1016/S0198-0149(10)80021-4), 1991.
- 829 Gallego-Torres, D., Martínez-Ruiz, F., De Lange, G. J., Jiménez-Espejo, F. J. and Ortega-Huertas, M.: Trace-  
830 elemental derived paleoceanographic and paleoclimatic conditions for Pleistocene Eastern Mediterranean  
831 sapropels, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 293(1–2), 76–89,  
832 <https://doi.org/10.1016/j.palaeo.2010.05.001>, 2010.
- 833 Ganopolski, A. and Rahmstorf, S.: Rapid changes of glacial climate simulated in a coupled climate model, *Nature*,  
834 409(6817), 153–158, 2001.
- 835 Garzanti, E., Andò, S., Vezzoli, G., Ali Abdel Megid, A. and El Kammar, A.: Petrology of Nile River sands  
836 (Ethiopia and Sudan): Sediment budgets and erosion patterns, *Earth Planet. Sci. Lett.*, 252(3–4), 327–341,  
837 <https://doi.org/10.1016/j.epsl.2006.10.001>, 2006.

- 838 Garzanti, E., Andò, S., Padoan, M., Vezzoli, G. and El Kammar, A.: The modern Nile sediment system: Processes  
839 and products, *Quat. Sci. Rev.*, 130, 9–56, <https://doi.org/10.1016/j.quascirev.2015.07.011>, 2015.
- 840 Garzanti, E., Vermeesch, P., Rittner, M. and Simmons, M.: The zircon story of the Nile: Time-structure maps of  
841 source rocks and discontinuous propagation of detrital signals, *Basin Res.*, 30(6), 1098–1117,  
842 <https://doi.org/10.1111/bre.12293>, 2018.
- 843 Gebco team: Gebco, Gebco Gen. Bathymetr. Chart Ocean <https://www.gebco.net/>, 2014.
- 844 Goudie, A. S. and Middleton, N. J.: Saharan dust storms: nature and consequences, *Earth-Sci. Rev.*, 56(1–4), 179–  
845 204, [https://doi.org/10.1016/S0012-8252\(01\)00067-8](https://doi.org/10.1016/S0012-8252(01)00067-8), 2001.
- 846 Grimm, R., Maier-Reimer, E., Mikolajewicz, U., Schmiedl, G., Müller-Navarra, K., Adloff, F., Grant, K. M.,  
847 Ziegler, M., Lourens, L. J. and Emeis, K.-C.: Late glacial initiation of Holocene eastern Mediterranean sapropel  
848 formation, *Nat. Commun.*, 6, 7099, <https://doi.org/10.1038/ncomms8099>, 2015.
- 849 Halim, Y., Guergues, S. K. and Saleh, H. H.: Hydrographic Conditions and Plankton in the South East  
850 Mediterranean During the Last Normal Nile Flood (1964), *Int. Rev. Gesamten Hydrobiol. Hydrogr.*, 52(3), 401–  
851 425, <https://doi.org/10.1002/iroh.19670520305>, 1967.
- 852 Hennekam, R., Jilbert, T., Schnetger, B. and de Lange, G. J.: Solar forcing of Nile discharge and sapropel S1  
853 formation in the early to middle Holocene eastern Mediterranean, *Paleoceanography*, 29(5), 343–356,  
854 <https://doi.org/10.1002/2013PA002553>, 2014.
- 855 Hennekam, R., Donders, T. H., Zwiep, K. and de Lange, G. J.: Integral view of Holocene precipitation and  
856 vegetation changes in the Nile catchment area as inferred from its delta sediments, *Quat. Sci. Rev.*, 130, 189–199,  
857 <https://doi.org/10.1016/j.quascirev.2015.05.031>, 2015.
- 858 Hopmans, E. C., Weijers, J. W. H., Schefuß, E., Herfort, L., Sinninghe Damsté, J. S. and Schouten, S.: A novel  
859 proxy for terrestrial organic matter in sediments based on branched and isoprenoid tetraether lipids, *Earth Planet.*  
860 *Sci. Lett.*, 224(1–2), 107–116, <https://doi.org/10.1016/j.epsl.2004.05.012>, 2004.
- 861 Jacob, J., Disnar, J.-R., Boussafir, M., Spadano Albuquerque, A. L., Sifeddine, A. and Turcq, B.: Pentacyclic  
862 triterpene methyl ethers in recent lacustrine sediments (Lagoa do Caçò, Brazil), *Org. Geochem.*, 36(3), 449–461,  
863 <https://doi.org/10.1016/j.orggeochem.2004.09.005>, 2005.
- 864 Jazi, S. D. and Wells, M. G.: Enhanced sedimentation beneath particle-laden flows in lakes and the ocean due to  
865 double-diffusive convection, *Geophys. Res. Lett.*, 43(20), 10,883–10,890, <https://doi.org/10.1002/2016GL069547>,  
866 2016.
- 867 Jilbert, T. and Slomp, C. P.: Iron and manganese shuttles control the formation of authigenic phosphorus minerals  
868 in the euxinic basins of the Baltic Sea, *Geochim. Cosmochim. Acta*, 107, 155–169,  
869 <https://doi.org/10.1016/j.gca.2013.01.005>, 2013.
- 870 Kasten, S., Zabel, M., Heuer, V. and Hensen, C.: Processes and Signals of Nonsteady-State Diagenesis in Deep-  
871 Sea Sediments and their Pore Waters, in *The South Atlantic in the Late Quaternary: Reconstruction of Material*

- 872 Budgets and Current Systems, edited by G. Wefer, S. Mulitza, and V. Ratmeyer, pp. 431–459, Springer, Berlin,  
873 Heidelberg, [https://doi.org/10.1007/978-3-642-18917-3\\_20](https://doi.org/10.1007/978-3-642-18917-3_20), , 2004.
- 874 Keeling, R. F., Körtzinger, A. and Gruber, N.: Ocean Deoxygenation in a Warming World, *Annu. Rev. Mar. Sci.*,  
875 2(1), 199–229, <https://doi.org/10.1146/annurev.marine.010908.163855>, 2009.
- 876 Kubin, E., Poulain, P.-M., Mauri, E., Menna, M. and Notarstefano, G.: Levantine Intermediate and Levantine Deep  
877 Water Formation: An Argo Float Study from 2001 to 2017, *Water*, 11(9), 1781,  
878 <https://doi.org/10.3390/w11091781>, 2019.
- 879 Kuhnt, T., Schmiedl, G., Ehrmann, W., Hamann, Y. and Andersen, N.: Stable isotopic composition of Holocene  
880 benthic foraminifers from the Eastern Mediterranean Sea: Past changes in productivity and deep water oxygenation,  
881 *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 268(1–2), 106–115, <https://doi.org/10.1016/j.palaeo.2008.07.010>, 2008.
- 882 Larrasoaña, J. C., Roberts, A. P., Hayes, A., Wehausen, R. and Rohling, E. J.: Detecting missing beats in the  
883 Mediterranean climate rhythm from magnetic identification of oxidized sapropels (Ocean Drilling Program Leg  
884 160), *Phys. Earth Planet. Inter.*, 156(3–4), 283–293, <https://doi.org/10.1016/j.pepi.2005.04.017>, 2006.
- 885 Le Houedec, S., Mojtahid, M., Bicchi, E., Lange, G. J. de and Hennekam, R.: Suborbital Hydrological Variability  
886 Inferred From Coupled Benthic and Planktic Foraminiferal-Based Proxies in the Southeastern Mediterranean  
887 During the Last 19 ka, *Paleoceanogr. Paleoclimatology*, 35(2), e2019PA003827,  
888 <https://doi.org/10.1029/2019PA003827>, 2020.
- 889 Liu, Q., Larrasoaña, J. C., Torrent, J., Roberts, A. P., Rohling, E. J., Liu, Z. and Jiang, Z.: New constraints on  
890 climate forcing and variability in the circum-Mediterranean region from magnetic and geochemical observations  
891 of sapropels S1, S5 and S6, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 333–334, 1–12,  
892 <https://doi.org/10.1016/j.palaeo.2012.02.036>, 2012.
- 893 Marlowe, I. T., Green, J. C., Neal, A. C., Brassell, S. C., Eglinton, G. and Course, P. A.: Long chain (n-C37–C39)  
894 alkenones in the Prymnesiophyceae. Distribution of alkenones and other lipids and their taxonomic significance,  
895 *Br. Phycol. J.*, 19(3), 203–216, <https://doi.org/10.1080/00071618400650221>, 1984.
- 896 Matthews, A., Azrieli-Tal, I., Benkovitz, A., Bar-Matthews, M., Vance, D., Poulton, S. W., Teutsch, N., Almogi-  
897 Labin, A. and Archer, C.: Anoxic development of sapropel S1 in the Nile Fan inferred from redox sensitive proxies,  
898 Fe speciation, Fe and Mo isotopes, *Chem. Geol.*, 475, 24–39, <https://doi.org/10.1016/j.chemgeo.2017.10.028>,  
899 2017.
- 900 MEDAR Group: MEDATLAS/2002 database. Mediterranean and Black Sea database of temperature salinity and  
901 bio-chemical parameters. Climatological Atlas. (4 CD-Roms), [http://www.ifremer.fr/medar/cdrom\\_database.htm](http://www.ifremer.fr/medar/cdrom_database.htm),  
902 2002.
- 903 Ménot, G., Pivot, S., Bouloubassi, I., Davtian, N., Hennekam, R., Bosch, D., Ducassou, E., Bard, E., Migeon, S.  
904 and Revel, M.: Timing and stepwise transitions of the African Humid Period from geochemical proxies in the Nile  
905 deep-sea fan sediments, *Quat. Sci. Rev.*, 228, 106071, <https://doi.org/10.1016/j.quascirev.2019.106071>, 2020.

- 906 Mercone, D., Thomson, J., Abu-Zied, R. H., Croudace, I. W. and Rohling, E. J.: High-resolution geochemical and  
907 micropalaeontological profiling of the most recent eastern Mediterranean sapropel, *Mar. Geol.*, 177(1), 25–44,  
908 [https://doi.org/10.1016/S0025-3227\(01\)00122-0](https://doi.org/10.1016/S0025-3227(01)00122-0), 2001.
- 909 Michaelis, W., Seifert, R., Nauhaus, K., Treude, T., Thiel, V., Blumenberg, M., Knittel, K., Gieseke, A.,  
910 Peterknecht, K., Pape, T., Boetius, A., Amann, R., Jørgensen, B. B., Widdel, F., Peckmann, J., Pimenov, N. V. and  
911 Gulin, M. B.: Microbial Reefs in the Black Sea Fueled by Anaerobic Oxidation of Methane, *Science*, 297(5583),  
912 1013–1015, <https://doi.org/10.1126/science.1072502>, 2002.
- 913 Mojtahid, M., Manceau, R., Schiebel, R., Hennekam, R. and de Lange, G. J.: Thirteen thousand years of  
914 southeastern Mediterranean climate variability inferred from an integrative planktic foraminiferal-based approach,  
915 *Paleoceanography*, 30(4), 402–422, <https://doi.org/10.1002/2014PA002705>, 2015.
- 916 Mologni, C., Revel, M., Blanchet, C., Bosch, D., Develle, A.-L., Orange, F., Bastian, L., Khalidi, L., Ducassou, E.  
917 and Migeon, S.: Frequency of exceptional Nile flood events as an indicator of Holocene hydro-climatic changes in  
918 the Ethiopian Highlands, *Quat. Sci. Rev.*, 247, 106543, <https://doi.org/10.1016/j.quascirev.2020.106543>, 2020.
- 919 Nixon, S. W.: Replacing the Nile: Are Anthropogenic Nutrients Providing the Fertility Once Brought to the  
920 Mediterranean by a Great River?, *AMBIO J. Hum. Environ.*, 32(1), 30–39, [https://doi.org/10.1579/0044-7447-](https://doi.org/10.1579/0044-7447-921)  
921 32.1.30, 2003.
- 922 Oeschies, A., Dietze, H. and Kähler, P.: Salt-finger driven enhancement of upper ocean nutrient supply, *Geophys.*  
923 *Res. Lett.*, 30(23), <https://doi.org/10.1029/2003GL018552>, 2003.
- 924 Pante, E. and Simon-Bouhet, B.: marmap: A Package for Importing, Plotting and Analyzing Bathymetric and  
925 Topographic Data in R, *PLOS ONE*, 8(9), e73051, <https://doi.org/10.1371/journal.pone.0073051>, 2013.
- 926 Parsons, J. D., Bush, J. W. M. and Syvitski, J. P. M.: Hyperpycnal plume formation from riverine outflows with  
927 small sediment concentrations, *Sedimentology*, 48(2), 465–478, [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-928)  
928 3091.2001.00384.x, 2001.
- 929 Pinardi, N., Zavatarelli, M., Adani, M., Coppini, G., Fratianni, C., Oddo, P., Simoncelli, S., Tonani, M., Lyubartsev,  
930 V., Dobricic, S. and Bonaduce, A.: Mediterranean Sea large-scale low-frequency ocean variability and water mass  
931 formation rates from 1987 to 2007: A retrospective analysis, *Prog. Oceanogr.*, 132, 318–332,  
932 <https://doi.org/10.1016/j.pocean.2013.11.003>, 2015.
- 933 Powley, H. R., Krom, M. D. and Van Cappellen, P.: Circulation and oxygen cycling in the Mediterranean Sea:  
934 Sensitivity to future climate change, *J. Geophys. Res. Oceans*, 121, 8230–8247, 2016.
- 935 Pujol, C. and Grazzini, C. V.: Distribution patterns of live planktic foraminifers as related to regional hydrography  
936 and productive systems of the Mediterranean Sea, *Mar. Micropaleontol.*, 25(2), 187–217,  
937 [https://doi.org/10.1016/0377-8398\(95\)00002-I](https://doi.org/10.1016/0377-8398(95)00002-I), 1995.
- 938 Pujol-Pay, M., Conan, P., Oriol, L., Cornet-Barthaux, V., Falco, C., Ghiglione, J.-F., Goyet, C., Moutin, T. and  
939 Prieur, L.: Integrated survey of elemental stoichiometry (C, N, P) from the western to eastern Mediterranean Sea,  
940 *Biogeosciences*, 8(4), 883–899, <https://doi.org/10.5194/bg-8-883-2011>, 2011.



- 941 R Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing,  
942 Vienna, Austria. <http://www.R-project.org/>, 2013.
- 943 Revel, M., Ducassou, E., Skonieczny, C., Colin, C., Bastian, L., Bosch, D., Migeon, S. and Mascle, J.: 20,000 years  
944 of Nile River dynamics and environmental changes in the Nile catchment area as inferred from Nile upper  
945 continental slope sediments, *Quat. Sci. Rev.*, 130, 200–221, <https://doi.org/10.1016/j.quascirev.2015.10.030>, 2015.
- 946 Riedinger, N., Brunner, B., Lin, Y.-S., Voßmeyer, A., Ferdelman, T. G. and Jørgensen, B. B.: Methane at the  
947 sediment–water transition in Black Sea sediments, *Chem. Geol.*, 274(1), 29–37,  
948 <https://doi.org/10.1016/j.chemgeo.2010.03.010>, 2010.
- 949 Robinson, A. R., Malanotte-Rizzoli, P., Hecht, A., Michelato, A., Roether, W., Theocharis, A., Ünlüata, Ü., Pinardi,  
950 N., Artegiani, A., Bergamasco, A., Bishop, J., Brenner, S., Christianidis, S., Gacic, M., Georgopoulos, D.,  
951 Golnaraghi, M., Hausmann, M., Junghaus, H.-G., Lascaratos, A., Latif, M. A., Leslie, W. G., Lozano, C. J., Oguz,  
952 T., Özsoy, E., Papageorgiou, E., Paschini, E., Rozentroub, Z., Sansone, E., Scarazzato, P., Schlitzer, R., Spezie,  
953 G.-C., Tziperman, E., Zodiatis, G., Athanassiadou, L., Gerges, M. and Osman, M.: General circulation of the  
954 Eastern Mediterranean, *Earth-Sci. Rev.*, 32(4), 285–309, [https://doi.org/10.1016/0012-8252\(92\)90002-B](https://doi.org/10.1016/0012-8252(92)90002-B), 1992.
- 955 Roether, W., Manca, B. B., Klein, B., Bregant, D., Georgopoulos, D., Beitzel, V., Kovačević, V. and Luchetta, A.:  
956 Recent Changes in Eastern Mediterranean Deep Waters, *Science*, 271(5247), 333–335,  
957 <https://doi.org/10.1126/science.271.5247.333>, 1996.
- 958 Roether, W., Klein, B., Manca, B. B., Theocharis, A. and Kioroglou, S.: Transient Eastern Mediterranean deep  
959 waters in response to the massive dense-water output of the Aegean Sea in the 1990s, *Prog. Oceanogr.*, 74(4), 540–  
960 571, <https://doi.org/10.1016/j.pocean.2007.03.001>, 2007.
- 961 Rogerson, M., Colmenero-Hidalgo, E., Levine, R. C., Rohling, E. J., Voelker, A. H. L., Bigg, G. R., Schönfeld, J.,  
962 Cacho, I., Sierro, F. J., Löwemark, L., Reguera, M. I., Abreu, L. de and Garrick, K.: Enhanced Mediterranean-  
963 Atlantic exchange during Atlantic freshening phases, *Geochem. Geophys. Geosystems*, 11(8),  
964 <https://doi.org/10.1029/2009GC002931>, 2010.
- 965 Rogerson, M., Rohling, E. J., Bigg, G. R. and Ramirez, J.: Paleocyanography of the Atlantic-Mediterranean  
966 exchange: Overview and first quantitative assessment of climatic forcing, *Rev. Geophys.*, 50(2),  
967 <https://doi.org/10.1029/2011RG000376>, 2012.
- 968 Rohling, E. J.: Review and new aspects concerning the formation of eastern Mediterranean sapropels, *Mar. Geol.*,  
969 122(1), 1–28, [https://doi.org/10.1016/0025-3227\(94\)90202-X](https://doi.org/10.1016/0025-3227(94)90202-X), 1994.
- 970 Rohling, E. J., Marino, G. and Grant, K. M.: Mediterranean climate and oceanography, and the periodic  
971 development of anoxic events (sapropels), *Earth-Sci. Rev.*, 143, 62–97,  
972 <https://doi.org/10.1016/j.earscirev.2015.01.008>, 2015.
- 973 Rossignol-Strick, M.: Mediterranean Quaternary sapropels, an immediate response of the African monsoon to  
974 variation of insolation, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 49(3–4), 237–263, [https://doi.org/10.1016/0031-0182\(85\)90056-2](https://doi.org/10.1016/0031-0182(85)90056-2), 1985.

- 976 Rossignol-Strick, M., Nesteroff, W., Olive, P. and Vergnaud-Grazzini, C.: After the deluge: Mediterranean  
977 stagnation and sapropel formation, *Nature*, 295(5845), 105–110, <https://doi.org/10.1038/295105a0>, 1982.
- 978 Rush, D., Talbot, H. M., Meer, M. T. J. van der, Hopmans, E. C., Douglas, B. and Sinninghe Damsté, J. S.:  
979 Biomarker evidence for the occurrence of anaerobic ammonium oxidation in the eastern Mediterranean Sea during  
980 Quaternary and Pliocene sapropel formation, *Biogeosciences*, 16(12), 2467–2479, [https://doi.org/10.5194/bg-16-](https://doi.org/10.5194/bg-16-2467-2019)  
981 2467-2019, 2019.
- 982 Rutten, A. and de Lange, G. J.: Sequential extraction of iron, manganese and related elements in S1 sapropel  
983 sediments, eastern Mediterranean, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 190, 79–101,  
984 [https://doi.org/10.1016/S0031-0182\(02\)00600-4](https://doi.org/10.1016/S0031-0182(02)00600-4), 2003.
- 985 Schimmelmann, A., Lange, C. B., Schieber, J., Francus, P., Ojala, A. E. K. and Zolitschka, B.: Varves in marine  
986 sediments: A review, *Earth-Sci. Rev.*, 159, 215–246, <https://doi.org/10.1016/j.earscirev.2016.04.009>, 2016.
- 987 Schlitzer, R.: Ocean Data View, Alfred-Wegener Institute AWI Bremerhaven. [odv.awi.de](http://odv.awi.de), 2018.
- 988 Schmiedl, G., Kuhnt, T., Ehrmann, W., Emeis, K.-C., Hamann, Y., Kotthoff, U., Dulski, P. and Pross, J.: Climatic  
989 forcing of eastern Mediterranean deep-water formation and benthic ecosystems during the past 22?000 years, *Quat.*  
990 *Sci. Rev.*, 29(23–24), 3006–3020, <https://doi.org/10.1016/j.quascirev.2010.07.002>, 2010.
- 991 Schmitt, R. W., Ledwell, J. R., Montgomery, E. T., Polzin, K. L. and Toole, J. M.: Enhanced Diapycnal Mixing by  
992 Salt Fingers in the Thermocline of the Tropical Atlantic, *Science*, 308(5722), 685–688,  
993 <https://doi.org/10.1126/science.1108678>, 2005.
- 994 Schouten, S., Wakeham, S. G. and Damsté, J. S. S.: Evidence for anaerobic methane oxidation by archaea in euxinic  
995 waters of the Black Sea, *Org. Geochem.*, 32(10), 1277–1281, [https://doi.org/10.1016/S0146-6380\(01\)00110-3](https://doi.org/10.1016/S0146-6380(01)00110-3),  
996 2001.
- 997 Schouten, S., Huguet, C., Hopmans, E. C., Kienhuis, M. V. M. and Sinninghe Damsté, J. S.: Analytical  
998 Methodology for TEX<sub>86</sub> Paleothermometry by High-Performance Liquid Chromatography/Atmospheric Pressure  
999 Chemical Ionization-Mass Spectrometry, *Anal. Chem.*, 79(7), 2940–2944, <https://doi.org/10.1021/ac062339v>,  
000 2007.
- 001 Sinninghe Damsté, J. S.: Spatial heterogeneity of sources of branched tetraethers in shelf systems: The  
002 geochemistry of tetraethers in the Berau River delta (Kalimantan, Indonesia), *Geochim. Cosmochim. Acta*, 186,  
003 13–31, <https://doi.org/10.1016/j.gca.2016.04.033>, 2016.
- 004 Sinninghe Damsté, J. S., Kuypers, M. M. M., Schouten, S., Schulte, S. and Rullkötter, J.: The lycopane/C31 n-  
005 alkane ratio as a proxy to assess palaeoacidity during sediment deposition, *Earth Planet. Sci. Lett.*, 209(1–2), 215–  
006 226, [https://doi.org/10.1016/S0012-821X\(03\)00066-9](https://doi.org/10.1016/S0012-821X(03)00066-9), 2003.
- 007 Tachikawa, K., Vidal, L., Cornuault, M., Garcia, M., Pothin, A., Sonzogni, C., Bard, E., Menot, G. and Revel, M.:  
008 Eastern Mediterranean Sea circulation inferred from the conditions of S1 sapropel deposition, *Clim. Past*, 11, 855–  
009 867, 2015.

- 010 Tang, C. M. and Stott, L. D.: Seasonal salinity changes during Mediterranean sapropel deposition 9000 years B.P.:  
011 Evidence from isotopic analyses of individual planktonic foraminifera, *Paleoceanography*, 8(4), 473–493,  
012 <https://doi.org/10.1029/93PA01319>, 1993.
- 013 Vadsaria, T., Ramstein, G., Dutay, J.-C., Li, L., Ayache, M. and Richon, C.: Simulating the Occurrence of the Last  
014 Sapropel Event (S1): Mediterranean Basin Ocean Dynamics Simulations Using Nd Isotopic Composition  
015 Modeling, *Paleoceanogr. Paleoclimatology*, 34(2), 237–251, <https://doi.org/10.1029/2019PA003566>, 2019.
- 016 Volkman, J. K.: Ecological and environmental factors affecting alkenone distributions in seawater and sediments,  
017 *Geochem. Geophys. Geosystems*, 1(9), <https://doi.org/10.1029/2000GC000061>, 2000.
- 018 Weldeab, S., Menke, V. and Schmiedl, G.: The pace of East African monsoon evolution during the Holocene,  
019 *Geophys. Res. Lett.*, 41(5), 1724–1732, <https://doi.org/10.1002/2014GL059361>, 2014.
- 020 Weltje, G. J. and Tjallingii, R.: Calibration of XRF core scanners for quantitative geochemical logging of sediment  
021 cores: Theory and application, *Earth Planet. Sci. Lett.*, 274(3–4), 423–438,  
022 <https://doi.org/10.1016/j.epsl.2008.07.054>, 2008.
- 023 Weltje, G. J., Bloemsa, M. R., Tjallingii, R., Heslop, D., Röhl, U. and Croudace, I. W.: Prediction of Geochemical  
024 Composition from XRF Core Scanner Data: A New Multivariate Approach Including Automatic Selection of  
025 Calibration Samples and Quantification of Uncertainties, in *Micro-XRF Studies of Sediment Cores*, vol. 17, edited  
026 by I. W. Croudace and R. G. Rothwell, pp. 507–534, Springer Netherlands, Dordrecht, [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-027-94-017-9849-5_21)  
027 94-017-9849-5\_21, , 2015.
- 028 Xiao, W., Wang, Y., Zhou, S., Hu, L., Yang, H. and Xu, Y.: Ubiquitous production of branched glycerol dialkyl  
029 glycerol tetraethers (brGDGTs) in global marine environments: a new source indicator for brGDGTs,  
030 *Biogeosciences*, 13(20), 5883–5894, <https://doi.org/10.5194/bg-13-5883-2016>, 2016.
- 031 Zirks, E., Krom, M. D., Zhu, D., Schmiedl, G. and Goodman-Tchernov, B. N.: Evidence for the Presence of  
032 Oxygen-Depleted Sapropel Intermediate Water across the Eastern Mediterranean during Sapropel S1, *ACS Earth  
033 Space Chem.*, <https://doi.org/10.1021/acsearthspacechem.9b00128>, 2019.
- 034 Zirks, E., Krom, M., Schmiedl, G., Katz, T., Xiong, Y., Alcott, L. J., Poulton, S. W. and Goodman-Tchernov, B.:  
035 Redox evolution and the development of oxygen minimum zones in the Eastern Mediterranean Levantine basin  
036 during the early Holocene, *Geochim. Cosmochim. Acta*, 297, 82–100, <https://doi.org/10.1016/j.gca.2021.01.009>,  
037 2021.
- 038 Zwiep, K. L., Hennekam, R., Donders, T. H., van Helmond, N. A. G. M., de Lange, G. J. and Sangiorgi, F.: Marine  
039 productivity, water column processes and seafloor anoxia in relation to Nile discharge during sapropels S1 and S3,  
040 *Quat. Sci. Rev.*, 200, 178–190, <https://doi.org/10.1016/j.quascirev.2018.08.026>, 2018.
- 041