

Extending and understanding the South West Western Australian rainfall record using a snowfall reconstruction from Law Dome, East Antarctica

Yaowen Zheng^{1,2}, Lenneke M. Jong^{3,4}, Steven J. Phipps^{1,5}, Jason L. Roberts^{3,4}, Andrew D. Moy^{3,4}, Mark A. J. Curran^{3,4}, and Tas D. van Ommen^{3,4}

¹Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

²Antarctic Research Centre, Victoria University of Wellington, Wellington, New Zealand

³Australian Antarctic Division, Kingston, Tasmania, Australia

⁴Australian Antarctic Program Partnership, Institute of Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

⁵Ikigai Research, Hobart, Tasmania, Australia

Correspondence: Yaowen Zheng (yaowen.zheng@vuw.ac.nz)

Abstract.

South West Western Australia (SWWA) has experienced a prolonged reduction in rainfall in recent decades, with associated reductions in regional water supply and residential and agricultural impacts. The cause of the reduction has been widely considered, but remains unclear. The relatively short length of the instrumental record limits long-term investigation. A previous proxy-based study used a statistically negative correlation between SWWA rainfall and snowfall from the Dome Summit South (DSS) ice core drilling site, Law Dome, East Antarctica and concluded that the anomaly of recent decades is unprecedented over the \sim 750 year period of the study (1250–2004 CE). Here we extend the snow accumulation record to cover the period 22 BCE–2015 CE and derive a rainfall reconstruction over this extended period. This extended record confirms that the recent anomaly is unique in the period since 1250 CE and unusual over the full \sim 2000 year period, with just two other earlier droughts of similar duration and intensity. The reconstruction shows that SWWA rainfall started to reduce around 1971 CE. Ensembles of climate model simulations are used to investigate the potential roles of natural variability and external climate drivers in explaining changes in SWWA rainfall. We find that anthropogenic greenhouse gases are likely to have contributed towards the SWWA rainfall drying trend after 1971 CE. However, natural variability may also have played a role in determining the timing and magnitude of the reduction in rainfall.

15 1 Introduction

South West Western Australia (SWWA) is characterized as a region with a Mediterranean-like climate (Yu and Neil, 1993; Siddique et al., 1999; Timbal et al., 2006). Most of the rainfall occurs during the winter and spring seasons, comprising around 70% of the annual total rainfall (Wright, 1974a; Hennessy et al., 1999; Ludwig and Asseng, 2006; Timbal et al., 2006). The seasonal concentration of rainfall makes May to October (approximately) the growing season for SWWA (Watson

20 and Lapins, 1969; French and Schultz, 1984; Anderson et al., 1996; Ward and Dunin, 2001). In comparison to other regions of
Western Australia (WA), SWWA has a more constant supply of water throughout the year for agriculture, industry and residents
(Wright, 1974a, b; Anderson et al., 1996; Pitman et al., 2004; Power et al., 2005; Ludwig and Asseng, 2006). Abundant rainfall
contributes to suitable conditions for dryland farming such as wheat with a yearly sowed area of more than 4,000,000 ha
(Ludwig and Asseng, 2006). It also directly provides water supply to the WA capital city Perth (Pitman et al., 2004; Power
25 et al., 2005). A number of studies have found the growing season rainfall in SWWA has decreased (Pittock, 1983; Li et al.,
2005; Cai and Cowan, 2006; Samuel et al., 2006; Timbal et al., 2006; Ludwig et al., 2009; Feng et al., 2010). Rainfall in this
region has decreased by 22% in August and 30% in September for the period 1946–1978 compared to the period 1913–1945
(Pittock, 1983), and has weakened by about 20% from the 1960s to the beginning of the 21st century (Cai and Cowan, 2006).
Winter rainfall has decreased by 15%-20% from the 1970s to the early 21st century (van Ommen and Morgan, 2010). This
30 sizable winter rainfall reduction contributes to decades of persistent drought since 1975 (Ansell et al., 2000; Cai and Cowan,
2006; Hope et al., 2006) causing water supply problems in WA (Pitman et al., 2004; Samuel et al., 2006). The SWWA rainfall
reduction occurring in the mid-20th century decreased the water supply to Perth by about 42% (Pitman et al., 2004). This has
required an additional investment of around \$300 million to develop alternative water sources (Pitman et al., 2004). This long-
term ongoing drought poses a potential threat to residential water supplies, industrial and agricultural production and makes
35 the study of this phenomenon and the determination of its driving factors urgent.

The large-scale climate drivers for both Australia and specifically the SWWA region have been investigated over the past
decades. The El Niño–Southern Oscillation (ENSO), specifically the equatorial Pacific Ocean sea surface temperature anomaly
in the Nino 3.4 region (Trenberth and Hoar, 1997), tends to be associated with interannual rainfall variations in Australia (Cai
et al., 2011), including dry conditions (Chiew et al., 1998). However, this teleconnection linking ENSO and Australian rainfall
40 is stronger in Eastern Australia than in SWWA. The Indian Ocean Dipole (IOD) shows a robust relationship during the growing
season from May to October (Fierro and Leslie, 2013) with the rainfall in southern and western regions of Australia (Ashok
et al., 2003), but not specifically to the SWWA region. The Southern Annular Mode (SAM), or the Antarctic Oscillation (AAO),
is a large-scale mode of climate variability that is correlated with rainfall in WA (Gong and Wang, 1999; Thompson et al.,
2011; Fierro and Leslie, 2013). The relationship between the SAM and SWWA rainfall is seasonal, with the SAM influencing
45 rainfall in June-July-August (JJA) but not in December-January-February (DJF) (Cai and Shi, 2005; Cai and Cowan, 2006).
The SAM has experienced a shift towards a more positive phase since the 1970s, which can be attributed to the depletion of
stratospheric ozone (Thompson and Solomon, 2002; Gillett and Thompson, 2003). This shift, in conjunction with the increase
in anthropogenic greenhouse gases over this period, may be responsible for at least part of the reduction in SWWA rainfall
(Cai and Shi, 2005).

50 Variability in the SAM is not the sole driver of SWWA rainfall anomalies; for example local Indian Ocean sea surface tem-
peratures can also play a role. In particular, negative sea surface temperature anomalies in the eastern Indian Ocean and positive
sea surface temperature anomalies in the central subtropical Indian Ocean are related to dry years in SWWA (Ummenhofer
et al., 2008). Some indication was found that mean sea level pressure anomalies over the Indian Ocean drive Indian Ocean sea
surface temperature anomalies and SWWA rainfall, but this link did not appear robust at the interannual time scale (Smith et al.,

55 2000). A link between SWWA rainfall extremes and large-scale Indian Ocean climate was found due to moisture advection onto the SWWA coast (England et al., 2006). This is subject to influence from the large-scale wind field over the eastern and southeastern Indian Ocean, which may contribute to SWWA rainfall extremes (England et al., 2006). This link between the Indian Ocean and SWWA rainfall may be influenced by the IOD–ENSO link (England et al., 2006). Thus, the tropical and subtropical Indian and Pacific Oceans are both likely to play a role in SWWA rainfall variations, but not the only role. Changes
60 of a similar magnitude to those observed can potentially also arise through natural multidecadal climate variability (Cai and Shi, 2005). Thus the drivers of the winter rainfall attenuation in SWWA are still unclear.

The relatively short length of the observational record in SWWA limits long-term studies of rainfall. Rainfall studies rely on Australian Government Bureau of Meteorology (BoM) instrumental station data. The earliest record goes back to around the 1880s, while most of the station records started at around 1900. The availability of 120 years of rainfall data is insufficient
65 to investigate the SWWA long-term rainfall evolution in history and makes it difficult to determine the uncertainties of the climate drivers in SWWA. Connections between mid-latitude climate and that of coastal East Antarctica have been reported for some time (Goodwin et al., 2004; van Ommen and Morgan, 2010). The strength and position of the southern hemisphere westerlies are important for coastal Antarctic snowfall rates, especially for cyclonically driven locations such as Law Dome, East Antarctica (Bromwich, 1988). The specific relationship between SWWA and snowfall recorded in the Dome Summit
70 South (DSS) ice core from Law Dome was reported by van Ommen and Morgan (2010) who found a statistically significant anticorrelation between winter (JJA) mean SWWA regional rainfall and Law Dome snow accumulation. This link, which accounts for up to 40% of the shared variance on interannual to decadal timescales is associated with simultaneous anomalous northward flow of relatively cool, dry air to SWWA and southward flow of relatively warm, dry air to the Law Dome region. This teleconnection pattern is characterised by a broadly zonal wave three pattern in 500 hPa geopotential field. The northward
75 flow to SWWA, while bringing showers to the southern coast, is distinct from the higher rainfall patterns which bring prefrontal rain from north/north-westerly directions (van Ommen and Morgan, 2010). More recently, van Ommen and Morgan (2010)'s work has been extended using a longer 2035-year accumulation record from the Law Dome core (Roberts et al., 2015). As with the earlier accumulation record, this was dated by determining annual layers in the seasonally varying water stable isotope ratios and trace ions from multiple ice cores drilled at the DSS site (66.7697°S, 112.8069°E, 1370 m elevation) (Roberts et al.,
80 2015). Taken together, the relationship between SWWA rainfall and DSS snowfall and the DSS long-term ice core record allows us to extend the SWWA rainfall record to span the past 2000 years.

In this study, we mainly focus on reconstructing growing season (May to October) SWWA rainfall over the past two thousand years and answering the questions about changes in SWWA rainfall since around 1970s and the drivers of these changes by comparing observational data with climate model simulations. We calculate and test the significance of the correlation
85 between growing season SWWA rainfall and DSS snow accumulation. We individually calculate and test the significance of the correlation based on 116 years of the Australian Water Availability Project (AWAP) gridded growing season rainfall data for 110 Local Government Areas (Australian Government, 2020) and combine the statistically significant ($p < 0.05$) areas to define the SWWA region of significance. We assume stationarity and build a linear model between the growing season rainfall in the significant region and the DSS snow accumulation and use this model to reconstruct the growing season rainfall in

90 SSWA from 2015 CE back to 22 BCE. We then compare the reconstruction with ensembles of simulations conducted using the Commonwealth Scientific and Industrial Research Organisation Mark 3L (CSIRO Mk3L) model (Phipps et al., 2011, 2012, 2013). This allows us to explore the drivers of the changes in rainfall in SSWA rainfall and place them in a longer-term context.

2 Data

2.1 DSS ice core record

95 DSS annual snow accumulation is calculated by dating the ice cores drilled at DSS (Roberts et al., 2015), East Antarctica. The length of this DSS snow accumulation record has been extended to 2035-years, from 2012 CE back to 22 BCE (Roberts et al., 2015).

Table 1. Observational data types include two DSS snow accumulation records, AWAP gridded rainfall and BoM stations data.

Data type	Time period	Reference
Law Dome Summit accumulation composite*	22 BCE–2012 CE	Roberts et al. (2015)
Law Dome Summit accumulation composite ^Δ	22 BCE–2015 CE	This study
AWAP Gridded rainfall	1900–2018 CE	Jones et al. (2009)
BoM station data	around 1900s–2019 CE	Lavery et al. (1997)

*DSS accumulation composite – DSS-main, DSS99, DSS97 and DSS1213.

^ΔDSS accumulation composite – DSS-main, DSS99, DSS97 and DSS1617.

In addition, the latest DSS snow accumulation data has been extended to 2015 CE using a recent new ice core (DSS1617). We use the annual layer thickness data from this new core (DSS1617) for the period 1990 to 2015 CE to extend and replace the 100 DSS1213 annual layer thickness data for the period 1990–2012 used in Roberts et al. (2015), and calculate the corresponding snow accumulation record using their Power-law vertical strain rate model. Therefore, the DSS snow accumulation record we use in this study will be a 2038-year long-term record from 22 BCE to 2015 CE.

2.2 SSWA instrumental rainfall record

The SSWA rainfall station records are monthly records which are obtained from BoM. We initially selected 16 stations, 105 which are as the same as van Ommen and Morgan (2010). Since the instrumental rainfall records for station Avondale Farm (ID: 10795, 116.87°E 32.12°S) are only available for 51 years (1965–2015), which is substantially less than the average of remaining 15 stations records length (around 118 years), we will not discuss the rainfall records of this station any further. The other 15 stations cover the 1900s to 2019 CE.

To investigate more completely the spatial variability and reconstruct SSWA rainfall, we use AWAP Australian Gridded Climate Data (AGCD) (Jones et al., 2009). This AWAP data is gridded data that has 0.05° longitude * 0.05° latitude geospatial resolution and 119 years' availability from 1900 CE to 2018 CE.

2.3 CSIRO Mk3L climate model simulations

The model simulations used for paleoclimate data-model comparison are from the CSIRO Mk3L climate model. CSIRO Mk3L is a reduced-resolution coupled general circulation model, comprising components that describe the atmosphere, ocean, sea ice and land surface (Phipps et al., 2011, 2012). The model is explicitly designed for studying climate variability and change on millennial time scales. The atmospheric component of CSIRO Mk3L is taken from the CSIRO Mk3 climate model (Gordon et al., 2002), but with reduced horizontal resolution. Both CSIRO Mk3 and CSIRO Mk3L produce credible simulations of large-scale precipitation, including over Australia (Cai et al., 2003; Cai and Shi, 2005; England et al., 2006; Phipps et al., 2011). The model is also skilful at capturing the dominant modes of large-scale variability in the Southern Hemisphere — ENSO and SAM — including the teleconnections between these modes and precipitation over Australia (Cai et al., 2003; Phipps et al., 2011, 2013; Abram et al., 2014; Barr et al., 2019).

In this study, we use four ensembles of simulations based on a combination of orbital forcing (O), greenhouse gases (G), solar irradiance (S) and volcanic aerosols (V) (Phipps et al., 2013). The four ensembles (O, OG, OGS and OGSV) are generated by combining each of these forcings (Table 2). Each ensemble contains three members, which differ only in that they are initialised from different years of a pre-industrial control simulation. As the initial states therefore differ, the internal variability will be different between each member but the responses to external forcings will be consistent.

Table 2. The forcing(s) and duration for each ensemble of the CSIRO Mk3L model simulations

Ensemble	Forcing(s)	Duration
O	Orbital	1 CE–2000 CE
OG	Orbital, greenhouse gases	1 CE–2000 CE
OGS	Orbital, greenhouse gases, solar irradiance	1 CE–2000 CE
OGSV	Orbital, greenhouse gases, solar irradiance, volcanic aerosols	501 CE–2000 CE

To compare the CSIRO Mk3L simulations with the SSWA rainfall reconstruction, we use all 12 simulations, three for each of the four ensembles. The simulations have a geospatial resolution of 5.625° longitude by 3.1857° latitude. The model has been run globally for 2000 years from 1 CE to 2000 CE and the simulations have been published (Phipps et al., 2013). For 9 simulations, three members for each of the O, OG and OGS ensembles, 2000 years of simulation is available covering the period from 1 to 2000 CE. For the three members of ensemble OGSV, 1500 years of data is available covering the period from 501 to 2000 CE. The model simulations consist of monthly mean data.

All the data we use in this study, including the stations rainfall, gridded rainfall, DSS record and model simulations, are for the growing season from May to October.

135 **3 Methods**

3.1 Normality testing

We use one-sample Kolmogorov-Smirnov tests (hereafter KS test) to assess the normality of the observational data. There is statistically significant evidence that the 15 BoM stations data (interpreted periods up to 2015 CE), AWAP gridded data and DSS snow accumulation data (interpreted the period 1900 CE to 2015 CE) all fit the normal distribution corresponding to their mean
140 and standard deviation, respectively. A visual validation, which consists of comparing the differences between the empirical and normal cumulative distribution functions (CDFs), is presented in Supplement Figure S1. The 15 BoM stations, AWAP gridded rainfall and DSS snow accumulation data empirical CDFs approximately fit their corresponded normal CDFs with minimal biases (Supplement Figure S1). With no irregularity between the empirical and normal CDFs, the visual validation for the observational data is consistent with the results of a one-sample KS test. Therefore, we are confident that the data is
145 described by a normal distribution.

3.2 Significance testing and region definition

Low-pass filtering (or smoothing) the data increases the correlation of the precipitation time-series between SWWA and Law Dome (van Ommen and Morgan, 2010). Annual-scale noise arises from site surface processes including local snow deposition variability which is ameliorated by the smoothing. In order to determine the optimal window size for running average smoothing, we test smoothing using windows sizes of 1–10 years on SWWA and DSS data and then calculate the Pearson correlation coefficients. As smoothing reduces the temporal degrees of freedom, we estimate the Effective Sample Sizes (ESS) or temporal degrees of freedom (Bretherton et al., 1999) by first calculating the lag-1 autocorrelation for each sample and then apply the following formula,
150

$$\text{ESS} = N \frac{1 - r_1 r_2}{1 + r_1 r_2} \quad (1)$$

155 where N is the sample size, r_1 and r_2 are the lag-1 autocorrelation coefficients corresponded to the two samples (Bretherton et al., 1999). Next, we calculate Student's t-statistic using:

$$t_S = r \sqrt{\frac{\text{ESS} - 2}{1 - r^2}} \quad (2)$$

160 where r is the Pearson correlation coefficient and then compute the corresponding p-value. We repeat the above calculations for both the gridded and stations rainfall data matching with DSS snow accumulation data. A six-year window (Supplement Figure S2) maximizes the area of the correlation between AWAP gridded rainfall and DSS snow accumulation with statistical significance. For the 15 BoM stations data, the number of the stations to show a statistically significant correlation is maximized

by a window size of 5-years (Supplement Figure S2). The rainfall changes and its drivers we are interested in this study is for the SWWA region instead of separate stations. For consistency with the AWAP gridded data, we also use a six-year window for the BoM stations data

165 To define a region with statistically significant correlation between observed (AWAP) and reconstructed (DSS) rainfall over SWWA, we independently calculate the correlation coefficient and test its significance for 110 Local Government Areas (Australian Goverment, 2020) (Supplement Table S1) in WA. There are 9 Local Government Areas smaller than the 0.05° longitude * 0.05° latitude geospatial resolution of the AWAP data grid. Therefore, we actually calculate and test 101 areas. There are 52 areas that are statistically significant (6-year window, $p < 0.05$). We combine these 52 areas to define the significant 170 region (Figure 1) over SWWA. For convenience, we name this significant region as "MASK".




Figure 1. The correlation map for the southwest part of WA region for 6-year window AWAP rainfall and DSS snow accumulation from 1900 CE to 2015 CE. The outline area (green line) is the MASK region where the correlation is statistically significant ($p < 0.05$). Red diagonal line connecting $115^\circ\text{E } 30^\circ\text{S}$ and $120^\circ\text{E } 35^\circ\text{S}$ is the boundary of SWWA (van Ommen and Morgan, 2010). Perth is the capital city of WA. Boyanup, Wonnaminta, Arthur River and Cranbrook are the four significant (6-year window, $p < 0.05$) stations.

175 The correlation map (Figure 1) shows the correlation coefficients for 6-year smoothed AWAP rainfall and DSS snow accumulation. The map shows that the strongly negatively correlated ($r \leq -0.5$) areas are mainly concentrated in the southwest corner. Areas around the top right show positive correlations with no statistical significance ($p > 0.05$). MASK covers almost the whole SWWA region apart from some coastal areas in the south. Southern coastal areas show no statistical significance ($p > 0.05$), consistent with the findings of van Ommen and Morgan (2010) who reported the south coast local rainfall related to onshore northward flow that weakens the anticorrelation with DSS that prevails across the rest of the region.

In order to quantify the MASK correlation coefficient and evaluate the statistical significance, we multiply the mask matrix (in the region has a value of 1 and outside the region has a value of 0) of the MASK with the AWAP gridded data to generate the MASK rainfall. Hereafter, the SSWA rainfall we are describing, is the MASK rainfall where the MASK region is delineated in Figure 1. Then we calculate the SSWA rainfall correlation coefficient with DSS and test its statistical significance.

Table 3. The Pearson correlation coefficients for the SSWA rainfall and the four BoM stations rainfall with the DSS snow accumulation. R^2 is the square of the correlation coefficient. All the correlations are statistically significant (6-year window, $p < 0.05$).

Sample	Correlation coefficient	R^2	Year (CE)
SSWA	-0.529	0.280	1900–2015
Arthur River	-0.548	0.300	1891–2015
Boyanup	-0.623	0.388	1898–2013
Cranbrook	-0.540	0.292	1891–2015
Wonnaminta	-0.546	0.298	1905–2015

180

Table 3 integrates the results of the significance testing and correlations. There is statistical significance ($p < 0.05$) that the SSWA rainfall and DSS snow accumulation are strongly negative correlated. BoM stations rainfall also show consistent results. From the 15 stations tested, there are statistically significant ($p < 0.05$) correlations with DSS for four stations. These 185 four stations are all geographically located in the MASK region (Figure 1) showing the consistency with the significance of MASK. Five other stations within the region have correlations of a similar magnitude, but these correlations are not significant at the 5% probability level (Supplement Figure S4). The squares of the correlation coefficients show that the explained variance is around 30-40%. We note that the tropics and subtropics can also play an important role in driving rainfall changes in SSWA (Smith et al., 2000; England et al., 2006; Ummenhofer et al., 2008). Using Law Dome accumulation to reconstruct SSWA rainfall therefore focuses on the SAM-related component of the rain-bearing systems, not the tropical or subtropical 190 components. However, our analysis shows that it makes a valuable contribution to our ability to reconstruct past climate and so we construct a linear model for SSWA rainfall and DSS snow accumulation.

3.3 Linear model construction

The time series and the scatter plot for SSWA rainfall and DSS snow accumulation are shown in Figure 2a and Figure 2b, respectively. The data show a generally linear distribution with a negative slope (Figure 2b). We use Ordinary Least Square 195 linear regression to construct a linear model to SSWA rainfall and DSS snow accumulation (for the four BoM stations, see Supplement Figure S3), and estimate the 95% confidence interval (CI) in the gradient and the intercept as 1.96 multiplied by the standard error of the model.




Figure 2. (a) The time series of anomalies for AWAP rainfall in the MASK region and DSS snow accumulation (6-year running averages) for the period 1900–2015 CE. (b) The scatter plot for AWAP rainfall in the MASK region and DSS snow accumulation (6-year running averages) for the period 1900–2015 CE with their linear model and 95% confidence interval. (c) The probability density function histogram for the model residuals. (d) The scatter plot for model residuals and fitted data with their linear fit and 95% confidence interval.

The model is Rain = Snow * (-0.389 ± 0.114) + 672 ± 82 mm/year in the growing season for the period 1900 CE to 2015 CE. The gradient interval $[-0.503, -0.274]$ is always negative, which is consistent with the result from the Pearson correlation test, 200 that the coefficient is negative and statistically significant ($p < 0.05$).

Figure 2c is a histogram of the raw residuals using probability density function scaling. Residuals broadly follow a normal distribution. Figure 2d is a scatter plot of fitted rainfall data and residuals. The distribution of residuals has no obvious regularity and trend, and is generally symmetric along $y = 0$. We furthermore assess the robustness of the model in Supplement Section 5. We calculate the autocorrelation length (where the autocorrelation coefficient is equal or below zero) of the SSWA rainfall for

205 the period 1900–2015 CE, and perform a jackknife (modified) analysis. The correlation coefficients of each individual jackknife ensemble member are all consistent with the period 1900–2015 CE (Supplement Table S4), and the 95% confidence intervals for the gradient and intercept are overlapped (Supplement Table S4), showing the statistically indistinguishable between each individual jackknife ensemble member. Taken together, there is no obvious evidence to reject the linear model. Thus, we use this linear model and the time series of DSS snow accumulation to reconstruct SWWA rainfall from 2015 CE back to 22 BCE.

210 **3.4 CUSUM analysis**

The Cumulative Summation (CUSUM) technique has been used to investigate rainfall data (Kampata et al., 2008; Chowdhury and Beecham, 2010). CUSUM is a method to sum the data anomalies using:

$$y_p = \sum_{k=1}^p x'_k \quad (3)$$

215 where x' is the anomaly relative to the mean of the whole series (Chowdhury and Beecham, 2010). The aim of CUSUM is to identify the step change in a time series by continuously accumulating anomalies. A step in the original data would be identified by the change of the gradient in the CUSUM.

Numerical integration enhances the signal-to-noise ratio. Using the CUSUM data to both pick the breakpoint (and evaluate its significance) and estimate the timing uncertainty potentially offer advantages compared to using the original data, especially at lower signal-to-noise ratios.

220 To specifically determine the breakpoint and evaluate its significance, we use the BREAKFIT analysis software (Mudelsee, 2009) on the CUSUM data.

3.5 Interpolation and model-data comparison

225 In light of the different spatial resolutions of the model simulations and AWAP gridded rainfall, we perform the interpolation for model simulations to be the same geospatial resolution as the AWAP gridded rainfall and multiply the MASK matrix with the interpolated model simulations to consistently capture the same region, same geospatial resolution data for both the rainfall reconstruction and simulation.

The rainfall reconstruction of SWWA back to 22 BCE is reconstructed by the linear model over MASK where the AWAP gridded rainfall and the DSS snow accumulation are statistically significantly correlated. The rainfall simulation is the precipitation simulation of the CSIRO Mk3L model run as four ensembles combining four climate forcings (Table 2). We use 230 the CUSUM analysis to compare the simulation of each ensemble and three members of CONTROL (a pre-industrial control simulation (Phipps et al., 2013)) to the rainfall reconstruction. Also, as each ensemble member represents the randomly different initialised weather states, we compare the mean of each ensemble member to the rainfall reconstruction to estimate the uncertainties in the simulated rainfall.

To statistically compare the rainfall reconstruction with the model simulations and test the differences, we use Welch's t-test
235 to calculate the adapted t-statistics using:

$$t_W = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}} \quad (4)$$

where \bar{x}_1 and \bar{x}_2 , s_1^2 and s_2^2 , n_1 and n_2 are sample mean values, standard deviations and sample sizes for two samples, respectively. We also estimate the adapted degrees of freedom (DF) using the Welch–Satterthwaite equation:

$$DF = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\frac{s_1^4}{n_1^2(n_1 - 1)} + \frac{s_2^4}{n_2^2(n_2 - 1)}} \quad (5)$$

240 Taken together, we calculate the p-value using t-statistics and DF.

To evaluate the specific time period we interested in, for example the rainfall before and after 1850 CE, we calculate the Root Mean Square Error (RMSE) between reconstructions with model simulations and also calculate the Pearson correlation coefficients, ESS (Equation 1), Student's t-statistic (Equation 2) and its adjusted p-value.

4 Results and discussion

245 4.1 Rainfall reconstruction

Figure 3a shows the reconstructed SSWA rainfall time series for the growing season (May to October) from 22 BCE to 2015 CE. We further validate the rainfall reconstruction in Supplement Section 5. We have found that the temporal stability of large-scale circulation over mid-latitude Australasian and higher latitude Law Dome regions is consistent, indicated by the correlation fields between Law Dome precipitation and mean sea level pressure in each member of the CSIRO Mk3L OG5V
250 ensemble (Supplement Figure S5). This rainfall reconstruction has shown that it is possible to use ice cores to investigate the longer-term context of rainfall variability over SSWA prior to the instrumental era. We should also be aware that the Law Dome—SSWA precipitation correlation is stronger during periods of enhanced (stronger/more frequent) meridional circulation, but weaker during periods of weaker meridional circulation (van Ommen and Morgan, 2010). The robustness of the reconstruction may therefore be reduced during periods of reduced meridional flow between southern Australia and Law
255 Dome (Supplement Section 5). Furthermore, the CSIRO Mk3L simulations exhibit variability in the correlation between Law Dome accumulation and SSWA precipitation on decadal to centennial timescales (Supplement Figure S6). Some caution is therefore needed in interpreting the reconstruction. We choose 1850 CE to be the year that separates before and after the Industrial Revolution consistent with other studies (e.g. Stocker et al., 2013).

Next, we assess whether the growing season rainfall in SSWA before and after 1850 CE belong to the same distribution. We
260 divide the rainfall reconstruction into two periods 22 BCE to 1849 CE and 1851 CE to 2015 CE. We plot the empirical distribu-




Figure 3. (a) The rainfall reconstruction in SWWA significant region (MASK region (Figure 1)) from 22 BCE to 2015 CE. The mean of this 2038-year time series (405 mm/year) is shown as a black horizontal line. (b) The rainfall reconstruction CUSUM (Equation 3). The black horizontal line equals zero. Red vertical dashed lines highlight the years 1850 CE and 1971 CE. (c) The 45-year running change in the rainfall reconstruction CUSUM series. Red horizontal dash line equals -1000 mm (accumulated change).

tion function (Figure 4a) and probability density function scaling histogram (Figure 4b) for each sample and its corresponding normal distribution (black dash line) with the same mean and standard deviation.

The distribution functions (Figure 4) for the period 1851 CE to 2015 CE are shifted left relative to the period 22 BCE to 1849 CE, indicating the rainfall after 1850 CE has reduced. To quantify the rainfall reduction and test its statistical significance, 265 we perform a two-sample KS test on the two samples and independently conduct Welch's t-test to validate two-sample KS nonparametric test results.

Results from two-sample KS test reject ($p < 0.01$) the null hypothesis that the samples are from same continuous distribution. Independently, Welch's t-test also allows us to reject ($p < 0.01$) the null hypothesis that the means are equal.

The mean of growing season rainfall reconstruction for 1851 CE to 2015 CE is 398 mm/year which is around 98% of the 270 rainfall mean during 22 BCE to 1849 CE which is 405 mm/year. The attenuation of the rainfall before and after 1850 CE is




Figure 4. (a) Empirical distribution function plot for rainfall reconstruction samples 22 BCE to 1849 CE (blue curve) and 1851 CE to 2015 CE (red curve). (b) Probability density function scaling histogram for two samples. Black dash curves are each sample's corresponding normal distribution.

statistically significant but the degree is around 2%. The 1851–2015 CE sample is missing the higher end of the distribution (i.e. there are no years with rainfall greater than 440 mm/year) and this lack of recharging events might have more of an impact than the shift in the mean (e.g., Gallant et al., 2013). Also the 1851 CE–2015 CE sample has a lower low-end of the distribution than 22 BCE–1849 CE, indicating more prevalent drier years. This suggests a possibility of an anthropogenic influence on the hydroclimate of this region.

The time series of the rainfall reconstruction gives us a clue that there might be another attenuation which happened after 1850 CE at around the late 20th century (Figure 3a,b). In order to accurately determine the changes in rainfall after 1850 CE, we calculate the CUSUM time series (Equation 3) for rainfall time series (Figure 3a) and use BREAKFIT analysis to identify any significant changes of the gradient. As CUSUM continuously accumulates anomalies of rainfall reconstruction along timeline, the variability of the rainfall becomes clearly visually observable (Figure 3b). The sign and magnitude of the gradient of CUSUM plot indicate rainfall anomalies. A positive (negative) gradient indicates a positive (negative) anomaly which is a rainfall increase (decrease) event. The larger the magnitude indicates the higher degree of the event. For consistency with the duration of the simulated rainfall which is available from 501 CE to 2000 CE, we take the rainfall reconstruction for the same period to calculate CUSUM time series and conduct BREAKFIT analysis.




Figure 5. CUSUM time series on rainfall reconstruction in period 1850–2000 CE. Yellow dotted lines are each interval's gradient calculated by using BREAKFIT analysis, with 1971 CE identified as the break-point. The gradient before 1971 CE is -1.40 mm/year. The gradient after 1971 CE is -37.16 mm/year.

285 Figure 5 shows the rainfall reconstruction CUSUM time series from 1850 CE to 2000 CE. Results from BREAKFIT show the break-point is at 1971 CE \pm 7 years (95% CI). The gradients for the intervals (1850, 1971) CE and (1971, 2000) CE are -1.4 mm/year [-3.2, -0.4] and -37.16 mm/year [-45.9, -28.8] respectively. Thus, the break in gradient at 1971 is statistically significant.

290 We test the findings of break-point analysis on the CUSUM time series by analysis of the original data. Breaking the rainfall into two samples, we found a shift of distribution to lower rainfall after 1971 CE in both empirical distribution function (Figure 6a) and probability density function scaling histogram (Figure 6b). The shift is around 30 mm/year showing a nearly 8% attenuation after 1971 CE, more than 4 times the reduction from 1850 CE. Two further statistical tests were preformed to verify this finding.

295 Results from the two-sample KS test allow us to reject the null hypothesis ($p < 0.01$) that the samples are from the same continuous distribution. Independently, it is statistically significant ($p < 0.01$) that the result from Welch's t-test can also allow us to reject the null hypothesis that two samples have equal means. The mean rainfall during the period 1972 CE to 2000 CE is 373 mm/year, which is 92% of the mean rainfall during the period 1850 CE to 1970 CE (406 mm/year). This change is more than four times bigger than the change around 1850 CE. Not only is there a large reduction in the mean rainfall, but also the 1972 CE to 2000 CE sample's distribution has largely shifted to lower values compared with 1850 CE to 1970 CE. There are




Figure 6. (a) Empirical distribution function plot for rainfall reconstruction samples 1850 CE to 1970 CE (blue curve) and 1972 CE to 2000 CE (red curve). (b) Probability density function scaling histogram for two samples. Black dash curves are each sample's corresponding normal distribution.

300 no years with rainfall over 410 mm/year, and nearly half of the years have rainfall less than 375 mm/year in the 1972 CE to 2000 CE sample (Figure 6b). This likely has a larger impact on agriculture than the rainfall shift before and after 1850 CE.

Results from the two individual statistical tests are very consistent with the BREAKFIT analysis on CUSUM time series that indicate 1971 CE marks the change in gradients from 1850 CE to 2000 CE. Comparison of Figure 6 with Figure 4 reveals that the shift in the rainfall distribution at around 1971 CE was much more pronounced than the shift at around 1850 CE. The 305 reduction in the mean rainfall at around 1971 CE resulted in a prolonged drought in SWWA. This drought continued during 2015–2020 (Supplement Figure S7). To highlight dry epochs of an equivalent duration to the observed drought to date, we calculate the 45-year running change in the CUSUM series (Figure 3c). Over the 45 years from 1971 CE to 2015 CE, this prolonged drought is expressed by an integrated rainfall reduction of more than 1000 mm (Figure 3c). We note there are two comparable prior epochs during the past two millennia, lasting from around 385–429 CE and 732–776 CE (Figure 3c).

310 **4.2 Data-model comparison**

Consistent with the analysis applied to the rainfall reconstruction, we analyze the simulated rainfall from CONTROL and the four forced ensembles. We divide the simulated time series for each simulation into two samples: 501 CE to 1849 CE and 1851 CE to 2000 CE. We further divide the industrial period into samples before and after the observed shift in 1971 CE. We

then independently perform the two-sample KS test and the Welch's t-test. We also use two different metrics to assess the
315 degree of agreement between the CUSUM time series for the various model simulations and the reconstruction: the differences
in the slope, and the Root Mean Square Error (RMSE).

Summary statistics for the model simulations are shown in Supplement Table S5. The model can be seen to have a dry
bias, but the simulated internal variability is of a similar magnitude (~ 80%) to the reconstruction. The simulated rainfall for
each individual ensemble member is shown in Supplement Figure S8, with the corresponding CUSUM time series shown in
320 Supplement Figure S9. Within each forced ensemble, there are considerable differences between the CUSUM time series for
individual ensemble members. This highlights the role of unforced internal variability in driving SWWA rainfall, consistent
with the findings of Cai and Shi (2005).

A number of prolonged droughts that approach the magnitude of the current integrated rainfall deficit are apparent in the
reconstruction (Figure 3c), as discussed in subsection 4.1. These prior drought epochs occurred during the pre-industrial era,
325 suggesting that they may have arisen through natural climate variability or natural forcings such as volcanoes. To explore
this within the model simulations, the 45-year changes in the CUSUM for the pre-industrial control simulation are shown in
Supplement Figure S10. A number of prolonged droughts are also apparent in all three ensemble members, for example at
years 589–633 CE in Control1 (Supplement Figure S10b). This supports the hypothesis that prolonged droughts of a similar
magnitude to the currently observed drought can arise through natural climate variability or natural forcings.

330 4.2.1 Industrial era

Table 4. The results for two independent statistical tests for different time periods from the rainfall time series. KS test is the two-sample KS test and W's t-test is the Welch's t-test. The tests results "same" means we cannot reject the null hypothesis (the two samples of data are from the same distribution), while "different" means we can reject the null hypothesis ($p < 0.05$).

	501–1849 CE VS 1851–2000 CE		1850–1970 CE VS 1972–2000 CE	
Sample	KS test	W's t-test	KS test	W's t-test
CONTROL	same	same	same	same
O	same	same	same	same
OG	different	different	same	same
OGS	different	different	same	same
OGSV	different	different	same	same
Rainfall reconstructions	different	different	different	different

We now examine the model ensembles and assess whether the simulated rainfall during the industrial era has a different
distribution from the simulated rainfall during the pre-industrial era. We cannot reject the null hypothesis, with either the
two-sample KS test or Welch's t-test performed on both CONTROL and O series, that the rainfall in 501–1849 CE and 1851–

2000 CE are from the same distribution (KS test) or equal mean (Welch's t-test) (Table 4). Conversely, the results from both
 335 two-sample KS test and Welch's t-test allow us to reject ($p < 0.05$) the null hypothesis for OG, OGS or OGSV and rainfall
 reconstruction (Table 4). These results give us confidence to say that orbital forcing is not the main driver for the rainfall
 changes after 1850 CE. Once the greenhouse gases are added to the model climate forcings, the rainfall simulations before and
 340 after 1850 CE belong to different distributions with different means (Table 4). There is no change when the solar irradiance
 and volcanic aerosols are added. Based on these statistical tests, we suggest that there is evidence for a role of anthropogenic
 greenhouse gases. However, there is no evidence of any additional impacts due to either solar irradiance or volcanic aerosols.




Figure 7. CUSUM time series for the rainfall reconstruction and CSIRO Mk3L model rainfall simulations in the MASK region from 501 CE to 2000 CE. The model simulations include CONTROL (pre-industrial control simulation (Phipps et al., 2013)), O, OG, OGS and OGSV. The black horizontal line equals zero. Red vertical dash lines highlight the year 1850 CE and 1971 CE.

Figure 7 shows the CUSUM time series for each of the rainfall reconstruction and CSIRO Mk3L model simulations from 501 CE to 2000 CE. For each ensemble including CONTROL scenario, the CUSUM time series is the mean of three members. As for the statistical tests (Table 4), the CUSUM series in Figure 7 show that the rainfall change around 1850 CE is not simulated by CONTROL or O ensembles, but is simulated by OG, OGS and OGSV ensemble series.

345 Examining each of the ensembles in turn, we see that CONTROL does not capture the key features of the rainfall reconstruction (Figure 7). This suggests either a role of internal variability, or that climate forcings might be driving the rainfall changes in SSWA. The time series of O varies slightly around zero with no obvious gradient changes (Figure 7), which suggests that orbital forcing alone cannot explain the changes in SSWA rainfall in recent decades. In contrast, ensembles OG, OGS

and OGSV are able to reproduce some of the key features of the rainfall reconstruction, particularly the decline after 1850 CE.
 350 These ensembles have approximately equal magnitudes especially after around 1900 CE and their gradient changes are broadly synchronous with large negative magnitudes.

4.2.2 Late 20th century

Figure 8 shows the CUSUM time series for the rainfall reconstruction and simulations in the period 1972–2000 CE. Ensemble O shows that the orbital forcing cannot explain the rainfall reduction after 1971 CE. However the reduction is reproduced by
 355 ensembles OG, OGS and OGSV. All three ensembles have the same sign of gradients and similar magnitudes. The magnitude of the gradient for the rainfall reconstruction is larger than any of the rainfall simulations.

We also perform a two-sample KS test and Welch's t-test on the rainfall simulations in the periods 1850–1970 CE and 1972–
 2000 CE. None of the model results from the statistical tests allow us to reject the null hypothesis with statistical significance.
 Unlike the rainfall reconstruction, none of the model simulations show that the rainfall in period 1972–2000 CE is different
 360 from the period 1850–1970 CE (Table 4).




Figure 8. The CUSUM time series on the rainfall reconstruction and simulations in the period 1972–2000 CE from the 501–2000 CE series. The black horizontal line equals zero.

As we can see in Figure 8, CONTROL and O have negligible slope. However, OG, OGS and OGSV do exhibit a negative slope and are therefore much closer to the rainfall reconstruction. We note that the amplitude of the internal variability in the underlying raw time series will influence the magnitude of the slopes in the CUSUM analysis. CONTROL and O are essentially indistinguishable as the time frame of orbital forcing is long compared to the 30 year period studied here. To determine which

Table 5. The RMSE and difference in slope calculated between the rainfall reconstruction and simulations in the period 1972–2000 CE. The difference in slope is the difference of the slope of the Ordinary Least Square linear fitted CUSUM time series data for each sample to the slope of the fitted CUSUM time series data for rainfall reconstruction in the period 1972–2000 CE.

Simulation	Difference in slope (mm/year)	RMSE (mm)
CONTROL	41.50	580
O	37.00	558
OG	20.09	296
OGS	23.00	287
OGSV	20.66	263

365 model simulation is in best agreement with the rainfall reconstruction, we calculate the difference in slope and the Root Mean Square Error (RMSE) (Table 5) between the CUSUM series for the rainfall reconstruction and each model ensemble.

Comparing CONTROL and the four ensembles, CONTROL and O have the largest differences in slope and RMSE, relative to the rainfall reconstruction, than any of the others. The forced ensembles OG, OGS and OGSV have much smaller differences in slope and much smaller RMSE. Therefore, both of difference in slope and RMSE analysis suggest that the simulations 370 including greenhouse gases are the closest to the rainfall reconstruction. Including solar irradiance and volcanic aerosols has negligible additional impact.

Although the model ensembles that include greenhouse gas forcing do simulate a drying trend after 1971 CE, the simulated trend is weaker than in the reconstruction. This may reflect the influence of stratospheric ozone depletion, which is not modelled in the climate model simulations that we analyse in this study. Alternatively, it may reflect the role of natural climate variability. 375 Because of its stochastic nature, the model simulations would not be expected to replicate the phase or magnitude of the specific internal variability encountered in the real world.

The largest improvement in agreement comes from adding greenhouse gases into the model. Therefore, our analysis suggests that greenhouse gases are the dominant climate forcing out of the four natural and anthropogenic forcing considered in this study driving the rainfall changes in SWWA during the 20th century.

380 **5 Conclusions**

We have used the DSS snow accumulation record to extend the previous published reconstruction of SWWA growing season (May to October) rainfall. Based on the statistically significant ($p < 0.05$) negative correlation between SWWA rainfall and DSS snow accumulation in the growing season, we built a linear model and reconstructed the rainfall for 2038 years (22 BCE to 2015 CE). Consistent with other studies (e.g. Stocker et al., 2013), we use 1850 CE to divide the reconstruction into two 385 periods: before and after the industrial revolution. We independently performed the two-sample KS test and Welch's t-test to test whether the two samples are from the equal means and distributions. The results from tests rejected ($p < 0.01$) the

390 null hypotheses, suggesting a detectable change in SSWA rainfall much earlier than the observed climate shift in SSWA in the 1970s. The CUSUM time series and the BREAKFIT analysis suggest a reduction rainfall after 1971 CE that was much larger in magnitude than the reduction after 1850 CE. The rainfall means and distributions in the periods 1850–1970 CE and 1972–2000 CE are independently tested and shown to be statistically significantly ($p < 0.01$) different.

395 With statistically significant ($p < 0.01$) rainfall reductions after 1850 CE and after 1971 CE, we compared the rainfall reconstructions with ensembles of CSIRO Mk3L climate model simulations driven by different natural and anthropogenic climate forcings. Comparing the periods 501–1849 CE and 1851–2000 CE, adding greenhouse gases significantly ($p < 0.05$) changed the rainfall means and distributions for the two periods. There is no detectable influence from solar irradiance and volcanic aerosols. However, the rainfall means and distributions for the periods 1850–1970 CE and 1972–2000 CE are statistically indistinguishable.

400 Examining the CUSUM for the period 1972–2000 CE, we do not find any drying trend in CONTROL simulation or in simulations driven only with orbital forcing. However, we do find a drying trend in simulations that include anthropogenic greenhouse gases. We again find that solar irradiance and volcanic aerosols have negligible additional impact. The model simulations therefore suggest that anthropogenic forcing has been the dominant driver of the rainfall trend in SSWA since 1971 CE.

405 Both the reconstruction and the climate model simulations suggest that the drying trend began earlier than the 1970s. However, the model simulations do not capture the acceleration in the reconstructed drying trend at around 1971 CE. This suggests that this acceleration cannot be attributed to external forcings, or at least not any of the forcings considered in this study. 410 Either natural variability or stratospheric ozone depletion are potential alternative explanations (e.g. Cai and Shi, 2005). A investigation into possible ozone forcing and a comparison against multiple climate models is a promising avenue for future work.

415 The reconstruction reveals that the rainfall reduction in SSWA in recent decades is not unprecedented. We note that two previous prolonged drought events of similar magnitude in SSWA have occurred during the past two millennia. Droughts of similar duration and magnitude also occur in an unforced pre-industrial control simulation. The model simulations therefore support the hypothesis that these pre-industrial drought events may have arisen through natural climate variability. However, forced climate model simulations indicate that anthropogenic greenhouse gases are the dominant driver of the rainfall reduction in SSWA since the early 1970s.

415 *Data availability.* The SSWA rainfall reconstruction data is available via <https://doi.org/10.25959/5f4c50b7b661f>. The DSS snow accumulation record is available via <https://www.clim-past.net/11/697/2015/>. The CSIRO Mk3L climate model simulations are available via <https://doi.org/10.5281/zenodo.3908926/>. The BoM instrumental rainfall records are available via <http://www.bom.gov.au/climate/data/>. The AWAP AGCD data is available via <http://www.bom.gov.au/metadata/catalogue/19115/ANZCW0503900567?template=full>.

Author contributions. YZ, SP, LJ and JR conceived the study. YZ performed the analysis. All authors contributed to the writing of the manuscript.

420 *Competing interests.* The authors declare that there are no competing interests.

Acknowledgements. This project received grant funding from the Australian Government as part of the Antarctic Science Collaboration Initiative program and was supported by (i) the Australian Research Council's Special Research Initiative for the Antarctic Gateway Partnership (Project ID SR140300001), (ii) the Centre for Southern Hemisphere Oceans Research, a joint research centre between QNLM and CSIRO, and (iii) the Australia's Antarctic program (AAS 4061, 4062 and 4537).

Abram, N. J., Mulvaney, R., Vimeux, F., Phipps, S. J., Turner, J., and England, M. H.: Evolution of the Southern Annular Mode during the past millennium, *Nature Climate Change*, 4, 564–569, <https://doi.org/10.1038/NCLIMATE2235>, 2014.

Anderson, W., Heinrich, A., and Abbotts, R.: Long-season wheats extend sowing opportunities in the central wheat belt of Western Australia, *Australian Journal of Experimental Agriculture*, 36, 203–208, 1996.

430 Ansell, T., Reason, C., Smith, I., and Keay, K.: Evidence for decadal variability in southern Australian rainfall and relationships with regional pressure and sea surface temperature, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 20, 1113–1129, 2000.

Ashok, K., Guan, Z., and Yamagata, T.: Influence of the Indian Ocean Dipole on the Australian winter rainfall, *Geophysical Research Letters*, 30, 2003.

435 Australian Government: Local Government Area, <https://data.gov.au/data/dataset/8a8e0037-e474-422f-8026-241c7c88551f>, 2020.

Barr, C., Tibby, J., Leng, M. J., Tyler, J. J., Henderson, A. C. G., Overpeck, J. T., Simpson, G. L., Cole, J. E., Phipps, S. J., Marshall, J. C., McGregor, G. B., Hua, Q., and McRobie, F. H.: Holocene El Niño-Southern Oscillation variability reflected in subtropical Australian precipitation, *Scientific Reports*, 9, 1627, <https://doi.org/10.1038/s41598-019-38626-3>, 2019.

Bretherton, C. S., Widmann, M., Dymnikov, V. P., Wallace, J. M., and Bladé, I.: The effective number of spatial degrees of freedom of a 440 time-varying field, *Journal of climate*, 12, 1990–2009, 1999.

Bromwich, D. H.: Snowfall in high southern latitudes, *Reviews of Geophysics*, 26, 149–168, 1988.

Cai, W. and Cowan, T.: SAM and regional rainfall in IPCC AR4 models: Can anthropogenic forcing account for southwest Western Australian winter rainfall reduction?, *Geophysical Research Letters*, 33, 2006.

Cai, W. and Shi, G.: Multidecadal fluctuations of winter rainfall over southwest Western Australia simulated in the CSIRO Mark 3 coupled 445 model, *Geophysical Research Letters*, 32, L12 701, <https://doi.org/10.1029/2005GL022712>, 2005.

Cai, W., Collier, M. A., Gordon, H. B., and Waterman, L. J.: Strong ENSO Variability and a Super-ENSO Pair in the CSIRO Mark 3 Coupled Climate Model, *Monthly Weather Review*, 131, 1189–1210, [https://doi.org/10.1175/1520-0493\(2003\)131<1189:SEVAAS>2.0.CO;2](https://doi.org/10.1175/1520-0493(2003)131<1189:SEVAAS>2.0.CO;2), 2003.

Cai, W., Van Rensch, P., Cowan, T., and Hendon, H. H.: Teleconnection pathways of ENSO and the IOD and the mechanisms for impacts on 450 Australian rainfall, *Journal of Climate*, 24, 3910–3923, 2011.

Chiew, F. H., Piechota, T. C., Dracup, J. A., and McMahon, T. A.: El Niño/Southern Oscillation and Australian rainfall, streamflow and drought: Links and potential for forecasting, *Journal of hydrology*, 204, 138–149, 1998.

Chowdhury, R. and Beecham, S.: Australian rainfall trends and their relation to the southern oscillation index, *Hydrological Processes: An International Journal*, 24, 504–514, 2010.

455 England, M. H., Ummenhofer, C. C., and Santoso, A.: Interannual rainfall extremes over southwest Western Australia linked to Indian Ocean climate variability, *Journal of Climate*, 19, 1948–1969, 2006.

Feng, J., Li, J., and Li, Y.: A monsoon-like southwest Australian circulation and its relation with rainfall in southwest Western Australia, *Journal of Climate*, 23, 1334–1353, 2010.

Fierro, A. O. and Leslie, L. M.: Links between central west Western Australian rainfall variability and large-scale climate drivers, *Journal of 460 climate*, 26, 2222–2246, 2013.

French, R. and Schultz, J.: Water use efficiency of wheat in a Mediterranean-type environment. II. Some limitations to efficiency, *Australian Journal of Agricultural Research*, 35, 765–775, 1984.

Gallant, A. J., Reeder, M. J., Risbey, J. S., and Hennessy, K. J.: The characteristics of seasonal-scale droughts in Australia, 1911–2009, *International Journal of Climatology*, 33, 1658–1672, 2013.

465 Gillett, N. P. and Thompson, D. W. J.: Simulation of Recent Southern Hemisphere Climate Change, *Science*, 302, 273–275, <https://doi.org/10.1126/science.1087440>, 2003.

Gong, D. and Wang, S.: Definition of Antarctic oscillation index, *Geophysical research letters*, 26, 459–462, 1999.

Goodwin, I., Van Ommen, T., Curran, M., and Mayewski, P.: Mid latitude winter climate variability in the South Indian and southwest Pacific regions since 1300 AD, *Climate Dynamics*, 22, 783–794, 2004.

470 Gordon, H. B., Rotstayn, L. D., McGregor, J. L., Dix, M. R., Kowalczyk, E. A., O'Farrell, S. P., Waterman, L. J., Hirst, A. C., Wilson, S. G., Collier, M. A., Watterson, I. G., and Elliott, T. I.: The CSIRO Mk3 Climate System Model, Technical Paper 60, CSIRO Atmospheric Research, available online at http://www.cmar.csiro.au/e-print/open/gordon_2002a.pdf, 2002.

Hennessy, K. J., Suppiah, R., and Page, C. M.: Australian rainfall changes, 1910–1995, *Australian meteorological magazine*, 48, 1–13, 1999.

Hope, P. K., Drosdowsky, W., and Nicholls, N.: Shifts in the synoptic systems influencing southwest Western Australia, *Climate Dynamics*, 475 26, 751–764, 2006.

Jones, D. A., Wang, W., and Fawcett, R.: High-quality spatial climate data-sets for Australia, *Australian Meteorological and Oceanographic Journal*, 58, 233, 2009.

Kampata, J. M., Parida, B. P., and Moalafhi, D.: Trend analysis of rainfall in the headstreams of the Zambezi River Basin in Zambia, *Physics and Chemistry of the Earth, Parts A/B/C*, 33, 621–625, 2008.

480 Lavery, B., Joung, G., and Nicholls, N.: An extended high-quality historical rainfall dataset for Australia, *Australian Meteorological Magazine*, 46, 27–38, 1997.

Li, Y., Cai, W., and Campbell, E.: Statistical modeling of extreme rainfall in southwest Western Australia, *Journal of climate*, 18, 852–863, 2005.

Ludwig, F. and Asseng, S.: Climate change impacts on wheat production in a Mediterranean environment in Western Australia, *Agricultural 485 Systems*, 90, 159–179, 2006.

Ludwig, F., Milroy, S. P., and Asseng, S.: Impacts of recent climate change on wheat production systems in Western Australia, *Climatic Change*, 92, 495–517, 2009.

Mudelsee, M.: Break function regression, *The European Physical Journal Special Topics*, 174, 49–63, 2009.

490 Phipps, S., Rotstayn, L., Gordon, H., Roberts, J., Hirst, A., and Budd, W.: The CSIRO Mk 3 L climate system model version 1. 0-Part 1: Description and evaluation, *Geoscientific Model Development*, 4, 483–509, 2011.

Phipps, S., Rotstayn, L., Gordon, H., Roberts, J., Hirst, A., and Budd, W.: The CSIRO Mk3L climate system model version 1.0-Part 2: Response to external forcings, *Geoscientific Model Development*, 5, 649–682, 2012.

Phipps, S. J., McGregor, H. V., Gergis, J., Gallant, A. J., Neukom, R., Stevenson, S., Ackerley, D., Brown, J. R., Fischer, M. J., and van Ommen, T. D.: Paleoclimate data–model comparison and the role of climate forcings over the past 1500 years, *Journal of Climate*, 26, 495 6915–6936, 2013.

Pitman, A. J., Narisma, G. T., Pielke Sr, R. A., and Holbrook, N.: Impact of land cover change on the climate of southwest Western Australia, *Journal of Geophysical Research: Atmospheres*, 109, 2004.

Pittock, A.: Recent climatic change in Australia: Implications for a CO 2-warmed earth, *Climatic Change*, 5, 321–340, 1983.

Power, S., Sadler, B., and Nicholls, N.: The influence of climate science on water management in Western Australia: Lessons for climate
500 scientists, *Bulletin of the American Meteorological Society*, 86, 839–844, 2005.

Roberts, J., Plummer, C., Vance, T., van Ommen, T., Moy, A., Poynter, S., Treverrow, A., Curran, M., and George, S.: A 2000-year annual
record of snow accumulation rates for Law Dome, East Antarctica, *Climate of the Past*, 11, 697–707, 2015.

Samuel, J. M., Verdon, D. C., Sivapalan, M., and Franks, S. W.: Influence of Indian Ocean sea surface temperature variability on southwest
Western Australian winter rainfall, *Water Resources Research*, 42, 2006.

505 Siddique, K., Loss, S., Regan, K., and Jettner, R.: Adaptation and seed yield of cool season grain legumes in Mediterranean environments of
south-western Australia, *Australian Journal of Agricultural Research*, 50, 375–388, 1999.

Smith, I., McIntosh, P., Ansell, T., Reason, C., and McInnes, K.: Southwest Western Australian winter rainfall and its association with Indian
Ocean climate variability, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 20, 1913–1930, 2000.

510 Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P. M., et al.: Climate
change 2013: The physical science basis, Contribution of working group I to the fifth assessment report of the intergovernmental panel on
climate change, 1535, 2013.

Thompson, D. W. and Solomon, S.: Interpretation of recent Southern Hemisphere climate change, *Science*, 296, 895–899, 2002.

Thompson, D. W., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of the Antarctic ozone hole in
Southern Hemisphere surface climate change, *Nature geoscience*, 4, 741–749, 2011.

515 Timbal, B., Arblaster, J. M., and Power, S.: Attribution of the late-twentieth-century rainfall decline in southwest Australia, *Journal of
Climate*, 19, 2046–2062, 2006.

Trenberth, K. E. and Hoar, T. J.: El Niño and climate change, *Geophysical Research Letters*, 24, 3057–3060, 1997.

Ummenhofer, C. C., Sen Gupta, A., Pook, M. J., and England, M. H.: Anomalous rainfall over southwest Western Australia forced by Indian
Ocean sea surface temperatures, *Journal of Climate*, 21, 5113–5134, 2008.

520 van Ommen, T. D. and Morgan, V.: Snowfall increase in coastal East Antarctica linked with southwest Western Australian drought, *Nature
Geoscience*, 3, 267–272, 2010.

Ward, P. and Dunin, F.: Growing season evapotranspiration from duplex soils in south-western Australia, *Agricultural Water Management*,
50, 141–159, 2001.

Watson, E. and Lapins, P.: Losses of nitrogen from urine on soils from south western Australia, *Australian Journal of Experimental Agricul-
525 ture*, 9, 85–91, 1969.

Wright, P. B.: Seasonal rainfall in southwestern Australia and the general circulation, *Monthly Weather Review*, 102, 219–232, 1974a.

Wright, P. B.: Temporal variations in seasonal rainfalls in southwestern Australia, *Monthly Weather Review*, 102, 233–243, 1974b.

Yu, B. and Neil, D.: Long-term variations in regional rainfall in the south-west of Western Australia and the difference between average and
high intensity rainfalls, *International Journal of Climatology*, 13, 77–88, 1993.