

Climate indices in historical climate reconstructions: A global state-of-the-art

David J. Nash^{a,b}, George C.D. Adamson^c, Linden Ashcroft^{d,e}, Martin Bauch^f, Chantal Camenisch^{g,h}, Dagomar Degrootⁱ, Joelle Gergis^{j,k}, Adrian Jusopović^l, Thomas Labbé^{f,m}, Kuan-Hui Elaine Lin^{n,o}, Sharon D. Nicholson^p, Qing Pei^q, María del Rosario Prieto^{r†}, Ursula Rack^s, Facundo Rojas^r and Sam White^t

^a School of Environment and Technology, University of Brighton, Brighton, United Kingdom

^b School of Geography, Archaeology and Environmental Studies, University of the Witwatersrand, Johannesburg, South Africa

^c Department of Geography, King's College London, London, United Kingdom

^d School of Earth Sciences, University of Melbourne, Melbourne, Australia

^e ARC Centre of Excellence for Climate Extremes, University of Melbourne, Melbourne, Australia

^f Leibniz Institute for the History and Culture of Eastern Europe, University of Leipzig, Leipzig, Germany

^g Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

^h Institute of History, University of Bern, Bern, Switzerland

ⁱ Department of History, Georgetown University, Washington DC, USA

^j Fenner School of Environment & Society, Australian National University, Canberra, Australia

^k ARC Centre of Excellence for Climate Extremes, Australian National University, Canberra, Australia

^l Institute of History, Polish Academy of Sciences, Warsaw, Poland

^m Maison des Sciences de l'Homme de Dijon, University of Burgundy, Dijon, France

ⁿ Research Center for Environmental Changes, Academia Sinica, Taipei, Taiwan

^o Graduate Institute of Environmental Education, National Taiwan Normal University, Taipei, Taiwan

^p Department of Earth, Ocean, and Atmospheric Science, Florida State University, Tallahassee, Florida, USA

^q Department of Social Sciences, Education University of Hong Kong, Hong Kong, Peoples Republic of China

^r Argentine Institute of Nivology, Glaciology and Environmental Sciences (IANIGLA-CONICET), Mendoza, Argentina

^s Gateway Antarctica, University of Canterbury, Christchurch, New Zealand

^t Department of History, Ohio State University, Columbus, Ohio, USA

[†] Deceased

Correspondence to: David J. Nash (d.j.nash@brighton.ac.uk). ORCID: 0000-0002-7641-5857

1 **Abstract.** Evidence contained within historical documents and inscriptions provides an important record of climate
2 variability for periods prior to the onset of systematic meteorological data collection. A common approach used by historical
3 climatologists to convert such qualitative documentary evidence into continuous quantitative proxy data is through the
4 generation of ordinal-scale climate indices. There is, however, considerable variability in the types of phenomena
5 reconstructed using an index approach and the practice of index development in different parts of the world. This review,
6 written by members of the PAGES CRIAS Working Group – a collective of climate historians and historical climatologists
7 researching Climate Reconstructions and Impacts from the Archives of Societies – provides the first global synthesis of the
8 use of the index approach in climate reconstruction. We begin by summarising the range of studies that have used indices for
9 climate reconstruction across six continents (Europe, Asia, Africa, the Americas, Australia) plus the world’s oceans. We then
10 outline the different methods by which indices are developed in each of these regions, including a discussion of the processes
11 adopted to verify and calibrate index series, and the measures used to express confidence and uncertainty. We conclude with
12 a series of recommendations to guide the development of future index-based climate reconstructions to maximise their
13 effectiveness for use by climate modellers and in multiproxy climate reconstructions.
14
15 **Keywords.** Climate reconstruction; temperature reconstruction; precipitation reconstruction; historical climatology; climate
16 history; documentary evidence

1. Introduction

Much of the effort of the palaeoclimatological community in recent decades has focussed on understanding long-term changes in climate, typically at millennial, centennial, or at best (in the case of dendroclimatology and palaeolimnology) sub-decadal to annual resolution. The results of this research have revolutionised our knowledge both of how climates have varied in the past and the potential drivers of such variability. However, as Pfister et al. (2018) identify, the results of palaeoclimate research are often at a temporal and spatial scale that is not suitable for understanding the short-term and local impacts of climate variability upon economies and societies. To this end, historical climatologists work to reconstruct high-resolution – annual, seasonal, monthly and in some cases daily – series of past temperature and precipitation variability from the archives of societies, as these are the scales at which weather impacts upon individuals and communities (e.g. Allan et al., 2016; Brönnimann et al., 2019).

The archives of societies, used here in a broad sense to refer to both written records and evidence preserved in the built environment (e.g. historic flood markers, inscriptions), contain extensive information about past local weather and its repercussions for the natural environment and on daily lives. Information sources include, but are not limited to, annals, chronicles, inscriptions, letters, diaries/journals (including weather diaries), newspapers, financial, legal and administrative documents, ships' logbooks, literature, poems, songs, paintings and pictographic and epigraphic records (Brázdil et al., 2005; Brázdil et al., 2010; Brázdil et al., 2018; Pfister, 2018; Rohr et al., 2018). Three main categories of information appear in these sources that can be used independently or in combination for climate reconstruction: (i) early instrumental meteorological data; (ii) records of recurring physical and biological processes (e.g. dates of plant flowering, grape ripening, the freezing of lakes and rivers); and (iii) narratives of short-term atmospheric processes and their impacts on environments and societies (Brönnimann et al., 2018).

The heterogeneity of the archives of societies – in time, space and in the types of information included in individual sources – raises conceptual and methodological challenges for climate reconstruction. Historical meteorological data can be quality-checked and analysed using standard climatological methods, while records of recurrent physical and biological phenomena provide proxy information that may be assessed using a variety of palaeoclimatological approaches (cf. Brönnimann et al., 2018). Descriptive and narrative accounts, however, require different treatment to make local observations of weather and its impacts compatible with the statistical requirements of climatological research.

A common approach used in historical climatology for the analysis of descriptive (or narrative) evidence is the generation of ordinal-scale indices as a bridge between raw weather descriptions and climate reconstructions. A simple unweighted index might, for example, employ a three-point classification, with months classed as –1 (cold or dry), 0 (normal) and 1 (warm or wet) depending upon the prevailing conditions described within historical sources. As Pfister et al. (2018) note, this “index” approach provides a means of converting “disparate documentary evidence into continuous quantitative proxy data... but without losing the ability to get back to the short-term local information for critical inspection and analysis” (p.116). Brázdil et al. (2010) provide a detailed account of the issues associated with the generation of indices.

The index approach to historical climate reconstruction over much of the world – an exception being China – has its roots in European scholarship. There is, however, considerable variability in the types of phenomena reconstructed using an index approach in different areas. There is also variability in practice, both in the way that historical evidence is treated to generate indices and in the number of ordinal categories in individual index series. Variability in the treatment of evidence arises, in part, from the extent to which analytical approaches have developed independently. In terms of categorisation, three-, five- and seven-point index series are most widely used but greater granularity (i.e. a greater number of index classes) may be achieved in different regions and for different climate phenomena depending upon the quantity, resolution and/or richness of the original historical evidence.

This study arises from the work of the PAGES (Past Global Changes) CRIAS Working Group, a cooperative of climate historians and historical climatologists researching Climate Reconstructions and Impacts from the Archives of Societies. The first meeting of the Working Group in Bern, Switzerland, in September 2018 identified the need to understand variability and – ideally – harmonise practice in the use of indices to maximise the utility of historical climate reconstructions for climate change investigations. This study, written by regional experts in historical climatology with contributions from other CRIAS members, is intended to address this need.

The ~~main~~ aims of ~~this~~ paper are ~~twofold, namely~~ to: (i) provide a global state-of-the-art review of the development and ~~application-use~~ of the index approach ~~as applied to descriptive evidence~~ in historical climate reconstruction; and (ii) identify best practice for future investigations. It does so through a continent-by-continent overview of practice, followed by a review of the use of indices in the reconstruction of climate variability over the oceans. Studies from northern polar regions are reviewed within sections 5 (the Americas) and 7 (the Oceans), as appropriate. To the knowledge of the authors, no studies of the climate history of Antarctica use an index approach.

Two caveats are necessary to frame the coverage of the review. First, the nature of documentary sources is well discussed in the climate history literature for most parts of the world. As such, we provide only limited commentary on sources for each continent, except for selected regions. These include China, where only a few overviews of documentary sources have been published (e.g. Wang, 1979; Wang and Zhang, 1988; Zhang and Crowley, 1989; Ge et al., 2018), and Japan and Russia where, to our knowledge, few no detailed descriptions are available for Anglophone audiences. Second, there are instances in the literature where quantifiable data in documentary sources (e.g. sea-ice cover, phenological phenomena) and even instrumental meteorological data are converted to indices for climate reconstruction purposes. This occurs mainly in studies where such data are integrated with information from narrative sources to generate longer, more continuous and homogenous series with a consistent (monthly or seasonal) resolution. We do not describe the generation of such index series in detail, but do provide examples in sections 2 to 7, as appropriate.

2. Climate indices in Europe

2.1. Origins of documentary-based indices in Europe


The use of climate indices has a long tradition in Europe, with the earliest studies published during the 1920s CE. As in any area, the start date for meaningful index-based reconstructions is determined by the availability of source material. Narrative sources in Central, Western and Mediterranean Europe, for example, are sufficiently dense from the 15th century CE onwards to enable seasonal index reconstruction for more than half of all covered years. Exceptionally, indices can be generated from the 12th century onwards, but with greatest confidence from the 14th century when serial sources join the available historiographic information (Wozniak, 2020). The number of index-based climate reconstructions for Europe is large; as such, this section of the review focusses mainly upon studies that include original published series based on primary sources and that reconstruct meteorological entities. This excludes climate modelling and other studies that synthesise or reanalyse previously published historical index series.

Due to the dominance of references to winter conditions in European documentary sources, early investigations centred primarily on winter severity (Pfister et al., 2018). The first use of the index approach was by the Dutch journalist, astronomer and later climatologist Cornelis Easton, who published his oeuvre on historical European winter severity in 1928 (Easton, 1928). In this monograph, Easton presented early instrumental data but also a catalogue of descriptions of winter conditions dating back to the 3rd century BCE derived from narrative sources. For the period prior to 1205 CE, this catalogue lists only

98 remarkable winter seasons; however, after this date every winter up to 1916 is attributed to a ten-point classification,
 99 including a quantifiable coefficient and a descriptive category. Easton's classification appears as an adapted graph in the
 100 second edition of Charles E. P. Brooks (1949) book on *Climate Through the Ages* (Pfister et al., 2018).

101 An isolated attempt to quantify the evaluation of weather diaries ([spanning 1182-1780 CE](#)) was proposed by the German
 102 meteorologist Fritz Klemm (1970), with a two-point scale for winter and summer temperature (cold/mild and mild/warm
 103 respectively) and precipitation (dry/wet). The Dutch meteorologist Folkert IJnsen also developed winter severity indices for
 104 the Netherlands ([1200-1916 CE](#)) but following a slightly different approach (IJnsen and Schmidt, 1974). However, one of
 105 the most important advances came in the late 1970s when British climatologist Hubert Horace Lamb published three-point
 106 indices of winter severity and summer wetness for Western Europe ([1100-1969 CE](#)) in his seminal book *Climate: Past,*
 107 *Present and Future* (Lamb, 1977). Lamb's methodology was more easily applicable compared to Easton's – a likely reason
 108 why successive studies refer to Lamb's method and why, in the aftermath of his publication, the index approach was applied
 109 in many different European regions.

110 In 1984, the Swiss historian Christian Pfister published his first temperature and precipitation indices for Switzerland in the
 111 volume *Das Klima der Schweiz von 1525-1860*, expanding his climate indices to cover all months and seasons of the year
 112 (Pfister, 1984). Pfister's work adapted Lamb's methods, extending Lamb's three-point scale into monthly seven-point
 113 ordinal-scale temperature and precipitation indices (Figure 1). Shortly after Pfister's initial study, Pierre Alexandre (1987)
 114 developed a comprehensive overview of the climate of the European Middle Ages ([1000-1425 CE](#)), also using indices. Over
 115 a decade later, Van Engelen et al. (2001) published a nine-point index-based temperature reconstruction for the Netherlands
 116 and Belgium ([764-1998 CE](#)). Most research groups investigating European climate history – including those led by Rüdiger
 117 Glaser (Freiburg, Germany) and Rudolf Brázdil (Brno, Czech Republic) – now adopt Pfister's approach as the standard
 118 method for index development, at least for temperature and precipitation reconstructions. This is described in more detail in
 119 section 8 as part of a global overview of approaches to index construction. [The opportunity to combine narrative sources](#)
 120 [with quantifiable information is one of the great advantages of the index-approach](#) (Pfister et al., 2018). [As a result, many](#)
 121 [index-based series for Europe incorporate some quantitative data. Many series also contain data gaps; the earlier the epoch,](#)
 122 [the more likely there are to be breaks in series – this is common to almost all index-based series globally.](#)



124 **Figure 1:** Monthly seven-point temperature indices for the Swiss Plateau (1680-1700), reconstructed using the Pfister index
 125 approach (data from Pfister, 1998). Zero values for specific months are indicated by a small green bar.

One area of Europe with a different research tradition is Russia (Jusupović and Bauch, 2020). Here, the earliest climate history research was by K.S. Veselovskij (1857), who compared historical information from various source types against early 19th century statistical climate data (for more details of Veselovskij's work, see Zhogova, 2013). M.A. Bogolepov later analysed climate-related information in published Cyrillic and Latin sources from the 10th century onwards (Bogolepov, 1907, 1908, 1911). Other studies have focused on accounts of anomalous weather in Russian sources (e.g. Borisenkov and Paseckij, 1983, 1988) and on reconstructing historical climate (Burchinskij, 1957; Liakhov, 1984; Borisenkov, 1988; Klimanov et al., 1995; Klimenko et al., 2001; Slepcev and Klimenko, 2005; Klimenko and Solomina, 2010), river flows (Oppokov, 1933) and famine years (Leontovich, 1892; Bozherianov, 1907).

The most important collection of Russian narrative sources is the 43-volume Полное Собрание Русских Летописей ('Complete Collection of Russian Chronicles', abbreviated to ПСРЛ; Borisenkov and Paseckij, 1988). These chronicles document events including infestations of insects, droughts, wet summers, wet autumns, unusual frost events, famine, floods, storms and earthquakes. The records have been used, in conjunction with other European sources, by Borisenkov and Paseckij (1988) to reconstruct a qualitative Russian climate history for the last 1000 years. More recent reconstructions have extended beyond historical sources to include a variety of other climate proxies (e.g. Klimenko and Solomina, 2010). The development of index-based series from narrative sources has yet to be attempted, although reconstructions of specific meteorological extremes, including wet/dry/warm/cold seasons and floods plus related socio-economic events such as famines, have been published by Shahgedanova (2002) (based on Borisenkov and Paseckij, 1983).

2.2. Temperature indices

Temperature is the most common meteorological phenomenon analysed ~~in~~ using an index approach over northern and central Europe ~~an indices~~. Authors who have developed temperature index series include Christian Pfister (1984, 1992, 1999), Pierre Alexandre (1987), Rudolf Brázdil (e.g. Brázdil and Kotyza, 1995, 2000; Brázdil et al., 2013a; spanning periods from 1000-1830 CE), Rüdiger Glaser (e.g. Glaser et al., 1999; Glaser, 2001; Glaser and Riemann, 2009; 1000-2000 CE), Astrid Ogilvie and Graham Farmer (1997; 1200-1439 CE), Gabriela Schwarz-Zanetti (1998; 1000-1524 CE), Lajos Rácz (1999; 16th century onwards), the Dutch working group around Aryan van Engelen (Van Engelen et al., 2001; Shabalova and van Engelen, 2003), Maria-João Alcoforado et al. (2000; 1675-1715 CE), Elena Xoplaki et al. (2001; 1675-1715 and 1780-1830 CE), Anita Bokwa et al. (2001; 16th and 17th centuries), Petr Dobrovolný et al. (2009), Dario Camuffo et al. (2010; 1500-2000 CE), Maria Fernández-Fernández et al. (2014; 2017; 1750-1840 CE), Laurent Lützenburger (2015; 1400-1530 CE) and Chantal Camenisch (2015a; 2015b; 15th century). The basis of these reconstructions is mainly narrative sources, or in the case of Brázdil and Kotyza (1995, 2000) and Fernández-Fernández et al. (2014), a single narrative source. However, depending on the epoch, narrative materials are supplemented by information from early weather diaries, administrative records and legislative sources. The majority of these studies (e.g. Pfister, 1984, 1992; Brázdil and Kotyza, 1995; Glaser et al., 1999; Pfister, 1999; Rácz, 1999; Brázdil and Kotyza, 2000; Glaser, 2001; Van Engelen et al., 2001; Shabalova and van Engelen, 2003; Dobrovolný et al., 2009; Glaser and Riemann, 2009; Camuffo et al., 2010) include an overlap with available instrumental data.

In Europe, different types of index scales have been used. As noted above, Christian Pfister (1984) developed a seven-point scale with a monthly resolution for temperature and precipitation (e.g., for temperature, -3: extremely cold, -2: very cold, -1: cold, 0: normal, 1: warm, 2: very warm, 3: extremely warm). Most historical climatologists follow this approach, though in some cases less granulated versions have had to be applied due to limited source density or quality. For instance, Glaser (2013) followed Pfister's indexing approach but used a three-point scale for the period 1000-1500 as information on weather appear only occasionally in documentary sources from this time. Schwarz-Zanetti (1998), Lützenburger (2015) and Camenisch (2015a) have also applied seven-point indices for the late Middle Ages, the latter two series at a seasonal resolution (Figure 2).




Figure 2: Comparison of seven-point winter temperature indices for Metz (Litzenburger, 2015) and the Low Countries (Belgium, Luxembourg and The Netherlands; Camenisch, 2015a) for the period 1420-1500, reconstructed using the Pfister index approach. Zero values for specific years are indicated by a small bar.

In addition to these studies, ~~three-four~~ other approaches exist for Europe. ~~First;~~ (i) IJnsen's temperature index (IJnsen and Schmidt, 1974) consists of a nine-point scale, which was also adopted by ~~Van Engelen et al. (2001).~~ ~~Second;~~ (ii) Alexandre (1987) used a five-point scale seasonal index, with categories from -2 (very warm) to +2 (very cold) and 0 being attributed to non-documented seasons; (iii) Fernández-Fernández et al. (2014; 2017) used a three-point-scale: (+1: warmer than usual; 0: normal; -1: colder than usual) and (iv) Domínguez-Castro et al. (2015) a five-point index (+2: very hot; +1: hot; 0: normal; -1: cold; -2: very cold). As noted in section 2.1, Klemm (1970) proposed a two-point index (warm/cold) for winter conditions.

2.3. Precipitation indices

Many of the authors mentioned in section 2.2 have also published precipitation indices. These reconstructions are usually based on the same source materials as the temperature indices (an exception being Dobrovolný et al., 2015). However, for certain regions, very specific source types exist that are more favourable for precipitation reconstructions than temperature – see, for example, the precipitation series for ~~southern Spain~~ the Mediterranean – based on the analysis of urban annals, religious chronicles and books of church and city archives (e.g. Rodrigo et al., 1994; Rodrigo et al., 1998; Rodrigo et al., 1999; Rodrigo and Barriendos, 2008; Fernández-Fernández et al., 2014; Domínguez-Castro et al., 2015; Fernández-Fernández et al., 2015). These series span various periods of the 16th to 20th centuries and, in some cases, overlap with instrumental data.

Often the same scale is applied for both temperature and precipitation indices; however, in certain regions, precipitation indices may show more gaps than their temperature counterparts as data may be seasonal or more sporadic. The studies by Van Engelen et al. (2001), ~~and~~ Alexandre (1987), ~~Fernández-Fernández et al. (2014; 2017) and~~ Domínguez-Castro et al. (2015) are exceptions, in that ~~both each~~ adopted a different or more rudimentary scale for precipitation compared to their temperature reconstructions. ~~—~~ Van Engelen et al. (2001) opted for a five-point scale for precipitation compared to a nine-point scale for temperature, and Alexandre (1987) a three-point rather than five-point index. Alexandre's (1987) precipitation index is also relatively simple and separates events by their nature (1: Snow; 2: Rain; 3: Dry conditions) rather than intensity. Fernández-Fernández et al. (2014; 2017) used a two-point scale (0: total absence of rain; 1: occurrence of rain) and Domínguez-Castro et al. (2015) a four-point scale.

197 Index series based on historical records of rogation ceremonies – closely linked to precipitation (or a lack thereof) – warrant
198 separate discussion. This source type is particularly valuable for western Mediterranean regions (e.g. Álvarez Vázquez,
199 1986; Martín-Vide and Vallvé, 1995; Barriendos, 1997; Piervitali and Colacino, 2001; Domínguez-Castro et al., 2008;
200 Barriendos, 2010; Domínguez-Castro et al., 2010; Garnier, 2010; Domínguez-Castro et al., 2012b; Fragoso et al., 2018;
201 Tejedor et al., 2018; Gil-Guirado et al., 2019; Bravo-Paredes et al., 2020). However, as most studies base their indices on the
202 type or cost of ceremonies – or the space within individual documents devoted to describing each ceremony – rather than a
203 meteorological entity, we do not go into further detail.

204 **2.4. Flood indices**

205 Flood events – the result of short periods of heavy precipitation and/or prolonged rainfall – can also be classified using
206 indices. The basis of European flood indices include narrative sources, administrative records such as bridge master's
207 accounts (e.g. those in Wels, Austria, which span the period 1350-1600 CE; Rohr, 2006, 2007, 2013), historic flood marks
208 and river profiles (Wetter et al., 2011; spanning 1268-present and overlapping with instrumental data). In some regions, the
209 availability and characteristics of sources may vary, and certain source types may be more important for flood reconstruction
210 than others. This is, for instance, the case in Hungary, where charters play a particularly important role in flood
211 reconstruction (Kiss, 2019; for the period 1001-1500 CE).

212 The scales used for flood reconstruction differ slightly from those used for the reconstruction of temperature and
213 precipitation. Drawing on Brázdil et al. (1999; which spans the 16th century), scholars mainly from Central Europe (e.g.
214 Sturm et al., 2001 [for the period 1500 CE-present]; Glaser and Stangl, 2003; 2004 [1000 CE-present]; Kiss, 2019) and
215 France (Litzenburger, 2015) have applied a three-point scale. In contrast, Pfister (1999), Wetter et al. (2011) and Salvisberg
216 (2017; 1550-2000 CE) used a five-point scale for floods of the River Rhine in Basel and the River Gürbe in the vicinity of
217 Bern. The French historian Emmanuel Garnier also developed a five-point scale to reconstruct flood time-series from 1500
218 to 1850 CE, taking into consideration the spatial extent and economic consequences of each event (Garnier, 2009, 2015). A
219 novel feature of the Garnier index is that it includes a -1 value for events where intensity cannot be estimated through
220 documentary sources. Rohr (2006, 2007, 2013) chose a four-point scale for his flood reconstruction of the river Traun in
221 Wels (Austria). In many cases, the index values express the amount of flood damage and/or the duration of flooding in
222 combination with the geographical extent (e.g. Pfister and Hächler, 1991 [covering the period 1500-1989 CE]; Salvisberg,
223 2017; Kiss, 2019). Comprehensive overviews of flood reconstruction, including the index method, are given in Glaser et al.
224 (2010), Brázdil et al. (2012) and recent work by the PAGES Floods Working Group synthesised in Wilhelm et al. (2018).

225 **2.5. Drought indices**

226 Drought events are closely linked to precipitation variability. As a result, many analyses of historical European droughts use
227 indices adapted from precipitation reconstructions. Evidence of past droughts can be found in administrative sources, diaries,
228 newspapers, religious sources and epigraphic evidence (see Brázdil et al., 2005; Brázdil et al., 2018; Erfurt and Glaser, 2019
229 [which spans the period 1800 CE-present]). Different approaches exist in historical climatology to express the severity of
230 droughts in index form. Brázdil and collaborators (2013b) proposed a three-point scale (-1: dry; -2: very dry; -3: extremely
231 dry) adapted from the precipitation indices described in section 2.3. Dry periods appear only in the drought index if they last
232 for at least two successive months. A similar approach is used by Pfister et al. (2006), Camenisch and Salvisberg (2020;
233 covering 1315-1715 CE) and Bauch et al. (2020; 1200-1400 CE). However, Garnier (2018) applies a five-point scale with an

additional sixth category for known drought-years with insufficient evidence for a more precise classification. Readers are referred to Brázdil et al. (2018) for a detailed discussion of the different types of drought indices.

2.6. Other indices

In Europe, the index method has only rarely been applied in contexts other than for temperature, precipitation, flood and drought reconstruction. Pichard and Roucaute (2009) developed, for example, an index for snowfall in the French Mediterranean region since 1715 [CE](#), including ordinal categories escalating from 1 to 3 depending on the event duration and quantity of snow fallen. This study is based on information from dairies and other urban narratives sources. Marie-Luise Heckmann (2008, 2015), coming from the field of historical seismology and seemingly unconnected to discussions in historical climatology, developed a combined temperature/precipitation index that differentiates winters and summers by weather description and phenological phenomena; this index was applied to documentary data from late-medieval Prussia and Livonia [\(1200-1500 CE\)](#).

Sea ice reconstructions for the seas around Iceland have been developed by Astrid Ogilvie, the pioneer of Icelandic climate history (Ogilvie, 1984, 1992; Ogilvie and Jónsson, 2001). She developed a monthly resolution sea-ice index based on historical observations in 37 sectors of the sea around Iceland (Ogilvie, 1996), including sightings of sea-ice in ships' logbooks, whalers' and sealers' charts, diaries, letters, books and newspapers. The index values hence vary from 1 to (theoretically) 37, with data weighed by source reliability. Pre-1900 [CE](#) records report single observations of icebergs and varying concepts of sea-ice have to be taken into consideration. The record is presented as a 5-year summarised value for the period 1600-1784 [CE](#), with monthly and annual values given from 1785 to present.

3. Climate indices in Asia

3.1. Origins of documentary-based indices in Asia

The use of the index approach in Asia is limited to research in China and India. With the exception of Japan, historical climatology research is either in its infancy or completely absent in other parts of the continent (Adamson and Nash, 2018). Very little work to reconstruct climate from documentary sources has occurred in southeast Asia, for example, and efforts to utilise records from the Byzantine Empire (Telelis, 2008; Haldon et al., 2014) and Muslim world (e.g. Vogt et al., 2011; Domínguez-Castro et al., 2012a) are only recently emerging. In Korea, only Kong and Watts (1992) have developed anything resembling climate indices, categorising individual years as warm/cold or dry/humid using information from diaries and histories.

Climate reconstruction work in China has developed largely independently from European historical climatology traditions. The Central Meteorological Bureau of China has published several fundamental works on Chinese wet/dry series. In 1981, a milestone work showed 120 cities with a five-point wet/dry series for the whole of China spanning the period 1470 to 1979 CE (Central Meteorological Bureau of China, 1981). Nowadays, most reconstructions (including coldness, drought, frost, hail and others) are based on the *Compendium of Chinese Meteorological Records of the Last 3,000 Years* edited by Zhang De'er (2004). This compendium provides details of a wide range of historical meteorological phenomena from across China at a daily level. However, due to an imbalance in population distribution, records are more abundant for eastern than western China (Ge et al., 2013). In India, the only study to use an index approach (Adamson and Nash, 2014) was developed from Nash and Endfield's work in southern Africa (see section 4); there were, however, several differences in approach, notably the inclusion of calibration tables.

One country where the field of historical climatology is relatively well-developed is Japan. Japan has weather data recorded in documents dating back to at least 55 CE (Ingram et al., 1981), and diaries in particular have been utilised to reconstruct

climate conditions (e.g. Mikami, 2008; Zaiki et al., 2012; Ichino et al., 2017; Shō et al., 2017). Access to documentary data on past weather phenomena is provided by detailed collections that evaluate historical sources (Mizukoshi, 2004-2014; Fujiki, 2007). However, Japanese historical climatology has no tradition of using indices, instead tending to use information in documentary sources to reconstruct units of meteorological measurement such as temperature and precipitation directly. For example, Mikami (2008) correlated mean monthly summer temperature with number of rain days. Mizukoshi (1993) and Hirano and Mikami (2008) used historical records to provide detailed reconstructions of weather patterns. Mizukoshi (1993) divided rainy seasons into three types: “heavy rain type”, “light rain type” and “clear rainy season type”, although these are not indices *per se*. In a similar way, Itō (2014) distinguished precipitation in categories such as “persisting rainfall” or “long downpour”, depending on seven keywords for each category. He used a similar approach to define indicators for cold spells, using keywords such as “cold”, “frost”, and “put on cotton [clothes]”. This keyword method for climatic conditions is also applied by Tagami (2016). There has also been much effort to reconstruct climate from climate-dependent phenomena such as cherry blossom or lake freezing dates (e.g. Aono and Kazui, 2008; Mikami, 2008; Aono and Saito, 2010).

3.2. Types of documentary evidence used to create index series

Historical climate index development in India has used a similar range of sources to those noted above for Europe – specifically ~~18th and 19th~~ newspapers and private diaries spanning the period 1781 to 1860 CE, supplemented by government records, missionary materials and some reports (Adamson and Nash, 2014). The sources used for the development of climate indices in China, however, are very different and require further explanation.

The earliest known written weather records in China, inscribed onto oracle bones, bronzes and wooden scripts, date to the Shang dynasty (~1600 BCE). These records were intended for weather forecasting, but later included actual weather observations (Wang and Zhang, 1988). Emperors of succeeding dynasties compiled more systematic records to allow them to better understand the weather, forecast harvests and hence maintain social stability (Tan et al., 2014). Some scholars use an old Chinese concept of *Tien* (or *Tian*, meaning Heaven) to explain the tradition. *Tien* was viewed as a medium used by gods and divinities to forward messages. Natural hazards (e.g. droughts and floods) were regarded as displaying *Tien*’s displeasure with the emperor and his court and were often followed by uprisings and rebellions (Perry, 2001; Pei and Forêt, 2018). To help them understand the long-term pattern of such hazards, imperial governments appointed specialists such as *Taishi* (imperial historians) or *Qintian Jian* (imperial astronomers) to record unusual and/or extreme weather events. Later, related environmental and socioeconomic events, such as early or late blossoming, agricultural conditions, famine, plagues and locust outbreaks, were also recorded (see Wang et al., 2018, for further details). This long tradition of chronicling has resulted in an exceptional range of materials for understanding and reconstructing past climates. It is worth noting, however that – due to a desire in imperial China to generalise details (Hansen, 1985) – phenomena were often only recorded as narrative descriptions with magnitude categorised as large, medium, or small.

The earliest official chronicle was *Han Shu* (‘The Book of Han’) written by Ban Gu (32-92 CE). However, many earlier historical books incorporate climate observations, including *Shi Ji* (‘Records of the Grand Historian’) by Sima Qian (145-86 BCE) and *Chun Qiu* (‘Spring and Autumn Annals’) compiled by Confucius (551-479 BCE) for the history of the *Lu* Kingdom (722- 481 BCE) (Wang and Zhang, 1988). Classic literature called *Jing Shi Zi Ji* was compiled in *Si Ku Quan Shu* (‘Complete Library in Four Branches of Literature’) published in 1787 (full-text digital versions are accessible at websites including Scripta Sinica: <http://hanchi.ihp.sinica.edu.tw/ihp/hanji.htm>). The *Shi* (meaning ‘history’) branch contains, but is not limited to, the ‘Twenty-Four Histories’ (later expanded to ‘Twenty-Five Histories’ by adding *Qing Shi Gao*, the ‘Draft History of Qing’), other historical books, documents of the central administration, local gazettes and private diaries (Ge et al., 2018).

While providing consistency in recording practices, the spatial coverage of official historical books was often limited to national capitals or other important locations. However, the writing of *Fang Zhi* – local chronicles or gazettes, popular in the Ming (1368-1643 CE) and especially Qing (1644-1911 CE) dynasties – substantively expanded the availability of documentary sources. Local gazettes contain unusual weather- and climate-related statements like those in the official chronicles, but incorporate additional details at provincial, prefectural, county or township levels depending on the local administrative unit. For more information, see Ge et al. (2018) and a database of local gazettes at <http://lcd.ccnu.edu.cn/#/index>.

In the 1980s, the Central Meteorological Bureau of China initiated a massive project for the compilation of weather- and climate-related records. The work resulted in the most influential publication in contemporary Chinese climate literature. The *Compendium of Chinese Meteorological Records of the Last 3,000 Years* edited by Zhang De'er (2004); this contains more than 150,000 records quoted from 7,930 historical documents, mostly local gazettes. To maximise the availability of the compendium, Wang et al. (2018) have digitised the records into the REACHES database (Figure 3). The quantity of records peaks in the last six hundred years, during the Ming and Qing dynasties. This is due to a large number of local gazettes spread across the country; however, only a few are available for the Tibetan Plateau and arid western regions. The Institute of Geographic Sciences and Natural Resources Research (Chinese Academy of Sciences) has also collated phenological records from historical documents (Zhu and Wang, 1973; Ge et al., 2003).


Two ~~series-sources~~ of documentary ~~sources-evidence~~ are of particular importance for historical climate reconstruction in China. Daily observations of sky conditions, wind directions, precipitation types and duration are recorded in *Qing Yu Lu* ('Clear and Rain Records') (Wang and Zhang, 1988). The records, however, are descriptive and only available for selected areas; these include Beijing (1724-1903 with six missing years), Nanjing (1723-1798), Suzhou (1736-1806), and Hangzhou (1723-1773). *Yu Xue Fen Cun* ('Depth of Rain and Snow') reported the measured depth of rainfall infiltration into the soil or depth of snow accumulation above ground in the Chinese units *fen* (~3.2mm) and *cun* (~3.2cm). From 1693 to the end of the Qing dynasty in 1911, these measurements were taken in eighteen provinces; however, many records include imprecise measurements and/or dates (Ge et al., 2005; Ge et al., 2011). Despite their descriptive and semi-quantitative nature, the two documentary sources are valuable for reconstructing past climate, especially for summer precipitation (Gong et al., 1983; Zhang and Liu, 1987; Zhang and Wang, 1989; Ge et al., 2011) and meiyu (or 'plum rains', marking the beginning of the rainy season; see Wang and Zhang, 1991) in different cities depending on the record length as described above. They are also useful for cross checking and/or validating climate indices derived from other documentary sources.

3.3. Temperature indices

The availability of documentary temperature indices for Asia is restricted to China. Zhu (1973) was the first Chinese scholar to use historical weather records and phenological evidence to identify temperature variability over the last 5,000 years (~3000 BCE to 1955 CE). He consulted a range of data sources for his reconstruction, including the dates of lake/river freezing/thawing, the start/end dates of snow and frost seasons, arrival dates of migrating birds, the distribution of plants such as bamboo, lychee and orange, the blossoming dates of cherry trees and harvest records. However, the study did not clearly indicate his methodology.

Winter temperature anomalies were initially regarded as key indicators of temperature changes in China (Zhang and Gong, 1979; Zhang, 1980; Gong et al., 1983; Wang and Wang, 1990a; Shen and Chen, 1993; Ge et al., 2003), as (i) there were more temperature-related descriptions in winter than in other seasons and (ii) winter temperatures have higher regional uniformity than summer temperatures (Wang and Zhang, 1992). However, this uniformity mainly reflects changes in the Siberian High system, so reconstructions of summer (and other season) temperature and precipitation anomalies to reflect

354 other aspects of monsoon circulation soon received increasing attention (see, for example, Zhang and Liu, 1987; Wang and
 355 Wang, 1990b; Yi et al., 2012).
 356



357
 358 **Figure 3:** Numbers of historical documentary records in the REACHES database for China. (a) Spatial distribution of
 359 records at 1,692 geographical sites across China. (b) Temporal evolution of the records in the database from 1 to 1911 CE
 360 (brown series); inset (blue series) shows the same data for 1 to 1350 CE but with an expanded vertical axis.

361 Zhu’s (1973) pioneering work has had a great influence upon the development of historical climatology in China. Successive
 362 studies used a similar approach to reconstruct winter temperature indices for every decade from the 1470s to 1970s by
 363 counting the frequency of years with cold- or warm-related records (Zhang and Gong, 1979; Zhang, 1980; Shen and Chen,
 364 1993; Zheng and Zheng, 1993). Zhang (1980) adopted binary (cold/warm) categories and further developed an equation to
 365 derive decadal temperature indices [for the period 1470-1970 CE](#) (see Section 8.2); this approach was applied in several
 366 studies (Gong et al., 1983; Wang and Wang, 1990b; Zheng and Zheng, 1993; Man, 1995).

The formal development of an ordinal-scale temperature index was first introduced by Wang and Wang (1990b) who used a four-point scale to build decadal winter cold index series [for the period 1470-1979 CE in eastern China](#) (0: no or light snow or no frost; 1: heavy snow over several days; 2: heavy snow over months; 3: heavy snow and frozen ground until the following spring). This approach was widely applied in subsequent series in different regions, for different seasons and at differing temporal resolutions (Wang and Wang, 1990a; Wang and Wang, 1990b; Wang et al., 1998; Wang and Gong, 2000; Tan and Liao, 2012; Tan and Wu, 2013). For example, Wang and Gong (2000) developed a fifty-year resolution winter cold index for eastern China spanning the period 800-2000 CE. Tan and colleagues adapted the approach to reconstruct decadal temperature index series (-2: rather cold; -1: cold; 0: normal; 1: warm) in the Ming_ (1368-1643 CE; Tan and Liao, 2012) and Qing dynasties_ (1644-1911 CE; Tan and Wu, 2013) in the Yangtze delta region.

3.4. Drought/flood and moisture indices

China has a particularly rich legacy of documents describing historical floods and droughts, and using such records to define drought-flood series has a long tradition. Zhu (1926) and Yao (1943) presented the earliest drought-flood series for all of eastern China [\(206 BCE-1911 CE\)](#), although their temporal and spatial resolutions are vague. Due to the higher number of available records for the last several hundred years, reconstructions using frequency counts were avoided in their series; instead the ratio between flood and drought events was used to build moisture indices (see section 8.2). Examples of other early studies include Yao (1982), Zhang and Zhang (1979), Zheng et al. (1977) and Gong and Hameed (1991).

Beginning in the 1970s, the Central Meteorological Administration initiated a project to reconstruct historic annual precipitation. This adopted a five-point ordinal scale (1: very wet; 2: wet; 3: normal; 4: dry; 5: very dry) to form drought-flood indices for 120 locations in China spanning the period 1470-1979 CE (Academy of Meteorological Science of China Central Meteorological Administration, 1981). The indices were compiled based on the evaluation of historical descriptions (section 8.2), with the series later extended to 2000 CE (Zhang and Liu, 1993; Zhang et al., 2003). Most reconstructions in China now use this five-point index (Zheng et al., 2006; Tan and Wu, 2013; Tan et al., 2014; Ge et al., 2018). For example, Zhang et al. (1997) used the approach to establish six regional series of drought-flood indices for eastern China (from the North China Plain to the Lower Yangtze Plain) spanning the period 960-1992 CE. Zheng et al. (2006) developed a dataset covering 63 stations across the North China Plain and the middle and lower reaches of the Yangtze Plain and reconstructed a drought-flood index series spanning 137 BCE to 1469 CE.

Adamson and Nash (2014) also adopted a five-point index series when reconstructing monsoon precipitation in western India (Figure 4). Where data quality allowed, indices were derived for individual ‘monsoon months’ (May/June, July, August and September/October) and summed to produce an index value for each entire monsoon season. Where monthly-level indices could not be constructed, indices pertaining to the whole monsoon were generated directly from narrative sources. The five-point index was chosen to correspond with the terminology currently used by the Indian Meteorological Department for their seasonal forecasts (from ‘deficient’ to ‘excess’ rainfall) and regular reports of rainfall conditions (a 4-point scale from ‘scanty’ to ‘excess’, with a fifth category ‘heavy’ added by the authors). As each of these correspond to percentage deviations from a rainfall norm, this allowed the generation of calibration tables within an instrumental overlap period, to assign descriptive terms to specific index points (e.g. the term ‘seasonable rain’ to the category +1 ‘excess’). This should allow the same methodology to be repeated elsewhere in India but limits the methodology to the subcontinent.




Figure 4: Five-point Western India Monsoon Rainfall reconstruction for 1780-1860. The reconstruction is a combination of separate series for Mumbai, Pune and the Gulf of Khambhat (see inset). Monsoon index categories map broadly onto Indian Meteorological Department (IMD) descriptors of seasonal monsoon rainfall (data for reconstruction from Adamson and Nash, 2014). Zero values are shown as small bars; years with insufficient data to generate an index value are left blank.

3.5. Other series

Several other studies have used weather descriptions within documentary records to reconstruct past climate series in China. These include reconstructed winter thunderstorm frequency (Wang, 1980, spanning 250 BCE-1900 CE), dust fall (Zhang, 1984, for the period 1860-1898 CE; Fei et al., 2009, for the past 1700 years) and typhoon series in Guangdong (Liu et al., 2001, 1000-1909 CE) and coastal China (Chen et al., 2019, 0-1911 CE). Many scholars have also used information in *Qing Yu Lu* and *Yu Xue Fen Cun* to count and build winter snowfall days series (Zhou et al., 1994; Ge et al., 2003), while Hao et al. (2012) have further used the series to regress annual winter temperatures over the middle and lower reaches of the Yangtze River since 1736.

Phenology-related phenomena have also been widely used in China to indicate past climate variability (Liu et al., 2014). Flower blossom dates in Hunan between 1888 and 1916 (Fang et al., 2005) and in the Yangtze Plain from 1450 to 1649 (Liu, 2017) were used to indicate temperature change. The date of the first recorded ‘song’ of the adult cicada has also been used to reconstruct precipitation change during the rainy season in Hunan from the late 19th to early 20th century (on the principle that cicada growth to adulthood requires sufficient humidity, and this coincides with the peak rainy season; Xiao et al., 2008). In recent years, researchers have been able to reconstruct various series including typhoons (Chen et al., 2019; Lin et al., 2019) and droughts (Lin et al., 2020) from the compendium of Chinese records compiled by Zhang (2004).

Using descriptions of agricultural outputs in the *Twenty-Four Histories* and *Qing History*, Yin et al. (2015) developed a grain harvest yield index and used this to infer temperature variations from 210 BCE to 1910 CE. Details of outbreaks of Oriental migratory locusts in these same histories have been used by Tian et al. (2011) to construct a 1910-year-long locust index through which precipitation and temperature variations can be inferred. *The History of Natural Disasters and Agriculture in*

427 *Each Dynasty of China*, published by the Chinese Academy of Social Science (1988), includes details of disasters such as
428 famines to reconstruct indices of climate variability during the imperial era.

429

430 **4. Climate indices in Africa**

431 **4.1. Origins of documentary-based indices in Africa**

432 Compared to the wealth of documentary evidence available for Europe and China, there are relatively few collections of
433 written materials through which to explore the historical climatology of Africa (Nash and Hannaford, 2020). The bulk of
434 written evidence stems from the late 18th century onwards, with a proliferation of materials for the 19th century following
435 the expansion of European missionary and other colonial activity.

436 Most historical rainfall reconstructions for Africa use evidence from one or more source type. A small number of studies are
437 based exclusively upon early instrumental meteorological data. Of these, some (e.g. the continent-wide analysis by
438 Nicholson et al., 2018) combine early rain gauge data with more systematically collected precipitation data from the 19th to
439 21st centuries, to produce quantitative time series. Others, such as Hannaford et al. (2015) for southeast Africa, use data
440 digitised from ships' logbooks to generate quantitative regional rainfall chronologies. Most climate reconstructions,
441 however, make use of narrative accounts to develop relative rainfall chronologies based on ordinal indices, either for the
442 whole continent or for specific regions.

443 While drawing upon European traditions and sharing many similar elements, methodologies for climate index development
444 in Africa have evolved largely in isolation from approaches in Europe (see section 8.3). The earliest work by Sharon
445 Nicholson, for example, was published around the same time that [Henry-Hubert Lamb](#) was developing his index approach
446 (Nicholson, 1978a, 1978b, 1979, 1980). Her early methodological papers on precipitation reconstruction (Nicholson, 1979,
447 1981, 1996) use a qualitative approach to identify broadly wetter and drier periods in African history. A seven-point index
448 (+3 to -3) integrating narrative evidence with instrumental precipitation data was introduced in Nicholson (2001) and
449 expanded in Nicholson et al. (2012a) and Nicholson (2018).

450 The many regional studies in southern Africa owe their approach to the work of Coleen Vogel (Vogel, 1988, 1989), who
451 drew on Nicholson's research but advocated the use of a five-point index to classify rainfall levels in the Cape region of
452 South Africa (+2: very wet, severe floods; +1: wet, good rains; 0: seasonal rains; -1: dry, months of no rain reported; -2: very
453 dry, severe drought). Subsequent regional studies, starting with Endfield and Nash (2002) and Nash and Endfield (2002),
454 have adopted the same five-point approach.

455 **4.2. Precipitation indices**

456 The main continent-wide index-based series for Africa originate from research undertaken by Sharon Nicholson (e.g.
457 Nicholson et al., 2012a). This series uses a seven-point scale and has been used to explore both temporal (Figure 5) and
458 spatial (Figure 6) variations in historical rainfall across Africa during the 19th century. One regional rainfall reconstruction is
459 available for West Africa, ~~spanning 1750-1800 and also using a seven-point scale~~ (Norrgård, 2015, spanning 1750-1800 CE
460 and using a seven-point scale) [and one for Kenya](#) (Mutua and Runguma, 2020, spanning 1845-1976 CE with a five-point
461 scale). The greatest numbers of regional reconstructions – all using a five-point scale – are available for southern Africa.
462 These include chronologies [covering all or part of the 19th century](#) for the Kalahari (Endfield and Nash, 2002; Nash and
463 Endfield, 2002, 2008) and Lesotho (Nash and Grab, 2010), and – most recently – Malawi (Nash et al., 2018) and Namibia
464 (Grab and Zumthurm, 2018). Several reconstructions are available for South Africa, including separate [19th century](#) series
465 for the Western and Eastern Cape, Namaqualand and present-day KwaZulu-Natal (Vogel, 1988, 1989; Kelso and Vogel,
466 2007; Nash et al., 2016). Most studies, including the continent-wide series, reconstruct rainfall at an annual level, but, where

information density permits, it has been possible to construct rainfall at seasonal scales (e.g. Nash et al., 2016). Regional studies from southern Africa have recently been combined with instrumental data and other annually-resolved proxies (including sea surface temperature data derived from analyses of fossil coral) to produce two multi-proxy reconstructions of rainfall variability (Neukom et al., 2014a; Nash et al., 2016).




Figure 5: Seven-point "wetness" index series for 1801 to 1840 for the 90 homogenous rainfall regions of Africa indicated across the x-axis. This series is reconstructed using documentary and instrumental data, with data gaps infilled using substitution and statistical inference (see section 8.3 and Nicholson et al., 2012a). From left to right, the regions approximately extend by latitude from the northern (region 1 – Northern Algeria/Tunisia) to southern (region 84 – western Cape, South Africa) extremes of the continent. Anomalies in the numbering sequence are regions 85, 86, 90 (all equatorial Africa), 87 (eastern Africa) and 88, 89 (Horn of Africa).





Figure 6: Rainfall anomaly patterns for 1835 and 1888 for the 90 homogenous rainfall regions of Africa delineated on the maps (modified after Nicholson et al., 2012b).

481 **4.3. Temperature indices**

482 To date, the only study exploring temperature variations in Africa using an index approach is an annually-resolved
483 chronology of cold season variability spanning 1833-1900 CE for the high altitude kingdom of Lesotho in southern Africa
484 (Grab and Nash, 2010). This uses a three-point index for winter severity (normal/mild; severe; very severe) and identifies
485 more severe and snow-rich cold seasons during the early- to mid-19th century (1833-1854) compared with the latter half of
486 the 19th century (Figure 7). A reduction in the duration of the frost season by over 20 days during the 19th century is also
487 identified.



488

489 **Figure 7:** Three-point “cold season severity” index for Lesotho and surrounding areas during the 19th century (top), with
490 major volcanic eruptions indicated. The occurrence of snowfall events (bottom) during the same period is also shown
491 (modified after Grab and Nash, 2010).

492 **5. Climate indices in the Americas**

493 **5.1. Origins of documentary-based indices in the Americas**

494 The use of the index approach in climate reconstruction is variable across the Americas. Although sufficient historical
495 records exist in some regions, particularly the north-eastern United States since the 18th century, few researchers have
496 generated climate indices for the USA or Canada (White, 2018). Mexico, in contrast, has produced pioneering studies in
497 climate history, especially on extreme droughts (see Prieto and Rojas, 2018; Prieto et al., 2019). In South America,
498 documentary evidence is overall lower in quality and quantity compared to Europe, so more complex indices have been
499 replaced by simpler ones, which extend to the 1500s CE.

500 **5.2. Temperature, precipitation and river-flow indices**

501 The only index-based temperature and precipitation reconstructions for the USA and Canada are those produced by William
502 Baron and collaborators. Although influenced by the work of Pfister, Baron (1980, 1982) used a distinct content analysis of
503 weather diaries (see section 8.4) to produce open-ended seasonal indices of New England temperature and precipitation for
504 1620-1800 CE-, a period overlapping with the first local instrumental temperature series (which began in the 1740s)from

~~weather diaries~~. He later combined seasonal indices, early instrumental records and phenological observations to create annual temperature and precipitation series and reconstruct frost-free periods (Baron et al., 1984; Baron, 1989, 1995).

There are a number of valuable compilations of extreme droughts in Mexico (e.g. Florescano, 1969; Jáuregui, 1979; Castorena et al., 1980; Endfield, 2007) and research that has identified climate trends across the country for 1450-1977 CE (Metcalf, 1987; Garza Merodio, 2002). Garza Merodio, ~~who was a student of Mariano Barriendas and the Pfister school~~, systemised the frequency and duration of climatic anomalies in the Basin of Mexico for 1530-1869 CE. García-Acosta et al. (2003) developed an unprecedented catalogue of historic droughts in central Mexico for 1450-1900 CE. Later work compared this information with a tree-ring series and found a significant correlation between major droughts and ENSO years over the same period (Mendoza et al., 2005). Mendoza et al. (2007) constructed a similar series of droughts on the Yucatan Peninsula for the 16th to 19th centuries. Garza Merodio (2017) improved this index and extended it back in time (see Hernández and Garza Merodio, 2010), based on the frequency and complexity of rain ceremonies (16th to 20th centuries). This approach identified droughts in bishoprics and towns of Mexico. Most recently, Dominguez-Castro et al. (2019) developed series for rainfall, temperature and other meteorological phenomena for Mexico City using information recorded in the books of Felipe de Zúñiga and Ontiveros; these volumes provide meteorological data with daily resolution for the twelve years spanning 1775 to 1786 CE.

In South America, the most detailed available historical information is on the scarcity or abundance of water. For investigations into historical rainfall and river flow rates, most studies construct 5-7 classes of data with annual or seasonal resolution. For example, a number of flood series have been compiled for rivers in Argentina (Prieto et al., 1999; Herrera et al., 2011; Prieto and Rojas, 2012, 2015; Gil-Guirado et al., 2016) – see Figure 8. ~~In contrast, temperature records are less reliable and generally begin with the earliest instrumental data in the late 1800s CE (Prieto and García-Herrera, 2009; Prieto and Rojas, 2018), but there are exceptions (Prieto, 1983). Most temperature-related indices use three classes.~~ In Bolivia, Gioda and Prieto (1999) and Gioda et al. (2000) developed a precipitation series for Potosí beginning in 1574 CE. In northern Chile, Ortlieb (1995) also compiled a detailed precipitation series for the 1800s CE in northern Chile. In Colombia, Mora Pacheco has developed a drought series for the Altiplano Cundiboyacense spanning the period 1778-1828 CE (Mora Pacheco, 2018). Finally, Dominguez-Castro et al. (2018) present a precipitation instrumental series from Quito (1891-2015 CE) and a series of wet and dry extremes from rogation ceremonies from 1600 to 1822 CE. ~~In contrast, temperature records are less reliable and generally begin with the earliest instrumental data in the late 1800s CE (Prieto and García-Herrera, 2009; Prieto and Rojas, 2018), but there are exceptions (e.g. Prieto, 1983, which covers the 17th and 18th centuries). Most temperature-related indices use three classes. Ortlieb (1994) compared a record of precipitation in central Chile to the occurrence of ENSO events in the tropical Pacific. Ortlieb (1995) also compiled a detailed precipitation series for the 1800s CE in northern Chile.~~

Some of the world's most important index-based chronologies of the El Niño Southern Oscillation (ENSO) derive from the analysis of ENSO-related impacts recorded in South American documentary evidence. This area of research was pioneered by William Quinn and colleagues (Quinn et al., 1987; Quinn and Neal, 1992), with Quinn's chronologies revised and improved by various authors using additional primary documentary sources (e.g. Ortlieb, 1994; Ortlieb, 1995, 2000; García-Herrera et al., 2008)~~compared a record of precipitation in central Chile to the occurrence of ENSO events in the tropical Pacific.~~




Figure 8: Six-point index series of historical flow in the Bermejo River (northern Argentina) between 1600 and 2008 CE based on documentary evidence. These annual-level data were used to create the decadal-scale flood series in Prieto and Rojas (2015). Zero values are indicated by short orange bars.

5.3. Sea-ice and snowfall indices

Relatively few studies have developed indices of winter conditions for the Americas. Building on their content analysis approach and that of Astrid Ogilvie in Iceland (see section 2.6), Catchpole and Faurer (1983) and Catchpole (1995) produced open-ended annual sea-ice indices for the western and eastern Hudson Bay, spanning the period 1751-1869 CE. A different type of three-class index was developed for snowfall in the Andes at 33°S spanning 1600-1900 CE, based on the number of months per year that the main mountain pass between Argentina and Chile was closed (Prieto, 1984).

6. Climate indices in Australia

6.1. Origins of documentary-based indices in Australia

Like Africa, Australia has a limited history of using documentary records for developing regional climate indices. Aside from early compilations of 19th century colonial documents and newspaper records (Jevons, 1859; Russell, 1877), or climate almanacs published by the Australian Bureau of Meteorology (Hunt, 1911, 1914, 1918; Watt, 1936; Warren, 1948), few attempts were made in the 20th century to use historical sources to develop climate indices ~~during the 20th century~~. Those that were developed focussed predominantly on drought conditions (see, for example, Foley, 1957; McAfee, 1981; Nicholls, 1988). However, considerable effort has been given in recent years to reconstruct climate variability in south-eastern Australia since British colonisation in 1788 CE using both historical documents and instrumental observations (e.g. Gergis et al., 2009; Fenby, 2012; Fenby and Gergis, 2013; Gergis and Ashcroft, 2013; Ashcroft et al., 2014a; Ashcroft et al., 2014b; Gergis et al., 2018; Ashcroft et al., 2019; Gergis et al., 2020). There have also been attempts to reconstruct storm and tropical cyclones along the east coast of Australia (e.g. Callaghan and Helman, 2008; Callaghan and Power, 2011, 2014; Power and Callaghan, 2016), although these are not index-based.

Documentary-based indices for Australia have focussed on regional rainfall histories using largely ~~using~~ material from previously published drought and/or rainfall compilations (Fenby and Gergis, 2013). These compilations contained a vast collection of primary source material including newspaper reports, unpublished diaries and letters, almanacs, observatory reports, 19th century Australian publications and official government reports. For example, the seminal 19th century sources

of Jevons (1859) and Russell (1877), that formed the foundation of the Fenby and Gergis (2013) analysis, contain 79 primary sources, including 40 accounts from personal diaries, letters and correspondence between a range of people in the colony with the authors. Most recently, Gergis et al. (2020) compiled colonial newspaper and government reports to ~~reconstruct~~ identify daily temperature extremes of snowfall and heatwaves from South Australia back to 1838. Although a temperature index from this material has not yet been developed, there is great potential to do so alongside recently homogenised 19th century instrumental temperature observations from the Adelaide region.

6.2. Precipitation and drought indices

The most extensive analysis of documentary records was compiled by Fenby (2012) and Fenby and Gergis (2013) as part of a large-scale project to reconstruct climate in south-eastern Australia using palaeoclimate, early instrumental and documentary data (Gergis et al., 2018). Fenby and Gergis (2013) used twelve secondary source compilations to collate monthly summaries of drought conditions experienced in five modern states in south-eastern Australia between 1788 and 1860 CE into a three-point index (wet, normal, drought). As explained in section 8.5, agreement between sources and several months of dry conditions was required before a period was considered a drought, rather than just ‘normal’ low summer rainfall. In coastal New South Wales, months of above average rainfall were only compiled where sufficiently detailed rainfall information was available (Fenby and Gergis, 2013). Given that Australian rainfall has high spatial variability, and many of the secondary sources only contained descriptions of localised floods or severe storm events, there were insufficient local reports from other regions to reconstruct larger-scale rainfall conditions using the sources considered.

To combine instrumental and documentary data into a single series spanning European settlement of Australia (1788 CE ~~to the present~~), Gergis and Ashcroft (2013) developed a three-point drought and wet year index based on instrumental rainfall observations from using a five-station network ~~of historical instrumental rainfall observations from in~~ the Sydney region ~~(for spanning 1832-1859) and, along with~~ a 45-station rainfall network from across south-eastern Australia ~~over the period (1860-2008). This~~ As with the “wetness” index for Africa (Figure 5), the instrumental data were converted to an index so they could be combined with the documentary-based index of Fenby and Gergis (2013) to create a single, complete rainfall reconstruction. was then combined with the documentary-based index of Fenby and Gergis (2013). Good agreement was found during the overlapping period between instrumental and documentary-derived indices (1832-1860), and between the eastern New South Wales index and the wider south-eastern Australian indices. This, providesing some confidence that the two indices could be combined, and that data from the very early period, when only eastern New South Wales ~~details records~~ are available, ~~can provide information on~~ are indicative of conditions experienced in the ~~wider broader~~ region.


Given the exploratory nature of this work in south-eastern Australia, the aim of these studies was to use documentary and instrumental data to simply identify the occurrence of wet and dry years in the first instance, rather than develop a more finely resolved scale of the magnitude of the rainfall anomalies. The recent digitisation and analysis of daily instrumental rainfall data from Sydney, Melbourne and Adelaide (Ashcroft et al., 2019) provides an excellent opportunity to develop ~~documentary~~ indices combining documentary and instrumental data from these regions in the future.

7. Climate indices and the world’s oceans

7.1. Challenges in generating documentary-based indices for the world’s oceans

The oceans constitute a challenging environment for historical climatologists. Written evidence of past weather at sea is generally local in scope, especially before the 17th century, and direct weather observations scarcely extend beyond the coast before the 15th century. Historical climatologists can use two categories of information to create reconstructions of past oceanic climate: (i) direct observations of weather, water, and sea ice conditions; and (ii) records of activities that were

610 influenced by weather and water conditions. Such information can be found in documents written at sea (on ships, boats or,
611 from the twentieth century, submarines; Figure 9), documents written on the coast within sight of the sea, and documents
612 written inland that record weather or activities at sea.



613
614 **Figure 9:** Journal written by a Dutch whaler during a voyage to the "Greenland Fishery," between Jan Mayen and Svalbard,
615 1615. Source: 0120 Oud archief stad Enkhuizen 1353-1815 (1872), Westfries Archief, Hoorn.


616 Between the 16th and 20th centuries, ships' logbooks are perhaps the most useful source type (see Wheeler, 2005a, 2005b;
617 Wheeler and García-Herrera, 2008; Ward and Wheeler, 2012; García-Herrera and Gallego, 2017; Degroot, 2018). Sailors
618 originally recorded the speed and direction of the wind in order to calculate their location, and their compass-aided
619 measurements of wind direction are often assumed to be true instrumental observations (Gallego et al., 2015). Yet naval
620 officers on different ships in the same fleet could record slightly different measurements, and they ~~However, there are~~
621 ~~challenges in using these sources~~ (Wilkinson, 2009; García-Herrera et al., 2018). For example, early modern mariners did
622 not always accurately estimate their longitude, or consistently describe whether recorded wind directions related to real or
623 magnetic north (Wilkinson, 2009; García-Herrera et al., 2018). Logs kept by flag officers – which survive in larger quantities
624 in early periods than logs kept by subordinate officers – may not include systematic weather observations. Ships did not sail
625 in sufficient numbers prior to the 18th and 19th centuries for scholars to use surviving logbooks for comprehensive regional
626 weather reconstructions, and, ~~indeed~~, many logbooks have been lost. Finally, logbooks written aboard British naval ~~some~~
627 ships copied ~~most~~ wind measurements earlier recorded in simple tables (~~called log-boards~~) and should therefore be
628 considered secondary sources for the purpose of climate reconstruction (Norrgård, 2017).

629 Ship logbooks of the 16th and 17th centuries, in particular, are most valuable when used alongside other documentary
630 evidence. Journals kept during exceptional voyages may provide similar environmental data but in a narrative format.
631 Accounts of the passage of ships through ports and tollhouses; the annual catch brought in by fishermen or whalers; or the

632 duration of voyages may provide evidence of changes in the distribution of sea ice or patterns of prevailing wind.
633 Correspondence, diary entries, intelligence reports, newspaper articles and chronicles may describe weather at sea, or
634 weather blown in from the sea, often at high resolution and occasionally for decades. Paintings, illustrations, and even
635 literature may provide insights into the changing frequency or severity of weather events at sea. These sources can
636 supplement other human records of the oceanic climate, including oral histories, or shipwrecks distributed in areas of heavy
637 trade (Chenoweth, 2006; Trouet et al., 2016).

638 7.2. Indices of wind direction and velocity

639 If carefully contextualised, information in written records of oceanic weather – especially ships’ logbooks and accounts of
640 naval voyages – can be quantified and entered into databases. The Climatological Database of the World’s Oceans
641 (CLIWOC; Figure 10), for example, quantified ~~data in~~ nearly 300,000 logbooks from 1750 to 1850 CE, and their data are
642 now among 456 million marine reports within the International Comprehensive Ocean-Atmosphere Data Set (ICOADS)
643 (García-Herrera et al., 2005b; Koek and Konnen, 2005; García-Herrera et al., 2006). By using such datasets, or by creating
644 databases of their own, ~~a small but growing group of~~ scholars ~~has attempted to have~~ reconstructed aspects of past climate at
645 sea, ~~or in many cases verifying or extending~~ verify reconstructions compiled by scientists using other means. ~~Some of the~~
646 ~~most important~~ High resolution reconstructions of regional trends in the frequency of winds from different directions, for
647 example, reveal broadscale atmospheric circulation changes associated with stratovolcanic eruptions, ENSO, the North
648 Atlantic Oscillation (NAO) or the monsoons of the Northern and Southern Hemispheres ~~are those of wind direction and~~
649 ~~velocity, which even when regional in scale can suggest broader changes in atmospheric circulation~~ (e.g. Garcia et al., 2001;
650 Küttel et al., 2010; Barriopedro et al., 2014; Barrett, 2017; Barrett et al., 2018; García-Herrera et al., 2018).



651
652 **Figure 10:** Plot of the position of all ships’ logbook entries in the CLIWOC database (Degroot and Ottens, 2020).

653 7.3. Indices of sea-ice extent

654 Records of sea ice in harbours and heavily trafficked waterways – or records of dues paid at ports and tollhouses – yield
655 easily quantified data. However, reports of sea ice at high latitudes in correspondence, logbooks or journals written before
656 the 19th century often give unclear descriptions of sea ice density, which makes it harder to determine how much sea ice
657 there might have been in different regions from year to year (Prieto et al., 2004). The resolution and precision of Arctic

index-based sea ice or iceberg reconstructions that rely on surviving early modern documents is accordingly quite low (Catchpole and Faurer, 1983; Catchpole and Halpin, 1987; Catchpole and Hanuta, 1989). An emerging way to circumvent this issue is to focus on particular regions where warm and cold ocean currents mixed, and that were sensitive to (a) changes in sea and air surface temperatures and (b) current strength, for example, around northern-western Svalbard or the Yugorsky Strait (Degroot, 2015). Logbook reports of the presence of sea ice in these target areas can be quantified, indexed, and used to develop reconstructions that suggest broadscale shifts in the strength of marine currents (Degroot, 2020).

7.4. Indices of precipitation and storms

Some ships' logbooks note the occurrence of precipitation at sea, and most record winds that must have influenced precipitation on land. Historical climatologists have therefore used logbooks to classify and graph precipitation at or near the sea (e.g. Wheeler, 2005b; Hannaford et al., 2015). Moreover, most documents that directly describe weather at sea or blown in from the sea faithfully report storms and at least approximately note their severity (Lamb, 1992; García-Herrera et al., 2004; García-Herrera et al., 2005a; Chenoweth and Divine, 2008; Wheeler et al., 2010). Reconstructions based on written evidence of damage inflicted along the coast, however, can be more problematic, as damage reflected both complex social conditions and environmental circumstances beyond the severity of storms (de Kraker, 2011; Degroot, 2018).

8. Methods for the derivation of climate indices

The preceding sections have highlighted the variable number of classes used in index-based climate reconstructions and hinted at the variety of different approaches to index development. This section summarises the main methodological approaches used to derive indices on the different continents, with an emphasis on temperature and rainfall series.

8.1. Climate index development in Europe – “Pfister indices”

In Europe, the most widely adopted approach to the reconstruction of temperature and rainfall variability for climatically-homogenous regions is through the development of seven-point indices (Pfister, 1984; Pfister et al., 2018), which the climate historian Franz Mauelshagen has termed “Pfister indices” (Mauelshagen, 2010). These indices are normally generated at a monthly level through the analysis of (bio)physically-based proxies and contemporary reports of climate and related conditions. This is not without its challenges, and requires a source-critical understanding of the evidence-base in addition to a knowledge of regional climates (Brázdil et al., 2010). To aid interpretation, any contemporary report should be accompanied by a range of information, including details of the date, time, location affected, author and source quality (see Brázdil et al., 2010; Pfister et al., 2018). The criteria used to allocate a specific month to a specific index category will vary from place to place according to regional climatic variability. Table 1, for example, illustrates the indicators used to classify individual months as either “warm” (+2/+3) or “cold” (-2/-3) in a temperature reconstruction for Switzerland (Pfister, 1992); these include regionally relevant phenomena such as the timing and duration of snowfall and various plant-phenological indicators. Pfister et al. (2018) recommend that monthly rankings of above +1 and below -1 should only be attributed based on proxy data such as phenological evidence, with values of -3 and +3 reserved only for exceptional months. An index value of 0 should only be used where reports of climate suggest normal conditions – an absence of data should be reported as a gap in the time series rather than a 0 value.

693
694

Table 1: Criteria used in the generation of seven-point temperature indices for “warm” (+2/+3) or “cold” (-2/-3) months in Switzerland (after Pfister, 1992; Pfister et al., 2018). *Italics* indicate criteria grounded in statistical analyses.

Month	“Cold” (index values of -2/-3)	“Warm” (index values of +2/+3)
Dec, Jan, Feb	Uninterrupted snow cover <i>Freezing of lakes</i>	Scarce snow cover Early vegetation activity
Mar	Long duration of snow cover Frequent snowfalls	Early sweet cherry flowering No snowfall
Apr	Several days of snow cover Frequent snowfalls	<i>Beech tree leaf emergence</i> <i>Early vine flower</i>
May	<i>Late grain and grape harvest</i> <i>Late vine flower</i>	<i>Early grain and grape harvest</i> <i>Start of barley harvest</i>
Jun	<i>Late vine flower</i> Several low altitude snowfalls	<i>Early grain and grape harvest</i> <i>High vine yields</i>
Jul	<i>Low vine yields</i> Snowfalls at higher altitudes	<i>High vine yields</i>
Aug	<i>Low tree ring density</i> Low sugar content of vine Snowfalls at higher altitudes	<i>High tree ring density</i> High sugar content of vine
Sep	Low sugar content of vine Snowfalls at higher altitudes	High sugar content of vine
Oct	Snowfalls, snow cover	Second flowering of spring plants
Nov	Long duration of snow cover	Second flowering of spring plants No snowfall

695

696 Once monthly index values have been generated, these can be summed to produce seasonal or annual classifications where
697 required. Three-month seasonal values can, as a result, fluctuate from -9 to +9 and annual values from -36 to +36 (see
698 Pfister, 1984). Implicit in this methodological approach is that runs of monthly indices are available with almost no gaps
699 (e.g. Litzenburger, 2015) or that, where gaps occur, there is a high probability that conditions during a given month reflect
700 the longer-term average for that month (e.g. Dobrovolný et al., 2009). Variations in source density, however, mean that it
701 may not always be possible to define indices at a monthly level. Such variations could simply be due to a scarcity of
702 available sources, or could be the product of seasonal variability that results in observations of a climate phenomenon being
703 concentrated in specific parts of the year (e.g. observations of rainfall in areas of Europe with a Mediterranean climate are
704 likely to be concentrated between September and April). Once monthly index values have been generated, these can be
705 summed to produce seasonal or annual classifications where required. Three-month seasonal values can, as a result, fluctuate
706 from -9 to +9 and annual values from -36 to +36 (see Pfister, 1984). In these cases situations where source density is
707 insufficient to permit the generation of monthly indices, researchers should (i) choose an appropriate temporal resolution (i.e.
708 seasonal or annual) based on the number and quality of available records, and (ii) develop specific seasonal- or annual-level
709 criteria – see, for example, the temperature and precipitation reconstructions for Belgium, Luxembourg and The Netherlands
710 generated by Camenisch (2015a) or the Mediterranean temperature series by Camuffo et al. (2010) (see, for example, the
711 temperature and precipitation reconstructions for Belgium, Luxembourg and The Netherlands generated by Camuffo et al.,
712 2010; Camenisch, 2015a). The methods used for verification and calibration are outlined in the following section.

713 In the development of his seven-point scale, Pfister assumed that temperature and precipitation followed a Gaussian
714 distribution. Initially, Pfister (1984) developed duodecile classes based on the frequency distribution of monthly
715 temperature/precipitation means for the sixty-year reference period 1901-1960 as the standard of comparison (Table 2). The
716 most extreme months (i.e. those given an index value of -3/+3) were those that fell into duodecile classes 1 and 12,
717 representing the 8.3% driest (or coldest) or 8.3% wettest (or warmest) months, respectively. Other index categories were
718 defined using 16.6% intervals. In the later version of his indices, Pfister (1999 and onwards) discontinued the use of
719 duodecile classes, using instead the standard deviation from the mean temperature/precipitation for the 1901-1960 reference

720 period to define index categories: $\pm 180\%$ (of the standard deviation from the reference period) for index values $-3/+3$, -
 721 $\pm 130\%$ for values $-2/+2$, and $\pm 65\%$ for values $+1/-1$.

722 **Table 2:** The definition of the weighted temperature and precipitation index values used in the creation of initial (pre-1999)
 723 seven-point ‘‘Pfister’’ indices (after Pfister, 1992).

	Lowest						Highest
	8.3%	16.6%	16.6%	16.6%	16.6%	16.6%	8.3%
Duodecile	1	2-3	4-5	6-7	8-9	10-11	12
Index	-3	-2	-1	0	1	2	3

724

725 8.2. Climate index development in Asia

726 In China, the quantification of historical records to reconstruct climate change originated with a Semantic Differential
 727 Method based on an analysis of each record’s content (see Central Meteorological Bureau of China, 1981; Su et al., 2014;
 728 Yin et al., 2015). Temperature series were traditionally established at a decadal scale only. In creating a series, each year was
 729 first defined as ‘cold’, ‘warm’ or ‘normal’ according to direct weather descriptions or environmental and phenological
 730 evidence. In contrast to the Pfister method (see section 8.1), ‘Normal’-normal was also used when there was insufficient
 731 information available to determine temperature abnormalities. This approach reflects the nature of most Chinese documents,
 732 where the primary mission of the recorders was to detail abnormal or extreme events; fewer descriptions of abnormal events
 733 are therefore interpreted as indicating conditions closer to normal. After each year had been defined as cold, warm or normal,
 734 an equation was then used to derive the decadal indices. The earliest example was published by Zhang (1980): $T_i = -[n_1 +$
 735 $0.3(10 - (\overline{n_1} + \overline{n_2}))]$, where T_i is the decadal winter temperature index, n_1 the number of cold years, n_2 the number of warm
 736 years, and 0.3 the empirical coefficient (see also Zhang and Crowley, 1989). The resulting value is always negative; the
 737 lower the value, the more severe the coldness.

738 A second approach to the construction of ordinal scale indices was developed by the Wangs in the 1990s (e.g. Wang and
 739 Wang, 1990a; Wang and Wang, 1990b; Wang et al., 1998). This used a four-point scale (0, 1, 2, 3) (Table 3). As in Europe,
 740 indices were generated through the analysis of phenological descriptions and contemporary reports of climate and related
 741 phenomena. Like Europe, criteria for individual index categories could also be adjusted for specific places at specific
 742 seasons according to geographical and climatic attributes. However, unlike the Pfister method, an index value of 0 could be
 743 used where there were missing data. The Wangs further introduced a statistical method to compare phenological evidence
 744 with modern (1951-1985 CE) and early instrumental data (1873-1972 CE in Shanghai) and allocate temperature ranges to
 745 ordinal scales (Wang and Wang, 1990b). An index value of -0.5 corresponded to a $-0.5\sim-0.9^\circ\text{C}$ temperature anomaly, a value
 746 of -1.0 to a $-1.0\sim-1.9^\circ\text{C}$ anomaly and a value of -2.0 to an anomaly of $\leq -2.0^\circ\text{C}$; values of 1.5 were added to indicate warm
 747 temperatures and -3.0 to capture extreme cold periods. These cold indices were then regressed with the decadal mean
 748 temperature (1873-1972 CE) to derive a coefficient through which the index value could be transferred into a ‘real’
 749 temperature.

750 **Table 3:** Criteria used in the development of temperature indices in China.

Cold index values				Temperature index values	
Wang, R. and Wang, S. (1990)		Wang, S. and Wang, R. (1990)		Tan and Wu (2013), adapted from Chen and Shi (2002)	
Index value	Criteria (winter)	Index value	Criteria (distinguishing four seasons; example of winter)	Index value	Criteria (winter and summer; example of winter)

0	No record of ice/frost; no snow; light snow	1.5	Warm records	1	Warm records such as 'winter warm as spring'
1	River/lake freezing; heavy snow over several days or several cm depth	-0.5	Heavy snow; freezing rain; ice glaze on trees	0	No specific records
2	River/lake frozen for weeks to allow human passage; heavy snow for months; snow frozen for months	-1.0	Frozen river or lake	-1	Heavy snow; freezing rain; ice glaze on trees
3	River/lake frozen for months to allow horse- drawn wagons or carriages to cross; heavy snow for months; ice melt in following spring	-2.0	Extreme cold; ocean water and large lakes or rivers frozen	-2	River/lake frozen for months to allow horse- drawn wagons or carriages to cross
		-3.0	River/lake frozen for months to allow horse- drawn wagons or carriages to cross		

751

752 Chen and Shi (2002) built upon Zhang (1980) and the Wangs' approaches in developing an equation to calculate decadal
753 temperature indices: $T_i = 10 - 2n_1 - n_2 + n_3$, where n_1 = number of extremely cold years, n_2 = number of cold years, n_3 =
754 number of warm years. A resulting decadal temperature index value of 10 denotes average conditions; <10 anomalous cold;
755 and >10 anomalous warm. Successive work (Tan and Liao, 2012; Tan and Wu, 2013) adopted the Chen and Shi (2002)
756 approach with a slight modification of the index criteria while retaining the four-point ordinal scale. The temperature series
757 generated using this approach have been incorporated into multi-proxy temperature reconstructions (e.g. Yi et al., 2012; Ge
758 et al., 2013). Zheng et al. (2007) and Ge et al. (2013) provide useful reviews of the approach used to generate temperature
759 indices in China.

760 As noted in section 3.2, drought-flood index reconstruction in China has a long tradition. Two main approaches are used.
761 Earlier studies adopted a proportionality index approach (Zhu, 1926; Yao, 1943). As explained by Gong and Hameed (1991),
762 Zhu used the equation $I = D/F$ to calculate the index, where D represents the number of droughts and F the number of
763 floods in a given time period. This equation is poorly defined if F or D is zero. Brooks (1949) modified this equation and
764 used the flood percentage, $I = 100 \times F/(F + D)$, to derive moisture conditions in Britain and some European regions from
765 100 BCE onwards at a 50-year resolution. Gong and Hameed (1991) further modified the equation as $I = 2F/(F + D)$ to
766 derive indices at a 5-year resolution. Their index takes the values $0 \leq I \leq 2$, with larger values reflecting wetter conditions.
767 Zhang and Zhang (1979) adopted a slightly different approach by counting the number of places with reported drought
768 events: $I_D = 2D/N$, where D represents the number of places having extreme drought (grade 5) and drought (grade 4) events
769 in a given year (see Table 4), and N is the total number of places.

770 The Academy of Meteorological Science of China Central Meteorological Administration (1981) adopted a five-point
771 ordinal scale approach to reconstruct annually resolved drought-flood indices in China. The key descriptors for each
772 classification (see Table 4) are mainly based on accounts of the onset, duration, areal extent and severity of each drought or
773 flood event in each location. They then assume a probability distribution of the five grades following a normal distribution: 1
774 (10%), 2 (25%), 3 (30%), 4 (25%), and 5 (10%). For the period of overlap between written and instrumental records (after
775 1950 CE), the graded series were compared against the observed May-September (major rainy season) precipitation and
776 regressed to transform the indices into numerical series (Table 4). Based on the five-point ordinal scale, Wang et al. (1993)
777 and Zheng et al. (2006) developed further formulae to calculate decadal drought-flood indices that can be applied to earlier
778 periods (i.e. before 1470) when less information is available.

779 **Table 4:** Criteria used in the generation of five-point drought-flood indices in China (Academy of Meteorological Science of
780 China Central Meteorological Administration, 1981). For more details, see Zhang and Crowley (1989), Zhang et al. (1997),
781 and Yi et al. (2012).

Index value	Norm	Transfer function for precipitation amount
1 (Very wet)	Prolonged heavy rain, continuous flood over two seasons, extensive flood, unusually heavy typhoon rain	$R_i > (\bar{R} + 1.17\sigma)$, where, \bar{R} is mean May-Sep precipitation, σ is standard deviation, R_i is precipitation in the i^{th} year
2 (Wet)	Spring or autumn prolonged rain with moderate damage, local flood	$(\bar{R} + 0.33\sigma) < R_i \leq (\bar{R} + 1.17\sigma)$
3 (Normal)	Favourable weather, usual case, or nothing special to be noted in records	$(\bar{R} - 0.33\sigma) < R_i \leq (\bar{R} + 0.33\sigma)$
4 (Dry)	Minor impacts of drought in a single season, local minor drought disaster	$(\bar{R} - 1.17\sigma) < R_i \leq (\bar{R} - 1.33\sigma)$
5 (Very dry)	Severe drought over a season, drought continued for several months, severe drought over an extensive area, or records describing extensive areas of barren land	$R_i \leq (\bar{R} - 1.17\sigma)$

782

783 8.3. Climate index development in Africa

784 Historical climate reconstructions for Africa use two different approaches to index development. The continent-wide rainfall
785 reconstruction by Nicholson et al. (2012a) is based upon 90 regions that are homogeneous with respect to interannual rainfall
786 variability. An underpinning assumption is that historical information for any location within a region – be it narrative or
787 instrumental – can be used to produce a precipitation time series representing that region. Instrumental rainfall data are
788 converted into seven “wetness” classes (-3 to +3) based on standard deviations from the long-term mean. A wetness index
789 value of zero corresponds to annual rainfall totals within +/-0.25 standard deviations of the mean. Index values of -1/+1 are
790 assigned to annual values between -0.25/+0.25 and -0.75/+0.75 standard deviations. Values of -2/+2 are given to annual
791 totals between -0.75/+0.75 and -1.25/+1.25 standard deviations, with more extreme departures classed as -/+3.

792 Documentary data are integrated by first assigning individual pieces of narrative evidence to a specific region; each piece of
793 evidence is then classified into one of the seven “wetness” categories. Like the approach used by Pfister, the presence of key
794 descriptors of climate conditions is used to distinguish these categories. The scores for each item of evidence for a specific
795 region/year are summed and averaged. Where there are several sources, a ‘0 index’ value represents an average of
796 conditions. Where only single sources are available, some contain so much climate-related information that, as in China,
797 absence of evidence for a specific season is taken to infer “normal” conditions; such cases are indicated in the original data
798 file accompanying the Nicholson et al. (2012a) reconstruction. Algorithms are then used to weight and combine
799 documentary and instrumental data for each region and year. These are defined subjectively according to the accuracy of the
800 quantitative versus qualitative indicators. For example, when one of each type is available, the qualitative indicator is
801 weighted twice as much as the gauge because of the inherent spatial variability within African rainfall. A second assumption
802 is that when the correlation between rainfall in two regions is >0.5 the regions are appropriate substitutes for each other
803 (Nicholson, 2001). In this way, classifications for regions without evidence for a given year can be derived by substitution.
804 Statistical inference (termed ‘spatial reconstruction’ by Nicholson) is then used to generate classifications for any remaining
805 regions. The cutoff of 0.5 was selected based on examination of time series that correlate with each other at various levels.
806 Those with a correlation of 0.5 showed marked similarity, though it should be noted that, in most cases, the correlation was
807 much higher, with the statistical significance being >0.001. Statistical inference is then used to generate classifications for
808 any remaining regions.

809

Regional rainfall reconstructions in southern Africa use an approach much closer to the Pfister method to classify documentary evidence into one of five rainfall classes (-2 to +2); these classes are ordinal rather than based on statistical distributions. Like the Pfister method, a '0 index' value is only awarded where narrative evidence suggests normal conditions – years with inconclusive or no data are left unclassified. Owing to the relatively paucity of documentary data for Africa compared to Europe, conditions for specific rainy seasons are categorised at a quarterly (e.g. Nash et al., 2016) or more commonly annual level. Again, key descriptors are used to distinguish the various index classes. The main point of divergence with the approach used by Nicholson is that – rather than assigning individual pieces of evidence to wetness classes and averaging – qualitative analysis is undertaken of all quotations describing weather and related conditions for an entire quarter/year (see Nash, 2017). These different methodological approaches, as well as the type of documentary evidence used, can introduce discrepancies between rainfall series for overlapping regions. Hannaford and Nash (2016) and Nash et al. (2018) note, for example, that the reconstructions in Nicholson et al. (2012a) for KwaZulu-Natal during the first decade of the 19th century and Malawi for the 1880s-1890s show generally drier conditions than overlapping series generated using different methods.

8.4. Climate index development in the Americas

Temperature, precipitation and phenological indices for North America have been based on a distinctive content analysis approach. This method was first applied to historical climatology in the 1970s to reconstruct freeze and break-up dates around Hudson Bay for the period 1714-1871 CE by quantifying the frequency and co-occurrence of key weather descriptors in Hudson's Bay Company records (Catchpole et al., 1970; Moodie and Catchpole, 1975). The resulting indices are open-ended, since more and stronger descriptors in the sources could generate indefinitely larger (positive or negative) values. Baron (1980) adapted content analysis to analyse historical New England diaries, by ranking and then numerically weighting descriptors of several types of weather found in those sources. In subsequent publications, he and collaborators adopted different scales for annual and seasonal temperature and precipitation depending on the level of detail in the underlying sources (e.g. Baron, 1995).

In Mexico, Mendoza et al. (2007) constructed a series of historical droughts for the Yucatan Peninsula using the method of Holmes and Lipo (2003). In this investigation, historical drought data were transformed into a series of pulse width modulation types (1 drought, 0 no drought) and linked to the Atlantic Multidecadal Oscillation and Southern Oscillation Index. Other studies have used key descriptors as the basis for index development. Garza Merodio (2017), for example, classified rain ceremonies into five ordinal levels based on Garza and Barriendos (1998), creating drought series for México, Puebla, Morelia, Guadalajara, Oaxaca, Durango, Sonora, Chiapas and Yucatán. Dominguez-Castro et al. (2019) generated binary series (presence or absence) for precipitation, frost, hail, fog, thunderstorm and wind in Mexico City. Temperature indices for Mexico have been developed using the applied content analysis approach of Baron (1982) and Prieto et al. (2005).

In South America, the methodology used to analyse historical sources for climate reconstruction initially followed Moodie and Catchpole's (1975) content analysis approach, but was later adapted in a number of papers by María del Rosario Prieto (e.g. Prieto et al., 2005). As noted in section 5, most historical rainfall and river flow index series use 5–7 annually- or seasonally-resolved classes based on key descriptors, while most temperature-related series use 3 classes. To date, all South American rainfall and temperature series are ordinal in nature and do not make background assumptions about the statistical distribution of climate-related phenomena. However, the method used to derive '0 index' values is not always clearly stated, and many series do not discriminate between 'no data' and 'normal' years (both of which are expressed as zero values). For example, in many studies that use rogation ceremonies as the basis for rainfall index development, months when there are no ceremonies are categorised as zero. There are exceptions, e.g. Dominguez-Castro et al. (2018), who explicitly identify an

absence of ceremonies as ‘no data’, and Prieto and Rojas (2015), who clearly differentiate between normal years and no data. A systematic reanalysis of many series would be useful to determine exactly how each was constructed. The approach used by Quinn et al. (1987) and Quinn and Neal (1992) to construct El Niño series over the past four and a half centuries is slightly different. The relative strengths of individual El Niño events were based on a range of subjective and objective measures in narrative sources from coastal Peru. These include descriptions of relative rainfall, the extent of flooding and the degree of physical damage and destruction associated with each event, alongside accounts of impacts on shipping (e.g. wind and current effects on travel times between ports), fisheries (e.g. changes to fish catches, changes in fish meal production), and marine life (e.g. mass mortality of endemic marine organisms and guano birds, extent of invasion by tropical nekton) (Quinn et al., 1987). This broad approach was continued in subsequent studies by Ortlieb (1994, 1995, 2000), García-Herrera et al. (2008) and others.

8.5. Climate index development in Australia

Australian efforts have largely been based on the Pfister approach (section 8.1) and regional-scale historical climatology investigations in southern Africa (section 8.3), although instrumental and documentary sources have been analysed separately. Fenby and Gergis (2013) and Gergis and Ashcroft (2013) converted documentary and instrumental data into a three-point scale of wet, normal and drought conditions. Historical data availability along with high spatial variability and known non-linearities in Australian rainfall meant that wet and dry conditions were assessed differently. Years were classified as ‘normal’ if they failed to reach either wet or dry criteria. To avoid introducing errors or biases in the record, years with missing data were marked as missing, rather than given a value of zero.

For droughts, agreement between a minimum of three of the twelve documentary sources used was required for drought conditions to be identified in a given month. Droughts were identified regionally in one of five modern southeastern Australian states. To avoid issues associated with exaggerated accounts of dry conditions and/or localised drought, a year was classified as a ‘drought year’ only when at least 40% of historical sources indicated dry conditions for at least six consecutive months during the May–April ‘ENSO’ year (the period with strongest association between south-eastern Australian rainfall variations and ENSO; Fenby and Gergis, 2013). Dry conditions were defined as times where a lack of rainfall was perceived as severe by society, or negatively impacted upon agriculture or water availability.

Months of above average rainfall in coastal New South Wales were identified using the annual rainfall summaries of Russell (1877), as this was the only source with consistent yearly information about rainfall events and impacts. Along with specific reports of good rainfall, monthly classifications of wet conditions were also based on accounts of flooding, abundant crops, excellent pasture and the occurrence of insect plagues (Fenby and Gergis, 2013). Six months of high rainfall were required for a year (May–April) to be defined as wet.

Combining the documentary-based indices with an instrumentally-derived index enabled the development of a single index of wet and dry conditions for eastern New South Wales from 1788 to 2008 CE. Each year of the instrumental rainfall datasets – the nine-station network for the Sydney region (1832–1860 CE) and a larger 45-station network representing the wider south-eastern Australian region – was assigned an index value of wet (1), normal (0) or dry (–1) based on normalised precipitation anomalies. Years with a normalised precipitation anomaly greater than the 70th percentile were counted as wet for that station, while those with an anomaly below the 30th percentile were counted as dry. Overall, a year was classified as wet or dry for the region if at least 40% of the stations with data available were in agreement, in line with the documentary classification of Fenby and Gergis (2013). Similar methods were employed by Ashcroft et al. (2014a) who used half a standard deviation above or below the 1835–1859 CE mean to build three-point indices of temperature, rainfall and pressure variability in southeastern Australia before 1860 CE using early instrumental data.

8.6. Climate index development in the oceans

The most common indices for marine climate reconstruction quantify shifts in prevailing wind direction. Most convert directional measurements from the 32-point system used by mariners in logbooks to one, four- or (very recently) eight-point indices. ~~This is, in – in part, because sailors were apparently-biased towards four, eight, and 16-point compass readings (Wheeler, 2004, 2005a). Scholars thereby create~~ These “directional indices” ~~that resemble the ordinal scales used to quantify qualitative temperature and rainfall observations on land. Influential publications also attempt to verify weather information in ships’ logbooks using other sources, including instrumental data (for more recent periods) or correspondence, intelligence reports, and diary entries (Degroot, 2018).~~ Few calculate error or confidence in their reconstructions, in part because those considerations are difficult to quantify (see García-Herrera et al., 2018). Recent studies have quantified the uncertainty involved in connecting logbook observations to broadscale circulation changes by using a calibration process that correlates wind directions in a target area traversed by ships to, for example, the strength of a monsoon (Gallego et al., 2015; Gallego et al., 2017).

Wind velocity and storm intensity or frequency indices, ~~meanwhile,~~ have ~~repeatedly also~~ made use of ~~wind measurements~~ observations recorded in logbooks. Beginning in the 19th century, mariners made these measurements using the 12-point Beaufort wind force scale. Before that, measurements ~~often~~ refer to the ~~state of a ship’s sails that mariners needed to furl or unfurl in winds of different velocity. The measurements are therefore more subjective than those of wind direction, yet they can still be roughly and can now be~~ translated into Beaufort indices (see García-Herrera et al., 2003; Koek and Konnen, 2005). It ~~has is been therefore~~ possible to use these indices to develop high-resolution reconstructions of trends in average wind velocities and storm frequency and intensity both to reconstruct annual wind velocities and storms at daily resolution, and to develop decadal reconstructions of wind force trends (Degroot, 2014). Yet because shifts in wind direction were more objectively recorded by sailors than changes in wind velocity, and are equally indicative of broadscale circulation changes, directional reconstructions are generally favoured by historical climatologists (Ordóñez et al., 2016).


9. Calibration, verification and dealing with uncertainty

9.1. Calibration and verification in index development

There are several approaches for calibrating and verifying index series used globally. Where overlapping meteorological data are available, long series of temperature and precipitation indices can be converted into quantitative meteorological units by using statistical climate reconstruction procedures; some of these have been inherited from fields such as dendroclimatology (see Brázdil et al., 2010, for a full discussion of statistical methods). For regions of the world lacking long instrumental records, simple cross-checking of climate indices against shorter periods of overlapping data is often used.

In Europe, Pfister (1984) was the first to use a calibration and verification process in the development of his indices. His approach – an example of best practice for regions where there is a lengthy period of instrumental overlap with the documentary record – is summarised by Brázdil et al. (2010) and Dobrovolný (2018) and illustrated in Figure 11. However, even where a period of overlap is lacking, indices from documentary sources can still be used to test reconstructions from proxy data (e.g. Bauch et al., 2020) or reconstruction, modelling results and observations (e.g. Bothe et al., 2019). The aim of calibration is to develop a transfer function between an index series and the measured climate variable, with verification against an independent period or subset of the overlapping meteorological data used to check the validity of this transfer function. In studies where there is a multi-decadal period of overlap, the instrumental data are normally divided into two subperiods; the index series is first calibrated to the earlier subperiod and then verified against the later subperiod (Dobrovolný, 2018). If only a short period of overlap is available, then cross-validation procedures are required.

932 The transfer function derived from a calibration period is normally evaluated by statistical measures (e.g. squared correlation
 933 r^2 , standard error of the estimate) before being applied in the verification period. During verification, index values are
 934 compared with the instrumental data and, again, evaluated statistically using r^2 , reduction of error and the coefficient of
 935 efficiency (see Cook et al., 1994; Wilson et al., 2006). If the calibrated data series, derived by applying the transfer function
 936 obtained for the calibration period, expresses the variability of the climate factor under consideration with satisfactory
 937 accuracy in the verification period, then the index series can be considered as useful for climate reconstruction back beyond
 938 the instrumental period (Brázdil et al., 2010). Caution is needed, however, as transfer functions, which are usually derived
 939 from relatively modern periods, may be non-stationary (e.g. where phenological series have been influenced by the
 940 introduction of new varieties or different harvesting technologies; Pfister, 1984; Meier et al., 2007).



941

942 **Figure 11:** The main steps in quantitative climate reconstruction based on temperature or precipitation indices derived from
 943 documentary evidence. Historical documentary sources are analysed to generate seven-point monthly indices (step 1), which
 944 are then summed to produce annual index series (step 2). Calibration and verification are carried out on periods of

945 overlapping instrumental data (step 3), with statistical results from verification used to define error bars for the final
946 reconstruction (step 4). Reprinted by permission from: Brázdil, R., Dobrovolný, P., Luterbacher, J., Moberg, A., Pfister, C.,
947 Wheeler, D., and Zorita, E.: European climate of the past 500 years: new challenges for historical climatology, *Climatic*
948 *Change*, 101, 7-40 (© Springer 2010).

949 Like the European approach, calibration and verification methods in China are applied to reconstructed temperature and
950 drought-flood indices by comparing the series overlap between instrumental and documentary periods. Shanghai has the
951 longest instrumental data coverage (1873 CE onwards), with Beijing, Suzhou, Nanjing, and Hangzhou also having century-
952 long data series (Chen and Shi, 2002; Zhang and Liu, 2002). As a result, most calibration is performed with reference to
953 these cities. Wang and Wang (1990a) compared their temperature series with these instrumental data to estimate correlation
954 coefficients and allocate corresponding values to their indices. A transfer function was also estimated between the number of
955 snow days (or number of lake freezing days) and observed temperatures by using multiple regression methods (Zhang, 1980;
956 Gong et al., 1983; Zhang and Liu, 1987; Wang and Gong, 2000; Ge et al., 2003). However, the statistical correlation reports
957 in these earlier studies appear incomplete.

958 The Academy of Meteorological Science of China Central Meteorological Administration (1981) have ~~also~~ used
959 ~~instrumental-precipitation data (1951-2000 CE) to calibrate-validate historical-drought-flood indices. However, the approach~~
960 ~~used focused on determining the probability distribution function of their five index classes to make the series comparable~~
961 ~~with instrumental data, rather than calibration per se (Yi et al., 2012; Shi et al., 2017), using chi-square tests, comparisons~~
962 ~~with the eigenvectors of instrumental precipitation series for 1951-2000 (Zhang and Liu, 1993; Zhang et al., 2003), and the~~
963 ~~standard error of the estimate to derive transfer functions (see Shi et al., 2017). These calculations assume that the~~
964 ~~distribution of the relative frequencies of the five classes is consistent (Yi et al., 2012).~~ A special feature of calibration and
965 verification in China is the utilisation of records in the *Qing Yu Lu* and *Yu Xue Fen Cun* (Hao et al., 2018; see section 3.2),
966 where comparisons can be made between reconstructed drought-flood indices and observed precipitation patterns (Zhang
967 and Wang, 1990). Such correlations can further be compared and calibrated using instrumental data, for example for Beijing
968 (Zhang and Liu, 2002), Suzhou, Nanjing and Hangzhou (Zhang and Wang, 1990).

969 Validation within the Nicholson et al. (2012a) rainfall reconstruction for continental Africa was carried out by comparing
970 time series based on those entries with instrumental rainfall data available for the same time and region. Quality control in
971 the final seven-class combined instrumental-historical reconstruction was provided by comparing the spread of estimates
972 from the various sources. If more than a two-class spread existed among the entries for an individual region and year, each of
973 those entries was re-evaluated. In most, it was found that an error was made in determining the location or year of a piece of
974 documentary evidence. Only eight “conflicts” in the Nicholson series could not be resolved in this way. The various regional
975 studies in southern Africa employ a simpler approach, using short periods of overlap with available instrumental data for
976 qualitative cross-checking/validation purposes (e.g. Nash and Endfield, 2002; Kelso and Vogel, 2007; Nash and Grab, 2010;
977 Nash et al., 2016).

978 The content analysis method developed for North American historical climatology uses replication by other researchers to
979 test the reliability of the quantification process and compared results from multiple independent sources to test validity
980 (Baron, 1980, pp.150-170). Subsequent studies have elaborated on this method, but many also draw on the Pfister index
981 approach as summarised in section 8.1. For South America, Neukom et al. (2009) created “pseudo-documentary” series to
982 quantify the relationship between document-derived precipitation indices and instrumental data (see also Mann and
983 Rutherford, 2002; Pauling et al., 2003; Xoplaki et al., 2005; Küttel et al., 2007). Following European conventions, index
984 series were transformed to instrumental units by linear regression with overlapping instrumental data. The skill measures
985 were quantified based on two calibration/verification intervals, using the first and second half of the overlap periods as

calibration and verification period, respectively and vice versa (Neukom et al., 2009). A similar approach has been used in southern Africa to integrate documentary-derived index series with other annually-resolved proxy data for the 19th century as part of multiproxy rainfall reconstructions (Neukom et al., 2014a; Nash et al., 2016).

Calibration and verification of indices in Australia (Figure 12) has been conducted using overlapping and largely independent instrumental data products, similar to approaches used in African reconstructions. In an example of good practice for future studies, independent high-resolution palaeoclimate reconstructions and records of water availability, such as lake levels, were also used for verification (Gergis and Ashcroft, 2013). Disagreements between these different sources were examined closely and often attributed to spatial variability in individual sources. For example, the 1820s in south-eastern Australia were identified as wetter than average in a regional palaeoclimate reconstruction (Gergis et al., 2012), but drier than average in a documentary-derived index and in historical information about water levels in Lake George, New South Wales (Gergis and Ashcroft, 2013). This was put down to geographical differences between the datasets – the palaeoclimate reconstruction was biased towards rainfall variability in southern parts of south-eastern Australia while the lake records and documentary index represented the east.




Figure 12: Wet and dry years for eastern New South Wales (Australia) identified using the nine-station network (1860–2008, purple) and a documentary index (1788–1860, grey). The median rainfall reconstruction (1788–1888) from Gergis et al. (2012) is also plotted as anomalies (mm) relative to a 1900–1988 base period. Note that 1841, 1844, 1846 and 1859 have been classified as wet, in accordance with a rainfall index derived from observations in the Sydney region (blue). Adapted from Gergis and Ashcroft (2013).

It is a long-standing best practice in marine historical climatology to verify weather observations by comparing different kinds of documentary evidence, or alternative different examples of the same evidence (e.g. multiple logbooks in the same fleet). Despite the very real challenges of interpreting measurements even in logbooks, there are indications that reconstructions that use these sources are reliable. There appears to be a high consistency and homogeneity both within wind measurements derived entirely from ships' logbooks, and between such measurements and data obtained from diverse sources that register the marine climate. Researchers have therefore linked documentary weather observations in, for example, the CLIWOC database, with datasets that homogenise and synthesise evidence from both textual and natural proxies, such as the National Oceanic and Atmospheric Administration's International Comprehensive Ocean-Atmosphere Data Set (ICOADS) (Jones and Salmon, 2005; Barriopedro et al., 2014).

9.2. Reporting confidence and uncertainty in index-based climate ~~series~~ reconstructions

Two forms of uncertainty are encountered when developing index-based climate series: (i) uncertainties related to the compilation of the index series themselves from documentary evidence; and (ii) uncertainties within any resulting index-based climate reconstruction. The first form of uncertainty relates mainly to the nature of information contained within specific source types. A detailed discussion is beyond the scope of this review. However, where indices are compiled from a unique documentary source – such as a private diary or diaries (e.g. Brázdil et al., 2008; Adamson, 2015; Domínguez-Castro et al., 2015), a series of correspondence (e.g. Rodrigo et al., 1998; Nash and Endfield, 2002; Fernández-Fernández et al., 2014) or a series of acts of municipal and ecclesiastical institutions for a location (e.g. Barriendos, 1997; Domínguez-Castro et al., 2018) – it is easier to identify and correct unexpected bias or homogeneity problems. Other index series draw together information from many different documentary sources (e.g. Camuffo et al., 2010; Nash and Grab, 2010; Fenby and Gergis, 2013; Brázdil et al., 2016), allowing the analysis of longer periods or larger regions but at the risk of incorporating homogeneities. Methodological differences – for example in the way in which ‘0 index’ values are derived (see section 8) – may also mask uncertainties introduced by data gaps. (Barriendos, 1997; Domínguez-Castro et al., 2018)

While compiling this review, it became apparent that relatively few index-based climate series investigations that adopt an index approach provide an assessment of the degree of uncertainty in the compilation of their index classification series/indices – in effect, something akin to the error bars used in quantitative climate reconstructions (e.g. Dobrovolný et al., 2010). Further, very few studies report directly on potential biases in their series due to the well-known tendency for documentary evidence to better record extreme events. The incorporation of statistical error is achievable where index-based series have been subject to full calibration and verification (section 9.1). However, it is less straightforward for climate reconstructions in regions (or for time periods) where a lack of overlapping instrumental data renders full calibration impossible.

To overcome this issue, Australian studies include some assessment of confidence by showing details of the number of sources in agreement, and the proportion of the study regions affected (see Fenby and Gergis, 2013). Independent high-resolution palaeoclimate and historical records were also used to verify each year of the reconstruction to assess confidence in the results (Fenby and Gergis, 2013; Gergis and Ashcroft, 2013).

One innovation from African historical climatology is the introduction by Clare Kelso and Coleen Vogel (2007) of a qualitative three-point ‘confidence rating’ (CR) for the classification of each rainy season in their climate history of Namaqualand (South Africa). The rating for each season (Figure 13) was derived from the number of sources consulted combined with the number of references to that particular climatological condition. CR=1 was awarded where there was only one source referring to the climatic condition. In contrast, years awarded CR=3 were those that had more than three date- and place-specific references describing climatic conditions. This approach has been adopted in subsequent studies in southern Africa by Nash et al. (2016), Nash et al. (2018) and Grab and Zumthurm (2018), with slight variations in the criteria used to award specific ratings according to source density.




Figure 13: Five-point index of rainfall variability in Namaqualand (South Africa) during the 1800s, including the first use of confidence ratings in relation to annual classifications in a documentary-derived index series (1 = low confidence, 3 = high confidence). Data from Kelso and Vogel (2007).

A similar approach was adopted by Quinn et al. (1987) and Quinn and Neal (1992) in their development of El Niño indices for Peru. El Niño events with a confidence rating of 1 were those that lacked a source reference or informational basis; these were not incorporated into the final list of reconstructed events. CR=2 was awarded when an event was based on limited or circumstantial evidence; CR=3, when additional information was needed to confirm the time of occurrence or intensity of an event; CR=4, when the occurrence time and intensity information was generally satisfactory, but additional evidence was needed to confirm the spatial extent of the event; and CR=5, when the available information concerning the occurrence and intensity of the event was considered to be satisfactory.

The second form of uncertainty relates specifically to index-based climate reconstruction. Where uncertainties can be quantified (either formally with statistics or less formally by comparison with other reconstructions), index-based reconstructions can be made fully comparable to natural proxy-based quantitative reconstructions. One example of this approach is the central Europe temperature series by Dobrovolný et al. (2010), the only documentary series used as part of the PAGES 2k Consortium (2013) continent-by-continent temperature reconstruction. Calibrated temperature series from China, including Zhang (1980) and Wang and Wang (1990a), are also incorporated into the PAGES 2k Consortium (2017) community-sourced database of temperature-sensitive proxy records.

10. Towards best practice in the use of climate indices for historical climate reconstruction

10.1. Regional variations in the development and application of climate indices

This review has shown that there are multiple approaches globally to the development and application of indices for historical climate reconstruction. Returning to the themes identified in the introduction, three categories of variability can be recognised. First, there is variability in the types of climate phenomena reconstructed in different regions (Table 5). Studies of the historical climatology of Europe and Asia span the greatest range of climate phenomena. This is partly a product of the range of climate zones present in these continents, and therefore the diversity of weather phenomena to which observers

might be exposed and document. However, it also reflects the relative abundance of documentary materials available for analysis and the richness of climate-related information they contain. Where smaller volumes of documentary evidence are available, reconstructions naturally tend to be skewed towards the climate parameters that were of sufficient importance to people that they captured them in writing or as artefacts – hence the emphasis on precipitation reconstructions for Africa and Australia and on winds and storm events over the oceans.

Table 5: Types of historical environmental phenomena reconstructed using an index approach in different parts of the world, with a qualitative indication of the relative emphasis of studies in each region (3 indicates a large number of studies, 1 a small number of studies, - indicates no studies).

Region	Temperature	Precipitation	Floods	Drought	Snow/ice	Wind/storms
Europe	***3	**3	**3	**2	*1	*1
Africa	*1	**2	-	*1	*1	*1
Americas	*1	*1	*1	-	-	*1
Asia	**2	**2	**2	*1	*1	*1
Australia	-	*1	*1	*1	-	-
Oceans	-	*1	-	-	-	**2

Second, there is variability in the way that historical evidence is treated to develop individual index series. Such variability arises, in part, from the extent to which analytical methods have developed independently. Thus, approaches to index-based climate reconstruction in parts of Asia are very different to those used in Europe. Chains of influence in practice can also be identified with, for example, elements of the ‘Pfister method’ from Europe being adopted by regional studies in southern Africa from the 1980s and then feeding into more recent precipitation reconstructions in Australia. There are common features of most historical treatments, regardless of tradition. These include the use of key descriptors or indicator criteria to match either individual observations (e.g. the continent-wide precipitation series for Africa developed by Nicholson) or sets of monthly, seasonal or annual observations (as per the Pfister method) to specific index classes. Most reconstructions are ordinal but, particularly where long runs of overlapping instrumental data are available, many are grounded in statistical distributions and present semi- or fully-quantified climate series.

The final source of variability across index-based investigations is in the number of index points used in individual reconstructions. A snapshot of this variability can be seen from investigations in Europe (Table 6). While most index-based reconstructions of European temperature and precipitation published since the 1990s employ the seven-point Pfister approach, some use up to nine classes. The number of classes used in European flood and drought reconstruction is usually smaller but, even here, may extend to seven-point classifications. There are also some commonalities. For example, most temperature and precipitation reconstructions use an odd number of classes – to allow the mid-point of the reconstruction to reflect ‘normal’ conditions – while open-ended unidirectional climate-related phenomena such as droughts and floods may be classified using either an even or odd number of classes. Similar patterns can be seen in other parts of the world (Table 7). In the rare instances where authors justify the number of index categories they use, most point to limitations in the quantity and/or richness of the historical evidence available for reconstruction as the reason for a smaller number of index categories.

Table 6: Variability in the number of index classes used in index-based historical climate reconstructions across Europe.

Climate phenomenon	Number of index classes used in climate reconstructions	Examples
--------------------	---	----------

Temperature	7-point most common (but also 2-, 3-, 5- and 9- point)	e.g. Pfister (1984), Alexandre (1987), Brázdil and Kotyza (1995, 2000), Van Engelen et al. (2001), Glaser (2013), Litzemberger (2015)
Precipitation	7-point most common (but also 3- and 5-point)	e.g. Alexandre (1987), Pfister (1992), Glaser et al. (1999), Van Engelen et al. (2001), Rodrigo and Barriendos (2008)
Floods	3-, 4- 5-point all common	e.g. Pfister (1999), Rohr (2006, 2013), Wetter et al. (2011), Brázdil et al. (2012), Garnier (2015), Kiss (2019)
Drought	3-point most common (but also 5- and 7-point)	e.g. Pfister et al. (2006), Brázdil et al. (2013b), Garnier (2018), Erfurt and Glaser (2019)

Table 7: Variability in the number of index classes used in index-based historical climate reconstructions in Africa, the Americas, Asia, Australia and over the oceans.

Region	Number of index classes used in climate reconstructions	Examples
Africa	3-point for temperature; 5- or 7-point for precipitation	e.g. Nicholson (2001), Nash and Endfield (2002), Kelso and Vogel (2007), Grab and Nash (2010), Nicholson et al. (2012a), Nash et al. (2016), Grab and Zumthum (2018)
Americas	3-point for temperature, 5- or 7-point for floods / precipitation; 3-point for snowfall	e.g. Baron et al. (1984), Prieto (1984), Baron (1989, 1995), Prieto et al. (1999), Prieto and Rojas (2015), Gil-Guirado et al. (2016)
Asia	4- or 5-point most common for temperature / precipitation and floods/drought	e.g. Zhu (1926), Zhang and Zhang (1979), Wang and Wang (1990a), Academy of Meteorological Science of China Central Meteorological Administration (1981), Wang and Wang (1990b), Wang et al. (1998), Tan and Wu (2013), Tan et al. (2014), Ge et al. (2018)
Australia	3-point for precipitation	e.g. Fenby and Gergis (2013), Gergis and Ashcroft (2013), Gergis et al. (2018)
Oceans	1-, 4- or 8-point for wind direction, 12-point for wind speed	e.g. Garcia et al. (2001), Prieto et al. (2005), Küttel et al. (2010), Barriopedro et al. (2014), Barrett et al. (2018), García-Herrera et al. (2018)

10.2. Guidelines for generating future documentary-based indices

The diversity of practice revealed in this review raises two issues. First, different approaches to index development make it harder for climate historians and historical climatologists working in different parts of the world to compare their climate indices directly, since each will include indices with differing climatological boundaries. Second, they make it harder for (palaeo)climatologists to use the resulting time series in synthesis and modelling studies without recourse to the methodology used in each original study. As noted in section 9.2, fully calibrated series have been included within global climate compilations such as the PAGES 2k Consortium (2013, 2017) [temperature syntheses](#). ~~These include, for example, the central Europe temperature series by Dobrovolný et al. (2010), the only documentary series used as part of the PAGES 2k Consortium (2013) continent by continent reconstruction. A number of more recent calibrated temperature reconstructions from Europe and China are now included in the PAGES 2k Consortium (2017) community sourced database of temperature sensitive proxy records.~~ Non-calibrated index series have also been incorporated into multi-proxy reconstructions using the “Pseudo proxy” approach of Mann and Rutherford (2002) – see, for example, Neukom et al. (2014a) and Neukom et al. (2014b) – but these reconstructions are relatively rare.

Having a standard approach to index-based climate reconstruction would clearly have its benefits. However, we recognise that a ‘one size fits all’ approach is not appropriate for all climate phenomena – the reconstruction of historical wind patterns

over the oceans from ships' logbooks, for example, already has well-developed methodologies and protocols. We further recognise that the most widely used approaches such as the Pfister method would require modification to be useful for temperature and/or rainfall reconstruction in all regions, since climates with strong seasonality may not have documentary evidence available year-round. ~~T-and~~ their use would, in some areas, also override the legacy of decades of methodological effort and require the reanalysis of enormous volumes of documentary evidence.

Rather than suggest a prescriptive method, we instead offer a series of guidelines as best practice for generating indices from collections of historical evidence. The guidelines are of greatest relevance to index-based reconstructions of temperature and precipitation but also have resonance for other climate phenomena (e.g. winter severity). The guidelines are based, in part, on the excellent reviews by Brázdil et al. (2010) and Pfister et al. (2018), but also incorporate insights from this study:

1. Researchers should be familiar with the climatology of their study region, as this may influence the temporal distribution of documentary evidence. Indices should, ideally, be based on collections of historical records that overlap with a climatically homogenous region with respect to the particular phenomena to be reconstructed.
2. Researchers should be familiar with the strengths and weaknesses of each of their historical sources prior to their use in climate reconstruction.
3. Researchers should select an appropriate temporal resolution for their index series according to the quantity, quality and richness (in terms of climate information) of available historical sources. This may be monthly, seasonal, annual or longer. ~~F, although~~ for information-rich areas, a monthly resolution is optimal as it offers the greatest potential for comparison with early instrumental series (which may be published as monthly averages prior to the wider availability of daily data) and the greatest flexibility for comparison with more coarsely-resolved sources, such as palaeoclimate reconstructions. For regions with marked variations in the quantity and quality of climate information across the year, the choice of resolution may be dictated by the length of period during the year when information is most sparse.
4. Whether to develop a three-, five- or seven- (or more) point index series ~~will also depend upon data quantity and quality but~~ may also be influenced by the legacy of previous studies in a region if direct comparisons are required; however, following guideline 3, researchers should only generate series with higher numbers of index classes if source density and richness permit.
5. Transforming the information in historical documents to numbers on a scale requires a high degree of expertise to minimise subjectivity and should, ideally, be undertaken by experienced researchers with a good knowledge of the climate of a region and an understanding of the language of the time period in which sources were written.
6. Historical records should ideally be sorted chronologically prior to analysis, with indices developed in a stepwise manner. Pfister et al. (2018, p.120) recommend that indexing begin with the most recent period (a process referred to by Brázdil et al., 2010, as 'hind-casting'), which for most studies will also be the period with the greatest volume of documentary evidence. This allows researchers to become familiar with the vagaries of their evidence during well-documented periods before working backwards to periods where information may be less complete.
7. For regions and periods where large volumes of historical information are available, indices should always be generated using evidence from more than one independent contemporary observer or record. Where reconstruction must rely on a single observer or record, or on secondary sources, appropriate levels of uncertainty should be noted in the final reconstruction (see ~~4-12~~).
- 7.8. It is advisable to sum-up index series – either in time (i.e. from monthly to seasonal or annual) or in space (i.e. by combining several index series from a climatologically homogeneous region). This approach may well approximate index series to natural climate variability.
- 8.9. Where possible, index series should be developed independently from the same set of historical sources by more than one researcher to minimise subjectivity. The final index series for southeast Africa produced by Nash et al.

(2016), for example, was first developed independently by two members of the research team who then met to agree the final series.

[9.10.](#) To maximise their wider usefulness, index series should, ideally, overlap with runs of local or regional instrumental data to permit calibration and verification. Where instrumental data are not available, overlaps with independent high-resolution palaeoclimate records may be used for calibration.

[10.11.](#) If fully calibrated, statistical measures of error should be incorporated into the presentation of any reconstruction.

[11.12.](#) Where insufficient overlapping instrumental data are available to permit full calibration and verification, some form of “Confidence Rating” (see section 9.2 and Kelso and Vogel, 2007) should be incorporated into the presentation of any reconstruction.

[12.13.](#) Finally, as Pfister et al. (2018, p.121) identify, the purpose and process of index development should be “fully transparent and open to critical evaluation”, with the method of index development described in detail and a source-critical evaluation of the underlying evidence included.

There remain vast collections of documentary evidence from all parts of the globe that have yet to be explored for information about past climate. We hope that, if such collections are scrutinised following these guidelines, they will lead to index-based reconstructions of climate variability that can be used to both extend climate records and contextualise studies of climate-society relationships to the wider benefit of humankind.

1181 **Dedication.** This paper is dedicated to the memory of María del Rosario Prieto, a pioneer in historical climatology and
 1182 active promoter of climate history studies in South America, who sadly passed away in 2020 during the preparation of the
 1183 first draft of this manuscript. Rest in peace, María.

1184 **Author contributions.** DJN, MB, CC and TL conceived the original study. Overall manuscript development was led by
 1185 DJN. All authors contributed to the writing of the first draft of the paper and to the preparation of the final manuscript.

1186 **Competing interest.** The authors declare that they have no conflict of interest.

1187 **Acknowledgements.** The authors would like to thank: PAGES (Past Global Changes) for supporting CRIAS Working
 1188 Group meetings in Bern (2018) and Leipzig (2019) that led to the conception and subsequent development of this
 1189 publication; the Leibniz Institute for the History and Culture of Eastern Europe (Leipzig, Germany), Oeschger Center for
 1190 Climate Change Research (Bern, Switzerland) and Education University of Hong Kong (Peoples Republic of China) for
 1191 supporting open access publication charges; and Lina Lerch (Leipzig) for help on Japanese climate-historical sources.

1192 **Financial support.** The meetings that underpinned this article were supported by PAGES (Past Global Changes). The article
 1193 processing charges for this open-access publication were covered by the Leibniz Institute for the History and Culture of
 1194 Eastern Europe, Oeschger Center for Climate Change Research and Education University of Hong Kong.

1195 References

- 1196 Academy of Meteorological Science of China Central Meteorological Administration: Yearly Charts of Dryness/Wetness in
 1197 China for the Last 500 Years, Cartographic Publishing House, Beijing, 1981.
- 1198 Adamson, G.C.D.: Private diaries as information sources in climate research, *Wiley Interdisciplinary Reviews: Climate*
 1199 *Change*, 6, 599-611, 2015.
- 1200 Adamson, G.C.D. and Nash, D.J.: Documentary reconstruction of monsoon rainfall variability over western India, 1781-
 1201 1860, *Climate Dynamics*, 42, 749-769, 2014.
- 1202 Adamson, G.C.D. and Nash, D.J.: Climate history of Asia (excluding China). In: *The Palgrave Handbook of Climate*
 1203 *History*, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 203-211, 2018.
- 1204 Alcoforado, M.J., Nunes, M.D., Garcia, J.C., and Taborda, J.P.: Temperature and precipitation reconstruction in southern
 1205 Portugal during the late Maunder Minimum (AD 1675-1715), *The Holocene*, 10, 333-340, 2000.
- 1206 Alexandre, P.: *Le climat en Europe au moyen âge: contribution à l'histoire des variations climatiques de 1000 à 1425,*
 1207 *d'après les narratives de l'Europe Occidentale, Recherches d'histoire et de sciences sociales 24*, Éditions de l'École des
 1208 hautes études en sciences sociales, Paris, 1987.
- 1209 Allan, R.J., Endfield, G.H., Damodaran, V., Adamson, G.C.D., Hannaford, M.J., Carroll, F., MacDonald, N., Groom, N.,
 1210 Jones, J., Williamson, F., Hendy, E., Holper, P., Arroyo-Mora, J.P., Hughes, L., Bickers, R., and Bliuc, A.M.: Toward
 1211 integrated historical climate research: the example of Atmospheric Circulation Reconstructions over the Earth, *Wiley*
 1212 *Interdisciplinary Reviews-Climate Change*, 7, 164-174, 2016.
- 1213 Álvarez Vázquez, J.A.: Drought and rainy periods in the Province of Zamora in the 17th, 18th, and 19th centuries. In:
 1214 *Quaternary Climate in Western Mediterranean*, López-Vera, F. (Ed.), Universidad Autonoma de Madrid, Madrid, 221-
 1215 233, 1986.
- 1216 Aono, Y. and Kazui, K.: Phenological data series of cherry tree flowering in Kyoto, Japan, and its application to
 1217 reconstruction of springtime temperatures since the 9th century, *International Journal of Climatology*, 28, 905-914, 2008.
- 1218 Aono, Y. and Saito, S.: Clarifying springtime temperature reconstructions of the medieval period by gap-filling the cherry
 1219 blossom phenological data series at Kyoto, Japan, *International Journal of Biometeorology*, 54, 211-219, 2010.
- 1220 Ashcroft, L., Gergis, J., and Karoly, D.J.: A historical climate dataset for southeastern Australia, 1788-1859, *Geoscience*
 1221 *Data Journal*, 1, 158-178, 2014a.
- 1222 Ashcroft, L., Karoly, D.J., and Gergis, J.: Southeastern Australian climate variability 1860-2009: a multivariate analysis,
 1223 *International Journal of Climatology*, 34, 1928-1944, 2014b.
- 1224 Ashcroft, L., Karoly, D.J., and Dowdy, A.J.: Historical extreme rainfall events in southeastern Australia, *Weather and*
 1225 *Climate Extremes*, 25, 100210, <https://doi.org/10.1016/j.wace.2019.100210>, 2019.
- 1226 Baron, W.R.: *Tempests, Freshets and Mackerel Skies; Climatological Data from Diaries Using Content Analysis*,
 1227 Unpublished PhD thesis, 1980. University of Maine at Orono, 1980.

- 1228 Baron, W.R.: The reconstruction of 18th-century temperature records through the use of content-analysis, *Climatic Change*,
1229 4, 384-398, 1982.
- 1230 Baron, W.R.: Retrieving American climate history - a bibliographic essay, *Agricultural History*, 63, 7-35, 1989.
- 1231 Baron, W.R.: Historical climate records from the northeastern United States, 1640 to 1900. In: *Climate since A.D. 1500*,
1232 Bradley, R.S. and Jones, P.D. (Eds.), Routledge, London, 74-91, 1995.
- 1233 Baron, W.R., Gordon, G.A., Borns, H.W., and Smith, D.C.: Frost-free record reconstruction for eastern Massachusetts,
1234 1733-1980, *Journal of Climate and Applied Meteorology*, 23, 317-319, 1984.
- 1235 Barrett, H.G.: El Niño Southern Oscillation from the pre-instrumental era: Development of logbook-based reconstructions;
1236 and evaluation of multi-proxy reconstructions and climate model simulations, Unpublished PhD thesis, 2017. University
1237 of Sheffield, 2017.
- 1238 Barrett, H.G., Jones, J.M., and Bigg, G.R.: Reconstructing El Nino Southern Oscillation using data from ships' logbooks,
1239 1815-1854. Part II: Comparisons with existing ENSO reconstructions and implications for reconstructing ENSO
1240 diversity, *Climate Dynamics*, 50, 3131-3152, 2018.
- 1241 Barriendos, M.: Climatic variations in the Iberian Peninsula during the late Maunder Minimum (AD 1675-1715): An
1242 analysis of data from rogation ceremonies, *The Holocene*, 7, 105-111, 1997.
- 1243 Barriendos, M.: Climate change in the Iberian Peninsula: Indicator of rogation ceremonies (16th-19th centuries), *Revue*
1244 *d'Histoire Moderne et Contemporaine*, 57, 131-159, 2010.
- 1245 Barriopedro, D., Gallego, D., Alvarez-Castro, M.C., Garcia-Herrera, R., Wheeler, D., Pena-Ortiz, C., and Barbosa, S.M.:
1246 Witnessing North Atlantic westerlies variability from ships' logbooks (1685-2008), *Climate Dynamics*, 43, 939-955,
1247 2014.
- 1248 Bauch, M., Labbé, T., Engel, A., and Seifert, P.: Prequel to the Dantean Anomaly: The Water Seesaw and droughts of 1302–
1249 1307 in Europe, *Climate of the Past*, 16, 2343-2358, 2020.
- 1250 Bogolepov, M.A.: О колебаниях климата Европейской России в историческую эпоху [Climate fluctuations in European
1251 Russia in the historical age], *Землеведение [Geology]*, Kushnerev and Co., Moscow, 1907.
- 1252 Bogolepov, M.A.: Колебания климата в Западной Европе с 1000 по 1500 г [Climate fluctuations in Western Europe from
1253 1000 to the year 1500], *Землеведение [Geology]*, Kushnerev and Co., Moscow, 1908.
- 1254 Bogolepov, M.A.: Колебания климата и истории [Climate fluctuations and history], Kushnerev and Co., Moscow, 1911.
- 1255 Bokwa, A., Limanówka, D., and Wibig, J.: Pre-instrumental weather observations in Poland in the 16th and 17th century. In:
1256 *History and Climate*, Jones, P.D. (Ed.), Springer, Boston, 9-27, 2001.
- 1257 Borisenkov, E.P. (Ed.): Колебания климата за последнее тысячелетие [Climate fluctuations over the past millennium],
1258 *Gidrometeoizdat*, Leningrad, 1988.
- 1259 Borisenkov, E.P. and Paseckij, V.M.: Экстремальные природные явления в русских летописях XI-XVII вв. [Extreme
1260 natural phenomena in the Russian chronicles of the 11-17 centuries], *Gidrometeoizdat*, Leningrad, 1983.
- 1261 Borisenkov, E.P. and Paseckij, V.M.: Тысячелетняя летопись необычайных явлений природы [Millennial chronicle of
1262 extraordinary natural phenomena], *Mysl'*, Moscow, 1988.
- 1263 Bothe, O., Wagner, S., and Zorita, E.: Inconsistencies between observed, reconstructed, and simulated precipitation indices
1264 for England since the year 1650 CE, *Climate of the Past*, 15, 307-334, 2019.
- 1265 Bozherianov, I.N.: Голодовки русского народа с 1024 по 1906 г [Starving of the Russian nations from 1024 to year 1906],
1266 *Gannibal*, Saint Petersburg, 1907.
- 1267 Bravo-Paredes, N., Gallego, M.C., Domínguez-Castro, F., García, J.A., and Vaquero, J.M.: Pro-pluvia rogation ceremonies
1268 in extremadura (Spain): Are they a good proxy of winter NAO?, *Atmosphere*, 11,
1269 <https://doi.org/10.3390/atmos11030282>, 2020.
- 1270 Brázdil, R. and Kotyza, O.: History of Weather and Climate in the Czech Lands I. Period 1000-1500, *Zürcher Geographische*
1271 *Schriften* 62, Zürich, 1995.
- 1272 Brázdil, R. and Kotyza, O.: History of Weather and Climate in the Czech Lands II. Utilisation of Economic Sources for the
1273 Study of Climate Fluctuation in the Louny Region in the Fifteenth-Seventeenth Centuries, Masaryk University, Brno,
1274 2000.
- 1275 Brázdil, R., Černušák, T., and Řezníčková, L.: Weather information in the diaries of the Premonstratensian Abbey at
1276 Hradisko, in the Czech Republic, 1693–1783, *Weather*, 63, 201-207, 2008.
- 1277 Brázdil, R., Pfister, C., Wanner, H., von Storch, H., and Luterbacher, J.: Historical climatology in Europe – the state of the
1278 art, *Climatic Change* 70, 363-430, 2005.

- Brázdil, R., Kotyza, O., Dobrovolný, P., Řezníčková, L., and Valášek, H.: Climate of the Sixteenth Century in the Czech Lands (History of Weather and Climate in the Czech Lands 10), Masaryk University, Brno 2013a.
- Brázdil, R., Kiss, A., Luterbacher, J., Nash, D.J., and Řezníčková, L.: Documentary data and the study of past droughts. A global state of the art, *Climate of the Past*, 14, 1915-1960, 2018.
- Brázdil, R., Kundzewicz, Z.W., Benito, G., Demarée, G., MacDonald, N., and Roald, L.A.: Historical floods in Europe in the past millennium. In: *Changes in Flood Risk in Europe*, Kundzewicz, Z.W. (Ed.), IAHS Press, Wallingford, 121-166, 2012.
- Brázdil, R., Dobrovolný, P., Luterbacher, J., Moberg, A., Pfister, C., Wheeler, D., and Zorita, E.: European climate of the past 500 years: new challenges for historical climatology, *Climatic Change*, 101, 7-40, 2010.
- Brázdil, R., Dobrovolný, P., Trnka, M., Kotyza, O., Řezníčková, L., Valášek, H., Zahradnický, P., and Štěpánek, P.: Droughts in the Czech Lands, 1090-2012 AD, *Climate of the Past*, 9, 1985-2002, 2013b.
- Brázdil, R., Dobrovolný, P., Trnka, M., Büntgen, U., Řezníčková, L., Kotyza, O., Valášek, H., and Štěpánek, P.: Documentary and instrumental-based drought indices for the Czech Lands back to AD 1501, *Climate Research*, 70, 103-117, 2016.
- Brázdil, R., Glaser, R., Pfister, C., Dobrovolný, P., Antoine, J.M., Barriendos, M., Camuffo, D., Deutsch, M., Enzi, S., Guidoboni, E., Kotyza, O., and Rodrigo, F.S.: Flood events of selected European rivers in the sixteenth century, *Climatic Change*, 43, 239-285, 1999.
- Brönnimann, S., Pfister, C., and White, S.: Archives of nature and archives of society. In: *The Palgrave Handbook of Climate History*, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 27-36, 2018.
- Brönnimann, S., Martius, O., Rohr, C., Bresch, D.N., and Lin, K.-H.E.: Historical weather data for climate risk assessment, *Annals of the New York Academy of Sciences*, 1436, 121-137, 2019.
- Brooks, C.E.P.: *Climate through the ages*, 2nd revised edition, Ernest Benn Ltd, London, 1949.
- Burchinskij, I.E.: О климате прошлого Русской равнины [On the climate of the past of the Russian Plain], *Gidrometeoizdat*, Leningrad, 1957.
- Callaghan, J. and Helman, P.: *Severe Storms on the East Coast of Australia, 1770-2008*, Griffith Centre for Coastal Management, Griffith University, Southport, 2008.
- Callaghan, J. and Power, S.B.: Variability and decline in the number of severe tropical cyclones making land-fall over eastern Australia since the late nineteenth century, *Climate Dynamics*, 37, 647-662, 2011.
- Callaghan, J. and Power, S.B.: Major coastal flooding in southeastern Australia 1860-2012, associated deaths and weather systems, *Australian Meteorological and Oceanographic Journal*, 64, 183-214, 2014.
- Camenisch, C.: Endless cold: a seasonal reconstruction of temperature and precipitation in the Burgundian Low Countries during the 15th century based on documentary evidence, *Climate of the Past*, 11, 1049-1066, 2015a.
- Camenisch, C.: Endlose Kälte: Witterungsverlauf und Getreidepreise in den Burgundischen Niederlanden im 15. Jahrhundert, *Wirtschafts-, Sozial- und Umweltgeschichte* 5, Schwabe, Basel, 2015b.
- Camenisch, C. and Salvisberg, M.: Droughts in Bern and in Rouen from the 14th to the beginning of the 18th century derived from documentary evidence, *Climate of the Past*, 16, 2173-2182, 2020.
- Camuffo, D., Bertolin, C., Barriendos, M., Dominguez-Castro, F., Cocheo, C., Enzi, S., Sghedoni, M., della Valle, A., Garnier, E., Alcoforado, M.J., Xoplaki, E., Luterbacher, J., Diodato, N., Maugeri, M., Nunes, M.F., and Rodriguez, R.: 500-year temperature reconstruction in the Mediterranean Basin by means of documentary data and instrumental observations, *Climatic Change*, 101, 169-199, 2010.
- Castorena, G., Sánchez Mora, E., Florescano, E., Padillo Ríos, G., and Rodríguez Viqueira, L.: *Análisis histórico de las sequías en México Documentación de la Comisión del Plan Nacional Hidráulico*, Secretaría de Agricultura y Recursos Hidráulicos (SARH), Comisión del Plan Nacional Hidráulico, Mexico, 1980.
- Catchpole, A.J.W.: Hudson's Bay Company ships' log-books as sources of sea ice data, 1751-1870. In: *Climate since A.D. 1500*, Bradley, R.S. and Jones, P.D. (Eds.), Routledge, London, 17-39, 1995.
- Catchpole, A.J.W. and Faurer, M.A.: Summer sea ice severity in Hudson Strait, 1751-1870, *Climatic Change*, 5, 115-139, 1983.
- Catchpole, A.J.W. and Halpin, J.: Measuring summer sea ice severity in Eastern Hudson Bay, 1751-1870, *Canadian Geographer-Geographe Canadien*, 31, 233-244, 1987.
- Catchpole, A.J.W. and Hanuta, I.: Severe summer sea ice in Hudson Strait and Hudson Bay following major volcanic eruptions, 1751-1889 A.D., *Climatic Change*, 14, 61-79, 1989.
- Catchpole, A.J.W., Moodie, D.W., and Kaye, B.: Content analysis: A method for the identification of dates of first freezing and first breaking from descriptive accounts, *Professional Geographer*, 22, 252-257, 1970.

1332 Central Meteorological Bureau of China: Atlas of Drought and Flood Distribution in China over the Last 500 Years, China
1333 Cartographic Publishing House, Beijing, 1981.

1334 Chen, H.F., Liu, Y.C., Chiang, C.W., Liu, X.Q., Chou, Y.M., and Pan, H.J.: China's historical record when searching for
1335 tropical cyclones corresponding to Intertropical Convergence Zone (ITCZ) shifts over the past 2 kyr, *Climate of the Past*,
1336 15, 279-289, 2019.

1337 Chen, J.Q. and Shi, Y.F.: The comparison between 1000-yr winter temperature change in the Yangtze river delta and ice
1338 core record of Guliya, *Journal of Glaciology and Geocryology*, 24, 32-39, 2002.

1339 Chenoweth, M.: A reassessment of historical Atlantic basin tropical cyclone activity, 1700-1855, *Climatic Change*, 76, 169-
1340 240, 2006.

1341 Chenoweth, M. and Divine, D.: A document-based 318-year record of tropical cyclones in the Lesser Antilles, 1690-2007,
1342 *Geochemistry Geophysics Geosystems*, 9, <https://doi.org/10.1029/2008GC002066>, 2008.

1343 Chinese Academy of Social Science: The history of natural disasters and agriculture in each dynasty of China, Agriculture
1344 Press, Beijing, 1988.

1345 Cook, E.R., Briffa, K.R., and Jones, P.D.: Spatial regression methods in dendroclimatology - a review and comparison of
1346 two techniques, *International Journal of Climatology*, 14, 379-402, 1994.

1347 de Kraker, A.M.J.: Reconstruction of storm frequency in the North Sea area of the pre-industrial period, 1400-1625 and the
1348 connection with reconstructed time series of temperatures, *History of Meteorology*, 2, 51-69, 2011.

1349 Degroot, D.: 'Never such weather known in these seas': Climatic fluctuations and the Anglo-Dutch wars of the seventeenth
1350 century, 1652-1674, *Environment and History*, 20, 239-273, 2014.

1351 Degroot, D.: Testing the limits of climate history: The quest for a northeast passage during the Little Ice Age, 1594-1597,
1352 *Journal of Interdisciplinary History*, 45, 459-484, 2015.

1353 Degroot, D.: The Frigid Golden Age: Climate Change, the Little Ice Age, and the Dutch Republic, 1560-1720, Cambridge
1354 University Press, New York, 2018.

1355 Degroot, D.: War of the whales: Climate change, weather and Arctic conflict in the early seventeenth century, *Environment*
1356 *and History*, 26, 549-577, 2020.

1357 Degroot, D. and Ottens, S.: Climatological Database of the World's Oceans, historicalclimatology.com,
1358 <https://www.historicalclimatology.com/cliwoc.html#>, last access: 18 June 2020.

1359 Dobrovolný, P.: Analysis and interpretation: Calibration-verification. In: The Palgrave Handbook of Climate History, White,
1360 S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 107-113, 2018.

1361 Dobrovolný, P., Brázdil, R., Trnka, M., Kotyza, O., and Valášek, H.: Precipitation reconstruction for the Czech Lands, AD
1362 1501–2010, *International Journal of Climatology*, 35, 1-14, 2015.

1363 Dobrovolný, P., Brázdil, R., Valášek, H., Kotyza, O., Macková, J., and Halíčková, M.: A standard paleoclimatological
1364 approach to temperature reconstruction in historical climatology: an example from the Czech Republic, AD 1718–2007,
1365 *International Journal of Climatology*, 29, 1478–1492, 2009.

1366 Dobrovolný, P., Moberg, A., Brázdil, R., Pfister, C., Glaser, R., Wilson, R., van Engelen, A., Limanówka, D., Kiss, A.,
1367 Halíčková, M., Macková, J., Riemann, D., Luterbacher, J., and Böhm, R.: Monthly, seasonal and annual temperature
1368 reconstructions for Central Europe derived from documentary evidence and instrumental records since AD 1500,
1369 *Climatic Change*, 101, 69-107, 2010.

1370 Dominguez-Castro, F., Garcia-Herrera, R., and Vicente-Serrano, S.M.: Wet and dry extremes in Quito (Ecuador) since the
1371 17th century, *International Journal of Climatology*, 38, 2006-2014, 2018.

1372 Dominguez-Castro, F., Gallego, M.C., Vaquero, J.M., Herrera, R.G., Pena-Gallardo, M., El Kenawy, A., and Vicente-
1373 Serrano, S.M.: Twelve years of daily weather descriptions in North America in the eighteenth century (Mexico City,
1374 1775-86), *Bulletin of the American Meteorological Society*, 100, 1531-1547, 2019.

1375 Domínguez-Castro, F., García-Herrera, R., and Vaquero, J.M.: An early weather diary from Iberia (Lisbon, 1631-1632),
1376 *Weather*, 70, 20-24, 2015.

1377 Domínguez-Castro, F., Santisteban, J.I., Barriendos, M., and Mediavilla, R.: Reconstruction of drought episodes for central
1378 Spain from rogation ceremonies recorded at the Toledo Cathedral from 1506 to 1900: A methodological approach,
1379 *Global and Planetary Change*, 63, 230–242, 2008.

1380 Domínguez-Castro, F., García-Herrera, R., Ribera, P., and Barriendos, M.: A shift in the spatial pattern of Iberian droughts
1381 during the 17th century, *Climate of the Past*, 6, 553–563, 2010.

1382 Domínguez-Castro, F., Vaquero, J.M., Marin, M., Cruz Gallego, M., and Garcia-Herrera, R.: How useful could Arabic
1383 documentary sources be for reconstructing past climate?, *Weather*, 67, 76-82, 2012a.

Domínguez-Castro, F., Ribera, P., García-Herrera, R., Vaquero, J.M., Barriendos, M., Cuadrat, J.M., and Moreno, J.M.: Assessing extreme droughts in Spain during 1750-1850 from rogation ceremonies, *Climate of the Past*, 8, 705-722, 2012b.

Easton, C.: *Les hivers dans l'Europe occidentale*, E. J. Brill, Leiden, 1928.

Endfield, G.H.: Climate and crisis in eighteenth century Mexico, *Medieval History Journal*, 10, 99-125, 2007.

Endfield, G.H. and Nash, D.J.: Drought, desiccation and discourse: missionary correspondence and nineteenth-century climate change in central southern Africa, *The Geographical Journal*, 168, 33-47, 2002.

Erfurt, M. and Glaser, R.: Changing impacts and societal responses to drought in southwestern Germany since 1800, *Regional Environmental Change*, 2019. <https://doi.org/10.1007/s10113-019-01522-7>, 2019.

Fang, X., Xiao, L., Ge, Q., and Zheng, J.: Changes of plants phenophases and temperature in spring during 1888-1916 around Changsha and Hengyang in Hunan Province, *Quaternary Sciences*, 25, 74-79, 2005.

Fei, J., Hu, H., Zhang, Z., and Zhou, J.: Research on dust weather in Beijing during 1860-1898: Inferred from the diary of Tonghe Weng, *Journal of Catastrophology (Zaihai Xue)*, 24, 116-122, 2009.

Fenby, C.D.: *Experiencing, understanding and adapting to climate in south-eastern Australia, 1788-1860*, Unpublished PhD Thesis, 2012. School of Earth Sciences and School of Historical and Philosophical Studies, University of Melbourne, Australia, 2012.

Fenby, C.D. and Gergis, J.: Rainfall variations in south-eastern Australia part 1: consolidating evidence from pre-instrumental documentary sources, 1788-1860, *International Journal of Climatology*, 33, 2956-2972, 2013.

Fernández-Fernández, M.I., Gallego, M.C., Domínguez-Castro, F., Trigo, R.M., and Vaquero, J.M.: The climate in Zafra from 1750 to 1840: precipitation, *Climatic Change*, 129, 267-280, 2015.

Fernández-Fernández, M.I., Gallego, M.C., Domínguez-Castro, F., Trigo, R.M., and Vaquero, J.M.: The climate in Zafra from 1750 to 1840: temperature indexes from documentary sources, *Climatic Change*, 141, 671-684, 2017.

Fernández-Fernández, M.I., Gallego, M.C., Domínguez-Castro, F., Trigo, R.M., García, J.A., Vaquero, J.M., Gonzalez, J.M.M., and Duran, J.C.: The climate in Zafra from 1750 to 1840: history and description of weather observations, *Climatic Change*, 126, 107-118, 2014.

Florescano, E.: *Precios del maíz y crisis agrícolas en México*, El Colegio de México, Mexico, 1969.

Foley, J.C.: Droughts in Australia: Review of records from earliest years of settlement to 1955, *Bulletin No. 43*, Bureau of Meteorology, Melbourne, 1957.

Fragoso, M., Carraça, M.D.G., and Alcoforado, M.J.: Droughts in Portugal in the 18th century: A study based on newly found documentary data, *International Journal of Climatology*, 38, 5522-5541, 2018.

Fujiki, H.: *Nihon chūsei saigaishi nenpyōkō* [Draft of a Chronological Timeline for the History of Japanese Medieval Catastrophes], Kōshi Shoin, Tokyo, 2007.

Gallego, D., García-Herrera, R., Peña-Ortiz, C., and Ribera, P.: The steady increase of the Australian Summer Monsoon in the last 200 years, *Scientific Reports*, 7, 16166, <https://doi.org/10.1038/s41598-017-16414-1>, 2017.

Gallego, D., Ordóñez, P., Ribera, P., Peña-Ortiz, C., and García-Herrera, R.: An instrumental index of the West African Monsoon back to the 19th century, *Quarterly Journal of the Royal Meteorological Society*, 141, 3166-3176, 2015.

García-Acosta, V., Pérez Zevallos, J.M., and Molina Del Villar, A.: *Desastres Agrícolas en México. Catálogo histórico, Tomo I: Épocas prehispánica y colonial (958-1822)*, Fondo de Cultura Económica (FCE), Centro de Investigaciones y Estudios Superiores en Antropología Social (CIESAS), Mexico, 2003.

García-Herrera, R. and Gallego, D.: Ship logbooks help to understand climate variability. In: *Advances in Shipping Data Analysis and Modeling*, Ducruet, C. (Ed.), Routledge, London, 37-51, 2017.

García-Herrera, R., Durán, F.R., Wheeler, D., Martín, E.H., Prieto, M.R., and Gimeno, L.: The use of Spanish and British documentary sources in the investigation of Atlantic hurricane incidence in historical times. In: *Hurricanes and Typhoons: Past, Present, and Future*, Murnane, R.J. and Liu, K.-B. (Eds.), Columbia University Press, New York, 149-176, 2004.

García-Herrera, R., Díaz, H.F., García, R.R., Prieto, M.R., Barriopedro, D., Moyano, R., and Hernández, E.: A chronology of El Niño events from primary documentary sources in Northern Peru, *Journal of Climate*, 21, 1948-1962, 2008.

García-Herrera, R., Prieto, L., Gallego, D., Hernández, E., Gimeno, L., Können, G., Koek, F.B., Wheeler, D., Wilkinson, C., Prieto, M.R., Báez, C., and Woodruff, S.: *CLIWOC Multilingual Meteorological Dictionary: An English-Spanish-Dutch-French dictionary of wind force terms used by mariners from 1750 to 1850*, Koninklijke Nederlands Meteorologisch Instituut, Den Haag, 2003.

1435 García-Herrera, R., Gimeno, L., Ribera, P., and Hernandez, E.: New records of Atlantic hurricanes from Spanish
1436 documentary sources, *Journal of Geophysical Research-Atmospheres*, 110, D03109,
1437 <https://doi.org/10.1029/2004JD005272>, 2005a.

1438 García-Herrera, R., Konnen, G.P., Wheeler, D.A., Prieto, M.R., Jones, P.D., and Koek, F.B.: CLIWOC: A climatological
1439 database for the world's oceans 1750-1854, *Climatic Change*, 73, 1-12, 2005b.

1440 García-Herrera, R., Können, G.P., Wheeler, D.A., Prieto, M.R., Jones, P.D., and Koek, F.B.: Ship logbooks help analyze
1441 pre-instrumental climate, *Eos, Transactions of the American Geophysical Union*, 87, 173-180, 2006.

1442 García-Herrera, R., Barriopedro, D., Gallego, D., Mellado-Cano, J., Wheeler, D., and Wilkinson, C.: Understanding weather
1443 and climate of the last 300 years from ships' logbooks, *Wiley Interdisciplinary Reviews-Climate Change*, 9, 2018.

1444 Garcia, R.R., Diaz, H.F., Herrera, R.G., Eischeid, J., Prieto, M.D., Hernandez, E., Gimeno, L., Duran, F.R., and Bascary,
1445 A.M.: Atmospheric circulation changes in the tropical Pacific inferred from the voyages of the Manila galleons in the
1446 sixteenth-eighteenth centuries, *Bulletin of the American Meteorological Society*, 82, 2435-2455, 2001.

1447 Garnier, E.: Le renversement des saisons. Climats et sociétés en France (vers 1500 – vers 1850), *Mémoire d'étude our*
1448 *l'obtention de l'Habilitation à diriger des recherches*, Université de Franche-Comté, 2009.

1449 Garnier, E.: Bassesses extraordinaires et grandes chaleurs. 500 ans de sécheresses et de chaleurs en France et dans les pays
1450 limitrophes, *Houille Blanche-Revue Internationale De L Eau*, 4, 26-42, 2010.

1451 Garnier, E.: At the risk of floodwaters: historical flood risk and its social impacts in the area of the Wash in eastern England
1452 (Cambridgeshire, Norfolk, Lincolnshire), mid 17th century – end of the 19th century, *Hydrology and Earth System*
1453 *Sciences Discussions*, 19, 1-33, 2015.

1454 Garnier, E.: Historic drought from archives. Beyond the instrumental record. In: *Drought. Science and Policy*, Iglesias, A.,
1455 Assimacopoulos, D., and Van Lanen, H.A.J. (Eds.), John Wiley & Sons, Hoboken, 45-67, 2018.

1456 Garza, G.M. and Barriendos, M.: El Clima en la historia, *Ciencias*, 51, 22-25, 1998.

1457 Garza Merodio, G.G.: Frecuencia y duración de sequías en la Cuenca de México de fines del siglo XVI a mediados del XIX,
1458 *Investigaciones Geográficas*, 2002, 106-115, 2002.

1459 Garza Merodio, G.G.: Variabilidad climática en México a través de fuentes documentales (siglos XVI al XIX), UNAM,
1460 Instituto de Geografía, Mexico City, 2017.

1461 Ge, Q.-S., Ding, L.-L., and Zheng, J.-Y.: Research on methods of starting date of pre-summer rainy season reconstruction in
1462 Fuzhou derived from Yu-Xue-Fen-Cun records, *Advances in Earth Science*, 26, 1200-1207, 2011.

1463 Ge, Q.-S., Hao, Z., Zheng, J., and Shao, X.: Temperature changes over the past 2000 yr in China and comparison with the
1464 Northern Hemisphere, *Climate of the Past*, 9, 1153-1160, 2013.

1465 Ge, Q.-S., Hao, Z.-X., Zheng, J.-Y., and Liu, Y.: China: 2000 years of climate reconstruction from historical documents. In:
1466 *The Palgrave Handbook of Climate History*, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan,
1467 London, 189-201, 2018.

1468 Ge, Q.-S., Zheng, J.Y., Hao, Z.X., Zhang, P.Y., and Wang, W.C.: Reconstruction of historical climate in China - High-
1469 resolution precipitation data from Qing dynasty archives, *Bulletin of the American Meteorological Society*, 86, 671-680,
1470 2005.

1471 Ge, Q.S., Zheng, J.Y., Fang, X.Q., Man, Z.M., Zhang, X.Q., Zhang, P.Y., and Wang, W.C.: Winter half-year temperature
1472 reconstruction for the middle and lower reaches of the Yellow River and Yangtze River, China, during the past 2000
1473 years, *The Holocene*, 13, 933-940, 2003.

1474 Gergis, J. and Ashcroft, L.: Rainfall variations in south-eastern Australia, Part 2: a comparison of documentary, early
1475 instrumental and palaeoclimate records, 1788-2008, *International Journal of Climatology* 33, 2973-2987, 2013.

1476 Gergis, J., Karoly, D.J., and Allan, R.J.: A climate reconstruction of Sydney Cove, New South Wales, using weather journal
1477 and documentary data, 1788-1791, *Australian Meteorological and Oceanographic Journal*, 58, 83-98, 2009.

1478 Gergis, J., Ashcroft, L., and Garden, D.: Recent developments in Australian climate history. In: *The Palgrave Handbook of*
1479 *Climate History*, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 237-245, 2018.

1480 Gergis, J., Ashcroft, L., and Whetton, P.: A historical perspective on Australian temperature extremes, *Climate Dynamics*,
1481 55, 843-868, 2020.

1482 Gergis, J., Gallant, A., Braganza, K., Karoly, D.J., Allen, K., Cullen, L., D'Arrigo, R.D., Goodwin, I., Grierson, P., and
1483 McGregor, S.: On the long-term context of the 1997-2009 "Big Dry" in south-eastern Australia: insights from a 206-year
1484 multi-proxy rainfall reconstruction, *Climatic Change*, 111, 923-944, 2012.

1485 Gil-Guirado, S., Espin-Sanchez, J.A., and Prieto, M.D.: Can we learn from the past? Four hundred years of changes in
1486 adaptation to floods and droughts. Measuring the vulnerability in two Hispanic cities, *Climatic Change*, 139, 183-200,
1487 2016.

1488 Gil-Guirado, S., Gómez-Navarro, J., and Pedro Montávez, J.: The weather behind words - new methodologies for integrated
1489 hydrometeorological reconstruction through documentary sources, *Climate of the Past*, 15, 1303–1325, 2019.

1490 Gioda, A. and Prieto, M.R.: Histoire des sécheresses andines. Potosí, El Niño et le Petit Âge Glaciaire. *La Météorologie*,
1491 *Revue de la Société Météorologique de France*, 8, 33-42, 1999.

1492 Gioda, A., Prieto, A.R., and Forenza, A.: Archival climate history survey in the Central Andes (Potosí, 16th -17th Centuries).
1493 In: *Prace Geograficzne, zeszyt 107*, Instytut Geografii UJ, Kraków, 107-112, 2000.

1494 Glaser, R.: *Klimageschichte Mitteleuropa. 1000 Jahre Wetter, Klima, Katastrophen*, Primus Verlag, Darmstadt, 2001.

1495 Glaser, R.: *Klimageschichte Mitteleuropas, 1200 Jahre Wetter, Klima, Katastrophen: Mit Prognosen für das 21. Jahrhundert*,
1496 3rd edition, Primus, Darmstadt, 2013.

1497 Glaser, R. and Stangl, H.: Historical floods in the Dutch Rhine Delta, *Natural Hazards and Earth System Sciences*, 3, 605-
1498 613, 2003.

1499 Glaser, R. and Stangl, H.: Climate and floods in Central Europe since AD 1000: Data, methods, results and consequences,
1500 *Surveys in Geophysics*, 25, 485-510, 2004.

1501 Glaser, R. and Riemann, D.: A thousand-year record of temperature variations for Germany and Central Europe based on
1502 documentary data, *Journal of Quaternary Science*, 24, 437-449, 2009.

1503 Glaser, R., Brazdil, R., Pfister, C., Dobrovolný, P., Vallve, M.B., Bokwa, A., Camuffo, D., Kotyza, O., Limanowka, D.,
1504 Racz, L., and Rodrigo, F.S.: Seasonal temperature and precipitation fluctuations in selected parts of Europe during the
1505 sixteenth century, *Climatic Change*, 43, 169-200, 1999.

1506 Glaser, R., Riemann, D., Schönbein, J., Barriendos, M., Brázdil, R., Bertolin, C., Camuffo, D., Deutsch, M., Dobrovolný, P.,
1507 van Engelen, A., Enzi, S., Halickova, M., Koenig, S.J., Kotyza, O., Limanowka, D., Mackova, J., Sghedoni, M., Martin,
1508 B., and Himmelsbach, I.: The variability of European floods since AD 1500, *Climatic Change*, 101, 235-256, 2010.

1509 Gong, G.F. and Hameed, S.: The variation of moisture conditions in China during the last 2000 years, *International Journal*
1510 *of Climatology*, 11, 271-283, 1991.

1511 Gong, G.F., Zhang, P.Y., and Zhang, J.Y.: A study on the climate of the 18th century of the Lower Changjiang Valley in
1512 China, *Geographic Research*, 2, 20-33, 1983.

1513 Grab, S.W. and Nash, D.J.: Documentary evidence of climate variability during cold seasons in Lesotho, southern Africa,
1514 1833-1900, *Climate Dynamics*, 34, 473-499, 2010.

1515 Grab, S.W. and Zumthurn, T.: The land and its climate knows no transition, no middle ground, everywhere too much or too
1516 little: a documentary-based climate chronology for central Namibia, 1845–1900, *International Journal of Climatology*, 38
1517 (Suppl. 1), e643-e659, 2018.

1518 Haldon, J., Roberts, N., Izdebski, A., Fleitmann, D., McCormick, M., Cassis, M., Doonan, O., Eastwood, W., Elton, H.,
1519 Ladstätter, S., Manning, S., Newhard, J., Nicoll, K., Telelis, I., and Xoplaki, E.: The climate and environment of
1520 Byzantine Anatolia: Integrating science, history, and archaeology, *Journal of Interdisciplinary History*, 45, 113-161,
1521 2014.

1522 Hannaford, M.J. and Nash, D.J.: *Climate, history, society over the last millennium in southeast Africa*, Wiley
1523 *Interdisciplinary Reviews-Climate Change*, 7, 370-392, 2016.

1524 Hannaford, M.J., Jones, J.M., and Bigg, G.R.: Early-nineteenth-century southern African precipitation reconstructions from
1525 ships' logbooks, *The Holocene*, 25, 379-390, 2015.

1526 Hansen, C.: Chinese language, Chinese philosophy, and "Truth", *Journal of Asian Studies* 44, 491-519, 1985.

1527 Hao, Z.X., Zheng, J.Y., Ge, Q.-S., and Wang, W.C.: Winter temperature variations over the middle and lower reaches of the
1528 Yangtze River since 1736 AD, *Climate of the Past*, 8, 1023-1030, 2012.

1529 Hao, Z.X., Yu, Y.Z., Ge, Q.-S., and Zheng, J.Y.: Reconstruction of high-resolution climate data over China from rainfall and
1530 snowfall records in the Qing Dynasty, *Wiley Interdisciplinary Reviews-Climate Change*, 9, 2018.

1531 Heckmann, M.-L.: Zwischen Weichseldelta, Großer Wildnis und Rigaischem Meerbusen. Ökologische Voraussetzungen für
1532 die Landnahme im spätmittelalterlichen Baltikum. In: *Von Nowgorod bis London. Studien zu Handel, Wirtschaft und*
1533 *Gesellschaft im mittelalterlichen Europa. Festschrift für Stuart Jenks zum 60. Geburtstag*, Heckmann, M.-L. and
1534 Röhrkasten, J. (Eds.), V&R Unipress, Göttingen, 255-295, 2008.

1535 Heckmann, M.-L.: Wetter und Krieg – im Spiegel erzählender Quellen zu Preußen und dem Baltikum aus dem 13. und 14.
1536 Jahrhundert. In: *Piśmienność pragmatyczna – Edytorstwo źródeł historycznych-archiwistyka. Studia ofiarowane*
1537 *Profesorowi Januszowi Tandeckiemu w sześćdziesiątą piątą rocznicę urodzin*, Czaia, R. and Kopiński, K. (Eds.), TNT,
1538 Toruń, 191-212, 2015.

1539 Hernández, M.E. and Garza Merodio, G.G.: Rainfall variability in Mexico's Southern Highlands (instrumental and
1540 documentary phases), 17th to 21st centuries. In: *Environmental quality in the large cities and industrial zones: problems*

- and management. Ecology and hydrometeorology of big cities and industrial zones (Russia-Mexico), Vol. I, Analysis of the environment, Karlin, N.L. and Shelutko, A.V. (Eds.), Russian State Hydrometeorology, University of St. Petersburg, 94-113, 2010.
- Herrera, R., Prieto, M.R., and Rojas, F.: Lluvias, sequías e inundaciones en el Chaco semiárido argentino entre 1580 y 1900, *Revista de la Junta de Estudios Históricos de Santa Fe*, LXIX, 173-200, 2011.
- Hirano, J. and Mikami, T.: Reconstruction of winter climate variations during the 19th century in Japan, *International Journal of Climatology*, 28, 1423-1434, 2008.
- Holmes, D.G. and Lipo, T.A.: Pulse width modulation for power converters: principles and practice, Wiley-IEEE Press, 2003.
- Hunt, H.A.: Results of rainfall observations made in Victoria during 1840-1910. Including all available annual rainfall totals from 1,114 stations; together with maps and diagrams, Bureau of Meteorology, Melbourne, 1911.
- Hunt, H.A.: Results of rainfall observations made in Queensland including all available annual rainfall totals from 1040 stations for all years of record up to 1913; together with maps and diagrams, Bureau of Meteorology, Melbourne, 1914.
- Hunt, H.A.: Results of rainfall observations made in South Australia and the Northern Territory, including all available annual rainfall totals from 829 stations for all years of record up to 1917, with maps and diagrams; also, appendices, presenting monthly and yearly meteorological elements for Adelaide and Darwin, Bureau of Meteorology, Melbourne, 1918.
- Ichino, M., Masuda, K., Kitamoto, A., Hirano, J., and Shō, K.: Experience of historical climatology as a material in Digital Humanities. In: Computers and the Humanities Symposium (December 2017), Information Processing Society of Japan, Tokyo, 139-146, 2017.
- IJnsen, F. and Schmidt, F.H.: Onderzoek naar het Optreden van Winterweer in Nederland, Scientific report, Royal Netherlands Meteorological Institute, de Bilt, 1974.
- Ingram, M.J., Farmer, G., and Wigley, T.M.L.: The use of documentary sources for the study of past climates. In: *Climate and History: Studies in Past Climates and their Impact on Man*, Wigley, T.M.L., Ingram, M.J., and Farmer, G. (Eds.), Cambridge University Press, Cambridge, 180-213, 1981.
- Itō, K.: Fujiki Hisashi nihon chūsei saigaishi nenpyōkō o riyō shita kikō hendō to saigai shiryō no kankei no kentō. Dai kikin no jiki o chūshin ni [Research on Historical Weather Sources Using Hisashi Fujiki's "Draft of a Chronological Timeline for the History of Medieval Japanese Catastrophes". Focussing on "Great Famine" Periods]. Kikō tekiōshi project. Kekka hōkokusho 1 [Historical Adaptation Project. Working Papers 1], 65-75, 2014.
- Jáuregui, E.: Algunos aspectos de las fluctuaciones pluviométricas en México en los últimos cien años, *Boletín del Instituto de Geografía*, 9, 39-64, 1979.
- Jevons, W.S.: Some data concerning the climate of Australia & New Zealand. In: *Waugh's Australian Almanac for the year 1859*, James William Waugh, Sydney, Australia, 47-98, 1859.
- Jones, P.D. and Salmon, M.: Preliminary reconstructions of the North Atlantic Oscillation and the Southern Oscillation Index from measures of wind strength and direction taken during the CLIWOC period, *Climatic Change*, 73, 131-154, 2005.
- Jusupović, A. and Bauch, M.: Surprising eastern perspectives: Historical climatology and Russian narrative sources, *PAGES News*, 28, <https://doi.org/10.22498/pages.28.2.16>, 2020.
- Kelso, C. and Vogel, C.H.: The climate of Namaqualand in the nineteenth century, *Climatic Change* 83, 257-380, 2007.
- Kiss, A.: *Floods and Long-Term Water-Level Changes in Medieval Hungary*, Springer, Cham, 2019.
- Klemm, F.: Witterungschronik des Barfüßerklosters Thann im Oberelsaß von 1182-1700, *Meteorologische Rundschau* 23/1, 15-18, 1970.
- Klimanov, V.A., Khotinskij, N.A., and Blagoveshchenskaia, N.V.: Колебания климата за исторический период в центре Русской равнины [Climate fluctuations over the historical period in the centre of the Russian Plain], *Известия Российской Академии Наук. Серия географическая* [Bulletin of the Russian Academy of Sciences: Geographic Series], 1, 89-96, 1995.
- Klimenko, V. and Solomina, O.: Climatic variations in the East European Plain during the last millennium: State of the art. In: *The Polish Climate in the European Context: An Historical Overview*, Przybylak, R., Majorowicz, J., Brázdil, R., and Kejan, M. (Eds.), Springer, Dordrecht, 71-101, 2010.
- Klimenko, V.V., Klimanov, V.A., Sirin, A.A., and Slepcev, A.M.: Изменения климата на западе европейской части России в позднем голоцене [Climate change in the west of European part of Russia in the late Holocene], *Доклады Российской Академии Наук* [Proceedings of the Russian Academy of Sciences], 376, 679-683, 2001.
- Koek, F.B. and Konnen, G.P.: Determination of wind force and present weather terms: The Dutch case, *Climatic Change*, 73, 79-95, 2005.

- 1595 Kong, W.S. and Watts, D.: A unique set of climatic data from Korea dating from 50 BC, and its vegetational implications,
1596 *Global Ecology and Biogeography Letters*, 2, 133-138, 1992.
- 1597 Küttel, M., Luterbacher, J., Zorita, E., Xoplaki, E., Riedwyl, N., and Wanner, H.: Testing a European winter surface
1598 temperature reconstruction in a surrogate climate, *Geophysical Research Letters*, 34,
1599 <https://doi.org/10.1029/2006GL027907>, 2007.
- 1600 Küttel, M., Xoplaki, E., Gallego, D., Luterbacher, J., Garcia-Herrera, R., Allan, R., Barriendos, M., Jones, P., Wheeler, D.,
1601 and Wanner, H.: The importance of ship log data: reconstructing North Atlantic, European and Mediterranean sea level
1602 pressure fields back to 1750, *Climate Dynamics*, 34, 1115-1128, 2010.
- 1603 Lamb, H.H.: *Climate. Past, Present and Future*, vol. 2, Methuen, London, 1977.
- 1604 Lamb, H.H.: *Historic Storms of the North Sea, British Isles and Northwest Europe*, Cambridge University Press, Cambridge,
1605 1992.
- 1606 Leontovich, F.I.: Голодовки в России до конца прошлого века [Famine in Russia until the end of the last century],
1607 *Северный Вестник [Northern Herald]*, March, 2–35, 1892.
- 1608 Liakhov, M.E.: Климатические экстремумы в центральной части Европейской территории СССР в XIII–XX вв.
1609 [Climatic extremes in the central part of the European territory of the USSR in the 13th-20th centuries], *Известия*
1610 *Академии Наук СССР: Серия географическая [Bulletin of the Academy of Sciences of the USSR: Geographic Series]*,
1611 6, 68-74, 1984.
- 1612 Lin, K.-H.E., Wang, P.K., Pai, P.L., Lin, Y.S., and Wang, C.W.: Historical droughts in the Qing dynasty (1644-1911) of
1613 China, *Climate of the Past*, 16, 911-931, 2020.
- 1614 Lin, K.-H.E., Hsu, C.T., Wang, P.K., Hsu, S.M., Lin, Y.S., Wan, C.W., Tseng, W.L., Wu, W.C., and Pan, W.:
1615 Reconstructing historical typhoon series and spatiotemporal characteristics from REACHES documentary records,
1616 *Journal of Geographical Sciences*, 93, 81-107, 2019.
- 1617 Litzenburger, L.: *Une ville face au climat. Metz à la fin du Moyen Age (1400–1530)*, Presses Universitaires de Nancy,
1618 Nancy, 2015.
- 1619 Liu, B.: Phenological change in Yangtze Plain during late Ming Dynasty (1450-1649), *Historical Geography*, 35, 22-33,
1620 2017.
- 1621 Liu, K.B., Shen, C.M., and Louie, K.S.: A 1,000-year history of typhoon landfalls in Guangdong, southern China,
1622 reconstructed from Chinese historical documentary records, *Annals of the Association of American Geographers*, 91,
1623 453-464, 2001.
- 1624 Liu, Y., Wang, H., Dai, J., Li, T.S., Wang, H., and Tao, Z.: The application of phenological methods for reconstructing past
1625 climate change, *Geographical Research*, 33, 603-613, 2014.
- 1626 Man, Z.M.: Some fundamentals in research on changes of warm and cold climate making use of historical records, *Historical*
1627 *Geography*, 12, 21-31, 1995.
- 1628 Mann, M.E. and Rutherford, S.: Climate reconstruction using “Pseudoproxies”, *Geophysical Research Letters*, 29,
1629 <https://doi.org/10.1029/2001GL014554>, 2002.
- 1630 Martín-Vide, J. and Vallvé, M.B.: The use of rogation ceremony records in climatic reconstruction: a case study from
1631 Catalonia (Spain), *Climatic Change*, 30, 201-221, 1995.
- 1632 Mauelshagen, F.: *Klimageschichte der Neuzeit 1500-1900 (Geschichte kompakt)*, Wissenschaftliche Buchgesellschaft,
1633 Darmstadt, 2010.
- 1634 McAfee, R.J.: *The fires of summer and the floods of winter: towards a climatic history for southeastern Australia, 1788–*
1635 *1860*, Macquarie University Library, Sydney, Australia, 1981.
- 1636 Meier, N., Rutishauser, T., Pfister, C., Wanner, H., and Luterbacher, J.: Grape harvest dates as a proxy for Swiss April to
1637 August temperature reconstructions back to AD 1480, *Geophysical Research Letters*, 34, L20705,
1638 <https://doi.org/10.1029/2007GL031381>, 2007.
- 1639 Mendoza, B., Garcia-Acosta, V., Velasco, V., Jauregui, E., and Diaz-Sandoval, R.: Frequency and duration of historical
1640 droughts from the 16th to the 19th centuries in the Mexican Maya lands, Yucatan Peninsula, *Climatic Change*, 83, 151-
1641 168, 2007.
- 1642 Mendoza, B., Jauregui, E., Diaz-Sandoval, R., Garcia-Acosta, V., Velasco, V., and Cordero, G.: Historical droughts in
1643 central Mexico and their relation with El Nino, *Journal of Applied Meteorology*, 44, 709-716, 2005.
- 1644 Metcalfe, S.E.: Historical data and climatic change in Mexico - a review, *Geographical Journal*, 153, 211-222, 1987.
- 1645 Mikami, T.: Climatic variations in Japan reconstructed from historical documents, *Weather*, 63, 190-193, 2008.
- 1646 Mizukoshi, M.: Climatic reconstruction in central Japan during the Little Ice Age based on documentary sources, *Journal of*
1647 *Geography (Chigaku Zasshi)*, 102, 152-166, 1993.

1648 Mizukoshi, M.: Kokiroku ni yoru jūroku/ jūgo/ jūyon/ jūsan/ jūni/ jūichi seiki no kikōkiroku [Weather Documentation in
1649 Historical Sources of the 16th/15th/14th/13th/12th/11th Century], 6 volumes, Tōkyōdō Shuppan, Tokyo, 2004-2014.

1650 Moodie, D.W. and Catchpole, A.J.W.: Environmental Data from Historical Documents by Content Analysis: Freeze-up and
1651 Break-up of Estuaries on Hudson Bay, 1714-1871, Department of Geography, University of Manitoba, 1975.

1652 Mora Pacheco, K.: Conmociones bajo un "cielo conspirador". Sequías en el Altiplano Cundiboyacense, 1778-1828. In: VII
1653 Simposio de Historia Regional y Local, Universidad Industrial de Santander, Colombia, 2018.

1654 Mutua, T.M. and Runguma, S.N.: Documentary driven chronologies of rainfall variability for Kenya, 1845-1976, Journal of
1655 Climatology and Weather Forecasting, 8, <https://doi.org/10.35248/2332-2594.2020.8.255>, 2020.

1656 Nash, D.J.: Changes in precipitation over southern Africa during recent centuries. In: Oxford Research Encyclopedia of
1657 Climate Science, Oxford Research Encyclopedias, 2017.

1658 Nash, D.J. and Endfield, G.H.: A 19th century climate chronology for the Kalahari region of central southern Africa derived
1659 from missionary correspondence, International Journal of Climatology, 22, 821-841, 2002.

1660 Nash, D.J. and Endfield, G.H.: 'Splendid rains have fallen': links between El Nino and rainfall variability in the Kalahari,
1661 1840-1900, Climatic Change, 86, 257-290, 2008.

1662 Nash, D.J. and Grab, S.W.: "A sky of brass and burning winds": documentary evidence of rainfall variability in the Kingdom
1663 of Lesotho, Southern Africa, 1824-1900, Climatic Change, 101, 617-653, 2010.

1664 Nash, D.J. and Hannaford, M.J.: Historical climatology in Africa: A state of the art, PAGES News, 28, 42-43, 2020.

1665 Nash, D.J., Pribyl, K., Endfield, G.H., Klein, J., and Adamson, G.C.D.: Rainfall variability over Malawi during the late 19th
1666 century, International Journal of Climatology, 38 (Suppl. 1), e629-e642, 2018.

1667 Nash, D.J., Pribyl, K., Klein, J., Neukom, R., Endfield, G.H., Adamson, G.C.D., and Kniveton, D.R.: Seasonal rainfall
1668 variability in southeast Africa during the nineteenth century reconstructed from documentary sources, Climatic Change,
1669 134, 605-619, 2016.

1670 Neukom, R., Prieto, M.D., Moyano, R., Luterbacher, J., Pfister, C., Villalba, R., Jones, P.D., and Wanner, H.: An extended
1671 network of documentary data from South America and its potential for quantitative precipitation reconstructions back to
1672 the 16th century, Geophysical Research Letters, 36, 2009.

1673 Neukom, R., Nash, D.J., Endfield, G.H., Grab, S.W., Grove, C.A., Kelso, C., Vogel, C.H., and Zinke, J.: Multi-proxy
1674 summer and winter precipitation reconstruction for southern Africa over the last 200 years, Climate Dynamics, 42, 2713-
1675 2716, 2014a.

1676 Neukom, R., Gergis, J., Karoly, D.J., Wanner, H., Curran, M., Elbert, J., Gonzalez-Rouco, F., Linsley, B.K., Moy, A.D.,
1677 Mundo, I., Raible, C.C., Steig, E.J., van Ommen, T., Vance, T., Villalba, R., Zinke, J., and Frank, D.: Inter-hemispheric
1678 temperature variability over the past millennium, Nature Climate Change, 4, 362-367, 2014b.

1679 Nicholls, N.: More on early ENSOs - evidence from Australian documentary sources, Bulletin of the American
1680 Meteorological Society, 69, 4-6, 1988.

1681 Nicholson, S.E.: Climatic variations in the Sahel and other African regions during the past five centuries, Journal of Arid
1682 Environments, 1, 3-24, 1978a.

1683 Nicholson, S.E.: Comparison of historical and recent African rainfall anomalies with late Pleistocene and early Holocene,
1684 Palaeoecology of Africa 10, 99-123, 1978b.

1685 Nicholson, S.E.: The methodology of historical climate reconstruction and its application to Africa, Journal of African
1686 History, 20, 31-49, 1979.

1687 Nicholson, S.E.: Saharan climates in historic times. In: The Sahara and the Nile, Williams, M.A.J., Faure, H. (Ed.), Balkema,
1688 Rotterdam, 173-200, 1980.

1689 Nicholson, S.E.: The historical climatology of Africa. In: Climate and History, Wigley, T.M.L., Ingram, M.J., and Farmer,
1690 G. (Eds.), Cambridge University Press, Cambridge, 249-270, 1981.

1691 Nicholson, S.E.: Environmental change within the historical period. In: The Physical Geography of Africa, Goudie, A.S.,
1692 Adams, W. M., and Orme, A. (Ed.), Oxford University Press, Oxford, 60-75, 1996.

1693 Nicholson, S.E.: A semi-quantitative, regional precipitation data set for studying African climates of the nineteenth century,
1694 part 1. Overview of the data set, Climatic Change, 50, 317-353, 2001.

1695 Nicholson, S.E.: A multi-century history of drought and wetter conditions in Africa. In: The Palgrave Handbook of Climate
1696 History, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 225-236, 2018.

1697 Nicholson, S.E., Klotter, D., and Dezfuli, A.K.: Spatial reconstruction of semi-quantitative precipitation fields over Africa
1698 during the nineteenth century from documentary evidence and gauge data, Quaternary Research 78, 13-23, 2012a.

1699 Nicholson, S.E., Dezfuli, A.K., and Klotter, D.: A two-century precipitation dataset for the continent of Africa, *Bulletin of*
1700 *the American Meteorological Society*, 93, 1219-1231, 2012b.

1701 Nicholson, S.E., Funk, C., and Fink, A.: Rainfall over the African continent from the 19th through the 21st century, *Global*
1702 *and Planetary Change*, 165, 114-127, 2018.

1703 Norrgård, S.: Practising historical climatology in West Africa: a climatic periodisation 1750-1800, *Climatic Change*, 129,
1704 131-143, 2015.

1705 Norrgård, S.: Royal Navy logbooks as secondary sources and their use in climatic investigations: introducing the log-board,
1706 *International Journal of Climatology*, 37, 2027-2036, 2017.

1707 Ogilvie, A.E.J.: The past climate and sea-ice record from Iceland. Part 1: Data to A.D. 1780, *Climatic Change*, 6, 131-152,
1708 1984.

1709 Ogilvie, A.E.J.: Documentary evidence for changes in the climate of Iceland, A.D. 1500-1800. In: *Climate since A.D. 1500*,
1710 Bradley, R.S. and Jones, P.D. (Eds.), Routledge, London, 92-117, 1992.

1711 Ogilvie, A.E.J.: Sea-ice conditions off the coasts of Iceland A.D. 1601-1850 with special reference to part of the Maunder
1712 Minimum period (1675-1715), *AmS-Varia* 25, 9-12, 1996.

1713 Ogilvie, A.E.J. and Farmer, G.: Documenting the Medieval climate. In: *Climates of the British Isles. Present, Past and*
1714 *Future*, Hulme, M. and Barrow, E. (Eds.), Routledge, London, 1997.

1715 Ogilvie, A.E.J. and Jónsson, T.: "Little Ice Age" research: A perspective from Iceland, *Climatic Change*, 48, 9-52, 2001.

1716 Oppokov, E.V.: Колебания водоносности рек в историческое время [Fluctuations in river flow in historical time]. In:
1717 *Исследования рек СССР [Research on Rivers of the USSR]*, vol. 4, State Institute of Hydrology, Leningrad, 1933.

1718 Ordóñez, P., Gallego, D., Ribera, P., Peña-Ortiz, C., and García-Herrera, R.: Tracking the Indian Summer Monsoon onset
1719 back to the pre-instrumental period, *Journal of Climate*, 29, 8115-8127, 2016.

1720 Ortlieb, L.: Las mayores precipitaciones históricas en Chile central y la cronología de eventos ENOS en los siglos XVI-XIX,
1721 *Revista Chilena de Historia Natural*, 67, 463-485, 1994.

1722 Ortlieb, L.: Eventos El Niño y episodios lluviosos en el desierto de Atacama: el registro de los dos últimos siglos, *Bulletin de*
1723 *l'Institut Français d'Études Andines*, 24, 519-537, 1995.

1724 Ortlieb, L.: The documentary historical record of El Niño events in Peru: An update of the Quinn record. In: *El Niño and the*
1725 *Southern Oscillation: Multiscale Variability and Global and Regional Impacts*, Diaz, H.F. and Markgraf, V. (Eds.),
1726 Cambridge University Press, Cambridge 207-297, 2000.

1727 PAGES 2k Consortium: Continental-scale temperature variability during the past two millennia, *Nature Geoscience*, 6, 339-
1728 346, 2013.

1729 PAGES 2k Consortium: A global multiproxy database for temperature reconstructions of the Common Era, *Scientific Data*,
1730 4, 170088, <https://doi.org/10.1038/sdata.2017.88>, 2017.

1731 Pauling, A., Luterbacher, J., and Wanner, H.: Evaluation of proxies for European and North Atlantic temperature field
1732 reconstructions, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2003GL017589>, 2003.

1733 Pei, Q. and Forêt, P.: Introduction to the climate records of Imperial China, *Environmental History*, 23, 863-871, 2018.

1734 Perry, E.J.: Challenging the Mandate of Heaven - Popular protest in modern China, *Critical Asian Studies*, 33, 163-180,
1735 2001.

1736 Pfister, C.: *Klimageschichte der Schweiz 1525-1860. Das Klima der Schweiz und seine Bedeutung in der Geschichte von*
1737 *Bevölkerung und Landwirtschaft*, Paul Haupt, Bern, 1984.

1738 Pfister, C.: Monthly temperature and precipitation patterns in Central Europe from 1525 to the present. A methodology for
1739 quantifying man-made evidence on weather and climate. In: *Climate Since A.D. 1500*, Bradley, R.S. and Jones, P.D.
1740 (Eds.), Routledge, London, 118-142, 1992.

1741 Pfister, C.: Raum-zeitliche Rekonstruktion von Witterungsanomalien und Naturkatastrophen 1496-1995. In cooperation with
1742 Daniel Brändli. Schlussbericht zum Projekt 4031-33198 des NFP 31, vdf Hochschulverlag AG and ETH Zürich, Zurich,
1743 1998.

1744 Pfister, C.: *Wetternachhersage. 500 Jahre Klimavariationen und Naturkatastrophen (1496-1995)*, Paul Haupt, Bern, Stuttgart,
1745 Wien, 1999.

1746 Pfister, C.: Evidence from the archives of societies: Documentary evidence - overview. In: *The Palgrave Handbook of*
1747 *Climate History*, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 37-47, 2018.

1748 Pfister, C. and Hächler, S.: Überschwemmungskatastrophen im Schweizer Alpenraum seit dem Spätmittelalter.
1749 Raumzeitliche Rekonstruktion von Schadensmustern auf der Basis historischer Quellen. In: *Historical Climatology in*

1750 Different Climatic Zones. Würzburger Geographische Arbeiten 80, Glaser, R. and Walsh, R.P.D. (Eds.), Institut für
1751 Geographie/Geographische Gesellschaft, Würzburg, 127-148, 1991.

1752 Pfister, C., Weingartner, R., and Luterbacher, J.: Hydrological winter droughts over the last 450 years in the Upper Rhine
1753 basin: a methodological approach, *Hydrological Sciences Journal-Journal Des Sciences Hydrologiques*, 51, 966-985,
1754 2006.

1755 Pfister, C., Camenisch, C., and Dobrovolný, P.: Analysis and Interpretation: Temperature and Precipitation Indices. In: *The*
1756 *Palgrave Handbook of Climate History*, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave-Macmillan, London,
1757 115-129, 2018.

1758 Pichard, G. and Roucaute, E.: Une déclinaison régionale du Petit Âge Glaciaire. Apport des archives historiques en
1759 Provence, *Archéologie du Midi Medieval* 27, 237-247, 2009.

1760 Piervitali, E. and Colacino, M.: Evidence of drought in western Sicily during the period 1565-1915 from liturgical offices,
1761 *Climatic Change*, 49, 225-238, 2001.

1762 Power, S.B. and Callaghan, J.: The frequency of major flooding in coastal southeast Australia has significantly increased
1763 since the late 19th century, *Journal of Southern Hemisphere Earth Systems Science*, 66, 2-11, 2016.

1764 Prieto, M.R.: El clima de Mendoza durante los siglos XVII y XVIII, *Meteorológica*, XIV, 165-174, 1983.

1765 Prieto, M.R.: Métodos para derivar información sobre precipitaciones nivales de fuentes históricas en la Cordillera de los
1766 Andes, *Zentralblatt für Geologie und Palaontologie*, 11/12, 1984.

1767 Prieto, M.R. and García-Herrera, R.: Documentary sources from South America: Potential for climate reconstruction,
1768 *Palaeogeography Palaeoclimatology Palaeoecology*, 281, 196-209, 2009.

1769 Prieto, M.R. and Rojas, F.: Documentary evidence for changing climatic and anthropogenic influences on the Bermejo
1770 Wetland in Mendoza, Argentina, during the 16th-20th century, *Climate of the Past*, 8, 951-961, 2012.

1771 Prieto, M.R. and Rojas, F.: Determination of droughts and high floods of the Bermejo River (Argentina) based on
1772 documentary evidence (17th to 20th century), *Journal of Hydrology*, 529, 676-683, 2015.

1773 Prieto, M.R. and Rojas, F.: Climate history in Latin America. In: *The Palgrave Handbook of Climate History*, White, S.,
1774 Pfister, C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 213-224, 2018.

1775 Prieto, M.R., Herrera, R., and Dussel, P.: Historical evidences of streamflow fluctuations in the Mendoza River, Argentina,
1776 and their relationship with ENSO, *The Holocene*, 9, 473-481, 1999.

1777 Prieto, M.R., Garcia-Herrera, R., and Hernández, E.: Early records of icebergs in the South Atlantic Ocean from Spanish
1778 documentary sources, *Climatic Change*, 66, 29-48, 2004.

1779 Prieto, M.R., Rojas, F., and Castillo, L.: La climatología histórica en Latinoamérica. Desafíos y perspectivas, *Bulletin de*
1780 *l'Institut français d'études andines*, 47, 141-167, 2019.

1781 Prieto, M.R., Gallego, D., Garcia-Herrera, R., and Calvo, N.: Deriving wind force terms from nautical reports through
1782 content analysis. The Spanish and French cases, *Climatic Change*, 73, 37-55, 2005.

1783 Quinn, W.H. and Neal, V.T.: The historical record of El Niño events. In: *Climate Since A.D. 1500*, Bradley, R.S. and Jones,
1784 P.D. (Eds.), Routledge, London, 623-648, 1992.

1785 Quinn, W.H., Neal, V.T., and Antunez de Mayolo, S.E.: El Nino occurrences over the past four and a half centuries, *Journal*
1786 *of Geophysical Research*, 92, 14449-14461, 1987.

1787 Rácz, L.: Climate history of Hungary since 16th Century. Past, present and future, Centre for Regional Studies of the
1788 Hungarian Academy of Sciences, Pécs, 1999.

1789 Rodrigo, F.S. and Barriendos, M.: Reconstruction of seasonal and annual rainfall variability in the Iberian peninsula (16th-
1790 20th centuries) from documentary data, *Global and Planetary Change*, 63, 243-257, 2008.

1791 Rodrigo, F.S., Estebanparra, M.J., and Castro-Diez, Y.: An attempt to reconstruct the rainfall regime of Andalusia (southern
1792 Spain) from 1601 ad to 1650 AD using historical documents, *Climatic Change*, 27, 397-418, 1994.

1793 Rodrigo, F.S., Esteban-Parra, M.J., and Castro-Diez, Y.: On the use of the Jesuit order private correspondence records in
1794 climate reconstructions: A case study from Castille (Spain) for 1634-1648 AD, *Climatic Change*, 40, 625-645, 1998.

1795 Rodrigo, F.S., Esteban-Parra, M.J., Pozo-Vazquez, D., and Castro-Diez, Y.: A 500-year precipitation record in Southern
1796 Spain, *International Journal of Climatology*, 19, 1233-1253, 1999.

1797 Rohr, C.: Measuring the frequency and intensity of floods of the Traun River (Upper Austria), 1441-1574, *Hydrological*
1798 *Sciences Journal-Journal Des Sciences Hydrologiques*, 51, 834-847, 2006.

1799 Rohr, C.: Extreme Naturereignisse im Ostalpenraum. Naturerfahrung im Spätmittelalter und am Beginn der Neuzeit,
1800 *Umwelthistorische Forschungen* 4, Böhlau, Cologne, Weimar, Vienna, 2007.

- Rohr, C.: Floods of the Upper Danube River and Its tributaries and their impact on urban economies (c. 1350-1600): The examples of the towns of Krems/Stein and Wels (Austria), *Environment and History*, 19, 133-148, 2013.
- Rohr, C., Camenisch, C., and Pribyl, K.: European Middle Ages. In: *The Palgrave Handbook of Climate History*, White, S., Pfister, C., and Mauelshagen, F. (Eds.), Palgrave-Macmillan, London, 247-263, 2018.
- Russell, H.C.: *Climate of New South Wales: Descriptive, historical, and tabular*, Charles Potter, Government Printer, Sydney, Australia, 1877.
- Salvisberg, M.: *Der Hochwasserschutz an der Gürbe. Eine Herausforderung für Generationen (1855-2010)*, Wirtschafts-, Sozial- und Umweltgeschichte 7, Schwabe, Basel, 2017.
- Schwarz-Zanetti, G.: *Grundzüge der Klima- und Umweltgeschichte des Hoch- und Spätmittelalters in Mitteleuropa*, Studentendruckerei, Zürich, 1998.
- Shabalova, M.V. and van Engelen, A.G.V.: Evaluation of a reconstruction of winter and summer temperatures in the low countries, AD 764-1998, *Climatic Change*, 58, 219-242, 2003.
- Shahgedanova, M. (Ed.): *The physical geography of northern Eurasia*, Oxford University Press, Oxford, 2002.
- Shen, X.Y. and Chen, J.Q.: Grain production and climatic variation in Taihu Lake Basin, *Chinese Geographical Science*, 3, 173-178, 1993.
- Shi, F., Zhao, S., Guo, Z.T., Goosse, H., and Yin, Q.Z.: Multi-proxy reconstructions of May-September precipitation field in China over the past 500 years, *Climate of the Past*, 13, 1919-1938, 2017.
- Shō, K., Shibuya, K., and Tominaga, A.: Examination of long-term changes in the rainy season by comparing diary weather records with meteorological observation data, *Journal of Hydrology and Water Resources*, 30, 294-306, 2017.
- Slepcev, A.M. and Klimenko, V.V.: Обобщение палеоклиматических данных и реконструкция климата восточной Европы за последние 2000 лет [Generalization of paleoclimatic data and reconstruction of the climate of Eastern Europe for the last 2000 years], *История и современность [History and Modernity]*, 1, 118-137, 2005.
- Sturm, K., Glaser, R., Jacobeit, J., Deutsch, M., Brázdil, R., Pfister, C., Luterbacher, J., and Wanner, H.: Hochwasser in Mitteleuropa seit 1500 und ihre Beziehung zur atmosphärischen Zirkulation, *Petermanns Geographische Mitteilungen*, 145, 14-23, 2001.
- Su, Y., Fang, X.Q., and Yin, J.: Impact of climate change on fluctuations of grain harvests in China from the Western Han Dynasty to the Five Dynasties (206 BC-960 AD), *Science China-Earth Sciences*, 57, 1701-1712, 2014.
- Tagami, Y.: Kohyōki chūki no nipponrettō no kikō hendō [Reconstruction of Climate Variation of Japan from 11th to 16th Century], *Ningen hattatsu kagakubu kiyō [Bulletin of the Faculty of Human Development, University of Toyama]*, 10, 205-219, 2016.
- Tan, L.C., Ma, L., Mao, R.X., and Tsai, Y.J.: Past climate studies in China during the last 2000 years from historical documents, *Journal of Earth Environment*, 5, 434-440, 2014.
- Tan, P.-H. and Liao, H.-M.: Reconstruction of temperature, precipitation and weather characteristics over the Yangtze River Delta Area in Ming Dynasty, *Journal of Geographical Science* 57, 61-87, 2012.
- Tan, P.-H. and Wu, B.-L.: Reconstruction of climatic and weather characteristics in the Shanghai area during the Qing dynasty, *Journal of Geographical Science* 71, 1-28, 2013.
- Tejedor, E., de Luis, M., Barriendos, M., Cuadrat, J.M., Luterbacher, J., and Saz, M.Á.: Rogation ceremonies: key to understand past drought variability in northeastern Spain since 1650, *Climate of the Past Discussions*, 2018. <https://doi.org/10.5194/cp-2018-67>, 2018.
- Telelis, I.G.: Climatic fluctuations in the Eastern Mediterranean and the Middle East AD 300–1500 from Byzantine documentary and proxy physical paleoclimatic evidence – A comparison, *Jahrbuch der Österreichischen Byzantinistik*, 58, 167-207, 2008.
- Tian, H., Stige, L.C., Cazelles, B., Kausrud, K.L., Svarverud, R., Stenseth, N.C., and Zhang, Z.: Reconstruction of a 1910-y-long locust series reveals consistent associations with climate fluctuations in China, *Proceedings of the National Academy of Sciences*, 108, 14521-14526, 2011.
- Trouet, V., Harley, G.L., and Dominguez-Delmas, M.: Shipwreck rates reveal Caribbean tropical cyclone response to past radiative forcing, *Proceedings of the National Academy of Sciences*, 113, 3169-3174, 2016.
- Van Engelen, A.F.V., Buisman, J., and IJnsen, F.: A millennium of weather, winds and water in the Low Countries. In: *History and Climate. Memories of the Future?*, Jones, P.D., Ogilvie, A.E.J., Davies, T.D., and Briffa, K.R. (Eds.), Kluwer Academic/Plenum Publishers, New York, 101-123, 2001.
- Veselovskij, K.S.: *O klimате Rossii [About Russian climate]*, Publishing House of the Imperial Academy of Sciences, Saint Petersburg, 1857.
- Vogel, C.H.: 160 years of rainfall in the Cape - has there been a change?, *South African Journal of Science*, 84, 1988.

- 1854 Vogel, C.H.: A documentary-derived climatic chronology for South Africa, 1820–1900, *Climatic Change*, 14, 291-307,
1855 1989.
- 1856 Vogt, S., Glaser, R., Luterbacher, J., Riemann, D., Al Dyab, G., Schönbein, J., and Garcia-Bustamente, E.: Assessing the
1857 Medieval Climate Anomaly in the Middle East: The potential of Arabic documentary sources, *PAGES News*, 19, 28-29,
1858 2011.
- 1859 Wang, P.K.: Meteorological records from ancient chronicles of China, *Bulletin of the American Meteorological Society*, 60,
1860 313-317, 1979.
- 1861 Wang, P.K.: On the relationship between winter thunder and the climatic change in China in the past 2200 years, *Climatic*
1862 *Change*, 3, 37-46, 1980.
- 1863 Wang, P.K. and Zhang, D.: An introduction to some historical governmental weather records of China, *Bulletin of the*
1864 *American Meteorological Society*, 69, 753-758, 1988.
- 1865 Wang, P.K. and Zhang, D.: A study on the reconstruction of the 18th century meiyu (plum rains) activity of Lower Yangtze
1866 region of China, *Science in China: Series B*, 34, 1237-1245, 1991.
- 1867 Wang, P.K. and Zhang, D.: Recent studies of the reconstruction of east Asian monsoon climate in the past using historical
1868 literature of China, *Meteorological Society of Japan*, 70, 423-446, 1992.
- 1869 Wang, P.K., Lin, K.-H.E., Liao, Y.C., Liao, H.M., Lin, Y.S., Hsu, C.T., Hsu, S.M., Wan, C.W., Lee, S.Y., Fan, I.C., Tan,
1870 P.H., and Ting, T.T.: Construction of the REACHES climate database based on historical documents of China, *Scientific*
1871 *Data*, 5, 2018.
- 1872 Wang, R.S. and Wang, S.W.: Reconstruction of winter temperature in Eastern China during the past 500 years using
1873 historical documents, *Acta Meteorologica Sinica*, 48, 379-386, 1990a.
- 1874 Wang, S.L., Ye, J.L., and Gong, D.Y.: Climate in China during the Little Ice Age, *Quaternary Sciences* 1, 54-64, 1998.
- 1875 Wang, S.W. and Wang, R.S.: Variations of seasonal and annual temperatures during 1470-1979 in eastern China,
1876 *Meteorological Bulletin*, 48, 26-35, 1990b.
- 1877 Wang, S.W. and Gong, D.Y.: Climate in China during the four special periods in Holocene, *Progress in Natural Science*, 10,
1878 379-386, 2000.
- 1879 Wang, S.W., Wang, K.S., Zhang, Z.M., and Ye, J.L.: The change of drought and flood disasters over the areas of Yangtze
1880 and Yellow rivers during 1380-1989. In: *Diagnosis Research of Frequency and Economic Effect for Drought and Flood*
1881 *Disasters over Yangtze and Yellow Rivers*, Wang, S.W. and Huang, Z.I. (Eds.), China Meteorological Press, Beijing, 41-
1882 54, 1993.
- 1883 Ward, C. and Wheeler, D.A.: Hudson's Bay Company ship's logbooks: a source of far North Atlantic weather data, *Polar*
1884 *Record*, 48, 165-176, 2012.
- 1885 Warren, H.N.: Results of rainfall observations made in New South Wales, Sections I - VI, Districts 46 - 75, including rainfall
1886 tables (monthly and annual), discussion of rainfall and its relation to primary industries, also temperature and humidity
1887 tables, records of floods, cyclones, and local storms, etc., Bureau of Meteorology, Canberra, 1948.
- 1888 Watt, W.S.: Results of rainfall observations made in Tasmania including all available annual rainfall totals from 356 stations
1889 for all years of record up to 1934, with maps and diagrams; and record of severe floods., Bureau of Meteorology,
1890 Melbourne, 1936.
- 1891 Wetter, O., Pfister, C., Weingartner, R., Luterbacher, J., Reist, T., and Trösch, J.: The largest floods in the High Rhine basin
1892 since 1268 assessed from documentary and instrumental evidence, *Hydrological Sciences Journal*, 56, 733-758, 2011.
- 1893 Wheeler, D.A.: Understanding seventeenth-century ships' logbooks: An exercise in historical climatology, *Journal for*
1894 *Maritime Research* 6, 21-36, 2004.
- 1895 Wheeler, D.A.: An examination of the accuracy and consistency of ships' logbook weather observations and records,
1896 *Climatic Change*, 73, 97-116, 2005a.
- 1897 Wheeler, D.A.: British naval logbooks from the late seventeenth century: New climatic information from old sources,
1898 *History of Meteorology* 2, 133-145, 2005b.
- 1899 Wheeler, D.A. and Garcia-Herrera, R.: Ships' logbooks in climatological research: Reflections and prospects. In: *Trends and*
1900 *Directions in Climate Research*, Gimeno, L., Garcia-Herrera, R., and Trigo, R.M. (Eds.), Annals of the New York
1901 Academy of Sciences, 1-15, 2008.
- 1902 Wheeler, D.A., Garcia-Herrera, R., Wilkinson, C., and Ward, C.: Atmospheric circulation and storminess derived from
1903 Royal Navy logbooks: 1685 to 1750, *Climatic Change*, 101, 257-280, 2010.
- 1904 White, S.: North American climate history (1500–1800). In: *The Palgrave Handbook of Climate History*, White, S., Pfister,
1905 C., and Mauelshagen, F. (Eds.), Palgrave Macmillan, London, 297-308, 2018.

1906 Wilhelm, B., Ballesteros Canovas, J.A., Corella Aznar, J.P., Kämpf, L., Swierczynski, T., Stoffel, M., Søren, E., and Tonen,
1907 W.: Recent advances in paleoflood hydrology: From new archives to data compilation and analysis, *Water Security* 3, 1-
1908 8, 2018.

1909 Wilkinson, C.: British logbooks in UK archives, 17th–19th centuries – a survey of the range, selection and suitability of
1910 British logbooks and related documents for climatic research, Climatic Research Unit, School of Environmental
1911 Sciences, University of East Anglia, Norwich, 2009.

1912 Wilson, R., Tudhope, A., Brohan, P., Briffa, K.R., Osborn, T., and Tett, S.F.B.: Two-hundred-fifty years of reconstructed
1913 and modeled tropical temperatures, *Journal of Geophysical Research*, 111, C10007,
1914 <https://doi.org/10.1029/2005JC003188>, 2006.

1915 Wozniak, T.: *Naturereignisse im frühen Mittelalter: Das Zeugnis der Geschichtsschreibung vom 6. bis 11 Jahrhundert*, De
1916 Gruyter, Berlin, 2020.

1917 Xiao, L., Fang, X., and Zhang, X.: Location of rainbelt of Meiyu during second half of 19th Century to early 20th Century,
1918 *Scientia Geographica Sinica*, 28, 385-389, 2008.

1919 Xoplaki, E., Maheras, P., and Luterbacher, J.: Variability of climate in Meridional Balkans during the periods 1675-1715 and
1920 1780-1830 and its impact on human life, *Climatic Change*, 48, 581-615, 2001.

1921 Xoplaki, E., Luterbacher, J., Paeth, H., Dietrich, D., Steiner, N., Grosjean, M., and Wanner, H.: European spring and autumn
1922 temperature variability and change of extremes over the last half millennium, *Geophysical Research Letters*, 32,
1923 <https://doi.org/10.1029/2005GL023424>, 2005.

1924 Yao, C.S.: A statistical approach to historical records of flood and drought, *Journal of Applied Meteorology*, 21, 588-594,
1925 1982.

1926 Yao, S.Y.: The geographical distribution of floods and droughts in Chinese history, 206 B.C.-A.D. 1911 *The Far Eastern*
1927 *Quarterly*, 2, 357-378, 1943.

1928 Yi, L., Yu, H.J., Ge, J.Y., Lai, Z.P., Xu, X.Y., Qin, L., and Peng, S.Z.: Reconstructions of annual summer precipitation and
1929 temperature in north-central China since 1470 AD based on drought/flood index and tree-ring records, *Climatic Change*,
1930 110, 469-498, 2012.

1931 Yin, J., Su, Y., and Fang, X.Q.: Relationships between temperature change and grain harvest fluctuations in China from 210
1932 BC to 1910 AD, *Quaternary International*, 355, 153-163, 2015.

1933 Zaiki, M., Grossman, M.J., and Mikami, T.: Document-based reconstruction of past climate in Japan, *PAGES News*, 20, 82-
1934 83, 2012.

1935 Zhang, D.: Winter temperature changes during the last 500 years in South China, *Chinese Science Bulletin*, 25, 497-500,
1936 1980.

1937 Zhang, D.: Preliminary analyses of the weather and climate during dust storms in the historical time, *Science in China B*, 24,
1938 278-288, 1984.

1939 Zhang, D.: *A Compendium of Chinese Meteorological Records of the Last 3,000 Years*, Jiangsu Education Press, Nanjing,
1940 2004.

1941 Zhang, D. and Liu, C.Z.: Reconstruction of summer temperature series (1724-1903) in Beijing, *Kexue Tongbao*, 32, 1046-
1942 1049, 1987.

1943 Zhang, D. and Wang, P.-K.: Reconstruction of the eighteenth century summer monthly precipitation series of Nanjing,
1944 Suzhou, and Hangzhou using the Clear and Rain Records of Qing Dynasty, *Journal of Meteorological Research*, 3, 261-
1945 278, 1989.

1946 Zhang, D. and Wang, P.-K.: Reconstruction of the 18th century summer monthly precipitation series of Nanjing, Suzhou and
1947 Hangzhou using Clear and Rain Records of Qing dynasty, *Quarterly Journal of Applied Meteorology*, 1, 260-270, 1990.

1948 Zhang, D. and Liu, C.J.: Continuation (1980-1992) to ‘Yearly Charts of Dryness/Wetness in China for the Last 500-year
1949 Period’, *Meteorological Monthly*, 19, 41-46, 1993.

1950 Zhang, D. and Liu, Y.: A new approach to the reconstruction of temporal rainfall sequences from 1724-1904 Qing dynasty
1951 weather records for Beijing, *Quaternary Sciences*, 22, 199-208, 2002.

1952 Zhang, D., Liu, C., and Jiang, J.: Reconstruction of six regional dry/wet series and their abrupt changes during the last 1000
1953 years in East China, *Quaternary Sciences*, 17, 1-11, 1997.

1954 Zhang, D., Lee, X.C., and Liang, Y.Y.: Continuation (1993-2000) to ‘Yearly Charts of Dryness/Wetness in China for the
1955 Last 500-year Period’, *Journal of Applied Meteorological Science*, 14, 379-384, 2003.

1956 Zhang, J.C. and Zhang, X.G.: Climatic fluctuations during the last 500 years in China and their interdependence, *Acta*
1957 *Meteorological Sinica*, 37, 49-57, 1979.

- 1958 Zhang, J.C. and Crowley, T.J.: Historical climate records in China and reconstruction of past climates, *Journal of Climate*, 2,
1959 833-849, 1989.
- 1960 Zhang, P.Y. and Gong, G.F.: Some characteristics of climate fluctuations in China since 16th century, *Acta Geographica*
1961 *Sinica*, 46, 238-247, 1979.
- 1962 Zheng, J.Y. and Zheng, S.Z.: An analysis on cold/warm and dry/wet in Shandong Province during historical times, *Acta*
1963 *Geographica Sinica*, 48, 348-357, 1993.
- 1964 Zheng, J.Y., Ge, Q.S., Fang, Z.Q., and Zhang, X.Z.: Comparison on temperature series reconstructed from historical
1965 documents in China for the last 2000 years, *Acta Meteorologica Sinica*, 65, 428-439, 2007.
- 1966 Zheng, J.Y., Wang, W.C., Ge, Q.-S., Man, Z.M., and Zhang, P.Y.: Precipitation variability and extreme events in eastern
1967 China during the past 1500 years, *Terrestrial Atmospheric and Oceanic Sciences*, 17, 579-592, 2006.
- 1968 Zheng, S.Z., Zhang, F.C., and Gong, G.F.: Preliminary analysis of moisture condition in southeastern China during the last
1969 two thousand years. In: *Proceedings of Symposium on Climatic Variations and Long-term Forecasting*, Science Press,
1970 Beijing, 1977.
- 1971 Zhogova, M.L.: Klimaticheskie zakonomernosti na territorii Rossii v Trudah K.S. Veselovskogo [Climate Regularities of
1972 Russia in the Works of K.S. Veselovsky], *Естественные науки: История естествознания* [Natural Sciences: History of
1973 Natural Science], 1, 160-167, 2013.
- 1974 Zhou, Q., Zhang, P., and Wang, Z.: Reconstruction of annual winter mean temperature series in Hefei area during 1973–
1975 1991, *Acta Geographica Sinica* 49, 332-337, 1994.
- 1976 Zhu, C.: Climate pulsations during historical times in China, *Geographical Review*, 16, 274-281, 1926.
- 1977 Zhu, C. and Wang, M.: *Phenology*, Science Press, Beijing, 1973.
- 1978 Zhu, K.: A preliminary study on climate change in China in the last 5000 years, *Scientia Sinica Mathematica*, 16, 168-189,
1979 1973.
- 1980