

# Signals of Holocene climate transitions amplified by anthropogenic land-use changes in the Westerly-Indian Monsoon realm and its anthropogenic impact

Nicole Burdanowitz<sup>1</sup>, Tim Rixen<sup>1,2</sup>, Birgit Gaye<sup>1</sup>, Kay-Christian Emeis<sup>1,3</sup>

5 <sup>1</sup>Institute for Geology, Universität Hamburg, Bundesstraße 55, 20146 Hamburg, Germany

<sup>2</sup>Leibniz-Zentrum für Marine Tropenforschung (ZMT), Fahrenheitstraße 6, 28359 Bremen, Germany

<sup>3</sup>Institute of Coastal Research, Helmholtz Center Geesthacht, Max-Planck-Straße 1, 21502 Geesthacht, Germany

*Correspondence to:* Nicole Burdanowitz (nicole.burdanowitz@uni-hamburg.de)

10 **Abstract.** The Indian Summer Monsoon (ISM) with its rainfall is the lifeline for people living on the Indian subcontinent today and possibly was the driver of the rise and fall of early agricultural societies in the past. Intensity and position of the ISM have shifted in response to orbitally forced thermal land-ocean contrasts. At the northwestern monsoon margins, interactions between the subtropical westerly jet (STWJ) and the ISM constitute a tipping element in the Earth's climate system, because their non-linear interaction may be a first-order influence on rainfall. We reconstructed marine sea surface  
15 temperature (SST), supply of terrestrial material and vegetation changes from a very well-dated sediment core from the northern Arabian Sea to reconstruct the STWJ-ISM interaction. The Holocene record (from 11,000 years) shows a distinct, but gradual, southward displacement of the ISM in the Early to Mid-Holocene, increasingly punctuated by phases of intensified STWJ events that are coeval with interruptions of North Atlantic overturning circulation (Bond events). Effects of the non-linear interactions culminate between 4.6-3 ka BP, marking a climatic transition period during which the ISM shifted  
20 southwards and the influence of SWTJ became prominent. The lithogenic matter input shows an up to 4-fold increase after this time period probably related to signaling the strengthened influence of agricultural activities of the Indus civilization with enhanced erosion of soils. This anthropogenic land-use change is amplifying the impact of Bond events and adding to the marine sedimentation rates adjacent to the continent.

## 1 Introduction

25 Changes in latitude and strength of the ISM were gradual over the course of the Holocene, but were punctuated by distinct climatic events (Herzschuh, 2006). The ISM was strong during the Early Holocene until approximately 6.5 ka ago, and influenced regions farther north than today (Herzschuh, 2006; Prasad and Enzel, 2006). A weakening of the ISM is reflected in many records after the Holocene Optimum Period, commonly attributed to declining summer insolation (Banerji et al., 2020; Herzschuh, 2006), latitudinal insolation gradients (Mohtadi et al., 2016; Ramisch et al., 2016) and feedbacks of climate  
30 anomalies such as the Arctic Oscillation (Zhang et al., 2018), the North Atlantic Oscillation (NAO) (Band et al., 2018; Banerji

et al., 2020; Kotlia et al., 2015; Lauterbach et al., 2014) or the El Niño Southern Oscillation (ENSO) (Banerji et al., 2020; Prasad et al., 2014; Srivastava et al., 2017).

Changes of the ISM system not only affected hydrological conditions and vegetation in this region, but were also an important constraint on the expansion and dispersal of the Indus civilization, also known as Harappan civilization, emerging around 5 ka ago (Durcan et al., 2019; Giosan et al., 2012, 2018; Possehl, 1997). Growing agricultural activities adjacent to the northwestern Arabian Sea increased soil erosion and, therefore, terrigenous sediment input (Gourlan et al., 2020). Further, the decrease of ISM precipitation favored the development of combined summer and winter cropping as well as drought-tolerant crops (Petrie and Bates, 2017). Winter precipitation intensity and interannual variability in the present-day northwestern region of the Indian Monsoon (IM) domain is governed by STWJ-induced events of strong precipitation, so called Western Disturbances (WD) (Anoop et al., 2013; Dimri et al., 2016; Hunt et al., 2018; Leipe et al., 2014; Munz et al., 2017; Zhang et al., 2018). Climatological data show a link between the NAO, the position of the STWJ and the strength of the WD during the winter over northwest India (Yadav et al., 2009). A positive NAO is associated with an intensified STWJ and stronger WD and, thus, above normal winter precipitation in this region (Yadav et al., 2009). The interaction, position and strength of the IM and the STWJ on millennial to Holocene timescales are yet poorly understood, because the pronounced signal of the ISM dominates sedimentary records, and high-quality records in the region where IM and STWJ interact are sparse.

The northeastern (NE) Arabian Sea (AS) is located in the region of interaction of IM and STWJ. Whereas the primary productivity during the warm ISM season is driven by lateral transport of nutrient-rich waters from the Oman upwelling area (Schulz et al., 1996), the NE winds deepen the mixed layer during the colder IWM season due to convective mixing leading to enhanced primary productivity and carbon exports to the deep sea (Banse and McClain, 1986; Madhupratap et al., 1996; Rixen et al., 2019). The NE AS holds high-quality archives due to excellently preserved and highly resolved sediment records deposited on the shelf and slope impinged by a strong oxygen minimum zone (OMZ) between 200 m and 1200 m water depth (von Rad et al., 1995; Schott et al., 1970; Schulz et al., 1996). Most existing records have poor temporal resolution or do not bracket the entire Holocene (Böll et al., 2014, 2015; Giosan et al., 2018; Lückge et al., 2001; Munz et al., 2015), but they do outline the interplay of the Indian Winter Monsoon (IWM) with the westerlies in the NE AS realm during the Holocene and suggest that winter monsoon activity intensified between 3.9 and 3.0 ka BP (Giesche et al., 2019; Giosan et al., 2018; Lückge et al., 2001) and intermittently during the last ca. 1.5 ka (Böll et al., 2014; Lückge et al., 2001; Munz et al., 2015).

For a detailed and high-resolution reconstruction, we analyzed lithogenic mass accumulation rates (LIT MAR), alkenone-based SST and *n*-alkane-based land vegetation from the NE AS (Figure 1) over the last 10.7 ka. They document the influence of the IWM and the westerlies on the marine and terrestrial environment in the NE AS region and show that the advent of agriculture in step with climate change significantly enhanced soil erosion.




Figure 1: Location of the study site SO90-63KA (red star), the nearby record Indus 11C (Giosan et al., 2018) (red circle) and Lake Qinghai (Hou et al., 2016) (blue dot in the inset map). The rivers (blue lines) important for the study site are shown. Prevailing wind pattern during the summer monsoon, here indicated as southwest (SW) monsoon, is shown with a black arrow. The prevailing wind pattern of the winter monsoon, here indicated as northeast (NE) monsoon, as well as the subtropical westerly jet and its associated rain bearing Western Disturbances, are shown as black dashed arrows. The map was created using ArcGIS v.10.8 (ESRI, 2019). The bathymetric data are from the General Bathymetric Chart of the Oceans (GEBCO2014; www.gebco.net).

65

## 2 Material and Methods

70 Records presented here were ~~raised~~obtained from the box core SO90-63KA (697 cm long), which was retrieved from the Arabian Sea off Pakistan (24°36.6'N, 65°59.0'E, 315 m water depth) during the RV SONNE cruise SO90 in 1993. With the upper 18 cm missing, SO90-63KA covers the last ca. 10.7 ka and encompasses the development of early Bronze Age agricultural societies (Staubwasser et al., 2002, 2003). The chronology is based on 80 <sup>14</sup>C dates of planktic foraminifers *Globigerinoides sacculifer* (Staubwasser et al., 2002, 2003) and belongs to the ~~most-well~~best dated cores from the Arabian  
75 Sea. ~~The bulk components, grain sizes and major and trace elements of SO90-63KA were published previously (Burdanowitz et al., 2019; Staubwasser and Sirocko, 2001). We used an extended data set for the grain size analyses of SO90-63KA sediments for the endmember modelling analyses. The procedure is described in the earlier published coarser resolution data set (Burdanowitz et al., 2019).~~ The major and trace elements of SO90-63KA were published previously (Burdanowitz et al., 2019; Staubwasser and Sirocko, 2001).

### 80 2.1 Lipid analyses (alkenones and *n*-alkanes)

For the lipid analyses, 2.0 to 13.4 g of freeze-dried and grounded sediment were extracted with a DIONEX Accelerated Solvent Extractor (ASE 200) at 75°C, 1000 psi for 5 min using dichloromethane (DCM) as solvent. The procedure was repeated three times. A known amount of an internal standard (14-heptacosanone, squalane) was added prior to extraction. An additional working sediment standard was extracted for each ASE running sequence. The total lipid extracts (TLEs) were concentrated  
85 using rotary evaporation. The TLEs were separated into a hexane-soluble and a hexane-insoluble fraction by Na<sub>2</sub>SO<sub>4</sub> column chromatography. The hexane-soluble fractions were saponified at 85°C for 2h using a solution of 5% potassium hydroxide in methanol (MeOH). Afterwards, the neutral fraction was extracted with hexane. Column chromatography using deactivated silica gel (5% H<sub>2</sub>O, 60 mesh) was carried out to separate the neutral fraction into an apolar- (containing *n*-alkanes), ketone- (containing alkenones) and polar-fraction by using hexane, DCM and DCM:MeOH (1:1), respectively. The apolar fraction  
90 was further cleaned by AgNO<sub>3</sub>-Si column chromatography using hexane as solvent. Quantification of the *n*-alkanes and the alkenones was carried out using an Agilent 6850 gas chromatograph (GC) equipped with an Optima1 column (30 m, 0.32 mm, 0.1 μm), split/splitless injector operating at 280°C and a flame ionization detector (FID, 310°C). H<sub>2</sub> was used as carrier gas with a flow rate of 1.5 ml/min. Samples were injected in hexane and duplicate measurements of each sample were carried out for the alkenones. For the *n*-alkanes, the GC temperature was programmed  
95 from 50°C (held 1 min) ramped at 8°C/min to 320°C (held 15 min). An external standard was used for quantification, containing *n*-C<sub>8</sub> to *n*-C<sub>40</sub> alkanes in known concentration. Repeated analyses of the external standard resulted in a quantification precision of 6 % (1<sub>SD</sub>).

The average chain length (ACL) of the homologues C<sub>27</sub>-C<sub>33</sub> was calculated using following equation:

$$ACL_{27-33} = \frac{27 * C_{27} + 29 * C_{29} + 31 * C_{31} + 33 * C_{33}}{C_{27} + C_{29} + C_{31} + C_{33}}$$

100 where  $C_x$  is the concentration of the  $n$ -alkane with  $x$  carbon atoms.

The carbon preference index (CPI) of the  $n$ -alkanes was calculated as:

$$CPI_{27-33} = 0.5 * \left( \frac{C_{27} + C_{29} + C_{31} + C_{33}}{C_{26} + C_{28} + C_{30} + C_{32}} + \frac{C_{27} + C_{29} + C_{31} + C_{33}}{C_{28} + C_{30} + C_{32} + C_{34}} \right)$$

where  $C_x$  is the concentration of the  $n$ -alkanes with  $x$  carbon atoms.

For the alkenones, the GC temperature was programmed to increase from 50°C (held 1 min) to 230°C at 20°C/min, then at  
105 4.5°C/min to 260°C and at 1.5°C/min to 320°C (held 15 min). The  $C_{37:2}$  and  $C_{37:3}$  alkenones were identified by comparing the retention time peaks of the samples and the known working sediment standard. Quantification was carried out by using a known amount of an external standard (14-heptacosanone and hexatriacontane).

We calculated the alkenone unsaturation index using the following equation (Prahl et al., 1988):

$$U_{37}^{k'} = \frac{C_{37:2}}{C_{37:2} + C_{37:3}}$$

110 and using the core top calibration of Indian Ocean sediments (Sonzogni et al., 1997) to convert the UK'37 index to SSTs:

$$SST = \frac{U_{37}^{k'} - 0.043}{0.033}$$

For each sample, at least one duplicate measurement was obtained with an average precision of 0.1°C (1<sub>SD</sub>). The precision of the replicate extractions of the working standard sediment (n=13) and duplicate measurements of each replicate was 0.5°C (1<sub>SD</sub>).

## 115 2.2 Lithogenic mass accumulation rates

The total mass accumulation rates (MAR) were calculated by multiplying the dry bulk density (DBD) with the linear sedimentation rate (LSR). The LSR was ~~derivedascertain by using age model depth and age interval valuesfrom depth increments of successive age model tie points~~. DBD was calculated with the following equation (Avnimelech et al., 2001):

$$DBD = \frac{\text{weight dry sample (Wd)}}{\text{total sample volume (Vt)}}$$

120 where

$$\begin{aligned} Vt &= \text{volume of solids} + \text{volume of water} \\ &= \frac{\text{weight dry sample}}{\text{particle density}} + (\text{weight wet sample} - \text{weight dry sample}) \end{aligned}$$

~~For calculating the sediment particle density, We we~~ assumed 2.65 g/cm<sup>3</sup> ~~as the density of for~~ inorganic sediment particles (Blake and Hartge, 1986; Boyd, 1995) and 1 g/cm<sup>3</sup> for water density. ~~We performed a density correction, and corrected~~ for  
125 organic matter contents assuming a density of 1.25 g/cm<sup>3</sup> for organic particles (Boyd, 1995):

$$\text{sediment particle density}_{\text{weighted average}} = 1.25 * \%OM + 2.65 * (100 - \%OM)$$

It has to be noted that we did not calculate the LIT MAR between 9.4 and 8.5 ka BP because the core was dried out in this section.

The wavelet power spectrum analyses of the LIT MAR record were carried out using a Matlab Wavelet script (Torrence and Compo, 1998). For this analysis we interpolated our data set to achieve a temporal resolution of about 15.6 years. This is the minimum time step between two samples and also covers the gap between 9.4 and 8.5 ka BP.

The bulk components, grain sizes and major and trace elements of SO90-63KA were published previously (Burdanowitz et al., 2019; Staubwasser and Sirocko, 2001).

### **2.3 Grain size analyses**

A coarser resolution data set of grain size analyses of SO90-63KA was previously published (Burdanowitz et al., 2019). Here, we used an extended data set of 146 sediment samples for the grain size analyses of SO90-63KA sediments and for the endmember modelling analyses. The grain size analyses were carried out at the Center for Tropical Marine Research (ZMT, Bremen). To remove inorganic carbon and organic matter 250 mg of bulk sediment were treated with muriatic acid (HCl) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) (e.g. Sun et al., 2002). Afterwards, the samples were treated with Calgon® (Na<sub>6</sub>P<sub>6</sub>O<sub>18</sub>) and boiled to avoid particle coagulation. The grain sizes were then measured with a Horiba LA-950V2 laser scattering particle size analyser. A triplicate measurement of each sample was carried out and results were averaged. The grain size distribution was calculated for 93 grain size classes between 10 and 3 μm and grain size distribution parameters were calculated with GRADISTAT V.8. The end-member modelling analysis (EMMA) Matlab script by (Dietze et al., 2012) was used for the end-member modelling.

The procedure is described in the earlier published coarser resolution data set (Burdanowitz et al., 2019).

## **3 Results & Discussion**

### **3.1 Alkenones and SST record of SO90-63KA**

SST estimates based on the UK'37 index reconstructed SST varies by about 2°C from 25.9°C to 27.9°C (UK'37: 0.896 to 0.963) throughout the Holocene at the core site (Figure 2). Periods of low SSTs were the Early Holocene until ca. 8.7 ka BP and the time interval between ca. 4.6 and 3.0 ka. A marked decrease of about 1°C occurred between 4.2 and 3.8 ka BP. High SSTs were registered between 8.1 and 5.2 ka BP during the Mid-Holocene. The last 3 ka show also high and, in addition, more variable SSTs than before at the core site.

### **3.2 n-Alkane record of SO90-63KA**

The n-alkane patterns of sediment samples (n=93) display a strong odd-over-even carbon number predominance with CPI<sub>27-33</sub> values of 2.1 to 7.6 (Fig. S1). The n-alkane C<sub>31</sub> is the most abundant component (mean: 25.5 ± 2.1 %) followed by n-C<sub>29</sub> (mean: 17.0 ± 1.8 %) and n-C<sub>33</sub> (mean: 12.2 ± 2.3 %) for most samples (Fig. S2). The ACL<sub>27-33</sub> ranges between 29.9 and 30.7 with highest values between 9.2 and 6.8 ka BP and lowest values from about 6.8 to 3.6 ka BP during the Mid-Holocene (Fig. S1).

The  $ACL_{27-33}$  values increase slightly during the last about 3.5 ka, but on average, do not reach the high values characteristic for the Early Holocene period.

### **3.3 Lithogenic mass accumulation rates (LIT MAR) of SO90-63KA**

The LIT MAR in our record range between 225  $g/(m^2*yr)$  and 3714  $g/(m^2*yr)$  (Figure 2). Periods of high LIT MAR occurred from 10.1 – 9.8 ka BP, at ca. 8.3 ka BP, around 3.4 ka BP, between 2.8 - 2.7 ka BP, 2.2 - 2.1 ka BP, 1.5 – 1.1 ka BP and over the last 300 years. The LIT MAR are relatively low in the Mid-Holocene period from 8.3 to 3.5 ka BP with only minor fluctuations compared to the rest of our record.

### **3.4 Grain size and end-member modelling analyses of SO90-63KA**

The grain size analyses of SO90-63KA ( $n=146$ ) show a bimodal pattern with modal grain sizes of 7.2  $\mu m$  and 0.5  $\mu m$  (Fig. S3). The median grain sizes of all samples range between 4.6  $\mu m$  and 10.0  $\mu m$  (mean:  $6.3 \pm 1.0 \mu m$ ) and the EMMA segregates three main end-members (Fig. S4). End-member 1 (EM1) presents a polymodal pattern with modal grain sizes of 5.5  $\mu m$ , 24.5  $\mu m$  and 0.4  $\mu m$ . End-member 2 (EM2) has a bimodal pattern with modal grain sizes of 16.27  $\mu m$  and 0.4  $\mu m$ , whereas end-member 3 (EM3) has a trimodal pattern with modal grain sizes of 0.4  $\mu m$ , 4.2  $\mu m$  and 42.1  $\mu m$  (Fig. S4).

## **4 Discussion**

### **4.1 Holocene climate change in the NE Arabian Sea realm**

The prime indicator for climate- and human-driven change on land adjacent to the NE AS is the lithogenic material archived in the sediment record. The lithogenic material at the core site is supplied by fluvial input from the Makran rivers (Forke et al., 2019; Lückge et al., 2002; von Rad et al., 2002; Staubwasser and Sirocko, 2001; Stow et al., 2002) and by aeolian input probably that originates from the Sistan Basin region in the border region of Afghanistan and Iran (Burdanowitz et al., 2019; Forke et al., 2019; Kaskaoutis et al., 2014; Rashki et al., 2012). At present dust from the Sistan region is transported to the NE AS by northerly Levant winds during summer and northeasterly and westerly winds during winter (Hussain et al., 2005; Kaskaoutis et al., 2015; Pease et al., 1998; Rashki et al., 2012; Sirocko et al., 1991; Tindale and Pease, 1999). Ti/Al ratios of SO90-63KA are an indicator for fluvial input by the Makran rivers (Burdanowitz et al., 2019) that helps to distinguish between aeolian and fluvial input. Ti along the Makran margin is enriched in terrestrial detritus mainly delivered from rivers (Lückge et al., 2001) and Ti/Al ratios are particularly high in fluvial deposited sediments along the Hingol River. In thick varves Ti/Al ratios are enriched compared to the thin varves indicating enhanced transport energy as a result of a more intense fluvial discharge (Lückge et al., 2001). Another useful proxy is end-member 3 (EM3) of the EMMA as an indicator for aeolian input (Burdanowitz et al., 2019). Dust storms over Karachi, located east of the Hab River mouth, exhibits median grain sizes ranging between 2.4 and 3.4  $\mu m$  (Figure S3) and EM3 has modal grain sizes of 4.2  $\mu m$  (Figure S4, this study) and 3.7  $\mu m$  (Burdanowitz

et al., 2019), respectively. The modal grain size of 4.2  $\mu\text{m}$  is similar to that of the dust over Karachi and can thus be used as an indicator for aeolian input.

190 Alkenone based SSTs reconstructed from our record range between 25.9 and 27.9°C throughout the Holocene. It has been pointed out that ~~Changes~~ changes in the alkenone producing coccolithophores communities *Emiliania huxleyi* and *Gephyrocapsa oceanica* in the NE AS are controlled by ~~changes in the~~ nutrient availability and ~~the~~ mean mixed layer depth (Andruleit et al., 2004). ~~This variation in~~ Although, the coccolithophores communities ~~in the NE AS varies~~ over the course of the year with ~~results in~~ higher alkenone fluxes during winter and spring ~~compared to other seasons. However, investigations of seasonal data on alkenone temperatures;~~ (Böll et al., 2014) ~~showed~~ found that the alkenone-based SST reconstructions ~~from sediments are is~~ representative for annual mean SST ~~in our study area. The lithogenic material at the core site is supplied by fluvial input from the Makran rivers (Forke et al., 2019; Lückge et al., 2002; von Rad et al., 2002; Staubwasser and Sirocko, 2001; Stow et al., 2002) and by aeolian input probably from the Sistan Basin region in the border region of Afghanistan and Iran (Burdanowitz et al., 2019; Forke et al., 2019; Kaskaoutis et al., 2014; Rashki et al., 2012). At present dust from the Sistan region is transported to the NE AS by northerly Levant winds during summer and northeasterly and westerly wind during winter (Hussain et al., 2005; Kaskaoutis et al., 2015; Pease et al., 1998; Rashki et al., 2012; Sirocko et al., 1991; Tindale and Pease, 1999).~~

195 ~~The SSTs reconstructed from our record range between 25.9 and 27.9°C throughout the Holocene. During waning glacial conditions in the region during of the Early Holocene resulted in consistently lower the SSTs remained low until ca. 8.7 ka BP in the NE AS (Böll et al., 2015; Gaye et al., 2018; Giosan et al., 2018) (Figure 2Figure 2). Between 8.7 and 8.1 ka BP SST increased from 26.0°C to 27.2°C and was accompanied by with rising LIT MAR around 8.4 to 8.3 ka BP. However, since there is a gap in our LIT MAR record, as mentioned before, we cannot exclude an onset of rising LIT MAR even prior to 8.4 ka BP. Coeval rapid changes have been identified across the entire IM realm (Dixit et al., 2014; Rawat et al., 2015) and mark the so-called 8.2 ka event, which was associated with a weakened ISM (Cheng et al., 2009; Dixit et al., 2014).~~

200

205

The North Atlantic hematite stained grain (HSG) records indicate eight pronounced cold events, so called Bond events, during the Holocene (Bond et al., 2001). Recent studies suggested that changes in the NAO and the Atlantic Meridional Overturning Circulation (AMOC) ~~may might~~ have played ~~an~~ important roles ~~and causing~~ at least some of these cold events ~~in the North Atlantic region~~ (Ait Brahim et al., 2019; Goslin et al., 2018; Klus et al., 2018). Whereas some Bond events (0, 1, 5, 7 & 8) occurred during negative NAO-like conditions, linked to a reduction in North Atlantic Deep Water formation, the other Bond events (mainly during the Mid-Holocene) occurred during positive NAO-like conditions (Ait Brahim et al., 2019). The weakened ISM of the 8.2 ka event has been linked to one of these North Atlantic cool phases (Bond event 5) and is interpreted to signal a decrease in ocean heat transport that induced a marked southward shift of the Intertropical Convergence Zone (ITCZ) (Cheng et al., 2009).


210


215

Although our SST record shows no evidence of the 8.2 ka cold event in the NE AS,  $\delta^{18}\text{O}$  data of *Globigerinoides ruber* of the same core ~~interval~~ signal ~~abruptly~~ declining Indus river discharge around 8.4 ka BP (Staubwasser et al., 2002) (Figure 2), accompanied by ~~declining Ti/Al ratios and~~ rising LIT MAR indicative for strong soil erosion ~~due to enhanced aridity on land. It is plausible that this cold event, observed in both the continental records and marine proxies, led to a short southward~~

220

225 migration of the ITCZ and drying on the adjacent continent, resulting in a decoupling of the oceanic vs. land proxies in our  
eore. High SSTs prevailed after the Early-Mid Holocene transition ~~at~~from ca. 8.1 ka BP until ca. 5.2 ka BP and are consistent  
with other Arabian Sea records (Böll et al., 2015; Gaye et al., 2018; Giosan et al., 2018). ~~The~~is warm period ~~encompasses~~  
the Mid-Holocene climate optimum period and is characterized by ~~low LIT MAR and increasing fluvial input (Figure 2) due~~  
~~to~~ high precipitation rates in the North Himalayan region (Dallmeyer et al., 2013; Herzschuh, 2006) ~~and a dense vegetation~~  
~~cover which lowered soil erosion. High Ti/Al ratios thus reflect increasing fluvial inputs, although low LIT MAR indicate low~~  
~~soil erosion (Figure 2). The decoupling probably was due to a dense vegetation cover during this Mid-Holocene wet phase~~  
~~which also explains why the LIT MAR record shows no correlation to the Bond events during the Mid-Holocene. The regional~~  
230 ~~vegetation damped their remote influence.~~





235 Figure 2: Environmental changes in the NE Arabian Sea. a) alkenone-based reconstructed SST of SO90-63KA, b) factor 1 based on  
DNA factor analyses of planktic DNA interpreted as SST/winter monsoon indicator in core Indus 11C (Giosan et al., 2018). c) Ti/Al  
ratios indicating fluvial sediment input, d) EM3 based on end-member modelling analyses of grain sizes as an indicator for aeolian  
input (Burdanowitz et al., 2019) and fe lithogenic mass accumulation rates (LIT MAR) of SO90-63KA. e)  $\delta^{18}\text{O}$  of *Globigerinoides*  
240 ruber of SO90-63KA (Staubwasser et al., 2003). gf latitudinal insolation gradients (LIG) of summer (JJA: 30°N – 0°, black line) and  
winter (DJF: 60°N – 30°N, blue line) and summer insolation at 30°N (dashed red line) (Laskar et al., 2004). hg stacked North  
Atlantic hematite stained grains (HSG) as representing drift-ice in record from core MC52-V29191 with so called “Bond events”  
(numbered red and grey bars) (Bond et al., 2001). Thick black lines in a) and c) indicates five-point running averages.

~~Thereafter, at~~From about 4.6 ka BP onwards a cooling period ~~occurred which was punctuated by~~began with a moderate initial  
cooling event at 4.4 ka BP and a rapid SST decrease of about 1°C between 4.2 and 3.8 ka BP. Foraminiferal analyses of the  
same core imply ~~lower-decreased~~ Indus river discharge and increased IWM mixing between 4.5 – 4.25 ka BP (Giesche et al.,  
2019; Staubwasser et al., 2003) which coincides with enhanced eolian input (EM3, Figure 2~~Figure 2~~). Giesche et al. (2019)  
245 ~~found-postulated~~ a weakening of the IWM strength from 4.1 to 3.9 ka BP and a period of cooling between 3.7 and 3.3 ka BP  
indicated by ~~using~~ salinity sensitive foraminera as proxies. This partly contradicts our SST record. However, the  $\delta^{18}\text{O}$  record  
of *G. ruber*, mainly interpreted as a salinity signal, is also affected by water temperature (Giesche et al., 2019; Staubwasser et  
al., 2003). If the  $\delta^{18}\text{O}$  signals were interpreted as a temperature rather than a salinity signal, Giesche et al. (2019) argued that  
the observed increase of the  $\delta^{18}\text{O}$  around 4.1 ka BP would be consistent with a surface water cooling of about 1°C. This  
250 interpretation matches our alkenone-based SST reconstruction at that time. Our record shows that lower SSTs ~~along with~~  
~~decreasing fluvial input~~ prevaileding until 3.0 ka BP and occurred along with decreased fluvial and aeolian input as shown by  
the Ti/Al ratios and EM3. The marked cooling between 4.6 and 3.0 is widespread in the Arabian Sea and on the Indian  
subcontinent (Gaye et al., 2018; Zhao et al., 2017) and coincides with the rise and fall of the Indus civilization (Wright et al.,  
2008).

255 Overall, long-term trends in the Ti/Al ratio and LIT MAR are driven by a strong ISM influence during the Early Holocene and  
a gradual increase (decrease) of STWJ (ISM) influence from the Mid- to Late Holocene. The aridification trend from the Mid-  
to Late Holocene led to a generally decreased fluvial input during the Late Holocene (Burdanowitz et al., 2019). The last ca.  
3.0 ka in the record are characterized by high and variable SSTs, LIT MAR, and increasing aeolian inputs, ~~while-whereas~~  
fluvial inputs of the Makran rivers decrease (Burdanowitz et al., 2019). In contrast to the generally smoothly decreasing river  
260 discharges, the higher SST and especially LIT MAR and aeolian inputs, reveal pronounced variations (Figure 2~~Figure 2~~).  
Furthermore, short episodes of high fluvial sediment discharge interrupt the general trend of decreasing fluvial input. The  
variations appear to be associated with a 1425 cycle, similar to the Bond events, that is superimposed on this long-term trend  
(Burdanowitz et al., 2019). We posit, that the combination of pronounced soil erosion, flood events due to precipitation events,  
and strong winds during arid phases led to the high LIT MAR in the Late Holocene part of our record. These changes are  
265 mainly associated with decreasing ISM activity as well as increasing wind in the source areas of the aeolian material and  
variable IWM strength (Böll et al., 2014, 2015; Burdanowitz et al., 2019; Ivory and Lézine, 2009; Lückge et al., 2001). The  
main forcing mechanisms have been suggested to be the southward shift of the ITCZ due to decreasing summer solar insolation,  
thermal land-ocean contrast and/or teleconnection to mid-high-latitude NH climate probably via the STWJ (Böll et al., 2014,


2015; Burdanowitz et al., 2019; Fleitmann et al., 2007; Giosan et al., 2018; Lückge et al., 2001; Mohtadi et al., 2016; Munz et al., 2015, 2017). A stronger impact of NH climate is indicated by high accumulation rates during Bond events 0-2 (Figure 2): The up to 4-fold increase of LIT MAR during the Late Holocene Bond events marks a system shift towards a strong response to mid-high latitude NH climate via the STWJ. This increased precipitation in winter, but decreased precipitation during the summer. Furthermore, this change in the seasonal precipitation pattern enhanced soil erosion due to stronger erosive forces of rivers in largely derelict farmlands exposed to desertification due to an overall increased aridification trend (Giosan et al., 2012, 2018; Wright et al., 2008).

#### 34.2 Transition period from low latitudinal to increased mid-high latitudinal influence on the Makran coast

The smooth and long-term decrease of the summer insolation at 30°N after ca. 11 ka BP is an important driver of climate and vegetation changes in the IM and westerlies realm (Fleitmann et al., 2007; Herzschuh, 2006). It does, however, not explain the marked cooling event between about 4.6 and 3 ka BP and the subsequently unstable warmer period seen in our record. The smoothly decreasing summer insolation cannot be the sole driver of these delayed and fast-paced environmental changes even if one considers that the maximum ISM intensity lags the maximum summer insolation by about 3 ka (Ansari and Vink, 2007; Fleitmann et al., 2007; Leipe et al., 2014; Reichert et al., 1998; Zhang et al., 2018; Zhao et al., 2017). A decisive influence instead may be the latitudinal insolation gradient (LIG) that triggers changes of the atmospheric pressure gradient, heat transport and determines the strength and position of regional atmospheric circulation (Bosmans et al., 2012; Davis and Brewer, 2009; Lee and Wang, 2014; Mohtadi et al., 2016; Wang et al., 2017). This which has so far received little attention in paleostudies (Clemens and Prell, 2003). The cross-equatorial insolation gradients play an important role for inter-hemispheric moisture transport and the extent of the Hadley cell and summer cross-equatorial insolation gradients may drive glacial-interglacial variability (Bosmans et al., 2015; Reichert, 1997). However, (Davis and Brewer, 2009) argued that intra-hemispheric summer LIG and the latitudinal temperature gradient (LTG) also influence the strength of the monsoon system and its poleward displacement and, respectively, that of the Hadley cell, because the LIG leads to differential heating of the polar and tropical regions. For instance, ISM variability over the Tibetan Plateau is found to be more sensitive to the summer LIG between 44°N and the Himalayan barrier at 30°N than to the cross-equatorial LIG (Ramisch et al., 2016). A study from the West African monsoon region shows a strong link of intra-hemispheric summer LIG and the West African monsoon (Kuechler et al., 2018). Further, different seasonal changes of the LIG (summer LIG varies with the obliquity and winter LIG with precession) have a strong response to climatic changes during the past (Davis and Brewer, 2009). Winter and summer LIG are equally important drivers for the strengths, positions and areal extents of the STWJ and the ISM (Mohtadi et al., 2016; Ramisch et al., 2016). As the catchment area of the core location is influenced by both, ISM (summer) and STWJ (winter), the winter LIG of the mid-high latitudes has to be considered. (Fallah et al., 2017) found that the STWJ over Iran ranged between 35°N and about 54°N during the last 6 ka. During the winter months (here defined as December – February), the winter LIG (here defined as the gradient between 60°N-30°N) determines the southernmost position of the STWJ and the frequency of the WDs (Fallah et al., 2017). The summer LIG (here defined as the gradient between 30°N – equator as the Himalayan range acts

~~an barrier, June-August) during the summer months (here defined as June-August)~~ affects the strength and extension of the ISM over ~~larger-centennial to millennial~~ timescales (Ramisch et al., 2016). The antagonism and relative magnitudes of the winter and summer LIGs influence the duration of the ISM, as well as STWJ and the frequency of the WDs. Therefore, changes  
305 in LIG have probably modulated seasonal length of the IM in the course of the Holocene.

The summer LIG reached its maximum between 8 and 7 ka BP, whereas the winter LIG peaked during the last 1 ka (~~Figure 2~~~~Figure 2~~). The net effect of ISM weakening and STWJ strengthening, in combination with dry periods on the adjacent continent to the north, resulted in more frequent dust storms that increased aeolian input to the Arabian Sea during the Late Holocene (Burdanowitz et al., 2019). We suggest that the time period between 4.6 and 3 ka BP marks a transition from an  
310 ISM-dominated climate system towards one which is more influenced by the STWJ. This strengthened the teleconnection between the Makran coastal ~~climate~~ and climate variability in the North Atlantic, which is ~~best illustrated~~~~most visible~~ by the link between the LIT MAR record and the Bond events ~~during the last 3.5 ka -since the end of the time period and associated fall the Indus civilization-~~(~~Figure 2~~~~Figure 2~~). A wavelet analysis of LIT MAR data (~~Figure 3~~) ~~reveals further evidence for~~~~illustrates the shift of influences during~~ the transition period with a ~~distinct~~ change of periodicities between ca 4.2 and 3.5  
315 ka BP (~~Figure 3~~). ~~The last 3.5 ka of the LIT MAR record is characterized by periodicities of about 512 to 1024 years, whereas periodicities of about 2048 to 3000 years were dominant between 7.5 to 3.5 ka BP. This shift indicates that the Makran region was more influenced by the mid-high latitude climate during the past 3.5 ka BP than during the Mid-Holocene. During the Mid-Holocene, in contrast, the ISM and the low latitude climate were dominant drivers in the Makran region and suppressed the influence of the Bond events.~~ These findings corroborate the hypothesis of a teleconnection between the mid-high-latitude  
320 NH climate via the SWTJ and IWM in the NE Arabian Sea region (Böll et al., 2015; Munz et al., 2015, 2017). On a more regional scale, increasing influence of the STWJ coinciding with the transition phase is documented in ~~several~~ marine and terrestrial records from the Arabian Sea and Indian subcontinent (Giosan et al., 2018; Hou et al., 2016). Climate simulations for the last 6 ka over Iran have shown a southward shift of the STWJ during the Late Holocene due to increasing winter insolation (Fallah et al., 2017), suggesting a “tipping point” around 3-4 ka BP in the interaction of ISM and STWJ with waxing  
325 influence of the STWJ towards the present day. The “tipping point” corresponds to so-called neoglacial anomalies in the Makran region, when a strong winter monsoon and weak interhemispheric temperature contrast (30°N-30°S) ~~were suggested to have been~~~~was~~ accompanied by a decrease in ISM precipitation (Giosan et al., 2018). Significant temperature oscillations over the last 3 ka were also recorded in the continental record of Lake Qinghai, but are commonly interpreted as an amplified response to volcanic and/or solar forcing (Hou et al., 2016). The manifestation of a meridional shift in STWJ is not restricted  
330 to the Arabian Sea region. A southward displacement of the STWJ during the Late Holocene has also been reported from the southern Alps in Europe, the Japan Sea and climate simulations for East Asia (Kong et al., 2017; Nagashima et al., 2013; Wirth et al., 2013), indicating teleconnections on a hemispheric spatial scale.



335 **Figure 3: Change of the periodicity in the lithogenic mass accumulation rates (LIT MAR) record. Wavelet power spectrum for LIT MAR of SO90-63KA using a Matlab Wavelet script (Torrence and Compo, 1998). The black line indicates the cone of influence, the dashed black line the 95% significance level. The periodicities clearly change in the time from 4.2 ka to around 3.5 ka BP and 4.2 ka BP from ca. 2360 years during the Early and Mid-Holocene to ca. 590-710 and 1400 years during the Late Holocene.**

Several studies have suggested ENSO as an important driver modulating the climate in the IM realm during the Holocene (Banerji et al., 2020; Munz et al., 2017; Prasad et al., 2014, 2020; Srivastava et al., 2017). In general, an ENSO event is associated with reduced ISM precipitation (Gadgil, 2003). Even though the ENSO events have apparently increased since-over 340 the last ca. 5000 years (Moy et al., 2002), roughly matching the timing of our observed transition period, we cannot find a correlation of our LIT MAR record with periods of strong ENSO events (not shown). Other studies suggested have shown that the relationship between IM precipitation and ENSO is not linear. Observations (1880 – 2005) imply suggest that less than half of the severe droughts occurred during El Niño years, while the amount of precipitation was normal or above normal during 345 other El Niño years (Rajeevan and Pai, 2007). A possible reason for this mismatch is the strong effect of the Indian Ocean Dipole (IOD). It can act as an amplifier or suppressor of the ENSO influence in the ISM realm (Ashok et al., 2001, 2004; Behera and Ratnam, 2018; Krishnaswamy et al., 2015; Ummenhofer et al., 2011). A positive IOD can counteract even a strong El Niño forcing leading to “normal” precipitation amounts (Ashok et al., 2004; Krishnaswamy et al., 2015; Ummenhofer et al., 2011). A recent study further suggested that a positive IOD shows a tripolar pattern over the IM realm, with above normal


350 precipitation in central India and below normal precipitation in northern and southern India (Behera and Ratnam, 2018). In contrast, a negative IOD creates a zonal pattern with above normal precipitation in western India and below normal precipitation in eastern India (Behera and Ratnam, 2018). In addition, these authors have linked a positive IOD to warmer SST in the northern AS and vice versa. However, ~~we can only speculate how strong its~~ impact ~~of the IOD was~~ during the transition period ~~is unknown~~ due to ~~the~~ lack of paleorecords ~~and reconstructions~~ ~~reconstructing the IOD~~.


### 355 **34.3 The environmental-anthropogenic interaction during the mid-Late Holocene**

Widespread anthropogenic land use, such as deforestation or rice irrigation, and associated environmental change is recorded during the Early-Mid Holocene transition period (Petrie et al., 2017; Petrie and Bates, 2017). Especially rice irrigation, starting around 6-4 ka ago, has been suggested to have increased the atmospheric CH<sub>4</sub> concentration since about 5 ka (Figure 4) and thus had a small, but emblematic impact on climate (Ruddiman and Thomson, 2001). The rise of the Indus civilization began  
360 around 5.2 ka BP when general aridification decreased river discharges (Giosan et al., 2012). Since ca. 4 ka BP, increasing winter precipitation enabled the development of agriculture along the Indus valley due to less intense floods favoring urbanization (Giosan et al., 2012; Wright et al., 2008). The urban Harappan civilization collapsed towards the end of the climatic transition period when less frequent annual flooding occurred, which was the basis of agriculture. Most of the people migrated from the Indus valley to the Himalaya plains, whereas the remaining society turned into a post-urban society (Clift  
365 and Plumb, 2008; Giosan et al., 2012, 2018; Possehl, 1997).

The average chain length (ACL) of plant-wax derived long-chain *n*-alkanes vary in response to vegetation type and/or climatic conditions (Bush and McInerney, 2013; Carr et al., 2014; Meyers and Ishiwatari, 1993; Rao et al., 2011; Vogts et al., 2009). For instance, African savanna plants produce, on average, longer chain *n*-alkanes than rainforest plants (Rommerskirchen et al., 2006; Vogts et al., 2009). In the region of Indus civilization, natural dominated until agriculture proliferated (Giosan et al.,  
370 2012) driven by the changing precipitation pattern due to the interplay of the position and strength of the STWJ and ISM (Ansari and Vink, 2007) (Figure 4). During the Mid-Holocene warm period, high SSTs are thought to have fueled the moisture transport to the adjacent continent, which is in line with our SST record (~~Figure 2~~~~Figure-2~~). High average summer precipitation and/or more uniformly distributed precipitation during the different seasons favored the development of grasslands in southern Pakistan (Ansari and Vink, 2007; Ivory and Lézine, 2009). This may ~~be~~ reflected in a decrease in the ~~ACL of *n*-alkane~~  
375 ~~homologues C<sub>27</sub>—C<sub>33</sub> (ACL<sub>27-33</sub>)~~ between ca. 7 to 5 ka BP (Figure 4). The ACL<sub>27-33</sub> increased after the transition period, although the amplitude of ACL<sub>27-33</sub> variation is not very high, and decreasing river discharges and increasing aeolian input (~~Figure 2~~~~Figure-2~~) indicate prevailing dry conditions after 3 ka BP. It is impossible to separate effects of anthropogenic modified vegetation or climatic driven vegetation changes on ACL<sub>27-33</sub> fluctuations ~~s~~ as the ACL<sub>27-33</sub> reflects to whole vegetation community. However, ~~but~~ the ACL<sub>27-33</sub> increase suggests that humans started to influence the vegetation in the  
380 sediment source region and led to the decoupling of the climate signal and the ACL<sub>27-33</sub>. This is in line with the coeval decoupling of the atmospheric CH<sub>4</sub> concentration and orbitally driven climate changes (Ruddiman and Thomson, 2001). These authors suggested that inefficient early rice farming caused atmospheric CH<sub>4</sub> concentration to increase at around 5 ka BP. Rice

agriculture began around 4 ka BP on the Indian subcontinent and near the Indus River region, adding to the atmospheric CH<sub>4</sub> concentration (Fuller et al., 2011). Further, climatic changes during the Mid-Holocene were suggested to have changed the agricultural system in the region to a “multi-cropping” (summer/winter crops) system (Petrie and Bates, 2017; Wright et al., 2008). This clearly shows that human activities not only modified the vegetation, but also started to impact climate during this time. In a modelling study land cover change associated with prehistoric cultures modified not only local climate, but climate on a broader hemispheric scale via teleconnections (Dallmeyer and Claussen, 2011). Since the transformation of natural into cultivated landscapes favors soil erosion, it is likely that this early human land-use change intensified the impact of the NAO and Bond events on the sedimentation in the NE AS during the late Holocene. The latter in turn documents a clear shift of the climatic system that was associated with the collapse of and deep crises of Late Bronze Age societies in the Mediterranean, Middle East and East Asia.





395 Figure 4: Environmental and anthropogenic interactions illustrated by proxy records. a) different phases of the Indus civilization (Giosan et al., 2012) (L=Late, M=Mature, E=Early) and time periods of ~~our~~ interpreted natural and anthropogenic modified  
 400 vegetation (green bar and arrow) of core SO90-83KA in comparison to b) average chain length (ACL) of the *n*-alkanes homologues 27-33 (black line, 5 point running mean) and c) LIT MAR of SO90-63KA as well as b. The ISM humidity index (Herzschuh, 2006) (blue), CH<sub>4</sub> concentration (red) reconstruction based on Greenland ice cores (Blunier et al., 1995) are also depicted in panel b and d) Bond events (number eds and grey bars) based on the HSG drift-ice record from the North Atlantic (Bond et al., 2001).

The climate transition period between 4.6 and 3 ka BP impacted not only the Indus civilization but also civilizations in Mesopotamia, the Eastern Mediterranean and China (Giosan et al., 2012, 2018; Kaniewski et al., 2013; Liu and Feng, 2012; Wang et al., 2016; Weiss et al., 1993). Today these regions are located slightly north of the northern border of the region influenced by the Asian monsoon (Wang et al., 2016), but where probably within the monsoon realm prior to the

405 ~~transition~~transition period. The series of climatically triggered collapses started at about 4.4 ka in China and reached Mesopotamia and the Eastern Mediterranean between 3.1 and 3.2 ka BP (Giosan et al., 2012, 2018; Kaniewski et al., 2013; Liu and Feng, 2012; Wang et al., 2016). This indicates that this transition period was characterized by a southward shift of the ISM realm, which in combination with socio-economic responses turned into catastrophic changes for Late Bronze Age societies that thrived in the ISM/ STWJ transition zone.

## 410 ~~4.5~~Conclusions

Our study of reconstructed SST, LIT MAR and vegetation changes at the Makran coastal region suggests a climatic transition period between 4.6 – 3 ka BP from a low latitude to increased mid-high latitude influence via the STWJ. Prior to this transition period the region was strongly influenced by the ISM and the ITCZ. Our LIT MAR record shows an up to 4-fold increase after the transition period coincident with Bond events in the North Atlantic. We argue that after this period the North Atlantic signals of NAO and Bond events are transmitted to the Makran coast region via the southward shifted STWJ. This supports an earlier modelling study (Fallah et al., 2017) suggesting a southward shift of the STWJ and more winter precipitation near our study area in Iran between 4 and 3 ka BP, which the authors mark as a “tipping point” in that region. Besides a long term drying trend since the Mid-Holocene, this transition period and the associated change in the precipitation pattern (winter vs. summer precipitation) affected the settlements and agriculture of the Indus civilization (Giosan et al., 2012, 2018; Petrie and Bates, 2017). The temporal coincidence gives rise to the hypothesis that humans themselves became environmental drivers prior to the onset of the transition period (Fuller et al., 2011; Ruddiman and Thomson, 2001), instead of only being passive victims of climate change. Whether human impacts suffice to influence the global climate is an ongoing debate, but it is very likely that land cover changes associated with prehistoric cultures modified at least local climate (Dallmeyer and Claussen, 2011). Consequences included a weakened local hydrological cycle that favored desertification, especially in semi-arid regions.

### Data availability

The alkenone-based SST, the LIT MAR, the extended grain size used for the EM3 and *n*-alkane data sets of core SO90-63KA ~~will be~~are available at PANGAEA data repository (<https://doi.pangaea.de/10.1594/PANGAEA.925909>). The previous grain size and elemental data of SO-90KA are available at PANGAEA data repository (<https://doi.pangaea.de/10.1594/PANGAEA.900973>).

## Author contribution

BG and KE designed the study. NB analyzed LIT MAR, alkenones and *n*-alkanes. TR analyzed grain sizes. NB, BG and TR interpreted the data. NB prepared the manuscript with input from all co-authors.

## Competing interests.

435 The authors declare that they have no conflict of interest.

## Acknowledgements

We thank F. Langenberg, M. Metzke, C. Staschok, L. Hilbig, L. Meiritz and Y. Akkul for analytical support. S. Beckmann is thanked for the technical support for *n*-alkane and alkenone analyses. We thank ~~three~~ an anonymous reviewer ~~of a previous version~~ for comments that helped to improve the manuscript. The study was supported by the German Federal Ministry of Education and Research (BMBF) as part of the CAHOL (Central Asian Holocene Climate, project number 03G0864A), a subproject of the research program CAME II (Central Asia: Climatic Tipping Points & Their Consequences, project number 03G0863G). This study is a contribution to the Cluster of Excellence 'CLICCS - Climate, Climatic Change, and Society', contribution to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg.

## References

- 445 Ait Brahim, Y., Wassenburg, J. A., Sha, L., Cruz, F. W., Deininger, M., Sifeddine, A., Bouchaou, L., Spötl, C., Edwards, R. L. and Cheng, H.: North Atlantic Ice-Rafting, Ocean and Atmospheric Circulation During the Holocene: Insights From Western Mediterranean Speleothems, *Geophys. Res. Lett.*, 46(13), 7614–7623, doi:10.1029/2019GL082405, 2019.
- Andrulleit, H., Rogalla, U. and Stäger, S.: From living communities to fossil assemblages: Origin and fate of coccolithophorids in northern Arabian Sea, *Micropaleontology*, 50, suppl.(1), 5–21, 2004.
- 450 Anoop, A., Prasad, S., Krishnan, R., Naumann, R. and Dulski, P.: Intensified monsoon and spatiotemporal changes in precipitation patterns in the NW Himalaya during the early-mid Holocene, *Quat. Int.*, 313(Supplement C), 74–84, doi:https://doi.org/10.1016/j.quaint.2013.08.014, 2013.
- Ansari, M. H. and Vink, A.: Vegetation history and palaeoclimate of the past 30 kyr in Pakistan as inferred from the palynology of continental margin sediments off the Indus Delta, *Rev. Palaeobot. Palynol.*, 145(3), 201–216, doi:https://doi.org/10.1016/j.revpalbo.2006.10.005, 2007.
- 455 Ashok, K., Guan, Z. and Yamagata, T.: Impact of the Indian Ocean dipole on the relationship between the Indian monsoon rainfall and ENSO, *Geophys. Res. Lett.*, 28(23), 4499–4502, doi:10.1029/2001GL013294, 2001.
- Ashok, K., Guan, Z., Saji, N. H. and Yamagata, T.: Individual and Combined Influences of ENSO and the Indian Ocean Dipole

- on the Indian Summer Monsoon, *J. Clim.*, 17(16), 3141–3155, doi:10.1175/1520-0442(2004)017<3141:IACIOE>2.0.CO;2,  
460 2004.
- Avnimelech, Y., Ritvo, G., Meijer, L. E. and Kochba, M.: Water content, organic carbon and dry bulk density in flooded sediments, *Aquac. Eng.*, 25(1), 25–33, doi:https://doi.org/10.1016/S0144-8609(01)00068-1, 2001.
- Band, S., Yadava, M. G., Lone, M. A., Shen, C. C., Sree, K. and Ramesh, R.: High-resolution mid-Holocene Indian Summer Monsoon recorded in a stalagmite from the Kotumsar Cave, Central India, *Quat. Int.*, 479, 19–24,  
465 doi:10.1016/j.quaint.2018.01.026, 2018.
- Banerji, U. S., Arulbalaji, P. and Padmalal, D.: Holocene climate variability and Indian Summer Monsoon: An overview, *The Holocene*, 1–30, doi:10.1177/0959683619895577, 2020.
- Banse, K. and McClain, C. R.: Winter blooms of phytoplankton in the Arabian Sea as observed by the Coastal Zone Color Scanner, *Mar. Ecol. Prog. Ser.*, 34, 201–211, doi:10.3354/meps034201, 1986.
- 470 Behera, S. K. and Ratnam, J. V.: Quasi-asymmetric response of the Indian summer monsoon rainfall to opposite phases of the IOD, *Sci. Rep.*, 8(1), 123, doi:10.1038/s41598-017-18396-6, 2018.
- Blake, G. R. and Hartge, K. H.: Bulk Density, *Methods Soil Anal.*, 363–375, doi:doi:10.2136/sssabookser5.1.2ed.c13, 1986.
- Blunier, T., Chappellaz, J., Schwander, J., Stauffer, B. and Raynaud, D.: Variations in atmospheric methane concentration during the Holocene epoch, *Nature*, 374(6517), 46–49, doi:10.1038/374046a0, 1995.
- 475 Böll, A., Lückge, A., Munz, P., Forke, S., Schulz, H., Ramaswamy, V., Rixen, T., Gaye, B. and Emeis, K. C.: Late Holocene primary productivity and sea surface temperature variations in the northeastern Arabian Sea: Implications for winter monsoon variability, *Paleoceanography*, doi:10.1002/2013PA002579, 2014.
- Böll, A., Schulz, H., Munz, P., Rixen, T., Gaye, B. and Emeis, K. C.: Contrasting sea surface temperature of summer and winter monsoon variability in the northern Arabian Sea over the last 25ka, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 426, 10–  
480 21, doi:10.1016/j.palaeo.2015.02.036, 2015.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M. N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I. and Bonani, G.: Persistent solar influence on North Atlantic climate during the Holocene., *Science*, 294, 2130–2136, doi:10.1126/science.1065680, 2001.
- Bosmans, J. H. C., Drijfhout, S. S., Tuenter, E., Lourens, L. J., Hilgen, F. J. and Weber, S. L.: Monsoonal response to mid-  
485 holocene orbital forcing in a high resolution GCM, *Clim. Past*, 8(2), 723–740, doi:10.5194/cp-8-723-2012, 2012.
- Bosmans, J. H. C., Hilgen, F. J., Tuenter, E. and Lourens, L. J.: Obliquity forcing of low-latitude climate, *Clim. Past*, 11, 1335–1346, doi:10.5194/cp-11-1335-2015, 2015.
- Boyd, C. E.: *Bottom Soils, Sediment, and Pond Aquaculture*, Springerl., 1995.
- Burdanowitz, N., Gaye, B., Hilbig, L., Lahajnar, N., Lückge, A., Rixen, T. and Emeis, K. C.: Holocene monsoon and sea level-  
490 related changes of sedimentation in the northeastern Arabian Sea, *Deep. Res. Part II Top. Stud. Oceanogr.*, 166, 6–18, doi:10.1016/j.dsr2.2019.03.003, 2019.
- Bush, R. T. and McInerney, F. A.: Leaf wax *n*-alkane distributions in and across modern plants: Implications for paleoecology

- and chemotaxonomy, *Geochim. Cosmochim. Acta*, 117, 161–179, doi:10.1016/j.gca.2013.04.016, 2013.
- 495 Carr, A. S., Boom, A., Grimes, H. L., Chase, B. M., Meadows, M. E. and Harris, A.: Leaf wax *n*-alkane distributions in arid zone South African flora: Environmental controls, chemotaxonomy and palaeoecological implications, *Org. Geochem.*, 67, 72–84, doi:10.1016/j.orggeochem.2013.12.004, 2014.
- Cheng, H., Fleitmann, D., Edwards, R. L., Wang, X., Cruz, F. W., Auler, A. S., Mangini, A., Wang, Y., Kong, X., Burns, S. J. and Matter, A.: Timing and structure of the 8.2 kyr B.P. event inferred from  $\delta^{18}\text{O}$  records of stalagmites from China, Oman, and Brazil, *Geology*, 37(11), 1007–1010, doi:10.1130/G30126A.1, 2009.
- 500 Clemens, S. C. and Prell, W. L.: A 350,000 year summer-monsoon multi-proxy stack from the Owen Ridge, Northern Arabian Sea, *Mar. Geol.*, 201(1–3), 35–51, doi:10.1016/S0025-3227(03)00207-X, 2003.
- Clift, P. D. and Plumb, R. A.: *The Asian Monsoon: Causes, History and Effects*, Cambridge University Press, Cambridge., 2008.
- Dallmeyer, A. and Claussen, M.: The influence of land cover change in the Asian monsoon region on present-day and mid-  
505 Holocene climate, *Biogeosciences*, 8(6), 1499–1519, doi:10.5194/bg-8-1499-2011, 2011.
- Dallmeyer, A., Claussen, M., Wang, Y. and Herzschuh, U.: Spatial variability of Holocene changes in the annual precipitation pattern: a model-data synthesis for the Asian monsoon region, *Clim. Dyn.*, 40(11–12), 2919–2936, doi:10.1007/s00382-012-1550-6, 2013.
- Davis, B. A. S. and Brewer, S.: Orbital forcing and role of the latitudinal insolation/temperature gradient, *Clim. Dyn.*, 32(2–  
510 3), 143–165, doi:10.1007/s00382-008-0480-9, 2009.
- Dietze, E., Hartmann, K., Diekmann, B., IJmker, J., Lehmkuhl, F., Opitz, S., Stauch, G., Wünnemann, B. and Borchers, A.: An end-member algorithm for deciphering modern detrital processes from lake sediments of Lake Donggi Cona, NE Tibetan Plateau, China, *Sediment. Geol.*, 243, 169–180, doi:http://dx.doi.org/10.1016/j.sedgeo.2011.09.014, 2012.
- Dimri, A. P., Yasunari, T., Kotlia, B. S., Mohanty, U. C. and Sikka, D. R.: Indian winter monsoon: Present and past, *Earth-  
515 Science Rev.*, 163, 297–322, doi:10.1016/j.earscirev.2016.10.008, 2016.
- Dixit, Y., Hodell, D. A., Sinha, R. and Petrie, C. A.: Abrupt weakening of the Indian summer monsoon at 8.2 kyr B.P., *Earth Planet. Sci. Lett.*, 391, 16–23, doi:http://dx.doi.org/10.1016/j.epsl.2014.01.026, 2014.
- Durcan, J. A., Thomas, D. S. G. G., Gupta, S., Pawar, V., Singh, R. N. and Petrie, C. A.: Holocene landscape dynamics in the Ghaggar-Hakra palaeochannel region at the northern edge of the Thar Desert, northwest India, *Quat. Int.*, 501, 317–327,  
520 doi:10.1016/j.quaint.2017.10.012, 2019.
- ESRI: ArcGIS Desktop: Release 10.8. Redlands, CA: Environmental Systems Research Institute., 2019.
- Fallah, B., Sodoudi, S., Russo, E., Kirchner, I. and Cubasch, U.: Towards modeling the regional rainfall changes over Iran due to the climate forcing of the past 6000 years, *Quat. Int.*, 429(Part B), 119–128, doi:https://doi.org/10.1016/j.quaint.2015.09.061, 2017.
- 525 Fleitmann, D., Burns, S. J., Mangini, A., Mudelsee, M., Kramers, J., Villa, I., Neff, U., Al-Subbary, A. A., Buettner, A., Hippler, D. and Matter, A.: Holocene ITCZ and Indian monsoon dynamics recorded in stalagmites from Oman and Yemen

- (Socotra), *Quat. Sci. Rev.*, 26(1–2), 170–188, doi:10.1016/j.quascirev.2006.04.012, 2007.
- Forke, S., Rixen, T., Burdanowitz, N., Ramaswamy, V., Munz, P., Wilhelms-dick, D., Vogt, C., Kasten, S. and Gaye, B.: Sources of laminated sediments in the northeastern Arabian Sea off Pakistan and implications for sediment transport mechanisms during the late Holocene, *The Holocene*, 29(1), 130–144, doi:10.1177/0959683618804627, 2019.
- Fuller, D. Q., van Etten, J., Manning, K., Castillo, C., Kingwell-Banham, E., Weisskopf, A., Qin, L., Sato, Y.-I. and Hijmans, R. J.: The contribution of rice agriculture and livestock pastoralism to prehistoric methane levels: An archaeological assessment, *The Holocene*, 21(5), 743–759, doi:10.1177/0959683611398052, 2011.
- Gadgil, S.: The Indian Monsoon and its variability, *Annu. Rev. Earth Planet. Sci.*, 31(1), 429–467, doi:10.1146/annurev.earth.31.100901.141251, 2003.
- Gaye, B., Böll, A., Segschneider, J., Burdanowitz, N., Emeis, K.-C., Ramaswamy, V., Lahajnar, N., Lückge, A. and Rixen, T.: Glacial–interglacial changes and Holocene variations in Arabian Sea denitrification, *Biogeosciences*, 15(2), 507–527, doi:10.5194/bg-15-507-2018, 2018.
- Giesche, A., Staubwasser, M., Petrie, C. A. and Hodell, D. A.: Indian winter and summer monsoon strength over the 4.2 ka BP event in foraminifer isotope records from the Indus River delta in the Arabian Sea, *Clim. Past*, 15(1), 73–90, doi:10.5194/cp-15-73-2019, 2019.
- Giosan, L., Clift, P. D., Macklin, M. G., Fuller, D. Q., Constantinescu, S., Durcan, J. A., Stevens, T., Duller, G. A. T., Tabrez, A. R., Gangal, K., Adhikari, R., Alizai, A., Filip, F., VanLaningham, S. and Syvitski, J. P. M.: Fluvial landscapes of the Harappan civilization, *Proc. Natl. Acad. Sci. U. S. A.*, 109(26), E1688–E1694, doi:10.1073/pnas.1112743109, 2012.
- Giosan, L., Orsi, W. D., Coolen, M., Wuchter, C., Dunlea, A. G., Thirumalai, K., Munoz, S. E., Clift, P. D., Donnelly, J. P., Galy, V. and Fuller, D. Q.: Neoglacial climate anomalies and the Harappan metamorphosis, *Clim. Past*, 14(11), 1669–1686, doi:10.5194/cp-14-1669-2018, 2018.
- Goslin, J., Fruergaard, M., Sander, L., Gałka, M., Menviel, L., Monkenbusch, J., Thibault, N. and Clemmensen, L. B.: Holocene centennial to millennial shifts in North-Atlantic storminess and ocean dynamics, *Sci. Rep.*, 8(1), 12778, doi:10.1038/s41598-018-29949-8, 2018.
- Gourlan, A. T., Albarede, F., Achyuthan, H. and Campillo, S.: The marine record of the onset of farming around the Arabian Sea at the dawn of the Bronze Age, *The Holocene*, 30(6), 878–887, doi:10.1177/0959683620902218, 2020.
- Herzschuh, U.: Palaeo-moisture evolution in monsoonal Central Asia during the last 50,000 years, *Quat. Sci. Rev.*, 25(1–2), 163–178, doi:10.1016/j.quascirev.2005.02.006, 2006.
- Hou, J., Huang, Y., Zhao, J., Liu, Z., Colman, S. and An, Z.: Large Holocene summer temperature oscillations and impact on the peopling of the northeastern Tibetan Plateau, *Geophys. Res. Lett.*, 43(3), 1323–1330, doi:10.1002/2015GL067317, 2016.
- Hunt, K. M. R., Turner, A. G. and Shaffrey, L. C.: The evolution, seasonality and impacts of western disturbances, *Q. J. R. Meteorol. Soc.*, 144(710), 278–290, doi:10.1002/qj.3200, 2018.
- Hussain, B. A., Mir, H. and Afzal, M.: Analysis of dust storms frequency over Pakistan during 1961–2000, *Pakistan J. Meteorol.*, 2(3), 2005.

- Ivory, S. J. and Lézine, A. M.: Climate and environmental change at the end of the Holocene Humid Period: A pollen record off Pakistan, *Comptes Rendus - Geosci.*, 341(8–9), 760–769, doi:10.1016/j.crte.2008.12.009, 2009.
- Kaniewski, D., Van Campo, E., Guiot, J., Le Burel, S., Otto, T. and Baeteman, C.: Environmental Roots of the Late Bronze Age Crisis, *PLoS One*, 8(8), 1–10, doi:10.1371/journal.pone.0071004, 2013.
- 565 Kaskaoutis, D. G., Rashki, A., Houssos, E. E., Goto, D. and Nastos, P. T.: Extremely high aerosol loading over Arabian Sea during June 2008: The specific role of the atmospheric dynamics and Sistan dust storms, *Atmos. Environ.*, 94, 374–384, doi:10.1016/j.atmosenv.2014.05.012, 2014.
- Kaskaoutis, D. G., Rashki, A., Francois, P., Dumka, U. C., Houssos, E. E. and Legrand, M.: Meteorological regimes modulating dust outbreaks in southwest Asia: The role of pressure anomaly and Inter-Tropical Convergence Zone on the 1–3 July 2014 case, *Aeolian Res.*, 18(Supplement C), 83–97, doi:https://doi.org/10.1016/j.aeolia.2015.06.006, 2015.
- 570 Klus, A., Prange, M., Varma, V., Bruno Tremblay, L. and Schulz, M.: Abrupt cold events in the North Atlantic Ocean in a transient Holocene simulation, *Clim. Past*, 14(8), 1165–1178, doi:10.5194/cp-14-1165-2018, 2018.
- Kotlia, B. S., Singh, A. K., Joshi, L. M. and Dhaila, B. S.: Precipitation variability in the Indian Central Himalaya during last ca. 4,000 years inferred from a speleothem record: Impact of Indian Summer Monsoon (ISM) and Westerlies, *Quat. Int.*,
- 575 371(Supplement C), 244–253, doi:https://doi.org/10.1016/j.quaint.2014.10.066, 2015.
- Krishnaswamy, J., Vaidyanathan, S., Rajagopalan, B., Bonell, M., Sankaran, M., Bhalla, R. S. and Badiger, S.: Non-stationary and non-linear influence of ENSO and Indian Ocean Dipole on the variability of Indian monsoon rainfall and extreme rain events, *Clim. Dyn.*, 45(1), 175–184, doi:10.1007/s00382-014-2288-0, 2015.
- Kuechler, R. R., Dupont, L. M. and Schefuß, E.: Hybrid insolation forcing of Pliocene monsoon dynamics in West Africa,
- 580 *Clim. Past*, 14(1), 73–84, doi:10.5194/cp-14-73-2018, 2018.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M. and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astron. Astrophys.*, 428(1), 261–285, doi:10.1051/0004-6361:20041335, 2004.
- Lauterbach, S., Witt, R., Plessen, B., Dulski, P., Prasad, S., Mingram, J., Gleixner, G., Hettler-Riedel, S., Stebich, M., Schnetger, B., Schwab, A. and Schwarz, A.: Climatic imprint of the mid-latitude Westerlies in the Central Tian Shan of
- 585 Kyrgyzstan and teleconnections to North Atlantic climate variability during the last 6000 years, *The Holocene*, 24(8), 970–984, doi:10.1177/0959683614534741, 2014.
- Lee, J.-Y. and Wang, B.: Future change of global monsoon in the CMIP5, *Clim. Dyn.*, 42(1), 101–119, doi:10.1007/s00382-012-1564-0, 2014.
- Leipe, C., Demske, D. and Tarasov, P. E.: A Holocene pollen record from the northwestern Himalayan lake Tso Moriri: Implications for palaeoclimatic and archaeological research, *Quat. Int.*, 348, 93–112, doi:https://doi.org/10.1016/j.quaint.2013.05.005, 2014.
- Liu, F. and Feng, Z.: A dramatic climatic transition at ~4000 cal. yr BP and its cultural responses in Chinese cultural domains, *The Holocene*, 22(10), 1181–1197, doi:10.1177/0959683612441839, 2012.
- Lückge, A., Dose-Rolinski, H., Khan, A. ., Schulz, H. and von Rad, U.: Monsoonal variability in the northeastern Arabian

- 595 Sea during the past 5000years: geochemical evidence from laminated sediments, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 167(3), 273–286, doi:10.1016/S0031-0182(00)00241-8, 2001.
- Lückge, A., Reinhardt, L., Andruleit, H., Doose-Rolinski, H., von Rad, U., Schulz, H. and Treppke, U.: Formation of varve-like laminae off Pakistan: decoding 5 years of sedimentation, *Geol. Soc. London, Spec. Publ.*, 195(1), 421–431, doi:10.1144/GSL.SP.2002.195.01.23, 2002.
- 600 Madhupratap, M., Kumar, S. P., Bhattathiri, P. M. A., Kumar, M. D., Raghukumar, S., Nair, K. K. C. and Ramaiah, N.: Mechanism of the biological response to winter cooling in the northeastern Arabian Sea, *Nature*, 384(6609), 549–552, doi:10.1038/384549a0, 1996.
- Meyers, P. A. and Ishiwatari, R.: Lacustrine organic geochemistry—an overview of indicators of organic matter sources and diagenesis in lake sediments, *Org. Geochem.*, 20(7), 867–900, doi:10.1016/0146-6380(93)90100-P, 1993.
- 605 Mohtadi, M., Prange, M. and Steinke, S.: Palaeoclimatic insights into forcing and response of monsoon rainfall, *Nature*, 533(7602), 191–199, doi:10.1038/nature17450, 2016.
- Moy, C. M., Seltzer, G. O., Rodbell, D. T. and Anderson, D. M.: Variability of El Niño/Southern Oscillation activity at millennial timescales during the Holocene epoch., *Nature*, 420(6912), 162–5, doi:10.1038/nature01194, 2002.
- Munz, P. M., Siccha, M., Lückge, A., Böll, A., Kucera, M. and Schulz, H.: Decadal-resolution record of winter monsoon intensity over the last two millennia from planktic foraminiferal assemblages in the northeastern Arabian Sea, *The Holocene*, doi:10.1177/0959683615591357, 2015.
- 610 Munz, P. M., Lückge, A., Siccha, M., Böll, A., Forke, S., Kucera, M. and Schulz, H.: The Indian winter monsoon and its response to external forcing over the last two and a half centuries, *Clim. Dyn.*, 49(5), 1801–1812, doi:10.1007/s00382-016-3403-1, 2017.
- 615 Pease, P. P., Tchakerian, V. P. and Tindale, N. W.: Aerosols over the Arabian Sea: geochemistry and source areas for aeolian desert dust, *J. Arid Environ.*, 39(3), 477–496, doi:10.1006/jare.1997.0368, 1998.
- Petrie, C. A. and Bates, J.: ‘Multi-cropping’, Intercropping and Adaptation to Variable Environments in Indus South Asia, *J. World Prehistory*, 30(2), 81–130, doi:10.1007/s10963-017-9101-z, 2017.
- Petrie, C. A., Singh, R. N., Bates, J., Dixit, Y., French, C. A. I., Hodell, D. A., Jones, P. J., Lancelotti, C., Lynam, F., Neogi, S., Pandey, A. K., Parikh, D., Pawar, V., Redhouse, D. I. and Singh, D. P.: Adaptation to Variable Environments, Resilience to Climate Change: Investigating Land, Water and Settlement in Indus Northwest India, *Curr. Anthropol.*, 58(1), 1–30, doi:10.1086/690112, 2017.
- 620 Possehl, G. L.: The transformation of the Indus Civilization, *J. World Prehistory*, 11(4), 425–472, doi:10.1007/BF02220556, 1997.
- 625 Prah, F. G., Muehlhausen, L. A. and Zahnle, D. L.: Further evaluation of long-chain alkenones as indicators of paleoceanographic conditions, *Geochim. Cosmochim. Acta*, 52(9), 2303–2310, doi:10.1016/0016-7037(88)90132-9, 1988.
- Prasad, S. and Enzel, Y.: Holocene paleoclimates of India, *Quat. Res.*, 66(3), 442–453, doi:http://dx.doi.org/10.1016/j.yqres.2006.05.008, 2006.

- Prasad, S., Anoop, A., Riedel, N., Sarkar, S., Menzel, P., Basavaiah, N., Krishnan, R., Fuller, D., Plessen, B., Gaye, B., Röhl, U., Wilkes, H., Sachse, D., Sawant, R., Wiesner, M. G. and Stebich, M.: Prolonged monsoon droughts and links to Indo-Pacific warm pool: A Holocene record from Lonar Lake, central India, *Earth Planet. Sci. Lett.*, 391, 171–182, doi:<https://doi.org/10.1016/j.epsl.2014.01.043>, 2014.
- Prasad, S., Marwan, N., Eroglu, D., Goswami, B., Mishra, P. K., Gaye, B., Anoop, A., Basavaiah, N., Stebich, M. and Jehangir, A.: Holocene climate forcings and lacustrine regime shifts in the Indian summer monsoon realm, *Earth Surf. Process. Landforms*, doi:10.1002/esp.5004, 2020.
- von Rad, U., Schulz, H., Khan, A. A., Ansari, M., Berner, U., Čepek, P., Cowie, G., Dietrich, P., Erlenkeuser, H., Geyh, M., Jennerjahn, T., Lückge, A., Marchig, V., Riech, V., Rösch, H., Schäfer, P., Schulte, S., Sirocko, F., Tahir, M. and Weiss, W.: Sampling the oxygen minimum zone off Pakistan: glacial-interglacial variations of anoxia and productivity (preliminary results, sonne 90 cruise), *Mar. Geol.*, 125(1), 7–19, doi:[http://dx.doi.org/10.1016/0025-3227\(95\)00051-Y](http://dx.doi.org/10.1016/0025-3227(95)00051-Y), 1995.
- von Rad, U., Khan, A. A., Berger, W. H., Rammelmair, D. and Treppke, U.: Varves, turbidites and cycles in upper Holocene sediments (Makran slope, northern Arabian Sea), *Geol. Soc. London, Spec. Publ.*, 195(1), 387–406, doi:10.1144/GSL.SP.2002.195.01.21, 2002.
- Rajeevan, M. and Pai, D. S.: On the El Niño-Indian monsoon predictive relationships, *Geophys. Res. Lett.*, 34(4), doi:10.1029/2006GL028916, 2007.
- Ramisch, A., Lockot, G., Haberzettl, T., Hartmann, K., Kuhn, G., Lehmkuhl, F., Schimpf, S., Schulte, P., Stauch, G., Wang, R., Wünnemann, B., Yan, D., Zhang, Y. and Diekmann, B.: A persistent northern boundary of Indian Summer Monsoon precipitation over Central Asia during the Holocene., *Sci. Rep.*, 6, 25791, doi:10.1038/srep25791, 2016.
- Rao, Z., Wu, Y., Zhu, Z., Jia, G. and Henderson, A.: Is the maximum carbon number of long-chain n-alkanes an indicator of grassland or forest? Evidence from surface soils and modern plants, *Chinese Sci. Bull.*, 56, 1714–1720, doi:10.1007/s11434-011-4418-y, 2011.
- Rashki, A., Kaskaoutis, D. G., Rautenbach, C. J. deW., Eriksson, P. G., Qiang, M. and Gupta, P.: Dust storms and their horizontal dust loading in the Sistan region, Iran, *Aeolian Res.*, 5, 51–62, doi:<http://dx.doi.org/10.1016/j.aeolia.2011.12.001>, 2012.
- Rawat, S., Gupta, A. K., Sangode, S. J., Srivastava, P. and Nainwal, H. C.: Late Pleistocene–Holocene vegetation and Indian summer monsoon record from the Lahaul, Northwest Himalaya, India, *Quat. Sci. Rev.*, 114, 167–181, doi:10.1016/j.quascirev.2015.01.032, 2015.
- Reichart, G.-J.: Late quaternary variability of the Arabian Sea monsoon and oxygen minimum zone, PhD thesis, Utrecht University., 1997.
- Reichart, G. J., Lourens, L. J. and Zachariasse, W. J.: Temporal variability in the northern Arabian Sea oxygen minimum zone (OMZ) during the last 225,000 years, *Paleoceanography*, 13(6), 607–621, doi:10.1029/98PA02203, 1998.
- Rixen, T., Gaye, B. and Emeis, K.-C.: The monsoon, carbon fluxes, and the organic carbon pump in the northern Indian Ocean, *Prog. Oceanogr.*, 175, 24–39, doi:<https://doi.org/10.1016/j.pocean.2019.03.001>, 2019.

- Rommerskirchen, F., Plader, A., Eglinton, G., Chikaraishi, Y. and Rullkötter, J.: Chemotaxonomic significance of distribution and stable carbon isotopic composition of long-chain alkanes and alkan-1-ols in C4 grass waxes, *Org. Geochem.*, 37(10), 665 1303–1332, doi:10.1016/j.orggeochem.2005.12.013, 2006.
- Ruddiman, W. F. and Thomson, J. S.: The case for human causes of increased atmospheric CH<sub>4</sub> over the last 5000 years, *Quat. Sci. Rev.*, 20(18), 1769–1777, doi:https://doi.org/10.1016/S0277-3791(01)00067-1, 2001.
- Schott, W., von Stackelberg, U., Eckhardt, F. J., Mattiat, B., Peters, J. and Zobel, B.: Geologische Untersuchungen an Sedimenten des indisch-pakistanischen Kontinentalrandes (Arabisches Meer), *Geol. Rundschau*, 60(1), 264–275, 670 doi:10.1007/BF01820944, 1970.
- Schulz, H., Von Rad, U. and Von Stackelberg, U.: Laminated sediments from the oxygen-minimum zone of the northeastern Arabian Sea, *Geol. Soc. London, Spec. Publ.*, 116(116), 185–207, doi:10.1144/GSL.SP.1996.116.01.16, 1996.
- Sirocko, F., Sarnthein, M., Lange, H. and Erlenkeuser, H.: Atmospheric summer circulation and coastal upwelling in the Arabian Sea during the Holocene and the last glaciation, *Quat. Res.*, 36(1), 72–93, doi:10.1016/0033-5894(91)90018-Z, 1991.
- 675 Sonzogni, C., Bard, E., Rostek, F., Lafont, R., Rosell-Mele, A. and Eglinton, G.: Core-top calibration of the alkenone index vs sea surface temperature in the Indian Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 44(6), 1445–1460, doi:10.1016/S0967-0645(97)00010-6, 1997.
- Srivastava, P., Agnihotri, R., Sharma, D., Meena, N., Sundriyal, Y. P., Saxena, A., Bhushan, R., Sawlani, R., Banerji, U. S., Sharma, C., Bisht, P., Rana, N. and Jayangondaperumal, R.: 8000-year monsoonal record from Himalaya revealing 680 reinforcement of tropical and global climate systems since mid-Holocene, *Sci. Rep.*, 7(1), 1–10, doi:10.1038/s41598-017-15143-9, 2017.
- Staubwasser, M. and Sirocko, F.: On the formation of laminated sediments on the continental margin off Pakistan: the effects of sediment provenance and sediment redistribution, *Mar. Geol.*, 172(1), 43–56, doi:10.1016/S0025-3227(00)00119-5, 2001.
- Staubwasser, M., Sirocko, F., Grootes, P. M. and Erlenkeuser, H.: South Asian monsoon climate change and radiocarbon in 685 the Arabian Sea during early and middle Holocene, *Paleoceanography*, 17(4), doi:10.1029/2000PA000608, 2002.
- Staubwasser, M., Sirocko, F., Grootes, P. M. and Segl, M.: Climate change at the 4.2 ka BP termination of the Indus valley civilization and Holocene south Asian monsoon variability, *Geophys. Res. Lett.*, 30(8), doi:10.1029/2002GL016822, 2003.
- Stow, D. A. V., Tabrez, A. R. and Prins, M. A.: Quaternary sedimentation on the Makran margin: turbidity current-hemipelagic interaction in an active slope-apron system, *Geol. Soc. London, Spec. Publ.*, 195(1), 219–236, 690 doi:10.1144/GSL.SP.2002.195.01.12, 2002.
- Sun, D., Bloemendal, J., Rea, D. K., Vandenberghe, J., Jiang, F., An, Z. and Su, R.: Grain-size distribution function of polymodal sediments in hydraulic and aeolian environments, and numerical partitioning of the sedimentary components, *Sediment. Geol.*, 152(3–4), 263–277, doi:10.1016/S0037-0738(02)00082-9, 2002.
- Tindale, N. W. and Pease, P. P.: Aerosols over the Arabian Sea: Atmospheric transport pathways and concentrations of dust 695 and sea salt, *Deep. Res. Part II Top. Stud. Oceanogr.*, 46(8–9), 1577–1595, doi:10.1016/S0967-0645(99)00036-3, 1999.
- Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bull. Am. Meteorol. Soc.*, 79(1), 61–78, 1998.

- Ummenhofer, C. C., Gupta, A. Sen, Li, Y., Taschetto, A. S. and England, M. H.: Multi-decadal modulation of the El Niño-Indian monsoon relationship by Indian Ocean variability, *Environ. Res. Lett.*, 6(3), 34006, doi:10.1088/1748-9326/6/3/034006, 2011.
- 700 Vogts, A., Moossen, H., Rommerskirchen, F. and Rullkötter, J.: Distribution patterns and stable carbon isotopic composition of alkanes and alkan-1-ols from plant waxes of African rain forest and savanna C3 species, *Org. Geochem.*, 40(10), 1037–1054, doi:10.1016/j.orggeochem.2009.07.011, 2009.
- Wang, J., Sun, L., Chen, L., Xu, L., Wang, Y. and Wang, X.: The abrupt climate change near 4,400 yr BP on the cultural transition in Yuchisi, China and its global linkage, *Sci. Rep.*, 6, 27723 [online] Available from:  
705 <https://doi.org/10.1038/srep27723>, 2016.
- Wang, P. X., Wang, B., Cheng, H., Fasullo, J., Guo, Z. T., Kiefer, T. and Liu, Z. Y.: The global monsoon across time scales: Mechanisms and outstanding issues, *Earth-Science Rev.*, 174(July 2016), 84–121, doi:10.1016/j.earscirev.2017.07.006, 2017.
- Weiss, H., Courty, M.-A., Wetterstrom, W., Guichard, F., Senior, L., Meadow, R. and Curnow, A.: The Genesis and Collapse of Third Millennium North Mesopotamian Civilization, *Science* (80-. ), 261(5124), 995–1004,  
710 doi:10.1126/science.261.5124.995, 1993.
- Wright, R. P., Bryson, R. A. and Schuldenrein, J.: Water supply and history: Harappa and the Beas regional survey, *Antiquity*, 82(315), 37–48, doi:10.1017/S0003598X00096423, 2008.
- Yadav, R. K., Rupa Kumar, K. and Rajeevan, M.: Increasing influence of ENSO and decreasing influence of AO/NAO in the recent decades over northwest India winter precipitation, *J. Geophys. Res. Atmos.*, 114(12), 1–12, doi:10.1029/2008JD011318,  
715 2009.
- Zhang, X., Jin, L., Chen, J., Lu, H. and Chen, F.: Lagged response of summer precipitation to insolation forcing on the northeastern Tibetan Plateau during the Holocene, *Clim. Dyn.*, 50(9), 3117–3129, doi:10.1007/s00382-017-3784-9, 2018.
- Zhao, J., An, C.-B., Huang, Y., Morrill, C. and Chen, F.-H.: Contrasting early Holocene temperature variations between monsoonal East Asia and westerly dominated Central Asia, *Quat. Sci. Rev.*, 178, 14–23,  
720 doi:<https://doi.org/10.1016/j.quascirev.2017.10.036>, 2017.