

# Hydroclimatic variability of opposing late Pleistocene climates in the Levant revealed by deep Dead Sea sediments

Yoav Ben Dor<sup>1</sup>, Francesco Marra<sup>1,2</sup>, Moshe Armon<sup>1</sup>, Yehouda Enzel<sup>1</sup>, Achim Brauer<sup>3,4</sup>, Markus Julius Schwab<sup>3</sup>, Efrat Morin<sup>1</sup>

<sup>1</sup> The Fredy and Nadine Herrmann Institute of Earth Sciences, The Hebrew University of Jerusalem, Jerusalem, 9190501, Israel

<sup>2</sup> Institute of Atmospheric Sciences and Climate, National Research Council of Italy, Bologna, 40129, Italy

<sup>3</sup> Section Climate Dynamics and Landscape Evolution, GFZ German Research Centre for Geosciences, Telegrafenberg, 14473, Potsdam, Germany

<sup>4</sup> Department of Earth Sciences, Potsdam University, Karl-Liebknecht-Straße 24/25, 14476, Potsdam, Germany

Correspondence to: Yoav Ben Dor ([Yoav.Bendor@mail.huji.ac.il](mailto:Yoav.Bendor@mail.huji.ac.il))

**Abstract.** Annual and decadal-scale hydroclimatic variability ~~is~~ describes key ~~characteristic~~ characteristics embedded into climate ~~insitu~~. ~~It is therefore crucial to, and is of specifically prime importance in subtropical regions. The study of hydroclimatic variability in order is therefore crucial~~ to understand its ~~effects on~~ manifestation as climate derivatives such as hydrological ~~processes~~ phenomena and water availability. However, the study of this variability from modern records is limited due to their relatively short span, whereas model simulations ~~relying~~ relying on modern dynamics could ~~miss~~ misrepresent some of its aspects. Here we study annual to decadal hydroclimatic variability in the Levant using two sedimentary sections covering ~700 years ~~deposited at 18 and 27 Ka retrieved each~~, from the depocenter of the Dead Sea, which has been continuously recording environmental conditions ~~throughout since~~ the late Pleistocene. We focus on ~~the two ~700-year-long series of annually-deposited laminated intervals (i.e., varves) representing two episodes of opposing mean climates, deposited during lake level rise and fall at 27 and 18 Ka, respectively. These two series comprise alternations of authigenic aragonite that precipitated during summer and flood-derived boric detrital laminae deposited during by winter floods. Within this record, aragonite laminae serve as form a proxy of annual inflow and the extent of epilimnion dilution, whereas detrital laminae comprise sub-laminae that record individual floods/flooding events. The two series depict distinct characteristics with increased mean and variance of annual inflow and flood frequency during “wetter”, with respect to the “drier”, conditions, reflected by opposite lake level changes. In addition, decades of intense flood frequency are identified (e.g., clusters), suggesting shifts between depicting the insitu impact of shifting centennial-scale climatic/climate regimes, which are particularly pronounced during wetter, lake-rising conditions. The combined application of multiple time series analyses indicate/suggests that the studied episodes of falling lake levels are characterized by multiple pronounced quasi-weak and non-significant periodic components with periodicities of 2-4, 6-8 and 12 years, whereas the rising lake level episode presents weaker, less persistent periodical components with similar periodicities, sub-decadal frequencies. Combining these observations with the modern synoptic-scale hydroclimatology indicates/suggests that Pleistocene climate changes resulted in shifts of the dominance~~

Formatted: English (United States)

35 of the key synoptic systems governing that govern rainfall, annual inflow and flood frequency in the eastern Mediterranean  
over centennial time-scales.

## 1 Introduction

40 Human activity relies on stable availability of freshwater, which in turn stems from the interaction of climatic, environmental  
and hydrologic conditions. Water availability is particularly crucial in drylands, which cover significant portions of the Earth's  
surface and host nearly 20% of the world's population (Safriel et al., 2005). In these areas, hydroclimatic variability of seasonal,  
annual and decadal scales bear significant effects on water availability that could lead to water stress as population grows  
(Luck et al., 2015; Luo et al., 2015). It is thus crucial to configure how hydroclimatic variability in drylands and Mediterranean  
regions could be affected by climatic changes, similar to those humanity faces in the upcoming decades (Peleg et al., 2015;  
Seager et al., 2019; Zappa et al., 2015). However, because the high-resolution measurements required to study hydroclimatic  
variability only span several decades, the interpretation and the quantification of relationships between trends, oscillations and  
transitional states, as well as their impact on the local water cycle, are still limited and debated (e.g., Morin, 2011; Serinaldi  
et al., 2018). Furthermore, the impacts of climate change on discrete and short-term phenomena such as individual storms and  
floods, is proving harder to determine from the available short records because the extent of available data does not adequately  
capture all possible diversity of hydroclimatic states (Armon et al., 2018; Greenbaum et al., 2010; Metzger et al., 2020; Tarolli  
et al., 2012). Palaeohydrologic records cover centennial and millennial intervals at various resolutions (e.g., Allen et al., 2020;  
Baker, 2008; Brauer et al., 2008; Redmond et al., 2002; Witt et al., 2016), and hence, could improve our understating of how  
climate change is locally manifested into short-term hydroclimatic variability (e.g., Ahlborn et al., 2018; Swierczynski et al.,  
2012). Nevertheless, this requires continuous high-resolution archives, which are rare, especially in sub-tropical arid regions  
(e.g., Zolitschka et al., 2015).

55 The subtropical Levant exhibits a sharp climatic gradient ranging from subhumid Mediterranean to hyperarid climatic zones,  
where slight modifications of global atmospheric circulation could severely impact the hydrological cycle and affect water  
availability (Fig. 1; e.g., Held and Soden, 2006; Luck et al., 2015; Shohami et al., 2011; Tamarin Brodsky and Kaspi, 2017).  
Precipitation in the Levant is limited to the mild and wet winters, whereas summers are dry and hot (Kelley et al., 2012;  
Kushnir et al., 2017; Seager et al., 2019). These properties further obscure the impacts of climatic changes on intra- and inter-  
annual hydroclimatic variability in the wetter Mediterranean regions, as they ultimately stem from the characteristics and  
frequency of discrete extra-tropical Mediterranean cyclones (i.e., Mediterranean/Cyprus Lows; e.g., Campins et al., 2011;  
Floacas et al., 2010; Saaroni et al., 2010; Ziv et al., 2004).

65 In this study, we use two high-resolution sequences of annually deposited laminations (i.e., varves) of the Dead Sea deep  
drilling project (DSDDP) obtained from the Dead Sea depoecenter within the framework of the international continental


scientific drilling program (ICDP; e.g., Neugebauer et al., 2014; Stein et al., 2011). These sequences are analyzed to determine hydroclimatic variability in the southern Levant under contrasting late Pleistocene global and regional climate changes, recorded by independently determined lake level trends (i.e., mean “wetter” vs. “drier” conditions; e.g., Bartov et al., 2003; Torfstein et al., 2013; Torfstein and Enzel, 2017). The segments were continuously sampled for thin-sections preparation, and analyzed using high-resolution microfacies and multiple time-series analyses in order to address the following questions: (a) Can any significant differences between the two studied episodes deposited during opposing mean climates be identified? (b) Do the series illustrate significant intra-series transitions indicative of hydroclimatic regime shifts? (c) Do the studied sequences (or parts of them) record any periodic components? (d) Do the identified periodic components (partially) resemble periodic of known global teleconnections (e.g., NAO, the North Atlantic Oscillation; Black, 2012; Seager et al., 2019)? i.e., can some aspects of the modern hydroclimatic variability assist in interpreting dominant aspects of hydroclimatic variability during past climate changes?

Human activity relies on continuous availability of freshwater, which in turn depends on the interaction of climatic, environmental and hydrologic processes. Water availability is particularly crucial in drylands and subtropical regions that cover significant portions of the Earth’s surface and host nearly 20% of the world’s population (e.g., Safriel et al., 2005). In these areas, hydroclimatic variability at seasonal, annual and decadal scales bear significant impact on water availability that could result in growing water stress as populations grow (e.g., Luck et al., 2015; Luo et al., 2015). It is therefore crucial to understand how hydroclimatic variability in drylands and Mediterranean regions could be affected by climate changes, as humanity faces in the upcoming decades (Peleg et al., 2015; Seager et al., 2019; Zappa et al., 2015; Ipcc, 2021). However, because the study of hydroclimatic variability requires high resolution measurements, that currently only cover several decades, the interpretation and the quantification of relationships between trends, oscillations and transition between climatic states, as well as their impact on the local water cycle insitu, are still limited and are therefore debated (e.g., Morin, 2011; Serinaldi et al., 2018). Furthermore, the impacts of climate change on short-term phenomena such as individual storms and floods, which bear substantial influence on the insitu water cycle in Mediterranean regions, are harder to determine from the available short historical records because the extent of available data does not adequately capture the full diversity of possible hydroclimatic states (Armon et al., 2018; Greenbaum et al., 2010; Tarolli et al., 2012; Metzger et al., 2020). Because palaeohydrologic archives often record centennial and millennial intervals at various resolutions (e.g., Allen et al., 2020; Baker, 2008; Brauer et al., 2008; Redmond et al., 2002; Witt et al., 2017), they have the potential to improve our understating of how climate change is manifested locally into short-term hydroclimatic variability (e.g., Ahlborn et al., 2018; Swierczynski et al., 2012). Nevertheless, this requires continuous high-resolution archives, which are rare, especially in sub-tropical arid terrestrial environments (e.g., Zolitschka et al., 2015).

The subtropical Levant exhibits a sharp climatic gradient ranging from subhumid Mediterranean in the north (Köppen-Geiger classification), where precipitation is focused in winter months (December-May), to hyperarid climate zones in the south (N-S; Kottek et al., 2006), and summers (June-September) are dry and hot . Because the main source of moisture to the region is

105 the Mediterranean Sea, the orographic effect of the central mountainous belt of Israel results in an abrupt W-E climatic gradient, where the eastern parts are substantially drier than its eastern parts (Fig. 1; Kushnir et al., 2017). Under these seasonal conditions, the region is sensitive to spatiotemporal changes in global circulation, where slight modifications could bear substantial impact on the hydrological cycle and water availability (Fig. 1; e.g., Held and Soden, 2006; Luck et al., 2015; Shohami et al., 2011; Tamarin-Brodsky and Kaspi, 2017; Drori et al., 2021). The interaction of substantial seasonality and the abovementioned climatic gradients (N-S and E-W) over the eastern Mediterranean further obscure the impacts of climatic changes on intra- and inter-annual hydroclimatic variability in the wetter Mediterranean regions, as they ultimately stem from the frequency and properties of discrete extra-tropical Mediterranean cyclones (i.e., Mediterranean/Cyprus Lows/Cyclones; e.g., Campins et al., 2011; Enzel et al., 2003; Flocas et al., 2010; Saaroni et al., 2010; Ziv et al., 2004; Armon et al., 2020).

110 In this study, two high-resolution sequences of annually-deposited laminations (i.e., varves) of the Dead Sea deep drilling project (DSDDP) obtained from the Dead Sea depocenter within the framework of the international continental scientific drilling program (ICDP; e.g., Neugebauer et al., 2014) are analyzed to determine hydroclimatic variability in the southern Levant under contrasting late Pleistocene global and regional climate changes, recorded by independently-determined lake level trends (i.e., mean “wetter” vs. “drier” conditions; Fig. 2; e.g., Bartov et al., 2003; Torfstein et al., 2013a; Torfstein and Enzel, 2017). Two laminated segments were continuously sampled for thin-sections preparation, and analyzed using microfacies analyses in high-resolution and multiple time-series analyses in order to address the following questions: (a) Can any significant differences between the two studied episodes recording opposing climatic trends be identified? (b) Do the series depict significant intra-series transitions indicative of hydroclimatic regime shifts? (c) Do the studied sequences (or parts of them) record any periodic components of spectral bands (e.g., interannual, decadal, multi-decadal components)? (d) Do the identified periodic components (partially) resemble the periodicity of known global teleconnection patterns (e.g., the North Atlantic Oscillation, NAO; Seager et al., 2019; Black, 2012)? i.e., can aspects of the modern hydroclimatic variability assist in interpreting dominant aspects of hydroclimatic variability during past climate changes such as periodic components?



125 **Figure 1** - a) Satellite image of the Mediterranean and its surroundings depicting the pivotal location of the ICDP-DSDDP site 5017-1 on the seam of global climate belts (extracted from The Blue Marble Next Generation, NASA's Earth Observatory; Stöckli et al., 2005). b) Satellite image of the eastern Mediterranean depicting the sharp climatic gradient between the Mediterranean north (mean annual precipitation  $>1000 \text{ mm} \cdot \text{yr}^{-1}$ ) and the hyper-arid south (mean annual precipitation  $<50 \text{ mm} \cdot \text{yr}^{-1}$ ; image extracted from Google Earth). c) The Dead Sea watershed and the extent of Lake Lisan during the last glacial maximum (lake level  $\sim 170 \text{ m bmsl}$ ;  $\sim 260 \text{ m}$  above modern level) with modern mean annual precipitation (after Enzel et al., 2003; Torfstein et al., 2013a).

130

## 2 Geological, hydrological, and climatological settings

135 The Dead Sea is the latest of a waterbodies series filling the deepest depression on earth, along the central part of the Dead  
Sea transform since the late Miocene (e.g., Garfunkel, 1981; Garfunkel and Ben-Avraham, 1996; Waldmann, 2017; Waldmann  
et al., 2017). Because level changes in the Dead Sea are directly tied with precipitation fluctuations over its watershed (e.g.,  
Enzel et al., 2003; Morin et al., 2019), its reconstructed past lake levels were utilized as a “mega-palaeo-rain-gauge” (Enzel et  
al., 2003), recording the integrated impacts of environmental conditions and climate on watershed hydrology (Bartov et al.,  
2003; Bookman et al., 2006; Machlus et al., 2000; Morin et al., 2019; Torfstein and Enzel, 2017). These impacts are propagated  
140 into the lake’s sedimentary record by lithological transitions from sequences of alternating aragonite and detrital laminae that  
accumulate during high lake stands and positive water budget (e.g., Ben-Dor et al., 2019; Neugebauer et al., 2014; Torfstein  
et al., 2013), to halite (e.g., Kiro et al., 2016; Palchan et al., 2017; Sirota et al., 2017) or gypsum—are deposited during  
drawdowns and droughts (Torfstein et al., 2008).

145 It was suggested that the climatic gradient over the Dead-Sea watershed has been similar since the late Pleistocene (Enzel et  
al., 2008). This has dictated water and sediment paths to the lake, where the main water source is the Mediterranean climate  
zone at the watershed’s northern region (Enzel et al., 2003; Ziv et al., 2006). In this northern part of the watershed, annual  
water supply largely depends on the spatiotemporal properties and frequency of extra-tropical winter cyclonic systems (i.e.,  
Mediterranean cyclones; Campins et al., 2011; Goldreich et al., 2004; Saaroni et al., 2010). Detrital sediments are delivered  
150 by flash floods (e.g., Nehorai et al., 2013) and debris flows (e.g., Ahlborn et al., 2018; David-Novak et al., 2004), which may  
result from both Mediterranean cyclones, active Red-Sea troughs (Dayan and Morin, 2006; Kahana et al., 2002; Tsvieli and  
Zangvil, 2005), and the less frequent disturbances to the subtropical jet stream also known as Tropical Plumes (Armon et al.,  
2018; Enzel et al., 2012; Tubi and Dayan, 2014; Ziv, 2001). Thus, the annual accumulation of alternating authigenic aragonite  
and detrital laminae throughout the majority of marine isotope stage 2 (MIS2) was attributed to seasonal deposition under high  
155 lake levels and wetter than current mean conditions (Begin et al., 1980; Ben-Dor et al., 2019; Haase-Schramm et al., 2004), as  
later confirmed by the comparison of laminae counting and radiometric dating (Prasad et al., 2004). These sequences dominate  
the record of Lake Lisan, which was the late Pleistocene predecessor of the Dead-Sea (Begin et al., 1980). In Lake Lisan,  
winter inflow was probably a dominant source for replenishing carbonate content in the carbonate-poor Dead-Sea after  
aragonite precipitation took place during summer (Kolodny et al., 2005; Neev, 1963; Neev and Emery, 1967; Stein et al.,  
160 1997), whilst delivering detrital material to the lake (Nehorai et al., 2013).

Until recently, studies of the sedimentary record of the Dead-Sea and its predecessors focused on available exposures  
surrounding the lake (e.g., Bartov et al., 2003, 2007; Prasad et al., 2004; Stein et al., 1997; Torfstein, 2008) and on short cores  
of Holocene sequences (e.g., Heim et al., 1997; Migowski et al., 2006). They were thus affected by local conditions and  
165 hiatuses when lake levels had fallen below the exposures’ altitude (Machlus et al., 2000; Torfstein et al., 2009, 2013) or by

170 wave erosion during episodes rising lake level. In addition, these exposures were accumulated on the lake's shelf, and they therefore reflect its shallow environment, whereas the deeper part of the lake remained nearly inaccessible except for short coring campaigns (e.g., Garber et al., 1987; Nissenbaum et al., 1972). The recently collected ICDP-DSDDP cores from the lake's depocenter, therefore, provide a new regional perspective on sedimentary, limnological and hydrological processes in the lake (e.g., Coianiz et al., 2019; Neugebauer et al., 2014). Detailed microfacies investigation of these cores revealed that the detrital laminae, which alternate with aragonite laminae, are comprised of multiple sub-laminae, each recording individual flood that reached the ICDP-DSDDP coring site from the eastern and western streams flowing to the lake (Ben Dor et al., 2018). Because the majority of these streams are ephemeral, and have relatively small watersheds (<1000 km<sup>2</sup>), flash floods may form both by regional precipitating storms and/or local convection. The laminated record of the deep ICDP-DSDDP core therefore provides a regional record of two independent hydroclimatological variables at annual resolution: (a) the thickness of aragonite laminae, and (b) the number of graded sub-laminae in the detrital layers. Although more recent investigations indicate that the aragonite laminae thickness are not linearly related to total annual inflow into the lake, as previously suggested by Stein et al., (1997), they are still correlated with dilution of the lake and bicarbonate content, and therefore still provide valuable insights on pacing, inherent periodicities and inter-annual inflow variability (Ben Dor et al., in review). The number of detrital sub-laminae in every detrital lamina, on the other hand, records the number of floods that managed to deliver detritus to the coring site, i.e., the floods exceeding some kind of threshold (Ben Dor et al., 2018).




Figure 1: a) Satellite image of the Mediterranean and its surroundings depicting the pivotal location of the ICDP-DSDDP site 5017-1 on the seam of global climate belts (extracted from The Blue Marble Next Generation, NASA's Earth Observatory; Stöckli et al., 2005). b) Top: Dead Sea lake level curve between 10 and 40 Ka showing the high stage of Lake Lisan, the Late Pleistocene predecessor of the Dead Sea during the last glacial (marine isotope stage, MIS2, 15–30 Ka) after Bartov et al., 2003 and Torfstein and Enzel, 2017. Bottom: The age-depth model of the ICDP-DSDDP 5017-1 core and the respective position of the studied segments along the core (after Kitagawa et al., 2017). c) The Dead Sea watershed and the geographic extent of Lake Lisan during the last glacial maximum plotted on modern precipitation isohyets (after Enzel et al., 2003 and Torfstein et al., 2013). d) The studied segments of the ICDP-DSDDP cores dated to 18Ka and 27Ka (MIS 2). e-h) Microscope images of alternating aragonite (summer-precipitate) and detritus (deposited during winter) varves depicting a detrital laminae composed of multiple sub-laminae (labeled 1-11) deposited by individual floods.

The Dead Sea is the recent modern member of a series of waterbodies filling the deepest terrestrial depression on earth along the central part of the Dead Sea transform since the late Miocene (e.g., Garfunkel, 1981; Garfunkel and Ben-Avraham, 1996; Waldmann et al., 2017). Because changes Dead Sea level are directly linked to precipitation fluctuations over its watershed (e.g., Morin et al., 2019), its reconstructed lake levels are utilized as a “mega palaeo-rain gauge” (Enzel et al., 2003), recording the integrated impacts of environmental and climatic conditions on watershed hydrology (Bartov et al., 2003; Bookman et al., 2006; Machlus et al., 2000; Torfstein and Enzel, 2017). These impacts are propagated into the lake’s sedimentary record and



200 are reflected by lithological transitions from sequences of alternating aragonite and detrital laminae that are deposited during high lake stands and positive water budget (Fig. 2; e.g., Ben Dor et al., 2019; Neugebauer et al., 2014; Torfstein et al., 2013b), to halite (e.g., Palchan et al., 2017; Sirota et al., 2017) or gypsum during drawdowns and droughts (Torfstein et al., 2008).

205 Although the full wet season lasts from October to May, ~65% of precipitation over the Dead Sea watershed is limited to the mild and wet winters (December-February). At the northern, and wetter part of the region, >90% of precipitation is delivered by low pressure systems from the Mediterranean Sea (i.e., Mediterranean Lows), whereas during summer (June-August), large-scale atmospheric subsidence results in stable dry and hot conditions (Goldreich, 2012; Kushnir et al., 2017; Tyrlis et al., 2013; Kelley et al., 2012). On average, the northern and western parts of the Dead Sea watershed are characterized by increased precipitation and lower temperatures, resulting in a pronounced climatic N-S gradient. The average temperatures during winter range between ~10 and ~20°C, whereas during summer, temperatures range from ~20 to ~35°C, in the northern and southern parts of the Dead Sea watershed, respectively (Israel Meteorological Service, 2021). Mean annual precipitation in the hyperarid region is under 50 mm·yr<sup>-1</sup>, delivered during less than 10 rainy days, whereas in the mountainous region at the north, it exceeds 1000 mm·yr<sup>-1</sup> and spread over more than 75 rainy days (Goldreich, 2012; Sharon and Kutiel, 1986). Precipitation over the southern part of the Levant is characterized by substantial spatiotemporal variability, depicting scattered, spotty, and intense convective rain (Sharon, 1972). The few observed precipitation events in the southern arid parts are separated by prolonged dry intervals ranging from weeks to months. The pronounced orographic effect of the Judean mountains, which separate the Lower Jordan Valley from the Mediterranean Sea, substantially decreases rainfall as storms progress eastward towards the Dead Sea basin, thus resulting in a sharp E-W precipitation gradient, which decreases from >500 mm·yr<sup>-1</sup> in Jerusalem to <100 mm·yr<sup>-1</sup> at the Dead Sea, over less than 50 km. East of the Dead Sea, the mountainous topography of the Jordanian escarpment increases annual precipitation to over 400 mm·yr<sup>-1</sup> at its highest parts (Fig. 1; Enzel et al., 2003).

215 Available records indicate that the modern climatic gradients over the Dead Sea watershed was similar during the late Pleistocene (Keinan et al., 2019; Enzel et al., 2008). These observations determine water and sediment transport paths into the lake, where the main water source is its Mediterranean climatic zone to the north (Fig. 1; e.g., Enzel et al., 2003; Ziv et al., 2006), where annual precipitation and water supply largely depend on the spatiotemporal properties and the frequency of extra-tropical cyclonic systems during winter (i.e. Mediterranean Lows; Campins et al., 2011; Goldreich, 2004; Saaroni et al., 2010). Detrital sediments are delivered to the ICDP 5017 coring site at the depocenter of the Dead Sea lake primarily by flash floods from adjacent streams (e.g., Nehorai et al., 2013; Ben Dor et al., 2018), and debris flows (Ahlborn et al., 2018). Whereas floods are generally associated with Mediterranean Lows (Dayan and Morin, 2006; Kahana et al., 2002; Tsvieli and Zangvil, 2005), debris flows, are associated with active Red Sea troughs (e.g., Ahlborn et al., 2018; Ben David-Novak et al., 2004). More rarely, subtropical jet stream disturbances can also generate flash floods (i.e., Tropical Plumes; Armon et al., 2018; Enzel et al., 2012; Tubi and Dayan, 2014; Ziv, 2001). Thus, the deposition of alternating authigenic aragonite and detrital laminae, which dominate the record of Lake Lisan the late Pleistocene predecessor of the Dead Sea (~70-14 Ka; Begin et al., 1980;


235 [Katz et al., 1977; Kaufman, 1971](#)), throughout the majority of marine isotope stage 2 (MIS2) was attributed to annual  
deposition under seasonal climate during high lake levels and wetter-than-modern conditions, which sufficed to replenish the  
lake with necessary bicarbonate to support aragonite precipitation during summer and deliver detrital sediments during winter  
([Begin et al., 1980; Ben Dor et al., 2019; Haase-Schramm et al., 2004](#)), as was further confirmed by the comparison of laminae  
counting and radiometric dating ([Prasad et al., 2004; Migowski et al., 2006](#)). In Lake Lisan, winter inflow from its northern  
part was likely the dominant source of water and alkalinity, replenishing carbonate content in the carbonate-poor Dead Sea  
240 after aragonite precipitation took place during summer ([Kolodny et al., 2005; Neev, 1963; Neev and Emery, 1967; Stein et al.,  
1997](#)), whereas local floods in adjacent (ephemeral) streams delivered detrital sediments to the lake's depocenter ([Nehorai et  
al., 2013](#)).

245 Until recently, studies of the sedimentary record of the Dead Sea and its predecessors focused on available outcrops  
surrounding the lake (e.g., [Bartov et al., 2003; Bartov et al., 2007; Stein et al., 1997; Prasad et al., 2004; Torfstein et al., 2008](#))  
as well as short cores of Holocene sequences (e.g., [Heim et al., 1997; Migowski et al., 2006](#)). The broad aspect of their  
implications was thus limited due to local conditions and depositional hiatuses that form when lake levels fell below the  
outcrop's altitude ([Machlus et al., 2000; Torfstein et al., 2013b; Torfstein et al., 2009](#)) or by wave erosion during episodes of  
250 rising lake level. In addition, these outcrops were accumulated on the lake's shelf, and they therefore reflect its shallow  
environment, whereas the deeper part of the lake remained nearly inaccessible except for short coring campaigns (e.g., [Garber  
et al., 1987; Nissenbaum et al., 1972](#)). The recently collected ICDP-DSDDP cores from the lake's depocenter (ICDP site 5017),  
therefore, provide a new regional perspective on sedimentary, limnological and hydrological processes in the lake (Fig. 1; e.g.,  
[Coianiz et al., 2019; Neugebauer et al., 2014](#)). The 5017 cores were dated by combining <sup>14</sup>C dating of macro organic debris  
including seeds and twigs ([Neugebauer et al., 2014](#)) with U-Th dating of primary aragonites ([Torfstein et al., 2015](#)), and by  
255 correlating their  $\delta^{18}\text{O}$  with cave deposits (e.g., [Lisiecki and Raymo, 2005; Bar-Matthews et al., 1999; Bar-Matthews et al.,  
2003](#)). The age-depth model of the sections used in this study was established through a Markov chain Monte Carlo procedure  
utilizing <sup>14</sup>C dates covering ~50 Ka (Fig. 2; [Kitagawa et al., 2017](#)).

260 More recently, a detailed microfacies investigation revealed that the detrital laminae of the alternating aragonite and detritus  
facies, are comprised of multiple sub-laminae, recording individual flooding events from adjacent streams that face the 5017  
ICDP coring site ([Ben Dor et al., 2018](#)). Because the majority of these streams are ephemeral, and have relatively small  
watersheds (<1000 km<sup>2</sup>), such flash-floods may be triggered either by regional storms, that substantially contribute to the  
lake's annual water budget, and/or by smaller-scale local convection that has negligible impact on annual inflow. The laminated  
265 sections of the ICDP-DSDDP core therefore provide a regional record of two (nearly) independent hydroclimatological  
variables at annual resolution: (a) the thickness of aragonite laminae (this study), and (b) the number of sub-laminae within  
detrital layers (Fig. 2; [Ben Dor et al., 2018](#)). Although more recent investigations indicate that aragonite laminae thickness is  
not linearly related to total annual inflow, as was previously suggested ([Stein et al., 1997](#)), it is correlated and monotonously

associated with carbonate inflow and lake dilution, where increased inflow would result in increased aragonite thickness, and may therefore still provide valuable insights on inherent periodicities and inter-annual inflow variability (Ben Dor et al., 2021).

The alkalinity required to support annual aragonite precipitation may be derived from dust settling directly on the lake and its watershed, as well as freshwater inflow. However, direct dust deposition is insufficient to support the deposition of aragonite laminae (e.g., Ganor and Foner, 1996; Kalderon-Asael et al., 2009), whereas the dissolution and remobilization of accumulated dust from the watershed, has the potential to supply bicarbonate that could increase inflow alkalinity (e.g., Crouvi et al., 2017; Belmaker et al., 2019). Although this cannot be directly addressed for MIS2, recent studies of the snow-affected Mt. Hermon region in Israel (Avni et al., 2018) and the denudation rates in the Judea region (Ryb et al., 2014), suggest that the dissolution of bedrock could not have increased alkalinity inflow by a factor greater than two, that would bear a more substantial impact on the northern part of the watershed. Thus the water and alkalinity budgets of the lake are ultimately dominated by rainfall over the northern part of the watershed (~75-85%), and subsurface inflow (~10-15%; Siebert et al., 2014; Levy et al., 2020), whereas the water (and alkalinity) contribution of floods is negligible (5-10%; Armon et al., 2019). Because the number of detrital sub-laminae, on the other hand, records the number of floods exceeding a threshold and reaching the coring site that deliver negligible water volumes (Ben Dor et al., 2018), the two series are addressed in this study independently in order to examine their properties and interactions. Thus, although floods over the small watersheds draining into the Dead Sea deliver both water and alkalinity into the Dead Sea (e.g., Belmaker et al., 2016; Golan et al., 2017), their overall negligible volume with respect to the annual inflow, makes the two studied proxies of aragonite thickness (this study) and the previously published number of detrital sub-laminae (Ben Dor et al., 2018) practically independent.



290 **Figure 2 - a) Top:** Dead Sea lake level depicting the highest stage of Lake Lisan during last glacial maximum, the Late Pleistocene  
 predecessor of the Dead Sea (Bartov et al., 2003; Torfstein and Enzel, 2017). **Bottom:** The age-depth model of the ICDP-DSDDP  
 5017-1 core and the position of the studied segments along the core (after Kitagawa et al., 2017). **b) The studied sections of the ICDP-  
 DSDDP** cores dated to 18 Ka and 27 Ka (marine isotope stage: MIS2, ~15-30 Ka). **c-f) Microscope images of alternating aragonite  
 (summer precipitate) and detritus (deposited during winter) varves depicting a detrital laminae composed of multiple sub-laminae  
 (labelled 1-11) deposited by individual floods (after Ben Dor et al., 2018).**

295

### 3 Materials and methods

#### 3.1 Microfacies analyses

Two chronologically-constrained segments of the ICDP-DSDDP 5017-1 core were selected for this study (ca. 18 and 27 Ka; Fig. 1; Neugebauer et al., 2014). These segments are coeval with opposing climatic trends, reflected by independently-determined rising (27 Ka) and falling (18 Ka) lake level trends (Fig. 1b; Bartov et al., 2002; Torfstein and Enzel, 2017). Continuous thin sections were prepared using a procedure adjusted for saline sediments (e.g., Brauer et al., 1999; Neugebauer et al., 2015), and analyzed using a Hirox RH 2000 optical microscope. A total of ~700 varves (years) were counted and described in each segment (Fig. S1). The segments comprise alternations of finely laminated varves (Ben Dor et al., 2019) and event-related deposits (ERDs; Neugebauer et al., 2014), attributed to debris flows (Ahlborn et al., 2018) and earthquakes (Kagan et al., 2018). This study focuses on two properties of the varves interpreted as hydrological proxies: (a) the thicknesses of aragonite laminae, and (b) the number of detrital sub-laminae in each detrital lamina (Figs. 1 and 2). Because the thickness of detrital laminae cannot be directly linked to specific hydroclimatic aspects, but instead depends on the interaction of multiple factors such as sediment availability and the location of the activated catchment, it is not analyzed here as a hydroclimatic proxy. The thickness of aragonite laminae was suggested to reflect annual inflow and lake dilution during the wet season (Kolodny et al., 2005; Stein et al., 1997), whereas individual sub-laminae in detrital laminae correspond to flood events capable of delivering siliciclastic sediments to the coring site (Ben Dor et al., 2018). These sub-laminae are attributed to floods that surpassed the threshold required to reach the coring site from their catchments. Missing data, such as were laminae are trimmed or distorted (~5%) were completed by median values prior to time-series analyses.

Two chronologically-constrained segments (each ~1.5 meter long) of the ICDP-DSDDP 5017-1 core deposited during ~18 and 27 Ka were selected for this study (Fig. 2; Neugebauer et al., 2014). These segments are coeval with opposing climatic trends, reflected by independently-determined rising (27 Ka) and falling (18 Ka) lake level trends (Bartov et al., 2002; Torfstein and Enzel, 2017). The two sections were sampled continuously with overlap to prepare sequence comprising 10 cm long thinsections using a dry-freeze procedure adjusted for saline sediments (e.g., Brauer et al., 1999; Neugebauer et al., 2015), and analyzed using a Hirox RH-2000 optical microscope (e.g., Ben Dor et al., 2018; Ben Dor et al., 2021). A total of ~700 varves (years) were counted and described in each segment. The segments comprise alternations of finely laminated varves (Ben Dor et al., 2019) and event-related deposits (Neugebauer et al., 2014), attributed to debris flows (Ahlborn et al., 2018) and earthquakes (Fig. 2; Kagan et al., 2018). The nature of these sediments and the way their mode of formation according based on the study of available exposures (Begin et al., 1980; Begin et al., 1974; Stein et al., 1997; Marco et al., 1996), and modern analogous lakes (e.g., Dean et al., 2015; Roeser et al., 2021) suggest that these detritus-aragonite couplets were deposited annually, thus forming varves. This is further supported by the uniformity of the sediments revealed by microfacies analyses carried out in this study, in which slight changes sediment properties are observed in detail and the agreement between laminae counting and independent radiometric dating such as  $^{14}\text{C}$  and U-Th (Prasad et al., 2009; Haase-Schramm et al., 2004; Neugebauer et al., 2015). Because no deposition of alternating aragonite and detritus sedimentary facies

330 [takes place under modern conditions in the Dead Sea \(e.g., Ben Dor et al., 2021\), the interpretation of alternating aragonite and detritus facies as annual deposits is a relatively solid assumption used within the framework of this study, as it cannot be directly determined for the studied interval Lake Lisan \(e.g., Prasad et al., 2004; Ben Dor et al., 2019; Ben Dor et al., 2020\).](#)

335 [This study focuses on two properties of the varves interpreted as hydrological proxies: \(a\) the number of detrital sub-laminae in each detrital lamina \(Ben Dor et al., 2018\), and \(b\) the thicknesses of aragonite laminae \(this study\). Because the thickness of detrital laminae \(Fig. 2\) is not directly linked to specific hydroclimatic processes, but instead depends on the interaction of multiple factors such as sediment availability and the geographical position of the activated catchment, it is not analyzed here as a hydroclimatic proxy. The thickness of aragonite laminae was suggested to reflect annual inflow and lake dilution during the wet season \(Kolodny et al., 2005; Stein et al., 1997\), whereas individual sub-laminae in detrital laminae correspond to flood events capable of delivering siliciclastic sediments to the coring site \(Ben Dor et al., 2018\). These sub-laminae are attributed to floods that surpassed the threshold required to reach the coring site from their catchments. Missing data where laminae are trimmed or distorted \(~5%\) were imputed using singular spectrum analysis \(e.g., Kondrashov and Ghil, 2006\).](#)

340 [Because](#) the typical number of sub-laminae is low, and the minimal number of detrital sub-laminae that can be counted is one, as no cases of “zero sub-laminae” are sedimentologically-distinguishable, varves comprising a single detrital lamina should be considered as varves recording either years with no floods or years with a single flood. The two series deposited during rising and falling lake levels were compared using key statistical properties: the Mann-Whitney-Wilcoxon ranksum test to determine if they were sampled from two populations of a statistically different median (MWW; Mann and Whitney, 1947), and the Ansari-Bradley (1960) ranksum dispersion test to determine if they were characterized by statistically significant dispersion. (MWW; Mann and Whitney, 1947), and the Ansari-Bradley ranksum dispersion test (AB; Ansari and Bradley, 1960) to determine if they were characterized by statistically significant dispersion.

### 3.2 Detecting intra-record regime shifts

355 [Regime shifts were identified using a non-parametric test that utilizes the normalized running sum as the test statistic was developed for identifying flood-rich episodes \(i.e., clusters; Appendix B\) and refined using a running MWW and Ansari-Bradley tests \(Figs. S2 and S3\). Following a visual inspection, the tests were applied with a symmetric window of 51 years width \(25 years in each direction\), that was adequate for order to identify regime shifts as part of this purpose study by visual inspection. First, comparing the observed running sum of sub-laminae was calculated for each series count normalized by its mean value using consecutive multiple equally-spaced window widths ranging from 10 to 300 years. The running sum was consequently normalized to its mean value for every window width, and peaks in the series were identified. In order to avoid overlaps, only local maximum values, separated by at least half window width, were further considered for the analysis \(Figs. S2-1 and S2-6\). The peaks were ordered according to their rank in descending order, where the largest value was assigned the](#)

Formatted: English (United States)

first position and the smallest value was assigned the last position. Each, against the surrogate series generated by point-wise permutations of the original data series was then randomly permuted 10,000 times without replacement and the same statistics (peaks of (Eq. 1; Figs. 3 and S1-S5)).

$$\text{Eq. 1: } y_{w(i)} = \begin{cases} i < \frac{w}{2} \\ \frac{w}{2} < i < N - \frac{w}{2} \\ i > N - \frac{w}{2} \end{cases} \frac{\sum n_j}{\widehat{y}_w}, \left[ i - \frac{w}{2} \leq j \leq i + \frac{w}{2} \right] \begin{cases} NaN \\ NaN \\ NaN \end{cases}$$


Equation 1 - the normalized running sum of sub-laminae counts,  $y_{w(i)}$  is the normalized running sum,  $i$  is the location index ranging between 1 and  $N$ ,  $N$  is the number of data points in the studied series,  $\sum n_j$  is the sum of sub-laminae count over the range between  $[i - \frac{w}{2} \leq j \leq i + \frac{w}{2}]$ ,  $w$  is the window width ranging from 10 to 300 by 10 years steps ( $w = [10, 20, 30, \dots, 290, 300]$ ), and  $\widehat{y}_w$  is the mean normalized running sum, allowing the calculation of their distributions, and the estimation of the likelihood of randomly of  $n$  for window width  $w$ . The resulting  $y_{w(i)}$  values larger than unity indicate flood-rich episodes, whereas  $y_{w(i)}$  values lower than unity indicate flood-poor episodes.

Because the aim of this analysis is to assess the probability of observing such values (Figs. S2-2 and S2-7). The observed value decadal to centennial intervals of increased flood frequency (i.e., clustering of floods) whilst avoiding a-priori assumptions and parametrization, and because the two series depicts low serial dependence (Fig. S6), the surrogate time series were produced by point-wise permutations. After the running sum was normalized running sum by its mean value for every peak rank window width, and local maxima values that are separated by at least half of the window width was were considered for the analysis (Fig. 3c and 3d). The peaks were ranked in descending order and compared to the distribution against a dataset derived from the 10,000 randomly permuted series (Figs. S2-3 and S2-8), 10,000 permutations without replacement of the respective series, revealing the significance level of observing intervals of the studied window width. Observed values larger than the 95<sup>th</sup> percentile are hereby of the permuted series are considered statistically significant with a false positive error of  $\alpha=0.05$  (Figs. S2-4 Fig. 3e and S2-93f). Because of the wide range significant peaks of applied different window widths for the statistical test, the significant peaks of different widths tend to overlap each other, so, the clusters were eventually evaluated by considering their summed probabilities (Figs. S2-5 Fig. 3g and S2-103h), and their edges were further refined by the a running MWW test and AB tests, in which the two halves of the window are compared against each other (Figs. S7-S8).

Recurrence and cross-recurrence analyses were applied to identify non-linear transitions and distinct regimes (Eckmann et al., 1987). These methods can be used to detect abrupt transitions as well as short-term periodic and quasi-periodic behaviors in non-stationary and noisy series by investigating their dynamics in the reconstructed phase space (Donges et al., 2011; Marwan et al., 2007). This approach is specifically useful in this study because the studied proxies record the convolved non-linear interaction of climatic, hydrological and limnogeological processes. Because the effect of their interactions could stem from non-linear dynamics that bear different implications on the studied sedimentary sections, the application of recurrence-based methods could prove more robust than specific harmonic or wavelet functions (Marwan et al., 2003; Marwan and Kurths,

395 2004). The analysis was carried out using the CRP toolbox for MATLAB® after normalizing the series to zero mean and unit  
standard deviation (ver. 5.22; Marwan et al., 2007). The embedding dimension and the time delay were determined using the  
false nearest neighbor approach (Kennel et al., 1992) and the mutual information method (Fig. S4; Marwan, 2011),  
respectively. An environment of uniform size was selected by a visual inspection of the plots ( $\epsilon=1\sigma$  for recurrence and  $\epsilon=1.5\sigma$   
400 for joint recurrence analyses). Recurrence quantification analysis (RQA) was used to calculate the recurrence rate (RR) and  
the transitivity coefficient, which describe the degree of regularity of the system's dynamics using the recurrence plot (Marwan  
et al., 2007), using a 30-years moving window with 20% overlap, which were found to be informative and not too noisy for  
this purpose.





405

**Figure 3:** the non-parametric test developed for identifying flood-rich episodes (i.e., flood clustering) a,b) the studied series of flood frequency analyzed using microfacies analyses of two DSDDP core sections. c,d) the running normalized sum calculated for identifying flood clusters and its peaks (triangle) using different window widths. e,f) binary diagrams indicating where observed  $y_{w(t)}$  values are higher than the calculated 95<sup>th</sup> percentile of the distributions derived from 10,000 randomly permuted series, and are thus statistically significant at  $\alpha=0.05$  (black pixels) or non-significant (white pixels). Missing values, where not enough peaks

410 were detected to calculate the 95th percentile are coded grey, g,h) the location of statistically significant peaks at  $\alpha=0.05$  depicted as semi-transparent shading for the top three highly ranked peaks of window widths ranging between 50 and 300. The red line depicts the probability of the summed identification of clusters by all window widths. The refined clusters are marked by dashed vertical lines.

415 Recurrence and joint-recurrence techniques and their analyses (*ROA* and *JROA*) were applied to identify non-linear transitions and distinct regimes within the series (Eckmann et al., 1987). These methods can be used to detect abrupt transitions as well as short-term periodic and oscillatory behaviours in non-stationary and noisy data by investigating their dynamics in the reconstructed phase space (Donges et al., 2011; Marwan et al., 2007). The following short introduction and included equations are based on the detailed review by Marwan et al. (2007), to which the keen reader is referred for further details. The recurrence of the system is computed by considering the pairwise similarity (proximity) of points along its phase space trajectory  
420 reconstructed using a time delay ( $\tau$ ) and an embedding dimension ( $m$ ). The recurrence plot (*RP*), which is used to efficiently visualize the recurrence of the studied series, is generated by calculating the  $R_{i,j(\epsilon)}$  matrix, through the binarization of the difference between the norm of the pairwise trajectories of steps in the reconstructed phase space using a predefined threshold  $\epsilon$  (Eq. 2; Marwan et al., 2007). When the trajectory in two points ( $i$  and  $j$ ) is smaller than  $\epsilon$ , the value of the corresponding point is set to one in  $R$ , and the pixel painted black in the *RP*. Otherwise the value is set to zero and the corresponding pixel in  
425 the *RP* is painted white.

$$\text{Eq. 2: } R_{i,j(\epsilon)} = \theta(\epsilon - \|\vec{x}_i - \vec{x}_j\|), \quad i, j = 1, 2, 3, \dots, N$$

430 Equation 2 (after Marwan et al., 2007) – the calculation of the recurrence matrix using a threshold  $\epsilon$ , where  $N$  is the number of points of the reconstructed  $\vec{x}_i$  trajectory, and  $\theta(\cdot)$  is the Heaviside function (step function), where  $\theta(x) = 0$ , if  $x < 0$  and  $\theta(x) = 1$  otherwise.

Similarly, the joint recurrence plot can be derived (*JRP*) for studying the dynamical relationship of two series or more by the product of recurrence plots (Eq. 3; Romano et al., 2004).

$$\text{Eq. 3: } J_{R_{i,j(\epsilon_{\vec{x}}, \epsilon_{\vec{y}})}} = \theta(\epsilon_{\vec{x}} - \|\vec{x}_i - \vec{x}_j\|) \cdot \theta(\epsilon_{\vec{y}} - \|\vec{y}_i - \vec{y}_j\|), \quad \vec{x}_i \in \mathbb{R}^m, \vec{y}_i \in \mathbb{R}^n, \quad i, j = 1, 2, 3, \dots, N$$

435 Equation 3 (after Marwan et al., 2007) – the calculation of the joint recurrence matrix as the product of two recurrence matrices calculated for two series with trajectories  $\vec{x}$  and  $\vec{y}$ , using the thresholds  $\epsilon_{\vec{x}}$  and  $\epsilon_{\vec{y}}$ , respectively.  $N$  is the number of points of the trajectories, and  $\theta(\cdot)$  is the Heaviside function (step function), where  $\theta(x) = 0$ , if  $x < 0$  and  $\theta(x) = 1$  otherwise.

440 The *RP* of a cyclic system would show diagonal lines parallel to the line of identity (*LOI*), which connects its lower left corner with its upper right one, whereas a noisy system would be characterized by spotty and random appearance of points throughout the *RP*. Once the *RP* is calculated, its properties and the time dependent behaviour of the system can be studied by quantifying its recurrence using recurrence quantification analysis (*ROA*) by computing different measures for each sub-matrix formed by sliding a square window along its *LOI* with a predefined step. In this study the recurrence rate (*RR*), determinism (*DET*), the

maximum length of a diagonal line within a sub-matrix ( $L_{max}$ ) and the laminarity ( $LAM$ ) metrics are used. The maxima of  $DET$  and  $L_{max}$  can be used to identify periodic-chaos and chaos-periodic transitions. Alongside those transitions in  $LAM$  (minima), it is capable of identifying chaos-chaos transitions as well (maxima). The recurrence rate is the percentage of points in which recurrence occurs within the studied section (Eq. 4).

$$\text{Eq. 4: } RR(\epsilon) = \frac{1}{N^2} \sum_{i,j=1}^N R_{i,j}(\epsilon)$$

The determinism ( $DET$ ) reflects the extent to which the system maintains an ordered, cyclic and deterministic behaviour. Because uncorrelated, weakly correlated, stochastic or chaotic processes form very short diagonals or none at all, whereas deterministic processes generate longer diagonals, the ratio of recurrence points that form diagonal lines (with length  $\geq l_{min}$ ) to all recurrence points can be used to estimate the main behaviour of the system within the studied period (Eq. 5). In this study the minimal length of  $l(l_{min})$ , was set to  $l_{min} = 2$ .

$$\text{Eq. 5: } DET = \frac{\sum_{l=l_{min}}^N l P(l)}{\sum_{l=1}^N l P(l)}$$

Similar to  $DET$ , the laminarity ( $LAM$ ) is computed as the portion of points forming a vertical line of minimal length  $v$  out of the entire set of recurrence points within the RP (or in its sub-matrices).

$$\text{Eq. 6: } LAM = \frac{\sum_{v=v_{min}}^N v P(v)}{\sum_{v=1}^N v P(v)}$$

In order to decrease the influence of the tangential motion  $LAM$  is computed using the points forming  $v$  that exceed a minimal length  $v_{min}$ , set in this study as  $v_{min} = 2$ . In cases where the RP consists of more single recurrence points than vertical structures  $LAM$  will decrease.

The length of the longest diagonal line ( $L_{max}$ ) is computed by identifying the longest diagonal line found within the RP (or in its sub-matrices; Eq. 7).

$$\text{Eq. 7: } L_{max} = \max(\{l_i\}_{i=1}^{N_i})$$

where  $N_i = \sum_{l \geq l_{min}} P(l)$  is the total number of diagonal lines.

This approach is specifically useful in this study because the studied proxies record the convolved non-linear interaction of climatic, hydrological and limnogeological processes, because the effect of their interactions could stem from non-linear dynamics that bear very different implications on sedimentary sections (Marwan and Kurths, 2004; Marwan et al., 2003). The analysis was carried out using the CRP toolbox for MATLAB© after normalizing the series to zero mean and unit standard deviation (ver. 5.22; Marwan et al., 2007). The embedding dimension and the time-delay were determined using the false nearest neighbour approach (Kennel et al., 1992) and the mutual information method (Fig. S6; Marwan, 2011). The threshold

$\varepsilon$  was set to fit a recurrence rate (RR) of 0.1 using the max norm method (Schinkel et al., 2008), and the RQA (and JRQA) measures were calculated with a window of 30 years with a time step of a single year.

### 3.3 Detecting periodical/periodic components

Hydrological data are usually not normally distributed, and may show short-term periodicities that would be hard to identify using standard Fourier-derived approaches such as the periodogram (Blackman and Tukey, 1958; e.g., Welch, 1967). Wavelet analyses provide a flexible tool for identifying irregular and non-stationary periodicities in short time series (Lau and Hengyi Weng, 1995; Torrence and Compo, 1998). We utilized the principles delineated by Torrence and Compo (1998) using the Morlet wavelet function after normalizing the data to zero mean and unit standard deviation (Grinsted et al., 2004). Significance testing was carried out against simulated red noise modeled by a lag-1 autoregressive process (AR(1); Grinsted et al., 2004; Torrence and Compo, 1998).

Singular spectrum analysis (SSA) is used to identify periodic and quasi-periodic signals in short, non-stationary and noisy time series (Broomhead and King, 1986). Unlike wavelet and Fourier-based procedures, where the signal is compared against a predefined function, SSA identifies key components that best explain the variance of the time-delayed series in an embedding matrix. This non-parametric approach reveals the dominant components that best explain data variance and defines their relative contribution for generating the observed signal (Vautard and Ghil, 1989). This, in turn, allows the breaking-down of the signal into its principle components and calculating their corresponding reconstructed components (RCs), which demonstrate the extent of contribution for each component to the signal (Ghil et al., 2002). Thus, the analyses of the individual RCs, and the determination of their dominant periodicities can point to dominant frequencies embedded in the observed series, which in turn could stem from teleconnections affecting hydroclimatic variability (e.g., NAO, SOI, etc.; Feliks et al., 2010; Le Mouél et al., 2019; Seager et al., 2019). To balance between the analyses complexity and usefulness, and avoid overfitting of the SSA, it was performed with an embedding dimension (window) of ten years after visually inspecting the effect of RC number of the eigenvalues. The first RC ( $RC^{(1)}$ ) was later used as the local trend component that was subtracted from the data as a high-pass filter to reveal short-term periodicities (Fig. S5). The SSA RCs were additionally analyzed using Welch's periodogram for evaluating the RCs power spectral density (Welch, 1967), and wavelet and cross-wavelet analyses (Grinsted et al., 2004) in order to identify their dominant periodicities in cluster and the background periods. The periodogram was calculated after applying a Hamming window of width 25 years and an overlap of 50%, applied to merge close periodical components and reduce spectral noise, after subtracting the mean from the series and zero-padding it to  $2^{14}$  points in order to minimize spectral leakage (Figs. S7-S11). Additionally, a multi-channel SSA (Groth and Ghil, 2015) followed with a cross-wavelet and Welch's periodogram analyses was performed (Figs. S12 and S13).

Hydrological data are commonly skewed, and may exhibit short-term periodic behaviour that is difficult to identify using standard Fourier-derived approaches such as the periodogram (e.g., Blackman and Tukey, 1958; Welch, 1967). Wavelet

505 analyses provide a set of flexible tool for identifying irregular and non-stationary periodicities in short time series (Lau and Weng, 1995; Torrence and Compo, 1998). The coherence of two series provides an estimate of the importance of areas with high common power in the two series, that reflects the localized correlation coefficient in time-frequency space (Grinsted et al., 2004). Wavelet analyses were carried out using the Morlet wavelet function after normalizing the data to zero mean and unit standard deviation. The significance of the wavelet power and coherence were determined against red noise simulated using a lag-1 autoregressive process (AR(1)) using the cumulative area-wise approach for detecting false positive periodicities at the  $\alpha=0.05$  (Schulte, 2019, 2016).

515 Singular spectrum analysis (SSA) is used to identify periodic and oscillatory signals in short, non-stationary and noisy time series (Broomhead and King, 1986). Unlike wavelet and Fourier-based procedures, where the signal is compared against a predefined function, SSA identifies principle components that best explain the variance of the time-delayed series through the embedded matrix. This non-parametric approach reveals the dominant components that best explain the variance and defines their relative contribution for generating the observed signal (Vautard and Ghil, 1989). This, in turn, allows the breaking-down of the signal into its principle components and calculating their corresponding reconstructed components (RCs), which demonstrate the extent of contribution for each component to the signal (Ghil et al., 2002). Generally, SSA complements wavelet and other spectral analyses because it provides increased flexibility, and may identify oscillatory components of different shapes that change in time. Thus, the analyses of the individual RCs, and the determination of their dominant periodic components can point at dominant frequencies embedded in the observed series, which in turn could stem from teleconnection patterns affecting hydroclimatic variability (e.g., NAO, SOI, etc.; Feliks et al., 2010; Le Mouél et al., 2019; Seager et al., 2019). In order to balance between the analyses complexity and usefulness, the SSA was performed with an embedding dimension (window) of ten years after visually inspecting the effect of the number of RCs on the relevant eigenvalues. The first RC ( $RC^{(1)}$ ) was later used as the local trend component that was subtracted from the data as a high-pass filter to reveal short-term periodicities (Fig. S9). The SSA RCs were additionally analyzed using the Welch periodogram for evaluating the RCs power spectral density (Welch, 1967), and wavelet analyses using the area-wise test for significance (Schulte, 2016, 2019) in order to compare their time-frequency properties during cluster and background periods. The Welch periodogram was calculated after applying a Hamming window of width 25 years and an overlap of 50%, applied to merge close periodic components and reduce spectral noise, after subtracting the mean from the series and zero-padding it to  $2^{11}$  points in order to minimize spectral leakage (Figs. S10-S13).

## 4 Results

### 4.1 Microfacies analyses

535 The studied segments are dominated by alternating laminae of aragonite and detritus with a greenish tint (Figs. 1, S1) characterized by a typical laminae thickness of <1 mm. The laminated segments are sporadically interrupted by massive or

graded ERDs attributed to abrupt mass wasting events (e.g., turbidities and slumps), likely triggered by earthquakes (Kagan et al., 2018), debris flows and slope instability (Ahlborn et al., 2018). Both laminae thickness and number of sub-laminae show statistically significant larger mean and variance during lake level rise (~27 Ka) than during lake level fall (~18Ka; Table 1; Fig. 2).

The studied segments are dominated by greenish alternating laminae of aragonite and detritus (Fig. 2) characterized by a typical laminae thickness of <1 mm. Detrital laminae are dominated by quartz, clay and organic matter, with some calcite, dolomite and feldspar grains, commonly showing graded bedding, a clay cap of and a thin lamina of amorphous organic matter (Fig. 2).

The laminated segments are interrupted by massive or graded ERDs attributed to abrupt mass wasting events (e.g., turbidities and slumps), triggered by earthquakes (Kagan et al., 2018), debris flows and slope instability (Ahlborn et al., 2018). Both laminae thickness and number of sub-laminae show statistically significant larger mean and variance during lake level rise (~27 Ka) than during lake level fall (~18 Ka; Table 1; Figs. 4 and 5).

Aragonite laminae thicknesses range between 51 and 22592.259  $\mu\text{m}$  ( $\mu=457\mu\text{m}$ ,  $\sigma=308\mu\text{m}$ ), and 53 and 6,020  $\mu\text{m}$  ( $\mu=642\mu\text{m}$ ,  $\sigma=529\mu\text{m}$ ) in the falling and rising segments, respectively. Detrital laminae thicknesses range from 27 to 13,136  $\mu\text{m}$  ( $\mu=312\mu\text{m}$ ,  $\sigma=638\mu\text{m}$ ), and 27 and 8,760  $\mu\text{m}$  ( $\mu=416\mu\text{m}$ ,  $\sigma=675\mu\text{m}$ ) in the falling and rising segments, respectively. The number of sub-laminae during the falling lake level ranges between 1 and 10 ( $\mu=1.2$ ,  $\sigma=0.7$ ), and between 1 and 17 ( $\mu=1.6$ ,  $\sigma=1.6$ ) in the rising lake level, respectively (Table 1). The MWW test indicates a significant difference between all studied proxies when comparing the two studied series (p value  $\ll 0.05$ ). The difference in dispersions of all variables between rising and falling lake levels are statistically significant (p value  $\ll 0.05$ ) except for detrital laminae, where the Ansari-Bradley test failed to reject the null hypothesis (p value  $\approx 0.12$ ). All three parameters show positive skewness of 2.1  $\mu\text{m}$  and 3.5  $\mu\text{m}$  for aragonite laminae thickness, 12.7  $\mu\text{m}$  and 7.6  $\mu\text{m}$  for detrital aragonite laminae, and 5.9 and 6.1 of the number of sub-laminae in sublaminae count during falling and rising lake levels, respectively. No substantially significant correlation exists between any two of the parameters in each time series, with the exception of a moderate correlation between the number of sub-laminae in the detrital laminae and laminae thickness (Fig. 34;  $r=0.52$  and  $0.61$ , for falling and rising lake levels, respectively; with p values  $\ll 0.005$ ).

Table 1 – Bulk statistical properties of the studied sedimentary series.

Rise/fall	Parameter	Unit	Min	Max	Median	Mean	Std. deviation	Skewness
fall	aragonite	$\mu\text{m}$	51	22592.259	388389	457458	308	2.1

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: Times New Roman, 10 pt, Font color: Auto, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: Times New Roman, 10 pt, Font color: Auto, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: Times New Roman, Font color: Auto

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: Times New Roman, 10 pt, Font color: Auto, English (United Kingdom)

Formatted: Font: Times New Roman, 10 pt, Font color: Auto, English (United Kingdom)

Formatted: Font: Times New Roman, 10 pt, Font color: Auto, English (United Kingdom)


Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: 10 pt, English (United Kingdom)

Formatted: Font: 10 pt, English (United Kingdom)
















Figure 4 – Time series of microfacies analysis of the studied varved sections. (a, b; blue) aragonite laminae thickness, (c, d; black) detritus laminae thickness, (e, f; red) number of floods identified by modifying the number of counted detrital sub-laminae as described in the text.

Formatted: Font: Not Bold






Figure 5 – Distributions and correlation scatter plots of the trivariate geological time series showing the distributions of (a) aragonite laminae thickness, (b) detritus laminae thickness and (c) number of sub-laminae counted in each detrital lamina. Plots d-f show the correlations between aragonite and detritus thickness (d,  $r^2=0.07$ ,  $0.03$ ), number of floods and detritus thickness (e,  $r^2=0.27$ ,  $0.37$ ), and number of floods and aragonite laminae thickness (f,  $r^2=0.03$ ,  $0.01$ )


580

#### 4.2 Detecting intra-record regime shifts

Two clusters of intense flood frequency were identified in each of the studied series (Fig. 3; Table 3). Although two clusters are identified, one of the clusters in each series appears more robust in comparison with the other, as it was identified using multiple window widths (Figs. S2-S5), and S2-10). The clusters are labeled as robust and less robust accordingly (Table 3). However, ~~we note it is noted~~ that this different robustness estimation could stem from an “edge effect” bias, that limits the analysis of segments close to the edges of the series ~~using a large range of window widths. The recurrence analyses demonstrate a substantial drop in both recurrence rate and the transitivity coefficient at the beginning of all clusters, although the first, less robust cluster, also depicts a rise in both parameters after the initial drop, which testifies to the reduced robustness of its identification as a cluster (Fig. 4 and 5)-when large window widths are applied. The RPs of aragonite series depict some short diagonal lines, whereas those of the sub-laminae show more spotty properties with blocky character (Figs. 6 and 7). An increase in both DET and L<sub>max</sub> is identified close to the clusters’ edges, although the RQA measures fluctuate throughout the series, and do not exhibit pronounced changes or trends with respect to the identified clusters. Both JRPs depict fairly low recurrence with spotty and patchy patterns, with some diagonal lines in the series of the falling lake level.~~

Table 3 – Identified clusters in the two studied series following the procedure Appendix B based on number of detrital sub-laminae.

	Level fall [index years]	Estimated robustness	Level rise [index years]	Estimated robustness
1 <sup>st</sup> cluster	47-138	Less robust	1-83	Less robust
2 <sup>nd</sup> cluster	<del>519487-</del> 662	Robust	<del>394363-</del> 534	Robust







Figure 6 – Recurrence analyses (a-c) Enlarged segments of the recurrence plots of the studied lake level fall depicting the recurrence plots for aragonite thickness (a), flood frequency (b) and the joint recurrence plot for the two proxies calculated using a maximum norm. Blue rectangle indicates (c). (d-f) Recurrence plots of the aragonite thickness (d), flood frequency (e) and the joint recurrence plot for the two proxies (f), with blue rectangles indicating the extent of data presented in (d-f). (d-f) depict a zoomed-in segment of corresponding plots a-c. Note the parallel diagonal lines that characterize the aragonite thickness (a,d), and indicate point on to an inherent quasi-periodic oscillatory behaviour, whereas the flood series (b,e) forms sporadic points and polygons that reflect a more chaotic or random behaviour. (g-i show the) The recurrence rate (RR) and the transitivity/determinism (DET) coefficient over using a running window size of 30 years and a single year step for aragonite thickness (g), floods (h) and the joint recurrence (i). (j-l) The laminarity (LAM) and the maximum length of the diagonal line within the sub-matrix ( $L_{max}$ ) coefficient using a running window size of 30 years and 20% overlap a single year step for aragonite thickness (j), floods (k) and the joint recurrence (l). The analyses were carried over after detrending the detrended data using the SSA  $RC^{(1)}$ . Selected parameters for the embedding are  $m$  (dimension),  $\tau$  (delay) and  $\epsilon$  (threshold) used for space phase reconstructions are presented in the figure. Red lines represent possible clusters. Pronounced non-stationarity is shown by a sharp decrease in recurrence and transitivity prior and during the possibly clusters identified independently of the recurrence analyses.







Figure 7—Recurrence analyses of— (a-c) Enlarged segments of the recurrence plots of the studied lake level rise depicting the recurrence-plots for aragonite thickness (a), flood frequency (b) and the joint recurrence plot for the two proxies calculated using a maximum norm. Blue rectangle indicates (c). (d-f) Recurrence plots of the aragonite thickness (d), flood frequency (e) and the joint recurrence plot for the two proxies (f), with blue rectangles indicating the extent of data presented in (d-f). d-f illustrate zoomed-in segment of the corresponding plots a-c. Note the parallel diagonal lines that characterize the aragonite thickness (a,d), and indicate point on to an inherent quasi-periodic oscillatory behaviour, whereas the flood series (b,e) forms sporadic points and polygons that reflect a more chaotic or random behaviour. g-i show the recurrence rate and the transitivity coefficient over (g-i) The recurrence rate (RR) and the determinism (DET) coefficient using a running window size of 30 years and a single year step for aragonite thickness (g), floods (h) and the joint recurrence (i). (j-l) The laminarity (LAM) and the maximum length of the diagonal line within the sub-matrix ( $L_{max}$ ) coefficient using a running window size of 30 years and 20% overlap, a single year step for aragonite thickness (j), floods (k) and the joint recurrence (l). The analyses were carried over after detrending the detrended data using the 1<sup>st</sup> SSA  $RC_{s(1)}$ . Selected parameters for the embedding are  $m$  (dimension),  $\tau$  (delay) and  $\epsilon$  (threshold) used for space phase reconstructions are presented in the figure. Red lines portray represent possible cluster. Pronounced non-stationarity is depicted by a sharp decrease in recurrence and transitivity prior and during the possibly clusters identified cluster independently of the recurrence analyses.




## 4.2 Detecting ~~periodical~~periodic components

635 Recurrence analyses of aragonite thicknesses indicate periodic or ~~quasi-periodic~~oscillatory behaviours, reflected by the ~~appearance of~~ diagonal lines ~~appearing throughout~~in the recurrence plot of the series (Figs. 4a,4d6 and 5a,5d7). Conversely, the flood frequency series present blocky and irregular patterns that indicate an underlying chaotic, random, or practically non-periodic process (Figs. 4b,4e6 and 5b,5e7). In particular, the segments identified as clusters demonstrate ~~decreased-recurrence rate and transitivity, suggesting some abrupt changes in the studied RQA measures that could indicate~~ a shift in the system's dynamics, both in aragonite thickness and in the number of sub-laminae. Although this ~~behavior~~behaviour is somewhat expected for the sub-laminae series that was used to determine clusters in the first place, it is also evident for the aragonite ~~data~~series.

645 Discontinuous short-term ~~increased power of~~ periodicities ~~with a period ranging between two and eight years of the annual-decadal band~~ appear sporadically throughout the entire dataset in both aragonite and flood frequency series (Fig. 6). ~~These~~8). ~~However, these short-term oscillations fail the significance test of the global wavelet spectrum against an AR(1) red noise process, but appear to be~~ ~~are not~~ significant in the analyses of the detrended data after subtracting  $RC^{(1)}$  (Figs. S14), which may indicate that their importance could be obscured by contrasting trend components. A statistically with an  $\alpha=0.05$ , and additionally do not pass the global significance test. Instead, a couple of the significant patches that appear within this spectral band appear within clusters as vertically stretched patches that range over a set of periodicities ranging from sub-decadal to a multidecadal (30-60 years) frequency bands, which indicate that these periodicities are attributed to the abrupt pulse of increased variance occurring during the identified clusters (Hochman et al., 2019). These several significant patches pass the global significance test after detrending using the  $RC^{(1)}$  of the SSA (Fig. S14). A centennial- to bicentennial-scale band (~120-250 years) periodic component is observed in both aragonite thickness series, although its ~~robustness is difficult to~~ ~~estimate~~significance cannot be determined as it falls largely outside the cone of influence, and it fails the cumulative area-wise significance test. A similar periodic component is also identified in the sub-laminae series during lake level rise, but not in the interval of lake level fall. ~~Altogether, the wavelet analyses do not suggest a robust detection of periodic components in any of the studied series.~~





660 Figure 8 – Wavelet and global-wavelet spectra of aragonite thickness and flood frequency during falling (left) and rising (right) episodes. Periodicities with significance level above 0.95 ( $\alpha=0.05$ ) of the cumulative area-wise significance test are depicted by a bold black line (Schulte et al., 2018; Schulte, 2016). Each triplot section shows the normalized data after normalization (a-d), the wavelet spectra (e-h) and the global wavelet spectra (blue) compared against a background red noise estimate the significance test of the global wavelet power spectra (dashed red/black, i-l). Significant peaks of arcs are marked in red (Schulte, 2019). Vertical dashed lines in (e-h) depict clusters identified as episodes of increased flood frequency. Note the loss of significance of that the short-term periodic component in the global wavelet spectra that is resolved, which fail the significance of the series, are significant in the analysis of the detrended series (Fig. S14).

670 The SSA RCs are characterized by multiple several periodic components (Figs. S6-S10-S13). The first two components are considered as the local trend component, and accordingly depict persistent “long-wave” periodic components with wave lengths >32 years. The following components show roughly three peaks characterized by periodicities of 2-3 years, 4-5 years, and 7-8 years. The multichannel SSA and its corresponding cross-wavelet analyses demonstrates that the periodic component

Formatted: Justified

with 4-5 years periodicity is relatively robust during lake level fall, but less persistent during lake level rise (Figs. S12 and S13)-periodic components of sub-decadal frequencies.

## 5 Discussion

### 5.1 The mechanistic link between mean properties and variability

Due to the unique hydroclimatic settings of the lake and its sedimentary record, the record cannot be directly interpreted from the perspective of a flood frequency analysis (e.g., Metzger et al., 2020), where the frequency and magnitude of floods in individual watersheds are studied. This is because the studied detrital sub-laminae in the ICDP-DSDDP core record the number of flood events (potentially lasting up to several days) that exceeded the threshold required to reach the coring site at the lake center (Ben-Dor et al., 2018), instead of recording discharge properties at individual watersheds (Nehorai et al., 2013). Furthermore, the bathymetry of the basin dictates that only large enough floods in tributaries, situated in front of the coring site on both sides of the escarpment, could have deposited detrital material at the coring site (Ben-Dor et al., 2018; Nehorai et al., 2013). Although this aspect masks some delicate features that could otherwise be extracted in the analyses of modern measurements, its length and established context with respect to regional climatic conditions testifies to its value.

By assuming that the dominant synoptic-scale circulation patterns during the late Pleistocene resembles modern conditions in the region (Enzel et al., 2008), it can be argued that flood events recorded as detrital sub-laminae at the depocenter of Lake Lisan can result from three main synoptic-circulation systems (e.g., Armon et al., 2019); the interplay between these three systems determines the mean climatic conditions (i.e., precipitation). Thus, the association of increased flood frequency with rising lake levels was initially interpreted as reflecting increased frequency and/or modulation of the track and depth of Mediterranean lows that deliver the majority of annual precipitation over the Dead-Sea watershed (Ben-Dor et al., 2018). However, a more detailed comparison of flood frequency with aragonite laminae thickness, that reflects annual inflow and dilution to some extent (Kolodny et al., 2005; Stein et al., 1997), reveals more subtle insights and a delicate interplay of hydroclimatic factors manifested through the watershed during opposing climatic regimes.

Both aragonite laminae thickness and flood frequency have larger mean and variance during lake level rise (Table 1). Because rising lake levels indicate a “wetter” climate on average, this relationship between mean and variance is similar to this relationship observed in modern precipitation (e.g., Morin et al., 2019). This relationship, thus, strengthens the interpretation of these sedimentary proxies as hydroclimatic proxies. Namely, the increased thickness of aragonite laminae and the increased number of detrital sub-laminae, in detrital laminae containing more than one sub-laminae, suggests that the episodes of rising lake level were characterized both by increased annual inflow and increased frequency of floods. As suggestions invoking increased flood frequency were raised both for wetter and drier conditions were made (e.g., Alpert et al., 2002; Rohling, 2013;

Yosef et al., 2019), this observation demonstrates that increased frequency of floods during wetter intervals in Mediterranean climate zone is the likely scenario. Furthermore, the coupling of increased observed thickness of aragonite laminae with increased flood frequency may support their interpretation by a common hydroclimatic mechanism, such as the increased occurrence of Mediterranean low pressure systems, suggested by earlier studies (Armon et al., 2019; Ben Dor et al., 2018; Enzel et al., 2008).

5.3 Periodical The poor correlation between aragonite thickness and the number of sub-laminae testifies to their low inter-dependence, and strengthen their interpretation as (nearly) independent proxies of the hydrological cycle in the Dead Sea watershed, in accordance with the previous interpretation of their roles in the hydrological cycle (Ben Dor et al., 2021; Ben Dor et al., 2018). Because of the unique hydroclimatic settings of the lake and its sedimentary record, the record cannot be directly interpreted from the perspective of a flood frequency analysis (e.g., Metzger et al., 2020), where the frequency and magnitude of floods in individual watersheds is analyzed. This is because detrital sub-laminae in the ICDP-DSDDP core record the number of flooding events, which potentially last up to several days (Nehorai et al., 2013), that exceeded the threshold required to reach the coring site at the lake depocenter (Ben Dor et al., 2018), instead of recording discharge properties at individual watersheds. Furthermore, the bathymetry of the basin dictates that only large-enough floods in tributaries situated in front of the coring site on both sides of the escarpment, could have deposited detrital material at the coring site (Fig. 1). Although this aspect masks some of the delicate features that could have been otherwise extracted such as in the analyses of modern measurements, its length and established context with respect to regional climatic conditions testifies to its value for interpreting hydroclimatic variability during episodes of climate change.

Available modern observations of sediment and water transport-paths into the lake, cannot unambiguously determine which synoptic systems affected the eastern Mediterranean when the studied sediment segments were deposited during the LGM. Nevertheless, considering available data that addresses this question, the likely suggestion that dominant weather regimes and synoptic circulation patterns during the LGM were similar to present is adopted for the sake of discussion in the framework of this work (Greenbaum et al., 2006; Amit et al., 2011; Enzel et al., 2008), with some possible modifications of their spatial properties (e.g., Goldsmith et al., 2017; Keinan et al., 2019). By considering that the dominant synoptic-scale circulation patterns during the late Pleistocene resemble modern observations in the region (Enzel et al., 2008), it can be argued that flooding events recorded as detrital sub-laminae at the depocenter of Lake Lisan can be attributed to the three key synoptic patterns that govern precipitation over the eastern Mediterranean (e.g., Armon et al., 2019). It is further suggested that the interplay and relative frequency of these three systems determine mean climatic conditions, that determine the Dead Sea lake level and ultimately propagate into its sedimentary record (Weber et al., 2021; Waldmann et al., 2017). Thus, the association of increased flood frequency with rising lake levels was interpreted as reflecting increased frequency and/or modulation of the Mediterranean Lows characteristics that deliver the majority of annual precipitation over the Dead Sea watershed, and hence determine annual inflow into the lake (Saaroni et al., 2010). However, a more detailed comparison of flood frequency with aragonite laminae thickness, that reflects annual inflow into the lake and the dilution of its epilimnion (Kolodny et al., 2005;


740 [Stein et al., 1997; Ben Dor et al., 2021](#)), reveals more subtle insights and a delicate interplay of hydroclimatic factors manifested through the watershed during opposing climatic regimes.

745 Both aragonite laminae thickness and flood frequency have larger mean and variance during lake level rise (Table 1). Because rising lake levels indicate a “wetter-on-average” climate, this relationship between mean and variance is similar to that observed in modern hydrologic parameters such as precipitation (e.g., [Morin et al., 2019](#)). The observed relationship between the two studied proxies, thus, strengthens the interpretation of these sedimentary proxies as hydroclimatic proxies. Namely, the increased thickness of aragonite laminae and the increased number of detrital sub-laminae suggests that the episodes of rising lake level were characterized both by increased annual inflow and increased flood frequency. As suggestions invoking increased flood frequency were raised both for wetter and drier conditions were made (e.g., [Alpert et al., 2002; Rohling, 2013; Yosef et al., 2019](#)), this observation demonstrates that increased frequency of floods during wetter intervals in Mediterranean climate zones is the likely scenario. Furthermore, the observed coupling of increased thickness of aragonite laminae with increased flood frequency may support their formation by a common hydroclimatic mechanism, such as the increased occurrence of Mediterranean low pressure systems ([Armon et al., 2019; Enzel et al., 2008; Ben Dor et al., 2018; Goldreich et al., 2004](#)).

### 750 **5.3 Periodic components and variability hydroclimatological pacing**

755 One of the goals in studying hydroclimatological proxies is identifying key periodical components that pace short-term variability under different mean climatic regimes (e.g., [Ghil et al., 2002; Grinsted et al., 2004](#)). Because (a) the thickness of aragonite laminae was suggested to reflect annual hydroclimatic conditions ([Kolodny et al., 2005](#)) One of the goals in studying hydroclimatological proxies is identifying key periodic components that pace short-term variability under different mean climatic regimes (e.g., [Ghil et al., 2002; Grinsted et al., 2004](#)). Because (a) the thickness of aragonite laminae was suggested to reflect hydroclimatic ([Kolodny et al., 2005](#)) and limnologic conditions ([Ben Dor et al., 2021](#)), and (b) periodicities attributed to solar cycles were previously identified by spectral analyses of laminae thickness at the exposed eo-eval MIS2 White Cliff Member of the Lisan Formation at Masada ([Prasad et al., 2004](#)), a similar attempt was made in this study as well. [Prasad et al., \(2004\)](#) report periodicities of 50-60 years, similar to those identified here (Figs. 7, S12 and S13). Additionally, wavelet analyses of eo-eval White Cliff Member outcrop the Lisan Formation at Masada deposited during MIS2 ([Prasad et al., 2004](#)), a similar attempt was made in this study identify similar periodic components as well. [Prasad et al., \(2004\)](#) report periodicities of 50-60 years, similar to those identified here (Figs. 8, S14). Additionally, previously reported  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  from ~200 aragonite laminae recovered from MIS2 exposures of the Lisan Formation at Masada and Perazim Valley outcrops were hereby conducted analyzed using wavelet analyses. Because the systematics of these isotopic compositions are distinctly different, these proxies bear different implication on environmental conditions in the lake and its surroundings. More specifically,  $\delta^{18}\text{O}$  is directly influenced by hydroclimatology, whereas  $\delta^{13}\text{C}$  is primarily affected by biological activity. More specifically,  $\delta^{18}\text{O}$  is directly influenced by hydroclimatology, whereas  $\delta^{13}\text{C}$  is primarily affected by biological activity ([Kolodny et al., 2005](#)).

775 Nevertheless, it appears that because the extent of biological activity in Lake Lisan depended on freshwater inflow that also replenished its surface water with required nutrients (Begin et al., 2004), the two proxies share similar pacing and are broadly characterized by non-persistent periodic band of annual to decadal scales that fail the cumulative area-wise significance test at the  $\alpha=0.05$  level, although the coherence of this band between the signals is significant, and shows similar phase relationship (Fig. 9).



780 **Figure 9** – Wavelet (a-d) and cross-wavelet (e-f) analyses of  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  in two segments of the Masada (30 Ka) and Perazim (25 Ka) exposures of the White Cliff Member (Lisan Formation) deposited during the last glacial over the lake's shelf. Areas with significance level above 0.95 ( $\alpha=0.05$ ) are marked by a thick black line. Data is from (Kolodny et al., 2005). Nevertheless, it appears that because the extent of biological activity in the saline Lake Lisan depended on freshwater inflow that also replenished its surface water with required nutrients (Begin et al., 2004), the two proxies seem to share similar pacing and are broadly characterized by non-persistent periodical components of 2-6 years (Fig. 7), similar to those observed in the deep core (Fig. 6).

Formatted: Caption, Don't keep with next




Figure 7—Wavelet (a-d) and cross-wavelet analyses of  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  in two segments of the Masada (30-Ka) and Perazim (25-Ka) exposures of the White Cliff Member (Lisan Formation) deposited during the last glacial over the lake's shelf. Areas with significance level above 0.9 ( $\alpha=0.1$ ) are marked by a thick black line. Data is from (Kolodny et al., 2005).

The North Atlantic (NAO) and the Eastern Atlantic (EA) oscillations are commonly considered as affecting interannual precipitation variability over the eastern Mediterranean (Feldstein, 2000; e.g., Feldstein and Dayan, 2008; Feliks et al., 2010; Krichak et al., 2002; Seager et al., 2019), primarily due to their reported effect on discrete precipitation-bearing synoptic patterns over the eastern Mediterranean (e.g., Black, 2012). Others, on the other hand, have proposed that, unlike western Turkey, the direct impact of NAO on precipitation in the Levant is minimal (Enzel et al., 2003; Seager et al., 2020; Ziv et al., 2006). Although such relationships are hard to identify in geological records, a North Atlantic impact over the Dead Sea hydrology during the last glacial was identified at lower frequency by lake level reconstructions, where abrupt lake level drops were identified during Heinrich events (Bartov et al., 2003). Additionally, a NAO-like periodic component was also identified in laminated halite sequences during the last interglacial (Palehan et al., 2017). However, modern observations (Sirota et al., 2017), may indicate that this pacing is working through temperature variations, which bear significant implications on halite deposition, rather than through precipitation and hydrologic forcing.

Century-long precipitation data from the Kfar Giladi and Jerusalem stations correlate well with pre-regulated modern Dead Sea levels, and are thus considered as recorders of mean hydrological conditions over the northern and central Dead Sea watershed in recent and past times (Enzel et al., 2003; Morin et al., 2019). Thus, available records of annual precipitation and the number of annual rainy days in both stations were also compared here to winter (DJF) NAO and EA indices using wavelet and cross-wavelet spectra in order to detect common pacing and periodic components (Fig. S15 and S16; NOAA, 2020).



810 Although these records are relatively short, spanning less than a century, they show little similarity with winter NAO and EA. However, short-term fluctuations with periodicity of 2-6 years were observed in the ICDP-DSDDP cores (Figs. 6, S6, S9-S14); in the isotopic composition of aragonite (Fig. 7; Kolodny et al., 2005), in modern precipitation records and in winter (DJF) NAO and EA (Figs. S15 and S16). This similarity, although not robust, could indeed point to a weak or non-linear effect of these teleconnections on precipitation in the eastern Mediterranean (e.g., Black, 2012). Alternatively, this can point to difficulties in extracting these delicate relationships from sedimentary archives that ultimately record a chain of processes and convoluted impacts of hydrological and limnological processes involved in laminae formation and post-depositional changes; for example, these may include non-homogenous spread of fine-grained sediments over the lake floor and the effects of inflows from different catchments of multiple sizes as well as temperature.

820 The North Atlantic oscillation (NAO) and the Eastern Atlantic (EA) patterns were considered to affect interannual precipitation variability over the eastern Mediterranean (Feldstein and Dayan, 2008; Feliks et al., 2010; Krichak et al., 2002; Seager et al., 2019), primarily due to their reported effect on discrete precipitation-bearing synoptic patterns over the eastern Mediterranean (e.g., Black, 2012). Other studies, on the other hand, have suggested that unlike western Turkey, NAO shows only minimal direct impact on precipitation in the Levant, and that it might affect temperature instead (Enzel et al., 2003; Seager et al., 2020; Ziv et al., 2006). Although such relationships are hard to identify in geological records, a North Atlantic impact over the Dead Sea hydrology during the last glacial was identified by lake level reconstructions, where abrupt lake level drops took place during Heinrich events (Bartov et al., 2003). Additionally, a periodic component of a sub-decadal to decadal frequency bands (4-5, 7-8 and ~11 years) was also identified in laminated halite sequences deposited during the last interglacial and was attributed to NAO (Palchan et al., 2017). However, modern observations, may indicate that this pacing is manifested by temperature variations, which bear determine halite deposition, rather than hydrologic forcing (Sirota et al., 2017, 2018).

830 Century-long precipitation data from the Kfar Giladi and Jerusalem stations correlate well with pre-regulated modern Dead Sea levels, and are thus considered as recorders of mean hydrological conditions over the northern and central parts of the Dead Sea watershed in recent and past times (Enzel et al., 2003; Morin et al., 2019). Thus, available precipitation records in both stations were also compared to the winter (DJF) NAO and EA indices using wavelet and wavelet coherence in order to detect common pacing and periodic components (Figs. S15 and S16; NOAA, 2020). These relatively short records are do not exhibit significant periodic components in the series, and no significant coherence as well. However, although the available records do not exhibit significant periodic behaviour (at  $\alpha=0.05$ ), increased non-significant spectral power of short-term fluctuations are evident in the ICDP-DSDDP cores (Figs. 8, S6, S10-S13), the isotopic composition of aragonite laminae (Fig. 9; Kolodny et al., 2005), and modern precipitation records and winter indices of teleconnection patterns (Figs. S15 and S16). This similarity that is not statistically significant, could either point at the lack of influence of NAO and EA on hydroclimate or at a weak and non-linear effect of these teleconnection patterns on precipitation in the eastern Mediterranean (Black, 2012). Alternatively, this can point at the inherent difficulties in extracting such delicate relationships from sedimentary archives that

ultimately record the interaction of hydrological and limnological processes involved in laminae formation and post depositional changes, such as the non-homogenous spread of fine-grained sediments over the lake floor and the effects of inflows from different catchments.

### 845 5.3 Clusters and regime transitions

855 The studied episodes demonstrate pronounced clustering that emerged by the application of the Monte Carlo-based method that we developed to identify clusters of intense flooding (Appendix B). Similar, non-Poisson and non-uniform flood frequency distributions were also noted in other high-resolution palaeo-hydrological records (e.g., Witt et al., 2016) as well as modern flood data (Metzger et al., 2020). When this is considered together with reduced recurrence rate during clusters, the increased frequency of floods during clusters could be indicative of a hydroclimatic regime shift into a less persistent, possibly less-ordered or chaotic regime (Figs. 6-7). In addition, the clusters observed during lake level fall are characterized by flood frequencies similar to those of background episodes during lake level rise. This suggests that the wetter episodes are characterized by increased mean precipitation and variability, as well as increased frequency of intense storms, which is in agreement with modern and more recent observations (Morin et al., 2019).

860 The studied episodes demonstrate pronounced centennial-scale clustering of flooding events (Fig. 3). Similarly, other high-resolution palaeo-hydrological records (e.g., Witt et al., 2017) and modern observations (e.g., Metzger et al., 2020) have demonstrated non-uniform and non-Poisson flood frequencies. In addition, the clusters observed during lake level fall are characterized by flood frequencies similar to those in background episodes during lake level rise (Figs. S17 and S18). This suggests that the wetter episodes are characterized by increased mean precipitation and variability, as well as increased frequency of intense storms, which is in agreement with modern and more recent observations (Morin et al., 2019).

865 The comparison of aragonite laminae thickness of the two clusters were identified in each of the studied late-Pleistocene intervals, reveal revealing opposing properties when their increased flood frequencies are compared with aragonite thicknesses. The opposing characteristics of the two clusters are, which become evident by when comparing their running mean (calculated as  $RC^{(1)}$ ) and dispersion, which is calculated as the running standard deviation of the residuals (after subtracting  $RC^{(1)}$ ) using a window of 30 years (Figs. S1710 and S1811). In each of the studied intervals, one cluster demonstrates increased mean and variance in both flood frequency and aragonite thickness, whereas the other cluster is characterized by increased mean flood frequency, but also by similar mean and variance of aragonite thickness (Figs. S19 and S20). Similarly, contrasting properties of the two clusters in each of the two intervals are revealed by their recurrence analyses, where one cluster shows a more pronounced drop in recurrence rate and transitivity than the other (Figs. 6-7). These two distinct approaches further demonstrate that the underlying processes responsible for each cluster are distinct, and in turn are also different from the similar to that of background processes. Altogether, episodes (Figs. S17 and S18). In order to address these observations call for a


unified explanation that can account for this discrepancy ~~in terms of~~ hereby suggested by considering the modern hydroclimatic ~~regimes~~regime and dominant synoptic circulation patterns.

Although modern records are relatively short, and the comparison of sparse hydrological measurements with the detailed framework of atmospheric circulation only permits cautious conclusions, the abovementioned observations can be explained using the modern synoptic framework by considering two distinct synoptic and hydroclimatic scenarios that govern flood formation and flood clustering over the catchments facing the ICDP-DSDDP coring site. Because most of the precipitation over the Dead Sea watershed is delivered by extra-tropical cyclones (Mediterranean lows) during winter months (e.g., Enzel et al., 2003; Saaroni et al., 2010; Ziv et al., 2006), when these cyclones are deep and characterized by a slightly southern track, they can pass slightly slower over the region and generate much more rainfall, and therefore they effectively deliver more precipitation to the watershed (e.g., Ben Dor et al., 2018; Ziv et al., 2006). Thus, they effectively increase annual precipitation, while increasing the chances of generating floods over the catchments that face the ICDP-DSDDP coring site (Armon et al., 2019; Belaehsen et al., 2017; Goldreich et al., 2004). As suggested above, this scenario could account for the observed increase in both mean and variance of flood frequency and aragonite laminae thickness, observed in one of the clusters of each studied series (Fig. S19 and S20).

During the other cluster, on the other hand, the increased flood frequency is uncoupled with a clear increase in the mean and/or variance of the aragonite laminae thickness. Studies of modern floods and their synoptic settings suggest that floods can also be generated by other two synoptic systems: active Red Sea troughs (ARST) and jet stream disturbances (i.e., Tropical Plumes). These synoptic conditions can generate significant floods over the small catchments surrounding the ICDP-DSDDP coring site, thus having the potential to deliver sediments, but because of their spatiotemporal properties and frequency, would have negligible contribution to total inflow into the lake (Armon et al., 2019). However, in present-day ARSTs are more frequent than Tropical Plumes, and are characterized by high peak discharge and relatively low volume floods (Armon et al., 2018; Shentsis et al., 2012) the second cluster may be thus explained by increased ARST frequency. This would lead to increased flood frequency (more sub-laminae) without drastically increasing annual inflow (similar aragonite thickness).

Although modern records are relatively short, and the comparison of sparse hydrological measurements with the detailed framework of atmospheric circulation only permits cautious suggestions at this stage, the abovementioned observations can be explained using the modern synoptic framework by considering two distinct synoptic and hydroclimatic scenarios that govern flood generation and their clustering over the catchments that face the ICDP-DSDDP coring site. Most of the precipitation over the Dead Sea watershed is delivered by extra-tropical cyclones (Mediterranean lows) during the winter months (e.g., Enzel et al., 2003; Saaroni et al., 2010; Ziv et al., 2006). When these cyclones are deep and have a southern track, their resulted precipitating clouds can pass impact the Dead Sea watershed for longer periods and generate more rainfall, and therefore effectively deliver more precipitation to the watershed (Ziv et al., 2006; Saaroni et al., 2010). Thus, under these conditions they increase annual precipitation, while increasing the chances of generating floods over the catchments that face

the ICDP-DSDDP coring site (Armon et al., 2019; Belachsen et al., 2017; Goldreich, 2004). As suggested above, this scenario could account for the observed coupled increase in both mean and variance of flood frequency and aragonite laminae thickness, observed in one of the clusters in each of studied series (Fig. S17 and S18).




**Figure 10** –running statistical properties of aragonite laminae (a-d) and flood frequency (e-h) for the studied lake level fall. (a,e) the measured data (b,f), the residuals (after subtracting the  $RC^{(1)}$ ) (c,g),  $RC^{(1)}$  and running std. deviation (30 years window), and the coefficient of variation calculated as  $\frac{std}{RC^{(1)}}$  (d,h).

The general understanding of precipitation patterns induced by different synoptic systems in the Dead Sea watershed depicts a “de-coupling” of annual inflow into the lake, which depends on annual precipitation over the northern parts of the watershed.

920 and floods reaching the coring site. This is because the frequency and intensity of (mainly *eastern Mediterranean Lows*  
determines annual precipitation over the watershed (Saaroni et al., 2010) and flood frequency in the relevant ephemeral streams  
(Goldreich et al., 2004), whereas the contribution of the other synoptic systems to annual precipitation by far, is less substantial  
(Armon et al., 2019; Marra et al., 2021). This is also evident by the low correlation ( $r^2 = 0.086$ ) of major floods (return period  
>5 years) in the Negev Desert (Kahana et al., 2002) and precipitation in Jerusalem (Fig. R1), which found to be closely  
correlated with Dead Sea lake level, and hence with annual inflow into the lake (Enzel et al., 2003). Thus, although this cannot  
be directly proven for the LGM, these modern observations are hereby considered as means to decipher the sedimentary record  
925 (e.g., Enzel et al., 2008; Goldsmith et al., 2017).

930 During the other cluster, on the other hand, the observed increased flood frequency is not coupled with increased mean and  
variance of aragonite laminae thickness. Studies of modern floods and their synoptic settings indicate that floods over the  
eastern Mediterranean are also generated by two other synoptic systems: the active Red Sea trough (ARST) and disturbances  
to the subtropical jet stream (or Tropical Plumes). These synoptic conditions can generate significant floods over the small  
catchments surrounding the ICDP-DSDDP coring site, thus having the potential to deliver sediments, but owing to their  
spatiotemporal properties and frequency, they provide negligible contribution to total inflow into the lake (Armon et al., 2019).  
935 Under current conditions ARSTs are more frequent than Tropical Plumes, and are characterized by high peak discharge and  
relatively low volume floods (e.g., Armon et al., 2018; Shentsis et al., 2012) the second cluster may be therefore explained by  
increased frequency of ARSTs during decades of decreased mean annual precipitation. This scenario would result in increased  
flood frequency, recorded by more sub-laminae, without substantially increasing annual inflow, thus exhibiting aragonite  
thicknesses similar to background periods.



940 **Figure 11 – running statistical properties of aragonite laminae (a-d) and flood frequency (e-h) for the studied lake level rise. (a,e) the measured data (b,f), the residuals (after subtracting the  $RC^{(1)}$ ) (c,g),  $RC^{(1)}$  and running std. deviation (30 years window), and the coefficient of variation calculated as  $\frac{std}{RC^{(1)}}$  (d,h).**

## 6 Conclusions

945 The short-term hydroclimatic variability of opposing climatic trends in the Levant was studied using multiple statistical in detail through several analyses of two annually-resolved varve sequences of the ICDP-DSDDP cores, representing opposing mean climates recorded by contrasting lake level trends. This unique sedimentary record complements the otherwise short and sparse modern climatic and hydrological records; it adds, and elucidates aspects and properties of longer time series of regional

hydroclimatology. on centennial time scale. By the analyses of two distinct sedimentary proxies that reflect annual inflow and flood frequency, and by their comparison with modern climatic data from a and synoptic perspective observations, new insights on hydroclimatic stationarity during late Pleistocene climate changes are/were revealed. These findings improve our understanding of short-term late Pleistocene hydroclimatic variability; moreover, they point to factors pacing hydroclimatic aspects and periodic components. Several

Key conclusions arise:

1. The dispersion of aragonite laminae thickness and flood frequency in the ICDP-DSDDP cores are fairly correlated/change with their corresponding mean, similar to as observed in modern hydrologic observations. These findings strengthen the interpretation of these proxies as recorders of hydroclimatic phenomena in the Dead Sea watershed. During the “wetter” interval, which is characterized by lake level rise, both the mean and the variance of these proxies are larger than during the studied episode of falling lake level.
2. Singular No significant periodic components were identified in the studied records using singular spectrum, wavelet and recurrence analyses reveal. However, it is suggested that this could stem from the interaction of climatic, hydrologic, and limnogeological processes that may have increased the noise/signal ratio of the studied proxies. Thus, the integration of these methods suggest that some short-term quasi-periodical/oscillatory components that affect/could have affected annual precipitation and flood frequency during the late Pleistocene. Although the inherent link between aragonite thickness and annual inflow is probably, but were not linear, the two proxies show similar periodical components of 2-3, 4-5, and 7-8 years, that could stem from the interaction of quasi-periodic 3-4 years components, possibly related/found to the North Atlantic Oscillation, be significant.
3. Flood frequencies over the studied series demonstrate strong hydro-climatological regime shifts operating at the centennial time scale. The clusters show Floods are clustered into episodes of two distinct characteristics in terms of the relationship/relationships between mean and dispersion of flood frequency and annual inflow. Namely, in each studied series one cluster is characterized by increased mean and variance of the two proxies, whereas the other one shows cluster is characterized by increased mean and variance only in dispersion of flood frequency, but not in annual precipitation. This implies that the clusters were generated by a distinct hydroclimatic regimes/regime, that was characterized by different relative dominance of various synoptic systems/circulation patterns that lead to/resulted in increased floods frequency at a decadal scale, but without increasing annual precipitation. More specifically, it may suggest that clusters of increased flood frequencies could either result from the increased frequency of Mediterranean Lows or of Active Red Sea Troughs. Such regime shifts could also affect modern and future conditions that would manifest as drastic hydroclimatic shifts at decadal to centennial scales.

**Formatted:** Indent: Before: 0.63 cm, Hanging: 0.63 cm,  
Line spacing: 1.5 lines

## 7 Appendices

See attached file

[Supplementary 1—figures](#)

[Supplementary 2—non-parametric test developed for identifying clusters](#)

## 980 **8-Code availability**

All code that was used for this research is available upon request from the corresponding author.

## **9-Data availability**

All the data that was used for this research is available in the appendices or upon request from the corresponding author.

## **10-Competing interests**

985 The authors declare that they have no conflict of interest.

## **Acknowledgements**

This study is a contribution to the PALEX project “Paleohydrology and Extreme Floods from the Dead Sea ICDP core”, funded by the DFG to A. Brauer, Y. Enzel, E. Morin, and Y. Erel (grant no. BR2208/13-1/-2). The authors acknowledge the support and contribution of laboratory staff and technicians in the GFZ, where preparation of thin-sections and photography were carried. We thank J. Mingram, N. Nowaczyk, B. Brademann, F. Ott, N. Dräger and M. Köppel for technical support and fruitful discussions. Y.B. is also grateful for a scholarship from the Advanced School of Environmental Studies, the Hebrew University of Jerusalem, and from the Rieger Foundation-Jewish National Fund program for environmental studies.

## **References**

- 995 [Ahlborn, M., Armon, M., Ben Dor, Y., Neugebauer, I., Schwab, M. J., Tjallingii, R., Shoqeir, J. H., Morin, E., Enzel, Y. and Brauer, A.: Increased frequency of torrential rainstorms during a regional late Holocene eastern Mediterranean drought, \*Quat. Res. \(United States\)\*, 89\(2\), doi:10.1017/qua.2018.9, 2018.](#)
- [Allen, K. J., Hope, P., Lam, D., Brown, J. R. and Wasson, R. J.: Improving Australia’s flood record for planning purposes—can we do better?, \*Aust. J. Water Resour.\*, 00\(00\), 1–10, doi:10.1080/13241583.2020.1745735, 2020.](#)
- 1000 [Alpert, P., Ben Gai, T., Baharad, A., Benjamini, Y., Yekutieli, D., Colacino, M., Diodato, L., Ramis, C., Homar, V., Romero, R., Michaelides, S. and Manes, A.: The paradoxical increase of Mediterranean extreme daily rainfall in spite of decrease in total values, \*Geophys. Res. Lett.\*, 29\(11\), 31–31–4, doi:10.1029/2001GL013554, 2002.](#)
- [Ansari, A. R. and Bradley, R. A.: Rank-Sum Tests for Dispersions, \*Ann. Math. Stat.\*, 31\(4\), 1174–1189 \[online\] Available](#)



from: <https://www.jstor.org/stable/pdf/2237814.pdf?refreqid=excelsior%3A3d25ddb377eccc2621e692b57882f61>, 1960.

1005 Armon, M., Dente, E., Smith, J. A., Enzel, Y. and Morin, E.: Synoptic-scale control over modern rainfall and flood patterns in the Levant drylands with implications for past climates, *J. Hydrometeorol.*, 19(6), 1077–1096, doi:10.1175/JHM-D-18-0013.1, 2018.

Armon, M., Morin, E. and Enzel, Y.: Overview of modern atmospheric patterns controlling rainfall and floods into the Dead Sea: Implications for the lake's sedimentology and paleohydrology, *Quat. Sci. Rev.*, 216, 58–73, doi:10.1016/j.quascirev.2019.06.005, 2019.

1010 Baker, V. R.: Paleoflood hydrology: Origin, progress, prospects, *Geomorphology*, 101(1–2), 1–13, doi:10.1016/j.geomorph.2008.05.016, 2008.

Bartov, Y., Stein, M., Enzel, Y., Agnon, A. and Reches, Z.: Lake levels and sequence stratigraphy of Lake Lisan, the late Pleistocene precursor of the Dead Sea, *Quat. Res.*, 57(1), 9–21, doi:10.1006/qres.2001.2284, 2002.

Bartov, Y., Goldstein, S. J., Stein, M. and Enzel, Y.: Catastrophic aid episodes in the Eastern Mediterranean linked with the North Atlantic Heinrich events, *Geology*, 31(5), 439–442, doi:10.1130/0091-7613(2003)031<0439:CAEITE>2.0.CO;2, 2003.

1015 Bartov, Y., Enzel, Y., Porat, N. and Stein, M.: Evolution of the Late Pleistocene Holocene Dead Sea Basin from Sequence Stratigraphy of Fan Deltas and Lake Level Reconstruction, *J. Sediment. Res.*, 77(9), 680–692, doi:10.2110/jsr.2007.070, 2007.

Begin, Z. B., Nathan, Y. and Ehrlich, A.: Stratigraphy and facies distribution in the Lisan Formation—new evidence from the area south of the Dead Sea, Israel, *Isr. J. Earth Sci.*, 29, 182–189, 1980.

1020 Begin, Z. B., Stein, M., Katz, A., Machlus, M., Rosenfeld, A., Buchbinder, B. and Bartov, Y.: Southward migration of rain tracks during the last glacial, revealed by salinity gradient in Lake Lisan (Dead Sea rift), *Quat. Sci. Rev.*, 23(14–15), 1627–1636, doi:10.1016/j.quascirev.2004.01.002, 2004.

Belachsen, I., Marra, F., Peleg, N. and Morin, E.: Convective rainfall in a dry climate: Relations with synoptic systems and flash flood generation in the Dead Sea region, *Hydrol. Earth Syst. Sci.*, 21(10), 5165–5180, doi:10.5194/hess-21-5165-2017, 2017.

1025 Black, E.: The influence of the North Atlantic Oscillation and European circulation regimes on the daily to interannual variability of winter precipitation in Israel, *Int. J. Climatol.*, 32(11), 1654–1664, doi:10.1002/joc.2383, 2012.

Blackman, R. B. and Tukey, J. W.: The Measurement of Power Spectra from the Point of View of Communications Engineering—Part II, *Bell Syst. Tech. J.*, 37(2), 485–569, doi:10.1002/j.1538-7305.1958.tb01530.x, 1958.

1030 Bookman, R., Bartov, Y., Enzel, Y. and Stein, M.: Quaternary lake levels in the Dead Sea basin: Two centuries of research, *Spec. Pap. Geol. Soc. Am.*, 401(10), 155–170, doi:10.1130/2006.2401(10), 2006.

Brauer, A., Endres, C., Günter, C., Litt, T., Stebich, M. and Negendank, J. F. W.: High resolution sediment and vegetation responses to Younger Dryas climate change in varved lake sediments from Meerfelder Maar, Germany, *Quat. Sci. Rev.*, 18(3), 321–329, doi:10.1016/S0277-3791(98)00084-5, 1999.

1035 Brauer, A., Mangili, C., Moseariello, A. and Witt, A.: Palaeoclimatic implications from micro-facies data of a 5900 varve time series from the Pianico interglacial sediment record, southern Alps, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 259(2–3), 121–

135, doi:10.1016/j.palaeo.2007.10.003, 2008.

Broomhead, D. S. and King, G. P.: Extracting qualitative dynamics from experimental data, *Phys. D Nonlinear Phenom.*, 20(2-3), 217–236, doi:10.1016/0167-2789(86)90031-X, 1986.

1040 Campins, J., Genovés, A., Picornell, M. A. and Jansà, A.: Climatology of Mediterranean cyclones using the ERA-40 dataset, *Int. J. Climatol.*, 31(11), 1596–1614, doi:10.1002/joc.2183, 2011.

Coianiz, L., Ben-Avraham, Z., Stein, M. and Lazar, M.: Spatial and temporal reconstruction of the late Quaternary Dead Sea sedimentary facies from geophysical properties, *J. Appl. Geophys.*, 160, 15–27, doi:10.1016/j.jappgeo.2018.11.002, 2019.

1045 David-Novak, H., Ben-Morin, E. and Enzel, Y.: Modern extreme storms and the rainfall thresholds for initiating debris flow on the hyperarid western escarpment of the Dead Sea, Israel, *Bull. Geol. Soc. Am.*, 116(5–6), 718–728, doi:10.1130/B25403.2, 2004.

Dayan, U. and Morin, E.: Flash flood—producing rainstorms over the Dead Sea: A review, *New Front. Dead Sea ...*, 2401(04), 53–62, doi:10.1130/2006.2401(04), 2006.

1050 Donges, J.-F., Donner, R. V., Trauth, M. H., Marwan, N., Schellnhuber, H. J. and Kurths, J.: Nonlinear detection of paleoclimate variability transitions possibly related to human evolution, *Proc. Natl. Acad. Sci. U. S. A.*, 108(51), 20422–20427, doi:10.1073/pnas.1117052108, 2011.

Ben-Dor, Y., Armon, M., Ahlborn, M., Morin, E., Erel, Y., Brauer, A., Schwab, M. J. M. J., Tjallingii, R. and Enzel, Y.: Changing flood frequencies under opposing late Pleistocene eastern Mediterranean climates, *Sci. Rep.*, 8(1), 1–10, doi:10.1038/s41598-018-25969-6, 2018.

1055 Ben-Dor, Y., Neugebauer, I., Enzel, Y., Schwab, M. J., Tjallingii, R., Erel, Y. and Brauer, A.: Varves of the Dead Sea sedimentary record, *Quat. Sci. Rev.*, 215, 173–184, doi:10.1016/j.quaseirev.2019.04.011, 2019.

Eckmann, J. P., Oliffson-Kamphorst, O. and Ruelle, D.: Recurrence plots of dynamical systems, *Epl*, 4(9), 973–977, doi:10.1209/0295-5075/4/9/004, 1987.

1060 Enzel, Y., Bookman (Ken-Tor), R., Sharon, D., Gvirtzman, H., Dayan, U., Ziv, B. and Stein, M.: Late Holocene climates of the Near East deduced from Dead Sea level variations and modern regional winter rainfall, *Quat. Res.*, 60(3), 263–273, doi:10.1016/j.yqres.2003.07.011, 2003.

Enzel, Y., Amit, R., Dayan, U., Crouvi, O., Kahana, R., Ziv, B. and Sharon, D.: The climatic and physiographic controls of the eastern Mediterranean over the late Pleistocene climates in the southern Levant and its neighboring deserts, *Glob. Planet. Change*, 60(3–4), 165–192, doi:10.1016/j.gloplacha.2007.02.003, 2008.

1065 Enzel, Y., Amit, R., Grodek, T., Ayalon, A., Lekaeh, J., Porat, N., Bierman, P., Blum, J. D. and Erel, Y.: Late Quaternary weathering, erosion, and deposition in Nahal Yael, Israel: An “impact of climatic change on an arid watershed”?, *Bull. Geol. Soc. Am.*, 124(5–6), 705–722, doi:10.1130/B30538.1, 2012.

Feldstein, S. B.: The Timescale, Power Spectra, and Climate Noise Properties of Teleconnection Patterns, *American Meteorological Society*, 2000.

1070 Feldstein, S. B. and Dayan, U.: Circumglobal teleconnections and wave packets associated with Israeli winter precipitation,

- Q.-J. R. Meteorol. Soc., 134(631), 455–467, doi:10.1002/qj.225, 2008.
- Feliks, Y., Ghil, M. and Robertson, A. W.: Oscillatory climate modes in the eastern Mediterranean and their synchronization with the North Atlantic Oscillation, *J. Clim.*, 23(15), 4060–4079, doi:10.1175/2010JCLI3181.1, 2010.
- Floecas, H. A., Simmonds, I., Kouroutzoglou, J., Keay, K., Hatzaki, M., Bricolas, V. and Asimakopoulos, D.: On cyclonic tracks over the Eastern Mediterranean, *J. Clim.*, 23(19), 5243–5257, doi:10.1175/2010JCLI3426.1, 2010.
- Garber, R. A., Levy, Y. and Friedman, G. M.: The sedimentology of the Dead Sea, Carbonates and Evaporites, 2(1), 43–57, doi:10.1007/BF03174303, 1987.
- Garfunkel, Z.: Internal structure of the Dead Sea leaky transform (rift) in relation to plate kinematics, *Tectonophysics*, 80(1–4), 81–108, doi:10.1016/0040-1951(81)90143-8, 1981.
- Garfunkel, Z. and Ben Avraham, Z.: The structure of the Dead Sea basin, *Tectonophysics*, 266(1–4), 155–176, doi:10.1016/S0040-1951(96)00188-6, 1996.
- Ghil, M., Allen, M. R., Dettinger, M. D., Ide, K., Kondrashov, D., Mann, M. E., Robertson, A. W., Saunders, A., Tian, Y., Varadi, F. and Yiou, P.: Advanced spectral methods for climatic time series, *Rev. Geophys.*, 40(1), 3–13–41, doi:10.1029/2000RG000092, 2002.
- Goldreich, Y., Mozes, H. and Rosenfeld, D.: Radar analysis of cloud systems and their rainfall yield in Israel, *Isr. J. Earth Sci.*, 53(2), 63–76, doi:10.1560/G68K-30MN-D5V0-KUHU, 2004.
- Greenbaum, N., Schwartz, U. and Bergman, N.: Extreme floods and short term hydroclimatological fluctuations in the hyper-arid Dead Sea region, Israel, *Glob. Planet. Change*, 70(1–4), 125–137, doi:10.1016/j.gloplacha.2009.11.013, 2010.
- Grinsted, A., Moore, J. C. and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, *Nonlinear Process. Geophys.*, 11(5/6), 561–566, doi:10.5194/npg-11-561-2004, 2004.
- Groth, A. and Ghil, M.: Monte Carlo singular spectrum analysis (SSA) revisited: Detecting oscillator clusters in multivariate datasets, *J. Clim.*, 28(19), 7873–7893, doi:10.1175/JCLI-D-15-0100.1, 2015.
- Haase-Schramm, A., Goldstein, S. L. and Stein, M.: U-Th dating of Lake Lisan (late Pleistocene dead sea) aragonite and implications for glacial east Mediterranean climate change, *Geochim. Cosmochim. Acta*, 68(5), 985–1005, doi:10.1016/j.gca.2003.07.016, 2004.
- Heim, C., Nowaczyk, N. R., Negendank, J. F. W., Leroy, S. A. G. and Ben Avraham, Z.: Near East desertification: Evidence from the Dead Sea, *Naturwissenschaften*, 84(9), 398–401, doi:10.1007/s001140050416, 1997.
- Held, I. M. and Soden, B. J.: Robust responses of the hydrological cycle to global warming, *J. Clim.*, 19(21), 5686–5699, doi:10.1175/JCLI3990.1, 2006.
- Kagan, E., Stein, M. and Marco, S.: Integrated Paleoseismic Chronology of the Last Glacial Lake Lisan: From Lake Margin Seismites to Deep-Lake Mass Transport Deposits, *J. Geophys. Res. Solid Earth*, 123(4), 2806–2824, doi:10.1002/2017JB014117, 2018.
- Kahana, R., Ziv, B., Enzel, Y. and Dayan, U.: Synoptic climatology of major floods in the Negev Desert, Israel, *Int. J. Climatol.*, 22(7), 867–882, doi:10.1002/joc.766, 2002.

- 1105 Kelley, C., Ting, M., Seager, R. and Kushnir, Y.: Mediterranean precipitation climatology, seasonal cycle, and trend as simulated by CMIP5, *Geophys. Res. Lett.*, 39(21), 1–7, doi:10.1029/2012GL053416, 2012.
- Kennel, M. B., Brown, R. and Abarbanel, H. D. I.: Determining embedding dimension for phase space reconstruction using a geometrical construction, *Phys. Rev. A*, 45(6), 3403–3411, doi:10.1103/PhysRevA.45.3403, 1992.
- Kiro, Y., Goldstein, S. L., Lazar, B. and Stein, M.: Environmental implications of salt facies in the Dead Sea, *Geol. Soc. Am. Bull.*, 128(5–6), 824–841, doi:10.1130/b31357.1, 2016.
- 1110 Kolodny, Y., Stein, M. and Machlus, M.: Sea-rain-lake relation in the Last Glacial East Mediterranean revealed by  $\delta^{18}\text{O}$ - $\delta^{13}\text{C}$  in Lake Lisan aragonites, *Geochim. Cosmochim. Acta*, 69(16), 4045–4060, doi:10.1016/j.gca.2004.11.022, 2005.
- Krichak, S. O., Kishcha, P. and Alpert, P.: Decadal trends of main Eurasian oscillations and the eastern Mediterranean precipitation, *Theor. Appl. Climatol.*, 72(3–4), 209–220, doi:10.1007/s007040200021, 2002.
- 1115 Kushnir, Y., Dayan, U., Ziv, B., Morin, E. and Enzel, Y.: Climate of the Levant, *Quat. Levant*, 31–44, doi:10.1017/9781316106754.004, 2017.
- Lau, K. M. and Hengyi-Weng: Climate signal detection using wavelet transform: how to make a time series sing, *Bull.—Am. Meteorol. Soc.*, 76(12), 2391–2402, doi:10.1175/1520-0477(1995)076<2391:CSDUWT>2.0.CO;2, 1995.
- Luck, M., Landis, M. and Gassert, F.: Aqueduct Water Stress Projections: Decadal projections of water supply and demand using CMIP5 GCMs., *World Resour. Inst. Technical Note*, (April), 1–20 [online] Available from: [wri.org/publication/aqueduct-water-stress-projections](http://wri.org/publication/aqueduct-water-stress-projections), 2015.
- 1120 Luo, T., Young, R. and Reig, P.: Aqueduct Projected Water Stress Country Rankings. Technical Note, Washington, D.C.: World Resources Institute, (August), 1–16, 2015.
- Machlus, M., Enzel, Y., Goldstein, S. L., Mareo, S. and Stein, M.: Reconstructing low levels of Lake Lisan by correlating fan-delta and lacustrine deposits, *Quat. Int.*, 73–74, 137–144, doi:10.1016/S1040-6182(00)00070-7, 2000.
- 1125 Mann, H. B. and Whitney, D. R.: On a Test of Whether one of Two Random Variables is Stochastically Larger than the Other, *Ann. Math. Stat.*, 18(1), 50–60 [online] Available from: <https://www.jstor.org/stable/pdf/2236101.pdf>, 1947.
- Marwan, N.: How to avoid potential pitfalls in recurrence plot-based data analysis, *Int. J. Bifurc. Chaos*, 21(4), 1003–1017, doi:10.1142/S0218127411029008, 2011.
- 1130 Marwan, N. and Kurths, J.: Cross-Recurrence Plots and Their Applications, *Math. Phys. Res. Cut. Edge*, 101–139, doi:10.1093/brain/aws023, 2004.
- Marwan, N., Trauth, M. H., Vuille, M. and Kurths, J.: Comparing modern and Pleistocene ENSO-like influences in NW Argentina using nonlinear time series analysis methods, *Clim. Dyn.*, 21(3–4), 317–326, doi:10.1007/s00382-003-0335-3, 2003.
- 1135 Marwan, N., Carmen Romano, M., Thiel, M. and Kurths, J.: Recurrence plots for the analysis of complex systems, *Phys. Rep.*, 438(5–6), 237–329, doi:10.1016/j.physrep.2006.11.001, 2007.
- Metzger, A., Marra, F., Smith, J. A. and Morin, E.: Flood frequency estimation and uncertainty in arid/semi-arid regions, *J. Hydrol.*, 590, 125254, doi:10.1016/j.jhydrol.2020.125254, 2020.

Migowski, C., Stein, M., Prasad, S., Negendank, J. F. W. and Agnon, A.: Holocene climate variability and cultural evolution in the Near East from the Dead Sea sedimentary record, *Quat. Res.*, 66(3), 421–431, doi:10.1016/j.yqres.2006.06.010, 2006.

Morin, E.: To know what we cannot know: Global mapping of minimal detectable absolute trends in annual precipitation, *Water Resour. Res.*, 47(7), doi:10.1029/2010WR009798, 2011.

Morin, E., Ryb, T., Gavrieli, I. and Enzel, Y.: Mean, variance, and trends of Levant precipitation over the past 4500 years from reconstructed Dead Sea levels and stochastic modeling, *Quat. Res. (United States)*, 91(2), 751–767, doi:10.1017/qua.2018.98, 2019.

Le Mouél, J.-L., Lopes, F. and Courtillot, V.: A Solar Signature in Many Climate Indices, *J. Geophys. Res. Atmos.*, 124(5), 2600–2619, doi:10.1029/2018JD028929, 2019.

Neev, D.: Recent precipitation of calcium salts in the Dead Sea, in *Israel Research Council Bulletin*, pp. 153–154., 1963.

Neev, D. and Emery, K. O.: The Dead Sea: depositional processes and environments of evaporites, *Isr. Geol. Surv. Bull.*, 41, 1–47, 1967.

Nehorai, R., Lensky, I. M., Hochman, L., Gertman, I., Brenner, S., Muskin, A. and Lensky, N. G.: Satellite observations of turbidity in the Dead Sea, *J. Geophys. Res. Ocean.*, 118(6), 3146–3160, doi:10.1002/jgre.20204, 2013.

Neugebauer, I., Brauer, A., Schwab, M. J., Waldmann, N. D., Enzel, Y., Kitagawa, H., Torfstein, A., Frank, U., Dulski, P., Agnon, A., Ariztegui, D., Ben-Avraham, Z., Goldstein, S. L. and Stein, M.: Lithology of the long sediment record recovered by the ICDP Dead Sea Deep Drilling Project (DSDDP), *Quat. Sci. Rev.*, 102, 149–165, doi:10.1016/j.quascirev.2014.08.013, 2014.

Neugebauer, I., Brauer, A., Schwab, M. J., Dulski, P., Frank, U., Hadzhiivanova, E., Kitagawa, H., Litt, T., Schiebel, V., Taha, N. and Waldmann, N. D.: Evidences for centennial dry periods at ~3300 and ~2800 cal. yr BP from micro-facies analyses of the Dead Sea sediments, Holocene, 25(8), 1358–1371, doi:10.1177/0959683615584208, 2015.

Nissenbaum, A., Baedeker, M. J. and Kaplan, I. R.: Organic geochemistry of Dead Sea sediments, *Geochim. Cosmochim. Acta*, 36(7), 709–727, doi:10.1016/0016-7037(72)90082-8, 1972.

NOAA: Climate Prediction Center, [online] Available from: <https://www.cpc.ncep.noaa.gov/data/teledoc/ea.shtml> (Accessed 29 July 2020), 2020.

Palehan, D., Neugebauer, I., Amitai, Y., Waldmann, N. D., Schwab, M. J., Dulski, P., Brauer, A., Stein, M., Erel, Y. and Enzel, Y.: North Atlantic controlled depositional cycles in MIS 5e layered sediments from the deep Dead Sea basin, *Quat. Res. (United States)*, 87(1), 168–179, doi:10.1017/qua.2016.10, 2017.

Peleg, N., Bartov, M. and Morin, E.: CMIP5-predicted climate shifts over the East Mediterranean: Implications for the transition region between Mediterranean and semi-arid climates, *Int. J. Climatol.*, 35(8), 2144–2153, doi:10.1002/joc.4114, 2015.

Prasad, S., Vos, H., Negendank, J. F. W., Waldmann, N., Goldstein, S. L. and Stein, M.: Evidence from Lake Lisan of solar influence on decadal to centennial-scale climate variability during marine oxygen isotope stage 2, *Geology*, 32(7), 581–584, doi:10.1130/G20553-1, 2004.

Redmond, K. T., Enzel, Y., House, P. K. and Biondi, F.: Climate variability and flood frequency at decadal to millennial time scales, *Anc. floods, Mod. hazards Princ. Appl. paleoflood Hydrol.*, 5, 21–45, 2002.

1175 Rohling, E. J.: Quantitative assessment of glacial fluctuations in the level of Lake Lisan, Dead-Sea rift, *Quat. Sci. Rev.*, 70, 63–72, doi:<https://doi.org/10.1016/j.quascirev.2013.03.013>, 2013.

Saaroni, H., Halfon, N., Ziv, B., Alpert, P. and Kutiel, H.: Links between the rainfall regime in Israel and location and intensity of Cyprus lows, *Int. J. Climatol.*, 30(7), 1014–1025, doi:[10.1002/joc.1912](https://doi.org/10.1002/joc.1912), 2010.

1180 Safriel, U., Adeel, Z., Niemeijer, D., Puigdefabregas, J., White, R., Lal, R., Winslow, M., Ziedler, J., Prince, S., Atcher, E., King, C., Shapiro, B., Wessels, K., Nielsen, T. T., Portnov, B., Reshef, I., Thornell, J., Lachman, E. and McNab, D.: Dryland systems, , 623–662 [online] Available from: <https://forskning.ruc.dk/en/publications/dryland-systems> (Accessed 27 August 2020), 2005.

Seager, R., Osborn, T. J., Kushnir, Y., Simpson, I. R., Nakamura, J. and Liu, H.: Climate variability and change of mediterranean-type climates, *J. Clim.*, 32(10), 2887–2915, doi:[10.1175/JCLI-D-18-0472.1](https://doi.org/10.1175/JCLI-D-18-0472.1), 2019.

1185 Seager, R., Liu, H., Kushnir, Y., Osborn, T. J., Simpson, I. R., Kelley, C. R. and Nakamura, J.: Mechanisms of Winter Precipitation Variability in the European Mediterranean Region Associated with the North Atlantic Oscillation, *J. Clim.*, 33(16), 7179–7196, doi:[10.1175/JCLI-D-20-0011.1](https://doi.org/10.1175/JCLI-D-20-0011.1), 2020.

Serinaldi, F., Kilsby, C. G. and Lombardo, F.: Untenable nonstationarity: An assessment of the fitness for purpose of trend tests in hydrology, *Adv. Water Resour.*, 111, 132–155, doi:[10.1016/j.advwatres.2017.10.015](https://doi.org/10.1016/j.advwatres.2017.10.015), 2018.

1190 Shentsis, I., Laronne, J. B. and Alpert, P.: Red Sea Trough flood events in the Negev, Israel (1964–2007), *Hydrol. Sci. J.*, 57(1), 42–51, doi:[10.1080/02626667.2011.636922](https://doi.org/10.1080/02626667.2011.636922), 2012.

Shohami, D., Dayan, U. and Morin, E.: Warming and drying of the eastern Mediterranean: Additional evidence from trend analysis, *J. Geophys. Res. Atmos.*, 116(22), 1–12, doi:[10.1029/2011JD016004](https://doi.org/10.1029/2011JD016004), 2011.

1195 Sirota, I., Enzel, Y. and Lensky, N. G.: Temperature seasonality control on modern halite layers in the Dead Sea: In situ observations, *Bull. Geol. Soc. Am.*, 129(9–10), 1181–1194, doi:[10.1130/B31661.1](https://doi.org/10.1130/B31661.1), 2017.

Stein, M., Starinsky, A., Katz, A., Goldstein, S. L., Machlus, M. and Schramm, A.: Strontium isotopic, chemical, and sedimentological evidence for the evolution of Lake Lisan and the Dead Sea, *Geochim. Cosmochim. Acta*, 61(18), 3975–3992, doi:[10.1016/S0016-7037\(97\)00191-9](https://doi.org/10.1016/S0016-7037(97)00191-9), 1997.

1200 Stein, M., Ben-Avraham, Z. and Goldstein, S. L.: Dead-Sea Deep Cores: A Window A Library of Georeferenced Into Past Climate and Seismicity, *Eos (Washington, DC)*, 92(49), 453–454, doi:[10.1111/j.2006.0906-7590.04596.x](https://doi.org/10.1111/j.2006.0906-7590.04596.x)Xiao, 2011.

Stöckli, R., Vermote, E., Saleous, N., Simmon, R. and Herring, D.: The Blue Marble Next Generation—A true color earth dataset including seasonal dynamics from MODIS, Publ. by NASA Earth Obs., 2005.

1205 Swierczynski, T., Brauer, A., Lauterbach, S., Martín-Puertas, C., Dulski, P., von Grafenstein, U. and Rohr, C.: A 1600-yr seasonally resolved record of decadal-scale flood variability from the Austrian Pre-Alps, *Geology*, 40(11), 1047–1050, doi:[10.1130/G33493.1](https://doi.org/10.1130/G33493.1), 2012.

Tamarin-Brodsky, T. and Kaspi, Y.: Enhanced poleward propagation of storms under climate change, *Nat. Geosci.*, 10(12),

908–913, doi:10.1038/s41561-017-0001-8, 2017.

Tarolli, P., Borga, M., Morin, E. and Delrieu, G.: Analysis of flash flood regimes in the North-Western and South-Eastern Mediterranean regions, *Nat. Hazards Earth Syst. Sci.*, 12(5), 1255–1265, doi:10.5194/nhess-12-1255-2012, 2012.

1210 Torfstein, A., Gavrieli, I., Katz, A., Kolodny, Y. and Stein, M.: Gypsum as a monitor of the paleo-limnological-hydrological conditions in Lake Lisan and the Dead Sea, *Geochim. Cosmochim. Acta*, 72(10), 2491–2509, doi:10.1016/j.gca.2008.02.015, 2008.

Torfstein, A.: Brine-freshwater interplay and effects on the evolution of saline lakes: The Dead Sea Rift terminal lakes by Brine-freshwater interplay and effects on the evolution of saline lakes: The Dead Sea Rift terminal lakes, (August), 2008.

1215 Torfstein, A. and Enzel, Y.: Dead Sea Lake Level Changes and Levant Palaeoclimate, in *Quaternary of the Levant: Environments, Climate Change, and Humans*, pp. 115–126., 2017.

Torfstein, A., Haase-Schramm, A., Waldmann, N., Kolodny, Y. and Stein, M.: U-series and oxygen isotope chronology of the mid-Pleistocene Lake Amora (Dead Sea basin), *Geochim. Cosmochim. Acta*, 73(9), 2603–2630, doi:10.1016/j.gca.2009.02.010, 2009.

1220 Torfstein, A., Goldstein, S. L., Kagan, E. J. and Stein, M.: Integrated multi-site U-Th chronology of the last glacial Lake Lisan, *Geochim. Cosmochim. Acta*, 104, 210–231, doi:10.1016/j.gca.2012.11.003, 2013.

Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bull. Am. Meteorol. Soc.*, 79(1), 61–78, doi:10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2, 1998.

Tsvieli, Y. and Zangvil, A.: Synoptic climatological analysis of “wet” and “dry” Red Sea Troughs over Israel, *Int. J. Climatol.*, 25(15), 1997–2015, doi:10.1002/joc.1232, 2005.

1225 Tubi, A. and Dayan, U.: Tropical Plumes over the Middle East: Climatology and synoptic conditions, *Atmos. Res.*, 145–146, 168–181, doi:10.1016/j.atmosres.2014.03.028, 2014.

Vautard, R. and Ghil, M.: Singular spectrum analysis in nonlinear dynamics, with applications to paleoclimatic time series, *Phys. D Nonlinear Phenom.*, 35(3), 395–424, doi:10.1016/0167-2789(89)90077-8, 1989.

1230 Waldmann, N.: The Stratigraphy and Chronology of the Samra Formation, in *Quaternary of the Levant: Environments, Climate Change, and Humans*, edited by O. Bar-Yosef and Y. Enzel, pp. 99–106, Cambridge University Press, Cambridge., 2017.

Waldmann, N., Neugebauer, I., Palchan, D., Hadzhiivanova, E., Taha, N., Brauer, A. and Enzel, Y.: Sedimentology of the Lacustrine Formations in the Dead Sea Basin, *Quat. Levant*, 83–90, doi:10.1017/9781316106754.009, 2017.

Welch, P. D.: Welch Periodogram.pdf, *IEEE Trans. Audio Electroacoust.*, 15, 70–73, 1967.

1235 Witt, A., Malamud, B. D., Mangili, C. and Brauer, A.: Analysis and Modelling of a 9.3 kyr Palaeoflood Record: Correlations, Clustering and Cycles, *Hydrol. Earth Syst. Sci. Discuss.*, (October), 1–60, doi:10.5194/hess-2016-470, 2016.

Yosef, Y., Aguilar, E. and Alpert, P.: Changes in extreme temperature and precipitation indices: Using an innovative daily homogenized database in Israel, *Int. J. Climatol.*, 39(13), 5022–5045, doi:10.1002/joc.6125, 2019.

1240 Zappa, G., Hawcroft, M. K., Shaffrey, L., Black, E. and Brayshaw, D. J.: Extratropical cyclones and the projected decline of winter Mediterranean precipitation in the CMIP5 models, *Clim. Dyn.*, 45(7–8), 1727–1738, doi:10.1007/s00382-014-2426-8,

2015.

Ziv, B.: A subtropical rainstorm associated with a tropical plume over Africa and the Middle-East, *Theor. Appl. Climatol.*, 69(1-2), 91-102, doi:10.1007/s007040170037, 2001.

1245

Ziv, B., Saaroni, H. and Alpert, P.: The factors governing the summer regime of the eastern Mediterranean, *Int. J. Climatol.*, 24(14), 1859-1871, doi:10.1002/joc.1113, 2004.

Ziv, B., Dayan, U., Kushnir, Y., Roth, C. and Enzel, Y.: Regional and global atmospheric patterns governing rainfall in the southern Levant, *Int. J. Climatol.*, 26(1), 55-73, doi:10.1002/joc.1238, 2006.

Zolitschka, B., Francus, P., Ojala, A. E. K. and Schimmelmann, A.: Varves in lake sediments—a review, *Quat. Sci. Rev.*, 117, 1-41, doi:10.1016/j.quaseirev.2015.03.019, 2015.

1250

[Ahlborn, M., Armon, M., Ben Dor, Y., Neugebauer, I., Schwab, M. J., Tjallingii, R., Shoqir, J. H., Morin, E., Enzel, Y., and Brauer, A.: Increased frequency of torrential rainstorms during a regional late Holocene eastern Mediterranean drought, \*Quaternary Research\*, 89, 425-431, 10.1017/qua.2018.9, 2018.](#)

1255

[Allen, K., Hope, P., Lam, D., Brown, J., and Wasson, R.: Improving Australia's flood record for planning purposes—can we do better?, \*Australasian Journal of Water Resources\*, 24, 36-45, 2020.](#)

1260

[Alpert, P., Ben-Gai, T., Baharad, A., Benjamini, Y., Yekutieli, D., Colacino, M., Diodato, L., Ramis, C., Homar, V., Romero, R., Michaelides, S., and Manes, A.: The paradoxical increase of Mediterranean extreme daily rainfall in spite of decrease in total values, \*Geophysical Research Letters\*, 29, 31-31-31-34, 10.1029/2001GL013554, 2002.](#)

1265

[Amit, R., Simhai, O., Ayalon, A., Enzel, Y., Matmon, A., Crouvi, O., Porat, N., and McDonald, E.: Transition from arid to hyper-arid environment in the southern Levant deserts as recorded by early Pleistocene cummulic Aridisols, \*Quaternary Science Reviews\*, 30, 312-323, 2011.](#)

1270

[Ansari, A. R. and Bradley, R. A.: Rank-sum tests for dispersions, \*The Annals of Mathematical Statistics\*, 31, 1174-1189, 1960.](#)

[Armon, M., morin, E., and Enzel, Y.: Overview of modern atmospheric patterns controlling rainfall and floods into the Dead Sea: Implications for the lake's sedimentology and paleohydrology, \*Quaternary Science Reviews\*, 216, 58-73, 2019.](#)

1275

[Armon, M., Dente, E., Smith, J. A., Enzel, Y., and Morin, E.: Synoptic-Scale Control over Modern Rainfall and Flood Patterns in the Levant Drylands with Implications for Past Climates, \*Journal of Hydrometeorology\*, 19, 1077-1096, 10.1175/jhm-d-18-0013.1, 2018.](#)

1280

[Armon, M., Marra, F., Enzel, Y., Rostkier-Edelstein, D., and Morin, E.: Radar-based characterisation of heavy precipitation in the eastern Mediterranean and its representation in a convection-permitting model, \*Hydrology and Earth System Sciences\*, 24, 1227-1249, 2020.](#)

[Avni, S., Joseph-Hai, N., Haviv, I., Matmon, A., Benedetti, L., and Team, A.: Patterns and rates of 103–105 yr denudation in carbonate terrains under subhumid to subalpine climatic gradient, Mount Hermon, Israel, \*Bulletin\*, 131, 899-912, 2018.](#)

1285

[Baker, V. R.: Paleoflood hydrology: Origin, progress, prospects, \*Geomorphology\*, 101, 1-13, 2008.](#)



\_\_\_\_\_  
[Bar-Matthews, M., Ayalon, A., Kaufman, A., and Wasserburg, G. J.: The Eastern Mediterranean paleoclimate as a reflection of regional events: Soreq cave, Israel, Earth Planet. Sci. Lett., 166, 85-95, 1999.](#)

1290 \_\_\_\_\_  
[Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A., and Hawkesworth, C. J.: Sea-land oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern Mediterranean region and their implication for paleorainfall during interglacial intervals, Geochim. Cosmochim. Acta, 67, 3181-3199, 2003.](#)

1295 \_\_\_\_\_  
[Bartov, Y., Enzel, Y., Porat, N., and Stein, M.: Evolution of the Late Pleistocene–Holocene Dead Sea Basin from sequence stratigraphy of fan deltas and lake-level reconstruction, Journal of Sedimentary Research, 77, 680-692, 2007.](#)

\_\_\_\_\_  
[Bartov, Y., Goldstein, S. L., Stein, M., and Enzel, Y.: Catastrophic arid episodes in the Eastern Mediterranean linked with the North Atlantic Heinrich events, Geology, 31, 439-442, 2003.](#)

1300 \_\_\_\_\_  
[Bartov, Y., Stein, M., Enzel, Y., Agnon, A., and Reches, Z.: Lake levels and sequence stratigraphy of Lake Lisan, the Late Pleistocene precursor of the Dead Sea, Quaternary Research, 57, 9-21, 2002.](#)

\_\_\_\_\_  
[Begin, Z., Ehrlich, A., and Nathan, Y.: Lake Lisan: the Pleistocene precursor of the Dead Sea, Ministry of Commerce and Industry, Geological Survey of Israel, Jerusalem61, 45, 1974.](#)

1305 \_\_\_\_\_  
[Begin, Z. B., Nathan, Y., and Ehrlich, A.: Stratigraphy and facies distribution in the Lisan Formation—new evidence from the area south of the Dead Sea, Israel, Israel Journal of Earth Sciences, 29, 182-189, 1980.](#)

1310 \_\_\_\_\_  
[Begin, Z. B., Stein, M., Katz, A., Machlus, M., Rosenfeld, A., Buchbinder, B., and Bartov, Y.: Southward migration of rain tracks during the last glacial, revealed by salinity gradient in Lake Lisan \(Dead Sea rift\), Quaternary Science Reviews, 23, 1627-1636, 10.1016/j.quascirev.2004.01.002, 2004.](#)

\_\_\_\_\_  
[Belachsen, I., Marra, F., Peleg, N., and Morin, E.: Convective rainfall in dry climate: relations with synoptic systems and flash-flood generation in the Dead Sea region, Hydrol. Earth Syst. Sci. Discuss, 2017.](#)

1315 \_\_\_\_\_  
[Belmaker, R., Taha, N., and Bookman, R.: Evidence for aragonite deposition in flood plumes of the Dead Sea, Goldschmidt Abstract 196.](#)

1320 \_\_\_\_\_  
[Belmaker, R., Lazar, B., Stein, M., Taha, N., and Bookman, R.: Constraints on aragonite precipitation in the Dead Sea from geochemical measurements of flood plumes, Quaternary Science Reviews, 221, 105876, 2019.](#)

\_\_\_\_\_  
[Ben David-Novak, H., Morin, E., and Enzel, Y.: Modern extreme storms and the rainfall thresholds for initiating debris flows on the hyperarid western escarpment of the Dead Sea, Israel, Geological Society of America Bulletin, 116, 718-728, 2004.](#)

1325 \_\_\_\_\_  
[Ben Dor, Y., Flax, T., Levitan, I., Enzel, Y., Brauer, A., and Erel, Y.: The paleohydrological implications of aragonite precipitation under contrasting climates in the endorheic Dead Sea and its precursors revealed by experimental investigations, Chemical Geology, 576, 10.1016/j.chemgeo.2021.120261, 2021.](#)

1330 \_\_\_\_\_  
[Ben Dor, Y., Neugebauer, I., Enzel, Y., Schwab, M. J., Tjallingii, R., Erel, Y., and Brauer, A.: Varves of the Dead Sea sedimentary record, Quaternary Science Reviews, 215, 173-184, 2019.](#)

1335 \_\_\_\_\_  
[Ben Dor, Y., Neugebauer, I., Enzel, Y., Schwab, M. J., Tjallingii, R., Erel, Y., and Brauer, A.: Reply to comment on Ben Dor Y. et al. "Varves of the Dead Sea sedimentary record." Quaternary Science Reviews 215 \(2019\): 173–184, Quaternary Science Reviews, 231, 10.1016/j.quascirev.2019.106063, 2020.](#)

1340 [Ben Dor, Y., Armon, M., Ahlborn, M., Morin, E., Erel, Y., Brauer, A., Schwab, M. J., Tjallingii, R., and Enzel, Y.: Changing flood frequencies under opposing late Pleistocene eastern Mediterranean climates, \*Sci Rep.\* 8, 8445, 10.1038/s41598-018-25969-6, 2018.](#)

[Black, E.: The influence of the North Atlantic Oscillation and European circulation regimes on the daily to interannual variability of winter precipitation in Israel, \*International Journal of Climatology\*, 32, 1654-1664, 2012.](#)

1345 [Blackman, R. B. and Tukey, J. W.: The measurement of power spectra, 1958.](#)

[Bookman, R., Bartov, Y., Enzel, Y., and Stein, M.: Quaternary lake levels in the Dead Sea basin: two centuries of research, \*Geological Society of America Special Papers\*, 401, 155-170, 2006.](#)

1350 [Brauer, A., Endres, C., and Negendank, J. F.: Lateglacial calendar year chronology based on annually laminated sediments from Lake Meerfelder Maar, Germany, \*Quaternary International\*, 61, 17-25, 1999.](#)

1355 [Brauer, A., Mangili, C., Moscarriello, A., and Witt, A.: Palaeoclimatic implications from micro-facies data of a 5900 yearve time series from the Piànico interglacial sediment record, southern Alps, \*Palaeogeography, Palaeoclimatology, Palaeoecology\*, 259, 121-135, 2008.](#)

[Broomhead, D. S. and King, G. P.: Nonlinear phenomena and chaos, \*On the Qualitative Analysis of Experimental Dynamical Systems\*, Malvern science series, Bristol, UK, 113-144, 1986.](#)

1360 [Campins, J., Genovés, A., Picornell, M., and Jansà, A.: Climatology of Mediterranean cyclones using the ERA-40 dataset, \*International Journal of Climatology\*, 31, 1596-1614, 2011.](#)

[Coianiz, L., Ben-Avraham, Z., Stein, M., and Lazar, M.: Spatial and temporal reconstruction of the late Quaternary Dead Sea sedimentary facies from geophysical properties, \*Journal of Applied Geophysics\*, 160, 15-27, 2019.](#)

1365 [Crouvi, O., Amit, R., Ben Israel, M., and Enzel, Y.: Loess in the Negev desert: sources, loessial soils, palaeosols, and palaeoclimatic implications, in: \*Quaternary of the Levant: Environments, Climate Change, and Humans\*, Cambridge University Press, Cambridge, edited by: Enzel, Y., and Bar-Yosef, O., 471-482, 2017.](#)

1370 [Dayan, U. and Morin, E.: Flash flood-producing rainstorms over the Dead Sea: A review, \*Geological Society of America Special Papers\*, 401, 53-62, 2006.](#)

1375 [Dean, J. R., Eastwood, W. J., Roberts, N., Jones, M. D., Yiğitbaşoğlu, H., Allcock, S. L., Woodbridge, J., Metcalfe, S. E., and Leng, M. J.: Tracking the hydro-climatic signal from lake to sediment: A field study from central Turkey, \*Journal of Hydrology\*, 529, 608-621, 2015.](#)

[Donges, J. F., Donner, R. V., Trauth, M. H., Marwan, N., Schellnhuber, H.-J., and Kurths, J.: Nonlinear detection of paleoclimate-variability transitions possibly related to human evolution, \*Proceedings of the National Academy of Sciences\*, 108, 20422-20427, 2011.](#)

1380 [Drori, R., Ziv, B., Saaroni, H., Etkin, A., and Sheffer, E.: Recent changes in the rain regime over the Mediterranean climate region of Israel, \*Climatic Change\*, 167, 15, 10.1007/s10584-021-03161-6, 2021.](#)

1385 [Enzel, Y., Amit, R., Dayan, U., Crouvi, O., Kahana, R., Ziv, B., and Sharon, D.: The climatic and physiographic controls of the eastern Mediterranean over the late Pleistocene climates in the southern Levant and its neighboring deserts, \*Global and Planetary Change\*, 60, 165-192, 2008.](#)

1390 \_\_\_\_\_ Enzel, Y., Bookman, R., Sharon, D., Gvirtzman, H., Dayan, U., Ziv, B., and Stein, M.: Late Holocene climates of the Near East deduced from Dead Sea level variations and modern regional winter rainfall, *Quaternary Research*, 60, 263-273, 2003.

\_\_\_\_\_ Enzel, Y., Amit, R., Grodek, T., Ayalon, A., Lekach, J., Porat, N., Bierman, P., Blum, J. D., and Erel, Y.: Late Quaternary weathering, erosion, and deposition in Nahal Yael, Israel: An “impact of climatic change on an arid watershed”?, *Bulletin*, 124, 705-722, 2012.

1395 \_\_\_\_\_ Feldstein, S. B. and Dayan, U.: Circumglobal teleconnections and wave packets associated with Israeli winter precipitation, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 134, 455-467, 2008.

1400 \_\_\_\_\_ Feliks, Y., Ghil, M., and Robertson, A. W.: Oscillatory climate modes in the Eastern Mediterranean and their synchronization with the North Atlantic Oscillation, *Journal of Climate*, 23, 4060-4079, 2010.

\_\_\_\_\_ Flocas, H. A., Simmonds, I., Kouroutzoglou, J., Keay, K., Hatzaki, M., Bricolas, V., and Asimakopoulos, D.: On cyclonic tracks over the eastern Mediterranean, *Journal of Climate*, 23, 5243-5257, 2010.

1405 \_\_\_\_\_ Ganor, E. and Foner, H.: The mineralogical and chemical properties and the behaviour of aeolian Saharan dust over Israel, in: *The impact of desert dust across the Mediterranean*, Springer, 163-172, 1996.

\_\_\_\_\_ Garber, R. A., Levy, Y., and Friedman, G. M.: The sedimentology of the Dead Sea, *Carbonates and Evaporites*, 2, 43-57, 1987.

1410 \_\_\_\_\_ Garfunkel, Z.: Internal structure of the Dead Sea leaky transform (rift) in relation to plate kinematics, *Tectonophysics*, 80, 81-108, 1981.

\_\_\_\_\_ Garfunkel, Z. and Ben-Avraham, Z.: The structure of the Dead Sea basin, *Tectonophysics*, 266, 155-176, 1996.

1415 \_\_\_\_\_ Ghil, M., Allen, M., Dettinger, M., Ide, K., Kondrashov, D., Mann, M., Robertson, A. W., Saunders, A., Tian, Y., and Varadi, F.: Advanced spectral methods for climatic time series, *Reviews of geophysics*, 40, 2002.

1420 \_\_\_\_\_ Golan, R., Lazar, B., Wurgaft, E., Lensky, N., Ganor, J., and Gavrieli, I.: Continuous CO<sub>2</sub> escape from the hypersaline Dead Sea caused by aragonite precipitation, *Geochim. Cosmochim. Acta*, 207, 43-56, 10.1016/j.gca.2017.02.031, 2017.

\_\_\_\_\_ Goldreich, Y.: Spatial Distribution of Mid-season Rainfall Date in Israel - a Review, *Horizons in Geography*, 177-182, 2004.

1425 \_\_\_\_\_ Goldreich, Y.: *The climate of Israel: observation, research and application*, Springer Science & Business Media 2012.

\_\_\_\_\_ Goldreich, Y., Mozes, H., and Rosenfeld, D.: Radar analysis of cloud systems and their rainfall yield in Israel, *Isr. J. Earth Sci.*, 53, 63-76, 2004.

1430 \_\_\_\_\_ Goldsmith, Y., Polissar, P., Ayalon, A., Bar-Matthews, M., and Broecker, W.: The modern and Last Glacial Maximum hydrological cycles of the Eastern Mediterranean and the Levant from a water isotope perspective, *Earth Planet. Sci. Lett.*, 457, 302-312, 2017.

1435 [Greenbaum, N., Schwartz, U., and Bergman, N.: Extreme floods and short-term hydroclimatological fluctuations in the hyper-arid Dead Sea region, Israel, \*Global and Planetary Change\*, 70, 125-137, 2010.](#)

[Greenbaum, N., Ben-Zvi, A., Haviv, I., and Enzel, Y.: The hydrology and paleohydrology of the Dead Sea tributaries, \*Geological Society of America Special Papers\*, 401, 63-93, 2006.](#)

1440 [Grinsted, A., Moore, J. C., and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, \*Nonlinear processes in geophysics\*, 11, 561-566, <https://doi.org/10.5194/npg-11-561-2004>, 2004.](#)

1445 [Haase-Schramm, A., Goldstein, S. L., and Stein, M.: U-Th dating of Lake Lisan \(late Pleistocene dead sea\) aragonite and implications for glacial east Mediterranean climate change, \*Geochim. Cosmochim. Acta\*, 68, 985-1005, <http://dx.doi.org/10.1016/j.gca.2003.07.016>, 2004.](#)

[Heim, C., Nowaczyk, N. R., Negendank, J. F., Leroy, S. A., and Ben-Avraham, Z.: Near East desertification: evidence from the Dead Sea, \*Naturwissenschaften\*, 84, 398-401, 1997.](#)

1450 [Held, I. M. and Soden, B. J.: Robust responses of the hydrological cycle to global warming, \*Journal of Climate\*, 19, 5686-5699, 2006.](#)

1455 [Hochman, A., Saaroni, H., Abramovich, F., and Alpert, P.: Artificial Detection of Lower-Frequency Periodicity in Climatic Studies by Wavelet Analysis Demonstrated on Synthetic Time Series, \*Journal of Applied Meteorology and Climatology\*, 58, 2077-2086, 2019.](#)

1460 [IPCC, Zhai, V. P., Pirani, A. S., Connors, L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, J. B. E., Matthews, R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B. \(Eds.\): \*Climate Change 2021: The Physical Science Basis\*, Cambridge University Press. In Press2021.](#)

1465 [Kagan, E., Stein, M., and Marco, S.: Integrated Paleoseismic Chronology of the Last Glacial Lake Lisan: From Lake Margin Seismites to Deep-Lake Mass Transport Deposits, \*Journal of Geophysical Research: Solid Earth\*, 10.1002/2017JB014117, 2018.](#)

[Kahana, R., Ziv, B., Enzel, Y., and Dayan, U.: Synoptic climatology of major floods in the Negev Desert, Israel, \*International Journal of Climatology\*, 22, 867-882, 2002.](#)

1470 [Kalderon-Asael, B., Erel, Y., Sandler, A., and Dayan, U.: Mineralogical and chemical characterization of suspended atmospheric particles over the east Mediterranean based on synoptic-scale circulation patterns, \*Atmospheric Environment\*, 43, 3963-3970, 2009.](#)

[Katz, A., Kolodny, Y., and Nissenbaum, A.: The geochemical evolution of the Pleistocene Lake Lisan-Dead Sea system, \*Geochim. Cosmochim. Acta\*, 41, 1609-1626, 10.1016/0016-7037\(77\)90172-7, 1977.](#)

1475 [Kaufman, A.: U-Series dating of Dead Sea Basin carbonates, \*Geochim. Cosmochim. Acta\*, 35, 1269-1281, \[http://dx.doi.org/10.1016/0016-7037\\(71\\)90115-3\]\(http://dx.doi.org/10.1016/0016-7037\(71\)90115-3\), 1971.](#)

1480 [Keinan, J., Bar-Matthews, M., Ayalon, A., Zilberman, T., Agnon, A., and Frumkin, A.: Paleoclimatology of the Levant from Zalmon Cave speleothems, the northern Jordan Valley, Israel, \*Quaternary Science Reviews\*, 220, 142-153, 2019.](#)

[Kelley, C., Ting, M., Seager, R., and Kushnir, Y.: Mediterranean precipitation climatology, seasonal cycle, and trend as simulated by CMIP5, \*Geophysical Research Letters\*, 39, 2012.](#)

1485 [Kennel, M. B., Brown, R., and Abarbanel, H. D.: Determining embedding dimension for phase-space reconstruction using a geometrical construction, \*Physical review A\*, 45, 3403, 1992.](#)

[Kitagawa, H., Stein, M., Goldstein, S. L., Nakamura, T., and Lazar, B.: Radiocarbon chronology of the DSDDP core at the deepest floor of the Dead Sea, \*Radiocarbon\*, 59, 383-394, 10.1017/RDC.2016.120, 2017.](#)

1490 [Kolodny, Y., Stein, M., and Machlus, M.: Sea-rain-lake relation in the Last Glacial East Mediterranean revealed by  \$\delta^{18}\text{O}\$ - \$\delta^{13}\text{C}\$  in Lake Lisan aragonites, \*Geochim. Cosmochim. Acta\*, 69, 4045-4060, 10.1016/j.gca.2004.11.022, 2005.](#)

1495 [Kondrashov, D. and Ghil, M.: Spatio-temporal filling of missing points in geophysical data sets, \*Nonlinear Processes in Geophysics\*, 13, 151-159, 2006.](#)

[Kottek, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F.: World map of the Köppen-Geiger climate classification updated, \*Meteorologische Zeitschrift\*, 15, 259-263, 2006.](#)

1500 [Krichak, S., Kishcha, P., and Alpert, P.: Decadal trends of main Eurasian oscillations and the Eastern Mediterranean precipitation, \*Theoretical and Applied Climatology\*, 72, 209-220, 2002.](#)

[Kushnir, Y., Dayan, U., Ziv, B., Morin, E., and Enzel, Y.: Climate of the Levant, in: \*Quaternary of the Levant\*, edited by: Enzel, Y., and Bar-Yosef, O., Cambridge University Press, Cambridge, 31-44, 10.1017/9781316106754.004, 2017.](#)

1505 [Lau, K. and Weng, H.: Climate signal detection using wavelet transform: How to make a time series sing, \*Bulletin of the American Meteorological Society\*, 76, 2391-2402, 1995.](#)

1510 [Le Mouél, J. L., Lopes, F., and Courtillot, V.: A Solar Signature in Many Climate Indices, \*Journal of Geophysical Research: Atmospheres\*, 2019.](#)

[Levy, Y., Burg, A., Yeichieli, Y., and Gvirtzman, H.: Displacement of springs and changes in groundwater flow regime due to the extreme drop in adjacent lake levels: The Dead Sea rift, \*Journal of Hydrology\*, 587, 10.1016/j.jhydrol.2020.124928, 2020.](#)

1515 [Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic  \$\delta^{18}\text{O}\$  records, \*Paleoceanography\*, 20, 2005.](#)

1520 [Luck, M., Landis, M., and Gassert, F.: Aqueduct water stress projections: decadal projections of water supply and demand using CMIP5 GCMs, Technical Note, Washington, DC: World Resources Institute. Available online at: <http://www.wri.org/publication/aqueduct-waterstress-projections>, 2015.](#)

[Luo, T., Young, R., and Reig, P.: Aqueduct Projected Water Stress Country Rankings, Technical Note, 2015.](#)

1525 [Machlus, M., Enzel, Y., Goldstein, S. L., Marco, S., and Stein, M.: Reconstructing low levels of Lake Lisan by correlating fan-delta and lacustrine deposits, \*Quaternary International\*, 73, 137-144, 2000.](#)

[Mann, H. B. and Whitney, D. R.: On a test of whether one of two random variables is stochastically larger than the other, \*The annals of mathematical statistics\*, 50-60, 1947.](#)

1530 [Marco, S., Stein, M., Agnon, A., and Ron, H.: Long-term earthquake clustering: A 50,000-year paleoseismic record in the Dead Sea Graben, \*Journal of Geophysical Research: Solid Earth\*, 101, 6179-6191, 1996.](#)

- 1535 [Marra, F., Armon, M., Adam, O., Zoccatelli, D., Gazal, O., Garfinkel, C. I., Rostkier-Edelstein, D., Dayan, U., Enzel, Y., and Morin, E.: Toward Narrowing Uncertainty in Future Projections of Local Extreme Precipitation, \*Geophysical Research Letters\*, 48, e2020GL091823, 2021.](#)
- 1540 [Marwan, N.: How to avoid potential pitfalls in recurrence plot based data analysis, \*International Journal of Bifurcation and Chaos\*, 21, 1003-1017, 2011.](#)
- [Marwan, N. and Kurths, J.: Cross recurrence plots and their applications, \*Mathematical physics research at the cutting edge\*, 101-139, 2004.](#)
- 1545 [Marwan, N., Romano, M. C., Thiel, M., and Kurths, J.: Recurrence plots for the analysis of complex systems, \*Physics reports\*, 438, 237-329, 2007.](#)
- [Marwan, N., Trauth, M. H., Vuille, M., and Kurths, J.: Comparing modern and Pleistocene ENSO-like influences in NW Argentina using nonlinear time series analysis methods, \*Climate Dynamics\*, 21, 317-326, 2003.](#)
- 1550 [Metzger, A., Marra, F., Smith, J. A., and Morin, E.: Flood frequency estimation and uncertainty in arid/semi-arid regions, \*Journal of Hydrology\*, 590, 125254, 2020.](#)
- [Migowski, C., Stein, M., Prasad, S., Negendank, J. F., and Agnon, A.: Holocene climate variability and cultural evolution in the Near East from the Dead Sea sedimentary record, \*Quaternary Research\*, 66, 421-431, 2006.](#)
- 1555 [Morin, E.: To know what we cannot know: Global mapping of minimal detectable absolute trends in annual precipitation, \*Water Resources Research\*, 47, 2011.](#)
- [Morin, E., Ryb, T., Gavrieli, I., and Enzel, Y.: Mean, variance, and trends of Levant precipitation over the past 4500 years from reconstructed Dead Sea levels and stochastic modeling, \*Quaternary Research\*, 91, 751-767, 2019.](#)
- 1560 [Neev, D.: Recent precipitation of calcium salts in the Dead Sea, \*Bull. Res. Council Israel, Sect. G\*, 11, 153-154, 1963.](#)
- [Neev, D. and Emery, K. O.: The Dead Sea: depositional processes and environments of evaporites, Ministry of Commerce and Industry, Geological Survey of Israel, Jerusalem, 1967.](#)
- 1565 [Nehorai, R., Lensky, I., Hochman, L., Gertman, I., Brenner, S., Muskin, A., and Lensky, N.: Satellite observations of turbidity in the Dead Sea, \*Journal of Geophysical Research: Oceans\*, 118, 3146-3160, 2013.](#)
- 1570 [Neugebauer, I., Brauer, A., Schwab, M. J., Dulski, P., Frank, U., Hadzhiivanova, E., Kitagawa, H., Litt, T., Schiebel, V., and Taha, N.: Evidences for centennial dry periods at~ 3300 and~ 2800 cal. yr BP from micro-facies analyses of the Dead Sea sediments, \*The Holocene\*, 0959683615584208, 2015.](#)
- 1575 [Neugebauer, I., Brauer, A., Schwab, M. J., Waldmann, N. D., Enzel, Y., Kitagawa, H., Torfstein, A., Frank, U., Dulski, P., and Agnon, A.: Lithology of the long sediment record recovered by the ICDP Dead Sea Deep Drilling Project \(DSDDP\), \*Quaternary Science Reviews\*, 102, 149-165, 2014.](#)
- 1580 [Nissenbaum, A., Baedeker, M. J., and Kaplan, I. R.: Organic geochemistry of Dead Sea sediments, \*Geochim. Cosmochim. Acta\*, 36, 709-727, 1972.](#)
- [NOAA Climate Prediction Center, teleconnection patterns, \[dataset\], 2020. <https://www.cpc.ncep.noaa.gov/data/teledoc/telecontents.shtml>](#)

1585 [Palchan, D., Neugebauer, I., Amitai, Y., Waldmann, N. D., Schwab, M. J., Dulski, P., Brauer, A., Stein, M., Erel, Y., and Enzel, Y.: North Atlantic controlled depositional cycles in MIS 5e layered sediments from the deep Dead Sea basin, \*Quaternary Research\*, 87, 168-179, 2017.](#)

1590 [Peleg, N., Bartov, M., and Morin, E.: CMIP5-predicted climate shifts over the East Mediterranean: implications for the transition region between Mediterranean and semi-arid climates, \*International journal of climatology\*, 35, 2144-2153, 2015.](#)

[Prasad, S., Negendank, J., and Stein, M.: Varve counting reveals high resolution radiocarbon reservoir age variations in palaeolake Lisan, \*Journal of Quaternary Science: Published for the Quaternary Research Association\*, 24, 690-696, 2009.](#)

1595 [Prasad, S., Vos, H., Negendank, J., Waldmann, N., Goldstein, S. L., and Stein, M.: Evidence from Lake Lisan of solar influence on decadal-to centennial-scale climate variability during marine oxygen isotope stage 2, \*Geology\*, 32, 581-584, 2004.](#)

[Redmond, K. T., Enzel, Y., House, P. K., and Biondi, F.: Climate variability and flood frequency at decadal to millennial time scales, \*Ancient floods, modern hazards: principles and applications of paleoflood hydrology\*, 5, 21-45, 2002.](#)

1600 [Roesser, P., Dräger, N., Brykała, D., Ott, F., Pinkerneil, S., Gierszewski, P., Lindemann, C., Plessen, B., Brademann, B., Kaszubski, M., Fojutowski, M., Schwab, M. J., Słowiński, M., Błaszczewicz, M., and Brauer, A.: Advances in understanding calcite varve formation: new insights from a dual lake monitoring approach in the southern Baltic lowlands, \*Boreas\*, 10.1111/bor.12506, 2021.](#)

1605 [Rohling, E. J.: Quantitative assessment of glacial fluctuations in the level of Lake Lisan, Dead Sea rift, \*Quaternary Science Reviews\*, 70, 63-72, 10.1016/j.quascirev.2013.03.013, 2013.](#)

1610 [Romano, M. C., Thiel, M., Kurths, J., and von Bloh, W.: Multivariate recurrence plots, \*Physics letters A\*, 330, 214-223, 2004.](#)

[Ryb, U., Matmon, A., Erel, Y., Haviv, I., Benedetti, L., and Hidy, A.: Styles and rates of long-term denudation in carbonate terrains under a Mediterranean to hyper-arid climatic gradient, \*Earth Planet. Sci. Lett.\*, 406, 142-152, 2014.](#)

1615 [Saaroni, H., Halfon, N., Ziv, B., Alpert, P., and Kutiel, H.: Links between the rainfall regime in Israel and location and intensity of Cyprus lows, \*International Journal of Climatology\*, 30, 1014-1025, 2010.](#)

[Safriel, U., Adeel, Z., Niemeijer, D., Puigdefabregas, J., White, R., Lal, R., Winslow, M., Ziedler, J., Prince, S., and Archer, E.: Dryland systems, in: \*Ecosystems and Human Well-being: Current State and Trends: Findings of the Condition and Trends Working Group\*, Island Press, 623-662, 2005.](#)

1620 [Schinkel, S., Dimigen, O., and Marwan, N.: Selection of recurrence threshold for signal detection, \*The european physical journal special topics\*, 164, 45-53, 2008.](#)

1625 [Schulte, J. A.: Cumulative areawise testing in wavelet analysis and its application to geophysical time series, \*Nonlinear Processes in Geophysics\*, 23, 45-57, 2016.](#)

[Schulte, J. A.: Statistical hypothesis testing in wavelet analysis: theoretical developments and applications to Indian rainfall, \*Nonlinear Processes in Geophysics\*, 26, 91-108, 2019.](#)

1630 [Schulte, J. A., Georgas, N., Saba, V., and Howell, P.: North Pacific influences on Long Island sound temperature variability, \*Journal of Climate\*, 31, 2745-2769, 2018.](#)

1635 \_\_\_\_\_ Seager, R., Osborn, T. J., Kushnir, Y., Simpson, I. R., Nakamura, J., and Liu, H.: Climate variability and change of Mediterranean-type climates, *Journal of Climate*, 32, 2887-2915, 2019.

\_\_\_\_\_ Seager, R., Liu, H., Kushnir, Y., Osborn, T. J., Simpson, I. R., Kelley, C. R., and Nakamura, J.: Mechanisms of Winter Precipitation Variability in the European–Mediterranean Region Associated with the North Atlantic Oscillation, *Journal of Climate*, 33, 7179-7196, 2020.

1640 \_\_\_\_\_ Serinaldi, F., Kilsby, C. G., and Lombardo, F.: Untenable nonstationarity: An assessment of the fitness for purpose of trend tests in hydrology, *Advances in Water Resources*, 111, 132-155, 2018.

\_\_\_\_\_ Israel Meteorological Service, Climatic data in Israel (1995-2009), [dataset], 2021. <https://ims.gov.il/he/ClimateAtlas>

1645 \_\_\_\_\_ Sharon, D.: The spottiness of rainfall in a desert area, *Journal of Hydrology*, 17, 161-175, 1972.

\_\_\_\_\_ Sharon, D. and Kutiel, H.: The distribution of rainfall intensity in Israel, its regional and seasonal variations and its climatological evaluation, *Journal of Climatology*, 6, 277-291, 1986.

1650 \_\_\_\_\_ Shentsis, I., Laronne, J. B., and Alpert, P.: Red Sea Trough flood events in the Negev, Israel (1964–2007), *Hydrological sciences journal*, 57, 42-51, 2012.

\_\_\_\_\_ Shohami, D., Dayan, U., and Morin, E.: Warming and drying of the eastern Mediterranean: Additional evidence from trend analysis, *Journal of Geophysical Research: Atmospheres*, 116, 2011.

1655 \_\_\_\_\_ Siebert, C., Rodiger, T., Mallast, U., Grabe, A., Guttman, J., Laronne, J. B., Storz-Peretz, Y., Greenman, A., Salameh, E., Al-Raggad, M., Vachtman, D., Zvi, A. B., Ionescu, D., Brenner, A., Merz, R., and Geyer, S.: Challenges to estimate surface- and groundwater flow in arid regions: the Dead Sea catchment, *Sci Total Environ*, 485-486, 828-841, 10.1016/j.scitotenv.2014.04.010, 2014.

1660 \_\_\_\_\_ Sirota, I., Enzel, Y., and Lensky, N. G.: Temperature seasonality control on modern halite layers in the Dead Sea: In situ observations, *Geological Society of America Bulletin*, B31661.31661, 2017.

1665 \_\_\_\_\_ Sirota, I., Enzel, Y., and Lensky, N. G.: Halite focusing and amplification of salt layer thickness: From the Dead Sea to deep hypersaline basins, *Geology*, 2018.

\_\_\_\_\_ Stein, M., Starinsky, A., Katz, A., Goldstein, S. L., Machlus, M., and Schramm, A.: Strontium isotopic, chemical, and sedimentological evidence for the evolution of Lake Lisan and the Dead Sea, *Geochim. Cosmochim. Acta*, 61, 3975-3992, 1997.

1670 \_\_\_\_\_ Stöckli, R., Vermote, E., Saleous, N., Simmon, R., and Herring, D.: The Blue Marble Next Generation-A true color earth dataset including seasonal dynamics from MODIS, Published by the NASA Earth Observatory, 2005.

1675 \_\_\_\_\_ Swierczynski, T., Brauer, A., Lauterbach, S., Martín-Puertas, C., Dulski, P., von Grafenstein, U., and Rohr, C.: A 1600 yr seasonally resolved record of decadal-scale flood variability from the Austrian Pre-Alps, *Geology*, 40, 1047-1050, 2012.

\_\_\_\_\_ Tamarin-Brodsky, T. and Kaspi, Y.: Enhanced poleward propagation of storms under climate change, *Nature Geoscience*, 10, 908, 2017.

1680 \_\_\_\_\_ Tarolli, P., Borga, M., Morin, E., and Delrieu, G.: Analysis of flash flood regimes in the North-Western and South-Eastern Mediterranean regions, *Natural Hazards and Earth System Sciences*, 12, 1255-1265, 2012.



- 1685 \_\_\_\_\_ Torfstein, A. and Enzel, Y.: Dead Sea Lake Level Changes and Levant Palaeoclimate, in: Quaternary of the Levant, edited by: Enzel, Y., and Bar-Yosef, O., Cambridge University Press, Cambridge, 115-126, 10.1017/9781316106754.013, 2017.
- 1690 \_\_\_\_\_ Torfstein, A., Goldstein, S. L., Kagan, E. J., and Stein, M.: Integrated multi-site U–Th chronology of the last glacial Lake Lisan, *Geochim. Cosmochim. Acta*, 104, 210-231, 2013a.
- \_\_\_\_\_ Torfstein, A., Goldstein, S. L., Stein, M., and Enzel, Y.: Impacts of abrupt climate changes in the Levant from Last Glacial Dead Sea levels, *Quaternary Science Reviews*, 69, 1-7, 2013b.
- 1695 \_\_\_\_\_ Torfstein, A., Gavrieli, I., Katz, A., Kolodny, Y., and Stein, M.: Gypsum as a monitor of the paleo-limnological–hydrological conditions in Lake Lisan and the Dead Sea, *Geochim. Cosmochim. Acta*, 72, 2491-2509, 2008.
- \_\_\_\_\_ Torfstein, A., Haase-Schramm, A., Waldmann, N., Kolodny, Y., and Stein, M.: U-series and oxygen isotope chronology of the mid-Pleistocene Lake Amora (Dead Sea basin), *Geochim. Cosmochim. Acta*, 73, 2603-2630, 2009.
- 1700 \_\_\_\_\_ Torfstein, A., Goldstein, S. L., Kushnir, Y., Enzel, Y., Haug, G., and Stein, M.: Dead Sea drawdown and monsoonal impacts in the Levant during the last interglacial, *Earth Planet. Sci. Lett.*, 412, 235-244, 2015.
- \_\_\_\_\_ Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bulletin of the American Meteorological society*, 79, 61-78, 1998.
- 1705 \_\_\_\_\_ Tsvieli, Y. and Zangvil, A.: Synoptic climatological analysis of ‘wet’ and ‘dry’ Red Sea troughs over Israel, *International journal of climatology*, 25, 1997-2015, 2005.
- \_\_\_\_\_ Tubi, A. and Dayan, U.: Tropical Plumes over the Middle East: Climatology and synoptic conditions, *Atmospheric Research*, 145, 168-181, 2014.
- 1710 \_\_\_\_\_ Tyrlis, E., Lelieveld, J., and Steil, B.: The summer circulation over the eastern Mediterranean and the Middle East: influence of the South Asian monsoon, *Climate Dynamics*, 40, 1103-1123, 2013.
- 1715 \_\_\_\_\_ Vautard, R. and Ghil, M.: Singular spectrum analysis in nonlinear dynamics, with applications to paleoclimatic time series, *Physica D: Nonlinear Phenomena*, 35, 395-424, 1989.
- \_\_\_\_\_ Waldmann, N., Neugebauer, I., Palchan, D., Hadzhiivanova, E., Taha, N., Brauer, A., and Enzel, Y.: Sedimentology of the Lacustrine Formations in the Dead Sea Basin, in: Quaternary of the Levant: Environments, Climate Change, and Humans, edited by: Enzel, Y., and Bar-Yosef, O., Cambridge University Press, Cambridge, 83-90, DOI: 10.1017/9781316106754.009, 2017.
- 1720 \_\_\_\_\_ Weber, N., Lazar, B., Gavrieli, I., Yechieli, Y., and Stein, M.: Gypsum Deltas at the Holocene Dead Sea linked to Grand Solar Minima, *Geophysical Research Letters*, 10.1029/2020gl091034, 2021.
- 1725 \_\_\_\_\_ Welch, P. D.: The use of fast Fourier transform for the estimation of power spectra: A method based on time averaging over short, modified periodograms, *IEEE Transactions on audio and electroacoustics*, 15, 70-73, 1967.
- 1730 \_\_\_\_\_ Witt, A., Malamud, B. D., Mangili, C., and Brauer, A.: Analysis and modelling of a 9.3 kyr palaeoflood record: correlations, clustering, and cycles, *Hydrology and Earth System Sciences*, 21, 5547-5581, 10.5194/hess-21-5547-2017, 2017.

1735 \_\_\_\_\_ Yosef, Y., Aguilar, E., and Alpert, P.: Changes in extreme temperature and precipitation indices: using an innovative daily homogenized database in Israel, *International Journal of Climatology*, 39, 5022-5045, 2019.

\_\_\_\_\_ Zappa, G., Hoskins, B. J., and Shepherd, T. G.: The dependence of wintertime Mediterranean precipitation on the atmospheric circulation response to climate change, *Environmental Research Letters*, 10, 104012, 2015.

1740 \_\_\_\_\_ Ziv, B.: A subtropical rainstorm associated with a tropical plume over Africa and the Middle-East, *Theoretical and Applied Climatology*, 69, 91-102, 2001.

\_\_\_\_\_ Ziv, B., Saaroni, H., and Alpert, P.: The factors governing the summer regime of the eastern Mediterranean, *International Journal of Climatology*, 24, 1859-1871, 2004.

1745 \_\_\_\_\_ Ziv, B., Dayan, U., Kushnir, Y., Roth, C., and Enzel, Y.: Regional and global atmospheric patterns governing rainfall in the southern Levant, *International Journal of Climatology*, 26, 55-73, 2006.

\_\_\_\_\_ Zolitschka, B., Francus, P., Ojala, A. E., and Schimmelmann, A.: Varves in lake sediments—a review, *Quaternary Science Reviews*, 117, 1-41, 2015.

1750