



## Sampling density and date influence spatial representation of tree ring reconstructions

Justin T. Maxwell<sup>1,2</sup>, Grant L. Harley<sup>3</sup>, Trevis J. Matheus<sup>4</sup>, Brandon M. Strange<sup>5</sup>, Kayla Van Aken<sup>6</sup>, Tsun  
5 Fung Au<sup>1</sup>, and Joshua C. Bregy<sup>1,7</sup>

1 Department of Geography, Indiana University

2 Harvard Forest, Harvard University

3 Department of Geography, University of Idaho

10 4 Department of Geography and the Environment, California State University, Fullerton

5 School of Natural Resources and the Environment, University of Arizona

6 School of Biological, Environmental and Earth Sciences, University of Southern Mississippi

7 Department of Earth and Atmospheric Sciences, Indiana University

15 *Correspondence to:* Justin T. Maxwell (maxweljt@indiana.edu)

**Abstract.** Our understanding of the natural variability of hydroclimate before the instrumental period (*ca.*  
1900 in the United States; US) is largely dependent on tree-ring-based reconstructions. Large-scale soil  
moisture reconstructions from a network of tree-ring chronologies have greatly improved our  
20 understanding of the spatial and temporal variability in hydroclimate conditions, particularly extremes of  
both drought and pluvial (wet) events. However, certain regions within these large-scale reconstructions  
in the US have a sparse network of tree-ring chronologies. Further, several chronologies were collected  
in the 1980s and 1990s, thus our understanding of the sensitivity of radial growth to soil moisture in the  
US is based on a period that experienced multiple extremely severe droughts and neglects the impacts of  
25 recent, rapid global change. In this study, we expanded the tree-ring network of the Ohio River Valley in  
the US, a region with sparse coverage. We used a total of 72 chronologies across 15 species to examine  
how increasing the density of the tree-ring network influences the representation of reconstructing the  
Palmer Meteorological Drought Index (PMDI). Further, we tested how the sampling date influenced the  
reconstruction models by creating reconstructions that ended in the year 1980 and compared them to  
30 reconstructions ending in 2010 from the same chronologies. We found that increasing the density of the  
tree-ring network resulted in reconstructed values that better matched the spatial variability of  
instrumentally recorded droughts and to a lesser extent, pluvials. By sampling tree in 2010 compared to  
1980, the sensitivity of tree rings to PMDI decreased in the southern portion of our region where severe  
drought conditions have been absent over recent decades. We emphasize the need of building a high-  
35 density tree-ring network to better represent the spatial variability of past droughts and pluvials. Further,  
chronologies on the International Tree-Ring Data Bank need updating regularly to better understand how  
the sensitivity of tree rings to climate may vary through time.

Keywords: Drought, Pluvial, Midwest United States, Dendrochronology, Palmer Meteorological Drought

40 Index



## 1 Introduction

Understanding the mechanisms that drive climate variability, particularly before the modern instrumental  
45 record (*ca.* 1900 in the United States; US), depends on proxy-based reconstructions. Precisely-dated tree-  
ring chronologies represent one of the primary proxies that can reconstruct inter-annual climate variability  
over recent centuries to millennia (Fritts, 1976). Tree rings provide robust historical context for  
instrumentally-recorded droughts and pluvials (wet periods) throughout the mid-latitudes (*e.g.*,Stahle and  
Cleaveland 1994 ; Woodhouse and Overpeck, 1998; Cook et al., 1999; Cook et al., 2010; Pederson et al.,  
50 2013; Maxwell and Harley, 2017; Oliver et al. 2019). Most of our understanding of past drought severity  
and variability in North America is the result of the North American Drought Atlas (NADA; Cook et al.,  
1999). The NADA comprises a network of tree-ring chronologies across North America from the  
International Tree-Ring Data Bank (ITRDB; [https://www.ncdc.noaa.gov/data-access/paleoclimatology-  
data/datasets/tree-ring](https://www.ncdc.noaa.gov/data-access/paleoclimatology-data/datasets/tree-ring)), creating a 2.5° x 2.5° reconstruction of summer (average June, July, and August;  
55 JJA) Palmer Drought Severity Index values (Palmer, 1965). The NADA produced multiple centuries of  
both spatial and temporal data of drought variability, providing an essential context to extreme soil-  
moisture conditions witnessed in the most recent centuries. More recently, the Living Blended Drought  
Atlas (LBDA; Cook et al., 2010) updated the NADA using additional tree-ring chronologies from the  
ITRDB and higher spatial-resolution climate data to calibrate models, creating a 0.5° x 0.5° reconstruction  
60 of the Palmer Meteorological Drought Index (PMDI; Palmer, 1965).

While the NADA and LBDA have provided invaluable information of past droughts and pluvials in North  
America, they were generated to compare large, regional events. Each gridded reconstruction uses tree-  
ring data that are within a 450-km radius from the center of each grid point. Therefore, the NADA and  
LBDA are excellent at representing large-scale extremes. However, these drought atlases may not  
65 represent local conditions in areas with sparse coverage of tree-ring chronologies, such as certain regions  
of the midwestern US (Maxwell and Harley, 2017; Strange et al., 2019). The tree-ring chronologies from  
the ITRDB can have biases in terms of tree species used and the spatial density of the tree-ring network



(Zhao *et al.*, 2019). When collecting tree-ring data for the purpose of reconstructing climate, the general goal is to target long-lived species that are sensitive to the climate variable to be reconstructed while also maximizing the length of the reconstruction. However, including multiple species can improve model performance and skill (Pederson *et al.*, 2001; Frank and Esper, 2005; Cook and Pederson, 2011; Maxwell *et al.*, 2011; Pederson *et al.*, 2012; Maxwell *et al.*, 2015). In the US, the ITRDB has excellent spatial replication in certain regions, such as the American Southwest, but other regions are poorly represented, such as the Ohio River Valley (ORV) (Zhao *et al.*, 2019). Due to changes in the density of the tree-ring network of the ITRDB and the use of a large radius (450 km) to reconstruct drought for the LBDA, soil moisture variability at small or local scales is potentially absent in areas that are underrepresented in the tree-ring network. Further, many of the chronologies that are available on the ITRDB were collected in the 1980s and have not been updated (Larson *et al.* 2013; Zhao *et al.*, 2019).

The wealth of climate information derived from tree rings is based on the key assertion that their physiological development is related to specific climatic conditions. An explicit relationship between climate and tree growth can be estimated during the instrumental period. Yet, developing a reconstruction assumes that this climate-tree-growth relationship is stationary over time. This assumption was generally true in the early development of the field of dendrochronology (*ca.* 1920s–1950s; Fritts, 1976). However, as human activities drive the Earth’s climate system into historically unprecedented, and potentially non-stationary and non-analogous conditions (Milly *et al.*, 2008), exceptions to this assumption have emerged. Changes in the drought signal recorded by tree rings have been established only recently (Larson *et al.*, 2013; Maxwell *et al.*, 2015, 2016, 2019; Helcoski *et al.*, 2019), making an investigation of its causes essential to ensuring the interpretability of tree-ring-based hydroclimate reconstructions. Of these recent studies, Maxwell *et al.* (2016) provided the first documentation of an apparent deteriorating relationship between radial tree growth and summer soil moisture that is not accompanied by an increase in signal strength during another season. The declining relationship—referred to as the “Fading Drought Signal”—was consistent across multiple species and sites within the Central Hardwoods Forest region of the Midwestern US. However, Maxwell *et al.* (2019) found that *Acer* (maple) species had a stable relationship, indicating that including species from this genus in reconstructions could improve model performance. In this paper, we test the hypothesis that increasing the spatial density of the tree-ring



network results in reconstructions that better replicate the local variation of the instrumental data. We also examine if the year when trees are sampled influences the climate reconstruction. We calibrate the reconstruction with recent (post-1980) radial growth and climate data and compare to reconstructions generated using data only from pre-1980. We test the hypothesis that including recent data could reduce the amount of variance explained in tree-ring reconstruction of soil moisture in the ORV.

## 2 Methods

### 2.2 *Living Blended Drought Atlas*

For the LBDA, Cook *et al.* (2010) created a gridded instrumental dataset of PMDI to calibrate tree-ring reconstruction models. The instrumental data were created using observations for temperature and precipitation from over 5,000 and 7,000 weather stations, respectively, which were spatially interpolated with a trivariate thin-plate spline in the ANUSPLIN program (Hutchinson, 1995). Cook *et al.* (2010) derived the reconstructions by gathering standardized tree-ring chronologies within 450 km of each instrumental grid point center. Chronologies that were significantly correlated with PMDI were retained and used in a principal component analysis (PCA). The resulting principal components (PCs) that had eigenvalues greater than one were then used as predictors in the reconstruction model. For the LBDA, we gathered both the instrumental and reconstructed  $0.5^\circ \times 0.5^\circ$  gridded PMDI data for the ORV region (Figure 1) from the National Oceanic and Atmospheric Administration, National Center for Environmental Information (<https://www.ncdc.noaa.gov/paleo-search/study/19119>; Cook *et al.*, 2010).

### 2.2 *Ohio River Valley Tree-Ring Network*

To examine how the density of the tree-ring network could impact the reconstruction, we gathered recently published chronologies and collected new chronologies across the ORV to fill the spatial gaps of the ITRDB (Figure 1; Supplemental Table 1). For the new chronologies, we either 1) updated existing chronologies from the ITRDB; 2) sampled new co-occurring species at an ITRDB site, or 3) created new chronologies from previously unsampled sites. For this study, we used a total of 72 chronologies across a variety of 15 species. Of these chronologies, 37 were published, three were newly updated ITRDB records, and 32 new collections (Figure 1; Supplemental Table 1). For the new ( $n=32$ ) and updated ( $n=$



3) chronologies, we used standard field methods to target at least 10 old growth trees for each species using morphological characteristics (Pederson, 2010). We used a hand-held 4.3-mm-diameter increment borer to extract two samples from each tree at breast height, from opposite sides of the tree (Stokes and Smiley, 1968). All newly collected samples were mounted and sanded with progressively finer sandpaper to reveal ring structure. We used the list method to visually crossdate all samples (Yamaguchi, 1991). Each sample was then measured using a Velmex stage with 0.001-mm precision using the program MeasureJ2X (Voor Tech 2008), and then statistically crossdated using the program COFECHA (Holmes, 1983). For the three updated chronologies, we crossdated the new sampled series with those previously sampled and available through the ITRDB.

### 2.3 *Detrending Tree-Ring Series*

For all chronologies, we removed both age-related growth trends and non-climatic influences of tree growth (*e.g.*, forest dynamics or insect outbreaks) by using signal-free standardization (Melvin and Briffa, 2008) with a two-thirds smoothing spline applied to each measured series (Cook and Peters, 1981). To ensure we achieved the desired spline flexibility of the two-thirds spline in the standardization, we used the approximation suggested by Bussberg *et al.* (2020) and used an 83% spline to account for end point adjustments. We stabilized the variance of the standardized chronologies using the data-adaptive power transformation (Cook and Peters, 1997). Signal-free standardization reduces “trend distortion” problems near the ends of the record (Melvin and Briffa, 2008). We trimmed each chronology to remove the portion of the record where low sample depth inflated the variance in standardized growth using an expressed population signal (EPS) value of 0.80 (Wigley *et al.*, 1984).

### 2.4 *Point-by-Point Regression*

We replicated the point-by-point regression procedure for the LBDA in Cook *et al.* (2010) and described in Cook *et al.* (1999) for the ORV tree-ring network. We developed a network of 0.5° x 0.5° grid points reconstructions ( $n = 181$ ) across the ORV region, defined as 37.75–42.25° N, 82.25–90.75° W (Figure 1). Similar to the LBDA, we produced PMDI reconstructions at each grid point by first screening standardized tree-ring chronologies through correlation analysis with PMDI. However, because we



wanted to examine how increasing the density of the tree-ring network influences the reconstruction, we gathered tree-ring chronologies within a 250-km radius from the center of each grid point instead of the 450-km radius used for LBDA. For each grid point, we built a reconstruction model by taking the screened standardized chronologies and conducting a PCA. Per the Kaiser-Guttman rule (Guttman, 1954, Kaiser, 1960), we then used the PCs with eigenvalues greater than one as predictors in a regression model to predict mean June–August (JJA) PMDI. We then used Pearson’s correlation to compare the reconstructed PMDI values from the LBDA to the ORV reconstruction at each grid point. We further chose well-known drought and pluvial years in the instrumental period to examine how the ORV and LBDA compared spatially. To compare the reconstructions with the instrumental data, we calculated the mean absolute error for each extreme event. To examine the species contribution to the overall ORV reconstruction, we gathered the absolute beta weights for each species from the reconstruction model (Frank and Esper, 2005).

## 2.5 Droughts and Pluvials

To determine if the ORV and LBDA reconstructions had differences in the amount of extreme hydroclimatic conditions, we calculated the number of years in each gridded reconstruction that had a JJA PMDI value of  $\geq 2.0$  or  $\leq -2.0$  to represent at least moderately wet and dry conditions, respectively. We further examined how the volatility in extreme conditions compared between the two reconstructions by calculating “flips” from one extreme to the other in consecutive years (Loecke *et al.* 2017; Oliver *et al.*, 2019). We specifically used an index developed by Loecke *et al.* (2017) to quantify large “whiplashes” (termed flips here) interannually. The flip index is defined as:

$$i = \text{PMDI}(t + 1) - t / \text{PMDI}(t + (t + 1))$$

where the index (i) equals the PMDI value of a given year (t) subtracted from the PMDI value of the following year (t + 1), divided by the sum of the PMDI values over the two-year period (t+(t+1)).

Positive index values indicate that conditions shifted from dry to wet over the two-year period.

Similarly, negative values represent a shift from wet to dry conditions. We used an index value  $> 75^{\text{th}}$  percentile to define an abnormally wet period and  $< 25^{\text{th}}$  percentile an extreme dry period. We then calculated wet flip events as years that were abnormally dry followed directly by extreme wet years.



175 Dry flips were calculated as abnormally wet years followed by extreme drought years. Lastly, we  
summed the wet and dry flips to calculate the total flips. These flips were calculated for each grid point  
in the ORV reconstruction where sample depth was determined by an EPS value of 0.80 to reproduce  
the variance in the instrumental data (Wigley *et al.*, 1984). We limited the calculation of flips to the  
period 1658–2005, which was the common period of overlap between the longest gridded ORV  
180 reconstruction and the LBDA.

## 2.6 Model Validation Comparisons

To examine the temporal stability of the relationship between tree growth and PMDI, we followed the  
same validation procedures used for the LBDA (Cook *et al.*, 2010). We used the early half of the common  
period (1901–1955) to calibrate a model between tree growth and PMDI to validate the late half (1956–  
185 2010). We used two tests of fit, the reduction of error statistic (RE) and the coefficient of efficiency (CE;  
Fritts, 1976; Cook *et al.*, 1999), to validate our calibration models. RE and CE both range from  $-\infty$  to +1,  
with positive values indicating robust predictive skill. However, RE is compared to the mean of the  
instrumental data, while CE relies on the verification period mean and therefore is a more conservative  
verification metric. We then compared the variance explained ( $R^2$ ), RE, and CE values between the LBDA  
190 and the ORV PMDI reconstructions for each grid point. We also mapped the gridded reconstructed PMDI  
values from extreme years in the observation period and well-known years in the historical record for  
both the LBDA and the ORV reconstructions to provide examples of spatial difference between the two  
reconstructions.

195 To examine how validation statistics may change based on when the trees were sampled, we created a  
second ORV reconstruction where the most recent year was 1980. This year was chosen because several  
chronologies available on the ITRDB were sampled in the 1980s, and this marked the beginning of a  
weakening relationship between radial growth and soil moisture in this region (Maxwell *et al.*, 2016). We  
used the same validation process described above except the early period was from 1901 to 1940 and the  
200 late period was from 1941 to 1980. We then calculated the difference between the 1980 and the 2010  
reconstruction for  $R^2$ , RE, and CE values for each grid point.



### 3 Results

#### 3.1 ORV vs. LBDA

Our first comparisons of chronologies distributed for the LBDA and ORV networks revealed broad spatial  
205 discrepancies. PMDI point-by-point regressions for the LBDA included 20 chronologies from six species  
over the study region, whereas the ORV network included 72 chronologies from 15 tree species. Not only  
is the spatial density of sites more sparse for the LBDA network, but it only included mostly single-  
chronology sites, whereas many ( $n = 18$ ) of the sites included in the ORV are multiple-chronology sites  
(2–6 co-occurring species) (Figure 1A, B). Although site coverage is sparse for both networks along the  
210 west-central, northwest, and southeast sectors, the ORV network included major spatial coverage  
improvements in other sectors (Figure 1). Particularly, the ORV increased spatial coverage in south-  
central Indiana where many of the sites included 4–6 co-occurring species chronologies ( $n = 27$  total  
chronologies). The PMDI reconstructions from the ORV network and the LBDA demonstrated strong  
and positive correlations, with  $r$ -values ranging from 0.50 to 0.90 (Figure 2). These correlations were  
215 calculated for the period of overlap between the two gridded reconstructions, 1830–2005 C.E. The highest  
correlations were found along the western portion of the gridded region, while the lowest agreement was  
found in the southeast (Figure 2).

The ORV reconstructions were shorter in length (maximum of 343 years) compared to the LBDA  
reconstructions (maximum of 2,006 years) due to each grid reconstruction having a smaller search radius  
220 (250 km vs 450 km) for chronology inclusion. The larger search radius allows the inclusion of longer  
chronologies in more of the gridded reconstructions. Secondly, we focused on increasing the spatial  
density of the network, which resulted in sampling younger sites (*e.g.*, the earliest years are in the early  
to late 19<sup>th</sup> century). While the ORV reconstructions were shorter, comparing certain well known extreme  
climatic years during the period of the overlap between the LBDA show some important differences.

#### 225 3.2 ORV and LBDA Extreme Year Comparisons

We chose a series of well-known drought and pluvial years (events) to compare the reconstructions  
between ORV and LBDA. Specifically, we examined the droughts of 1988, 1954, 1936, 1816, and 1774





and the pluvial periods of 1945–1951, 1882–1883, and 1811. In general, the increased spatial density of tree-ring chronologies used in the ORV reconstruction displayed more local variation in the reconstructions of extreme climatic events (Figure 3). However, in a few examples, such as 1774 and 1816, the spatial pattern of where extreme drought was located changed between the two reconstructions (Figure 3). Using extreme events in the observed record (three droughts and one pluvial), both the ORV and LBDA underestimated wet and dry extremes. However, the ORV reconstruction better matched the distribution of soil moisture values and the spatial patterns of the instrumental data compared to the LBDA reconstruction (Figure 4; Supplemental Figs 1–3). For droughts, the ORV consistently had lower mean absolute errors (differences ranging for 0.21 to 0.41) compared to the LBDA (Figure 4; Supplemental Figs 1–3). However, for the pluvial event the two reconstructions had similar mean absolute errors (difference of 0.03) with the LBDA being slightly smaller (Supplemental Fig 3).

In general, the probability distribution function (PDF) of the ORV reconstruction had a lower occurrence (densities of 0.17 compared to 0.23) of near-average years but higher densities (differences ranging from 0.01 to 0.05) for extremes, particularly drought, compared to the LBDA (Figure 5). Similarly, the ORV had a larger number of reconstructed drought (median difference of 32 years) and pluvial (difference of 7 years) conditions compared to the LBDA (Figure 5). Due to the larger number of extreme years, the ORV reconstructions had more frequent flips according to the flip index values compared to the LBDA (Figure 6). The central and southeastern portions of the region in particular showed a greater number of wet, dry, and total flips, resulting in ~30 more wet and dry flips and ~60 more total flips (Figure 6).

### 3.3 Species Contributions

With the highest average beta-weight values, *Quercus* spp. chronologies demonstrated consistently to be the strongest contributors to reconstruction models (Figure 7). The chestnut oak (*Q. montana*) chronology from Dale and Jackie Riddle State Nature Preserve in Ohio had the highest single-forest model beta weight at 0.68, though as a species, *Q. alba* were the strongest species contributor (Figure 7). The Lincoln's New Salem *Q. alba* collection demonstrated the strongest correlation to JJA PMDI, with  $r = 0.75$  during the period 1900–2014. *Q. alba* beta values had a distribution with the highest median and smallest



interquartile range. In addition to *Quercus* spp., *L. tulipifera* was also strong contributors to drought  
255 models, with median beta values of 0.08 (Figure 7).

### 3.2 ORV and LBDA Validation Statistics

Comparing how well each reconstruction model represented the instrumental data, we find that the  
variance explained ( $R^2$ -values) in the calibration and verification periods match well for the northern  
portion of the network, with values ranging from 40 to 60 percent variance explained (Figure 8). However,  
260 the ORV models for the southern half of the region generally explain less variance compared to the LBDA  
(Figure 8). Interestingly, the RE- and CE-values between the two reconstructions are generally more  
similar, with the ORV having poorer validation statistics in the southernmost portion of the region and  
the LBDA having weaker statistics in the central portion of the region (Figure 8).

Previous work has shown that radial growth from trees in the south-central portion of the region are  
265 becoming less sensitive to soil moisture compared to earlier time periods (Maxwell *et al.*, 2016). The  
comparison between a point-by-point reconstruction that ended in 1980 to a reconstruction that ended in  
2010 demonstrates that while the calibration  $R^2$ -values are similar, the 2010 verification models explain  
much less variance in the southern portion of the ORV (Figure 9). These are the same regions in the ORV  
reconstruction that explain less variance compared to the LBDA. Importantly, the ORV 1980 and 2010  
270 reconstructions used the same tree-ring chronologies (Figure 9). Therefore, our results indicate that tree  
rings in the southern portion of our study region have become less responsive to soil moisture.

## 4 Discussion

### 4.1 ORV and LBDA Extreme Year Comparisons

Tree rings have long been used to provide an historical context to past hydroclimatic extremes (Stahle  
275 and Cleaveland 1994; Woodhouse and Overpeck, 1998; Cook *et al.*, 1999; Cook *et al.*, 2010; Pederson *et al.*,  
*et al.*, 2013). However, in some regions in the US, the tree-rings sites are sparsely distributed, and it is  
unknown what kind of impact that has on the representation of past climate. Due to the higher density of  
tree-ring chronologies and the smaller search radius (250 km for the ORV compared to 450 km for LBDA)  
of the PC regression models when determining the pool of predictors, the ORV better replicates the spatial



280 variability of the instrumental data compared the LBDA (Figure 4; Supplemental Figures 1–3). By using  
a 450-km radius for potential tree-ring chronologies, the LBDA was successful at reconstructing soil  
moisture even in areas that have a limited number of tree-ring chronologies. However, this approach  
results in the use of the same tree-ring chronologies in multiple gridded reconstructions, spatially  
smoothing the variability of the reconstructed PMDI compared to the instrumental data. The same is true  
285 of the ORV; however, the increase in the spatial density of the chronologies allows a smaller search radius  
and therefore, can increase the spatial variability in the ORV. The increase in spatial variability in PMDI  
values of the ORV better matches the instrumental data while still providing a statistically valid  
reconstruction model (Figure 4; Supplemental Figures 1–3). These findings have important implications,  
particularly in regions with a sparse tree-ring network where the LBDA likely underestimates localized  
290 droughts and pluvials. Increasing the spatial density of the tree-ring network will allow a more accurate  
spatial representation of extreme events nearly anywhere where trees are sensitive to climate.

In addition to the increase in spatial variability of extremes that we find, previous work suggests  
increasing the density of the tree-ring network can result in the discovery of previously unknown droughts  
295 and pluvials at more local scales (Maxwell and Harley, 2017). Here, we find support of better localized  
representations of extremes by increasing the density of the tree-ring network with the ORV having a  
larger number of droughts and pluvials compared to the LBDA (Figure 5). The increase in extremes have  
important implications on the long-term variability of past hydroclimate and to the interannual volatility  
of PMDI. Recent work has shown increases in interannual volatility has important impacts on agriculture  
300 (Locke *et al.*, 2017), and social and ecological systems (Casson *et al.*, 2019). Our finding suggests that in  
areas with a sparse tree-ring network, such as in the ORV, tree-ring reconstructions underestimate  
extremes and therefore, volatility in extremes is also underestimated. We find a higher number of flips by  
increasing the tree-ring network (Figure 6) and therefore, provide a better representation of past volatility  
to put current and future projected change into context.

#### 305 4.2 *Species Contributions*



Historically, soil moisture reconstructions from tree rings in the eastern US have been dominated by a few species, e.g. *Quercus alba*, *Taxodium distichum*, *Tsuga canadensis* (Zhao *et al.*, 2019). In addition to increasing the spatial density of the network, the ORV reconstruction has increased the number of species  
310 used, many of which are co-occurring. The use of multiple species has been shown to increase model performance (Pederson *et al.*, 2001; Frank and Esper, 2005; Cook and Pederson, 2011; Maxwell *et al.*, 2011; Pederson *et al.*, 2012, Maxwell *et al.* 2015). Examining the beta values of the species used in the reconstructions models, *Quercus* (oak) species in general contribute more to the models (Figure 7), which is part of the reason why they have been traditionally used so frequently. However, we find that several  
315 species, including *Liriodendron tulipifera* (tuliptree), make strong contributions to the model as well (Figure 7). These findings agree with recent studies that suggest less commonly used species can increase the representativeness of tree-ring reconstructions of climate (Pederson *et al.*, 2012; Maxwell, 2016; Maxwell and Harley, 2017).

#### 4.3 ORV and LBDA Validation Statistics

320 While increasing the spatial density of the tree-ring network allowed the reconstructions to more accurately capture the spatial variability of extreme conditions, the reconstruction models of the ORV have less predictive skill compared to those of the LBDA, especially during the verification period (Figure 8). The two networks have some overlap in chronologies, but while the ORV has a higher density of chronologies within the Ohio River Valley region, the LBDA can draw from more chronologies across a  
325 larger region. While the larger radius increases the number of samples in the model and could lead to more explained variance for the LBDA, the ORV reconstruction better spatially replicates extremes in the instrumental period (Figures 4; Supplemental Figures 1–3).

330 Interestingly, the decrease in variance explained in the southern portion of the region may not attribute from differences of sample depth in the tree-ring network. When using the same chronologies while ending the calibration period at 1980 instead of 2010 for the ORV reconstruction, the validation statistics compare very well with the LBDA. However, by updating the chronologies to 2010, the  $R^2$  and the validation statistics drop dramatically for the grid reconstructions in the southern portion of the region



(Figure 9). These findings support Maxwell *et al.* (2016), where they found trees in in this region to have  
335 a weakening signal to soil moisture, termed the “Fading Drought Signal.” The recent decrease in  
sensitivity of tree growth to soil moisture has also been documented outside of the ORV, in the Mid-  
Atlantic US (Helcoski *et al.*, 2019), indicating the impact of a changing climate could influence the  
representation of tree rings to climate in mid-latitude locations. Drought in the Midwest during the  
instrumental period (1901–2010) was temporally clustered in the 1930s and 1950s. The only recent  
340 droughts in the study period were in 1988 and 2002. In both cases, the northern portions of the region  
experienced severe drought (in excess of -4.0 PMDI values for 1988), but the southern portion of the  
region only experienced moderate dryness (PMDI values of  $\sim -2.0$ ). Maxwell *et al.* (2016) attributed the  
weakening signal to a recent period without severe drought; however, Helcoski *et al.* (2019) discussed  
the possibility of increases in carbon dioxide concentrations in addition to a long period of wetness  
345 interacting to weaken tree growth responses to soil moisture. However, recent works examining the  
simultaneous influence of water availability, carbon dioxide concentrations, and acidic deposition found  
that water availability was the leading influence on tree growth (Levesque *et al.*, 2017; Maxwell *et al.*,  
2019), suggesting a wet period is likely driving the weakening signal. The decreasing performance of the  
southern reconstructions support these findings as this region has been generally wet and absent of severe  
350 drought. While Maxwell *et al.* (2019) found that *Acer* species had a more stable relationship with soil  
moisture, the inclusion of multiple, co-occurring *A. saccharum* records did not dramatically influence the  
performance of the reconstruction models in the southern portion of the region. Our findings demonstrate  
the complexity of tree species interactions with environmental variability in the midwestern US with  
regard to rapidly changing climate regimes and stress the need to better understand species responses to  
355 changing climate and what impact that could have on reconstructions of soil moisture.

## 5 Conclusions

By increasing the density of the tree-ring network in a region that is poorly represented in the LBDA, we  
created a gridded PMDI reconstruction for the ORV region. We compared our gridded reconstruction  
with the LBDA and found that increasing the density of the tree-ring network resulted in an increase in  
360 localized hydroclimatic extremes that better match the spatial and temporal patterns of the instrumental



data. However, calibrating our models with more recent data (up to the year 2010) resulted in a decrease in variance explained for the southern portion of the region. This region has not experienced extreme droughts recently, which is likely driving the decrease in model performance. Increasing spatial density of the tree-ring network is important to better represent localized extremes in the past, indicating that  
365 researchers should continue to target previously unsampled old-growth forests. Similarly, the time in which the trees are sampled is also important to model performance. Long periods without extreme hydroclimate variability can result in reconstruction models that are less representative of climatic conditions. We stress the need to update previously-sampled chronologies to the current period so that longer calibration models can have the chance to better represent the range of sensitivity of trees rings to  
370 climate. Overall, we find that a higher spatial density of the tree-ring network will improve the local representation of reconstructed climate. However, more work is needed to better quantify how the strength of the relationship between tree growth and climate varies through time.

Data Availability: All reconstructions will be uploaded onto the NOAA paleoclimate page. All tree ring  
375 chronologies used in this manuscript will but uploaded to the International Tree-Ring Databank.

Author Contributions: JTM and GLH designed the methods of manuscript. JTM preformed analyses with feedback from GLH. TJM, BMS, KVK, and TFA helped develop tree ring chronologies with assistance from JTM and GLH. All authors contributed to data collection and the preparation of the manuscript.  
380

Acknowledgments: The collection of tree ring samples was partially funded by a USDA Agriculture and Food Research Initiative grant 2017-67013-26191 and from the Indiana University Vice Provost for Research Faculty Research Program. We would like to thank James Dickens, James McGee, Josh Oliver, Karly Schmidt-Simard, Brynn Taylor, Michael Thornton, Senna Robeson, Matt Wenzel, and Luke Wylie  
385 for assistance in the field and laboratory.



## References

- 390 Bussberg, N. W., Maxwell, J. T., Robeson, S. M. and Huang, C.: The effect of end-point adjustments on smoothing splines used for tree-ring standardization, *Dendrochronologia*, 60, 125665, doi:10.1016/j.dendro.2020.125665, 2020.
- 395 Casson, N. J., Contosta, A. R., Burakowski, E. A., Campbell, J. L., Crandall, M. S., Creed, I. F., Eimers, M. C., Garlick, S., Lutz, D. A., Morison, M. Q., Morzillo, A. T. and Nelson, S. J.: Winter Weather Whiplash: Impacts of Meteorological Events Misaligned With Natural and Human Systems in Seasonally Snow-Covered Regions, *Earth's Future*, 7(12), 1434–1450, doi:10.1029/2019EF001224, 2019.
- 400 Coats, S., Smerdon, J. E., Cook, B. I., Seager, R., Cook, E. R. and Anchukaitis, K. J.: Internal ocean-atmosphere variability drives megadroughts in Western North America, *Geophysical Research Letters*, 43(18), 9886–9894, doi:10.1002/2016GL070105, 2016.
- 405 Cook, E. R. and Pederson, N.: Uncertainty, Emergence, and Statistics in Dendrochronology, in *Dendroclimatology: Progress and Prospects*, edited by M. K. Hughes, T. W. Swetnam, and H. F. Diaz, pp. 77–112, Springer Netherlands, Dordrecht., 2011.
- Cook, E. R. and Peters, K.: The Smoothing Spline: A New Approach to Standardizing Forest Interior Tree-Ring Width Series for Dendroclimatic Studies, [online] Available from: <https://repository.arizona.edu/handle/10150/261038> (Accessed 21 February 2020), 1981.
- 410 Cook, E. R. and Peters, K.: Calculating unbiased tree-ring indices for the study of climatic and environmental change, *The Holocene*, 7(3), 361–370, doi:10.1177/095968369700700314, 1997.
- Cook, E. R., Meko, D. M., Stahle, D. W. and Cleaveland, M. K.: Drought Reconstructions for the Continental United States, *J. Climate*, 12(4), 1145–1162, doi:10.1175/1520-0442(1999)012<1145:DRFTCU>2.0.CO;2, 1999.



415 Cook, E. R., Seager, R., Heim, R. R., Vose, R. S., Herweijer, C. and Woodhouse, C.: Megadroughts in  
North America: placing IPCC projections of hydroclimatic change in a long-term palaeoclimate context,  
Journal of Quaternary Science, 25(1), 48–61, doi:10.1002/jqs.1303, 2010.

Frank, D., Wilson, R. and Esper, J.: Synchronous variability changes in Alpine temperature and tree-ring  
420 data over the past two centuries, Boreas, 34(4), 498–505, doi:10.1080/03009480500231443, 2005.

Fritts, H.: Tree Rings and Climate, Academic Press. New York., 1976.

Guttman, L.: Some necessary conditions for common-factor analysis, Psychometrika, 19(2), 149–161,  
425 doi:10.1007/BF02289162, 1954.

Helcoski, R., Tepley, A. J., Pederson, N., McGarvey, J. C., Meakem, V., Herrmann, V., Thompson, J. R.  
and Anderson-Teixeira, K. J.: Growing season moisture drives interannual variation in woody  
productivity of a temperate deciduous forest, New Phytologist, 223(3), 1204–1216,  
430 doi:10.1111/nph.15906, 2019.

Holmes, R. L.: COMPUTER -ASSISTED QUALITY CONTROL IN TREE -RING DATING AND  
MEASUREMENT, , 11, n.d.

435 Hutchinson, M. F.: Interpolating mean rainfall using thin plate smoothing splines, International Journal  
of Geographical Information Systems, 9(4), 385–403, doi:10.1080/02693799508902045, 1995.

Kaiser, H. F.: The Application of Electronic Computers to Factor Analysis, Educational and  
Psychological Measurement, 20(1), 141–151, doi:10.1177/001316446002000116, 1960.

440





- Larson, E. R., Allen, S., Flinner, N. L., Labarge, S. G. and Wilding, T. C.: The Need and Means To Update Chronologies In A Dynamic Environment, *trre*, 69(1), 21–27, doi:10.3959/1536-1098-69.1.21, 2013.
- 445 Levesque, M., Andreu-Hayles, L. and Pederson, N.: Water availability drives gas exchange and growth of trees in northeastern US, not elevated CO<sub>2</sub> and reduced acid deposition, *Sci Rep*, 7(1), 1–9, doi:10.1038/srep46158, 2017.
- Loecke, T. D., Burgin, A. J., Riveros-Iregui, D. A., Ward, A. S., Thomas, S. A., Davis, C. A. and Clair,  
450 M. A. St.: Weather whiplash in agricultural regions drives deterioration of water quality, *Biogeochemistry*, 133(1), 7–15, doi:10.1007/s10533-017-0315-z, 2017.
- Maxwell, J. T.: The Benefit of Including Rarely-Used Species in Dendroclimatic Reconstructions: A Case Study Using *Juglans nigra* in South-Central Indiana, USA, *Tree-Ring Research*, 72(1), 44–52,  
455 doi:10.3959/1536-1098-72.01.44, 2016.
- Maxwell, J. T. and Harley, G. L.: Increased tree-ring network density reveals more precise estimations of sub-regional hydroclimate variability and climate dynamics in the Midwest, USA, *Clim Dyn*, 49(4), 1479–1493, doi:10.1007/s00382-016-3396-9, 2017.
- 460 Maxwell, J. T., Harley, G. L. and Matheus, T. J.: Dendroclimatic reconstructions from multiple co-occurring species: a case study from an old-growth deciduous forest in Indiana, USA, *International Journal of Climatology*, 35(6), 860–870, doi:10.1002/joc.4021, 2015.
- 465 Maxwell, J. T., Harley, G. L. and Robeson, S. M.: On the declining relationship between tree growth and climate in the Midwest United States: the fading drought signal, *Climatic Change*, 138(1), 127–142, doi:10.1007/s10584-016-1720-3, 2016.



- Maxwell, J. T., Harley, G. L., Mandra, T. E., Yi, K., Kannenberg, S. A., Au, T. F., Robeson, S. M.,  
470 Pederson, N., Sauer, P. E. and Novick, K. A.: Higher CO<sub>2</sub> Concentrations and Lower Acidic Deposition  
Have Not Changed Drought Response in Tree Growth But Do Influence iWUE in Hardwood Trees in the  
Midwestern United States, *Journal of Geophysical Research: Biogeosciences*, 124(12), 3798–3813,  
doi:10.1029/2019JG005298, 2019.
- 475 Maxwell, R. S., Hessl, A. E., Cook, E. R. and Pederson, N.: A multispecies tree ring reconstruction of  
Potomac River streamflow (950–2001), *Water Resources Research*, 47(5), doi:10.1029/2010WR010019,  
2011.
- Melvin, T. M. and Briffa, K. R.: A “signal-free” approach to dendroclimatic standardisation,  
480 *Dendrochronologia*, 26(2), 71–86, doi:10.1016/j.dendro.2007.12.001, 2008.
- Milly, P. C. D., Betancourt, J., Falkenmark, M., Hirsch, R. M., Kundzewicz, Z. W., Lettenmaier, D. P.  
and Stouffer, R. J.: Stationarity Is Dead: Whither Water Management?, *Science*, 319(5863), 573–574,  
doi:10.1126/science.1151915, 2008.
- 485 Palmer, W. C.: *Meteorological Drought*, U.S. Department of Commerce, Weather Bureau., 1965.
- Pederson, N.: External Characteristics of Old Trees in the Eastern Deciduous Forest, *naar*, 30(4), 396–  
407, doi:10.3375/043.030.0405, 2010.
- 490 Pederson, N., Jacoby, G. C., D’Arrigo, R. D., Cook, E. R., Buckley, B. M., Dugarjav, C. and Mijiddorj,  
R.: Hydrometeorological Reconstructions for Northeastern Mongolia Derived from Tree Rings: 1651–  
1995, *J. Climate*, 14(5), 872–881, doi:10.1175/1520-0442(2001)014<0872:HRFNMD>2.0.CO;2, 2001.
- Pederson, N., Bell, A. R., Knight, T. A., Leland, C., Malcomb, N., Anchukaitis, K. J., Tackett, K., Scheff,  
495 J., Brice, A., Catron, B., Blozan, W. and Riddle, J.: A long-term perspective on a modern drought in the  
American Southeast, *Environ. Res. Lett.*, 7(1), 014034, doi:10.1088/1748-9326/7/1/014034, 2012.



Pederson, N., Bell, A. R., Cook, E. R., Lall, U., Devineni, N., Seager, R., Eggleston, K. and Vranes, K. P.: Is an Epic Pluvial Masking the Water Insecurity of the Greater New York City Region?, *J. Climate*, 500 26(4), 1339–1354, doi:10.1175/JCLI-D-11-00723.1, 2013.

Routson, C. C., Woodhouse, C. A. and Overpeck, J. T.: Second century megadrought in the Rio Grande headwaters, Colorado: How unusual was medieval drought?, *Geophysical Research Letters*, 38(22), doi:10.1029/2011GL050015, 2011.

505

Stahle, D. W. and Cleaveland, M. K.: Tree-Ring Reconstructed Rainfall Over the Southeastern U.S.A. During the Medieval Warm Period and Little Ice Age, in *The Medieval Warm Period*, edited by M. K. Hughes and H. F. Diaz, pp. 199–212, Springer Netherlands, Dordrecht., 1994.

510 Stahle, D. W., Cook, E. R., Cleaveland, M. K., Therrell, M. D., Meko, D. M., Grissino-Mayer, H. D., Watson, E. and Luckman, B. H.: Tree-ring data document 16th century megadrought over North America, *Eos, Transactions American Geophysical Union*, 81(12), 121–125, doi:10.1029/00EO00076, 2000.

Strange, B. M., Maxwell, J. T., Robeson, S. M., Harley, G. L., Therrell, M. D. and Ficklin, D. L.: 515 Comparing three approaches to reconstructing streamflow using tree rings in the Wabash River basin in the Midwestern, US, *Journal of Hydrology*, 573, 829–840, doi:10.1016/j.jhydrol.2019.03.057, 2019.

Swain, D. L., Langenbrunner, B., Neelin, J. D. and Hall, A.: Increasing precipitation volatility in twenty-first-century California, *Nature Clim Change*, 8(5), 427–433, doi:10.1038/s41558-018-0140-y, 2018.

520

Voor Tech Consulting, 2008. Measure J2X



525 Wigley, T. M. L., Briffa, K. R. and Jones, P. D.: On the Average Value of Correlated Time Series, with  
Applications in Dendroclimatology and Hydrometeorology, *J. Climate Appl. Meteor.*, 23(2), 201–213,  
doi:10.1175/1520-0450(1984)023<0201:OTAVOC>2.0.CO;2, 1984.

530 Woodhouse, C. A. and Overpeck, J. T.: 2000 Years of Drought Variability in the Central United States,  
*Bull. Amer. Meteor. Soc.*, 79(12), 2693–2714, doi:10.1175/1520-  
0477(1998)079<2693:YODVIT>2.0.CO;2, 1998.

535 Yamaguchi, D. K.: A simple method for cross-dating increment cores from living trees, *Can. J. For. Res.*,  
21(3), 414–416, doi:10.1139/x91-053, 1991.

540 Zhao, S., Pederson, N., D’Orangeville, L., HilleRisLambers, J., Boose, E., Penone, C., Bauer, B., Jiang,  
545 Y. and Manzanedo, R. D.: The International Tree-Ring Data Bank (ITRDB) revisited: Data availability  
and global ecological representativity, *Journal of Biogeography*, 46(2), 355–368, doi:10.1111/jbi.13488,  
2019.

540

545

550



555

560

Figures:





Figure 1: Map of  $0.5^\circ \times 0.5^\circ$  PMDI grid points ( $n = 181$ ) across the Ohio River Valley (ORV) region, 565 Midwest US—defined as  $37.75\text{--}42.25^\circ$  N,  $90.75\text{--}82.25^\circ$  W—plotted with tree-ring chronology sites included from the (A) ITRDB and (B) ORV networks. Sites with single-species and multiple-species are denoted by symbol shape and color (see Supplemental Table 1). Note: most ITRDB sites consist of single species in the LBDA but multiple species are represented in the ORV.



570

Figure 2: Map of correlation values between the LBDA and ORV reconstruction during the period of 1830–2005. The correlations of each grid shown in the map are all significant at the 0.05-level. The white grids represent locations over the Great Lakes and therefore, no data is available for correlation analysis.

575




Figure 3: Spatial comparison of the ORV (left column) and the LBDA (right column) of reconstructed PMDI during years that experienced hydroclimatic extremes. Red cells represent below average PMDI and blue cells represent above average PMDI. White cells represent no data either due to being over water or from no chronologies being old enough to create a reconstruction.






Figure 4: A map of PMDI values for the instrumental data, ORV, and LBDA reconstructions for the year 1954. The histogram represents frequency of PMDI values for the instrumental, ORV, and LBDA PMDI values. The mean absolute error values show that the ORV reconstruction better matches the instrumental data compared to the LBDA reconstruction. White grids represent areas over water and therefore, no data.




Figure 5: A) Probability distribution functions for all gridded reconstructed PMDI values for the ORV and LBDA reconstructions. B) Boxplot of the number of droughts (PMDI  $\le -2.0$ ) years between LBDA and ORV. C) Boxplot of the number of pluvials (PMDI  $\ge 2.0$ ) years between LBDA and ORV.




Figure 6: Maps of the number of wet flips (top row), dry flips (middle row), and total flips (bottom row), for the ORV (left column) and the LBDA (right column). White grids represent values over water and therefore, no data.




Figure 7: Beta values from the PMDI reconstruction models for each species. The “x” represents the mean  
605 beta weight for the species. QUMO=*Quercus montana*, QUAL=*Q. alba*, QURU=*Q. rubra*, QUMA=*Q.*  
*macrocarpa*, LITU=*Liriodendron tulipifera*, TSCA=*Tsuga canadensis*, FAGR=*Fagus grandifolia*,  
FRNI=*Fraxinus nigra*, ACSA=*Acer saccharum*, PIST=*Pinus strobus*, CAOV=*Carya ovata*,  
JUNI=*Juglans nigra*, and QUVE= *Q. velutina*. The species are ranked by their mean beta values from  
highest to lowest.

610





Figure 8: Comparison of the calibration and validation statistics between the ORV (left column) and LBDA (right column) reconstructions.



615 Figure 9: Maps of the difference between the ORV reconstruction when ending the calibration period in 2010 compared to 1980 for calibration  $R^2$ , verification  $R^2$ , RE, and CE.