

1 **Last 2400 yrs. of environmental changes and human activity recorded in the**
2 **gyttja-type bottom sediments of the Lake Mlynek (Warmia and Masuria**
3 **Region, northern Poland)**
4

5 Fabian Welc (1), Jerzy Nitychoruk (2), Leszek Marks (3), Krzysztof Bińka (3), Anna Rogóż-
6 Matyszczyk (2), Milena Obremska (4), Abdelfattah Zalut (5)

7
8 1. Institute of Archaeology, Cardinal Stefan Wyszyński University in Warsaw, e-mail: f.welc@uksw.edu.pl.

9 2. Faculty of Economic and Technical Sciences, Pope John Paul II State Higher School of Education, e-mail:
10 jerzy.nitychoruk@pswbp.pl, annarogoz@interia.pl

11 3. Faculty of Geology, University of Warsaw, e-mail: k.binka@uw.edu.pl, leszek.marks@uw.edu.pl

12 4. Polish Academy of Sciences, Institute of Geological Sciences, e-mail: mobremska@twarda.pan.pl

13 5. Tanta University, Faculty of Science, Tanta University, e-mail: abzalut@science.tanta.edu.eg
14

15 **Abstract**

16 *In the densely forested Warmia and Masuria region (north Poland) there are many endorheic lakes*
17 *characterized by small size and ~~slow~~ **sedimentation**, which makes them excellent Holocene*
18 *environmental and palaeoclimatic archives. One of them - the Lake Mlynek, located near the*
19 *village of Janiki Wielkie, has been selected for multi-faceted palaeoenvironmental research based*
20 *on a precise radiocarbon ~~scale~~. Sediments of this lake also contain unique information about*
21 *human impact on the environment, because a stronghold has been operating on its northern shore*
22 *since the early Iron Age to the early Medieval period, ~~which gives~~ opportunity to correlate*
23 *palaeoenvironmental data with phases of the human activity in the last 2400 years. During 3rd and*
24 *2nd century BC ~~in~~ the lake was surrounded by a dense deciduous forest. From the 1st century BC to*
25 *2nd century AD the forest around the lake was much reduced, what can be associated with the first*
26 *pre roman (La Tene) and Roman occupation phase ~~attested on~~ the stronghold located close to the*
27 *lake. From the 2nd up to 9th century AD ~~gradual~~ restoration of the forest and ~~decline~~ of human*
28 *activity ~~is attested~~, along with lake deepening and onset of colder and humid climatic phase which*
29 *corresponded to ~~global~~ cooling episode known as ~~the~~ Bond 1 Event (1.5 ka BP). The next intensive*
30 *forest clearing around the lake occurred in the 9th – 13th century AD as result of human activity*
31 *(Middle Age settlement phase on the stronghold). This period is marked by a ~~climate change~~*
32 *towards ~~warming~~, which is confirmed by a gradual shallowing of the lake (Middle Age Warm*

33 *Period). In latter time, the strong human activity which transformed the landscape caused that*
34 *possible climate-induced natural environmental changes are not so clear. Only cooling during the*
35 *Little Ice Age is indicated by the changing sedimentation rate in the lake.*

36
37 **Keywords:** late Holocene, lake sediments, Lake Młynek, environmental change, human impact, Late Holocene, Iron
38 Age, Middle Ages, north Poland.

40 1. Introduction

41 Lake sediments are a useful source of proxies of past environmental and climate changes
42 in the Holocene (see Brauer, 2004; Zolitschka, 2007; Wanner et al., 2008; Francus et al., 2013;
43 Ojala et al., 2013; Welc, 2017). The main advantage of lake sediment archives is usually continuous
44 and uninterrupted accumulation. Well-dated lake sediment columns let to trace both long- and
45 short-term Holocene palaeoclimate changes (Smol et al., 2001; Tiljander et al., 2002; Valpola and
46 Ojala, 2006; Czymzik et al., 2010; Elbert et al., 2012; Tylmann et al., 2012; Welc, 2017).
47 Particularly valuable for palaeoclimate reconstructions are sequences from lakes, without river
48 spring inflow and outflow (Wetzel, 2001; Stankevica et al., 2015). In northeastern Poland eutrophic
49 lakes are common, similarly as in northeastern Europe. They are typical for their substantial
50 primary production (algae and aquatic macrophytes), because of the predominance of nutrient input
51 over mineralization processes (Cooke et al., 2005). Such intensive bio-productivity results in a
52 deposition of thick organic sedimentary sequences, mostly of organic gyttja composed of remains
53 of aquatic plants, plankton and benthic organisms transformed by activity of bacteria and mixed
54 with mineral components supplied from the lake basin (Kurzo et al., 2004; Stankevica et al., 2015).
55 There are ca. 1000 freshwater lakes of different size in the Warmia and Mazury Region in north
56 Poland (Fig. 1). Most of them are located within past glacial tunnel valleys formed by meltwater
57 erosion at the termination of the Vistulian (Weichselian) Glaciation (ca. 115-12 ka BP). After
58 deglaciation at the end of the Pleistocene these tunnel valleys were partly filled with deposits and
59 water, and persisted in the Holocene. Such lake basins have steep slopes and their bottom deposits
60 are underlain by glaciofluvial sand, gravel and silt or glacial till (Kondracki, 2002; Gałązka, 2009).
61 Many lakes in the Warmia and Mazury Region are small (<1 ha), with stable sedimentation rate
62 and without river inflow and outflow. It is among the reasons that palaeoclimatic investigations,
63 based mainly on pollen analysis are undertaken in this area (e.g., Kupryjanowicz, 2008; Kołaczek
64 et al., 2013).

65 Lake Mlynek is located near the village of Janiki Wielkie and it was selected for multi-
66 faceted palaeoenvironmental research (pollen analysis, diatom, chrysophyte cysts, geochemistry-
67 and other) based on a precise radiocarbon scale, as it is hypothesized that the bottom sediments of
68 this lake contain a unique record of human impact on environment, as a result of the location of an
69 Iron Age stronghold on the northern shore, which was active (though not continuously) up until
70 the early Middle Ages (Fig. 1). Performed analysis provided an opportunity to reconstruct the
71 transformation of the vegetation around the lake and the changes in the reservoir that occurred
72 under the influence of the climate (regional significance) and as a result of human activity. Our
73 results were correlated with geoarchaeological data to determine mutual relations between
74 environmental and climatic changes with development of human settlement phases in the Warmia
75 and Mazury Region during the last 2000 years.

76

77 2. Study area

78 The Lake Mlynek is a small water body that has occupied a glacial tunnel valley in the Holocene.
79 The lake is located in the Iława Lakeland in northern Poland, maintains the NNE-SSW course and
80 it is about 720 m long and 165 m wide. The lake occupies 7.5 ha in area, its water surface rises to
81 about 101 m a.s.l. and the maximum depth is just over 2 m. The lake Mlynek is surrounded by a
82 morainic plateau at 120-130 m a.s.l and is surrounded by dense forest (Fig. 1). A large part of the
83 Iława Lake District is covered with forest, whereas meadows and synanthropic communities which
84 have a smaller share. ~~The forest covers 41.5% of the area.~~ Among the habitats, a highly-productive
85 mixed coniferous forest prevails. The basic components of the Iława forest are pine, oak, beech,
86 alder, birch, in smaller amounts there are spruce, larch, ash, hornbeam, maple and linden. Currently,
87 the Lakeland is characterized by a transitional climate that shapes the influence of the continental
88 and maritime climate circulation. The vegetation period lasts about 206 days, and the snow cover
89 remains for 70-90 days. Average temperature values range from approximately -4.0°C in February
90 to above 17.0°C in July, maximum from -1.0° to 22.0°C, minimum from approximately -7.0° to
91 12.0°C. Due to the greater proportion of the polar air masses and a large number of natural water
92 reservoirs, air humidity is relatively high, ranging from 72% to 89%. The precipitation sums from
93 500 to 550 mm a year. Throughout the year, SW winds predominate. The westerly winds are
94 stronger in winter. The highest wind speeds are recorded in the winter months (from 2 to 4 m/s)
95 and the lowest in the summer (from 2.0 to 3.0 m/s) (Jutrzenka-Trzebiatowski and Polakowski,

96 1997; Stopa-Boryczka et al., 2013). It is important to note, that from the north, a small stream flows
97 into the Lake Młynek, which is active in winter and dries up almost completely in summer (Fig. 1:
98 D). Most probably it is an effect of irrigation works related to the construction of the mill in the
99 15th century, somewhere in the vicinity of the medieval stronghold located on the shore of the lake
100 (Semrau, 1935, Bińka et al., 2020).

101

102 3. Material and methods

103 3.1. Bathymetry

104 Determination of lake bathymetry and thickness of bottom sediments are extremely
105 important in palaeolimnological research to help locate coring sites. This can be achieved through
106 the use of GPR sounding (Lin et al., 2009; Sambuelli et al., 2009; Sambuelli and Silvia, 2012). In
107 Poland winter is a particularly convenient season when ice cover of a lake makes GPR profiling
108 much easier and improves access and speed of data collection (Hunter et al., 2003). Measurements
109 along and across the lake were carried out in 2017, directly on a lake ice and a snow cover (Fig. 2).
110 We used the radar system ProEx of the Malå Geoscience. A radar pulse was generated at a regular
111 distance interval of 0.02 m (900 samples were recorded from a single pulse). The time window of
112 recording was between 250 and 300 ns. Prospection was done with use of a shielded monostatic
113 antenna with 250 MHz nominal frequency of the electromagnetic wave.

114

115 3.2. Coring and sampling

116 Based on the results of the GPR 4 drillings were done at ca 2 m water depth (Fig. 3) to
117 collect cores according to Givelet et al. (2004) collecting protocol. Sediment cores were film-
118 wrapped in 1 m plastic tubes and transported to the laboratory. These cores (M1-4) were then
119 subjected to magnetic susceptibility measurements which enabled to select M1, the longest and
120 most continuous core, to carry out detailed analysis. Samples from the 3.5 m long core M-1
121 (geographic coordinates: 53.82486 N, 19.72419 E) were sub-sampled at 5 cm interval and used for
122 multi-proxy laboratory analyses.

123

124 3.3. Age-depth model

125 Radiocarbon dating was performed on 4 bulk samples from the core M-1, collected either
126 from organic-rich gyttja or gyttja with dispersed organic matter (Table 1). The organic matter

127 seems to have been derived both from aquatic and terrestrial sources. AMS dating was done in the
 128 Poznań Radiocarbon Laboratory in Poland, where ^{14}C measurements were performed in **graphite**
 129 **targets** (Goslar et al., 2004). Construction of proper and correct age-depth model required an
 130 assessment of several agents that could disturb constant accumulation of bottom deposits of the
 131 Lake Młynek. Disturbances could result both from sedimentary and post-sedimentary processes
 132 (varied rate of deposition and compaction, impact of bioturbation). The varied influx of material
 133 delivered to the lake from the adjacent area is a very important factor of disturbance. Therefore, a
 134 Bayesian age-depth ~~routine mode~~ was chosen as it takes into account the sedimentation rate and its
 135 variability (Blaauw and Christen, 2005, 2011; Blaauw et al., 2007) (Fig. 4). The model was based
 136 on default settings, except for section thickness which was set at 0.05 cm given the length of this
 137 core. The Bacon model uses the IntCal3 curve (Reimer et al., 2013) to calibrate the radiocarbon
 138 data.

139

140 Table 1. List of radiocarbon determinations.

No.	Depth in m	Lab. reference	^{14}C yr. BP	Age calibrated 95% probability	Material dated
1	0.95-1.00	S/JW 1/2015/A	435 ± 30	1418 – 1494 AD	Bulk of gyttja
2	1.65-1.70	S/JW 1/2015/B	1015 ± 30	971 – 1048 AD	Bulk of gyttja
3	2.40-2.45	S/JW 1/2015/C	1730 ± 30	236 – 386 AD	Bulk of gyttja
4	3.45-3.50	S/JW 1/2015/D	2275 ± 30	401 – 351 BC	Bulk of gyttja

141

142 3.4. Palaeobotanical analysis

143

144 3.4.1 Pollen

145 The core M-1 was sampled every 5 cm for pollen analysis. 70 samples (ca. 10 g each) were
 146 treated with 5% HCl, boiled in 5% KOH and hot 30% HF. They were washed with 15% HCl and
 147 treated by the standard Erdtman's acetolysis. In each sample about 1000 pollen grains were counted
 148 using an optical microscope at 400x magnification.

149

150 3.4.2 Diatom and Chrysophyte cysts analysis

151 70 samples were prepared for the analysis of diatoms and chrysophyte cysts. They were
 152 extracted from 1 g of dry sediment of each sample using the disintegration method in HCl and
 153 H_2O_2 , according to the technique proposed by Zalat and Servant-Vildary (2007). For slide

154 preparation, 0.1 ml of the final suspension was dried on coverslips and then mounted onto slides
155 using Naphrax. Diatoms were identified to species level using a Leica photomicroscope with a
156 digital camera and equipped with differential interference contrast (DIC) optics at 1000x
157 magnification with oil immersion. Identification and ecological information of the diatom species
158 were based primarily upon the published literature (e.g. Kilham et al., 1986; Douglas and Smol,
159 1999; Witkowski et al., 2000; Hofmann et al., 2011). Recent taxonomic advances split many
160 diatom taxa of the former genus *Fragilaria sensu lato* into several new genera, including
161 *Fragilaria*, *Pseudostaurosira*, *Staurosira* and *Staurosirella* spp. (Williams and Round, 1987);
162 these new names herein collectively referred to as *Fragilaria sensu lato*. Chrysophyte cysts were
163 described and enumerated following Duff et al. (1995, 1997), Pla (2001) and Wilkinson et al.
164 (2002). Preliminary results of the diatom studies based on the core M-1 were already published by
165 Zalat et al. (2018).

166

167 3.5. Geochemical analysis

168 ICP-OES spectrometer was used for determination of basic (Al, Ca, Mg, Na, K, Fe, P) and
169 trace elements (As, Cd, Mn, Th, Ti, U, V, Zn) ~~in the analysed samples~~. Powdered samples were
170 mineralized in a closed microwave Anton Paar Multiwave PRO reaction system. Mineralization
171 procedure was based on the procedure of Lacort & Camarero. Characteristics of lake sediments
172 ~~was~~ done with the extraction method of elements soluble in aquaregia (according to European
173 Standard CEN/TC 308/WG 1/TG 1, slightly modified). Dry samples of about 0.2 g weight were
174 transferred to the PTFE vessel and HNO₃, and HCL Merck Tracepur® was added. The vessels
175 were placed in a rotor and loaded to a microwave. Finally, the samples were analysed in the Spectro
176 Blue ICP OES spectrometer at Regional Research Centre for Environment, Agricultural and
177 Innovative Technologies, Pope John II State School of Higher Education in Biała Podlaska. Berndt
178 Kraft Spectro Genesis ICAL solution and VHG SM68-1-500 Element Multi Standard 1 in 5%
179 HNO₃ were used.

180

181 3.6. Total organic carbon (TOC)

182 Analyses were done after sample acidification to remove carbonates in the SHIMADZU
183 SSM 5000A analyser with a solid sample combustion unit. The method was the catalytically aided
184 combustion oxidation at 900°C with pre-acidification and oven temperature 200°C. A measuring

185 range TC was 0.1 mg to 30 mg carbon. Sample amount was 1 g and aqueous content <0.5 g.
186 Repeatability: S.D. $\pm 1\%$ of full scale range (www.ssi.shimadzu.com/products/toc-analyzers/ssm-
187 5000a).

188

189 3.7. *Magnetic susceptibility (MS)*

190 The cores from the Lake Młynek were subjected to MS measurements using SM-30
191 magnetic susceptibility meter (ZH Instruments). Due to very high sensitivity (1×10^{-7} SI units) this
192 device was provided with 8 kHz LC oscillator and its pick-up coil sensor was large enough to
193 measure sufficiently high volume of sediments with very low MS. The measurements were done
194 at every 5 cm along each core (M1-4).

195

196 3.8. *SEM microscopic analysis*

197 All selected samples were analysed using a scanning electron microscope (SEM) HITACHI
198 TM3000 with an energy dispersive spectrometer (EDS) SWIFT ED 3000 Oxford Instruments. The
199 samples were not covered with any conductive material. Magnification range was x 20 to x 30 000.
200 This method was used to perform basic microscopic observations of samples of the core M-1 with
201 point determination of their chemical composition of major elements.

202

203 4. Results

204 4.1. *Archaeological records*

205 Archaeological data were collected from the stronghold Janiki Wielkie, located on a hill at
206 the north-eastern shore of the Młynek Lake. During the archaeological research carried out in 2013
207 and 2016, a total of 143 stratigraphic units were distinguished on this site, which were divided into
208 seven main settlement phases: phase I – early Iron Age, phase II –stronghold abandoned after the
209 early Iron Age, phase III – early Middle Ages, phase IV – stronghold abandoned in the early Middle
210 Ages, phase V – settlement activity on the stronghold in the 11th-13th century, phase VI –
211 stronghold definitely abandoned in the 14th century (Rabiega et al., 2017, Nitychoruk and Welc,
212 2017).

213 4.2 *Bathymetry*

214 A georadar transect across the lake reflects both its bathymetry and lithologic variety of its
215 bottom (Figs 2 - 3). The superficial layer of the transect is represented by lake ice, ca 25 cm thick

216 and although it is almost not visible on radar images due to its thickness being smaller than a
217 vertical resolution of measurements, beneath there are abundant horizontal multiple reflections of
218 energy from the bottom of the ice. Two narrow and vertical zones with small diffraction hyperboles
219 at 23 and 29 m of the transect indicate upward deformation of bottom sediments at the location
220 sites of the sounding core and the core M-1 (Fig. 2a). The top of the underlying mineral deposits
221 (so-called hard bottom) is indicated as a distinct downward-deflected reflection surface (Fig. 2b).
222 In a central part of the lake it occurs at 2.6 m depth (two-way travel time 290 ns) and indicates the
223 top of the Holocene organic sediments. Unfortunately, beneath there is a signal-absorption zone
224 (Fig. 2d), resulting from the fact that most sediments are composed of fine-grained organic material
225 (gyttja). However, thickness of this layer was determined by drillings to about 5 m. A relief of the
226 lake bottom in the GPR image reflects a cross-section of a buried glacial tunnel valley that was
227 eroded mainly in sandy and sandy-gravel deposits. Close to the lake shore (0 to 20 m in the
228 northwest and 110 to 140 m in the southeast), there are numerous oblique and chaotically parallel
229 reflection surfaces dipping towards the channel axis. They reflect bedding of the Pleistocene sandy-
230 gravel series that partly filled a subglacial channel (Fig. 2c).

231 4.3. *Age-depth model*

232 The age-depth model of the core M-1 from the Lake Młynek is shown in Fig. 4. Grey
233 stippled lines show 95% confidence intervals and the red curve shows the ‘best’ model based on
234 the weighted mean age for each depth. Good runs of a stationary distribution are shown in the upper
235 left panel, green curves and grey histograms in the upper middle panel present distributions for the
236 sediment accumulation rate. The main bottom panel shows the calibrated ^{14}C dates (transparent
237 blue) and the age-depth model (darker grey areas) which are indicating calendar ages.

238 4.4. *Lithology*

239 Bottom deposits of the Młynek Lake are organic-rich. The core M-1 is composed of grey-
240 brown gyttja at 1.8 – 3.6 m depth (Fig. 5). At 1.45 – 1.80 m depth there is grey-brown gyttja-
241 detritus and at 1.10 – 1.45 m depth algal gyttja is recorded. The uppermost part of the core is
242 composed of grey-brown (depth 0.4 – 1.1 m) and detritus gyttja (0.0 – 0.4 m).

243 4.5 *Sedimentation rate*

244 The sedimentation rate was calculated based on the age-depth model. Results reflect quite
245 a stable sedimentary environment with a general rate of 1.5 mm a year. The rate is stable at 3.40 –
246 1.77 m depth and equal ca 1.5 mm a year. It drops to 1 mm, then rises to 1.3 – 1.8 mm a year at

247 1.77 – 0.30 m. At 0.0 – 0.30 m the sedimentary rate is the highest and equal ca 3 mm a year (Fig.
248 5).

249 4.6. *Magnetic susceptibility and total organic carbon*

250 MS is highly dependent on lithology and grain size of deposits (Dearing, 1994; Sandgren
251 and Snowball, 2001). It reflects presence and size of ferromagnetic particles in a sample (Verosub
252 and Roberts, 1995). Increased content of ferromagnetic minerals such as magnetite, Fe-Ti oxides
253 or pyrrhotite generates higher MS whereas biotite, pyrite, carbonates and organics result in its lower
254 values. Total volume of magnetic minerals in lake sediments reflects mostly climatic changes in a
255 catchment (Bloemdal and deMenocal, 1989; Snowball, 1993; Peck et al., 1994).

256 MS in the core M-1 is varied but due to organic character of the sediments, its values are
257 relatively low, from 0.002 to 0.034×10^{-7} units SI. In grey-brown gyttja with organic matter at 3.50
258 – 2.58 m depth, MS rises and drops in turn from 0.01 to 0.02×10^{-7} SI. MS drops from 2.60 m depth,
259 reaching a minimum at 1.63 m. Higher up, MS rises again, with the highest value at 1.35 m, then
260 there is a minimum at 1.05 m and the next maximum at 0.69 m depth (Fig. 6).

261 MS is generally low in biogenic sediments as gyttja, which is composed mainly of
262 microfossil skeletons e.g. diatoms and radiolarians (Thompson and Oldfield, 1986). In the Lake
263 Młynek there is an apparent negative relationship between TOC and MS. Several intervals show
264 both higher percentages of TOC and lower MS values. At 1.40 m depth, TOC suddenly drops,
265 probably due to deforestation and then, MS significantly rises due to increasing input of terrestrial
266 (non-organic) material to the lake. Such coincidence clearly indicates that TOC is both of
267 autochthonous and allochthonous derivation (Fig. 6).

268 Changes of MS in the Lake Młynek sediments record most probably an input of clay into
269 the lake and diagenetic conditions in bottom sediments. Iron oxides are presumably of detrital
270 origin and were delivered to the basin through deep valleys incised at the north-western shore.
271 Concentration of ferromagnetic minerals is connected with periodical intensive soil erosion around
272 the lake. Their higher content depends also on diagenetic processes in bottom sediments. Oxidation
273 of organic matter in anoxic conditions (by iron-oxide-reducing bacteria) results usually in increased
274 content of ferromagnetic particles (small particles are removed first). In opposite, oxygenation by
275 heavy floods stops this process and small magnetic particles are preserved (Jelinowska et al., 1997).

276 4.7. *Water-soluble ions*

277 Various factors influence distribution and accumulation of geochemical elements in lake
278 sediments. Most important are texture, mineral composition, oxidation/reduction state,
279 absorption/desorption and physical transportation processes (Ma et al., 2016). Curves of
280 representative elements are generally used to characterize sedimentary environments. Most
281 analysed elements do not indicate any clear trend with depth in the Lake Młynek. The curves of S
282 and TOC show significant rises at 2.0 – 1.4 m depth that are slightly correlated with decreased
283 contents of Al, Fe, K, Ca, Mg and MS (Fig. 6).

284 Sulphur content is correlated with existence of iron sulphides. In the studied core, Fe is
285 positively correlated with Al and Ti (Fig. 7). Fe-Ti oxides are noted in SEM EDS analysis. They
286 are resistant to surface weathering and carry trace elements (Bauer and Velde, 2014). At ca. 3 m,
287 high frequency peaks of Al, K, Ca, Na, Mg, Fe and S occur (Fig. 6). Such occasional high intensity
288 events leave a stronger geochemical imprint, because of sedimentation in shallow water (Ivanić et
289 al., 2018). The highest contents of detrital elements like Al, K, Ca and Mg are to be associated with
290 sudden delivery of clastic material to the lake e.g. during increasing flood or rainfall (Wirth, et al.,
291 2013). Especially Al is extremely immobile, that is why it should be regarded as a typical lithogenic
292 element (Price et al., 1999). Additionally, Al is a major constituent of soils and other sediments as
293 a structural element of clays. It has a strong positive correlation with many major elements (Fig.
294 7). The association between Al and other elements can be therefore used as the basis to compare
295 natural elemental contents in sediments and soils. Ca is well correlated with Al, originated mainly
296 from terrigenous bicarbonate inputs and was deposited in a lake as a solid carbonate (Miko et al.,
297 2003). Calcium is evidently more easily removed in solution from a mineral material and it is
298 highly concentrated in highly erosional periods (Mackereth, 1965).

299 The Fe/Ca ratio is considered as a eutrophication proxy. The highest ratio points out to low
300 oxygenation, eutrophic or dystrophic reservoirs (i.e. Kraska and Piotrowicz, 2000; Holmes and De
301 Decker, 2012), whereas the low Fe/Ca ratio in bottom sediments indicates oligotrophic character
302 of a lake. In the studied core sediments, Fe/Ca ratio varies from 0.808 (depth 3.05 m) to 3.677 (1.2
303 m). The ratio is low, indicating oligotrophic conditions in bottom sediments which gives conflicting
304 results with other data. The Fe/Ca ratio can be disturbed by detrital input to the lake (Fig. 6). The
305 dysaerobic conditions in the lake are confirmed with Th/U ratios (0.03 – 0.41) which are lower
306 than the critical value of 2 as indicated by Myers and Wignall (1987) and Wignall (1994). The ratio
307 of total Fe to total P ranges from 13.91 (1.6 m) to 30.82 (0.55 m). The values are typical for other

308 lakes in northern Poland, which vary from 3 to 180 according to Bojakowska (2016). The release
309 of P follows in reducing conditions. According to Ahlgren et al. (2011) is even up to ten times
310 greater than in aerobic conditions. However, there is a poor correlation with other redox proxies
311 i.e. Th/U (R=0.08). It can be caused by presence of Al which forms Al(OH)₃. In such systems even
312 though the redox state favours release of P from iron minerals, the P is immobilized by binding
313 with hydroxides. Thus, the presence of Al(OH)₃ can stop release of P even in an anoxic
314 hypolimnion (Hupfer and Lewandowski, 2008). It can be a case in the studied sediments as Al
315 shows positive correlation with P content (R=0.49). Except for Fe/Ca, all counted ratios point out
316 to anoxic conditions in all studied samples which is typical to eutrophic lakes. Nevertheless, as all
317 proxies are characterized by extreme values at the 3.05 m depth, they seem to depend on external
318 load of terrigenous material. It is confirmed with very good positive correlation between Fe and Al
319 (0.95), Fe and Ti (0.64) Mn and Al (0.46) or Mn and Ti (0.78).

320 4.8. Diatoms and chrysophyte cysts

321 Studies of the Lake Mlynek bottom sediments revealed presence of more than 200 diatom
322 taxa belonging to 54 genera (Zalat et al., 2018) (Fig. 8). Diatoms were generally abundant and well
323 to moderately preserved in most samples, although with admixture of mechanically broken valves,
324 especially in the topmost part of the core. Results of the diatom analysis and relative abundance of
325 the most dominant taxa enabled subdivision of the M-1 core section into 11 diatom assemblage
326 zones (Fig. 8) that reflected six phases of lake development (Zalat et al., 2018). Moreover, changes
327 in chrysophyte cysts distributions along with variation in diatom composition could be related to
328 changes in pH, climate and trophic status. Stomatocysts can be used as the index of lake-level
329 changes, habitat availability, metal concentrations and salinity. The periphytic diatom species
330 dominate the planktonic ones throughout the core. A high proportion of periphyton to plankton
331 assemblages was reported as indicative for a long-lasting ice-cover (Karst-Riddoch et al., 2005)
332 whereas a shift from benthic to planktonic diatom taxa is considered for an ecological indicator
333 that is generally interpreted in high-altitude lakes as record of shorter winter and increased in
334 temperatures. Common occurrence of benthic forms represented by *Staurosira venter/Staurosirella*
335 *pinnata* diatom assemblage indicates circumneutral to slightly alkaline shallow water with
336 lowering lake levels and prolonged ice cover. However, *Aulacoseira* is the most dominant
337 planktonic genus followed by *Cyclotella* and low frequency of *Cyclostephanos*.

338 Diatom preservation in the upper part of the core (depth 1.40 – 0.15 m) is moderate to
 339 relatively poor and the recognized assemblage was represented by the occurrence of some
 340 dissolved and teratological diatoms valves, in particular the topmost part of the core section (0.30-
 341 0.15 m) (Zalat et al., 2018).

342 4.9. Pollen

343 Based on percentage contents of main trees and terrestrial herbs five local pollen
 344 assemblage zones (LPAZ M1-M5) were established in the pollen sequence of the Lake Młynek.
 345 The Pollen levels were determined on the basis of changes in the percentage of individual taxa,
 346 confirmed by a cluster analysis (Fig. 9):

Zone	Depth [m]	Main features of pollen spectra
LPAZ M-1	345÷315 cm	Pollen grains of <i>Carpinus</i> reached 33.5% and <i>Alnus</i> 25%, <i>Pinus</i> and <i>Betula</i> are <20%. A top border of this zone is indicated by decline of <i>Carpinus</i> .
LPAZ M-2	315÷265 cm	The share of <i>Carpinus</i> drops significantly (<10%), contents of <i>Betula</i> , <i>Quercus</i> and <i>Corylus</i> are slightly raised. The percentages of Gramineae significantly increased up to 7.5%. There are continuous curves of <i>Cannabis/Humulus</i> , Chenopodiaceae, <i>Plantago lanceolata</i> , <i>Rumex acetosella</i> and <i>Secale cereale</i> and a top boundary is indicated by decline of Gramineae.
LPAZ M-3	195÷265 cm	At the beginning the curve of <i>Betula</i> raises to 24% but then drops <10%. The share of <i>Carpinus</i> and <i>Fagus</i> rises to 19% and 27%, respectively. Content of Gramineae decreased <2% and the curves of <i>Secale cereale</i> , <i>Plantago lanceolata</i> and <i>Rumex acetosella</i> disappear. There are only single pollen grains of Chenopodiaceae and <i>Cannabis/Humulus</i> . A top boundary is indicated by a rise of Gramineae.
LPAZ M-4	195÷145 cm	Content of <i>Fagus</i> pollen began gradually decrease. The share of pollen grains of <i>Betula</i> increases and becomes stable at 22-27%. Content of the Gramineae pollen grains increases again to 7%. Curves of <i>Cannabis/Humulus</i> , <i>Plantago lanceolata</i> , <i>Rumex acetosella</i> and <i>Secale</i> raise and a top boundary is marked by a rapid rise of <i>Cannabis/Humulus</i> .
LPAZ M-5	145÷15 cm	Curves of main deciduous trees decline: <i>Carpinus</i> <9%, <i>Fagus</i> <5%, <i>Quercus</i> <5%, <i>Alnus</i> <15%, <i>Betula</i> <14% and content of <i>Pinus</i> increases to 40%. There is significant rise of Gramineae up to 15%. Percentages of <i>Cannabis/Humulus</i> reached absolute maxima (25%) but close to middle part of this zone their strong decline is observed (below 2-3%). The continuous curves of <i>Cerealia undiff.</i> , <i>Centaurea cyanus</i> , <i>Plantago lanceolata</i> , <i>Rumex acetosella</i> , <i>Rumex acetosella</i> appeared, and single pollen grains of <i>Polygonum dumetorum</i> , <i>Polygonum aviculare</i> and <i>Urtica</i> were present.

347

348 5. Discussion

349 5.1. Phases of the lake Mlynek development during last 2400 years

350 Based on results of performed analysis and archaeological data, five main environmental
351 phases of the Lake Mlynek development were distinguished (Fig. 10):

352 5.1. Phase 1: ~2300 – 2100 cal. BP (ca. 4 – 2/1 c. BC). Depth: 3.45 – 3.15 m

353 This phase is recorded in LPAZ M-1 which represents closed forest communities dominated
354 by hornbeam and alder, which colonized marshland near lake shores. Plants of open spaces are
355 only rarely noted as well as indicators of anthropogenic activity (e.g. *Plantago lanceolata*). There
356 is therefore every reason to conclude that vegetation at that time was relatively natural and not
357 disturbed. A diatom assemblage at the beginning of the diatom subzone DZ1 at 3.45 – 3.40 m depth
358 (Fig. 8) indicates a shallow and slightly alkaline lake, followed (3.35 – 3.15 m) by a rising lake
359 level. Common occurrence and domination of *A. granulata* suggests a high trophic status of slightly
360 alkaline freshwater environment with high silica concentration (Zalat et al., 2018). MS is high and
361 corresponds to high content of Fe, Ti and Al, indicating increased influx of terrigenous material,
362 presumably activated by intensive rainfalls.

363 5.2. Phase 2: 2100 – 1830 cal. BP (ca. 1 c. BC – 2 c. AD). Depth: 3.15 – 2.65 m

364 A vicinity of the lake began to change and major changes in the environment were caused
365 by significant human impact. This phase corresponds with the LPAZ M2, characterized by
366 reduction and fragmentation of the hornbeam-dominated forest. Birch, pine and hazel expanded
367 under better lighting conditions in a partly open forest whereas higher oak values resulted from
368 higher production of pollen. Mid-forest pastures occupied rather small-scale open areas, as can be
369 notified from higher percentages of *Plantago lanceolata* and other herbaceous plants, e.g.
370 Gramineae, *Artemisia* and *Rumex acetosa/acetosella*. Cultivated plants as *Cannabis t.* and *Secale*
371 are rare, however their occurrence is entirely consistent with other indicators present during this
372 phase. Human occupation is attested by presence of *Cannabis/Humulus*, *Plantago lanceolata*,
373 *Rumex acetosella*, *Secale* and cereals undiff. This phase is commonly noted and similarly
374 expressed in numerous palynological sequences in neighbouring areas (see for example
375 Noryśkiewicz, 1982, 1987, 2013; Bińka et al., 1991; Ralska-Jasiewiczowa et al., 1998). Pollen data
376 indicate that societies of that time cultivated rye and probably hemp. It is the oldest settlement
377 phase at Janiki Wielkie stronghold and corresponds to a termination of the La Tène and the early
378 Roman period. Human communities in the vicinity of the lake can be connected with settlements

379 of the East-Baltic Kurgan Culture (Rabiega et al., 2017). During this phase, planktonic diatoms
380 were replaced by benthic taxa, accompanied by *Gyrosigma acuminatum* indicating a lower lake
381 level and dominance of mesotrophic alkaline freshwater environment. The lower stands were
382 interrupted by a short rise of water level at 2.90 – 2.85 m (Zalat et al., 2018). During this phase
383 climatic conditions were still similar to ones in the previous phase, but it was drier what is reflected
384 by shallowing of the lake. Described phase can be correlated with the so-called Roman Climatic
385 Optimum (see McCormick et al., 2012).

386 5.3. Phase 3: 1830 – 1150 cal. BP (ca. 2 – 9 c. AD). Depth: 2.65 – 1.95 m

387 This phase is recorded by the LPAZ M3 when a forest restoration occurred. Absence of human
388 indicator plants suggest that the settlement in the catchment was abandoned. There are also no traces
389 of human activity nearby (Rabiega et al., 2017). Reduction of human impact and human-generated
390 semi-open habitats, allowed for a short-term expansion of birch into empty, open areas, and then
391 replaced by hornbeam that rebuilt its position to the level as in the LPAZ M1. Also, elm expanded
392 again in a riparian forest. This restoration of the natural forest was followed by abrupt expansion of
393 beech in the second half of the LPAZ M3. The area of open herbaceous plants communities,
394 previously widespread, was limited.

395 Abundant planktonic diatoms occurred in the lake (Fig. 8) what indicates its deepening,
396 enhanced thermal stratification, reduced mixing and increased thermal stability (Zalat et al., 2018).
397 Intensified development of a vegetation cover and higher lake water level are indicated by
398 geochemical indices. A gradual drop of MS corresponds with decreased content of detrital elements
399 as Fe, Ti, Al and K, accompanied by gradual increase of TOC and the proxy ratio Fe/Ca. Lower MS
400 and content of Al (acting as a major constituent of soils) accompanied by the higher TOC suggest
401 limited erosion, in spite of gradually higher precipitation in the lake catchment and therefore, a rise
402 of its water level (Fig. 6). The climate in this phase has become more humid. Increased rainfall and
403 decreased evaporation are reflected in lake sedimentation as the lake got deeper, resulting in reduced
404 deposition, lower productivity and greater stability. This phase could be associated with a global
405 cooling of the Bond Event 1 (1.5 ka BP) (Bond et al., 1997; Welc, 2019).

406 5.4 Phase 4: 1150 – 780 cal. BP (ca. 9 – 13 c. AD). Depth: 1.95-1.45 m.

407 This phase is correlated with the LPAZ M4 and was divided into two subphases 4a and 4b
408 (Fig. 10). The subphase 4a marks the onset of another settlement phase, resulting in forest clearing,
409 similarly as in the LPAZ M2. Disturbances took place firstly in a beech forest and less in a

410 hornbeam-dominated one. The anthropogenic activity was only slightly smaller than in the LPAZ
411 M2 ~~and is~~ reflected by presence of Gramineae, *Artemisia*, *Cannabis/Humulus*, *Plantago*
412 *lanceolata*, *Rumex acetosella*, *Secale* and cerealia undiff. Diatom assemblages prove deepening of
413 the lake (Zalat et al., 2018) as indicated by abundance of *Aulacoseira* associated with *Puncticulata*
414 *radiosa* in the upper part of the diatom zone 5 at 1.85 – 1.70 m depth. The diatom assemblage
415 suggests a rising lake level, higher trophic and stronger turbulent mixing conditions. Moreover, the
416 greatest reduction of abundant *Fragilaria sensu lato* accompanied by abundant *A. granulata*, could
417 resulted from forest clearing around the lake. Higher TOC corresponds with lower content of
418 detrital material (Fe, Ti, Al and K) and lower MS, and it can reflect a progressing humidity (Fig.
419 6). This phase can be correlated with the Migration Period and the early Middle Ages. A wooden-
420 loamy defence rampart was raised at the end of the phase in a settlement close to the lake
421 (archaeological phase III), after removal of a natural soil developed during abandonment of the
422 site in the early Roman Period. After a short period, this stronghold was destroyed. A charcoal
423 from a fired wall that represents destruction at the end of the archaeological phase IIIA, was dated
424 at 1245 ± 25 cal. BP i.e. 682-870 AD (95.4% probability) and 1090 ± 30 cal. BP i.e. 892-1014 AD
425 (95.4% probability) (Rabiega et al., 2017).

426 Human impact declines during the subphase 4b (1.70 – 1.45 m depth) onwards. At this time
427 birch and less intensively poplar occupied temporarily abandoned open areas, especially toward
428 the end of the zone, when a human activity was less intensive. Alder became more abundant,
429 probably expanding into exposed marginal areas of the lake. The subphase 4b, corresponds to the
430 diatom zone 6 which is characterized by abundant benthic *Fragilaria sensu lato* with sporadic
431 occurrence of planktonic taxa. A diatom assemblage reflects lowering water level and slight
432 alkaline freshwater, lower nutrient concentrations and low silica content (Zalat et al., 2018). In the
433 stronghold at the lake shore, the next phase of human activity took place at the end of the 11th
434 century AD when a new rampart was raised. Wooden constructions were also built, traces of which
435 were excavated in the gate passage. The settlement was finally abandoned presumably in the first
436 half of the 13th century and then, its ramparts were strongly eroded, with their material moving
437 towards a yard and the moat (Rabiega et al., 2017). Previous climatic conditions continue in this
438 phase. The subphase 4b is characterized by ~~climate change towards warming~~, which confirms
439 gradual shallowing of the lake and increased rate of sedimentation. Human impact on the
440 environment in this subphase is already so great that reconstruction of a climate change is not

441 clear. There is no doubt, however, that this is a warm period, which should be correlated with the
442 Medieval Warm Period (Mann et al., 2009).

443 5.5. Phase 5: 780 – 0 cal. BP (13 c. AD – present time). Depth: 1.45 - 0 m

444 This phase starts about 1200 AD and is connected with the early Modern Period of the
445 Middle Ages. Cultivation and treatment of hemp has been ~~terminated but cultivation of~~ cereals and
446 presence of synanthropic plants indicate human activity ~~in~~ near the lake. The water level is not high
447 and slightly changes. There is a drop of TOC and rise of MS caused by increasing input of terrestrial
448 material at 1.4 m depth, resulting presumably from human deforestation. The small watercourse
449 which enters the Lake from the north – east appeared most probably during this phase and had the
450 strong impact on the its water environment (see, Bińka et al, 2020). ~~How it was mentioned,~~ in 15
451 c. AD a mill was built near the lake using water from ~~the newly created~~ stream. Damming of the
452 water in the mill reservoir probably contributed to periodical blooms of dinoflagellate populations
453 in the Lake Młynek. Major blooms of *Tetraedron* which usually preceding blooms of the
454 dinoflagellate, ~~was~~ most probably main factor that contributed to the decline of settlement on the
455 stronghold near the shore of the lake (Bińka et al, 2020). ~~Described~~ zone is also characterized by
456 increased precipitation which is reflected by significantly more intensive terrestrial inflow to the
457 lake and is confirmed by quasi-linear correlation of MS with contents of Fe and Ti in sediments
458 (Fig. 6). The modern lake is shallow (2-3 m) and gradually overgrowing. Summing up, the phase
459 5 is marked by intensive human activity around the lake and therefore, most environmental and
460 climate changes are obliterated.

461 5.2 Development of the Lake Młynek on regional bedground

462 The above scenario seems to be confirmed by earlier palaeoenvironmental research carried
463 out in the south-western part of the Warmia-Masuria Lake District (Kupryjanowicz, 2008;
464 Kołaczek et al., 2013). Previous studies of the lake and paleolake sediments in this region were
465 based mainly on pollen analysis and enables to compare the Lake Młynek record with other
466 sequences. The closest site Woryty (Pawlikowski et al., 1982, Noryśkiewicz and Ralska-
467 Jasiewiczowa, 1989, Ralska-Jasiewiczowa and Latałowa, 1996), just 35 km to the east, is a
468 reference one. Palaeoenvironmental records delivered by the Lake Młynek core are very similar to
469 the Woryty palynological succession with distinctive human impact during the Roman Period and
470 the Medieval Ages. More detailed comparison is impossible, because of low resolution of the
471 pollen spectrum obtained in Woryty. The second site is the Lake Drużno, located in the Vistula

472 Delta, 35 km to the north of Młynek Lake (Zachowicz et al., 1982; Zachowicz and Kępińska, 1987;
473 Miotk-Szpiganowicz et al., 2008). Unfortunately, low resolution and lack of the age-depth model
474 from this lake makes comparison also very difficult. Despite of this and habitat differences between
475 the Lake Družno and the Lake Młynek, a pollen records obtained in both sites are very similar and
476 comprises human indicators during the Roman Period and human impact during the Medieval time.
477 Differences in natural vegetation are local and especially exposed in higher share of alder in a
478 pollen diagram from the Lake Družno, most probably caused by wet habitats in the Vistula Delta.
479 The pollen spectrum from the Lake Łańskie (Madeja, 2013), located 55 km to the south-east from
480 the Lake Młynek, shows higher content of pine and lower share of beech than in the case of the
481 Lake Młynek. Such divergences are probably not only due to different location and environmental
482 conditions in the lake vicinity but also depend on different size of these lakes. The Lake Młynek is
483 a very small (0.7 km²) mid-forest basin, whereas the Lake Łańskie is over 10 km² large and contains
484 mostly a regional pollen record. Based on periodical appearances of human plant indicators and
485 archaeological data between 300 BC and 800 AD, three human phases of West Baltic Barrow,
486 Wielbark and Prussian cultures were distinguished (Madeja, 2013). In the pollen diagram from the
487 Lake Młynek (phase 2), the first culture is indicated, including termination of the La Tene and the
488 Roman Period. Significant growth of human indicators from the beginning of 11th century is visible
489 in diagrams from both sites. A more local record from the Lake Młynek is marked especially by
490 high content of *Humulus/Cannabis* (to 25%) in 13-15th centuries AD. In the sediments of the Lake
491 Łańskie, hemp occurred discontinuously and was <1%.

492 Numerous pollen data are available from the area adjacent in the south-west in the Brodnica
493 Lake District, including the Strażym Lake (Noryśkiewicz, 1987; Noryśkiewicz and Ralska-
494 Jasiewiczowa, 1989), the Oleczno Lake (Filbrandt-Czaja, 1999; Filbrandt-Czaja et al., 2003) and
495 the Chełmno Lakeland (Noryśkiewicz, 2013). Pollen record from this region suggests settlements
496 during La Tene, Roman and Medieval periods. Pollen record from other sites located to the east of
497 the Lake Młynek indicates differences in a beech content. The *Fagus sylvatica* content changes to
498 the north-east and its significantly high content in the Lake Młynek sediments represents a very
499 local record in a small lake. Decline of *Fagus sylvatica* depend on a continental climate and is
500 noted in pollen diagrams from the lakes: Salet (Szal et al., 2014a), Mikołajki (Ralska-
501 Jasiewiczowa, 1989), Żabińskie (Wacnik et al., 2016) and Wigry (Kupryjanowicz, 2007). A decline
502 of beech is accompanied by a rise of *Picea abies*. A record of human activity in pollen spectra from

503 eastern Poland was noted at many sites. The pollen record from the Lake Młynek are similar to the
504 ones from the Masurian Lakes: Wojnowo, Miłkowskie and Jędzelek, located over 100 km to the
505 east (Wacnik et al., 2014). Recorded episodes of human impact on vegetation during the Roman
506 Period and Medieval time are separated by 500-600 years long intervals without cultivation and
507 with natural reforestation (indicated by strong share of birch which is a pioneer tree). Similar lasting
508 of human withdrawal in the Lake Młynek section began and terminated earlier than recorded in the
509 lakes Wojnowo and Miłkowskie. Another history of human activity is represented in a record from
510 the Lake Sałęt (Szał et al., 2014b). Pollen grains of cultivated and ruderal plants are noted
511 continuously from the early Iron Age to the early Medieval time. In opposite to the pollen record
512 from the lakes Młynek, Wojnowo and Miłkowskie, the suggested constant settlement in the
513 neighbourhood of the Lake Sałęt was interrupted by a single very short decline of human impact
514 at 880-980 AD (Szał et al., 2014a).

515 6. Conclusions

516 6. 1. Based on results of lithological, geochemical, palynological and diatomological analysis,
517 supplemented with archaeological data, five main environmental phases of the Lake Młynek
518 development were distinguished (Fig. 10). Radiocarbon ages enabled detailed chronology whereas
519 pollen data and stratigraphy of the stronghold to the north-east of the lake made correlation of
520 human activity with environmental data possible for the last 2300 years. From the 1 century BC to
521 2nd century AD the forest around the lake was much reduced, what can be associated with pre
522 roman and Roman occupation phase (attested also on the stronghold located close to the lake).
523 From the 2nd to 9th century AD is attested gradual restoration of the forest and decline of human
524 activity along with lake which is deepening due to the advent of more wet climatic conditions. This
525 colder and humid phase corresponded to the Bond 1 Event (1.5 ka BP) cooling episode. Intensive
526 forest clearing around the lake occurred in the 9th – 13th century AD as result of next phase human
527 activity. This period is marked by warming confirmed by a gradual shallowing of the lake (Middle
528 Age Warm Period). In next 5 strong human impact transformed the local landscape, especially
529 construction and activity small mill since 15 c. AD. This caused that possible climate-induced
530 natural environmental changes are not so clear.

531 6. 2. Environmental transformations recorded in bottom lake sediments of the Lake Młynek were
532 highly dependent on human activity and were especially intensive in the Roman and Middle Age
533 periods due to favourable climatic conditions.

534 6. 3. Human colonisation deduced from a pollen record of the Lake Młynek is coincident with
535 archaeological data, including existence of a stronghold and in spite of a local character, it
536 correlates well with data from other, more regionally significant palynological sites.

537 6. 4. Transformations of the Młynek lake reconstructed based on diatom analysis, not only indicate
538 changes of the lake water level and correspond with a human impact but also determine episodes
539 of more humid climate during coolings.

540

541 **Acknowledgments**

542 The research was funded by the National Science Centre in Poland in the project UMO-
543 2016/21/B/ST10/03059: *Correlation of prehistoric and early medieval settlement phases in north*
544 *-east Poland with the changes of the natural environment in the light of lacustrine sediments study.*

545

546 **References**

547 Ahlgren, J., Reitzel, K., De Brabandere, H., Gogoll, A., Rydin, E.: Release of organic P forms from
548 lake sediments, *Water Research*, 45, 565-72. 2011.

549 Bauer, A., Velde, B.: *Geochemistry at the Earth's Surface Movement of Chemical Elements.*
550 Springer – Verlag, Berlin-Heidelberg, 2014.

551 Bińka, K., Cieśla, A., Łącka, B., Madeyska, T., Marciniak, B., Szeroczyńska, K., Więckowski, K.:
552 The development of Błędowo Lake (Central Poland) - A palaeoecological study, *Studia*
553 *Geologica Polonica*, 100, 1-83, 1991.

554 Bińka, K., Welc, F., Nitychoruk, J., Sieradz, D., Lewczuk, A: Unique finds in palynological
555 spectra: acetolyze resistant vegetative forms of freshwater dinofagellate based on the Lake
556 Młynek record from northeastern Poland, *Studia Quaternaria* 37/2, 59-67. 2020.

557 Blaauw, M., Christen, J.A.: Flexible Palaeoclimate Age-Depth Models Using an Autoregressive
558 Gamma Process, *Bayesian Analysis* 6/3, 457–474. 2011

559 Blaauw, M., Christen, J.A., Mauquoy, D., van der Plicht, J., Bennett, K.D.: Testing the timing of
560 radiocarbon dated events between proxy archives, *The Holocene* 17, 283-288, 2007.

561 Bloemdal, J., deMenocal, P.: Evidence for a change in the periodicity of tropical climate cycles at
562 2.4 Myr from whole – core magnetic susceptibility measurements, *Nature* 342, 897- 900, 1989.

563 Bojakowska, I.: Phosphorous in lake sediments of Poland – results of monitoring research,
564 *Limnological Review*, 16, 15-25, 2016.

565 Brauer, A.: Annually laminated lake sediments and their palaeoclimatic relevance. In: The Climate
566 in Historical Times. Towards a Synthesis of Holocene Proxy Data and Climate Models. GKSS
567 School of Environmental Research, 111-129, edited by: Fischer, H., Kumke, T., Lohmann, G.,
568 Flöser, G., Miller, H., von Storch, H., Negendank, J.F.W., Springer Verlag, 2004.

569 Brauer, A., Dulski, P., Mangili, C., Mingram, J., Liu, J.: The potential of varves in high-resolution
570 palaeolimnological studies, *PAGES News* 17 (3), 96-98, 2009.

571 Brenner M., Whitmore T.J., Curtis J.H., Hodell D.A., Schelske, C.L.: Stable isotope ($\delta^{13}\text{C}$ and
572 $\delta^{15}\text{N}$) signatures of sedimented organic matter as indicators of historic lake trophic state, *J.*
573 *Paleolimnol.*, 22, 205-221, 1999.

574 Czymzik, M., Dulski, P., Plessen, B., von Grafenstein, U., Naumann, R., Brauer, A.: A 450-year
575 record of spring-summer flood layers in annually laminated sediments from Lake Ammersee
576 (southern Germany), *Water Resour. Res.*, 46, W11528, 2010.

577 Dearing, J. A.: Environmental magnetic susceptibility: using the Bartington MS2, 1994.

578 Douglas, M.S.V., Smol, J.P.: Freshwater diatoms as indicators of environmental change in the High
579 Arctic, 227–244, in: *The Diatoms: Applications for the Environmental and Earth Sciences*,
580 edited by: Stoermer, E.F., Smol, J.P., Cambridge Univ. Press, Cambridge, 1999.

581 Duff, K.E., Zeeb, B.A., Smol, J.P.: *Atlas of Chrysophycean Cysts*, 2. Kluwer Academic Publishers,
582 Dordrecht-Boston-London, 1995.

583 Duff, K.E., Zeeb, B.A., Smol, J.P.: Chrysophyte cyst biogeographical and ecological distributions:
584 a synthesis, *Journal Biogeography* 24, 791–812, 1997

585 Elbert, J., Grosjean, M., von Gunten, L., Urrutia, R., Fischer, D., Wartenburger, R., Ariztegui, D.,
586 Fujak, M., Hamann, Y.: Quantitative high-resolution winter (JJA) precipitation reconstruction
587 from varved sediments of Lago Plomo 47°S, Patagonian Andes, AD 1530-2001, *Holocene* 22
588 (4), 465-474, 2012.

589 Filbrandt-Czaja A.: Zmiany szaty roślinnej okolic jeziora Oleczno w późnym holocenie pod
590 wpływem czynników naturalnych i antropogenicznych, in: *Studia nad osadnictwem*
591 *średniowiecznym ziemi chełmińskiej* 3, 61–68, edited by: W. Chudziak, Toruń, 1999.

592 Filbrandt-Czaja A., Noryśkiewicz B., Piernik A.: Intensification gradient of settlement processes
593 in pollen diagrams from Dobrzyńsko-Olsztyńskie Lake District, *Ecol. Quest.*, 3, 125-137, 2003.

594 Snowball, I.: Mineral magnetic properties of Holocene lake sediments and soils from the Karsa
595 Valley, Lappland, Sweden, and their relevance to paleoenvironmental reconstruction, *Terra*
596 *Nova* 5, 258-270, 1993.

597 Francus, P., von Suchodoletz, H., Dietze, M., Donner, R.V., Bouchard, F., Roy, A.-J., Fagot, M.,
598 Verschuren, D., Kröopelin, S.: Varved sediments of Lake Yoa (Ounianga Kebir, Chad) reveal
599 progressive drying of the Sahara during the last 6100 years, *Sedimentology* 60 (4), 911-934,
600 2013.

601 Gałązka, D.: Szczegółowa mapa geologiczna Polski 1:50 000, ark. Iława (210). *Centr. Arch. Geol.*
602 *Państw. Inst. Geol.*, [Detailed Geological Map of Poland, scale 1:50 000, Iława sheet (210).
603 Warsaw, 2009.

604 Givélet, N., Le Roux, G., Cheburkin, A., Chen, B., Frank, J., Goodsite, M. E., Kempter, H.,
605 Krachler, M., Noernberg, T., Rausch, N., Rheinberger, S., Roos-Barraclough, F., Sapkota, A.,
606 Scholzb, Ch., Shotyk, W.: Suggested protocol for collecting, handling and preparing peat cores
607 and peat samples for physical, chemical, mineralogical and isotopic analyses. *J. Environ. Monit.*,
608 6, 481–492, 2004.

609 Hofmann, G., Werum, M., Lange-Bertalot, H.: Diatomeen im Süßwasser-Benthos von
610 Mitteleuropa. A.R.G. GantnerVerlag, Rugell, Liechtenstein, 1–908, 2011.

611 Holmes J.A., De Decker P.: The chemical composition of ostracod shells: application in Quaternary
612 paleoclimatology, in: *Ostracoda as proxies for Quaternary climate change, Developments in*
613 *Quaternary Science* 12, 131-140, edited by: Horne D., Holmes J.A., Rodriguez-Lazaro J. &
614 Viehberg F., 2012.

615 Hunter, L. E., Delaney, A. J., Lawson, D. E. Davis, L.; Downhole GPR for high-resolution analysis
616 of material properties near Fairbanks, Alaska, in: *Ground Penetrating Radar in Sediments,*
617 *Geological Society, Special Publications* 211, 275-285, edited by: Bristow, C. S., Jol, H. M.,
618 London, 2003.

619 Ivanić, M., Lojen, S., Grozić, D., Jurina, I., Škapin, S.D., Troskot-Čorbić, T., Mikac, N., Juračić,
620 M.: Geochemistry of sedimentary organic matter and trace elements in modern lake sediments
621 from transitional karstic land–sea environment of the Neretva River delta (Kuti Lake, Croatia),
622 *Quaternary International* 494, 286-299, 2018.

623 Jelinowska, A., Tucholka, P., Wieckowski, K.: Magnetic properties of sediments in a Polish lake:
624 evidence of a relation between the rock-magnetic record and environmental changes in Late
625 Pleistocene and Holocene sediments, *Geophys. J. Int.*, 129,727-736, 1997.

626 Karst-Riddoch, T.L., Pisaric, M.F.J., Smol, J.P.: Diatom responses to 20th century climate-related
627 environmental changes in high elevation mountain lakes of the northern Canadian Cordillera,
628 *Journal of Paleolimnology* 33, 265–282, 2005.

629 Kilham, P., Kilham, S.S., Hecky, R.E.: Hypothesized resource relationships among African
630 planktonic diatoms, *Limnology and Oceanography* 31, 1169–1181, 1986.

631 Jutrzenka – Trzebiatowski, A., Hołdyński, A.: *Roślinność rzeczywista Parku Krajobrazowego*
632 *Pojezierza Iławskiego*. Akademia Rolniczo-Techniczna. Olsztyn: 1–36, 1997.

633 Kołaczek, P., Kuprjanowicz, M., Karpińska-Kołaczek, M., Szal, M., Winter, H., Danel, W.,
634 Pochocka-Szwarc, K., Stachowicz-Rybka, R.: The Late Glacial and Holocene development of
635 vegetation in the area of a fossil lake in the Skaliska Basin (north-eastern Poland) inferred from
636 pollen analysis and radiocarbon dating, *Acta Palaeobot.*, 53(1), 23–52, 2013.

637 Kondracki, J.: *Geografia regionalna Polski*. Wyd. Nauk. PWN, Warszawa, 2002.

638 Kraska M., Piotrowicz R.: Lobelia lakes: specificity, trophy, vegetation and protection, in:
639 protection of beds and wetlands of Pomerania region 3, 48-52, edited by: Malinowski B., 2000.

640 Kuprjanowicz, M.: *Badania palinologiczne w Polsce północno-wschodniej*, in: *Człowiek i jego*
641 *środowisko (Polska północno-wschodnia w holocenie)*, Botanical Guidebooks 30, 77–95, edited
642 by: Wacnik A., Madeyska, 2008.

643 Kupryjanowicz, M.: Postglacial development of vegetation in the vicinity of the Wigry Lake,
644 *Geochronometria* 27, 53-66, 2007.

645 Lin, Y.T., Schuettpehl, C.C., Wu, C.H., and Fratta, D.: A combined acoustic and electromagnetic
646 wave-based techniques for bathymetry and subbottom profiling in shallow waters, *Journal of*
647 *Applied Geophysics*, 68, 203–218, 2009.

648 Ma, L., Wu, J., Abuduwaili, L., Liu, W.: Geochemical Responses to Anthropogenic and Natural
649 Influences in Ebinur Lake Sediments of Arid Northwest China, *Plos One*, 13/11(5):e0155819:
650 doi: 10.137, 2016.

651 Madeja J.: Vegetation changes and human activity around Lake Łańskie (Olsztyn Lake District,
652 NE Poland) from the mid Holocene, based on palynological study, *Acta Palaeobotanica* 53(2),
653 235–261, 2013.

654 Mann, M. E., Zhang, Z., Rutherford, S., et al.: (2009). Global Signatures and Dynamical Origins
655 of the Little Ice Age and Medieval Climate Anomaly, *Science* 326 (5957), 1256–60, 2009.

656 McCormick, M., Büntgen, U., Cane, M.A., Cook, E.R., Harper, K., Huybers, P., Litt, T., Manning,
657 S.W., Mayewski, P. A., More, A.F.M., Nicolussi, K., Tegel, W.: Climate Change during and
658 after the Roman Empire: Reconstructing the Past from Scientific and Historical Evidence,
659 *Journal of Interdisciplinary History* 43:2 (Autumn, 2012), 169–220.

660 Miotk-Szpiganowicz, G., Zachowicz, J., Uścińowicz, S.: Review and reinterpretation of the pollen
661 and diatom data from the deposits of the Southern Baltic lagoons, *Polish Geological Institute*
662 *Special Papers*, 23, 45–70, 2008.

663 Myers, K.J., Wignall, P.: Understanding Jurassic Organic-rich Mudrocks—New Concepts using
664 Gamma-ray Spectrometry and Palaeoecology: Examples from the Kimmeridge Clay of Dorset
665 and the Jet Rock of Yorkshire, in: *Marine Clastic Sedimentology*, 172-189, edited by: J.K.,
666 Legett, G., Zauffa, 1987.

667 Nitychoruk, J. Welc, F.: Janiki Wielkie. Środowisko fizyczno-geograficzne, in: *Katalog Grodzisk*
668 *Warmii i Mazur*, tom 2, 153 – 155, edited by: Kobyliński, Z., Warszawa, 2017.

669 Noryskiewicz, B.: Lake Steklin – a reference site for the Dobrzyń-Chełmno Lake District, N
670 Poland, Report on palaeoecological studies for the IGCP-Project No. 158B, *Acta Palaeobot.*,
671 22(1), 65-83, 1982.

672 Noryskiewicz A. M.: *Historia roślinności i osadnictwa Ziemi Chełmińskiej w późnym holocenie*,
673 *Studium palinologiczne*, Toruń, 2013.

674 Noryskiewicz, B.: History of vegetation during the Late-Glacial and Holocene in Brodnica Lake
675 District in light of pollen analysis of Lake Strażym deposits, *Acta Palaeobot.*, 27(1), 283-304,
676 1987.

677 Noryskiewicz, B., Ralska-Jasiewiczowa, M.: Type region P-w: Dobrzyń-Olsztyn lake District,
678 *Acta Palaeobotanica* 29(2), 85-93, 1989.

679 Ojala, A.E.K., Kosonen, E., Weckström, J., Korkonen, S., Korhola, A.: Seasonal formation of
680 clastic-biogenic varves: the potential for palaeoenvironmental interpretations, *GFF*, Special
681 issue: *Varve Genesis Chronol.*, *Paleoclimate* 135 (3/4), 237-247, 2013.

682 Pawlikowski M., Ralska-Jasiewiczowa M., Schönborn W., Stupnicka E., Szeroczyńska K.:
683 Woryty near Gietrzwałd, Olsztyn Lake District, NE Poland – history and lake development during
684 the last 12000 years, *Acta Palaeobotanica* 22(1), 85-116, 1982.

685 Peck, J. A., King, J. W., Colman, S. M., Kravchinsky, V. A.:1994. A rock magnetic record from
686 lake Baikal, Syberia: Evidence for Late Quaternary climate change, *Earth. Planet. Sci. Lett.*,
687 122, 221-238, 1994.

688 Rabięga, K., Rutyna, M., Wach, D.: Janiki Wielkie. Badania Archeologiczne, w: Katalog Grodzisk
689 Warmii i Mazur, tom 2. Warszawa, 155–174, edited by: Kobyliński, Z., Warszawa, 2017.

690 Ralska-Jasiewiczowa M.: Type region P-x: Masurian Great Lake Dystrykt. *Acta Palaeobotanica*
691 29(2), 95-100, 1989.

692 Ralska-Jasiewiczowa M., Latałowa A.: Poland, in: *Palaeoecological events during the Last 15,000*
693 *years: Regional Synthesis of Palaeoecological Studies of Lakes and Mires in Europe*, 403-472,
694 edited by: Berglund, B.E., Birks H.J.B., Ralska-Jasiewiczowa, M., Wright H.E., John Wiley &
695 Sons: Chichester, 1996.

696 Ralska-Jasiewiczowa M, Goslar T, Madeyska T, Starkel L.: Lake Gościąg, Central Poland, a
697 monographic study, W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków, 1998.

698 Sambuelli, L., Silvia, S.: Case study: A GPR survey on a morainic lake in northern Italy for
699 bathymetry, water volume and sediment characterization, *Journal of Applied Geophysics* 81,
700 48-56, 2012.

701 Sambuelli, L., Calzoni, C., Pesenti, M.: Case history: Waterborne GPR survey for estimation
702 bottom-sediment variability: A survey on the Po River, Turin, Italy, *Geophysics* 74, 95–102,
703 2009.

704 Sandgren, P., Snowball, I.: Application of mineral magnetic techniques to Paleolimnology, in:
705 *Tracking environmental change using lake sediments, Physical and geochemical methods 2*,
706 edited by: Last, W., Smol, J., Kluwer Academic Publishers. Netherlands, 2001.

707 Semrau, A.: *Mitteilungen des Copernicus-Vereins für Wissenschaft und Kunst zu Thorn. Heft 42. Elbing.*
708 1939.

709 Smol, J.P., Birks, J.B., Last, W.M.: *Tracking Environmental Change Using Lake Sediments,*
710 *Terrestrial, Algal and Siliceous Indicators* 3, 371, Springer Verlag, 2001.

711 Snowball, I. F.: Mineral magnetic properties of Holocene lake sediments and soils from the Karsa
712 valley, Lapland, Sweden, and their relevance to palaeoenvironmental reconstruction, *Terra*
713 *Nova* 5, 258 – 270, 1993.

714 Stankevica, K., Kalnina, L., Klavins, M., Cerina, A., Ustupe, L., Kaup, E.: Reconstruction of the
715 Holocene Palaeoenvironmental Conditions Accordingly to the Multiproxy Sedimentary
716 Records from Lake Pilvelis, Latvia, *Quaternary International* 386, 102-15, 2015.

717 Stopa-Boryczka, M., Boryczka, J., Wawer, J., Grabowska, K., Dobrowolska, M., Osowiec, M.,
718 Błażek, E., Skrzypczuk, J., Grzęda.: Climate of north–eastern Poland based on J. Kondracki and
719 J. Ostrowski`s Physiographic division. Atlas of interdependence of meteorological and
720 geographical parameters in Poland. Warsaw University, Warsaw, 2013.

721 Szal M., Kupryjanowicz M., Wyczółkowski M.: Late Holocene changes in vegetation of the
722 Mrągowo Lakeland (NE Poland) as registered in the pollen record from Lake Salet, *Studia*
723 *Quaternaria*, 31(1), 51–60, 2014a.

724 Szal M., Kupryjanowicz M., Wyczółkowski M., Tylmann W.: The Iron Age in the Mrągowo Lake
725 District, Masuria, NE Poland: the Salet settlement microregion as an example of long-lasting
726 human impact on vegetation. *Veget. Hist. Archaeobot.*, 23, 419–437, 2014b.

727 Thompson, R., Oldfield, F.: *Environmental magnetism*, Allen and Unwin London, 1986.

728 Tiljander, M., Ojala, A.E.K., Saarinen, T., Snowball, I.: Documentation of the physical properties
729 of annually laminated (varved) sediments at a sub-annual to decadal resolution for
730 environmental interpretation, *Quat. Int.*, 88 (1), 5-12, 2002.

731 Tylmann, W., Szpakowska, K., Ohlendorf, C., Woszczyk, M., Zolitschka, B.: Conditions for
732 deposition of annually laminated sediments in small meromictic lakes: a case study of Lake
733 Suminko (northern Poland), *J. Paleolimnol.*, 47 (1), 55-70, 2012.

734 Valpola, S.E., Ojala, A.E.K.: Post-glacial sedimentation rate and patterns in six lakes of the
735 Kokemäenjoki upper watercourse, Finland, *Boreal Environ. Res.*, 11, 195-211, 2006.

736 Verosub, K. L., Roberts, A. P.: *Environmental Magnetism: past, present and future*, *Journal of*
737 *Geophysical Research* 100, 2175-2192, 1995.

738 Wacnik A, Tylmann W, Bonk A et al., Determining the responses of vegetation to natural processes
739 and human impacts in north-eastern Poland during the last millennium: Combined pollen,
740 geochemical and historical data, *Vegetation History and Archaeobotany*. Epub ahead of print
741 17 March. DOI: 10.1007/s00334-016-0565-z, 2016.

742 Wacnik A., Kupryjanowicz M., Mueller-Bieniek A., Karczewski M., Cywa K.: The environmental
743 and cultural contexts of the late Iron Age and medieval settlement in the Mazurian Lake District,

744 NE Poland: combined palaeobotanical and archaeological data, *Veget. Hist. Archaeobot.*, 23,
745 439–459, 2014.

746 Welc, F.: Lake sediments and geoarchaeology (editorial), *Studia Quaternaria* 34(1), 3-8, 2017.

747 Welc, F.: Geoarchaeological evidence of late and post-Antiquity (5th-9th c. AD) climate changes
748 recorded at the Roman site in Plemići Bay (Zadar region, Croatia), *Studia Quaternaria* 36 (1),
749 3–17, 2019.

750 Wetzel, R.G.: Past productivity: paleolimnology, in: *Limnology. Lake and River Ecosystems*, third
751 ed., edited by: Wetzel, R.G., Elsevier, Oxford, 785-804, 2001.

752 Wilkinson, A.N., Zeeb, B.A., Smol, J.P.: *Atlas of chrysophycean cysts*, Kluwer Academic
753 Publishers, Dordrecht, 2002.

754 Williams, D.M., Round, F.E.: Revision of the genus *Fragilaria*, *Diatom Research* 2, 267–288, 1987.

755 Wirth S.B., Gilli A., Niemann H., Dahl T.W., Ravasi D., Sax N., Hamann Y., Peduzzi R., Peduzzi
756 S., Tonolla M., Lehmann M.F., Anselmetti F.S.: Combining sedimentological, trace metal (Mn,
757 Mo) and molecular evidence for reconstructing past water-column redox conditions: the
758 example of meromictic Lake Cadagno (Swiss Alps), *Geochimica et Cosmochimica Acta*, 120,
759 220-238, 2013.

760 Witkowski, A., Lange-Bertalot, H., Metzeltin, D.: Diatom flora of marine coasts, I. *Iconographia*
761 *Diatomologica* 7, 1–925, 2000.

762 Zachowicz J., Kępińska U.: The palaeoecological development of Lake Družno (Vistula Deltaic
763 Area), *Acta Palaeobotanica* 27(1), 227-249, 1987.

764 Zachowicz J., Przybyłowska-Lange W., Nagler J.: The Late-glacial and Holocene vegetational
765 history of the Żuławy region, N Poland, A. Biostratigraphic study of Lake Družno sediments,
766 *Acta Palaeobotanica* 22(1), 141-161, 1982.

767 Zalat, A., Welc, F., Nitychoruk, J., Marsk, L., Chodyka, M., Zbucki, Ł.: Last two millennia water
768 level changes of the Młynek Lake (Northern Poland) inferred from diatoms and chrysophyte
769 cysts record, *Studia Quaternaria* 35 (2), 77 – 89, 2018.

770 Zolitschka, B.: Varved lake sediments, in: *Encyclopedia of Quaternary Science*, edited by: Elias,
771 S.A., Elsevier, Amsterdam, 3105-3114, 2007.


772

773


774

775
776
777
778
779
780
781
782
783
784


ILLUSTRATIONS




785
786 Fig. 1. A - location of the Młynek Lake in the Warmia and Mazury Region (north-eastern Poland) (Drawing; Fabian
787 Welc). B – view of the Młynek Lake from the north-west (Photo: Fabian Welc), C – satellite image of the lake (open
788 source: ©Google Earth : www.google.com/intl/pl/earth). D – LIDAR image of the lake: a – lake basin, b – Janiki
789 Wielkie archaeological site established in early Iron Age (open source: ©Geoportal Poland : www.geoportal.gov.pl).
790
791
792




793
 794 Fig. 2. GPR reflection profile across the Młynek Lake (cf. Fig. 2), a – drilling M-1, b – upper boundary of the so-called
 795 hard bottom, c –stratified glaciofluvial sandy-gravel beds beneath the lake sediments, d – attenuation zone of
 796 electromagnetic waves connected with occurrence of organic sediment (gyttja) (measurements, processing and
 797 interpretation: Fabian Welc).
 798




799
 800 Fig. 3. Młynek Lake: A – location of drillings M 1-4 and transect of GPR sounding (open source: Google Earth©:
 801 www.google.com/intl/pl/earth/).
 802




803
 804 Fig. 4. Age-depth model of the core M-1 from the Młynek Lake. Good runs of a stationary distribution are shown in
 805 the upper left panel, green curves and grey histograms in the upper right panel present distributions for the sediment
 806 accumulation rate. The main bottom panel shows the calibrated ^{14}C dates (transparent blue) and the age-depth model
 807 (darker gray areas) which are indicating calendar ages. Grey stippled lines show 95% confidence intervals and the red
 808 curve shows the 'best' model based on the weighted mean age for each depth. The model was created by F. Welc using
 809 the open Bacon software (Blaauw and Christen, 2011).




810
 811 Fig. 5. Lithology of the M-1 borehole with radiocarbon determinations with 95% confidence, close up - photo of the
 812 log at 2.6 - 3.0 m depth and sedimentary rate (mm/year) estimated based on the age/depth model. Description of LOG:
 813 1 - hydrated – detritus type gyttja, 2 - very plastic - algal gyttja, 3 - gray-brown peaty - detritus gyttja, 4 - gray-brown
 814 gyttja (Photo and drawing: Fabian Welc).
 815
 816




817
 818 Fig. 6. Concentration depth curves for selected elements and TOC in the core M-1 of the Mlynek Lake sediments.
 819 Description of LOG: 1 - hydrated – detritus type gytja, 2 - very plastic - algal gytja, 3 - gray-brown peaty - detritus
 820 gytja, 4 - gray-brown gytja (Drawing: Fabian Welc).



821
 822 Fig. 7. Scatter plot showing the correlation in the core M-1 between S and TOC, Al and TOC, Ti and Fe, and Ti and
 823 Fe. (Drawing: Anna Rogóż-Matyszcak)




824
 825 Fig. 8. Diatom stratigraphy of the core M-1, showing diatom zones and lake phases and relative water level changes
 826 estimated on relation between planktonic and benthonic diatom taxa (Interpretation and drawing: Abdelfattah Zalat).
 827
 828



829
830
831
832

Fig. 9. Percentage pollen diagram from core M-1 – selected taxa.



833
 834 Fig. 10. Diagram with selected palaeoenvironmental proxies including lithology (1 - hydrated – detritus type gyttja, 2
 835 - very plastic - algal gyttja, 3 - gray-brown peaty - detritus gyttja, 4 - gray-brown gyttja) with phases of human activity
 836 in the vicinity of the Mlynek Lake, supplemented by archaeological chronology for Poland (Drawing: Fabian Welc).
 837