



# Reconstructing the Evolution of Ice Sheets, Sea Level and Atmospheric CO<sub>2</sub> During the Past 3.6 Million Years

Constantijn J. Berends<sup>1</sup>, Bas de Boer<sup>2</sup>, Roderik S. W. van de Wal<sup>1,3</sup>

<sup>1</sup> Institute for Marine and Atmospheric research Utrecht, Utrecht University, Utrecht, The Netherlands

5 <sup>2</sup> Earth and Climate Cluster, Faculty of Science, Vrije Universiteit Amsterdam, The Netherlands

<sup>3</sup> Faculty of Geosciences, Department of Physical Geography, Utrecht University, Utrecht, The Netherlands

*Correspondence to:* Constantijn J. Berends (c.j.berends@uu.nl)

**Abstract.** Understanding the evolution of, and the interactions between, ice sheets and the global climate over geological time is important for being able to constrain earth system sensitivity. However, direct observational evidence of past CO<sub>2</sub> concentrations only exists for the past 800,000 years. Records of benthic δ<sup>18</sup>O date back millions of years, but contain signals from both land ice volume and ocean temperature. In recent years, inverse forward modelling has been developed as a method to disentangle these two signals, resulting in mutually consistent reconstructions of ice volume, temperature and CO<sub>2</sub>. We use this approach to force a hybrid ice-sheet – climate model with a benthic δ<sup>18</sup>O stack, reconstructing the evolution of the ice sheets, global mean sea level and atmospheric CO<sub>2</sub> during the late Pliocene and the Pleistocene, from 3.6 million years (Myr) ago to the present day. During the warmer-than-present climates of the Late Pliocene, reconstructed CO<sub>2</sub> varies widely, from 320 – 440 ppmv for warm periods such as Marine Isotope Stage (MIS) KM5c, to 235 – 250 ppmv for the MIS M2 glacial excursion. Sea level is relatively stable during this period, with a high stand of 6 – 14 m, and a drop of 12 – 26 m during MIS M2. Both CO<sub>2</sub> and sea level are within the wide ranges of values covered by available proxy data for this period. Our results for the Pleistocene agree well with the ice-core CO<sub>2</sub> record, as well as with different available sea-level proxy data. During the early Pleistocene, 2.6 – 1.2 Myr ago, we simulate 40 kyr glacial cycles, with interglacial CO<sub>2</sub> decreasing from 280 – 300 ppmv at the beginning of the Pleistocene, to 250 – 280 ppmv just before the Mid-Pleistocene Transition (MPT). Peak glacial CO<sub>2</sub> decreases from 220 – 250 ppmv to 205 – 225 ppmv during this period. After the MPT, when the glacial cycles change from 40 kyr to 80/120 kyr cyclicity, the glacial-interglacial contrast increases, with interglacial CO<sub>2</sub> varying between 250 – 320 ppmv, and peak glacial values decreasing to 170 – 210 ppmv.

## 25 1 Introduction

Understanding the response of ice sheets and the global climate as a whole to changes in the concentrations of atmospheric CO<sub>2</sub>, is important for understanding the future evolution of the climate system. Since large-scale changes in ice sheet geometry typically occur over thousands to tens of thousands of years, sources of information other than direct observational evidence are required. In order to gain more insight in the relation between these components of the Earth system, studying their evolution during the geological past is useful.




One particularly rich source of information is presented by ice cores. The chemical and isotopical content of the ice itself can provide valuable information on the state of the world's climate, and in particular on the temperature, at the time the ice was formed (Dansgaard, 1964; Jouzel et al., 1997; Alley, 2000; Kindler et al., 2014). Air bubbles, trapped when the snow compresses first into firm and ultimately into ice, contain tiny samples of the atmosphere as it was at the time the air got trapped. The oldest ice core presently available, the EPICA Dome C core, contains ice and air bubbles dating back 800 kyr (Bereiter et al., 2015). The information obtained from these ice cores has greatly improved our understanding of the dynamics of the glacial cycles of the Late Pleistocene.

Different methods of relating chemical and isotopic properties of ocean sediments to the atmospheric CO<sub>2</sub> concentration have been used to create proxy data extending back further in time than the ice core record. Many studies have measured the  $\delta^{11}\text{B}$  of different fossil foraminifera, using the observed relation to seawater pH to calculate atmospheric CO<sub>2</sub> concentrations (Hönisch et al., 2009; Seki et al., 2010; Bartoli et al., 2011; Martínez-Botí et al., 2015; Foster and Rae, 2016; Stap et al., 2016; Chalk et al., 2017; Dyez et al., 2018; Sosdian et al., 2018). Another line of evidence has focused on the ratio of different types of alkenones in fossil foraminifera (Seki et al., 2010; Badger et al., 2013; Zhang et al., 2013), although the reliability of this proxy for CO<sub>2</sub> concentrations lower than ~400 ppmv has recently been called into question (Badger et al., 2019).

A different line of work relates the density of stomata on fossil plant leaves to atmospheric CO<sub>2</sub> concentration, providing data throughout the Cenozoic (Kürschner et al., 1996; Wagner et al., 2002; Finsinger and Wagner-Cremer, 2009; Beerling and Royer, 2011; Stults et al., 2011; Bai et al., 2015; Hu et al., 2015). This proxy has been the subject of some discussion regarding its reliability (Indermühle et al., 1999; Jordan, 2011; Porter et al., 2019), based on its discrepancies with the ice-core record, and presently unresolved problems due to the influence of other effects such as evolution, extinction, changes in local environment, and immigration of species.

Information about the global glacial state is obtained from the  $\delta^{18}\text{O}$  of fossil benthic foraminifera, which is influenced by both total ice volume and deep-sea temperature. Ocean sediment cores containing foraminiferal shells have been used to create stacks of benthic  $\delta^{18}\text{O}$  records dating back 65 Myr (Lisiecki and Raymo, 2005; Zachos et al., 2001, 2008). Fig. 1 shows the LR04 stack of benthic  $\delta^{18}\text{O}$  records (Lisiecki and Raymo, 2005), together with the EPICA Dome C CO<sub>2</sub> record (Lüthi et al., 2008) and the Earth's orbital parameters and Northern hemisphere summer insolation (Laskar et al., 2004), during the last 800 kyr.



**Figure 1: Earth's orbital parameters (obliquity, eccentricity and precession) and the resulting Northern summer insolation (Laskar et al., 2004), benthic  $\delta^{18}\text{O}$  (Lisiecki and Raymo, 2005) and ice core  $\text{CO}_2$  (Lüthi et al., 2008) during the past 800 kyr.**

Since benthic  $\delta^{18}\text{O}$  contains both an ice-based and a climate-based signal, using it to understand the evolution of either of  
5 these two Earth system components is only possible after disentangling the two contributions. Several studies have aimed at  
performing this separation by deriving either of the two signals from independent other proxies. One approach has been to  
derive deep-water temperatures from foraminiferal Mg/Ca ratios (Sosdian et al., 2009; Elderfield et al., 2012; Shakun et al.,  
2015). Global mean sea level is then solved as a closure term. Alternatively, proxies for global mean temperature have been  
obtained from records of oxygen and hydrogen isotope abundances from ice cores from Greenland (Alley, 2000) and  
10 Antarctica (Jouzel et al., 2007). In that case ocean water temperature can be resolved as closure term from the benthic signal.  
Focusing on the signal from ice volume rather than temperature, Rohling et al. (2014) used planktic  $\delta^{18}\text{O}$  in the  
Mediterranean Sea as a proxy for sea-level at the Strait of Gibraltar, translating the result into a global mean sea-level record



spanning the last 5.5 Myr. Grant et al. (2014) applied the same method to the Red Sea, producing a record of sea-level at the Bab-el-Mandab Strait with a higher accuracy, but going back only 500 kyr.

These studies are generally viewed as “data” studies; observed variables (chemical concentrations, isotope ratios, etc.) are used to derive climate parameters (ice volume, global mean temperature, pCO<sub>2</sub>, etc.) using relatively simple, time-  
5 independent concepts. Discussions about the validity of the results generally focus less on the physical processes described by these simple models, and more on the properties of the data, such as sample contamination and diagenesis, statistical limitations and measurement uncertainty.

A different family are the “model” studies, where physical-based models are used to describe the evolution of the entire climate system (atmosphere, ocean, cryosphere, carbon cycle, etc.) through time. Typically, the aim of a model study is to  
10 determine if our understanding of a physical process is good enough to explain the observations, or to use that understanding to make predictions of that process in the future. Several recent studies have aimed to reproduce the evolution of global climate and the cryosphere throughout the Pleistocene, using nothing but reconstructions of insolation as forcing (Brovkin et al., 2012; Willeit et al., 2015, 2019). By studying the differences between model results and observations from proxy data, such studies can help interpret these data, providing insight into the physical processes that govern the relation between  
15 observed and derived variables. In turn, the differences between data and model results can help to assess the importance of physical processes that are not included in the model.

Inverse modelling is a hybrid method, using an approach that combines elements from both these families of research. This method aims to derive the evolution of the entire global climate-cryosphere system through time, based on observations of benthic δ<sup>18</sup>O (Bintanja et al., 2005; Bintanja and van de Wal., 2008; de Boer et al., 2010, 2012, 2013, 2014, 2017; van de  
20 Wal et al., 2011; Stap et al., 2017, 2017; Berends et al., 2018, 2019). This is done by using ice-sheet models, with climate forcing components of varying complexity, to describe the known relations between atmospheric CO<sub>2</sub>, global temperature and climate, and ice-sheet evolution. Such models can be used to calculate the contributions to the observed benthic δ<sup>18</sup>O signal from both ice volume and deep-water temperature over time. By using a tool called an “inverse routine”, the model can be forced to (almost) exactly reproduce the benthic δ<sup>18</sup>O signal, providing a reconstruction of global climate *as it should*  
25 *have evolved* in order to explain the observations. The advantage of this approach over other proxy-based reconstructions is that the simulated changes in global climate and ice-sheet evolution are mutually consistent, building on the physical equations within the model framework. Earlier studies adopting this approach used simple, one-dimensional ice-sheet models to represent all ice on Earth or on a single hemisphere, and different parameterisations of the relation between ice volume and climate (e.g. de Boer et al., 2010; Stap et al., 2016). More recent studies have used more elaborate 3-D ice-sheet  
30 models covering different regions of the Earth, using more comprehensive mass balance parameterisations and representations of the global climate (e.g. Bintanja and van de Wal, 2008; de Boer et al., 2013, 2017; Berends et al., 2018, 2019). The most recent of these studies, by Berends et al. (2019), used a 3-D ice-sheet-shelf model, forced with output from several different GCM simulations by a so-called “matrix method” of model coupling, to simulate the last four glacial cycles.




They showed that their results agreed with geomorphological and proxy-based evidence of ice-sheet volume and extent, benthic  $\delta^{18}\text{O}$ , deep-water temperature, ice-sheet temperature and atmospheric  $\text{CO}_2$ .

The work presented here builds on the work by Berends et al. (2019), extending their results to 3.6 Myr ago to produce a time-continuous, self-consistent reconstruction of atmospheric  $\text{CO}_2$ , global climate, ice-sheet geometry and global mean sea-level, over the period where ice core records of  $\text{CO}_2$  are not available. The inverse modelling approach that was adopted to force this model with a benthic  $\delta^{18}\text{O}$  record is described in Sect. 2.1. The ice-sheet and climate models are described in Sects. 2.2 and 2.3, respectively, while the matrix method used to couple these two models is described in Sect. 2.4. The resulting reconstructions of  $\text{CO}_2$ , ice volume, temperature and sea-level over the past 3.6 Myr are presented in Sect. 3 and discussed in Sect. 4.

## 10 2 Methodology

### 2.1 Inverse modelling

In order to disentangle the contributions to the benthic  $\delta^{18}\text{O}$  signal from global land ice volume and deep-water temperature changes, we use the inverse modelling method proposed by Bintanja and van de Wal (2008) and refined by de Boer et al. (2013, 2014, 2017) and Berends et al. (2019). The two contributions are not independent; both result from changes in the Earth's climate, driven by changes in insolation and  $\text{CO}_2$ . The inverse modelling method determines how  $\text{CO}_2$  should have evolved over time in order to affect the global climate (in terms of temperature and precipitation, which control the evolution of the ice sheets) in such a way that the observed changes in benthic  $\delta^{18}\text{O}$  are reproduced. In the inverse modelling method, the modelled value for  $\text{CO}_2$  is used, together with the ice sheets simulated by the ice-sheet model, to determine the global climate by using the matrix method of model forcing. This method was presented by Berends et al. (2018) and refined by Berends et al. (2019), and is described in more detail in Sect. 2.4. The resulting climate is used to determine the surface mass balance over the ice sheets, which forces the ice-sheet model. The resulting global ice volume and deep-water temperature anomaly (which is derived from the global annual mean surface temperature, using a moving average) are used to calculate a modelled value of benthic  $\delta^{18}\text{O}$ . This approach is visualized conceptually in Fig. 2.



**Figure 2: A conceptual visualization of the inverse forward modelling approach. The model is forced externally by an insolation reconstruction and a benthic δ<sup>18</sup>O record (black boxes). pCO<sub>2</sub> is changed based on the difference between observed and modelled δ<sup>18</sup>O. The climate matrix interpolates between the pre-calculated GCM snapshots, based on the prescribed pCO<sub>2</sub> value and the modelled state of the cryosphere (ice thickness and albedo), to determine the climate that is used to calculate the surface mass balance over the ice sheets. Figure adapted from Berends et al. (2019).**

The modelled value δ<sup>18</sup>O<sub>mod</sub> is compared to the observed value δ<sup>18</sup>O<sub>obs</sub>. A modelled value that is too low indicates that the modelled climate is too warm, or the ice sheets are too small. pCO<sub>2</sub> is then decreased in the next time-step, leading to both a cooling and a subsequent increase in ice volume. This relationship is quantified by the following equation:

$$pCO_2 = \overline{pCO_2} + \alpha(\delta^{18}O_{mod} - \delta^{18}O_{obs}). \quad (1)$$

Here,  $\overline{pCO_2}$  is the mean modelled pCO<sub>2</sub> over the preceding 8.5 kyr, and  $\alpha$  is a scaling parameter. In order to calculate δ<sup>18</sup>O<sub>mod</sub>, the spatially variable isotope content of the individual ice-sheets is tracked through time, with the surface isotope balance based on the observed present-day relation between precipitation rates and isotope content according to Zwally and Giovinetto (1997). Benthic δ<sup>18</sup>O is assumed to be linearly dependent on the global mean deep-water temperature anomaly, which is calculated by temporally smoothing the global mean surface temperature anomaly over the preceding 3 kyr. The values of 8.5 kyr for the length of the CO<sub>2</sub> averaging window, 3 kyr for the deep-water temperature averaging window, and 120 ppmv ‰<sup>-1</sup> for the δ<sup>18</sup>O-CO<sub>2</sub> scaling parameter, were determined experimentally to accurately reproduce the observed benthic δ<sup>18</sup>O record.


The combination of this inverse modelling method to reconstruct pCO<sub>2</sub> and the matrix method to determine the global climate has been shown to accurately reproduce changes in benthic δ<sup>18</sup>O and the individual contributions from both global



ice volume and deep-water temperature, as well as ice-sheet volume and extent, ice-sheet surface temperature and  $p\text{CO}_2$  during the last four glacial cycles (Berends et al., 2019).

## 2.2 Ice-sheet model

The evolution of the ice sheets is simulated using ANICE, a coupled 3-D ice-sheet-shelf model (de Boer et al., 2013; Berends et al., 2018). ANICE uses a combination of the shallow ice approximation (SIA; Morland and Johnson, 1980) for grounded ice and the shallow shelf approximation (SSA; Morland, 1987) for floating ice to solve the ice mechanical equations. Internal ice temperatures, used to calculate ice viscosity, are calculated using a coupled thermodynamical module. The SIA and SSA ice velocities are averaged near the grounding line according to the approach by Winkelmann et al. (2011), as explained in detail by de Boer et al. (2013). The surface mass balance is parameterized based on monthly mean surface temperature and precipitation, where ablation is calculated using the surface temperature-albedo-insolation parameterization, as explained in more detail by Berends et al. (2018). The reconstruction by Laskar et al. (2004) is used to prescribe time- and latitude-dependent insolation at the top of the atmosphere. A combination of the temperature-based formulation by Martin et al. (2011) and the glacial-interglacial parameterization by Pollard & DeConto (2009), is used to calculate sub-shelf melt underneath the Antarctic ice shelves, calibrated by de Boer et al. (2013) to produce realistic present-day Antarctic shelves and grounding lines. A more detailed explanation is provided by de Boer et al. (2013) and references therein. A simple threshold thickness of 200 m is used to describe ice calving, where any shelf ice below this threshold thickness is removed. The model is run on four separate grids simultaneously, covering North America, Eurasia, Greenland and Antarctica, as shown in Fig. 3. The horizontal resolution is 40 km for Antarctica, North America and Eurasia, and 20 km for Greenland.



20

**Figure 3: The areas of the world covered by the four model domains of ANICE. In the North America and Eurasia domains, Greenland is omitted. Figure adapted from Berends et al. (2018).**



### 2.3 Climate model

HadCM3 is a coupled atmosphere-ocean GCM (Gordon et al., 2000; Valdes et al., 2017), which has been shown to accurately reproduce the present-day climate heat budget (Gordon et al., 2000). It has been used for future climate projections in the IPCC AR4 (Solomon et al., 2007), and paleoclimate reconstructions such as PlioMIP (Haywood and Valdes, 2003; Dolan et al., 2011, 2015; Haywood et al., 2013) and PMIP2 (Braconnot et al., 2007). Atmospheric circulation is calculated at a resolution of 2.5 ° latitude by 3.75 ° longitude. The ocean is modelled at a horizontal resolution of 1.25 ° by 1.25 °, with 20 vertical layers. We use results from several different steady-state time slice simulations with HadCM3 of different climate states to force our ice-sheet model, using the matrix method explained in Sect. 2.4.

### 2.4 Matrix method

According to Pollard (2010), a climate matrix is a collection of pre-calculated output data from several steady-state GCM simulations, called “snapshots”. These snapshots differ from each other in one or more key parameters, such as orbital configuration, prescribed atmospheric pCO<sub>2</sub>, or ice-sheet configuration. Each of these constitutes a separate dimension of the matrix. When using an ice-sheet model to simulate the evolution of an ice sheet over time, the prescribed climate is determined in every model time-step by combining the GCM snapshots according to position of the ice-sheet model state in the climate matrix. This approach occupies the middle ground between offline forcing and fully coupled ice-sheet – climate models. The different GCM snapshots contain the key feedback effects of the altitude-temperature and albedo-temperature. In addition, the matrix method captures the effect of ice sheet geometry on large-scale atmospheric circulation and precipitation. The matrix method creates a spatially variable linear interpolation of these snapshots providing a first order approach to the strength of the feedback and the effects of ice-sheet geometry on atmospheric circulation and precipitation.

In this study, we use the matrix method developed by Berends et al. (2018), where temperature fields from the different climate states are combined based on a prescribed value for pCO<sub>2</sub> and on the internally modelled ice-sheets. The feedback of the ice sheets on the climate is calculated via the effect on absorbed insolation through changes in surface albedo. This interpolation is carried out separately for all four ice sheets. The altitude-temperature feedback is parameterized by a constant lapse-rate derived from the GCM snapshots. Precipitation fields are combined based on changes in surface elevation, reflecting the orographic forcing of precipitation and plateau desert effect caused by the presence of a large ice-sheet. The equations describing these calculations are explained in more detail by Berends et al. (2018), who demonstrated the viability of this method by simulating the evolution of the North American, Eurasian, Greenland and Antarctic ice-sheets throughout the entire last glacial cycle at the same time. They showed that model results agree well with available data in terms of ice-sheet extent, sea-level contribution, ice-sheet surface temperature and contribution to benthic δ<sup>18</sup>O. Here, we apply this matrix method to the climate matrix created by Berends et al. (2019), consisting of eleven pre-calculated GCM snapshots, created with HadCM3. Two of these, produced by Singarayer and Valdes (2010), respectively represent the pre-industrial period (PI) and the last glacial maximum (LGM). The other nine, produced by Dolan et al. (2015), represent the






global climate during Marine Isotope Stage (MIS) M2 (3.3 My ago), for four different possible ice-sheet geometries and two different pCO<sub>2</sub> concentrations, plus one Pliocene control run. The total set of eleven snapshots allows the climate matrix to disentangle the effects on climate off changes in pCO<sub>2</sub> and changes in ice-sheet extent, and provides information on climates that are both colder and warmer than present-day.

### 5 3 Experimental set-up and results

Here, we describe a set of simulations from 3.6 Myr ago to the present day. The choice for this starting point aims to include the end of the Late Pliocene and the inception of the Pleistocene glacial cycles. The model was initialized with the same PRISM3 ice sheets that were used as boundary conditions in several of the HadCM3 simulations by Dolan et al. (2015) that were included in the climate matrix. The model was forced with the LR04 stack of benthic  $\delta^{18}\text{O}$  records (Lisiecki and Raymo, 2005). We chose here to perform three simulations: one “default” run, one with the  $\delta^{18}\text{O}$  forcing adjusted upwards by 0.1 ‰ and one with the forcing adjusting downwards by 0.1 ‰, as this uncertainty dominates the total uncertainty in the reconstructed CO<sub>2</sub> and global mean sea-level. Since simulating such long periods of time is very computationally demanding, we decided that the added value of including other simulations (as in the ensemble by Berends et al., 2019) did not outweigh the computational cost. As a result, uncertainties are conservative in this study.

The resulting reconstructions of CO<sub>2</sub> and global mean sea-level, together with the  $\delta^{18}\text{O}$  forcing, are shown in Fig. 4. The 40 kyr glacial cycles of the Early Pleistocene, between 2.6 and 1.2 Myr ago, are clearly visible, changing to 80/120 kyr cycles after the MPT. The visibly larger uncertainty for higher CO<sub>2</sub> concentrations, occurring during the warm periods of the late Pliocene, is due to the relative sparsity of the climate matrix for warmer-than-present states (three snapshots only) compared to the colder-than-present part (seven snapshots).




**Figure 4:** Observed benthic  $\delta^{18}\text{O}$  (black; Lisiecki and Raymo, 2005) and reconstructed  $\text{CO}_2$  (brown) and global mean sea level (blue) for the entire 3.6 Myr simulation period. The present-day values for the three variables (pre-industrial value of 280 ppmv for  $\text{CO}_2$ ) are shown by horizontal dashed lines.

- 5 The modelled sea-level is compared to several other reconstructions in Fig. 5. Shown are the reconstructions based on planktic  $\delta^{18}\text{O}$  in the Red Sea (Grant et al., 2014) and in the Mediterranean  $\delta^{18}\text{O}$  (Rohling et al., 2014), which respectively cover the last 500 kyr and 5.5 Myr. The reconstruction by de Boer et al. (2014) is based on an ice-sheet model-based decomposition of the benthic  $\delta^{18}\text{O}$  signal, very similar to the work presented here. The main difference is the climate forcing applied to the ice-sheet model, which by de Boer et al. (2014) was described by a simple globally uniform temperature
- 10 offset, and which was replaced by the climate matrix approach plus inversely modelled  $\text{CO}_2$  in our work. Despite this improvement in the climate forcing, which resulted in a simulated ice-sheet at LGM which agrees better with geomorphological evidence (Berends et al., 2018), the sea-level reconstructions are virtually indistinguishable. The reconstruction by Willeit et al. (2019) shown in Fig. 5 is based on a fully coupled ice-sheet – climate – carbon cycle model,



forced only with insolation. However, their ice-sheet model only simulated the Northern Hemisphere, so any possible contribution from Antarctica to changes in sea-level, especially during the warm Late Pliocene, is not accounted for in their reconstruction. The reconstruction by Elderfield et al. (2012) was made using a Mg/Ca-based reconstruction of sea surface temperature, which was then used to disentangle the ice volume and deep-sea temperature signals in the benthic  $\delta^{18}\text{O}$  record. Lastly, the reconstruction by Naish et al. (2009), adjusted by Miller et al. (2012) to match results from geological backstripping in New Zealand, suggests a relatively stable sea-level during the late Pliocene, 3.4 – 2.58 Myr ago, which agrees well with our results. Rohling et al. (2014), arguing that geological backstripping can provide information on relative changes in sea-level but not on absolute values, added a +20 m offset to the results by Miller et al. (2012), which made those results agree better with their own reconstruction from the Mediterranean Sea. However, our own results are reasonably close those by Miller et al. (2011) over the period between 3.4 and 2.6 Myr ago without requiring such a correction. The strong variability in global mean sea-level visible in the results of Rohling et al. (2014) during this period, which suggests the repeated disappearance and reappearance of most of the East Antarctic ice sheet, is not visible in the data of Miller et al. (2012), nor in our own results.




15 **Figure 5: Reconstructed global mean sea-level for the entire simulation period (black), compared to reconstructions based on Red Sea  $\delta^{18}\text{O}$  (Grant et al., 2014; red), Mediterranean  $\delta^{18}\text{O}$  (Rohling et al., 2014; yellow), an ice-sheet model-based inversion of the global benthic  $\delta^{18}\text{O}$  similar to this study (de Boer et al., 2014; blue), an insolation-forced, fully coupled ice-sheet – climate –**



carbon-cycle model (Willeit et al., 2019; green), a separation of the ice- and temperature-induced  $\delta^{18}\text{O}$  signals based on Mg/Ca ratios (Elderfield et al., 2012; purple), and a direct scaling of benthic  $\delta^{18}\text{O}$  scaled to match results from geological backstripping (Miller et al. 2012; Naish et al. 2009; dark red). The present-day value of zero is shown by a dashed line.

The  $\text{CO}_2$  reconstruction for the past 800 kyr is compared to the EPICA Dome C ice core record (Lüthi et al., 2008) in Fig. 6, as well as to several different model reconstructions and proxy-based reconstructions. The three other model-based reconstructions shown were created by decoupling the benthic  $\delta^{18}\text{O}$  signal using a 1-D ice-sheet model (van de Wal et al., 2011; Stap et al., 2017), or by using an insolation-forced, fully coupled ice-sheet – climate – carbon cycle model (Willeit et al., 2019). The geological proxies are based either on alkenones (Seki et al., 2010; Badger et al., 2013; Zhang et al., 2013) or  $\delta^{12}\text{B}$  ratios (Hönisch et al. 2009; Seki et al., 2010; Bartoli et al., 2011; Martínez-Botí et al., 2015; Stap et al., 2016; Chalk et al., 2017; Dyez et al., 2018; Sosdian et al., 2018) derived from benthic foraminifera, or based on stomata (Kürschner et al., 1996; Stults et al., 2011; Bai et al., 2015; Hu et al., 2015).



15 **Figure 6: Reconstructed atmospheric  $\text{CO}_2$  for the entire simulation period, compared to the EPICA Dome C ice core record (Lüthi et al., 2008), three different reconstructions based on ice-sheet-climate models (van de Wal et al., 2011; Stap et al., 2017; Willeit et al., 2019), as well as to chemical proxies based on  $^{11}\text{B}$  ratios (Seki et al., 2010; Martínez-Botí et al., 2015; Stap et al., 2016; Chalk et al., 2017; Dyez et al., 2018; Sosdian et al., 2018), alkenones (Hönisch et al., 2009; Seki et al., 2010; Bartoli et al., 2011; Badger et al., 2013; Zhang et al., 2013) and stomata (Kürschner et al., 1996; Stults et al., 2011; Bai et al., 2015; Hu et al., 2015). The pre-industrial value of 280 ppmv is shown by a dashed line.**



Our results broadly match those of van de Wal et al. (2011), who reconstructed CO<sub>2</sub> using 1-D ice models, forced with a simple spatially uniform temperature offset (assuming the strength of all feedbacks in the CO<sub>2</sub>-temperature relation to be constant in time) calculated from benthic δ<sup>18</sup>O using a similar inverse forward modelling approach. Both our results and those of van de Wal et al. (2011) show values during the late Pliocene that were higher than pre-industrial, but lower than present-day (412 ppmv at the date of writing), though the values by van de Wal et al. (2011) are at the low end of the uncertainty of our own results, as can be seen in Fig. 6. Stap et al. (2017) used a 1-D ice-sheet model set-up very similar to that by van de Wal et al. (2011), but used a more elaborate energy-balance model to represent the global climate. Their results differ markedly from ours, and those of van de Wal et al. (2011). The reconstruction by Willeit et al. (2019), who used a coupled ice-sheet – climate – carbon cycle model, agrees with ours and that by van de Wal et al. (2011) in terms of the glacial-interglacial amplitude both before and after the MPT. However, the downward linear trend in pCO<sub>2</sub>, which they prescribed manually to induce the inception of northern hemisphere glaciation at 2.6 Myr ago, and the resulting high values during the Early Pleistocene, are not visible in the other reconstructions.

None of the model-based reconstructions agrees better than the others with the available proxy data. A recent study by Badger et al. (2019) showed that the alkenone proxy has only a very low sensitivity to atmospheric CO<sub>2</sub> at low to moderate CO<sub>2</sub> concentrations, as is clearly visible in the top panel of Fig. 6, where the alkenone proxy data show a nearly constant CO<sub>2</sub> concentration throughout the last glacial cycle. They state that the reliability of this proxy for values lower than ~350 ppmv (the entire Pleistocene and large parts of the Late Pliocene) is doubtful. In addition, several studies (Indermühle et al., 1999; Jordan, 2011; Porter et al., 2019) have questioned the reliability of the stomatal proxies. However, even when we disregard both of these proxies, the level of disagreement between the different boron-based proxies is such that using them to choose one model-based reconstruction over another is not possible.

The only available data record which is reliable enough to allow such a comparison is the ice-core record. In Table 1, we compare the performance of the different model-based studies at reproducing this record. Our results show the best agreement, both in terms of the linear correlation coefficient  $R^2 = 0.68$  and of the root-mean-square-error RMSE = 15.3 ppmv (both taken over the entire 800 kyr ice-core period), of all four studies.


25

**Table 1: Statistical comparison of the different model-based CO<sub>2</sub> reconstructions (this study; van de Wal et al., 2011; Stap et al., 2017; Willeit et al., 2019) to the EPICA Dome C ice core record (Lüthi et al., 2008), as well as the glacial-interglacial amplitude in CO<sub>2</sub> both before and after the MPT for the different reconstructions.**

	EPICA	This study	Wal2011	Stap2017	Willeit2019
model-EPICA R <sup>2</sup>	-	0.68	0.62	0.25	0.25
model-EPICA RMSE (ppmv)	-	15.3	16.2	30.5	24.7
pre-MPT amplitude (ppmv)	-	46	31	75	38
post-MPT amplitude (ppmv)	86	85	74	103	72



Fig. 7 illustrates the difference between our modelled CO<sub>2</sub> values for the Early and Late Pleistocene. Interglacial CO<sub>2</sub> decreases from 280 – 300 ppmv at the beginning of the Early Pleistocene, to 250 – 280 ppmv just before the MPT. Glacial CO<sub>2</sub> decreases from 220 – 250 ppmv to 205 – 225 ppmv during this period, indicating a background trend of about -20 ppmv in 1.5 Myr. After the MPT, interglacial CO<sub>2</sub> varies between 250 – 320 ppmv, and glacial values decrease to 170 – 210 ppmv, with neither range showing a clear trend. The reduced glacial-interglacial CO<sub>2</sub> difference of 46 ppmv (averaged over the entire period) that we find in our results agrees with the value of 43 ppmv observed in the boron isotope data by Chalk et al. (2017). The values for all four model-based reconstructions, both before and after the MPT, are shown in Table 1. After the MPT, we find a glacial-interglacial difference of 85 ppmv, which agrees very well with the value of 86 ppmv over the past 800 kyr from the EPICA Dome C ice core record (Lüthi et al., 2008).




10

**Figure 7: Reconstructed atmospheric CO<sub>2</sub> during the Pleistocene. The colored regions indicate the glacial and interglacial ranges for the Early (2.6 – 1.2 Myr ago; blue) and Late (1.2 – 0.011 Myr ago; green) Pleistocene.**

The non-linear, time-dependent relation between CO<sub>2</sub> and sea level is visualised in Fig. 8, both for the different proxy data (panel A) and the different model reconstructions (panels B-F). While the lag between changes in CO<sub>2</sub> and changes in global climate is not included in any of the models except the one by Willeit et al. (2019), it is too short to be visible in the figure. Instead, the spread of the data points is due to both the slow response of the ice sheets to climate change, and the intrinsic hysteresis of ice sheet volume under a changing climate.

15



**Figure 8:** A: Sea level vs atmospheric CO<sub>2</sub> from proxy data. Blue: sea-level data from Rohling et al. (2014; last 5 Myr) vs boron isotope CO<sub>2</sub> data. Red: sea-level data from Elderfield et al. (2012; last 1.5 Myr) vs boron isotope CO<sub>2</sub> data. Gold: sea-level data from Miller et al. (2012) and Naish et al. (2009; 3.4 – 2.3 Myr ago) vs boron isotope CO<sub>2</sub> data. purple: sea-level data from Grant et al. (2014; last 500 kyr), vs the EPICA Dome C ice core record (Lüthi et al., 2008). B-F: reconstructed sea level and atmospheric CO<sub>2</sub> from different modelling studies (B: this study; C: van de Wal et al., 2011; D: de Boer et al., 2012; E: Stap et al., 2017; F: Willeit et al., 2019). de Boer et al. (2012; panel D) did not explicitly model CO<sub>2</sub>, calculating only a global mean temperature change. We converted this to CO<sub>2</sub>, using the logarithmic relation from van de Wal et al. (2011).

The combination of ice core CO<sub>2</sub> data (Lüthi et al., 2008) and Red Sea sea-level data (Grant et al., 2014), regarded as the two most reliable proxy reconstructions for the colder-than-present worlds of the Late Pleistocene and shown by the purple data points in Panel A, displays a pattern that is very similar to the model results by de Boer et al. (2012), and by this study. The reconstructions by van de Wal et al. (2011) and Willeit et al. (2019) show a smaller spread in data points, indicating that their ice sheets respond faster to changes in climate. The reconstruction by Stap et al. (2017) shows a vast spread, indicating that their ice sheets respond very slowly. The cause of this is not known (*personal communication with authors*). The results by van de Wal et al. (2011), de Boer et al. (2012), Willeit et al. (2019) and this study all show a CO<sub>2</sub> “threshold” for glaciation and sea-level drop around 250 ppmv, slightly lower than what is indicated by the Grant/EPICA data.

The much weaker response of sea level to changes in CO<sub>2</sub> for concentrations higher than 280 ppmv, which is displayed by all models except Stap et al. (2017), seems to be confirmed by the sea-level data by Miller et al. (2012) and the boron isotope



CO<sub>2</sub> data. The sea-level data by Rohling et al. (2014) show much more variation, especially during the early Pleistocene. The results by van de Wal et al. (2011) show a very clear threshold around 375 ppmv CO<sub>2</sub>, beyond which sea level rises rapidly with increasing CO<sub>2</sub>. As indicated by the colours, they only find these warm greenhouse worlds during the first half of the Miocene (20 – 12 Myr ago), which explains why none of the other models, nor the proxy data, display this behaviour.

5 Although some of the boron isotope proxies suggest CO<sub>2</sub> concentrations higher than this 375 ppmv threshold value, the sea level data do not show the associated strong rise.

## 5 Conclusions and discussion

We have presented a new, time-continuous, self-consistent reconstruction of atmospheric CO<sub>2</sub>, ice sheet evolution and global mean sea-level of the last 3.6 Myr, based on the benthic δ<sup>18</sup>O glacial proxy. This reconstruction was created by using a

10 hybrid ice-sheet – climate model (Berends et al., 2018) to decouple the contributions from ice volume and deep-water temperature to the benthic δ<sup>18</sup>O record (Lisiecki and Raymo, 2005). Our sea-level reconstruction agrees well with similar model-based results (de Boer et al., 2014), the reconstruction based on Red Sea δ<sup>18</sup>O by Grant et al. (2014) and the combined results of benthic δ<sup>18</sup>O (Naish et al., 2009) and geological backstripping (Miller et al., 2012). Our results agree less well with the reconstruction based on Mediterranean Sea δ<sup>18</sup>O by Rohling et al. (2014), which goes back further in time but has a

15 lower signal-to-noise ratio, and with the reconstruction based on benthic δ<sup>18</sup>O and Mg/Ca ratios by Elderfield et al. (2012). Although previous studies (van de Wal et al., 2011; de Boer et al., 2014; Stap et al., 2017) used more simple, parameterized ice-sheet and climate models, the resulting sea-level reconstructions are very similar.

Our CO<sub>2</sub> reconstruction agrees well with the EPICA Dome C ice core record (Lüthi et al., 2008), showing the strongest correlation of all four model-based reconstructions. Our results do not agree well with different chemical proxy-based

20 reconstructions based on foraminiferal alkenones (Seki et al., 2010; Badger et al., 2013; Zhang et al., 2013), foraminiferal δ<sup>12</sup>B ratios (Seki et al., 2010; Martínez-Botí et al., 2015; Stap et al., 2016; Chalk et al., 2017; Dyez et al., 2018; Sosdian et al., 2018) or fossil plant stomata (Kürschner et al., 1996; Stults et al., 2011; Bai et al., 201; Hu et al., 2015). However, the strong spread between those different proxies and the large uncertainty of each individual proxy record do not allow to conclude whether the difference between those proxies, and our model reconstructed values is significant.

25 Between 2.8 and 1.2 Myr ago, during the early Pleistocene, we simulate 40 kyr glacial cycles with glacial-interglacial sea-level changes of 25 – 50 m. During this period, CO<sub>2</sub> varies between 270 – 280 ppmv during interglacials and 210 – 240 ppmv during glacial maxima, with both values decreasing by about 20 ppmv over the 1.4 Myr course of the Early Pleistocene. After the Mid-Pleistocene Transition (MPT), when the glacial cycles change from 40 kyr to 80/120 kyr cyclicity, their sea-level amplitude increases to 70 – 120 m, with CO<sub>2</sub> varying between 250 – 320 ppmv during interglacials

30 and 170 – 210 ppmv during glacial maxima. This implies a glacial interglacial contrast of about 45 ppmv pre MPT and 85 ppmv post MPT. The two most reliable proxy records, the sea-level reconstruction by Grant et al. (2014) and the EPICA Dome C ice core record (Lüthi et al., 2008), display a relation between sea-level and CO<sub>2</sub> for colder-than-present world that





matches our model results. For warmer-than-present worlds, the large spread and uncertainty in available proxy data for both sea-level and CO<sub>2</sub> prevent us from drawing conclusions on the reliability of our model results in this regard. We believe that the inverse modelling method of using benthic δ<sup>18</sup>O, combined with coupled or hybrid ice-sheet – climate models, as a proxy for past CO<sub>2</sub> and sea-level, can add valuable insights into the evolution of the Earth system during these warm episodes.

5

*Data availability.* The reconstructed global mean sea-level and atmospheric CO<sub>2</sub> concentration over the past 3.6 Myr are available online at doi.org/10.5281/zenodo.3534940

*Author contributions.* CJB, BdB, and RSWvdW designed the study. CJB created the model set-up and carried out the simulations, with support from RSWvdW. CJB drafted the paper, and all authors contributed to the final version.

10

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* The Ministry of Education, Culture and Science (OCW), in the Netherlands, provided financial support for this study via the program of the Netherlands Earth System Science Centre (NESSC). B. de Boer was funded by NWO Earth and Life Sciences (ALW), project 863.15.019. This work was sponsored by NWO Exact and Natural Sciences for the use of supercomputer facilities. Model runs were performed on the LISA Computer Cluster, we would like to acknowledge SurfSARA Computing and Networking Services for their support. Special thanks go to B. Hönisch for advice on paleo-CO<sub>2</sub> data usage.

15

## 20 **References**

- Alley, R. B.: The Younger Dryas cold interval as viewed from central Greenland, *Quaternary Science Reviews* 19, 213-226, 2000
- Badger, M. P. S., Schmidt, D. N., Mackensen, A., and Pancost, R. D.: High-resolution alkenone palaeobarometry indicates relatively stable pCO<sub>2</sub> during the Pliocene (3.3-2.8 Ma), *Philosophical Transactions of the Royal Society A* 371, 2013
- 25 Badger, M. P. S., Chalk, T. B., Foster, G. L., Brown, P. R., Gibbs, S. J., Sexton, P. F., Schmidt, D. N., Pälike, H., Mackensen, A., and Pancost, R. D.: Insensitivity of alkenone carbon isotopes to atmospheric CO<sub>2</sub> at low to moderate CO<sub>2</sub> levels, *Climate of the Past* 15, 539-554, 2019
- Bai, Y.-J., Chen, L.-Q., Ranhotra, P. S., Wang, Q., Wang, Y.-F., and Li, C.-S.: Reconstructing atmospheric CO<sub>2</sub> during the Plio-Pleistocene transition by fossil *Typha*, *Global Change Biology* 21, 874-881, 2015
- 30 Bartoli, G., Hönisch, B., and Zeebe, R. E.: Atmospheric CO<sub>2</sub> decline during the Pliocene intensification of Northern Hemisphere glaciations, *Paleoceanography* 26, 2011
- Beerling, D. J. and Royer, D. L.: Convergent Cenozoic CO<sub>2</sub> history, *Nature Geoscience* 4, 418-420, 2011



- Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H., Kipfstuhl, S., and Chappellaz, J.: Revision of the EPICA Dome C CO<sub>2</sub> record from 800 to 600 kyr before present, *Geophysical Research Letters* 42, 542-549, 2015
- Berends, C. J., de Boer, B., and van de Wal, R. S. W.: Application of HadCM3@Bristolv1.0 simulations of paleoclimate as forcing for an ice-sheet model, ANICE2.1: set-up and benchmark experiments, *Geoscientific Model Development* 11, 4657-4675, 2018
- Berends, C. J., de Boer, B., Dolan, A. M., Hill, D. J., and van de Wal, R. S. W.: Modelling ice sheet evolution and atmospheric CO<sub>2</sub> during the Late Pliocene, *Climate of the Past* 15, 1603-1619, 2019
- Bintanja, R., van de Wal, R. S. W., and Oerlemans, J.: Global ice volume variations through the last glacial cycle simulated by a 3-D ice-dynamical model, *Quaternary International* 95-96, 11-23, 2002
- Bintanja, R., van de Wal, R., and Oerlemans, J.: Modelled atmospheric temperatures and global sea levels over the past million years, *Nature* 437, 125-128, 2005
- Bintanja, R. and van de Wal, R. S. W.: North American ice-sheet dynamics and the onset of 100,000-year glacial cycles, *Nature* 454, 869-872, 2008
- Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J.-Y., Abe-Ouchi, A., Crucifix, M., Driesschaert, E., Fichet, T., Hewitt, C. D., Kageyama, M., Kitoh, A., Laîné, A., Loutre, M.-F., Marti, O., Merkel, U., Ramstein, G., Valdes, P. J., Weber, S. L., Yu, Y., and Zhao, Y.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum - Part 1: experiments and large-scale features, *Climate of the Past* 3, 261-277, 2007
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus* 16, 436-468, 1964
- de Boer, B., van de Wal, R. S. W., Bintanja, R., Lourens, L. J., and Tuenter, E.: Cenozoic global ice-volume and temperature simulations with 1-D ice-sheet models forced by benthic δ<sup>18</sup>O records, *Annals of Glaciology* 51, 23-33, 2010
- de Boer, B., van de Wal, R. S. W., Lourens, L. J., and Bintanja, R.: Transient nature of the Earth's climate and the implications for the interpretation of benthic δ<sup>18</sup>O records, *Palaeogeography, Palaeoclimatology, Palaeoecology* 335-336, 4-11, 2012
- de Boer, B., van de Wal, R., Lourens, L. J., Bintanja, R., and Reerink, T. J.: A continuous simulation of global ice volume over the past 1 million years with 3-D ice-sheet models, *Climate Dynamics* 41, 1365-1384, 2013
- de Boer, B., Stocchi, P., and van de Wal, R.: A fully coupled 3-D ice-sheet-sea-level model: algorithm and applications, *Geoscientific Model Development* 7, 2141-2156, 2014
- de Boer, B., Lourens, L. J., and van de Wal, R. S. W.: Persistent 400,000-year variability of Antarctic ice volume and the carbon cycle is revealed throughout the Plio-Pleistocene, *Nature Communications* 5, 2014
- de Boer, B., Dolan, A. M., Bernales, J., Gasson, E., Goelzer, H., Gollledge, N. R., Sutter, J., Huybrechts, P., Lohmann, G., Rogozhina, I., Abe-Ouchi, A., Saito, F., and van de Wal, R. S. W.: Simulating the Antarctic ice sheet in the Late-Pliocene warm period: PLISMIP-ANT, an ice-sheet model intercomparison project, *The Cryosphere* 9, 881-903, 2015



- de Boer, B., Haywood, A. M., Dolan, A. M., Hunter, S. J., and Prescott, C. L.: The Transient Response of Ice Volume to Orbital Forcing During the Warm Late Pliocene, *Geophysical Research Letters* 44, 10,486-10,494, 2017
- Brovkin, V., Ganopolski, A., Archer, D., and Munhoven, G.: Glacial CO<sub>2</sub> cycle as a succession of key physical and biogeochemical processes, *Climate of the Past* 8, 251-264, 2012
- 5 Chalk, T. B., Hain, M. P., Foster, G. L., Rohling, E. J., Sexton, P. F., Badger, M. P. S., Cherry, S. G., Hasenfratz, A. P., Haug, G. H., Jaccard, S. L., Martínez-García, A., Pälike, H., Pancost, R. D., and Wilson, P. A.: Causes of ice age intensification across the Mid-Pleistocene Transition, *Proceedings of the National Academy of Sciences* 114, 13114-13119, 2017
- Dolan, A. M., Haywood, A. M., Hill, D. J., Dowsett, H. J., Hunter, S. J., Lunt, D. J., and Pickering, S. J.: Sensitivity of Pliocene ice sheets to orbital forcing, *Palaeogeography, Palaeoclimatology, Palaeoecology* 309, 98-110, 2011
- 10 Dolan, A. M., Haywood, A. M., Hunter, S. J., Tindall, J. C., Dowsett, H. J., Hill, D. J., and Pickering, S. J.: Modelling the enigmatic Late Pliocene Glacial Event - Marine Isotope Stage M2, *Global and Planetary Change* 128, 47-60, 2015
- Dyez, K. A., Hönisch, B., and Schmidt, G. A.: Early Pleistocene Obliquity-Scale pCO<sub>2</sub> Variability at ~1.5 Million Years Ago, *Paleoceanography and Paleoclimatology* 33, 1270-1291, 2018
- 15 Elderfield, H., Ferretti, P., Greaves, M., Crowhurst, S., McCave, I. N., Hodell, D. A., and Piotrowski, A. M.: Evolution of Ocean Temperature and Ice Volume Through the Mid-Pleistocene Climate Transition, *Science* 337, 704-711, 2012
- Finsinger, W. and Wagner-Cremer, F.: Stomatal-based inference models for reconstruction of atmospheric CO<sub>2</sub> concentration: a method assessment using a calibration and validation approach, *The Holocene* 18, 757-764, 2009
- Foster, G. L. and Rae, J. W. B.: Reconstructing Ocean pH with Boron Isotopes in Foraminifera, *Annu. Rev. Earth Planet. Sci.* 44, 207-237, 2016
- 20 Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments, *Climate Dynamics* 16, 147-168, 2000
- Grant, K. M., Rohling, E. J., Bronk Ramsey, C., Cheng, H., Edwards, R. L., Florindo, F., Heslop, D., Marra, F., Roberts, A. P., Tamisiea, M. E., and Williams, F.: Sea-level variability over five glacial cycles, *Nature Communications* 5, 2014
- 25 Haywood, A. M., Hill, D. J., Dolan, A. M., Otto-Bliesner, B., Bragg, F., Chan, W.-L., Chandler, M. A., Contoux, C., Dowsett, H. J., Jost, A., Kamae, Y., Lohmann, G., Lunt, D. J., Abe-Ouchi, A., Pickering, S. J., Ramstein, G., Rosenbloom, N. A., Salzmann, U., Sohl, L., Stepanek, C., Ueda, H., Yan, Q., and Zhang, Z.: Large-scale features of Pliocene climate: results from the Pliocene Model Intercomparison Project, *Climate of the Past* 9, 191-209, 2013
- 30 Haywood, A. M. and Valdes, P. J.: Modelling Pliocene warmth: contribution of atmosphere, oceans and cryosphere, *Earth and Planetary Science Letters* 218, 363-377, 2003
- Hönisch, B., Hemming, N. G., Archer, D., Siddall, M., and McManus, J. F.: Atmospheric carbon dioxide concentration across the mid-Pleistocene transition, *Science* 324, , 2009



- Hu, J.-J., Xing, Y.-W., Turkington, R., Jacques, F. M. B., Su, T., Huang, Y.-J., and Zhou, Z.-K.: A new positive relationship between pCO<sub>2</sub> and stomatal frequency in *Quercus guyavifolia* (Fagaceae): a potential proxy for palaeo-CO<sub>2</sub> levels, *Annals of Botany* 115, 777-788, 2015
- Indermühle, A., Stauffer, B., Stocker, T. F., Raynaud, D., and Barnola, J.-M.: Technical Comment: Early Holocene  
5 Atmospheric CO<sub>2</sub> Concentrations, *Science* 286, 1999
- Jordan, G. J.: A critical framework for the assessment of biological palaeoproxies: predicting past climate and levels of atmospheric CO<sub>2</sub> from fossil leaves, *New Phytologist* 192, 29-44, 2011
- Jouzel, J., Alley, R. B., Cuffey, K. M., Dansgaard, W., Grootes, P., Hoffmann, G., Johnsen, S. J., Koster, R. D., Peel, D., Shuman, C. A., Stievenard, M., Stuiver, M., and White, J.: Validity of the temperature reconstruction from water isotopes in  
10 ice cores, *Journal of Geophysical Research* 102, 26,471-26,487, 1997
- Jouzel, J., Masson-Delmote, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S., Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J. P., Stenni, B., Stocker, T. F., Tison, J. L., Werner, M., and Wolff, E. W.: Orbital and Millennial Antarctic Climate Variability  
15 over the Past 800,000 Years, *Science* 317, 793-797, 2007
- Kürschner, W. M., van der Burgh, J., Visscher, H., and Dilcher, D. L.: Oak leaves as biosensors of late Neogene and early Pleistocene paleoatmospheric CO<sub>2</sub> concentrations, *Marine Micropaleontology* 27, 299-312, 1996
- Kindler, P., Guillevic, M., Baumgartner, M., Schwander, J., Landais, A., and Leuenberger, M.: Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Climate of the Past* 10, 887-902, 2014
- 20 Laskar, J., Robutel, P., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astronomy & Astrophysics* 428, 261-285, 2004
- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic delta-18-O records, *Paleoceanography* 20, , 2005
- Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fisher, H.,  
25 Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000-800,000 years before present, *Nature Letters* 452, 379-382, 2008
- Martin, M. A., Winkelmann, R., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) - Part 2: Dynamic equilibrium simulation of the Antarctic ice sheet, *The Cryosphere* 5, 727-740, 2011
- 30 Martínez-Botí, M. A., Foster, G. L., Chalk, T. B., Rohling, E. J., Sexton, P. F., Lunt, D. J., Pancost, R. D., Badger, M. P. S., and Schmidt, D. N.: Plio-Pleistocene climate sensitivity evaluated using high-resolution CO<sub>2</sub> records, *Nature* 518, 49-54, 2015



- Miller, K. G., Wright, J. D., Browning, J. V., Kulpecz, A., Kominz, M., Naish, T., Cramer, B. S., Rosenthal, Y., Peltier, W. R., and Sostdian, S.: High tide of the warm Pliocene: Implications of global sea level for Antarctic deglaciation, *Geology* 40, 407-410, 2012
- Morland, L. W. and Johnson, I. R.: Steady motion of ice sheets, *Journal of Glaciology* 25, 229-246, 1980
- 5 Morland, L. W.: Unconfined ice-shelf flow, in *Dynamics of the West Antarctic Ice Sheet*, edited by C. J. van der Veen, and J. Oerlemans, pp. 99–116, Kluwer Acad., Dordrecht, Netherlands, 1987
- Naish, T., Powell, R., Levy, R., Wilson, G., Scherer, R., Talarico, F., Krissek, L., Niessen, F., Pompilio, M., Wilson, T., Carter, L., DeConto, R. M., Huybers, P., McKay, R. M., Pollard, D., Ross, J., Winter, D., Barrett, P., Browne, G., Cody, R., Cowan, E., Crampton, J., Dunbar, G., Dunbar, N., Florindo, F., Gebhardt, C., Graham, I., Hannah, M., Hansaraj, D.,
- 10 Harwood, D., Helling, D., Henrys, S., Hinnov, L., Kuhn, G., Kyle, P., Laufer, A., Maffioli, P., Magens, D., Mandernack, K., McIntosh, W., Millan, C., Morin, R., Ohneiser, C., Paulsen, T., Persico, D., Raine, I., Reed, J., Riesselman, C. R., Sagnotti, L., Schmitt, D., Sjunneskog, C., Strong, P., Taviani, M., Vogel, S., Wilch, T., and Williams, T.: Obliquity-paced Pliocene West Antarctic ice sheet oscillations, *Nature* 458, 322-328, 2009
- Pollard, D.: A retrospective look at coupled ice sheet-climate modeling, *Climatic Change* 100, 173-194, 2010
- 15 Pollard, D. and DeConto, R. M.: Modelling West Antarctic ice sheet growth and collapse through the past five million years, *Nature* 458, 329-332, 2009
- Porter, A. S., Evans-FitsGerald, C., Yiotis, C., Montañez, I. P., and McElwain, J. C.: Testing the accuracy of new paleoatmospheric CO<sub>2</sub> proxies based on plant stable isotopic composition and stomatal traits in a range of simulated paleoatmospheric O<sub>2</sub>:CO<sub>2</sub> ratios, *Geochimica et Cosmochimica Acta* 259, 69-90, 2019
- 20 Rohling, E. J., Foster, G. L., Marino, G., Roberts, A. P., Tamisiea, M. E., and Williams, F.: Sea-level and deep-sea-temperature variability over the past 5.3 million years, *Nature* 508, 477-492, 2014
- Seki, O., Foster, G. L., Schmidt, D. N., Mackensen, A., Kawamura, K., and Pancost, R. D.: Alkenone and boron-based Pliocene pCO<sub>2</sub> records, *Earth and Planetary Science Letters* 292, 201-211, 2010
- Shakun, J. D., Lea, D. W., Lisiecki, L. E., and Raymo, M. E.: An 800-kyr record of global surface ocean δ<sup>18</sup>O and
- 25 implications for ice volume-temperature coupling, *Earth and Planetary Science Letters* 426, 58-68, 2015
- Singarayer, J. S. and Valdes, P. J.: High-latitude climate sensitivity to ice-sheet forcing over the last 120 kyr, *Quaternary Science Reviews* 29, 43-55, 2010
- Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., and Averyt, K. B.: *Climate Change 2007: the Physical Science Basis*, Contribution Of Working Group 1 To The Fourth Assessment Report Of The Intergovernmental Panel On Climate
- 30 Change, 2007
- Sostdian, S. and Rosenthal, Y.: Deep-sea temperature and ice volume change across the Pliocene-Pleistocene climate transitions, *Science* 325, 306-310, 2009



- Sosdian, S. M., Greenop, R., Hain, M. P., Foster, G. L., Pearson, P. N., and Lear, C. H.: Constraining the evolution of Neogene ocean carbonate chemistry using the boron isotope pH proxy, *Earth and Planetary Science Letters* 498, 362-376, 2018
- Stap, L. B., de Boer, B., Ziegler, M., Bintanja, R., Lourens, L. J., and van de Wal, R.: CO<sub>2</sub> over the past 5 million years: Continuous simulation and new δ<sup>11</sup>B-based proxy data, *Earth and Planetary Science Letters* 439, 1-10, 2016
- 5 Stults, D. Z., Wagner-Cremer, F., and Axsmith, B. J.: Atmospheric paleo-CO<sub>2</sub> estimates based on *Taxodium distichum* (Cupressaceae) fossils from the Miocene and Pliocene of Eastern North America, *Palaeogeography, Palaeoclimatology, Palaeoecology* 309, 327-332, 2011
- Valdes, P. J., Armstrong, E., Badger, M. P. S., Bradshaw, C. D., Bragg, F., Crucifix, M., Davies-Barnard, T., Day, J. J., Farnsworth, A., Gordon, C., Hopcroft, P. O., Kennedy, A. T., Lord, N. S., Lunt, D. J., Marzocchi, A., Parry, L. M., Pope, V., Roberts, W. H. G., Stone, E. J., Tourte, G. J. L., and Williams, J. H. T.: The BRIDGE HadCM3 family of climate models: HadCM3@Bristol v1.0, *Geoscientific Model Development* 10, 3715-3743, 2017
- 10 Wagner, F., Aaby, B., and Visscher, H.: Rapid atmospheric CO<sub>2</sub> changes associated with the 8,200-years-B.P. cooling event, *Proceedings of the National Academy of Sciences* 99, 12011-12014, 2002
- 15 van de Wal, R. S. W., de Boer, B., Lourens, L. J., Köhler, P., and Bintanja, R.: Reconstruction of a continuous high-resolution CO<sub>2</sub> record over the past 20 million years, *Climate of the Past* 7, 1459-1469, 2011
- Willeit, M., Ganopolski, A., Calov, R., Robinson, A., and Maslin, M.: The role of CO<sub>2</sub> decline for the onset of Northern Hemisphere glaciation, *Quaternary Science Reviews* 119, 22-34, 2015
- Willeit, M., Ganopolski, A., Calov, R., and Brovkin, V.: Mid-Pleistocene transition in glacial cycles explained by declining CO and regolith removal, *Science Advances* 5, 2019
- 20 Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) - Part 1: Model description, *The Cryosphere* 5, 715-726, 2011
- Zachos, J. C., Pagani, M., Sloan, L., Thomas, E., and Billups, K.: Trends, Rhythms, and Aberrations in Global Climate 65 Ma to Present, *Science* 292, 686-694, 2001
- 25 Zachos, J. C., Dickens, G. R., and Zeebe, R. E.: An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics, *Nature* 451, 279-283, 2008
- Zhang, Y. G., Pagani, M., Liu, Z., Bohaty, S. M., and DeConto, R. M.: A 40-million year history of atmospheric CO<sub>2</sub>, *Philosophical Transactions of the Royal Society A* 371, 2013
- Zwally, H. J. and Giovinetto, M. B.: Areal distribution of the oxygen-isotope ratio in Greenland, *Annals of Glaciology* 25, 208-213, 1997
- 30