

1 **Younger Dryas ice-margin retreat in Greenland, new evidence** 2 **from Southwest Greenland**

3 Svend Funder¹, Anita H.L. Sørensen², Nicolaj K. Larsen¹, Anders A. Bjørk³, Jason P. Briner⁴,
4 Jesper Olsen⁵, Anders Schomacker⁶, Laura B. Levy⁷ Kurt H. Kjær¹

5 1Globe Institute, University of Copenhagen, 1350 Copenhagen K, Denmark

6 2Geosyd, 2730 Copenhagen, Denmark

7 3Department of Geoscience and Natural Resource Management, University of Copenhagen, 1350 Copenhagen K,
8 Denmark

9 4Department of Geology, University at Buffalo, Buffalo, NY 14260, USA

10 5Department of Physics and Astronomy, Aarhus University, 8000 Aarhus C, Denmark

11 6Department of Geosciences, UiT, the Arctic University of Norway, N-9037 Tromsø, Norway

12 7Department of Geology, Humboldt State University, 1 Harpst St Arcata, CA 95521, USA

13

14 **Abstract.** To date the final stage in deglaciation of the Greenland shelf, when a contiguous ice sheet margin on the
15 inner shelf transitioned to outlet glaciers in troughs with intervening ice-free areas, we generated cosmogenic ¹⁰Be dates
16 from bedrock knobs on six outlying islands along a stretch of 300 km of the Southwest Greenland coast. Despite ¹⁰Be
17 inheritance influencing some dates, the ages generally support a Greenland Ice Sheet (GrIS) margin that retreated off
18 the inner shelf during the middle Younger Dryas (YD) period. Published ¹⁰Be and ¹⁴C-dated records show that this
19 history of the GrIS margin is seen in other parts of Greenland, but with large variations in extent and speed of retreat,
20 sometimes even between neighbouring areas. Areas with a chronology extending into the Allerød period show no
21 marked ice margin change at the Allerød/YD transition except in northernmost Greenland. In contrast, landforms on the
22 shelf (moraines and grounding zone wedges) have been suggested to indicate YD readvances or long-lasting ice-margin
23 stillstands on the middle shelf. However, these features have been dated primarily by correlation with cold periods in
24 the ice core temperature records. Ice-margin retreat during the middle and late YD is explained by advection of warm
25 subsurface water at the ice-margin, and by increased seasonality. Our results therefore point to the complexity of the
26 climate/ice-margin relation, and to the urgent need for direct dating of the early deglaciation history of Greenland.

27 **Keywords:** Younger Dryas, Greenland ice sheet, Climate change, Cosmogenic exposure dating.


28

29 **1 Introduction**

30 The Younger Dryas (YD) cold climate oscillation from 12.8 to 11.7 ka BP (thousand years Before Present) began after
31 the Allerød warm period with a 200-yr-long period of cooling, and culminated with a 60-yr-long period of abrupt
32 warming, as recorded in Greenland ice cores (Steffensen et al., 2008). Over the Greenland Ice Sheet (GrIS), annual
33 mean temperatures dropped between 5 and 9°C (Buizert et al., 2014), when both summer insolation (65° N) and
34 atmospheric CO₂ were increasing (e.g. Buizert et al., 2014). YD climate changes were especially concentrated around

35 the north-eastern North Atlantic in the areas of Atlantic Meridional Overturning Circulation (AMOC) (Carlson, 2013).
36 Similar to present climate change, the YD oscillation was a result of perturbations in the Earth's climate system and,
37 with a view to the future, it is of great interest to study the effect of these climate changes on the margin of the GrIS.
38 During the YD, it seems that the GrIS in most areas had its margin on the shelf, and earlier work has concentrated on
39 the behaviour of ice streams in transverse troughs on the shelf (e.g. Larsen et al., 2016), and newer references discussed
40 below.

41 In this study, we present 18 new cosmogenic ^{10}Be exposure ages from six localities from the inner shelf spanning
42 300 km of Southwest Greenland. Our purpose is to shed light on ice-margin behaviour during the final phase of
43 deglaciation of the shelf when a contiguous GrIS margin transformed into outlet glaciers in transverse troughs feeding
44 the shelf (Fig. 1). Despite field observations that coastal islands experienced warm-based glacial scouring, ^{10}Be
45 inheritance from episodes of earlier exposure influences some samples in our chronology. Still, clustered ages suggest
46 that the GrIS margin generally retreated during the middle YD. These results are discussed in the context of previous
47 studies elsewhere in Greenland, indicating a mismatch between temperature records and ice margin behaviour. Possible
48 mechanisms which may overrule or mute the effect of temperature change in this environment, are discussed.
49



50

51

52

53

54

55

Figure 1: Location of study area (A), Estimated deglaciation ages and ^{14}C minimum constraints from this work and previous studies (B); Details of cosmogenic dates from this work, and shelf bathymetry. For references to previous results see text. Background map of Greenland and surrounding seas from BedMachine Greenland v.3 (Morlighem et al., 2017).

56 **2. Background**

57 **2.1. Setting**

58 The continental shelf in the study area in Southwest Greenland narrows from a width of c. 70 km in the north to c. 50
59 km in the south (Fig. 1). It is composed of an inner c. 25-km-wide and up to 500-m-deep trough running along the coast
60 and dissected by glacial erosion in Proterozoic orthogneiss bedrock (Henriksen, 2008). On the outer shelf, a belt of
61 shallow banks with an gently undulating surface are composed of younger stratified marine and fluvial sediments.
62 These banks are dissected by 400- to 500-m-deep transverse troughs that are a continuation of the major fjords inland
63 (Holtedahl, 1970; Henderson, 1975; Sommerhoff, 1975; Roksandic, 1979; Sommerhoff, 1981; Ryan et al., 2016). At a
64 distance of 10-15 km beyond the coast, the inner trough forms an archipelago with a multitude of small glacially
65 sculptured rocky islands and skerries, reflecting intensive, but uneven glacial erosion. From these rocky islands we
66 collected our samples (Fig. 2).

67 **2.2. Deglaciation history**

68 Although there is little evidence for glacier overriding, it is likely that the ice sheet covered the rather narrow shelf
69 during LGM evidence from a marine core in the Davis Strait outside the Fiskebanke trough suggests that the ice-
70 margin here stood at the shelf break until deglaciation began at c. 18.6 cal. ka BP (Winsor et al., 2015a). By c. 11 ka,
71 the retreating ice-margin reached the present coastline, and the subsequent deglaciation of the fjords and land began, as
72 summarised by Winsor et al. (2015b). This leaves a period of c. 7000 years with the ice-margin inland of the shelf edge,
73 but otherwise unaccounted for. Possible evidence for prolonged GrIS margin during this interval is a series of lobate
74 moraines that run along the troughs and impinge on the inner side of the banks (Fig. 1) (Sommerhoff, 1975; Winsor et
75 al., 2015a). From their setting, these moraines were correlated with the Fiskebanke moraine system to the north (Funder
76 et al., 2011), where they were thought to date from a YD readvance on the shelf (van Tatenhove et al., 1996; Roberts et
77 al., 2009). In our area a limited YD readvance on the inner shelf, the Neria stade, was postulated by Weidick et al.
78 (2004), based on weathering limits on coastal mountains. A YD readvance in this part of the GrIS was also suggested
79 by modelling, which indicated that the ice-margin in SW Greenland retreated from the shelf edge to the present
80 coastline in the Bølling-Allerød period, but then returned to the shelf during YD (Simpson et al., 2009; Lecavalier et al.,
81 2014). A grounding zone wedge in the Fiskebanke trough points to a stillstand or readvance of the glacier front at an
82 unknown time during deglaciation (Fig. 1, Ryan et al., 2016). The significance of these features is discussed below in
83 the light of our new chronology.

84




Figure 2: Sampling localities: (a) Buksefjord (Sample X1526, 12.0 ka), (b) Fiskenæsset (Sample X1521, 13.0 ka), (c) Ravens Storø (Sample X1520, 17.0 ka, inheritance), (d) Avigaat (Sample X1518, 10.3 ka), (e) Pamiut (Sample X1515, 12.0 ka), (f) Sermiligarsuk (Sample X1507, 10.9 ka).

85
86
87
88
89
90
91
92

93

94 **3. Field and laboratory methods**

95

96 Samples were collected from the summits of bedrock knobs in glacially sculpted islands along the inner shelf in
97 Southwest Greenland. Unfortunately, erratic boulders on the bare rock surfaces were largely absent. This potentially
98 represents a problem because while boulders were ideally incorporated in the ice in a pristine condition without
99 previous exposure to cosmic radiation, the glacial erosion of the bedrock surface may not have been deep enough to
100 remove inherited isotopes from older exposures, which may result in overestimation of the age (Briner et al., 2006;
101 Corbett et al., 2013; Larsen et al., 2014). To minimize the risk for inheritance, we selected sites in the lowland where
102 the overlying ice would have been thickest and most erosive, but above the marine limit to avoid the risk of shielding of
103 the rock surface by the sea. From each site we collected 3-4 samples within a small radius to be sure that all samples
104 from each locality had been deglaciated at the same time.

105 Contrary to inheritance, other geologic processes may yield young ages that are younger than true time of
106 exposure. This may occur if a surface has been partially shielded from cosmic radiation by vegetation, snow cover or a
107 veneer of glacial sediments for some time (Gosse and Phillips, 2001). However, it is unlikely that the rocky summits
108 were vegetated, as soil would have been washed into the depressions of the glacial sculpture, which was not observed
109 (Fig. 2). Also, long-lasting and deep snow cover over sample sites is unlikely in the stormy and maritime climate at the
110 outer coast. Indeed, we experienced heavy snowfall during the sampling, with thick snow accumulating in hollows,
111 while the tops were left free of snow (Fig. 2). Topographic shielding from nearby mountains was checked with a
112 clinometer in the field.

113 The laboratory work comprised sample preparation at the University of Buffalo and measurement of ^{10}Be -
114 concentrations at the AMS facility at Aarhus University. The laboratory procedure for the preparation followed the
115 University at Buffalo's protocols (Briner, 2015). Samples were crushed and sieved to 250–500 μm , then exposed to a
116 magnetic separator to remove the more magnetic minerals and facilitate the subsequent froth flotation. In addition to
117 flotation, some samples (X1509, X1513, X1521) had to undergo heavy mineral separation to obtain sufficient amounts
118 of quartz. Before the next step the samples were examined under a microscope to see if they had been substantially
119 purified. Finally, the samples were etched by hydrogen chloride (HCl) and a mixture of hydrofluoric and nitric acid
120 (HF/HNO_3) in order to further isolate pure quartz from remaining minerals. Quartz purity was then verified by
121 inductively coupled plasma optical emission spectroscopy at the University of Colorado. Pure quartz samples were fully
122 dissolved with a ^9Be carrier and $\text{Be}(\text{OH})_2$ was isolated through column separation. The $^{10}\text{Be}/^9\text{Be}$ ratios were measured at
123 Aarhus AMS Centre (AARAMS) and all samples were blank corrected (Olsen et al., 2016). Nuclide concentrations
124 were normalized to the Beryllium standard 07KNSTD (Nishiizumi et al., 2007).

125 The ages were calculated with the CRONUS-Earth online calculator (Balco et al., 2008), using the $^{10}\text{Be}/^9\text{Be}$ -
126 ratio measured by the AMS subtracted the processed blank ratio. The processed blank ratio was 2.10×10^{-15} and the
127 blank-corrected sample ratios ranged from 0.76×10^{-13} to 2.58×10^{-13} . The Arctic ^{10}Be production rate (Young et al.
128 (2013) and the time-invariant scaling scheme for spallation processes given by (Lal, 1991) and Stone (2000) were
129 applied. The time-invariant scaling scheme does not incorporate variations in past geomagnetic field strength, but these
130 usually only affect younger samples, at c. 10 ka, by 1% (Nishiizumi et al., 2007). The maximum deviation between
131 different scaling schemes in this material is c. 1%, so they generally provide consistent ages and do not affect the

132 relative chronology. We used a rock density of 2.65 g cm^3 and made no correction for potential surface erosion or
133 snow/vegetation cover. The study area has undergone glacioisostatic uplift since the deglaciation, and this may
134 potentially influence the ^{10}Be ages. However, as the production rate calibration dataset probably experienced a similar
135 uplift history at our sample sites, no correction for glacioisostatic uplift is applied (cf. Young et al. 2020). Accordingly,
136 we present ^{10}Be ages without correcting for glacioisostatic uplift, similar to most other ^{10}Be studies from Greenland.
137 Individual ^{10}Be ages are presented with their 1-sigma analytical uncertainties, which include the uncertainty in the blank
138 correction, the “internal” uncertainty (Table 1). When we compare our ^{10}Be ages with ^{14}C ages or climate records we
139 include the production rate uncertainty, the “external” uncertainty (Balco et al., 2008).

140 Previously published ^{14}C ages have been re-calibrated using the Intcal20 calibration programme (Reimer et
141 al., 2013). Following the procedure adopted for dates on marine shells from Greenland, ages on marine shells have been
142 corrected with a ΔR of 0 for western Greenland and with a ΔR of -150 yr for eastern Greenland, based on dating
143 modern pre-bomb shells (e.g. Mörner and Funder, 1990), acknowledging that significant, but unknown, changes in the
144 reservoir effect may potentially have affected the ages especially in the turbulent millennia during the early deglaciation
145 phases (e.g. Andrews et al., 2018).

146

Table 1. Summary of ^{10}Be data from Southwest Greenland.

Sample ID	Latitude (N)	Longitude (W)	Elevation (m a.s.l.)	Sample type†	Shielding factor	Thickness (cm)	Quartz (g)	^9Be carrier weight (g)	^{10}Be conc. (atoms/g)×10 ⁴	^{10}Be uncert. (atoms/g)×10 ⁴	^{10}Be age (ka) internal (external) uncertainties††
Buksefjord											
X1524	63.83957	51.73826	118	bedrock	1	4.5	40.45	0.6067	6.88	0.36	14.43 ± 0.76 (1.03)
X1525	63.83970	51.73851	117	bedrock	1	5.5	33.19	0.6082	5.90	0.30	<i>12.48 ± 0.64 (0.88)</i>
X1526	63.83967	51.73839	102	bedrock	1	6	40.13	0.6086	5.59	0.26	<i>12.06 ± 0.57 (0.82)</i>
Calculated average (number of samples out of total)											<i>12.3 ± 0.4 (n=3)</i>
Fiskenæsset											
X1521	63.04961	50.99505	76	bedrock	0.999962	4.5	21.16	0.6068	5.93	0.43	<i>12.98 ± 0.95 (1.14)</i>
X1522	63.05008	50.99449	75	bedrock	0.999969	5.5	26.75	0.6074	5.85	0.31	<i>12.93 ± 0.68 (0.93)</i>
X1523	63.05016	50.99454	76	bedrock	0.999969	5.5	36.35	0.6049	6.31	0.36	<i>13.92 ± 0.81 (1.05)</i>
Calculated average (number of samples out of total)											<i>13.3 ± 0.6 (n=3)</i>
Ravns Storo											
X1519	62.71573	50.40947	193	bedrock	0.999986	7	35.09	0.6074	6.95	0.38	13.71 ± 0.76 (1.01)
X1520	62.71573	50.40947	189	bedrock	0.999986	6	45.21	0.6083	8.66	0.87	17.03 ± 1.72 (1.91)
X9364	62.71799	50.41719	209	bedrock	1	4.5	34.47	0.6092	6.26	0.37	<i>11.91 ± 0.70 (0.91)</i>
X9365	62.71770	50.41629	208	boulder	1	4.5	39.87	0.613	6.05	0.43	<i>11.52 ± 0.81 (0.99)</i>
Calculated average (number of samples out of total)											<i>11.7 ± 0.4 (n=2/4)</i>
Avigaat											
X1516	62.17882	49.80153	47	bedrock	1	7	45.06	0.6062	5.94	0.48	<i>13.68 ± 1.11 (1.29)</i>
X1517	62.17888	49.80107	44	bedrock	1	6	45.08	0.6089	5.23	0.24	<i>11.98 ± 0.56 (0.81)</i>
X1518	62.17894	49.80064	42	bedrock	1	4.5	45.26	0.608	4.55	1.10	<i>10.31 ± 2.49 (2.54)</i>
Calculated average (number of samples out of total)											<i>12.0 ± 1.7 (n=3/3)</i>
Paamiut											
X1513	61.85744	49.53121	65	bedrock	1	6	32.31	0.6111	5.57	0.29	<i>12.45 ± 0.66 (0.89)</i>
X1514	61.85734	49.53098	61	bedrock	1	6.5	25.48	0.6086	5.34	0.56	<i>12.05 ± 1.28 (1.40)</i>
X1515	61.85708	49.53045	60	bedrock	1	5.5	35.44	0.607	5.36	0.62	<i>12.01 ± 1.40 (1.52)</i>
Calculated average (number of samples out of total)											<i>12.2 ± 0.3 (n=3/3)</i>
Sermiligarsuk											
X1507	61.32122	48.86104	57	boulder	0.999672	6	33.35	0.6086	4.81	1.02	<i>10.86 ± 2.32 (2.38)</i>
X1509	61.32136	48.86013	61	bedrock	0.999704	5.5	24.76	0.5672	6.55	0.37	<i>14.66 ± 0.84 (1.10)</i>
Best estimate for deglaciation age											<i>10.9 ± 2.3 (n=1/1)</i>

†: All samples are coarse grained orthogneiss

††: Italics: used in average/best estimate (see text)

147
148
149150 **4. Results and Interpretations**

151

152

153

154

155


156

157

158

159

As discussed below we consider a spread of old ages as “inheritance outliers”, while the mean of clustered younger ages gives the most reliable deglaciation age. Where there is no overlap between the uncertainties, we regard the youngest age as a maximum age for deglaciation. At each site our new ages are compared to previously published cosmogenic dates of deglaciation or thinning of ice streams at fjord mouths. In addition, we also show ^{14}C results on dating marine molluscs or onset of organic sedimentation in coastal lakes. Although not dating deglaciation, these dates serve as minimum constraints for local deglaciation. Much of this information has recently been reviewed by Sinclair et al. (2016). The six sites are described below, and the results are shown in Table 1 and Fig. 3.



160

161

162

163

164

165

166

167

Figure 3: Normal kernel density plots for the ^{10}Be ages from six coastal sites in SW Greenland. The mean age (black) is calculated based on the individual samples (blue) after excluding statistical outliers (red) (see table 1).

168

4.1 Buksefjord

169

170

171

172

173

This site is located at the outer margin of the strandflat, c. 15 km from the main coastline and midway between the mouth of Ameralikfjorden and Buksefjorden (Fig. 1). The three bedrock samples from this locality were collected between 102 and 118 m a.s.l. and yielded ages of 14.4 ± 0.8 ka (X1524), 12.5 ± 0.6 ka (X1525) and 12.1 ± 0.6 ka (X1526). We interpret the oldest age as an outlier. The two youngest ages have overlapping internal uncertainty and average 12.3 ± 0.4 ka, which we interpret as the time of deglaciation at this site.

174

175

176

177

178

179

180

On the coastal mountains c. 10 km to the east, ^{10}Be dates of boulders from between 82 and 360 m a.s.l. gave an average deglaciation age of 10.7 ± 0.6 ka (Larsen et al., 2014). At the mouth of the Nuuk Fjord Complex, 30 km to the north, marine shells on the outer coast gave a minimum constraint for the deglaciation of 11.4 cal. ka BP (Weidick, 1976a), while ^{10}Be ages close to Nuuk showed deglaciation at c. 11 ka (Winsor et al., 2015b). This may imply that our dates here are some centuries too old, although the dates from around Nuuk Fjord indicate that the outer coast became ice free while an ice stream still occupied the Nuuk trough.

181 **4.2. Fiskenæsset**

182 Three bedrock samples were collected in the outer archipelago c. 6 km from the coast, from a small ice-scoured island
183 c. 15 kilometres west of the Fiskenæsset settlement (Fig. 1). The samples were collected from 75-76 m a.s.l and yielded
184 ages of 13.0 ± 1.0 ka (X1521), 12.9 ± 0.7 ka (X1522), and 13.9 ± 0.8 ka (X1523). The average, 13.3 ± 0.6 ka, is the oldest
185 deglaciation date of our sites (Fig. 1).

186 These ages imply that the GrIS margin here was close to the coast prior to the YD. This result should be
187 substantiated by other sources, but there is no available evidence from the adjacent coast to support or oppose this
188 timing of deglaciation. Farther north, at Sermilik Fjord, ^{14}C dates of marine molluscs show that the initial marine
189 transgression and retreat of the GrIS from the outer coast probably did not begin until a short time before 10.5 cal ka
190 BP, and on coastal mountains nearby, ^{10}Be ages from 450 m a.s.l. show that the GrIS surface had thinned at c. 10.6 ka
191 (Larsen et al. 2014). Even though these results come from a different trough, the difference in dates on deglaciation of
192 the coast of 2000 years warrants confirmation. However, it should be noted that Weidick (1976b) considered the
193 Sermilik glacier to be the last to retreat from the shelf in this part of Greenland, while the ice sheet margin both to the
194 north and south had already been ice free for several millennia. In areas to the south; although no direct chronology
195 exists to support this idea.

196

197 **4.3. Ravns Storø**

198 Four samples were collected on the island of Ravns Storø, in the middle of the archipelago, c. 5 km from the coast (Fig.
199 1). The samples were collected between 189 and 209 m a.s.l. within a radius of 200 m. The ages show a spread of more
200 than 5000 years: 13.7 ± 0.8 ka (X1519), 17.0 ± 1.7 ka (X1520), 11.9 ± 0.7 ka (X9364) and 11.5 ± 0.8 ka (X9365). The two
201 youngest ages, including our only boulder sample (sample X9365), have overlapping internal uncertainties, and we
202 consider their average, 11.7 ± 0.4 ka, as the best estimate for the time of deglaciation at this site, while the oldest ages are
203 outliers. From this area there is no supporting information on deglaciation history.

204

205 **4.3. Avigaat**

206 Three samples from the bedrock surface were collected from an islet in the inner archipelago, c. 3 km from the coast
207 and the abandoned Avigaat settlement (Fig. 1). The samples were taken between 42 and 47 m a.s.l. and yield ages of
208 13.7 ± 1.1 ka (X1516), 12.0 ± 0.5 ka (X1517) and 10.3 ± 2.5 ka (X1518). The variable ages and their uncertainties are
209 large. Because these ages overlap, we consider the average of 12.0 ± 1.7 ka as the best estimate for deglaciation at this
210 site. Some support that this might be generally correct comes from a ^{14}C age of c. 11.3 cal ka BP from basal gyttja in a
211 lake in coastal Nerutussoq fjord to the south (Fig. 1), giving a minimum age for deglaciation at this site (Kelly and
212 Funder, 1974).

213

214 **4.4. Paamiut**

215 This site is located on a small ice-scoured island on the inner archipelago, c. 5 km from the coast and close to the mouth
216 of Kuanersoq fjord and the town of Paamiut (Fig. 1). Here, three bedrock samples between 60 and 65 m a.s.l., are dated

217 to 12.5 ± 0.7 (X1513), 12.1 ± 1.3 (X1514) and 12.0 ± 1.4 ka (X1515). All three ages overlap within the internal
 218 uncertainty and average 12.2 ± 0.3 ka.

219 Around Paamiut and Kuanersoq several studies have supplied both ^{10}Be and ^{14}C deglaciation dates for the
 220 outer fjord. As expected, these deglaciation dates from farther inland are somewhat younger than ours. ^{10}Be dates from
 221 Kuanersoq indicate thinning of the ice margin in the fjord beginning c. 11.7 ka, and, by extrapolation, retreat from the
 222 fjord mouth at c. 11.2 ka (Winsor et al., 2015b). From a ^{14}C age of 11.0 cal ka BP for basal gyttja in a lake 8 km from
 223 our samples and well below the local marine limit, Woodroffe et al. (2014) suggested that deglaciation could not have
 224 been much earlier than c. 11 ka. These results from nearby coastal localities therefore indicate deglaciation c. 1000
 225 years later than at our site. Much of this work concerned the ice stream in Kuanersoq, while our samples come from the
 226 open coast to the south, and we suggest that an ice stream in the Kuanersoq trough remained at the inner shelf while the
 227 adjacent coastal areas became ice free.

228

229 **4.5. Sermiligaarsuk**

230 From a small island in the inner archipelago, c. 2 km from the coast and 12 km south of the Sermiligaarsuk fjord, we
 231 collected two samples from 56 and 61 m a.s.l. One sample is from bedrock (X1509) and one is from a 1-m-diameter
 232 boulder (X1507). The two samples have widely scattered ages of 10.9 ± 2.3 ka (X1507, boulder) and 14.7 ± 0.8 ka
 233 (X1509, bedrock) with a large uncertainty, particularly in the boulder age. The oldest age is unrealistic for deglaciation
 234 and interpreted as an inheritance outlier. The age of 10.9 ± 2.3 ka (sample X1507), one of our few boulder dates, is
 235 interpreted as a closer approximation for deglaciation at this site. This age is the youngest for deglaciation of the inner
 236 shelf, but the island is also closer to the coast than any of the other sites.

237 Marine shells below the marine limit in the nearby outer Sermiligaarsuk Fjord have an ^{14}C age of 9.7 cal ka
 238 BP, providing a minimum for deglaciation at this site (Weidick et al., 2004).

239

240 **5. Discussion**

241

242 **5.1. Overview of results from Southwest Greenland**

243 According to the criteria outlined above, two of the sites, Paamiut (12.2 ± 0.2 ka; $n=3$) and Fiskenæsset (13.3 ± 0.6 ka;
 244 $n=3$), contain no obvious outliers, hence no obvious inheritance, implying deglaciation ages during the middle YD
 245 (Paamiut) and prior to the YD (Fiskenæsset). Also, Avigaat has overlapping uncertainties, indicating deglaciation in late
 246 YD times, but with a large uncertainty. At two sites, Buksefjord and Ravns Storø, one or two samples are interpreted as
 247 being influenced by inheritance, but the remaining clusters indicate deglaciation during middle to late YD. Finally, at
 248 Sermiligarssuk only one sample is considered free of inheritance, yielding a best estimate for deglaciation in the Early
 249 Holocene. From this, the results, although affected by inheritance, would point to deglaciation on the inner shelf in this
 250 part of Greenland at varying times between the late Allerød and the early Holocene. However, at some sites the ages are
 251 significantly older than expected from a comparison with previous dating of deglaciation at the adjacent coast.

252 At Paamiut, Fiskenæsset and Buksefjord, our ages are up to 2000 years older than deglaciation ages obtained
 253 at nearby fjord mouths. A possible explanation may be that while the coastal areas became ice free, ice lingered in the

254 major troughs, not reaching the inner shelf until the Early Holocene, as shown previously for ice streams in Disko Bugt
255 (Jennings et al., 2014) and suggested by Weidick (1976b) for our area in Southwest Greenland.


256 However, the offset in deglaciation ages, especially the oldest from Fiskenæsset, could also be impacted by a
257 small amount of uniform inheritance in the bedrock, as demonstrated at Utsira, Norway (Briner et al., 2016). Uniform
258 inheritance may influence the mean age from a cluster, meaning that several bedrock samples from adjacent sites could
259 all have experienced a similar amount of inheritance. This effect is considered to particularly affect bedrock and
260 boulders in areas that experienced long ice-free periods between brief maximum glacial phases (Briner et al. 2016). The
261 landscape in the coastal archipelago is the result of intense erosion by warm based ice, probably back through several
262 glaciations and during the better part of the last Ice Age (e.g. Nielsen and Kuijpers, 2013; Seidenkrantz et al., 2019),
263 and we consider the type of deep, uniform inheritance as described by Briner et al. (2016) to be unlikely in our samples.

264 A possibly more likely type of uniform inheritance could be if the ice margin readvanced, but failed to erode
265 the bedrock deeply enough to remove the ^{10}Be signal from previous exposure. This could have happened during a YD
266 readvance, as suggested for this area by Weidick et al. (2004) and Lecavalier et al. (2014). However, independent dating
267 is required to show if any of these potential errors have affected the ages, especially those from Fiskenæsset.

268 In summary, the ^{10}Be dates from the coast of Southwest Greenland – although affected by inheritance –
269 suggest that the ice sheet margin retreated on the inner shelf close to the coast at least since mid/late YD times. Some
270 dates are very old compared to deglaciation dates on the coast. This could be due to differential ice margin behaviour in
271 and away from troughs or, in the case of the oldest age, to ice margin readvance over ice free land.

272

273



274
 275
 276
 277
 278
 279

Figure 4: Deglacial ice-margin features in Greenland discussed in the text. (Question marks at grounding zone wedges apply to age and come from the original literature). Sources Larsen et al. (2016) and references discussed in the text. Background map of Greenland and surrounding seas from BedMachine Greenland v.3 (Morlighem et al. 2017)

280 5.2. YD ice-margins in Greenland

281 From a recent review of YD ice-margins in Greenland, Larsen et al. (2016) concluded that ice-margin retreat indeed
 282 characterised most areas with a dated record going back through or at least into the YD. This is well constrained by ^{10}Be
 283 dating in coastal areas and ^{14}C -dated marine sediment cores in and outside major cross-shelf troughs, and applies to
 284 areas in western, eastern and southernmost Greenland (Fig 4). More recently, GrIS retreat during the YD has been
 285 corroborated from the Disko Bugt shelf (Hogan et al., 2016; Oksman et al., 2017), and from East and South Greenland
 286 (Levy et al., 2016; Andrews et al., 2018; Dyke et al., 2018; Rainsley et al., 2018).

287 In available records, the most dramatic and studied retreat occurred on the shelf at Disko Bugt, where the ice
 288 stream apparently retreated over more than 200 km from an Allerød-early YD position near the shelf break (Fig. 4), but
 289 the retreat was punctuated by periods of topographically conditioned stillstand and a spectacular, but brief, readvance
 290 (e.g. O'Cofaigh et al., 2013; Hogan et al., 2016). This is the only readvance dated to YD in West Greenland, and
 291 deserves a closer look. The readvance/retreat is recorded by till on the outer shelf and debris flows at the shelf edge.
 292 Surprisingly, ^{14}C dates on reworked shell fragments below, within and above the till, as well as in debris flows at the
 293 shelf edge, all give overlapping ages. Moreover, mid-shelf *in situ* shells, which postdate the retreat, also gives an
 294 overlapping age. This shows that both advance and retreat took place over a very short period. Using the median ages as
 295 indication Hogan et al. (2016) suggested that retreat from the shelf edge began at c. 12.24 cal. ka BP and proceeded
 296 until c. 12.1 cal. ka BP, when the ice margin stabilised in late YD times, pinned on mid-shelf topography c. 150 km
 297 from the shelf edge. This would imply an average retreat rate of c. 1 km/yr, including decade-long stops at several
 298 GZWs on the way. Even considering the large uncertainties in the dates, the retreat rate would be comparable to the
 299 highest retreat rates recorded for tidewater glaciers in SE Greenland in recent times (Bjørck et al., 2012), and, as noted
 300 by O'Cofaigh et al. (2013), seem irreconcilable with YD temperatures. As a possible explanation O'Cofaigh et al. (2013)
 301 tentatively suggested that the advance was a glacio-dynamic surge-like event, when a thin and mobile ice stream
 302 confined in the trough advanced to the shelf edge. Whatever the explanation, this singular event is without parallel
 303 anywhere in Greenland, and is hardly significant for YD climate change.

304 On the east side of Greenland, in Kangerlussuaq trough and fjord (Fig. 4), not only the shelf, but also most of
 305 the fjord became ice free during YD (e.g. Andrews et al., 2018). It is noteworthy that in these areas, as well as other
 306 areas with a record going back to the Allerød (Scoresby Sund, southernmost Greenland), there is no evidence for
 307 marked ice margin response to the initial YD cooling. In these two areas, the GrIS had retreated behind the present
 308 coastline before YD (e.g. Björck et al., 2002; Larsen et al., 2016; Levy et al., 2016; Levy et al., 2020). Only on the north
 309 coast of Greenland did glaciers apparently advance/retreat at the beginning/end of YD (Larsen et al., 2016).

310 In areas where the dated record goes back only to the middle YD, such as most of our area, or areas where
 311 the ice remained on the shelf until early Preboreal times, as shown by recent cosmogenic dates from coastal Southeast
 312 Greenland and the Sisimiut area (Fig. 4) (Dyke et al., 2018; Rainsley et al., 2018; Levy et al., 2020; Young et al., 2020),
 313 the ice margin retreat may be seen as a response to the slow warming in the latter portion of the YD (Vacca et al., 2009;
 314 Buizert et al., 2018). Therefore, while the dated records going back to Allerød times do not show evidence of ice margin
 315 readvance/stillstand at the initial YD cooling, a readvance may have occurred in other areas, considering the large
 316 variation in YD ice margin behaviour seen in the dated records. Below we discuss evidence, (e.g. landforms on the
 317 shelf), which has been attributed to such a YD ice margin readvance/stillstand.
 318

319 **5.3 YD readvance on the shelf? – moraines and GZWs**

320 Since their first discovery, the Hellefisk and Fiskebanke moraines on the West Greenland shelf have played a prominent
 321 role in the discussion of early deglaciation history (Fig.4) (e.g. Kelly, 1985; Funder et al., 2011; Hogan et al., 2016).
 322 The outermost and oldest, the Hellefisk moraine system, runs along the shelf break for 200 km at a depth of c. 200 m, c.
 323 120 km from the coast and consists of swarms of up to 100-m-high ridges (Brett and Zarudski, 1979). To the east of
 324 this, halfway towards the coast, the younger Fiskebanke moraines impinge on the inner side of the fishing banks c. 40
 325 km from the coast. These are composed of single ridges, which occur intermittently, on the inner banks and along the
 326 sides of transverse troughs for a distance of c. 500 km along the coast (Fig. 4). Although undated, the two moraine
 327 systems have generally been regarded as climate signals for two distinct periods of cooling – either Saalian and LGM
 328 (Funder et al., 2011) or LGM and YD (van Tatenhove et al., 1996; Roberts et al., 2009; Simpson et al., 2009; Lecavalier
 329 et al., 2014).

330 Recently, for the first time, absolute ages have been supplied for parts of the Hellefisk moraines, where
 331 Hogan et al. (2016), from ^{14}C dates in marine cores, found that in their study area south of Disko Bugt, these moraines
 332 represented a topographically controlled calving bay, dated to c. 12.2 cal. ka BP. In contrast to this, from a marine core
 333 at the shelf edge off Nuuk, Seidenkrantz et al. (2019) found that the outer Hellefisk moraine here dated to c. 60 ka.
 334 These results imply that the outer Hellefisk moraines are not synchronous, but have widely different ages, and are to
 335 some extent controlled not by climate but by topography.

336 Extending for 500 km along the coast the younger Fiskebanke moraines would be the most compelling
 337 evidence for ice margin response to YD cooling in Greenland, if they can be dated to YD. The results from Hellefisk
 338 moraines may also cast some doubt on the climatic significance of the Fiskebanke moraines. Their affinity to the inner
 339 shelf trough and transverse troughs could indicate topographic, rather than climatic, control. As noted above our dates
 340 from Fiskebæset, well behind the moraines, could be interpreted either in favour of or against a YD readvance to the
 341 moraines. This stresses the need for climate-independent dating of this important event.

342 Other geomorphic evidence for YD readvance or long-lasting stillstand has recently been suggested from
 343 several major transect troughs. This is based on high-resolution multibeam bathymetry, revealing a large variety of
 344 glacial landforms that formed during retreat of major ice streams. Notably, the occurrence of large GZWs has been
 345 suggested to reflect long lasting stillstand on mid-shelf (see references below). GZWs are wedge-shaped sediment
 346 accumulations deposited at the front of an ice stream during a period of stability (e.g. Dowdeswell and Fugelli, 2012).
 347 GZWs have now been observed in most investigated troughs around Greenland, and, although not dated, prominent
 348 GZWs have tentatively been assigned to the YD based on the assumption that they correlate with cold periods in the ice
 349 core temperature record (Sheldon et al., 2016; Slabon et al., 2016; Arndt et al., 2017; Newton et al., 2017; Arndt, 2018).

350 In the Uummannaq trough deglaciation of the shelf began before 15 ka and by c. 11.5 cal. ka BP the large ice
 351 stream in the trough had disintegrated into fjord glaciers with their front close to the present ice-margin (e.g. Jennings et
 352 al., 2017). However, there are two very different views on what happened in the intervening 3500 years. Based on
 353 exposure ages on coastal mountains and ^{14}C dates in the fjords Roberts et al. (2013) found that the large Uummannaq
 354 ice stream had retreated from the trough and into the fjords during YD, controlled by topography and bathymetry (Fig.
 355 4). In contrast, Sheldon et al. (2016), from a series of marine cores and a prominent GZW in the transect trough,
 356 suggested that the ice stream was stabilised for 2000 years since Allerød times on the outer shelf, 150 km further away

357 from the coast (Fig. 4). This was based on extrapolation from a ^{14}C age and correlation with the ice core temperature
 358 record.

359 An even larger discrepancy between the two dating approaches is seen in Northeast and East Greenland,
 360 where Arndt et al. (2017) and Arndt (2018), used multibeam bathymetry to interpret lineaments, mid shelf GZWs and a
 361 moraine at the mouth of Scoresby Sund as evidence for readvance of fast flowing ice streams in major troughs along the
 362 northern east coast of Greenland (Fig. 4). These features were attributed to the YD by climatic inference. This overlooks
 363 earlier work from land, especially in Scoresby Sund. Here Greenland's highest concentration of ^{10}Be and ^{14}C dates
 364 show that the outlet glaciers in this fjord system had retreated into the fjord during the Allerød, forming a swarm of
 365 moraines dating from Allerød through YD and into the Preboreal (Denton et al., 2005; Kelly et al., 2008; Hall et al.,
 366 2010; Vasskog et al., 2015; Levy et al., 2016), ruling out the possibility of a major YD readvance to or on the shelf,

367 Similar outbursts of fast flowing ice streams, reaching mid shelf GZWs were recorded also farther north in
 368 transect troughs at Kong Oscar Fjord, Kejser Franz Joseph Fjord and the wide shelf of Northeast Greenland, and dated
 369 by the same means to YD (Arndt et al., 2017; Arndt, 2018). Also here it has been overlooked that mid fjord moraines,
 370 100 km behind the mid shelf GZWs, previously have been dated to late YD/earliest Preboreal – after calibration of the
 371 ^{14}C ages (Hjort, 1979). Reconciling these two datasets would imply an extraordinarily dynamic behaviour of the ice-
 372 margin along the East Greenland seaboard, with both advances and retreats of more than 100 km within YD, in a period
 373 with increased sea ice along the coast (Flückiger et al., 2008; Buizert et al., 2018),.

374 In Northwest Greenland, mid-shelf GZWs with a length of more than 100 km have been recorded in transect
 375 troughs on the shelf in Melville Bugt (Slabon et al., 2016; Newton et al., 2017). From analogy with the “climate-
 376 correlated” Uummannaq GZW they were tentatively referred to YD, although non-climatic, bathymetric conditions may
 377 also have determined their position (Newton et al., 2017).

378 In summary, some of the landforms on the shelf, which, on climatic grounds, have been attributed to YD ice
 379 margin readvance/stillstand apparently do not date from YD, and surprisingly, none of the ^{14}C and ^{10}Be dated records
 380 show evidence for ice margin response to initial YD cooling (Fig. 4). This highlights the need for climate-independent
 381 dating of the submarine landforms and GZWs, to exploit this rich source of information, and get a better understanding
 382 of the ice sheet/climate relation.

383

384 **5.5. Ice-margin retreat during the YD cold oscillation?**

385 To explain the mismatch between YD cooling, and apparent ice-margin retreat, two agents have especially been called
 386 on: advection of warm oceanic subsurface water to the ice-margin, and increased climatic seasonality.

387 In both cases the sequence of events begins with increased production of meltwater around the North Atlantic
 388 during the Allerød warm period. The fresher and lighter water eventually sealed off the Atlantic surface circulation
 389 from the atmosphere and impeded Atlantic Meridional Overturning (AMOC). However, warm water from the
 390 subtropical areas was still driven into the North Atlantic, but now as subsurface currents (Marcott et al., 2011; Ezat et
 391 al., 2014). The subsurface water followed the path of the present North Atlantic surface circulation in the Irminger
 392 Current running south along Southeast Greenland, continuing around Greenland's southern tip, and heading northwards
 393 as the West Greenland Current (Fig. 1). Along the Greenland shelf the warm Atlantic subsurface water was present and
 394 caused ice-margin retreat in Southeast and West Greenland at 15-16 cal. ka BP, and it was continuously present at the

395 Southeast Greenland shelf edge through Bølling-Allerød and YD times (Kuijpers et al., 2003; Knutz et al., 2011;
396 Jennings et al., 2017; Andrews et al., 2018).

397 Today warm subsurface water from these currents, below a cap of fresher water, causes extensive melting of
398 floating outlet glaciers in Greenland (e.g. Mayer et al., 2000; Motyka et al., 2011), and during the early phase of
399 deglaciation when the GrIS had its entire margin on the shelf it was especially sensitive to the advection of warm
400 subsurface water, causing ice-margin retreat even when temperatures were dropping as discussed extensively in the
401 literature (Kuijpers et al., 2003; Jennings et al., 2006; Knutz et al., 2011; Rinterknecht et al., 2014; Winsor et al., 2015b;
402 Sheldon et al., 2016; Sinclair et al., 2016; Jennings et al., 2017; Oksman et al., 2017; Andrews et al., 2018; Dyke et al.,
403 2018; Rainsley et al., 2018).

404 Crucial to the impact of the warm water is the depth of the grounding line at the ice-margin, and the
405 accessibility for the warm subsurface water. This is again dependent on local bathymetry and – in the troughs – of the
406 type of connection to the open ocean, which in each area may control the impact of the warm water on the ice-margin,
407 as well as the impact from changing sea level and presence or absence of buttressing sea ice. This may explain why the
408 deglaciation of the shelf and troughs had such a different character even between neighbouring troughs, such as
409 between the rapid deglaciation of the Kangerlussuaq trough and the much slower deglaciation along the adjacent coast
410 to the south (Dyke et al., 2018). It may also explain the differences in the timing of onset of deglaciation in the
411 neighbouring Nuussuaq and Disko troughs in West Greenland as discussed by Jennings et al. (2017).

412 Increased seasonality is also connected to advection of meltwater over the North Atlantic, and reduced
413 AMOC. The fresher meltwater-diluted water seals off the ocean surface and, especially in winter, cuts off the
414 ocean/atmosphere exchange (Denton et al., 2005; Hall et al., 2008; Vacco et al., 2009; 2010; Buizert et al., 2014; Levy
415 et al., 2016; Buizert et al., 2018). This results in very cold and arid winters and increase in extent and duration of sea
416 ice, while summer temperatures – which primarily determines a glacier’s mass balance - are less affected and may even
417 warm up (Björck et al., 2002).

418 Vacco et al. (2009) presented a glaciological model for warm based glaciers on land, tuned with a
419 temperature record from a Greenland ice core, showing that for areas with high amplitude change between Allerød and
420 YD moraine deposition would be expected in the beginning of YD followed by recession of the ice margin in the mid
421 and late YD. This may explain why major YD moraines in Greenland have been seen only in northernmost Greenland
422 and Scoresby Sund – two areas where the glaciers had become landlocked already before YD, while the ice margin in
423 most other areas retreated on the shelf, controlled to a large extent by warm subsurface ocean circulation.

424 The importance of changing seasonality has recently been investigated in a model where the deglacial ice-
425 core temperature records in three ice cores are combined with simulated seasonal air temperatures for the whole of
426 Greenland, enabling assessment of variations in seasonality in time and space (Buizert et al., 2018). The varied
427 seasonality model deviates from previous models (e.g. Lecavalier et al., 2014), indicating that the dramatic temperature
428 changes in the ice core records at the beginning and end of YD, are muted in the varied seasonality model.

429 In summary, the apparent contradiction between ice core temperature records, where temperatures dropped
430 dramatically at the onset of YD, and the dated glacial record where glaciers in many parts retreated, may be explained
431 by the effect of warm subsurface water on the ice-margin, which was, all around Greenland, located on the shelf during
432 the early phase of deglaciation. Local topographic and bathymetric conditions controlled the access of warm water to
433 the ice-margin when on the shelf. Increased seasonality owes to increase in distribution and duration of winter sea ice,

434 and the YD temperature drop in the ice cores was due to a large extent to lowering of winter temperatures with little
 435 impact on the ice-margin, but a large effect on distribution, thickness and duration of sea ice. This may also explain why
 436 neither the rapid initial YD cooling nor the abrupt warming at the end left a clear signal in the dated records.
 437

438 **6. Conclusions**

439 ^{10}Be dates on bedrock surfaces in the glacially eroded archipelago on the inner shelf of southwestern Greenland are
 440 affected by ^{10}Be inherited from earlier exposure, but clustering of ages from each site suggest that the ice-margin here
 441 was retreating and close to the coast at least since mid YD times.

442 A survey of ^{10}Be and ^{14}C dated records, going back through YD elsewhere in Greenland - south, east and
 443 west - shows that also here the ice-margin was retreating, but with large differences in speed and extent even between
 444 neighbouring basins, probably controlled by local topography and trough geometry.

445 While the retreat in our and in other part dates from mid or late YD and may be seen as retreat from an initial
 446 YD readvance, areas with a record that goes back into the Allerød show no evidence for ice margin readvance at the
 447 initial YD transition. Only in northernmost Greenland did glaciers from a local ice cap apparently advance/retreat at the
 448 beginning/end of YD.

449 Moraines and GZWs on the shelf, which have been attributed to YD readvance, are dated by climatic
 450 inference, and need direct age control to distinguish climate- from non-climatic factors.

451 The apparent mismatch between the ice core temperature record and the ice-margin may be explained by the
 452 circumstance that during LGM the GrIS, contrary to other large ice sheets, around the whole perimeter was standing on
 453 the shelf and especially sensitive to changes in ocean currents, sea level, and sea ice distribution and thickness.

454 Recently, high resolution bathymetry has supplied a wealth of data on ice stream dynamics during
 455 deglaciation. However, this evidence is essentially dated only by climate-inference. To tap this rich source of
 456 information and get a better understanding of the ice sheet/climate relation, climate-independent dating of the
 457 submarine features is badly needed.

458 We subscribe to the contention by Andrews et al. (2018, p. 16): “The use of the GrIS’s isotopic records as a
 459 one-to-one template for coeval changes in glacier and ocean response potentially ignores the different response
 460 timescales between the atmosphere, oceans and cryosphere” – with a bearing also on the future.

461
 462 **Author contributions.** SF, KK and AB conceptualised the project and carried out the work in the field. AS did the
 463 laboratory work, critical assessment and compilation of data under supervision of and according to methodologies
 464 developed by JB, NL, AS, LL and JO. Visualization owes to NL and AB. The writing and editing was made by SF in
 465 close cooperation with NL and AB. KK was responsible for the funding acquisition.

466

467 **Acknowledgement**

468 This study was made possible by grants from Danish Council of Natural Sciences (FNU) and Danish National Science
 469 Foundation to carry-out field and laboratory work. It would not have been possible without generous support from the
 470 Danish Navy and the ship “HDMS Knud Rasmussen” with its skipper and his extremely help- and joyful crew. We are

471 grateful for the constructive and insightful comments offered by the referees David Ullman, David Roberts and Nicolas
472 Adams, as well as discussion with Richard Alley.
473

474

475 **References**

- 476 Andrews, J. T., Cabedo-Sanz, P., Jennings, A. E., Olafsdottir, S., Belt, S. T., and Geirsdottir, A.: Sea ice, ice-
477 rafting, and ocean climate across Denmark Strait during rapid deglaciation (similar to 16-12 cal ka BP) of the
478 Iceland and East Greenland shelves, *Journal of Quaternary Science*, 33, 112-130, 2018.
- 479 Arndt, J. E.: Marine geomorphological record of Ice Sheet development in East Greenland since the Last
480 Glacial Maximum, *Journal of Quaternary Science*, 33, 853-864, 2018.
- 481 Arndt, J. E., Jokat, W., and Dorschel, B.: The last glaciation and deglaciation of the Northeast Greenland
482 continental shelf revealed by hydro-acoustic data, *Quaternary Sci Rev*, 160, 45-56, 2017.
- 483 Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J.: A complete and easily accessible means of calculating
484 surface exposure ages or erosion rates from Be-10 and Al-26 measurements, *Quaternary Geochronology*, 3,
485 174-195, 2008.
- 486 Björck, S., Bennike, O., Rosen, P., Andresen, C. S., Bohncke, S., Kaas, E., and Conley, D.: Anomalously mild
487 Younger Dryas summer conditions in southern Greenland, *Geology*, 30, 427-430, 2002.
- 488 Bjørk, A. A., Kjær, K. H., Korsgaard, N. J., Khan, A., Kjeldsen, K. K., Andresen, C., Box, J. E., Larsen, N. K., and
489 Funder, S.: An aerial view of 80 years of climate-related glacier fluctuations in southeast Greenland, *Nature*
490 *Geoscience*, 5, 427-432, 2012.
- 491 Brett, C. P. and Zarudski, E. F. K.: Project Westmar, a shallow marine geophysical survey on the West
492 Greenland shelf, *Geological Survey of Greenland, Reports*, 87, 1-27, 1979.
- 493 Briner, J.: UB Quartz Cleaning Procedure, University at Buffalo, 1-20 pp., 2015.
- 494 Briner, J. P., Goehring, B. M., Mangerud, J., and Svendsen, J. I.: The deep accumulation of Be-10 at Utsira,
495 southwestern Norway: Implications for cosmogenic nuclide exposure dating in peripheral ice sheet
496 landscapes, *Geophysical Research Letters*, 43, 9121-9129, 2016.
- 497 Briner, J. P., Miller, G. H., Davis, P. T., and Finkel, R. C.: Cosmogenic radionuclides from fiord landscapes
498 support differential erosion by overriding ice sheets, *Geological Society of America Bulletin*, 118, 406-420,
499 2006.
- 500 Buizert, C., Gkinis, V., Severinghaus, J. P., He, F., Lecavalier, B. S., Kindler, P., Leuenberger, M., Carlson, A. E.,
501 Vinther, B., Masson-Delmotte, V., White, J. W. C., Liu, Z. Y., Otto-Bliesner, B., and Brook, E. J.: Greenland
502 temperature response to climate forcing during the last deglaciation, *Science*, 345, 1177-1180, 2014.
- 503 Buizert, C., Keisling, B. A., Box, J. E., He, F., Carlson, A. E., Sinclair, G., and DeConto, R. M.: Greenland-Wide
504 Seasonal Temperatures During the Last Deglaciation, *Geophysical Research Letters*, Research Letter, doi:
505 10.1002/2017GL075601, 2018. 1-10, 2018.
- 506 Carlson, A. E.: The Younger Dryas Climate Event, *Encyclopedia of Quaternary Science*, 3, 126-134, 2013.
- 507 Corbett, L. B., Bierman, P. R., Graly, J. A., Neumann, T. A., and Rood, D. H.: Constraining landscape history
508 and glacial erosivity using paired cosmogenic nuclides in Upernavik, northwest Greenland, *Geological*
509 *Society of America Bulletin*, 125, 1539-1553, 2013.
- 510 Denton, G. H., Alley, R. B., Comer, G. C., and Broecker, W. S.: The role of seasonality in abrupt climate
511 change, *Quaternary Sci Rev*, 24, 1159-1182, 2005.
- 512 Dowdeswell, J. A. and Fugelli, E. M. G.: The seismic architecture and geometry of grounding-zone wedges
513 formed at the marine margins of past ice sheets, *Geol Soc Am Bull*, 124, 1750-1761, 2012.
- 514 Dyke, L. M., Hughes, A. L., Andresen, C. S., Murray, T., Hiemstra, J. F., Bjørk, A. A., and Rodés, Á.: The
515 deglaciation of coastal areas of southeast Greenland, *The Holocene, Reports*, 2018.
- 516 Ezat, M. M., Rasmussen, T. L., and Groeneveld, J.: Persistent intermediate water warming during cold
517 stadials in the southeastern Nordic seas during the past 65 k.y., *Geology*, 42, 663-666, 2014.
- 518 Flückiger, J., Knutti, R., White, J. W. C., and Renssen, H.: Modeled seasonality of glacial abrupt climate
519 events, *Clim Dynam*, 31, 633-645, 2008.
- 520 Funder, S., Kjeldsen, K. K., Kjaer, K. H., and Cofaigh, C. O.: The Greenland Ice Sheet During the Past 300,000
521 Years: A Review, *Quaternary Glaciations - Extent and Chronology: A Closer Look*, 15, 699-713, 2011.

- 522 Gosse, J. C. and Phillips, F. M.: Terrestrial in situ cosmogenic nuclides: theory and application, *Quaternary*
523 *Sci Rev*, 20, 1475-1560, 2001.
- 524 Hall, B. L., Baroni, C., and Denton, G. H.: The most extensive Holocene advance in the Stauning Alper, East
525 Greenland, occurred in the Little Ice Age Polar Research, 27, 128-134, 2008.
- 526 Hall, B. L., Baroni, C., and Denton, G. H.: Relative sea-level changes, Schuchert Dal, East Greenland, with
527 implications for ice extent in late-glacial and Holocene times, *Quaternary Sci Rev*, 29, 3370-3378, 2010.
- 528 Henderson, G.: New bathymetric maps covering offshore West Greenland 59° 30' N, 1975, 761-774.
- 529 Henriksen, N.: Geological History of Greenland - Four billion years of earth evolution, Geological Survey of
530 Denmark and Greenland, Copenhagen, 2008.
- 531 Hjort, C.: Glaciation in Northern East Greenland during the Late Weichselian and Early Flandrian, *Boreas*, 8,
532 281-296, 1979.
- 533 Hogan, K. A., Cofaigh, C. O., Jennings, A. E., Dowdeswell, J. A., and Hiemstra, J. F.: Deglaciation of a major
534 palaeo-ice stream in Disko Trough, West Greenland, *Quaternary Sci Rev*, 147, 5-26, 2016.
- 535 Holtedahl, O.: On Morphology of West Greenland Shelf with General Remarks on Marginal-Channel
536 Problem, *Mar Geol*, 8, 155-+, 1970.
- 537 Jennings, A. E., Andrews, J. T., Cofaigh, C. O., St Onge, G., Sheldon, C., Belt, S. T., Cabedo-Sanz, P., and
538 Hillaire-Marcel, C.: Ocean forcing of Ice Sheet retreat in central west Greenland from LGM to the early
539 Holocene, *Earth Planet Sc Lett*, 472, 1-13, 2017.
- 540 Jennings, A. E., Hald, M., Smith, M., and Andrews, J. T.: Freshwater forcing from the Greenland Ice Sheet
541 during the Younger Dryas: evidence from southeastern Greenland shelf cores, *Quaternary Sci Rev*, 25, 282-
542 298, 2006.
- 543 Jennings, A. E., Walton, M. E., Cofaigh, C. O., Kilfeather, A., Andrews, J. T., Ortiz, J. D., De Vernal, A., and
544 Dowdeswell, J. A.: Paleoenvironments during Younger Dryas-Early Holocene retreat of the Greenland Ice
545 Sheet from outer Disko Trough, central west Greenland, *Journal of Quaternary Science*, 29, 27-40, 2014.
- 546 Kelly, M.: A review of the Quaternary geology of western Greenland. In: *Quaternary environments eastern*
547 *Canadian Arctic, Baffin Bay and western Greenland*, Andrews, J. T. (Ed.), Allen & Unwin, Boston, 1985.
- 548 Kelly, M. and Funder, S.: The pollen stratigraphy of late Quaternary lake sediments of South West
549 Greenland, *Geological Survey of Greenland, Reports*, 64, 1-26, 1974.
- 550 Kelly, M. A., Lowell, T. V., Hall, B. L., Schaefer, J. M., Finkel, R. C., Goehring, B. M., Alley, R. B., and Denton,
551 G. H.: A Be-10 chronology of lateglacial and Holocene mountain glaciation in the Scoresby Sund region, east
552 Greenland: implications for seasonality during lateglacial time, *Quaternary Sci Rev*, 27, 2273-2282, 2008.
- 553 Knutz, P. C., Sicre, M. A., Ebbesen, H., Christiansen, S., and Kuijpers, A.: Multiple-stage deglacial retreat of
554 the southern Greenland Ice Sheet linked with Irminger Current warm water transport, *Paleoceanography*,
555 26, 2011.
- 556 Kuijpers, A., Troelstra, S. R., Prins, M. A., Linthout, K., Akhmetzhanov, A., Bouryak, S., Bachmann, M. F.,
557 Lassen, S., Rasmussen, S., and Jensen, J. B.: Late Quaternary sedimentary processes and ocean circulation
558 changes at the Southeast Greenland margin, *Mar Geol*, 195, 109-129, 2003.
- 559 Lal, D.: Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models, *Earth*
560 *Planet Sc Lett*, 104, 424-439, 1991.
- 561 Larsen, N. K., Funder, S., Kjaer, K. H., Kjeldsen, K. K., Knudsen, M. F., and Linge, H.: Rapid early Holocene ice
562 retreat in West Greenland, *Quaternary Sci Rev*, 92, 310-323, 2014.
- 563 Larsen, N. K., Funder, S., Linge, H., Moller, P., Schomacker, A., Fabel, D., Xu, S., and Kjaer, K. H.: A Younger
564 Dryas re-advance of local glaciers in north Greenland, *Quaternary Sci Rev*, 147, 47-58, 2016.
- 565 Lecavalier, B. S., Milne, G. A., Simpson, M. J. R., Wake, L., Huybrechts, P., Tarasov, L., Kjeldsen, K. K., Funder,
566 S., Long, A. J., Woodroffe, S. A., Dyke, A. S., and Larsen, N. K.: A model of Greenland ice sheet deglaciation
567 based on observations of ice extent and relative sea-level, *Quaternary Sci Rev*, 102, 54-84, 2014.
- 568 Levy, L. B., Kelly, M. A., Lowell, T. V., Hall, B. L., Howley, J. A., and Smith, C. A.: Coeval fluctuations of the
569 Greenland ice sheet and a local glacier, central East Greenland, during late glacial and early Holocene time,
570 *Geophysical Research Letters*, 43, 1623-1631, 2016.

571 Levy, L. B., Larsen, N. K., Knudsen, M. F., Egholm, D. L., Bjørk, A. A., Kjeldsen, K. K., Kelly, M. A., Howley, J. A.,
572 Olsen, J., Tikhomirov, D., Zimmerman, S. R. H., and Kjær, K. H.: Multi-phased deglaciation of south and
573 southeast Greenland controlled by climate and topographic setting, *Quaternary Sci Rev*, 242, 106454, 2020.
574 Marcott, S. A., Clark, P. U., Padman, L., Klinkhammer, G. P., Springer, S. R., Liu, Z. Y., Otto-Bliesner, B. L.,
575 Carlson, A. E., Ungerer, A., Padman, J., He, F., Cheng, J., and Schmittner, A.: Ice-shelf collapse from
576 subsurface warming as a trigger for Heinrich events, *Proceedings of the National Academy of Sciences of
577 the United States of America*, 108, 13415-13419, 2011.
578 Mayer, C., Reeh, N., Jung-Rothenhausler, F., Huybrechts, P., and Oerter, H.: The subglacial cavity and
579 implied dynamics under Nioghalvfjærdssjøen Glacier, NE-Greenland, *Geophysical Research Letters*, 27,
580 2289-2292, 2000.
581 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauche, N.,
582 Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M.,
583 Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noel, B. P. Y., O'Cofaigh, C., Palmer, S., Rysgaard, S.,
584 Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and
585 Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland
586 From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44,
587 11051-11061, 2017.
588 Motyka, R. J., Truffer, M., Fahnestock, M., Mortensen, J., Rysgaard, S., and Howat, I.: Submarine melting of
589 the 1985 Jakobshavn Isbrae floating tongue and the triggering of the current retreat, *J Geophys Res-Earth*,
590 116, 2011.
591 Mörner, N.-A. and Funder, S.: C-14 dating of samples collected during the NORQUA 86 expedition, and
592 notes on the marine reservoir effect., *Meddelelser om Grønland*, 22, 57-59, 1990.
593 Newton, A. M. W., Knutz, P. C., Huuse, M., Gannon, P., Brocklehurst, S. H., Clausen, O. R., and Gong, Y.: Ice
594 stream reorganization and glacial retreat on the northwest Greenland shelf, *Geophysical Research Letters*,
595 44, 7826-7835, 2017.
596 Nielsen, T. and Kuijpers, A.: Only 5 southern Greenland shelf edge glaciations since the early Pliocene,
597 *Scientific reports*, 3, 1875-1875, 2013.
598 Nishiizumi, K., Imamura, M., Caffee, M. W., Southon, J. R., Finkel, R. C., and McAninch, J.: Absolute
599 calibration of Be-10 AMS standards, *Nucl Instrum Meth B*, 258, 403-413, 2007.
600 O'Cofaigh, C., Dowdeswell, J. A., Jennings, A. E., Hogan, K. A., Kilfeather, A., Hiemstra, J. F., Noormets, R.,
601 Evans, J., McCarthy, D. J., Andrews, J. T., Lloyd, J. M., and Moros, M.: An extensive and dynamic ice sheet on
602 the West Greenland shelf during the last glacial cycle, *Geology*, 41, 219-222, 2013.
603 Oksman, M., Weckstrom, K., Miettinen, A., Juggins, S., Divine, D. V., Jackson, R., Telford, R., Korsgaard, N. J.,
604 and Kucera, M.: Younger Dryas ice margin retreat triggered by ocean surface warming in central-eastern
605 Baffin Bay, *Nat Commun*, 8, 2017.
606 Olsen, J., Tikhomirov, D., Grosen, C., Heinemeier, J., and Klein, M.: Radiocarbon Analysis on the New
607 AARAMS 1MV Tandetron, *Radiocarbon*, 59, 905-913, 2016.
608 Rainsley, E., Menviel, L., Fogwill, C. J., Turney, C. S. M., Hughes, A. L. C., and Rood, D. H.: Greenland ice mass
609 loss during the Younger Dryas driven by Atlantic Meridional Overturning Circulation feedbacks, *Scientific
610 Reports*, 8, 11307, 2018.
611 Reimer, P. J., Bard, E., Bayliss, J. W., Beck, P. G., Blackwell, C., Bronk Ramsey, C. E., Buck, H., Cheng, R. L.,
612 Edwards, M., Friedrich, P. M., Grootes, T. P., Guilderson, H., Hafliðason, I., Hajdas, C., Hatté, T. J., Heaton, D.
613 L., Hoffmann, A. G., Hogg, K. A., Hughen, K. F., Kaiser, B., Kromer, S. W., Manning, M., Niu, R. W., Reimer, D.
614 A., Richards, E. M., Scott, J. R., Southon, R. A., Staff, C., Turney, S. M., and van der Plicht, J.: IntCal13 and
615 Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP *Radiocarbon*, 55, 1869-1887, 2013.
616 Rinterknecht, V., Jomelli, V., Brunstein, D., Favier, V., Masson-Delmotte, V., Bourles, D., Leanni, L., and
617 Schlappy, R.: Unstable ice stream in Greenland during the Younger Dryas cold event, *Geology*, 42, 759-762,
618 2014.

- 619 Roberts, D. H., Long, A. J., Schnabel, C., Davies, B. J., Xu, S., Simpson, M. J. R., and Huybrechts, P.: Ice sheet
620 extent and early deglacial history of the southwestern sector of the Greenland Ice Sheet, *Quaternary Sci*
621 *Rev*, 28, 2760-2773, 2009.
- 622 Roberts, D. H., Rea, B. R., Lane, T. P., Schnabel, C., and Rodes, A.: New constraints on Greenland ice sheet
623 dynamics during the last glacial cycle: Evidence from the Uummannaq ice stream system, *J Geophys Res-*
624 *Earth*, 118, 519-541, 2013.
- 625 Roksandic, M. M.: Geology of the continental shelf off West Greenland between 61 15'N and 64 00'N,
626 Geological Survey of Greenland, Reports, 92, 1-15, 1979.
- 627 Ryan, J. C., Dowdeswell, J. A., and Hogan, K. A.: Three cross-shelf troughs on the continental shelf of SW
628 Greenland from Olex data, *Atlas of Submarine Glacial Landforms: Modern, Quaternary and Ancient*, 46,
629 167-168, 2016.
- 630 Seidenkrantz, M. S., Kuijpers, A., Olsen, J., Pearce, C., Lindblom, S., Ploug, J., Przybylo, P., and Snowball, I.:
631 Southwest Greenland shelf glaciation during MIS 4 more extensive than during the Last Glacial Maximum,
632 *Sci Rep*, 9, 15617, 2019.
- 633 Sheldon, C., Jennings, A., Andrews, J. T., Cofaigh, C. O., Hogan, K., Dowdeswell, J. A., and Seidenkrantz, M.
634 S.: Ice stream retreat following the LGM and onset of the west Greenland current in Uummannaq Trough,
635 west Greenland, *Quaternary Sci Rev*, 147, 27-46, 2016.
- 636 Simpson, M. J. R., Milne, G. A., Huybrechts, P., and Long, A. J.: Calibrating a glaciological model of the
637 Greenland ice sheet from the Last Glacial Maximum to present-day using field observations of relative sea
638 level and ice extent, *Quaternary Sci Rev*, 28, 1631-1657, 2009.
- 639 Sinclair, G., Carlson, A. E., Mix, A. C., Lecavalier, B. S., Milne, G., Mathias, A., Buizert, C., and DeConto, R.:
640 Diachronous retreat of the Greenland ice sheet during the last deglaciation, *Quaternary Sci Rev*, 145, 243-
641 258, 2016.
- 642 Slabon, P., Dorschel, B., Jokat, W., Myklebust, R., Hebbeln, D., and Gebhardt, C.: Greenland ice sheet retreat
643 history in the northeast Baffin Bay based on high-resolution bathymetry, *Quaternary Sci Rev*, 154, 182-198,
644 2016.
- 645 Sommerhoff, G.: Geomorphologische Prozesse in der Labrador - und Imingersee. Ein Beitrag zur
646 submarinen Geomorphologie einer subpolaren Meeresregion, *Polarforschung*, 51, 175-191, 1981.
- 647 Sommerhoff, G.: Versuch einer geomorphologischen Gliederung des südwestgrönländischen
648 Kontinentalabhangs, *Polarforschung*, 45, 87-101, 1975.
- 649 Steffensen, J. P., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Fischer, H., Goto-Azuma, K.,
650 Hansson, M., Johnsen, S. J., Jouzel, J., Masson-Delmotte, V., Popp, T., Rasmussen, S. O., Rothlisberger, R.,
651 Ruth, U., Stauffer, B., Siggaard-Andersen, M. L., Sveinbjornsdottir, A. E., Svensson, A., and White, J. W. C.:
652 High-resolution Greenland Ice Core data show abrupt climate change happens in few years, *Science*, 321,
653 680-684, 2008.
- 654 Stone, J. O.: Air pressure and cosmogenic isotope production, *Journal of Geophysical Research* 105 (B10),
655 23753-23759, 2000.
- 656 Vacco, D. A., Alley, R. B., and Pollard, D.: Modeling dependence of moraine deposition on climate history:
657 the effect of seasonality, *Quaternary Sci Rev*, 28, 639-646, 2009.
- 658 van Tatenhove, F. G. M., van der Meer, J. J. M., and Koster, E. A.: Implications for deglaciation chronology
659 from new AMS age determinations in central west Greenland, *Quaternary Research*, 45, 245-253, 1996.
- 660 Vasskog, K., Langebroek, P. M., Andrews, J. T., Nilsen, J. E. O., and Nesje, A.: The Greenland Ice Sheet during
661 the last glacial cycle: Current ice loss and contribution to sea-level rise from a palaeoclimatic perspective,
662 *Earth-Science Reviews*, 150, 45-67, 2015.
- 663 Weidick, A.: C-14 datings of survey material, carried out in 1975, *Rapport Grønlands Geologiske*
664 *Undersøgelse*, 80, 136-144, 1976a.
- 665 Weidick, A.: Observations on the Quaternary geology in the Fiskeneset area during the summer of 1973,
666 *Rapport Greenlands Geologiske Undersøgelse*, 73, 96-99, 1976b.
- 667 Weidick, A., Kelly, M., and Bennike, O.: Late Quaternary development of the southern sector of the
668 Greenland Ice Sheet, with particular reference to the Qassimiut lobe, *Boreas*, 33, 284-299, 2004.

- 669 Winsor, K., Carlson, A. E., Caffee, M. W., and Rood, D. H.: Rapid last-deglacial thinning and retreat of the
670 marine-terminating southwestern Greenland ice sheet, *Earth Planet Sc Lett*, 426, 1-12, 2015b.
- 671 Winsor, K., Carlson, A. E., Welke, B. M., and Reilly, B.: Early deglacial onset of southwestern Greenland ice-
672 sheet retreat on the continental shelf, *Quaternary Sci Rev*, 128, 117-126, 2015a.
- 673 Woodroffe, S. A., Long, A. J., Lecavalier, B. S., Milne, G. A., and Bryant, C. L.: Using relative sea-level data to
674 constrain the deglacial and Holocene history of southern Greenland, *Quaternary Sci Rev*, 92, 345-356, 2014.
- 675 Young, N. E., Briner, J. P., Miller, G. H., Lesnek, A. J., Crump, S. E., Thomas, E. K., Pendleton, S. L., Cuzzone, J.,
676 Lamp, J., Zimmerman, S., Caffee, M., and Schaefer, J. M.: Deglaciation of the Greenland and Laurentide ice
677 sheets interrupted by glacier advance during abrupt coolings, *Quaternary Sci Rev*, 229, 106091, 2020.
- 678 Young, N. E., Schaefer, J. M., Briner, J., and . A ^{10}Be production-rate calibration for the Arctic, *Journal of*
679 *Quaternary Science* 28, 516-526, 2013.

680

681