

A data-model approach to interpreting speleothem oxygen isotope records from monsoon regions on orbital timescales

Sarah E. Parker¹, Sandy P. Harrison¹, Laia Comas-Bru¹, Nikita Kaushal², Allegra N. LeGrande³, Martin Werner⁴

5 ¹School of Archaeology, Geography and Environmental Science, Reading University, Reading, UK

²Asian School of the Environment, Nanyang Technological University, Singapore

³NASA Goddard Institute for Space Studies and Center for Climate Systems Research, Columbia University, New York, USA

⁴Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

Correspondence to: Sarah E. Parker (s.parker@pgr.reading.ac.uk)

10 **Abstract.** Reconstruction of past changes in monsoon climate from speleothem oxygen isotope ($\delta^{18}\text{O}$) records is complex because $\delta^{18}\text{O}$ signals can be influenced by multiple factors including changes in precipitation, precipitation recycling over land, temperature at the moisture source and changes in the moisture source region and transport pathway. Here, we analyse >150 speleothem records from version 2 of the Speleothem Isotopes Synthesis and Analysis (SISAL) database to produce composite regional trends in $\delta^{18}\text{O}$ in monsoon regions; compositing minimises the influence of site-specific karst and cave processes that can influence individual site records. We compare speleothem $\delta^{18}\text{O}$ observations with isotope-enabled climate model simulations to investigate the specific climatic factors causing these regional trends. We focus on differences in $\delta^{18}\text{O}$ signals between interglacial (mid-Holocene and Last Interglacial) and glacial (Last Glacial Maximum) states, and on $\delta^{18}\text{O}$ evolution through the Holocene. Differences in speleothem $\delta^{18}\text{O}$ between the mid-Holocene and Last Interglacial in the East Asian and Indian monsoons are small, despite the larger summer insolation values during the Last Interglacial. Last Glacial Maximum $\delta^{18}\text{O}$ values are significantly less negative than interglacial values. Comparison with simulated glacial-interglacial $\delta^{18}\text{O}$ shows that changes are principally driven by global shifts in temperature and regional precipitation. Holocene speleothem $\delta^{18}\text{O}$ records show distinct and coherent regional trends. Trends are similar to summer insolation in India, China and southwestern South America, but different in the Indonesian-Australian region. Redundancy analysis shows that 37% of Holocene variability can be accounted for by latitude and longitude, supporting the differentiation of records into individual monsoon regions. Regression analysis of simulated precipitation $\delta^{18}\text{O}$ and climate variables show that global Holocene monsoon $\delta^{18}\text{O}$ trends are driven by changes in precipitation, atmospheric circulation and (to a lesser extent) source area temperature, whilst precipitation recycling is non-significant. However, there are differences in regional scale mechanisms; there are clear relationships between changes in precipitation and in $\delta^{18}\text{O}$ for India, southwestern South America and the Indonesian-Australian regions, but not for the East Asian monsoon. Changes in atmospheric circulation contributes to $\delta^{18}\text{O}$ trends in the East Asian, Indian and Indonesian-Australian monsoons, and a weak source area temperature effect is observed over southern and central America and Asia. Precipitation recycling is influential in southwestern South America and southern

15
20
25
30

Africa. Overall, our analyses show that it is possible to differentiate the impacts of specific climatic mechanisms influencing precipitation $\delta^{18}\text{O}$ and use this analysis to interpret changes in speleothem $\delta^{18}\text{O}$.

1 Introduction

35 The oxygen isotopic ($\delta^{18}\text{O}$: $^{18}\text{O}/^{16}\text{O}$ ratio relative to a standard, in permil, ‰) composition of speleothems is widely used to infer past regional climates (Bar-Matthews et al., 1997; McDermott, 2004; Wang et al., 2008). Speleothem oxygen isotope ($\delta^{18}\text{O}_{\text{spel}}$) signals are inherited from $\delta^{18}\text{O}$ in precipitation ($\delta^{18}\text{O}_{\text{precip}}$) above the cave, which in turn is determined by the initial $\delta^{18}\text{O}$ of water vapour as it evaporates at the oceanic moisture source region, the degree of rainout and evaporation from source to cave site and air temperature changes encountered throughout the moisture transport pathway (Fairchild and Baker, 2012; 40 Lachniet, 2009). Understanding the effects and contribution of each of these climate processes to $\delta^{18}\text{O}_{\text{precip}}$ (and therefore $\delta^{18}\text{O}_{\text{spel}}$) is essential to inferring palaeoclimate from speleothem $\delta^{18}\text{O}$ records.

Initial $\delta^{18}\text{O}$ is determined by oceanic $\delta^{18}\text{O}$ at the evaporative moisture source region (Craig and Gordon, 1965), which varies spatially (LeGrande and Schmidt, 2006) and through time (e.g. Waelbroeck et al., 2002). During evaporation from the moisture source, ^{16}O is preferentially incorporated into the vapour, whilst subsequent fractionation during atmospheric transport occurs 45 by Rayleigh distillation; As air masses cool and moisture condenses, heavier ^{18}O is enriched in the liquid phase and removed by precipitation. With progressive rainout along a moisture pathway, precipitation becomes gradually more depleted (Dansgaard, 1964). Within this framework, $\delta^{18}\text{O}_{\text{precip}}$ is controlled by two variables: temperature and the amount of precipitation along a moisture pathway. The temperature effect stems from the cooling required for progressive rainout during Rayleigh distillation (Dansgaard, 1964; Rozanski et al., 1993). The temperature- $\delta^{18}\text{O}$ impact is dominant at mid to high 50 latitudes, whilst observations suggest that changes in upstream and local precipitation dominate changes in the $\delta^{18}\text{O}_{\text{precip}}$ signal at tropical latitudes. The negative relationship between local precipitation and $\delta^{18}\text{O}_{\text{precip}}$, often referred to as the “amount effect” (Bailey et al., 2018; Dansgaard, 1964), results from the re-evaporation and diffusive exchange between precipitation and water vapour during deep convective precipitation (Risi et al., 2008). However, Rayleigh distillation is complicated by changes in atmospheric circulation and moisture recycling. Changes in the area from which the moisture is sourced will modify $\delta^{18}\text{O}_{\text{precip}}$ 55 because the initial $\delta^{18}\text{O}$ values differ between sources (Cole et al., 1999; Friedman et al., 2002), whilst changes in the moisture transport pathway and/or distance between source and cave site can result in differing degrees of fractionation associated with condensation and evaporation (Aggarwal et al., 2012; Bailey et al., 2018). The isotopic composition of atmospheric water vapour may also be modified by precipitation recycling over land, since evapotranspiration returns moisture from precipitation back to the atmosphere thereby reducing the $\delta^{18}\text{O}_{\text{precip}}$ /distance gradient along an advection path that occurs with Rayleigh 60 distillation (Gat, 1996; Salati et al., 1979).

Speleothem $\delta^{18}\text{O}$ records from monsoon regions show multi-millennial variability that has been interpreted as documenting the waxing and waning of the monsoons in response to changes in summer insolation, manifested as either a change in the

absolute amount of precipitation (Cai et al., 2012; Cheng et al., 2006) or a change in the ratio of more negative $\delta^{18}\text{O}$ summer precipitation to less negative $\delta^{18}\text{O}$ winter precipitation (Cruz et al., 2005; Dong et al., 2010; Wang et al., 2001). However, the multiplicity of processes that influence $\delta^{18}\text{O}$ before incorporation in the speleothem make it difficult to attribute the climatic causes of changes in individual speleothem records unambiguously. In the East Asian monsoon, for example, speleothem $\delta^{18}\text{O}$ records have been interpreted as a summer monsoon signal, manifested either as a change in the amount of water vapour removed along the moisture trajectory (Yuan et al., 2004), and/or as a change in the contribution of summer precipitation to annual totals (Cheng et al., 2006, 2009, 2016; Wang et al., 2001) based on the relationship between modern $\delta^{18}\text{O}_{\text{precip}}$ and climate. Other interpretations of Chinese monsoon $\delta^{18}\text{O}_{\text{spel}}$ have included rainfall source changes (Tan 2009, 2011, 2014) or local rainfall changes (Cai et al., 2010; Tan et al., 2015). Maher (2008) interpreted $\delta^{18}\text{O}_{\text{spel}}$ as reflecting changes in moisture source area, based on differences between $\delta^{18}\text{O}_{\text{spel}}$ and loess/palaeosol records of rainfall and the strong correlation between East Asian and Indian monsoon speleothems. Maher and Thompson (2012) used a mass balance approach to show that the changes in precipitation (either local or upstream) or rainfall seasonality required to reproduce $\delta^{18}\text{O}_{\text{spel}}$ trends would be unreasonably large. They therefore argued that changes in moisture source were required to explain shifts in $\delta^{18}\text{O}$ both on glacial/interglacial time scales and during interglacials. There are also multiple interpretations of the causes of $\delta^{18}\text{O}_{\text{spel}}$ variability in other monsoon regions. In the Indonesian-Australian monsoon region, for example, $\delta^{18}\text{O}_{\text{spel}}$ variability has been interpreted as a precipitation amount signal (Carolin et al., 2016; Krause et al., 2019) or a precipitation seasonality signal (Ayliffe et al., 2013; Griffiths et al., 2009), based on modern $\delta^{18}\text{O}_{\text{precip}}$ and climate observations (Cobb et al., 2007; Moerman et al., 2013), and/or as a moisture source/trajectory signal (Griffiths et al., 2009; Wurtzel et al., 2018). South American speleothem records have been interpreted as records of monsoon intensity, due to changes in the amount of precipitation over the region (Cruz et al., 2006; Wang et al., 2006; Cheng et al., 2013), changes in the degree of upstream precipitation and evapotranspiration (Cheng et al., 2013) or changes in the ratio of precipitation sourced from the low-level jet versus the Atlantic (Cruz et al., 2005; Wang et al., 2006). In the Indian monsoon region, speleothem $\delta^{18}\text{O}$ records are interpreted primarily as an amount effect signal (Berkelhammer et al., 2010; Fleitmann et al., 2004), supported by $\delta^{18}\text{O}_{\text{precip}}$ /climate observations (e.g. Battacharya et al., 2003). However, other studies have suggested that $\delta^{18}\text{O}_{\text{precip}}$ changes in this region are driven primarily by large-scale changes in monsoon circulation and hence, Indian monsoon $\delta^{18}\text{O}_{\text{spel}}$ should be interpreted as a moisture source/trajectory signal (Breitenbach et al., 2010; Sinha et al., 2015).

These interpretations generally rely on modern $\delta^{18}\text{O}_{\text{precip}}$ -climate observations, which may not have remained constant through time. The sources of $\delta^{18}\text{O}$ variability can also be explored using isotope-enabled climate models (e.g. Hu et al., 2019), which incorporate known isotope effects and therefore provide plausible explanations for $\delta^{18}\text{O}_{\text{spel}}$ trends. Modelling studies suggest that changes in East Asian monsoon $\delta^{18}\text{O}_{\text{precip}}$ (during Heinrich events: Lewis et al., 2010; Pausata et al., 2010, and on orbital timescale: Battisti et al., 2014; LeGrande and Schmidt, 2009) do not reflect local rainfall variability but instead reflect changes in $\delta^{18}\text{O}$ of vapour delivered to the region. Variability in the $\delta^{18}\text{O}$ of vapour delivered to East Asia on orbital timescales has been diagnosed as due to changes in precipitation upstream of the region (Battisti et al., 2014), changes in moisture source

location (Hu et al., 2019; Tabor et al., 2018) or changes in the strength of monsoon winds (LeGrande and Schmidt, 2009; Liu et al., 2014). $\delta^{18}\text{O}_{\text{precip}}$ variability in the East Asian monsoon during Heinrich events has also been attributed to non-local isotope fractionation (Lewis et al., 2010; Pausata et al., 2011). Modelling results suggest that changes in precipitation amount are the predominant source of $\delta^{18}\text{O}$ variability in the Indian monsoon during the Holocene (LeGrande and Schmidt, 2009) and in the glacial (Lewis et al., 2010), and in the South American and Indonesian/Australian regions during Heinrich events (Lewis et al., 2010) and the Last Interglacial (Sjolte and Hoffman, 2014).

In this study, we combine speleothem $\delta^{18}\text{O}$ records from version 2 of the Speleothem Isotopes Synthesis and Analysis (SISAL) database with isotope-enabled palaeoclimate simulations from two climate models to investigate the plausible mechanisms driving changes in $\delta^{18}\text{O}$ in monsoon regions through the Holocene (last 11,700 years) and between interglacial (mid-Holocene and Last Interglacial) and glacial (Last Glacial Maximum) states. We compare $\delta^{18}\text{O}_{\text{spel}}$ signals across geographically separated cave sites to extract a regional signal, thus minimising the influence of karst and in-cave processes, such as the mixing of groundwaters from different precipitation events or changes in cave ventilation, that can be important for the $\delta^{18}\text{O}_{\text{spel}}$ of individual records. We use Principal Coordinate Analysis (PCoA) to identify regions with geographically coherent $\delta^{18}\text{O}_{\text{spel}}$ records, and then examine how these regions behave on glacial-interglacial time scales and through the Holocene. We use isotope-enabled model simulations, to investigate the main drivers of $\delta^{18}\text{O}_{\text{spel}}$ variability in regions where the models reproduce the large-scale $\delta^{18}\text{O}$ changes shown by observations. We exploit the fact that models produce internally physically consistent changes to explore potential and plausible causes of the trends observed in speleothem records across specific monsoon regions, using multiple regression analysis.

2. Methods

2.1 Speleothem oxygen isotope data

Speleothem $\delta^{18}\text{O}$ records were obtained from the SISAL (Speleothem Isotopes Synthesis and Analysis) database (Atsawaranunt et al., 2018; Comas-Bru et al., 2020a, 2020b). Records were selected based on the following criteria:

- They are located in monsoon regions, between 35°S and 40°N;
- The mineralogy is known and does not vary (i.e. between calcite and aragonite) through time, because oxygen isotope fractionation during speleothem precipitation is different for calcite and aragonite;
- For the interglacial-glacial analyses, the records contain samples in the periods used for the model simulations: mid-Holocene (MH, 6,000±500 years BP, 6ka), Last Glacial Maximum (LGM, 21,000±1,000 years BP, 21ka) or the Last Interglacial (LIG, 125,000±1,000 years BP, 125 ka), where BP (before present) is 1950 CE;
- For the PCoA, the records have a temporal coverage of at least 4,000 years in the Holocene;
- For Holocene trend analyses, speleothems have a record of the period from 7,000 to 3,000 years BP;
- They are the most recent update of the record from a site available in version 2 of the SISAL database.

This resulted in the selection of 125 records from 44 sites for the PCoA analysis, 64 records from 38 sites for the analysis of MH, LGM and LIG signals and 79 records from 40 sites for the Holocene trend analysis (Fig. 1). Although the SISALv2 database contains multiple age models for some sites, we use the published age models given by the original authors for all records.

2.2 Climate model simulations

There are relatively few paleoclimate simulations made with models that incorporate oxygen isotope tracers, and the available simulations do not necessarily focus on the same periods or use the same modelling protocols. Here, we use simulations of opportunity from two isotope-enabled climate models: ECHAM5 (version 5 of the European Centre for medium range weather forecasting model in HAMBURG) and GISS E-R (Goddard Institute for Space Studies Model version E-R). The ECHAM5 simulations provide an opportunity to examine large-scale changes between glacial and interglacial states, using simulations of the MH, LGM and LIG. The GISS Model E-R Ocean-Atmosphere Coupled General Circulation Model was used to investigate the evolution of $\delta^{18}\text{O}$ evolution during the Holocene, using eight time slices (9 ka, 6 ka, 5 ka, 4 ka, 3 ka, 2 ka, 1 ka and 0 ka) experiments. Although simulations of the MH 6ka time slice are available with both models, there are differences in the protocol used for the two experiments which preclude direct comparison of the simulations.

The ECHAM5-wiso MH experiment (Wackerbarth et al., 2012; Werner, 2019) was forced by orbital parameters (based on Berger and Loutre, 1991) and greenhouse gas (GHG) concentrations ($\text{CO}_2 = 280$ ppm, $\text{CH}_4 = 650$ ppb, $\text{N}_2\text{O} = 270$ ppb) appropriate to 6 ka. Changes in sea-surface temperature (SST) and sea-ice were derived from a transient Holocene simulation (Varma et al., 2012). The control simulation for the MH experiment was an ECHAM-wiso simulation of the period 1956-1999 (Langebroek et al., 2011), using observed SSTs and sea-ice cover. This control experiment was forced by SSTs and sea-ice only, with atmospheric circulation free to evolve. The ECHAM5-wiso LGM experiment (Werner, 2019; Werner et al., 2018) was forced by orbital parameters (Berger and Loutre, 1991), GHG concentrations ($\text{CO}_2 = 185$ ppm, $\text{CH}_4 = 350$ ppb, $\text{N}_2\text{O} = 200$ ppb), land-sea distribution and ice sheet height and extent appropriate to 21 ka; SST and sea-ice cover were prescribed from the GLAMAP dataset (Schäfer-Neth and Paul, 2003). Sea surface water and sea-ice $\delta^{18}\text{O}$ were uniformly enriched by 1 % at the start of the experiment. The control simulation for the LGM experiment used present-day conditions, including orbital parameters and GHG concentrations set to modern values, and SSTs and sea-ice cover from the last 20 years (1979-1999). Both the MH and LGM simulations were run at T106 horizontal grid resolution, approximately 1.1° by 1.1° . Comparison of the MH and LGM simulations with speleothem data globally (Comas-Bru et al., 2019; Fig. S1 and S2) show that the ECHAM model reproduces the broadscale spatial gradients and the sign of isotopic changes at the majority of cave sites (MH: 72%; LGM: 76%). However, the changes compared to present are generally more muted in the simulations than shown by the speleothem records.

The LIG experiment (Gierz et al., 2017b, 2017a) was run using the ECHAM5/MPI-OM Earth System Model, with stable water isotope diagnostics included in the ECHAM5 atmosphere model (Werner et al., 2011), the dynamic vegetation model JSBACH

(Haese et al., 2012) and the MPI-OM ocean/sea-ice module (Xu et al., 2012). This simulation was run at a T31L19 horizontal
160 grid resolution, approximately 3.75° by 3.75°. The LIG simulation was forced by orbital parameters derived from Berger and
Loutre (1991) and GHG concentrations (CO₂ = 276 ppm, CH₄ = 640 ppb, N₂O = 263 ppb) appropriate to 125 ka, but it was
assumed that ice sheet configuration and land-sea geography is unchanged from modern and therefore no change was made to
the isotopic composition of sea water. The LIG simulation is compared to a pre-industrial (PI) control with appropriate
insolation, GHG and ice sheet forcing for 1850 CE. The sign of simulated isotopic changes in the LIG is in good agreement
165 with ice core records from Antarctica and Greenland and speleothem records from Europe, the Middle East and China (Gierz
et al., 2017b) although, as with the MH and LGM, the observed changes tend to be larger than the simulated changes (Fig. S3).

There are GISS ModelE-R (LeGrande and Schmidt, 2009) simulations for eight time slices during the Holocene (9 ka, 6 ka, 5
ka, 4 ka, 3 ka, 2 ka, 1 ka and 0 ka). The 0 ka experiment is considered as the pre-industrial control (ca 1880 CE). Orbital
parameters were based on Berger and Loutre (1991) and GHG concentrations were adjusted based on ice core reconstructions
170 (Brook et al., 2000; Indermühle et al., 1999; Sowers, 2003) for each time slice. A remnant Laurentide ice sheet was included
in the 9 ka simulation, following Licciardi et al. (1998), and the corresponding adjustment was made to mean ocean salinity
and ocean water δ¹⁸O to account for this (Carlson et al., 2008). The ice sheet in all the other experiments was specified to be
the same as modern, and therefore no adjustment was necessary. The simulations were run using the M20 version of GISS
ModelE-R, which has a horizontal resolution of 4° by 5°. Each experiment was run for 500 years and we use the last 100
175 simulated years for the analyses. Comparison of the simulated trends in δ¹⁸O show good agreement with Greenland ice core
records, marine records from the tropical Pacific and Chinese speleothem records (LeGrande and Schmidt, 2009). However,
as is the case with the ECHAM simulations, the model tends to produce changes less extreme than shown by the observations
(Fig S4, S5 and S6).

2.3 Principal Coordinate Analysis and Redundancy Analysis

180 We used PCoA to identify regionally coherent patterns in the speleothem δ¹⁸O records for the Holocene. PCoA is a multivariate
ordination technique that uses a distance/dissimilarity matrix to represent inter-object (dis)similarity in reduced space (Gower,
1966; Legendre and Legendre, 1998). Speleothem records from individual sites are often discontinuous; missing data is
problematic for many ordination techniques. PCoA is more robust to missing data than other methods (Kärkkäinen and
Saarela, 2015; Rohlf, 1972). We used a correlation matrix of speleothem records as the (dis)similarity measure. The temporal
185 resolution of speleothem records was first standardised by calculating a running average mean with non-overlapping 500-year
windows. This procedure produces a single composite record when there are several records for a given site. PCoA results
were displayed as a biplot, where sites ordinated close to one another (i.e., with similar PCoA scores) show similar Holocene
trends and sites ordinated far apart have dissimilar trends. We used the ‘broken stick’ model (Bennett, 1996) to identify which
PCoA axes were significant. We used redundancy analysis (RDA: Legendre and Legendre, 1998; Rao, 1964) with latitude and
190 longitude as predictor variables to identify if PCoA (dis)similarities were related to geographical location, and Principal

Components Analysis (PCA) to identify the main patterns of variation. As these explanatory variables are not dimensionally homogeneous, they were centred on their means and standardised to allow direct comparison of the gradients. PCoA and RDA analyses were carried out using the ‘vegan’ package in R (Oksanen et al., 2019).

2.4 Glacial-interglacial changes in $\delta^{18}\text{O}$

195 We examined glacial-interglacial shifts in $^{18}\text{O}_{\text{spel}}$ observations and in annual precipitation-weighted mean $\delta^{18}\text{O}_{\text{precip}}$ from ECHAM-wiso in regions influenced by the monsoon. We focus on regional differences between MH, LGM and LIG with respect to the present-day for speleothems or the control simulation experiment for model outputs. Comas-Bru et al. (2019) have shown that differences in speleothem $\delta^{18}\text{O}$ data between the 20th century and the pre-industrial period (i.e. 1850±15 CE) are within the temporal and measurement uncertainties of the data, and thus the use of different reference periods (i.e. PI for
200 the ECHAM LIG experiment, 20th century for ECHAM MH, LGM experiments) should have little effect on our analyses. We used mean site $\delta^{18}\text{O}_{\text{spel}}$ values for each period for the regions identified in the PCoA analysis. Where there are multiple speleothem $\delta^{18}\text{O}$ records for a site in a time period, they were averaged to calculate mean $\delta^{18}\text{O}_{\text{spel}}$. Three sites above 3500m were excluded from the calculation of the means because high elevation sites have more negative $\delta^{18}\text{O}$ values than their low-elevation counterparts and their inclusion would distort the regional estimates.

205 There are relatively few speleothems covering both the present-day and the period of interest (i.e., MH, LGM or LIG), precluding the calculation of $\delta^{18}\text{O}_{\text{spel}}$ anomalies from the speleothem data. We therefore calculated anomalies with respect to modern (1960-2017 CE) using as reference the Online Isotopes in Precipitation Calculator (OIPC: Bowen, 2018; Bowen and Revenaugh, 2003), a global gridded dataset of interpolated mean annual precipitation-weighted $\delta^{18}\text{O}_{\text{precip}}$ data. This dataset combines data from 348 stations from the Global Network of Isotopes in Precipitation (IAEA/WMO, 2018), covering part or
210 all of the period 1960-2014, and other records available at the Water Isotopes Database (Waterisotopes Database, 2017).

OIPC $\delta^{18}\text{O}_{\text{precip}}$ was converted to its speleothem equivalent assuming that: (i) precipitation-weighted mean annual $\delta^{18}\text{O}_{\text{precip}}$ is equivalent to mean annual drip-water $\delta^{18}\text{O}$ (Yonge et al., 1985) and (ii) precipitation of calcite is consistent with the empirical speleothem-based kinetic fractionation factor of Tremaine et al. (2011) and precipitation of aragonite follows the fractionation factor from Grossman and Ku (1986), as formulated by Lachniet (2015):

$$215 \quad \delta^{18}\text{O}_{\text{calcite_SMOW}} = w\delta^{18}\text{O}_{\text{precip_SMOW}} + \left(\left(\frac{16.1 \cdot 1000}{T} \right) - 24.6 \right) \quad (\text{T in K}) \quad (1)$$

$$\delta^{18}\text{O}_{\text{aragonite_SMOW}} = w\delta^{18}\text{O}_{\text{precip_SMOW}} + \left(\left(\frac{18.34 \cdot 1000}{T} \right) - 31.954 \right) \quad (\text{T in K}) \quad (2)$$

where $\delta^{18}\text{O}_{\text{calcite_SMOW}}$ and $\delta^{18}\text{O}_{\text{aragonite_SMOW}}$ are the speleothem isotopic composition for calcite and aragonite speleothems with reference to the V-SMOW standard (in permil); $w\delta^{18}\text{O}_{\text{precip}}$ is the OIPC precipitation-weighted annual mean isotopic
220 composition of precipitation with respect to the V-SMOW standard and T is the mean annual cave temperature (in degrees Kelvin). We used the long-term (1960-2016) mean annual surface air temperature from the CRU-TS4.01 database (Harris et

al., 2014) at each site as a surrogate for mean annual cave air temperature. The resolution of the gridded data means that $w\delta^{18}\text{O}_{\text{precip_SMOW}}$ and T may be the same for nearby sites.

We use the V-SMOW to V-PDB conversion from Coplen et al. (1983), which is independent of speleothem mineralogy:

$$225 \quad \delta^{18}\text{O}_{\text{PDB}} = 0.97001 \cdot \delta^{18}\text{O}_{\text{SMOW}} - 29.29 \quad (3)$$

where $\delta^{18}\text{O}_{\text{PDB}}$ is relative to the V-PDB standard and $\delta^{18}\text{O}_{\text{SMOW}}$ is relative to V-SMOW standard.

Average uncertainties in the speleothem age-depth models are ~50 years during the Holocene. This interval is smaller than the time windows used in this analysis, and age uncertainty is therefore expected to have a negligible impact on the results. We investigated the influence of age uncertainties on the LGM and LIG $\delta^{18}\text{O}_{\text{spel}}$ anomalies by examining the impact of using
230 different window widths (± 500 , ± 700 , ± 1000 , ± 2000 years) on the regional mean $\delta^{18}\text{O}_{\text{spel}}$ anomalies.

We used anomalies of $w\delta^{18}\text{O}_{\text{precip}}$, mean annual surface air temperature (MAT) and mean annual precipitation (MAP) from the ECHAM5-wiso simulations to investigate the drivers of glacial-interglacial $\delta^{18}\text{O}_{\text{spel}}$ variability. Values were calculated from land grid cells (>50% land) $\pm 3^\circ$ around each speleothem site. This distance was chosen with reference to the coarsest resolution simulation (LIG, ca. 3.75 x 3.75 $^\circ$). Gridded values of MAT and MAP were weighted by the proportion of each grid cell that
235 lies within $\pm 3^\circ$ of the site and linear distance-weighted means were calculated for each site and time slice. We only considered regions with at least one speleothem record for each of the three time periods, although these were not required to be the same sites, and where the observed shifts in $\delta^{18}\text{O}_{\text{spel}}$ were in the same direction and of a similar magnitude to the simulated $w\delta^{18}\text{O}_{\text{precip}}$.

2.5 Holocene and Last Interglacial regional trends

Regional speleothem $\delta^{18}\text{O}$ changes through the Holocene were examined by creating composite time-series for each region
240 identified in the PCoA analysis with at least four Holocene records (> 5000 years long). Regional composites were constructed using a 4-step procedure, modified from Marlon et al. (2008): (i) the $\delta^{18}\text{O}$ data for individual speleothems were transformed to z-scores, so all records have a standardised mean and variance:

$$z\text{-score}_i = (\delta^{18}\text{O}_i - \overline{\delta^{18}\text{O}}_{(\text{base period})}) / s\delta^{18}\text{O}_{(\text{base period})} \quad (4)$$

Where $\overline{\delta^{18}\text{O}}$ is the mean and $s\delta^{18}\text{O}$ is the standard deviation of $\delta^{18}\text{O}$ for a common base period. A base period of 7,000 to
245 3,000 years BP was chosen to maximise the number of records included in each composite. (ii) the standardised data for a site were re-sampled by applying a 100-year non-overlapping running mean with the first bin centred at 50 years BP, in order to create a single site time series while ensuring that highly resolved records do not dominate the regional composite; (iii) each regional composite was constructed using locally weighted regression (Cleveland and Devlin, 1988) with a window width of 3,000 years and fixed target points in time; and (iv) confidence intervals (5th and 95th percentiles) for each composite were
250 generated by bootstrap resampling by site over 1,000 iterations. There are too few sites to construct regional composites for the LIG and thus the trends in $\delta^{18}\text{O}_{\text{spel}}$ were examined using records from individual sites covering the period 130-116 ka BP. We calculated Holocene regional composites from annual precipitation-weighted mean $\delta^{18}\text{O}_{\text{precip}}$ anomalies simulated by the GISS model. Simulated $\delta^{18}\text{O}_{\text{precip}}$ trends were calculated using linear distance-weighted mean $\delta^{18}\text{O}_{\text{precip}}$ values from land grid

cells (>50% land) within $\pm 4^\circ$ around each site. This distance was determined by the grid resolution of the model. Regional
255 composites were then produced using bootstrap resampling in the same way as for the speleothem data. The simulated
anomalies are relative to the control run rather than the specified base period used for the speleothem-based composites, so
absolute values of simulated and observed Holocene trends are expected to differ. Preliminary analyses showed that neither
the mean values nor trends in $\delta^{18}\text{O}_{\text{precip}}$ were substantially different if the sampled area was reduced to match the sampling
used for the ECHAM-based box plot analysis, or was increased to encompass the larger regions shown in Fig. 1 and used in
260 the multiple regression analysis.

2.6 Multiple regression analysis

We investigate the drivers of regional $\delta^{18}\text{O}_{\text{precip}}$, and by extension $\delta^{18}\text{O}_{\text{spel}}$, through the Holocene using multiple linear
regression (MLR) of annual precipitation-weighted mean $\delta^{18}\text{O}_{\text{precip}}$ anomalies and climate variables from GISS model E-R.
Climate variables were chosen to represent the four potential large-scale drivers of regional changes in the speleothem $\delta^{18}\text{O}$
265 records. Specifically, we use changes in mean precipitation and precipitation recycling over the monsoon regions, and changes
in mean surface air temperature and surface wind direction over the moisture source regions. Whereas the influence of changes
in precipitation, recycling and temperature are relatively direct measures, the change in surface wind direction over the
moisture source region is used as an index of potential changes in the moisture source region and transport pathway. The
boundaries of each monsoon region (Fig. 1) were defined to include all the speleothem sites used to construct the Holocene
270 $\delta^{18}\text{O}_{\text{spel}}$ composites. Moisture source area limits (Fig. 1) were defined based on moisture tracking studies (Bin et al., 2013;
Breitenbach et al., 2010; D'Abreton and Tyson, 1996; Drumond et al., 2008, 2010; Durán-Quesada et al., 2010; Kennett et al.,
2012; Nivet et al., 2018; Wurtzel et al., 2018) and GISS simulated summer surface winds. All climate variables were extracted
for the summer months, defined as May to September (MJJAS) for northern hemisphere regions and November to March
(NDJFM) for southern hemisphere regions (Wang and Ding, 2008). Only grid cells with >50% land were used to extract
275 variables over monsoon regions and only grid cells with <50% land were used to extract variables over moisture source regions.
The inputs to the MLR for each time interval were calculated as anomalies from the control run.

Precipitation recycling was calculated as the ratio of locally sourced precipitation versus total precipitation. Although the GISS
E-R mid-Holocene experiment explicitly estimates recycling using vapour source distribution tracers (Lewis et al., 2014), this
was not done for all the Holocene time slice simulations. Therefore, we calculate a precipitation recycling index (RI), following
280 Brubaker et al. (1993):

$$RI = \frac{P_R}{P} = \frac{E}{2Q_H + E} \quad (5)$$

Where locally sourced (recycled) precipitation (P_R) is estimated using total evaporation over a region (E) and total precipitation
(P) is estimated as the sum of total evaporation and net incoming moisture flux integrated across the boundaries of the region
(Q_H). RI therefore expresses the change in the contribution of local, recycled precipitation independently of any overall change
285 in precipitation amount.

We use pseudo- R^2 to determine the goodness-of-fit for the overall MLR model, and t values (the regression coefficient divided by its standard error) to determine the strength of each relationship. Partial residual plots were used to show the relationship between each predictor variable and $\delta^{18}\text{O}_{\text{precip}}$ when the effects of the other variables are held constant.

All statistical analyses were performed in R (R Core Team, 2019) and plots were generated using ggplot (Wickham, 2016).

290 **3 Results**

3.1 Principal Coordinate Analysis and Redundancy Analysis

PCoA shows the (dis)similarity of Holocene $\delta^{18}\text{O}_{\text{spel}}$ evolution across individual records, and thus allows an objective regionalisation of these records. The first two PCoA axes are significant, according to the broken stick test, and account for 65% and 20% of $\delta^{18}\text{O}_{\text{spel}}$ variability respectively (Table 1). The PCoA scores differentiate records geographically (Fig. 2a):
295 southern hemisphere monsoon regions such as the southwestern South American Monsoon (SW-SAM) and South African Monsoon (SAfM) are characterised by low PCoA1 scores, whilst northern hemisphere monsoons such as the Indian Summer Monsoon (ISM) and the East Asian Monsoon (EAM), are characterised by higher PCoA1 scores. This indicates that regions can be differentiated based on their temporal evolution as captured by the first PCoA axis. Most southern hemisphere regions also have lower PCoA2 scores although this is not consistent over time. Speleothem records from Central America (CAM)
300 and Indonesian-Australian monsoon (IAM) have PCoA scores intermediate between the northern and southern hemisphere regions. PCoA clearly separates the South American records into a northeastern region (NE-SAM) with scores similar to other northern hemisphere monsoon regions and a southwestern region (SW-SAM), with scores similar to other southern hemisphere regions. The RDA supports a geographical control on the (dis)similarity of speleothem $\delta^{18}\text{O}$ records over the Holocene (Fig. 2b). RDA1 explains 37% of the variability and is significantly correlated with both latitude and longitude (Table 2).

305 **3.2 Regional interglacial-glacial differences**

To investigate the causes of glacial-interglacial shifts in $\delta^{18}\text{O}$, we compare simulated and observed regional $\delta^{18}\text{O}$ signals during the LIG, LGM and MH with shifts in climate variables (precipitation and temperature). Only the ISM, EAM and IAM regions have sufficient speleothem data (i.e. at least one record from every time period) to allow comparisons across the MH, LGM and LIG (Fig. 3) and have similar shifts in observed $\delta^{18}\text{O}_{\text{spel}}$ and simulated $\delta^{18}\text{O}_{\text{precip}}$. The regional mean $\delta^{18}\text{O}_{\text{spel}}$ anomalies
310 calculated for different time windows (± 500 , ± 700 , ± 1000 , ± 2000 years) vary by less than 0.35‰ for the LGM (ISM: $<0.16\text{‰}$, EAM: $<0.35\text{‰}$, IAM: $<0.22\text{‰}$) and 0.48 ‰ for the LIG (ISM: $<0.16\text{‰}$, EAM: $<0.48\text{‰}$, IAM: $<0.11\text{‰}$), indicating that age uncertainties have a minimal impact on these mean values. The most positive $\delta^{18}\text{O}_{\text{spel}}$ anomalies in all three regions occur at the LGM, with more negative anomalies for the MH and LIG. The simulated $\delta^{18}\text{O}_{\text{precip}}$ anomalies show a similar pattern, more positive anomalies during the LGM than during the MH or the LIG. Although the mean values of the simulated
315 $\delta^{18}\text{O}_{\text{precip}}$ anomalies differ from the observed $\delta^{18}\text{O}_{\text{spel}}$ anomalies, they are consistent with the observations within the range of their uncertainties. The differences in regional $\delta^{18}\text{O}_{\text{spel}}$ anomalies between MH and LIG differ across the three regions. In both

the ISM and the EAM, differences in $\delta^{18}\text{O}_{\text{spel}}$ values between the MH and LIG are small (Fig. 3a, 3b), although ISM LIG $\delta^{18}\text{O}_{\text{spel}}$ values are slightly more negative than MH values. In the IAM, MH values are less negative than the LIG (Fig. 3c). However, there are only a limited number of speleothem records from the ISM and IAM during the LIG, so the apparent differences between the two intervals in these regions may not be meaningful. The glacial-interglacial changes in $\delta^{18}\text{O}_{\text{precip}}$ are consistent with the simulated temperature and precipitation changes, with warmer and wetter conditions during interglacials and cooler and drier conditions during the LGM in all three regions. Differences in simulated precipitation between the MH and the LIG could help explain the differences between $\delta^{18}\text{O}_{\text{spel}}$ in the ISM and IAM, since the LIG is wetter than the MH in the ISM and drier than the MH in the IAM. However, the LIG is also drier than the MH in the EAM, a feature that appears inconsistent with the lack of differentiation between the $\delta^{18}\text{O}$ signals in this region.

3.3 Regional-scale interglacial $\delta^{18}\text{O}$ evolution

There are four monsoon regions with sufficient data to examine regional Holocene $\delta^{18}\text{O}$ trends: EAM, ISM, SW-SAM and IAM (Fig. 4). The IAM region has the fewest records ($n=7$) whilst the EAM has the largest number ($n=14$). The regional composites are expressed as z-scores, i.e. changes with respect to the mean and variance of $\delta^{18}\text{O}$ for the base period (3000-7000 yr BP). The confidence intervals on the regional composites are small for all regions, except SW-SAM in the early Holocene. The EAM and ISM regions (Fig. 4 a-e) show the most positive $\delta^{18}\text{O}_{\text{spel}}$ z-scores around 12 ka followed by a rapid decrease towards their most negative values at ~ 9.5 ka and ~ 9 ka, respectively. The $\delta^{18}\text{O}_{\text{spel}}$ z-scores in the EAM are relatively constant from 9.5 to ~ 7 ka, whereas this plateau is present but less marked in the ISM. There is a gradual trend towards more positive $\delta^{18}\text{O}_{\text{spel}}$ z-scores towards the present in both regions thereafter. The SW-SAM records (Fig. 4i) have their most positive $\delta^{18}\text{O}_{\text{spel}}$ z-scores in the early Holocene with a gradual trend to more negative scores towards the present. By contrast, the IAM z-scores (Fig. 4g) are most positive at 12ka, gradually decreasing until ca 5 ka and are relatively flat thereafter.

There are insufficient data to create composite curves for the LIG, but individual records from the four regions (Fig. 5) show similar features to the Holocene trends. Records from the ISM and EAM (Fig. 5 left), for example, are characterised by an initial sharp decrease in $\delta^{18}\text{O}_{\text{spel}}$ values of about 4 ‰ between 130-129 ka and then most of the records (Dykoski et al., 2005; Kathayat et al., 2016; Wang et al., 2008) show little variability for several thousand years. Despite the fact that the Tianmen record (Cai et al., 2010, 2012) shows considerable variability between 123-127 ka, there is nevertheless a similar plateau in the average observed value before the rapid change to less negative values after 127 ka. Similar to the Holocene, the SW-SAM record (Cheng et al., 2013) shows increasingly negative $\delta^{18}\text{O}_{\text{spel}}$ values through the LIG. The trend shown for Whiterock cave (Carolin et al., 2016) also shows similar features to the IAM Holocene composite, with a gradual trend towards more negative values initially and a relatively complacent curve towards the end of the interglacial (Fig. 5 right).

3.4 Multiple regression analysis of Holocene $\delta^{18}\text{O}_{\text{precip}}$

The MLR analyses of simulated $\delta^{18}\text{O}_{\text{precip}}$ trends identify the impact of an individual climate variable on $\delta^{18}\text{O}_{\text{precip}}$ in the absence of changes in other variables. The global model for the Holocene (1 to 9ka) $\delta^{18}\text{O}_{\text{precip}}$ trends has a pseudo- R^2 of 0.80 and shows statistically significant relationships between the anomalies in $\delta^{18}\text{O}_{\text{precip}}$ and anomalies in regional precipitation, temperature and surface wind direction (Table 3). The partial residual plots (Fig. 6) show there is a strong negative relationship with regional precipitation (t value = -8.75) and a strong positive relationship (t value = 8.03) with surface wind direction over the moisture source region, an index of changes in either source area or moisture pathway. This indicates that increases in regional precipitation alone will lead to a decrease in $\delta^{18}\text{O}$ while changes in source area/moisture pathway, in the absence of changes in other variables, will lead to significant change in $\delta^{18}\text{O}$. The relationship with temperature over the moisture source region is weaker, but positive (t value = 2.05), i.e. an increase in temperature over the moisture source region will lead to an increase in $\delta^{18}\text{O}$ if there are no changes in other climate variables. Precipitation recycling is not significant in this global analysis.

There are too few data points to make regressions for individual monsoon regions, but the distribution of data points for each region in the partial residual plots (Fig. 6) is indicative of the degree of conformity to the global model. Data points from the ISM, SW-SAM, IAM and SAfM are well aligned with the global relationship with regional precipitation (Fig. 6a), indicating that precipitation is an important control on changes in $\delta^{18}\text{O}_{\text{precip}}$ in these regions. The NE-SAM, EAM and CAM values deviate somewhat from the global relationship and, although there are relatively few points, this suggests that changes in precipitation are a less important influence on $\delta^{18}\text{O}_{\text{precip}}$ changes in these regions. The impact of temperature changes (Fig. 6b) in the ISM, EAM and SW-SAM is broadly consistent with the global model. The slope of the relationship with temperature is negative for the IAM and NE-SAM, and since this is physically implausible it suggests that some factor not currently included in the MLR is influencing these records. However, the inconsistencies between the regional signals helps to explain why the global relationship between anomalies in temperature and $\delta^{18}\text{O}_{\text{precip}}$ is weak (Fig. 6b) and probably reflects the fact that tropical temperature changes during the Holocene are small. Data points from the EAM, ISM and IAM are well aligned with the global relationship between changes in $\delta^{18}\text{O}_{\text{precip}}$ and changes in wind direction (Fig. 6c), indicating that changes in source area or moisture pathway are an important control on changes in $\delta^{18}\text{O}_{\text{precip}}$ in these regions. However, values for CAM, SW-SAM, NE-SAM and SAfM deviate strongly from the global relationship. Recycling does not appear to be an important contributor to changes in $\delta^{18}\text{O}_{\text{precip}}$ except in SW-SAM and SAfM (Fig. 6d).

4 Discussion

We have shown that it is possible to derive an objective regionalisation of speleothem records based on PCoA of the oxygen-isotope trends through the Holocene (Fig. 2). This approach separates out regions with a distinctive northern hemisphere signal (e.g. ISM, EAM, NE-SAM) from regions with a distinctive southern hemisphere signal (e.g. SW-SAM, SAfM), reflecting the fact that the evolution of regional monsoons in each hemisphere follows, to some extent, insolation forcing. It also identifies

regions that have an intermediate pattern (e.g. IAM). The robustness of the regionalisation is borne out by the fact that Holocene composite trends from each region have tight confidence intervals (Fig. 4), showing that the signals of individual records across a region show broad similarities. The monsoon regions identified by PCoA are consistent with previous studies (Wang et al., 2014). The tracking of northern hemisphere insolation is a recognised feature of monsoon systems in India and China (see reviews by Kaushal et al., 2018; Zhang et al., 2019). The separation of speleothem records from NE-SAM from those in SW-SAM is consistent with the precipitation dipole that exists between northeastern Brazil (Nordeste) and the continental interior (Berbery and Barros, 2002; Boers et al., 2014). The anti-phasing of speleothem records from the two regions during the Holocene has been recognised in previous studies (Cruz et al., 2009; Deininger et al., 2019). The intermediate nature of the records from the maritime continent is consistent with the fact that the Indonesian-Australian (IAM) summer monsoon is influenced by cross-equatorial air flow and hence can be influenced by northern hemisphere conditions (Trenberth et al., 2000). Palaeoenvironmental records from this region show mixed signals for the Holocene: some have been interpreted as showing enhanced (Beaufort et al., 2010; Mohtadi et al., 2011; Quigley et al., 2010; Wyrwoll and Miller, 2001) and others reduced precipitation (Kuhnt et al., 2015; Steinke et al., 2014) during the early and mid-Holocene. Modelling studies have shown that this region is highly sensitive to SST changes in the Indian Ocean and South China Sea, which in turn reflect changes in the northern hemisphere winter monsoons. Although most climate models produce a reduction in precipitation across the IAM during the mid-Holocene in response to orbital forcing, this is less than might be expected in the absence of ocean feedbacks associated with changes in the Indian Ocean (Zhao and Harrison, 2012).

The separation of northern and southern monsoon regions is consistent with the idea that changes in monsoon rainfall are primarily driven by changes in insolation (Ding and Chan, 2005; Kutzbach et al., 2008). Indeed, regional $\delta^{18}\text{O}_{\text{spei}}$ composites from the EAM, ISM and SW-SAM show a clear relationship with the long-term trends in local summer insolation (Fig. 4). Similar patterns are seen in individual speleothem records from each region confirming that the composite trends are representative. However, the composite trends are not an exact mirror of the insolation signal over the Holocene. For example, the ISM and EAM composites show a more rapid rise during the early Holocene than implied by the insolation forcing. The maximum wet phase in these two regions lasts for ca 3,000 years, again contrasting with the gradual decline in insolation forcing after its peak at ca 11 ka. Both the rapid increase and the persistence of wet conditions for several thousand years is also observed in other palaeohydrological records across southern and central China, including pollen (Zhao et al., 2009; Li et al., 2018) and peat records (Hong et al., 2003; Zhou et al., 2004). These features are also characteristic of lake records from India (Misra et al., 2019). The lagged response to increasing insolation is thought to be due to the presence of northern hemisphere ice sheets in the early Holocene (Zhang et al., 2018). The persistence of wetter conditions through the early and mid-Holocene is thought to reflect the importance of land-surface and ocean feedbacks in sustaining regional monsoons (Dallmeyer et al., 2010; Kutzbach et al., 1996; Marzin and Braconnot, 2009; Rachmayani et al., 2015; Zhao and Harrison, 2012). The evolution of regional monsoons during the LIG shows patterns similar to those observed during the Holocene, including the lagged response to insolation and the persistence of wet conditions after peak insolation. This is again consistent

with the idea that internal feedbacks play a role in modulating the monsoon response to insolation forcing. We have also shown that there is little difference in the isotopic values between the MH and the LIG in the ISM and EAM regions, which is also observed in individual speleothem records (Kathayat et al., 2016; Wang et al., 2008). Given that the increase in summer insolation is much larger during the LIG than the MH, this finding indicates that other factors play a role in modulating the monsoon response to insolation forcing and may reflect the importance of global constraints on the externally-forced expansion of the tropical circulation (Biasutti et al., 2018).

Global relationships between $\delta^{18}\text{O}_{\text{precip}}$ and climate variables (precipitation amount, temperature and surface wind direction; Fig. 6) are consistent with existing studies: a strong relationship with precipitation and a weaker temperature effect has been widely observed at tropical and sub-tropical latitudes in modern observations (Dansgaard, 1964; Rozanski et al., 1993). The significant global relationship between $\delta^{18}\text{O}_{\text{precip}}$ and surface winds supports the idea that changes in moisture source and pathway are also important for explaining $\delta^{18}\text{O}$ variability over the Holocene. The multiple regression analysis also provides insights into the relative importance of different influences at a regional scale. In the ISM, the results support existing speleothem studies that suggest changes in precipitation amount (Cai et al., 2015; Fleitmann et al., 2004) and to a lesser extent moisture pathway (Breitenbach et al., 2010) drive $\delta^{18}\text{O}_{\text{spel}}$ variability. The $\delta^{18}\text{O}$ variability in the IAM region through the Holocene also appears to be strongly driven by changes in precipitation and moisture pathway, consistent with the interpretation of Wurtzel et al. (2018). Changes in regional precipitation (where the cave sites are located) do not seem to explain the observed changes in $\delta^{18}\text{O}_{\text{spel}}$ in the EAM during the Holocene, where Holocene $\delta^{18}\text{O}_{\text{precip}}$ evolution is largely driven by changes in atmospheric circulation (indexed by changes in surface winds). This is consistent with existing studies that emphasise changes in moisture source and/or pathway rather than local precipitation changes (Maher, 2016; Maher and Thompson, 2012; Tan, 2014; Yang et al., 2014). Speleothem $\delta^{18}\text{O}$ records in the SW-SAM clearly reflect regional-scale changes in precipitation, consistent with interpretations of individual records (Cruz et al., 2009; Kanner et al., 2013). However, this is a region where changes in precipitation recycling also appears to be important, perhaps unsurprisingly given that recycling presently contributes ca 25-35% of the precipitation over the Amazon (Brubaker et al., 1993; Eltahir and Bras, 1994).

The LGM is characterised by lower northern hemisphere summer insolation, globally cooler temperatures, expanded global ice volumes and lower GHG concentrations than either the MH or the LIG. The $\delta^{18}\text{O}_{\text{spel}}$ anomalies are more positive during the LGM than the MH or LIG, suggesting drier conditions in the ISM, EAM and IAM, supported by simulated changes in $\delta^{18}\text{O}_{\text{precip}}$ and precipitation (Fig. 3). Cooler SSTs of approximately 2°C (relative to the MH and LIG) in the ISM and EAM and of approximately 3°C in IAM source areas, together with a ca 5% decrease in relative humidity (Yue et al., 2011) would result in a water vapour $\delta^{18}\text{O}$ signal at the source ca 1 ‰ more depleted than seawater. This depletion results from the temperature dependence of equilibrium fractionation during evaporation and kinetic isotope effects related to humidity (Clark and Fritz, 1997). This fractionation counteracts any impact from enriched seawater $\delta^{18}\text{O}$ values during the LGM (ca. +1 ‰ relative to the MH or LIG; Waelbroeck et al., 2002). Enriched $\delta^{18}\text{O}_{\text{precip}}$ and $\delta^{18}\text{O}_{\text{spel}}$ values during the LGM must therefore be caused by a significant decrease in atmospheric moisture and precipitation that resulted from the cooler conditions.

We have used version 2 of the SISAL database (Atsawawaranunt et al., 2018; Comas-Bru et al., 2020a) in our analyses. Despite
445 the fact that SISALv2 includes more than 70% of known speleothem isotope records, there are still too few records from some
regions (e.g. Africa, the Caribbean) to make meaningful analyses. The records for older time periods are also sparse. There are
only 14 records from monsoon regions covering the LIG in SISALv2, for example. Nevertheless, our analyses show that there
are robust and explicable patterns for most monsoon regions during the Holocene and sufficient records to make meaningful
analyses of the LGM and LIG. Whilst there is a need for the generation of new speleothem records from key regions such as
450 northern Africa, further expansion of the SISAL database will certainly provide additional opportunities to analyse the
evolution of the monsoons through time.

The impact of age uncertainties, included in SISALv2, are not taken into account in our analyses. Age uncertainties during the
Holocene are smaller than the interval used for binning records and the width of the time windows used, and thus should not
have a significant effect on our conclusions. The mean age uncertainty at the LGM and LIG is ca 430 and 1140 years,
455 respectively. However, varying the window length for the selection of LGM and LIG samples from ± 500 to ± 2000 years,
thereby encompassing this uncertainty, has a negligible effect (<0.5 ‰) on the average $\delta^{18}\text{O}$ values. Thus, the interglacial-
glacial contrast in regional $\delta^{18}\text{O}_{\text{spel}}$ is also robust to age uncertainties.

Isotope-enabled climate models are used in this study to explain observed regional-scale trends in $\delta^{18}\text{O}_{\text{spel}}$. There is a limited
number of isotope-enabled models, and there are no simulations of the same time period using the same experimental protocol.
460 Although there are simulations of the MH from both ECHAM5-wiso and GISS, for example, these models have different grid
resolutions and used different boundary conditions. This could help to explain why the two models yield different estimates
of the change in regional $\delta^{18}\text{O}_{\text{precip}}$ (of 0.5 ‰) at the MH. However, both models show trends in $\delta^{18}\text{O}_{\text{precip}}$ that reproduce the
observed changes in regional $\delta^{18}\text{O}_{\text{spel}}$ (Figs 3 and 4), and this provides a basis for using these models to explore the causes of
these trends on different timescales. The failure to reproduce the LGM $\delta^{18}\text{O}_{\text{spel}}$ signal in SW-SAM in the ECHAM5-wiso
465 model, which precluded a consideration of interglacial-glacial shifts in this region, is a common feature of other isotope-
enabled simulations (Caley et al., 2014; Risi et al., 2010).

This study illustrates a novel data-model approach to investigate the drivers of $\delta^{18}\text{O}_{\text{spel}}$ under past conditions, by comparing
composite regional records and then using multiple linear regression of isotope-enabled palaeoclimate simulations to determine
the role of individual climate drivers of these trends. This obviates the need to use modern $\delta^{18}\text{O}_{\text{precip}}$ -climate relationships to
470 explain changes under conditions considerably different from today or to rely on coherency between different
palaeohydrological archives which may respond to different climate variables. This model interrogation approach could be
employed to address questions about the regional drivers of speleothem records outside the monsoon regions.

5 Conclusions

Geographically distributed speleothem $\delta^{18}\text{O}$ records and isotope-enabled climate models can be used together to understand the drivers of $\delta^{18}\text{O}_{\text{spel}}$ on orbital timescales. Speleothem records, objectively grouped into monsoon regions by record correlation and multivariate ordination techniques, show regional trends that are consistent with changes in summer insolation but modulated by land-surface and ocean feedbacks. Glacial $\delta^{18}\text{O}_{\text{spel}}$ signals are best explained by a large decrease in precipitation, as a consequence of lower atmospheric moisture content driven by global cooling. The evolution of $\delta^{18}\text{O}_{\text{spel}}$ through the Holocene across the global monsoon domain is driven by changes in precipitation, atmospheric circulation and temperature. At the regional scale, our analyses support the increasing number of studies suggesting that East Asian monsoon speleothem $\delta^{18}\text{O}$ evolution through the Holocene relates to changes in atmospheric circulation (i.e. changes in moisture pathway and/or source). Changes in regional precipitation are the predominant driver of Holocene $\delta^{18}\text{O}_{\text{spel}}$ evolution in the Indian, southwestern South American and Indonesian-Australian monsoons, although changes in atmospheric circulation also contribute in the Indian and Indonesian-Australian monsoon regions and changes in precipitation recycling in southwestern South America.

Code and data availability

The SISAL (Speleothem Isotopes Synthesis and AnaLysis) database version 2 is available through the University of Reading Research Archive at <http://dx.doi.org/10.17864/1947.256>. The ECHAM5-wiso MH and LGM simulations are available at <https://doi.org/10.1594/PANGAEA.902347>. The ECHAM LIG simulation is available at <https://doi.pangaea.de/10.1594/PANGAEA.879229>. The OIPC mean annual $\delta^{18}\text{O}_{\text{precip}}$ data are available at http://wateriso.utah.edu/waterisotopes/pages/data_access/ArcGrids.html. CRU-TS4.01 mean annual temperature data are available at <http://doi.org/10/gcmcz3>. The GISS simulations and code used to generate the figures in this paper are available at <https://doi.org/10.5281/zenodo.3875496>.

495 Author contributions

The study was designed by SP, SPH, LCB and NK. MW and ALG provided climate model outputs. SP ran the analyses. The first draft of the manuscript was written by SP, SPH and LCB and all authors contributed to the final draft.

Competing interests

The authors declare no competing interests.

500 **Funding**

SP, LCB and SPH acknowledge funding support from the ERC-funded project GC2.0 (Global Change 2.0: Unlocking the past for a clearer future, grant number 694481). SPH also acknowledges support from the JPI-Belmont Forum project "Palaeoclimate Constraints on Monsoon Evolution and Dynamics (PaCMEDy)" funded through NERC (NE/P006752/1).

Acknowledgements

505 Ideas in this paper were developed at a meeting of the SISAL (Speleothem Isotopes Synthesis and Analysis) working group of the Past Global Changes (PAGES) programme. We thank PAGES for their support for this meeting and our colleagues in SISAL for the useful discussions. We thank Gabriele Messori for help with the code for calculating precipitation recycling.

510 **References**

- Aggarwal, P. K., Alduchov, O. A., Froehlich, K. O., Araguas-Araguas, L. J., Sturchio, N. C. and Kurita, N.: Stable isotopes in global precipitation: A unified interpretation based on atmospheric moisture residence time, *Geophys. Res. Lett.*, 39(11), <https://doi.org/10.1029/2012GL051937>, 2012.
- 515 Atsawawaranunt, K., Comas-Bru, L., Amirnezhad Mozhdehi, S., Deininger, M., Harrison, S. P., Baker, A., Boyd, M., Kaushal, N., Masood Ahmad, S., Ait Brahim, Y., Arienzo, M., Bajo, P., Braun, K., Burstyn, Y., Chawchai, S., Duan, W., Hatvani, I. G., Hu, J., Kern, Z., Labuhn, I., Lachniet, M., Lechleitner, F. A., Lorrey, A., Pérez-Mejías, C., Pickering, R., Scroton, N. and SISAL working group members: The SISAL database: A global resource to document oxygen and carbon isotope records from speleothems, *Earth Syst. Sci. Data*, 10, 1687–1713, <https://doi.org/10.5194/essd-10-1687-2018>, 2018.
- 520 Ayliffe, L.K., Gagan, M.K., Zhao, J.X., Drysdale, R.N., Hellstrom, J.C., Hantoro, W.S., Griffiths, M.L., Scott-Gagan, H., St Pierre, E., Cowley, J.A. and Suwargadi, B.W.: Rapid interhemispheric climate links via the Australasian monsoon during the last deglaciation, *Nat. Commun.*, 4(1), 1-6, <https://doi.org/10.1038/ncomms3908>, 2013
- Bailey, A., Posmentier, E. and Feng, X.: Patterns of evaporation and precipitation drive global isotopic changes in atmospheric moisture, *Geophys. Res. Lett.*, 45(14), <https://doi.org/10.1029/2018GL078254>, 2018.
- 525 Bar-Matthews, M., Ayalon, A. and Kaufman, A.: Late Quaternary paleoclimate in the eastern Mediterranean region from stable isotope analysis of speleothems at Soreq Cave, Israel, *Quat. Res.*, 47(2), 155–168, <https://doi.org/10.1006/qres.1997.1883>, 1997.
- Battacharya, S.K., Froehlich, K., Aggarwal, P.K., and Kulkarni, K.M.: Isotopic variation in Indian Monsoon precipitation: Records from Bombay and New Delhi, *Geophys. Res. Lett.*, 30(24), <http://doi.org/10.1029/2003GL018453>, 2003.
- 530 Battisti, D. S., Ding, Q. and Roe, G. H.: Coherent pan-Asian climatic and isotopic response to orbital forcing of tropical insolation, *J. Geophys. Res. Atmos.*, 119(21), 11997-12020, <https://doi.org/10.1002/2014JD021960>, 2014.

- Beaufort, L., Van Der Kaars, S., Bassinot, F. C. and Moron, V.: Past dynamics of the Australian monsoon: precession, phase and links to the global monsoon concept, *Clim. Past*, 6, 695–706, <https://doi.org/10.5194/cp-6-695-2010>, 2010.
- Bennett, K. D.: Determination of the number of zones in a biostratigraphical sequence, *New Phytol.*, 132(1), 155–170, <https://doi.org/10.1111/j.1469-8137.1996.tb04521.x>, 1996.
- 535 Berbery, E. H. and Barros, V. R.: The hydrologic cycle of the La Plata basin in South America, *J. Hydrometeorol.*, 3(6), 630–645, [https://doi.org/10.1175/1525-7541\(2002\)003<0630:THCOTL>2.0.CO;2](https://doi.org/10.1175/1525-7541(2002)003<0630:THCOTL>2.0.CO;2), 2002.
- Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quat. Sci. Rev.*, 10(4), 297–317, [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.
- 540 Berger, A. L.: Long-term variations of daily insolation and Quaternary climatic changes, *J. Atmos. Sci.*, 35(12), 2362–2367, [https://doi.org/10.1175/1520-0469\(1978\)035<2362:ltvodi>2.0.co;2](https://doi.org/10.1175/1520-0469(1978)035<2362:ltvodi>2.0.co;2), 1978.
- Berkelhammer, M., Sinha, A., Mudelsee, M., Cheng, H., Edwards, R.L. and Cannariato, K.: Persistent multidecadal power of the Indian Summer Monsoon, *Earth Planet. Sci. Lett.*, 290(1-2), 166–172, <http://doi.org/10.1016/j.epsl.2009.12.017>, 2010.
- Biasutti, M., Voigt, A., Boos, W., Braconnot, P., Hargreaves, J.C., Harrison, S.P., Kang, S.M., Mapes, B.E., Scheff, J., Schumaker, C., Sobel, A.H. and Xie, S.P.: Global energetics and local physics as drivers of past, present and future monsoons, *Nat. Geosci.*, 11(6), 392–400, <https://doi.org/10.1038/s41561-018-0137-1>, 2018.
- 545 Bin, C., Xiang-De, X. and Tianliang, Z.: Main moisture sources affecting lower Yangtze River Basin in boreal summers during 2004–2009, *Int. J. Climatol.*, 33(4), 1035–1046, <https://doi.org/10.1002/joc.3495>, 2013.
- Boers, N., Rheinwalt, A., Bookhagen, B., Barbosa, H. M. J., Marwan, N., Marengo, J. and Kurths, J.: The South American rainfall dipole: A complex network analysis of extreme events, *Geophys. Res. Lett.*, 41(20), 7397–7405, <https://doi.org/10.1002/2014GL061829>, 2014.
- 550 Bowen, G.: Gridded maps of the isotopic composition of meteoric waters, [online] Available from: <http://waterisotopes.org> (Accessed 15 October 2018), 2018.
- Bowen, G. J. and Revenaugh, J.: Interpolating the isotopic composition of modern meteoric precipitation, *Water Resour. Res.*, 39(10), <https://doi.org/10.1029/2003WR002086>, 2003.
- 555 Breitenbach, S. F. M., Adkins, J. F., Meyer, H., Marwan, N., Kumar, K. K. and Haug, G. H.: Strong influence of water vapor source dynamics on stable isotopes in precipitation observed in Southern Meghalaya, NE India, *Earth Planet. Sci. Lett.*, 292(1–2), 212–220, <https://doi.org/10.1016/j.epsl.2010.01.038>, 2010.
- Brook, E. J., Harder, S., Severinghaus, J., Steig, E. J. and Sucher, C. M.: On the origin and timing of rapid changes in atmospheric methane during the Last Glacial Period, *Global Biogeochem. Cycles*, 14(2), 559–572, <https://doi.org/10.1029/1999GB001182>, 2000.
- 560 Brubaker, K. L., Entekhabi, D. and Eagleson, P. S.: Estimation of continental precipitation recycling, *J. Clim.*, 6(6), 1077–1089, [https://doi.org/10.1175/1520-0442\(1993\)006<1077:EOCPR>2.0.CO;2](https://doi.org/10.1175/1520-0442(1993)006<1077:EOCPR>2.0.CO;2), 1993.
- Cai, Y., Cheng, H., An, Z., Lawrence Edwards, R., Wang, X., Tan, L. and Wang, J.: Large variations of oxygen isotopes in precipitation over south-central Tibet during Marine Isotope Stage 5, *Geology*, 38(3), 243–246, <https://doi.org/10.1130/G30306.1>, 2010.
- 565

- Cai, Y., Zhang, H., Cheng, H., An, Z., Lawrence Edwards, R., Wang, X., Tan, L., Liang, F., Wang, J. and Kelly, M.: The Holocene Indian monsoon variability over the southern Tibetan Plateau and its teleconnections, *Earth Planet. Sci. Lett.*, 335, 135–144, <https://doi.org/10.1016/j.epsl.2012.04.035>, 2012.
- 570 Cai, Y., Fung, I. Y., Edwards, R. L., An, Z., Cheng, H., Lee, J. E., Tan, L., Shen, C. C., Wang, X., Day, J. A., Zhou, W., Kelly, M. J. and Chiang, J. C. H.: Variability of stalagmite-inferred Indian monsoon precipitation over the past 252,000 y, *Proc. Natl. Acad. Sci. U. S. A.*, 112(10), 2954–2959, <https://doi.org/10.1073/pnas.1424035112>, 2015.
- Caley, T., Roche, D. M., Waelbroeck, C. and Michel, E.: Oxygen stable isotopes during the Last Glacial Maximum climate: Perspectives from data-model (iLOVECLIM) comparison, *Clim. Past*, 10(6), 1939–1955, <https://doi.org/10.5194/cp-10-1939-2014>, 2014.
- 575 Carlson, A. E., LeGrande, A. N., Oppo, D. W., Came, R. E., Schmidt, G. A., Anslow, F. S., Licciardi, J. M. and Obbink, E. A.: Rapid early Holocene deglaciation of the Laurentide ice sheet, *Nature Geosci.*, 1(9), 620–624, <https://doi.org/10.1038/ngeo285>, 2008.
- 580 Carolin, S. A., Cobb, K. M., Lynch-Stieglitz, J., Moerman, J. W., Partin, J. W., Lejau, S., Malang, J., Clark, B., Tuen, A. A. and Adkins, J. F.: Northern Borneo stalagmite records reveal West Pacific hydroclimate across MIS 5 and 6, *Earth Planet. Sci. Lett.*, 439, 182–193, <https://doi.org/10.1016/j.epsl.2016.01.028>, 2016. Cheng, H., Edwards, R. L., Wang, Y., Kong, X., Ming, Y., Kelly, M. J., Wang, X., Gallup, C. D. and Liu, W.: A penultimate glacial monsoon record from Hulu Cave and two-phase glacial terminations, *Geology*, 34(3), 217–220, <https://doi.org/10.1130/G22289.1>, 2006.
- 585 Cheng, H., Fleitmann, D., Edwards, R. L., Wang, X., Cruz, F. W., Auler, A. S., Mangini, A., Wang, Y., Kong, X., Burns, S. J. and Matter, A.: Timing and structure of the 8.2 kyr B.P. event inferred from $\delta^{18}\text{O}$ records of stalagmites from China, Oman, and Brazil, *Geology*, 37(11), 1007–1010, <https://doi.org/10.1130/G30126A.1>, 2009.
- Cheng, H., Sinha, A., Cruz, F. W., Wang, X., Edwards, R. L., d’Horta, F. M., Ribas, C. C., Vuille, M., Stott, L. D. and Auler, A. S.: Climate change patterns in Amazonia and biodiversity, *Nat. Commun.*, 4(1), 1411, <https://doi.org/10.1038/ncomms2415>, 2013.
- Clark, I. and Fritz, P.: *Environmental Isotopes in Hydrology*, Lewis Publishers, New York., 1997.
- 590 Cleveland, W. S. and Devlin, S. J.: Locally weighted regression: An approach to regression analysis by local fitting, *J. Am. Stat. Assoc.*, 83(403), 596–610, <https://doi.org/10.1080/01621459.1988.10478639>, 1988.
- Cobb, K.M., Adkins, J.F., Partin, J.W. and Clark, B.: Regional-scale climate influences on temporal variations of rainwater and cave dripwater oxygen isotopes in northern Borneo, *Earth Planet. Sci. Lett.*, 263(3-4), 207–220, <https://doi.org/10.1016/j.epsl.2007.08.024>, 2007.
- 595 Cole, J. E., Rind, D., Webb, R. S., Jouzel, J. and Healy, R.: Climatic controls on interannual variability of precipitation $\delta^{18}\text{O}$: Simulated influence of temperature, precipitation amount, and vapor source region, *J. Geophys. Res. Atmos.*, 104(D12), 14223–14235, <https://doi.org/10.1029/1999JD900182>, 1999.
- 600 Comas-Bru, L., Harrison, S. P., Werner, M., Rehfeld, K., Scroxton, N. and Veiga-Pires, C.: Evaluating model outputs using integrated global speleothem records of climate change since the last glacial, *Clim. Past.*, 15(4), 1557–1579, <https://doi.org/10.5194/cp-15-1557-2019>, 2019.
- Comas-Bru, L., Atsawawanunt, K., Harrison, S. P. and SISAL working group members: SISAL (Speleothem Isotopes Synthesis and AnaLysis Working group) database version 2.0, <http://dx.doi.org/10.17864/1947.242>, 2020a.

- Comas-Bru, L., Rehfeld, K., Roesch, C., Amirnezhad-Mozhdehi, S., Harrison, S. P., Atsawawaranunt, K., Ahmad, S. M., Ait Brahim, Y., Baker, A., Bosomworth, M., Breitenbach, S. F., Burstyn, Y., Columbu, A., Deininger, M., Demény, A., Dixon, B., Fohlmeister, J., Hatvani, I. G., Hu, J., Kaushal, N., Kern, Z., Labuhn, I., Lechleitner, F. A., Lorrey, A., Martrat, B., Novello, V. F., Oster, J., Pérez-Mejías, C., Scholz, D., Scroxton, N., Sinha, N., Ward, B. M., Warken, S., Zhang, H. and SISAL working group members: SISALv2: A comprehensive speleothem isotope database with multiple age-depth models, *Earth Syst. Sci. Data*, 12(4), <https://doi.org/10.5194/essd-2020-39>, 2020b.
- 605 Coplen, T. B., Kendall, C. and Hopple, J.: Comparison of stable isotope reference samples, *Nature*, 302(5905), 236–238, <https://doi.org/10.1038/302236a0>, 1983.
- 610 Cruz, F. W., Burns, S. J., Karmann, I., Sharp, W. D., Vuille, M., Cardoso, A. O., Ferrari, J. A., Silva Dias, P. L. and Viana, O.: Insolation-driven changes in atmospheric circulation over the past 116,000 years in subtropical Brazil, *Nature*, 434(7029), 63–66, <https://doi.org/10.1038/nature03365>, 2005.
- Cruz, F.W., Burns, S.J., Karmann, I., Sharp, W.D. and Vuille, M.: Reconstruction of regional atmospheric circulation features during the late Pleistocene in subtropical Brazil from oxygen isotope composition of speleothems, *Earth Planet. Sci. Lett.*, 248(1-2), 495-507, <https://doi.org/10.1016/j.epsl.2006.06.019>, 2006.
- Cruz, F. W., Vuille, M., Burns, S. J., Wang, X., Cheng, H., Werner, M., Lawrence Edwards, R., Karmann, I., Auler, A. S. and Nguyen, H.: Orbitally driven east-west antiphasing of South American precipitation, *Nat. Geosci.*, 2(3), 210–214, <https://doi.org/10.1038/ngeo444>, 2009.
- 620 D’Abreton, P. C. and Tyson, P. D.: Three-dimensional kinematic trajectory modelling of water vapour transport over Southern Africa, *Water SA*, 22(4), 297–306, 1996.
- Dallmeyer, A., Claussen, M. and Otto, J.: Contribution of oceanic and vegetation feedbacks to Holocene climate change in monsoonal Asia, *Clim. Past*, 6(2), 195–218, <https://doi.org/10.5194/cp-6-195-2010>, 2010.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16(4), 436–468, <https://doi.org/10.3402/tellusa.v16i4.8993>, 1964.
- 625 Deininger, M., Ward, B. M., Novello, V. F. and Cruz, F. W.: Late Quaternary variations in the South American monsoon system as inferred by speleothems—New perspectives using the SISAL database, *Quaternary*, 2(1), 6, <https://doi.org/10.3390/quat2010006>, 2019.
- Ding, Y. and Chan, J. C. L.: The East Asian summer monsoon: an overview, *Meteorol. Atmos. Phys.*, 89(1–4), 117–142, <https://doi.org/10.1007/s00703-005-0125-z>, 2005.
- 630 Dong, J., Wang, Y., Cheng, H., Hardt, B., Lawrence Edwards, R., Kong, X., Wu, J., Chen, S., Liu, D., Jiang, X. and Zhao, K.: A high-resolution stalagmite record of the Holocene East Asian monsoon from Mt Shennongjia, central China, *Holocene*, 20(2), 257–264, <https://doi.org/10.1177/0959683609350393>, 2010.
- Drumond, A., Nieto, R., Gimeno, L. and Ambrizzi, T.: A Lagrangian identification of major sources of moisture over Central Brazil and la Plata Basin, *J. Geophys. Res. Atmos.*, 113(D14), <https://doi.org/10.1029/2007JD009547>, 2008.
- 635 Drumond, A., Nieto, R., Trigo, R., Ambrizzi, T., Souza, E. and Gimeno, L.: A Lagrangian identification of the main sources of moisture affecting northeastern Brazil during its pre-rainy and rainy seasons, *PLoS One*, 5(6), <https://doi.org/10.1371/journal.pone.0011205>, 2010.
- Durán-Quesada, A. M., Gimeno, L., Amador, J. A. and Nieto, R.: Moisture sources for Central America: Identification of moisture sources using a Lagrangian analysis technique, *J. Geophys. Res. Atmos.*, 115(5), 1–15,

- 640 <https://doi.org/10.1029/2009JD012455>, 2010.
- Dykoski, C. A., Edwards, R. L., Cheng, H., Yuan, D., Cai, Y., Zhang, M., Lin, Y., Qing, J., An, Z. and Revenaugh, J.: A high-resolution, absolute-dated Holocene and deglacial Asian monsoon record from Dongge Cave, China, *Earth Planet. Sci. Lett.*, 233(1–2), 71–86, <https://doi.org/10.1016/J.EPSL.2005.01.036>, 2005.
- Eltahir, E. A. B. and Bras, R. L.: Precipitation recycling in the Amazon basin, *Q. J. R. Meteorol. Soc.*, 120(518), 861–880, 645 <https://doi.org/10.1002/qj.49712051806>, 1994.
- Fairchild, I. J. and Baker, A.: *Speleothem Science: From Process to Past Environments*, 1st ed., Wiley-Blackwell, Chichester., 2012.
- Fleitmann, D., Burns, S. J., Neff, U., Mudelsee, M., Mangini, A. and Matter, A.: Palaeoclimatic interpretation of high-resolution oxygen isotope profiles derived from annually laminated speleothems from Southern Oman, *Quat. Sci. Rev.*, 23(7–8), 935–945, <https://doi.org/10.1016/j.quascirev.2003.06.019>, 2004. 650
- Friedman, I., Harris, J. M., Smith, G. I. and Johnson, C. A.: Stable isotope composition of waters in the Great Basin, United States 1. Air-mass trajectories, *J. Geophys. Res. Atmos.*, <https://doi.org/10.1029/2001JD000565>, 2002.
- Gat, J. R.: Oxygen and hydrogen isotopes in the hydrological cycle, *Annu. Rev. Earth Planet. Sci.*, 24(1), 225–262, <https://doi.org/10.1146/annurev.earth.24.1.225>, 1996.
- 655 Gierz, P., Werner, M. and Lohmann, G.: Simulating climate and stable water isotopes during the last interglacial using a coupled climate-isotope model, link to model results, PANGAEA, <https://doi.org/10.1594/PANGAEA.879229>, 2017a.
- Gierz, P., Werner, M. and Lohmann, G.: Simulating climate and stable water isotopes during the Last Interglacial using a coupled climate-isotope model, *J. Adv. Model. Earth Syst.*, 9(5), 2027–2045, <https://doi.org/10.1002/2017MS001056>, 2017b.
- 660 Goldschneider, N., Chen, Z., Auler, A.S., Bakalowicz, M., Broda, S., Drew, D., Hartmann, J., Jiang, G., Moosdorf, N., Stevanovic, Z. and Veni, G.: Global distribution of carbonate rocks and karst water resources, *Hydrogeol. J.*, 28(5), 1661–1677, <https://doi.org/10.1007/s10040-020-02139-5>, 2020.
- Gower, J. C.: Some distance properties of latent root and vector methods used in multivariate analysis, *Biometrika*, 53(3–4), 325–338, <https://doi.org/10.2307/2333639>, 1966.
- 665 Griffiths, M. L., Drysdale, R. N., Gagan, M. K., Zhao, J. X., Ayliffe, L. K., Hellstrom, J. C., Hantoro, W. S., Frisia, S., Feng, Y. X., Cartwright, I., Pierre, E. S., Fischer, M. J. and Suwargadi, B. W.: Increasing Australian-Indonesian monsoon rainfall linked to early Holocene sea-level rise, *Nat. Geosci.*, 2(9), 636–639, <https://doi.org/10.1038/ngeo605>, 2009.
- Grossman, E. L. and Ku, T. L.: Oxygen and carbon isotope fractionation in biogenic aragonite: Temperature effects, *Chem. Geol. Isot. Geosci. Sect.*, 59(1), 59–74, [https://doi.org/10.1016/0168-9622\(86\)90057-6](https://doi.org/10.1016/0168-9622(86)90057-6), 1986.
- 670 Haese, B., Werner, M. and Lohmann, G.: Stable water isotopes in the coupled atmosphere-land surface model ECHAM5-JSBACH, *Geosci. Model Dev.*, 6, 3375–3418, <https://doi.org/10.5194/gmd-6-1463-2013>, 2012.
- Hardy, D. R., Vuille, M. and Bradley, R. S.: Variability of snow accumulation and isotopic composition on Nevado Sajama, Bolivia, *J. Geophys. Res. D Atmos.*, 108(D22), <https://doi.org/10.1029/2003jd003623>, 2003.
- Harris, I., Jones, P. D., Osborn, T. J. and Lister, D. H.: Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset, *Int. J. Climatol.*, 34(3), 623–642, <https://doi.org/10.1002/joc.3711>, 2014.

- 675 Hong, Y. T., Hong, B., Lin, Q. H., Zhu, Y. X., Shibata, Y., Hirota, M., Uchida, M., Leng, X. T., Jiang, H. B., Xu, H., Wang, H. and Yi, L.: Correlation between Indian Ocean summer monsoon and North Atlantic climate during the Holocene, *Earth Planet. Sci. Lett.*, 211(3–4), 371–380, [https://doi.org/10.1016/S0012-821X\(03\)00207-3](https://doi.org/10.1016/S0012-821X(03)00207-3), 2003.
- Hu, C. Y., Henderson, G. M., Huang, J. H., Xie, S. C., Sun, Y. and Johnson, K. R.: Quantification of Holocene Asian monsoon rainfall from spatially separated cave records, *Earth Planet. Sci. Lett.*, 266(3–4), 221–232, <https://doi.org/10.1016/j.epsl.2007.10.015>, 2008.
- 680 Hu, J., Emile-Geay, J., Tabor, C., Nusbaumer, J. and Partin, J.: Deciphering oxygen isotope records from Chinese speleothems with an isotope-enabled climate model, *Paleoceanogr. Paleoclimatology*, 43(12), 2098–2112, <https://doi.org/10.1029/2019PA003741>, 2019.
- IAEA/WMO: Global Network of Isotopes in Precipitation, The GNIP Database, available at: <http://www.iaea.org/water> (last access: 20 January 2019), 2018.
- 685 Indermühle, A., Stocker, T. F., Joos, F., Fischer, H., Smith, H. J., Wahlen, M., Deck, B., Mastroianni, D., Tschumi, J., Blunier, T., Meyer, R. and Stauffer, B.: Holocene carbon-cycle dynamics based on CO₂ trapped in ice at Taylor Dome, Antarctica, *Nature*, 398(6723), 121–126, <https://doi.org/10.1038/18158>, 1999.
- Kanner, L. C., Burns, S. J., Cheng, H., Edwards, R. L. and Vuille, M.: High-resolution variability of the South American summer monsoon over the last seven millennia: Insights from a speleothem record from the central Peruvian Andes, *Quat. Sci. Rev.*, 75, 1–10, <https://doi.org/10.1016/j.quascirev.2013.05.008>, 2013.
- 690 Kärkkäinen T. and Saarela M.: Robust Principal Component Analysis of Data with Missing Values. In: Perner P. (eds) *Machine Learning and Data Mining in Pattern Recognition. MLDM 2015. Lecture Notes in Computer Science*, vol 9166. Springer, Cham, 2015
- 695 Kathayat, G., Cheng, H., Sinha, A., Spötl, C., Edwards, R. L., Zhang, H., Li, X., Yi, L., Ning, Y., Cai, Y., Lui, W. L. and Breitenbach, S. F. M.: Indian monsoon variability on millennial-orbital timescales, *Sci. Rep.*, 6, 24374, <https://doi.org/10.1038/srep24374>, 2016.
- Kaushal, N., Breitenbach, S. F. M., Lechleitner, F. A., Sinha, A., Tewari, V. C., Ahmad, S. M., Berkelhammer, M., Band, S., Yadava, M., Ramesh, R. and Henderson, G. M.: The Indian Summer Monsoon from a Speleothem $\delta^{18}\text{O}$ Perspective—A Review, *Quaternary*, 1(3), 29, <https://doi.org/10.3390/quat1030029>, 2018.
- 700 Kennett, D. J., Breitenbach, S. F. M., Aquino, V. V., Asmerom, Y., Awe, J., Baldini, J. U. L., Bartlein, P., Culleton, B. J., Ebert, C., Jazwa, C., Macri, M. J., Marwan, N., Polyak, V., Pruffer, K. M., Ridley, H. E., Sodemann, H., Winterhalder, B. and Haug, G. H.: Development and disintegration of Maya political systems in response to climate change, *Science*, 338(6108), 788–791, <https://doi.org/10.1126/science.1226299>, 2012.
- 705 Krause, C. E., Gagan, M. K., Dunbar, G. B., Hantoro, W. S., Hellstrom, J. C., Cheng, H., Edwards, R. L., Suwargadi, B. W., Abram, N. J. and Rifai, H.: Spatio-temporal evolution of Australasian monsoon hydroclimate over the last 40,000 years, *Earth Planet. Sci. Lett.*, 513, 103–112, <https://doi.org/10.1016/j.epsl.2019.01.045>, 2019.
- Kuhnt, W., Holbourn, A., Xu, J., Opdyke, B., De Deckker, P., Röhl, U. and Mudelsee, M.: Southern Hemisphere control on Australian monsoon variability during the late deglaciation and Holocene, *Nat. Commun.*, 6(1), 1–7, <https://doi.org/10.1038/ncomms6916>, 2015.
- 710 Kutzbach, J., Bonan, G., Foley, J. and Harrison, S. P.: Vegetation and soil feedbacks on the response of the African monsoon to orbital forcing in the early to middle Holocene, *Nature*, 384(6610), 623–626, <https://doi.org/10.1038/384623a0>, 1996.

- Kutzbach, J. E., Liu, X., Liu, Z. and Chen, G.: Simulation of the evolutionary response of global summer monsoons to orbital forcing over the past 280,000 years, *Clim. Dyn.*, 30(6), 567–579, <https://doi.org/10.1007/s00382-007-0308-z>, 2008.
- 715 Lachniet, M. S.: Climatic and environmental controls on speleothem oxygen-isotope values, *Quat. Sci. Rev.*, 28(5–6), 412–432, <https://doi.org/10.1016/j.quascirev.2008.10.021>, 2009.
- Lachniet, M. S.: Are aragonite stalagmites reliable paleoclimate proxies? Tests for oxygen isotope time-series replication and equilibrium, *Bull. Geol. Soc. Am.*, 127(11–12), 1521–1533, <https://doi.org/10.1130/B31161.1>, 2015.
- 720 Langebroek, P. M., Werner, M. and Lohmann, G.: Climate information imprinted in oxygen-isotopic composition of precipitation in Europe, *Earth Planet. Sci. Lett.*, 311(1–2), 144–154, <https://doi.org/10.1016/j.epsl.2011.08.049>, 2011.
- Legendre, P. and Legendre, L.: *Numerical Ecology*, 2nd edition, Elsevier B.V, Amsterdam., 1998.
- LeGrande, A. N. and Schmidt, G. A.: Sources of Holocene variability of oxygen isotopes in paleoclimate archives, *Clim. Past*, 5(3), 441–455, <https://doi.org/10.5194/cp-5-441-2009>, 2009.
- 725 Lewis, S. C., LeGrande, A. N., Kelley, M. and Schmidt, G. A.: Water vapour source impacts on oxygen isotope variability in tropical precipitation during Heinrich events, *Clim. Past*, 6(3), 325–343, <https://doi.org/10.5194/cp-6-325-2010>, 2010.
- Lewis, S. C., LeGrande, A. N., Schmidt, G. A. and Kelley, M.: Comparison of forced ENSO-like hydrological expressions in simulations of the preindustrial and mid-Holocene, *J. Geophys. Res. Atmos.*, 119(12), 7064–7082, <https://doi.org/10.1002/2013JD020961>, 2014.
- 730 Licciardi, J. M., Clark, P. U., Jenson, J. W. and Macayeal, D. R.: Deglaciation of a soft-bedded Laurentide Ice Sheet, *Quat. Sci. Rev.*, 17(4–5), 427–448, [https://doi.org/10.1016/S0277-3791\(97\)00044-9](https://doi.org/10.1016/S0277-3791(97)00044-9), 1998.
- Maher, B. A.: Holocene variability of the East Asian summer monsoon from Chinese cave records: A re-assessment, *Holocene*, 18(6), 861–866, <https://doi.org/10.1177/0959683608095569>, 2008.
- Maher, B. A.: Palaeoclimatic records of the loess/palaeosol sequences of the Chinese Loess Plateau, *Quat. Sci. Rev.*, 154, 23–84, <https://doi.org/10.1016/j.quascirev.2016.08.004>, 2016.
- 735 Maher, B. A. and Thompson, R.: Oxygen isotopes from Chinese caves: Records not of monsoon rainfall but of circulation regime, *J. Quat. Sci.*, 27(6), 615–624, <https://doi.org/10.1002/jqs.2553>, 2012.
- Marlon, J. R., Bartlein, P. J., Carcaillet, C., Gavin, D. G., Harrison, S. P., Higuera, P. E., Joos, F., Power, M. J. and Prentice, I. C.: Climate and human influences on global biomass burning over the past two millennia, *Nat. Geosci.*, 1(10), 697–702, <https://doi.org/10.1038/ngeo313>, 2008.
- 740 Marzin, C. and Braconnot, P.: Variations of Indian and African monsoons induced by insolation changes at 6 and 9.5 kyr BP, *Clim. Dyn.*, 33(2–3), 215–231, <https://doi.org/10.1007/s00382-009-0538-3>, 2009.
- McDermott, F.: Palaeo-climate reconstruction from stable isotope variations in speleothems: A review, *Quat. Sci. Rev.*, 23(7–8), 901–918, <https://doi.org/10.1016/j.quascirev.2003.06.021>, 2004.
- 745 Misra, P., Tandon, S. K. and Sinha, R.: Holocene climate records from lake sediments in India: Assessment of coherence across climate zones, *Earth-Science Rev.*, 190, 370–397, <https://doi.org/10.1016/j.earscirev.2018.12.017>, 2019.
- Moerman, J.W., Cobb, K.M., Adkins, J.F., Sodemann, H., Clark, B. and Tuen, A.A.: Diurnal to interannual rainfall $\delta^{18}\text{O}$

- variations in northern Borneo driven by regional hydrology. *Earth Planet. Sci. Lett.*, 369, 108–119, <https://doi.org/10.1016/j.epsl.2013.03.014>, 2013.
- 750 Mohtadi, M., Oppo, D. W., Steinke, S., Stuut, J. B. W., De Pol-Holz, R., Hebbeln, D. and Lückge, A.: Glacial to Holocene swings of the Australian-Indonesian monsoon, *Nat. Geosci.*, 4(8), 540–544, <https://doi.org/10.1038/ngeo1209>, 2011.
- Nivet, F., Bergonzini, L., Mathé, P. E., Noret, A., Monvoisin, G., Majule, A. and Williamson, D.: Influence of the balance of the intertropical front on seasonal variations of the isotopic composition in rainfall at Kisiba Masoko (Rungwe Volcanic Province, SW, Tanzania), *Isotopes Environ. Health Stud.*, 54(4), 352–369, <https://doi.org/10.1080/10256016.2018.1443923>, 2018.
- 755 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. and Wagner, H.: *vegan: Community Ecology Package*. R package version 2.5-2, Cran R, 2019.
- Pausata, F. S. R., Battisti, D. S., Nisancioglu, K. H. and Bitz, C. M.: Chinese stalagmite $\delta^{18}\text{O}$ controlled by changes in the Indian monsoon during a simulated Heinrich event, *Nat. Geosci.*, 4(7), 474–480, <https://doi.org/10.1038/ngeo1169>, 2011.
- 760 Quigley, M. C., Horton, T., Hellstrom, J. C., Cupper, M. L. and Sandiford, M.: Holocene climate change in arid Australia from speleothem and alluvial records, *Holocene*, 20(7), 1093–1104, <https://doi.org/10.1177/0959683610369508>, 2010.
- R Core Team: *R: A Language and Environment for Statistical Computing*, [online] Available from: <https://www.r-project.org/>, 2019.
- 765 Rachmayani, R., Prange, M. and Schulz, M.: North African vegetation-precipitation feedback in early and mid-Holocene climate simulations with CCSM3-DGVM, *Clim. Past*, 11(2), <https://doi.org/10.5194/cp-11-175-2015>, 2015.
- Rao, C. R.: The use and interpretation of principal component analysis in applied research, *Sankhya Indian J. Stat. Ser. A*, 329–358, 1964.
- 770 Risi, C., Bony, S. and Vimeux, F.: Influence of convective processes on the isotopic composition ($\delta^{18}\text{O}$ and δD) of precipitation and water vapor in the tropics: 2. Physical interpretation of the amount effect, *J. Geophys. Res. Atmos.*, 113(D19), <https://doi.org/10.1029/2008JD009943>, 2008.
- Risi, C., Bony, S., Vimeux, F. and Jouzel, J.: Water-stable isotopes in the LMDZ4 general circulation model: Model evaluation for present-day and past climates and applications to climatic interpretations of tropical isotopic records, *J. Geophys. Res. Atmos.*, 115(D12), <https://doi.org/10.1029/2009JD013255>, 2010.
- 775 Rohlf, F.J.: An Empirical Comparison of Three Ordination Techniques in Numerical Taxonomy, *Syst. Biol.*, 21, 271–280, <https://doi.org/10.1093/sysbio/21.3.271>, 1972.
- Rozanski, K., Araguás-Araguás, L. and Gonfiantini, R.: Isotopic patterns in modern global precipitation, *Geophys Monogr.*, 78, 1–36, <https://doi.org/10.1029/GM078p0001>, 1993.
- Salati, E., Dall'Olio, A., Matsui, E. and Gat, J. R.: Recycling of water in the Amazon Basin: An isotopic study, *Water Resour. Res.*, 15(5), 1250–1258, <https://doi.org/10.1029/WR015i005p01250>, 1979.
- 780 Schäfer-Neth, C. and Paul, A.: The Atlantic Ocean at the Last Glacial Maximum: 1. Objective mapping of the GLAMAP sea-surface conditions, in *The South Atlantic in the Late Quaternary*, edited by G. Wefer, S. Mulitza, and V. Ratmeyer, pp. 531–548, Springer, Berlin, Heidelberg., 2003.

- 785 Sinha, A., Kathayat, G., Cheng, H., Breitenbach, S.F., Berkelhammer, M., Mudelsee, M., Biswas, J. and Edwards, R.L.: Trends and oscillations in the Indian summer monsoon rainfall over the last two millennia, *Nat. Commun.*, 6(1), 1-8, <https://doi.org/10.1038/ncomms7309>, 2015.
- Sowers, T.: Ice core records of atmospheric N₂O covering the last 106,000 years, *Science*, 301(5635), 945–948, <https://doi.org/10.1126/science.1085293>, 2003.
- Steinke, S., Mohtadi, M., Prange, M., Varma, V., Pittauerova, D. and Fischer, H. W.: Mid- to Late-Holocene Australian-Indonesian summer monsoon variability, *Quat. Sci. Rev.*, 93, 142–154, <https://doi.org/10.1016/j.quascirev.2014.04.006>, 2014.
- 790 Tan, M.: Circulation effect: climatic significance of the short term variability of the oxygen isotopes in stalagmites from monsoonal China, *Quat. Sci.*, 29(5), 851–862, 2009.
- Tan, M.: Tread-wind driven inverse coupling between stalagmite $\delta^{18}\text{O}$ from monsoon region of China and large scale temperature-circulation effect on decadal to precessional timescales, *Quat. Sci.*, 31(6), 1086–1097, 2011.
- 795 Tan, M.: Circulation effect: Response of precipitation $\delta^{18}\text{O}$ to the ENSO cycle in monsoon regions of China, *Clim. Dyn.*, 42(3–4), 1067–1077, <https://doi.org/10.1007/s00382-013-1732-x>, 2014.
- Tan, L.C., Cai, Y., Cheng, H., Edwards, R.L., Shen, C., Gao, Y. and An, Z.: Climate significance of speleothem $\delta^{18}\text{O}$ from central China on decadal timescale, *J. Asian Earth Sci.*, 106, 150–155, <https://doi.org/10.1016/j.jseaeas.2015.03.008>, 2015.
- 800 Tremaine, D. M., Froelich, P. N. and Wang, Y.: Speleothem calcite formed in situ: Modern calibration of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ paleoclimate proxies in a continuously-monitored natural cave system, *Geochim. Cosmochim. Acta*, 75(17), 4929–4950, <https://doi.org/10.1016/j.gca.2011.06.005>, 2011.
- Trenberth, K. E., Stepaniak, D. P. and Caron, J. M.: The global monsoon as seen through the divergent atmospheric circulation, *J. Clim.*, 13(22), 3969–3993, [https://doi.org/10.1175/1520-0442\(2000\)013<3969:TGMAST>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<3969:TGMAST>2.0.CO;2), 2000.
- 805 Varma, V., Prange, M., Merkel, U., Kleinen, T., Lohmann, G., Pfeiffer, M., Renssen, H., Wagner, A., Wagner, S. and Schulz, M.: Holocene evolution of the Southern Hemisphere westerly winds in transient simulations with global climate models, *Clim. Past*, 8, 391–402, <https://doi.org/10.5194/cp-8-391-2012>, 2012.
- Wackerbarth, A., Langebroek, P. M., Werner, M., Lohmann, G., Riechelmann, S., Borsato, A. and Mangini, A.: Simulated oxygen isotopes in cave drip water and speleothem calcite in European caves, *Clim. Past*, 8(6), 1781–1799, <https://doi.org/10.5194/cp-8-1781-2012>, 2012.
- 810 Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J. C., McManus, J. F., Lambeck, K., Balbon, E. and Labracherie, M.: Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records, *Quat. Sci. Rev.*, 21(1–3), 295–305, [https://doi.org/10.1016/S0277-3791\(01\)00101-9](https://doi.org/10.1016/S0277-3791(01)00101-9), 2002.
- Wang, P. X., Wang, B., Cheng, H., Fasullo, J., Guo, Z. T., Kiefer, T. and Liu, Z. Y.: The global monsoon across timescales: Coherent variability of regional monsoons, *Clim. Past*, 10(6), 2007–2052, <https://doi.org/10.5194/cp-10-2007-2014>, 2014.
- 815 Wang, X., Auler, A. S., Edwards, R. L., Cheng, H., Ito, E. and Solheid, M.: Interhemispheric anti-phasing of rainfall during the last glacial period, *Quat. Sci. Rev.*, 25(23–24), 3391–3403, <https://doi.org/10.1016/j.quascirev.2006.02.009>, 2006.
- Wang, Y., Cheng, H., Edwards, R. L., Kong, X., Shao, X., Chen, S., Wu, J., Jiang, X., Wang, X. and An, Z.: Millennial- and orbital-scale changes in the East Asian monsoon over the past 224,000 years, *Nature*, 451(7182), 1090–1093, <https://doi.org/10.1038/nature06692>, 2008.

- 820 Wang, Y. J., Cheng, H., Edwards, R. L., An, Z. S., Wu, J. Y., Shen, C. C. and Dorale, J. A.: A high-resolution absolute-dated late Pleistocene monsoon record from Hulu Cave, China, *Science*, 294(5550), 2345–2348, <https://doi.org/10.1126/science.1064618>, 2001.
- Waterisotopes Database: <http://waterisotopes.org>, last access: 15 April 2017.
- Werner, M.: ECHAM5-wiso simulation data – present-day, mid-Holocene, and Last Glacial Maximum, PANGAEA, <https://doi.org/https://doi.pangaea.de/10.1594/PANGAEA.902347>, 2019.
- 825 Werner, M., Langebroek, P. M., Carlsen, T., Herold, M. and Lohmann, G.: Stable water isotopes in the ECHAM5 general circulation model: Toward high-resolution isotope modeling on a global scale, *J. Geophys. Res. Atmos.*, 116(D15), <https://doi.org/10.1029/2011JD015681>, 2011.
- 830 Werner, M., Haese, B., Xu, X., Zhang, X., Butzin, M. and Lohmann, G.: Glacial-interglacial changes in H₂¹⁸O, HDO and deuterium excess—results from the fully coupled ECHAM5/MPI-OM Earth system model, *Geosci. Model Dev.*, 9(2), 647–670, <https://doi.org/10.5194/gmd-9-647-2016>, 2016.
- Werner, M., Jouzel, J., Masson-Delmotte, V. and Lohmann, G.: Reconciling glacial Antarctic water stable isotopes with ice sheet topography and the isotopic paleothermometer, *Nat. Commun.*, 9(1), 1–10, <https://doi.org/10.1038/s41467-018-05430-y>, 2018.
- 835 Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag, New York. [online] Available from: <https://ggplot2.tidyverse.org>, 2016.
- Wurtzel, J. B., Abram, N. J., Lewis, S. C., Bajo, P., Hellstrom, J. C., Troitzsch, U. and Heslop, D.: Tropical Indo-Pacific hydroclimate response to North Atlantic forcing during the last deglaciation as recorded by a speleothem from Sumatra, Indonesia, *Earth Planet. Sci. Lett.*, 492, 264–278, <https://doi.org/10.1016/j.epsl.2018.04.001>, 2018.
- 840 Wyrwoll, K. H. and Miller, G. H.: Initiation of the Australian summer monsoon 14,000 years ago, *Quat. Int.*, 83, 119–128, [https://doi.org/10.1016/S1040-6182\(01\)00034-9](https://doi.org/10.1016/S1040-6182(01)00034-9), 2001.
- Xu, X., Werner, M., Butzin, M. and Lohmann, G.: Water isotope variations in the global ocean model MPI-OM, *Geosci. Model Dev.*, 5(3), 809–818, <https://doi.org/10.5194/cpd-8-4885-2012>, 2012.
- 845 Yang, X., Liu, J., Liang, F., Yuan, D., Yang, Y., Lu, Y. and Chen, F.: Holocene stalagmite $\delta^{18}\text{O}$ records in the East Asian monsoon region and their correlation with those in the Indian monsoon region, *Holocene*, 24(12), 1657–1664, <https://doi.org/10.1177/0959683614551222>, 2014.
- Yonge, C. J., Ford, D. C., Gray, J. and Schwarcz, H. P.: Stable isotope studies of cave seepage water, *Chem. Geol. Isot. Geosci. Sect.*, 58(1–2), 97–105, [https://doi.org/10.1016/0168-9622\(85\)90030-2](https://doi.org/10.1016/0168-9622(85)90030-2), 1985.
- 850 Yuan, D., Cheng, H., Edwards, R. L., Dykoski, C. A., Kelly, M. J., Zhang, M., Qing, J., Lin, Y., Wang, Y., Wu, J., Dorale, J. A., An, Z. and Cai, Y.: Timing, duration, and transitions of the Last Interglacial Asian monsoon, *Science*, 304(5670), 575–578, <https://doi.org/10.1126/science.1091220>, 2004.
- Yue, X., Wang, H., Liao, H. and Jiang, D.: Simulation of the Direct Radiative Effect of Mineral Dust Aerosol on the Climate at the Last Glacial Maximum, *J. Clim.*, 24(3), 843–858, <https://doi.org/10.1175/2010JCLI3827.1>, 2011.
- Zhang, Brahim, Li, Zhao, Kathayat, Tian, Baker, Wang, Zhang, Ning, Edwards and Cheng: The Asian Summer monsoon: Teleconnections and forcing Mechanisms—A review from Chinese speleothem $\delta^{18}\text{O}$ records, *Quaternary*, 2(3), 26,


855 <https://doi.org/10.3390/quat2030026>, 2019.

Zhang, X., Jin, L., Chen, J., Lu, H. and Chen, F.: Lagged response of summer precipitation to insolation forcing on the northeastern Tibetan Plateau during the Holocene, *Clim. Dyn.*, 50(9–10), 3117–3129, <https://doi.org/10.1007/s00382-017-3784-9>, 2018.


860 Zhao, Y. and Harrison, S. P.: Mid-Holocene monsoons: A multi-model analysis of the inter-hemispheric differences in the responses to orbital forcing and ocean feedbacks, *Clim. Dyn.*, 39(6), 1457–1487, <https://doi.org/10.1007/s00382-011-1193-z>, 2012.

Zhou, W., Yu, X., Jull, A. J. T., Burr, G., Xiao, J. Y., Lu, X. and Xian, F.: High-resolution evidence from southern China of an early Holocene optimum and a mid-Holocene dry event during the past 18,000 years, *Quat. Res.*, 62(1), 39–48, <https://doi.org/10.1016/j.yqres.2004.05.004>, 2004.

865




870 **Figure 1: Spatial distribution of speleothem records used in this study. Colours indicate the sites used in Principal Coordinates**
Analysis and Redundancy Analysis (PCoA, RDA) to separate monsoon regions, and sites not used in PCoA and RDA but used in
subsequent analyses. The individual regional monsoons are shown by boxes: CAM = Central American Monsoon latitude: 10 to
33°; longitude: -115 to -58°, SW-SAM = southwestern South American Monsoon latitude: -10° to 0°; longitude: -80° to -64° and
latitude: -30° to -10°; longitude: -68° to -40°), NE-SAM = northeastern South American Monsoon (latitude: -10° to 0°; longitude: -
875 **60° to -30°), SAFm = southern African Monsoon (latitude: -30° to -17°; longitude: 10° to 40°), ISM = Indian Summer Monsoon**
(latitude: 11° to 32°; longitude: 50° to 95°), EAM = East Asian Monsoon (latitude: 20° to 39°; longitude: 100° to 125°), IAM =
Indonesian-Australian Monsoon (latitude: -24° to 5°; longitude: 95° to 135°). Source region limits used in the multiple linear
regression analysis are also shown. The background carbonate lithology is from the World Karst Aquifer Mapping (WOKAM)
project (Goldschneider et al., 2020).




880 (a) PCoA biplot showing the loadings of each site on the first 2 axes, which represent 85% of the total variance. Shapes indicate the Holocene coverage of each site, where
sites with a coverage ≥ 8000 years represent most or all of the Holocene (Hol). Sites with a temporal coverage of < 8000 years are
885 coded to show whether they represent the early to mid-Holocene (EH to MH, record midpoint $> 8,000$ years BP), the mid Holocene
(MH, record midpoint between 8,000 and 5,000 years BP), or the mid to late-Holocene (LH to MH, midpoint $< 5,000$ years BP). (b)

RDA triplot, where the response variables are the PCoA1 and PCoA2 axes explained by latitude and longitude. The direction of the PCoA axes have been fixed so that they align with the explanatory variables.




890 Figure 3: Speleothem $\delta^{18}\text{O}$ anomalies compared to anomalies of $\delta^{18}\text{O}_{\text{precip}}$, precipitation and temperature from the ECHAM simulations for the (a) East Asian (EAM), (b) Indian (ISM) and (c) Indonesian-Australian (IAM) monsoons. The boxes show the median value (line) and the interquartile range, and the whiskers shown the minimum and maximum values, with outliers represented by grey dots. Note that the isotope axes are reversed, so that the most negative anomalies are at the top of the plot, to be consistent with the assumed relationship with the direction of change in precipitation and temperature.


895



900 **Figure 4: Evolution of regional speleothem $\delta^{18}\text{O}$ signals through the Holocene compared to $\delta^{18}\text{O}_{\text{precip}}$ simulated by the GISS model.**
 The left panel shows northern hemisphere monsoons (EAM = East Asian Monsoon; ISM = Indian Summer Monsoon) and summer
 (May through September) insolation at 30°N (Berger, 1978). The right panel shows southern hemisphere monsoons (SW-SAM =
 southwest South American Monsoon; IAM = Indonesian-Australian Monsoon) and summer (November through March)
 905 insolation for 20°S (Berger, 1978). The speleothem $\delta^{18}\text{O}$ changes are expressed as z-scores, with a smoothed loess fit (3,000 year window), and
 confidence intervals obtained by bootstrapping by site. $\delta^{18}\text{O}_{\text{precip}}$ values are expressed as anomalies from the pre-industrial control
 simulation. Note that the isotope axes are reversed, so that the most negative anomalies are at the top of the plot, to be consistent
 with the assumed relationship with the changes in insolation.



910 **Figure 5: Comparison of changes in summer insolation and Last Interglacial $\delta^{18}\text{O}_{\text{spe}}$ records from the (b,c) East Asian Monsoon (EAM), (d,e) Indian Summer Monsoon (ISM), (g) southwest South American Monsoon (SW-SAM) and (h) Indonesian-Australian Monsoon (IAM) regions. The U/Th dates and uncertainties are shown for each record. The summer insolation curves (Berger, 1978) are for May through September at 30°N in the northern hemisphere (a) and for November through March for 20°S in the southern hemisphere (f). Note that the isotope axes are reversed, so that the most negative anomalies are at the top of the plot, to be consistent with the assumed relationship with the changes in insolation.**



915

Figure 6: Partial residual plots from the multiple linear regression analysis, showing the relationship between anomalies in simulated $\delta^{18}\text{O}_{\text{precip}}$ and the four predictor variables, after taking account of the fitted partial effects of all the other predictors. The simulated $\delta^{18}\text{O}_{\text{precip}}$ are anomalies relative to the pre-industrial control simulation, and are annual values weighted by precipitation amount. The predictor variables are: precipitation in the delineated monsoon region (mm/d), temperature in the source region ($^{\circ}\text{C}$), surface wind direction over the source region ($^{\circ}$) as an index of potential changes in source region and the ratio of precipitation recycling to total precipitation over the monsoon region (RI, unitless). The predictor variables are summer mean values, where summer is defined as May to September for northern hemisphere monsoons and November to March for southern hemisphere monsoons.

920

	PCoA1	PCoA2	PCoA3	PCoA4	PCoA5
Eigenvalue	269.06	85.22	16.81	10.25	5.55
Explained (%)	64.87	20.55	4.054	2.47	1.34
Cumulative (%)	64.87	85.42	89.48	91.95	93.27

925

Table 1: Results of the Principal Coordinates Analysis (PCoA). Significant axes, as determined by the broken stick method (Bennett, 1996), are shown in bold.

	RDA1	RDA2
Latitude	0.88	-0.47
Longitude	0.75	0.67
Eigenvalue	0.73	0.04
Explained (%)	36.7	2.2

Table 2: Results of the redundancy analysis (RDA). Variables that are significantly correlated ($P < 0.01$) with the RDA axes are shown in bold.

930

	Regression coefficient	T value
Regional precipitation	-0.16	-8.75
Source area temperature	0.39	2.05
Wind direction	0.06	8.03
Precipitation recycling	4.34	1.92

Table 3: Results of the multiple linear regression analysis. Significant relationships ($P > 0.01$) are shown in bold.

935