

Holocene wildfire regimes in forested peatlands in western Siberia: interaction between peatland moisture conditions and the composition of plant functional types

5 Angelica Feurdean*^{1,2,3}, Andrei-Cosmin Diaconu⁴, Mirjam Pfeiffer², Mariusz Gałka⁵, Simon M. Hutchinson⁶, Geanina Butiseaca², Natalia Gorina⁷, Spassimir Tonkov⁸, Aidin Niamir,² Ioan Tantau⁴, Sergey Kirpotin^{9,10}

¹Department of Physical Geography, Goethe University, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

10 ²Senckenberg Biodiversity and Climate Research Centre (BiK-F), Senckenberganlage, 25, 60325, Frankfurt am Main, Germany

³ STAR-UBB Institute, Babeş-Bolyai University, Kogălniceanu 1, 400084, Cluj-Napoca, Romania

⁴Department of Geology, Babeş-Bolyai University, Kogălniceanu 1, 400084, Cluj-Napoca, Romania

15 ⁵Department of Biogeography, Paleoecology and Nature Conservation, Faculty of Biology and Environmental Protection, University of Lodz, Banacha 1/3, Lodz, Poland

⁶School of Science, Engineering and Environment, University of Salford, Greater Manchester M5 4WT, Salford, UK

⁷ Department of Ecology, Natural Use and Environmental Engineering, National Tomsk State University, Lenina Pr., 36, 634050, Tomsk, Russia

20 ⁸Laboratory of Palynology, Faculty of Biology, Sofia University St. Kliment Ohridski, Dragan Tsankov 8, 1164, Sofia, Bulgaria

⁹Tuvan State University, Lenina 36, 667000, Kyzyl, Russia

¹⁰Bio-Clim-Land Center of Excellence, National Research, Tomsk State University, Lenina Pr., 36, Tomsk, 634050, Russia

Correspondence to: Feurdean@em.uni-frankfurt.de; angelica.feurdean@gmail.com

25

30

35 **Abstract.** Wildfire is the most common disturbance type in boreal peatland forests and can trigger significant changes in forest composition. Waterlogging in peatlands determines the degree of tree cover and the depth of the burnt horizon associated with wildfires. However, interactions between peatland moisture, vegetation composition and flammability, and fire regime in forested peatland in Eurasia remain largely unexplored, despite their huge extent in boreal regions. To address this knowledge gap, we reconstructed the Holocene fire regime, vegetation composition and peatland hydrology at two sites in Western Siberia
40 in the Tomsk Oblast, Russia. The palaeoecological records originate from forested peatland areas in predominantly light taiga (*Pinus sylvestris*-*Betula*) with an increase in dark taiga communities (*Pinus sibirica*, *Picea obovata*, *Abies sibirica*) towards the east. We found that past water levels fluctuated between 8 and 30 cm below the peat surface. Wet peatland conditions promoted deciduous trees (i.e., *Betula*) and *Sphagnum*, whereas dry peatland conditions favoured conifers and a higher dark-to-light-taiga ratio, thus perhaps greater forest density. The frequency and severity of fire increased with a declining water
45 table and enhanced fuel dryness and flammability, and an intermediate dark-to-light-taiga ratio. We found that the probability of greater fire severity increased when the water level declined below 20 cm suggesting a tipping point in peatland hydrology at which the wildfire regime intensifies. On a Holocene scale, we found two periods with contrasting moisture-vegetation-fire interactions: (1) severe fires were recorded between 7.5 and 4.5 ka BP with lower water level and an increased proportion of dark taiga and fire avoiders (*Pinus sibirica* at Rybnaya and *Abies sibirica* at Ulukh-Chayakh) with a predominantly light taiga
50 and fire-resister community characterized by *Pinus sylvestris*; (2) severe fires occurred over the last 1.5 ka BP and were associated with fluctuating water tables, a declining abundance of fire avoiders, and an expansion of fire invaders (*Betula*). These findings suggest that frequent high-severity fires can lead to compositional and structural changes in forests when trees fail to reach reproductive maturity between fire events or where extensive forest gaps limit seed dispersal. This study also shows prolonged periods of synchronous fire activity across the sites, particularly during the early to mid-Holocene, suggesting
55 a regional imprint of centennial to millennial-scale Holocene climate variability on wildfire activity. Increasing human presence in the region of the Ulukh-Chayakh Mire near Teguldet over the last four centuries drastically enhanced ignitions compared to natural background levels. Frequent warm and dry spells predicted by climate change scenarios in Siberia for the future will enhance peatland drying and may convey a competitive advantage to conifer taxa. However, dry conditions, particularly a water table decline below the threshold of 20 cm, will probably exacerbate the frequency and severity of wildfire,
60 disrupt conifers' successional pathway and accelerate shifts towards more fire-adapted deciduous broadleaf tree cover. Furthermore, climate-disturbance-fire feedbacks will accelerate changes in the carbon balance of forested boreal peatlands and affect their overall future resilience to climate change.

1 Introduction

Wildfire is the most common type of disturbance in boreal forests and forested peatlands (Kharuk et al., 2021 and refs therein).
65 It can change tree community composition and accelerate climate warming via carbon release into the atmosphere and

alteration of the radiative balance due to changes in land surface albedo (Rogers et al., 2015; Kharuk et al., 2021). However, the impacts of wildfire on vegetation and climate strongly depend on the fire regime. High-intensity crown fires kill most trees and alter species composition for an extended period of time. In the short term, such fires release black carbon aerosols (Rogers et al., 2015). When these aerosols persist in the atmosphere, this leads to a medium-term increase in albedo and ultimately to regional cooling. Contrastingly, surface fires typically do not kill mature trees or trigger stand-scale forest replacement and have little effect on albedo. Fire types in the Siberian boreal forests and forested peatlands are often litter-fuelled surface fires that infrequently transition to the crown, depending on forest composition (Sannikov and Goldammer, 1996; Grooth et al., 2013; Kharuk et al., 2021). A high diversity of fire-related plant functional types (PFT), including resisters (*Pinus sylvestris*, *Larix sibirica*, *L. gmelinii*), avoiders (*Abies sibirica*, *Picea obovata*, *Pinus sibirica*), invaders (*Betula pubescens*, *B. pendula*) and endurers (*Populus tremula*, *B. pubescens*) dominate these forest communities (Rowe, 1983; Agee, 1998; Wirth, 2005). Fire resisters, invaders, and endurers are commonly associated with a high-frequency but low-intensity surface fires, whereas avoiders are associated with a low-frequency but high-intensity crown fires (Goldammer and Furyaev, 1996; Wirth, 2005). As these two types of fire can have a different net effect on tree community composition and climate, it is essential to understand the patterns and drivers behind each fire regime. This understanding will also improve the prediction of the impact of changes on the extent, frequency, and severity of fire in the future.

A fire regime emerges from the combination of ignition sources, climatic conditions, fuel properties and human activities. The interactions between these factors are spatiotemporally complex (Moritz et al., 2014; Andela et al., 2017). The spatial complexity of wildfire patterns is particularly accentuated in open and forested peatlands, where the variability in the local peat moisture content, vegetation composition and structure results in pronounced small-scale heterogeneity (Camill et al., 2009; Magnan et al., 2012; Kuosmanen et al., 2014; Remy et al., 2018; Barhoumi et al., 2019; Stivrints et al., 2019; Feurdean et al., 2020a). The water-logged conditions that prevail in peatlands can limit the depth to which the fuel is burnt. However, this attenuation effect decreases with water table decline and increasing tree cover (Whitman et al., 2018). Unlike drained peatlands with shrub and tree-dominated communities, well-hydrated peatlands composed of a dense *Sphagnum* cover remain wet despite droughts and exhibit only slight to medium fire damage (Whitman et al., 2018; Gewin, 2020). This pattern in fire dynamics is particularly concerning because climate warming accelerates woody encroachment in peatlands and increases fuel availability for peatland forest fires (Kharuk et al., 2021). Intensification of fire severity may also lead to a shift from conifer towards deciduous tree dominance (Kelly et al., 2013; Mekonnen et al., 2019). Understanding how hydrological conditions in peatlands are influenced by climate change and how this may interact with peatland forest composition and fuel flammability is a key research priority that remains largely unexplored (Page et al., 2009; Walker et al., 2009; Kharuk et al., 2021).

Reducing uncertainty in predicting the future trajectories of fire regimes and the impact of changing fire activity on the functioning of boreal ecosystems requires approaches with a broad spatial and temporal scope (Kasischke et al., 2010). Tree-dominated ecosystems have long successional cycles ranging between decades and centuries. Palaeoecological records capture long time scales and are, therefore, particularly suitable to track the fire dynamics of forest ecosystems (Whitlock et al., 2018). However, studies of past millennial-scale variability in fire regimes in Siberian boreal forests, forested peatlands and forest

100 steppe started to emerge only recently (Feurdean et al., 2020a; Rudaya et al., 2020; Glückner et al., 2021; Barhoumi et al., 2021). Although Siberia is covered by extensive forested peatlands, particularly to the west (Vompersky et al., 1994; Liss et al. 2001; Kirpotin et al., 2021), no studies have explicitly explored the interactions between peatland moisture, vegetation composition, and fire regime in this vast region.

In this study, we used multi-proxy analyses (pollen, non-pollen palynomorphs, microcharcoal, macrocharcoal morphologies, 105 testate amoeba, geochemistry) on two new peat records from Tomsk Oblast to explore the patterns and drivers of the western Siberian fire regime throughout the Holocene. Specifically, we have explored how hydrological conditions in peatlands affect fuel dryness and how peat hydrology interact with tree community composition with regard to plant functional types and fuel flammability in driving the frequency and type of fires. Combining our new data with our two other published records from the region (Feurdean et al., 2020a) allowed a quantification of the regional changes in biomass burning and fire frequency.

110 2 Material and Methods

2.1 Geographical location and site selection

The study areas stretch along a west-east transect in a permafrost-free area in Tomsk Oblast, Siberia (Bleuten and Filippov, 2008; Kirpotin et al., 2021). The region has a continental boreal climate. The mean annual temperature is 0°C, the mean temperature of the coldest/warmest season is -17°C/17°C and mean annual precipitation is ~ 525 mm falling predominantly 115 in the warm season (www.climexp.knmi.nl). Regionally forest is made up of both light (*Pinus sylvestris*, *Betula pubescens*, *B. pendula*, *Populus tremula*) and dark (*Pinus sibirica*, *Picea obovata*, and *Abies sibirica*) taiga, the latter in greater proportions towards the east (Laschinsky and Koroliuk, 2014). Fires, at present primarily ignited by humans (80%), occur predominantly between spring and early autumn, with summer and autumn fires being usually the most severe (Kukavskaya et al., 2014, 2016; Kharuk et al., 2021).

120 We cored two sites, Rybnaya Mire (N57,27566800°, E84,48758900°) located near the Rybnaya river, a tributary of the Ob river, and Ulukh-Chayakh Mire (N57,34326197°, E88,32306189°) located on a terrace of the Chulym river, near the village of Teguldet, about 30-40 km from the border of the Krasnoyarsk Krai (Fig. 1). The local peatland vegetation at both coring sites includes mesotrophic open sedge-*Sphagnum* communities with small birch and pine trees at Rybnaya Mire and standing dead *Pinus* tree trunks at Ulukh-Chayakh Mire. These dead trees reflect a rise in water levels across the mire due to road 125 construction in the 1960s. Peat cores (50 cm length, 5 cm diameter) totalling 400 cm in length at Rybnaya Mire and 348 cm at Ulukh-Chayakh Mire were extracted with a Russian-type corer in 2017 and 2019, respectively.

2.2 Chronology

We established the chronology of both cores based on AMS radiocarbon measurements performed at Isotoptech, Debrecen, 130 Hungary (File S1). The ¹⁴C AMS age estimates were converted to calendar years BP using the IntCal20 data set of Reimer et al. (2020) and we constructed the age-depth models using the smooth spline method implemented in CLAM software (Blaauw,

2010). In the age-depth models, we assigned a surface age of -67 cal yr BP (coring year 2017 for Rybnaya) and -69 cal yr BP (coring year 2019 for Ulukh-Chayakh) to the surface samples of each sequence.

135 2.3 Fire history

2.3.1 Charcoal-based reconstructions of fire history

We inferred changes in local-scale fire regime based on charcoal particles identified in peat samples of 2 cm³ extracted at 1 cm contiguous intervals (Whitlock and Larsen, 2001). Sample preparation, identification and the categorisation of charcoal morphotypes followed Feurdean et al. (2020a) and Feurdean (2021). To deduce the predominant fire type, we pooled the charcoal morphologies into two categories: non-woody (graminoids, forbs) and woody types (deciduous leaves, wood, and resins) and calculated their ratio. These two categories represent the dominant fuel sources for surface and crown fires, respectively (Enache et al., 2006; Jensen et al., 2007; Courtney-Mustaphi and Pisaric, 2014; Feurdean et al., 2017; 2020a; Feurdean, 2021). At Ulukh-Chayakh, we additionally tested the aspect ratio (length:width) of charcoal particles in selected samples to discriminate between the relative dominance of graminoids versus leaves and wood morphologies (Feurdean, 2021; Vachula et al., 2021). We grouped the charcoal fractions into two size classes (150-500 µm and >500 µm) to deduce the approximative proximity of fires, i.e., local versus on-site fire events (Adolf et al., 2018). We calculated the influx of charcoal morphotypes and size classes (particles cm⁻² yr⁻¹) by dividing their respective concentration (particles/cm³) by the sediment deposition rate determined from the age-depth models (yr/cm).

We estimated the frequency and severity of fire episodes using charcoal peaks identified from macrocharcoal particles using the method of Higuera et al. (2009). This methodology involved interpolating CHAR values to constant time steps (30 yr), decomposing the record into a background and a peak component (local fire episodes) using a robust LOWESS smoother (900 yr), and evaluating charcoal peaks using the 95th percentile of the modelled noise distribution. We used the signal-to-noise index (SNI) to assess the suitability of the record for peak detection (Kelly et al. 2011; File S2).

To integrate fire history across the study area, we created composite records of CHAR and fire frequency (FF) by combining the two new records with our two published profiles from Plotnikovo Mire (Feurdean et al., 2019; 2020a; Fig. 1). Composite CHAR was created using the Power et al. (2010) protocol implemented in the R palaeofire package version 4.0 (Blarquez et al., 2014) with the following setup for Z score transformation: 8.5 ka to present as the base period, and LOWESS smoothing to a 25-yr mean. The composite FF was created by pooling the identified fire events from all four records and applying a 150-yr smoothing.

160 2.3.2 Fire type identification from satellite images and forest statistics

To evaluate the ability of charcoal records to capture the occurrence and severity of fires near the two sites, we used satellite imagery of vegetation cover from LANDSAT 4-5 TM, LANDSAT-7 ETM+, and LANDSAT 8 OLI/TIRS (<https://earthexplorer.usgs.gov/>). We also used observation-based fire data from the Forest Department of the Tomsk Region

from 1950 to 2020 CE and data from MODIS sensors onboard the NASA Terra-1 and Aqua-1, which detect forest fires
165 (<http://fires.ru/>). All maps and images were processed using ESRI ArcGIS version 10.8.

2.4 Pollen-based reconstruction of vegetation dynamics

To reconstruct past vegetation dynamics, we analysed pollen and non-pollen palynomorphs (NPPs) mainly at 4 cm intervals in both profiles following the protocol of Bennett and Willis (2001). A pollen sum exceeding 500 terrestrial pollen grains, excluding Cyperaceae, was counted for most samples and converted into percentages based on the total terrestrial pollen sum.
170 The percentages of Cyperaceae, ferns and mosses were calculated relative to the terrestrial pollen sum and their respective individual sum. The bottom part of the Ulukh-Chayakh profile i.e., from 300-348 cm (age >6500 cal yr BP) contained extremely poorly preserved pollen and was excluded from further analysis. Microscopic charcoal particles longer than 10 µm were also counted alongside pollen, and their influx was estimated relative to the added *Lycopodium* marker and the sediment deposition rate.
175 We assigned pollen-derived tree taxa to one of the fire-related functional PFT groups - resisters (*Pinus sylvestris*, *Larix* spp.), avoiders (*Picea obovata*, *Abies sibirica*, *Pinus sibirica*), or invaders (*Betula* spp.). The two tree birch species (*B. pubescens* and *B. pendula*) and the dwarf shrub (*B. nana*) could not be differentiated by pollen analysis and were considered as *Betula* spp. However, at present day, *B. nana* tends to be found only sparsely in this region. Due to its poor pollen preservation, *Populus*, a fire endurer, is mostly absent from our pollen records. The necessity to combine the *Betula* pollen signal and the
180 paucity of pollen of *Populus* may have resulted in the endurer PFT group failing to be represented in our records. The PFT-specific fire coping adaptations relate to a taxon's strategy to complete its life cycle in the context of a given fire regime (Gill, 1981; Wirth, 2005). We also separately grouped the tree pollen into taxa associated with dark taiga (*Pinus sibirica*, *Abies sibirica*, *Picea obovata*) and light taiga (*Pinus sylvestris*, *Betula* spp., *Larix* spp.) and created an index for dark-to-light taiga composition, an approximation of forest density, by calculating the ratio between the relative abundance of dark and light taiga
185 tree taxa (Higuera et al., 2014). Additionally, we pooled the non-arboreal pollen types as shrubs, herbs, ferns and mosses. To determine the regional changes in tree community composition, we created composite records of PFTs. This was done by averaging the Z-scores and smoothing them to 300 years with a locally weighted regression and a 95% confidence envelope on the same sites used for composite biomass burning reconstructions in the R palaeofire package.

190 2.5 Climate reconstruction based on testate amoebae and water table depth

To determine changes in peatland hydrology at each site, we used testate amoeba-based water table reconstructions. The testate amoebae preparation and identification were conducted at 4 cm intervals following the methods of Hendon and Charman (1997) and Charman et al. (2000). Based on the available literature (Grospietsch, 1958; Ogden and Hedley, 1980; Mazei and Tsyganov, 2006), we identified and counted a minimum of 150 testate amoebae per sample. No testate amoebae were present
195 in core segments below 285 cm (age > 6000 ka BP) in the Ulukh-Chayakh sequence. We used the transfer function developed

for the pan-European region (Amesbury et al., 2016) to derive the water depth. The sample errors were generated using 1000 bootstrapping cycles (Line et al., 1994). For the composite water table reconstruction, we used the same setting as for pollen.

2.6 Elemental geochemistry

200 The concentration of the geochemical element Ti was measured using a non-destructive Niton XL3t 900 X-Ray Fluorescence analyser (fpXRF) to determine the potential influence of water influx (i.e., floods) on mire water table. Sedimentary Ti NCS DC73308 was used as a Certified Reference Material (CRM). Measurements followed the procedure described by Hutchinson et al. (2016).

205 2.6 Numerical methods

To quantify the relationship between peatland moisture conditions, biomass burning, and the relative abundances of PFTs and dark-to-light taiga composition, we performed a correlation analysis. We used microcharcoal (CHAR particles <150 µm) and machrocharcoal (CHAR particles >150 µm) influx to determine regional and local biomass burning, pollen percentages as indicators of the three PFTs (resisters, invaders, avoiders, and others) at local to regional scales, and testate amoebae-based water table depth as a proxy of local peatland moisture. The group “others” includes pollen and spores of shrubs, herbs, ferns and moss. Before the analysis, we interpolated all datasets fed into the model to a 100- year interval using linear interpolation. We also developed generalised linear models (GLMs) to explore the response of biomass burning (CHAR particles >150 µm) to changes in peatland moisture conditions (DTW) and to test for the existence of thresholds in the water level at which biomass burning may increase.

215

3. Results

3.1 Chronology

The age-depth model of the Rybnaya sequence spanned ~ the last 8400 years and showed a mean peat accumulation rate of 25 yr/cm (ranging between 6-36 yr/cm). The Ulukh-Chayakh sequence may cover ~ the last 8500 years, but the chronology of the bottom part of this site (>6000 years) relied on linear extrapolation and it is therefore highly uncertain (File S1). The subsequence covering the last 6000 years had an average temporal resolution of 21 yr/cm (ranging between 3-37 yr/cm).

220

3.2 Fire type identification from charcoal, satellite images and forest statistics

The comparison of sedimentary charcoal from sub-recent samples with satellite imagery of the Plotnikovo Mire (sites SP and SD) confirmed that the marked increase in charcoal particles towards the present day at SD location closely agrees with the occurrence of high-severity fires (Feurdean et al., 2020a). Satellite images and official forest statistics show no evidence of recent fires in the vicinity of the Rybnaya and Ulukh-Chayakh sites, except for a small fire documented in 1993 near the Ulukh-Chayakh site (Fig.1). This aligns with the scarcity of charcoal pieces from sub-surface samples in both cores (Fig. 2; File S3). A high-severity fire event occurred in 2015 some ca. 30 km from Ulukh-Chayakh, which is not documented in our surface

225

230 charcoal record and corroborates the localised origin of charcoal found in this sedimentary profile. Samples from the Ulukh-Chayakh site show a minor increase in charcoal (2-7 pieces; up to $0.6 \text{ cm}^{-2} \text{ yr}^{-1}$) ca. fifty years ago, possibly related to a local fire event in 1993. The first clear charcoal peak at this site (250 pieces; $7.3 \text{ cm}^{-2} \text{ yr}^{-1}$), with numerous wood fragments, was found ca. 200 years ago, whereas the preceding charcoal peak (400 pieces; $9.2 \text{ cm}^{-2} \text{ yr}^{-1}$) was ca. 900 years ago (Fig. 2; File S3). These two peaks therefore demonstrate the intermittent occurrence of infrequent, high-severity local fires that produce large quantities of charcoal, embedded in what is otherwise predominately a surface fire regime generating low levels of charcoal. It also indicates that the charcoal peak extraction method applied here reliably reflects the occurrence of high-severity local fires.

3.3 Site-specific and composite record of biomass burning and fire frequency

240 Both records displayed a high signal-to-noise index (File S2) with almost all peaks well above 3 (a mean of 4.98 at Rybnaya and 6.45 at Ulukh-Chayakh), the theoretical minimum value for justification of peak analysis (Kelly et al., 2011). At Rybnaya, CHAR was highest between 7.5 and 6 ka BP and around 4.5 ka BP, whereas the frequency of high-severity fires inferred from the charcoal peak magnitude and abundance of woody morphologies ranged between 1 and 2 fires/900 yr (Figs. 2, 4; Files S2, S3). At Ulukh-Chayakh, CHAR was highest >6 and 3.5 ka BP, and the fire frequency varied between 2 and 5 fires/900 yr.

245 Both profiles showed a decline towards a minimum in CHAR and frequency of severe fires, i.e., low peak magnitude and reduced occurrence of woody charcoal morphologies, approximately between 4 and 1.5 ka BP (3-1.5 ka BP at Ulukh-Chayakh). The CHAR increased towards the present at both sites, but the charcoal peak amplitude was considerably higher at Ulukh-Chayakh.

The composite CHAR record showed augmented regional biomass burning between 7.5 and 4 ka BP (positive Z-scores) with peaks centred at 7.5 and 4.5 ka BP (Fig. 5). We found a pattern of CHAR decline towards a minimum in the composite profiles between 4 and 2 ka BP (negative Z-scores), followed by another increase between 2 ka BP to the present (positive Z-scores). The composite fire frequency showed an increase until ca. 5-4.5 ka BP and then again from 2 ka to the present.

3.4 Site-specific and composite record of vegetation composition and forest density

255 Early Holocene data is only available from the Rybnaya site (8.5 and 7.5 ka BP), indicating that light taiga and the fire-invader *Betula* were most abundant (up to 60%) during that period, whereas other light taiga fire-resisters (*Pinus sylvestris*) and dark taiga and fire avoiders (mainly *Picea obovata*), had an abundance of 10% and 5%, respectively (Fig. 3; File S4a). The high proportion of light taiga resulted in a low dark-to-light taiga ratio of 0.1, and perhaps a low forest density. An increased dark-to-light taiga ratio (> 0.2), was identified between 7.5 and 4 ka BP at Rybnaya and between 5 and 2.5 ka BP at Ulukh-Chayakh (Fig. 5; File S4a,b). The light taiga was composed of up to 70% fire-resisters (*Pinus sylvestris* and *Larix spp.*), while the dark taiga, with its fire avoiders (*Pinus sibirica*, *Picea obovata*, and *Abies sibirica*), made up to 25% of the forest. A decline in fire-resisters (down to 50%), avoiders (below 10%) and dark-to-light taiga ratio and increasing invader abundance (up to 60%) occurred after 4.5 ka BP at Rybnaya and after 2.5 ka BP at Ulukh-Chayakh. Between-site variability was highest within the

avoider group. *Picea obovata* had higher values at Rybnaya (up to 10%) than at Ulukh-Chayakh (3%), whereas *Abies sibirica* was more abundant at Ulukh-Chayakh (15%) than at Rybnaya (7%; Fig. 4). The composite vegetation record showed a higher proportion of fire-resisters between 7.5 and 5 ka BP, fire avoiders between 7.5 and 3 ka BP, and fire invaders over the last 4.5 ka BP (Fig. 5).

Understory vegetation primarily reflects local peatland development. It was predominantly composed of herbs/sedges (Cyperaceae; up to 20%) and ferns (Polypodiaceae, up to 20%) and *Equisetum* (up to 25%) at both sites until ca. 4.5 ka BP. An increase in the abundance of shrubs (Ericaceae; up to 10%) and moss (*Sphagnum*; highly fluctuating between 10 and 100%) at Rybnaya, and Ericaceae (up to 5%) and moss (*Sphagnum*; up to 100%) at Ulukh-Chayakh was noted over the last 4.5 ka BP (Fig. 3; File S4a,b). The sum and individual levels of herbaceous taxa, including those potentially related to human impact (*Artemisia*, Asteraceae, Chenopodiaceae, Urticaceae, Brassicaceae, *Plantago lanceolata*), were constant throughout the profiles. However, their abundance increased slightly over the last 1.5 ka BP (Fig. 3; File S4a,b). Coprophilous dung spores, *Podospora* and *Sporormiella*, were present throughout both records, although they became more abundant between 4.5 and 3.5 ka BP and 3.2 and 1.0 ka BP at Ulukh-Chayakh and over the past millennium at Rybnaya (File S4a,b).

3.5 Site-specific and composite records of peatland moisture conditions

At Rybnaya, the testate amoeba-based reconstruction of the water table indicated relatively high levels (around 15 cm below surface) between 8.5 and 7.5 ka BP, 4 and 2.5 ka BP, and over the last 1 ka BP (Fig. 4). We reconstructed low levels (20-30 cm below surface) for the time interval between 7.5 and 4.5 ka BP and 2.5 and 1 ka BP (Fig. 4). At Ulukh-Chayakh, the water level was high (around 15 cm below surface) between 3.5 and 2.5 ka BP, 2 and 0.5 ka BP, and towards the present day. There, times of low water table (34-20 cm below surface) occurred between 6 and 3.5 ka BP, 2.5 and 2 ka BP, and around 0.5 ka BP. The composite record of the water table showed high levels approximately between 8.5 to 7.5 ka BP, 3.5 and 3 ka BP and during the last 1.5 ka BP, and low levels between 7.5 and 4.5 ka BP, and around 2.5 ka BP (Fig. 5).

3.6 Numerical analysis

The correlation analysis at Rybnaya showed that micro- and macrocharcoal were highly correlated and positively associated with the proportion of fire avoiders and light-to-dark taiga index and negatively with resisters and invaders (Fig. 6a). However, not all these relationships were statically significant (Appendix A1). At Ulukh-Chayakh, micro- and macrocharcoal were weakly correlated. Microcharcoal was positively associated with fire avoiders, resisters, others and light-to-dark taiga index and negatively with invaders (mostly statically significant; Fig. 6a, b). All other relationships i.e., between PFTs and macrocharcoal and depth to water table were statistically non-significant (Fig. 6a; Appendix A1) The generalised linear model showed no significant influence of water levels above 20 cm on the probability of fire occurrence and severity at both sites (Fig. 6c,d). However, a drop in water level below 20 cm contributed statistically significantly to the increase in fire occurrence and severity at Rybnaya but had no statistically significant influence at Ulukh-Chayakh.

4 Discussion

4.1 Changes in fuel, fire type and fire frequency over the Holocene

300 Stand-replacing fires combust substantial amounts of biomass because they burn entire trees. Such fires manifest themselves
in a greater abundance of charcoal particles that exceed the quantities typically produced by surface fires (van Marle et al.,
2017). This feature allows a distinct separation of charcoal peaks from charcoal background levels, which yields a higher
signal-to-noise index (Higuera et al., 2005; 2009; Courtney Mustaphi and Pisaric 2014). The comparison of sedimentary
305 charcoal from sub-recent samples with satellite imagery of the Plotnikovo Mire (Feurdean et al., 2020a) and the vicinity of the
current study sites (Rybnaya and Ulukh-Chayakh) demonstrates the intermittent occurrence of high-severity local fires, that
produce large quantities of charcoal, embedded in what is otherwise predominately a surface fire generated low level of
charcoal.

We reason that the prevalence of a high charcoal influx with well-defined charcoal peaks and abundant woody morphotypes
found between ca. 7.5 and 4 (3) ka BP at both sites, and over the past 1.5 ka BP at Ulukh-Chayakh, is indicative of the
310 predominance of high-severity local fires (Fig. 2; File S3). Between 4 (3) and 1.5 ka BP, the observed low charcoal influx and
peak magnitude were coupled with a lower abundance of woody morphologies at both sites, although the latter feature was
more pronounced at Ulukh-Chayakh (Fig. 2; File S3). For this period, we inferred predominately low-temperature surface fires
that mostly burned understory biomass. Experimental production of charcoal additionally provides evidence that graminoids
and *Sphagnum* retain a high charred mass only during low-temperature fires (Hudspith et al., 2017; Feurdean, 2021).
315 Measurements of aspect ratio ($L:W$) and surface area on selected charcoal samples at Ulukh-Chayakh show that the aspect
ratio only partially agrees with the predominant fuel type inferred from charcoal morphologies. A lower aspect ratio of charcoal
particles, typical for wood and leaf morphotypes, was what we expected at times of increased woody morphologies, and an
increased aspect ratio with a rise in the relative proportion of graminoids (Feurdean, 2021; Vachula et al., 2021). This is
probably the result of a strongly mixed morphology of charcoal types in the same sample (Fig. 2). Our measurements of
320 charcoal surface area indicate high values at times of an increased abundance of leaves (Fig. 2; File S3) with a typically high
surface area (Crawford and Belcher, 2014; Feurdean, 2021).

Our reconstructed fire return interval is ~ 450 yr (2 fires/900 yr) for both sites throughout most of the Holocene (Fig. 4).
However, fire frequency (at Rybnaya) and severity (at Ulukh-Chayakh) increased over the last 1.5-2 ka BP compared to the
long-term averages of both records suggesting an intensification of fire activity during the most recent two millennia. The
325 composite fire frequency derived from all four records also shows increasing biomass burning and fire frequency over the last
2-1.5 ka BP (Fig. 5). The reconstructed fire return interval (FRI) in the study area is in line with other sites located in forested
peatlands (*Pinus-Betula* dominated, *Picea obovata*, *Abies sibirica*) in Russia, which report an FRI range between 100 and 600
yr (Barhoumi et al., 2019, 2020; 2021). In contrast to a study in the northern Ural region (Barhoumi et al., 2019), we have not
found a gradual increase in fire frequency from the early Holocene towards the present, but rather two distinct periods of
330 enhanced activity, between 7.5 and 4 ka BP, and approximately over the last 2 ka BP (Fig. 5), a pattern more similar to sites
in the West Siberian Plain (Rudaya et al., 2020) and Lake Baikal area (Bourhami et al., 2021).

4.2 Drivers of vegetation and fire regime change

Climate conditions influence peatland hydrology, fuel dryness and flammability as well as vegetation composition with regard to moisture- and fire-related PFTs. The relative abundances of the fire-related PFTs, in turn, influence the prevalence of specific fire regime types, e.g., low-intensity vs. high-intensity regimes. Below, we discuss how hydrological conditions in peatlands affect fuel dryness and how peat hydrology interact with peatland tree community composition and fuel flammability.

4.2.1 The influence of climate on peatland moisture and fuel flammability

Waterlogging in peatlands and the occurrence of a dense *Sphagnum* cushion can limit fire severity and reduce the depth of the burnt horizon. Moreover, a waterlogged horizon provides a substrate that serves as a seedbed for post-fire regeneration (Whitman et al., 2018; Gewin 2020). However, increasingly dry peatland conditions and consequently greater tree cover can reduce the limiting effect of waterlogging on fire severity (Magnan et al., 2012; Kettridge et al., 2015; Whitman et al., 2018; 2019; Loisel et al., 2020). Deep burns can even smoulder over winter and re-ignite the following spring (Scholten et al., 2021). Our water table reconstructions indicate that the water level fluctuated between 8 and 30 cm below the peat surface and was around 20 cm at both sites during large parts of the Holocene (Fig. 4). We found that the probability of fire occurrence and intensification of fire severity was greater at times when the water level dropped below 20 cm (drier conditions) and lower at water levels between 10 and 20 cm below the surface (wetter conditions), a pattern that was more evident at Rybnaya (Fig. 6 c,d). The stronger moisture-fire relationship at Rybnaya may be related to the higher amplitude of changes in local hydrological conditions and the overall drier conditions at this site (8-30 cm) compared to Ulukh-Chayakh (10-23 cm).

Water levels of mesotrophic mires are co-regulated by the balance between precipitation (P) and evapotranspiration (Et), and external water inflow (Chambers and Charman, 2004). The detrital element Ti, a possible indicator of water influx in minerogenic mires, increased notably between 4 and 3 ka BP and 1.5 and 0.1 ka BP at Ulukh-Chayakh, but only slightly over the last 1 ka at Rybnaya (File S5). This suggests increasing fluvial input, reflecting possible flooding or channel position change as a transport mechanism for the delivery of such material at Ulukh-Chayakh (File S5; Leshchinskiy et al., 2011). Mire type development and changes in peat plant communities could have also influenced peatland hydrology (Galka et al., 2016; Kurina and Li, 2018; Blyakharchuk and Kurina, 2021). At Rybnaya, the minerogenic stage of the mire occurred from 8 to 4.5 ka BP and the water level was more stable. With the meso-oligotrophisation of the mire, indicated by the appearance of Ericaceae at 4.5 ka BP, and the dominance of *Sphagnum* from 3.6 ka BP to the present, the water level and the amplitude of its fluctuation rose and the occurrence of peat fires declined (Figs. 3, 4; File S4a,b). At the Ulukh-Chayakh site, Ericaceae developed at 4.5 ka BP and remained dominant up to the present, whereas *Sphagnum* established around 2 ka BP and became dominant only in the past decades (Fig. S4). Further, mire margins with thinner peat depth burn more frequently than thicker peat (Turunen et al., 2001). Thus the location of the Ulukh-Chayakh sampling point close to the mineral margin of the mire (ca. 160 m), as opposed to a few kilometres at Rybnaya, could have sustained a high fire occurrence at this site until recently. Lastly, fire itself can influence peatland hydrology, with field measurements indicating an age-dependent water table lowering

after burning (Holden et al., 2015). It is beyond this study to discuss all the details of the drivers behind peatland hydrology, however, given our main focus is the link between state changes in peat hydrology and fire occurrence and severity.

At a Holocene scale, the intensity of fires and/or fire size was greatest between 7.5 and 6.5 ka BP and 5 and 4 ka BP, but the timing was not fully synchronous between the sites (Figs. 4, 5). From 9 to 4.5 ka BP, annual temperatures in the Northern Hemisphere were up to 3.5°C warmer than at present (Fig. 5) and exhibited a more pronounced seasonality, i.e., higher summer and lower winter temperatures (Kauffman et al., 2020; Bova et al., 2021). Proxy records from Siberia attest to warmer and drier-than-present climate conditions between 9 to 6 ka BP with a temperature and moisture optimum between 6 and 4.5 ka BP (Borisova et al., 2011; Groisman et al., 2012). Although variable at a wide regional scale in Siberia, multiproxy records from West Siberia including testate amoebae show strikingly similar hydrological conditions to ours i.e., moist between 8.1 and 7.4 ka BP, followed by dry conditions between 7.4 and 5.1 ka BP, associated with a diminished North Atlantic influence at this latitudinal band (see review of Mikhailova et al., 2021). Other testate amoebae-based hydrological reconstructions in the study region show partial agreement to ours, however, there are coeval shifts to dry conditions between 7.5 and 5.5 ka BP in all these records (Kurina et al., 2018; 2021), the latter study primarily linked changes in the surface wetness of mires to those in precipitation. It is possible that warm summer temperatures enhanced evapotranspiration, which, combined with overall reduced precipitation, contributed to a lowering of peatland water levels, leading to drier surface conditions from 8 to 5 ka BP (Fig. 5).

Contrastingly, the interval of lowest biomass burning and fire severity approximately between 4 (3) and 1.5 ka BP coincides with one of the periods of wettest peatland conditions during the Holocene (at Rybnaya) or a highly fluctuating water table (at Ulukh-Chayakh; Fig. 4). Annual and summer temperatures declined after 4.5 ka BP to values below the present day (Kauffman et al., 2020; Bova et al., 2021). Our reconstructed hydrological conditions align with published records from western Siberia in revealing wet conditions between 5.1 and 1.4 ka BP with maximum wetness between 4.5 and 3 ka BP (Mikhailova et al., 2021; Blyakharchuk and Kurina, 2021). Regionally, this wet and cool trend (Groisman et al., 2012) led to high fuel moisture and reduced fuel flammability. Although biomass burning and the frequency of severe fires show more centennial-scale variability over the past 1.5 ka BP, this matches only partially known climatic fluctuations, such as the Medieval Climate Anomaly (MCA1200-650 cal yr BP) and the Little Ice Age (LIA; 700-400 cal yr BP). Notably, an increased charcoal peak amplitude occurred during the second part of the LIA (400-200 cal yr BP), at a time of extremely wet conditions (Fig.4; Feurdean et al., 2019). We hypothesise that the climatic linkage between the cool and wet conditions, that imply high fuel moisture conditions and typically low fire activity, was likely disrupted by anthropogenic activity towards the present day (Fig 3; File S4a, b), as document by other pollen records in the region (Blyakharchuk et al., 2019). Historical records indicate the widespread colonisation of the region by Russian (i.e., from the west) at this time that was associated with the conversion of forest to arable land and pastures, and likely increased anthropogenic ignitions (Naumov, 2006). Most recently, local fire suppression near the village of Teguldet near Rybnaya mire may have hampered contemporary fire spread on the mire leading to the recent pattern of low fire activity.

Nevertheless, humans may have altered the frequency of fires earlier than the widespread Russian colonisation of the region. A GIS survey of archaeological finds in the Tomsk region reveals a high density of sites, but no distinctions between cultural phases of these sites exist (Zolnikov et al., 2020). In parallel with modern settlement distribution, most of these ancient sites were situated near rivers favouring the floodplain terraces of river valleys (Zolnikov et al., 2020). Another study in the Lower Tom basin attests to the presence of tools that have the characteristics of sites commonly associated with the Neolithic and Bronze Age close to Rybnaya (Idimishev et al., 2018). There is little evidence of pollen indicative of human impact at Rybnaya until 400 years ago, but coprophilous-based evidence of herbivores (*Podospora*) is present from 7 ka onward and increased over the past 400 years, whereas fire activity was high from 7 to 5 ka BP (Fig. 4, S4). At the Ulukh-Chayakh site, coprophilous spores became more abundant between 4.5-3.5 ka BP (*Sporormiela*) and 3.2 and 1.0 ka BP (*Podospora* and *Sporormiela*) while fire activity increased between 4 and 3 ka and over the last millennium. Given the locations of our sites near rivers, the archaeological evidence suggests the possibility of some low level of human impact on vegetation and fire. Although, the effect of herbivores on forest composition and density is well known (Morales-Molino et al., 2019), it is difficult to say from our records whether herbivores contributed to the opening of the forest and facilitated the fire spread, and whether the coprophilous remains originated from wild or domesticated animals.

4.3 Feedbacks between peatland moisture, community PFT composition and fire regime

Peatland moisture conditions affect the relative abundances of PFTs. Fire-related PFTs, in turn, create positive or negative feedbacks for fire regimes via specific fire-related traits (Feurdean et al., 2020a; Kharuk et al., 2021). We found a dominance of open light taiga species and fire invaders (*Betula* spp) at times of higher water table (wet conditions) between 8.5 and 7.5 ka BP and after 4.5 and 3.5 ka BP (Figs. 2, 4). However, *Betula* spp. also remained dominant during the periods of water table decline recorded between 2.5 and 1.5 ka BP. *B. pubescens* grows on wet peatlands, whereas *B. pendula* grows more frequently on drier sandy soils and peatlands with higher fire incidence (Wirth, 2005; Groisman et al., 2012; Blyakharchuk, 2003). Unfortunately, pollen analysis cannot differentiate the prevalence of these birch species and the prevalence of their contrasting hydrological and fire requirements. An increased dark-to-light taiga ratio shows that denser peatland forests of light taiga and fire-resisters (*Pinus sylvestris*), interspaced with dark taiga taxa and fire avoiders tree taxa, primarily *Pinus sibirica* at Rybnaya and *Abies sibirica* at Ulukh Chayakh (only from 5.5 ka BP), prevailed between 7.5 and 4.5 ka BP (Fig.3; File S4a,b). This forest composition coincides with drier peatland conditions and an increased fire severity. Although not statistically significant, our correlations suggest a greater tolerance of deciduous trees to wetter conditions (high-water levels) compared to conifers (Fig. 6 a,b). Visual trends and correlation reveal that the biomass burning increase at intermediate dark-to-light taiga index (ranging from 0.15 to 0.3) was connected to dry peat conditions (Figs. 4, 6). This finding is consistent with emerging evidence on fire-fuel relationships in boreal forests, which points to a higher fire hazard in less dense as opposed to denser forests, which allow fuel to dry (Scheffer et al., 2012; Feurdean et al., 2020b). However, it should also be noted that the spatial scale of our

proxy reconstructions differs. Testate amoebae reveal local changes in peatland hydrology, whereas pollen and microcharcoal reflect local to regional changes in vegetation composition and fire activity (Bennet and Willis, 2001; Whitlock and Larsen, 2001; Lamentowicz et al., 2015; Blyakharchuk and Kurina, 2021). Nonetheless, although hydrological conditions may vary across a peatland locally, general trends in peat moisture should be consistent on a larger, regional spatial scale. It is also not possible to differentiate pollen originating from forested peatlands and that produced in adjacent (non-peatland) areas, which complicates the link between peat moisture and PFTs. However, given the vast extent of forested peatlands in the study area, we assume that most of the pollen rain reflects the tree composition of such peatlands.

The presence of the fire-resisters (i.e., *Pinus sylvestris*), in Siberia is associated with a light surface fire regime (mean FRI of 28 years) that is occasionally interrupted by longer-term (mean FRI of 200 yr) stand-replacing crown fire events (Ivanova, 2005; Kukavskaya et al., 2016). Fire avoiders typically experience severe stand-replacing crown fires at relatively long return intervals (mean FRI 150 years; range 99 and 300 years; Goldammer and Furyaev, 1996; Wirth, 2005; Kharuk et al., 2021).

The post-fire regeneration pathway of conifer species involves recruitment directly after burning (*P. sylvestris*) or after an early successional phase initiated by *Betula* (*Pinus sibirica*, *Picea obovata*, and *Abies sibirica*). A mixed taiga peatland forest with deciduous trees and dark taiga conifer taxa typically develops between 60-120 years after a stand-replacing fire, followed by the dominance of conifers as a late-successional stage (Tautenhahn et al., 2016; Coop et al., 2020; Kharuk et al., 2021). We reason that the prevailing periodicity of severe fires (every 180 to 450 years) between 7.5 and 4.5 ka BP provided fire-free periods that were long enough for the fire avoiders *Pinus sibirica*, *Picea obovata* and *Abies sibirica* to reach reproductive age. Alternatively, burning may have been too patchy to create sufficiently large forest gaps to limit seed dispersal and inhibit post fire recovery. *Pinus sibirica* is typically classified as a fire avoider but behaves like a fire resister when reaching old age, which could also explain its increased presence at times of high fire activity. Fire also allows dark taiga species to compete with *P. sylvestris* and become established beyond poor soils and boggy areas (Kharuk et al., 2021).

A different fire regime and altered vegetation feedbacks emerged over the last two millennia at Ulukh-Chayakh, the site that experienced more severe fire episodes. The intensification of local fire severity at Ulukh-Chayakh paralleled a decline in the proportion of fire avoiders (mostly *Abies sibirica* and *Picea obovata*), an increase in fire invaders (*Betula*) and the abundant occurrence of heathland shrubs (Figs. 3, 4). Shrubs burn hotter than other surface fuel types and reach temperatures that kill most conifer seedlings (Tautenhahn et al., 2016). High-severity burned areas have also been found to be more prone to repeated burning and have a more negative impact on conifers (Kukavskaya et al., 2016; Whitmann et al., 2019; Coop et al., 2020). We propose that the intensification in fire severity at Ulukh-Chayakh may have eliminated mature *Abies sibirica* and *Picea obovata*, limited their seed dispersal and disrupted their successional pathways. These dynamics resulted in a shift in tree community composition towards more post-fire-adapted invader communities with better dispersal capabilities, recruitment strategies for burned areas and rapidly maturing taxa. Our results support emerging findings of increases in deciduous trees (*Betula* or *Populus*) at the expense of evergreen conifers that are associated with contemporary warming and increasing fire severity in boreal forests and forested peatlands in Siberia and North America (Kelly et al., 2013; Tautenhahn et al., 2016; Mekkonen et al., 2019; Whitman et al., 2019; Kharuk et al., 2021). At Rybnaya, the modest increase in biomass burning over

the past two millennia coincided with wet conditions, the occurrence of *Sphagnum*, and a peatland forest composition that remained dominated by *Betula*, indicating that fire has played a more limited role in tree community dynamics at this site. The post-disturbance (fire, insect outbreak) forest regeneration pathway in Siberia also involves *Populus tremula*. Eurasian aspen forest communities can reach hundreds of years in age and change forest composition to *Populus* dominance where regeneration of dark taiga conifers is absent (Tautenhahn et al., 2016; Kharuk et al., 2021). Although we have not encountered *Populus* in our pollen record due to its poor pollen preservation, it cannot be discounted that *Populus* may have been a part of post fire forest succession pathways in the past.

475

Conclusion

This study provides novel insights into past fire regimes based on Holocene records from forested peatlands in western Siberia. It demonstrates that peatland hydrology is a critical determinant of fuel dryness, plant community composition (in terms of peat plant composition and fire-related plant functional types) and fire regime. Wet peatland conditions promoted open tree communities with deciduous trees typically classified as light-taiga taxa and fire invaders (*Betula*), whereas dry peatland conditions facilitated conifers and therefore a high dark-to-light-taiga ratio. Although the probability of fire occurrence in these peatlands increased with drier conditions and a high dark-to-light-taiga ratio, the greatest fire incidence occurred when the water level declined below 20 cm and at an intermediate forest density. At a Holocene timescale, we found two periods with contrasting moisture-vegetation-fire interactions. During the first period, an enhanced fire severity episode occurred between 7.5 and 4.5 ka BP and was associated with a low water level and a higher proportion of dark taiga/fire avoider taxa (mainly *Pinus sibirica* at Rybnaya and *Abies sibirica* at Ulukh Chayakh) in a light taiga and fire resister (*Pinus sylvestris*) dominant community. However, the second period of increased fire severity (i.e., last ca. 1.5 ka BP) coincided with a reduced abundance of fire avoiders (mostly *Abies sibirica* and *Picea obovata*) and an expansion of fire invaders (*Betula*). These community changes demonstrate that frequent fires of higher severity can lead to compositional changes in forest tree communities, either because the trees were unable to reach their reproductive age between burning events, or the fires created substantial forest gaps that hindered seed dispersal. This study also shows certain prolonged periods of synchronous fire activity across the sites, suggesting that the magnitude of centennial to millennial-scale climate variability over the Holocene was marked enough to drive fire regimes on a regional scale.

Based on our findings from fossil records, the frequent warm and dry spells predicted by climate change scenarios in Siberia for the future (IPCC, 2021) will likely enhance peatland drying and may convey a competitive advantage to some conifers, especially *Pinus sylvestris*. But dry peatland conditions will also exacerbate the likelihood of greater fire frequency, increase fire severity (i.e., stand replacing fires) and likely disrupt the successional pathway of other typical taiga conifers, notably that of fire avoiders. Such a fire regime change may accelerate a shift towards communities dominated by deciduous fire invader-type trees. Our results indicate an increased sensitivity of mesotrophic peatlands to burning at water levels below 20 cm, however, more regional studies are necessary to determine thresholds in peatland moisture to fire. Future climate-disturbance-


500

fire feedbacks will not only accelerate changes in boreal forest structure and composition, but also impact the carbon balance of the forested peatlands and ultimately lead to albedo-mediated feedbacks on the regional climate system.

Appendices

505 Appendix A1.

Correlation coefficients for the two sequences. Note that forest density refers to dark-to-light taiga ratio.



510

Data availability

515 All data generated for this paper will be deposited in the Neotoma database following publication.

Author contribution

Conceptualization: AF; Methodology: AF, ACD, MG, SMH, GB, NG, ST, IT, SK; Investigation and writing: AF prepared the manuscript with contribution from MP, and SMH; Editing: ACD, MP, MG, GB, NG, SMH, ST, AN, IO, SK; Funding acquisition: AF.

Competing interests: The authors declare that they have no conflict of interest.

Acknowledgements

We thank Rosa Rytönen, University of Manchester for her help during fieldwork, and the reviewers for useful comments to improve the manuscript. Sergey Loiko also provided archaeological literature from the regions.

Funding

This research was supported by the German Research Foundation (grant no. FE-1096/6-1). SK is grateful for support from the Russian Science Foundation (No. 20-67-46018) and the Tomsk State University Development Programme («Priority-2030»).

References

- Amesbury, M.J., Swindles, G.T., Bobrov, A., Charman, D.J., Holden, J., Lamentowicz, M., Mallon, G., Mazei, Y., Mitchell, E.A.D., Payne, R.J., Roland, T.P., Turner, T.E., and Warner, B.G.: Development of a new pan-European testate amoeba transfer function for reconstructing peatland palaeohydrology. *Quat. Sci. Rev.* 152, 132–151, <https://doi.org/10.1016/j.quascirev.2016.09.024>, 2018.
- Adolf, C., Wunderle, S., Colombaroli, D., Weber, H., Gobet, E., Heiri, O., van Leeuwen, J.F., Bigler, C., Connor, S.E., Gafka, M. and La Mantia, T., Tinner, W.: The sedimentary and remote sensing reflection of biomass burning in Europe. *Global Ecology and Biogeography*, 27, 199–212. <https://doi.org/10.1111/geb.12682>, 2018.
- Agee, J.K.: Fire and Pine Ecosystems. In: Richardson, D.M. (Ed.): *Ecology and Biogeography of Pinus*. Cambridge Univ Press, Cambridge, pp. 193–218, 1998.
- Amesbury, M.J., Swindles, G.T., Bobrov, A., Charman, D.J., Holden, J., Lamentowicz, M., Mallon, G., Mazei, Y., Mitchell, E.A.D., Payne, R.J., Roland, T.P., Turner, T.E., and Warner, B.G.: Development of a new pan-European testate amoeba transfer function for reconstructing peatland palaeohydrology. *Quat. Sci. Rev.* 152, 132–151, <https://doi.org/10.1016/j.quascirev.2016.09.024>, 2018.
- Andela, N., Morton, D. C., Giglio, L., Chen, Y., van der Werf, G. R., Kasibhatla, P. S., DeFries, R. S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Yue, C., and Randerson, J.T.: A human-driven decline in 10 global burned area, *Science*, 356, 1356–1362, <https://doi.org/10.1126/science.aal4108>, <http://science.sciencemag.org/content/356/6345/1356>, 2017.
- Bennett, K.D., Willis, K.J.: Pollen. In: Smol, J.S., Birks, H.J.B., Last, W.M. (Eds.): *Tracking Environmental Change Using Lake Sediments*. Kluwer Academic Publishers, Dordrecht, pp. 5–32, 2001.

- Blaauw, M.: Methods and code for ‘classical’ age-modelling of radiocarbon sequences. *Quaternary Geochronology* 5, 512–518, <https://doi.org/10.1016/j.quageo.2010.01.002>, 2010.
- Blarquez, O., Vanni re, B., Marlon, J.R., Daniau, A.-L., Power, M.J., Brewer, S., and Bartlein, P.J.: Paleofire An R package to analyse sedimentary charcoal records from the Global Charcoal Database to reconstruct past biomass burning. *Computers & Geosciences*, 72, 255-261, <https://doi.org/10.1016/j.cageo.2014.07.020>, 2014.
- Barhoumi, C., Peyron, O., Joannin, S., Subetto, D., Kryshen, A., Drobyshhev, I., Girardin, M.P., Brossier, B., Paradis, L., Pastor, T., and Alleaume, S.: Gradually increasing forest fire activity during the Holocene in the northern Ural region (Komi Republic, Russia). *The Holocene*, 29906-20, <https://doi.org/10.1177/0959683619865593>, 2019.
- Barhoumi, C., Ali, A.A., Peyron, O., Dugerdil, L., Borisova, O., Golubeva, Y., Subetto, D., Kryshen, A., Drobyshhev, I., Ryzhkova, N., and Joannin, S.: Did long-term fire control the coniferous boreal forest composition of the northern Ural region (Komi Republic, Russia) *J. Biogeogr.*, 47, 2426–2441. <https://doi.org/10.1111/jbi.13922>, 2020.
- Barhoumi, C., Vogel, M., Dugerdil, L., Limani, H., Joannin, S., Peyron, O., Ali, A.A.: Holocene Fire Regime Changes in the Southern Lake Baikal Region Influenced by Climate-Vegetation-Anthropogenic Activity Interactions. *Forests*, 12, 978, <https://doi.org/10.3390/f12080978>, 2021.
- Blyakharchuk, T. A.: Four new pollen sections tracing the Holocene vegetational development of the southern part of the West Siberian Lowland. *The Holocene* 13, 715–731. <https://doi.org/10.1191/0959683603hl658rp>, 2003.
- Blyakharchuk, T.A., Kurina I.V., and Pologova N.N.: Late-Holocene dynamics of vegetation cover and humidity of climate in the southeastern sector of the West Siberian Plain according to the data of palynological and rhizopod research of peat deposits. *Tomsk State University Journal of Biology*, 45, 164-189. doi: 10.17223/19988591/44/9, In Russian, English Summary, 2019.
- Blyakharchuk, T.A., and Kurina, I.: Late Holocene environmental and climatic changes in the Western Sayan Mountains based on high-resolution multi-proxy data. *Boreas*, 50, 919– 934. <https://doi.org/10.1111/bor.12493>, 2021.
- Borisova, O.K., Novenko, E.Y., Zelikson, E.M., and Kremenetski, K.V.: Lateglacial and Holocene vegetational and climatic changes in the southern taiga zone of West Siberia according to pollen records from Zhukovskoye peat mire. *Quaternary International*, 237, 65-73. <https://doi.org/10.1016/j.quaint.2011.01.015>, 2011.
- Bleuten, W., and Filippov, I.: Hydrology of mire ecosystems in central West Siberia: the Mukhrino field station. in: Glagolev, M.V., Lapshina, E.D. (Eds.), *Transactions of UNESCO Department of Yugorsky State University Dynamics of Environment and Global Climate Change*, <https://doi.org/10.17816/edgcc11S208-224>, 2008.
- Bova, S., Rosenthal, Y., Liu, Z., Godat, S.P., and Yan, M.: Seasonal origin of the thermal maxima at the Holocene and the last interglacial. *Nature* 589, 548–553, <https://doi.org/10.1038/s41586-020-03155-x>, 2021.
- Camill, P., Barry, A., Williams, E., Andreassi, C., Limmer, J., and Solick, D.: Climate vegetation–fire interactions and their impact on long-term carbon dynamics in a boreal peatland landscape in northern Manitoba, Canada. *Journal of Geophysical Research*, 114, G04017, <https://doi.org/10.1029/2009JG001071>, 2009.

- 585 Chambers, F.M. and Charman, D.J.: Holocene environmental change: contributions from the peatland archive. *The Holocene* 14, 1-6. doi:10.1191/0959683604hl684ed, 2014.
- Charman, D.J., Hendon, D., and Woodland, W.A.: *The Identification of Testate Amoebae (Protozoa: Rhizopoda) in Peats. Technical Guide No. 9.* Quaternary Research Association, London, 2000.
- Coop, J.D., Parks, S.A., Stevens-Rumann, C.S., Crausbay, S.D., Higuera, P.E., Hurteau, M.D., Tepley, A., Whitman, E., Assal, T., Collins, B.M., and Davis, K.T.: Wildfire-driven forest conversion in western North American landscapes. *BioScience*, 590 70, 659-73, <https://doi.org/10.1093/biosci/biaa061>, 2020.
- Courtney-Mustaphi, C.J., and Pisaric, M.F.: A classification for macroscopic charcoal morphologies found in Holocene lacustrine sediments. *Progress in Physical Geography*, 38, 734-754, <https://doi.org/10.1177/0309133314548886>, 2014.
- Dieleman, C.M., Rogers, B.M., Potter, S., Veraverbeke, S., Johnstone, J.F., Laflamme, J., Solvik, K., Walker, X.J., Mack, M.C., and Turetsky, M.R.: Wildfire combustion and carbon stocks in the southern Canadian boreal forest: implications 595 for a warming world. *Global Change Biol.* 26, 6062–6079, <https://doi.org/10.1111/gcb.15158>, 2020.
- Enache, M.D., and Cumming, B.F.: Tracking recorded fires using charcoal morphology from the sedimentary sequence of Prosser Lake, British Columbia (Canada). *Quat. Res.*, 65, 282-292, <https://doi.org/10.1016/j.yqres.2005.09.003>, 2006
- Evseeva, N.S.: Geography of the Tomsk Region (Natural conditions and resources). Tomsk. West Siberian Plain Great Soviet Encyclopedia: in A. M. Prokhorov (Eds): - 3rd ed. - M.: Soviet Encyclopedia. 223, 2001. 600
- Feurdean, A., Veski, S., Florescu, G., Vannière, B., Pfeiffer, M., O'Hara, R.B., Stivrins, N., Amon, L., Heinsalu, A., Vassiljev, J., and Hickler, T.: Broadleaf deciduous forest counterbalanced the direct effect of climate on Holocene fire regime in hemiboreal/boreal region (NE Europe). *Quaternary Science Reviews*, 169, 378-390, <https://doi.org/10.1016/j.quascirev.2017.05.024>, 2017.
- 605 Feurdean, A., Gałka, M., Tantau, Florescu, G., I., Hutchinson, S.M., Diaconu, A., and Kirpotin, S.: 2000 years of variability in hydroclimate and carbon accumulation in western Siberia and the relationship with large scale atmospheric circulation: A multiproxy peat record. *Quaternary Science Review*, 226, 105948, <https://doi.org/10.1016/j.quascirev.2019.105948>, 2019.
- Feurdean, A., Florescu, G., Tanțău, I., Vannière, B., Diaconu, A.C., Pfeiffer, M., Warren, D., Hutchinson, S.M., Gorina, N., Gałka, M., and Kirpotin, S.: 2020a. Recent fire regime in the southern boreal forests of western Siberia is unprecedented 610 in the last five millennia. *Quaternary Science Reviews*, 244, 106495, <https://doi.org/10.1016/j.quascirev.2020.106495>, 2020.
- Feurdean, A., Vannière, B., Finsinger, W., Warren, D., Connor, S. C., Forrest, M., Liakka, J., Panait, A., Werner, C., Andrič, M., Bobek, P., Carter, V. A., Davis, B., Diaconu, A.-C., Dietze, E., Feeser, I., Florescu, G., Gałka, M., Giesecke, T., Jahns, S., Jamrichová, E., Kajukalo, K., Kaplan, J., Karpińska-Kołaczek, M., Kołaczek, P., Kuneš, P., Kupriyanov, D., Lamentowicz, M., Lemmen, C., Magyari, E. K., Marcisz, K., Marinova, E., Niamir, A., Novenko, E., Obremska, M., Pędziszewska, A., Pfeiffer, M., Poska, A., Rösch, M., Słowiński, M., Stančikaitė, M., Szal, M., Święta-Musznicka, J., Tanțău, I., Theuerkauf, M., Tonkov, S., Valkó, O., Vassiljev, J., Veski, S., Vincze, I., Wacnik, A.,

- 620 Wiethold, J., and Hickler, T., 2020. Fire risk modulation by long-term dynamics in land cover and dominant forest type in Eastern and Central Europe. *Biogeosciences*, 17, 1–18. <https://doi.org/10.5194/bg-17-1-2020b>.
- Feurdean, A.: Experimental production of charcoal morphologies to discriminate fuel source and fire type: an example from Siberian taiga, *Biogeosciences*, 18, 3805–3821, <https://doi.org/10.5194/bg-18-3805-2021>, 2021.
- 625 Galka, M., Tanțău, I., Ersek, V., and Feurdean, A.: A 9000 year record of cyclic vegetation changes identified in a montane peatland deposit located in the Eastern Carpathians (Central-Eastern Europe): Autogenic succession or regional climatic influences? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 449, 52–61, <https://doi.org/10.1016/j.palaeo.2016.02.007>, 2016.
- Gewin, B.: How peat could protect the planet. *Nature*, 578, 7794, 204–208, DOI: 10.1038/d41586-020-00355-3, 2020.
- Gill, A.M.: Fire adaptive traits of vascular plants. Fire regimes and ecosystem properties: proceedings of the conference, vol WO-26. USDA Forest Service, Honolulu 208–230.
- 630 Grospietsch, T., 1958. Wechseltierchen (Rhizopoden). Kosmos Verlag, Stuttgart.
- Grooth, W.J., Cantin, A.S., Flannigan, M.D., Soja, A.J., Gowman, L.M., and Newbery, A.: A comparison of Canadian and Russian boreal forest fire regimes. *Forest Ecology Management*, 294, 23–34, <https://doi.org/10.1016/j.foreco.2012.07.033>, 2013.
- 635 Groisman, P.Y., Blyakharchuk, T.A., Chernokulsky, A.V., Arzhanov, M.M., Marchesini, L.B., Bogdanova, E.G., Borzenkova, II, Bulygina, O.N., Karpenko, A.A., Karpenko, L.V., and Knight, R.W.: Climate changes in Siberia, in: Regional environmental changes in Siberia and their global consequences Springer, Dordrecht, 2012.
- Glückler, R., Herzschuh, U., Kruse, S., Andreev, A., Vyse, S. A., Winkler, B., Biskaborn, B. K., Pestrykova, L., and Dietze, E.: Wildfire history of the boreal forest of south-western Yakutia (Siberia) over the last two millennia documented by a lake-sediment charcoal record, *Biogeosciences*, 18, 4185–4209, <https://doi.org/10.5194/bg-18-4185-2021>, 2021.
- 640 Goldammer J.G., and Furyaev V.V.: Fire in Ecosystems of Boreal Eurasia: Ecological Impacts and Links to the Global System. In: Goldammer J.G., Furyaev V.V. (Eds.): Fire in Ecosystems of Boreal Eurasia. Forestry Sciences, vol 48. Springer, Dordrecht. https://doi.org/10.1007/978-94-015-8737-2_1, 1996.
- Hendon, D., and Charman, D.J.: The preparation of testate amoebae (Protozoa: rhizopoda) samples from peat. *Holocene* 7, 199–205. <https://doi.org/10.1177/095968369700700207>, 1997.
- 645 Higuera, P.E., Sprugel, D.G., and Brubaker, L.B.: Reconstructing fire regimes with charcoal from small-hollow sediments: a calibration with tree-ring records of fire. *The Holocene* 15, 238–51, 10.1191/0959683605hl789rp, 2005.
- Higuera, P., Brubaker, L., Anderson, P., Hu, F., and Brown, T.: Vegetation mediated the impacts of postglacial climate change on fire regimes in the south-central Brooks Range, Alaska. *Ecological Monographs* 79, 201–219, <https://doi.org/10.1890/07-2019.12009>, 2009.
- 650 Higuera, P.E., Briles, C.E., and Whitlock, C.: Fire-regime complacency and sensitivity to centennial-through millennial-scale climate change in Rocky Mountain subalpine forests, Colorado, USA. *Journal of Ecology*, 102, 1429–1441, <https://doi.org/10.1111/1365-2745.12296>, 2014.

- Holden, J., Palmer, S. M., Johnston, K., Wearing, C., Irvine, B., and Brown, L.E.: Impact of prescribed burning on blanket peat hydrology, *Water Resour. Res.*, 51, 6472– 6484, doi:10.1002/2014WR016782, 2015.
- 655
- Hudspith, V. A., Hadden, R. M., Bartlett, A. I., and Belcher, C. M.: Does fuel type influence the amount of charcoal produced in wildfires? Implications for the fossil record, *Palaeontology*, 61, 159–171. <https://doi.org/10.1111/pala.12341>, 2018.
- Hutchinson, S.M., Akinyemi, F.O., Mîndrescu, M., Begy, R. and Feurdean, A.: Recent sediment accumulation rates in contrasting lakes in the Carpathians (Romania): impacts of shifts in socio-economic regime. *Regional environmental change*, 16, 501-513. <https://doi.org/10.1007/s10113-015-0764-7>, 2016.
- 660
- Idimeshev, A.A., Bychkov, D.A., and Asochakova, E.M.: Stone industry of Samuska III settlement based on the results of the statistic analysis. *Tomsk Journal of Linguistics and Anthropology*, 3, 115-127, 10.23951/2307-6119-2020-3-115-127, 2020.
- IPCC, 2021: Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press.
- 665
- Ivanova, G.A.: Vegetation zone-specific characteristics of scots pine forest fires in central Siberia *Dr Biol. Krasnoyarsk: VN Sukachev Institute of Forest Publishing*, 2005.
- 670
- Ivanova, G.A., Kukavskaya, E.A., Ivanov, V.A., Conrad, S.G., MvRae, D.: Fuel characteristics, loads and consumption in Scots pine forests of central Siberia. *J. For. Res.* 31, 2507–2524, <https://doi.org/10.1007/s11676-019-01038-0>, 2020.
- Jensen, K., Lynch, E., Calcote, R., and Hotchkiss, S.C.: Interpretation of charcoal morphotypes in sediments from Ferry Lake, Wisconsin, USA: do different plant fuel sources produce distinctive charcoal morphotypes? *Holocene* 17, 907-915, <https://doi.org/10.1177/0959683607082405>, 2007.
- 675
- Kaufman, D., McKay, N., Routson, C. et al. Holocene global mean surface temperature, a multi-method reconstruction approach. *Sci Data* 7, 201, <https://doi.org/10.1038/s41597-020-0530-7>, 2020.
- Kasischke, E.S.: Boreal ecosystems in the global carbon cycle. In: Kasischke ES, Stocks BJ (eds) *Fire, climate change, and carbon cycling in the boreal forest*. Springer, 2010.
- 680
- Kettridge, N., Turetsky, M., Sherwood, J.H., Thompson, D.K., Miller, C.A., Benscoter, B.W., Flannigan, M.D., Wotton, B.M., and Waddington, J.M.: Moderate drop in water table increases peatland vulnerability to post-fire regime shift. *Scientific Reports*, 5, 8063. <https://doi.org/10.1038/srep08063>, 2015.
- Kharuk, V.I., Ponomarev, E.I., Ivanova, G.A., Dvinskaya, M.L., Coogan, S.C., and Flannigan, M.D.: Wildfires in the Siberian taiga. *Ambio* 1-22, <https://doi.org/10.1007/s13280-020-01490-x>, 2021.
- 685
- Kelly, R.F., Higuera, P.E., Barrett, C.M., and Hu, F.S.: A signal-to-noise index to quantify the potential for peak detection in sediment-charcoal records. *Quaternary Research* 75, 11-17, <https://doi.org/10.1016/j.yqres.2010.07.011>, 2011.

- Kelly, R., Chipman, M.L., Higuera, P.E., Stefanova, I., Brubaker, L.B., and Hu, F.S.: Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National Academy of Sciences*, 110, 13055-13060, <https://doi.org/10.1073/pnas.1305069110>, 2013.
- 690 Kirpotin, S.N., Antoshkina, O.A., Berezin, A.E., Elshehawi, S., Feurdean, A., Lapshina, E.D., Pokrovsky, O.S., Peregon, A.M., Semenova, N.M., Tanneberger, F., and Volkov, I.V.: Great Vasyugan Mire: How the world's largest peatland helps addressing the world's largest problems. *Ambio*, 7:1-2. <https://doi.org/10.1007/s13280-021-01520-2>, 2021.
- Korovin, G.N.: Analysis of distribution of forest fires in Russia. In: Goldammer J.G., and V.V. Furyaev, V.V.(Eds.): *Fire in Ecosystems of Boreal Eurasia*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 2016.
- 695 Kuosmanen, N., Fang, K., Bradshaw, R.H., Clear, J.L., and Seppä, H.: Role of forest fires in Holocene stand-scale dynamics in the unmanaged taiga forest of northwestern Russia. *The Holocene* 24, 1503–1514, <https://doi.org/10.1177/0959683614544065>, 2014.
- Kukavskaya, E.A., Ivanova, G.A., Conard, S.G., McRae, D.J., and Ivanov, V.A.: Biomass dynamics of central Siberian Scots pine forests following surface fires of varying severity. *Int J Wildland Fire* 23, 872–886, <https://doi.org/10.1071/WF13043>, 2014.
- 700 Kukavskaya, E.A., Buryak, L.V., Shvetsov, E.G., Conard, S.G., and Kalenskaya, O.P.: The impact of increasing fire frequency on forest transformations in southern Siberia. *Forest ecology and management*, 15, 382:225-35, [10.1016/j.foreco.2016.10.015](https://doi.org/10.1016/j.foreco.2016.10.015), 2016.
- Kurina, I.V., Veretennikova, E. E., Golovatskaya, E.A., Blyakharchuk, T.A., and Smirnov, S.V.: Dynamics of the surface wetness of mires in the southern taiga subzone of Western Siberia in the middle and late Holocene. *Tomsk State University Journal of Biology*, 42, 218-241, doi: 10.17223/19988591/42/12, 2018.
- 705 Kurina, I.V., and Li, H. Why Do Testate Amoeba Optima Related to Water Table Depth Vary? *Microb Ecol*, 77, 37–55, <https://doi.org/10.1007/s00248-018-1202-4>, 2019.
- Laschinsky, N.N., and Koroliuk, Y, A.: Syntaxonomy of zonal dark-coniferous forests of southern taiga of the West Siberian plain and of humid low-mountains of the Altai-Sayan mountain region. *Vegetation of Russia*, 26, [10.31111/vegus/2015.26.85](https://doi.org/10.31111/vegus/2015.26.85), 2015.
- 710 Liss, O.L., L.I. Abramova, N.A. Avetov, N.A. Berezina, L.I. Inisheva, T.V., Kurnishkova, Z.A., Sluka, T.Yu, Tolpycheva, et al. Mire systems of Western Siberia and their environmental importance. Tula: Grifi K (in Russian).
- Lamentowicz, M., Słowinski, M., Marcisz, K., Zielinska, M., Kaliszan, K., Lapshina, E., Gilbert, D., Buttler, A., Fiałkiewicz-Kozieł, B., Jassey, V.E., and Laggoun-Defarge, F.: Hydrological dynamics and fire history of the last 1300 years in western Siberia reconstructed from a high-resolution, ombrotrophic peat archive. *Quat. Res.* 84, 312-325, 2015, <https://doi.org/10.1016/j.yqres.2015.09.002>, 2015.
- 715 Mazei, Y., Tsyganov, A.: *Freshwater Testate Amoebae*. KMK, Moscow, 2006.
- Magnan, G.M., Lavoie, M., and Payette, S.: Impact of fire on long-term vegetation dynamics of ombrotrophic peatlands in northwestern Quebec, Canada. *Quaternary Research*, 77, 110–121, 11, doi:10.1016/j.yqres.2011.10.006, 2012.
- 720



- Mekonnen, Z.A., Riley W.J., Randerson, J.T., Grant R.F, and. Rogers, B.M.: 2019. Expansion of high-latitude deciduous forests driven by interactions between climate warming and fire. *Nature Plants*. 5, 952–958. <https://doi.org/10.1038/s41477-019-04958>, 2019.
- 725 Mikhailova, A.B., Grenaderova, A.V., Kurina, I.V., Shumilovskikh, L. and Stojko, T.G.: Holocene vegetation and hydroclimate changes in the Kansk forest steppe, Yenisei River Basin, East Siberia. *Boreas*, 50, 948-966. <https://doi.org/10.1111/bor.12542>, 2021.
- Moore, P.D., and Web, J.A., Collinson, M.E.: Pollen analysis. Blackwell Science, 1991.
- Morales-Molino, C., Tinner. W., Perea. R., Carrión. J.S., Colombaroli. D., Valbuena-Carabaña. M., Zafra. E., and Gil. L.: Unprecedented herbivory threatens rear-edge populations of *Betula* in southwestern Eurasia. *Ecology*, 100, 2833, doi: 10.1002/ecy.2833, 2019.
- 730 Moritz, M.A., Batllori, E., Bradstock, R.A., Gill, A.M., Handmer, J., Hessburg, P.F., Leonard, J., McCaffrey, S., Odion, D.C., Schoennagel, T., and Syphard, A.D.: Learning to coexist with wildfire. *Nature*, 515, 58–66, <https://doi.org/10.1038/nature13946>, 2014.
- Marlon, J. R., Kelly, R., Daniau, A.-L., Vanni re, B., Power, M. J., Bartlein, P., Higuera, P., Blarquez, O., Brewer, S., Br cher, T., Feurdean, A., Romera, G. G., Iglesias, V., Maezumi, S. Y., Magi, B., Courtney Mustaphi, C. J., and Zhihai, T.: Reconstructions of biomass burning from sediment-charcoal records to improve data–model comparisons, *Biogeosciences*, 13, 3225–3244, <https://doi.org/10.5194/bg-13-3225-2016>, 2016.
- 735 Naumov, I. V.: The history of Siberia. London, New York: Routledge, 2006.
- Page, S., Hoscilo, A., Langner, A., Tansey, K., Siegert, F., Limin S., Rieley, J.: Tropical peatland fires in Southeast Asia. In: *Tropical Fire Ecology*. Springer Praxis Books. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-540-77381-8_9, 2009.
- 740 Ogden, C.G., and Hedley, R.H.: An Atlas of Freshwater Testate Amoebae. Oxford University Press, London.
- Pausas J.G., and Paula, S.: Fuel shapes the fire–climate relationship: evidence from Mediterranean ecosystems *Global Ecology and Biogeography*, 21, 1074–82, <https://doi.org/10.1111/j.1466-8238.2012.00769.x>, 2012.
- 745 Power, M.J., Marlon, J.R., Bartlein, P.J., Harrison, S.P.: Fire history and the Global Charcoal Database: a new tool for hypothesis testing and data exploration. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 291, 52-59, <https://doi.org/10.1016/j.palaeo.2009.09.014>, 2010.
- Remy, C.C., Fouquemberg, C., Asselin, H., Andrieux, B., Magnan, G., Brossier, B., Grondin, P., Bergeron, Y., Talon, B., Girardin, M.P., Blarquez, O.: Guidelines for the use and interpretation of palaeofire reconstructions based on various archives and proxies. *Quat. Sci. Rev.* 193, 312-322. <https://doi.org/10.1016/j.quascirev.2018.06.010>, 2018.
- 750 Reimer, P., Austin, W., Bard, E., Bayliss, A., Blackwell, P., Bronk Ramsey, C., Talamo, S.: The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon*, 62, 725-757. doi:10.1017/RDC.2020.41, 2020.
- Rogers, B.M., Soja, A.J., Goulden, M.L., and Randerson, J.T.: Influence of tree species on continental differences in boreal fires and climate feedbacks. *Nature Geosciences*, 8, 228-234. <https://doi.org/10.1038/ngeo2352>, 2015.

- 755 Rowe, J.S.: Concepts of fire effects on plant individuals and species. In: Wein RW, McLean DA (eds) *The role of fire in northern circumpolar ecosystems*, 18th edn. Wiley, Chichester, 1983.
- Rudaya, N., Krivonogov, S., Słowiński, M., Cao, S., and Zhilich, S: 2020. Postglacial history of the Steppe Altai: Climate, fire and plant diversity. *Quaternary Science Reviews*, 249, p.106616. <https://doi.org/10.1016/j.quascirev.2020.106616>, 2020.
- 760 Sannikov, S.N., and Goldammer, J.G.: Fire ecology of pine forests of northern Eurasia. In: Goldammer JG, Furyaev VV (eds) *Fire in ecosystems of boreal Eurasia*, Forestry Sciences vol 48. Kluwer, Dordrecht, pp 151–167. . https://doi.org/10.1007/978-94-015-8737-2_1, 1996.
- Scholten, R.C., Jandt, R., Miller, E.A. Rogers, B.M., Veraverbeke, S. Overwintering fires in boreal forests. *Nature* 593, 399–404, <https://doi.org/10.1038/s41586-021-03437>, 2021.
- 765 Stivrins, N., Aakala, T., Ilvonen, L., Pasanen, L., Kuuluvainen, T., Vasander, H., ... Seppä, H., 2019. Integrating fire-scar, charcoal and fungal spore data to study fire events in the boreal forest of northern Europe. *The Holocene*, 29, 1480–1490. <https://doi.org/10.1177/0959683619854524>.
- Scheffer, M., Hirota, M., Holmgren, M., Van, Nes E.H., and Chapin, III F.S.: Thresholds for Boreal Biome Transitions. *PNAS*, 109, 21384–21389, <https://doi.org/10.1073/pnas.1219844110>, 2012
- 770 Tautenhahn, S., Lichstein, J.W., Jung, M., Kattge, J., Bohlman, S.A., Heilmeyer, H., Prokushkin, A., Kahl, A. and Wirth, C., 2016. Dispersal limitation drives successional pathways in Central Siberian forests under current and intensified fire regimes. *Global Change Biology*, 22 2178–2197, <https://doi.org/10.1111/gcb.13181>, 2016.
- Turunen, J., Tahvanainen, T., Tolonen, K., and Pitkänen, A.: Carbon accumulation in West Siberian Mires, Russia Sphagnum peatland distribution in North America and Eurasia during the past 21,000 years, *Global Biogeochem. Cycles*, 15(2), 285– 296, doi:10.1029/2000GB001312, 2001.
- 775 van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015). *Geosci. Model Dev.*, 10, 3329–3357, [https://doi.org/10.5194/gmd-10-3329-](https://doi.org/10.5194/gmd-10-3329-2017)
- 780 2017, 2017.
- Vachula, R.S., Sae-Lim, J., and Li, R.: A critical appraisal of charcoal morphometry as a paleofire fuel type proxy. *Quaternary Science Reviews*, 262, 106979. <https://doi.org/10.1016/j.quascirev.2021.106979>, 2021.
- Vompersky, S.E., Ivanova, A.I., Tsyganova, O.P., Valiaeva, N.A., Glukhova, T.V., Dubinin, F.I., L.G. Markelova.: Wet soilsand mires in Russia and their carbon pool. *Pochvovedenie*, 12, 17–25 (in Russian), 1994.
- 785 Walker, X.J., Baltzer, J.L., Cumming, S.G., Day, N.J., Ebert , C., Goetz, S., Johnstone, J.F., Potter, S., Rogers, B.M., Schuur, E.A. G., Turetsky, M.R., and Mack, M. C.: Increasing wildfires threaten historic carbon sink of boreal forest soils. *Nature*, 572, 520–523, <https://doi.org/10.1038/s41586-019-1474-y>, 2019.

- Wirth, C.: Fire regime and tree diversity in boreal forests: implications for the carbon cycle. In: Forest Diversity and Function. pp 309-344. Springer, Berlin, Heidelberg, 2005.
- 790 Whitlock, C., and Larsen, C.: Charcoal as a fire proxy. In Smol, JP, Birks, HJB and Last, WM. (Eds.): Tracking environmental change using lake sediments. Volume 3: terrestrial, algal, and siliceous indicators. Kluwer Academic Publishers, 75-97, 10.1007/0-306-47668-1, https://doi.org/10.1007/0-306-47668-1_5, 2002.
- Whitlock, C., Colombaroli, D., Conedera, M., Tinner, W.: Land-use history as a guide for forest conservation and management. *Conservation Biology*, 32, 84-97, <https://doi.org/10.1111/cobi.12960>, 2018.
- 795 Whitman, E., Parisien, M.A., Thompson, D.K., and Flannigan, M.D.: Short-interval wildfire and drought overwhelm boreal forest resilience. *Scientific Reports*, 9, <https://doi.org/10.1038/s41598-019-55036-7>, 2019.
- Whitman, E., Parisien, M.A., Thompson, D.K., and Flannigan, M.D.: Topoedaphic and forest controls on post-fire vegetation assemblies are modified by fire history and burn severity in the northwestern Canadian boreal forest. *Forests*, 9, 151. <https://doi.org/10.3390/f9030151>, 2018.
- 800 Zolnikov, I.D., Nikulina, A.V., Pavlenok, K.K., Vybornov, A.V., Postnov, A.V., Bychkov, D.A., Glushkova, and N.V.: Regularities in the spatial location of archaeological objects in Timks. *Rossiiskaia arkheologiya*, 17959, 22, 31, 10.31857/S086960630008251-5, 2020.

805 **Figures**

810 **Figure 1.** Location of the study area in Eurasia, Russia and the Tomsk region (A). Satellite based images showing the location and spatial extent of vegetation types and the fire event at the previously published sites (B; Feurdean et al., 2020a) and the new study sites: Rybnaya (C) and Ulukh-Chayakh (D).



Legend

- ▲ Location of the study sites
- ~ River
- Gravel road

Vegetation

- mixed (deciduous and coniferous) forest
- deciduous forest
- boggy forest

- floodplain forest and meadow
- mire
- burnt area

815 **Figure 2.** The ratio of charcoal morphotype influx ($\#/\text{cm}^{-2}\text{yr}^{-1}$) of the two main categories: woody (wood, deciduous leaves, resins) and non-woody type (forbs, grass) in the Rybnaya and Ulukh-Chayakh sequences. The aspect ratio ($L:W$) at Ulukh-Chayakh is used for fuel type identification. A high $L:W$ ratio is typical for a higher abundance of graminoids charcoal, whereas a lower $L:W$ ratio is typical for charcoal from wood and leaves. The ratio of charcoal influx ($\#/\text{cm}^{-2}\text{yr}^{-1}$) of the two main size classes: small 150–500 μm and large $> 500\text{ }\mu\text{m}$ in the two sequences and the charcoal surface area (μm^2) at Ulukh-Chayakh (see File S3 for a full range of size classes). Bullets at Ulukh-Chayakh represent the extra-large charcoal fraction ($>1000\text{ }\mu\text{m}$) identified during routine plant macrofossils analysis of sediment volumes of ca. 20 cm^3 . The grey time window highlights the period with an uncertain chronology.






Figure 3. Temporal trends in the pollen and spore percentages of individual tree taxa and lumped group-wise for shrub, herb, fern and moss at Rybnaya and Ulukh-Chayakh. Periods with substantial changes in fire regime are indicated by the dotted vertical lines.


840



845 **Figure 4.** Temporal trends in the pollen percentages of the fire-related plant functional groups (invaders, resisters and avoiders)
 and the forest type determined from the ratio between pollen percentages of dark and light taiga tree taxa in the Rybnaya and
 Ulukh-Chayakh sequences. Hydrological conditions on the peatlands were derived from testate amoeba-based estimates of the
 water table depth (DTW) where values range from 8 cm (high water level) to 30 cm (low water level). The fire metrics include
 burned biomass (CHAR), fire frequency (number of fires /900 years) and charcoal peak magnitude (the higher the values, the
 850 greater the fire severity and/or closer to the site) derived from macro-charcoal particles >150 µm. Colours in the background
 denote periods with marked changes in fire regime and vegetation composition, where a change from blue to yellow indicates
 an intensification of the fire episodes.



Figures 5. Charcoal based fire frequency at here locations in *Pinus-Betula* dominated boreal forests Russia: northern Ural region (Bourhami et al., 2019), Lake Baikal (Bourhami et al., 2021) and western Siberian Plain (Rudaya et al., 2020) (a). Composite record of biomass burning (n=4) based on Z-score charcoal influx where positive/negative Z-score values represent greater-than-mean/lower-than-mean charcoal influx over the base period (b). Composite record of fire frequency (n=4) (c). Composite record of peatland hydrology (n=4) from testate amoeba where positive/negative Z-score values represent lower/higher-than mean water level (d). Annual temperature (anomalies) for 30-60 °N (Kaufman et al., 2020) (e). Composite record (n=4) of the relative abundance of avoiders, resisters, and invaders determined from pollen percentages (f). Grey curves represent confidence intervals. Colours highlight the periods with higher biomass burning in the composite record.



870

Figure 6. Correlation analysis between water table position, the main plant functional types, and the light-to dark taiga index (forests density) and micro and microcharcoal for Rybnaya (a) and Ulukh-Chayakh (b). Generalised linear models (GLMs) showing the response of biomass burning (CHAR) to changes in peatland moisture conditions (higher values in depth to water table=DTW representing drier conditions) for Rybnaya (c) and Ulukh-Chayakh (d).

