

## Reassessing Ireland's Hottest Temperature Record

Katherine Dooley<sup>1,2</sup>, Ciaran Kelly<sup>1</sup>, [Natascha Seifert<sup>1</sup>](#), Therese Myslinski<sup>1</sup>, Sophie O'Kelly<sup>1</sup>, Rushna Siraj<sup>1</sup>, [Natascha Seifert<sup>1</sup>](#), Ciara Crosby<sup>1</sup>, Jack Kevin Dunne<sup>1</sup>, Kate McCauley<sup>1</sup>, James Donoghue<sup>1</sup>, Eoin Gaddren<sup>1</sup>, Daniel Conway<sup>1</sup>, Jordan Cooney<sup>1</sup>, Niamh McCarthy<sup>1</sup>, Eoin Cullen<sup>1</sup>, Simon Noone<sup>1</sup>, Conor Murphy<sup>1</sup>, Peter Thorne<sup>1</sup>

5 <sup>1</sup> [The Irish Climate Analysis and Research UnitS \(ICARUS\), National University of Ireland Maynooth Maynooth University, Maynooth, Co. Kildare, Ireland.](#)

<sup>2</sup> [Environmental Protection Agency, PO Box 3000, Johnstown Castle Estate, County Wexford, Ireland, Y35 W821.](#)

Correspondence to: Katherine Dooley (katherine.dooley.2020@mumail.ie)

### Abstract

10 The highest currently recognised air temperature (33.3°C) ever recorded in the Republic of Ireland was logged at Kilkenny Castle in 1887. ~~This~~ ~~How~~ ~~the original observational record~~ ~~however~~ ~~s no longer exists,~~ ~~an issues which appears to be reflected globally.~~ Given that Ireland is now the only country in Europe to have a national heat record ~~that was~~ ~~that was~~ set in the 19th century, a reassessment of the verity of this record is both timely and valuable. The present analysis undertakes a fundamental reassessment of the plausibility of the 1887 temperature record using ~~methods~~ similar ~~methods~~ to those used to assess various  
15 weather extremes under WMO auspices over recent years. Specifically, we undertake an inter-station reassessment using sparse available records and make recourse to the new and improved 20CRv3 sparse-input reanalysis product. Neither surrounding available stations nor the reanalysis offer substantive support for the Kilkenny record of 33.3°C being correct. ~~Moreover,~~ ~~r~~Recent data rescue efforts have uncovered several earlier ~~extreme~~ values, one of which exceeds the Kilkenny value (~~XX~~33.5°C on 16<sup>th</sup> July 1876 ~~DATE~~ recorded at ~~PP~~the Phoenix Park). ~~However,~~ ~~the Given~~ sparsity of early observational  
20 ~~networks,~~ a distinct lack of synoptic support from 20CRv3 ~~for many of the extreme heat values,~~ ~~for a number of these values,~~ and the fact that these ~~ee~~ measurements ~~will have~~ ~~were used~~ ~~obtained using~~ non-standard exposures ~~leads us to conclude that,~~ ~~while these temperatures may well be correct~~ there is grossly insufficient evidence to support ~~any of~~ these 19<sup>th</sup> Century ~~extremes~~ as robust national heat record candidates. Data from the early 20<sup>th</sup> Century onwards benefits from a denser network of stations undertaking measurements in a more standardised manner, many under the direct auspices of Met Éireann and its  
25 predecessors, adhering to WMO guidance and protocols. This enables more robust cross-checking of records. ~~We argue that~~ ~~The~~ Met Éireann recognised 20<sup>th</sup> Century heat record from Boora in 1976 verifies as ~~the most plausia~~ ~~reasonable~~ ~~robust national temperature record~~ based upon the synoptic situation and comparisons with nearby neighbouring stations ~~and,~~ ~~has not been surpassed in the 21<sup>st</sup> Century.~~ This measurement of 32.5°C thus likely constitutes the highest ~~ever~~ reliably recorded temperature measurement in the Republic of Ireland. ~~This work was undertaken as a masters class assignment and u~~ Ultimately,  
30 the formal decision on any reassessment and reassignment of the national record rests with the national meteorological service,

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Superscript

Met Éireann. Finally, this analysis highlights the potential value in reanalysing other national meteorological records around the world using the techniques pioneered by WMO record assessment teams.

Formatted: English (United Kingdom)

## 1. Introduction and Context

National meteorological records of heat, cold, precipitation and other meteorological parameters play a key role in the communication of weather events and climate change at both national and international scales. Over the past decade or so the WMO has instigated teams of experts to assess or reassess global and regional records for a range of phenomena. These have either recognised (Quetlard et al., 2009, Courtney et al., 2012, Purevjav et al., 2015, Lang et al., 2017, Cervený et al., 2017, Lásková et al., 2018, Merlone et al., 2020, Peterson et al., 2020, Weidner et al., 2021) or occasionally removed and replaced (El Fadli et al., 2013) various global or regional records for meteorological phenomena including heat and cold records. The latter revocation by El Fadli et al. was of particular note as it removed a long-standing global all-time heat record which had appeared e.g. in the Guinness book of world records, encyclopedias and other similar reference materials. In comparison, national records have, at least in the vast majority of cases, received scant attention. Yet the tools and approaches employed by the WMO are eminently scaleable to a national context. Here, by use of Ireland's reported all-time heat record as an example, we make the case for assessing anew national records using the tools and techniques pioneered by these WMO expert teams. Such investigations are particularly valuable for early records that occurred prior to standardisation of instrumentation and methods of observation when the biases in the original records may be substantial (Böhm et al. 2010, Brunet et al. 2011, Murphy et al. 2020).

The highest recognised shaded air temperature recorded in Ireland was observed at Kilkenny Castle on 26th June 1887 (Rohan, 1986). Kilkenny Castle is located at 52.6505° N, 7.2493° W, 60 metres above sea level. In 1841, the population of Kilkenny City was approximately 19,071, decreasing to 15,257 in 1851 (McDill, 2021). According to the 2016 census, Kilkenny City now has a population of 25,512 (City Population, 2021). The reported temperature was observed from a "standard site," reading 33.3°C (Rohan, 1986:133). Unfortunately, the original records from Kilkenny Castle for the period are missing and unavailable for scrutiny (Mary Curley, pers comm). Curiosity surrounds this temperature record as Ireland remains the only country in Europe to have an existing heat record from the 19th century, with almost all other European countries having had their heat records set much more recently (Table 1). Given this, and the relative paucity of metadata supporting the Irish record, it is appropriate to re-examine this record using modern investigative techniques.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Country	Temperature (°C)	Date	Location
Republic of Ireland	33.3	26 <sup>th</sup> Jun 1887	Kilkenny Castle (Kilkenny)
England	38.7	25 <sup>th</sup> Jul 2019	Cambridge Botanic Garden
Wales	35.2	2 <sup>nd</sup> Aug 1990	Hawarden Bridge (Flintshire)
Scotland	32.9	9 <sup>th</sup> Aug 2003	Greycrook (Scottish Borders)
<u>Northern Ireland</u>	<u>31.34<sup>*</sup></u>	<u>21<sup>st</sup> July 2021</u>	<u>Armagh Castle (County Tyrone)</u>
<del>France</del> <del>Northern Ireland</del>	<del>46.030.8</del>	<del>28<sup>th</sup> Jun 2019</del> <del>30<sup>th</sup> Jun 1976</del>	<del>Veragues, Herault</del> <del>Knockarevan (County Fermanagh)</del>
<del>Belgium</del> <del>Northern Ireland</del>	<del>41.830.8</del>	<del>25<sup>th</sup> Jul 2019</del> <del>14<sup>th</sup> July 1983</del>	<del>Begijnendijk</del> <del>Flemish Brabant</del> <del>Shaw's Bridge, Belfast (County Antrim)</del>
<del>Netherlands</del> <del>France</del>	<del>40.746.0</del>	<del>25<sup>th</sup> July 2019</del> <del>28<sup>th</sup> Jun 2019</del>	<del>Gilze en Rijen</del> <del>Veragues, Herault</del>
<del>Norway</del> <del>Belgium</del>	<del>35.641.8</del>	<del>20<sup>th</sup> June 1976</del> <del>25<sup>th</sup> Jul 2019</del>	<del>Nesbyen, Buskerud</del> <del>Begijnendijk</del> <del>Flemish Brabant</del>
<del>Iceland</del> <del>Netherlands</del>	<del>30.540.7</del>	<del>22<sup>nd</sup> June 1939</del> <del>25<sup>th</sup> July 2019</del>	<del>Teigarhorn, Djupivogur</del> <del>Gilze-en-Rijen</del>
<del>Norway</del>	<del>35.6</del>	<del>20<sup>th</sup> June 1976</del>	<del>Nesbyen, Buskerud</del>
<del>Iceland</del>	<del>30.5</del>	<del>22<sup>nd</sup> June 1939</del>	<del>Teigarhorn, Djupivogur</del>

Table 1: ~~Table showing highest daily maximum temperature records of several NW European countries surrounding Ireland including constituent countries of the United Kingdom sourced from various National Meteorological Service websites at the time of writing (\* record yet to be ratified as of 21/07/21).~~

Formatted: Superscript

Commented [PWT1]: I'd check this is still the case and if so note this with the updated change – otherwise modify the table.

- 70 The national temperature record was logged at Kilkenny Castle using a Negretti and Zambia thermometer, a standard thermometer in use at the time. This thermometer was tested by the Meteorological Council in 1890 which showed the instrument required no corrections, thus reporting the true value ([Report of the Meteorological Council, 1890](#))([Kilkennyweather, 2019](#)). During the 1880s the Stevenson Screen (Stevenson, 1864) became the preferred standard screen to shield thermometers from sunlight and radiation. Dolores Gaffney, the collections officer at Kilkenny Castle, believes the temperature in question was recorded within a Stevenson Screen (pers. comm.). The original Kilkenny Castle

site was surveyed in 1890 but not considered a 'second order' ([maintained by trained staff \(UCAR, 2021\)](#)) station ([RMS, 1890](#)) and eventually closed [in 1933 \(Niall Dollard, pers. comm.\)](#). The exact location of the weather station in Kilkenny Castle at the time of the record is unknown. Station siting is known to have substantial potential impacts upon the representativity of resulting measurements and modern guidance from WMO specifies criteria for siting (WMO, 2018).

80 Kilkenny Castle is in the centre of Kilkenny [Town](#) beside the River Nore (Figure 1). Given ample grounds it is possible that the instrument was well sited but, equally, there are many potential sitings which would today be considered to yield the possibility of biased records. Met Éireann explained that a station inspection report from approximately 1911 stated that Kilkenny Castle only kept the records for four years before destroying them (Mary Curley, pers. comm.). This both means that the original record from which the national temperature record arises is lost for perpetuity and that necessary metadata to

85 understand the measurement context [are](#) also unavailable. Unfortunately, the timeseries of daily recordings around the record heat event [were](#) not saved and so the single 'observation' of record heat must be analysed in isolation.

[This added challenge of missing data is not an isolated occurrence. The crossover of manual to digitised recordings, early inconsistencies in logs before standardisation and the flippant destroying of records viewed as 'no longer useful' has been reported on at both European and global scales \(Brunet and Jones, 2011\). Data rescue efforts are being](#)

90 [recognised as imperative tools in the understanding of early climate variability and change \(Allan et al., 2011, Brönnimann et al 2019\), but are of limited utility for stations that had a policy of actively disposing of their records as is the case at Kilkenny Castle.](#)

**Commented [KD2]:** 90% sure that the official record stating that kilkenny was not considered of second order status is available at <https://doi.org/10.1002/qj.4970167403> but I do not have access to RMetS unfortunately. Peter/Conor/Simon, if you do would you check? It should be under Appendix VII. Sentence should read "Kilkenny, visited August 30. - The instruments were in good order, but there is no prospect of the station becoming one of Second Order."

Strong proof of closure is proving difficult. It appears that the station closed in 1900 but there is not much to directly state this. <https://www.kilkennyweather.com/index.php/1957-met-station-opened> states that the Met Eireann station opened in 1957 after a '57 year gap'.

**Commented [KD3R2]:** I now have a pers comm. with Niall Dollard (who runs Kilkennyweather.com). This is his reply:


*I believe daily observations were made up to 1900, however I am not sure if there were many between 1901 and 1910. Records exist for period 1911 to 1920. Also for 1921 to 1933 excluding the months of May 1923 and Oct to Dec 1933. The last readings, as far as I know, were completed in Sept 1933. It seems the Butlers (the owners) were winding things down at that stage.*

Next email reply:

*I believe now that there were records also in the period 1900 to 1910. So the only gaps I am sure about are the ones I mentioned in my previous email.*

*So it was an almost continuous record from 1886 to 1933.*

**Formatted:** Not Highlight



95 **Figure 1.** Aerial Image of Kilkenny Castle and its' surrounding area (top) and the grounds of the castle (bottom) ([Google Earth v. 9.152.0.1 \(2021\)](#)).

Met Éireann also recognises a 20<sup>th</sup> Century highest air temperature of 32.5°C recorded on 29<sup>th</sup> June 1976 [in Boora, County Offaly](#). This reading came in the midst of the long heatwave and summer drought which affected the British and Irish Isles



Oak Park, Carlow	14/07/1983				32.2	Met Éireann
Killarney, Kerry	12/07/1921				32.2	Mateus et al. (2020)
Markree, Sligo	29/06/1854	54°10'30" N	08°27'20" W	34m	32.2	Mateus et al. (2020)
Scattery Island, Clare	07/08/1851				32.2	Mateus et al. (2020)
Shannon Airport, Clare	28/06/2018				32.0	Met Éireann
Dooks, Kerry	18/07/2006				32.0	Met Éireann
Ballybrittas, Laois	29/06/2006				32.0	Met Éireann
Boora, Offlay	02/07/1976				32.0	Met Éireann

Table 2. Notable **apparent** heat extremes in available records from the Republic of Ireland over the instrumental record sourced from Met Éireann's website and available digital records, including the recently rescued early holdings from Mateus et al. (2020). The table includes all records that exceed 32°C at least one of which must plausibly constitute the robust national record. **Blank spaces** **Grey fill** **are** **is** **present** **where** **the** **exact** **location/height** **of** **the** **stations** **could** **not** **be** **found**.

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Times New Roman, 11 pt, Font color: Auto, Not Highlight

Formatted: Font color: Auto

Formatted: Font: Times New Roman, 11 pt, Font color: Auto, Not Highlight

Formatted: Font color: Auto, Not Highlight

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Highlight

Formatted: Highlight

**Commented [PWT5]:** Blank or colour block them or add N/A? I'd be tempted to grey fill so its visually obvious they are intentionally blank and modify this caption accordingly

110 Given the uncertainty associated with the Irish national high temperature record, together with its novelty in being the earliest national temperature record in Europe, the 2019-20 class of the MSc Climate Change at Maynooth University were set a classgroup assignment to re-evalaute the record. This paper represents the outcomes of that reassessment. Having introduced in the present section the national context and aspects of the recognised all-time and 20<sup>th</sup> Century national heat records, as well as selected additional heat extremes in available records, the rest of the paper is structured as follows. Section 2 introduces key methodological approaches and results from past WMO extremes verification efforts which this ~~exercise novel masters assignment as part of the MSc in Climate Change at Maynooth University~~ was designed to mimic. Section 3 goes on to apply these approaches to propose that the long-standing Kilkenny Castle national heat record should be rejected. Section 4 then considers additional candidates arising from Table 2, until alighting upon a recommendation for the most plausible highest reliable heat record in the available observations. Section 5 ~~provides reflections and~~ Section 6 concludes.

## 120 2. Assessments and Certifications of Climatic Extremes

The WMO has instigated a formal process to verify and certify a broad range of extremes. A rapporteur, when advised of a candidate record extreme that is desired to be assessed convenes a team of domain area and regional experts to investigate the event and recommend either acceptance or rejection as a bona fide observation. If the event is rejected the team is asked to



125 recommend an alternative record if they can do so. Teams have been instigated for temperature (El Fadli et al., 2013, Laska et al., 2018, Merlone et al., 2020, Weidner et al., 202), precipitation (Quetelard et al., 2009), winds (Courtney et al., 2012) and pressure (Purevjav et al., 2015) amongst other records. The WMO teams consider global or WMO region records and have not, to date, assessed national records.

130 Teams assess all aspects of the potential record including available metadata, data, the synoptic situation and, if possible, the instrumentation, to come to a conclusion. The extent of the investigation depends upon the nature and timing of the event being analysed. More recent events can benefit from more thorough analyses which may include instrument characterisation and site visits (Merlone et al., 2020). Whereas revisiting old records, as is the case herein, has to rely more on event characterisation and reference to nearby instrumental records (e.g. El Fadli et al., 2013). Increasingly, recourse is made to meteorological reanalysis products, including sparse-input centennial scale reanalyses (Slivinski et al., 2019) that can provide information on  
135 the evolving synoptic meteorological situation surrounding the record event.

Several analyses of various meteorological records worldwide have been undertaken over recent years leading to their removal (WMO, 2021). Perhaps most famously, what was cited to be the world's highest recorded temperature for over 90 years was invalidated following a careful WMO-team reappraisal. A temperature of 58°C was recorded at El Azizia (modern day Libya)  
140 but was disproved due to various issues brought to light after an in-depth investigation of the record. The issues surrounding the reading included problematic instrumentation, siting, observer interpretation and, notably in the context of the present study, that the record did not correspond to other nearby locations (El Fadli et al., 2013). The South African Weather Service (SAWS, 2019) reported a temperature of 50.1°C from the Vioolsdrif weather station on the 28th of November 2019 (Austral summer). This reading exceeded all previous heat records for South Africa. However, the temperature sensor in the Vioolsdrif  
145 weather station was replaced two days prior to the record temperature and upon tracking the behaviour of the station over the following days, it was decided that the behaviour of the temperature sensor was questionable. This resulted in the 50.1 °C record being deemed invalid. A temperature of 42.9 °C was recorded at the Deelen weather station, in the Netherlands, on the 25<sup>th</sup> of July 2019 (Schildkamp, 2019). The sensors at the Deelen station were inspected and any technical faults were ruled out. However, another instrument on the same site, a few hundred metres away, didn't record the same high values, which  
150 resulted in the record being dismissed. Lastly, a high temperature of 53°C recorded in Cloncurry, Queensland, Australia in 1889 was disregarded due to faults in the measurement and was therefore not included in Australia's Bureau of Meteorology's list of recognised extremes (Smith, 2012).

Of those assessments of possible records, the most analogous to the present analysis is that of El Fadli et al. (2013) in that the original instrumentation has long since disappeared and the metadata is imperfect. Like El Fadli et al. (2013), we therefore  
155 must make recourse to nearby stations and sparse-input centennial timescale reanalyses to reassess the Kilkenny record.



### 3. A Critical Reassessment of the Kilkenny Castle National Heat Record

160 Figure 2 displays the location of the Kilkenny station as well as the [six](#) closest stations with continuously available daily records over the period of the national heat record. There are very few stations with continuous daily records over this period, with only [5](#) other stations on the island of Ireland for which records had been digitised and made available [at the time of the present analysis](#). Data was sourced from holdings under preparation for the Copernicus Climate Change Service that arise from Met Éireann and the UK Met Office.

165 Additional records may exist in the Met Éireann archives that are yet to be digitised or those of the UK Met Office, and the analysis [largely](#) preceded the availability of the very recently rescued holdings from Mateus et al. (2020) ([Table 2](#)). The stations on the island of Ireland all consist of long-running sites that have been well maintained either by Met Éireann [and](#) its [predecessors](#) [or](#) the Armagh Observatory. To augment the records on the island of Ireland, recourse is also made to records from Sheffield in the UK. There

170 are [other](#) British [stations](#) for the same time but they are even further from the Kilkenny site (e.g. Oxford, Kew Gardens). With the notable exception of a station situated south-[east](#) of Kilkenny, the stations provide a reasonable geographical spread sufficient to infer spatial temperature gradient behaviour on the day of the record.




Figure 2. Location of the seven stations in Ireland and England used in this study to assess the reliability of the reported Kilkenny record. The green marker shows the location of the Kilkenny station. Station selection is based upon availability over the period of the Kilkenny record high temperature event.

Figure 3 illustrates observed surface temperatures for the month of June 1887 for the six comparator weather stations and the reported record temperature from Kilkenny. All station series show similar temporal evolution in their maximum daily temperatures through June of 1887 (although there is some ambiguity as to the timing of daily observations at several sites). A period of sustained heat builds during the second half of June peaking on or around the day of the record, before breaking the next day. The break in heat appears to progress from north to south. Geographically, the nearest recording to the Kilkenny record on the 26<sup>th</sup> was Birr, which reported 29.3°C, followed by Roches Point which reached 22.2°C. These high temperatures are backed up by contemporary records of the summer of 1887 being hot and dry. Noone et al. (2017) note that 1887 is one of the most intense drought years in Ireland in precipitation records spanning the past 250 years. Barrington (1888) assessed the impacts of the previous year's drought on agriculture, describing the drought of 1887 as being most extreme in the south and south east,

Formatted: Superscript

Formatted: Not Highlight

particularly in [Counties](#) Kilkenny, Wexford and Cork where April to June precipitation was as little as 30 percent of normal. Barrington also notes widespread crop failure across Ireland in 1887, while newspaper articles from the time indicate reduced harvests and crop failure throughout the country. [Industrial](#) activity, particularly the Linen industry in Northern Ireland was also adversely affected (Noone et al., 2017). [Although](#) the main focus of Barrington (1888) is on precipitation deficits, he does [also](#) note [the](#) temperature “On Sunday, June 5th, the temperature rose, and for the remainder of the month we had a combination of heat and drought, which lasted until July 10th. No record exists of such a hot and dry June in the south and southeast of Ireland”. Barrington (1888) goes on to [mention](#) that “... in June and July 1887, the S. of Ireland suffered more from excess of heat than any part of England or Scotland.” [further cementing the plausibility that the Island of Ireland was experiencing extreme hot and dry conditions at the time of the Kilkenny record high.](#)





Figure 3. The maximum [air](#) surface temperatures of the [six](#) selected weather stations for the month of June 1887. The [red point](#) marker displays the maximum surface temperature recorded at Kilkenny on the 26<sup>th</sup> June 1887.

The 20CRv3 reanalysis product (Slivinski et al., 2019) extends back to the early 19<sup>th</sup> Century using surface pressure and prescribed [sea surface temperature](#) boundary conditions to create physically consistent reconstructions of climate. Examination of 20CRv3 fields [over the period around](#) 26<sup>th</sup> June 1887 highlights that there was a high pressure system in place throughout the examined period (Figure 4). Prior to the 26<sup>th</sup> June the high pressure is centred to

the north of Ireland advecting air from the near continent. On 26<sup>th</sup> June the high pressure collapses and is subsequently replaced by a building high pressure from the near Atlantic initially advecting cooler air from the north which becomes centred over Ireland by the end of the month. The evolving pressure situation is **highly** consistent with the timing and relative phasing of changes reported from each site in Figure 3.

NOAA 20CRv3 MSLP & 850-hPa Temperatures



210 **Figure 4.** [NOAA 20CRv3 MSLP and 850-hPa Temperatures over an 8 day period centred upon the 26th June 1887. 20th Century Reanalysis V3 data provided by the NOAA/OAR/ESRL PSL, Boulder, Colorado, USA, from their Web site at https://psl.noaa.gov/data/gridded/data.20thC\\_ReanV3.html. The domain of latitude 35N-70N and longitude 45W-30E was chosen https://psl.noaa.gov/data/gridded/data.20thC\\_ReanV3.html. The domain of latitude 35N-70N and longitude 45W-30E was chosen to provide a synoptic view, with Ireland centred to provide an overview of the surrounding regions. 850hPa temperatures are given in colours while MSLP is given as contours.](https://psl.noaa.gov/data/gridded/data.20thC_ReanV3.html)

On the basis of the synoptic situation (Figure 4) and the contiguous station series (Figure 3) it is certain that the period was indeed a notable heatwave event that could, plausibly, be consistent with a record high temperature. However, such a synoptic situation is not uncommon in the summer season over Ireland. In isolation this evidence is insufficient to determine the validity of the record.

220 With the exception of the Kilkenny Castle site, all sites used in Figure 3 have [nearly complete](#) records through to present, [many of which have been digitised](#). Since 2010, data from Kilkenny Greenshill (also referred to as Green's Hill or Green Hills), which is very close to the location of the original Kilkenny Castle site, is available via Met Éireann. [The Greenshill site is at a distance of 1.6 km NNW from Kilkenny castle \(Figure 5\). The elevation differs by approximately 12 m with Kilkenny castle lying at 58m asl while the Greenhills station lies 46m asl. Both sites are reasonably proximal to the River Nore, and both are surrounded by some urban infrastructure. Thus, the Kilkenny Castle and Greenshill stations are likely to have been broadly comparable. Although clearly the lack of a period of overlap leaves some irreducible uncertainty in this interpretation.](#) The availability of several years of modern data using modern instrumental techniques both near the site of the original Kilkenny record and at the sites of the historical comparator sites permits statistically based comparisons. [Summer season](#) timeseries from each station were [matched](#) to account for missing data and then modern differences between station pairs analysed to provide context for the evident offsets between Kilkenny Castle and remaining sites in Figure 3 [on 26<sup>th</sup> June 1887.](#) [For Birr, the modern data was only available for the 2010 summer which would have been insufficient to analyse the frequency distributions. As a result, Birr station was excluded from the modern-day frequency distribution analysis.](#)

**Commented [KD8]:** Review comment: The application of 20CR to show the synoptic situation seems not so persuasive, I feel. It indicates a favorable weather condition for the extreme hot and dry event, but when the authors said that the condition is not uncommon in the summer season over Ireland, they did not exhibit an example of the synoptic situation at present. Is there any similar or even more extreme high pressure system over the study region during the same length of periods in modern time?

**Commented [KD9R8]:** As above

**Commented [CM10]:** Need to look at phrasing here

**Formatted:** Not Highlight

**Formatted:** Superscript




Figure 5: Illustration of the relative positioning of Kilkenny Castle and Greenshill, Kilkenny (Google Earth v. 9.154.0.1 (2022)).

Formatted: Not Highlight

240

When looking at the modern-day differences between Markree and Kilkenny Greenshill 90% of the distribution lies between a 0°C = 5°C temperature difference (Kilkenny warmer than Markree on average, Figure 6). The difference recorded between Markree and Kilkenny Castle on 26<sup>th</sup> June 1887 was 9.9°C which is a substantial outlier. The bulk of the differences (95%) between Kilkenny Greenshill and Sheffield (2010-2017) lie between -5°C and 5°C (broader because of the far greater distance between the sites). Despite this increased dispersion of modern era differences, the difference between Sheffield and Kilkenny Castle on 26<sup>th</sup> June 1887 of 9.4°C again indicates a substantial outlier relative to modern inter-site characteristics. Only a handful of higher differences have been reached within the recent era. For Kilkenny Greenshill and Phoenix Park (2013-2019), 95% of the data lies between -2.5°C and 2.5°C with Kilkenny Greenshill generally measuring slightly higher temperatures than Phoenix Park. The difference on 26<sup>th</sup> June 1887 between Kilkenny Castle and Phoenix Park at 9°C, lies entirely outside the

245


250

Commented [CM12]: ditto



distribution of modern inter-site behaviour. The highest recorded temperature difference in the modern era was 7.5°C, indicating the heat record to be an extreme anomaly. Most (95%) of differences between Kilkenny Greenshill and Roches Point (2010 -2019) lie within a range between 0°C - 5°C, 95%) of

255 differences between Kilkenny Greenshill and Roches Point (2010 -2019) lie within a range between 0°C - 5°C, showing that here again Kilkenny Greenshill generally measures higher temperatures than Roches Point. The temperature record of the 26<sup>th</sup> of June 1887 was 6.6°C warmer at Kilkenny Castle than Roches Point and lies within the upper 2% of the differences. Similarly, the difference between Armagh and Kilkenny Greenshill (2010 - 2018) shows that 90% of the data lies within -2.5°C and 3°C. Here, the temperature difference of 6.4°C between Kilkenny Castle and Armagh also lies within the upper 2%, showing that such a temperature difference is unlikely to occur but not impossible.



265 **Figure 6.** Modern day distribution for the JJA season differences between the maximum temperature of Kilkenny Greenshill and the four stations where the data was most available. Shown in red is the difference reported on June 26<sup>th</sup> 1887 between each station and Kilkenny Castle.

The June 26<sup>th</sup> 1887 difference between Kilkenny Castle and all surrounding sites is highly anomalous being in the extreme upper tail at Markree, Sheffield, Roches Point and Armagh and entirely outside the distribution for Phoenix Park. Differences overall are highly anomalous in a broad arc from SSW through N to due E. While there is a change in circulation 270 Differences overall are highly anomalous in a broad arc from SSW through N to due E. While there is a change in circulation via advection of a northerly airmass this occurred on June 27<sup>th</sup> (Figures 3 and 4). there is no active frontal system that may support a multi-degree thermal gradient between Kilkenny and its neighbours that would support such an extreme departure from all neighbouring sites. The slack pressure gradient also does 275 not support such a strong temperature gradient between Kilkenny and neighbouring sites. While many of the Irish stations are coastal, Birr (for which unfortunately a modern data comparison was not possible) is not and would have been less impacted by any sea breezes that may have set up than Kilkenny. Overall, therefore, while June 1887 undoubtedly was an unusually hot month and there is strong support for a build-up of heat breaking on or around June 26<sup>th</sup>/27<sup>th</sup> the very anomalous differences compared to modern behaviour argue strongly against the validity of the reported 280 Kilkenny Castle record value. The question therefore is what value may constitute the ‘true’ Irish national heat record and when did it occur?

#### 4. Consideration of Alternative Candidate Record Events

Table 2 noted a number of other candidate dates and locations in which the reported temperature exceeded 32°C. Unlike the 285 Kilkenny Castle value, many of these observations have been made as part of long-term series that are digitally available. These permit, in addition to the analyses performed for Kilkenny Castle, a consideration of the evolution of the station series around the event, including for some cases the hour-by-hour evolution of temperatures on the day. Also, at least for more recent events there is a denser neighbour series network available to perform the neighbour-based comparison and the difference series can be calculated directly rather than via a replacement site as is the case in Kilkenny. Their 290 recency also means there is better metadata, and the records are generally made in a more standardised and uniform manner consistent with WMO guidance on methods of observation (WMO, 2018). However, first it is necessary to consider the viability of the other records unearthed by Mateus et al. (2020) many of which pre-date the Kilkenny Castle record.

##### 4.1 Early period candidate record high temperatures

Mateus et al. (2020) have digitised a range of early Irish meteorological stations and made these data available. In constructing 295 Table 2 the majority of the remaining observations above 32°C arise from this source, including an observation of 33.5°C from Phoenix Park, Dublin in July of 1876 which is even warmer than the Kilkenny Castle observations, and a set of observations made in the 1850s.

#### 4.1.1 1850s candidates

The suite of observations made in the 1850s preceded the formation of the UK Met Office (at the time the Republic of Ireland had not gained independence from the UK). Metadata on who exactly took these observations is incomplete but suggests they were undertaken by military engineers to whom it is assumed some degree of training had been given. This period precedes efforts to shield instruments using a Stevenson Screen (Stevenson, 1864) and instruments may typically have been housed on north facing walls or placed in ventilated rooms (Parker, 1994). There is substantial literature pointing to potential biases in these early records with a tendency for summertime maxima to be overestimated (Parker, 1994, Camuffo, 2002, Böhm et al., 2010, Trewin, 2010, Brunet et al., 2011). Specifically, modern-day comparisons either of original instruments and exposures (Böhm et al., 2010) or reconstructed instruments and exposures (Brunet et al., 2011) with standard Stevenson Screens highlight important potential biases in summertime daily maximum temperatures. These are particularly marked for mid-latitude north wall exposures that would, especially if not oriented to true north, catch significant solar radiation in mid-summer (Parker, 1994, Böhm et al., 2010).

A further concern over the validity of several of these 1850s high temperature records relates to their geographical situation. A reading of 32.2°C on Scattery Island, Clare, an Island off the west coast of less than 1km by 1km seems implausible when climatological sea surface temperatures are in the high teens. Markree in Sligo also appears multiple times. As a coastal location in the NW of Ireland it is difficult to envisage how temperatures as warm as 32°C or higher could be attained on such a regular basis. They also are very much bunched around late June which may indicate the presence of radiative effects around the solstice upon this measurement series or something as simple as annual leave measurements being taken incorrectly, by another observer. Kiltrish, in Clare is similarly coastal and similar questions would pertain around how plausible such extreme heat was so close to the Atlantic Ocean. The same concerns pertain to Dunmore East which is on the Irish Sea coast of SE Ireland. The Royal College of Surgeons in Dublin is relatively close to the quay so, again, it is questionable whether such a warm temperature could be attained even allowing for urban heat island effects.

The 1850s candidates also are sometimes associated with implausible synoptic situations according to NOAA 20CRv3 MSLP and 850-hPa Temperature reanalysis reconstructions (Figure 7). Even in the 1850s the available pressure constraint is sufficiently robust over NW Europe to provide a robust synoptic scale reconstruction. The three synoptic charts in 1851 (Figure 7) are associated with high pressure and a slack flow of air from the continent which would, generally, be associated with climatologically hot conditions. Of note is that over 28<sup>th</sup>/29<sup>th</sup> of June 1851, Kiltrish, Markree and Dunmore East all reported temperatures in excess of 32°C. Combined with the synoptic chart reconstructed from 20CRv3 there is little doubt that 28<sup>th</sup>/29<sup>th</sup> June 1851 was, indeed, a heatwave event across the island of Ireland. Newspapers at the time reflected these conditions reporting on the ‘hot’ and ‘oppressive’ weather that was occurring in Ireland (Freemans Journal, 1851; Leinster Express, 1851).

**Commented [PWT13]:** Conor may be able to furnish some details.

**Commented [314R13]:** Don't know much about history of temp observations. This is likely the case though. What does Mateus have to say on this? Even if you look at their summer and spring mean series for max temps in the plots there is an obvious warm bias prior to 1880, which suggests such problems.

**Commented [SN15R13]:** See the PP metadata file which names some of the observer and provides details.

**Commented [KD16R13]:** 1855- 56: Sergeant George Brown, Thomas Rhodes (Royal Sappers and Miners)  
1856-58: Sergeant George Brown. Computations by C.A. John Kelly/Sapper John Meade/ Gerard Liddy (who is down as a civil assistant)/ Thomas A. Dennann  
1860-62: Sergeant George Brown and Sapper Edgar White/  
Computation by Thomas Dennann/William brown

Military Engineers basically

**Formatted:** Superscript

**Formatted:** Superscript

**Formatted:** Superscript

The remaining dates [in the 1850s](#) are associated synoptically with conditions that would tend to be climatically normal or even below normal temperatures. The 27/06/1852 chart has a low pressure to the north of Ireland with a slack north-westerly flow which would be an onshore wind at [Markree](#), Sligo, and completely inconsistent with a temperature in excess of 32°C.


335 The 26/06/1853 event, again for a temperature at [Markree](#), Sligo, is represented by a southwesterly flow, which is grossly inconsistent with temperatures as high as the low 30s in this location. The situation on 29/06/1854 is of a low pressure centred over England with a northerly flow, again inconsistent with temperatures in excess of 32°C at [Markree](#). The slack pressure gradient on 03/08/1856 leads to potentially a slight easterly onshore wind inconsistent with a high temperature at the Royal College of Surgeons in Dublin.

340

Taken together, the uncertainty relating to early instrumental bias, climatological locations, and in many cases lack of substantive synoptic situation support mean that the 1850s values contained in Table 2 almost certainly should be precluded as robust candidates for record temperatures. That is not to say that, at least in some cases, the values may [not](#) be correct or at the very least indicative of, climatologically speaking, extreme heat. But [Markree](#) in particular, which occurs the most frequently, has values that are often not supported at all by the synoptic situation.

345

NOAA 20CRv3 MSLP & 850-hPa Temperatures



350 Figure 7. [NOAA 20CRv3 MSLP and 850-hPa Temperatures](#) on the dates of the possible heat records in the 1850s arising from Mateus et al. (2020) recent data rescue and as documented in Table 2. [20th Century Reanalysis V3 data provided by the NOAA/OAR/ESRL PSL, Boulder, Colorado, USA, from their Web site at https://psl.noaa.gov/data/gridded/data.20thC\\_ReanV3.html](#). The domain of latitude 35N-70N and longitude 45W-30E was chosen to provide a synoptic view, with Ireland centred to provide an overview of the surrounding regions.


#### 4.1.2 The Phoenix Park value of 33.5°C

355 The Phoenix Park site is the longest running contiguous site in the Republic of Ireland. Originally run by the Ordnance Survey, then the military, it is now maintained by Met Éireann. It is a little unclear exactly how the potential record on 16<sup>th</sup> July 1876 was measured. Mateus et al. (2020) mentions “*Drawings when no station photographs are available for the early 19th century, for example Cameron (1856) of the thermometer screen at Phoenix Park Dublin, are furnished*”. In 1879 there is the first mention of measurements by a Stevenson Screen “*quality of the record improved and since 1879/1880, when the thermometers*  
360 *were housed in a Stevenson screen, the data may be considered to have a high level of accuracy and reability [sic]*” (Irish Meteorological Service, 1983: 4). It is hence probable, but not certain, that the measurement was made by a thermometer housed in something other than a Stevenson Screen, although exactly what is hard to ascertain precisely. The site is about 5km inland from Dublin Bay such that, in the absence of a sea-breeze, climatologically speaking, a temperature in the 30s could potentially be attained. The synoptic situation as reconstructed by [NOAA 20CRv3 MSLP and 850-hPa Temperatures](#) is supportive of very warm conditions with a high pressure of 1025hPa centred over the British and Irish isles (Figure 8). Preceding days had had a gentle flow of air from the near continent. There is thus clear synoptic support for very high temperatures.

Formatted: Font: Italic

Formatted: Font: Italic

NOAA 20CRv3 MSLP & 850-hPa Temperatures





370 Figure 8. NOAA 20CRv3 MSLP and 850-hPa Temperatures evolution around the Phoenix Park observed value of 33.5°C in 1876 on 16<sup>th</sup> July. 20th Century Reanalysis V3 data provided by the NOAA/OAR/ESRL PSL, Boulder, Colorado, USA, from their Web site at [https://psl.noaa.gov/data/gridded/data.20thC\\_ReanV3.html](https://psl.noaa.gov/data/gridded/data.20thC_ReanV3.html) . The domain of latitude 35N-70N and longitude 45W-30E was chosen to provide a synoptic view, with Ireland centred to provide an overview of the surrounding regions.

375 Having discounted the long-standing Kilkenny Castle national record in Section 3 and, in Section 4.1, ruled out as sufficiently robust any values from the 1850s, this leaves the Phoenix Park temperature a full 1°C warmer than any other remaining candidate value in Table 2. Most of these remaining candidate values are from far further inland where such high temperatures a priori would be more easily attainable. When analysed against other Dublin city temperature observations for the month of June 1876 (Figure 9) it can be seen that, while a similar temperature evolution is mostly followed throughout the month, the Phoenix Park station is consistently hotter than the other Dublin stations (except for the 3<sup>rd</sup> and 26<sup>th</sup> day). Dixon (1953) reported a sudden and rapid temperature high just a few years earlier, on the 21<sup>st</sup> July 1868; “the temperature in Phoenix Park one day during July shot up to 88.4 degrees F. (31.3°C), a record which has yet to be beaten”. At this time, the Botanic Gardens similarly recorded a very hot July, however the temperature recorded on the 21<sup>st</sup> of July is 81 degrees F (27.2°C), several degrees below the Phoenix Park recording despite their close proximity.

385 It is clear from the 20CRv3 reconstruction and the surface temperature recordings shown in Figure 8 that the synoptic situation in July 1876 supports very warm conditions. However, the near constant systematic elevation of the temperature readings in Phoenix Park relative to other stations casts doubt on whether it was reporting the actual surface temperature at the time. Phoenix Park is in an elevated position so on lapse rate basis alone would be expected to record slightly lower and not higher temperatures than nearby sites. While the value at the Phoenix Park cannot be discounted there is equally no obvious way to sufficiently robustly confirm the While the value at the Phoenix Park cannot be discounted there is equally no obvious way to sufficiently robustly confirm the While the value at the Phoenix Park cannot be discounted there is equally no obvious way to sufficiently robustly confirm the observation for it to be adequately verified as a national heat record.

395

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Font: Italic

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Superscript

**Commented [SN17]:** I have sent the sheet containing the PP comparison with Fitzwilliam sq, Trinity and Botanic gardens which show large differences in tmax for 1876. Should note these findings here.




Figure 8: The maximum surface temperatures of all available stations in Dublin City during June 1876.

400


#### 4.2 Candidates since the beginning of the 20th Century

The remaining candidate events in which the reported temperature attained or exceeded 32°C are all associated with synoptically broadly similar conditions. High pressure is situated over or to the north / north east of the island of Ireland, often with weak advection of air from the near continent (Figure 10). For all cases the 850hPa temperatures are elevated. The synoptic situations thus do not call into immediate question any of the remaining candidate values realism.


405

410

NOAA 20CRv3 MSLP & 850-hPa Temperatures



ERA5 MSLP & 850-Temperature



415 Figure 10. As Figure 4 but for the dates of post-1900 possible record heat events as detailed in Table 2. Data provided by the NOAA/OAR/ESRL PSL, Boulder, Colorado, USA, from their Web site at [https://psl.noaa.gov/data/gridded/data.20thC\\_ReanV3.html](https://psl.noaa.gov/data/gridded/data.20thC_ReanV3.html) (for NOAA 20CRv3 MSLP and 850-hPa Temperatures) and Hershbach et al. (2018) (for ERA5 during its period of record). The domain of latitude 35N-70N and longitude 45W-30E was chosen to provide a synoptic view, with Ireland centred to provide an overview of the surrounding regions.

420 Boora is the hottest value of the se remaining candidates at 32.5°C, attained on 29/06/1976. Boora is situated in the centre of the island of Ireland, which means it is highly unlikely to suffer from marine influences which may cap diurnal maxima closer to the coast. The summer of 1976 was famous across the British and Irish Isles for its drought and hot summer following on from 18 dry months previously (Noone et al. 2017; O’Laoghog, 1979). Therefore, the partitioning of heat between sensible (temperature) and latent (evaporation) terms would have been climatologically skewed toward sensible heating. The value at Boora furthermore is situated within a string of very warm days with many of the daily maximum observations exceeding

425 30°C.

430 The hot and dry conditions were widely reported in newspapers at the time; The Irish Press reported on 24<sup>th</sup> June 1976 ‘The Irish have been greeted by a heatwave which saw the temperature hover around the 88 degree (31.1°C) mark for a long time’ (Redmond, 1976). On 28<sup>th</sup> June 1976, the Irish Press reported on the weather in Britain and Europe: ‘you cannot expect to share the 90 degree (32.2°C) heatwave that has had Britain sweltering for days. But there is a chance you will soon be basking in a purely Irish hot spell, with temperatures reaching 85 degrees F (29.4°C). Britain, because of its proximity to the Continent and its distance from the Atlantic, is enjoying the effects of an anti-cyclone which is almost stationary over Europe...According to the Dublin Met Office, this could mean an Irish heatwave within the next few days, with temperatures rising from the present 60 - 70 degrees F. (15.5 – 21.1°C) to 86 degrees F (30°C)’ (Cahill, 1976).

435 A comparison to neighbouring sites for this candidate event benefits from the standardisation of meteorological observations during the 20<sup>th</sup> Century and the much denser network of Met Éireann sites since the mid-20<sup>th</sup> Century. Comparisons can now be made to sites at tens of kilometres distance rather than some hundreds of kilometres.

440 These elevated temperatures, experienced at Boora, were also experienced at the nearby stations which sit within tens of kms elevated temperatures, experienced at Boora, were also experienced at the nearby stations which sit within tens of kms of the Boora site (Table 3).

ID	Station name	Distance from Boora (KM)	Latitude	Longitude	Elevation (M)	Observed Maximum Temperature (°C)	Diff (°C)	Data Source

Formatted Table

2322	Boora	-	53.22	-7.72	58	32.5		Met Éireann
875	Mullingar	41	53.31	-7.21	101	28.8	-3.7	Met Éireann
1614	Clonsast Bord Na Mona	35	53.11	-7.12	73	30.8	-1.7	Met Éireann
3431	Derrygreenagh	36	53.21	-7.15	90	28.7	-3.8	Met Éireann
4414	Ballybrittas	41	53.06	-7.08	90	32	-0.5	Met Éireann

Table 3. Comparison between Boora, Offaly maximum temperature on 29/06/1976 and neighbouring stations within 50 km radius.

Formatted: Font color: Text 1

445

A comparison to neighbouring sites benefits from the standardisation of meteorological observations during the 20<sup>th</sup> Century and the much denser network of Met Éireann sites since the mid-20<sup>th</sup> Century. Comparisons can now be made to sites at tens of kilometres distance rather than some hundreds of kilometres. Commensurately the distribution of statistical differences in summertime maximum temperatures are considerable smaller. Repeating the methodology of the Kilkenny neighbour-based analysis but using Boora and nearby stations for a period of up to 20 years either side of the record, the observation in question sits well within the distributions of expected differences in daily maxima over summer (Figure 911).

450

In the modern-day differences between Boora and its four neighbouring stations – Mullingar, Clonsast, Derrygreenagh and Ballybrittas, most of the inter-site distribution lies between  $\pm 1.5^{\circ}\text{C}$  – 1.5°C temperature difference. On 29<sup>th</sup> June 1976 the difference recorded between Boora and Mullingar was 3.7°C which falls within the upper quartile 2% of its distribution. Differences between Boora and Clonsast are generally between -2°C and 2.5°C and the heat record difference on 29<sup>th</sup> June 1976 was 1.7°C, falling lower on the range, within the upper 10% interquartile range. For Boora and Derrygreenagh, most of the data lies between -1.5°C – 2°C with Boora typically measuring slightly higher temperatures than Derrygreenagh. The difference on 29<sup>th</sup> June 1976 between the two sites at 3.8°C, lies entirely outside the interquartile range but still falling within the upper quartile of the differences within the upper reaches 2% of the expected differences but has still been exceeded on several occasions. Lastly, Boora and Ballybrittas have very strong similarities in the recorded temperature with most of the differences between -2°C and 2°C. The June 29<sup>th</sup> 1976 difference between these stations was 0.5°C completely within the distribution of modern inter-site behaviour.


455

460

Formatted: Not Highlight

Commented [PWT18]: Just check this – visually may be in upper quartile?

Formatted: Not Highlight



465 **Figure 11.** Modern day distribution for the JJA season differences between the maximum temperature of Boora, Offlay and the four stations where the data was most available. Shown in red is the difference reported on June 29<sup>th</sup> 1976 between each station and Boora.

470 A combination of geographical location, the synoptic situation, antecedent conditions, neighbour comparisons and use of  
 475 standard modern-day meteorological observations leads us to conclude the Boora observation is *likely to be a credible reading*.  
 We would thus conclude logically that this represents the highest reliably recorded temperature in the Republic of Ireland since  
 observations began. If the observation were to be called into question any of several candidates between 32°C and 32.3°C  
 since 1900 based upon the synoptic situation on each date would be viable alternatives, but we do not consider these in greater  
 depth herein.

## 5. Conclusions

*Ireland is now highly anomalous in having its recognised national heat record set in the 19<sup>th</sup> Century. The record is from  
 a long-closed site in Kilkenny Castle and the associated series is not available, presumed lost, as is the bulk of relevant  
 metadata. The Kilkenny record of 33.3°C in June 1887 exceeds that reached since the beginning of the 20<sup>th</sup> Century by 0.8°C.*

480 *Results from our inter-station reassessment and reanalysis comparisons leads us to question if this truly represents the highest  
 reliably measured national temperature for Ireland. Based upon the few contemporaneous station records available on the  
 island of Ireland and at Sheffield and their modern differences to a site close to the original Kilkenny Castle site, our findings  
 cast very considerable doubt upon the record. Differences on the 26<sup>th</sup> June 1887 are implausibly large and not consistent with  
 the synoptic situation as reanalysed by NOAA 20CRv3.*

485 A range of similar concerns *including basic physical considerations* preclude a number of other 19<sup>th</sup> Century values in excess

Formatted: Font: Italic

A range of similar concerns including basic physical considerations preclude a number of other 19<sup>th</sup> Century values in excess of 32°C, including an observation of 33.5°C at Phoenix Park in 1876.

A search of candidate records since 1887 yields that the highest likely defensible observation is 32.5°C recorded at Boora on 29<sup>th</sup> June 1976.

This analysis only considered the national all-time heat record. Met Éireann maintains all time annual and monthly records, several of which date back to similarly early, pre-standardisation, measurements (Met Éireann, 2020). Most notably the all time cold record is -19.1°C, recorded in Markree, Sligo in January 1881 which is close to the Atlantic coast in NW Ireland. Reassessment of these records using a range of techniques such as those used here would likely be valuable.

It is probable that Ireland is far from alone in having national meteorological heat, cold, rainfall and other records that may benefit from a reassessment using these new techniques pioneered by WMO record assessment teams. Given the importance of national all-time meteorological records in media and the general public discourse around both weather and climate change it is important that the verity of these records be assured. The techniques and approaches used herein are broadly transferable to any similar reconsideration of meteorological records in other jurisdictions.

**Code availability.** The data and code that support the findings of this study are available through: <https://github.com/katherinedooley/Reassessing-Ireland-s-Hottest-Temperature-Record>.

**Competing interests.** The authors declare that they have no conflict of interest.

**Author contributions.** SOK, CC, EC and KM conducted a primary investigation and provided background information on the study. KD, CK, NS, and JKD conducted an inter-station series assessment to assess the plausibility of the reported record. DC, NM, JD, TM created and analysed reanalysis weather maps for the top recorded hottest years 1887, 1995, 2006 and 2018. RS, EG, JC explored the plausible national heat records if Kilkenny 1887 record was to fail. SN and CM provided data and support along with contributions from Met Éireann. KD, CK and PT prepared the paper with contributions from all the co-authors.

**Acknowledgements.** The authors would like to extend our gratitude to ICARUS and the Department of Geography in Maynooth University for their help on this assignment for the MSc in climate change.

Formatted: Not Highlight

Formatted: Font: 10 pt



## References

520 [Allan, R., Brohan, P., Compo, G.P., Stone, R., Luterbacher, J. and Brönnimann, S., \(2011\). The international atmospheric circulation reconstructions over the earth \(ACRE\) initiative. \*Bulletin of the American Meteorological Society\*, 92\(11\), pp.1421-1425 <https://doi.org/10.1175/2011BAMS3218.1>.](#)

525 Barrington, R. M. (1888). The drought of 1887, and some of its effects on Irish agriculture. *Journal of the Statistical and Social Inquiry Society of Ireland*, 9, 223. <https://search.proquest.com/docview/911736931?pq-origsite=gscholar&fromopenview=true>

530 Böhmer, R., et al. (2010). The early instrumental warm-bias: a solution for long central European temperature series 1760–2007 *Climatic Change*, 101: 41. <https://doi.org/10.1007/s10584-009-9649-4>

535 [Brönnimann, S., Allan, R., Ashcroft, L., Baer, S., Barriendos, M., Brázdil, R., Brugnara, Y., Brunet, M., Brunetti, M., Chimani, B. and Cornes, R., \(2019\). Unlocking pre-1850 instrumental meteorological records: A global inventory. \*Bulletin of the American Meteorological Society\*, 100\(12\), pp.ES389-ES413. <https://doi.org/10.1175/BAMS-D-19-0040.1>.](#)

535 [Brunet, M. and Jones, P., \(2011\). Data rescue initiatives: bringing historical climate data into the 21st century. \*Climate Research\*, 47\(1-2\), pp.29-40. <https://doi.org/10.3354/cr00960>.](#)

540 [Brunet, M., Asin, J., Sigró, J., Bañón, M., García, F., Aguilar, E., Palenzuela, J.E., Peterson, T.C. and Jones, P. \(2011\). The minimization of the screen bias from ancient Western Mediterranean air temperature records: an exploratory statistical analysis. \*Int. J. Climatol.\*, 31: 1879-1895, <https://doi.org/10.1002/joc.2192>](#)

Cahill, A., (1976) 'How the Irish keep their cool', *Irish Press*, 28<sup>th</sup> June 1976. Available at: <https://archive-irishnewsarchive-com.jproxy.nuim.ie/Olive/APA/INA.Edu/SharedView.Article.aspx?href=IPR%2F1976%2F06%2F28&id=Ar00101&sk=74D555A7> (accessed 22 March 2021).

545 [Cameron, H.J. \(1856\) \*Meteorological observations take during the year 1829 to 1852 at the Ordnance Survey Office, Phoenix Park Dublin: to which is added a series of similar observations made at the principal trigonometrical stations, and at other places, in Ireland.\* Dublin, Ireland: Alexander Thom and Sons.](#)

550 [Camuffo, D. \(2002\). Errors in Early Temperature Series Arising from Changes in Style of Measuring Time, Sampling Schedule and Number of Observations. \*Climatic Change\*, 53, 331–352. <https://doi.org/10.1023/A:1014962623762>](#)

Formatted: Font: Times New Roman, 10 pt

Field Code Changed

Formatted: Font: Italic

Formatted: Font: Times New Roman, 10 pt

Field Code Changed

City Population (2021) KILKENNY (Ireland) [online]. Available at:  
[https://www.citypopulation.de/en/ireland/towns/KK\\_kilkenny/](https://www.citypopulation.de/en/ireland/towns/KK_kilkenny/) (accessed 08th December 2021).

Cerveny, R.S., Bessemoulin, P., Burt, C.C., Cooper, M.A., Cunjie, Z., Dewan, A., Finch, J., Holle, R.L., Kalkstein, L., Kruger, A. and Lee, T.C., (2017). WMO assessment of weather and climate mortality extremes: lightning, tropical cyclones, tornadoes, and hail. *Weather, climate, and society*, 9(3), pp.487-497.  
<https://doi.org/10.1175/WCAS-D-16-0120.1>  
<https://doi.org/10.1175/WCAS-D-16-0120.1>  
<https://doi.org/10.1175/WCAS-D-16-0120.1>  
<https://doi.org/10.1175/WCAS-D-16-0120.1>

Courtney, J., Buchan, S., Cerveny, R., Bessemoulin, P., Peterson, TC, Rubiera Torres, JM, Beven, J, King, J, Trewin, B & Rancourt, K (2012). 'Documentation and verification of the world extreme wind gust record: 113.3 m s<sup>-1</sup> on Barrow Island Australia, during passage of tropical cyclone Olivia', *Australian Meteorological and Oceanographic Journal*, vol. 62, no. 1, pp. 1-9, <https://doi.org/10.22499/2.6201.001>

Curley, M., (2019). Interviewed by Conor Murphy, Peter Thorne & Sophie O'Kelly. Email . 14 October 2019.

Dixon, F.E., (1953). *Weather in old Dublin*. *Dublin Historical Record*, 13(3/4), pp.94-107.

Dollar, N., (2021). Email correspondence with Katherine Dooley, 20<sup>th</sup> April 2021.

El Fadli, K., Cerveny, R., Burt, C., Eden, P., Parker, D., Brunet, M., Peterson, T., Mordacchini, G., Pelino, V., Bessemoulin, P., Stella, J., Driouech, F., Wahab, M. and Pace, M., (2013). World Meteorological Organization Assessment of the Purported World Record 58°C Temperature Extreme at El Azizia, Libya (13 September 1922). *Bulletin of the American Meteorological Society*, 94(2), pp.199-204. <http://www.jstor.org/stable/26219496>.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N. (2018): ERA5 hourly data on pressure levels from 1979 to

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS). (Accessed 20<sup>th</sup> January 2022).  
[10.24381/cds.bd0915c6](https://cds.clm.copernicus.com/cds/details/?tableId=10.24381/cds.bd0915c6)

Google Earth v. 9.152.0.1 (2021). *Kilkenny Castle, grounds and surrounds*, 52°38'43.37"N 7°13'32.19"W Elevation 4084 m and 52°39'00.78"N 7°14'40.78"W Elevation 876 m, [Online] Available at: <https://earth.google.com/web/> (Accessed 14 December, 2021).

Google Earth v. 9.154.0.1 (2022). *Kilkenny Castle to Green's Hill*, 52°39'00"N 7°14'51"W Elevation 64 m and 52°39'40"N 7°15'15"W Elevation 46 m. [Online] Available at: <https://earth.google.com/web/> (Accessed 20 January 2022).

Irish Meteorological Service. (1983). *The climate of Dublin*. Dublin: Irish Meteorological Service.  
Irish Meteorological Service. (1983). *The climate of Dublin*. Dublin: Irish Meteorological Service.  
Irish Meteorological Service. (1983). *The climate of Dublin*. Dublin: Irish Meteorological Service.

Khan, F., (1976) 'This heat wave, it's just TANTastic', Irish Independent, 29th June 1976. Available at: <https://archive-irishnewsarchive-com.jproxy.nuim.ie/Olive/APA/INA.Edu/SharedView.Article.aspx?href=IND%2F1976%2F06%2F29&id=Ar00102&sk=242CE860> (accessed 22 March 2021).

Lang, T.J., Pédeboy, S., Rison, W., Cervený, R.S., Montanyà, J., Chauzy, S., MacGorman, D.R., Holle, R.L., Ávila, E.E.,  
Lang, T.J., Pédeboy, S., Rison, W., Cervený, R.S., Montanyà, J., Chauzy, S., MacGorman, D.R., Holle, R.L., Ávila, E.E.,  
Láska, K., King, J., Bromwich, D., Jones, P., Solomon, S., Renwick, J., Lazzara, M., de los Milagros Skansi, M., Brunet, M.,

'London Markets. Corn Exchange - Monday', *Leinster Express*, 28<sup>th</sup> June 1851. Available at: <https://archive-irishnewsarchive-com.jproxy.nuim.ie/Olive/APA/INA.Edu/SharedView.Article.aspx?href=LEE%2F1851%2F06%2F28&id=Ar00301&sk=C5F36D51> (accessed 29 April 2021).

Mateus, C, Potito, A, Curley, M. (2020) Reconstruction of a long-term historical daily maximum and minimum air temperature  
McDill, Z. (2021) A City in Co. Kilkenny in Ireland [online]. Available at:

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: Italic

Formatted: Font: (Default) Times New Roman

Merlone, A., Al-Dashti, H., Faisal, N., Cervený, R.S., AlSarmi, S., Bessemoulin, P., Brunet, M., Driouech, F., Khalatyan, Y., Peterson, T.C. and Rahimzadeh, F., (2019). Temperature extreme records: World Meteorological Organization metrological and meteorological evaluation of the 54.0 C observations in Mitribah, Kuwait and Turbat, Pakistan in 2016/2017. *International Journal of Climatology*, 39(13), pp.5154-5169.

DOI: 10.1002/joc.6132

DOI: 10.1002/joc.6132

DOI: 10.1002/joc.6132

DOI: 10.1002/joc.6132

Met Éireann (2020). *Weather Extreme Records for Ireland*. [online] Available at: <https://www.met.ie/climate/weather-extreme-records> (accessed 13th August 2021).

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Murphy, C, Wilby, R.L, Matthews, T.K.R., Thorne P., Broderick C., Fealy R., Hall J., Harrigan S., Jones P., McCarthy G., MacDonald N., Noone S., Ryan C. (2020) Multi-century trends to wetter winters and drier summers in the England and Wales precipitation series explained by observational and sampling bias in early records. *Int J Climatol.*; 40: 610– 619. <https://doi.org/10.1002/joc.6208>

NOAA Physical Science Laboratory (2020). *The Twentieth Century Reanalysis Project* [online]. Available at: NOAA Physical Science Laboratory (2020). *The Twentieth Century Reanalysis Project* [online]. Available at:

Noone, S., Broderick, C., Duffy, C., Matthews, T., Wilby, R.L. and Murphy, C., (2017). A 250-year drought catalogue for the Island of Ireland (1765–2015). *International Journal of Climatology*, 37, pp.239-254. <https://doi.org/10.1002/joc.4999>

Formatted: Font: Italic

O'Laoghog S.S. (1979). The Dry Period October 1974 to August 1976 [report], Met Éireann. Internal Memorandum, 88/79, 1979, 1979-01.

Parker, D. E. (1994). Effects of changing exposure of thermometers at land stations. *International Journal of Climatology*, 14, pp. 1-31. <https://doi.org/10.1002/joc.3370140102>

Peterson, M.J., Lang, T.J., Bruning, E.C., Albrecht, R., Blakeslee, R.J., Lyons, W.A., Pédeboy, S., Rison, W., Zhang, Y., Brunet, M. and Cervený, R.S., (2020). New WMO certified megaflash lightning extremes for flash distance (709 km) and duration (16.73 seconds) recorded from space. *Geophysical Research Letters*, 47,

DOI: 10.1029/2020GL088888.

655 Purevjav, G., Balling Jr, R.C., Cerveny, R.S., Allan, R., Compo, G.P., Jones, P., Peterson, T.C., Brunet, M., Driouech, F.,  
Stella, J.L. and Svoma, B.M.  
(2015). The Tosontsengel Mongolia  
world record sea-level pressure extreme: spatial analysis of elevation bias in adjustment-to-sea-level pressures *Int. J.  
Climatology* 35 (10) 2968-2977; DOI: 10.1002/joc.4186

660

Quetelard, H., Bessemoulin, P., Cerveny R.S., Peterson, T.C., Burton, A., and Y. Boodhoo,  
(2007). World record rainfalls (72-hour and four-day accumulations) at Cratère Commerson, Réunion Island, during the  
passage of Tropical Cyclone Gamede, *Bulletin of the American Meteorological Society*, 90(5): 603-608

665

Redmond, J., (1976). 'Sun Sets a Poser', *Irish Press*, 24<sup>th</sup> June 1976, Available at: [https://archive-irishnewsarchive-  
com.jproxy.nuim.ie/Olive/APA/INA.Edu/SharedView.Article.aspx?href=IPR%2F1976%2F06%2F24&id=Ar01400&sk=E0  
C59558](https://archive-irishnewsarchive-com.jproxy.nuim.ie/Olive/APA/INA.Edu/SharedView.Article.aspx?href=IPR%2F1976%2F06%2F24&id=Ar01400&sk=E0C59558) (accessed 22 March 2021).

670

Report of the Meteorological Council (1890). Report of the Meteorological Council to the Royal Society For the Year ending  
31<sup>st</sup> of March 1890. Q.J.R. Meteorol. Soc., 16: 73. Available at:  
[https://play.google.com/books/reader?id=YnEMAAAYAAJ&pg=GBS.PA72&hl=en\\_GB](https://play.google.com/books/reader?id=YnEMAAAYAAJ&pg=GBS.PA72&hl=en_GB) (accessed 30<sup>th</sup> July 2021).

675

RMS (1890). Report of the council for the year 1889. Q.J.R. Meteorol. Soc., 16: 86-  
119. <https://doi.org/10.1002/qj.4970167403>

Rohan, P.K. (1986). *The Climate of Ireland. 2<sup>nd</sup> Ed. Dublin: Meteorological Service.*

680

Schildkamp, V. (2019). So it was 42.9 degrees, where did the mysterious heat in Deelen come from? *Algemeen Dagblad*  
[online] 3 December 2019. Available at:  
[https://www.ad.nl/binnenland/het-was-dus-wel-42-9-graden-  
https://www.ad.nl/binnenland/het-was-dus-wel-42-9-graden-  
waar-kwam-die-mysterieuze-hitte-in-deelen-vandaag-br~a07dfb22/?referrer=https%3A%2F%2Fen.wikipedia.org%2F](https://www.ad.nl/binnenland/het-was-dus-wel-42-9-graden-waar-kwam-die-mysterieuze-hitte-in-deelen-vandaag-https://www.ad.nl/binnenland/het-was-dus-wel-42-9-graden-waar-kwam-die-mysterieuze-hitte-in-deelen-vandaag-br~a07dfb22/?referrer=https%3A%2F%2Fen.wikipedia.org%2F)  
(Accessed 20 October 2020)

Formatted: Font: Not Italic

Formatted: Font: Italic, Italian (Italy)

Formatted: Italian (Italy)

Formatted: Font: (Default) Times New Roman

Formatted: Hyperlink, Font: (Default) Times New Roman

685 Slivinski, L.C., Compo, G.P., Whitaker, J.S., et al. (2019) Towards a more reliable historical reanalysis: Improvements for  
version 3 of the Twentieth Century Reanalysis system. Quarterly Journal of the Royal Meteorological Society,  
145: 2876– 2908. <https://doi.org/10.1002/qj.3598>

690 ‘Smithfield Market – London, Friday’, *Freemans Journal*, 30<sup>th</sup> June 1851. Available at: [https://archive-irishnewsarchive-  
com.jproxy.nuim.ie/Olive/APA/INA.Edu/SharedView.Article.aspx?href=FMJ%2F1851%2F06%2F30&id=Ar00312&sk=73  
6004A9](https://archive-irishnewsarchive-com.jproxy.nuim.ie/Olive/APA/INA.Edu/SharedView.Article.aspx?href=FMJ%2F1851%2F06%2F30&id=Ar00312&sk=736004A9) (accessed 29 April 2021).

Smith B., (2012), ‘Hottest temperature on earth revised’. *The Sunday Morning Herald*, September 14<sup>th</sup>. [online] Available at:  
695 <https://www.smh.com.au/environment/hottest-temperature-on-earth-revised-20120914-25wji.html> (Accessed 16 October  
<https://www.smh.com.au/environment/hottest-temperature-on-earth-revised-20120914-25wji.html> (Accessed 16  
October 2020).

700 [South African Weather Station \(2019\), \*Violsdrif temperature on 28 November 2019\* \[online\]. Available at:  
https://www.weath  
ersa.co.za/Documents/Corporate/Medrel2Dec2019b%20Violsdrif%20temperature.pdf](https://www.weath) (Accessed 20 October 2020).

[Stevenson, T.C.E. \(1864\). New Description of Box for Holding Thermometers. \*Journal of the Scottish Meteorological  
Society\*, 1, pp 122, June 1864.](#)

705 [Stubbs, M.W. \(1977\). EXCEPTIONAL EUROPEAN WEATHER IN 1976\\*. \*Weather\*, 32: 457-  
463. https://doi.org/10.1002/j.1477-8696.1977.tb04504.x](https://doi.org/10.1002/j.1477-8696.1977.tb04504.x)

Trewin, B. (2010). Exposure, instrumentation, and observing practice effects on land temperature measurements, *WIREs  
Clim. Change*, 1, 490–506. <https://doi.org/10.1002/wcc.46>" <https://doi.org/10.1002/wcc.46>

710 [UCAR \(2021\). 3. Conventional Meteorological Station Data. \*An Introduction to Atmospherice and Oceanic Datasets\* \[online\].  
Available at: https://www.cgd.ucar.edu/cas/tn404/text/tn404\\_6.html \(Accessed 01 July 2021\).](https://www.cgd.ucar.edu/cas/tn404/text/tn404_6.html)

715 [Weidner, G., King, J., Box, J.E., Colwell, S., Jones, P., Lazzara, M., Cappelen, J., Brunet, M. and Cerveny, R.S., 2021. WMO  
evaluation of northern hemispheric coldest temperature:– 69.6° C at Klinck, Greenland, 22 December 1991. \*Quarterly Journal  
of the Royal Meteorological Society\*, 147\(734\), pp.21-29  
. <https://doi.org/10.1002/qj.3901>](https://doi.org/10.1002/qj.3901)

Formatted: Font: (Default) Times New Roman

Formatted: Hyperlink, Font: (Default) Times New Roman,  
Font color: Auto

Formatted: Font: (Default) Times New Roman

Formatted: Hyperlink, Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, English  
(Ireland)

Formatted: Font: (Default) Times New Roman, Font color:  
Auto

. <https://doi.org/10.1002/qj.3901>

720 WMO (2018). *Guide to Instruments and Methods of Observation, Volume I – Measurement of meteorological variables.*  
WMO (2018). *Guide to Instruments and Methods of Observation, Volume I – Measurement of meteorological variables.*

WMO (2021). *Main | World Meteorological Organization's World Weather & Climate Extremes Archive.* [online]  
Available at: <https://wmo.asu.edu/#global> "<https://wmo.asu.edu/#global>", [Accessed 9 April 2021].

725

730

735

**Formatted:** Hyperlink, Font: (Default) Times New Roman,  
Font color: Auto

**Formatted:** Font: (Default) Times New Roman