

## Sclerochronological evidence of pronounced seasonality from the late Pliocene of the southern North Sea Basin, and its implications

Andrew L.A. Johnson<sup>1</sup>, Annemarie M. Valentine<sup>2</sup>, Bernd R. Schöne<sup>3</sup>, Melanie J. Leng<sup>4</sup>, Stijn Goolaerts<sup>5</sup>

<sup>1</sup>*School of Built and Natural Environment, University of Derby, Derby DE22 1GB, UK*

<sup>2</sup>*School of Geography and Environmental Science, Nottingham Trent University, Southwell NG25 0QF, UK*

<sup>3</sup>*Institute of Geosciences, University of Mainz, 55128 Mainz, Germany*

<sup>4</sup>*National Environmental Isotope Facility, British Geological Survey, Keyworth NG12 5GG, UK*

<sup>5</sup>*OD Earth & History of Life and Scientific Heritage Service, Royal Belgian Institute of Natural Sciences, 1000 Brussels, Belgium*

Correspondence to: Andrew L.A. Johnson (a.l.a.johnson@derby.ac.uk)

**Abstract.** Oxygen isotope ( $\delta^{18}\text{O}$ ) sclerochronology of benthic marine molluscs provides a means of reconstructing the seasonal range in seafloor temperature, subject to use of an appropriate equation relating shell  $\delta^{18}\text{O}$  to temperature and water  $\delta^{18}\text{O}$ , reasonably accurate estimation of water  $\delta^{18}\text{O}$ , and due consideration of growth-rate effects. Taking these factors into account,  $\delta^{18}\text{O}$  data from late Pliocene bivalves of the southern North Sea Basin (Belgium and the Netherlands) indicate a seasonal seafloor range ~~at times larger~~ ~~a little smaller~~ -than now in the area. Microgrowth-increment data from *Aequipecten opercularis*, together with the species-composition of the bivalve assemblage and aspects of preservation, suggest a setting below the summer thermocline for all but the latest material investigated. This implies a higher summer temperature at the surface than on the seafloor and consequently a greater seasonal range. A ~~conservative-reasonable~~ (3 °C) estimate of the difference between maximum seafloor and surface temperature under circumstances of summer stratification points to seasonal surface ranges in excess of the present value (12.4 °C nearby). Using ~~a model-constrained estimates~~ ~~derived estimate~~ of water  $\delta^{18}\text{O}$  (0.0 ‰), summer surface temperature was initially in the cool temperate range (< 20 °C) and then (during the Mid-Piacenzian Warm Period; MPWP) increased into the warm temperate range (> 20 °C) before reverting to cool temperate values (in conjunction with shallowing and a loss of summer stratification). This pattern is in agreement with biotic-assemblage evidence. Winter temperature was firmly in the cool temperate range (< 10 °C) throughout, contrary to previous interpretations. Averaging of summer and winter surface temperatures for the MPWP provides a figure for ~~mean~~-annual sea

35 -surface temperature that is 2–3 °C higher than the present value (10.9 °C nearby) and in close agreement with a figure obtained by averaging alkenone- and TEX<sub>86</sub>-temperatures for the MPWP from the Netherlands. These proxies, however, respectively underestimate summer temperature and overestimate winter temperature, giving an incomplete picture of seasonality. A higher ~~mean~~ annual temperature than now is consistent with the notion of global warmth in 40 the MPWP, but a low winter temperature in the southern North Sea Basin suggests regional reduction in oceanic heat supply, contrasting with other interpretations of North Atlantic oceanography during the interval. Carbonate clumped isotope ( $\Delta_{47}$ ) and biomineral unit thermometry offer means of checking the  $\delta^{18}\text{O}$ -based temperatures.

## 1 Introduction

45 The foraminiferal  $\delta^{18}\text{O}$  record from the deep sea indicates that the global volume of land ice was generally lower than now during the Pliocene Epoch (Lisiecki and Raymo, 2005), and that global mean surface temperature (GMST) was therefore generally higher. The late Pliocene saw the last mainly-warm interval before the change to the typically cooler-than-present conditions of the Pleistocene. During this interval, the Mid-Piacenzian Warm Period (MPWP; 50 3.28–3.03 Ma; Dowsett et al., 2019), ‘warm peak average’ temperature was 2–3°C higher than now, similar to the GMST predicted for the end of the present century (Dowsett et al., 2013). As evident under current global warming, the mid-Piacenzian temperature anomaly was not uniform, being for instance greater than the global average figure at mid-latitudes in the oceans according to results from both proxies and modelling (Lunt et al., 2010). Despite general 55 agreement, strong discrepancies between proxy and model estimates of mean annual sea\_- surface temperature (ASST) (~~MASST~~) have been identified in some regions (Dowsett et al., 2011). Those formerly recognized in the northern North Atlantic Ocean have been reduced by limiting proxy estimates to one source—alkenone index—and adjusting model boundary conditions (Dowsett et al., 2019). It is, however, widely considered (e.g. Robinson, 2009; Bova et al., 2021) that alkenone index reflects temperature in the warmer part of the year, and the same is now thought to be generally the case for another commonly utilized geochemical proxy, the Mg/Ca ratio of foraminiferal calcite (Bova et al., 2021). The species-composition of assemblages of pelagic micro-organisms (particularly Foraminifera) has been extensively used to derive both summer and winter sea\_-surface temperatures for the Pliocene (e.g. Dowsett et 60 al., 2010). The methodology, employing information on the seasonal temperatures associated with modern representatives and relatives, assumes constancy of niche (‘ecological

uniformitarianism’; Vignols et al., 2019) and, furthermore, that both summer and winter temperatures exert an influence on modern occurrence. This is questionable for the many forms that ‘bloom’ in summer, and those (dinoflagellates) that survive winter as cysts (dinocysts).

70

It would be possible to obtain a more accurate estimate of regional mean MASST for comparison with model outputs by combining temperatures from a summer proxy with those from a winter proxy, if such existed. Dearing Crampton-Flood et al. (2020) obtained TEX<sub>86</sub> estimates about 6 °C lower than from alkenones for sea temperature during the MPWP in the Netherlands. They took the former data to reflect conditions during winter, when the source-organisms (archaea) of the lipids concerned may have bloomed. Given that alkenones (produced by haptophyte algae) seem to reflect summer conditions, the mid-point between the two figures is probably close to mean MASST. However, we cannot be sure in the absence of information on the precise times during winter and summer that are represented, and for the same reason we cannot say whether the figures give an accurate picture of seasonality. In this paper we use sclerochronology (investigation of the chemical and physical nature of accretionary mineralized skeletons) to obtain estimates of extreme summer and winter sea temperatures in individual years over a late Pliocene interval spanning the MPWP in Belgium and the Netherlands. The information substantially supplements initial sclerochronological estimates (Valentine et al., 2011) from these countries on the eastern side of the southern North Sea Basin (SNSB), and complements a sizeable body of equivalent data relating to the early Pliocene (Zanclean) sequence of eastern England, on the western side of the SNSB (Johnson et al., 2009, 2021b; Vignols et al., 2019). The results serve to: (1) test estimates of seasonality and annual (average) temperature obtained from other proxies; (2) expand and refine the proxy evidence of temperature available for testing models of Pliocene climate; (3) provide an insight into the controls on regional marine climate.

80

85

90

## 2 Sclerochronology and seasonality

The majority of sclerochronological studies of environment have been conducted on accretionary calcium carbonate skeletons, principally those of corals and molluscs in the marine realm. Trace element (Sr/Ca) profiles from shallow-water corals have been found to mirror seasonal changes in surface temperature (e.g. DeLong et al., 2007, 2011) but no such close relationship exists in molluscs (e.g. Gillikin et al., 2005; Markulin et al., 2019). In view of the absence of corals (at least long-lived, colonial forms) from extra-tropical shallow-water

95

environments and general inutility of trace (and minor) element data from molluscs for  
100 reconstructing seasonal temperature variation, sclerochronological investigations of  
palaeoseasonality in temperate and polar settings have been largely based on the  $\delta^{18}\text{O}$  of  
molluscan carbonate. Pelagic belemnites supplement benthic molluscs as a provider of  
information on Jurassic and Cretaceous conditions (e.g. Mettam et al., 2014) but after the  
extinction of the former at the end of the Cretaceous the latter become the sole source of data  
105 (e.g. Bice et al., 1996; Williams et al., 2010; Surge and Barrett, 2012; Johnson et al., 2009,  
2017, 2019; Vignols et al., 2019; de Winter et al., 2020a, b). There is no doubt that temperature  
exerts an influence on the  $\delta^{18}\text{O}$  of molluscan carbonate, but values are also affected by the  $\delta^{18}\text{O}$   
of the fluid from which the material was precipitated (usually taken to be equivalent to that of  
ambient water) and by kinetic and more obscure ‘vital’ effects (e.g. Owen et al., 2002a, b;  
110 Fenger et al., 2007; Garcia-March et al., 2011). At present it is possible only to constrain (not  
specify) the  $\delta^{18}\text{O}$  of ambient water in ancient settings so, although precise, the temperatures  
from  $\delta^{18}\text{O}$  thermometry are not necessarily accurate—i.e. they are questionable as absolute  
temperatures. Nevertheless, assuming that kinetic and vital effects do not vary with season or  
age, an assumption which is certainly valid for some molluscs (e.g. Fenger et al., 2007; Garcia-  
115 March et al., 2011), and that water  $\delta^{18}\text{O}$  is constant, ontogenetic profiles are, at least in  
principle, a true reflection of relative temperature and hence (from the difference between  
summer and winter  $\delta^{18}\text{O}$  values) of seasonality.


Unfortunately, molluscan growth is often discontinuous, and interruptions are frequently  
120 associated with seasonal temperature extremes (Schöne, 2008), so in such cases the shell  $\delta^{18}\text{O}$   
record does not fully reflect the range of temperatures experienced (e.g. Hickson et al., 2000;  
Peharda et al., 2019a). However, because of their typical manifestation as ‘growth lines’,  
interruptions can be recognized and instances of likely truncation of the  $\delta^{18}\text{O}$  record inferred  
(e.g. Johnson et al., 2017, 2019, 2021b). The increasing occurrence and/or duration of growth  
125 interruptions with age is part of the reason for the commonly observed reduction in amplitude  
of  $\delta^{18}\text{O}$  profiles through ontogeny, but general slowing of growth (and consequent greater time-  
averaging within samples) is also contributory (Goodwin et al., 2003; Ivany et al., 2003).  
Increasing sample resolution can potentially offset this effect (Schöne, 2008) but the most  
accurate indication of seasonal temperature variation is likely to be obtained from early  
130 ontogenetic data (Goodwin et al. 2003; Ivany et al., 2003). An exception to this rule is provided  
by *Magallana* (formerly *Crassostrea*) *gigas*, which exhibits substantial oxygen isotope  
disequilibrium in early ontogeny (Huyghe et al., 2020, 2022).

A problem as important as growth cessation/reduction for inference of seasonal temperature range is the choice of equation relating temperature to water and shell  $\delta^{18}\text{O}$ . Various equations exist for both aragonite and calcite, and for the same range in shell  $\delta^{18}\text{O}$  these yield different temperature ranges. Thus for a water value of 0.0 ‰ and summer and winter shell values of 0.0 ‰ and +2.0 ‰, respectively, the widely employed aragonite equation of Grossman and Ku (1986) yields seasonal temperatures of 19.4 °C and 10.7 °C—i.e. a seasonal range of 8.7°C. However, for the same  $\delta^{18}\text{O}$  values the *Glycymeris glycymeris*-specific aragonite equation of Royer et al. (2013) yields seasonal temperatures of 17.4 °C and 12.1 °C—i.e. a seasonal range of only 5.3 °C. Since both equations are linear, like the LL (low-light) calcite equation of Bemis et al. (1998), used by Johnson et al. (2021b), neither the absolute values specifying a summer-winter difference of 2.0 ‰ in shell  $\delta^{18}\text{O}$ , nor the value of water  $\delta^{18}\text{O}$ , affect the calculated seasonal temperature range. However, the non-linear calcite equations of O'Neil et al. (1969; as reformulated by Shackleton et al., 1974) and Kim and O'Neil (1997) not only yield different temperature ranges for a given range in shell  $\delta^{18}\text{O}$ , but the temperature range in each case varies with the absolute shell values concerned, and with water  $\delta^{18}\text{O}$ . Thus for a water value of 0.0 ‰ and summer and winter shell values of 0.0 ‰ and + 2.0‰, respectively, the calcite equation of O'Neil et al. (1969) yields a seasonal temperature range of 8.2 °C (summer 15.7 °C, winter 7.5 °C) and that of Kim and O'Neil (1997) a seasonal temperature range of 8.9 °C (summer 13.7°C, winter 4.8 °C), but for a water value of +0.4 ‰ and summer and winter shell values of +1.5 ‰ and +3.5 ‰ (i.e. the same 2.0 ‰ range but at higher absolute values), the equation of O'Neil et al. (1969) yields a seasonal temperature range of 7.8 °C (summer 11.1 °C, winter 3.3 °C) and that of Kim and O'Neil (1997) a seasonal temperature range of 8.6 °C (summer 8.8 °C, winter 0.2 °C). The differences in seasonal temperature range due to different water  $\delta^{18}\text{O}$  and absolute shell  $\delta^{18}\text{O}$  values are not great but the differences due to different equations are fairly significant for calcite (up to almost 1 °C for the water and shell  $\delta^{18}\text{O}$  values specified above) and quite major for aragonite (over 3 °C). Clearly, therefore, the choice of equation must be given careful consideration.

Modelling (e.g. Williams et al., 2009) and carbonate clumped isotope ( $\Delta_{47}$ ) analysis (e.g. Briard et al. 2020; Caldarescu et al., 2021) are techniques that have been used to constrain water  $\delta^{18}\text{O}$ . The studies cited in relation to the latter approach employ it to resolve seasonal fluctuations, and de Winter et al. (2021) discuss the best sampling strategy to achieve this end. In nearshore settings affected by major seasonal influxes of freshwater (normally isotopically light), and

which exhibit concomitant reductions in salinity, variation in water  $\delta^{18}\text{O}$  may be quite high. Lloyd (1964) documented change of more than 1 ‰ over a few months in part of Florida Bay and Ivany et al. (2004) inferred seasonal variation of 2.5 ‰ in an Eocene nearshore setting in the south-eastern USA. However, in more offshore settings the effects of freshwater influx are much less. Thus in the modern North Sea seasonal variation in salinity is in most places only 0.25 PSU (Howarth et al., 1993), which translates to a seasonal variation in water  $\delta^{18}\text{O}$  of just 0.07 ‰ using the salinity–water  $\delta^{18}\text{O}$  relationship for the North Sea of Harwood et al. (2008). Within a few tens of kilometres of the mouth of the Rhine seasonal variation in salinity rises to 0.75 PSU and hence calculated variation in water  $\delta^{18}\text{O}$  to 0.21 ‰. At 20–30 m depth in the eastern part of the central North Sea, Schöne and Fiebig (2009) identified variation in salinity of up to 2 PSU in certain years, which translates to a variation in water  $\delta^{18}\text{O}$  of 0.55 ‰. If minimum and maximum water  $\delta^{18}\text{O}$  values differing by this amount coincided respectively with the times of maximum and minimum water temperature it would increase the temperature range calculated from shell  $\delta^{18}\text{O}$  assuming constant water  $\delta^{18}\text{O}$  by an amount in the order of 2.6 °C (figure for calcite using the LL equation mentioned above). However, the data of Schöne and Fiebig (2009) provide no evidence of a negative correlation between salinity/water  $\delta^{18}\text{O}$  and temperature, and near the eastern shore of the central North Sea there is a very strong positive correlation between water  $\delta^{18}\text{O}$  and temperature over the seasonal cycle (Ullmann et al., 2011; de Winter et al., 2021). This presumably reflects relatively high evaporation in summer, combined with relatively low freshwater input, a common pattern in mid-latitude settings and one suggesting that seasonal variation in shell  $\delta^{18}\text{O}$  of marine organisms at mid-latitudes is more likely to be damped than enhanced by variation in water  $\delta^{18}\text{O}$ .

Even if there were a negative correlation between water  $\delta^{18}\text{O}$  and temperature it would be confined to nearshore waters (more susceptible to freshwater influx), hence the effect on calculated temperature range could be mitigated by use of offshore shells. This approach introduces the possibility of underestimation of the surface range as a result of life positions below the summer thermocline (typically at 25–30 m depth in shelf settings). However, shells from sub-thermocline settings may be recognized from the associated sediments and biota, and, in the case of the scallop *Aequipecten opercularis*, from microgrowth-increment patterns (Johnson et al., 2009, 2021b; Fig. 1).




200 **Figure 1:** Pictorial demonstration of microgrowth-increment patterns in left valves of *Aequipeecten*  
*opercularis*. (a) Typical supra-thermocline specimen (mesotidal setting, 23 m depth, La Coruña,  
 Galicia, Spain) showing small increments early and late in ontogeny. (b, c) Typical sub-thermocline  
 specimens (b: microtidal setting, 50 m depth, Gulf of Tunis, Tunisia; c: microtidal setting, 38 m depth,  
 205 Adriatic Sea, Pula, Croatia) showing large increments early in ontogeny. (d) Inferred sub-thermocline  
 specimen (Ramsholt Member, Coralline Crag Formation, Broom Pit, Suffolk, UK) showing large  
 increments early in ontogeny and a transition from large to small increments late in ontogeny. Scale  
 bars for whole-shell images (upper) and enlargements (lower) = 10 mm. Major growth breaks (gb)  
 identified in enlargements of (b, d). (a) = University of Derby, Geological Collections (UD) 53424; (b)  
 = National Museum of Natural History, Paris, IM-2008-1542 (one of seven specimens in this lot); (c)  
 210 = UD 53423 (one of 48 specimens in this lot, coded S3A29); (d) = UD 53425. See Johnson et al. (2009,  
 2021b) for numerical data and discussion of microgrowth-increment patterns in *A. opercularis*. Modern  
 supra-thermocline specimens show a difference of < 0.3 mm between the maximum and minimum  
 values of smoothed increment-height profiles, while the majority of sub-thermocline specimens show  
 a difference of > 0.3 mm.

215 While  $\delta^{18}\text{O}$  sclerochronology is potentially informative about seasonality, it should be clear  
 from the foregoing that results from the technique need to be interpreted carefully. The  
 reliability of the information is of course also dependent on preservation of the original shell  
 $\delta^{18}\text{O}$  signature. [Other issues are considered by Ivany and Judd \(2022\).](#)

### 3 Setting and material

220 In the Pliocene the marine area of the SNSB was somewhat greater than now (Fig. 2), partly  
 due to higher global sea-level and partly to subsequent regional uplift (Westaway et al., 2001).  
 To the west, onshore marine deposits exist in eastern England, and to the east in Belgium and  
 the Netherlands, those in the last two countries passing eastwards into essentially fluvial non-  
 marine deposits of the proto-Rhine/Meuse/Scheldt river system (Louwye et al., 2020;  
 225 Munsterman et al., 2020). The Eridanos river system, draining the Baltic area, had its exit into

the SNSB in the area of the present German Bight, some 400 km north-east of the proto-Rhine/Meuse/Scheldt exit (Gibbard and Lewin, 2016). While at certain times a link may have existed between the SNSB and the Channel Basin during the Pliocene (either at the present position or across southern England; Funnell, 1996; Westaway et al., 2002; van Vliet-Lanoë et al., 2002; Gibbard and Lewin, 2016), at others the basins were separated by the Weald–Artois land-bridge, as shown in Fig. 2. Water depth in the southern North Sea is now less than 40 m in most places but seismic stratigraphy indicates that it was greater in the Pliocene, at least in areas of low sediment accumulation (Overeem et al., 2001).



235 **Figure 2:** Pliocene palaeogeography in the vicinity of the SNSB, the location of sites in the Lillo (1) and Oosterhout (2) formations from which shells were obtained, and the area of onshore outcrop of the Coralline Crag Formation in eastern England (the partly Pleistocene Red Crag Formation occurs over a larger area). Adapted from Valentine et al. (2011, fig. 1), itself based on Murray (1992, map NG1).


In eastern England there is a large stratigraphic gap between the Zanclean Coralline Crag Formation and the late Piacenzian basal unit ('Walton Crag') of the Red Crag Formation, but 240 the Piacenzian is better recorded in northern Belgium by the Lillo Formation and in the south-



west Netherlands by the Oosterhout Formation (Fig. 3). The last two formations essentially comprise marine sands, the Oosterhout Formation at Ouwerkerk (Zeeland) probably deposited in deeper water than the Lillo Formation at Antwerp from the evidence of fish otoliths (Gaemers and Schwarzahns, 1973) and a position farther from the inferred shoreline (Fig. 2).  
245 Nevertheless, Slupik et al. (2007) inferred a depth of deposition above storm wavebase for ~~the~~ most of the Oosterhout Formation at Schelphoek, 15 km north-west of Ouwerkerk. In the Antwerp area depth estimates based on the fauna have varied between authors according to the group studied, most of them hardly taking into account the marked variation in sediment and sedimentary structures within members of the Lillo Formation (see Deckers et al., 2020, figs. 4–6). According to Gaemers (1975), the otolith assemblage indicates a depth of at least 10–20 m for the ‘Kallo Sands’ (= Lillo Formation, Oorderen Member; Marquet and Herman, 2009) but less than 10 m for the overlying Kruisschans Member. This indication of upward shallowing is supported by assemblage evidence from dinocysts (Louwye et al., 2004; De Schepper et al., 2009), Foraminifera (Laga, 1972) and bivalves (Marquet, 2004), but statistical data from the last group suggest greater absolute depths: 35–45 m for the Oorderen Member and 15–55 m for the Kruisschans Member by the common overlap in depth-range of extant species; 40–50 m for the former and 20–50 m for the latter by the medial depth of extant species. The articulated preservation of the semi-infaunal bivalve *Atrina fragilis*, locally in life position,  
260 within the Oorderen Member (Marquet and Herman, 2009) is difficult to reconcile with the 10–20 m minimum depth estimate of Gaemers (1975), since specimens would have been subject to fair-weather processes after death. It is more likely that they lived at the depth suggested by Marquet (2004) and were killed by rapid burial (and permanently interred) in storms. A somewhat greater depth still was inferred from the bivalve assemblage of the underlying Luchtbal Member: 40–50 m by ‘common overlap’; 40–60 m by ‘medial depth’. The low diversity of the bivalve fauna of the Merksem Member (overlying the Kruisschans Member) precluded the same statistical treatment but Marquet and Herman (2009) inferred from this impoverishment a depth of less than 15 m, an estimate consistent with the foraminiferal assemblage (Laga, 1972) and the high proportion of terrestrial palynomorphs (De Schepper et al., 2009).  
270

Dinocyst assemblages indicate surface temperatures within the warm temperate range (but possibly only with respect to summer; Sect. 1) during deposition of most of the Oorderen Member, but punctuated by cool intervals and preceded by continuously cool conditions during deposition of the Luchtbal Member (De Schepper et al., 2009). The dinocysts of the

280 Kruisschans and Merksem members mainly indicate a continuation of the warm conditions of the Oorderen Member but provide a few hints of cooling (Louwye et al. 2004; De Schepper et al., 2009). Other evidence of this is provided by bivalves, fish and pollen (Hacquaert, 1961; Vandenberghe et al., 2000; Marquet, 2005), and Wood et al. (1993) determined a 5–6 °C decrease in summer surface temperature from the ostracods of a contemporaneous part of the Oosterhout Formation.



295 **Figure 3:** Stratigraphy and correlation of marine mid–late Pliocene and early Pleistocene units of the southern North Sea Basin, with the general stratigraphic positions of shells sampled for the present study (specific positions in Table 1). Age (Ma) of the Red Crag Formation and constituent members (including the unofficial Walton Crag unit) according to Wood et al. (2009); of the Coralline Crag, Kattendijk and Lillo formations and constituent members according to De Schepper et al. (2009) and Louwye and De Schepper (2010); and of the Oosterhout (O.F.) and Maassluis formations according to Dearing Crampton-Flood et al. (2020) and Wesselingh et al. (2020). An additional small hiatus, of uncertain age, is present in the lower part of the Oosterhout Formation (Dearing Crampton-Flood et al., 2020). The Maassluis Formation includes a number of non-marine horizons (Slupik et al., 2007). Names of Lillo Formation members are in accordance with recent practice (Louwye et al., 2020; Wesselingh et al., 2020), omitting ‘Sands’/’Sands’, as included by previous authors. Wesselingh et al. (2020) found evidence of an additional layer (Broechem Unit) between the Kattendijk Formation and Luchtbal Member of the Lillo Formation. De Meuter and Laga (1976) designated an additional, uppermost division of the latter formation (Zandvliet Member), but this may be no more than the decalcified top of the Merksem Member (Louwye et al., 2020). Geographic provenance of shells and the location of the Coralline Crag Formation shown in Fig. 2. MPWP = Mid-Piacenzian Warm Period.

310

315

**Table 1:** Basic information for the investigated specimens (all single valves). Order stratigraphic within formations (by member, then by borehole-depth or bed, with entries for the Oosterhout Formation inserted immediately above those, if any, for the equivalent member in the Lillo Formation). Entries in square brackets are interpretations (see footnotes). Latitudes and longitudes are for the location indicated in the adjacent column and do not necessarily specify the exact place of collection (see Fig. 2 for the positions of Ouwerkerk and Antwerp). UD = University of Derby, Geological Collections; IRSNB = Royal Belgian Institute of Natural Sciences, Brussels (MSNB was used for specimens discussed in Valentine et al., 2011).


Formation	Member or equivalent (in quotes)	Borehole-depth (b-d) or bed	Location	Latitude, longitude	Genus and species	Repository and number	Code herein	Valve height (mm)	General physical condition	Mineralogy of sampled layer	Number of isotope samples
Oosterhout	'Merksem'	b-d: 89.75–91 m	Ouwerkerk	51.626° N, 3.983° E	<i>Aequipecten opercularis</i>	UD 53362	AO10	56	Incomplete	Calcite	42
Oosterhout	'Merksem'	b-d: 93.5–94.5 m	Ouwerkerk	51.626° N, 3.983° E	<i>Aequipecten opercularis</i>	UD 53363	AO9	46	Incomplete	Calcite	30
Oosterhout	'Oorderen'	b-d: 98.5–99.5 m	Ouwerkerk	51.626° N, 3.983° E	<i>Aequipecten opercularis</i>	UD 53347	AO8	34	Incomplete, abraded	Calcite	31
Lillo	Oorderen	<i>Atrina fragilis</i> bed	Vrasenedok, Kallo, Antwerp	51.263° N, 4.238° E	<i>Aequipecten opercularis</i>	IRSNB Invert-29710-10	AO7	42	Complete	Calcite	31
Lillo	Oorderen	<i>Atrina fragilis</i> bed	Vrasenedok, Kallo, Antwerp	51.263° N, 4.238° E	<i>Aequipecten opercularis</i>	IRSNB Invert-29710-09	AO6	51	Complete	Calcite	39
Lillo	Oorderen	Base <i>Atrina fragilis</i> bed	Deurganckdok, Doel, Antwerp	51.291° N, 4.257° E	<i>Aequipecten opercularis</i>	IRSNB Invert-D2-8	AO5	31	Complete	Calcite	23
Lillo	Oorderen	<i>Atrina fragilis</i> bed	Vrasenedok, Kallo, Antwerp	51.263° N, 4.238° E	<i>Pygocardia rustica</i>	IRSNB Invert-29710-04	PR	62	Complete	Aragonite	37
Lillo	Oorderen	<i>Atrina fragilis</i> bed	[Antwerp] <sup>a</sup>	51.217° N, 4.421° E	<i>Arctica islandica</i>	IRSNB Invert-18201-01	AI	64	Complete	Aragonite	32
Oosterhout	'Luchtbal'	b-d: 106–107.5 m	Ouwerkerk	51.626° N, 3.983° E	<i>Aequipecten opercularis</i>	UD 53364	AO4	44	Incomplete, abraded	Calcite	36
Lillo	Luchtbal	<i>Palliohum gerardi</i> bed	Deurganckdok, Doel, Antwerp	51.291° N, 4.257° E	<i>Aequipecten opercularis</i>	IRSNB Invert-29710-13	AO3	42	Complete	Calcite	28
Lillo	Luchtbal	<i>Palliohum gerardi</i> bed	Deurganckdok, Doel, Antwerp	51.291° N, 4.257° E	<i>Aequipecten opercularis</i>	IRSNB Invert-29710-12	AO2	47	Complete	Calcite	30
Lillo	Luchtbal	<i>Palliohum gerardi</i> bed	Deurganckdok, Doel, Antwerp	51.291° N, 4.257° E	<i>Aequipecten opercularis</i>	IRSNB Invert-29710-11	AO1	54	Complete	Calcite	28
Lillo	Luchtbal	[lower bed] <sup>b</sup>	Deurganckdok, Doel, Antwerp	51.291° N, 4.257° E	<i>Glycymeris radiolyrata</i>	IRSNB 7698	GR2	77	Broken in storage	Aragonite	74
Lillo	Luchtbal	[lower bed] <sup>b</sup>	Deurganckdok, Doel, Antwerp	51.291° N, 4.257° E	<i>Glycymeris radiolyrata</i>	IRSNB Invert-29710-0062	GR1	92	Broken in storage	Aragonite	42

- 320 a no specific location indicated within Belgium, but highly likely to be Antwerp  
 320 b species indicated as 'special' to the lower bed of the Luchtbal Member in the Deurganckdok (Marquet, 2002)

Previous sclerochronological investigation of late Pliocene temperatures in Belgium and the Netherlands focussed on the Oorderen Member and an equivalent horizon in the Oosterhout Formation, and was restricted to  $\delta^{18}\text{O}$  data from two bivalve species, *Aequipecten opercularis* and *Atrina fragilis* (Valentine et al., 2011). Here we supplement the existing  $\delta^{18}\text{O}$  data from *A. opercularis* with microgrowth-increment data from the same specimens to gain an insight into their hydrographic setting (sub- or supra-thermocline) and also supply  $\delta^{18}\text{O}$  data from two further bivalve species (*Arctica islandica* and *Pygocardia rustica*) from the Oorderen Member, and another (*Glycymeris radiolyrata*) from the Luchtbal Member. In addition, we provide *A. opercularis* data from the Luchtbal Member and horizons equivalent to the Luchtbal and Merksem members in the Oosterhout Formation. Values for  $\delta^{13}\text{C}$  (obtained alongside  $\delta^{18}\text{O}$ ) are reported for all species. Details of the provenance of the specimens are given in Table 1, together with alphanumeric codes (AO = *A. opercularis*; AI = *A. islandica*; PR = *P. rustica*; GR = *G. radiolyrata*) and sundry basic descriptive information. Note that all the five specimens

335

from the Oorderen Member *sensu stricto* come from the *Atrina fragilis* bed, a horizon with the ~~warm~~ warm temperate dinoflagellate-biotadinocyst assemblage found at most levels in the member. Illustrations of species other than *A. opercularis* (Fig. 1) are provided in Fig. 4. Most, if not all, of the material from the Lillo Formation was obtained from temporary exposures created during harbour works in the Antwerp area, while all the material from the Oosterhout Formation was obtained from a borehole (Rijkswaterstaat-Deltadienst, afdeling Waterhuishouding, 42H19-4/42H0039) at Ouwerkerk, Zeeland. Interpretation of positions (depths) within the Ouwerkerk borehole in terms of members within the Lillo Formation follows Gaemers and Schwarzans (1973) except in the case of AO8, for which we have accepted the opinion of F. Wesselingh (in Valentine et al., 2011) that the position is equivalent to the Oorderen Member. Gaemers and Schwarzans (1973) considered that strata of this age ('Kallo Sands') were missing at Ouwerkerk but they appear to be well represented at Schelphoek, only 15 km away (Slupik et al., 2007).



**Figure 4:** Right valves of (a) *Arctica islandica*, (b) *Pygocardia rustica* and (c) *Glycymeris radiolyrata* from the Lillo Formation, Antwerp. (a): probably Oorderen Member, Verrebroekdok (IRSNB 7699); (b): Oorderen Member, Verrebroekdok (IRSNB 7700); (c): Luchtbal Member, Deurganckdok (IRSNB 7701). Growth lines/breaks are evident in all three specimens—e.g. c. 10 major growth breaks in (b). Scale bar = 10 mm.

According to the latest chronostratigraphy (Fig. 3), the material investigated is largely or entirely Piacenzian (3.60–2.59 Ma) in age, the oldest (from the Luchtbal Member of the Lillo Formation) being possibly as old as 3.71 Ma (latest Zanclean) and the youngest (from horizons in the Oosterhout Formation equivalent to the Merksem Member of the Lillo Formation) being no younger than 2.76 Ma (De Schepper et al., 2009). The MPWP is probably represented by material from the Oorderen Member and the equivalent level in the Oosterhout Formation (Valentine et al., 2011). The Luchtbal and Oorderen members are separated by an unconformity interpreted by De Schepper et al. (2009) as a product of the sea-level fall associated with

Marine Isotope Stage (MIS) M2 (c. 3.3 Ma), which marks a glacial episode. The Luchtbal Member was therefore probably deposited before MIS M2 under interglacial conditions.

All the specimens come from stratigraphic intervals with a fully marine associated biota (e.g. Marquet, 2002, 2005; Gaemers and Schwarzahns, 1973), in conformity with modern occurrences in the case of the extant species *A. opercularis* and *A. islandica* (Tebble, 1976) and other fossil occurrences in the case of the extinct species *P. rustica* and *G. radiolyrata* (Norton, 1975; Buchardt and Simonarson, 2003; Marquet, 2002, 2005). Investigation of specimens of *A. islandica*, *P. rustica* and *G. radiolyrata* added information from infaunal, slow-growing taxa to that derived from fast-growing, epifaunal *A. opercularis*, hence serving to mitigate any 'ecological' bias in the results. We could not sample as many specimens of the infaunal, slow-growing species as of *A. opercularis* due to the limited availability of material (perforce from museums, in the lack of extant stratal exposures in the area of study). However, we sampled multiple years in the infaunal, slow-growing species so the combined number of seasonal cycles investigated was similar to that in *A. opercularis*. We nevertheless expected some imbalance in the data because modern examples of *Glycymeris* species, from both cool- and warm-temperate settings, show winter cessation or slowing of growth and thus supply (or would supply) underestimates of the seasonal temperature range from  $\delta^{18}\text{O}$  sclerochronology (Peharda et al., 2012, 2019a, b; Royer et al., 2013; Reynolds et al., 2017; Featherstone et al., 2020; Alexandroff et al., 2021). Various equations have been used to express the precise relationship between  $\delta^{18}\text{O}$  and temperature in modern *Glycymeris* (Royer et al., 2013; Peharda et al., 2019a, b) but species of this genus certainly exhibit something at least close to equilibrium isotopic incorporation. The same is true of *A. opercularis* (Hickson et al., 1999; Johnson et al., 2021b) and *A. islandica* (Schöne, 2013; Mette et al., 2018; Trofimova et al., 2018). Because of the similarity of  $\delta^{18}\text{O}$  values from seemingly well-preserved *P. rustica* from the Pliocene of Iceland to those from co-occurring, similarly preserved *A. islandica* (Buchardt and Simonarson, 2003) it is reasonable to assume equilibrium fractionation in the former (extinct) species. The specimens analysed showed no ~~obvious-physical~~ signs of alteration and they are ~~unlikely to have suffered significant~~ likely to have been heated by more than 10 °C heating through burial as the thickness of overlying sediments was probably never much more than 100 m (the depth below the present surface of the lowermost shell from the Ouwerkerk borehole). Examples of both calcitic *A. opercularis* (including AO7 herein) and aragonitic *A. fragilis* from the Lillo Formation were shown by Valentine et al. (2011) to exhibit the original shell microstructure. Similarly (Valentine et al., 2011), as do examples of a good preservation

400 has been demonstrated in a variety of calcitic and aragonitic species from the slightly earlier  
Ramsholt Member of the Coralline Crag Formation in eastern England (Johnson et al., 2009;  
Vignols et al., 2019). We therefore considered it reasonable to proceed with isotopic analysis  
of our material (both calcitic and aragonitic; see Sect. 4 and Table 1) without detailed  
investigation of its preservation. Moon et al. (2021) have recently shown that good  
mineralogical and microstructural preservation does not necessarily guarantee good  
405 preservation of original shell  $\delta^{18}\text{O}$ . They heated shell material to 200 °C and identified  
consistent negative shifts in  $\delta^{18}\text{O}$  (1.5 ‰ after two weeks at this temperature) over an annual  
cycle, but no significant mineralogical or microstructural changes. As our specimens  
experienced only minimal heating through burial, similar alteration of shell  $\delta^{18}\text{O}$  is unlikely.


## 4 Methods

### 4.1 Laboratory procedures

410 The exterior of *A. opercularis* shells was coated with a sublimate of ammonium chloride and  
digitally photographed for the purpose of measuring microgrowth increments and the position  
of growth breaks. The coating was washed off with tap-water and the shells then underwent  
the further cleaning procedure adopted by Valentine et al. (2011) for removal of any surficial  
organic matter, in preparation for isotopic sampling of the outer shell layer from the exterior,  
415 as in other such investigations of *A. opercularis* (e.g. Hickson et al., 1999, 2000; Johnson et  
al., 2009, 2021b; Vignols et al., 2019). The infaunal species were sampled in cross-section  
along the line of maximum growth, in accordance with universal practice for *A. islandica* (e.g.  
Schöne et al., 2005) and common practice for *Glycymeris* species (e.g. Royer et al., 2013). For  
this purpose shells were stabilized in resin before sectioning—by the partial-encasement  
420 method of Schöne et al. (2005) for *A. islandica* and *P. rustica*, and the total-encasement method  
of Johnson et al. (2021a) for *G. radiolyrata* (fragments bonded beforehand). Use of vacuum  
impregnation in the latter method resulted in resin penetration into the outer part of the outer  
shell layer.

425 Extraction of isotope samples from *A. opercularis* shells was by drilling a dorsal to ventral  
series of shallow commarginal grooves (depth and width < 1 mm; cf. Hickson et al., 1999, fig.  
2; 2000, fig. 3) in the external surface, with the sample sites more closely spaced towards the  
ventral margin in an attempt to maintain temporal resolution in the face of declining growth  
rate with age. Details of the procedure are given in Johnson et al. (2019) with respect to another

430 scallop species. Mean sample spacing for individuals—the average distance between the  
centres of grooves along the dorso-ventral (= maximum-growth/shell-height) axis—was 0.93  
(AO8)–1.35 (AO9) mm. Sampling of the infaunal species was by drilling a series of holes  
(depth and width < 1 mm; Fig. 5) in the outer shell layer as seen in cross-section, the curved  
path being located about midway between the external surface and the boundary between the  
435 outer and inner shell layers in *A. islandica* and *P. rustica*, but somewhat closer to the latter  
boundary in *G. radiolyrata* to avoid resin-contaminated material (Fig. 5). Sample spacing was  
more constant than for *A. opercularis*, although significantly reduced late in the long series  
from *G. radiolyrata* GR2, again to maintain temporal resolution. Mean sample spacing for  
440 individuals—the average distance between the centres of holes, measured in terms of the  
difference in straight-line distance from the origin of growth—was 0.69 mm for *A. islandica*,  
0.57 mm for *P. rustica*, and 0.54 (GR2) and 0.57 (GR1) mm for *G. radiolyrata*. Note that in  
these relatively convex species the straight-line distance from the origin of growth is not a  
measurement of shell height as normally defined (a distance from the umbo, which protrudes  
dorsal of the origin of growth in these forms; e.g. Fig. 5) and that the plane in which it was  
445 measured (along the line of maximum growth) arguably does not include the shell height axis  
in the prosogyrate species *A. islandica* and *P. rustica* (dependent on the point at the shell margin  
that is regarded as ventral). The lines of measurement and the values obtained are, however,  
regarded as ‘heights’ for all four species considered herein, for the sake of simplicity. The *A.*  
*opercularis* shells were relatively small (Table 1) and were sampled from near the origin of  
450 growth (dorsal margin) to a point at or close to the ventral margin (maximum sample height  
53.0 mm in AO10). The shells of the infaunal species were larger (Table 1) and not sampled to  
the end of ontogeny (maximum sample height 54.7 mm in GR2). Furthermore, the thinness of  
the outer layer close to the origin of growth meant that sampling had to start relatively far from  
this point (minimum sample height 15.4 mm in GR2).



455 **Figure 5:** Cross-section of *Glycymeris radiolyrata* specimen GR2 showing the origin of growth (og),  
position of sample holes (relatively far from the external surface in this species to avoid the darker,  
resin-contaminated material) and a major growth break (pale diagonal band in enlargement) at shell  
height 35.4 mm. Scale bars = 10 mm. Black spot in enlargement is a marker to assist sample numbering.

460 The cross-sections of the infaunal species were digitally photographed for the purpose of  
measuring the positions of sample holes and growth breaks, as seen on the shell exterior (cf.  
Fig. 4) and projected or traced (Fig. 5) into the isotope sample path. Distances from the origin  
of growth were determined from the images using the bespoke measuring software Panopea©  
(2004, Peinl and Schöne). Panopea was also used to measure the position of growth breaks and  
465 the height of microgrowth increments in the shell-exterior images of *A. opercularis* (cf. Fig.  
1). As in the case of isotope sample positions, measurements were made along the dorso-ventral  
axis or (where this was impossible due to abrasion or encrustation) lateral to this line, the  
measurements then being mathematically adjusted as described by Johnson et al. (2019) to  
correspond to ones made along the dorso-ventral axis. All the microgrowth-increment  
470 measurements were made by the same person (AMV), thus assuring a reasonably uniform  
approach given the subjective element in increment identification (Johnson et al., 2021b).  
Growth breaks were classified as major (incorporating ‘moderate’) or minor in all species  
dependent on their external prominence (cf. Figs. 1, 4).

475 Samples (typically 50–100 µg) were analysed for their stable oxygen and carbon isotope  
composition (given as  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$ ) at the stable isotope facility, British Geological Survey,  
Keyworth, UK (*A. opercularis*, *A. islandica*, *P. rustica*) and the Institute of Geosciences,  
University of Mainz, Germany (*G. radiolyrata*). At Keyworth, samples were analysed using  
an Isoprime dual inlet mass spectrometer coupled to a Multiprep system; powder samples were  
480 dissolved with concentrated phosphoric acid in borosilicate Wheaton vials at 90°C. At Mainz,  
samples were analysed using a Thermo Finnigan MAT 253 continuous flow–isotope ratio mass  
spectrometer coupled to a Gasbench II; powder samples were dissolved with water-free  
phosphoric acid in helium-flushed borosilicate exetainers at 72°C. Both laboratories  
calculated  $\delta^{13}\text{C}$ —and  $\delta^{18}\text{O}$  against VPDB and calibrated data against NBS-19 (preferred  
485 values: +1.95 ‰ for  $\delta^{13}\text{C}$ , –2.20 ‰ for  $\delta^{18}\text{O}$ ) and their own Carrara Marble standard  
(Keyworth: +2.00 ‰ for  $\delta^{13}\text{C}$ , –1.73 ‰ for  $\delta^{18}\text{O}$ ; Mainz: +2.01 ‰ for  $\delta^{13}\text{C}$ , –1.91 ‰  
for  $\delta^{18}\text{O}$ ). Values were consistently within  $\pm 0.05$  ‰ of the values for  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  in NBS-  
19. This confirms the comparability of results from each laboratory established in earlier work  
(Johnson et al., 2019). Note that  $\delta^{18}\text{O}$  of shell aragonite was not corrected for different acid-  
490 fractionation factors of aragonite and calcite (for further explanation see Füllenbach et al.,  
2015).


## 4.2 Calculation of temperatures



In previous work on late Pliocene bivalves from Belgium and the Netherlands, minimum and maximum estimates of global average seawater  $\delta^{18}\text{O}$  ( $-0.5\text{‰}$  and  $-0.2\text{‰}$ ), and minimum and maximum modelled values for the early Pliocene in the western part of the SNSB ( $+0.1\text{‰}$  and  $+0.5\text{‰}$ ), all adjusted downwards by  $0.1\text{‰}$  to allow for the input of isotopically light freshwater into the eastern SNSB, were used to calculate sets of temperatures from shell  $\delta^{18}\text{O}$  (Valentine et al., 2011). It seems appropriate to apply the adjusted modelled values more widely to late Pliocene material from Belgium and the Netherlands. ~~T, but the adjusted global values are~~  
495 ~~unreasonably low (they supply implausibly cold temperatures of  $0.1\text{ °C}$  and  $1.6\text{ °C}$ , respectively, from AO6, a specimen from a horizon with a warm temperate dinocyst assemblage) and are not used here.~~

~~— re probably unreasonably low and hence not worth applying here (Johnson et al.,~~  
Valentine et al. (2011) employed the calcite equation of O’Neil et al. (1969) for calculation of  
505 temperatures from *A. opercularis* but there are grounds for thinking that this provides slightly inaccurate figures (Hickson et al., 1999; Vignols et al. 2019). The LL calcite equation of Bemis et al. (1998) seems to provide more accurate figures (i.e. for modern shells, a better fit with directly measured temperatures) and certainly yields a larger estimate for seasonal range (Johnson et al., 2021b). Both equations have therefore been employed herein to generate  
510 ‘minimum’ and ‘maximum’ seasonal ranges from *A. opercularis*. Note that the calcite equation of Kim and O’Neil (1997) yields an intermediate estimate but the absolute temperatures obtained from modern *A. opercularis* are too low (Johnson et al., 2021b).

Just as there is some uncertainty as to the best equation for calculation of temperatures from *A.*  
515 *opercularis* calcite, so different equations have been favoured for use with aragonitic *Glycymeris glycymeris*. Royer et al. (2013) advocated use of a species-specific equation developed by them, while Reynolds et al. (2017) provide grounds for using the general aragonite equation of Grossman and Ku (1986). The former yields a smaller estimate of seasonal range than the latter so again both have been employed herein in relation to *G.*  
520 *radiolyrata*. The equation of Grossman and Ku (1986) is generally used in relation to aragonitic *A. islandica* and supplies similar temperatures from co-occurring (also aragonitic) *P. rustica* specimens (Buchardt and Simonarson, 2003). This, and no other equation, has therefore been used herein in relation to these species. In calculating temperatures appropriate adjustments were made to allow for the different scales used in measurement of water (VSMOW) and shell  
525 (VPDB)  $\delta^{18}\text{O}$  values (Coplen et al., 1983; Vignols et al., 2019).



**Figure 6:** Ontogenetic profiles of  $\delta^{18}\text{O}$ ,  $\delta^{13}\text{C}$  and microgrowth-increment height from Luchtbal Member (and equivalent) *A. opercularis* (a–d) and *G. radiolyrata* (e, f). Note that the isotopic axis has been reversed in each part such that lower values of  $\delta^{18}\text{O}$  (corresponding to higher temperatures) plot towards the top. While the axis range is 4‰ throughout, the minimum and maximum values for *A. opercularis* (calcitic; pale yellow background) have been set 0.5‰ lower than for *G. radiolyrata* (aragonitic; pale pink background) to facilitate comparison, given the different fractionation factors applying for  $\delta^{18}\text{O}$  (Kim et al., 2007). The criteria for recognition of reliable and unreliable summer and winter  $\delta^{18}\text{O}$  values are given in Sect. 6.1.1. The fairly large single-point  $\delta^{18}\text{O}$  excursion at height 18.5 mm in (d) is matched by a negative one in  $\delta^{13}\text{C}$  and probably reflects contamination. Smaller interruptions of the large-scale cyclical pattern of  $\delta^{18}\text{O}$  variation in this and other profiles can be regarded as represent ‘noise’ (unexplained variability), (like smaller excursions in other  $\delta^{18}\text{O}$  profiles) because it is associated with an aberrant  $\delta^{13}\text{C}$  value.

## 540 5 Basic results and analysis


The isotopic, microgrowth-increment and growth-break data are shown in Figs. 6 (Luchtbal Member and equivalent) and 7 (Oorderen Member and equivalent; Merksem-Member equivalent). Read top to bottom, left to right (i.e. in the alphabetical order of parts), the sequence in each figure is as in Table 1, read top to bottom. The raw data is available online  
545 (Johnson et al., 2021c).

### 5.1 $\delta^{18}\text{O}$ values and growth breaks

~~Disregarding~~ Apart from departures representing probable contamination or 'noise' (e.g. see Fig. 6e,d and caption), all profiles show cyclical patterns of  $\delta^{18}\text{O}$  variation, from less than half a cycle in *A. opercularis* profiles starting near the origin of growth and terminating at a height  
550 of about 30 mm (AO8, AO5—Fig. 7c, f, respectively) to between two and three in a profile terminating at 53 mm (AO10—Fig. 7a), but from between two and three cycles to substantially more over smaller height intervals in *G. radiolyrata* (between three and four from 25–49 mm in GR1—Fig. 6f; between eight and nine from 15–55 mm in GR2—Fig. 6e), and in *P. rustica* and *A. islandica* (between two and three from 27–48 mm in each case—Fig. 7g, h,  
555 respectively). In *A. opercularis* profiles extending beyond one  $\delta^{18}\text{O}$  cycle, the amplitude commonly shows a clear ontogenetic decrease. This pattern is less pervasive and pronounced amongst the other species, and the *A. islandica* specimen shows an ontogenetic increase in amplitude. However, the lack of early ontogenetic data for comparison from these species should be noted. The maximum amplitudes from *G. radiolyrata* specimens are less than from  
560 most *A. opercularis* specimens but those from *P. rustica* and *A. islandica* are similar to *A. opercularis*. Growth breaks, albeit sometimes only minor, are associated with (< 1 mm from the sample sites of) nearly all  $\delta^{18}\text{O}$  maxima and a few  $\delta^{18}\text{O}$  minima from *G. radiolyrata*, but with none of the maxima or minima from *A. opercularis*. Growth breaks are associated with two of the three maxima and two of the three minima from the *A. islandica* specimen, and with  
565 two of the three minima from the *P. rustica* specimen.

Taking the  $\delta^{18}\text{O}$  cycles to reflect seasonal temperature variation and hence intervals of a year, the much smaller number over a given height interval from *A. opercularis* confirms that this species grew a great deal faster than the others (more than twice as fast as *A. islandica* and *P. rustica*, and three to five times faster than *G. radiolyrata*). In *A. opercularis* profiles spanning  
570 two or more years (AO10, AO6 —Fig. 7a, e, respectively), there is an ontogenetic decrease in

wavelength as well as amplitude—i.e. growth was fastest in early ontogeny. ~~This~~  
decline in growth rate has been widely documented in *A. opercularis* from both  $\delta^{18}\text{O}$  and other  
evidence (e.g. Johnson et al., 2021b), and in the present instances (in which  $\delta^{18}\text{O}$  maxima and  
575 minima are not associated with growth breaks) the ontogenetic decrease in amplitude of  $\delta^{18}\text{O}$   
cycles is probably a consequence of the general slowing of growth with age, leading to time-  
averaging in samples. Whatever the explanation, seasonal temperature variation is likely to be  
most faithfully reflected by the first  $\delta^{18}\text{O}$  cycle in *A. opercularis* profiles. The profiles from *G.*  
*radiolyrata*, *P. rustica* and *A. islandica* undoubtedly omit several early ontogenetic cycles and  
580 given the short wavelength of the later cycles represented it may be that the amplitude of these  
is reduced by time-averaging, as inferred in *A. opercularis*. Even if the closer spacing of  
samples from *G. radiolyrata*, *P. rustica* and *A. islandica* may have been sufficient in principle  
for resolution of seasonal  $\delta^{18}\text{O}$  extremes, the association of growth breaks with maxima,  
minima or both suggests that some recorded extremes are not representative of the most  
585 extreme temperatures experienced by the organism in the season concerned—i.e.  $\delta^{18}\text{O}$   
variation may not fully reflect seasonal temperature variation.



590 **Figure 7:** Isotopic, microgrowth-increment and growth-break data from Merksem-equivalent *A. opercularis* (a, b) and Oorderen Member (and equivalent) *A. opercularis* (c-f), *P. rustica* (g) and *A. islandica* (h). Format and symbols as in Fig. 6.

595 Compared to  $\delta^{18}\text{O}$  values from the same specimen,  $\delta^{13}\text{C}$  values generally show much less variation, ~~usually substantially so, particularly within the span of  $\delta^{18}\text{O}$  cycles. Nevertheless, in~~  
~~some specimens~~ ~~While in some specimens~~ there are intervals ~~of ontogeny~~ exhibiting covariation between  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  ~~(~~ moderate ~~–~~ strong positive covariation in AO10 (Fig. 7a), AO6 (Fig. 7e) and PR (Fig. 7g) between shell heights 25 and 53 mm ( $r^2 = 0.61$ ), 18 and 46 mm ( $r^2 = 0.84$ ), and 31 and 46 mm ( $r^2 = 0.34$ ), respectively; ~~AO6, PR—Fig. 7a, e, g, respectively;~~  
600 ~~moderate–strong negative covariation in GR2 (Fig. 6e) and GR1 (Fig. 6e, (Fig. 6–f, respectively))~~ between shell heights 25 and 43 mm ( $r^2 = 0.68$ ), and 26 and 42 mm ( $r^2 = 0.40$ ), respectively. However, the general picture is of fluctuations (if any) in  $\delta^{13}\text{C}$  that are independent of  $\delta^{18}\text{O}$ . The *A. opercularis* specimens show a marginal to clear overall decrease in  $\delta^{13}\text{C}$  through ontogeny, while the *P. rustica* specimen shows little change and the *A. islandica* and *G. radiolyrata* specimens show clear overall increases. The mean values from  
605 the *A. opercularis* specimens are very similar—from  $+0.31 \pm 0.22$  ‰ ( $\pm 1\sigma$ ) in AO8 to  $+0.77 \pm 0.24$  ‰ in AO9—and comparable to the mean from the *P. rustica* specimen ( $+0.98 \pm 0.18$  ‰), but much lower than the means from the *A. islandica* ( $+2.44 \pm 0.35$  ‰) and *G. radiolyrata* (GR1,  $+2.42 \pm 0.40$  ‰; GR2,  $+2.69 \pm 0.32$  ‰) specimens. The data from *A. opercularis* and *A. islandica* compare closely with those from early Pliocene examples of these species from  
610 eastern England (Johnson et al., 2009; Vignols et al., 2019). The difference between the means from early Pliocene *A. opercularis* (calcitic) and *A. islandica* (aragonitic) was ascribed principally to the mineralogical difference (Vignols et al., 2019). This interpretation is supported by the mean values from the present *G. radiolyrata* (aragonitic) specimens, which are similar to those from *A. islandica*, but not by the *P. rustica* (also aragonitic) mean value,  
615 which is only a little outside the range of mean values from *A. opercularis*. The different pattern of overall ontogenetic change in *G. radiolyrata* and *A. islandica* (increase, unlike in *A. opercularis* and *P. rustica*) also remains to be explained, as does the unusual negative covariation between  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  in *G. radiolyrata*.

### 5.3 Microgrowth-increment patterns (*A. opercularis*)

620 Even in smoothed (5-point moving average) profiles of microgrowth-increment size from *A. opercularis*, substantial high-frequency variation is present in nearly all cases. However, amongst those profiles long enough to show a low-frequency pattern, in a number of cases a fairly clear and complete major cycle proceeding from small to large to small increments is

discernible over about the first 40 mm of shell height. Such a cycle is evident in three of the  
625 four Luchtbal-Member (and equivalent) profiles, in each case with an amplitude (difference  
between the maximum and minimum of the smoothed profile) of more than 0.30 mm. The  
exception (AO4—Fig. 5a) is a profile too short to show this pattern. Only one (AO6—Fig. 6g)  
of the four Oorderen-Member (and equivalent) profiles has an amplitude greater than 0.30 mm,  
but a second (AO5—Fig. 6f) has an amplitude only fractionally less and a third (AO8—Fig.  
630 6c) is too short to show equivalent ('high amplitude') variation. Despite their considerable  
length the Merkssem-equivalent profiles exhibit an amplitude well below 0.30 mm ('low  
amplitude'). The prevalent high-amplitude pattern from Luchtbal-Member (and equivalent)  
shells corresponds to that in modern sub-thermocline shells, and the occurrence of the pattern  
in an Oorderen-Member shell is at least inconsistent with a supra-thermocline setting (Johnson  
635 et al., 2009, 2021b). The low-amplitude pattern in the two Merkssem-equivalent shells is  
consistent with a supra-thermocline setting—~~however, but~~ given the occasional occurrence of  
such a pattern in sub-thermocline shells, ~~it~~ is not inconsistent with the latter setting.

## 6 Interpretation

### 6.1 Temperatures

#### 6.1.1 Derivation, comparison and evaluation of seasonal seafloor values

The equations and water  $\delta^{18}\text{O}$  values that were employed to calculate summer and winter  
temperatures from shell  $\delta^{18}\text{O}$  are explained in Sect. 4.2. Following the reasoning of Johnson et  
al. (2017), the shell  $\delta^{18}\text{O}$  values used were the extreme ones at inflection points in profiles,  
supplemented in the present case by those profile-end values possibly corresponding to  
645 inflection points on the evidence of similar (true) inflection-point values from the same and  
other co-occurring shells of the species concerned. The profile-end values probably provide  
slight underestimates of summer temperature and slight overestimates of winter temperature in  
some cases—i.e. had the profiles extended farther, slightly lower or higher  $\delta^{18}\text{O}$  values,  
respectively, might have been identified. As well as errors from this source, others of the same  
650 type no doubt exist in the case of data from locations close to growth breaks (as a result of an  
incomplete record) and, for at least *A. opercularis*, in the case of late ontogenetic data (as a  
result of time-averaging), although no significant error is likely where the temperatures  
concerned are higher (for summers) or lower (for winters) than a reliable estimate from another  
time in ontogeny. Alongside the profile-end and near-growth break (unreliable) winter  $\delta^{18}\text{O}$   
655 data from *G. radiolyrata* is one inflection-point value (the first, unaccompanied by a growth

break, at a height of 17.9 mm in GR2; Fig. 6e) which appears acceptable as a source of reliable temperature information. The  $\delta^{18}\text{O}$  value is, however, lower than any from this taxon regarded as an unreliable source of winter temperatures. Hence, it too must be regarded as suspect, possibly recording a downward temperature fluctuation in spring rather than true winter conditions.

All the calculated temperatures are represented in Fig. 8, those based on equations providing 'minimum' seasonal ranges for *G. radiolyrata* and *A. opercularis* combined in Fig. 8a, those providing 'maximum' seasonal ranges for these species combined in Fig. 8b, and those providing a 'minimum' seasonal range for *G. radiolyrata* and a 'maximum' for *A. opercularis* (a 'hybrid' set) combined in Fig. 8c. Unreliable values (probable underestimates for summers and overestimates for winters) are identified by use of an open symbol, whereby we have applied the above reasoning uniformly except to *G. radiolyrata*, *A. islandica* and *P. rustica*: in the absence of early ontogenetic data for comparison in these species, we have assumed that the late ontogenetic data represented are free from time-averaging effects.

Figure 8 shows a general similarity in seasonal temperatures within and between stratigraphic members, with the exception of winter values supplied by *G. radiolyrata* from the Luchtbal Member, which are markedly higher than those from Luchtbal-Member *A. opercularis*. Since the specimens of each species come from different (~~although-although~~ immediately adjacent), conceivable that the data reflect environmental change. However, given the fact that the summer temperatures supplied by the species are close or identical (dependent on the method of calculation) and that all of the 12 sets of winter temperatures from *G. radiolyrata* are probable overestimates, it seems much more likely that the change is apparent rather than real. If this is accepted then it is sensible to view those Luchtbal-Member (and equivalent) summer temperatures represented in Fig. 8a and Fig. 8c (very similar from *G. radiolyrata* and *A. opercularis* for the same water  $\delta^{18}\text{O}$  value) as more accurate than those in Fig. 8b (somewhat lower from *A. opercularis* than from *G. radiolyrata* for the same water  $\delta^{18}\text{O}$  value). The Oorderen-Member winter temperatures from *A. islandica* and *P. rustica* are closer to those from Oorderen-Member *A. opercularis* (from the same bed) in Fig. 8a than those in Fig. 8c (and Fig. 8b), suggesting that the data in Fig. 8a are the most accurate overall. However, if it is untrue that the winter data from *A. islandica* and *P. rustica* are free from time-averaging effects (i.e. if the data are unreliable) there is no reason for favouring the remaining data in Fig. 8a over those in Fig. 8c.



690

695

700

705

**Figure 8:** Winter and summer seafloor temperatures, calculated from the seasonal extreme  $\delta^{18}\text{O}$  values indicated in Figs. 6 and 7, using water  $\delta^{18}\text{O}$  values of 0.0 ‰ and +0.4 ‰ and various equations. **(a):** equation of Royer et al. (2013) for GR, of O'Neil et al. (1969) for AO, and of Grossman and Ku (1986) for AI and PR; **(b):** equation of Grossman and Ku (1986) for GR, AI and PR, and of Bemis et al. (1998) for AO; **(c):** equation of Royer et al. (2013) for GR, of Bemis et al. (1998) for AO, and of Grossman and Ku (1986) for AI and PR. Interval means are for reliable seasonal temperatures (see Sect. 6.1.1) from the Luchtbal Member and equivalent, Oorderen Member and equivalent, and Merksem-equivalent strata. The indicated present-day seasonal seafloor temperature range (4.7–16.9 °C) is for 25 m depth at 53° N, 03° E.

**Figure 8:** Winter and summer seafloor temperatures calculated from the seasonal extreme  $\delta^{18}\text{O}$  values (SFR: SSFT minus WSFT) and annual average temperature (ASFT; midpoint between SSFT and WSFT) for 'members', based on the reliable data indicated in Fig. 8. **Table 2:** Mean seasonal seafloor temperatures (°C;  $\pm 1\sigma$ ) and seafloor temperature range (summer minus winter) for 'members', based on the reliable data indicated in Fig. 8.

	Member and/or equivalent	Water $\delta^{18}\text{O} = 0.0 \text{ ‰}$				Water $\delta^{18}\text{O} = +0.4 \text{ ‰}$			
		SSFT	WSFT	SFR	ASFT	SSFT	WSFT	SFR	ASFT
Fig. 8a	Merksem	15.1 $\pm$ 1.1	6.2 $\pm$ 0.0	8.9	10.7	16.8 $\pm$ 1.1	7.8 $\pm$ 0.0	9.0	12.3
	Oorderen	17.3 $\pm$ 1.6	6.7 $\pm$ 1.0	10.6	12.0	19.1 $\pm$ 1.6	8.3 $\pm$ 1.1	10.8	13.7
	Luchtbal	16.2 $\pm$ 0.4	6.4 $\pm$ 0.6	9.8	11.3	17.5 $\pm$ 0.5	8.0 $\pm$ 0.6	9.5	12.8
Fig. 8b	Merksem	14.5 $\pm$ 1.2	4.1 $\pm$ 0.0	10.4	9.3	16.4 $\pm$ 1.2	6.0 $\pm$ 0.0	10.4	11.2
	Oorderen	17.2 $\pm$ 1.6	5.6 $\pm$ 2.0	11.6	11.4	19.0 $\pm$ 1.6	7.4 $\pm$ 1.9	11.6	13.2
	Luchtbal	17.1 $\pm$ 1.1	4.2 $\pm$ 0.7	12.9	10.7	18.9 $\pm$ 1.0	6.2 $\pm$ 0.7	12.7	12.6
Fig. 8c	Merksem	14.5 $\pm$ 1.2	4.1 $\pm$ 0.0	10.4	9.3	16.4 $\pm$ 1.2	6.0 $\pm$ 0.0	10.4	11.2
	Oorderen	17.2 $\pm$ 1.6	5.6 $\pm$ 2.0	11.6	11.4	19.0 $\pm$ 1.6	7.4 $\pm$ 1.9	11.6	13.2
	Luchtbal	16.1 $\pm$ 0.5	4.2 $\pm$ 0.7	11.9	10.2	17.4 $\pm$ 0.4	6.2 $\pm$ 0.7	11.2	11.8

710

715

The interval mean values for each member (and its equivalent) shown in Fig. 8 are listed in Table 2. Table 2 gives the interval mean values for seasonal temperatures based on reliable data, as depicted in Fig. 8. Unlike other changes, the Luchtbal to Oorderen increases and the Oorderen to Merksem decreases in mean summer temperature evident in Fig. 8a and Fig. 8c are statistically significant for both water  $\delta^{18}\text{O}$  values (one-tailed *t*-tests;  $p < 0.05$ ). The fact that these changes correspond at least qualitatively to the changes in summer temperature inferred from dinoflagellates dinocysts and ostracods (Sect. 3) cements the impression that the data in Fig. 8a and Fig. 8c provide a more accurate picture of seasonal temperatures, and hence of seasonal range, than the data in Fig. 8b, in which only the Oorderen to Merksem decrease in summer temperature (again statistically significant) is evident.

**Table 3** Summer (SSFT) and winter (WSFT) seafloor temperatures (°C) for the year showing the greatest range (SFR: SSFT minus WSFT) from each shell, together with the annual average seafloor temperature (ASFT; midpoint between SSFT and WSFT). Unreliable as well as reliable seasonal

720 temperatures used (Fig. 8). AO8 (no winter data) has been included to supplement the summer data.

Member and/or equiv.	Shell	Temperatures using the equations of O'Neil et al. (1969) for AO, Royer et al. (2013) for GR, Grossman and Ku (1986) for AI and PR								Temperatures using the equations of Bemis et al. (1998) for AO, Grossman and Ku (1986) for GR							
		Water $\delta^{18}\text{O} = 0.0 \text{ ‰}$				Water $\delta^{18}\text{O} = +0.4 \text{ ‰}$				Water $\delta^{18}\text{O} = 0.0 \text{ ‰}$				Water $\delta^{18}\text{O} = +0.4 \text{ ‰}$			
		SSFT	WSFT	SFR	ASFT	SSFT	WSFT	SFR	ASFT	SSFT	WSFT	SFR	ASFT	SSFT	WSFT	SFR	ASFT
M.	AO10	13.6	6.2	7.4	9.9	15.3	7.8	7.5	11.6	12.8	4.1	8.7	8.5	14.7	6.0	8.7	10.4
	AO9	15.9	6.2	9.7	11.1	17.6	7.8	9.8	12.7	15.4	4.1	11.3	9.8	17.3	6.0	11.3	11.7
Oorderen	AO8	16.6				18.4				16.2				18.1			
	AO7	18.0	5.9	12.1	12.0	19.8	7.4	12.4	13.6	17.7	3.6	14.1	10.7	19.6	5.6	14.0	12.6
	AO6	18.3	6.8	11.5	12.6	20.1	8.4	11.7	14.3	18.0	4.8	13.2	11.4	19.9	6.7	13.2	13.3
	AO5	18.6	5.8	12.8	12.2	20.4	7.3	13.1	13.9	18.3	3.5	14.8	10.9	20.3	5.4	14.9	12.9
	PR	18.4	7.6	10.8	13.0	20.1	9.4	10.7	14.8								
	AI	18.2	6.0	12.2	12.1	19.9	7.7	12.2	13.8								
Luchtbal	AO4	16.2	6.6	9.6	11.4	18.0	8.2	9.8	13.1	15.8	4.5	11.3	10.2	17.7	6.5	11.2	12.1
	AO3	16.0	6.4	9.6	11.2	17.8	8.0	9.8	12.9	15.5	4.3	11.2	9.9	17.5	6.3	11.2	11.9
	AO2	15.9	7.0	8.9	11.5	17.6	8.6	9.0	13.1	15.4	5.0	10.4	10.2	17.3	6.9	10.4	12.1
	AO1	16.3	5.4	10.9	10.9	18.0	7.0	11.0	12.5	15.8	3.1	12.7	9.5	17.8	5.0	12.8	11.4
	GR2	16.4	11.1	5.3	13.8	17.5	12.1	5.4	14.8	17.9	9.1	8.8	13.5	19.6	10.8	8.8	15.2
	GR1	16.7	11.5	5.2	14.1	17.8	12.5	5.3	15.2	18.3	9.8	8.5	14.1	20.0	11.5	8.5	15.8

In conclusion, the data in Fig. 8a are probably the most accurate but the data in Fig. 8c should not be excluded from consideration, especially as evidence from modern *A. opercularis* (Johnson et al., 2021b) suggests that the equation of Bemis et al. (1998), used for calculation of temperatures from this species in Fig. 8c, provides more accurate temperatures than the equation of O'Neil et al. (1969), used in Fig. 8a.

### 6.1.2 Seasonal seafloor ranges

With the exception of the Luchtbal values resulting from the questionable combination of data employed in Fig. 8b, all the seasonal ranges for members (~~differences between mean summer and winter values for the various combinations of data in Fig. 8~~irrespective of how data have been combined and temperatures calculated; Table 2) are lower than the current seafloor range at offshore locations in the southern North Sea—e.g. at 53° N, 03° E, where the ~~range difference between mean maximum and minimum temperatures~~ is 12.2 °C (Johnson et al., 2021b, fig. 1A). However, ~~they are only slightly lower (compare the separations of the dotted/dashed lines in Fig. 8 with the width of the grey bar showing the current range at 53° N, 03° E) and~~ not all individual specimens indicate a lower range (Table 3): the *A. islandica* specimen (Oorderen Member) shows the same range as currently at 53° N, 03° E, and even a water  $\delta^{18}\text{O}$  value of 0.0 ‰ used with the equation of O'Neil et al. (1969) yields a range (12.8 °C) from the Oorderen *A. opercularis* specimen AO5 that is higher than at present. The latter case is based on a winter temperature that is not reliable in the sense of Sect. 6.1.1, but the true (most extreme) winter temperature can only have been less and the seasonal range hence greater. Using a water  $\delta^{18}\text{O}$  value of +0.4 ‰ with the equation of O'Neil et al. (1969) yields a range of 12.4 °C from the Oorderen specimen AO7, and using the equation of Bemis et al.

(1998) yields a range of 13.2 °C from the Oorderen specimen AO6 and 12.8 °C from the  
745 Luchtbal specimen AO1 (independent of water  $\delta^{18}\text{O}$ ), as well as higher ranges than with the  
equation of O'Neil et al. (1969) from the Oorderen specimens AO5 (14.89 °C) and AO7 (14.04  
°C). It can therefore be said that at times during the period of deposition of the Oorderen  
Member, and possibly of the Luchtbal Member, -the seasonal range in seafloor temperature  
was higher than the current typical range at present. One should, however, bear in mind the  
750 variation of about 2 °C either side of the mean maximum and minimum temperatures in the  
North Sea at present (Lane and Prandle, 1996), so the 'high' individual Oorderen and Luchtbal  
ranges do not necessarily manifest significantly greater seasonality than now and possibly  
substantially so, and that the range may sometimes have been higher than at present during the  
period of deposition of the Luchtbal Member. Individual specimens from the equivalent of the  
755 Merksem Member provide no some evidence of a higher-lower seafloor range than now  
(maximum range 11.3 °C from AO9, calculated with the equation of Bemis et al., 1998), but  
the small sample size (two) should be noted.

The calculations leading to the above figures for seasonal range assume constant water  $\delta^{18}\text{O}$   
760 during the intervals of ontogeny concerned. If at the times of maximum temperature the actual  
water  $\delta^{18}\text{O}$  value was lower than assumed the calculated temperatures would be overestimates;  
similarly, if at the times of minimum temperature the actual water  $\delta^{18}\text{O}$  value was higher than  
assumed the calculated temperatures would be underestimates. Each of these situations, or the  
two together, would yield an overestimate of seasonal range. While these circumstances are  
765 possible, they are improbable. As noted in Sect. 2, water  $\delta^{18}\text{O}$  is relatively high (not low) during  
summer and relatively low (not high) during winter in coastal waters of the North Sea at  
present. The calculated seasonal ranges are thus more likely to be underestimates.

**Table 4** Mean summer (SSST) and winter (WSST) sea surface temperatures (°C;  $\pm 1\sigma$ ), seasonal range (SSR: SSST minus WSST) and annual average temperature (ASST; midpoint between SSST and WSST) for 'members', based on the reliable data indicated in Fig. 8, with a 3 °C addition to Luchtbal and Oorderen summer seafloor temperatures in recognition of summer stratification (see text, Section 6.1.3, for explanation).

	Member and/or equivalent	Water $\delta^{18}\text{O} = 0.0 \text{ ‰}$				Water $\delta^{18}\text{O} = +0.4 \text{ ‰}$			
		SSST	WSST	SSR	ASST	SSST	WSST	SSR	ASST
Fig. 9a	Merksem	15.1 ± 1.1	6.2 ± 0.0	8.9	10.7	16.8 ± 1.1	7.8 ± 0.0	9.0	12.3
	Oorderen	20.3 ± 1.6	6.7 ± 1.0	13.6	13.5	22.1 ± 1.6	8.3 ± 1.1	13.8	15.2
	Luchtbal	19.2 ± 0.4	6.4 ± 0.6	12.8	12.8	20.5 ± 0.5	8.0 ± 0.6	12.5	14.3
Fig. 9b	Merksem	14.5 ± 1.2	4.1 ± 0.0	10.4	9.3	16.4 ± 1.2	6.0 ± 0.0	10.4	11.2
	Oorderen	20.2 ± 1.6	5.6 ± 2.0	14.6	12.9	22.0 ± 1.6	7.4 ± 1.9	14.6	14.7
	Luchtbal	20.1 ± 1.1	4.2 ± 0.7	15.9	12.2	21.9 ± 1.0	6.2 ± 0.7	15.7	14.1
Fig. 9c	Merksem	14.5 ± 1.2	4.1 ± 0.0	10.4	9.3	16.4 ± 1.2	6.0 ± 0.0	10.4	11.2
	Oorderen	20.2 ± 1.6	5.6 ± 2.0	14.6	12.9	22.0 ± 1.6	7.4 ± 1.9	14.6	14.7
	Luchtbal	19.1 ± 0.5	4.2 ± 0.7	14.9	11.7	20.4 ± 0.4	6.2 ± 0.7	14.2	13.3

### 6.1.3 Seasonal surface ranges

775 The water depths indicated by the bivalve mollusc assemblages of the Luchtbal and Oorderen members (respective minimum depths 40 and 35 m; Sect. 3) are greater than the typical depth of the summer thermocline in shelf settings (25–30 m). Microgrowth-increment evidence from *A. opercularis* (Sect. 5.3) is consistent with a sub-thermocline setting for both members, and the higher frequency of such evidence from the Luchtbal Member is consistent with the indication from assemblage analysis that this was deposited at a greater depth than the Oorderen Member. Microgrowth-increment evidence of a supra-thermocline setting from Merksem-equivalent shells in the Oosterhout Formation at Ouwerkerk is similarly consistent with the shallow depth of deposition (maximum 15 m) indicated by the biota of the Merksem Member itself at Antwerp (note that supra-thermocline settings exist now in the southern North Sea at a distance from the shore well beyond that of Ouwerkerk from the Pliocene shoreline; Fig. 2). Given a sub-thermocline setting for the Luchtbal and Oorderen members we must add a ‘stratification factor’ to summer seafloor temperatures to derive estimates of summer surface temperature and hence seasonal surface range (winter surface temperature is likely to have been the same as on the seafloor; Johnson et al., 2021b). There is a difference of 2.6 °C between the mean annual seafloor and surface temperature maxima at a seasonally stratified location (depth 59 m) in the central North Sea some 600 km north of the sites of the Pliocene shells (Johnson et al., 2021b). At this location the mean maximum surface temperature is only 13.7 °C, a figure substantially exceeded in the (unstratified) southern North Sea at present (e.g. 17.1 °C at 53° N, 03° E; Johnson et al., 2021b, fig. 1A) and with little doubt also in the Pliocene at warm times during deposition of the Oorderen Member –(see Sect. 3). From the evidence of dinocysts (De Schepper et al., 2009) it is conceivable that at cool times, and during deposition of the Luchtbal Member, summer surface temperature was not much higher than the present central North Sea

figure. A stratification factor of 3 °C is therefore a reasonable estimate for these cool intervals and a serviceable (conservative) one for warm intervals during deposition of the Oorderen Member. Adding 3 °C to the interval mean values for Luchtbal and Oorderen summer seafloor temperature yields stratification-adjusted figures for seasonal surface range (Table 4) that are in all cases higher than the present typical range in the southern North Sea. Therefore it can be scarcely contested that a stratification factor larger than 2.6 °C should be employed. The northern Adriatic is probably a better modern analogue than the central North Sea for the Pliocene SNSB. At a stratified location there (depth 38 m) annual seafloor and surface temperature maxima differ by between 3.2 and 9.9 °C (mean difference 7.7 °C; Johnson et al., 2021b), indicating that a stratification factor of 3 °C is the very least that should be applied to derive estimates of seasonal surface range from Luchtbal and Oorderen seafloor temperature data. Adding this figure to the seafloor ranges in Table 3 yields estimates of surface range that are larger than the present value in the southern North Sea (12.4 °C at 53° N, 03° E; Johnson et al., 2021b). Adding the same to individual values also yields figures for seasonal surface range (Table 5) that are higher than the present typical range, except in the cases of from all individuals apart from the Luchtbal specimens GR1, GR2 and AO2. And a lower range is only obtained from the last when benthic temperature is calculated using the equation of O'Neil et al. (1969), and the lower Oorderen ranges are especially large. The low ranges from Luchtbal specimens GR1 and GR2 reflect the high winter seafloor temperatures recorded, all of which are unreliable. While the Luchtbal and Oorderen (and respective equivalent) shells provide clear evidence of a higher surface range than now (see Fig. 9, which depicts the interval mean and individual information from Tables 4 and 5), a lower range is indicated by the two Merksem-equivalent shells, and AO2 would provide estimates of surface range no less than present with a stratification factor of 3.5 °C, a far from unreasonable figure. In contrast to the evidence of a higher surface range than now at most times during the periods of deposition of the Luchtbal and Oorderen members, the two Merksem equivalent shells show a lower seasonal range (Table 3), assuming they have been correctly interpreted as from a supra-thermocline setting individuals. It may of course be the case that they provide an unrepresentative picture of conditions during this interval picture for their time.

**Table 5** Summer (SSST) and winter (WSST) seafloor temperatures (°C) for the year showing the greatest range (SSR; SSST minus WSST) from each shell, together with the annual average seafloor

temperature (ASST; midpoint between SSST and WSST). SSST = SSFT for Merksem-equivalent (M.) shells; = SSFT + 3°C for Luchtbal and Oorderen (and equivalent) shells (see text, Section 6.1.3. for explanation). Unreliable as well as reliable seasonal temperatures used (Fig. 9). AO8 (no winter data) has been included to supplement the summer data.

Member and/or equiv.	Shell	Temperatures using the equations of O'Neil et al. (1969) for AO, Royer et al. (2013) for GR, Grossman and Ku (1986) for AI and PR								Temperatures using the equations of Bemis et al. (1998) for AO, Grossman and Ku (1986) for GR							
		Water $\delta^{18}\text{O} = 0.0 \text{ ‰}$				Water $\delta^{18}\text{O} = +0.4 \text{ ‰}$				Water $\delta^{18}\text{O} = 0.0 \text{ ‰}$				Water $\delta^{18}\text{O} = +0.4 \text{ ‰}$			
		SSST	WSST	SSR	ASST	SSST	WSST	SSR	ASST	SSST	WSST	SSR	ASST	SSST	WSST	SSR	ASST
M.	AO10	13.6	6.2	7.4	9.9	15.3	7.8	7.5	11.6	12.8	4.1	8.7	8.5	14.7	6.0	8.7	10.4
	AO9	15.9	6.2	9.7	11.1	17.6	7.8	9.8	12.7	15.4	4.1	11.3	9.8	17.3	6.0	11.3	11.7
Oorderen	AO8	19.6				21.4				19.2				21.1			
	AO7	21.0	5.9	15.1	13.5	22.8	7.4	15.4	15.1	20.7	3.6	17.1	12.2	22.6	5.6	17.0	14.1
	AO6	21.3	6.8	14.5	14.1	23.1	8.4	14.7	15.8	21.0	4.8	16.2	12.9	22.9	6.7	16.2	14.8
	AO5	21.6	5.8	15.8	13.7	23.4	7.3	16.1	15.4	21.3	3.5	17.8	12.4	23.3	5.4	17.9	14.4
	PR	21.4	7.6	13.8	14.5	23.1	9.4	13.7	16.3								
	AI	21.2	6.0	15.2	13.6	22.9	7.7	15.2	15.3								
Luchtbal	AO4	19.2	6.6	12.6	12.9	21.0	8.2	12.8	14.6	18.8	4.5	14.3	11.7	20.7	6.5	14.2	13.6
	AO3	19.0	6.4	12.6	12.7	20.8	8.0	12.8	14.4	18.5	4.3	14.2	11.4	20.5	6.3	14.2	13.4
	AO2	18.9	7.0	11.9	13.0	20.6	8.6	12.0	14.6	18.4	5.0	13.4	11.7	20.3	6.9	13.4	13.6
	AO1	19.3	5.4	13.9	12.4	21.0	7.0	14.0	14.0	18.8	3.1	15.7	11.0	20.8	5.0	15.8	12.9
	GR2	19.4	11.1	8.3	15.3	20.5	12.1	8.4	16.3	20.9	9.1	11.8	15.0	22.6	10.8	11.8	16.7
	GR1	19.7	11.5	8.2	15.6	20.8	12.5	8.3	16.7	21.3	9.8	11.5	15.6	23.0	11.5	11.5	17.3


#### 6.1.4 Absolute surface temperatures

As pointed out in Sect. 6.1.1, there is a Luchtbal–Oorderen increase and an Oorderen–Merksem decrease in mean summer seafloor temperature from reliable individual summer values calculated using the equation of Royer et al. (2013) for *G. radiolyrata*, the equation of Grossman and Ku (1986) for *A. islandica* and *P. rustica*, and the equations of both O'Neil et al. (1969) and Bemis et al. (1998) for *A. opercularis*. These changes are evident whether a water  $\delta^{18}\text{O}$  value of 0.0 ‰ or +0.4 ‰ is applied, are statistically significant, and correspond qualitatively to changes in summer temperature inferred from assemblages of ostracods and dinocysts. These proxies (and organic biomarkers—see below) provide estimates that are time-averaged to an extent similar to the interval mean temperatures quoted herein, making comparison fair. The data from ostracods and probably from dinocysts relate to summer sea surface temperature (SSST) so quantitative comparison must involve the equivalent data, derived as indicated in Sect. 6.1.3. The relevant interval mean information is that for Fig. 9a and 9c in Table 4, showing Oorderen SSST above 20 °C and Merksem SSST below 20 °C irrespective of the water  $\delta^{18}\text{O}$  value used in calculation, and Luchtbal SSST above 20 °C with a water  $\delta^{18}\text{O}$  value of +0.4 ‰ and below 20 °C with a water  $\delta^{18}\text{O}$  value of 0.0 ‰. All the Oorderen SSST estimates are consistent with dinocyst evidence (Sect. 3) specifying mainly warm temperate conditions (SSST > 20 °C; Johnson et al., 2021b) and the Luchtbal SSST estimates with a water value of 0.0 ‰ are consistent with dinocyst evidence specifying cool temperate conditions (SSST < 20 °C; Johnson et al., 2021b). This suggests that temperatures calculated with a water  $\delta^{18}\text{O}$  value of 0.0 ‰ may be the most accurate for all intervals. The



860 Merksem interval mean SSST calculated with this value is 14.5 or 15.1 °C dependent on the  
equation used (Table 4), both figures being consistent with the Oorderen–Merksem SSST  
decrease of 5–6 °C indicated by ostracod assemblage data (Sect. 3). The statements made in  
respect of interval means also apply in general to individual SSST values calculated with the  
same equations (Table 5). The only exceptions are the temperatures derived using a water  $\delta^{18}\text{O}$   
value of 0.0 ‰ from Oorderen specimen AO8, which are slightly below the warm temperate  
range for summer. Such temperatures are entirely to be expected from some individuals in a  
865 regime with a mean SSST only a few degrees above the cool temperate range (cf. Sect. 6.1.2).

The similarity of  $\delta^{18}\text{O}$ -derived estimates of summer sea surface temperature, particularly those  
using a water  $\delta^{18}\text{O}$  value of 0.0 ‰, to assemblage-based estimates lends credence to the  
equivalent winter sea surface temperatures (WSST). With the exception only of unreliable  
870 individual data from *G. radiolyrata*, all WSST estimates (Tables 4, 5) are firmly in the cool  
temperate range (< 10 °C; Johnson et al., 2021b). We can take the midpoint between interval  
mean SSST and WSST estimates as a figure for annual (‘average’) sea surface temperature  
(ASST) and compare this with ASST estimates based on other information. Robinson et al.  
(2018) derived an ASST of 13.6 °C for the mid-Piacenzian North Sea using bivalve  $\delta^{18}\text{O}$  data  
875 from the Coralline Crag Formation (UK). It is, however, questionable whether the Coralline  
Crag is of this age (see Fig. 3). Dearing Crampton-Flood et al. (2020) obtained a SSST of about  
16 °C (cool temperate) from alkenone index and a WSST of about 10 °C (boundary cool/warm  
temperate) from  $\text{TEX}_{86}$  for part of the Oosterhout Formation in the Netherlands (Noord-  
Brabant) representing the MPWP. These figures specify an ASST of 13 °C. This is very similar  
880 to the  $\delta^{18}\text{O}$ -derived ASST estimates (Table 4) of 12.9 and 13.5 °C obtained using a water  $\delta^{18}\text{O}$   
value of 0.0 ‰ from Oorderen and equivalent shells, which represent the same interval. Such  
figures for ASST (2–3 °C higher than the modern ASST of 10.9 °C specified by an SSST of  
17.1 °C and WSST of 4.7 °C at 53° N, 03° E; Johnson et al., 2021b, fig. 1A) are entirely  
consistent with general expectations for the MPWP, but the  $\delta^{18}\text{O}$ -derived data reveal that they  
885 result from substantially warmer summer conditions than at present and winter conditions  
much the same as now, opposite to the picture provided by alkenone and  $\text{TEX}_{86}$  data. The  
interval mean Luchtbal figures for SSST obtained using a water  $\delta^{18}\text{O}$  value of 0.0 ‰ are  
likewise above the present SSST, although less markedly so, and in combination with WSST  
figures similar to present also specify a higher ASST than now (Table 4). By contrast, the  
890 equivalent Merksem figures for SSST are below the present value and in combination with  
WSST figures similar to present specify a lower ASST than now.



**Figure 9:** Winter and summer sea-surface temperatures, calculated as in Fig. 8, using the same equations for (a), (b) and (c), but with a 3 °C supplement to Luchtbal and Oorderen summer temperatures in recognition of thermal stratification (see text for explanation). Interval means calculated as in Fig. 8. The indicated present-day seasonal sea-surface temperature range (4.7–17.1 °C) is for 53° N, 03° E. Note that the Luchtbal and Oorderen ranges, as indicated by the separation of dotted or dashed lines for each interval, are larger than the present-day range (see text for more detailed discussion).

895



900 The ontogenetic decline in  $\delta^{13}\text{C}$  shown by *A. opercularis* is as seen in modern examples of the species from diverse settings (Johnson et al., 2021b), and probably reflects increasing output of isotopically light respiratory carbon with increasing body size alongside slower shell secretion—i.e. reduced ‘demand’ for carbon (Lorrain et al., 2004). Short-term fluctuations paralleling changes in  $\delta^{18}\text{O}$  might similarly reflect variation in respiratory output determined by seasonal variation in the availability of food (Chauvaud et al., 2011). The opposite ontogenetic trend shown by *G. radiolyrata* and *A. islandica* can hardly be explained by a reduction in respiratory output, and the opposite short-term pattern shown by *G. radiolyrata* is unlikely to reflect a reversal in the timing of maximum food availability from summer to winter. These changes might reflect variation in the  $\delta^{13}\text{C}$  of dissolved inorganic carbon (DIC), the main source of carbon in shells (Marchais et al., 2015). Preferential uptake of  $^{12}\text{C}$  by photosynthesizers is a major influence on the  $\delta^{13}\text{C}$  of DIC, but high photosynthetic fixation of carbon in summer is a doubtful cause of the high summer  $\delta^{13}\text{C}$  values in *G. radiolyrata* because it would require (cf. Arthur et al., 1983) that the shells lived above the thermocline, which other evidence argues against. The anomalously low  $\delta^{13}\text{C}$  values from *P. rustica* (for the aragonite mineralogy) cannot be explained as a consequence of the incorporation of isotopically light carbon from porewaters (cf. Krantz et al., 1987) because the other infaunal species, *G. radiolyrata* and *A. islandica*, exhibit high values. Conceivably, the *P. rustica* values reflect a food source with a particularly low value (Marchais et al., 2015).

920 In view of the multiple potential ‘local’ controls on shell  $\delta^{13}\text{C}$  it is questionable whether the relatively low values from late Pliocene *A. opercularis*, compared to pre-industrial Holocene examples (Hickson et al., 2000), are a reflection of relatively high atmospheric  $\text{CO}_2$ , as was suggested to explain the similarly low values from early Pliocene forms (Johnson et al., 2009; Vignols et al., 2019).

## 925 **7 Implications of temperature data**

We have shown that by adopting certain equations relating shell  $\delta^{18}\text{O}$  to temperature, selecting a particular modelled value for water  $\delta^{18}\text{O}$ , and making a reasonable allowance for summer stratification (where indicated by independent evidence), it is possible to generate summer surface temperatures from shell  $\delta^{18}\text{O}$  data that are consistent with assemblage-derived estimates (cool or warm temperate according to interval) for the late Pliocene of Belgium and

the Netherlands. The corresponding winter surface temperatures are cool temperate in each of the three (Luchtbal, Oorderen and Merksem) intervals studied, and in conjunction with the summer temperatures demonstrate a higher seasonal surface range than at present in the area during the Luchtbal and Oorderen intervals, ~~and a particularly high range during the latter.~~

935 There is no reason to suppose that the Luchtbal and Oorderen data reflect glacial episodes, when seasonality appears to be enhanced (Hennissen et al., 2015, Crippa et al., 2016)—i.e. the different seasonality from now was under equivalent overall conditions (interglacial).

940 The ~~same approach~~ approach used herein has also revealed high seasonal surface ranges in the early Pliocene of the SNSB and the early and late Pliocene of the eastern seaboard of the USA (Johnson et al., 2017, 2019, 2021b; Vignols et al. 2019). Southward-flowing cool currents, as exist now (north of Cape Hatteras), were probably influential in the latter area, but no such current exists at present in the North Sea or is likely to have done during the Pliocene. Presently, winter temperature is raised somewhat in the North Sea by offshoots of the warm North Atlantic Current, principally entering from the north (Winther and Johannessen, 2006). Reduction of this oceanic heat supply, in conjunction with global (atmospheric) warmth, might perhaps have led to the seasonal surface temperatures of the Luchtbal and Oorderen intervals (similar to now in winter, warmer than now in summer) that account for the enhanced seasonal ranges. Raffi et al. (1985) made essentially the same suggestion to explain evidence of high

950 Pliocene seasonality in the southern North Sea from the bivalve assemblage.—Fluctuations in the strength and position of the North Atlantic Current during the Pliocene are certainly recognized from proxy evidence, and episodes of reduced oceanic heat supply could correspond to the Luchtbal and Oorderen intervals (e.g. Bachem et al., 2017; Panitz et al., 2018). For the latter (i.e. the MPWP), ~~when seasonality was greatest in the SNSB,~~ most models

955 indicate an increase in ocean heat transport in the North Atlantic compared to now, but some indicate a decrease (Zhang et al., 2021). However, even if the times of high seasonality in the SNSB correspond to periods of reduced oceanic heat supply (relative to the Pliocene norm or to the present) it is not yet clear that this is a sufficient explanation for the low winter temperatures contributing to pronounced seasonality. The Merksem decline in summer temperature (post-dating the MPWP) can be more certainly attributed to the global cooling which presaged the onset of northern hemisphere glaciation. The lack of a decline in winter temperature is consistent with Mg/Ca evidence from North Atlantic *Globigerina bulloides* (Foraminifera) suggesting that minimum temperatures did not start falling until the very end of the Pliocene, after the time of deposition of the Lillo Formation (Hennissen et al., 2015, 2017).

965

The similarity of the alkenone/TEX<sub>86</sub>-based estimate of ~~M~~ASST for the MPWP in the eastern SNSB to the sclerochronologically derived estimates both corroborates the latter and suggests that alkenone temperatures for other areas could be usefully supplemented by information from TEX<sub>86</sub>, even if this would be unlikely to specify the full seasonal range. Combined (average) figures would probably be lower than alkenone-only estimates and the relatively close alignment of proxy with model temperatures recently achieved for the northern North Atlantic (Sect.1) might be lost, with implications for model adequacy. Alkenone ~~temperatures can also be checked against sclerochronologically derived estimates for the~~ temperatures for the MPWP in the US Middle Atlantic Coastal Plain are, like those for this interval in the eastern SNSB, similar to summer surface ~~Here, as in the eastern SNSB, summer surface~~ figures based on sclerochronology (Dowsett et al., 2021) and very much higher than winter temperatures derived by this means (Johnson et al., 2017, 2019). The need for ‘moderation’ by winter proxy data in order to facilitate meaningful proxy–model comparison is therefore underlined ~~projected from benthic isotopic temperatures (Johnson et al., 2017, 2019) are several degrees higher than alkenone temperatures, which are typically 25–26 °C (Dowsett et al., 2019). However, again like the eastern SNSB, winter isotopic temperatures are very much cooler (at least from scallop species) and specify a MASST in the region of 20 °C. Thus sclerochronological evidence from the Middle Atlantic Coastal Plain also suggests that alkenone temperatures should be ‘moderated’ by combination with winter proxy data in order to facilitate meaningful comparison with model results.~~

The summer seafloor temperatures recorded from a late Pliocene *A. islandica* specimen herein (18.2 and 19.9 °C using water δ<sup>18</sup>O values of 0.0 and +0.4 ‰, respectively) corroborate evidence supplied by an early Pleistocene example from Italy (20.3 °C from specimen ACG254-1 using a water δ<sup>18</sup>O value of +0.5 ‰; Crippa et al., 2016, table 1) of a thermal range greater in the past than at present (upper limit 16 °C; Witbaard and Bergman, 2003). This indication of change in realized thermal niche supplements evidence of the same in several other bivalve taxa from the early Pliocene of the UK, although in most of these cases it is manifested by tolerance then of winter conditions cooler than are experienced by modern representatives or relatives, which are restricted to Mediterranean locations (Vignols et al., 2019). This information raises some doubts about the use of assemblage evidence to interpret past environments by means of ecological uniformitarianism, the very widely applied approach

1000 which assumes that ancient examples of taxa occupied the same niche as modern. Certainly the accuracy of this methodology for times beyond the recent past deserves reconsideration.

## 8 Further work

1005 Even if the analysed shells of Luchtbal and Oorderen age were to be from supra- rather than sub-thermocline settings, the  $\delta^{18}\text{O}$  data from them would specify a seasonal surface temperature range much higher than previously inferred—for example, ~~most the majority of~~ shells of Oorderen age indicate a range more than double the  $6^\circ\text{C}$  suggested by organic proxies (see Sect. 6.1.4 and the SFR data in Table 3). We argued in Sect. 2 and Sect. 6.1.2 that seasonal variation in shell  $\delta^{18}\text{O}$  is unlikely to have been enhanced by variation in water  $\delta^{18}\text{O}$ , and it can be added here that fluvial input (the means by which variation in water  $\delta^{18}\text{O}$  might have been brought about) may have been less in the Pliocene than now due to a smaller catchment area of the Rhine/Meuse/Scheldt system (Dearing Crampton-Flood et al., 2020), making enhancement of shell  $\delta^{18}\text{O}$  variation by water  $\delta^{18}\text{O}$  variation even more improbable. Notwithstanding these arguments, actual evidence of water  $\delta^{18}\text{O}$  would be very welcome, both as a check on the stability of values through the year and as a means of deriving accurate absolute temperatures. At present, substitution of independently derived temperatures (e.g. from carbonate clumped isotope or biomineral unit thermometry; Briard et al., 2020, Höche et al. 2020, 2021; Caldarescu et al., 2021; de Winter et al., 2021) into equations relating shell  $\delta^{18}\text{O}$  to temperature and water  $\delta^{18}\text{O}$  seems the best approach to determining the last parameter. However, the existence of fluid inclusions in bivalve shell carbonate (Nooitgedacht et al., 2021) raises the possibility that these might serve as a source of direct insight into the isotopic composition of ambient water during life.

1025 It would, of course, also be useful to have additional shell  $\delta^{18}\text{O}$  (and increment) data, more to confirm the declines in ~~summer sea surface temperature~~SSST, ~~annual sea surface temperature~~ASST and ~~sea surface range~~SSR indicated by the limited information from Merksem-age shells than to expand the already substantial evidence of high values for these parameters from shells of Luchtbal and Oorderen age. Notwithstanding some doubts about the reliability of the approach (Johnson et al., 2017), it would also be worth obtaining independent evidence of seasonality from variation in the size of zooids within bryozoan colonies. Using

1030 this technique, Knowles et al. (2009, table 3) obtained locality-specific seasonality estimates  
of  $8.08 \pm 1.38$  and  $8.15 \pm 1.30$  °C for the early Pliocene Ramsholt Member of eastern England,  
figures comparing closely with the value-estimate of  $7.77 \pm 1.12$  °C obtained through isotope  
sclerochronology of *A. opercularis* from this unit (mean difference between the winter and  
summer seafloor temperatures in Johnson et al., 2021b, table 3).

1035 In addition to the above refinements of proxy evidence, further modelling efforts are required  
to see whether the low winter temperatures indicated by sclerochronological evidence for the  
mid-Piacenzian SNSB can be reproduced, and whether reduced oceanic heat supply is a  
feasible cause. Haywood et al. (2000) modelled mid-Piacenzian summer surface temperatures  
1040 2–4 °C higher than present for the area, similar to those determined herein. However, they  
modelled winter temperatures 4–6 °C higher than present, approaching or into the warm  
temperate range and markedly above the firmly cool temperate values indicated by  
sclerochronology. Haywood et al. (2000) ascribed the reduced seasonality specified by their  
results to greater westerly wind stress and strength in the North Atlantic compared to now, with  
1045 a resultant increase in heat transport by the Gulf Stream and North Atlantic Current. More  
recent modelling of mid-Piacenzian seasonal sea-surface temperatures at higher northern  
latitudes indicates greater warming in summer than winter (de Nooijer et al., 2020)—i.e. higher  
seasonality than now, as inferred for the SNSB. Moreover, as already noted (Sect. 7), oceanic  
heat supply may not have been greater in the mid-Piacenzian. Possibly, use of up-to-date  
1050 models with revised boundary conditions may yield results conforming with the evidence of  
low winter temperatures and high seasonality from the SNSB.

## 9 Conclusions

Sclerochronological evidence from bivalves indicates that for most of the late Pliocene  
(including the MPWP) the seasonal range in surface temperature in the SNSB was higher than  
1055 now. This was probably a consequence of higher summer temperatures associated with global  
(atmospheric) warmth. The apparently similar winter temperatures to now may reflect partial  
withdrawal of oceanic heat supply to the region through a change in strength and/or position  
of the North Atlantic Current.

1060 Averaging sclerochronologically derived summer and winter temperatures yields an annual sea  
surface temperature ~~–MASST~~ 2–3°C higher than now in the SNSB, as does averaging

temperatures from alkenone and TEX<sub>86</sub> thermometry. However, alkenones provide underestimates of extreme summer temperature and TEX<sub>86</sub> provides overestimates of extreme winter temperature, hence these proxies do not specify the full seasonal range. The  
1065 sclerochronologically derived temperatures are based on shell δ<sup>18</sup>O and dependent on estimates of water δ<sup>18</sup>O that require testing. Back-calculation from temperatures obtained by carbonate clumped isotope or biomineral unit thermometry from the same shells constitutes a potential means.

**Data availability.** The raw isotope and increment data, and the corresponding shell heights,  
1070 are available in the open-source online repository Zenodo (<https://doi.org/10.5281/zenodo.5585630>; Johnson et al. 2021c).

**Author contributions.** ALAJ conceived the study, obtained financial support and access to isotope analytical facilities, conducted some of the isotopic sampling and treatment of results, and drafted the manuscript. AMV conducted the rest of the isotopic sampling and treatment of  
1075 results, and obtained the increment data, within the context of a PhD project supervised by ALAJ. BRS and MJL facilitated the isotopic analysis. SG provided shell photographs together with detailed information concerning the stratigraphy and environments of the Belgian and Dutch Pliocene, supplied as comments on the first draft of the manuscript. BRS and MJL also commented on the first draft.

1080 **Competing interests.** The authors declare that they have no conflict of interest.

**Acknowledgements.** We thank the following for facilitating study of museum specimens in their care: Annelise Folie, Robert Marquet and Etienne Steurbaut (Institut royal des Sciences naturelles de Belgique, Brussels, Belgium); Frank Wesselingh (Naturalis Biodiversity Center, Leiden, The Netherlands); Serge Gofas and Virginie Héros (Muséum National d'Histoire  
1085 Naturelle, Paris, France). Melita Peharda (Institut za oceanografiju i ribarstvo, Split, Croatia) and Guillermo Roman (Instituto Español de Oceanografía, La Coruña, Spain) generously supplied live-collected individuals. Mark Dean and Matthew Hunt (Geoscience, University of Derby, UK) kindly assisted with specimen preparation, and Michael Maus (Institut für Geowissenschaften, Universität Mainz, Germany) and Hilary Sloane (National Environmental  
1090 Isotope Facility, British Geological Survey, Keyworth, UK) with isotopic analysis. We are

grateful to the NERC Isotope Geosciences Facilities Steering Committee for granting analytical services (IP-1108-0509, IP-1155-1109), the British Geological Survey for a PhD studentship award to AMV (BUFI S157), and the Alexander von Humboldt-Stiftung for support of a research stay at Mainz by ALAJ.

## 1095 References

- Alexandroff, S. J., Butler, P. G., Hollyman, P. R., Schöne, B. R. and Scourse, J. D.: Late Holocene seasonal temperature variability of the western Scottish shelf (St Kilda) recorded in fossil shells of the bivalve *Glycymeris glycymeris*, *Palaeogeogr. Palaeoclimatol.*, 562, <https://doi.org/10.1016/j.palaeo.2020.110146>, 2021.
- 1100 Arthur, M. A., Williams, D. F. and Jones, D. S.: Seasonal temperature-salinity changes and thermocline development in the mid-Atlantic Bight as recorded by the isotopic composition of bivalves, *Geology*, 11, 655–659, [https://doi.org/10.1130/0091-7613\(1983\)11<655:STCATD>2.0.CO;2](https://doi.org/10.1130/0091-7613(1983)11<655:STCATD>2.0.CO;2), 1983.
- Bachem, P. E., Risebrobakken, B., De Schepper, S. and McClymont, E. L.: Highly variable Pliocene sea surface conditions in the Norwegian Sea, *Clim. Past*, 13, 1153–1168, <https://doi.org/10.5194/cp-13-1153-2017>, 2017.
- 1105 Bemis, B. E., Spero, H. J., Bijma, J. and Lea, D. W.: Reevaluation of the oxygen isotopic composition of planktonic foraminifera: Experimental results and revised paleotemperature equations, *Paleoceanography*, 13, 150–160, <https://doi.org/10.1029/98PA00070>, 1998.
- 1110 Bice, K. L., Arthur, M. A. and Marincovich, L.: Late Paleocene Arctic Ocean shallow-marine temperatures from mollusc stable isotopes, *Paleoceanography*, 11, 241–249, <https://doi.org/10.1029/96PA00813>, 1996.
- Bova, S., Rosenthal, Y., Liu, Z., Godad, S. P. and Yan, M.: Seasonal origin of the thermal maxima at the Holocene and the last interglacial, *Nature*, 589, 548–553, <https://doi.org/10.1038/s41586-020-03155-x>, 2021.
- 1115 Briard, J., Pucéat, E., Vennin, E., Daëronc, M., Chavagnac, V., Jaillet, R., Merle, D. and de Raféllis, M.: Seawater paleotemperature and paleosalinity evolution in neritic environments of the Mediterranean margin: Insights from isotope analysis of bivalve shells, *Palaeogeogr. Palaeoclimatol.*, 543, <https://doi.org/10.1016/j.palaeo.2019.109582>, 2020.
- 1120 Buchardt, B. and Simonarson, L. A.: Isotopic palaeotemperatures from the Tjörnes Beds in Iceland: evidence of Pliocene cooling, *Palaeogeogr. Palaeoclimatol.*, 189, 71–95, [https://doi.org/10.1016/S0031-0182\(02\)00594-1](https://doi.org/10.1016/S0031-0182(02)00594-1), 2003.

- 1125 Caldarescu, D.E., Sadatzki, H., Andersson, C., Schäfer, P., Fortunato, H. and Meckler, A. N.:  
Clumped isotope thermometry in bivalve shells: A tool for reconstructing seasonal  
upwelling, *Geochim. Cosmochim. Ac.*, 294, 174–191,  
<https://doi.org/10.1016/j.gca.2020.11.019>, 2021.
- 1130 Chauvaud, L., Thébault, J., Clavier, J., Lorrain, A. and Strand, Ø.: What's hiding behind  
ontogenetic  $\delta^{13}\text{C}$  variations in mollusk shells? New insights from the Great Scallop (*Pecten*  
*maximus*), *Estuar.ies Coast\_s*, 34, 211–220, <https://doi.org/10.1007/s12237-010-9267-4>,  
2011.
- Coplen, T. B., Kendall, C. and Hopple, J.: Comparison of stable isotope reference samples,  
*Nature*, 302, 236–238, <https://doi.org/10.1038/302236a0>, 1983.
- 1135 Crippa, G., Angiolini, L., Bottini, C., Erba, E., Felletti, F., Frigerio, C., Hennissen, J. A. I.,  
Leng, M. J., Petrizzo, M. R., Raffi, I., Raineri, G. and Stephenson, M. H.: Seasonality  
fluctuations recorded in fossil bivalves during the early Pleistocene: Implications for  
climate change, *Palaeogeogr. Palaeoclim.*, 446, 234–251,  
<https://doi.org/10.1016/j.palaeo.2016.01.029>, 2016.
- 1140 Dearing Crampton-Flood, E., Noorbergen, L. J., Smits, D., Boschman, R. C., Donders, T. H.  
and Muns, D. K.: A new age model for the Pliocene of the southern North Sea basin: a  
multi-proxy climate reconstruction, *Clim. Past* 16, 523–541, <https://doi.org/10.5194/cp-16-523-2020>, 2020.
- Deckers, J., Louwey, S. and Goolaerts, S.: The internal division of the Pliocene Lillo  
Formation: correlation between Cone Penetration Tests and lithostratigraphic type sections,  
*Geol. Belg.*, 23, 333–343, <https://doi.org/10.20341/gb.2020.027>, 2020.
- 1145 DeLong, K. L., Quinn, T. M. and Taylor, F. W.: Reconstructing twentieth-century sea surface  
temperature variability in the southwest Pacific: A replication study using multiple  
coral Sr/Ca records from New Caledonia, *Paleoceanography*, 22, PA4212,  
<https://doi.org/10.1029/2007PA001444>, 2007.
- 1150 DeLong, K. L., Flannery, J. A., Maupin, C. R., Poore, R. Z. and Quinn, T.M.: A coral Sr/Ca  
calibration and replication study of two massive corals from the Gulf of Mexico,  
*Palaeogeogr. Palaeoclim.*, 307, 117–128, <https://doi.org/10.1016/j.palaeo.2011.05.005>,  
2011.
- 1155 De Meuter, F. and Laga, P.: Lithostratigraphy and biostratigraphy based on benthonic  
Foraminifera of the Neogene deposits of northern Belgium, *Bulletin van de Belgische*  
*Vereniging voor Geologie*, 85, 133–152, 1976.



- 1160 de Nooijer, W., Zhang, Q., Li, Q., Zhang, Q., Li, X., Zhang, Z., Guo, C., Nisancioglu, K. H.,  
Haywood, A.M., Tindall, J. C., Hunter, S. J., Dowsett, H. J., Stepanek, C., Lohmann, G.,  
Otto-Bliesner, B. L., Feng, R., Sohl, L. E., Chandler, M. A., Tan, N., Contoux, C.,  
1165 Ramstein, G., 13, Baatsen, M. L. J., von der Heydt, A. S., Chandan, D., Peltier, W. R., Abe-  
Ouchi, A., Chan, W.-L., Kamae, Y. and Brierley, C. M.: Evaluation of Arctic warming in  
mid-Pliocene climate simulations, *Clim. Past*, 16, 2325–2341, <https://doi.org/10.5194/cp-16-2325-2020>, 2020.
- De Schepper, S., Head, M. J. and Louwye, S.: Pliocene dinoflagellate cyst stratigraphy,  
palaeoecology and sequence stratigraphy of the Tunnel-Canal Dock, Belgium. *Geol. Mag.*,  
1165 146, 92–112, <https://doi.org/10.1017/S0016756808005438>, 2009.
- de Winter, N. J., Ullmann, C. V., Sorensen, A. M., Thibault, N., Goderis, S., Van Malderen, S.  
J. M., Snoeck, C., Goolaerts, S., Vanhaecke, F. and Claeys, P.: Shell chemistry of the boreal  
Campanian bivalve *Rastellum diluvianum* (Linnaeus, 1767) reveals temperature  
seasonality, growth rates and life cycle of an extinct Cretaceous oyster, *Biogeosciences*, 17,  
1170 2897–2922, <https://doi.org/10.5194/bg-17-2897-2020>, 2020a.
- de Winter, N. J., Vellekoop, J., Clark, A. J., Stassen, P., Speijer, R. P. and Claeys, P.: The giant  
marine gastropod *Campanile giganteum* (Lamarck, 1804) as a high-resolution archive of  
seasonality in the Eocene greenhouse world, *Geochem. Geophys. Geosy.*, 21,  
e2019GC008794, <https://doi.org/10.1029/2019GC008794>, 2020b.
- 1175 de Winter, N. J., Agterhuis, T. and Ziegler, M.: Optimizing sampling strategies in high-  
resolution paleoclimate records, *Clim. Past* 17, 1315–1340, <https://doi.org/10.5194/cp-17-1315-2021>, 2021.
- Dowsett, H., Robinson, M., Haywood, A., Salzmann, U., Hill, D., Sohl, L., Chandler, M.,  
Williams, M., Foley, K. and Stoll, D.: The PRISM3D paleoenvironmental reconstruction,  
1180 *Stratigraphy*, 7, 123–129, 2010.
- Dowsett, H. J. Haywood, A. M., Valdes, P. J., Robinson, M. M., Lunt, D. J., Hill, D., Stoll, D.  
K. and Foley, K. M.: Sea surface temperatures of the mid-Piacenzian Warm Period: A  
comparison of PRISM3 and HadCM3, *Palaeogeogr. Palaeoclimatol.*, 309, 83–91,  
| <https://doi.org/10.1016/j.palaeo.2011.03.016>, 2011.
- 1185 Dowsett, H. J., Robinson, M. M., Stoll, D. K., Foley, K. M., Johnson, A. L. A., Williams, M.  
and Riesselman, C. R.: The PRISM (Pliocene palaeoclimate) reconstruction: time for a  
paradigm shift, *Philos. T. R. Soc. A*, 371, 20120524, 24 pp.,  
<https://doi.org/10.1098/rsta.2012.0524>, 2013.

1190 Dowsett, H. J., Robinson, M. M., Foley, K. M., Herbert, T. D., Otto-Bliesner, B. L. and Spivey,  
W.: The mid-Piacenzian of the North Atlantic Ocean, *Stratigraphy*, 16, 119–144,  
<https://doi.org/10.29041/strat.16.3.119-144>, 2019.

1195 Dowsett, H. J., Robinson, M. M., Foley, K. M. and Herbert, T. D.: The Yorktown Formation:  
Improved stratigraphy, chronology, and paleoclimate interpretations from the U.S. Mid-  
Atlantic Coastal Plain, *Geosciences*, 11, 486,  
<https://doi.org/10.3390/geosciences11120486>, 2021.

1200 Featherstone, A. M., Butler, P. G., Schöne, B. R., Peharda, M. and Thébault, J.: A 45-year sub-  
annual reconstruction of seawater temperature in the Bay of Brest, France, using the shell  
oxygen isotope composition of the bivalve *Glycymeris glycymeris*, *Holocene*, 30, 3–12,  
<https://doi.org/10.1177/0959683619865592>, 2020.

1205 Fenger, T., Surge, D., Schöne, B. R. and Milner, N.: Sclerochronology and geochemical  
variation in limpet shells (*Patella vulgata*): A new archive to reconstruct coastal sea surface  
temperature, *Geochem. Geophys. Geosy.*, 8, Q07001,  
<https://doi.org/10.1029/2006GC001488>, 2007.

1210 Füllenbach, C. S., Schöne, B. R. and Mertz-Kraus., R.: Strontium/lithium ratio in shells of  
*Cerastoderma edule* (Bivalvia) - A new potential temperature proxy for brackish  
environments, *Chem. Geol.*, 417, 341–355,  
<https://doi.org/10.1016/j.chemgeo.2015.10.030>, 2015.

1215 Funnell, B. M.: Plio-Pleistocene palaeogeography of the southern North Sea Basin (3.75–0.60  
Ma), *Quaternary Sci. Rev.*, 15, 391–405, [https://doi.org/10.1016/0277-3791\(96\)00022-4](https://doi.org/10.1016/0277-3791(96)00022-4),  
1996.

1220 Gaemers, P. A. M.: Enkele paleo-ecologische opmerkingen over de Pliocene afzettingen in de  
tunnelput nabij Kallo, België, provincie Oost-Vlaanderen. Deel 2, Mededelingen van de  
Werkgroep voor Tertiare en Kwartaire Geologie, 12, 43–49, 1975.

1225 Gaemers, P. A. M. and Schwarzahns, W.: Fisch-Otolithen aus dem Pliozän von Antwerpen  
(Belgien) und Ouwerkerk (Niederlande) und aus dem Plio-Pleistozän der Westerschelde  
(Niederlande), *Leidse Geologische Mededelingen*, 49, 207–257, 1973.

Garcia-March, J. R., Surge, D., Lees, J. M. and Kersting, D. K.: Ecological information and  
water mass properties in the Mediterranean recorded by stable isotope ratios in *Pinna  
nobilis* shells, *J. Geophys. Res.-Biogeo.*, 116, G02009,  
<https://doi.org/10.1029/2010JG001461>, 2011.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Field Code Changed

Formatted: Hyperlink, Font: (Default) Times New Roman, 12 pt

- Gibbard, P. L. and Lewin, J.: Filling the North Sea Basin: Cenozoic sediment sources and river styles (André Dumont medallist lecture 2014), *Geol. Belg.*, 19, 201–217, <https://doi.org/10.20341/gb.2015.017>, 2016.
- 1225 Gillikin, D. P., Lorrain, A., Navez, J., Taylor, J. W., Keppens, E., Baeyens, W. and Dehairs, F.: Strong biological controls on Sr/Ca ratios in aragonitic marine bivalve shells, *Geochem. Geophys. Geosy.*, 6, Q05009, <https://doi.org/10.1029/2004GC000874>, 2005.
- 1230 Goodwin, D. H., Schöne, B. R. and Dettman, D. L.: Resolution and fidelity of oxygen isotopes as paleotemperature proxies in bivalve mollusk shells: Models and observations, *Palaios*, 18, 110–125, [https://doi.org/10.1669/0883-1351\(2003\)18<110:RAFOOI>2.0.CO;2](https://doi.org/10.1669/0883-1351(2003)18<110:RAFOOI>2.0.CO;2), 2003.
- Grossman, E. L. and Ku, T.: Oxygen and carbon isotope fractionation in biogenic aragonite: Temperature effects, *Chem. Geol. (Isotope Geoscience Section)*, 59, 59–74, [https://doi.org/10.1016/0009-2541\(86\)90044-6](https://doi.org/10.1016/0009-2541(86)90044-6), 1986.
- 1235 Hacquaert, N.: Palynologisch onderzoek van de cenozoïsche mariene zanden (Scaldisien en Merxemian) van het Hansadok te Antwerpen, *Natuurwetenschappelijk Tijdschrift*, 42, 65–112, 1960.
- Harwood, A. J. P., Dennis, P. F., Marca, A. D., Pilling, G. M. and Millner, R. S.: The oxygen isotope composition of water masses within the North Sea, *Estuar. Coast. Shelf Sci.*, 78, 353–359, <https://doi.org/10.1016/j.ecss.2007.12.010>, 2008.
- 1240 Haywood, A. M., Sellwood, B. W. and Valdes, P. J.: Regional warming: Pliocene (3 Ma) paleoclimate of Europe and the Mediterranean, *Geology*, 28, 1063–1066, 2000.
- Hennissen, J. A. I., Head, M. J., De Schepper, S. and Groeneveld, J.: Increased seasonality during the intensification of Northern Hemisphere glaciation at the Pliocene–Pleistocene transition ~2.6 Ma, *Quaternary Sci. Rev.*, 129, 321–332, <http://doi.org/10.1016/j.quascirev.2015.10.010>, 2015.
- 1245 Hennissen, J. A. I., Head, M. J., De Schepper, S. and Groeneveld, J.: Dinoflagellate cyst paleoecology during the Pliocene–Pleistocene climatic transition in the North Atlantic, *Palaeogeogr. Palaeoclimatol.*, 470, 81–108, <https://doi.org/10.1016/j.palaeo.2016.12.023>, 2017.
- 1250 Hickson, J. A., Johnson, A. L. A., Heaton, T. H. E. and Balson, P. S.: The shell of the Queen Scallop *Aequipecten opercularis* (L.) as a promising tool for palaeoenvironmental reconstruction: evidence and reasons for equilibrium stable-isotope incorporation, *Palaeogeogr. Palaeoclimatol.*, 154, 325–337, [https://doi.org/10.1016/S0031-0182\(99\)00120-0](https://doi.org/10.1016/S0031-0182(99)00120-0), 1999.

- 1255 Hickson, J. A., Johnson, A. L. A., Heaton, T. H. E. and Balson, P. S.: Late Holocene environment of the southern North Sea from the stable isotopic composition of Queen Scallop shells, *Palaeontol. Electron.*, 3 (2), art. 3, 11 pp., [https://palaeo-electronica.org/2000\\_2/scallop/issue2\\_00.htm](https://palaeo-electronica.org/2000_2/scallop/issue2_00.htm), 2000.
- 1260 Höche, N., Peharda, M., Walliser, E. O. and Schöne, B. R.: Morphological variations of crossed-lamellar ultrastructures of *Glycymeris bimaculata* (Bivalvia) serve as a marine temperature proxy, *Estuar. Coast. Shelf Sci.*, 237, 106658, <https://doi.org/10.1016/j.ecss.2020.106658>, 2020.
- 1265 Höche, N., Walliser, E. O., de Winter, N. J., Witbaard, R. and Schöne, B. R.: Temperature-induced microstructural changes in shells of laboratory-grown *Arctica islandica* (Bivalvia), *Plos One* 16, e0247968, <https://doi.org/10.1371/journal.pone.0247968>, 2021.
- Howarth, M. J., Dyer, K. R., Joint, I. R., Hydes, D. J., Purdie, D. A., Edmunds, H., Jones, J. E., Lowry, R. K., Moffat, T. J., Pomroy, A. J. and Proctor, R.: Seasonal cycles and their spatial variability, *Philos. T. R. Soc. A*, 343, 383-403, <https://doi.org/10.1098/rsta.1993.0054>, 1993.
- 1270 Huyghe, D., Emmanuel, L., de Rafelis, M., Renard, M., Ropert, M., Labourdette, N. and Lartaud, F.: Oxygen isotope disequilibrium in the juvenile portion of oyster shells biases seawater temperature reconstructions, *Estuar. Coast. Shelf Sci.*, 240, article 106777, <https://doi.org/10.1016/j.ecss.2020.106777>, 2020.
- 1275 Huyghe, D., Daëron, M., de Rafelis, M., Blamart, D., Sébilo, M., Paulet, Y.-M., and Lartaud, F.: Clumped isotopes in modern marine bivalves, *Geochim. Cosmochim. Ac.*, 316, 41–58, <https://doi.org/10.1016/j.gca.2021.09.019>, 2022.
- Ivany, L. C. and Judd, E. J.: Deciphering temperature seasonality in Earth’s ancient oceans, *Annu. Rev. Earth Planet. Sci.* 50:123–52, <https://doi.org/10.1146/annurev-earth-032320-095156>, 2022.
- 1280 Ivany, L. C., Wilkinson, B. H. and Jones, D. S.: Using stable isotope data to resolve rate and duration of growth throughout ontogeny: An example from the surf clam, *Spisula solidissima*, *Palaios*, 18, 126–137, [https://doi.org/10.1669/0883-1351\(2003\)18<126:USIDTR>2.0.CO;2](https://doi.org/10.1669/0883-1351(2003)18<126:USIDTR>2.0.CO;2), 2003.
- 1285 Ivany, L. C., Wilkinson, B. H., Lohmann, K. C., Johnson, E. R., McElroy, B. J. and Cohen, G. J.: Intra-annual isotopic variation in *Venericardia* bivalves: Implications for early Eocene temperature, seasonality, and salinity on the US Gulf Coast, *J. Sediment. Res.*, 74, 7–19, <https://doi.org/10.1306/052803740007>, 2004.

Field Code Changed

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Field Code Changed

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Field Code Changed

Formatted: Hyperlink, Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: Font color: Auto, Pattern: Clear

- 1290 Johnson, A. L. A., Hickson, J. A., Bird, A., Schöne, B. R., Balson, P. S., Heaton, T. H. E. and  
Williams, M.: Comparative sclerochronology of modern and mid-Pliocene (c. 3.5 Ma)  
1295 *Aequipecten opercularis* (Mollusca, Bivalvia): an insight into past and future climate  
change in the north-east Atlantic region, *Palaeogeogr. Palaeoclimatol.*, 284, 164–179,  
<https://doi.org/10.1016/j.palaeo.2009.0.022>, 2009.
- Johnson, A. L. A., Valentine, A., Leng, M. J., Sloane, H. J., Schöne, B. R. and Balson, P. S.:  
1295 Isotopic temperatures from the Early and Mid-Pliocene of the US Middle Atlantic Coastal  
Plain, and their implications for the cause of regional marine climate change, *Palaios*, 32,  
250–269, <https://doi.org/10.2110/palo.2016.080>, 2017.
- Johnson, A. L. A., Valentine, A. M., Leng, M. J., Schöne, B. R. and Sloane, H. J.: Life history,  
environment and extinction of the scallop *Carolinapecten eboreus* (Conrad) in the Plio-  
Pleistocene of the US eastern seaboard, *Palaios*, 34, 49–70,  
1300 <https://doi.org/10.2110/palo.2018.056>, 2019.
- Johnson, A. L. A., Harper, E. M., Clarke, A., Featherstone, A. C., Heywood, D. J., Richardson,  
K. E., Spink, J. O. and Thornton, L. A. H.: Growth rate, extinction and survival amongst  
late Cenozoic bivalves of the North Atlantic, *Hist. Biol.*, 33, 802–813,  
<https://doi.org/10.1080/08912963.2019.1663839>, 2021a.
- 1305 Johnson, A. L. A., Valentine, A. M., Schöne, B. R., Leng, M. J., Sloane, H. J. and Janeković,  
I.: Growth-increment characteristics and isotopic ( $\delta^{18}\text{O}$ ) temperature record of sub-  
thermocline *Aequipecten opercularis* (Mollusca:Bivalvia): evidence from modern Adriatic  
forms and an application to early Pliocene examples from eastern England, *Palaeogeogr.*  
*Palaeoclimatol.*, 561, <https://doi.org/10.1016/j.palaeo.2020.110046>, 2021b.
- 1310 Johnson, A. L. A., Valentine, A. M., Schöne, B. R., Leng, M. J., Sloane, H. J. and Goolaerts,  
S.: Raw data for “Sclerochronological evidence of pronounced seasonality from the late  
Pliocene of the southern North Sea Basin, and its implications (Version 1)”, Zenodo,  
<https://doi.org/10.5281/zenodo.5585630>, 2021c.
- Kim, S.-T. and O’Neil, J. R.: Equilibrium and nonequilibrium oxygen isotope effects in  
1315 synthetic carbonates, *Geochim. Cosmochim. Ac.*, 61, 3461–3475,  
[https://doi.org/10.1016/S0016-7037\(97\)00169-5](https://doi.org/10.1016/S0016-7037(97)00169-5), 1997.
- Kim, S.-T., O’Neil, J. R., Hillaire-Marcel, C. and Mucci, A.: Oxygen isotope fractionation  
between synthetic aragonite and water: Influence of temperature and  $\text{Mg}^{2+}$  concentration,  
*Geochim. Cosmochim. Ac.*, 71, 4704–4715, <https://doi.org/10.1016/j.gca.2007.04.019>,  
1320 2007.

Knowles, T., Taylor, P. D., Williams, M., Haywood, A. M. and Okamura, B.: Pliocene seasonality across the North Atlantic inferred from cheilostome bryozoans, *Palaeogeogr. Palaeoclimatol.*, 77, 226–235, <https://doi.org/10.1016/j.palaeo.2009.04.006>, 2009.

Krantz, D. E., Williams, D. F. and Jones, D. S.: Ecological and paleoenvironmental information using stable isotope profiles from living and fossil molluscs, *Palaeogeogr. Palaeoclimatol.*, 58, 249–266, [https://doi.org/10.1016/0031-0182\(87\)90064-2](https://doi.org/10.1016/0031-0182(87)90064-2), 1987.

Laga, P.: Stratigrafie van de mariene Plio-Pleistocene afzettingen uit de omgeving van Antwerpen met een bijzondere studie van de foraminiferen, Ph.D, thesis, Katholieke Universiteit Leuven, Belgium, 252 pp., 1972.

Lane, A. and Prandle, D.: *Inter-annual variability in the temperature of the North Sea, Cont. Shelf Res.*, 16, 1489–1507, [https://doi.org/10.1016/0278-4343\(96\)00001-5](https://doi.org/10.1016/0278-4343(96)00001-5), 1996.

Lisiecki, L. E. and Raymo, M. E.: A Pliocene–Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records, *Paleoceanography*, 20, 522–533, <https://doi.org/10.1029/2004PA001071>, 2005.

Lloyd, R. M.: Variations in the oxygen and carbon isotope ratios of Florida Bay mollusks and their environmental significance, *J. Geol.*, 72, 84–111, 1964..

Lorrain, A., Paulet, Y.-M., Chauvaud, L., Dunbar, R., Mucciarone, D. and Fontugne, M.:  $\delta^{13}\text{C}$  variation in scallop shells: Increasing metabolic carbon contribution with body size? *Geochim. Cosmochim. Ac.*, 68, 3509–3519, <https://doi.org/10.1016/j.gca.2004.01.025>, 2004.

Louwyse, S. and De Schepper, S.: *The Miocene-Pliocene hiatus in the southern North Sea Basin (northern Belgium) revealed by dinoflagellate cysts. Geol. Mag.* 147, 760–776, <https://doi.org/10.1017/S0016756810000191>, 2010.

Louwyse, S., Head, M. J. and De Schepper, S.: Dinoflagellate cyst stratigraphy and palaeoecology of the Pliocene in northern Belgium, southern North Sea Basin, *Geol. Mag.*, 141, 353–378, <https://doi.org/10.1017/S0016756804009136>, 2004.

Louwyse, S., Deckers, J. and Vandenberghe, N.: The Pliocene Lillo, Poederlee, Merksplas, Mol and Kieseloolite Formations in northern Belgium: a synthesis, *Geol. Belg.*, 23, 297–313, <https://doi.org/10.20341/gb.2020.016>, 2020.

Lunt, D. J., Haywood, A. M., Schmidt, G. A., Salzmann, U., Valdes, P. J. and Dowsett, H. J.: Earth system sensitivity inferred from Pliocene modelling and data, *Nat. Geosci.*, 3, 60–64, <https://doi.org/10.1038/NGEO706>, 2010.

Marchais, V., Richard, J., Jolivet, A., Flye-Sainte-Marie, J., Thébault, J., Jean, F., Richard, P., Paulet, Y.-M., Clavier, J. and Chauvaud, L.: Coupling experimental and field-based

Field Code Changed

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Field Code Changed

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: Not Italic

Field Code Changed

- 1355 approaches to decipher carbon sources in the shell of the great scallop, *Pecten*  
*maximus* (L.), Geochim. Cosmochim. Ac., 168, 58–69,  
<https://doi.org/10.1016/j.gca.2015.07.010>, 2015.
- Markulin, K., Peharda, M., Mertz-Kraus, R., Schöne, B. R., Uvanović, H., Kovač, Z. and  
1360 Janeković, I.: Trace and minor element records in aragonitic bivalve shells as environmental  
proxies, Chem. Geol., 507, 120–133, <https://doi.org/10.1016/j.chemgeo.2019.01.008>,  
2019.
- Marquet, R.: The Neogene Amphineura and Bivalvia (Protobranchia and Pteriomorphia) from  
Kallo and Doel (Oost-Vlaanderen, Belgium), Palaeontos, 2, 1–99, 2002.
- Marquet, R.: Ecology and evolution of Pliocene bivalves from the Antwerp Basin, Bull. Inst.  
1365 R. Sc. N. B-S, 74 suppl., 205–212, 2004.
- Marquet, R.: The Neogene Bivalvia (Heterodonta and Anomalodesmata) and Scaphopoda from  
Kallo and Doel (Oost-Vlaanderen, Belgium), Palaeontos, 6, 1–142, 2005.
- Marquet, R. and Herman, J.: The stratigraphy of the Pliocene in Belgium, Palaeofocus, 2, 1–  
39, 2009.
- 1370 Mettam, C., Johnson, A. L. A., Nunn, E. V. and Schöne, B. R. Stable isotope ( $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$ )  
sclerochronology of Callovian (Middle Jurassic) bivalves (*Gryphaea (Bilobissa) dilobotes*)  
and belemnites (*Cylindroteuthis pusoziana*) from the Peterborough Member of the Oxford  
Clay Formation (Cambridgeshire, England): evidence of palaeoclimate, water depth and  
belemnite behaviour, Palaeogeogr. Palaeoclimatol., 399, 187–201,  
1375 <https://doi.org/10.1016/j.palaeo.2014.01.010>, 2014.
- Mette, M. J., Whitney, N. M., Ballew, J. and Wanamaker, A. D.: Unexpected isotopic  
variability in biogenic aragonite: A user issue or proxy problem?, Chem. Geol., 483, 286–  
294, <https://doi.org/10.1016/j.chemgeo.2018.02>, 2018.
- 1380 Moon, L. R., Judd, E. J., Thomas, J., and Ivany, L. C.: Out of the oven and into the fire:  
Unexpected preservation of the seasonal  $\delta^{18}\text{O}$  cycle following heating experiments on shell  
carbonate, Palaeogeogr. Palaeoclimatol., 562, <https://doi.org/10.1016/j.palaeo.2020.110115>,  
2021.
- Munsterman, D. K., ten Veen, J. H., Menkovic, A., Deckers, J., Witmans, N., Verhaegen, J.,  
Kersthold-Boegehold, S. J., van de Ven, T. and Busschers, F.: An updated and revised  
1385 stratigraphic framework for the Miocene and earliest Pliocene strata of the Roer Valley  
Graben and adjacent blocks, Neth. J. Geosci., 98, e8, <https://doi.org/10.1017/njg.2019.10>,  
2020.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Field Code Changed

- Murray, J. W.: Palaeogene and Neogene, in: Atlas of Palaeogeography and Lithofacies, edited by: Cope, J. C. W., Ingham, J. K. and Rawson, P. F., Memoir 13, The Geological Society, London, UK, 141–147.  
1390
- Nooitgedacht, C. W., van der Lubbe, H. J. L., Ziegler, M. and Staudigel, P. T.: Internal water facilitates thermal resetting of clumped isotopes in biogenic aragonite, *Geochem. Geophys. Geosy.*, 22, e2021GC009730, <https://doi.org/10.1029/2021GC009730>, 2021.
- Norton, P. E. P.: Paleocology of the Mollusca of the Tjörnes sequence, Iceland, *Boreas*, 4, 97–110, 1975.  
1395
- O’Neil, J. R., Clayton, R. N. and Mayeda, T. K.: Oxygen isotope fractionation in divalent metal carbonates, *J. Chem. Phys.*, 51, 5547–5558, 1969.
- Overeem, I., Weltje, G. J., Bishop-Kay, C. and Kroonenberg, S. B.: The Late Cenozoic Eridanos delta system in the Southern North Sea Basin: a climate signal in sediment supply?, *Basin Res.*, 13, 293–312, <https://doi.org/10.1046/j.1365-2117.2001.00151.x>, 2001.  
1400
- Owen, R., Kennedy, H. and Richardson, C.: Isotopic partitioning between scallop shell calcite and seawater: Effect of shell growth rate, *Geochim. Cosmochim. Acta.*, 66, 1727–1737, [https://doi.org/10.1016/S0016-7037\(01\)00882-1](https://doi.org/10.1016/S0016-7037(01)00882-1), 2002a.  
1405
- Owen, R., Kennedy, H. and Richardson, C.: Experimental investigation into partitioning of stable isotopes between scallop (*Pecten maximus*) shell calcite and sea water, *Palaeogeogr. Palaeoclimatol.*, 185, 163–174, [https://doi.org/10.1016/S0031-0182\(02\)00297-3](https://doi.org/10.1016/S0031-0182(02)00297-3), 2002b.
- Panitz, S., Salzmann, U., Risebrobakken, B., De Schepper, S., Pound, M. J., Haywood, A. M., Dolan, A. M. and Lunt, D. J.: Orbital, tectonic and oceanographic controls on Pliocene climate and atmospheric circulation in Arctic Norway, *Global Planet. Change*, 161, 183–193, <https://doi.org/10.1016/j.gloplacha.2017.12.022>, 2018.  
1410
- Peharda, N., Crnčević, M., Bušelić, I., Richardson, C. A. and Ezgeta-Balić, D.: Growth and longevity of *Glycymeris nummaria* (Linnaeus, 1758) from the eastern Adriatic, Croatia, *J. Shellfish Res.*, 31, 947–950, <https://doi.org/10.2983/035.031.0406>, 2012.
- Peharda, M., Thébault, J., Markulin, K., Schöne, B. R., Janeković, I. and Chauvaud, L.: Contrasting shell growth strategies in two Mediterranean bivalves revealed by oxygen-isotope ratio geochemistry: The case of *Pecten jacobaeus* and *Glycymeris pilosa*, *Chem. Geol.*, 526, 23–35, <https://doi.org/10.1016/j.chemgeo.2017.09.029>, 2019a.  
1415
- Peharda, M., Walliser, E. O., Markulin, K., Purroy, A., Uvanović, H., Janeković, I., Župan, I., Vilibić, I. and Schöne, B. R.: *Glycymeris pilosa* (Bivalvia)—A high-potential geochemical  
1420



- archive of the environmental variability in the Adriatic Sea, *Mar. Environ. Res.*, 150, 104759, <https://doi.org/10.1016/j.marenvres.2019.104759>, 2019b.
- 1425 [Raffi, S., Stanley, S. M. and Marasti, R.: Biogeographic patterns and Plio-Pleistocene extinction of \*Bivalvia\* in the Mediterranean and southern North Sea, \*Paleobiology\*, 11, 368-388, <https://doi.org/10.1017/S0094837300011684>, 1985.](#)
- Reynolds, D. J., Hall, I. R., Slater, S. M., Scourse, J. D., Halloran, P. R. and Sayer, M. D. J.: Reconstructing past seasonal to multicentennial-scale variability in the NE Atlantic Ocean using the long-lived marine bivalve mollusk *Glycymeris glycymeris*, *Paleoceanography*, 32, 1153–1173, <https://doi.org/10.1002/2017PA003154>, 2017.
- 1430 Robinson, M. M.: New quantitative evidence of extreme warmth in the Pliocene Arctic, *Stratigraphy*, 6, 265–275, 2009.
- Robinson, M. M., Dowsett, H. J., Foley, K. M. and Riesselman, C. R.: PRISM marine sites: The history of PRISM sea surface temperature estimation, U.S. Geological Survey Open-File Report, 2018–1148, 49 pp., <https://doi.org/10.3133/ofr20181148>, 2018.
- 1435 Royer, C., Thébaud, J., Chauvaud, L. and Olivier, F.: Structural analysis and paleoenvironmental potential of dog cockle shells (*Glycymeris glycymeris*) in Brittany, northwest France. *Palaeogeogr. Palaeoclimatol.*, 373, 123–132, <https://doi.org/10.1016/j.palaeo.2012.01.033>, 2013.
- Schöne, B. R.: The curse of physiology—challenges and opportunities in the interpretation of geochemical data from mollusk shells, *Geo-Mar. Lett.*, 28, 269–285, <https://doi.org/10.1007/s00367-008-0114-6>, 2008.
- 1440 Schöne, B. R.: *Arctica islandica* (Bivalvia): A unique paleoenvironmental archive of the northern North Atlantic Ocean, *Global Planet. Change*, 111, 199–225, <https://doi.org/10.1016/j.gloplacha.2013.09.013>, 2013.
- 1445 Schöne, B. R., Fiebig, J., Pfeiffer, M., Gleß, R., Hickson, J., Johnson, A. L. A., Dreyer, W. and Oschmann, W.: Climate records from a bivalved Methuselah (*Arctica islandica*, Mollusca; Iceland), *Palaeogeogr. Palaeoclimatol.*, 228, 130–148, <https://doi.org/10.1016/j.palaeo.2005.03.049>, 2005.
- Schöne, B. R. and Fiebig, J.: Seasonality in the North Sea during the Allerød and Late Medieval Climate Optimum using bivalve sclerochronology, *Int. J. Earth Sci.*, 98, 83–98, <https://doi.org/10.1007/s00531-008-0363-7>, 2009.
- 1450 Shackleton, N. J.: Attainment of isotopic equilibrium between ocean water and the benthonic foraminifera genus *Uvigerina*: Isotopic changes in the ocean during the last glacial,

**Field Code Changed**

**Formatted:** Hyperlink, Font: (Default) +Body (Calibri), 11 pt

**Formatted:** Hyperlink, Font: (Default) Times New Roman, 12 pt, Font color: Auto, Pattern: Clear

**Formatted:** Font: (Default) Times New Roman, 12 pt

- Colloques Internationaux du Centre National de la Recherche Scientifique, 219, 203–209,  
1455 1974.
- Slupik, A. A., Wesselingh, F. P., Janse, A. C. and Reumer, J. W. F.: The stratigraphy of the  
Neogene–Quaternary succession in the southwest Netherlands from the Schelphoek  
borehole (42G4-11/42G0022)—a sequence stratigraphic approach, *Neth. J. Geosci.*, 86,  
317–332, <https://doi.org/10.1017/S0016774600023556>, 2007,
- 1460 Surge, D. and Barrett, J. H.: Marine climatic seasonality during medieval times (10th to 12th  
centuries) based on isotopic records in Viking Age shells from Orkney, Scotland,  
*Palaeogeogr. Palaeoclimatol.*, 350, 236–246, <https://doi.org/10.1016/j.palaeo.2012.07.003>,  
2012.
- Tebble, N.: *British Bivalve Seashells*, second ed., Her Majesty's Stationary Office, Edinburgh,  
1465 UK, 212 pp., 1976.
- Trofimova, T., Milano, S., Andersson, C., Bonitz, F. G. W. and Schöne, B. R.: Oxygen isotope  
composition of *Arctica islandica* aragonite in the context of shell architectural organization:  
Implications for paleoclimate reconstructions, *Geochem. Geophys. Geosy.*, 19, 453–470,  
<https://doi.org/10.1002/2017GC007239>, 2018.
- 1470 Ullmann, C. V., Wiechert, U. and Korte, C.: Oxygen isotope fluctuations in a modern North  
Sea oyster (*Crassostrea gigas*) compared with annual variations in seawater temperature:  
Implications for palaeoclimate studies, *Chem. Geol.*, 270, 170–176,  
<https://doi.org/10.1016/j.chemgeo.2010.07.019>, 2010.
- Valentine, A., Johnson, A. L. A., Leng, M. J., Sloane, H. J. and Balson, P. S.: Isotopic evidence  
1475 of cool winter conditions in the mid-Piacenzian (Pliocene) of the southern North Sea Basin,  
*Palaeogeogr. Palaeoclimatol.*, 309, 9–16. <https://doi.org/10.1016/j.palaeo.2011.05.015>, 2011.
- Vandenbergh, N., Herman, J., Laga, P., Louwe, S., De Schepper, S., Vandenbergh, J.,  
Bohncke, S. and Konert, W.: The stratigraphic position of a Pliocene tidal clay deposit at  
Grobbendonk (Antwerp Province, Belgium), *Geol. Belg.*, 3, 405–17,  
1480 <https://doi.org/10.20341/gb.2014.040>, 2000.
- Van-Vliet-Lanoë, B., Vandenbergh, N., Laurent, M., Laignel, B., Lauriat-Rage, A., Louwey,  
S., Mansy, J.-L., Mercier, D., Hallégouët, B., Laga, P., Laquement, F., Melliez, F., Michel,  
Y., Mougédet, G. and Villier, J.-P.: Palaeogeographic evolution of northwestern Europe  
during the Upper Cenozoic, in: *Messinia event: palaeobiological and  
1485 palaeoecological approaches*: edited by Néraudeau, D. and Goubert, É.,  
*Geodiversitas*, 24, 511–541, 2002.

- Vignols, R. M., Valentine, A. M., Finlayson, A. G., Harper, E. M., Schöne, B. R., Leng, M.J., Sloane, H. J. and Johnson, A.L.A.: Marine climate and hydrography of the Coralline Crag (early Pliocene, UK): isotopic evidence from 16 benthic invertebrate taxa, *Chem. Geol.*, 1490 536, 62–83, <https://doi.org/doi:10.1016/j.chemgeo.2018.05.034>, 2019.
- Wesselingh, F. P., Busschers, F. S. and Goolaerts, S.: Observations on the Pliocene sediments exposed at Antwerp International Airport (northern Belgium) constrain the stratigraphic position of the Broechem fauna, *Geol. Belg.*, 23, 315–321, <https://doi.org/10.20341/gb.2020.026>, 2020.
- 1495 Westaway, R., Maddy, D. and Bridgland, D.: Flow in the lower continental crust as a mechanism for the Quaternary uplift of south-east England: constraints from the Thames terrace record, *Quaternary Sci. Rev.*, 21, 559–603, [https://doi.org/10.1016/S0277-3791\(01\)00040-3](https://doi.org/10.1016/S0277-3791(01)00040-3), 2002.
- Williams, M., Haywood, A. M., Harper, E. M., Johnson, A. L. A., Knowles, T., Leng, M. J., 1500 Lunt, D. J., Okamura, B., Taylor, P. D. and Zalusiewicz, J.: Pliocene climate and seasonality in North Atlantic shelf seas, *Philos. T. R. Soc. A*, 367, 85–108, <https://doi.org/10.1098/rsta.2008.0224>, 2009.
- Williams, M., Nelson, A. E., Smellie, J. L., Leng, M. J., Johnson, A. L. A., Jarram, D. R., Haywood, A. M., Peck, V. L., Zalusiewicz, J., Bennett, C. and Schöne, B. R.: Sea ice extent 1505 and seasonality for the Early Pliocene northern Weddell Sea determined from fossil *Austrochlamys* bivalves, *Palaeogeogr. Palaeoclimatol.*, 292, 306–318, <https://doi.org/10.1016/j.palaeo.2010.04.003>, 2010.
- Winther, N. G. and Johannessen, J. A.: North Sea circulation: Atlantic inflow and its destination, *J. Geophys. Res.-Oceans*, 111, C12018, 1510 <https://doi.org/10.1029/2005JC003310>, 2006.
- Witbaard, R. and Bergman, M. J. N.: The distribution and population structure of the bivalve *Arctica islandica* L. in the North Sea: what possible factors are involved?, *J. Sea Res.*, 50, 11–25, [https://doi.org/10.1016/S1385-1101\(03\)00039-X](https://doi.org/10.1016/S1385-1101(03)00039-X), 2003.
- Wood, A. M., Wilkinson, I. P., Maybury, C. A. and Whatley, R.C.: Neogene, in: *Ostracods in British Stratigraphy*, edited by: Whittaker, J. E. and Hart, M. B., Spec. Publ., The Micropalaeontological Society, The Geological Society, London, UK, 411–446, 2009.
- Zhang, Z., Li, X., Guo, C., Otterå, O. H., Nisancioglu, K. H., Tan, N., Contoux, C., Ramstein, G., Feng, R., Otto-Bliesner, B. L., Brady, E., Chandan, D., Peltier, W. R., Baatsen, M.L.J., von der Heydt, A. S., Weiffenbach, J. E., Stepanek, C., Lohmann, G., Zhang, Q., Li, Q., 1520 Chandler, M. A., Sohl, L. E., Haywood, A. M., Hunter, S. J., Tindall, J. C., Williams, C.,

Lunt, D. J., Chan, W.-L. and Abe-Ouchi, A.: Mid-Pliocene Atlantic Meridional Overturning Circulation simulated in PlioMIP2, *Clim. Past*, 17, 529-543, <https://doi.org/10.5194/cp-17-529-2021>, 2021.