

1 Influence of long-term changes in solar irradiance forcing on the 2 Southern Annular Mode

3 Nicky M. Wright^{1,2,*}, Claire E. Krause^{1,3,+}, Steven J. Phipps⁴, Ghyslaine Boschat⁵, Nerilie J. Abram^{1,2}

4 ¹ Research School of Earth Sciences, Australian National University, Canberra ACT 2601, Australia

5 ² ARC Centre of Excellence for Climate Extremes, Australian National University, Canberra ACT 2601, Australia

6 ³ ARC Centre of Excellence for Climate System Science, Australian National University, Canberra ACT 2601, Australia

7 ⁴ Ikigai Research, Sandy Bay, Tasmania 7006, Australia

8 ⁵ Bureau of Meteorology and ARC Centre of Excellence for Climate Extremes, Melbourne, Victoria 3001, Australia

9

10 Correspondence to: Nicky M. Wright (nicky.wright@sydney.edu.au)

11 *Current address: School of Geosciences, University of Sydney, NSW 2006, Australia

12 +Current address: Geoscience Australia, Canberra ACT, Australia

13

14

15 **Abstract.** The Southern Annular Mode (SAM) is the leading mode of climate variability in the extratropical
16 Southern Hemisphere, with major regional climate impacts. Observations, reconstructions, and historical climate
17 simulations all show positive trends in the SAM since the 1960s; however, earlier trends in palaeoclimate SAM
18 reconstructions cannot be reconciled with last millennium simulations. There are also large differences in the
19 magnitude of solar irradiance change between various solar reconstructions, although most last millennium
20 climate simulations have relied on a low-amplitude solar forcing scenario. Here we investigate the sensitivity of
21 the SAM to solar irradiance variations using simulations with a range of constant solar forcing values, and last
22 millennium transient simulations with varying amplitude solar forcing scenarios. We find the mean SAM state can
23 be significantly altered by solar irradiance changes, and that transient last millennium simulations using a high-
24 amplitude solar scenario have an improved and significant agreement with proxy-based SAM reconstructions.
25 Our findings suggest that the effects of solar forcing on high-latitude climate may not be adequately incorporated
26 in most last millennium simulations, due to solar irradiance changes that are too small and/or the absence of
27 interactive atmospheric chemistry in the global climate models used for these paleoclimate simulations.

28

29

30 1 Introduction

31 The evolution of climate over the last millennium provides a unique setting for determining how modes of climate
32 variability respond to natural and anthropogenic forcing. The temporal evolution of external forcings (such as
33 atmospheric greenhouse gas concentrations, volcanic eruptions and changes in solar irradiance) is reasonably
34 well understood over the last millennium (Schmidt et al., 2011; Schmidt et al., 2012; Jungclaus et al., 2017),
35 allowing their effects on climate to be explored using global climate models. Such simulations have primarily
36 been compared to proxy-based reconstructions of global or hemispheric mean temperature (PAGES 2k-PMIP3

37 group; Neukom et al., 2018; Neukom et al., 2019), or analysed for changes in tropical or Northern Hemisphere
38 modes of climate variability (Ortega et al., 2015; Otto-Bliesner et al., 2016). There is considerably less
39 palaeoclimate data available in the Southern Hemisphere (Emile-Geay et al., 2017), and investigations into the
40 influence of external forcings on Southern Hemisphere climate variability are scarce. Despite this, evidence exists
41 that major changes in Southern Hemisphere climate variability occurred during the last millennium, including via
42 the Southern Annular Mode (SAM) (Abram et al., 2014; Dätwyler et al., 2018).

43 The SAM, also known as the Antarctic Oscillation (AAO), is the leading pattern of atmospheric variability
44 in the extratropical Southern Hemisphere. SAM variability describes changes in the strength and position of the
45 westerly wind belt (or mid-latitude westerly jet) over the Southern Ocean, and is represented by zonally opposing
46 geopotential height anomalies between the mid (~40°S) and high (~65°S) latitudes (Thompson and Wallace, 2000;
47 Marshall, 2003). A positive phase of the SAM is characterised by negative pressure anomalies over Antarctica
48 compared to positive pressure anomalies at mid-latitudes (Marshall, 2003), and a poleward contraction of the
49 westerly jet. Changes in the SAM have important impacts on temperature and precipitation across the Southern
50 Hemisphere, with particularly strong influences on weather across Australia, New Zealand, South America,
51 southern Africa and Antarctica (Gillett et al., 2006; Sen Gupta and England, 2006; Hendon et al., 2007). For
52 example, a positive SAM is associated with cool and wet conditions across most of southern Australia (excluding
53 Tasmania) (Gillett et al., 2006; Sen Gupta and England, 2006; Hendon et al., 2007; Fogt and Marshall, 2020), cool
54 and dry conditions across the Antarctic continent contrasted with warm and wet conditions along the Antarctic
55 Peninsula (Fogt and Marshall, 2020), and warm and dry conditions in New Zealand, Tasmania, and southern
56 South America (Gillett et al., 2006).

57 Characterisation of the SAM over the past century has relied on observational records and/or reanalysis
58 modelling (Marshall, 2003; Fan and Wang, 2004; Fogt et al., 2009; Visbeck, 2009). Short and sparse Antarctic
59 climate observations mean that SAM variability is directly measured only since 1957 (Marshall, 2003). Seasonal
60 SAM reconstructions from limited observations, primarily in the mid-latitudes, extend the instrumental record
61 back to 1865 for austral summer/autumn, and 1905 for winter (Fogt et al., 2009; Jones et al., 2009). Observations
62 and reanalyses have shown a robust positive trend in the SAM since the mid-20th Century that is most
63 pronounced in summer (Marshall, 2003; Fogt and Marshall, 2020), and has been primarily linked to the depletion
64 of stratospheric ozone (Thompson and Solomon, 2002; Gillett and Thompson, 2003; Son et al., 2009; Polvani et
65 al., 2011a; Thompson et al., 2011; Grise et al., 2013; Jones et al., 2016; Banerjee et al., 2020), with contributions
66 from increasing atmospheric greenhouse gases and internal variability/tropical decadal variability (e.g., Fyfe et al.,
67 1999; Kushner et al., 2001; Shindell and Schmidt, 2004; Arblaster and Meehl, 2006; Yang et al., 2020). During all
68 other seasons, the positive SAM trends have been mainly attributed to increasing greenhouse gases (year-round
69 trends) (Polvani et al., 2011b; Thompson et al., 2011). Analysis of historical simulations from the fifth Coupled
70 Model Intercomparison Project (CMIP5) has also found a significant response of the SAM in all seasons to solar
71 forcing, however the amplitude of this response is small compared to internal variability and anthropogenic
72 forcing, and unlikely to be identifiable in observations (Gillett and Fyfe, 2013). Future climate simulations suggest
73 that increasing greenhouse gases will cause further positive trends in the SAM (e.g., Wang and Cai, 2013; Gillett

74 and Fyfe, 2013; Goyal et al., 2021). In summer the trends are more uncertain as the changes in SAM will depend
75 on the opposing forcing from ozone recovery (Banerjee et al., 2020) and increasing greenhouse gases (Arblaster
76 et al., 2011; Meehl et al., 2012; Barnes and Polvani, 2013). The positive trend in austral summer SAM has paused
77 since the 2000s due to stratospheric ozone recovery resulting from the Montreal Protocol (Banerjee et al., 2020).

78 Palaeoclimate proxies (i.e., ice cores, tree-rings, stalagmites, corals, lake records, etc.) have been used
79 to reconstruct the SAM back through time, presenting a long-term picture of the natural variability of the SAM
80 prior to recent anthropogenic forcing (Abram et al., 2014; Dätwyler et al., 2018; Saunders et al., 2018; Villalba et
81 al., 2012). Proxy-based reconstructions indicate that the SAM experiences a large amount of natural variability
82 that may be intrinsic (unforced) or a response to natural external forcing, while SAM variability may also feedback
83 to force climate changes through modulating CO₂ outgassing from the Southern Ocean (Saunders et al., 2018)
84 and Antarctic sea ice extent (Crosta et al., 2021). During the last millennium, a minimum in the SAM index (i.e.,
85 negative SAM) occurred during the 1400s and a positive trend in the SAM (with superimposed interannual to
86 century-scale variability) is evident since this time (Fig. 1a). Anthropogenic forcing of the positive SAM trend since
87 the mid-20th Century has now moved the mean state of the SAM to its most positive state over at least the last
88 1000 years (Abram et al., 2014).

89 Climate simulations of the SAM response to rising atmospheric greenhouse gas levels and stratospheric
90 ozone depletion over the last century compare reasonably well to observations and proxy data (Miller et al., 2006;
91 Raphael and Holland, 2006; Swart and Fyfe, 2012; Gillett and Fyfe, 2013; Zheng et al., 2013). However, further
92 back in time, models are unable to reproduce the structure or magnitude of pre-industrial SAM trends that are
93 reconstructed from proxies (Abram et al., 2014). This could be due to errors in the reconstructions, or a
94 systematic issue in the way the SAM is forced or represented in current climate models (Abram et al., 2014;
95 Gillett and Fyfe, 2013). Visually, the temporal evolution of the reconstructed SAM over the last millennium
96 resembles some of the characteristics of long-term changes in solar irradiance over this time (Fig. 1a–c).

97 Variations in the solar constant (i.e., the rate at which energy reaches the Earth's surface from the sun)
98 have previously been suggested as an important driver of the SAM on observational time scales (Kuroda et al.,
99 2007; Kuroda and Kodera, 2005; Kuroda and Shibata, 2006; Roscoe and Haigh, 2007; Lu et al., 2011). The solar
100 constant varies naturally in an 11-year solar cycle, and recent work using changes in radiocarbon (¹⁴C) from
101 annually resolved and accurately dated tree rings confirms that the 11-yr cycle was present throughout the last
102 millennium (Brehm et al., 2021) (Fig. 1d). Other proxies (e.g., cosmogenic isotopes such as beryllium-10, ¹⁰Be)
103 also indicate that longer-term trends in solar forcing occur on century and millennial scales (Steinhilber et al.,
104 2009; Gray et al., 2010). During the last millennium, solar modulation reached a minimum during the Spörer
105 (1388–1558) and Maunder (1621–1718) Grand Solar Minima events.

106 Precise observations of solar irradiance (via satellites) are only available from 1978, and so
107 reconstructions of past changes in the magnitude of solar irradiance rely directly or indirectly on the relationship
108 between proxies (e.g., ¹⁴C, ¹⁰Be, sunspot properties) and solar irradiance, combined with processing and
109 calibration. Consequently, there are different published magnitudes for solar irradiance changes during the last
110 millennium (Fig. 1b), although the reconstructions are broadly similar in their temporal history of changes in solar

111 forcing. Many solar reconstructions (including those outlined in the Paleoclimate Modelling Intercomparison
112 Project, phase 3 (PMIP3) v1.0 protocol; Schmidt et al., 2011) have a $\sim 1.5 \text{ W m}^{-2}$ peak to peak variation in total
113 solar irradiance (TSI) between present day and the Maunder Minimum period (Judge et al., 2012). A high-
114 amplitude ($\sim 6 \text{ W m}^{-2}$ peak-to-peak variation) solar reconstruction (Shapiro et al. 2011) was also included as part
115 of the PMIP3 v1.1 protocol (Schmidt et al., 2012), but it has been argued that this reconstruction overestimates
116 the changes in TSI by a factor of two due to the model used in their methodology (Judge et al., 2012). Phase 4 of
117 the Paleoclimate Modelling Intercomparison Project (PMIP4) provides an alternative solar irradiance scenario
118 ('PMOD') derived using a similar approach to Shapiro et al. (2011) and Judge et al. (2012), with a Maunder
119 Minimum to present amplitude of $\sim 3.4 \text{ W m}^{-2}$ (Jungclaus et al., 2017): almost half the amplitude of Shapiro et al.
120 (2011), though still considerably larger than alternative solar reconstructions for the last millennium. This PMOD
121 reconstruction is currently considered to be the 'upper limit' on the magnitude of solar irradiance change
122 (Jungclaus et al., 2017). Existing last millennium climate simulations have almost exclusively been run using the
123 low amplitude solar forcing scenarios (e.g., Steinhilber et al., 2009) and in model set ups that do not
124 accommodate solar-relevant atmospheric chemistry or wavelength specification. This raises the possibility that
125 the effects of solar forcing may not have been adequately included in last millennium simulations, potentially
126 accounting for data-model SAM discrepancies.

127 Here we explore the sensitivity of the SAM to variations in solar forcing, in an attempt to understand if
128 variations in solar irradiance may have affected SAM variability over the last millennium (Fig. 1). We explore
129 simulations with constant solar forcing values that correspond to the range of total solar irradiance values from
130 the high amplitude solar reconstruction of Shapiro et al. (2011), as well as additional extreme solar forcing values.
131 In addition, we investigate transient simulations for the last millennium using intermediate and high amplitude
132 solar forcing scenarios that complement existing low amplitude transient solar forcing experiments. Our findings
133 demonstrate that the mean state of the SAM can be significantly altered by changes in solar irradiance, and that
134 transient solar forcing of a magnitude equivalent to high amplitude scenarios for the last millennium are sufficient
135 for a significant solar effect on the SAM to become evident despite the large magnitude of internal SAM
136 variability. Last millennium simulations using high amplitude solar forcing show an improved agreement with
137 proxy-based SAM reconstructions, suggesting that the effects of solar forcing may not be adequately
138 represented in current last millennium climate model simulations.

139

140

141 **2 Methods**

142 **2.1 Reconstructions**

143 The last millennium reconstructions that we use in this study are for the annual mean SAM index. These
144 are based on (i) a multiproxy network spanning Antarctica and South America where the temperature anomalies
145 caused by SAM variations are strong (Abram et al., 2014), and (ii) an extensive network of proxies from across the
146 Southern Hemisphere, using a long calibration period and a correlation plus stationarity criterion for proxy
147 selection (Dätwyler et al., 2018). Hereafter, these SAM reconstructions are referred to as A14 (Abram et al., 2014)

148 and D18 (Dätwyler et al., 2018). The A14 and D18 reconstructions share similar features in their long-term trends
149 (Fig. 1a), despite potential regional biases in the proxy networks used for the reconstructions and uncertainty
150 related to non-stationary proxy-SAM relationships (Huiskamp and McGregor, 2021; Hessel et al., 2017). Both
151 reconstructions indicate that the most negative phase of SAM conditions occurred during the 1400s prior to a
152 progressive, multi-century positive trend in the SAM since the 1400s including the rapid 20th Century increase in
153 the SAM. Both reconstructions also record strong interannual to centennial variability on top of these long-term
154 trends, with this characteristic particularly evident in the A14 reconstruction.

155 The A14 and D18 reconstructions were both developed using the instrumental SAM index as a
156 calibration target (i.e., Marshall, 2003, and Fogt et al., 2009, respectively); however, the A14 reconstruction
157 displays a larger magnitude of variability (Fig. 1a). This difference in magnitude is due to differences in the way
158 the annual SAM index can be calculated from instrumental data (Fig. 2). For example, the instrumental SAM index
159 (Marshall, 2003; <http://www.nerc-bas.ac.uk/icd/gjma/sam.html>) is publicly available in both monthly and annual
160 resolutions. Here, the annual resolution SAM index is calculated directly using the differences of annual means of
161 mean sea level pressure (MSLP) data at 40°S and 65°S. Alternatively, the monthly resolution SAM data
162 (calculated from differences of monthly means of MSLP data at 40°S and 65°S) can be used to then calculate
163 annual averages of the SAM. The two approaches result in similar trends and interannual variability of the SAM;
164 however, the magnitude of the directly calculated annual SAM index is 2.7-times larger than the annual mean
165 SAM derived from the monthly SAM index (Fig. 2).

166 The A14 and D18 SAM reconstructions both use an annual SAM index as their calibration target, but A14
167 used a calibration annual mean SAM index calculated from annual data (i.e., red line in Fig. 2), while D18 used a
168 calibration annual mean SAM index calculated from monthly data (e.g., orange line in Fig. 2). Consequently, while
169 the two reconstructions produce similar patterns and trends of annual SAM variability during the last millennium,
170 they have markedly different magnitudes of change due to differences in the instrumental calibration data used
171 (Fig. 3a). To confirm this, we recalculate the A14 reconstruction using the alternate calibration target (i.e.,
172 calibrated to the annual SAM index derived from monthly SAM data, as in the D18 reconstruction). This rescaled
173 reconstruction (referred to hereafter as A14-rescaled) has interannual to centennial variability and trends in the
174 last millennium that are of similar magnitude as the D18 reconstruction (Fig. 3b), confirming that the source of
175 apparent discrepancy between the A14 and D18 reconstructions is primarily due to the different instrumental
176 targets used to calibrate the reconstructions.

177 In this study we use the A14, D18 and A14-rescaled SAM reconstructions as indicators of temporal
178 changes in the mean state of SAM during the last millennium. These are used in data-model comparisons to
179 determine if different solar-forced model simulations are able to reproduce characteristics of reconstructed SAM
180 changes during the last millennium.

181

182 **2.2 Model Simulations**

183 The model simulations in this study were performed using the Commonwealth Scientific and Industrial Research
184 Organisation Mark 3L (CSIRO Mk3L) coupled climate system model, version 1.2 (Phipps et al., 2011, 2012,

185 2013). CSIRO Mk3L is a fully coupled general circulation model that includes components describing the
186 atmosphere, ocean, sea ice, and land surface. The horizontal resolution of the atmosphere, sea ice, and land
187 surface models is $5.6^\circ \times 3.2^\circ$ in the longitudinal and latitudinal dimensions, respectively, with 18 vertical levels.
188 The horizontal resolution of the ocean model is $2.8^\circ \times 1.6^\circ$, with 21 vertical levels. CSIRO Mk3L was used within
189 this study as it is computationally efficient, allowing for many multi-century to millennia-scale experiments to be
190 performed.

191 We investigate how the SAM changes in a series of ‘solar constant’ experiments and transient
192 experiments. Specifically, we perform:

193 (a) Seven solar constant experiments, where the solar constant (i.e., TSI, with no wavelength
194 dependence) was changed to capture the range of proxy-based realistic solar values in the Shapiro
195 reconstruction (-7 , -3 , $+1$ and $+3 \text{ W m}^{-2}$ anomalies relative to a 1365 W m^{-2} control), as well as unrealistic
196 extreme solar forcings (e.g., -15 , $+7$ and $+35 \text{ W m}^{-2}$ anomalies) to test the response of the model (Fig. 1c). These
197 experiments use preindustrial CO_2 (280 ppm) and we run each experiment to equilibrium. In our analysis of the
198 model output we primarily focus on the transient response of the SAM to the changed solar constant within the
199 first 200 years of each run (Fig. 4), where the climate has not yet equilibrated to the new solar forcing. These
200 experiments are referred to using their solar constant anomaly value (e.g., S-7, S+3, etc.)

201 (b) Three-member ensembles of transient experiments using three solar forcing scenarios for the last
202 millennium. These transient experiments were initialised from the control used in Phipps et al. (2013) (refer to
203 (Phipps et al., 2013) for further details), and all include orbital, greenhouse gas, and solar forcings. The first
204 ensemble is the orbital-greenhouse gases-solar (‘OGS’) ensemble first published in Phipps et al. (2013), which
205 uses the Steinhilber et al. (2009) solar forcing. We build upon this ‘OGS’ ensemble by performing additional
206 experiments with modified solar forcing (Fig. 1b). The second ensemble of transient experiments uses an
207 amplified Steinhilber et al. (2009) forcing, where the magnitude of the transient solar forcing anomaly relative to
208 the long-term mean is doubled (hereafter referred to as ‘OGS-x2’). The third ensemble of transient experiments
209 uses the Shapiro et al. (2011) high amplitude solar forcing that was included as a last millennium forcing option in
210 the PMIP3 v1.1 protocol (Schmidt et al., 2012) and is hereafter referred to as ‘OGS-Shapiro’. The CSIRO Mk3L
211 model does not include interactive chemistry, nor do we prescribe ozone variations scaled to the solar forcing in
212 our experiments in an attempt to replicate the response of atmospheric chemistry during the last millennium.
213 Each experiment was run for 1–2000 CE. Here, we focus our analysis to 850–1900 CE, to avoid the influence of
214 strong anthropogenic greenhouse gas forcings after 1900.

215 We supplement our transient experiments with CSIRO Mk3L using previously published simulations
216 using the HadCM3 model (‘Euroclim500’; Schurer et al., 2014). These simulations include one full-forcing
217 ensemble member for 800–2000 CE, three full-forcing ensemble members for 1400–2000 CE, four ‘weak solar’
218 only ensemble members for 1400–2000 CE, and one ‘solar Shapiro’ only run for 800–2000 CE using the solar
219 reconstruction from Shapiro et al. (2011). Forcings used in the HadCM3 experiments follow the PMIP3 protocol
220 (Schmidt et al., 2011; Schmidt et al., 2012); specifically, the solar reconstruction used in the full-forcing and
221 ‘weak solar’ runs is based on a combination of Steinhilber et al. (2009) (for times prior to 1810) and Wang et al.

222 (2005) (for 1810–2000) (see Schurer et al., 2014, for further details). We note that both our CSIRO Mk3L
223 experiments and these HadCM3 experiments do not have an interactive ozone.

224 The SAM index was calculated for each model run following the definition from Gong and Wang (1999):

225 $SAM = P^*_{40^{\circ}S} - P^*_{65^{\circ}S}$, where P^* is the normalised annual zonal mean MSLP anomaly in the model relative to

226 climatology. For the solar constant experiments, we use the solar constant (1365 W m^{-2}) control run as our

227 climatology. This allows us to assess the effect of modifying the solar constant anomaly on the SAM mean state

228 in the other solar constant experiment relative to the 1365 W m^{-2} control. We use a Welch's t -test and

229 Kolmogorov–Smirnov test to assess the difference between our solar constant experiments and the control run.

230 In the transient experiments we use the 1900–1999 period as our climatology to assess pre-industrial changes in

231 the SAM mean state over the last millennium, and use a Wilcoxon rank-sum test, linear least-squares regression,

232 and bootstrapping approach to compare our transient experiments with its radiative forcing and the SAM

233 reconstructions.

234

235

236 **3 Results**

237 **3.1 Solar constant experiments**

238 The solar constant experiments demonstrate how changes in the intensity of solar forcing influence the SAM

239 mean and extreme states (Fig. 5, Fig. 6). The distribution of the mean annual SAM index is significantly different

240 from the control in the S–15, S–7, and S+35 experiments (based on Welch's t -test, $P < 0.05$, see Fig. 5a). In

241 particular, changing the solar constant results in a mean shift in the SAM but no significant difference in the

242 magnitude of SAM variability about this mean shift (using a Kolmogorov–Smirnov test, $P > 0.05$) (Fig. 5a). This

243 mean shift in the SAM results in a change in the number of extreme events (outside $\pm 2\sigma$ of the control run),

244 whereby a reduced solar forcing (i.e., S–15, S–7, and S–3) results in an increase in extreme negative SAM events

245 and decrease in extreme positive SAM events, and vice versa for the increased solar forcing experiments (Fig.

246 5b).

247 The spatial patterns of anomalies in the solar constant experiments demonstrate the consistent

248 influences that changes in solar intensity have on the SAM and Southern Hemisphere climate (Fig. 5c–h).

249 Negative solar forcing anomalies result in a decrease in MSLP over the Southern Hemisphere mid-latitudes and

250 an increase in MSLP over the Antarctic, and thus a reduced meridional pressure gradient between these zones

251 (Fig. 5c, f–g). This is associated with an enhancement of the surface temperature gradient between the mid-

252 latitudes and Antarctica (Fig. 5d; i.e., Antarctica cools more than the mid-latitudes) and decreased westerly wind

253 anomalies in the Southern Ocean jet (Fig. 5e), leading to a mean negative shift in the SAM index and an increase

254 in the number of extreme negative SAM events relative to extreme positive events. The opposite is also true with

255 positive solar forcing, which leads to increased MSLP over the mid-latitudes, decreased MSLP over Antarctica,

256 and consequently strengthens the meridional pressure gradient and the westerly wind jet. This results in a more

257 positive mean SAM with an increase in the frequency of extreme positive SAM events. We note that there is an

258 asymmetry in the effect of changing the solar constant, with a larger response seen in the negative solar constant

259 experiments (i.e., S-3, S-7) than the positive solar constant anomalies (i.e., S+3, S+7) (Fig. 5a-e). The solar
260 constant experiments also show that the magnitude of MSLP change over the Antarctic (65°S) is larger than over
261 the mid-latitudes (40°S) in response to changes in solar forcing (Fig. 5c).

262

263 **3.2 Transient experiments for the last millennium**

264 The transient simulations build upon our findings from the solar constant experiments by modelling the time
265 evolution of the SAM index during the last millennium based on different amplitudes of transient solar forcing (Fig.
266 7, Fig. 8). The simulations also include transient orbital and greenhouse gas forcing, and so we express the
267 results using radiative forcing (i.e., the combined radiative forcing of orbital, greenhouse gas, and solar; instead of
268 solar irradiance only) and focus on pre-industrial times (i.e., prior to 1900). The ensemble mean SAM index from
269 the low solar amplitude OGS experiments (Phipps et al., 2013) is not significantly correlated with the radiative
270 forcing. This indicates that the low amplitude solar forcing in these simulations is not large enough to modulate
271 the mean SAM state in a way that is detectable beyond the magnitude of unforced internal SAM variability. We
272 additionally find that the OGS-ensemble mean is largely within the range of unforced internal SAM variability
273 during the last millennium (Fig. 7c), where our internal variability is represented using the $\pm 2\sigma$ range from the
274 orbital ('O') only simulations (Fig. 7a). We further assess this relationship by applying a bootstrapping approach to
275 randomly reorder the OGS ensemble mean SAM ($N = 10000$), and find that the OGS SAM index is not
276 significantly correlated with its radiative forcing any more than could be explained by a random distribution of the
277 model data. However, the ensemble mean SAM index from the intermediate solar amplitude OGS-x2
278 experiments is positively correlated with radiative forcing for pre-industrial times ($r = 0.43$, $P < 0.05$, effective
279 sample size $N_{\text{eff}} = 15.8$; based on 70-yr moving averages stepped by 35 years, with effective sample size taking
280 into account lag-1 autocorrelation of the time series; Fig. 8a-b). The OGS-x2 ensemble mean also exceeds the
281 range of internal variability during some intervals of the last millennium, in particular, during the 15th Century,
282 where the SAM is more negative than can be explained by internal variability alone (Fig. 7d). However, for
283 individual ensemble members with intermediate amplitude solar forcing, we cannot reject the null hypothesis that
284 there is no correlation. Using a bootstrapping approach ($N = 10000$) to our OGS-x2 ensemble mean SAM index
285 further finds a significant correlation with its radiative forcing ($P < 0.05$), relative to a random distribution of the
286 model data.

287 In contrast, the SAM index for the high amplitude OGS-Shapiro simulations is significantly correlated
288 with the radiative forcing anomaly (Fig. 8c). The correlation coefficient between radiative forcing and the SAM
289 index in the ensemble mean is 0.64 ($P < 0.05$, $N_{\text{eff}} = 14.4$) for pre-industrial times (i.e., 850-1900). The correlation
290 between the OGS-Shapiro ensemble mean SAM index and its radiative forcing remains significant ($P < 0.05$;
291 relative to a random distribution of the model data) when tested against a bootstrapping approach ($N = 10000$).
292 Each of the individual ensemble members in the high amplitude solar experiments also displays a significant
293 long-term SAM-radiative forcing relationship over the last millennium, demonstrating a forced signal detectable
294 beyond the large range of internal SAM variability (Fig. 7e).

295 We explore this relationship further by binning the model results across all of the transient experiments
296 based on the magnitude of radiative forcing (Fig. 9). Increasingly negative radiative forcing anomalies result in an
297 increasingly negative SAM index (Fig. 9a–b). Binning across all ensemble members based on radiative forcing
298 anomalies further illustrates the linear relationship between reduced radiative forcing and a more negative mean
299 SAM (Fig. 9b). The zonal mean MSLP and temperature profiles (Fig. 9c–d) and spatial structure of MSLP
300 anomalies (Fig. 9e–h) are consistent with the anomalies produced in the solar constant experiments (Fig. 5). This
301 suggests a consistent response of mid to high-latitude Southern Hemisphere climate to changes in solar forcing
302 within the CSIRO Mk3L experiments, that is also robust across different experiment designs.

303 Overall, our experiments show that a decrease in solar radiative forcing (in both the constant solar and
304 transient solar forcing runs) results in a mean negative SAM shift. Reducing solar forcing by $1.5\text{--}1.0\text{ W m}^{-2}$ is
305 roughly equivalent to a $\sim 7\text{ W m}^{-2}$ reduction in the solar constant (based on scaling the change in TSI by 0.7/4;
306 Lean and Rind, 1998). Both the S–7 fixed solar constant experiment and high amplitude transient radiative
307 forcing of -1.5 to -1.0 W m^{-2} result in a statistically significant negative anomaly in the SAM that is detectable
308 despite the large magnitude of unforced internal variability of the SAM. These simulation results with the CSIRO
309 Mk3L model suggest that reconstructed negative SAM conditions during the last millennium could have been the
310 result of reduced solar forcing at this time.

311

312

313 **4 Comparison with reconstructions**

314 Previous work using low amplitude solar forcing experiments has not found any significant relationship between
315 solar forcing and reconstructed long-term changes in the annual SAM during the last millennium (Abram et al.,
316 2014; Dätwyler et al., 2018). However, extreme changes in solar forcing in our model experiments that are
317 comparable with high amplitude estimates of solar irradiance anomalies during the last millennium (Shapiro et al.,
318 2011) are able to produce a significant change in the mean SAM index, where a -7 W m^{-2} change in solar
319 irradiance (or a roughly -1.23 W m^{-2} change in radiative forcing) results in a significant negative shift in the mean
320 SAM index. To explore further whether solar forcing may help to explain reconstructed trends in the SAM during
321 the last millennium, we compare our model results with proxy-based SAM reconstructions (see Section 2.1).

322 To test the significance of changes in the SAM during the last millennium, we use each of the SAM
323 reconstructions to assess whether 70-yr sliding windows of the annual SAM reconstructions are significantly
324 ($P < 0.05$; Wilcoxon rank-sum test) more negative than a 70-yr reference window of the reconstruction between
325 1831–1900. This reference interval was chosen as it is prior to strong positive SAM trends caused by
326 anthropogenic forcing during the 20th Century, and is longer than the standard 50-year preindustrial period so as
327 to improve the robustness of the distribution testing. These tests show that across all three reconstructions the
328 SAM index was significantly more negative between approximately 1390 and 1715 CE compared with the 1831–
329 1900 reference interval (Fig. 10). The A14 and A14-rescaled reconstructions also indicate significant negative
330 SAM distributions prior to around 1140 CE (Fig. 10).

331 Carrying out the same distribution tests on the CSIRO Mk3L transient simulations (Fig. 11) shows that
332 there are no significant negative shifts of the SAM index during the last millennium in the OGS ensemble mean
333 with low amplitude solar forcing. By comparison, the OGS-Shapiro ensemble mean shows a strong and
334 sustained negative SAM shift that peaked at approximately 1460 CE. This is in good agreement with the interval
335 where all SAM reconstructions also display a significant negative shift in the SAM. The OGS-Shapiro ensemble
336 mean also shows earlier intervals where the SAM index is significantly more negative than the 1831–1900
337 reference interval. Exact matches in the start and end times of significant negative shifts caused by solar forcing
338 in the SAM simulations and reconstructions are not expected due to the additional effect of large unforced
339 internal variability in the SAM (i.e., as seen in differences between ensemble members runs with the same solar
340 forcing). However, we do find that the maximum significance in negative SAM distribution shifts in the OGS-
341 Shapiro ensemble mean is around 1460 CE (Fig. 11d), which matches the timing of maximum significant shifts in
342 all of the last millennium SAM reconstructions (Fig. 10d; approximately 1415–1560 CE). We also find a strongly
343 significant negative shift in the simulated SAM index peaking at around 1025 CE (Fig. 11d), which coincides with
344 the reconstructed significant shift in the A14 and A14–rescaled reconstructions prior to 1140 CE (Fig. 10).

345 Direct correlation of the reconstructions and transient simulations further shows that the A14 proxy-
346 based SAM reconstruction shares significant ($P < 0.05$) variance during the pre-industrial last millennium (1000–
347 1900 CE) with the ensemble mean SAM index of the transient solar scenario simulations run with CSIRO Mk3L
348 (Fig. 12a). The increasing strength of the correlations with increasing magnitude of solar forcing indicates
349 improved coherence between the multi-decadal variability and long-term trends of the reconstructed and
350 modelled SAM when strong solar forcing is used over the last millennium. There are differences in the shorter-
351 term details of the modelled and reconstructed SAM indices, such as the timing during the 1400s when the most
352 negative SAM conditions are reached. However, these differences are of a comparable magnitude to the internal
353 variability between ensemble members of the same experiment (Fig. 8) and so may simply represent differences
354 between realisations (including the reconstructed single real-world realisation) in unforced variability of the SAM
355 on top of the solar-forced variability and trends.

356 We further explore the robustness of the correlation of the simulated SAM and the A14 proxy-based
357 SAM reconstruction by using a bootstrapping approach to randomly reorder the ensemble mean SAM indices
358 from our transient solar forcing experiments ($N = 10000$). Based on this, we find that the OGS simulation is not
359 significantly correlated with the A14 reconstruction any more than could be explained by a random distribution of
360 simulated data. However, we find that both the OGS-x2 and OGS-Shapiro ensemble mean SAM index are
361 significantly correlated to the A14 reconstruction ($P < 0.05$; relative to a random distribution of model data).

362 The D18 SAM reconstruction during the pre-industrial last millennium is not significantly correlated with
363 the ensemble mean SAM index of any of the CSIRO Mk3L transient solar forcing experiments (Fig. 13a). Based
364 on a bootstrapping approach ($N = 10000$), the OGS-Shapiro ensemble mean SAM index is significantly correlated
365 ($P < 0.05$; relative to a random distribution of simulated data) to the D18 reconstruction, while there is no
366 significant correlation between the D18 reconstruction and the OGS and OGS-x2 ensemble mean SAM index.
367 This appears to be mostly related to differences between the modelled and reconstructed SAM indices prior to

368 1400. This also corresponds to an apparent increase in the magnitude of noise within the D18 reconstruction
369 prior to 1400 (Fig. 10c), and a reduction of reconstruction skill (negative reduction of error [RE]) for the D18
370 reconstruction prior to 1400 (Dätwyler et al., 2018), and may reflect reduced reconstruction fidelity due to the
371 sparse proxy network during the early stages of the last millennium. Visually, the maximum negative SAM
372 anomaly during the 1400s and the long-term positive trend since that time appears to correspond well between
373 the transient simulations and the D18 reconstruction, particularly for the strong amplitude solar forcing
374 experiments (Fig. 13a).

375

376 **5 Discussion**

377 There have so far been few studies exploring the influence of high amplitude changes in solar forcing during the
378 last millennium. Research by Schurer et al. (2014) found little influence of stronger solar variability on Northern
379 Hemisphere temperature reconstructions of the last millennium using the HadCM3 model. However, a
380 comparison of the Shapiro et al. (2011)-solar forced run in HadCM3 (Schurer et al., 2014) and our OGS-Shapiro
381 simulation show some similarities in the long-term trends of the SAM index (Fig. 12). In particular, both CSIRO
382 Mk3L and HadCM3 show a large negative excursion in the SAM index around 1450 CE in their strong solar
383 forcing simulations (Fig. 12, purple and red time series in Fig. 12a and 12b, respectively). The transient strong-
384 solar forcing simulation from HadCM3 has a significant ($P < 0.05$) correlation with both proxy-based SAM
385 reconstructions ($r = 0.48$ for A14, and $r = 0.73$ for D18). In contrast, last millennium correlations are not significant
386 for the SAM reconstructions with the ensemble mean SAM index in the HadCM3 weak-solar forcing only
387 simulation and full-forcing (including weak-solar forcing) simulations (Fig. 12b, Fig. 13b). The HadCM3 strong
388 solar forcing experiment thus corroborates our findings using CSIRO Mk3L, indicating that high amplitude solar
389 forcing of last millennium simulations has a detectable effect on the annual SAM that improves the agreement
390 between modelled realisations of the SAM index and proxy-based SAM reconstructions.

391 Solar activity is thought to influence climate, the SAM, and its Northern Hemisphere equivalent—the
392 Northern Annular Mode (NAM)—through either ‘top-down’ (e.g., via changes associated with stratospheric-
393 tropospheric coupling due to ozone and UV-related stratospheric temperature and wind variations; Gray et al.,
394 2010) or ‘bottom-up’ mechanisms (e.g., through associated changes in sea surface temperature [SST] and ocean
395 heat uptake; Gray et al., 2010; Meehl et al., 2008). Notably, resolving the former mechanism requires climate
396 model simulations using interactive stratospheric chemistry that are computationally expensive to run (and not
397 currently feasible for the last millennium), while ‘bottom-up’ drivers do not require a well-resolved stratosphere or
398 changes in stratospheric ozone (Meehl et al., 2008). Studies based on observations and/or chemistry-enabled
399 climate models have previously suggested that the SAM (and the NAM) is sensitive to changes in solar activity
400 associated with the 11-year solar cycle (Kuroda et al., 2007; Kuroda and Kodera, 2005; Kuroda and Shibata,
401 2006; Gray et al., 2010; Gray et al., 2013; Ma et al., 2018; Kuroda, 2018; Arblaster and Meehl, 2006). These
402 studies link variations in the SAM/NAM to changes in stratospheric temperatures and/or stratospheric-
403 tropospheric coupling. For example, during years of higher solar activity, there is a stratospheric extension of the
404 SAM signal (Kuroda and Kodera, 2005), related to a corresponding increase in stratospheric-tropospheric

405 coupling (Kuroda et al., 2007). A similar process is seen for the NAM, though there may be a 2–4 year lagged
406 response of positive NAM following a solar high (Gray et al., 2013; Ma et al., 2018). Solar activity incites variations
407 in stratospheric temperature and winds related to changes in UV irradiance and ozone production, while
408 associated variations in stratospheric-tropospheric coupling result in changes in surface climate (e.g., a ‘top-
409 down’ forcing) (Gray et al., 2010). This varies from proposed ‘bottom-up’ mechanisms for solar variations
410 influencing surface climate, which involve changes in SST and ocean heat uptake during periods of increased
411 solar activity, resulting in an increase in latent heat flux and evaporation, which ultimately leads to intensified
412 precipitation along convergence zones and stronger trade winds (Meehl et al., 2008; Meehl et al., 2003; Gray et
413 al., 2010), as well as less heating over the ocean than land (Meehl et al., 2003). Solar forcing may also influence
414 climate through a combination of both top-down and bottom-up mechanisms (Rind et al., 2008), and simulations
415 with both mechanisms working together result in a stronger tropical SST response more similar to observations
416 than simulations with only a single mechanism (Meehl et al., 2009).

417 As the models examined in this study do not have a well-resolved stratosphere or incorporate interactive
418 stratospheric ozone, we suggest our simulated changes in the SAM are caused by a ‘bottom-up’ mechanism. We
419 find comparable changes in our increased solar constant experiments to previous studies invoking a bottom-up
420 mechanism, such as an increase in equatorial evaporation and precipitation, as well as a greater increase in land
421 surface temperatures than over the ocean (Fig. 6). Within the atmosphere, our experiments show an increase in
422 solar forcing causes an increase in temperature throughout the tropics, with a larger increase in the upper
423 troposphere than surface, as well as cooling in the high-latitude upper troposphere and a reduced warming in the
424 lower troposphere along 40°–60°S (Fig. 14b). This increase in temperature is combined with a westerly wind
425 anomaly that spans from the tropical upper troposphere to the high latitudes (~50°S) and extends into the lower
426 troposphere (Fig. 14a)—all of which are similar to the climate effects simulated from an increase in radiative
427 forcing by greenhouse gases (e.g., Kushner et al., 2001; Lim and Simmonds, 2009; Butler et al., 2011). The
428 poleward contraction in the westerly jets around 55°S is particularly pronounced in the S+7 and S+35 scenarios,
429 as is the increased meridional temperature gradient (e.g., tropics warming faster than the Southern Ocean),
430 leading to the development of a mean positive SAM state. A decrease in solar forcing results in an approximately
431 inverse pattern, with cooling in the tropical upper troposphere, warming in the high-latitude upper troposphere,
432 and more cooling in the tropical lower troposphere, with minimal cooling in the high latitudes (Fig. 14b), as well as
433 a decrease in the zonal wind anomaly across the tropical–high latitude upper troposphere and into the high
434 latitude lower troposphere (Fig. 14). Overall, this indicates a weakening of the westerly jet (i.e., negative zonal
435 wind anomaly around ~50°S). Solar forcing is primarily shortwave and varies seasonally and spatially, with
436 greater influence in the tropics, while greenhouse gas forcing is more spatially uniform (Meehl et al., 2003). While
437 solar forcing affects climate differently to greenhouse gas forcing, it is possible that we find in our solar
438 experiments a broadly similar tropospheric response as expected from greenhouse gases due to the extremely
439 large magnitude changes in our solar forcing experiments combined with exploring transient mean state changes
440 over only the first 200 years (Fig. 4) of the solar constant experiments.

441 Overall, our findings do suggest that the effects of solar forcing on the SAM are not adequately
442 represented in current last millennium climate simulations. It is possible that the reconstructed minimum in the
443 SAM during the 15th Century was a response to a minimum in solar irradiance at this time, and that this solar
444 response is not reproduced in last millennium simulations that are forced with low amplitude solar forcing.
445 However, our results do not necessarily imply that solar forcing of the last millennium involved the large
446 amplitude changes of the Shapiro et al. (2011) forcing scenario. The large amplitude solar forcing used in our
447 transient experiments was a plausible forcing scenario provided as an option for last millennium experiment
448 design of the Coupled Model Intercomparison Project (Schmidt et al., 2012). But this strong solar forcing
449 scenario (i.e., Shapiro et al., 2011) has rarely been applied in last millennium simulations, and it has been argued
450 that the magnitude of TSI change in Shapiro et al. (2011) have been overestimated by a factor of two (Judge et
451 al., 2012). Instead, it may be that in climate models that do not have the interactive atmospheric chemistry
452 needed to permit solar-impacts on stratospheric ozone, a large amplitude of solar forcing is instead needed to
453 reproduce the last millennium climate impacts on the SAM that were caused by more modest solar forcing
454 changes. While we cannot rule out the possibility that the strength of the forcing allows our experiments to
455 reproduce changes in the SAM without a realistic representation of all the forcings involved, our strong solar
456 forcing experiments produce a SAM response that better replicates reconstructed changes in the SAM during the
457 last millennium.

458

459

460 **6 Conclusion**

461 Palaeoclimate reconstructions indicate large changes in the SAM during pre-industrial times that are not
462 replicated in current last millennium climate simulations. We explore changes in solar forcing on the SAM using
463 solar constant and last millennium transient simulations that cover large amplitude solar changes and find that
464 the SAM index significantly decreases (increases) with a decrease (increase) in solar forcing. The magnitude of
465 solar forcing change required for a significant change in the SAM index is much greater than the most commonly
466 used solar forcing scenarios for the last millennium, with an approximately 7 W m^{-2} decrease in total solar
467 irradiance (or 1.5 W m^{-2} decrease in radiative forcing) required before the effect of solar forcing on the SAM can
468 be distinguished from the large range of unforced SAM variability. Transient simulations of the last millennium
469 with strong solar forcing result in an improved and significant agreement with proxy-based reconstructions of the
470 SAM. It is plausible that solar forcing may have been an important driver in long-term SAM trends prior to the
471 strong anthropogenic forcing of the 20th and 21st centuries, and that current climate model simulations of the
472 last millennium do not adequately represent the effect of solar variability on mid to high latitude Southern
473 Hemisphere climate. This may be due to a higher magnitude of solar irradiance changes than is usually applied in
474 last millennium simulations, or (more likely) due to the absence of important physical and chemical processes in
475 coupled global climate models that would allow more moderate changes in solar forcing to have a discernible
476 impact on high latitude climate.

477

478 **7 Acknowledgements**

479 We thank the Australian Research Council for support of this research through the Centre of Excellence for
480 Climate Extremes (CE170100023)), Discovery Project (DP140102059), Future Fellowship (FT160100029) and the
481 Australian Centre for Excellence in Antarctic Science (SR200100008). This work was also supported by the
482 Climate Systems Hub of the Australian Government's National Environmental Science Program, and was made
483 possible by computational resources provided by the Australian Government through the National Computational
484 Infrastructure, including a grant (xf4) through the ANU merit allocation scheme.

485

486

487 **8 Author contributions**

488 N.J.A. devised the study. C.E.K. performed the solar constant experiments, and N.M.W. and S.J.P. performed the
489 transient simulations. N.M.W. and N.J.A. performed the data analyses with help from G.B., and N.M.W. made the
490 figures. N.M.W. wrote the manuscript, with contributions from all co-authors.

491

492

493 **9 Competing interests**

494 The authors declare no competing interests.

495

496

497 **10 Data availability**

498 Data will be archived in a public repository upon acceptance.

499 **References**

- 500 Abram, N. J., Mulvaney, R., Vimeux, F., Phipps, S. J., Turner, J., and England, M. H.: Evolution of the Southern
501 Annular Mode during the past millennium, *Nature Climate Change*, 4, 564-569, 2014.
- 502 Arblaster, J. M. and Meehl, G. A.: Contributions of external forcings to southern annular mode trends, *Journal of*
503 *climate*, 19, 2896-2905, 2006.
- 504 Arblaster, J. M., Meehl, G. A., and Karoly, D. J.: Future climate change in the Southern Hemisphere: Competing
505 effects of ozone and greenhouse gases, *Geophysical Research Letters*, 38, 2011.
- 506 Banerjee, A., Fyfe, J. C., Polvani, L. M., Waugh, D., and Chang, K.-L.: A pause in Southern Hemisphere
507 circulation trends due to the Montreal Protocol, *Nature*, 579, 544-548, 2020.
- 508 Barnes, E. A. and Polvani, L.: Response of the midlatitude jets, and of their variability, to increased greenhouse
509 gases in the CMIP5 models, *Journal of Climate*, 26, 7117-7135, 2013.
- 510 Brehm, N., Bayliss, A., Christl, M., Synal, H.-A., Adolphi, F., Beer, J., Kromer, B., Muscheler, R., Solanki, S. K.,
511 and Usoskin, I.: Eleven-year solar cycles over the last millennium revealed by radiocarbon in tree rings,
512 *Nature Geoscience*, 14, 10–15, 2021.
- 513 Butler, A. H., Thompson, D. W., and Birner, T.: Isentropic slopes, downgradient eddy fluxes, and the extratropical
514 atmospheric circulation response to tropical tropospheric heating, *Journal of the atmospheric sciences*, 68,
515 2292-2305, 2011.
- 516 Crosta, X., Etourneau, J., Orme, L.C., Dalaiden, Q., Campagne, P., Swingedouw, D., Goosse, H., Massé, G.,
517 Miettinen, A., McKay, R.M., Dunbar, R.B., Escutia, C., and Ikehara, M.: Multi-decadal trends in Antarctic
518 sea-ice extent driven by ENSO–SAM over the last 2,000 years, *Nature Geoscience*, 14, 156–160,
519 10.1038/s41561-021-00697-1, 2021.
- 520 Dätwyler, C., Neukom, R., Abram, N. J., Gallant, A. J. E., Grosjean, M., Jacques-Coper, M., Karoly, D. J., and
521 Villalba, R.: Teleconnection stationarity, variability and trends of the Southern Annular Mode (SAM) during
522 the last millennium, *Climate Dynamics*, 51, 2321-2339, 10.1007/s00382-017-4015-0, 2018.
- 523 Emile-Geay, J., McKay, N. P., Kaufman, D. S., Von Gunten, L., Wang, J., Anchukaitis, K. J., Abram, N. J.,
524 Addison, J. A., Curran, M. A., and Evans, M. N.: A global multiproxy database for temperature
525 reconstructions of the Common Era, *Scientific data*, 4, 170088, 2017.
- 526 Fan, K. and Wang, H.: Antarctic oscillation and the dust weather frequency in North China, *Geophysical Research*
527 *Letters*, 31, 2004.
- 528 Fogt, R. L. and Marshall, G. J.: The Southern Annular Mode: variability, trends, and climate impacts across the
529 Southern Hemisphere, *Wiley Interdisciplinary Reviews: Climate Change*, e652, 2020.
- 530 Fogt, R. L., Perlwitz, J., Monaghan, A. J., Bromwich, D. H., Jones, J. M., and Marshall, G. J.: Historical SAM
531 variability. Part II: Twentieth-century variability and trends from reconstructions, observations, and the
532 IPCC AR4 models, *Journal of Climate*, 22, 5346-5365, 2009.
- 533 Fyfe, J., Boer, G., and Flato, G.: The Arctic and Antarctic Oscillations and their projected changes under global
534 warming, *Geophysical Research Letters*, 26, 1601-1604, 1999.

535 Gillett, N. and Fyfe, J.: Annular mode changes in the CMIP5 simulations, *Geophysical Research Letters*, 40,
536 1189-1193, 2013.

537 Gillett, N. P. and Thompson, D. W.: Simulation of recent Southern Hemisphere climate change, *Science*, 302,
538 273-275, 2003.

539 Gillett, N. P., Kell, T. D., and Jones, P.: Regional climate impacts of the Southern Annular Mode, *Geophysical
540 Research Letters*, 33, 2006.

541 Gong, D. and Wang, S.: Definition of Antarctic oscillation index, *Geophysical research letters*, 26, 459-462, 1999.

542 Goyal, R., Sen Gupta, A., Jucker, M., and England, M. H.: Historical and projected changes in the Southern
543 Hemisphere surface westerlies, *Geophysical Research Letters*, 48, e2020GL090849, 2021.

544 Gray, L. J., Beer, J., Geller, M., Haigh, J. D., Lockwood, M., Matthes, K., Cubasch, U., Fleitmann, D., Harrison, G.,
545 and Hood, L.: Solar influences on climate, *Reviews of Geophysics*, 48, 2010.

546 Gray, L. J., Scaife, A. A., Mitchell, D. M., Osprey, S., Ineson, S., Hardiman, S., Butchart, N., Knight, J., Sutton, R.,
547 and Kodera, K.: A lagged response to the 11 year solar cycle in observed winter Atlantic/European weather
548 patterns, *Journal of Geophysical Research: Atmospheres*, 118, 13,405-413,420, 2013.

549 Grise, K. M., Polvani, L. M., Tselioudis, G., Wu, Y., and Zelinka, M. D.: The ozone hole indirect effect: Cloud-
550 radiative anomalies accompanying the poleward shift of the eddy-driven jet in the Southern Hemisphere,
551 *Geophysical Research Letters*, 40, 3688-3692, 2013.

552 Hendon, H. H., Thompson, D. W., and Wheeler, M. C.: Australian rainfall and surface temperature variations
553 associated with the Southern Hemisphere annular mode, *Journal of Climate*, 20, 2452-2467, 2007.

554 Hessler, A., Allen, K. J., Vance, T., Abram, N. J., & Saunders, K. M.: Reconstructions of the southern annular mode
555 (SAM) during the last millennium. *Progress in Physical Geography*, 41(6), 834-849. 2017.

556 Huiskamp, W. and McGregor, S.: Quantifying Southern Annular Mode paleo-reconstruction skill in a model
557 framework, *Clim. Past*, 17, 1819–1839, 10.5194/cp-17-1819-2021, 2021.

558 Jones, J. M., Fogt, R. L., Widmann, M., Marshall, G. J., Jones, P. D., and Visbeck, M.: Historical SAM variability.
559 Part I: Century-length seasonal reconstructions, *Journal of Climate*, 22, 5319-5345, 2009.

560 Jones, J. M., Gille, S. T., Goosse, H., Abram, N. J., Canziani, P. O., Charman, D. J., Clem, K. R., Crosta, X., de
561 Lavergne, C., and Eisenman, I.: Assessing recent trends in high-latitude Southern Hemisphere surface
562 climate, *Nature Climate Change*, 6, 917-926, 2016.

563 Judge, P. G., Lockwood, G. W., Radick, R. R., Henry, G. W., Shapiro, A. I., Schmutz, W., and Lindsey, C.:
564 Confronting a solar irradiance reconstruction with solar and stellar data, *A&A*, 544, A88, 2012.

565 Jungclaus, J. H., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L. P., Egorova, T., Evans, M., González-
566 Rouco, J. F., Goosse, H., Hurtt, G. C., Joos, F., Kaplan, J. O., Khodri, M., Klein Goldewijk, K., Krivova, N.,
567 LeGrande, A. N., Lorenz, S. J., Luterbacher, J., Man, W., Maycock, A. C., Meinshausen, M., Moberg, A.,
568 Muscheler, R., Nehrbass-Ahles, C., Otto-Bliesner, B. I., Phipps, S. J., Pongratz, J., Rozanov, E., Schmidt,
569 G. A., Schmidt, H., Schmutz, W., Schurer, A., Shapiro, A. I., Sigl, M., Smerdon, J. E., Solanki, S. K.,
570 Timmreck, C., Toohey, M., Usoskin, I. G., Wagner, S., Wu, C. J., Yeo, K. L., Zanchettin, D., Zhang, Q., and
571 Zorita, E.: The PMIP4 contribution to CMIP6 – Part 3: The last millennium, scientific objective, and

572 experimental design for the PMIP4 past1000 simulations, *Geosci. Model Dev.*, 10, 4005-4033,
573 10.5194/gmd-10-4005-2017, 2017.

574 Kuroda, Y.: On the Origin of the Solar Cycle Modulation of the Southern Annular Mode, *Journal of Geophysical*
575 *Research: Atmospheres*, 123, 1959-1969, 10.1002/2017JD027091, 2018.

576 Kuroda, Y. and Kodera, K.: Solar cycle modulation of the Southern Annular Mode, *Geophysical Research Letters*,
577 32, 10.1029/2005GL022516, 2005.

578 Kuroda, Y. and Shibata, K.: Simulation of solar-cycle modulation of the Southern Annular Mode using a
579 chemistry-climate model, *Geophysical Research Letters*, 33, 10.1029/2005GL025095, 2006.

580 Kuroda, Y., Deushi, M., and Shibata, K.: Role of solar activity in the troposphere-stratosphere coupling in the
581 Southern Hemisphere winter, *Geophysical Research Letters*, 34, 10.1029/2007GL030983, 2007.

582 Kushner, P. J., Held, I. M., and Delworth, T. L.: Southern Hemisphere atmospheric circulation response to global
583 warming, *Journal of Climate*, 14, 2238-2249, 2001.

584 Lean, J. and Rind, D.: Climate forcing by changing solar radiation, *J. Climate*, 11, 3069-3094, 10.1175/1520-
585 0442(1998)011%3C3069%3ACFBCSR%3E2.0.CO;2, 1998.

586 Lim, E.-P. and Simmonds, I.: Effect of tropospheric temperature change on the zonal mean circulation and SH
587 winter extratropical cyclones, *Climate dynamics*, 33, 19-32, 2009.

588 Lu, H., Jarvis, M. J., Gray, L. J., and Baldwin, M. P.: High- and low-frequency 11-year solar cycle signatures in
589 the Southern Hemispheric winter and spring, *Quarterly Journal of the Royal Meteorological Society*, 137,
590 1641-1656, 10.1002/qj.852, 2011.

591 Ma, H., Chen, H., Gray, L., Zhou, L., Li, X., Wang, R., and Zhu, S.: Changing response of the North
592 Atlantic/European winter climate to the 11 year solar cycle, *Environmental Research Letters*, 13, 034007,
593 10.1088/1748-9326/aa9e94, 2018.

594 Marshall, G. J.: Trends in the Southern Annular Mode from Observations and Reanalyses, *Journal of Climate*, 16,
595 4134-4143, 10.1175/1520-0442(2003)016<4134:Titsam>2.0.Co;2, 2003.

596 Meehl, G. A., Arblaster, J. M., Branstator, G., and van Loon, H.: A Coupled Air-Sea Response Mechanism to
597 Solar Forcing in the Pacific Region, *Journal of Climate*, 21, 2883-2897, 10.1175/2007jcli1776.1, 2008.

598 Meehl, G. A., Arblaster, J. M., Matthes, K., Sassi, F., and van Loon, H.: Amplifying the Pacific climate system
599 response to a small 11-year solar cycle forcing, *Science*, 325, 1114-1118, 2009.

600 Meehl, G. A., Washington, W. M., Wigley, T., Arblaster, J. M., and Dai, A.: Solar and greenhouse gas forcing and
601 climate response in the twentieth century, *Journal of Climate*, 16, 426-444, 2003.

602 Meehl, G. A., Washington, W. M., Arblaster, J. M., Hu, A., Teng, H., Tebaldi, C., Sanderson, B. N., Lamarque, J.-
603 F., Conley, A., and Strand, W. G.: Climate system response to external forcings and climate change
604 projections in CCSM4, *Journal of Climate*, 25, 3661-3683, 2012.

605 Miller, R. L., Schmidt, G. A., and Shindell, D. T.: Forced annular variations in the 20th century Intergovernmental
606 Panel on Climate Change Fourth Assessment Report models, *Journal of Geophysical Research:*
607 *Atmospheres*, 111, 10.1029/2005JD006323, 2006.

608 Neukom, R., Schurer, A. P., Steiger, N. J., and Hegerl, G. C.: Possible causes of data model discrepancy in the
609 temperature history of the last Millennium, *Scientific Reports*, 8, 7572, 10.1038/s41598-018-25862-2,
610 2018.

611 Neukom, R., Barboza, L. A., Erb, M. P., Shi, F., Emile-Geay, J., Evans, M. N., Franke, J., Kaufman, D. S., Lücke,
612 L., Rehfeld, K., Schurer, A., Zhu, F., Brönnimann, S., Hakim, G. J., Henley, B. J., Ljungqvist, F. C., McKay,
613 N., Valler, V., von Gunten, L., and Consortium, P. k.: Consistent multidecadal variability in global
614 temperature reconstructions and simulations over the Common Era, *Nature Geoscience*, 12, 643-649,
615 10.1038/s41561-019-0400-0, 2019.

616 Ortega, P., Lehner, F., Swingedouw, D., Masson-Delmotte, V., Raible, C. C., Casado, M., and Yiou, P.: A model-
617 tested North Atlantic Oscillation reconstruction for the past millennium, *Nature*, 523, 71-74,
618 10.1038/nature14518, 2015.

619 Otto-Bliesner, B. L., Brady, E. C., Fasullo, J., Jahn, A., Landrum, L., Stevenson, S., Rosenbloom, N., Mai, A., and
620 Strand, G.: Climate Variability and Change since 850 CE: An Ensemble Approach with the Community
621 Earth System Model, *Bulletin of the American Meteorological Society*, 97, 735-754, 10.1175/bams-d-14-
622 00233.1, 2016.

623 PAGES 2k-PMIP3 group: Continental-scale temperature variability in PMIP3 simulations and PAGES 2k regional
624 temperature reconstructions over the past millennium, *Clim. Past*, 11, 1673–1699,
625 <https://doi.org/10.5194/cp-11-1673-2015>, 2015.

626 Phipps, S. J., Rotstayn, L. D., Gordon, H. B., Roberts, J. L., Hirst, A. C., and Budd, W. F.: The CSIRO Mk3L
627 climate system model version 1.0 – Part 1: Description and evaluation, *Geosci. Model Dev.*, 4, 483-509,
628 10.5194/gmd-4-483-2011, 2011.

629 Phipps, S. J., Rotstayn, L. D., Gordon, H. B., Roberts, J. L., Hirst, A. C., and Budd, W. F.: The CSIRO Mk3L
630 climate system model version 1.0 – Part 2: Response to external forcings, *Geosci. Model Dev.*, 5, 649-682,
631 10.5194/gmd-5-649-2012, 2012.

632 Phipps, S. J., McGregor, H. V., Gergis, J., Gallant, A. J. E., Neukom, R., Stevenson, S., Ackerley, D., Brown, J. R.,
633 Fischer, M. J., and van Ommen, T. D.: Paleoclimate Data-Model Comparison and the Role of Climate
634 Forcings over the Past 1500 Years, *Journal of Climate*, 26, 6915-6936, 10.1175/jcli-d-12-00108.1, 2013.

635 Polvani, L. M., Waugh, D. W., Correa, G. J., and Son, S.-W.: Stratospheric ozone depletion: The main driver of
636 twentieth-century atmospheric circulation changes in the Southern Hemisphere, *Journal of Climate*, 24,
637 795-812, 2011a.

638 Polvani, L. M., Waugh, D. W., Correa, G. J. P., and Son, S.-W.: Stratospheric Ozone Depletion: The Main Driver of
639 Twentieth-Century Atmospheric Circulation Changes in the Southern Hemisphere, *Journal of Climate*, 24,
640 795-812, 10.1175/2010jcli3772.1, 2011b.

641 Raphael, M. N. and Holland, M. M.: Twentieth century simulation of the southern hemisphere climate in coupled
642 models. Part 1: large scale circulation variability, *Climate Dynamics*, 26, 217-228, 10.1007/s00382-005-
643 0082-8, 2006.

644 Rind, D., Lean, J., Lerner, J., Lonergan, P., and Leboissitier, A.: Exploring the stratospheric/tropospheric
645 response to solar forcing, *Journal of Geophysical Research: Atmospheres*, 113, 2008.

646 Roscoe, H. K. and Haigh, J. D.: Influences of ozone depletion, the solar cycle and the QBO on the Southern
647 Annular Mode, *Quarterly Journal of the Royal Meteorological Society*, 133, 1855-1864, 10.1002/qj.153,
648 2007.

649 Saunders, K.M., Roberts, S.J., Perren, B. *et al.*: Holocene dynamics of the Southern Hemisphere westerly winds
650 and possible links to CO₂ outgassing. *Nature Geoscience*, 11, 650–655, 10.1038/s41561-018-0186-5,
651 2018

652 Schmidt, G., Jungclaus, J. H., Ammann, C., Bard, E., Braconnot, P., Crowley, T., Delaygue, G., Joos, F., Krivova,
653 N., and Muscheler, R.: Climate forcing reconstructions for use in PMIP simulations of the Last Millennium
654 (v1. 1), *Geoscientific Model Development*, 185-191, 2012.

655 Schmidt, G. A., Jungclaus, J. H., Ammann, C., Bard, E., Braconnot, P., Crowley, T., Delaygue, G., Joos, F.,
656 Krivova, N., and Muscheler, R.: Climate forcing reconstructions for use in PMIP simulations of the last
657 millennium (v1. 0), 2011.

658 Schurer, A. P., Tett, S. F., and Hegerl, G. C.: Small influence of solar variability on climate over the past
659 millennium, *Nature Geoscience*, 7, 104-108, 2014.

660 Sen Gupta, A. and England, M. H.: Coupled ocean–atmosphere–ice response to variations in the southern
661 annular mode, *Journal of Climate*, 19, 4457-4486, 2006.

662 Shapiro, A., Schmutz, W., Rozanov, E., Schoell, M., Haberleiter, M., Shapiro, A., and Nyeki, S.: A new approach
663 to the long-term reconstruction of the solar irradiance leads to large historical solar forcing, *Astronomy &*
664 *Astrophysics*, 529, A67, 2011.

665 Shindell, D. T. and Schmidt, G. A.: Southern Hemisphere climate response to ozone changes and greenhouse
666 gas increases, *Geophysical Research Letters*, 31, 2004.

667 Son, S. W., Tandon, N. F., Polvani, L. M., and Waugh, D. W.: Ozone hole and Southern Hemisphere climate
668 change, *Geophysical Research Letters*, 36, 2009.

669 Steinhilber, F., Beer, J., and Fröhlich, C.: Total solar irradiance during the Holocene, *Geophysical Research*
670 *Letters*, 36, 2009.

671 Swart, N. and Fyfe, J. C.: Observed and simulated changes in the Southern Hemisphere surface westerly wind-
672 stress, *Geophysical Research Letters*, 39, 2012.

673 Thompson, D. W. and Solomon, S.: Interpretation of recent Southern Hemisphere climate change, *Science*, 296,
674 895-899, 2002.

675 Thompson, D. W. and Wallace, J. M.: Annular modes in the extratropical circulation. Part I: Month-to-month
676 variability, *Journal of climate*, 13, 1000-1016, 2000.

677 Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of
678 the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nature Geoscience*, 4, 741-749,
679 10.1038/ngeo1296, 2011.

680 Villalba, R., Lara, A., Masiokas, M. H., Urrutia, R., Luckman, B. H., Marshall, G. J., Mundo, I. A., Christie, D. A.,
681 Cook, E. R., and Neukom, R.: Unusual Southern Hemisphere tree growth patterns induced by changes in
682 the Southern Annular Mode, *Nature geoscience*, 5, 793-798, 2012.

683 Visbeck, M.: A station-based southern annular mode index from 1884 to 2005, *Journal of Climate*, 22, 940-950,
684 2009.


685 Wang, G. and Cai, W.: Climate-change impact on the 20th-century relationship between the Southern Annular
686 Mode and global mean temperature, *Scientific reports*, 3, 1-6, 2013.

687 Wang, Y.-M., Lean, J., and Sheeley Jr, N.: Modeling the Sun's magnetic field and irradiance since 1713, *The*
688 *Astrophysical Journal*, 625, 522, 2005.

689 Yang, D., Arblaster, J. M., Meehl, G. A., England, M. H., Lim, E.-P., Bates, S., and Rosenbloom, N.: Role of
690 tropical variability in driving decadal shifts in the Southern Hemisphere summertime eddy-driven jet,
691 *Journal of Climate*, 33, 5445-5463, 2020.

692 Zheng, F., Li, J., Clark, R. T., and Nnamchi, H. C.: Simulation and projection of the Southern Hemisphere annular
693 mode in CMIP5 models, *Journal of Climate*, 26, 9860-9879, 2013.

694




696


697 **Fig. 1. Reconstructed SAM and solar variability over the last millennium.** (a) Observation-based mean annual SAM
 698 indices (normalised units) of Marshall (2003) (red) and Fogt et al. (2009) (orange), and the reconstructed SAM index of
 699 Abram et al. (2014) (A14, blue) and Dätwyler et al. (2018) (D18, black; see Section 2.1 for an explanation of the
 700 difference in magnitude between the SAM reconstructions). The reconstructions are plotted as a 7-yr moving average
 701 for the annual SAM index (thin lines) with a 70-yr loess filter (thick lines), while 7-yr moving averages are shown for the
 702 observational data. (b) Total solar irradiance reconstructions for the last millennium, including Steinhilber et al. (2009)
 703 (blue), and a modified version of Steinhilber et al. (2009) (green) where the amplitude of the variations relative to the
 704 long-term mean has been increased by a factor of two and Shapiro et al. (2011) (purple). For comparison with the

705 current high amplitude scenario, we also include the PMOD reconstruction from PMIP4 (Jungclaus et al., 2017)
706 (orange). Note that PMOD has been normalised to 1365 W m^{-2} , to be comparable to the other solar reconstructions
707 presented here. (c) Solar irradiance values for our solar constant experiments, with the experiment name provided in
708 parentheses. (d) Solar modulation parameter reconstruction for the last millennium, based on radiocarbon from annually
709 resolved and dated tree rings (Brehm et al., 2021), shown as monthly resolution (thin lines) and as a 70-yr moving
710 average (thick lines).

711
712
713
714
715
716
717
718
719




720
721 **Fig. 2.** Difference in the observed annual SAM index when calculated using monthly MSLP anomalies (i.e., using
722 normalised monthly MSLP anomalies at 40°S and 65°S , and then averaging this monthly SAM index into an annual
723 timeseries; orange line) and annual MSLP (i.e., normalised MSLP anomalies at 40°S and 65°S ; red line). Data from
724 Marshall (2003), and the updated index is publicly available (<http://www.nerc-bas.ac.uk/icd/gjma/sam.html>).



725

726 **Fig. 3. The effect of calibration target on scaling of last millennium SAM reconstructions.**

727 (a) observational-based annual SAM index (Marshall, 2003) calculated using annual MSLP anomalies (red line; shown as
 728 7-yr moving average), which forms the calibration target for the A14 SAM reconstruction (blue line; shown as 7-yr
 729 moving average and 70-yr loess filter by thin and thick lines, respectively). (b) observational-based annual SAM index
 730 (Marshall, 2003) calculated using monthly MSLP anomalies (orange line; shown as 7-yr moving average) and the result
 731 this alternate calibration target has on the scaling of A14 SAM reconstruction (purple line; shown as 7-yr moving
 732 average and 70-yr loess filter by thin and thick lines, respectively). For comparison, the D18 SAM reconstruction is
 733 shown in black (7-yr moving average and 70-yr loess filter by thin and thick lines, respectively) on both (a) and (b), and
 734 is based on a monthly-derived annual SAM index as the calibration target.



735

736 **Fig. 4. 70-yr moving average for the annual SAM index for each solar constant simulation.** Period of interest in this

737 study (first 200 years) is indicated with a green box.




Fig. 5. SAM index and events for the first 200 years of fixed solar forced model runs. (a) Violin plots of the distribution of mean annual SAM index calculated for each solar constant run. Coloured horizontal lines refer to the 25th percentile, mean, and 75th percentile. (b) Number of extreme SAM events (outside $\pm 2\sigma$ of the control run) in 200 years, with a second order line of best fit for positive and negative events. Blue circles refer to extreme negative SAM events, and red crosses refer to extreme positive SAM events. (c–e) Zonal mean anomalies for the first 200 years of the experiments for: (c) mean sea level pressure, (d) surface air temperature, and (e) surface zonal wind stress (positive values indicate westerly anomalies); where the anomaly is relative to the control run. Colours for each experiment correspond to those used in panel (a). Dashed lines at 40°S and 65°S are the latitudes used to calculate the SAM index. (f–h): MSLP anomalies relative to the control for the first 200 years in the (f) S–15, (g) S–7, and (h) S+35 experiments. Regions shown are significantly different to the control run (based on Welch’s t-test, $P < 0.05$); areas where $P \geq 0.05$ have been masked out.




Fig. 6. Surface anomalies (relative to the control run) for the first 200 years of the different solar constant experiments. (a) MSLP anomaly (shaded) and surface wind anomalies (vectors). (b) Surface temperature anomaly (shaded), with temperature contours additionally shown at 0.5 °C increments (to assist in interpreting S+35). Regions that are significantly different (based on a Welch's t -test, $P < 0.05$) are shown, while areas where $P \geq 0.05$ have been masked out.




Fig. 7: SAM index anomaly based on the different Mk3L simulations. SAM index anomaly is calculated relative to 1900–1999 mean and shown as 70-yr moving averages. (a) orbital (‘O’) only simulation; (b) orbital and greenhouse gases (‘OG’) simulation; (c) orbital, greenhouse gases, and low amplitude solar forcing (‘OGS’) simulation; (d) orbital, greenhouse gases, and the intermediate amplitude solar forcing (‘OGS-x2’) simulation; (e) orbital, greenhouse gases, and high amplitude solar (‘OGS-Shapiro’) simulation. Thick lines refer to the ensemble mean, while thin lines denote the individual ensemble members. Dashed lines on all subplots show the $\pm 2\sigma$ range based on the orbital (‘O’) only simulations, representing internal variability.




Fig. 8. SAM index response to different amplitudes of transient solar forcing during the last millennium. Time series (left-side panels) of the SAM index (coloured lines; ensemble mean: thick line; ensemble members: thin lines) and radiative forcing anomalies (black) in the transient experiments with (a) low, (b) intermediate and (c) large amplitude solar forcing. Time series shown for the SAM index are calculated relative to 1900–1999 climatology, and long-term changes are represented as 70-yr moving averages stepped by 35 years. Scatter plots (right-side panels) show the 850–1900 CE correlation between 70-yr moving averages stepped by 35 years of the SAM index and radiative forcing anomalies in the ensemble mean (black, first r -value), and ensemble members (coloured, r -values in square brackets give range across ensemble members); r -values in bold text are where $P < 0.05$, while r -values in grey text are where there is no significant relationship. Correlations are limited to 850–1900 CE to avoid the influence of recent anthropogenic greenhouse gas influences on the radiative forcing–SAM relationship.




Fig. 9: SAM index response to different levels of radiative forcing compiled across solar transient experiments.

(a) Violin plots of the 70-yr rolling mean SAM index between 850–1900 CE; lines refer to the 25th percentile, mean, and 75th percentile. (b) Scatter plot of 70-yr rolling mean for radiative forcing anomaly and the SAM index; $r = 0.78$ ($P < 0.05$). (c–d) Zonal mean anomalies for: (c) mean sea level pressure (MSLP), and (d) temperature, where the anomaly is relative to the climatological (1900–1999) mean. Coloured lines refer to the radiative forcing bins for -1.5 to -1.0 $W m^{-2}$ (purple), -1.0 to -0.5 $W m^{-2}$ (blue); -0.5 to 0.0 $W m^{-2}$ (orange); 0.0 to 0.5 $W m^{-2}$ (red). (e–h) MSLP anomalies (relative to the climatological mean based on radiative forcing bins). Regions shown are significantly different to the OGS ensemble mean (based on Welch’s t-test, $P < 0.05$); areas where $P \geq 0.05$ have been masked out. Dashed lines at $40^{\circ}S$ and $65^{\circ}S$ are the latitudes used to calculate the SAM index.




Fig. 10. Last millennium SAM reconstructions (a–c) and the significance of negative shifts in 70-year distributions through time relative to a 70-yr preindustrial reference period (1831–1900) (d). Annual mean SAM reconstructions (thin curves) and 70-y loess filters (thick curves) for (a) the A14 SAM reconstruction (blue); (b) the A14-rescaled SAM reconstruction (purple; See Fig. 3); and (c) the D18 SAM reconstruction. SAM reconstructions shown relative to 1900–1999 climatology (dashed horizontal lines) (d) Distribution tests using a Wilcoxon rank-sum test of moving 70-yr windows of each SAM reconstruction shown in parts (a–c), relative to its 70-year preindustrial reference period (1831–1900 CE; yellow vertical shading). Timeseries show the probability for the distribution of sliding test windows being more negative than the distribution in the reference interval; $P < 0.05$ are shown with a white background, while $P \geq 0.05$ are shown with a grey background. All three reconstructions show a significant ($P < 0.05$) negative shift in the SAM index compared with the 1831–1900 reference interval during the period from approximately 1390 to 1715 CE. Both versions of the A14 SAM reconstruction also have a significant negative SAM shift prior to around 1140 CE.




Fig. 11. SAM index in the CSIRO Mk3L transient last millennium simulations (a–c) and the significance of negative shifts in 70-year windows of annual SAM distributions through time relative to a 70-yr preindustrial reference period (1831–1900) (d). 70-yr loess filter of the SAM index of (a) the OGS simulations; (b) OGS-x2 simulations; and (c) OGS-Shapiro simulations (as in Fig. 8). The ensemble mean is shown as a thick line, and individual ensemble members are shown in thin lines. SAM indices are shown relative to their 1900–1999 climatology (dashed horizontal lines). (d) Distribution tests on ensemble means using a Wilcoxon rank-sum test of sliding 70-yr windows relative to the preindustrial reference period (1831–1900 CE; vertical yellow shading), where colours match the SAM simulations shown in parts a–c. Timeseries show the probability for the distribution of sliding test windows being more negative than the distribution in the reference interval. $P < 0.05$ are shown with a white background, while $P \geq 0.05$ are shown with a grey background.




Fig. 12. Comparison of simulated SAM index and the A14 SAM reconstruction between 1000 and 1900 CE. Time series (left-side plots) of the SAM index (70-yr moving averages stepped by 35 years, calculated relative to 1900–1999 historical mean) from ensemble means of transient simulations and SAM reconstructions (black line; A14) (a) for transient experiments from CSIRO Mk3L using low (blue, OGS), intermediate (green, OGSx2), and high (purple, OGS-Shapiro) amplitude solar forcing. (b) for simulations from HadCM3 (Schurer et al., 2014), including the full forcing (blue), weak solar (orange) and strong solar (“Solar Shapiro” runs using Shapiro et al., 2011; red) simulations. Note that there is only a single member for the strong solar scenario, and that the solar forcing simulations from HadCM3 do not include greenhouse gases whereas the full forcing simulation does. This results in the last millennium SAM index from the full-forcing simulation having a lower mean than the solar forcing-only experiments when calculated relative to the 1900–1999 historical mean. Scatter plots (right-side plots) show the correlation between the simulated ensemble mean and reconstructed SAM indices, as 70-yr moving averages stepped by 35 years, and r -values between the simulated and reconstructed SAM index are shown in bold text where $P < 0.05$.




Fig. 13. Comparison of simulated SAM index and the D18 SAM reconstruction between 1000 and 1900 CE. Time series of the SAM index (70-yr moving averages stepped by 35 years; ensemble mean: thick line; ensemble members: thin lines) from transient simulations and SAM reconstruction (70-yr moving averages stepped by 35 years) from D18 (black line) for (a) solar transient experiments from CSIRO Mk3L, and (b) simulations from HadCM3 (Schurer et al., 2014), including the full forcing (blue), weak solar (orange) and strong solar (i.e., Shapiro et al., 2011; red) simulations. Note that the solar forcing simulations from HadCM3 do not include GHGs. Scatter plots show the correlation between 70-yr moving averages of the simulated SAM index and 70-yr moving average of the reconstructed SAM index for the ensemble mean, and r values between the simulated and reconstructed SAM index where $P < 0.05$ are shown in bold text.




Fig. 14: Atmospheric profiles for the Southern Hemisphere mean anomalies for the first 200 years of each solar constant experiment. Anomalies are calculated relative to the long-term control. (a) Zonal wind anomalies (shaded), and contours of the control zonal wind (lines). (b) Temperature anomalies (shaded); contour lines of the temperature anomaly are additionally shown in 0.5 °C increments, to assist in interpreting S-15 and S+35.