

Wet season rainfall characteristics and temporal changes for Cape Town South Africa, 1841-2018

Nothabo E. Ndebele¹, Stefan Grab², Herbert Hove¹

¹School of Statistics and Actuarial Science, University of the Witwatersrand, South Africa

5 ²School of Geography, Archaeology and Environmental Studies, University of the Witwatersrand, South Africa

Correspondence to: Nothabo E. Ndebele (Nothabo.Ndebele@wits.ac.za)

Abstract. Wet seasons may be characterized by the frequency of wet/dry days, duration of wet/dry spells, and season length. These properties are investigated for Cape Town using rainfall data from four weather stations in the Cape Town metropolitan area located at the South African Astronomical Observatory (SAAO), Maitland, Kirstenbosch and Cape Town International airport. The primary focus is on the long SAAO daily rainfall record dating back to 1841, with the specific aim to statistically assess attributes of the wet season (April to October) and its temporal variability over the period 1841-2018. The decade 1950-1959 had significantly high frequencies of wet days, but there was a subsequent significant decline in wet days at the SAAO (-1day/decade) and Maitland (-1.1days/decade) during the period 1950-2018. A significant decline in wet days also occurred at the SAAO between 1880 and 1940 (-3.3 days/decade, $p = 0.005$). Dry spells longer than 5 days have become more prevalent since the beginning of the 20th century. A rain-based definition for the onset and termination of the wet season is presented using 5-day running sums and pentad means; these were applied to each year containing adequate daily data, so as to track changes during the wet season. We identify a recent decadal decline in mean wet season length (-1.4 days/decade) since c. 1940, particularly between 1941-1994 (-4.2 days/decade), such that wet seasons since 2000 have only averaged 182 days in length while the long-term (1841-2018) average is 188 days. This decline is attributed to an increased incidence of late onsets (after 15 April) and early terminations (earlier than 18 October) of the wet season, or a combination of both, particularly since the year 2000. Interannual variability in wet season characteristics is associated with solar (sunspot) cycles and fluctuations in the Southern Oscillation Index and Southern Annular Mode.

1. Introduction

Temporal characteristics of seasonal rainfall vary across regions and climates of the world (Knoben et al., 2019). In Africa, wet seasons are either unimodal (concentrated over only one peak rainfall period per 12 month cycle: e.g. southern Africa) or bimodal (two temporally separated wet seasons per 12 month cycle: e.g. south coast of South Africa and large parts of East and West Africa) (Camberlin et al., 2009; Amekudzi et al., 2015). Understanding the nature and potential changes to such wet seasons is important to several sectors, most particularly agriculture, water resources planning and health (Hachigonta et al., 2008; Singh and Ranade, 2010). To this end, considerable scientific interest has focused on establishing the onset/termination dates of the annual wet season(s) and how these may have changed over recent decades (e.g. Mupangwa et al., 2011; Owusu

and Waylen, 2013; Oguntunde et al., 2014; Obarein and Amanambu, 2019; Giráldez et al., 2020), or might change in future through climate modeling projections (e.g. Kusunoki et al., 2011; Sarr, 2012; Saeed et al., 2018). Defining seasonal rainfall onset/termination dates depends on location, climate and the study purpose; hence defining attributes have been variable across disciplines (climatology, hydrology, agronomy) and regions. Rainfall based variables are most commonly used to determine rainfall onset/termination dates; these include the number of wet days, rainfall amounts (mm) and wet/dry spells (days)(e.g. Hachigonta et al., 2008; Ngetich et al., 2014). However, in some instances, other variables such as temperature, wind and humidity have also been used to compute potential evapotranspiration and soil water balance, particularly to help determine suitable rainfall thresholds required for specific crop growth (Mugalavai et al., 2008; Moron and Robertson, 2014).

Establishing wet season length usually requires rainfall based empirical methods. For instance, Boyard-Micheau et al. (2013) determine the rainfall onset date in Kenya and northern Tanzania based on the first wet day of N consecutive days, receiving a minimum of P_{min} of rain, without a dry spell lasting more than N_d number of days, and additionally specify a minimum P_{min} value for subsequent days. Each of these parameter values is determined empirically to address a set of requirements, most typically associated with crop growth. So for example, understanding the rainfall requirements for maize to germinate and grow has helped establish an appropriate onset date, based on a threshold of 25mm, but over a 10 day period followed by at least 20mm over the subsequent 20 days (Tadross et al., 2005; Hachigonta et al., 2008). In this case, rainfall data were divided into pentads (5 day periods) so as to identify the pentad that best satisfied the criteria for maize growth (Tadross et al., 2005). The use of pentads is also common practice for identifying other characteristics of the wet season, such as wet season length (e.g. Hachigonta et al., 2008; Marengo et al., 2017; Byakatonda et al., 2018).

Alternative approaches to establish seasonal rainfall onset/termination dates include using the cumulative distribution of rainfall amounts (Nicholls, 1984; Amekudzi et al., 2015) or accumulated rainfall anomalies (Liebmann and Marengo, 2001; Camberlin et al., 2009; Liebmann et al., 2012; Dunning et al., 2016). Liebmann and Marengo (2001) delineate onset/termination dates by obtaining absolute minimum/maximum of annual accumulated anomalies, calculated by subtracting the long-term mean rainfall from the actual rainfall for each day of the year. This approach is less sensitive to those using specified thresholds and may thus be applied at local and regional scales. The method by Liebmann et al. (2012) has more recently been extended to accommodate regions that have two wet seasons per annum, such as parts of West and East Africa (e.g. Dunning et al., 2016).

Also of importance, is establishing rainfall characteristics (daily frequency, distribution, amount; wet and dry spells) during the wet season, as these may have an influence on crop production, biosystems, hydrology etc. Wet/dry days are defined by either using a fixed value or using a parameter such as mean daily rainfall, for the area of interest. Given that the usual precision of rain gauges is 0.1mm, at times this is taken to be the threshold value (e.g. Martin-Vide and Gomez, 1999). However, many prefer a 1mm threshold to define a wet day given the assumption that an amount less than 1mm is likely returned to the atmosphere through evapotranspiration (e.g. Kruger, 2006; Nastos and Zerefos, 2009; Froidurot and Diedhiou, 2017; Valdes-Abellan et al., 2017; Byakatonda et al., 2019). Most recently, Rivoire et al. (2019) argued that since evapotranspiration is a variable (depending on the season), a fixed threshold may not be suitable, particularly for the dry season. When a wet day

65 threshold has been defined, wet/dry spells are identified as consecutive days of wet/dry days, and these may be used to further investigate rainfall characteristics.

Several previous studies have examined winter rainfall and attributes of the wet season over southernmost Africa (e.g. Reason et al., 2002; Reason and Rouault, 2005; Dieppois et al., 2016; du Plessis, 2017; Mahlalela et al., 2019). Identified associations between rainfall and climate modes such as the El Niño Southern Oscillation (ENSO) and Southern Annular Mode (SAM) are now well established for the region (Tyson, 1981; Mason and Tyson, 1992; Reason and Rouault, 2002; Philippon et al., 2012). Variability in rainfall at interannual and decadal time scales is influenced by SAM in the winter rainfall region in South Africa (Reason and Rouault, 2005; Dieppois et al., 2016; Ndebele et al., 2020). Links between ENSO and rainfall are significant with longer wet spells and higher rainfall associated with El Niño events (Philippon et al., 2012). The association of solar activity with rainfall and other climate variables has also been confirmed for the region (Dyer, 1975; Mason and Tyson, 1992; Ndebele et al., 2020). Using 20th century data, Mason and Tyson (1992) found that an increase in solar activity during the early part of the winter season (April to June) corresponded with higher precipitation over the winter rainfall region of South Africa. More recently, Ndebele et al. (2020) confirmed positive correlations between annual rainfall and solar activity at decadal time scales. Similar associations between rainfall and solar activity have been found for other regions globally (e.g. Laurenz et al., 2019; Nitka and Burnecki, 2019). When investigating solar activity and rainfall in Europe using data since 1900, Laurenz et al., (2019) observed correlations between rainfall and the 11 year solar cycle. After detecting significant correlations between solar activity and precipitation in the U.S.A, Nitka and Burnecki (2019) found that precipitation forecasts improved when adding solar activity as a variable.

Despite previous scientific effort, to date, and to our knowledge, relatively few studies have investigated long-term (> a century) wet season characteristics in Africa, despite its potential importance for ongoing and future agricultural planning at sub-regional scales. To this end, we use one of the worlds' longest single station rainfall records (178 years), namely that of the South African Astronomical Observatory (SAAO) in Cape Town, South Africa, to statistically establish the nature and temporal changes of wet season characteristics through the period 1841-2018. The work expands on that recently describing the trends and variability of rainfall in Cape Town since 1841 (Ndebele et al., 2020). While the SAAO record is the primary focus of this paper given its long record and historic value, additional station records over the Cape Town metropolitan region are also analyzed for imputation and comparative purposes. Specific objectives of the current paper include the quantification of wet season length, the occurrence of dry/wet spells, frequency of wet days, and to establish the temporal variation of these attributes. Such rainfall characteristics for Cape Town, which previous studies have not yet examined, provide important long-term perspectives relevant to, in particular, agricultural production, hydrology, and ecosystem functioning. In addition, given the considerable length of our dataset, we statistically test for any associations between these wet season rainfall characteristics and their potential drivers (i.e. El Niño Southern Oscillation, Southern Annular Mode, and solar activity [sunspots]).

2. Site context and data

2.1. Site context

Cape Town is the second largest city in South Africa and has a coastal setting in the southwestern portion of the Western Cape Province (Fig. 1). This region is characterised by a complex topography with mountain ranges exceeding 1000m in elevation within a few km of the coast, but also includes the Cape flats, a low lying and flat relief bounded by the Atlantic Ocean on southern and western sides, and where Cape Town International Airport is located (Goodness and Anderson, 2013). The eastern side of the Cape Flats and False Bay also have mountain ranges which exceed 1500m in some places. Cape Town has a Mediterranean climate with typically cool, wet austral winters (May-Sept) and dry, warm summers (Ndebele et al., 2020). According to Mahlalela et al. (2019), the southwestern and eastern coastal regions, and also some windward facing mountain slopes, receive on average >500mm of rainfall per annum, yet parts of the SW Cape region, particularly northwards and inland of Cape Town, may receive as little as 200-400mm per annum. Rainfall quantity/distribution is thus highly variable and influenced by both topography and direction of rain-bearing weather systems. Rainfall over the southwestern Cape region is primarily associated with overpassing cold fronts connected to mid-latitude cyclones, but as much as 11% may be owing to cut-off lows which occasionally bring exceptionally heavy rainfall (Abba Omar and Abiodun, 2020).

Cape Town city centre is located between Table Mountain and Table Bay (Fig. 1). Rainfall stations selected for this study are all within 20km of the city centre, and despite receiving variable rainfall amounts, they are influenced by a common rainfall seasonal regime and rain bearing systems. Rainfall records from four weather stations across the Cape Town metropolitan region are investigated; these include Cape Town International Airport, Kirstenbosch, Maitland, and the SAAO (Fig. 1). The SAAO, founded in 1820, is located *c.* 5km east of central Cape Town and *c.* 2km inland of Table Bay, on the eastern side of Table Mountain. Daily rainfall records began in January 1841 and continue to this day, making it the longest continuous single station daily record in the southern hemisphere (Ndebele et al., 2020). Daily readings have been manually read from a rain gauge located in the gardens of the SAAO over the duration of this period. Given that our aim is to examine long-term (> one century) rainfall changes, for which reliable data become largely unavailable with distance from central Cape Town, and because the nature of rainfall regimes change over relatively small distances (few 10s of km) from this region, our focus of investigation is constrained to the Cape peninsula only.

2.2. Data

As is often the case with early long-term instrumental weather registers, the SAAO record (1841-2018) has some data gaps, owing mainly to the absence of recordings over weekends and public holidays, and in some cases due to instrument failure or replacement. Although annual rainfall totals are available for all years, daily records are completely absent for the years 1878 and 1879. For the remainder of the record, missing daily rainfall values fall into two categories; namely days recorded as completely absent of data, or as days (e.g. weekends and public holidays) absent of data but with an accumulated total rainfall

value on a subsequent day. About 3.2% of missing days have an accumulated value on a subsequent day and only *c.* 2% of the record is completely missing. Changes in the rainfall recording practice and instruments used to record precipitation did not change much over the years and should thus not have any significant impact on our results. However, we note that missing but accumulated data are more prevalent since the early 1960s. Methods of recording precipitation changed in the 2000s, from manually read rain gauges to digital gauges, which might account for slight (yet insignificant) differences in measured rainfall amounts in the most recent decades. For further comments on data quality and data cleaning, please refer to Ndebele et al., 2020. Imputation of missing values for the SAAO record was performed using data from neighbouring stations at Maitland (1906-2018), Kirstenbosch (1915-2018), and Cape Town International Airport (1950-2018) (Ndebele et al., 2020). The imputation involves a two-step approach enabling the estimation of wet days and rainfall amounts on those days. One method was applied to days with completely missing records, while a second method was applied to those with accumulated totals for each month separately as the frequency of wet days and daily rainfall amounts vary with each month (see Ndebele et al., 2020 for further details).

Data for the climate modes: Southern Oscillation Index-SOI (Ropelewski and Jones, 1987), Southern Annular Mode-SAM (Gong and Wang, 1999), and solar (sunspot number) (Vanlommel et al., 2004) were used for examining associations with interannual variability in wet season characteristics. Data are available as either monthly or bimonthly indices for the periods 1866-2018 (SOI), 1851-2011 (SAM), and 1841-2018 (solar) (Table 1). Solar data are based on the sunspot number obtained from the Royal Observatory of Belgium in Brussels. The Twentieth Century Reanalysis version 2 (20CR.v2) was used to extend SAM back to 1851.

2.3. Defining rainfall variables and season length

Please refer to Table 2 for terms used to define rainfall variables and the wet season. In this paper, a ‘wet day’ is defined as one having received ≥ 1 mm of rain (as per Camberlin et al., 2009; Singh and Ranade, 2010; Boyard-Micheau et al., 2013; Muthuwatta et al., 2017; Byakatonda et al., 2018). The number of years with complete daily data is represented by n_d . The frequency of days with rainfall amounts > 0 mm experienced on each calendar day ($d = 1, 2, \dots, 365$) over the complete years is given by f_{w_1} . The frequency of wet days (≥ 1 mm) over the complete years is given by f_{w_2} . These frequencies are used to estimate the probability of a wet day (> 0 mm) P_{WD}^1 and (> 1 mm) P_{WD}^2 calculated for each calendar day (Table 2). The average daily rainfall amount is calculated for each calendar day (\bar{R}_d), for each month (\bar{R}_m), and for each year (\bar{R}_Y). The long-term daily average is calculated using all days in the record and is represented by \bar{R}_L (Table 2).

A wet (dry) spell is defined as a period of consecutive days with daily rainfall amounting to ≥ 1 mm (< 1 mm) preceded and followed by a dry (wet) day during the wet season. The term ‘one day spell’ is used to refer to a wet (or dry) day preceded and followed by a dry (wet) day. On this basis, the following are established: a) frequency of wet/dry spells, b) amount of rainfall during wet spells, c) duration and timing of spells, and d) statistical distributions associated with the lengths of wet/dry spells.

Estimates of wet season onset/termination dates are obtained by applying a combination of methods including the sum of rainfall values over a five day period (pentads) (after Odekunle, 2004; Amekudzi et al., 2015), and by comparing with the long-term pentad average \bar{D}_l . Initially, data are divided into 73 non overlapping pentads (5 day periods of consecutive days) over 365 days each year, starting on 1 January each year. The 29th February is not included in affected years and should not adversely impact results as this is a dry time of year. Average rainfall over a 5-day range (\bar{D}_p) and long-term pentad average (\bar{D}_l) are used to determine the lower and upper limits for wet season onset/termination dates.

For all the stations, days between pentad 19 (starting on Julian day 91) and pentad 61 (ending on Julian day 305), potentially represent the wet season (i.e. 1 April to 31 October). Specific rainfall onset dates (W_o) for a calendar year are defined as the first day that the 5-day running sum is above the long-term pentad average (\bar{D}_l) occurring after Julian day 90. Rainfall termination dates (W_E) are defined as the last day with a 5-day running sum above the long-term pentad average \bar{D}_l occurring before Julian day 306. The wet season (l_{ws}) length is then quantified on the basis of onset/termination dates. To confirm that the period between onset and termination dates actually encapsulates the majority of rainfall received in any given year, the accumulated rainfall during the defined wet season for that year is taken as a proportion of the annual rainfall for that year (P_{ws}).

175

2.4. Statistical methods and tests

2.4.1. Mann Kendall tests for trends

The Mann Kendall test (Mann, 1945 and Kendall, 1948) is used to test the null hypothesis that there is no trend under the assumption that observations in a series are independent. Given a time series y_1, y_2, \dots, y_n , the following steps were undertaken:

180

$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n 1 \times \text{sign}(y_j - y_i) \quad (1)$$

where n is the number of data points in the time series y_1, y_2, \dots, y_n ; $i, = 1, 2, \dots, n - 1$ and $j = 2, 3, \dots, n$ for $j > i$. The mean of S , $E[S] = 0$ and the variance of S is given by:

$$\text{var}(S) = \frac{1}{18} [n(n-1)(2n+5)] - \sum_{p=1}^q t_p(t_p-1)(2t_p+5) \quad (2)$$

where q is the number of tied groups and t_p is the number of data points in the p th group.

185 The test statistic is then calculated as follows:

$$z = \begin{cases} \frac{S - 1}{\sqrt{\text{var}(S)}}, & \text{if } S > 0 \\ 0, & \text{if } S = 0 \\ \frac{(S + 1)}{\sqrt{\text{var}(S)}}, & \text{if } S < 0 \end{cases} \quad (3)$$

The z statistic follows a standard normal distribution and therefore if $|z| > z_{1-\frac{\alpha}{2}}$, the null is rejected and it is concluded that there is a trend present in the time series. Rainfall time series often exhibit serial correlation violating the assumptions of the test, thus adjustments have been made to the Mann Kendall test statistic that allow for dependence in the observations. In this study, modifications include variance correction (Hamed and Rao, 1998), variance correction using lag-1 correlation (Yue and Wang, 2004), bias corrected pre-whitening (Hamed, 2009) and a bootstrapped optional bias approach (Lacombe et al., 2012). The trend is considered to be significant if at least two of the tests have p-values less than 5% or 10%. Details on these modifications are provided as supplementary information. A multi temporal trend approach is used to trace the change in trend by considering test statistics, p-values and Sen's slopes to measure trend over all possible sub-periods of lengths ranging from 30 to 100 years.

195

2.4.2. Bivariate wavelet analysis

Bivariate wavelet analysis may be used to test for joint periodicity of two time series. Assuming a continuous time series $y(t)$, a wavelet function is defined as (Joshi et al., 2016):

$$\varphi(\omega) = \varphi(s, \tau) = s^{-\frac{1}{2}} \varphi\left(\frac{(t - \tau)}{s}\right) \quad (4)$$

where t is time, τ is the time step in which the window function is iterated, and s is the wavelet scale $(0, \infty)$. The chosen $\varphi(\omega)$ function must have a zero mean and be localised in both time and Fourier space. The cross wavelet transform is given by the convolution of $y(t)$, scaled and translated to give the function $W(s, \tau)$:

200

$$W(s, \tau) = s^{-\frac{1}{2}} \int_{-\infty}^{\infty} y(t) \varphi^*\left(\frac{(t - \tau)}{s}\right) dt \quad (5)$$

The cross wavelet spectrum of two series $y_1(t)$ and $y_2(t)$ is defined as:

$$W_{y_1, y_2}(s, \tau) = W_{y_1}(s, \tau)W_{y_2}^*(s, \tau) \quad (6)$$

where $W_{y_1}(s, \tau)$ and $W_{y_2}(s, \tau)$ are the continuous wavelet transforms of $y_1(t)$ and $y_2(t)$ respectively and $*$ denotes the complex conjugate. Two time series may also be analysed for wavelet coherence by the extraction of common periods of high power. This enables the identification of phase relationships between the two series. The wavelet coherence (R^2) of two time series is defined as:

$$R^2(s) = \frac{\left(\left| S \left(s^{-1} W_{y_1, y_2}(s, \tau) \right) \right|^2 \right)}{S \left(s^{-1} |W_{y_1}(s, \tau)|^2 \right) \cdot S \left(s^{-1} |W_{y_2}(s, \tau)|^2 \right)} \quad (7)$$

where S is a smoothing operator and is analogous to a localised correlation coefficient in time frequency space. The Morlet wavelet transform is recommended for cross wavelet analysis (Grinsted et al., 2004). Relationships between the wet season variables (wet day frequencies, wet spells, dry spells, season onset and season length) and climate modes including SOI, SAM, and the solar cycles (sunspots), were investigated using bivariate wavelet analysis. The time series were detrended to remove the effect of linear trend, and partial wavelet coherence (Ng and Chan, 2012) was calculated where there were correlations between the climate mode indices.

3. Results

3.1. Overview of dry and wet season characteristics

Average rainfall during the wet season (April to October) ranges from 385mm to 1085mm per annum (Table 3) for the four stations, constituting on average 75% of total annual rainfall. For the common period of 1950-2018, stations on average record 484mm pa at Maitland, 531mm pa at Cape Town International (CPT Int.), 608mm pa at the SAAO and 1422mm pa at Kirstenbosch. Such large spatial rainfall variability is a function of topographic influence. Although the SAAO wet season rainfall record is lower than the average across all stations (av =761mm pa), it is closest to this average value. The number of wet season rain days average 57.4 days for all stations over the period 1950-2018, while the SAAO records on average 55.6 rain days during this period and is thus relatively close to this average (Table 4). The dry summer months of November to March receive daily rainfall averaging below 0.7mm/day at the SAAO, Maitland and CPT Int., and *c.* 1.5mm/day at Kirstenbosch. Consequently, there are many dry days (<1mm) during the summer months with long dry spells at all stations.

225 In contrast, the cooler seasons (April to October) receive on average *c.* 2mm/day at Maitland and CPT Int., 2.4mm/day at the SAAO and 5.6mm/day at Kirstenbosch, usually peaking in June (Fig. 2).

The number of wet days per annum (January to December), wet days for the wet season (April to October), and the proportion of wet days during the wet season, relative to the total annual number of wet days, are presented for available records in Fig. 3. Over the period 1841-2018, high frequencies of wet days are recorded for the late 19th century (SAAO) and during the 1940s
230 and 1950s (all stations) (Table 4). In contrast, the period 1925-1940 records low frequencies of wet days (3 stations). Mann Kendall trend test results indicate a significant decline in the number of wet days during the wet season at the SAAO (by -0.2 days/decade over the period 1841-2018) and at Maitland (by -0.7 days/decade over the period 1906-2018) (Table 4). However, such declines are more pronounced in annual rain day records, which includes the dry season. The frequency of high intensity rainfall ($\geq 10\text{mm/day}$) constitutes on average 5% of days during a calendar year at the SAAO, Maitland and Cape Town Int.,
235 and 11.2% of days at Kirstenbosch (see Fig. 3) for the lengths of the records available at each station. Below (above) average frequencies of high intensity rainfall occurred in several years during the 1930s and 1970s (1950s).

A multi temporal trend analysis was performed to identify changes in the linear trends for wet day frequency records at 30-, 50-, 70- and 90-year time scales. The test statistics and Sen's slope at the end of the sub period tested for trend were plotted
240 for all possible overlapping sub-periods observed for each station record (Fig. 4). At the SAAO, significant positive 30-year trends and 50-year trends are observed for the 1890s and 1900s respectively, mainly owing to the 1880s and early 1890s having had a particularly high frequency of wet days compared to previous decades. In addition, several years between 1926 and 1936 recorded low frequencies compared to the period of high wet day frequencies in the 1940s and 1950s, resulting in significant positive 30-year trends in the latter periods. Outputs from other stations confirm these 30-year trends (Fig. 4). Negative trends
245 are observed for sub-periods ending in the last 30 years over 50- and 70-year sub-periods, but these are only significant at the SAAO and Maitland.

3.2. Wet and dry spells during the wet season

The average number of wet spells during the wet season ranges between 28 and 31 for all stations over the years 1950-2018,
250 and average 1.8 days in duration, apart from Kirstenbosch which records a longer average length of 2.3 days. There was a prolonged period of below average frequencies of wet spells and average spell lengths at the SAAO and Maitland stations in the 1920s and 1930s, as well as since 1975 for the SAAO (Fig. 5). Average dry spell lengths are between 4.4 and 5.6 days (Fig. 5), with maximum spell lengths of between 32 and 48 days. The early 20th century and 1940s had below average dry spell lengths which correspond with periods of higher rainfall (Ndebele et al., 2020). Above average dry spell lengths between 1925
255 and 1938 at Maitland are consistent with low rainfall during that period.

A finer-scale assessment of wet/dry spell frequencies at the SAAO is presented in Fig. 6. Wet/dry spell lengths below (and above) the mean spell lengths (≈ 2 days (wet)/ 5days (dry)) are considered and examined over the full study period. The total

number of wet spells significantly decreases over the full study period (1841-2018), and most strongly so during the period 1880-1940. One-day events generally have the highest frequencies, averaging 16.5 spells (std dev = 4.17) over the full study period. The average number of such spells was higher during the period 1941-1957 (av. = 17.2) and lower during the last c. 60 years (1958-2018: av. = 15.9), although this becomes higher again during the 21st century (2000-2018; av. = 17.3). There is a significant decline (-1.4 spells/decade) in one-day wet events during the wet season between 1958 and 1999. Two-day wet spells averaged 8.7 spells (std dev = 2.8) per wet season between 1841 and 2018 but decreased substantially over the period 1880-1957 (-0.52 spells/decade). After having increased during the period 1958-1999 (0.87 spells/decade; av. = 9 spells/wet season), two-day wet spells decreased to average 8.5 spells/wet season since 2000. Longer wet spells (> 2 days) averaged 6 spells/wet season between 1841 and 2018, but have progressively decreased in frequency since the 1960s (2000-2018: av. = 4.65 spells/wet season).

Dry spells lasting ≤ 5 days (short dry spells) average 21.7 spells per wet season (Fig. 6) and range between 9 spells (1872) and 36 spells (1892). A significant decline (-0.2 spells/decade) in short dry spells is detected for the full period 1841-2018, but is most pronounced during the period 1880-1941 (-1.0 spells/decade). Long dry spells (lasting ≥ 6 days) average 10 per wet season (Fig. 6), with fewest recorded in 1880 and 1892 (5 spells each), and most in 1930 and 2003 (16 spells each). Long dry spells on average represent 32% of all dry spells during the wet season for the period 1841-2018, but increase to on average 36% since the new millennium, and hence a tendency toward longer dry spells. A significant negative trend (-0.3 spells/decade) is evident prior to 1900, which subsequently changes to a significant increase (0.1 spells/decade) after 1900. Years with the highest proportion of shorter dry spells (≤ 5 days) include 1883 and 1892 (> 87% of all dry spells), while the highest proportion of dry spells ≥ 6 days (40% of all dry spells) occurred in 2010. The period 1880-1920 is notable for its high number of short dry spells (av. = 24.3 spells/yr) and below normal number of longer (≥ 6 days) dry spells (av. = 9 spells/yr) (Fig. 6). In contrast, since c. 1974, longer dry spells have for the most part become more frequent (av. since 1974 = 10.7 spells/yr).

280 3.3. Extreme wet and dry spells during the wet season (SAAO)

The geometric distribution and compound geometric distribution (see Supplementary information) were used to model wet and dry spell lengths, and to estimate the 95th and 99th percentiles for extreme wet and dry spells. The distribution parameters for the SAAO, Maitland and Cape Town Int. records are similar, and the 95th and 99th percentiles for wet spell lengths of 4 and 5 days respectively. Several years during the late 19th century (1883, 1890, 1892 and 1893) had high frequencies of long wet spells (≥ 4 days) at the SAAO, which coincidentally also recorded high rainfall amounts (see Ndebele et al., 2020). The 1940s and early 1950s had the highest recorded number of such extreme wet spells, during which time seven years experienced six or more wet spells of ≥ 4 days during the wet season at the SAAO. Long wet spells (≥ 4 days) are also noted during the 1940s and 1950s at Maitland and Cape Town International. The 95th percentile for wet spell lengths at Kirstenbosch = 6 days. The 1940s and 1950s stand out as a period with high frequencies of wet spells ≥ 6 days at Kirstenbosch.

290 The 95th and 99th percentiles for dry spell lengths are 14 and 20 days respectively at the SAAO. Long dry spells during the months of either April/May or August/September are associated with the onset/termination of wet seasons respectively, and at times reflect a late/early onset/termination to the wet season respectively. Long dry spells (19-28 days in length) during the mid-wet season (i.e. June/July; Julian days 160-210) were a particular concern in 1929, 1934 and 1935, and coincided with the driest period (1926-1936) since records began in 1841 (see Ndebele et al., 2020). Also noteworthy is the occurrence of long
295 (17 day) dry spells during June/July of recent years (2004, 2005 and 2014).

3.4. Pentad profiles

Decadal pentad profiles for the SAAO are provided so as to obtain a decadal cross-sectional view of the onset, end and distribution of wet season rainfall. Average total values for each pentad over decadal periods (only 9 years for the most recent
300 decade: 2010-2018) were calculated and plotted (Fig. 7). Pentads considered for the wet season are those between pentad 19 (April 1) and pentad 61 (ending 1 November), as these (and 95% confidence intervals) exceed the long-term pentad average (\bar{D}_l) (Fig. 7). Pentads with average values above that of the long-term pentad average are provided in Table 5 and represented in Fig. 8 and Fig. 9. The earliest pentads with mean values above the \bar{D}_l range between pentad 19 (1-5 April) and pentad 30 (26 -30 May); these provide an indication of the average wet season onset during each decade. The latest pentads with mean
305 values above \bar{D}_l range between pentad 51 (8-12 August) and pentad 61 (28 October to 1 November). Coincidentally, the most recent decade had the latest average wet season onset date, while the first decade had the earliest average wet season termination date. Decades recording the highest number of dry pentads (< 5mm) include 1920-1929 (10 pentads) and 2010-2018 (13 pentads); these are also the driest decades with peak pentad averages below 20mm. Decades with the most pentads above \bar{D}_l (32-33 pentads), early wet season onsets and late wet season termination dates, correspond with decades that had
310 high rainfall at the end of the 19th century and the 1940s and 1950s. The most recent decade (2010-2018) had the least number of pentads above \bar{D}_l (22 pentads). During most decades, rainfall peaks between pentads 28 (1-5 May) and 38 (5-9 July). The peak pentad ranges from an average of 18mm (1930-1939; 2010-2018) to 40mm (1860-1869; 1960-1969). Pentad profiles for the other stations are provided in the supplementary information. Similar observations are made for the period 2010-2018, which records a tendency toward an increased number of dry pentads, low peak pentad average rainfall, and shortened wet
315 seasons.

3.5. Rainfall onset/ termination dates

Rainfall onset/termination dates were calculated for each year across the four weather stations, as shown in Fig. 10. Onset days at the four stations are positively correlated (correlations range between 0.53 and 0.77; Table 6) and have similar averages for
320 the common period 1950-2018, ranging between Julian day 103 (13 April at the SAAO) and 106 (16 April at Cape Town Int.

and Kirstenbosch). The latest recorded onset date was at Maitland on Julian day 150 (30 May 1925). Early onsets were common in the 1950s and 1980s (averaging Julian day 100), while prominent late onsets occurred during the mid-1930s (av. = day 112) and early 1970s (av. = day 117). The average termination dates were Julian day 291/292 (18/19 October) across the four stations, resulting in an average wet season length of 185-188 days. The occurrence of early termination days (before 7 October) has become more prevalent during the most recent 40 years (av = day 284) and constitute 80% of termination dates before Julian day 280 during the last 100 years. Rain season lengths were notably shorter during the period 1970-1994 (av = 181 days) for all stations.

For the SAAO, the average rain season onset and termination dates for the period 1841-2018 are Julian days 103 (13 April) and 291 (18 October) respectively. The latest onset date was 23 May 1933, while the earliest termination date was 26 August 1993. During more recent decades (1958-2018) the mean onset and termination dates have shifted to Julian days 105 (15 April) and 288 (14 October) respectively. Early rainfall onsets (before Julian day 100; 10 April) occurred during the late 1840s to early 1860s, 1940-1960, during the 1980s and 2011-2014. Late rainfall onsets (after Julian day 105; 15 April) occurred during the decade 1930-1939, the latter part of the 1960s to early 1970s, and many years post 1997. Wet seasons have typically ended early (on av. before Julian day 290; 17 October) during the most recent decades, starting in 1970. There has been a recent decadal decline in wet season length (-1.4 days/decade) since *c.* 1940, especially between 1941-1994 (-4.2 days/decade), such that wet seasons since 2000 have only averaged 182 days in length while the long-term (1841-2018) average is 188 days.

To test our method of identifying wet seasons, the proportions of annual rainfall accumulated between onset and termination dates were calculated and compared to the totals for January to December (P_{WS}). The average proportion of annual rainfall at the SAAO during the identified wet seasons is 81% (1841-2018) (for days ≥ 1 mm). Over the common period 1950- 2018, the other three stations recorded a combined average proportion of 81% of annual rainfall during wet seasons. The most recent four decades (starting 1978) have had the highest proportion (*c.* 83%) of annual rainfall concentrated in the wet season at the SAAO. In addition, five years since 1988 have recorded $> 90\%$ of annual precipitation during the wet season, which suggests strong inter-seasonal contrasts in rainfall distribution during recent decades.

3.6. Associations with climate modes

Continuous bivariate wavelet analysis was used to investigate the relationships between wet season rainfall variables and possible climate drivers including solar cycles (sunspots), the Southern Annular Mode (SAM), and El Niño Southern Oscillation using the Southern Oscillation Index (SOI). Wet season variables included are wet day frequency, total number of wet spells, short wet spells (< 3 days), long wet spells (> 2 days), short dry spells (< 6 days), long dry spells (> 5 days), season onset day and season length. The averages for indices for months between March and October and subsets of these months were used for SAM, SOI and sunspots. Wavelet coherence between wet season variables and climate indices were analysed in pairs at multiple time scales; these are represented in plots indicating significant coherence at a 5% significance level (Fig. 11). Partial wavelet coherence was considered for SOI and SAM since these indices are correlated.

Figure 11 shows wavelet coherence plots for variables that had significant coherence. Black contour lines in the plots show areas of significance (at 5 % level) of the wavelet coherence. The cone of influence is the blocked area (U shaped) with a white foreground indicating the region where coherence may be inaccurate due to edge effects and zero padding. Significant relationships are observed between the 9-14 year solar cycle and short (≤ 5 days) dry spells (1908-1964), and also long (≥ 6 days) dry spells (1905-1998). The longer 32-40 year solar cycle significantly correlates with wet season onset days and season length between 1920 and 1975. Although the relationship has persisted into the 2000s, it is not significant after 1975, as this time frame is in the cone of influence. Similarly, at time scales between 32-36 years, there is significant partial coherence between rainfall onset days and the SOI. Wet days, season length and short dry spells show evidence of significant partial coherence with the SAM over 6-12 year time scales from 1890-1912, 1930-1962 and 1918-1938 respectively. [The supporting documents](#) provide further details on the dominant time scales for each combination of variables.

4. Discussion

Previous studies examining rainfall onset/termination dates, numbers of rain days, wet/dry spells and wet season lengths over various parts of southern Africa, have done so using composite datasets but over relatively short periods of investigation (< 60 years) (e.g. Tadross et al., 2005; Tadross et al., 2009; Moeletsi et al., 2011; Philippon et al., 2012; MacKellar et al., 2014). Although our datasets cannot represent a wide geographic region given that it is based on only four station records, these offer a substantially longer-term (up to 178 years) perspective on wet season rainfall characteristics than do records spanning only a few decades. A long record such as we have presented here, is particularly valuable as it can identify trends and interannual variability that may not be detected over shorter (< 60 years) periods of investigation, and in addition, offers an opportunity to test associations between such rainfall characteristics and potential causative mechanisms such as ENSO, SAM, and solar cycles.

While there is a general positive trend in wet day frequencies over Cape Town through the 19th century, as reflected in the SAAO data, trends during the 20th and early 21st centuries are broadly negative. Such trends are similar to those observed by Burls *et al.* (2019) who detected a significant decline in wet days since 1900 using data with more than 100 years from weather stations in the wider western Cape region (including SAAO). We identify two periods with significantly high wet day frequencies; 1881-1905 at the SAAO (av. = 63.6 days) and during the 1940s (av = 66.8 days at 3 stations) and 1950s for all stations (av. = 63.2 days). A decline in wet day frequencies is observed during the common period from 1950 to 2018 for all stations, although this is only significant at Maitland (-1.0 days/decade) and the SAAO (-1.0 days/decade) during the wet season. Declines are also evident over the 103-year period starting in 1915, for three stations (excluding CPT Int.). However, this is only significant for the SAAO during the wet season. The longer SAAO record indicates that for the period 1880-1940, there was a significant decline in wet day frequencies for both the wet season (-0.2 days/decade) and annual series (-0.5 days/decade). The average wet day frequencies during wet seasons of most recent decades (1958-2018, av. = 54.9) is similar to that for the period 1841-1877 (av. = 56.3) for the SAAO. However, the average frequency of wet days for the full year

between these periods has changed more significantly from 74.5 days and 69.1 days respectively, accounting for an overall decrease over the period 1841-2018. The annual number of wet days during summer dry seasons has thus decreased more significantly than during wet seasons over Cape Town. Long-term trends presented here for Cape Town are also consistent with observations made for the more recent period (1960-2010) by MacKellar et al. (2014), who identified declining trends in
390 rain days over large parts of South Africa, including the western Cape region. Studies have also projected declining rain days and an increase in dry spells for future decades (e.g. Abiodun et al., 2017).

Our methodology determined the onset and termination of wet seasons by establishing the first and last days on which the 5-day running sum of rainfall amounts exceed the long-term 5-day average rainfall amounts between April and October each year. This method effectively delineates the wet season [length] as supported by the fact that, on average, it represents 81% of
395 rainfall ($\geq 1\text{mm/day}$) captured across all years using daily data from the SAAO. Justification for limiting the season onset and end dates to days between 1 April and 31 October is further demonstrated by establishing mean pentad values and their confidence intervals using available daily data for all four stations. These limits prevent the occurrence of false start/end dates and thus the consequent recording of potential long dry spells at the beginning/end of each wet season. These limits can also be adjusted to suit the local climatology by considering the confidence intervals of mean pentad values. Our method is thus
400 not reliant on the growing season for specific crops, as used by Tadross *et al.* (2005) and Hachigonta *et al.* (2008). This method is reinforced by patterns observed in the pentad profiles across all decades for the period 1841-2018 at the SAAO. In addition, mid-season dry spells do not affect our defining of the wet season, unlike the method employed by Dunning *et al.* (2016).

Our study demonstrates that wet seasons in Cape Town were longest during the periods 1851-1865, 1890-1913 and 1940-1959 (av. > 195 days), and shortest over the periods 1841-1849, 1930-1939, 1963-1990 and 1998-2010 (av. < 181 days), with most
405 recent trends indicating a shortening of the wet season by 1.4 days/decade since 1941. The identified recent shortening of the wet season is consistent with findings from other studies for the southwestern Cape region (e.g. Tadross *et al.*, 2009; du Plessis, 2017; Sousa et al., 2018; Ndebele et al., 2020). But shortening wet seasons during recent decades have also been observed for several other sub-Saharan regions of Africa (e.g. Sarr, 2012; Oguntunde et al., 2014; Dunning et al., 2018; Seregina et al., 2019; Sibanda et al., 2020), and is thus a concern for both summer and winter rainfall regions of the continent. Later onset
410 dates for Cape Town are prevalent during the periods 1924-1939, 1961-1980 and 1999-2010, and are projected to become even further delayed in years to come (Mahlalela et al., 2019). Recent decadal shortening of the Cape peninsular wet season is thus a consequence of both later onset (after Julian day 105; 15 April) and earlier termination (before Julian day 291; 19 October) dates. The pentad profiles provide a decadal overview that combines wet season onset and termination dates, the distribution of rainfall during the wet season, and wet spell lengths. Our data show that some years over the period 2010-2018
415 had incidences of uncharacteristically high rainfall events during summer months, as also observed by de Kock et al. (2021). Although the most recent decade (2010-2018) experienced contracted rainfall seasons, the years 2015-2017 mark an outstanding drought, which was not particularly different to the situation over the period 1930-1939. To this end, it is important to place recent drought perspectives in context with both recent decades and longer-term historic events. The implication from

the longer record is that past periods of shortened rain seasons with drought were followed again by wetter periods, and that
420 patterns in variability continue into the future given their associations with a variety of climate modes.
Several previous studies have considered the association between wet season characteristics and climate modes, such as ENSO
and SAM (e.g. Reason et al., 2005; Tadross et al., 2009; Moeletsi et al., 2011; Raut et al., 2014; Mahlalela et al., 2019), but
have been temporally constrained in their analyses. Given our 178-year rainfall record, we have now ascertained interannual
and interdecadal variability sustained over longer time frames associated with such climate modes. Results demonstrate that
425 quasi-decadal variability (7-10 years) in onset days and season lengths is associated with SAM, while variability in dry spell
lengths (9-14 years) is connected to the solar cycle. Interdecadal variability (32 – 40 years) in wet season onsets and lengths
significantly correlate with long solar cycles of 32-38 years during the 20th century, starting from 1920, as also with SAM.
Should such associations still operate to greater or lesser extent during the 21st century, then some of the recent late onset dates
and shorter seasons might be attributed to the effects of SOI and SAM variability. Such associations suggest that these climate
430 modes, in combination, have influenced seasonal characteristics of rainfall over Cape Town and adjacent regions during the
recent past, and are likely to continue to exercise an influence in future (to a greater or lesser extent).

5. Conclusions

435 Discussions concerning recent, current and future (forthcoming years) wet season characteristics should take cognisance of
the underlying interannual variability associated with climate modes, as discussed in this paper. This has implications for
appropriate forecasting and planning of forthcoming wet seasons in the region, not least because of it being a water-stressed
region and one that is agriculturally (fruit and grain agriculture) important. Agricultural planning and production are strongly
influenced and sensitive to such rainfall patterns given the semi-arid climate. For instance, late onsets of the wet season may
affect the planting phase in crop production, with short wet seasons potentially resulting in low production yields. Critical
440 phases in crop growth may also be negatively impacted by long dry spells during the wet season. To this end, an improved
understanding of regional rainfall patterns associated with the combined role of influencing factors, as demonstrated in this
paper, is thus of value for short- to medium- term precipitation forecasts and associated societal (e.g. agricultural) planning.

Acknowledgements

445 We thank the South African Weather Service for providing rainfall data.

References

Abba Omar, S. and Abiodun, B. J.: Characteristics of cut-off lows during the 2015–2017 drought in the Western Cape, South
Africa, *Atmos. Res.*, 235, 104772, <https://doi.org/10.1016/j.atmosres.2019.104772>, 2020.

- Abiodun, B. J., Adegoke, J., Abatan, A. A., Ibe C. A., Egbebiyi, T. S., Engelbrecht, F., and Pinto, I.: Potential impacts of climate change on extreme precipitation over four African coastal cities, *Clim. Change*, 143(3), 399–413, <https://doi.org/10.1007/s10584-017-2001-5>, 2017.
- Amekudzi, L. K., Yamba, E. I., Preko, K., Asare, E. O., Aryee, J., Baidu, M., and Codjoe, S. N. A.: Variabilities in Rainfall Onset, Cessation and Length of Rainy Season for the Various Agro-Ecological Zones of Ghana, *Climate*, 3(2), 416–434, <https://doi.org/10.3390/cli3020416>, 2015.
- 455 Boyard-Micheau, J., Camberlin, P., Philippon, N., and Moron, V.: Regional-Scale Rainy Season Onset Detection: A New Approach Based on Multivariate Analysis, *J. Clim.*, 26(22), 8916–8928, <https://doi.org/10.1175/JCLI-D-12-00730.1>, 2013.
- Burls, N. J., Blamey, R. C., Cash, B. A., Swenson, E. T., Fahad, A. al, Bopape, M-J. M., Straus, D. M., and Reason, C. J. C.: The Cape Town “Day Zero” drought and Hadley cell expansion, *NPJ Clim. Atmos. Sci.*, 2(1), 1–8, <https://doi.org/10.1038/s41612-019-0084-6>, 2019.
- 460 Byakatonda, J., Parida, B. P., Kenabatho, P. K., and Moalafhi, D. B.: Prediction of onset and cessation of austral summer rainfall and dry spell frequency analysis in semiarid Botswana. *Theor. Appl. Climatol.*, 135(1), 101–117, <https://doi.org/10.1007/s00704-017-2358-4>, 2019.
- Camberlin, P., Moron, V., Okoola, R., Philippon, N., and Gitau, W.: Components of rainy seasons’ variability in Equatorial East Africa: onset, cessation, rainfall frequency and intensity, *Theor. Appl. Climatol.*, 98(3), 237–249, <https://doi.org/10.1007/s00704-009-0113-1>, 2009.
- 465 De Kock, W. M. D., Blamey, R. C., and Reason, C. J. C.: Large Summer Rainfall Events and Their Importance in Mitigating Droughts over the South Western Cape, South Africa, *J. Hydrometeorol.*, 22(3), 587–599, <https://doi.org/10.1175/JHM-D-20-0123.1>, 2021.
- Dieppois, B., Pohl B., Rouault, M., New, M., Lawler, D., and Keenlyside, N.: Interannual to interdecadal variability of winter and summer southern African rainfall, and their teleconnections, *J. Geophys. Res. Atmos.*, 121(11), 6215–6239, <https://doi.org/10.1002/2015JD024576>, 2016.
- 470 du Plessis, J. A.: An investigation into the evidence of seasonal rainfall pattern shifts in the Western Cape, South Africa, *J. S. Afr. Inst. Civ. Eng.*, 59(4), 47–55, <https://doi.org/10.17159/2309-8775/2017/v59n4a5>, 2017.
- Dunning, C. M., Black, E., and Allan, R. P.: Later Wet Seasons with More Intense Rainfall over Africa under Future Climate Change, *J. Clim.*, 31(23), 9719–9738, <https://doi.org/10.1175/JCLI-D-18-0102.1>, 2018.
- 475

- Dunning, C. M., Black, E. C. L., and Allan, R. P.: The onset and cessation of seasonal rainfall over Africa, *J. Geophys. Res. Atmos.*, 121(19), 11,405–11,424, <https://doi.org/10.1002/2016JD025428>, 2016.
- Dyer, T. G. J.: Solar Activity and Rainfall Variation over Southern Africa, *S. Afr. J. Sci.*, 71(12), 369, 1975.
- 480 Froidurot, S. and Diedhiou, A.: 2017. Characteristics of wet and dry spells in the West African monsoon system, *Atmos. Sci. Lett.*, 18(3), 125–131, <https://doi.org/10.1002/asl.734>, 2017.
- Giráldez, L., Silva, Y., Zubieta R., and Sulca J.: Change of the Rainfall Seasonality Over Central Peruvian Andes: Onset, End, Duration and Its Relationship With Large-Scale Atmospheric Circulation, *Climate*, 8(2), 23, <https://doi.org/10.3390/cli8020023>, 2020.
- Gong, D. and Wang, S.: Definition of Antarctic Oscillation index, *Geophys. Res. Lett.*, 26(4), 459–462. 485 <https://doi.org/10.1029/1999GL900003>, 1999.
- Grinsted, A., Moore, J. C., and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, *Nonlinear Process. Geophys.*, 11(5/6), 561–566, 2004.
- Hachigonta, S., Reason, C. J. C., and Tadross, M.: An analysis of onset date and rainy season duration over Zambia, *Theor. Appl. Clim.*, 91(1), 229–243, <https://doi.org/10.1007/s00704-007-0306-4>, 2008.
- 490 Hamed, K. H.: Enhancing the effectiveness of prewhitening in trend analysis of hydrologic data, *J. Hydrol*, 368(1), 143–155, <https://doi.org/10.1016/j.jhydrol.2009.01.040>, 2009.
- Hamed, K. H. and Ramachandra Rao, A.: A modified Mann-Kendall trend test for autocorrelated data., *J. Hydrol.*, 204(1), 182–196, [https://doi.org/10.1016/S0022-1694\(97\)00125-X](https://doi.org/10.1016/S0022-1694(97)00125-X), 1998.
- Joshi, N., Gupta, D., Suryavanshi, S., Adamowski, J., and Madramootoo, C. A.: Analysis of trends and dominant 495 periodicities in drought variables in India: A wavelet transform based approach, *Atmos. Res.*, 182: 200–220., <https://doi.org/10.1016/j.atmosres.2016.07.030>, 2016.
- Kendall, M. G.: 1948. *Rank correlation methods*, Griffin: Oxford, England, 1948.
- Knoben, W. J. M., Woods, R. A., and Freer, J. E.: Global bimodal precipitation seasonality: A systematic overview, *Int. J. Climatol.*, 39(1), 558–567, <https://doi.org/10.1002/joc.5786>, 2019.
- 500 Kruger, A. C.: Observed trends in daily precipitation indices in South Africa: 1910–2004, *Int. J. Climatol.*, 26(15), –2285, <https://doi.org/10.1002/joc.1368>, 2006.

- Kusunoki, S., Mizuta, R., and Matsueda, M.: Future changes in the East Asian rain band projected by global atmospheric models with 20-km and 60-km grid size, *Clim. Dyn.*, 37(11), 2481–2493, <https://doi.org/10.1007/s00382-011-1000-x>, 2011.
- 505 Lacombe, G., McCartney, M., and Forkuor, G.: Drying climate in Ghana over the period 1960–2005: evidence from the resampling-based Mann-Kendall test at local and regional levels, *Hydrol. Sci. J.*, 57(8), 1594–1609, <https://doi.org/10.1080/02626667.2012.728291>, 2012.
- Laurenz, L., Lüdecke, H-J., and Lüning, S.: Influence of solar activity changes on European rainfall, *J. Atmos. Sol-Terr. Phys.*, 185, 29–42, <https://doi.org/10.1016/j.jastp.2019.01.012>, 2019.
- Liebmann, B., Bladé, I., Kiladis, G. N., Carvalho, L. M. V., B. Senay, G., Allured, D., Leroux, S., and Funk, C.: Seasonality 510 of African Precipitation from 1996 to 2009, *J. Clim.*, 25(12), 4304–4322, <https://doi.org/10.1175/JCLI-D-11-00157.1>, 2012.
- Liebmann, B. and Marengo, José A.: Interannual Variability of the Rainy Season and Rainfall in the Brazilian Amazon Basin. *J. Clim.*, 14(22), 4308–4318, [https://doi.org/10.1175/1520-0442\(2001\)014<4308:IVOTRS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<4308:IVOTRS>2.0.CO;2), 2001.
- MacKellar, N., New, M., and Jack, C.: Observed and modelled trends in rainfall and temperature for South Africa: 1960-2010, *S. Afri. J. Sci.*, 110(7–8), 1–13, 2014.
- 515 Mahlalela, P. T., Blamey, R. C., and Reason, C. J. C.: Mechanisms behind early winter rainfall variability in the southwestern Cape, South Africa, *Clim. Dyn.*, 53(1–2), 21–39, <https://doi.org/10.1007/s00382-018-4571-y>, 2019.
- Mann, H. B.: Nonparametric Tests Against Trend, *Econometrica*, 13(3), 245–259, <https://doi.org/10.2307/1907187>, 1945.
- Marengo, J. A., Liebmann, B., Kousky, V. E., Filizola, N. P., and Wainer, I. C.: Onset and End of the Rainy Season in the Brazilian Amazon Basin, *J. Clim.*, 14(5), 833–852, [https://doi.org/10.1175/1520-0442\(2001\)014<0833:OAEOTR>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<0833:OAEOTR>2.0.CO;2), 520 2001.
- Martin-Vide, J. and Gomez, L.: Regionalization of Peninsular Spain based on the length of dry spells, *Int. J. Climatol.*, 19(5), 537–555, [https://doi.org/10.1002/\(SICI\)1097-0088\(199904\)19:5<537::AID-JOC371>3.0.CO;2-X](https://doi.org/10.1002/(SICI)1097-0088(199904)19:5<537::AID-JOC371>3.0.CO;2-X), 1999.
- Mason, S. J. and Tyson, P. D.: The modulation of sea surface temperature and rainfall associations over southern Africa with solar activity and the quasi-biennial oscillation, *J. Geophys. Res. Atmos.*, 97(D5), 5847–5856, 525 <https://doi.org/10.1029/91JD02189>, 1992.
- Moeletsi, M. E., Walker, S., and Landman, W. A.: ENSO and implications on rainfall characteristics with reference to maize production in the Free State Province of South Africa, *Phys. Chem. Earth*, 36(14), 715–726, <https://doi.org/10.1016/j.pce.2011.07.043>, 2011.

- Moron, V. and Robertson, A. W.: Interannual variability of Indian summer monsoon rainfall onset date at local scale, *Int. J. Climatol.*, 34(4), 1050–1061, <https://doi.org/10.1002/joc.3745>, 2014.
- 530 Mugalavai, E. M., Kipkorir, E. C., Raes, D., and Rao M. S.: Analysis of rainfall onset, cessation and length of growing season for western Kenya, *Agric. For. Meteorol.*, 148(6), 1123–1135, <https://doi.org/10.1016/j.agrformet.2008.02.013>, 2008.
- Mupangwa, W., Walker, S., and Twomlow, S.: Start, end and dry spells of the growing season in semi-arid southern Zimbabwe, *J. Arid Environ.*, 75(11), 1097–1104, <https://doi.org/10.1016/j.jaridenv.2011.05.011>, 2011.
- 535 Muthuwatta, L., Perera, H. P. T. W., Eriyagama, N., Surangika, K. B. N. U., and Premachandra, W. W.: Trend and variability of rainfall in two river basins in Sri Lanka: an analysis of meteorological data and farmers' perceptions, *Water Int.*, 42(8), 981–999, <https://doi.org/10.1080/02508060.2017.1406784>, 2017.
- Nastos, P. T. and Zerefos, C. S.: Spatial and temporal variability of consecutive dry and wet days in Greece, *Atmos. Res.*, 94(4), 616–628, <https://doi.org/10.1016/j.atmosres.2009.03.009>, 2009.
- 540 Ndebele, N. E., Grab, S., and Turasie, A.: Characterizing rainfall in the south-western Cape, South Africa: 1841–2016, *Int. J. Climatol.*, 40(4), 1992–2014, <https://doi.org/10.1002/joc.6314>, 2020.
- Ng, E. K. W. and Chan, J. C. L.: Geophysical Applications of Partial Wavelet Coherence and Multiple Wavelet Coherence, *J. Atmos. Ocean. Technol.*, AMS, 29(12), 1845–1853, <https://doi.org/10.1175/JTECH-D-12-00056.1>, 2012.
- Ngetich, K. F., Mucheru-Muna, M., Mugwe, J. N., Shisanya, C. A., Diels, J., and Mugendi, D. N.: Length of growing season, rainfall temporal distribution, onset and cessation dates in the Kenyan highlands, *Agric. For. Meteorol.*, 188, 24–32, <https://doi.org/10.1016/j.agrformet.2013.12.011>, 2014.
- 545 Nicholls, N.: A system for predicting the onset of the north Australian wet-season, *J. Climatol.*, 4(4), 425–435. <https://doi.org/10.1002/joc.3370040407>, 1984.
- Nitka, W. and Burnecki, K.: Impact of solar activity on precipitation in the United States, *Phys. A: Stat. Mech. Appl.*, 527, 121387, <https://doi.org/10.1016/j.physa.2019.121387>, 2019.
- 550 Obarein, O. A. and Amanambu, A. C.: Rainfall timing: variation, characteristics, coherence, and interrelationships in Nigeria, *Theor. Appl. Climatol.*, 137(3–4): 2607–2621, <https://doi.org/10.1007/s00704-018-2731-y>, 2019.
- Odekunle, T. O.: Rainfall and the length of the growing season in Nigeria., *Int. J. Climatol.*, 24(4), 467–479, <https://doi.org/10.1002/joc.1012>, 2004.

- 555 Oguntunde, P. G., Lischeid, G., Abiodun, B. J., and Dietrich, O.: Analysis of spatial and temporal patterns in onset, cessation and length of growing season in Nigeria, *Agric. For. Meteorol.*, 194, 77–87, <https://doi.org/10.1016/j.agrformet.2014.03.017>, 2014.
- Owusu, K. and Waylen, P. R.: The changing rainy season climatology of mid-Ghana, *Theor. Appl. Climatol.*, 112(3), 419–430, <https://doi.org/10.1007/s00704-012-0736-5>, 2013.
- 560 Philippon, N., Rouault, M., Richard, Y., and Favre, A.: The influence of ENSO on winter rainfall in South Africa, *Int. J. Climatol.*, 32(15), 2333–2347, <https://doi.org/10.1002/joc.3403>, 2012.
- Raut, B. A., Jakob, C., and Reeder, M. J.: Rainfall Changes over Southwestern Australia and Their Relationship to the Southern Annular Mode and ENSO, *J. Clim.*, 27(15), 5801–5814, <https://doi.org/10.1175/JCLI-D-13-00773.1>, 2014.
- Reason, C. J. C., Hachigonta, S., and Phaladi, R. F.: Interannual variability in rainy season characteristics over the Limpopo region of southern Africa, *Int. J. Climatol.*, 25(14), 1835–1853, <https://doi.org/10.1002/joc.1228>, 2005.
- 565 Reason, C. J. C. and Jagadheesha, D.: Relationships between South Atlantic SST Variability and Atmospheric Circulation over the South African Region during Austral Winter, *J. Clim.*, A. M S., 18(16), 3339–3355, <https://doi.org/10.1175/JCLI3474.1>, 2005.
- Reason, C. J. C. and Rouault, M.: ENSO-like decadal variability and South African rainfall, *Geophys. Res. Lett.*, John Wiley & Sons, Ltd, 29(13), 16–1, <https://doi.org/10.1029/2002GL014663>, 2002.
- 570 Reason C. J. C. and Rouault M.: Links between the Antarctic Oscillation and winter rainfall over western South Africa, *Geophys. Res. Lett.*, 32(7), <https://doi.org/10.1029/2005GL022419>, 2005.
- Reason, C. J. C., Rouault, M., Melice, J-L., and Jagadheesha, D.: Interannual winter rainfall variability in SW South Africa and large scale ocean–atmosphere interactions, *Meteorol. Atmos. Phys.*, 80(1), 19–29, <https://doi.org/10.1007/s007030200011>, 2002.
- 575 Rivoire, P., Trambly, Y., Neppel, L., Hertig, E., and Vicente-Serrano, S. M.: Impact of the dry-day definition on Mediterranean extreme dry-spell analysis, *Nat. Haz. Earth Sys. Sci.*, 19(8), 1629–1638, <https://doi.org/info:doi:10.5194/nhess-19-1629-2019>, 2019.
- Saeed, F., Bethke, I., Fischer, E., Legutke, S., Shiogama, H., Stone, D. A., and Schleussner, C-F. 2018: Robust changes in tropical rainy season length at 1.5 °C and 2 °C, *Environ. Res. Lett.*, 13(6), 064024, <https://doi.org/10.1088/1748-9326/aab797>, 2018.
- 580

- Sarr, B.: Present and future climate change in the semi-arid region of West Africa: a crucial input for practical adaptation in agriculture, *Atmos. Sci. Lett.*, 13(2), 108–112, <https://doi.org/10.1002/asl.368>, 2012.
- 585 Seregina, L. S., Fink, A. H., van der Linden, R., Elagib, N. A., and Pinto, J. G.: A new and flexible rainy season definition: Validation for the Greater Horn of Africa and application to rainfall trends, *Int. J. Climatol.*, 39(2), 989–1012, <https://doi.org/10.1002/joc.5856>, 2019.
- Sibanda, S., Grab, S. W., and Ahmed, F.: Long-term rainfall characteristics in the Mzingwane catchment of south-western Zimbabwe, *Theor. Appl. Climatol.*, 139(3), 935–948, <https://doi.org/10.1007/s00704-019-03020-6>, 2020.
- 590 Singh, N. and Ranade, A.: The Wet and Dry Spells across India during 1951–2007, *J. Hydrometeorol.*, 11(1), 26–45, <https://doi.org/10.1175/2009JHM1161.1>, 2010.
- Tadross, M., Suarez, P., Lotsch, A., Hachigonta, S. P., Mdoka, M. L., Unganai, L. S., Lucio F., Kamdonyo, D., and Muchinda, M.: Growing-season rainfall and scenarios of future change in southeast Africa: implications for cultivating maize, *Clim. Res.*, 40(2-3), 147-161, <https://doi.org/10.3354/cr00821>, 2009
- 595 Tadross, M. A., Hewitson, B. C., and Usman, M. T.: The Interannual Variability of the Onset of the Maize Growing Season over South Africa and Zimbabwe, *J. Clim.*, 18(16), 3356–3372, <https://doi.org/10.1175/JCLI3423.1>, 2005
- Tyson, P. D.: Atmospheric circulation variations and the occurrence of extended wet and dry spells over Southern Africa, *Int. J. Climatol.*, 1(2), 115–130, <https://doi.org/10.1002/joc.3370010203>, 1981.
- Valdes-Abellan, J., Pardo, M. A., and Tenza-Abril, A. J.: Observed precipitation trend changes in the western Mediterranean region, *Int. J. Climatol.*, 37(S1), 1285–1296, <https://doi.org/10.1002/joc.4984>, 2017.
- 600 Vanlommel, P., Cugnon, P., van der Linden, R.A.M., Berghmans, D., and Clette, F: The Sidc: World Data Center for the Sunspot Index, *Sol. Phys.*, 224(1), 113–120, <https://doi.org/10.1007/s11207-005-6504-2>, 2004.
- Yue, S. and Wang, C.: The Mann-Kendall Test Modified by Effective Sample Size to Detect Trend in Serially Correlated Hydrological Series, *Water Resour. Manag.*, 18(3), 201–218, <https://doi.org/10.1023/B:WARM.0000043140.61082.60>, 2004.

605

Tables

Table 1: Climate indices

Index	Dates	Frequency	References
Southern Oscillation Index	1866 -2018	Monthly	Ropelewski and Jones (1987)
Southern Annular Mode	1851-2011	Monthly	Gong and Wang (1999)
Solar (sunspot number)	1841-2018	Monthly	Vanlommel et al. (2004)

Table 2: Definitions of terms

Term	Symbol	Calculation
Proportion of wet days > 0mm (per Calendar day)	P_{WD}^1	$P_{WD}^1 = \frac{f_{w1}}{n_d}$
Proportion wet days \geq 1mm (per Calendar day)	P_{WD}^2	$P_{WD}^2 = \frac{f_{w2}}{n_d}$
Average daily rainfall (per Calendar day)	\bar{R}_d	$\bar{R}_d = \frac{\sum_{i=1}^{n_d} R_d^i}{n_d} \text{ for } d = 1, 2, \dots, 365$
Average daily rainfall (per month)	\bar{R}_m	$\bar{R}_m = \frac{\sum_{i=1}^{n_m} R_m^i}{n_m} \text{ for } m = 1, 2, \dots, 12$
Average daily rainfall (per year)	\bar{R}_Y	$\bar{R}_Y^i = \frac{\sum_{d=1}^{365} R_d^i}{365} \text{ for } i = 1, 2, \dots, n_d$
Long term daily average (full record)	\bar{R}_L	$\bar{R}_L = \frac{\sum_{i=1}^{n_d} \bar{R}_Y^i}{n_d}$
Pentad rainfall totals	D_p	$D_p^i = \sum_{d=5(p-1)+1}^{5p} R_d^i \text{ for } p = 1, 2, \dots, 73; \\ i = 1, 2, \dots, n_d$
Pentad averages	\bar{D}_p	$\bar{D}_p = \frac{\sum_{i=1}^{n_d} D_p^i}{5n_d} \text{ for } p = 1, 2, \dots, 73$
Long term pentad average	\bar{D}_1	$\bar{D}_1 = \frac{\sum_{p=1}^{73} \bar{D}_p}{73}$
Wet season length	l_{ws}^i	$l_{ws}^i = W_E - W_o$
Proportion of wet season rainfall	P_{ws}	$P_{ws} = \frac{\sum_{d=ws} R_d}{\sum_{k=1}^{12} R_m}$

Table 3: Information on rainfall stations. The mean total annual rainfall for the rainfall series and mean seasonal rainfall (Apr–Oct) is presented for each station. R = correlation between the Observatory daily rainfall record and that for other stations over common recording periods.

Name	Location	Years	Length	Annual	Apr-Oct	R
Observatory	-33.9330; 18.4770	1841 – 2018	178	619.19	483.6	1
Maitland	-33.9200; 18.5060	1906 – 2018	112	493.96	385.3	0.70
Kirstenbosch	-33.9830; 18.4330	1915 – 2018	104	1384.63	1184	0.63
Cape Town Int.	-33.9670; 18.6000	1950 – 2018	69	531.33	414.4	0.74

625 Table 4: Wet day frequency (days) for various stations over the Cape Town Metropol. Significant (Sig) trends have Mann Kendall (MK) test p-values < 0.10 (at 10% significance level) and p-values < 0.05 (at 5% significance level) for at least 2 of the modified MK tests.

Station (years)	Months	Mean (std. dev)	Min (year)	Max (year)	Sen's slope	MK p-value	Sig. (Y/N)
SAAO (1841-2018)	April to October	57.85 (8.9)	39 (1880)	83 (1892)	-0.2	< 0.10	Y
	Annual	74.59 (10.5)	51 (1880)	101 (1883)	-0.5	< 0.05	Y
Maitland (1906-2018)	April to October	50 (9.9)	31 (1936)	69 (1957)	-0.7	< 0.05	Y
	Annual	62.1 (8.3)	42 (1931)	87 (1913)	-1.0	< 0.05	Y
1915-2018							
SAAO	April to October	56.9 (8.8)	40 (1927)	80 (1941)	-0.74	< 0.05	Y
	Annual	72.7 (10.4)	52 (2000)	100 (1941)	-1.25	< 0.05	Y
Kirstenbosch	April to October	71.4 (8.7)	56 (1985)	91 (1941)	-0.30	> 0.10	N
	Annual	92.5 (10.1)	72 (1979)	117 (1941)	-0.40	> 0.10	N
Maitland	April to October	49.5 (8.3)	31 (1936)	69 (1957)	-0.60	> 0.10	N
	Annual	61.2 (9.7)	42 (1931)	86 (1957)	-0.72	< 0.10	Y
1950-2018							
SAAO	April to October	55.6 (8.3)	42 (1963)	75 (1951)	-1.010	< 0.10	Y
	Annual	70.5 (9.8)	52 (2000)	94 (1951)	-1.43	< 0.05	Y
Kirstenbosch	April to October	71.0 (8.2)	56 (1985)	91 (1970)	-0.83	> 0.10	N
	Annual	92.0 (9.7)	72 (1979)	116 (1970)	-0.87	> 0.10	N
Maitland	April to October	48.9 (7.5)	35 (2015)	69 (1957)	-1.11	< 0.05	Y
	Annual	60.5 (9.4)	42(2000)	86 (1957)	-1.23	< 0.05	Y

Cape Town Int.	April to October	54.1 (8.8)	39 (2015)	76 (1951)	0.00	> 0.10	N
	Annual	67.9 (7.9)	49 (1999)	88 (1951)	-0.30	> 0.10	N

630 **Table 5:** Pentad profiles per decade at the SAAO. Pentad long-term mean = 8.44mm and dry pentads have < 5mm of rain. Pentad 19-24 (1-30 April); Pentad 25-30 (1-30 May); Pentad 51-55 (8 September -2 October); and Pentad 56-61(3 October – 1 November).

Decade	Lowest pentad above mean	Highest pentad above mean	N° of Pentads above mean	N° of dry pentads	Peak pentad	Peak Pentad rainfall
1841 – 1849	20	51	27	7	32	27.7
1850 – 1859	19	57	29	6	28	33.8
1860 – 1869	23	61	30	1	36	35.5
1870 – 1877	22	57	29	4	31	28.4
1880 – 1889	21	57	31	2	34	25.2
1890 – 1899	23	61	32	4	43	30
1900 – 1909	19	60	30	3	33	29.8
1910 – 1919	19	55	30	8	39	29.5
1920 – 1929	24	56	25	10	35	25.8
1930 – 1939	20	56	25	8	44	18.8
1940 – 1949	19	61	32	7	30	33.3
1950 – 1959	20	60	32	3	28	30.1
1960 – 1969	21	58	23	8	32	35.4
1970 – 1979	24	59	27	6	33	27.7
1980 – 1989	20	56	30	8	48	32.6
1990 – 1999	20	59	30	6	35	27.3
2000 – 2009	20	56	33	6	38	31.9
2010 – 2018	30	61	22	13	42	17.4

635 **Table 6:** Wet season characteristic correlations for the four stations: 1950-2018. O = onset day, T = termination day, L= wet season length. 1- Observatory, 2- Maitland, 3 – Kirstenbosch and 4 – Cape Town International. Shaded correlations are for similar characteristics.

	O1	T1	L1	O2	T2	L2	O3	T3	L3	O4	T4	L4
O1	1.000											
T1	0.019	1.000										
L1	-0.616	0.776	1.000									
O2	0.532	0.058	-0.289	1.000								
T2	0.045	0.445	0.322	0.062	1.000							
L2	-0.326	0.306	0.447	-0.631	0.735	1.000						
O3	0.647	-0.021	-0.425	0.603	-0.185	-0.553	1.000					
T3	0.072	0.526	0.369	0.141	0.539	0.323	-0.017	1.000				
L3	-0.380	0.399	0.554	-0.300	0.519	0.607	-0.682	0.743	1.000			
O4	0.685	0.016	-0.419	0.578	-0.075	-0.451	0.765	0.002	-0.511	1.000		
T4	0.003	0.687	0.540	0.013	0.432	0.327	-0.093	0.681	0.560	-0.105	1.000	
L4	-0.439	0.470	0.648	-0.363	0.351	0.519	-0.558	0.475	0.721	-0.718	0.768	1.000





Figure 1: Location of the SAAO and other stations in Cape Town, southwestern Cape of South Africa (Ndebele et al., 2020).



645

Figure 2: Average daily mean for each Julian day calculated over the full record available at each station – Observatory (1841-2018), Maitland (1906-2018), Kirstenbosch (1915-2018) and Cape Town Int. (1950-2018). Day 91 (1 April) and day 305 (31 October) represented the start and end of the wet season (WS) respectively. The minimum for wet day (min. wet day) is 1mm.

650






Figure 3: The annual and wet season (April to October) frequency of wet days ($\geq 1\text{mm}$) and the corresponding proportion relative to the total number of days (365) per year (right vertical axis).



660 Figure 4: The test statistic values for the multi-temporal trends calculated over overlapping 30-, 50-, 70- and 90-year sub-periods for the 4 stations. The end year for each sub-period is given as the horizontal axis and the z test statistic for the modified Mann Kendall trend test applied is plotted against the last year included in the sub-period. The statistic $z = -1.96$ and $z = 1.96$ corresponds with the 5% level of significance (red horizontal dashed lines). Test statistic values above (below) $z = 1.96$ ($z = -1.96$) represent significant trends.



665

Figure 5: Total number of wet and dry spells per wet season (frequency) and mean spell lengths (MSL) in days for wet and dry spells during the wet season.




Figure 6: Wet/dry spell frequencies at the SAO.






Figure 7: Mean pentad rainfall (MR) and 95% confidence intervals. Horizontal dashed blue lines indicate threshold for wet days (5mm) and long-term mean of all pentads respectively. Vertical dashed blue lines the start (pentad 19) and end (pentad 61) of the wet season.



680

Figure 8: Pentad rainfall profiles: mean rainfall (MR) pentad totals for each decade at SAAO.

Observatory Pentad Profiles



685

Figure 9: The horizon plots are plotted for difference in the pentad mean relative to the overall long term pentad mean for the series. Positive values (blue) represent pentad mean values above the overall long term pentad mean and negative values indicate pentad mean values below the overall long term mean. Lighter shades are closer to the long term mean and darker

shades further away from the mean. The contrast between wet and dry season is observed and allows comparison between these seasons across decades. Pentad 19 is the approximate wet season onset and pentad 57 approximate termination.




Figure 10: Onset/termination and length of wet seasons during the wet season.




Figure 11: Wavelet coherence plots: seasonal rainfall characteristics with solar [sunspots]. The scale for shaded colours in the plot represents wavelet coherence and is scaled from lowest (blue) to highest (red). Black contour lines show areas of significance (at 5 % level) of the wavelet coherence. The cone of influence is the blocked area (U shaped) with a white foreground.