

5 kyr of fire history in the High North Atlantic Region: natural variability and ancient human forcing

Delia Segato^{1,2}, Maria Del Carmen Villoslada Hidalgo^{1,2,3}, Ross Edwards⁴, Elena Barbaro^{1,2}, Paul Vallelonga^{5,6}, Helle Astrid Kjær⁵, Marius Simonsen⁵, Bo Vinther⁵, Niccolò Maffezzoli^{1,2}, Roberta Zangrando^{1,2}, Clara Turetta^{1,2}, Dario Battistel^{1,2}, Orri Vésteinsson⁷, Carlo Barbante^{1,2}, and Andrea Spolaor^{1,2}

¹Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University of Venice, Via Torino 155, 30170 Mestre, Italy

²CNR-Institute of Polar Sciences (ISP-CNR), Via Torino 155, 30170 Mestre, Italy

³CIC nanoGUNE BRTA, Tolosa Hiribidea 76, 20018 San Sebastian, Spain

⁴Curtin University, Kent St, Bentley WA 6102, Australia

⁵Centre for Ice and Climate, Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

⁶UWA Oceans Institute, University of Western Australia, Crawley WA 6009, Australia

⁷University of Iceland, Archaeology Department, University of Iceland, Archaeology Department, Sæmundargata 2, 101 Reykjavík, Iceland

Correspondence: Andrea Spolaor (andrea.spolaor@cnr.it)

Abstract. Biomass burning influences global atmospheric chemistry by releasing greenhouse gases and climate-forcing aerosols. There is controversy about the magnitude and timing of Holocene changes in biomass burning emissions from millennial to centennial time scales and, in particular, ~~on~~ about the possible impact of ancient civilizations. Here we present a 5 kyr record of fire activity proxies levoglucosan, black carbon and ammonium measured in the RECAP ice core, drilled in the coastal East Greenland and therefore affected by processes occurring in the High North Atlantic Region. Levoglucosan and ammonium fluxes ~~show high levels~~ are high from 5 to 4.5 kyr (thousand years) followed by an abrupt decline, possibly due to monotonic decline in Northern Hemisphere summer insolation. Levoglucosan and black carbon show an abrupt decline at 1.1 kyr BP (before 2000 AD), suggesting a decline in wildfire regime in ~~the Icelandic territory~~ Iceland due to the extensive land clearing caused by Viking colonizers. ~~A minimum is reached at 0.5 kyr BP for all fire proxies~~ All fire proxies reach a minimum during the half of the last century, after which levoglucosan and ammonium fluxes increase again, in particular over the last 200 years. We find that the fire regime reconstructed from RECAP fluxes seems mainly related to climatic changes, however over the last millennium human activities might ~~have had a substantial influence controlling the occurrence of fire~~ influenced wildfire frequency/occurrence substantially.

1 Introduction

15 Extensive ~~forest fires~~ wildfires, also at high latitudes, have recently generated worldwide attention and raised concerns about the impacts of humans and climate change on ~~wildfire regimes~~ fire regime. During summer 2020 wildfires over the Arctic Circle

emitted 35% more CO₂ than the previous year, with a significant contributor being peatland fires (Witze, 2020). However, little is known about the patterns and driving forces of fire activity in the past. Quantitative observations of wildfires are severely limited both in time and space, and coverage in global datasets based on satellite observations began ~~only in the last decades~~ (Xiao-rui et al., 2005; van der Werf et al., 2006) in 1974 (Chuvieco et al., 2019). In order to understand the climate – human – fire relationship over long timescales, proxy records of biomass burning are invaluable.

Fire is influenced by ~~both human activities~~ human activities, vegetation and climate, and ~~climate and~~ is a key Earth system process (Bowman et al., 2009; Keywood et al., 2013). As a major component of the carbon cycle, fire interacts with the climate system by releasing particulates, greenhouse gases, including CO₂, CO, CH₄, NO_x, and black carbon (Bowman et al., 2009). Climatic conditions (temperature, insolation changes, atmospheric CO₂, precipitation) are the fundamental drivers for the ignition and spread of fire (Andela et al., 2017; Marlon et al., 2008, 2013; Power et al., 2008; Molinari et al., 2018) ~~and understanding the~~. Fuel load and vegetation type also influence fire behaviour, with high levels of biomass burning/fire severity coinciding with the dominance of fire invaders, and lower biomass burning/severity coinciding with the dominance of resisters (Feurdean et al., 2020a). Understanding the causes and consequences of fires is critical for assessing the state of the Earth system, because of the close relationship between fire, vegetation, and climate.

Anthropogenic activities have influenced the environment well before the Industrial revolution, as several studies suggest (Ruddiman, 2003; Doughty, 2013). All the continents (except Antarctica) were settled by the beginning of the Holocene, thus this period provides crucial context for fire-human-climate interactions and an opportunity to disentangle climate and human contributions in changing fire regimes (Marlon et al., 2013). It has been argued that human activities have influenced fire activity for millennia (Marlon et al., 2008; Power et al., 2008). In the earliest phase of agricultural development, farmers used fire to clear land through the slash and burn technique (Ruddiman and Ellis, 2009). With the establishment of the first agricultural societies in Europe in the mid-Holocene, humans substantially altered the European landscape (Price, 2000). The attribution of changes in fire regime to human impact is largely based on the synchronicity of these changes and indicators of human activity, such as changes in erosion rates, vegetation and land use (Marlon et al., 2013). Ruddiman (2003) explicitly argued that land use changes in Eurasia during the early to mid-Holocene could explain increases in CH₄ and CO₂ atmospheric concentrations and that these changes had a significant impact on climate. Fire variability inferred from the NEEM ice core (Greenland) was associated with droughts as well as temperature and summer insolation variability, however from 4 kyr BP fire trends could not be explained without considering human influence in altering vegetation distribution especially in Europe (Zennaro et al., 2015). Sapart et al. (2012) found an increase in pyrogenic CH₄ emissions at the times of the Roman Empire (~~100 BC~~ 2.1 – 300 AD 1.7 kyr BP), the Medieval Climate anomaly (~~800~~ 1.2 – 1200 AD 0.8 kyr BP) and the Little Ice Age (~~1300~~ 0.7 – 1600 AD 0.4 kyr BP) from the analysis of CH₄ isotopic composition of the air trapped in the Greenlandic EUROCORE and NEEM ice cores. The pyrogenic CH₄ increase associated with the Roman Empire was attributed to an increase of charcoal use from metal production and to the contemporary civilizations in China and India. On the other hand, the increase during Medieval time has been associated with both extended droughts in Northern Europe and accelerating deforestation in both Europe and Asia, while the increase detected during the Little Ice Age has been explained by natural wildfires and by rapid land clearance in the Northern Hemisphere (Sapart et al., 2012).

A diverse range of paleo-tracers has been used to reconstruct biomass burning from different climate archives such as fire scars on tree rings, ice core records of gases and aerosol-borne chemicals (Legrand et al., 2016; Rubino et al., 2015) and sedimentary charcoal records (Marlon et al., 2008, 2016; Power et al., 2008). These records reflect a wide range of different regions of fire locations, frequency, distribution and intensity integrated over a wide span of temporal scales (Grieman et al., 2018; Battistel et al., 2018; Lim et al., 2016). Since charcoal particles settle rapidly, one record is only locally representative (in the order of tens of kilometers) and several charcoal records need to be assembled in order to obtain a regional reconstruction of fire activity (Marlon et al., 2008; Blarquez et al., 2014). Furthermore, several regions, like Siberia, are under-represented in the available charcoal reconstructions, reducing their potential on the reconstruction of fire history on wide spatial areas.

Ice cores from polar regions, on the other hand, are extensively used to reconstruct past climate conditions and can provide forest-fire records (Legrand et al., 2016). Although being geographical point measurements, ice cores represent a record of air, moisture and aerosols sourced from regional or even hemispheric scales with up to annual and sub-annual temporal resolution (Zennaro et al., 2015; Legrand et al., 2016; Simonsen et al., 2019). They also have the great advantage to catch rapid events such as volcanoes-volcanic eruptions and wildfires.

Historically, the ice concentrations of some impurities like ammonium (NH_4^+) and K^+ were suggested to be partially influenced by forest fire emissions. However, these compounds are not specific proxies as their background variations also reflect continuous biogenic emissions from vegetation and soils, while only peak values can be associated with biomass burning events (Fischer et al., 2015). K^+ , additionally, has been found to be highly sensitive to contamination, making it difficult to measure its species in ice (Legrand et al., 2016). Recently, however, most of the attention has been given to two specific fire proxies levoglucosan and black carbon (BC), the latter being a specific tracer of biomass burning in the pre-industrial times (Osmont et al., 2019). Despite that NH_4^+ is influenced by biogenic emissions from Greenland ice-free areas, it has been included in this study as further-for comparison.

Levoglucosan (1,6-anhydro- β -D-glucopyranose) is the most abundant monosaccharide anhydride released when cellulose combustion occurs at temperatures $> 300^\circ\text{C}$ (Simoneit, 1999). It is injected in the atmosphere in convective smoke plumes and deposited on glacier surfaces through wet and dry deposition (Gambaro et al., 2008) with a residence time of up to 2 weeks (Bhattarai et al., 2019). Since levoglucosan is strongly water-soluble, it may be leached during the melt-refreeze process, reshaping the post-depositional distribution /signal-(You et al., 2016). Due to its high emission factors and relatively high concentrations in the ambient aerosols it is an ideal marker compound for biomass burning (Hoffmann et al., 2010).

BC is the light-absorbing refractory carbonaceous matter emitted during incomplete combustion of fossil and biofuels in fires ignited by both natural and human sources (McConnell et al., 2007). BC is an important indicator of biomass burning for paleoclimate reconstructions, as it is a specific fire proxy for pre-Industrial times and can be measured in high resolution with Continuous Flow Analysis (McConnell et al., 2007). BC does not refer to a single well-defined compound because carbonaceous aerosols are emitted in the form of a continuum of compounds with different physical and chemical properties. BC has very low chemical reactivity in the atmosphere and its residence time is about one week; its primary removal process is wet deposition, with dry deposition contributing to 15%-40% of the total removal (Bond et al., 2013; Cape et al., 2012; Barrett et al., 2019).

Previous studies found levoglucosan and BC to have similar trends in Greenland ice cores (Legrand et al., 2016), giving further hint that ice core archives provide a complementary tool to charcoal records in examining the link between climate, human influence and fire activity. By now, several records of fire proxies from Greenland ice cores reconstruct fire history for the last few centuries, however very few extend over the past millennia. In this work we present 5 kyr long records of levoglucosan, BC and NH_4^+ from the Renland ice core (71° 18' 18" N, 26° 43' 24" W). The main objective of this paper is to elucidate the role of climate and human activities on fire activity in the almost unstudied area of the High North Atlantic before the Industrial era.

2 Materials and methods

The RECAP (The REnland ice CAP) project retrieved in 2015 a 584-meter ice core drilled to the bedrock on the Renland ice cap. The Renland ice cap is situated in Eastern Greenland on a high elevation plateau on the Renland peninsula in the Scoresbysund fjord (71° 18' 18" N, 26° 43' 24" W). The ice cap is constrained by surrounding topography and its eastern plateau reaches an elevation of 2340 m at its summit. Its coastal location provides important geographic climate information that can be compared with central Greenland ice cores as well as providing a sensitive indicator of changes at the margins of the Greenland ice sheet.

The core was stored frozen and shipped to Europe, where it was cut at AWI (Alfred-Wegener Institute, Bremerhaven, Germany) and processed at the Centre for Ice and Climate (Niels Bohr Institute, University of Copenhagen, Denmark). The samples analyzed for levoglucosan were collected discretely every 55 cm from a continuous ice core melting system (Bigler et al., 2011) as part of the Continuous Flow Analysis (CFA) campaign conducted at the University of Copenhagen in autumn 2015 (Maffezzoli et al., 2019; Simonsen et al., 2019), while BC and NH_4^+ were measured continuously. After collection, the discrete samples were immediately frozen at -20°C and kept in the dark until analysis. In this work we consider the top 482 m of the core, corresponding to the last 5 kyr BP. The chronology is achieved by annual layer counting down to 458.3 m and below this point by volcanic matching to the GICC05 timescale (Simonsen et al., 2019).

2.1 Levoglucosan, BC and NH_4^+ analysis

Levoglucosan was determined using liquid chromatography/negative ion electrospray ionization – tandem mass spectrometry (HPLC/(-)ESI-MS/MS). This analytical method allows the direct injection of melted samples spiked with $^{13}\text{C}_6$ -labelled internal standard into the HPLC instrument, avoiding contamination during pre-analytical steps (Gambaro et al., 2008; Zennaro et al., 2014; Battistel et al., 2018). All pre-analytical steps were performed under a Class-100 clean bench located in a Class-100 clean room at Ca' Foscari University of Venice. Purelab Ultra system (Elga, High Wycombe, U.K.) was used to produce the ultrapure water (18.2 M Ω cm, 0.01 TOC) utilized in all analytical and pre-analytical procedures (i.e. cleaning and decontamination procedures, standard solutions preparation and chromatographic analysis) (Gambaro et al., 2008).

BC analysis was conducted using a BC analyzer (SP2, Droplet Measurement Technologies, Boulder, Colorado) connected to the CFA system, following the method of McConnell et al. (2007). The SP2 measures mass of individual BC particles using

laser-induced incandescence. BC particles absorb sufficient energy as they pass through the laser beam and reach a temperature at which they incandesce. Intensity of the incandescence is measured with a photomultiplier tube and recorded. The mass of an individual BC particle is proportional to the area of the incandescence signal (McConnell et al., 2007).

The analysis of NH_4^+ was performed by fluorescence within the CFA setup (Bigler et al., 2011). The melt water stream flowing at $1 \text{ mL}\cdot\text{min}^{-1}$ was added to a reagent made from 1.29 g O-phthaldehyde ($\text{C}_8\text{H}_6\text{O}_2$), 60 mL Ethanol, 900 mL purified water (MilliQ) and a buffer made from 35.8 g $\text{Na}_2\text{HPO}_4\cdot 12\text{H}_2\text{O}$, 1000 mL MilliQ, 600 μL NaOH (>32%), 100 μL HCHO (>37%) and 0.8 g Na_2SO_3 , following a 1 m-of-meter mixing coil at 80°C and a 0.2 m-meter mixing coil at room temperature. The mixture was excited at 365 nm and detected at 400 nm by means of a photomultiplier based detector (PMT-FL, FIAlab instruments). Three standards were used for calibration based on an $100 \text{ mg}\cdot\text{L}^{-1}$ IC multielement standard (VII, Certipur, Merck) and diluted to 24, 50 and 200 ppb NH_4^+ . The melt rate of the CFA was kept between 4 and $5 \text{ cm}\cdot\text{min}^{-1}$ and with a response time of 12 seconds; the equivalent depth resolution of the NH_4^+ dataset is less than 1 cm.

Based on the age scale released in Simonsen et al. (2019), the levoglucosan record covers the period from 0.089 to 5 kyr BP and a depth of 482 m, the BC record covers the period from 0.4 to 5 kyr BP and a depth of 482 m and the NH_4^+ covers the entire period back to 5 kyr.

2.2 Potential source regions of RECAP fluxes

To identify the potential forest fires regions able to influence the RECAP site, backward trajectories have been computed using the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPPLIT) model (Stein et al., 2015) with GDAS1 meteorological data. GDAS1 dataset is available at $1 \times 1^\circ$ resolution. The HYSPPLIT model was run using the PySPPLIT Python package (Warner, 2018). Trajectories were calculated every 6 h in backward mode at 3 different altitudes (500, 1000 and 2000 m above Renland elevation, 2315 m a.s.l.) for the entire dataset available, that is the period 2006-2019. A 7-days run time was chosen based on estimations of BC maximum residence time (Ramanathan and Carmichael, 2008; Cape et al., 2012).

HYSPPLIT modeling tool is useful to reconstruct the most likely transport areas over recent period for recent periods and the information derived can be extended over a longer timescales timescale (Pre-industrial or Holocene), assuming that the main atmospheric circulation mode has not changed significantly during those periods (Rubino et al., 2015). Stable isotope data ($\delta^{18}\text{O}$) suggest that the circulation pattern of air masses reaching the Greenland Ice Sheet did not significantly change over the last 10 000 years (Vinther et al., 2006) which supports extending to the past back trajectory analysis based on modern conditions.

2.3 Statistical approach

To investigate possible similarities with available time series, we estimated pairwise Pearson correlations among available records. Correlation coefficients were computed after interpolating all series to obtain a 20-years time resolution. 20-year bins have been chosen in order to maximize time resolution among all time series, limited by Northern Hemisphere temperature (Marcott et al., 2013). We investigated the correlation between our series with NEEM levoglucosan flux (Zennaro et al., 2014), Northern Hemisphere temperatures as tracer of main climate variability (Marcott et al., 2013), RECAP $\delta^{18}\text{O}$ (Hughes et al., 2020)

and sedimentary charcoal influx composites as regional fire reconstruction calculated over North Europe, North America, Siberia and more generally for the entire HLNH (High Latitude Northern Hemisphere, lat > 55°), available in the Global Charcoal Database (Blarquez et al., 2014).

155 To understand the mechanisms behind fire regime changes, we examine fire flux step changes and link them with climate and human history. To determine mean changes in time series we conduct the off-line change point analysis using the ruptures Python package (Truong et al., 2019). Three optimal breakpoints were found for levoglucosan flux using the Pelt method; Dynamic programming was then applied setting three breakpoints. As previously discussed, climate and especially summer temperatures have a central role in explaining the trends observed in RECAP fire reconstruction. However, other processes
160 might be of relevance and in order to disentangle the contribution of climate in changing the fire regime, we calculate ratios between levoglucosan and BC fluxes with Northern Hemisphere reconstructed temperature from Marcott et al. (2013): a stable ratio suggests that climate is the main driver, significant variations in the ratio suggest additional processes to be in place. We then perform change point analysis on ratios between normalized levoglucosan and BC fluxes and Northern Hemisphere temperature (Levo/T and BC/T).

165 3 Results

3.1 Results from back-trajectory analysis

Seven day long back trajectories calculated with HYSPLIT model over the period 2006-2019 suggest that the higher frequency of trajectories come from the nearby areas with respect to the Renland site (Figure 2), as also evidenced by Simonsen et al. (2019) and Maffezzoli et al. (2019) studying insoluble dust and sea ice respectively. In addition to the Renland ice cap surrounding area, the coastal area immediately on the North of the Renland site appears to be the most probable source region.
170 The South-East and South-West coastal areas of Greenland and Iceland are also ice-free source areas which provide the densest and shortest trajectories considering the density of gridded points and travelling time. We define the region ranging from the longitudes 60°W – 0° and the latitudes 90°N – 55° N as the region with the most likely source areas of materials arriving at Renland (~~red-black~~ box in Figure 2) and we hereafter call it High North Atlantic Region (HNAR).

175 Other contributors of air masses are North America, Northern Europe and Siberia. Boreal forests of North America and Siberia are also ~~a possible source~~ possible sources of impurities, as well as Northern Europe (Schüpbach et al., 2018), however they require a longer travel path for fire emissions to reach the Renland site compared to coastal Greenland and Iceland and ~~thus are~~ are thus expected to carry only a minor contribution.

3.2 Levoglucosan, BC and NH₄⁺ results

180 RECAP levoglucosan, BC and NH₄⁺ profiles display distinct variability on centennial timescales and do not exhibit a clear trend over the past 5 kyr (Figure S1). The levoglucosan concentrations vary from 0.006 to 0.1 ng·g⁻¹ with a mean (considering

the whole 5 kyr record) of $0.033 \pm 0.002 \text{ ng}\cdot\text{g}^{-1}\text{ng}\cdot\text{g}^{-1}$, BC concentration profile varies from 0.4 to $1.7 \text{ ng}\cdot\text{g}^{-1}$ with a mean of $0.942 \pm 0.006 \text{ ng}\cdot\text{g}^{-1}$ and NH_4^+ varies from 0.09 to $17 \text{ ng}\cdot\text{g}^{-1}$ with a mean of $5.0 \pm 0.2 \text{ ng}\cdot\text{g}^{-1}$.

Since Levoglucosan, BC and NH_4^+ can be deposited in snow and glaciers by both wet and dry deposition (Stohl et al., 185 2007), fluxes were calculated as the product of concentration with annual accumulation and expressed in $\mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$. Renland accumulation (Hughes et al., 2020; Corella et al., 2019; Simonsen et al., 2019) shows a stable profile, indicating that wet deposition ~~didn't~~ did not undergo drastic changes over the period covered by our record (Vinther et al., 2009). Levoglucosan flux (Figure 3a) exhibits the same major features observed in the concentration record, indicating that variations in the levoglucosan series reflect changes in the atmospheric concentration, rather than a change in snow accumulation (Grieman et al., 2018). Levoglucosan flux shows high levels from 5 to 4.5 kyr BP with $31.5 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$ on average. A rather stable deposition flux is determined between 4 to 2 kyr BP while higher variability is determined during the period 2 to 1 kyr BP, with an average of $17 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$. Lower fluxes are found in the period 1 to 0.5 kyr BP with an average of $10 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$. BC flux is stable with an average of $464 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$ over the period 5 to 1 kyr BP (Figure ~~2b~~ 3b). During this period some oscillations are detected with a higher depositional flux determined at ca. 3.6, 3, 2.5, 2 kyr BP. A decrease in BC flux is determined, similarly to levoglucosan, after 1.1 kyr BP with a value of $372 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$. NH_4^+ flux shows higher values from 5 to 4.6 kyr BP with an average of $2889 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$ and a rather stable profile from 4 to 1.25 kyr BP ($2387 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$ on average) and an abrupt decline at 2.7 kyr BP (Figure 3c). Low values ($2037 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$) are detected over the period 0.99 – 0.42 kyr BP with an abrupt increase during the last 200 years, with a maximum flux of $7915 \mu\text{g}\cdot\text{m}^{-2}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$. 190 195

3.3 ~~Statistical Results from statistical~~ analysis

~~To investigate possible similarities with available time series, we estimated pairwise Pearson correlations among available records (Table ??). Correlation coefficients were computed after interpolating all series to obtain a 20-years time resolution. 20-year bins have been chosen in order to maximize time resolution among all time series, limited by Northern Hemisphere temperature (Marcott et al., 2013). We investigated the correlation between our series with NEEEM levoglucosan flux (Zennaro et al., 2014), Northern Hemisphere temperatures as tracer of main climate variability (Marcott et al., 2013), RECAP $\delta^{18}\text{O}$ (Hughes et al., 2020) and sedimentary charcoal influx composites as regional fire reconstruction calculated over North Europe, North America, Siberia and more generally for the entire HLNH (High Latitude Northern Hemisphere, lat $> 55^\circ$), available in the Global Charcoal Database (Blarquez et al., 2014). For the RECAP core, the correlation between Figure 1 shows the correlation matrix computed for the considered time series. The correlation between RECAP levoglucosan and BC fluxes is low but significant ($r = 0.21$, $p < 0.01$), as well as between BC and NH_4^+ ($r = 0.27$, $p < 0.01$). RECAP levoglucosan flux is significantly correlated with Northern Hemisphere reconstructed temperature ($r = 0.46$, $p < 0.001$) as well as with RECAP $\delta^{18}\text{O}$ ($r = 0.33$, $p < 0.01$); RECAP BC flux is also significantly correlated with NH temperature ($r = 0.32$, $p < 0.001$) and with RECAP $\delta^{18}\text{O}$ ($r = 0.29$, $p < 0.01$). No statistically significant correlation has been determined between RECAP biomass burning fluxes and the regional sedimentary charcoal influx composites (Table 1).~~ 200 205 210

~~To understand the mechanisms behind fire regime changes, we examine fire flux step changes and link them with climate and human history. To determine mean changes in time series we conduct the off-line Breakpoints resulting from change point~~ 215

analysis using the ruptures Python package (Truong et al., 2019). Three optimal breakpoints were found for levoglucosan flux using the Pelt method; Dynamic programming was then applied setting three breakpoints. The breakpoints of each time series are showed in Figure 3-4 by a change of color. As for levoglucosan flux, two main step changes are found at 4.45 and 1.15 kyr, where the profile significantly decreases, while at 0.2 kyr BP it increases; regarding the BC flux, one strong step decline is found at 1.03 kyr BP, while for the NH_4^+ flux the main step changes are found at 4.57, 4.37 and 0.12 kyr BP (Figure 4a,c,e). It was not possible to detect the concomitant shift at 1.15 kyr BP on NH_4^+ flux due to absence of data.

As previously discussed, climate and especially summer temperatures have a central role in explaining the trends observed in RECAP fire reconstruction. However, other processes might be of relevance and in order to disentangle the contribution of climate in changing the fire regime, we calculate ratios between levoglucosan and BC fluxes with Northern Hemisphere reconstructed temperature from Marcott et al. (2013): a stable ratio suggests that climate is the main driver, significant variations in the ratio suggest additional processes to be in place. We perform change point analysis also on ratios between normalized levoglucosan and BC fluxes and Northern Hemisphere temperature (Levo/T and BC/T in Figure 4b,d). The RECAP ratio Levo/T ratio presents step changes at 4.49, 1.19 and 0.37 kyr BP, while the ratio BC/T at 4.41, 1.11 and 0.71 kyr BP (Figure 3b,d). The mean shifts at around 1.1 kyr BP of the Levo/T and BC/T differ by 100 years, however while the Levo/T ratio strongly decreases at 4.5 kyr BP, the BC/T flux increases.

4 Discussion

4.1 Climate influence on the fire regime in the High North Atlantic Region

Compared As opposed to other Greenlandic ice cores such as NEEM, the RECAP core is retrieved in at a coastal site, thus it is speculated to be more influenced by local sources in the Holocene (Simonsen et al., 2019). From back trajectory analysis (Figure 2) and available literature (Corella et al., 2019; Simonsen et al., 2019) we infer that impurities arriving at the Renland site mostly originate in the HNAR, as defined in Section 2.2. Our record is therefore a useful archive to evaluate the fire history at the high latitudes of the North Atlantic.

Several studies suggest that climate is the main driver of global biomass burning (Marlon et al., 2008). Elevated summer temperatures and sustained droughts can affect fuel flammability and lead to increased global fire activity over seasonal to centennial timescales (Daniau et al., 2012). Climate conditions also determined forest expansion (and glacier retreat), with positive effects on fuel moisture and a dampening effect on biomass burning (Feurdean et al., 2020b). Significant correlations of RECAP RECAP fire proxies with NH temperature during the past 5 kyr period (Table ?? Figure 1) suggest that climatic conditions play an important role in controlling forest fire activity in the HNAR. Fire reduced throughout the late Holocene in parallel with progressively decreasing summer insolation in the Northern Hemisphere through the late Holocene (Power et al., 2008). Greater-than-present summer insolation resulted in warmer and drier summers in the Northern Hemisphere with a consequent increasing of fire increasing fire activity as showed by records from North America and Europe. Renland, being located at a high latitude, receives minimal or no insolation throughout the winter, meaning that summer insolation dominates

(Hughes et al., 2020). Thus, summer solar input ~~along with temperatures may have the major influence in controlling and~~ temperature strongly influences fire activity in the HNAR.

250 High levels of levoglucosan and NH_4^+ fluxes from 5 to 4.5 kyr BP could be linked to the NH warm temperatures of the ~~early-to-mid Holocene~~ mid-Holocene. In fact, Marcott et al. (2013) find warmer temperatures from 9.5 to 5.5 kyr B.P, followed by a cooling trend from 5.5 kyr BP onwards. Such a cooling trend could explain levoglucosan and NH_4^+ ~~trends decrease~~ from 5 to 4.5 kyr BP. An additional source of NH_4^+ fluxes, however, could also be soil emission from Greenland ice-free areas, especially during summer when the seasonal ice melts (Fischer et al., 2015). RECAP levoglucosan flux ~~shows high fluxes~~ is high at 1.5 kyr BP, ~~with a declining trend in concomitance with elevated RECAP summer $\delta^{18}\text{O}$ (Hughes et al., 2020), and successively declines~~ until 1 kyr BP, where values remain low until ~~0.2 kyr~~ 200 years BP (Figure 3). This agrees with higher RECAP summer $\delta^{18}\text{O}$ trend (Hughes et al., 2020). Marcott et al. (2013) and (Mann et al., 2008) document a cooling trend from a warm interval (~~~1500-1.5 - 1000-1~~ kyr BP) to a cool interval (~~~500-0.5 - 100 years~~ 0.1 kyr BP). The increase in levoglucosan flux over the last 200 years is most likely connected with the increase in NH temperatures and green-house gases. Charcoal

260 records from the NH show a striking decrease in biomass burning from the late nineteenth century to mid-to-late twentieth century (Marlon et al., 2008), however this is below the resolution of the RECAP fire tracer fluxes.

RECAP levoglucosan and BC fluxes do not correlate with NEEM levoglucosan flux (~~Table ??~~ Figure 1) suggesting a different influence area. While Zennaro et al. (2014) suggest that NEEM fire sources are mainly North American and Eurasian boreal sources, RECAP is a coastal core and local sources like coastal Greenland and Iceland likely have a major influence.

265 Furthermore, the 15 kyr NEEM levoglucosan records shows a maximum in biomass burning at ~3.5 to 1.5 kyr BP, associated with high fire levels in North America and Europe and to wetter conditions in North America (Zennaro et al., 2015). Such a trend is not found for RECAP proxies. For the last 2 kyr, a declining trend starting at ~1.5 kyr BP is ~~in common with shared~~ by NEEM and RECAP ~~fore fire~~ proxies, however, during the ~~MWP between 1000-1300 AD (the Medieval Warm Period (1.2 - 0.8 kyr BP))~~ NEEM levoglucosan increases again while RECAP fire proxies ~~keep low values. The little similarities stay low.~~ The weak similarity between NEEM and RECAP fire proxies ~~give further hint~~ further suggests that the two sites ~~likely~~ have different source areas.

270

Based on the Global Charcoal Database, fire regime reconstructions in the Northern Hemisphere have been compiled for North America, North Europe and Siberia regions, as well as for the whole High Latitude Northern Hemisphere ($\text{lat} > 55^\circ$) and some of these could contribute to RECAP levoglucosan flux. Our statistical analysis suggests weak correlations between

275 RECAP biomass burning fluxes and charcoal composites from North America (~~Table ??~~ Figure 1). RECAP levoglucosan and BC are significantly but negatively correlated with HLNH and Northern Europe composite charcoal records. The negative correlation does not explain the trend observed in the RECAP records but suggests instead that different fire regimes might exist between Northern Europe and the HNAR. General circulation patterns and storm tracks mostly West to East likely move ~~the European fires~~ European fire proxies away from Renland, ~~giving further hint~~ further supporting the hypothesis that Northern

280 Europe fire history is not ~~be~~ captured by RECAP ice.

Two regions that might influence fire ~~proxies~~ proxy deposition in the Renland site ~~are yet remain~~ to be considered: Siberia and Iceland. Past information of fire activity in the Siberia region come from few charcoal datasets and ~~the Belukha ice core~~

nitrate and potassium records (Eichler et al., 2011). Comparing all these available proxies over the last 750 years show that the forest fire activity increased between 1600 and 1680 AD (Eichler et al., 2011) ice core records. From increasing abundance of charcoal morphologies of all types in Plotnikovo Mire, frequent local fires are found between 4.8 and 3.9 cal yr BP and 2.8 and 1.5 kyr BP (Feurdean et al., 2020a), while Eichler et al. (2011) find increased fire activity between 400 and 320 BP from ice core nitrate and potassium and charcoal. Similar to the regions of North America and North Europe, no correlation is found between RECAP ice core fire records and the composite charcoal record for the Siberia region. For Iceland, however, no fire history record is available to our knowledge. The contribution from Greenland can be only considered in warm periods of Holocene, when the ice sheet retreats and ~~the coasts are subject to the smoldering of peats.~~ vegetation of grass and shrublands along the coasts.

~~Based~~ We suggest that the main source of fire markers that are deposited at Renland, together with Eastern Greenland, is Iceland. Our hypothesis is based on the absence of correlation between fire tracer fluxes and any charcoal record from the regions investigated as possible sources, the low correlation with NEEM levoglucosan flux and the results from back-trajectory analysis ~~we suggest that the main source of fire markers that are deposited to Renland might be the Icelandic region.~~ Iceland has a high aeolian activity driven by climate, volcanic activity and glacial sediment supply. Atmospheric low-pressure systems are common, sometimes referred as the "Icelandic low", which frequently result in relatively high wind speeds (Arnalds et al., 2016). It is thus likely that RECAP fire proxies ~~respond more to a local influence~~ reflect fire emissions from Iceland rather than a hemispheric trend ~~and are affected by fire emissions from Iceland.~~ The fire history of Iceland has never been documented and the RECAP ice core might represent the first record preserving past Icelandic fire changes. The significant link between levoglucosan and BC with temperature reconstruction as well as RECAP $\delta^{18}\text{O}$ suggests that the change of Icelandic fire regime ~~might have been was~~ mostly driven by climate fluctuations. ~~The presence of birch forests covering~~ Also, although the Icelandic territory is ~~well documented for the last 2.5 kyr (Streeter et al., 2015) and the expansion of vegetation is strongly dependent on climate (Haraldsson and Ólafsdóttir, 2003).~~

~~Although the Icelandic territory is~~ characterized by high humidity and frequent precipitation (Ólafsson et al., 2007) that could limit fire propagation, natural fires might be triggered more easily than in other regions due to the large number of volcanoes and associated eruptions (Butwin et al., 2018).

4.1.1 Additional processes explaining fire proxy variability

Global climate alone may not be the only driver of fire activity. Additional processes are suggested to have an impact on the fire regime (Marlon et al., 2008), such as ~~the~~ anthropogenic activities including land clearance, ice sheet advance/retreat and peat fires accentuated by permafrost thawing, altering the temporal and spatial structure of fuel and the frequency of ignitions since the early Holocene (Pfeiffer et al., 2013; Vanni re et al., 2016; Andela et al., 2017).

From change point analysis conducted on fire fluxes and on ratios with temperature we identify three main changes at approximately 4.5, 1.1 and 0.2 kyr BP, found to be common to more than one time series. In the following sections we formulate hypotheses about possible processes inducing fire regime variability.

4.1.2 ~~Period~~ The period 5 – 4.5 kyr BP: ~~Decline of~~ in summer insolation

One possible process explaining decreasing levoglucosan and NH_4^+ fluxes ~~decline~~ during the period 5 – 4.5 kyr BP is the monotonic Holocene decline in Northern Hemisphere summer insolation. Cooler conditions caused cryosphere expansion and progressively a lowered Equilibrium Line Altitude in the ~~highlands in Iceland~~ Iceland highlands, as well as in the ~~eastern~~ Eastern Greenland, the Baffin Islands, Western Svalbard and Western Norway (Geirsdóttir et al., 2018). High-resolution lacustrine records (Geirsdóttir et al., 2013) indicate that, despite the monotonic decline in summer insolation, Iceland's landscape changes and ice cap expansions were nonlinear with abrupt changes occurring at 5, 4.5 – 4.0, 3, and 1.5 kyr BP. It is possible that the abrupt decline in levoglucosan and NH_4^+ fluxes at 4.5 kyr BP was the result of consistent glacier advance and consequent reduction of vegetation. Evidence of vegetation decline starting at 5 kyr BP is found both in Eastern Greenland and Iceland. Pollen records show the presence of dense dwarf shrubs of low Arctic *Betula nana* in the Scoresby Sund area (Funder, 1978) and Basaltsø (Wagner et al., 2000) from 8 to 5 kyr BP, indicating warm and dry conditions. As regards to Iceland, Eddudóttir (2016) found decreased ~~*Betula pubescens*~~ *Betula pubescens* pollen counts in a lake record from the ~~northwest~~ Northwest highlands, giving further evidence of the retreat of woodland from 6 to 4 kyr BP, which might have caused a reduction in fuel availability. ~~Cooler summers and the reduced availability of fuel~~ Feurdean et al. (2020a) find an association between high levels of biomass burning and the presence of *Betula* species, which are classified as fire invaders (Wirth, 2005). Cooler summers and vegetation type change may thus be the cause of the observed reduction in fire regime in the HNAR.

RECAP levoglucosan and BC fluxes have different trends in the period 5 – 4.5 kyr BP (Figure ~~2,3,3,4~~). One possible explanation is that during the ~~mid-Holocene~~ early Holocene (8 - 5 kyr BP) larger areas of the Greenland coasts and Iceland were dry (Wagner et al., 2000) and ice-free (Geirsdóttir et al., 2009) and thus subject to wildfires linked to smoldering of peats. Such wildfires are fueled by the organic matter contained in the soil and are accentuated by the melting of permafrost. Peat Levoglucosan, being emitted by low-temperature fires, might also capture burning grasslands and shrubs (Kehrwald et al., 2020), that grow densely in the HNAR during the early Holocene. Peat and grass fires along the Greenland coasts ~~might could~~ explain the higher fluxes of levoglucosan and ammonium ~~at this stage in this period~~. BC, instead, is mostly emitted under flaming conditions (Legrand et al., 2016) and there is uncertainty regarding ~~BC as a good~~ the quality of BC proxy to trace peat fires (Jayarathne et al., 2017). Further studies, however, are necessary to understand the mechanisms explaining the diverging behavior of levoglucosan and BC.

The latter part of the Holocene is characterized by unstable conditions in the HNAR. Eastern Greenland vegetation becomes progressively more sparse after 5 kyr BP, with a rise of *Salix* and *Cassiope* dominated poor dwarf shrub heaths, indicating moist and cool conditions (Wagner et al., 2000). Environmental instability starting from 4.2 kyr BP is reported in northwest Iceland, indicated by a sparse presence of *Betula pubescens* (Eddudóttir, 2016). Evidence of *Sorbus cf. aucuparia* pollen suggests that shade-intolerant herbs were replacing birch woodlands (Hallsdóttir and Caseldine, 2005), resulting in more sparse and open vegetation. Some woodland regenerations are seen throughout the late Holocene, the most apparent one at 1.6 kyr BP (Hallsdóttir and Caseldine, 2005). Stable fire fluxes may thus reflect cooler climate conditions with more sparse vegetation in the HNAR.

350 4.1.3 Change at 1.1 kyr BP: Viking colonization of Iceland

The step change ~~on~~of levoglucosan and BC fluxes, Levo/T and BC/T at 1.1 kyr BP may be explained ~~considering~~by human activities in Iceland (Figure 34). Iceland was colonized ~~AD~~from~~from~~ Scandinavia and Britain in the late 9th century AD (Vésteinnsson and McGovern, 2012) and since then it experienced a rapid and comprehensive deforestation in a matter of decades, during which large areas completely lost their woodland cover after the arrival of the first settlers (Erlendsson and Edwards, 2009; Streeter et al., 2015). The early settlers cleared land mainly through ~~felling compared to burning, since charcoal layers have been identified only in a small number of sites~~tree-felling, as inferred from the absence of charcoal layers which would indicate the use of fire either through forest clearance or application of slash-and-burn techniques (Trbojević, 2016). The Viking colonization of Iceland resulted in the loss of more than 25% of the ~~vegetation~~total vegetation cover and to catastrophic soil erosion (Haraldsson and Ólafsdóttir, 2003; Erlendsson and Edwards, 2009)(~~Greipsson, 2012~~)(~~Streeter et al., 2015~~), leading to a reduction of the areas which could support vegetation and that would be burned in case of the triggering of a fire. The decline of RECAP fire proxies at 1.1 kyr BP would thus be a consequence of the reduction of vegetation in Iceland. Although the climatic cooling, strengthened from year 700 BP onwards, ~~might have had the greatest effect on reducing fire regime in the HNAR, this might not be completely explained without considering~~can to a great extent explain the observed reduction in forest fires, the human contribution must also be accounted for. It is likely that the inappropriate human management of natural resources acted in parallel to climatic cooling, causing what is recognized to be one of the first environmental disasters.

4.1.4 The last 200 years: global increase in temperatures and land conversion rates

Levoglucosan flux and Levo/T shift over the last few centuries (Figure 4a,b). Levoglucosan flux changes 200 years BP while Levo/T ~~change is anticipated by~~~170 years~~changes earlier at 370 BP~~. The shift of levoglucosan to higher values is probably related to ~~increases in land cover conversion rates in boreal regions (Zennaro et al., 2014), which act together with~~increases in~~increasing~~ global temperatures and ~~green-house-gases concentrations~~is associated with increased dwarf shrub pollen percentages at the sediment surface in East Greenland (Wagner et al., 2000). The last decades of the nineteenth century were the period of the maximum ~~expansion of population and agriculture~~population expansion and land cover conversion for agricultural purposes in many areas of the Northern Hemisphere (Marlon et al., 2008). Zennaro et al. (2014) found relatively high levoglucosan values in the NEEM ice core during the last two centuries, although a clear assessment was not possible due to lack of data. Several charcoal records from North America, Siberia and Europe evidence an increase since ~~~1800~~200 BP (as also reported in Figure 4f,g,i) and ~~have~~has been attributed both to climate and ~~to the~~ increasing use of fire for land clearance (Marlon et al., 2008, 2016).

The abrupt increase of NH₄⁺ ~~in~~over the last 120 years BP may be attributed to anthropogenic emissions. Wendl et al. (2015) argued that trends in NH₄⁺ concentrations in the Lomonosovfonna-09 ice core (Svalbard) after ~~1940 AD~~60 BP indicate a strong anthropogenic influence mainly from NH₃ emissions from agriculture and livestock in Eurasia. No BC data are available for the last 500 years;~~;~~ however, high values of BC flux are expected since several shallow cores from Svalbard and Greenland

show broad concentration maxima starting from ~~1860 AD~~ 140 BP and are attributed to anthropogenic emissions (Osmont et al., 2018).

5 Conclusions

385 This paper provides 5 kyr records of fire proxies levoglucosan, BC and NH_4^+ in the RECAP ice core in Greenland.

From back trajectory analysis and the comparison with regional fire reconstructions based on charcoal records, we find that the most likely source area of impurities arriving at Renland is the High North Atlantic Region (ranging from the longitudes $60^\circ\text{W} - 0^\circ$ and the latitudes $90^\circ\text{N} - 55^\circ\text{N}$) and comprehends the Greenland Ice Sheet, the coasts of Greenland and Iceland. Iceland, in particular, ~~could have~~ might have had a great influence in driving fire regime changes in RECAP fire proxies due to
390 its position with respect to the Renland site and to the presence of large forested areas throughout the Holocene. In the Northern Hemisphere, Iceland is also the only region which is not up until now covered by fire reconstructions, and RECAP fire proxies ~~might represent~~ could enable the first reconstruction of past Icelandic fire regime.

We find that climate variability is the main control of changes in the High North Atlantic fire regime, and especially temperature driven by summer insolation. A downturn of levoglucosan and NH_4^+ fluxes at 4.5 kyr BP may be associated with the
395 monotonic Holocene decline in Northern Hemisphere summer insolation that resulted in cooler conditions, cryosphere expansion and vegetation reduction in the HNAR. BC shows a different trend in this period, possibly explained by wildfires linked to smoldering of peats along the Greenland coasts due to warmer temperatures and melting of permafrost. While levoglucosan and NH_4^+ can record peat fires, BC is not a good proxy because is emitted during flaming conditions when biomass burns. Further studies are necessary to confirm our hypothesis. During the last millennium the fire regime may have been influenced by
400 the human impact in the Icelandic environment, which was uninhabited before the arrival of the Vikings. ~~The~~ We hypothesize that the massive land clearing and the active fire suppression probably led to a reduction of wildfires. The human impact on the fire regime probably accentuated over the last centuries due to population expansion and increases in land cover conversion rates for agricultural purposes.

Data availability. The dataset will be published in the Pangaea database.

405 *Author contributions.* DS and AS designed the research. MCVH, RZ and EB performed the levoglucosan analysis, HAK, MS, BV and PV processed the ice core samples, run the ammonium analysis and helped in the data interpretation, RE performed black carbon analysis, DS and NM performed and run the back-trajectory analysis, DB, CB, CT, OV, AS and DS contributed to the data interpretation. DS and AS wrote the manuscript with inputs from all the authors.

Competing interests. The authors declare no conflict of interest.

410 *Acknowledgements.* The RECAP ice coring effort was financed by the Danish Research Council through a Sapere Aude grant, the NSF through the Division of Polar Programs, the Alfred Wegener Institute, and the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement 610055 through the Ice2Ice project. This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 689443 via project iCUPE (Integrative and Comprehensive Understanding on Polar Environments). The research leading to these results has received funding from the
415 European Union's Seventh Framework programme ("Ideas" Specific Programme, ERC Advanced Grant) under grant agreement no. 267696 "EARLYhumanIMPACT". We acknowledge the help of ELGA LabWater which provides the ultrapure water used for levoglucosan analysis at Ca' Foscari University of Venice. We thank the two anonymous reviewers whose comments/suggestions helped improve and clarify this manuscript.




Figure 1. Correlation matrix of 5 kyr records with resolution of 20 years. *** indicates $p < 0.01$, ** indicates $p < 0.05$ and * indicates $p < 0.1$. The colorbar represents the correlation from -1 to 1.




Figure 2. Seven-days back trajectories of years 2006-2019 at 500, 1000 and 2000 meters above Renland elevation, indicated as number of endpoints divided by number of trajectories. When more than one point of the same trajectory fall in the same grid cell, they are counted as one. The black box indicates the High North Atlantic Region (HNAR), extending between 60°W – 0° in longitude and 90°N – 55° N in latitude. The red star indicates the Renland site.




Figure 3. Normalized 5 kyr time series of: (a) RECAP levoglucosan flux; (b) RECAP BC flux; (c) RECAP NH_4^+ flux; (d) NEEEM levoglucosan flux (Zennaro et al., 2015); (e) NH temperature reconstruction ($90^\circ - 30^\circ$ latitude) (Marcott et al., 2013); (f-g-h-i) Charcoal composite record of North America, North Europe, Siberia and Northern Hemisphere ($\text{lat} > 55^\circ$), respectively (red lines) with 0.05 and 0.95 confidence levels (gray areas). The black line indicates the time of the Viking colonization of Iceland. Age scale is indicated as 1000 years before 2000.




Figure 4. Change point analysis of: (a) levoglucosan flux; (b) ratio of normalized levoglucosan and NH temperature; (c) BC flux; (d) ratio of normalized levoglucosan and NH temperature; (e) NH_4^+ flux. The change of background color indicates the mean shift. The significant breakpoints are indicated by a change of color. The changes in common to more than one time series are indicated by an arrow.

References

- 420 Andela, N., Morton, D. C., Giglio, L., Chen, Y., van der Werf, G. R., Kasibhatla, P. S., DeFries, R. S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Yue, C., and Randerson, J. T.: A human-driven decline in global burned area, *Science*, 356, 1356–1362, <https://doi.org/10.1126/science.aal4108>, 2017.
- Arnalds, O., Dagsson-Waldhauserova, P., and Olafsson, H.: The Icelandic volcanic aeolian environment: Processes and impacts — A review, *Aeolian Research*, 20, 176 – 195, <https://doi.org/https://doi.org/10.1016/j.aeolia.2016.01.004>, 2016.
- 425 Barrett, T., Ponette-González, A., Rindy, J., and Weathers, K.: Wet deposition of black carbon: A synthesis, *Atmospheric Environment*, 213, <https://doi.org/10.1016/j.atmosenv.2019.06.033>, 2019.
- Battistel, D., Kehrwald, N. M., Zennaro, P., Pellegrino, G., Barbaro, E., Zangrando, R., Pedeli, X. X., Varin, C., Spolaor, A., Vallelonga, P. T., Gambaro, A., and Barbante, C.: High-latitude Southern Hemisphere fire history during the mid- to late Holocene (6000–750 BP), *Climate of the Past*, 14, 871–886, <https://doi.org/10.5194/cp-14-871-2018>, 2018.
- 430 Bhattarai, H., Saikawa, E., Wan, X., Zhu, H., Ram, K., Gao, S., Kang, S., Zhang, Q., Zhang, Y., Wu, G., Wang, X., Kawamura, K., Fu, P., and Cong, Z.: Levoglucosan as a tracer of biomass burning: Recent progress and perspectives, *Atmospheric Research*, 220, 20 – 33, <https://doi.org/https://doi.org/10.1016/j.atmosres.2019.01.004>, 2019.
- Bigler, M., Svensson, A., Kettner, E., Vallelonga, P., Nielsen, M., and Steffensen, J.: Optimization of High-Resolution Continuous Flow Analysis for Transient Climate Signals in Ice Cores, *Environmental Science & Technology*, 45, 4483–9, <https://doi.org/10.1021/es200118j>,
435 2011.
- Blarquez, O., Vannière, B., Marlon, J. R., Daniau, A.-L., Power, M. J., Brewer, S., and Bartlein, P. J.: *paleofire*: An R package to analyse sedimentary charcoal records from the Global Charcoal Database to reconstruct past biomass burning, *Computers & Geosciences*, 72, 255 – 261, <https://doi.org/https://doi.org/10.1016/j.cageo.2014.07.020>, 2014.
- Bond, T., Doherty, S., Fahey, D., Forster, P., Berntsen, T., DeAngelo, B., Flanner, M., Ghan, S., Kärcher, B., Koch, D., Kinne, S.,
440 Kondo, Y., Quinn, P., Sarofim, M., Schultz, M., Michael, S., Venkataraman, C., Zhang, H., Zhang, S., and Zender, C.: Bounding the role of black carbon in the climate system: A Scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D’Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A., Prentice, I. C.,
445 Roos, C. I., Scott, A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S. J.: Fire in the Earth System, *Science*, 324, 481–484, <https://doi.org/10.1126/science.1163886>, 2009.
- Butwin, M., Löwis, S., Pfeffer, M., and Thorsteinsson, T.: The Effects of Volcanic Eruptions on the Frequency of Particulate Matter Suspension Events in Iceland, *Journal of Aerosol Science*, 128, 99–113, <https://doi.org/10.1016/j.jaerosci.2018.12.004>, 2018.
- Cape, J., Coyle, M., and Dumitrescu, P.: The atmospheric lifetime of black carbon, *Atmospheric Environment*, 59, 256 – 263,
450 <https://doi.org/https://doi.org/10.1016/j.atmosenv.2012.05.030>, 2012.
- Chuvieco, E., Mouillot, F., van der Werf, G. R., San Miguel, J., Tanase, M., Koutsias, N., García, M., Yebra, M., Padilla, M., Gitas, I., Heil, A., Hawbaker, T. J., and Giglio, L.: Historical background and current developments for mapping burned area from satellite Earth observation, *Remote Sensing of Environment*, 225, 45–64, <https://doi.org/https://doi.org/10.1016/j.rse.2019.02.013>, 2019.

- Corella, J., Maffezzoli, N., Cuevas, C., Vallelonga, P., Spolaor, A., Cozzi, G., Müller, J., Vinther, B., Barbante, C., Kjær, H., Edwards, R., and Alfonso, S.-L.: Holocene atmospheric iodine evolution over the North Atlantic, *Climate of the Past*, 15, 2019–2030, <https://doi.org/10.5194/cp-15-2019-2019>, 2019.
- Daniau, A., Bartlein, P. J., Harrison, S. P., Prentice, I. C., Brewer, S., Friedlingstein, P., Harrison-Prentice, T. I., Inoue, J., Izumi, K., Marlon, J. R., Mooney, S., Power, M. J., Stevenson, J., Tinner, W., Andrič, M., Atanassova, J., Behling, H., Black, M., Blarquez, O., Brown, K. J., Carcaillet, C., Colhoun, E. A., Colombaroli, D., Davis, B. A. S., D’Costa, D., Dodson, J., Dupont, L., Eshetu, Z., Gavin, D. G., Genries, A., Haberle, S., Hallett, D. J., Hope, G., Horn, S. P., Kassa, T. G., Katamura, F., Kennedy, L. M., Kershaw, P., Krivonogov, S., Long, C., Magri, D., Marinova, E., McKenzie, G. M., Moreno, P. I., Moss, P., Neumann, F. H., Norström, E., Paitre, C., Rius, D., Roberts, N., Robinson, G. S., Sasaki, N., Scott, L., Takahara, H., Terwilliger, V., Thevenon, F., Turner, R., Valsecchi, V. G., Vannièrè, B., Walsh, M., Williams, N., and Zhang, Y.: Predictability of biomass burning in response to climate changes, *Global Biogeochemical Cycles*, 26, <https://doi.org/10.1029/2011GB004249>, 2012.
- Doughty, C. E.: Preindustrial Human Impacts on Global and Regional Environment, *Annual Review of Environment and Resources*, 38, 503–527, <https://doi.org/10.1146/annurev-environ-032012-095147>, 2013.
- Eddudóttir, S.: Holocene Environmental Change in Northwest Iceland, Ph.D. thesis, University of Iceland., 2016.
- Eichler, A., Tinner, W., Brüttsch, S., Olivier, S., Papina, T., and Schwikowski, M.: An ice-core based history of Siberian forest fires since AD 1250, *Quaternary Science Reviews*, 30, 1027–1034, <https://doi.org/10.1016/j.quascirev.2011.02.007>, 2011.
- Erlendsson, E. and Edwards, K. J.: The timing and causes of the final pre-settlement expansion of *Betula pubescens* in Iceland, *The Holocene*, 19, 1083–1091, <https://doi.org/10.1177/0959683609341001>, 2009.
- Feurdean, A., Florescu, G., Tanțău, I., Vannièrè, B., Diaconu, A.-C., Pfeiffer, M., Warren, D., Hutchinson, S. M., Gorina, N., Gałka, M., and Kirpotin, S.: Recent fire regime in the southern boreal forests of western Siberia is unprecedented in the last five millennia, *Quaternary Science Reviews*, 244, 106495, <https://doi.org/https://doi.org/10.1016/j.quascirev.2020.106495>, 2020a.
- Feurdean, A., Vannièrè, B., Finsinger, W., Warren, D., Connor, S. C., Forrest, M., Liakka, J., Panait, A., Werner, C., Andrič, M., Bobek, P., Carter, V. A., Davis, B., Diaconu, A.-C., Dietze, E., Feeser, I., Florescu, G., Gałka, M., Giesecke, T., Jahns, S., Jamrichová, E., Kajukalo, K., Kaplan, J., Karpińska-Kołodziej, M., Kołodziej, P., Kuneš, P., Kupriyanov, D., Lamentowicz, M., Lemmen, C., Magyari, E. K., Marcisz, K., Marinova, E., Niamir, A., Novenko, E., Obremska, M., Pędziszewska, A., Pfeiffer, M., Poska, A., Rösch, M., Słowiński, M., Stančikaitė, M., Szal, M., Święta Musznicka, J., Tanțău, I., Theuerkauf, M., Tonkov, S., Valkó, O., Vassiljev, J., Veski, S., Vincze, I., Wacnik, A., Wiethold, J., and Hickler, T.: Fire hazard modulation by long-term dynamics in land cover and dominant forest type in eastern and central Europe, *Biogeosciences*, 17, 1213–1230, <https://doi.org/10.5194/bg-17-1213-2020>, 2020b.
- Fischer, H., Schüpbach, S., Gfeller, G., Bigler, M., Röthlisberger, R., Erhardt, T., Stocker, T. F., Mulvaney, R., and Wolff, E. W.: Millennial changes in North American wildfire and soil activity over the last glacial cycle, *Nature Geoscience*, 8, 723–727, <https://doi.org/https://doi.org/10.1038/ngeo2495>, 2015.
- Funder, S.: Holocene stratigraphy and vegetation history in the Scoresby Sund area, East Greenland, *Bulletin - Groenlands Geologiske Undersoegelse*, 129, 1–76, https://inis.iaea.org/search/search.aspx?orig_q=RN:10462964, 1978.
- Gambaro, A., Zangrando, R., Gabrielli, P., Barbante, C., and Cescon, P.: Direct Determination of Levoglucosan at the Picogram per Milliliter Level in Antarctic Ice by High-Performance Liquid Chromatography/Electrospray Ionization Triple Quadrupole Mass Spectrometry, *Analytical chemistry*, 80, 1649–1655, <https://doi.org/10.1021/ac701655x>, 2008.

- 490 Geirsdóttir, A., Miller, G. H., Axford, Y., and Sædís Ólafsdóttir: Holocene and latest Pleistocene climate and glacier fluctuations in Iceland, *Quaternary Science Reviews*, 28, 2107–2118, <https://doi.org/https://doi.org/10.1016/j.quascirev.2009.03.013>, holocene and Latest Pleistocene Alpine Glacier Fluctuations: A Global Perspective, 2009.
- Geirsdóttir, A., Miller, G., Larsen, D., and Olafsdottir, S.: Abrupt Holocene climate transitions in the northern North Atlantic region recorded by synchronized lacustrine records in Iceland, *Quaternary Science Reviews*, 70, 48–62, <https://doi.org/10.1016/j.quascirev.2013.03.010>,
495 2013.
- Geirsdóttir, A., Miller, G., Andrews, J., Harning, D., Anderson, L., and Thordarson, T.: The onset of Neoglaciation in Iceland and the 4.2 ka event, *Climate of the Past Discussions*, pp. 1–33, <https://doi.org/10.5194/cp-2018-130>, 2018.
- Greipsson, S.: Catastrophic soil erosion in Iceland: Impact of long-term climate change, compounded natural disturbances and human driven land-use changes, *CATENA*, 98, 41 – 54, <https://doi.org/https://doi.org/10.1016/j.catena.2012.05.015>, 2012.
- 500 Grieman, M. M., Aydin, M., Isaksson, E., Schwikowski, M., and Saltzman, E. S.: Aromatic acids in an Arctic ice core from Svalbard: a proxy record of biomass burning, *Climate of the Past*, 14, 637–651, <https://doi.org/10.5194/cp-14-637-2018>, 2018.
- Hallsdóttir, M. and Caseldine, C. J.: 14. The Holocene vegetation history of Iceland, state-of-the-art and future research, in: Iceland — Modern Processes and Past Environments, vol. 5 of *Developments in Quaternary Sciences*, pp. 319–334, Elsevier, [https://doi.org/https://doi.org/10.1016/S1571-0866\(05\)80016-8](https://doi.org/https://doi.org/10.1016/S1571-0866(05)80016-8), 2005.
- 505 Haraldsson, H. and Ólafsdóttir, R.: Simulating vegetation cover dynamics with regards to long-term climatic variations in sub-Arctic landscapes, *Global and Planetary Change*, 38, 313–325, [https://doi.org/10.1016/S0921-8181\(03\)00114-0](https://doi.org/10.1016/S0921-8181(03)00114-0), 2003.
- Hoffmann, D., Tilgner, A., Iinuma, Y., and Herrmann, H.: Atmospheric stability of levoglucosan: a detailed laboratory and modeling study, *Environmental Science & Technology*, 44, 694–699, <https://doi.org/10.1021/es902476f>, 2010.
- Hughes, A., Jones, T., Vinther, B., Gkinis, V., Stevens, C., Morris, V., Vaughn, B., Holme, C., Markle, B., and White, J.: High-frequency
510 climate variability in the Holocene from a coastal-dome ice core in East-Central Greenland, *Climate of the Past*, 16, 1369–1386, <https://doi.org/10.5194/cp-16-1369-2020>, 2020.
- Jayarathne, T., Stockwell, C., Gilbert, A., Daugherty, K., Cochrane, M., Ryan, K., Putra, E., Saharjo, B., Dwi Nurhayati, A., Albar, I., Yokelson, R., and Stone, E.: Chemical characterization of fine particulate matter emitted by peat fires in Central Kalimantan, Indonesia, during the 2015 El Niño, *Atmospheric Chemistry and Physics Discussions*, pp. 1–47, <https://doi.org/10.5194/acp-2017-608>, 2017.
- 515 Kehrwald, N. M., Jasmann, J. R., Dunham, M. E., Ferris, D. G., Osterberg, E. C., Kennedy, J., Havens, J., Barber, L. B., and Fortner, S. K.: Boreal blazes: biomass burning and vegetation types archived in the Juneau Icefield, *Environmental Research Letters*, 15, 085 005, <https://doi.org/10.1088/1748-9326/ab8fd2>, 2020.
- Keywood, M., Kanakidou, M., Stohl, A., Dentener, F., Grassi, G., Meyer, C. P., Torseth, K., Edwards, D., Thompson, A. M., Lohmann, U., and Burrows, J.: Fire in the Air: Biomass Burning Impacts in a Changing Climate, *Critical Reviews in Environmental Science and
520 Technology*, 43, 40–83, <https://doi.org/10.1080/10643389.2011.604248>, 2013.
- Legrand, M., McConnell, J., Fischer, H., Wolff, E. W., Preunkert, S., Arienzo, M., Chellman, N., Leuenberger, D., Maselli, O., Place, P., Sigl, M., Schüpbach, S., and Flannigan, M.: Boreal fire records in Northern Hemisphere ice cores: a review, *Climate of the Past*, 12, 2033–2059, <https://doi.org/10.5194/cp-12-2033-2016>, 2016.
- Lim, S., Faïn, X., Ginot, P., Mikhalenko, V., Kutuzov, S., Paris, J.-D., Kozachek, A., and Laj, P.: Black carbon variability since preindustrial
525 times in Eastern part of Europe reconstructed from Mt Elbrus, Caucasus ice cores, *Atmospheric Chemistry and Physics Discussions*, pp. 1–30, <https://doi.org/10.5194/acp-2016-804>, 2016.

- Maffezzoli, N., Vallelonga, P., Edwards, R., Saiz-Lopez, A., Turetta, C., Kjær, H. A., Barbante, C., Vinther, B., and Spolaor, A.: A 120 000-year record of sea ice in the North Atlantic?, *Climate of the Past*, 15, 2031–2051, <https://doi.org/10.5194/cp-15-2031-2019>, 2019.
- 530 Mann, M. E., Zhang, Z., Hughes, M. K., Bradley, R. S., Miller, S. K., Rutherford, S., and Ni, F.: Proxy-based reconstructions of hemispheric and global surface temperature variations over the past two millennia, *Proceedings of the National Academy of Sciences*, 105, 13 252–13 257, <https://doi.org/10.1073/pnas.0805721105>, 2008.
- Marcott, S., Shakun, J., Clark, P., and Mix, A.: A Reconstruction of Regional and Global Temperature for the Past 11,300 Years, *Science* (New York, N.Y.), 339, 1198–201, <https://doi.org/10.1126/science.1228026>, 2013.
- 535 Marlon, J. R., Bartlein, P. J., Carcaillet, C., Gavin, D. G., Harrison, S. P., Higuera, P. E., Joos, F., Power, M. J., and Prentice, I. C.: Climate and human influences on global biomass burning over the past two millennia, *Nature Geoscience*, 1, 697–702, <https://doi.org/https://doi.org/10.1038/ngeo313>, 2008.
- Marlon, J. R., Bartlein, P. J., Daniiau, A.-L., Harrison, S. P., Maezumi, S. Y., Power, M. J., Tinner, W., and Vanni re, B.: Global biomass burning: a synthesis and review of Holocene paleofire records and their controls, *Quaternary Science Reviews*, 65, 5 – 25, <https://doi.org/https://doi.org/10.1016/j.quascirev.2012.11.029>, 2013.
- 540 Marlon, J. R., Kelly, R., Daniiau, A.-L., Vanni re, B., Power, M. J., Bartlein, P., Higuera, P., Blarquez, O., Brewer, S., Br ucher, T., Feurdean, A., Romera, G. G., Iglesias, V., Maezumi, S. Y., Magi, B., Courtney Mustaphi, C. J., and Zhihai, T.: Reconstructions of biomass burning from sediment-charcoal records to improve data-model comparisons, *Biogeosciences*, 13, 3225–3244, <https://doi.org/10.5194/bg-13-3225-2016>, 2016.
- 545 McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J. R., Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th-Century Industrial Black Carbon Emissions Altered Arctic Climate Forcing, *Science*, 317, 1381–1384, <https://doi.org/10.1126/science.1144856>, 2007.
- Molinari, C., Lehsten, V., Blarquez, O., Carcaillet, C., Davis, B. A. S., Kaplan, J. O., Clear, J., and Bradshaw, R. H. W.: The climate, the fuel and the land use: Long-term regional variability of biomass burning in boreal forests, *Global Change Biology*, 24, 4929–4945, <https://doi.org/10.1111/gcb.14380>, 2018.
- 550  lafsson, H., Furger, M., and Br ummer, B.: The weather and climate of Iceland, *Meteorologische Zeitschrift*, 16, 5–8, <https://doi.org/10.1127/0941-2948/2007/0185>, 2007.
- Osmont, D., Wendl, I., Schmidely, L., Sigl, M., Vega Riquelme, C., Isaksson, E., and Schwikowski, M.: An 800 year high-resolution black carbon ice-core record from Lomonosovfonna, Svalbard, *Atmospheric Chemistry and Physics Discussions*, pp. 1–30, <https://doi.org/10.5194/acp-2018-244>, 2018.
- 555 Osmont, D., Sigl, M., Eichler, A., Jenk, T., and Schwikowski, M.: A Holocene black carbon ice-core record of biomass burning in the Amazon Basin from Illimani, Bolivia, *Climate of the Past*, 15, 579–592, <https://doi.org/10.5194/cp-15-579-2019>, 2019.
- Pfeiffer, M., Spessa, A., and Kaplan, J.: A model for global biomass burning in preindustrial time: LPJ-LMfire (v1.0), *Geoscientific Model Development*, 6, 643–685, <https://doi.org/10.5194/gmd-6-643-2013>, 2013.
- 560 Power, M., Marlon, J., Ortiz, N., Bartlein, P., Harrison, S., Mayle, F., Ballouche, A., Bradshaw, R., Carcaillet, C., Cordova, C., Mooney, S., Moreno, P., Prentice, I., Thonicke, K., Tinner, W., Whitlock, C., Zhang, Y., Zhao, Y., Ali, A., Anderson, R., Beer, R., Behling, H., Briles, C., Brown, K., Brunelle, A., Bush, M., Camill, P., Chu, G., Clark, J., Colombaroli, D., Connor, S., Daniiau, A., Daniels, M., Dodson, J., Doughty, E., Edwards, M., Finsinger, W., Foster, D., Frechette, J., Gaillard, M., Gavin, D., Gobet, E., Haberle, S., Hallett, D., Higuera, P., Hope, G., Horn, S., Inoue, J., Kaltenrieder, P., Kennedy, L., Kong, Z., Larsen, C., Long, C., Lynch, J., Lynch, E., McGlone, M., Meeks, S., Mensing, S., Meyer, G., Minckley, T., Mohr, J., Nelson, D., New, J., Newnham, R., Noti, R., Oswald, W., Pierce, J., Richard, P., Rowe, C.,

- 565 Sanchez Goñi, M., Shuman, B., Takahara, H., Toney, J., Turney, C., Urrego-Sanchez, D., Umbanhowar, C., Vandergoes, M., Vanniere, B., Vescovi, E., Walsh, M., Wang, X., Williams, N., Wilmshurst, J., and Zhang, J.: Changes in fire regimes since the Last Glacial Maximum: An assessment based on a global synthesis and analysis of charcoal data, *Climate Dynamics*, 30, 887–907, <https://doi.org/10.1007/s00382-007-0334-x>, 2008.
- Price, T. D.: *Europe's First Farmers*, Cambridge University Press, <https://doi.org/10.1017/CBO9780511607851>, 2000.
- 570 Ramanathan, V. and Carmichael, G.: Global and Regional Climate Changes Due to Black Carbon, *Nature Geoscience*, 1, 221–227, <https://doi.org/10.1038/ngeo156>, 2008.
- Rubino, M., D'Onofrio, A., Seki, O., and Bendle, J.: Ice-core records of biomass burning, *The Anthropocene Review*, 3, <https://doi.org/10.1177/2053019615605117>, 2015.
- Ruddiman, W.: The Anthropogenic Greenhouse Era Began Thousands of Years Ago., *Climatic change*, 61, 261–293, <https://doi.org/https://doi.org/10.1023/B:CLIM.0000004577.17928.fa>, 2003.
- 575 Ruddiman, W. and Ellis, E.: Effect Of Per-Capita Land Use Changes On Holocene Forest Clearance And CO₂ Emissions, *Quaternary Science Reviews*, 28, 3011–3015, <https://doi.org/10.1016/j.quascirev.2009.05.022>, 2009.
- Sapart, C., Monteil, G., Prokopiou, M., Wal, R., Kaplan, J., Sperlich, P., Krumhardt, K., Veen, C., Houweling, S., Krol, M., Blunier, T., Sowers, T., Martinerie, P., Witrant, E., Dahl-Jensen, D., and Röckmann, T.: Natural and anthropogenic variations in methane sources
- 580 during the past two millennia, *Nature*, 490, 85–8, <https://doi.org/10.1038/nature11461>, 2012.
- Schüpbach, S., Fischer, H., Bigler, M., Erhardt, T., Gfeller, G., Leuenberger, D., Mini, O., Mulvaney, R., Abram, N., Fleet, L., Frey, M., Thomas, E., Svensson, A., Dahl-Jensen, D., Kettner, E., Kjær, H., Seierstad, I., Steffensen, J., Rasmussen, S., and Wolff, E.: Greenland records of aerosol source and atmospheric lifetime changes from the Eemian to the Holocene, *Nature Communications*, 9, <https://doi.org/10.1038/s41467-018-03924-3>, 2018.
- 585 Simoneit, B. R. T.: A review of biomarker compounds as source indicators and tracers for air pollution, *Environmental Science and Pollution Research*, 6, 159–169, <https://doi.org/https://doi.org/10.1007/BF02987621>, 1999.
- Simonsen, M. F., Baccolo, G., Blunier, T., Borunda, A., Delmonte, B., Frei, R., Goldstein, S., Grinsted, A., Kjær, H. A., Sowers, T., Svensson, A., Vinther, B., Vladimirova, D., Winckler, G., Winstrup, M., and Vallelonga, P.: East Greenland ice core dust record reveals timing of Greenland ice sheet advance and retreat, *Nature Communications*, 10, <https://doi.org/https://doi.org/10.1038/s41467-019-12546-2>, 2019.
- 590 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Stohl, A., Berg, T., Burkhart, J. F., Fjærraa, A. M., Forster, C., Herber, A., Hov, Ø., Lunder, C., McMillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth, K., Treffeisen, R., Virkkunen, K., and Yttri, K. E.: Arctic smoke - record high air
- 595 pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006, *Atmospheric Chemistry and Physics*, 7, 511–534, <https://doi.org/10.5194/acp-7-511-2007>, 2007.
- Streeter, R., Dugmore, A., Lawson, I., Erlendsson, E., and Edwards, K.: The onset of the palaeoanthropocene in Iceland: Changes in complex natural systems, *The Holocene*, DOI: 10.1177/0959683615594468, <https://doi.org/10.1177/0959683615594468>, 2015.
- Trbojević, N.: *The Impact of Settlement on Woodland Resources in Viking Age Iceland*, Ph.D. thesis, The School of Humanities, University of Iceland, 2016.
- 600 Truong, C., Oudre, L., and Vayatis, N.: Selective review of offline change point detection methods, *Signal Processing*, 167, 107 299, <https://doi.org/10.1016/j.sigpro.2019.107299>, 2019.

- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmospheric Chemistry and Physics*, 6, 3423–3441, <https://doi.org/10.5194/acp-6-3423-2006>, 2006.
- Vanni re, B., Blarquez, O., Rius, D., Doyen, E., Br ucher, T., Colombaroli, D., Connor, S., Feurdean, A., Hickler, T., Kaltenrieder, P., Lemmen, C., Leys, B., Massa, C., and Olofsson, J.: 7000-year human legacy of elevation-dependent European fire regimes, *Quaternary Science Reviews*, 132, 206 – 212, <https://doi.org/https://doi.org/10.1016/j.quascirev.2015.11.012>, 2016.
- Vinther, B., Buchardt, S. L., Clausen, H., Dahl-Jensen, D., Johnsen, S., Fisher, D., Koerner, R., Raynaud, D., Lipenkov, V., Andersen, K., Blunier, T., Rasmussen, S., Steffensen, J., and Svensson, A.: Holocene thinning of the Greenland ice sheet, *Nature*, 461, 385–8, <https://doi.org/10.1038/nature08355>, 2009.
- Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M., Steffensen, J. P., Svensson, A., Olsen, J., and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006921>, 2006.
- V steinsson, O. and McGovern, T. H.: The Peopling of Iceland, *Norwegian Archaeological Review*, 45, 206–218, <https://doi.org/10.1080/00293652.2012.721792>, 2012.
- Wagner, B., Melles, M., Hahne, J., Niessen, F., and Hubberten, H.-W.: Holocene climate history of Geographical Society  , East Greenland — evidence from lake sediments, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 160, 45–68, [https://doi.org/https://doi.org/10.1016/S0031-0182\(00\)00046-8](https://doi.org/https://doi.org/10.1016/S0031-0182(00)00046-8), 2000.
- Warner, M.: Introduction to PySPLIT: A Python toolkit for NOAA ARL’s HYSPLIT model, *Computing in Science & Engineering*, 20, 47–62, <https://doi.org/10.1109/MCSE.2017.3301549>, 2018.
- Wendl, I., Eichler, A., Isaksson, E., Martma, T., and Schwikowski, M.: 800-Year ice-core record of nitrogen deposition in Svalbard linked to ocean productivity and biogenic emissions, *Atmospheric Chemistry and Physics*, 15, 7287–7300, <https://doi.org/10.5194/acp-15-7287-2015>, 2015.
- Wirth, C.: Fire Regime and Tree Diversity in Boreal Forests: Implications for the Carbon Cycle, pp. 309–344, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/3-540-26599-6_15, 2005.
- Witze, A.: The Arctic is burning like never before — and that’s bad news for climate change, *Nature*, 585, 336–337, <https://doi.org/https://doi.org/10.1038/d41586-020-02568-y>, 2020.
- Xiao-rui, T., Mcrae, D., Li-fu, S., Ming-yu, W., and Hong, L.: Satellite remote-sensing technologies used in forest fire management, *Journal of Forestry Research*, 16, 73–78, <https://doi.org/https://doi.org/10.1007/BF02856861>, 2005.
- You, C., Yao, T., Xu, B., Xu, C., Zhao, H., and Song, L.: Effects of sources, transport, and postdepositional processes on levoglucosan records in southeastern Tibetan glaciers, *Journal of Geophysical Research: Atmospheres*, 121, 8701–8711, <https://doi.org/10.1002/2016JD024904>, 2016.
- Zennaro, P., Kehrwald, N., McConnell, J. R., Sch pbach, S., Maselli, O. J., Marlon, J., Vallelonga, P., Leuenberger, D., Zangrando, R., Spolaor, A., Borrotti, M., Barbaro, E., Gambaro, A., and Barbante, C.: Fire in ice: two millennia of boreal forest fire history from the Greenland NEEM ice core, *Climate of the Past*, 10, 1905–1924, <https://doi.org/10.5194/cp-10-1905-2014>, 2014.
- Zennaro, P., Kehrwald, N., Marlon, J., Ruddiman, W. F., Br ucher, T., Agostinelli, C., Dahl-Jensen, D., Zangrando, R., Gambaro, A., and Barbante, C.: Europe on fire three thousand years ago: Arson or climate?, *Geophysical Research Letters*, 42, 5023–2033, <https://doi.org/10.1002/2015GL064259>, 2015.