

# Northern Hemisphere atmospheric history of carbon monoxide since preindustrial times reconstructed from multiple Greenland ice cores

Xavier Faïn<sup>1</sup>, Rachael H. Rhodes<sup>2</sup>, Philip Place<sup>3</sup>, Vasilii V. Petrenko<sup>3</sup>, Kévin Fourteau<sup>4</sup>,  
Nathan Chellman<sup>5</sup>, Edward Crosier<sup>3</sup>, Joseph R. McConnell<sup>5</sup>, Edward J. Brook<sup>6</sup>, Thomas Blunier<sup>7</sup>,  
Michel Legrand<sup>8</sup>, and Jérôme Chappellaz<sup>1</sup>

<sup>1</sup>Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

<sup>2</sup>Department of Earth Sciences, University of Cambridge, Cambridge, CB2 3EQ, UK

<sup>3</sup>Department of Earth and Environmental Sciences, University of Rochester, Rochester, NY, USA

<sup>4</sup>Univ. Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Études de la Neige, Grenoble, France

<sup>5</sup>Division of Hydrologic Sciences, Desert Research Institute, Reno, NV 89512, USA

<sup>6</sup>College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR 97331, USA

<sup>7</sup>Centre for Ice and Climate, Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

<sup>8</sup>LISA, UMR CNRS 7583, Université Paris-Est Créteil, Université de Paris, Institut Pierre Simon Laplace (IPSL), Créteil, France

**Correspondence:** Xavier Faïn (xavier.fain@univ-grenoble-alpes.fr)

**Abstract.** Carbon monoxide (CO) is a regulated pollutant and one of the key components determining the oxidizing capacity of the atmosphere. Obtaining a reliable record of atmospheric CO mixing ratios since pre-industrial times is necessary to evaluate climate-chemistry models in conditions different from today and to constrain past CO sources. We present high-resolution measurements of CO mixing ratios from ice cores drilled at five different sites on the Greenland ice sheet which experience a range of snow accumulation rates, mean surface temperatures, and different chemical compositions. An optical-feedback cavity-enhanced absorption spectrometer (OF-CEAS) was coupled to continuous melter systems and operated during four analytical campaigns conducted between 2013 and 2019. Overall, continuous flow analyses (CFA) of CO were carried out on over 700 m of ice. The CFA-based CO measurements exhibit excellent external precision (ranging 3.3–6.6 ppbv,  $1\sigma$ ), and achieve consistently low blanks (ranging from  $4.1 \pm 1.2$  to  $12.6 \pm 4.4$  ppbv), enabling paleo-atmospheric interpretations. However the five CO records all exhibit variability too large and rapid to reflect past atmospheric mixing ratio changes. Complementary tests conducted on discrete ice samples demonstrate that these variations are not artifacts of the analytical method (i.e., production of CO from organics in the ice during melting), but very likely are related to in situ CO production within the ice before analysis. Evaluation of signal resolution and co-investigation of high-resolution records of CO and TOC suggest that past atmospheric CO variations can be extracted from the records' baselines with accumulation rates higher than 20 cm water equivalent per year ( $\text{weq yr}^{-1}$ ). Consistent baseline CO records from four Greenland sites are combined to produce a multisite average ice core reconstruction of past atmospheric CO for the Northern Hemisphere high latitudes, covering the period from 1700 to 1957 CE. Such reconstruction should be taken as an upper bound of past atmospheric CO abundance. From 1700 to 1875 CE, the record reveals stable or slightly increasing values in the 100–115 ppbv range. From 1875 to 1957 CE, the record indicates a monotonic increase from  $114 \pm 4$  ppbv to  $147 \pm 6$  ppbv. The ice-core multisite CO record exhibits an excellent overlap with

20 the atmospheric CO record from Greenland firn air which spans the 1950-2010 time period. The combined ice-core and firn air CO history, spanning 1700-2010 CE provides useful constraints for future model studies of atmospheric changes since the preindustrial period.

## 1 Introduction

Carbon monoxide (CO) is a reactive trace gas that plays a crucial role in the interactions between climate and atmospheric chemistry. CO strongly affects the global oxidative capacity of the atmosphere by acting on the budgets of both hydroxyl radical (OH) and ozone ( $O_3$ ). CO is the principal sink for the tropospheric OH, with up to 40% of the OH radicals reacting with CO in the modern troposphere (Lelieveld et al., 2016). Thus, CO indirectly impacts the lifetime of numerous atmospheric constituents such as methane ( $CH_4$ ), non-methane hydrocarbons (NMHCs), and hydrofluorocarbons (HCFCs). Oxidation of CO by OH ultimately leads to  $CO_2$  production, and in the presence of high levels of nitrogen oxides ( $NO_x$ ) can also result in significant production of tropospheric ozone (Crutzen, 1973). Since CO impacts the atmospheric budgets of greenhouse gases such as  $CO_2$ ,  $CH_4$  and tropospheric ozone, its present-day emissions lead to an indirect radiative forcing of about  $0.23 \pm 0.05 \text{ W m}^{-2}$  (Myhre et al. 2013). Furthermore, CO has another indirect climatic impact through the formation of submicron secondary organic aerosol produced by oxidation of volatile organic compounds (VOCs).

CO is emitted by various surface processes and produced by atmospheric oxidation of different gaseous precursors. Atmospheric oxidation of  $CH_4$  and VOCs represents about half of the sources (Duncan et al., 2007). The surface sources include incomplete combustion of anthropogenic fossil fuels and biofuels (Hoesly et al., 2018), biomass burning (van der Werf et al., 2017), and minor contributions from plant leaves (Tarr et al., 1995; Bruhn et al., 2013) and the ocean (Conte et al., 2019). The oxidation by OH is the dominant sink of CO, which results in a mean global CO tropospheric lifetime of about 2 months (Khalil et al., 1999). However, the CO lifetime strongly varies with latitude and season, ranging from 20–40 days in the tropics and up to 3 months in polar areas (Duncan et al., 2007).

The understanding of the past CO trends is crucial for monitoring how anthropogenic activities have impacted CO emissions, atmospheric chemistry and composition. Over the last decades, analyses of past CO trends have been made possible by a growing number of direct and indirect observations. Monitoring of atmospheric CO, initiated by NOAA (National Oceanic and Atmospheric Administration) in 1988, revealed a global decrease of tropospheric CO by  $\sim 0.5 \text{ ppbv yr}^{-1}$  between 1988 and 1996 (Novelli et al., 1998). The space-born MOPITT (Measurements of Pollution in The Troposphere instrument) has monitored the atmospheric CO burden since 2000 (Deeter et al., 2017), and reveals a declining trend. This trend is more pronounced in the Northern Hemisphere (Zheng et al., 2019) which concentrates more sources related to the global economic activity such as fossil fuel use and CO emission from combustion processes.

Ground based and satellite-derived CO data are only available, however, for the last three decades. Ancient air preserved in glacial ice and firn is thus a unique archive for reconstructing the past atmospheric CO record prior to the 1990s. The firn is the upper layer of an ice sheet where snow is slowly transformed into ice. A large amount of air can be sampled from the interconnected open pores. Mean ages of atmospheric gas increase with firn depth. CO has only been measured in firn air at a

few Northern Hemisphere sites, including Summit, the North Greenland Ice Core Project (NGRIP), and the North Greenland Eemian drilling site (NEEM) (Petrenko et al., 2013). Measurements performed on the Devon Island (Canadian Arctic) firn air showed CO artefacts precluding atmospheric reconstruction (Clark et al., 2007). The three central Greenland firn records allowed for a 60 yrs reconstruction of atmospheric CO and revealed that atmospheric CO mixing ratios in the Arctic following industrialization increased from 145 ppbv in 1950 CE (Common Era) until around the late 1970s with a peak of 155 ppbv, and decreased to 140 ppbv in the late 1990s. This decreasing trend in atmospheric CO mixing ratio reconstructed from firn archive is in agreement with direct observations (ground and satellite based) for the last decades. Isotopic measurements of the NEEM firn air suggested this pattern was driven mainly by a change in CO emissions derived from fossil fuel combustion (Wang et al., 2012).

Analysis of air trapped in bubbles in solid ice below the firn layer is required to extend such reconstruction further back in time, beyond 1950 CE. The pioneering measurements conducted on the Eurocore ice archive (Haan et al., 1996; Haan and Raynaud, 1998) found that Northern Hemisphere (NH) [CO] increased gradually from ~90 ppbv in 1850 CE to 110 ppbv in ~1950 CE and that CO levels were stable at ~90 ppbv from 1625 to 1850 CE. Air older than 1600 CE (below 167 m depth) exhibited higher variability and elevated CO levels (100–180 ppbv) with parallel anomalies in the CO<sub>2</sub> record. This section of the Greenland Eurocore record is thought to reflect in situ CO production rather than an atmospheric signal. Haan and colleagues determined CO concentrations along the Eurocore archive using gas chromatography, including a mercuric oxide detector, combined with wet extractions of the trapped gases from discrete pieces of ice. This protocol was affected by a relatively large blank (typically 10 ppbv), was time consuming and resulted in limited resolution reconstructions.

Over the last decade, Continuous Flow Analyses (CFA) of ice core CH<sub>4</sub> concentrations utilising laser spectroscopy (Stowasser et al., 2012) has become a widely-used tool in palaeoclimatology (e.g., Rhodes et al., 2015). CFA was applied for the first time to CO with the measurement from the NEEM-2011-S1 core (Faïn et al., 2014), as an attempt to reconstruct an atmospheric history for NH CO over the past 1800 yrs. The ultra-high resolution of CFA analysis means it is a powerful tool to distinguish between depth sections impacted by in situ CO production and sections preserving the atmospheric record. Faïn et al. (2014) report stable measurements of CO mixing ratios with an external precision of 7.8 ppbv (1 $\sigma$ ), but a poorly constrained procedural blank and poor accuracy because absolute calibration was not yet possible. The NEEM-2011-S1 CO mixing ratios reported are highly variable throughout the ice core with high frequency (at the annual scale), high amplitude spikes characterizing the record. These CO signals likely result from in situ production occurring within the ice itself, with patterns too abrupt and rapid to reflect atmospheric variability. In situ production coupled with the procedural blank and accuracy problems largely prevented interpretation of the record in terms of atmospheric CO variations.

In this study, we expand on this exploratory investigation by reporting continuous CO data measured on a set of five additional Greenland ice cores. By combining the analysis of these new ice core records, and separating the high frequency CO signals driven by in situ production from baseline concentrations, we reveal in the Arctic and for the last three centuries (i) an upper bound estimate of past atmospheric CO abundance and (ii) atmospheric [CO] trends. Climate-chemistry models and/or Earth System Models can produce simulated atmospheric [CO] at ground level in Central Greenland, from the preindustrial era to present-day. Such models are presently intercompared within the AerChemMIP exercise (Collins et al., 2017). The compar-

ison of the past evolution of Arctic atmospheric CO mixing ratio extracted from Greenland ice archives in the frame of this study with AerChemMIP model outputs is out of the scope of this paper. Such comparison, which should also allow to better  
90 constrain CO emissions inventories of emissions factors, will be addressed in a future study.

## 2 Methods

### 2.1 Sample description

Five ice cores extracted from Greenland (Table 1) were investigated in this study (Fig. S1). High resolution CO mixing ratio data were measured continuously along with those of methane. The methane data for the NEEM-SC, D4, North Greenland  
95 Ice Core Project (NGRIP), and Tunu13 sites have been reported previously (Rhodes et al., 2016). The NEEM-SC section was chosen to extend the existing NEEM-2011-S1 record (Faïn et al., 2014; Rhodes et al., 2013) further back in time, and a composite NEEM signal including the NEEM-2011-S1 and new NEEM-SC sections is presented here. The calibration of the NEEM-2011-S1 CO data was revisited before building this composite (see Supplementary Information, SI Sect. 2.1). The NEEM-SC and NEEM-2011-S1 archives are different ice cores, but drilled at the same location.

100 The fifth ice core in this study is a new core, retrieved from central Greenland during June 2015: the PLACE ice core. The PLACE core was drilled 1 km away from the location of Eurocore site(72.58°N and 37.64°W, drilled 1989; Schwander and Rufli, 1994), which in prior work was suggested to contain an intact atmospheric CO signal for air ages younger than 1650 CE (Haan et al., 1996; Haan and Raynaud, 1998). Care was taken when selecting the PLACE site to avoid drilling in areas that were disturbed during the Eurocore and later GRIP ice core drilling operations. The core sections were drilled with the Blue  
105 Ice Drill (BID) (Kuhl et al., 2014) and immediately placed in the shade in a clean snow area after drilling, before packing in ice core boxes. These boxes were then stored in a snow cave (at -20°C or colder). At the end of the season, the cores were removed from the snow cave at the field site and returned to Summit Station for transport to the National Ice Core Laboratory (NICL), Denver, CO USA. Core temperature was verified to not have exceeded -15°C during transport by including temperature loggers in some of the ice core boxes.

110 The five sites investigated experience accumulation rates ranging from 8 to 41 cm weq yr<sup>-1</sup>, mean annual surface temperature between -24 and -32°C, and different chemical composition (Table 1, Sect. 3.3). Description of the ice and gas chronologies for the five ice cores is reported in Table S1.

### 2.2 CO continuous flow analyses

Since the first continuous, high resolution, CO measurements were performed along the NEEM-S1 ice core in 2011 (Faïn et al.,  
115 2014), CFA-based CO analyses have greatly improved, including lowering the CO blank and characterizing how CO is preferentially dissolved during the CFA process so as to establish an absolute calibration. The excellent precision of CO analyses has been confirmed, and the designs of CFA setups themselves have been optimized to limit the instrumental smoothing and

**Table 1.** Locations, site characteristics and other relevant information for ice cores featured in this study.

Ice core and location	Drilling year	Depth interval (m)	Gas age interval (yrs CE)	Accum. rate (cm ice yr <sup>-1</sup> )	Mean annual Temp. (°C)
D4 Central Greenland 71.40°N, 43.08°W 2713 m elevation	2001	63-145	1825;1961	41 <sup>a</sup>	-24 <sup>a</sup>
Tunu13 NE Greenland 78.03°N, 33.88°W 2200 m elevation	2013	61-213	836 ; 1893	8-14	-29 <sup>b</sup>
NGRIP Central Greenland 75.10°N, 42.32°W 2917 m elevation	2000	74-108 207-254 519-569 1215-1265	1780 ; 1926 980 ; 1237 -929 ; -616 -5545 ; -5899	19 <sup>c</sup>	-31.5 <sup>c</sup>
PLACE Central Greenland 72.58°N, 37.64°W 3200 m elevation	2015	80-153	1447; 1957	20.5	-32
NEEM-2011-S1 NW Greenland 77.45°N, 51.06°W 2450 m elevation	2011	71-409	270 ; 1961	22 <sup>d</sup>	-28.9 <sup>d</sup>
NEEM-2011-SC NW Greenland 77.45°N, 51.06°W 2450 m elevation	2009	399-573	-682 ; 322	22 <sup>d</sup>	-28.9 <sup>d</sup>

<sup>a</sup>Rhodes et al. (2016); <sup>b</sup>Butler et al. (1999); <sup>c</sup>community members (2004); <sup>d</sup>NEEM community members (2012).

improve signal resolution. This section reviews the operation and recent improvements of CFA-based CO measurements. More details can also be found in Supplementary Information (SI Sect. 1).

120 **2.2.1 System operation**

The ice cores listed in Table 1 were analysed using continuous ice core melter systems coupled with online gas measurements. The general principles of this analytical approach are provided by (Stowasser et al., 2012). Briefly, ice core sticks are cut at a 34 mm x 34 mm cross section and processed on a melterhead located in a cold room. The melterhead is composed of inner and outer collection areas with the inner area dedicated to sample collection. To prevent contamination, an overflow from the inner 125 to the outer melterhead areas of >10% is created by lowering the sample pumping speed. The water and gas bubble mixture is continuously pumped via a debubbler into a temperature-controlled gas extraction unit maintained at 30°C. The gas/water volume ratio of the sample is about 10% before the debubbler, and 50% after the debubbler. A fully degassed fraction of the melted sample is thus also available at the debubbler for complementary chemical analyses in the liquid phase. The gas is recovered from the sample line after the debubbler by applying a pressure gradient across a gas-permeable membrane, either 130 Transfer-Line (Idex) or Micromodule 0.5×1 (Membrana GmbH) degassers. Then, the gas is dried by a custom-made Nafion (Perma Pure) dryer. Finally, CO (and/or CH<sub>4</sub>) mixing ratios are continuously monitored along the gas sample flow by a laser spectrometer.

Our samples were all analysed at the Desert Research Institute (DRI), Reno (NV, USA) during two analytical campaigns (Tunu13, D4, NEEM-SC, and NGRIP in 2013; PLACE in 2015). The NEEM-2011-S1 core was analysed at DRI in 2011, 135 as described by Faïn et al. (2014). The PLACE core was also analysed at the Institute of Environmental Geosciences (IGE, Grenoble, France) in 2017. Specific descriptions of the DRI and IGE setups were already provided by Rhodes et al. (2015) and Fourteau et al. (2017), respectively, and a comparison of the two setups is reported in the SI Sect. 1.2.

A unique spectrometer, using optical feedback cavity enhanced absorption spectrometry (SARA, developed at Laboratoire Interdisciplinaire de Physique, University Grenoble Alpes, France; Morville et al., 2005) was used to analyse carbon monoxide 140 (and simultaneously, methane) at both DRI and IGE and during the three analytical campaigns. Detailed description of this instrument, which was used for CO measurements along the NEEM-2011-S1 core in 2011 at DRI, is reported by Faïn et al. (2014). The OF-CEAS instrument was always carefully calibrated against standard gases (Table S3) before melting ice cores (see SI Sect. 1.4).

**2.2.2 Calibration Loop**

145 The degassing membranes operated so far in gas-CFA systems extract bubbles from the sample flow but do not recover dissolved gases from the water phase efficiently. Carbon monoxide (or methane) have higher solubility than N<sub>2</sub> or O<sub>2</sub>. Consequently, mixing ratios of CO in the gas phase of the sample flow exhibit lower values than initially existing in the ice bubbles. These offsets need to be corrected for, and an attempt to replicate the conditions experienced by the ice core water-gas mixture 150 between the melterhead and the laser spectrometer was reported originally by Stowasser et al. (2012). As melting ice contains 10% air by volume, a 10:90 mixture of synthetic air and degassed deionized (DI) water can be introduced into the system via a 4-port valve located directly after the melterhead. The water is sourced from a 2 L reservoir degassed by constantly bubbling ultra-pure He through it. The air–water mixture follows the same path through the system as the ice core sample before being

analysed by the laser spectrometer. However, it is not completely identical as it includes more components such as an additional peristaltic pump or extra lines. This pathway for synthetic standard analysis will be referred to as “calibration loop” here, but 155 has been designated as “full loop” in previous studies (e.g., Rhodes et al., 2013; Faïn et al., 2014). Note that this “calibration loop” should not be confused with the “Internal Loop” (Rhodes et al., 2013) a system to isolate the gas extraction system from the remainder of the melting system, while keeping a continuous gas extraction (see Fig. 1 from Rhodes et al., 2013).

### 2.2.3 CO blank of CFA-based analyses

The CO procedural blank was evaluated for each analytical campaign. Two different evaluation approaches were applied, which 160 are described in SI Sect. 1.5. A relatively high CO blank of  $35 \pm 7$  ppbv was observed at DRI in 2013, when using a Membrana Micromodule degasser for analysing the Tunu13, D4, and a fraction of the NEEM-SC cores. Lower blank values, ranging  $4.1 \pm 1.2$  to  $12.6 \pm 4.4$  ppbv were observed when using an Idex Transfer-Line degasser (cores PLACE, NGRIP, a fraction of NEEM-SC).

### 2.2.4 Internal precision and stability

165 Internal precision and stability of gas-CFA measurements can be evaluated by Allan variance tests (Allan, 1966) applied to the calibration loop dataset. Laser spectrometers produce many measurements per minute, with a data acquisition rate of 6 Hz for our OF-CEAS instrument. Observed optimal integration time (i.e., time of lowest Allan deviation) was determined for each analytical campaign, and is now larger than 500 s for DRI, and 1000 s for IGE CFA setups (SI), which is much longer than the 5 s reported by Faïn et al. (2014) and testament to increased stability. However, to maximize the depth resolution, CO data 170 can be averaged over shorter integration time (IT). There is a trade-off between internal precision and IT. Abrupt, non-climatic signals in gas records, such as in situ CO production (Faïn et al., 2014), cannot be fully resolved without reducing the IT. Internal precision, defined as twice the Allan deviation at chosen integration times, ranged from 1.2 to 1.6 ppbv depending on the analytical campaign (SI Sect. 1.6 and Table S2).

### 2.2.5 External precision

175 External precision of the continuous CO measurements (i.e., including all sources of errors or bias) was investigated by melting replicate ice sticks on different days during each analytical campaign. Specifically, we defined the external precision as the pooled standard deviation calculated on the differences of CO concentrations from main and replicate analysed ice sticks, averaging continuous CO data over few cm long intervals (SI Sect. 1.7). This approach estimates an external precision of 5.7 ppbv for the DRI gas-CFA setup in 2011 (NEEM-2011-S1 campaign, 18 m long replicated section, using 9 cm intervals). We 180 reproduced this analysis for the 2015 DRI campaign, and established an external precision of 6.6 ppbv (19.6 m long replicated section, 10-cm long intervals, Fig. S3). With a similar methodology, but a shorter replicate section analysed, we evaluated the external precision of PLACE continuous CO measurements conducted at IGE in 2017 to 3.3 ppbv (2.3 m long replicated section, 1-cm intervals, Fig. S3). No replicate ice core sections were available and measured during the 2013 DRI campaign.

For the 2013 campaign, we consider the external precision to be 5.7 ppbv for the ice cores analyzed in 2013 with a Micromodule

185 and 6.6 ppbv using the Idex Transfer-Line degasser, by extrapolating results from similar CFA setups.

## 2.2.6 Absolute calibration and accuracy

CFA-based gas records must be corrected for under-recovery of gases dissolved in the water stream to obtain absolute values on the WMO-CO X2014A calibration scale. The magnitude of gas dissolution is monitored using the calibration loop (Sect.

2.2.2). In this study, a 73 m long ice core (PLACE) was analysed on two different CFA setups (at IGE and DRI), for CO but

190 also for CH<sub>4</sub> mixing ratios. This gas CFA laboratory intercomparison revealed that the fraction of CH<sub>4</sub> not recovered at IGE was larger than at DRI (14% and 10%, respectively). Note that the gas extraction unit and OF-CEAS detection instrument

were identical, and only the melter and lines upstream of the Idex Transfer-line degasser had different geometries (Sect. 2.2.3).

This intercomparison demonstrates that the solubility-based calibrations are required to transfer gas CFA datasets into absolute concentration scales, and are dependent on the geometry and operation of a CFA setup. This observation was further confirmed

195 at IGE by measuring the CH<sub>4</sub> mixing ratio at IGE when melting replicate ice sticks with varying CFA system geometries (e.g., length of lines) and operation of the CFA setup (e.g., melting speed, pumping rate). For each analytical campaign, the geometry and operational protocol of the CFA setup were thus kept unchanged during the measurements. However, these observations

rule out universal modelling of the preferential CO dissolution by using Henry's Law coefficient, and demonstrate that CFA setups are dynamic systems, not at solubility equilibrium.

200 In this study, we used the calibration loop operated with standard gases (Table S3) for absolute calibration of the CO dataset.

Such an approach has been successfully applied for CH<sub>4</sub> (e.g., Rhodes et al., 2015). However, calibration loops need to be carefully set up and operated, to closely reproduce solubility losses occurring in the CFA sample pathway. In the case of CH<sub>4</sub>, discrete datasets often allow for an external validation of this CFA internal calibration, but such validation was not available for Greenland CO. We hypothesise that CO and CH<sub>4</sub> dissolution follow the same physical laws: consequently, if a calibration

205 loop is able to reproduce methane preferential dissolution, it should also reproduce CO losses related to dissolution. Thus, calibration loop experiments reproducing the expected methane preferential dissolution were conducted for each analytical campaign. We extracted from such calibration loop experiments solubility calibration (SC) factors for CO. Overall, CO losses driven by preferential dissolution ranged from 4 to 9%. Replicate measurements show that the fraction of CO not recovered at

210 the outlet of the CFA system was very stable during analytical campaigns, both at DRI and IGE. Based on repeated calibration loop measurements throughout the campaigns we conservatively estimate the uncertainty of the SC factor to be  $\pm 1\%$  (2 sigma).

To further evaluate the robustness of the CFA procedure and this absolute calibration approach, five discrete samples of the PLACE core were analysed with a discrete CO setup (SI Sect. 1.8.4, Fig. S7). The agreement between discrete and CFA-based CO dataset (both IGE and DRI) was excellent, with differences in CO mixing ratio ranging from 0.4 to 2.8%. The depth intervals of the discrete samples encompassed the entire span of the PLACE core, from 84 to 146 m depth. More details about

215 absolute CO calibration are reported in Supplementary Information (SI Sect. 1.8).

### 2.2.7 Signal Smoothing

The mixing of gases and meltwater during the sample transfer from the melt head to the laser spectrometer induces a CFA experimental smoothing of the signal. The extent of the CFA-based damping was determined for each analytical campaign by performing step tests, i.e., switches between two synthetic mixtures of degassed DI water and synthetic air standards of 220 different CH<sub>4</sub> concentrations (Stowasser et al., 2012, see SI Sect. 1.9). A cut-off wavelength can be defined as the wavelength of a sine signal experiencing a 50% attenuation in amplitude. Cut-off wavelengths ranged from 15.0 cm during the DRI 2013 campaign (when using Membrana Micromodule degasser) to 1.6 cm during IGE 2017 campaign (Table S2). Lower cut-off wavelengths were obtained by limiting system dead volume, most notably by reducing the debubbler internal volume and by 225 using an optimized Idex Transfer-Line degasser instead of Membrana Micromodule degasser (1 and 5 ml internal volume, respectively)

### 2.2.8 Data processing

CO and ice chemistry data collected at DRI were mapped onto depth scales using high-resolution (0.1–0.5 Hz acquisition rate) liquid conductivity data and time–depth relationships recorded by system operators. A constant melt rate for each metre length of core is assumed. DRI depth scale uncertainties are estimated to be  $\pm 1$  cm ( $1\sigma$ ). 230 Occasional entry of ambient air into the analytical system as breaks in the core were encountered caused contamination. The OF-CEAS spectrometer simultaneously measures CO and CH<sub>4</sub> mixing ratios, and such contamination was characterised by a sharp increase in CH<sub>4</sub> concentration (ca. 1900 ppbv) followed by an exponential decrease. Data were manually screened for these ambient air contamination events.

Uncertainty on CO mixing ratios is established as  $1\sigma$  and calculated for each calibrated CO dataset. It combined uncertainties 235 evaluated specifically for each analytical setup on CO blanks, solubility calibration factors, and internal precision of CO CFA measurements (see SI and previous sections).

Finally, we used high resolution water isotope datasets to match DRI and IGE PLACE depth scales. Water isotopic ratios were measured simultaneously with gas concentrations in both laboratories using laser spectroscopy (Gkinis et al., 2011; Maselli et al., 2013).

## 240 2.3 Investigation of possible rapid CO production from trace organics in the ice during melting

To investigate if elevated or highly variable carbon monoxide levels observed in Greenland ice core could originate from the melting process (i.e., “in extractu” CO production from trace organics in the ice), CO concentrations were determined in tests with 20 discrete ice and firn samples (sized between 272 - 1089 g, average size of 690 g). The experimental protocol is reported in detail in the SI. Briefly, ice samples collected below close-off depth (fully closed porosity) are completely grated to 245 fine powder to remove trapped air. Firn samples do not contain a significant amount of trapped air; however some firn samples were still grated to powder to verify that the grating process did not increase the [CO] blank. The samples are then introduced in a glass vessel, evacuated and flushed with ultrapure air overnight. Gas standard of known concentration is then introduced

within the vessel. Consequently, for both firn and ice samples CO mixing ratio within the vessel prior to melting is equal to that of the gas standard introduced in the vessel. If CO in extractu production occurs during melting, it should cause an increase in  
250 CO mixing ratio above this known value.

The CO concentration in the gas available in the vessel headspace is determined by gas chromatography combined with a mercuric oxide reduction detector (RGD; Peak Performer 1 from Peak Laboratories), at three different stages: before melting, during melting, and after melting. CO concentrations are calibrated using four synthetic air gas standards with nominal CO concentrations ranging from 50 to 500 ppbv, and data are reported on the WMO-CO X2014A scale.


255 Three large diameter (27 cm) cores collected with the Blue Ice Drill (BID) were used for these tests: the PLACE ice core (Central Greenland, Table 1), the C14 firn core (Central Greenland, drilled in June 2015, 72°39.62' N, 38°34.85' W), and a Taylor Glacier ice core (Antarctica, drilled in December 2015, 77°45.69' S, 161°43.18' E). BID cores allowed for up to four true depth replicates to be measured from the same depth interval. CO concentrations were determined (before, during, and after melting) for the following cases: (i) Greenland C14 firn ( 45 m depth), (ii) Greenland PLACE ice from depths ranging  
260 110-125 m and showing low and stable [CO] on the continuous CFA record, (iii) Greenland PLACE ice from depths ( 111 m) showing elevated and highly variable [CO] on the continuous CFA, (iv) Oldest Dryas ( 15 kyrs BP) and Holocene ( 9.5 kyrs) Antarctica ice, and (v) gas-free ice made from degassed Milli-Q 18.2MΩ.

## 2.4 Chemistry data

During the two DRI analytical campaigns, melted ice core samples were analysed continuously by inductively coupled plasma  
265 mass spectrometry (ICP-MS) and CFA for chemical species. These analytical methods have been reported previously (Mc-  
Connell and Edwards, 2008; McConnell et al., 2007, 2002). High-resolution measurements of total organic carbon (TOC)  
were obtained at DRI by coupling a Sievers 900 TOC analyzer to the DRI ice core melter (Legrand et al., 2016).

## 3 Results and Discussion

The new CO records available from the PLACE, NGRIP, NEEM, D4, and Tunu13 ice-cores are reported in Fig. 1 and plotted  
270 for the period spanning 1650 to 1960 CE (gas age, see Table S1). Only the NEEM, NGRIP and Tunu13 datasets significantly  
extend beyond the last three centuries and the full records are reported in Fig. S17. Over the 1650-1960 CE period, all Greenland  
CO ice records exhibit highly variable CO values. A minimum mixing ratio of 92 ppbv is observed at NEEM and PLACE,  
and a maximum mixing ratio of 1191 ppbv at Tunu13. Considering all records, some 46 events revealed mixing ratios higher  
than 250 ppbv. In Sect. 3.1 we discuss the baseline CO levels of the different ice-core records, and the characteristics of abrupt  
275 CO spikes detected in all records. Tests and discussions presented in Sect 3.2. demonstrate that in situ production drives these  
abrupt CO spikes. The key question of whether an atmospheric signal can be extracted from the low frequency variability of  
the CO ice-records' baselines is discussed in Sect. 3.3. Comparison of our new CO records with existing data is presented in  
Sect. 3.4. Finally, a multi-site reconstruction of past atmospheric CO levels at northern latitudes is presented in Sect. 3.5.



**Figure 1.** Upper panels: Greenland continuous CO records available from the PLACE (DRI), NGRIP, NEEM, D4, and Tunu13 ice-cores and plotted for the period spanning 1650-1960 CE. Lower panel : CO baseline levels from each ice-record defined as the 5<sup>th</sup> percentile of data every 4 years over a moving window of 15 yrs, with shaded envelopes reported on each baseline representing the uncertainty.

### 3.1 Continuous CO records along Greenlandic ice cores

#### 280 3.1.1 CO baseline levels


We identified the CO baseline levels from each ice-record by calculating the 5<sup>th</sup> percentile of data every 4 years over a moving window of 15 yrs. A window of 15 yrs corresponds to a time interval shorter than the full width at half maximum of the CO age distribution at these sites (Table 1). We chose to use 5<sup>th</sup> percentile CO baselines because they can be calculated identically for all datasets and are not based on the subjective operator decisions involved in identifying and removing peaks  
285 to derive a baseline level. The temporal trends in records calculated using 10th percentiles are similar. 5<sup>th</sup> percentiles were not computed when more than 50% of the data were missing within the windows. Figure 1 (lower panel) reports 5<sup>th</sup> percentile CO baselines extracted over the last 300 years from the D4, Tunu13, NGRIP, NEEM, and PLACE ice-records (all DRI dataset). The shaded envelope reported on each baseline represents the uncertainty (1 $\sigma$ ) calculated by propagating CO external precision and uncertainties on CO blank and solubility calibration factor (see Sect. 2). The CO baseline records from the five ice cores  
290 are always within 20 ppbv of each other. From 1700 to 1875 CE, the ice-records indicate rather unchanged or slightly enhanced levels with time, whereas from 1875 to 1950 CE, all the ice-records reveal a steady increase.

#### 3.1.2 High frequency non atmospheric CO signals

Figure 1 shows that the five Greenlandic sites investigated in this study all exhibit highly variable CO mixing ratios. Referring to previous data reported by Faïn et al. (2014), such a pattern was expected when measuring the NEEM-SC samples and extending the NEEM-2011-S1 record. However, new records from central Greenland (PLACE, NGRIP), north-eastern Greenland  
295 (Tunu13) and southern Greenland (D4) appear to be also affected by variability in CO mixing ratios that is too large and rapid to reflect past atmospheric mixing ratio changes (Figs. 1 and S17).

The true extent of high CO variability may be masked, however, by CFA analytical smoothing. Figure 2 (upper panel) reports a 1.5 m long section of the PLACE ice-core, analysed with both DRI and IGE CFA setups. The higher CFA setup resolution  
300 at IGE (Sect. 2.2.7) reveals more features, but may still miss even higher frequency variability and dampen extrema values. Similarly, a CO record from a core drilled at a relatively low snow accumulation site will appear more smoothed out by CFA analysis than one from a relatively high snow accumulation rate site.

High frequency variability within CO records can be quantitatively characterized using MAD (Median Absolute Deviation). MAD calculated for all CO records collected at DRI every 4 yrs over a moving window of 15 yrs (similarly to baseline signals,  
305 see Sect. 3.1.1) are reported on Fig. S18. We observe for the 1700-1950 CE time period that NEEM, NGRIP, Tunu13 and PLACE records exhibit stable and similar MAD values, with mean MAD values over the entire 1700-1950 CE period ranging 9.9-14.2 ppbv, depending on the records. The D4 MAD values are larger, varying from 10 to 30 ppbv. The NEEM, NGRIP and Tunu13 datasets significantly extend beyond the last three centuries and reveal different MAD patterns for age older than 1700 CE. The Tunu13 record, spanning the last 1300 yrs, reveals that CO variability remains stable as depth increases while  
310 NEEM and NGRIP show increasing MAD with depth, clearly indicating that high variability in CO is more pronounced for



**Figure 2.** High-resolution TOC,  $\text{NH}_4^+$ , CO from the PLACE core. CO mixing ratios were measured with both the DRI and IGE CFA systems.  $\text{NH}_4^+$  shows a well-marked seasonal cycle characterized by a summer maximum. PLACE snow accumulation is about  $20 \text{ cm weq yr}^{-1}$

older ice. These increases are significant, with MAD values reaching  $\sim 60 \text{ ppbv}$  for the deeper sections of the analyzed NEEM and NGRIP ice-core sections.

### 3.2 Do drilling or analytical processes cause CO production?

The observation of high frequency CO variability along the NEEM-2011-S1 core led Faïn et al. (2014) to discuss if such 315 patterns could be driven by the processes involved in sample collection and analysis. Here we consider our new data to assess in more detail if such processes can drive CO production.

#### 3.2.1 Ice core drilling

We observe that abrupt CO spikes of similar magnitude occur in ice-cores extracted by dry drillings (e.g., PLACE, D4, Tunu13) and when a drilling fluid was used (e.g., NEEM, NGRIP) (Fig. 1). This clearly rules out contamination from the drilling fluid 320 as a cause for high frequency CO variability, confirming the previous conclusion from Faïn et al. (2014).

Haan et al. (2001) observed CO production within an alpine snowpack in daylight. Drillings are usually conducted at high-altitude sites in summer where UV radiation levels are high. One could thus hypothesise that CO is photochemically produced

within ice-cores just after drilling, when ice-cores are handled at the drilling site before being packed in ice-core boxes. During the PLACE and NEEM-SC drillings, specific care was taken to never expose freshly drilled cores to direct sunlight. For 325 PLACE, a shaded area was set up for core processing. NEEM 2011-S1 and SC core drillings were conducted in a tent and a trench, respectively. In spite of that, all these ice-cores reveal CO spikes. In contrast, no specific care was taken during the historical Eurocore drilling when handling cores, with respect to sunlight exposure, and Haan and Raynaud (1998) report that this archive does not exhibit highly variable CO concentration in the upper section. Based on this, it appears that brief direct exposure to sunlight after drilling is unlikely to be the cause of the large CO spikes

### 330 **3.2.2 Impact of long-term core storage**

By comparing the storage time of different ice-cores (storage without light exposure) prior to CO analysis with the same CFA setup (NEEM-2011-S1 and D4), Faïn et al. (2014) concluded that CO production in ice during core storage was unlikely. We extend this observation by comparing central Greenland archives: NGRIP and PLACE records reveal similar abrupt CO spikes and CO baselines (Fig. 1) while PLACE and NGRIP cores were analysed 3 months and 15 years, respectively, after drilling. 335 Furthermore, we measured at IGE a 10 m long replicate section of the PLACE core nearly two years after the main analytical campaign (respectively December 2018 and February 2017) and no change in the CO values was observed (Fig. S14).

### **3.2.3 Impact of CFA process**


In 2013 at DRI, a 4 m long replicate section of the D4 ice-core was melted with all laboratories kept in darkness. Similar in situ CO production patterns were observed under light and dark melting conditions. These results, already reported by Faïn et al. 340 (2014, Fig. S2), reveal that CO variability observed along the D4 core are not related to photochemically driven CO production occurring in the sampling lines between the melter head and the CFA extraction box. To further evaluate if the CFA analytical system could somehow induce CO production within the melted ice, a melted D4 sample was collected downstream of the degassing membrane and re-circulated in the calibration loop mode (Sect. 2.2.2) for one hour as degassed blank water. We observed similar CO levels to the deionised water blank.

### 345 **3.2.4 Results of investigation of possible rapid CO production from trace organics in the ice during melting**

As described above, CO analyses of 20 discrete ice and firn core samples (including the PLACE core) were carried out at the University of Rochester, with the goals of investigating the possibility of in extractu production of CO from trace organics in the ice during melting of ice core samples (Sect. 2.3 and SI Sect. 1.11, Table S4).

All of the different ice and firn sample types, including samples of different mass and a gas-free ice sample, showed a consistent 350 excess CO growth occurring during the melting step, with excess CO increasing by 6.4 ppbv on average (Fig. 3) - a much smaller CO enhancement than the spikes observed in the CFA records. Since the excess CO produced appears stable after melting, and consistent between the sample types, the excess CO observed is likely a systematic extraction system blank.

If the CO production mechanism involves organic compounds present in the ice lattice, it would be expected that variations in



**Figure 3.** Average excess in CO mixing ratios produced during melting of discrete samples measured by gas chromatography combined with a mercuric oxide reduction detector. The CO values shown have been corrected for a small ( $\sim 3$  ppbv) CO excess observed in equivalent tests where the ice samples were not melted (see SI Sect. 1.11.3). The following samples are investigated : gas free ice, firn from Central Greenland (including low mass samples), PLACE ice samples exhibiting high CFA [CO] and low CFA [CO], Taylor Glacier (Antarctica) ice. Samples description is reported in Table S4. Error bars represent one standard deviation for all sample runs of a given type and points without error bars represent single measurements

the concentrations of organics from samples collected at different sites / from different depth levels, as well as differences in  
 355 the melted ice mass would result in significant differences in excess CO. Some of the PLACE ice samples were specifically selected for having exhibited low and stable [CO] on the continuous CFA record while others were selected for exhibiting elevated and spiky [CO] behavior suggesting the possibility for instantaneous CO production during melting (Fig. 3, Table S4). The other samples, the two Taylor Glacier and gas-free ice were also selected as they should have very different trace organic loadings to the Greenland ice, especially the gas-free ice sample which have no significant organic loading (< 2 ppbC). Further,  
 360 replicate firn samples of different mass would be expected to have a total organic loading that scales with the firn mass. Because there are no significant differences observed in the excess CO produced from these very different sample types (Fig. 3), these

results indicate that the melt-extraction process itself does not result in significant CO production from trace organics found within the ice samples. Instead, these results lend further support to the idea that the elevated and highly variable [CO] values observed in Greenland ice samples are due to excess CO produced from in situ production within the ice itself (Faïn et al., 365 2014; Haan and Raynaud, 1998).

### 3.3 Atmospheric CO history retrieved from Greenland ice cores

We now explore if past atmospheric concentrations of CO can be extracted from the low frequency variability of the CO ice-records' baselines (Fig. 1). The key question is: does the in situ production present in all five cores still impact the 5<sup>th</sup> percentile of data that we adopt as a baseline?


370 In our earlier study on the NEEM-S1-2011 core (Fain et al., 2014), we reported that 68% of the CO spikes were observed in ice layers enriched with pyrogenic aerosols (i.e., exhibiting ammonium levels above 18 ppb). Within the 60 cm length core analysed in that study, elevated ammonium ( $\text{NH}_4^+$ ) and CO were found together with dissolved organic carbon (DOC) (>100 ppbC) representing an important reservoir of carbon, which could potentially be oxidized into CO. The oxidation processes involved, however, remain unidentified. Here we focus on the two cores for which we have obtained reliable total organic 375 carbon (TOC) data (Sect. 2.4). These are interesting cores to compare because they have very different snow accumulation rates: Tunu13 has the lowest accumulation rate, which also varies over time between 8 to 13 cm weq yr<sup>-1</sup>, while PLACE is more typical of a Late Holocene Greenland core with a stable accumulation rate of ~20 cm weq yr<sup>-1</sup>.

#### 3.3.1 The importance of data resolution

Typical CO, TOC, and  $\text{NH}_4^+$  patterns observed along the PLACE core are reported along a 1.6 m section on Fig. 2. Maximum 380 and minimum levels of the three species occur at similar depths.  $\text{NH}_4^+$  is commonly used in Greenland ice to identify summer layers that are characterized by a well-marked summer maximum (Legrand and Mayewski, 1997). Investigating summertime and wintertime distributions of [CO] and [TOC] over the full PLACE record confirms that both species share coherent seasonality (Fig. S22 and S26): CO and TOC show lower levels in winter-deposited ice layers, suggesting that at the PLACE site the CO signal preserved in winter layers is more suitable for paleo-atmospheric reconstructions.

385 Typical CO, TOC, and  $\text{NH}_4^+$  variations along the Tunu13 core appear different to those at PLACE (Fig. 4). The seasonal cycle in  $\text{NH}_4^+$  is still detectable but the TOC and CO records show much less co-variation than at PLACE. PLACE and Tunu13 TOC were analyzed with the same setup (i.e impacted by the same analytical smoothing), but the lower accumulation rate of Tunu13 results in a larger smoothing of the TOC signal. The higher degree of smoothing in CO relative to PLACE results 390 in part from Tunu13 being analysed with the 2013 DRI CFA setup (Micromodule degasser), which shows a larger response time in the gas phase (Table S2). Similarly to TOC, the effect is compounded by the much lower snow accumulation rate of Tunu13 relative to PLACE - the section of core in Fig. 4 has only 8 cm weq yr<sup>-1</sup>. Step tests (Sect. 2.2.7) suggest that for CO a 60% attenuation would be expected for a sine signal of 10 cm wavelength, a depth interval representing >1 yr in this case. The impact of analytical smoothing on Tunu13 CO patterns was further investigated by convoluting a Tunu13-like synthetic CO signal with the Green function of the 2013 DRI CFA setup (Fig. S8, middle panel; Fig. S12). A 5<sup>th</sup> percentile baseline was

395 extracted from the smoothed synthetic signal produced by this convolution. The baseline obtained is found up to 8 ppbv above the original synthetic signal for depth intervals spanning few meters (Fig. S13), confirming that CFA analytical smoothing can positively bias the Tunu13 baseline CO level. More details about this conceptual experiment are available in Sect. SI 1.9.2.



**Figure 4.** High-resolution TOC,  $\text{NH}_4^+$ , CO from the Tunu13 core. Snow accumulation is about  $8 \text{ cm weq yr}^{-1}$  at this depth.


The significant analytical smoothing for the Tunu13 ice-core suggests that annual winter minima in CO (associated with winter minima in TOC) may be unresolvable, which would lead to an overestimation of the baseline even when using the 5<sup>th</sup> percentile values. In contrast, repeated analyses of the higher snow accumulation PLACE core on two different gas CFA setups (DRI and IGE, Table S2) show that despite the DRI setup (2015 configuration) exerting a greater smoothing effect than the IGE one (SI Sect. 1.9, Fig. S10 and S11), the CO minima values are well-resolved in both cases (Fig. 2). Furthermore, the PLACE CO baseline extracted from the IGE continuous record is similar to the one based on the DRI dataset (SI section 2.4), suggesting that analytical smoothing is not a limitation on isolating CO levels in the winter ice layers at this site. Convoluting a PLACE-like synthetic CO signal with either DRI 2015 and IGE 2017 CFA setup Green function (e.g., SI Sect. 1.9.2) confirms this result. This suggests that the 5<sup>th</sup> percentile baseline calculated at PLACE and other high snow accumulation sites is suitable for investigating past evolutions of the atmospheric CO burden.

### 3.3.2 TOC patterns and in situ CO production

The high resolution TOC measurements collected along the PLACE core exhibit a stable TOC background value of 13.5 ppbC (baseline defined as the 5<sup>th</sup> percentile), with 17 spikes above 40 ppbC (70 m long record, Fig. S24). TOC levels observed at Tunu13 are lower, with a baseline value of 7.9 ppbC (1 $\sigma$ ) and only 10 spikes above 40 ppbC for a longer record (143 m long record, Fig. S25). Tunu13 mean TOC (10 meters window averaging) exhibits a low frequency variability with values ranging from 4 to 12 ppbC (Fig. 5). The low frequency variability in Tunu13 mean TOC is correlated to changes in the snow accumulation rate (Fig. 5) which varies from 8 to 13 cm weq yr<sup>-1</sup> (annual average). Along most of the core, this linear relationship between TOC and snow accumulation rate remains unchanged, although it becomes steeper below 160 m depth (Fig. S27). Such correlation is not observed at PLACE where the snow accumulation is constant, with annual mean values ranging 20-21.5 cm weq yr<sup>-1</sup>.

In situ CO production within Greenland ice archives is very often co-located in ice layers where TOC levels are high (e.g., Fig. 2, S22, and S26) as observed previously on the NEEM-2011-S1 core (Faïn et al., 2014), suggesting that in situ CO production may be related to the organic carbon (OC) availability. The importance of organic carbon in the processes driving CO in situ production is further supported by the long-term averaged mean CO record (running windows of 10 m) which is significantly correlated to mean TOC at Tunu13 ( $r^2 = 0.56$ ,  $p < 0.01$ , Fig. 5) for preindustrial times (gas age prior to 1850 CE, i.e. below 83 m depth), with increases in mean CO when TOC and snow accumulation are higher. However, comparison of the Tunu13 and PLACE datasets suggest that higher TOC levels does not always imply higher CO concentrations: over the 1640-1950 CE gas age time period, the mean CO mixing ratio is almost identical for Tunu13 and PLACE (137 and 136 ppbv, respectively) while mean TOC is higher for PLACE compared to Tunu13 (20 and 7 ppbC, respectively). This may suggest that the required amount of OC to lead to significant in situ CO production is already reached at the Tunu13 site.

Characterizing the fate of TOC in Greenland ice is a first requirement for a better understanding of in situ CO production processes. Figure 5 suggests that the fate of TOC deposited at Tunu13 is related to the rate of snow accumulation at the surface. Comparing Tunu13 and PLACE TOC records suggests that the lower the accumulation rate, the lower the TOC baseline concentrations in the deep ice. A snowpit study of recent snow layers deposited at Summit (Hagler et al., 2007) showed that a large fraction of water soluble and insoluble organic carbon may be lost at the surface during post-deposition processes, such as photochemical reactions. We hypothesize that such processes may be enhanced at lower accumulation rates when the snow layers get exposed to UV radiation for longer. However, the relationships between TOC and ammonium suggest that other processes can impact TOC after deposition at Tunu13. Forest fire debris reaching Central Greenland snow during spring or summer mainly consists of ammonium formate, and organic carbon is mostly made of formate at Summit (Legrand et al., 2016). Legrand and De Angelis (1996) reported a formate-to-ammonium molar ratio close to unity in elevated ammonium events recorded at Summit both during the last 200 years and the Holocene. The relationship between TOC and ammonium at PLACE (Fig. S28) exhibits a similar slope close to unity. A completely different pattern is observed at Tunu13 (Fig. S29), with unexpectedly low TOC observed along ammonium peaks. We suggest that ammonium formate has been remobilized after deposition in the snow at sites with snow accumulation



**Figure 5.** Running averages (10 m windows) of TOC and CO concentrations for the Tunu13 core, plotted along with annually averaged surface snow accumulation rate.


rates lower than  $12 \text{ cm weq yr}^{-1}$ . Ammonium remains when deposited due to acidic snow layers. In contrast to ammonium, formate evolves to a different form (presumably gaseous formic acid) that can be released into the free atmosphere above the snowpack or be remobilized to neighbouring winter layers. Such a process could contribute to explain lower TOC levels observed under lower snow accumulation rates, and also the smoothed shape of the TOC peaks at Tunu13 (Fig. 4). This process can also promote in situ CO production in winter layers where some organic substrates can be remobilized.

### 3.3.3 Summary

Investigating the CO and TOC PLACE records did not reveal any evidence that the CO baseline could be influenced by in situ production at that site. In contrast, Tunu13 has a lower snow accumulation rate than other sites (Table 1) and thus the analytical CFA smoothing of its CO record is more pronounced. This effect likely introduces a positive bias into the Tunu CO baseline record for ice sections impacted by in situ production, with some of the CO baseline signal likely incorporating in situ produced CO from summer ice layers. This limits the interpretation of Tunu13 baseline as an atmospheric signal in this study. While the mean CO mixing ratio is significantly correlated to the mean TOC concentrations in ice at Tunu13, this is not the case for baseline CO level which does not exhibit significant correlations with mean or baseline TOC concentrations. But we can not rule out that a redistribution of organic carbon along depth driven by OC post-deposition process (shifting the ammonium-formate equilibrium with the gas phase) impacts specifically the Tunu13 CO record by providing some additional organic substrates in winter layers.

### 3.4 Comparison of PLACE and Eurocore datasets

Haan et al. (1996, 1998) published a pioneering CO dataset based on discrete sampling of the Eurocore ice core. This dataset 460 revealed a smoothed CO signal, interpreted as not being affected by in situ production over the last 300 yrs (26 samples, depth range: 78-154 m). However, Haan et al.'s CO data for the period preceding 1700 CE displayed a relatively elevated and scattered signal with levels fluctuating from 90 to 180 ppbv, likely impacted by in situ production. Through our study, we were 465 unable to identify a Greenland site where the ice archive provides a low and stable CO record similar to Haan et al.'s over the last few centuries. Specifically, the CO record retrieved from the PLACE ice-core, which was drilled less than 1 km away from the Eurocore borehole, shows high frequency variability along the entire CFA record (80-153 m depth), and this observation is supported by discrete measurements conducted on the PLACE core with an analytical process in principle similar to the one 470 deployed by Haan et al. (Sect. 2.2.6) (Fig. 6).



**Figure 6.** CO mixing ratio collected (i) along the Place ice core: continuous IGE CFA record, with 5<sup>th</sup> percentile baseline (black line), and discrete dataset (green dot); (ii) along the Eurocore archive : historical Eurocore discrete CO data (blue dot, Haan et al 1998), and CFA-based dataset measured in 2019 at IGE (red).

Over the 1700-1900 CE period, Eurocore discrete CO data are about 10 ppbv lower than the baseline of the PLACE CFA-based record (Figs. 6 and S23). Furthermore, the trend and rate of change of the two records are similar from 1700 to 1905

470 CE period but diverge over the 1925-1950 CE interval, with an increasing offset between datasets, and a sharp increase in CO concentrations observed only in the Eurocore record for the 1935-1950 CE time period.

To further investigate this apparent offset between our new Greenland CO records and the Eurocore CO record of Haan et al., we analysed six 50 cm-long sections of the 1989 Eurocore ice core with the gas CFA IGE setup in January 2019, at depths spanning 84 to 153 m. A fraction of the Eurocore core has been stored as an archive for 30 yrs at -20°C. We were able to 475 sample the Eurocore archive at exactly the same depths that Haan et al. did in the 1990s (the gaps left by the discrete sampling conducted by Haan and colleagues could be seen). Figure 6 shows Eurocore data (both from Haan et al. and our new CFA dataset), along with the PLACE IGE CFA record: to do so, an offset of +6 m was applied to the Eurocore depth scale, so as to take into account 26 yrs of snow accumulation that occurred between the Eurocore and PLACE drillings. Our CFA-based CO analyses reveal high variability along the Eurocore archive. We systematically observed higher CO mixing ratios 480 than previously reported by (Haan and Raynaud, 1998) including at the exact depths sampled by them. We established that such discrepancies were not related to differences in CO calibration scales by reanalyzing in 2015 the standard gases used by Haan et al. for calibrating their Eurocore dataset in the 90s. Overall, our CFA-based Eurocore data are in contradiction with the historical Eurocore dataset but are in good agreement with the PLACE record, in terms of both absolute levels and high variability. The discrete PLACE CO data, measured with a methodology similar to the one applied by Haan et al., also exhibit 485 higher values than Haan et al. dataset (Fig. 6), while they show an excellent agreement with the CFA CO dataset (Fig. S7).

To summarize, we have not been able to reconcile the historical Eurocore dataset of Haan et al. with a new Greenland ice-core CO dataset (PLACE). This is despite analysis of remaining Eurocore ice from depths co-located with the published Haan data and the fact that PLACE ice core was collected just 1 km away from Eurocore. The reasons for the relatively low and stable CO measurements reported in the 1990s for the Eurocore ice core remain unknown and our results bring the validity of the 490 Haan et al. (1996, 1995) results into question.


### 3.5 Constraining past atmospheric CO in the northern hemisphere

#### 3.5.1 CO baseline records over the last 300 yrs

Extracting atmospheric information from a single CO record retrieved from a Greenland ice archive affected by in situ production features is challenging (e.g., Faïn et al., 2014). However, we show here that the PLACE, NGRIP, NEEM, and D4 records, 495 reconstructed from ice archives originating from different Greenland locations and drilled over a time period spanning more than a decade, exhibit common patterns. We conservatively exclude Tunu13 from paleoatmospheric interpretations, knowing that the Tunu13 baseline CO record could be positively bias by analytical smoothing or redistribution of organic carbon along depth (see Sect. 3.3). Fig. 1 reports 5<sup>th</sup> percentile CO baselines extracted over the last 300 years from the D4, NGRIP, NEEM, and PLACE records (DRI dataset). The four CO baseline records all lie within an envelope of maximum 20 ppbv range. From 500 1700 to 1875 CE, the records reveal stable or slightly increasing values remaining in the 100-115 ppbv range. From 1875 to 1950 CE, the records indicate a monotonic increase from 100-120 ppbv to 135-150 ppbv (i.e., a mean rate of increase of ~0.3 ppbv yr<sup>-1</sup>). Interestingly, in 1835 CE, PLACE, NGRIP, NEEM, and D4 baselines exhibit a common maximum in CO

concentration but at a level just slightly higher than concentrations observed during the 1700-1820 CE period.

A multisite composite CO baseline spanning 1700-1957 CE was generated by considering D4, NEEM, NGRIP, and PLACE 505 records and excluding Tunu13. For consistency, we only consider CO records collected on the same CFA gas setup (DRI). The multisite composite, reported in Fig. 7, was obtained by averaging CO baselines values interpolated on a common gas age scale. An uncertainty envelope ( $1\sigma$ ) obtained by considering the independent uncertainty of individual records is plotted along with the composite. Overall, Fig. 7 reports a  $\sim 30\%$  increase in CO concentration at high latitudes of the northern hemisphere atmosphere from pre-industrial to 1950 CE.



**Figure 7.** Past atmospheric CO mixing ratios for the Northern Hemisphere high latitudes and spanning 1700-2018 CE. Black line : multisite average obtained by combining baselines ( $5^{th}$  percentiles) of continuous CO records collected at four Greenland sites (PLACE, NEEM, NGRIP and D4), representing an upper bound estimate of past atmospheric CO levels. Light blue : firn air records obtained by combining samples from NEEM, NGRIP, and Summit; blue : average firn air record (Petrenko et al., 2013) ; Red line : NOAA atmospheric monitoring at Barrow (Alaska, USA).

510 The fact that the CO baseline records from the four different Greenland ice cores are so consistent gives us confidence that our multisite reconstruction provides a reliable reconstruction of past atmospheric changes. An additional argument for that conclusion comes from the good overlap between our multisite reconstruction and CO data from firn air Greenland measurements

(section 3.5.2). However, we cannot exclude the possibility that past atmospheric CO levels could have been lower than shown in Fig. 7 and recommend that our multisite average should be taken as an upper bound of past atmospheric CO levels. If in situ 515 CO production drives the high frequency variability seen throughout the CFA records of these four ice cores, as we hypothesize (see Sect. 3.2), it is impossible to exclude the possibility that minimum CO levels may also be slightly affected (Sect 3.3).

### 3.5.2 Comparison with firn air reconstruction

An independent means to assess the validity of our atmospheric CO reconstruction is to compare it with a reconstruction of northern hemisphere high-latitude atmospheric CO mixing ratios from Greenland firn air, available for the period 1950-2008 520 CE (Petrenko et al., 2013). This reconstruction includes air samples collected in the open porosity of the firn at three sites in Greenland (NGRIP in 2001, Summit in 2006, and NEEM in 2008), and shows that CO records from these three sites agree well with each other as well as with recent atmospheric measurements from Barrow, Alaska. However, a small contribution from in situ CO production (up to 5 ppbv) within the firn open porosity itself could not be ruled out (Petrenko et al., 2013). This implies that the in situ CO production observed in ice core records which exhibit [CO] enhancement much larger than 5 ppbv 525 (Fig. 1) is a process happening within the ice matrix, after bubble closure. The reconstructed firn air history suggests that Arctic CO mixing ratio in 1950 CE was 138–148 ppbv, rose by 10–15 ppbv from 1950 to the 1970s, peaked in the 1970s or early 1980s, and finally declined by about 30 ppbv to present-day levels. The average firn air-based CO history spanning the last six 530 decades (Petrenko et al., 2013) is shown on Fig. 7 along with our ice core CO composite reconstruction. The uncertainty in this scenario is captured by the variability of a set of 61 possible scenarios (Petrenko et al., 2013), and is shown as an envelope on Fig. 7.

PLACE and D4 high resolution CO baseline records overlap with the firn record. The firn air data suggest that CO concentration spanned 138-148 ppbv in 1950 CE for the Arctic atmosphere Fig. 7). The 1950 CE CO concentration range as defined by PLACE and D4 baselines is 135-150 ppbv, indicating good agreement between firn air and ice core reconstructions.

### 3.5.3 Spatial representativity of the record

535 The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP; Lamarque et al., 2013) investigated the long-term changes in atmospheric composition between 1850 and 2100, with the goal of documenting atmospheric composition changes. To investigate the spatial representativity of the ice core record, we compared the CO mixing ratios simulated in the frame of the ACCMIP exercice for the years 1980 and 2000 and averaged over three different areas: the 45-90°N latitudinal band, an area encompassing the Greenland ice sheet (20-60°W; 60-84°N), and the single grid point including the PLACE 540 drilling location. Average [CO] over Greenland and [CO] at the PLACE drilling location are identical. However, atmospheric [CO] simulated for the 45-90°N latitudinal band is about 12% higher than [CO] simulated over Greenland. These results support our approach to combine ice core reconstructions obtained from different Greenlandic locations, but indicate that the absolute CO concentrations derived from our datasets are representative of the Greenlandic, Arctic, atmosphere, and may underestimate the average CO concentrations for the entire 45-90°N latitudinal band. On the other hand, the temporal changes depicted by 545 our composite record probably have a larger spatial significance than the Greenlandic context alone.

#### 4 Summary and conclusions

New continuous profiles of CO mixing ratio have been measured along five Greenlandic ice cores, the PLACE, NGRIP, D4, Tunu13 and NEEM-SC archives. We also revisited the NEEM-2011-S1 dataset (Faïn et al., 2014). By coupling ice core melter systems with online measurements (SARA spectrometer), 700 m of ice were analyzed at high resolution for CO mixing ratio.

550 This methodology has greatly improved over the last decade, with excellent external precision, low and well-constrained blank values, and a good accuracy. All investigated records revealed high and variable concentrations that can not be interpreted as changes in atmospheric CO, but are most likely related to in situ production occurring within the ice archives themselves after bubble closure. Additional measurements conducted on individual discrete samples allowed us to rule out the possibility of rapid CO production from organics in the ice during melting or other large CFA system artifacts. We have not been able to  
555 reconcile the prior Eurocore dataset of Haan et al. (1996, 1998) with any of the new Greenland ice core CO measurements made by both discrete and continuous analytical methods. Our results therefore question the accuracy of the relatively low and stable CO values reported by Haan et al. (1996, 1998) for the Eurocore ice core.

We have presented a multisite average ice core reconstruction of past atmospheric CO for the northern hemisphere high latitudes, covering the period from 1700 to 1950 CE, providing an upper bound estimate of past atmospheric CO abundance. This  
560 signal of paleo-atmosphere was extracted from the low frequency variability of the CO records' baselines of the PLACE, D4, NEEM, and NGRIP archives which all exhibit consistent trends despite different surface accumulation snow rates and chemical compositions. From 1700 to 1875 CE, the multisite average record is stable with a slight increase with time from 103 to 114 ppbv. Our upper-bound value for CO mixing ratio at preindustrial times in the Arctic is 110 ppbv. From 1875 to 1950 CE, the CO increases monotonically from 114 to 142 ppbv, with a rate of increase of about  $0.3 \text{ ppbv yr}^{-1}$ . Finally, an excellent agreement  
565 between our ice core based record and previously published firn air reconstruction is observed in the 1950s (Petrenko et al., 2013). The ice-core and firn [CO] histories, spanning 1700-2010 CE, exhibit similar patterns to an up-to-date combined inventory of anthropogenic and open burning CO emissions (van Marle et al., 2017; Hoesley et al., 1998). Our CFA-based multisite Greenland ice core CO reconstruction provides an indication of the timing and magnitude of past variations in the high northern latitude CO burden that can provide a benchmark for future atmospheric chemistry model studies. Specifically,  
570 a natural extension of this work should be the comparison of our ice-core-based atmospheric CO reconstruction with model outputs from the AerChemMIP exercise (Collins et al., 2017).

*Acknowledgements.* This work was supported by the following programs: the French ANR projects RPD-COCLICO (#10-RPDOC-002-01, X.F.) and NEEM (#07-VULN-09-001), the EU grants FP7-IP #ENV-2010/265148 (project Pegasos, X.F.) and FP7 ERC #291062 (project Ice&Lasers, J.C.), the US NSF awards #1406236 (V.V.P.), #0221515 (J.R.M.), #0909541 (J.R.M.), #1204176 (J.R.M.), #1406219 (J.R.M.), and #0968391 (E.B., Partnerships in International Research and Education, project PIRE), the Packard Fellowship for Science and Engineering (V.V.P.), and finally the French national program LEFE/INSU (project GreenCO, X.F.). Grateful thanks go to Olivia Maselli, Larry Layman, Daniel Pasteris, Michael Sigl, and other members of the DRI team who assisted with the measurement campaigns. We thank Frederic Prié, Elise Fourre, and Amaelle Landais for their help in the continuous measurement of water isotopes along the PLACE core

580 at IGE, in 2017. We are grateful to Ray Langenfelds for the measurements of gas standards at CSIRO, in 2015. We thank Sophie Szopa  
581 and Kostas Tsigaridis for our useful discussions. Polar Field Services and the 109<sup>th</sup> New York Air National Guard, and the French Polar  
582 Institute (IPEV) provided logistical support for ice core drilling. We are grateful to drillers and field teams. The NEEM project is directed  
583 by the Center for Ice and Climate at the Niels Bohr Institute, Copenhagen, and the US NSF OPP. It is supported by funding agencies and  
584 institutions in Belgium (FNRS-CFB and FWO), Canada (NRCan/GSC), China (CAS), Denmark (FIST), France (IPEV, CNRS/INSU, CEA  
585 and ANR), Germany (AWI), Iceland (RannIs), Japan (NIPR), Korea (KOPRI), The Netherlands (NWO/ALW), Sweden (VR), Switzerland  
586 (SNF), United Kingdom (NERC), and the USA (US NSF, OPP). NGRIP is a multinational research program funded by participating institutions  
587 in Denmark, France, Germany, Japan, Sweden, Switzerland Belgium, Iceland, and the United States. This work also benefited from data  
588 available in the public NOAA ESRL database. We thank the people involved in the acquiring, analysis and accessibility of these data.  
589 This manuscript was largely improved by constructive reviews from Murat Aydin and Maria Elena Popa.

*Data availability.* The high resolution carbon monoxide datasets are available on the PANGEA database.

590 *Author contributions.* This scientific project was designed by XF, JC, VVP, RR, EB, and TB. The high resolution carbon monoxide measurements  
591 were carried out by XF and RR, with support of KF, TB, JC and EB. Discrete CO analyses were carried out by PP and EC, and  
592 chemistry analyses were carried out by NC and JRM. XF, PP, and VVP participated in the PLACE drilling. RR participated in the Tunu13  
593 drilling. The codes for data processing and modelling were developed by KF, JRM, and XF. All authors contributed to the interpretation of  
594 the data. The manuscript was written by XF with the help of all co-authors.

595 *Competing interests.* The authors declare having no competing interests

## References

Allan, D.: Statistics of atomic frequency standards, *Proceedings of the IEEE*, 54, 221–230, <https://doi.org/10.1109/PROC.1966.4634>, 1966.

Butler, J. H., Battle, M. O., Bender, M., Montzka, S. A., Clarke, A. D., Saltzman, E. S., M., S. C., Severinghaus, J., and Elkins, J. W.: A record of atmospheric halocarbons during the twentieth century from polar firn air, *Nature*, 399, 749–755, 1999.

600 Clark, I. D., Henderson, L., Chappellaz, J., Fisher, D., Koerner, R., Worthy, D. E. J., Kotzer, T., Norman, A. L., and Barnola, J. M.: CO<sub>2</sub> isotopes as tracers of firn air diffusion and age in an Arctic ice cap with summer melting, Devon Island, Canada, *Journal of Geophysical Research-Atmospheres*, 112, <https://doi.org/10.1029/2006jd007471>, 2007.

Collins, W. J., Lamarque, J.-F., Schulz, M., Boucher, O., Eyring, V., Hegglin, M. I., Maycock, A., Myhre, G., Prather, M., Shindell, D., and Smith, S. J.: AerChemMIP: quantifying the effects of chemistry and aerosols in CMIP6, *Geoscientific Model Development*, 10, 585–607, <https://doi.org/10.5194/gmd-10-585-2017>, 2017.

605 community members, N.: High-resolution record of Northern Hemisphere climate extending into the last interglacial period, *Nature*, <https://doi.org/10.1038/nature02805>, 2004.

Conte, L., Szopa, S., Séférian, R., and Bopp, L.: The oceanic cycle of carbon monoxide and its emissions to the atmosphere, *Biogeosciences*, 16, 881–902, <https://doi.org/10.5194/bg-16-881-2019>, 2019.

610 Crutzen, P. J.: A discussion of the chemistry of some minor constituents in the stratosphere and troposphere, *Pure and Applied Geophysics*, 106-108, 1385–1399, <https://doi.org/10.1007/BF00881092>, 1973.

Deeter, M. N., Edwards, D. P., Francis, G. L., Gille, J. C., Martínez-Alonso, S., Worden, H. M., and Sweeney, C.: A climate-scale satellite record for carbon monoxide: the MOPITT Version 7 product, *Atmospheric Measurement Techniques*, 10, 2533–2555, <https://doi.org/10.5194/amt-10-2533-2017>, 2017.

615 Duncan, B., Logan, J. A., Bey, I., Megretskia, I. A., Yantosca, R. M., Novelli, P. C., Jones, N. B., and Rinsland, C. P.: Global budget of CO, 1988–1997: Source estimates and validation with a global model, *Journal of Geophysical Research-Atmospheres*, 112, <https://doi.org/10.1029/2007jd008459>, 2007.

Faïn, X., Chappellaz, J., Rhodes, R. H., Stowasser, C., Blunier, T., McConnell, J. R., Brook, E. J., Preunkert, S., Legrand, M., Debois, T., and Romanini, D.: High resolution measurements of carbon monoxide along a late Holocene Greenland ice core: evidence for in situ production, *Climate of the Past*, 10, 987–1000, <https://doi.org/10.5194/cp-10-987-2014>, 2014.

620 Fourteau, K., Faïn, X., Martinerie, P., Landais, A., Ekyakin, A. A., Lipenkov, V. Y., and Chappellaz, J.: Analytical constraints on layered gas trapping and smoothing of atmospheric variability in ice under low-accumulation conditions, *Climate of the Past*, 13, 1815–1830, <https://doi.org/10.5194/cp-13-1815-2017>, 2017.

Gkinis, V., Popp, T. J., Blunier, T., Bigler, M., Schupbach, S., Kettner, E., Johnsen, S. J., and Schüpbach, S.: Water isotopic ratios from a 625 continuously melted ice core sample, *Atmospheric Measurement Techniques*, 4, 2531–2542, <https://doi.org/10.5194/amt-4-2531-2011>, 2011.

Haan, D. and Raynaud, D.: Ice core record of CO variations during the last two millennia: atmospheric implications and chemical interactions within the Greenland ice, *Tellus Series B-Chemical and Physical Meteorology*, 50, 253–262, 1998.

Haan, D., Martinerie, P., and Raynaud, D.: Ice core data of atmospheric carbon monoxide over Antarctica and Greenland during the last 200 630 years, *Geophysical Research Letters*, 23, 2235–2238, 1996.

Haan, D., Zuo, Y., Gros, V., and Brenninkmeijer, C. A. M.: Photochemical production of carbon monoxide in snow, *Journal of Atmospheric Chemistry*, 40, 217–230, 2001.

Hagler, G. S. W., Bergin, M. H., Smith, E. a., Dibb, J. E., Anderson, C., and Steig, E. J.: Particulate and water-soluble carbon measured in recent snow at Summit, Greenland, *Geophysical Research Letters*, 34, [https://doi.org/https://doi.org/10.1029/2007GL030110](https://doi.org/10.1029/2007GL030110), 2007.

635 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-i., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.

Khalil, M., Pinto, J., and Shearer, M.: Atmospheric carbon monoxide, *Chemosphere - Global Change Science*, 1, ix–xi, 640 [https://doi.org/10.1016/S1465-9972\(99\)00053-7](https://doi.org/10.1016/S1465-9972(99)00053-7), 1999.

Kuhl, T. W., Johnson, J. A., Shturmakov, A. J., Goetz, J. J., Gibson, C. J., and Lebar, D. A.: A new large-diameter ice-core drill: the Blue Ice Drill, *Annals of Glaciology*, 55, 1–6, <https://doi.org/10.3189/2014AoG68A009>, 2014.

Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. a., Nagashima, T., Naik, V., Plummer, 645 D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, a., and Zeng, G.: The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): overview and description of models, simulations and climate diagnostics, *Geoscientific Model Development*, 6, 179–206, <https://doi.org/10.5194/gmd-6-179-2013>, 2013.

Legrand, M. and De Angelis, M.: Light carboxylic acids in Greenland ice: A record of past forest fires and vegetation emissions from the boreal zone, *Journal of Geophysical Research*, 101, 4129, <https://doi.org/10.1029/95JD03296>, 1996.

650 Legrand, M. and Mayewski, P.: Glaciochemistry of polar ice cores: A review, *REVIEWS OF GEOPHYSICS*, 35, 219–243, 1997.

Legrand, M., McConnell, J. R., Fischer, H., Wolff, E. W., Preunkert, S., Arienzo, M., Chellman, N., Leuenberger, D., Maselli, O. J., Place, P., Sigl, M., Schupbach, S., and Flannigan, M.: Boreal fire records in Northern Hemisphere ice cores: A review, *Climate of the Past*, 12, 2033–2059, <https://doi.org/10.5194/cp-12-2033-2016>, 2016.

Lelieveld, J., Gromov, S., Pozzer, A., and Taraborrelli, D.: Global tropospheric hydroxyl distribution, budget and reactivity, *Atmospheric 655 Chemistry and Physics*, 16, 12 477–12 493, <https://doi.org/10.5194/acp-16-12477-2016>, 2016.

Maselli, O. J., Fritzsche, D., Layman, L., McConnell, J. R., and Meyer, H.: Comparison of water isotope-ratio determinations using two cavity ring-down instruments and classical mass spectrometry in continuous ice-core analysis., *Isotopes in environmental and health studies*, 49, 387–98, <https://doi.org/10.1080/10256016.2013.781598>, 2013.

660 McConnell, J. R. and Edwards, R.: Coal burning leaves toxic heavy metal legacy in the Arctic., *Proceedings of the National Academy of Sciences of the United States of America*, 105, 12 140–12 144, <https://doi.org/10.1073/pnas.0803564105>, 2008.

McConnell, J. R., Lamorey, S., and Taylor, K. C.: Continuous ice-core chemical analyses using inductively coupled plasma mass spectrometry, *Environmental Science and Technology*, 36, 7–11, 2002.

McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J. R., Pasteris, D. R., Carter, M. M., and Kahl, J. D. W.: 20th-century industrial black carbon emissions altered arctic climate forcing, *Science*, 317, 1381–1384, 2007.

665 Morville, J., Kassi, S., Chenevier, M., and Romanini, D.: Fast, low-noise, mode-by-mode, cavity-enhanced absorption spectroscopy by diode-laser self-locking, *Applied Physics B-Lasers and Optics*, 80, 1027–1038, <https://doi.org/10.1007/s00340-005-1828-z>, 2005.

NEEM community members: Eemian interglacial reconstructed from a Greenland folded ice core, *Nature*, 493, 489, <https://doi.org/10.1038/nature11789>, 2012.

670 Novelli, P. C., Masarie, K. A., and Lang, P. M.: Distributions and recent changes of carbon monoxide in the lower troposphere, *Journal of Geophysical Research*, 103, 19 015–19 033, 1998.

Petrenko, V. V., Martinerie, P., Novelli, P. C., Etheridge, D. M., Levin, I., Wang, Z., Blunier, T., Chappellaz, J., Kaiser, J., Lang, P., Steele, L. P., Hammer, S., Mak, J., Langenfelds, R. L., Schwander, J., Severinghaus, J., Witrant, E., Petron, G., Battle, M. O., Forster, G., Sturges, W. T., Lamarque, J.-F., Steffen, K., and White, J. W. C.: A 60 yr record of atmospheric carbon monoxide reconstructed from Greenland firn air, *Atmospheric Chemistry and Physics*, 13, 7567–7585, <https://doi.org/10.5194/acp-13-7567-2013>, 2013.

675 Rhodes, R. H., Faïn, X., Stowasser, C., Blunier, T., Chappellaz, J., McConnell, J. R., Romanini, D., Mitchell, L. E., and Brook, E. J.: Continuous methane measurements from a late Holocene Greenland ice core: Atmospheric and in-situ signals, *Earth and Planetary Science Letters*, 368, 9–19, <https://doi.org/10.1016/j.epsl.2013.02.034>, 2013.

Rhodes, R. H., Brook, E. J., Chiang, J. C. H., Blunier, T., Maselli, O. J., McConnell, J. R., Romanini, D., and Severinghaus, J.: Enhanced tropical methane production in response to iceberg discharge in the North Atlantic, *Science*, 348, 1016–9, 680 <https://doi.org/10.1126/science.1262005>, 2015.

Rhodes, R. H., Faïn, X., Brook, E. J., McConnell, J. R., Maselli, O. J., Sigl, M., Edwards, J. S., Buzert, C., Blunier, T., Chappellaz, J., and Freitag, J.: Local artifacts in ice core methane records caused by layered bubble trapping and in situ production: a multi-site investigation, *Climate of the Past*, 12, 1061–1077, <https://doi.org/10.5194/cp-12-1061-2016>, 2016.

Schwander, J. and Rufli, H.: Electromechanical drilling of a 300-m core in a dry hole at Summit, *Mem. Natl Inst. Polar Res.*, 49, 93–94, 1994.

685 Stowasser, C., Buzert, C., Gkinis, V., Chappellaz, J., Schüpbach, S., Bigler, M., Faïn, X., Sperlich, P., Baumgartner, M., Schilt, a., Blunier, T., and Schupbach, S.: Continuous measurements of methane mixing ratios from ice cores, *Atmospheric Measurement Techniques*, 5, 999–1013, <https://doi.org/10.5194/amt-5-999-2012>, 2012.

van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, 690 <https://doi.org/10.5194/essd-9-697-2017>, 2017.

Wang, Z., Chappellaz, J., Martinerie, P., Park, K., Petrenko, V. V., Witrant, E., Emmons, L. K., Blunier, T., Brenninkmeijer, C. A. M., and Mak, J.: The isotopic record of Northern Hemisphere atmospheric carbon monoxide since 1950: implications for the CO budget, *Atmospheric Chemistry and Physics*, 12, 4365–4377, <https://doi.org/10.5194/acp-12-4365-2012>, 2012.

Zheng, B., Chevallier, F., Yin, Y., Ciais, P., Fortems-Cheiney, A., Deeter, M. N., Parker, R. J., Wang, Y., Worden, H. M., and Zhao, Y.: Global atmospheric carbon monoxide budget 2000–2017 inferred from multi-species atmospheric inversions, *Earth System Science Data*, 11, 1411–1436, <https://doi.org/10.5194/essd-11-1411-2019>, 2019.