

# The unidentified eruption of 1809: a climatic cold case

Claudia Timmreck<sup>1</sup>, Matthew Toohey<sup>2</sup>, Davide Zanchettin<sup>3</sup>, Stefan Brönnimann<sup>4</sup>, Elin Lundstad<sup>4</sup>, and Rob Wilson<sup>5</sup>

<sup>1</sup>The Atmosphere in the Earth System, Max Planck Institute for Meteorology, Bundesstr. 53, 20146 Hamburg, Germany

<sup>2</sup>Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, Canada

<sup>3</sup>Department of Environmental Sciences, Informatics and Statistics, University Ca' Foscari of Venice, Mestre, Italy

<sup>4</sup>Institute of Geography Climatology and Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

<sup>5</sup>School of Earth & Environmental Sciences, University of St. Andrews, United Kingdom

**Correspondence:** Claudia Timmreck (claudia.timmreck@mpimet.mpg.de)

**Abstract.** The “1809 eruption” is one of the most recent unidentified volcanic eruptions with a global climate impact. Even though the eruption ranks as the 3rd largest since 1500 with a sulfur emission strength estimated to be two times that of the 1991 eruption of Pinatubo, not much is known of it from historic sources. Based on a compilation of instrumental and reconstructed temperature time series, we show here that tropical temperatures show a significant drop in response to the ~1809 eruption, similar to that produced by the Mt. Tambora eruption in 1815, while the response of Northern Hemisphere (NH) boreal summer temperature is spatially heterogeneous. We test the sensitivity of the climate response simulated by the MPI Earth system model to a range of volcanic forcing estimates constructed using estimated volcanic stratospheric sulfur injections (VSSI) and uncertainties from ice-core records. Three of the forcing reconstructions represent a tropical eruption with approximately symmetric hemispheric aerosol spread but different forcing magnitudes, while a fourth reflects a hemispherically asymmetric scenario without volcanic forcing in the NH extratropics. Observed and reconstructed post-volcanic surface NH summer temperature anomalies lie within the range of all the scenario simulations. Therefore, assuming the model climate sensitivity is correct, the VSSI estimate is accurate within the uncertainty bounds. Comparison of observed and simulated tropical temperature anomalies suggests that the most likely VSSI for the 1809 eruption would be somewhere between 12 -19 Tg of sulfur. Model results show that NH large-scale climate modes are sensitive to both volcanic forcing strength and its spatial structure. While spatial correlations between the N-TREND NH temperature reconstruction and the model simulations are weak in terms of the ensemble mean model results, individual model simulations show good correlation over North America and Europe, suggesting the spatial heterogeneity of the 1810 cooling could be due to internal climate variability.

## 1 Introduction

The early 19th century (~1800-1830 CE), at the tail end of the Little Ice Age, marks one of the coldest period of the last millennium (e.g. Wilson et al., 2016; PAGES 2k Consortium, 2019) and is therefore of special interest in the study of inter-decadal climate variability (Jungclauss et al., 2017). It was influenced by strong natural forcing: a grand solar minimum (Dalton Minimum, ~1790-1820 CE) and simultaneously a cluster of very strong tropical volcanic eruptions that includes the widely

known Mt. Tambora eruption in 1815, an unidentified eruption estimated to have occurred in 1808 or 1809, and a series of eruptions in the 1820s and 1830s. Brönnimann et al. (2019a) point out that this sequence of volcanic eruptions influenced the last phase of the Little Ice Age by not only leading to global cooling but also by modifying the large-scale atmospheric circulation through a southward shift of low-pressure systems over the North Atlantic related to a weakening of the African monsoon and the Atlantic–European Hadley cell (Wegmann et al., 2014).

The Mt. Tambora eruption in April 1815 was the largest in the last 500 years and had substantial global climatic and societal effects (e.g., Oppenheimer, 2003; Brönnimann and Krämer, 2016; Raible et al., 2016). In contrast to the Mt. Tambora eruption, little is known about the 1809 eruption. Although there is no historical source reporting a strong volcanic eruption in 1809, its occurrence is indubitably brought to light by ice-core sulfur records, which clearly identify a peak in volcanic sulfur in 1809/1810 (Dai et al., 1991). Simultaneous signals in both Greenland and Antarctic ice-cores with similar magnitude are consistent with a tropical origin, and analysis of sulfur isotopes in ice-cores supports the hypothesis of a major volcanic eruption with stratospheric injection (Cole-Dai et al., 2009).

Based on ice-core sulfur records from Antarctica and Greenland, the 1809 eruption is estimated to have injected  $19.3 \pm 3.54$  Tg of sulfur (S) into the stratosphere (Toohey and Sigl, 2017). This value is roughly 30% less than the estimate for the 1815 Mt. Tambora eruption, and roughly twice that of the 1991 Pinatubo eruption. Accordingly, the 1809 eruption produced the 2nd largest volcanic stratospheric sulfur injection (VSSI) of the 19th century, and the 6th largest of the past 1000 years. For comparison, the Ice-core Volcanic Index 2 (IVI2) database (Gao et al., 2008) estimates that the 1809 eruption injected 53.7 Tg of sulfate aerosols, which corresponds to 13.4 Tg S. While smaller than the estimate of Toohey and Sigl (2017), the IVI2 value lies within the reported 2-sigma uncertainty range. Uncertainties in VSSI and related uncertainties in the radiative impacts of the volcanic aerosol could be relevant for the interpretation of post-volcanic climate anomalies, as recently discussed for the 1815 Mt. Tambora eruption and the year without summer in 1816 (Zanchettin et al., 2019; Schurer et al., 2019).

While the location and the magnitude, of the 1809 eruption are unknown, its exact timing is also uncertain. A detailed analysis of high-resolution ice-core records points to an eruption in February 1809  $\pm$  4 months (Cole-Dai, 2010) which is consistent with the timing implied by other high-resolution ice-core records (Sigl et al., 2013, 2015; Plummer et al., 2012). Observations from South America of atmospheric phenomena consistent with enhanced stratospheric aerosol (Guevara-Murua et al., 2014) suggest a possible eruption in late November or early December 1808 ( $4.12.1808 \pm 7$  days), although there is no direct link between these observations and the ice-core sulfate signals. Chenoweth (2001) proposed an eruption date of March-June 1808 based on a sudden cooling in Malaysian temperature data and maximum cooling of marine air temperature in 1809. Such uncertainty in the eruption date has implications for the associated spatio-temporal pattern of aerosol dispersal and hemispheric and global climate impacts (Toohey et al., 2011; Timmreck, 2012). The climatic impacts of the 1809 eruption have been mostly studied in the context of the early 19th century volcanic cluster (e.g., Cole-Dai et al., 2009; Zanchettin et al., 2013, 2019; Anet et al., 2014; Winter et al., 2015; Brönnimann et al., 2019a) or of multi-eruption investigations (e.g., Fischer et al., 2007; Rao et al., 2017). Lesser attention has been placed on characterizing and understanding the short-term climatic anomalies that specifically followed the 1809 eruption. Available observations and reconstructions indicate ambiguous signals in NH land-mean summer temperatures reconstructed from tree-ring data for this period. For example, Schneider et al. (2017) found that,

among the ten largest eruptions of the past 2500 years, the 1809 event was one of two that did not produce a significant “break” in the temperature time series. While the temperature reconstruction reports cooling in 1809/1810, Schneider et al. (2017) note that reconstructed temperatures did not return to their climatological mean after the initial drop and remained low until the Mt. Tambora eruption in 1815. Hakim et al. (2016) presented multivariate reconstructed fields for the 1809 volcanic eruption from the last millennium climate reanalysis (LMR) project. They found abrupt global surface cooling in 1809 which was reinforced in 1815. The post volcanic global-mean 2m temperature anomalies show however a wide spread of up to 0.3 °C in the LMR between ensemble members and experiments using different combinations of calibration data for the proxy system models and prior data in the reconstruction. Using the LMR paleoenvironmental data assimilation framework, Zhu et al. (2020) demonstrate that some of the known discrepancies between tree-ring data and paleoclimate models can partly be resolved by assimilating tree-ring density records only and focusing on growing-season temperatures instead of annual temperature while performing the comparison at the proxy locales. However differences remain for large events like the Mt. Tambora 1815 eruption.

In this study, we investigate the climate impact of the 1809 eruption by using Earth system model ensemble simulations and by analyzing new and existing observational and proxy based data sets. We explore how uncertainties in the magnitude and spatial structure of the forcing propagate to the magnitude and ensemble variability of post-eruption regional and hemispheric climate anomalies.

In section 2, we briefly describe the applied methods, model, experiments and data sets. Section 3 provides an overview of the reconstructed and observed climate effects of the 1809 eruption, while section 4 presents the main results of the model experiments including a model-data intercomparison. The results are discussed in section 5. The paper ends with a summary and conclusions (section 6).

## **2 Methods and Data**

### **2.1 Methods**

#### **2.1.1 Model**

We use the latest low-resolution version of the Max-Planck-Institute Earth-System-Model (MPI-ESM1.2-LR, Mauritsen et al. (2019)), an updated version of the MPI-ESM used in the Coupled Model Intercomparison Project Phase 5 CMIP5 (Giorgetta et al., 2013). The applied MPI-ESM1.2 configuration is one of the two reference versions used in the Coupled Model Intercomparison Project Phase 6 (CMIP6, see Eyring et al., 2016). It consists of four components: the atmospheric general circulation model ECHAM6 (Stevens et al., 2013), the ocean-sea ice model MPIOM (Jungclaus et al., 2013), the land component JSBACH (Reick et al., 2013) and the marine biogeochemistry model HAMOCC (Ilyina et al., 2013). JSBACH is directly coupled to the ECHAM6.3 model and includes dynamic vegetation, whereas HAMOCC is directly coupled to the MPIOM. ECHAM6 and MPIOM are in turn coupled through the OASIS3-MCT coupler software. In MPI-ESM1.2, ECHAM6.3 is used, which is run with a horizontal resolution in the spectral space of T63 (~200 km) and with 47 vertical levels up to 0.01 hPa with 13 model levels above 100 hPa. In ECHAM6.3 aerosol microphysical processes are not included. The radiative forcing of the volcanic

90 aerosol is prescribed by monthly and zonal mean optical parameters which are generated with the Easy Volcanic Aerosol forcing generator (EVA, Toohey et al., 2016), see section 2.1.2


The MPIOM, which is run in its GR15 configuration with a nominal resolution of  $1.5^\circ$  around the equator and 40 vertical levels, has remained largely unchanged with respect to the CMIP5 version. Several revisions with respect to the MPI-ESM CMIP5 version have however been made for the for the atmospheric model including a new representation of radiation transfer, 95 land physics and biogeochemistry components and the ocean carbon cycle. A detailed description of all updates is given in Mauritsen et al. (2019). Previous studies have successfully shown the MPI-ESM is especially well-suited for paleo-applications and has been widely tested and employed in the context of the climate of the last millennium (e.g., Jungclaus et al., 2014; Zanchettin et al., 2015; Moreno-Chamarro et al., 2017).

### 2.1.2 Forcing

100 The applied volcanic forcing is compiled with the Easy Volcanic Aerosol (EVA) forcing generator (Toohey et al., 2016). EVA provides an analytic representation of volcanic stratospheric aerosol forcing, prescribing the aerosol's radiative properties and primary modes of their spatial and temporal variability. Although EVA represents an idealized forcing approach, its forcing estimates lie within the multi model range of global aerosol simulations for the Tambora eruption (Zanchettin et al., 2016; Clyne et al., 2021). This permits the compilation of physically consistent forcing estimates also for historic eruptions. EVA 105 uses sulfur dioxide ( $\text{SO}_2$ ) injection time series as input and applies a parameterized three-box model of stratospheric transport to reconstruct the space–time structure of sulfate aerosol evolution. Simple scaling relationships serve to construct stratospheric aerosol optical depth (SAOD) at  $0.55 \mu\text{m}$  and aerosol effective radius from the stratospheric sulfate aerosol mass, from which wavelength-dependent aerosol extinction, single scattering albedo, and scattering asymmetry factors are derived for pre-defined wavelength bands and latitudes. Volcanic stratospheric sulfur injection (VSSI) values for the simulations performed in this 110 work are taken from the eVolv2k reconstruction based on sulfate records from various ice cores from Greenland and Antarctica (Toohey and Sigl, 2017). Compared to prior volcanic reconstructions, eVolv2k includes improvements of the ice-core records in terms of synchronization and dating, as well as in the methods used to estimate VSSI from them.

Consistent with the estimated range given by Cole-Dai (2010) and the convention for unidentified eruptions used by Crowley and Unterman (2013), the eruption date of the unidentified 1809 eruption is set to occur on the 1st of January 1809 and located 115 at the equator. The eVolv2k best estimate for the VSSI of the 1809 eruption is  $19.3 \text{ Tg S}$  with a  $1\sigma$  uncertainty of  $\pm 3.54 \text{ Tg S}$ , based on the variability between individual ice-core records and model-based estimates of error due to the limited hemispheric sampling provided by ice sheets. To incorporate this uncertainty into climate model simulations, we constructed aerosol forcing time series using the central (or best) VSSI estimate, as well as versions which perturbed the central estimate by adding and subtracting two times the estimated uncertainty ( $\pm 2\sigma$ ) from the central VSSI estimate. These three forcing sets are hereafter 120 termed “Best”, “High”, and “Low”, respectively. Constructed in this manner, the range from “Low” to “High” forcing should roughly span a 95% confidence interval of the global-mean aerosol forcing.

There are other important sources of uncertainty in the reconstruction of stratospheric aerosol, further to that related to the magnitude of the sulfur deposition. For example, the transport of aerosol from the tropics to each hemisphere has been



**Figure 1.** Volcanic radiative forcing. Global stratospheric aerosol optical depth (SAOD) at  $0.55 \mu\text{m}$  based on eVolv2k VSSI estimates (Toohey and Sigl, 2017) and calculation with the volcanic forcing generator EVA (Toohey et al., 2016) for the four different forcing scenarios: “Best”, “Low”, “High” and “nNHP” for the 1809 eruption and the “Best” forcing scenario for the Mt. Tambora eruption. Bottom: Spatial and temporal distribution of a zonal mean stratospheric SAOD for the four experiments.

seen to be quite variable for the tropical eruptions of Pinatubo in June 1991, El Chichón in April 1982 and Agung in March  
 125 1963, which likely arises due to the particular meteorological conditions at the time of the eruption (Robock, 2000). While  
 the 1991 Pinatubo eruption produced an aerosol cloud that spread relatively evenly to each hemisphere, the aerosol from the  
 1982 El Chichón eruption and the 1963 Agung eruption were heavily biased to one hemisphere (Suppl. Fig. S2). Furthermore,  
 the lifetime, evolution and spatial structure of aerosol properties may vary significantly as a result of the injection height of  
 the volcanic plume (Toohey et al., 2019; Marshall et al., 2019). Recently, Yang et al. (2019) pointed out that an accurate  
 130 reconstruction of the spatial forcing structure of volcanic aerosol is important to get a reliable climate response. Motivated in  
 large part by the post-1809 surface temperature anomalies to be discussed below, which include strong cooling in the tropics  
 and a muted NH mean temperature signal, we constructed a fourth forcing set, which is identical to the “Best” forcing in the  
 tropics and in the Southern Hemisphere (SH), but has the aerosol mass in the NH extratropics completely removed, creating  
 a strongly asymmetric forcing structure. This forcing scenario, which we call “no-NH plume” or “nNHP” in the following,  
 135 should be interpreted as a rather extreme “end-member” in terms of NH forcing. The lack of aerosol for the NH in this  
 constructed forcing is clearly inconsistent with the polar ice-core records of sulfate deposition from the 1809 eruption, which  
 suggests roughly equal deposition between Greenland and Antarctica over a similar duration to other typical tropical eruptions,  
 indicating a long-lasting and global aerosol spread. Due to uncertainties in the conversion of ice-core sulfate to hemispheric

aerosol burden and radiative forcing (Toohey and Sigl, 2017; Marshall et al., 2020), it is not impossible that the radiative  
140 forcing from the 1809 eruption aerosol was characterized by some degree of hemispheric asymmetry in reality. Still, the nNHP  
forcing presented here should be interpreted as a rather unlikely scenario for the 1809 eruption. We explore here the impact  
of this forcing scenario as an extreme, idealized form of hemispheric asymmetry that might conceivably play some role in the  
response to the 1809 eruption, and is directly applicable to “unipolar” tropical eruptions like Agung (1963) and El Chichon  
(1982).

145 Time series of global-mean and zonal-mean SAOD at  $0.55 \mu\text{m}$  for the different 1809 aerosol forcing scenarios discussed  
above are shown in Fig. 1, together with the “Best” scenario after the Mt. Tambora eruption. Peak global-mean SAOD values  
following the 1809 eruption range from 0.17 to 0.33 from the “Low” to “High” scenarios, roughly corresponding to forcing  
from a little stronger than that from the 1991 Pinatubo eruption to a little weaker than the 1815 Mt. Tambora eruption, re-  
spectively. The “nNHP” scenario produces a global-mean SAOD that peaks at a value of 0.21, i.e., in between the “Low” and  
150 “Best” scenarios, and decays in a manner very similar in magnitude to the “Low” scenario. The latitudinal spread of aerosol  
is relatively evenly split between the NH and SH in the “Best”, “Low”, and “High” scenarios, with offsets in the timing of  
the peak hemispheric SAOD resulting from the parameterized seasonal dependence of stratospheric transport in EVA. After  
the removal of aerosol mass from the NH extratropics in the construction of the “nNHP” scenario, the SAOD is predictably  
negligible in the NH extratropics and a strong gradient in SAOD is produced at  $30^\circ\text{N}$ .

### 155 2.1.3 Experiments

We have performed ensemble simulations of the early 19th century with the MPI-ESM1.2-LR for each of the four forcing  
scenarios for the 1809 eruption (*Best*, *High*, *Low* and *nNHP*). All simulations also include the eVolv2k “Best” forcing estimate  
for the Mt. Tambora eruption from 1815 onwards. Related experiments using a range of different forcing estimates for the 1815  
Mt. Tambora eruption were used in Zanchettin et al. (2019) and Schurer et al. (2019) to investigate the role of volcanic forcing  
160 uncertainty in the climate response to the 1815 Mt. Tambora eruption, in particular the “year without summer” in 1816. For  
each experiment we have produced 10 realizations branched off every 100 to 200 years from an unperturbed 1,200-year-long  
pre-industrial control run (constant forcing, excluding background volcanic aerosols) to account for internal climate variability.  
All simulations were initialized on 1st January 1800 with constant preindustrial forcing except for stratospheric aerosol forcing.

## 2.2 Data

### 165 2.2.1 Temperature reconstructions

#### Tropical temperature reconstructions

In our study, we compare three different sea-surface temperature (SST) reconstructions with the MPI-ESM simulations.  
The temperature reconstruction TROP is a multi-proxy tropical ( $30^\circ\text{N}$ – $30^\circ\text{S}$ ,  $34^\circ\text{E}$ – $70^\circ\text{W}$ ) annual SST reconstruction between  
1546–1998 (D’Arrigo et al., 2009). TROP consists of 19 coral, tree-ring and ice-core proxies located between  $30^\circ\text{N}$  and  $30^\circ\text{S}$ .  
170 The records were selected on the basis of data availability, dating certainty, annual or higher resolution, and a documented

relationship with temperature. It shows annual-to multi decadal-scale variability and explains 55% of the annual variance in the most replicated period 1897–1981. Further, 400-year long spatially resolved tropical SST reconstructions for four specific regions: the Indian Ocean (20°N–15°S, 40–100°E), the western (25°N–25°S, 110–155°E) and eastern Pacific (10°N–10°S, 175°E–85°W), and the western Atlantic (15–30°N, 60–90°W) were compiled by Tierney et al. (2015) based on 57 published and publicly archived marine paleoclimate data sets. The four regions were selected based on the availability of nearby coral sampling sites and an analysis of spatial temperature covariance. An even more regionally specific SST reconstruction was developed by D'Arrigo et al. (2006) for the Indo-Pacific warm pool region (15°S–5°N, 110–160°E) using annually resolved teak ring width and coral  $\delta^{18}\text{O}$  records. This September–November mean SST reconstruction dates from A.D. 1782–1992 and explains 52% of the SST variance in the most replicated period. This record was used in the D'Arrigo et al. (2009) TROP reconstruction.

### **Northern Hemisphere extratropical temperature reconstruction**

We compare our climate simulations of the early 19th century with four near-surface air temperature (SAT) reconstruction, which have all been used to assess the impacts of volcanic eruptions on surface temperature. The N-TREND (Northern Hemisphere Tree-Ring Network Development) reconstructions (Wilson et al., 2016; Anchukaitis et al., 2017) are based on 54 published tree-ring records and use different parameters as proxies for temperature. Eleven of the records are derived from ring-width (RW), 18 from maximum latewood density (MXD), and 25 are mixed records which consist of a combination of RW, MXD, and blue intensity (BI) data (see, Wilson et al., (2016) for details). The N-TREND database domain covers the NH midlatitudes between 40° and 75°N with at least 23 records extending back to at least AD 978. Two versions of the N-TREND reconstructions are used herein. N-TREND (N), detailed in Wilson et al. (2016), is a large-scale mean composite May-August temperature reconstruction derived from averaging the 54 tree-ring records weighted to 4 longitudinal quadrats, with separate nested calibration and validation performed as each shorter record is removed back in time. N-TREND (S), detailed in Anchukaitis et al. (2017), is a spatial reconstruction of the same season, derived by using point-by-point multiple regression (Cook et al., 1994) of the tree-ring proxy records available within 1000 to 2000 kms of the centre point of each 5x5 degree instrumental grid cell. For each gridded reconstruction a similar nesting procedure was used as Wilson et al. (2016). Herein, we use the average of all the grid point reconstructions for the periods where the validation Reduction of Error (RE - Wilson et al. 2006) was greater than zero. The NVOLC reconstruction (Guillet et al., 2017) is a NH summer temperature reconstruction over land (40–90°N) composed of 25 tree-ring chronologies (12 MXD, 13 TRW) and 3 isotope series from Greenland ice-cores (DYE3, GRIP, Crete). NVOLC was generated using a nested approach and includes only chronologies which encompass the full time period between today and the 13th century. The temperature reconstruction by Schneider et al. (2015) is based on 15 MXD chronologies distributed across the NH extratropics. All the temperature reconstructions show distinct short-time cooling after the largest eruptions of the Common Era. However, Schneider et al. (2017) point to a notable spread in the post-volcanic temperature response across the different reconstructions. This has various possible explanations, including the different parameters used, the spatial domain of the reconstruction, the method(s) used for detrending, and choices made in the network compilations.

**Surface air temperature from English East India Company ship logs**

Brohan et al. (2012) compiled an early observational data set of weather and climate between 1789 and 1834 from records of the English East India Company (EEIC), which are archived in the British Library. The records include 891 ships' logbooks of voyages from England to India or China and back containing daily instrumental measurements of temperature and pressure, as well as wind-speed estimates. Several thousands of weather observations could be gained from these ship voyages across the Atlantic and Indian Oceans providing a detailed view of the weather and climate in the early 19th century. Brohan et al. (2012) found that mean temperatures expressed a modest decrease in 1809 and 1816 as a likely consequence of the two large tropical volcanic eruptions during the period. Following Brohan et al. (2012), here we calculate temperature anomalies from the SAT measurements recorded in the EEIC logs. We account for the relatively sparse and irregular spatial and temporal sampling by computing for each measurement its anomaly from the HadNMAT2 night marine air temperature climatology (Kent et al., 2013). The SAT anomalies were then binned according to month/year and location and averaged. We present the data as mean temperature anomalies for the tropics (20°S to 20°N) in monthly or annual means. To quantify the impact of the 1809 eruption, anomalies are referenced to the 1800-1808 time period.

**Station data**

Climate model output is compared with monthly temperature series from land stations that cover the period 1806-1820 from a number sources, as compiled in Brönnimann et al. (2019b). The sources include data available electronically from the German Weather Service (DWD), the Royal Dutch Weather service (KNMI), the International Surface Temperature Initiative (Rennie et al., 2014), and the Global Historical Climatology Network (Lawrimore et al., 2011). In addition, we added nine series digitized from the compilation of Friedrich Wilhelm Dove that were not contained in any of the other sources (Dove, 1838, 1839, 1842, 1845). Of the 73 series obtained, 20 had less than 50% data coverage within the period 1806 and 1820 and were thus not further considered. The remaining 53 time series (see appendix table A1) were deseasonalized based on the 1806-1820 mean seasonal cycle and grouped by region (see appendix table A2).

**2.3 Analysis of model output**

Post-eruption climatic anomalies in the volcanically-forced ensembles are compared with both anomalies from the control run (describing the range of intrinsic climate variability), and with anomalies from a set of proxy-based reconstructions and instrumental observations, providing a reference/target to evaluate the simulation under both volcanically-forced and unperturbed conditions. Comparison between the volcanically-forced ensembles and the control run is based on the generation of signals in the control simulation analogous to the post-eruption ensemble-mean and ensemble-spread anomalies. In practice, a large number (1000) of surrogate ensembles is sampled from the control run, each identified by a randomly chosen year as reference for the eruption. Ensemble means and spreads (defined by 5th and 95th percentiles) of such surrogate ensembles provide an empirical probability distribution that is used to determine the range of intrinsic variability, which is illustrated with the associated 5th-95th percentile ranges. Differences between the volcanically-forced ensembles and the surrogate ensembles are tested




statistically through the Mann-Whitney U test (following, e.g., Zanchettin et al., 2019). When the ensembles are compared with a one-value target, either an anomaly from reconstructions/observations or a given reference (e.g., zero), significance of the difference between the ensemble and the target is determined based on whether the latter exceeds a given percentile range from the ensemble (e.g., the interquartile or the 5th-95th percentile range) or, alternatively, based on a t-test.

Integrated spatial analysis between the simulations and the N-TREND (S) gridded reconstruction is performed through a combination of the Root Mean Square Error (RMSE) and spatial correlation. Both metrics are calculated by including grid points in the reconstructions that correspond to the proxy locations and interpolating the model output to those locations with a nearest-neighbour algorithm. The relative contribution of each location is weighted by the cosine of its latitude to account for differences in the associated grid-cell area.

### 3 The 1809 eruption in climatic observations and proxy records

In proxy and instrumental records of tropical temperatures, cooling in the years 1809-1811 is generally on par with that after the 1815 Mt. Tambora eruption. Based on annually-resolved temperature-related records from corals, TRs and ice-cores, D'Arrigo et al. (2009) report peak tropical cooling of  $-0.77^{\circ}\text{C}$  in 1811, compared to  $-0.84^{\circ}\text{C}$  in 1817 (Fig. 2a). Tropical SST variability is modulated by El-Nino Southern Oscillation (ENSO) variability such as neutral to La Nina-like conditions in 1810 and El Nino-like ones in 1816 (Li et al., 2013; McGregor et al., 2010). The lagged response to the 1809 and 1815 eruptions in the TROP reconstruction is therefore most likely a result of an overlaying El Nino signal. Removing the ENSO signal from the TROP reconstructions led to a shift of maximum post volcanic cooling from 2 years after the eruption to one year (D'Arrigo et al., 2009). A clear signal is found in reconstructed Indo-Pacific warm pool SST anomalies from the post-1809 period 1809–1812, with values of  $-0.28^{\circ}\text{C}$ ,  $-0.73^{\circ}\text{C}$ ,  $-0.76^{\circ}\text{C}$  and  $-0.79^{\circ}\text{C}$ ; compared to  $-0.30^{\circ}\text{C}$ ,  $-0.51^{\circ}\text{C}$  and  $-0.51^{\circ}\text{C}$  for the post-Tambora period 1815 – 1817 (D'Arrigo et al., 2006). Chenoweth (2001) reports pronounced tropical cooling from ship-based marine SAT measurements in 1809 ( $-0.84^{\circ}\text{C}$ ), similar to that in 1816 ( $-0.81^{\circ}\text{C}$ ). More recent analysis of a larger set of ship-based marine SAT records from the EEIC by Brohan et al. (2012) suggests a more modest cooling for the two early 19th century eruptions, of about  $0.5^{\circ}\text{C}$  (Fig. 2b). However, the cooling is again found to be of comparable magnitude after the 1809 and 1815 eruptions, and therefore hints to a tropical location of the 1809 eruption in agreement with the ice-core data.

Tree-ring records capture volcanically-forced summer cooling very well (e.g., Briffa et al., 1998; Hegerl et al., 2003; Schneider et al., 2015; Stoffel et al., 2015). However, in the NH extratropics, SAT anomalies after 1809 are more spatially and temporally complex compared to the typical post-eruption pattern with broad NH cooling. In tree-ring-based temperature reconstructions for interior Alaska/Yukon (Briffa et al., 1994; Davi et al., 2003; Wilson et al., 2019), 1810 is one of the coldest summers identified over recent centuries. In earlier reconstructions of summer SAT in different regions of the western United States (Schweingruber et al., 1991; Briffa et al., 1992), 1810 was shown to be the third coldest summer in the British Columbia/Pacific Northwest region. Likewise, European tree-ring records show cooling after 1809 (e.g., Briffa et al., 1992; Wilson et al., 2016). In contrast, tree-ring networks in certain regions such as Eastern Canada show minimal response after 1809 (Gennaretti et al., 2018).



**Figure 2.** Observed and reconstructed temperature anomalies around the 1809 volcanic eruption. a) Reconstructed tropical ( $30^{\circ}\text{N}$ – $30^{\circ}\text{S}$ ,  $34^{\circ}\text{E}$ – $70^{\circ}\text{W}$ ) sea surface temperature (TROP, D’Arrigo et al., 2009), measured tropical marine surface air temperatures from EEIC ship logs (Brohan et al., 2012) and Indo-Pacific warm pool data (D’Arrigo et al., 2006). b) NH summer land temperatures from four tree-ring based reconstructions (Wilson et al. 2016 (N-TREND(N)), Anchukaitis et al., 2017 (N-TREND (S)), Guillet et al., 2017 (NVOLC), Schneider et al., 2015 (SCH15)). c-d) Monthly mean NH winter (c) and summer (d) temperature anomalies ( $^{\circ}\text{C}$ ) from 53 station data averaged over different European regions (Central Europe (CEUR:  $46.1$ – $52.5^{\circ}\text{N}$ ,  $6$ – $17.8^{\circ}\text{E}$ ), Eastern Europe (EEUR:  $47$ – $57^{\circ}\text{N}$ ,  $18$ – $32^{\circ}\text{E}$ ), Northern Europe (NEUR:  $55$ – $66^{\circ}\text{N}$ ,  $10$ – $31^{\circ}\text{E}$ ), Southern Europe ( $38$ – $46^{\circ}\text{N}$ ,  $7$ – $13.5^{\circ}\text{E}$ ), Western Europe (WEUR:  $48.5$ – $56^{\circ}\text{N}$ ,  $6^{\circ}\text{W}$ – $6^{\circ}\text{O}$ ) and New England (NENG:  $41$ – $44^{\circ}\text{N}$ ,  $73$ – $69^{\circ}\text{W}$ ). e-g) Mean surface temperature anomalies ( $^{\circ}\text{C}$ ) for boreal summers of 1809 (e), 1810 (f) and 1811 (g) in NH tree-ring data N-TREND (S) (Anchukaitis et al., 2017). Pink dots in panel e) illustrate the location of the tree-ring proxies used in the N-TREND reconstructions.

Based on compilations of regional records, tree-ring-based reconstructions of NH-mean land summer SAT show a large spread in hemispheric cooling after the 1809 eruption (Fig. 2b), with anomalies of  $-0.87^{\circ}\text{C}$ ,  $-0.77^{\circ}\text{C}$ ,  $-0.21^{\circ}\text{C}$  and  $-0.15^{\circ}\text{C}$  in 1810 for the N-TREND (S), NVOLC, N-TREND (N) and SCH15 reconstructions, respectively). Although using the same data set, the spatial N-TREND (S) and the nested N-TREND (N) reconstructions show a quite different behaviour. In N-TREND (S), the nature of the spatial multiple regression modelling biases the input records to those that correlate most strongly with local temperatures which, when available, were are likely MXD data. In all four reconstructions, NH temperature does not return to the climatological mean after an initial drop in 1810, but remains low, or even exhibits a continued cooling trend until


the Mt. Tambora eruption in 1815 (Schneider et al., 2015). The spatial variability of the reconstructed NH extratropical temperature response to the 1809 eruption is illustrated in Fig. 2e,f,g based on the spatially resolved N-TREND (S) reconstruction (Anchukaitis et al., 2017), displaying zonal oscillations consistent with a “wave-2” structure, that are especially evident in 1810 but already appreciable in 1809. This hemispheric structure is in contrast with the relatively uniform cooling seen in tree-ring records for Tambora (Fig. S3) and indeed for many of the largest eruptions of the past millennium (Hartl-Meier et al., 2017).

Information about regional and seasonal mean NH temperature anomalies in the early 19th century can be obtained from different station data across Europe and from New England (Fig. 2c,d). In NH winter the measurements reflect the high variability of local scale weather (Fig. 2c). Warm anomalies, an indication for post-eruption “NH winter warming”, are clearly visible in 1816/1817 in the 2nd winter after the Mt. Tambora eruption in 1815. Northern Europe shows the largest warm anomaly for all regions (about 3 °C). Warm NH winter anomalies between 1.5°C and 2 °C are seen in the winter 1809/1810 over Northern and Eastern Europe and over New England. Strong cooling is however found for the 1808/1809 winter in Northern and Central Europe. NH summer temperature anomalies are less variable than in winter (Fig. 2d). A local distinct cooling is found in the “year without summer” in 1816 over all regions except Northern Europe, where it occurs a year later. The cooling after the 1809 eruption is not so pronounced as after the Mt. Tambora eruption in 1815. In general the station data support the spatial distribution of the reconstructed near surface temperature anomalies derived from tree-ring data. They show a local minimum over Northern, Eastern and Southern Europe in NH summer 1810, which does not appear over Western and Central Europe and New England. The warm anomalies in the order of 2°C, which are found in summer 1811 over Eastern Europe are not however captured by the N-TREND spatial reconstruction although some slight warming is seen in the data over East Poland, Belarus and the Baltic States.

## 4 Results

### 4.1 Simulations

Firstly, we compare the simulated evolutions of monthly mean near-surface (2m) air temperature anomalies between the four experiments globally, in the tropics and NH extratropics (Fig. 3). Ensemble mean global mean temperature anomalies grow through 1809, and reach peak values through 1810 in all experiments, before decaying towards climatological values (Fig. 3a). Peak cooling reaches around 1.0 °C in the *High* experiment, compared to 0.5 °C in the *Low* and *nNHP* experiments. Peak temperature anomalies across the experiments correlates with the magnitude of prescribed AOD (Fig. 1a), and the responses are qualitatively consistent with expectations, with the AOD for the *Low* and *nNHP* experiments, which is similar in magnitude to that from the observed 1991 Pinatubo eruption, leading to global mean temperature anomalies also similar to those observed after Pinatubo. Global ensemble mean near-surface temperature anomalies are close together in *Low* and *nNHP* over boreal summer but differ for boreal winter when the intrinsic variability is higher. *Low* is the only experiment for which large-scale temperatures return within the 5th-95th percentile range percentile interval of the control run before the Mt. Tambora eruption in 1815. Global mean temperature anomalies of the other three experiments return only within the 5th-95th percentile range of unperturbed variability by 1815. As expected, almost no significant near-surface temperature anomalies are found for the




**Figure 3.** Global, tropical and extratropical temperature anomalies. Left: Simulated ensemble-mean monthly anomalies of (a) global, b) extra-tropical northern hemisphere and c) tropical averages of near-surface air temperature with respect to the pre-eruption (1800-1808) climatology. All data are deseasonalized using the respective annual average cycle from the control run. Thick (thin) black dashed lines are the 5th–95th percentile intervals for signal occurrence in the control run for the ensemble mean (ensemble spread). Bottom bars indicate periods when an ensemble member’s monthly mean temperature (color code as for the time series plots) is significantly different ( $p = 0.05$ ) from the control run according to the Mann-Whitney U test. Right: ensemble distributions (median, 25th–75th and 5th–95th percentile ranges) of seasonal mean anomalies for the first post-eruption winter (1809–1810, DJF) and summer (1810, JJA) following the 1809 eruption as well as for the pre-eruption period (1800–1808).

*nNHP* simulation in the NH extratropics except a few months in spring and autumn 1813 (Fig. 3b). The *nNHP* ensemble-mean values stay within the interquartile range of the control run but show a slight negative trend between 1809 and 1815. *nNHP* is also the only experiment where the NH extratropical summer of 1814 is colder than the summer of 1809. Internal variability is

relatively high in the NH extratropics in particular in NH winter spanning more than 1.5 °C. So, even the ensemble mean near  
 315 surface temperature anomalies for *Best* and *High* almost reach the 5th-95th percentile range of the control run in the 1st post  
 volcanic winters. Peak cooling appears for all experiments except *nNHP* in the summer 1810. In the tropics, *Best*, *High* and  
*nNHP* are outside the 5th-95th percentile range in the first four -post volcanic years while *Low* exceeds the 5th-95th percentile  
 range only for two years (Fig. 3c).

The ensemble distributions for the seasonal mean of winter 1809/1810 and summer 1810 illustrate the differences between  
 320 the four experiments not only in the mean anomaly but also for the ensemble spread (Fig. 3 d-f). While for example in summer  
 1810 the global and tropical ensemble mean of *Low* and *nNHP* are quite close, the ensemble spread is much larger in *Low*  
 compared to *nNHP*. *Low* has in general the largest ensemble spread independent of season and hemispheric scale. The clearest  
 separation between the experiments appears in the NH extratropics in summer 1810 (Fig. 3f) in line with Zanchettin et al.  
 (2019), who could show with a k mean cluster analysis on a large ensemble that forcing uncertainties can overwhelm initial-  
 condition spread in boreal summer.




**Figure 4.** Simulated ensemble-mean zonal-mean near-surface air temperature anomalies (°C) for the four MPI-ESM experiments. Only anomalies exceeding one standard deviation of the control run are shown. Anomalies are calculated with respect to the pre-eruption (1800-1808) climatology.

325

A more detailed spatial distribution of the simulated temporal evolution of post-volcanic surface temperature anomalies is seen in the Hovmöller diagram in Fig. 4. It shows that in all four experiments a multiannual surface temperature response is found in the tropics (30°S-30°N). In the inner tropics, the cooling disappears after one and a half years in *Low* and two to three years later in *Best*, *High* and *nNHP*. In the subtropics, a significant surface cooling signal is found over the ocean until 1815

330 in *Best* and *High*, while over land no significant cooling appears in 1814 (Fig. S4). A strong cooling signal is found in the NH extra tropics in *Best*, *Low* and *High* in summer 1810 and in *High* and to a small extent also in *Best* in summer 1811. In *nNHP* no surface cooling is detectable over the NH extratropics in the 1st four years after the eruption, consistent with the prescribed volcanic forcing (see Fig. 1). However, a cooling anomaly is apparent around 60°N in summer 1813, which is seen in the zonal mean over the ocean (Fig. S4) and likely due to decreased poleward ocean heat transport. Significant cooling south of 30°S appears only in austral spring 1809.




**Figure 5.** Simulated ensemble-mean near-surface air temperature anomalies for the 1st winter (1809/1810) and the second summer (1810) after the 1809 eruption for the four different MPI-ESM simulations. Shaded regions are significant at the 95% confidence level according to a t-test. Anomalies are calculated with respect to the period 1800-1808.

335

Fig. 5 shows the spatial near-surface air temperature anomalies for the first boreal winter (1809/1810) and the second boreal summer (1810) after the 1809 eruption for the four experiments. In general, the cooling is strongest over the NH

continents in all experiments revealing a strong cooling pattern over Alaska, Yukon and the Northwest Territories in the first post-eruption winter. In the *Best* and *High* experiments, relatively strong cold anomalies are found over the Central Asian dry highland regions around 40°N from the Hindukush in the west to the Pacific, while *Low* and *nNHP* show a small yet significant cooling over India and Southeast China. In boreal winter a significant warming is visible over Eurasia in all experiments except for *Low*, where warming anomalies are instead found over the polar ocean, and it is most pronounced in *High* but also quite extensive in *nNHP*. Such NH winter warming pattern is known to be induced by atmospheric circulation changes (e.g., Wunderlich and Mitchell, 2017; DallaSanta et al., 2019) and can occur in post-eruption winters as a dynamic response to the enhanced stratospheric aerosol layer, when it displays highly variable amplitude of local anomalies (Shindell et al., 2004). Accordingly, in our simulations the Eurasian winter warming pattern consists of one or two areas with positive temperature anomalies centered over various locations between Fennoscandia and the Central Siberian Plateau in the different simulations. Significant cooling, albeit of different strength, is found in the NH extratropics in boreal summer in the three symmetric-forcing experiments (*Best*, *High*, *Low*). However, while all of them show significant negative temperature anomalies over the North America continent with a local maximum over California and also cooling over Greenland, no significant anomalies are seen in *Low* over Fennoscandia. Except for some small regions (Finland, the Kola Peninsula and West Alaska), no significant cooling is found in *nNHP* in the NH extra tropics in boreal summer. The spatial distribution of the forcing can impact the latitudinal position of peak surface cooling, which in turn can lead to a shift of the Intertropical Convergence Zone (e.g., Haywood et al., 2013; Pausata et al., 2020). This is clearly visible in the cold anomaly belt over the Sahel region in the asymmetric forcing experiment *nNHP*. Significant warm anomalies are detectable in a small band that extends from the Caspian sea to the west to Japan to the East. Cooling over the ocean is weaker and mostly confined in the tropical belt between 30°S and 30°N. *High* is the only experiment where a significant El Nino type anomaly is seen over the Pacific Ocean in boreal summer 1810, while in the other three experiments a slight but non-significant warming appears off the coast of South-American. Looking to the relative SST anomalies as calculated after Khodri et al. (2017) an El Nino type anomaly is seen for all 4 scenarios in boreal summer 1810, while in winter 1809/1810 a significant warming anomaly appears in the central tropical Pacific in all experiments except *Best* (Fig. S5).

The substantial differences found in the post-eruption evolution of continental and subcontinental climates reflect the variety of climate responses produced by different combinations of internal climate variability and forcing structure. In this regard, post-eruption anomalies of selected dominant modes of large-scale atmospheric circulation in the Northern Hemisphere and the tropics, including the Pacific/North American pattern, the North Atlantic Oscillation, the North Pacific Index and the Southern Oscillation, yield a spread of responses within individual ensembles that is often as large as the range of pre-eruption variability. Further, response distributions generated by different forcings in some cases do not overlap (see Supplementary Figure S1).



**Figure 6.** Comparison of tropical temperatures anomalies. Comparison of the MPI-ESM simulations with a) tropical and annual mean (30°N-30°S) surface air temperature from shipbourne measurements of the English East India Company (EEIC, Brohan et al., 2012), b) annual mean tropical sea surface temperature (SST) reconstruction (TROP, D’Arrigo et al., 2009) over the tropical IndoPacific (30°N-30°S, 34°E-70°W), c) seasonal mean (Sep-Nov) SST reconstruction (D’Arrigo et al., 2006) anomalies over the Indonesian warmpool (WP, 15°S–5°N, 110–160°E). The black line represents in all panels the observed or reconstructed data while the colored lines represent ensemble means of the respective model simulations. The grey shaded regions in b) indicate the 95% confidence interval of the reconstruction. Anomalies are taken with respect to the years 1800 to 1808.




## 4.2 Model-data comparison

### 4.2.1 Tropics

370 A multiannual cooling signal is found in the MPI-ESM simulations in the tropical region after the unidentified 1809 eruption (Figs. 3 and 4). The same signature is detected in the English East India Company (EEIC) ship-based surface air temperature anomaly annual means (Brohan et al., 2012) and in tropical SSTs reconstruction (TROP, D'Arrigo et al., 2009) and the Indo-Pacific warm pool (D'Arrigo et al., 2006) in Fig. 6. The simulated ensemble mean temperatures (Fig. 6a) bracket the observed anomaly in the EEIC data in 1809, with the observed value falling between the results of the *Low* and *nNHP* forcing experiments. In 1810-1812, the cooling in the *Best*, *High* and *nNHP* experiments is stronger than that observed, and therefore the results from the *Low* experiment are generally most consistent with the ship-borne measurements (Fig. 6a). When the model results are sampled at the locations and times of the EEIC measurements (Fig. S6), the mean negative temperature anomalies in 1809 are 10-30% smaller, with *Best*, *High* and *nNHP* experiments all producing anomalies similar to that of the EEIC measurements. For the 1810-1812 period, the sampling makes little difference compared to the full tropical average, with *Best*, *High* and *nNHP* experiments all showing larger negative temperature anomalies than the EEIC measurements. A comparison of TROP with our four experiments reveals that all experiments lie within the 5th-95th percentile interval of the TROP reconstruction although the reconstructed SST response appears to be dampened in comparison to the model experiments (Fig. 6b). Although the long term trends of TROP and the model experiments are in general agreement, the dampened post volcanic cooling could reflect autocorrelative biases in the proxies (Lücke et al., 2019). Detailed scrutiny of high resolution tropical SST proxies and their potential biases to robustly reflect volcanically forced cooling has not been made in the same way that has been performed for tree-ring archives over the last decade (Anchukaitis et al., 2012; D'Arrigo et al., 2013; Esper et al., 2015; Franke et al., 2013; Lücke et al., 2019). A similar behavior is found for the Indonesian warm pool (Fig. 6c). However, in contrast to the whole tropics, the differences between the different forcing experiments are much smaller for the warm pool region compared to the wider tropical regions and the volcanic signal is more pronounced in the reconstructed SST at least for the unidentified 1809 eruption.


Tierney et al. (2015) provided coral-based reconstructions of tropical SSTs for four different ocean regions: the Indian Ocean, the West and the East Pacific, and the West Atlantic. Comparison of our four experiments with the coral-based reconstructions reveals quite different behaviour and simulation-reconstruction agreement across the various regions (Fig. 7). For the East Pacific region, the reconstruction and the MPI-ESM simulations are not inconsistent with each other over the 1809 period showing a substantial high variability (Fig. 7a) reflecting the influence of both ENSO and volcanic cooling. A clear volcanic signal is therefore found for the four experiments only for the Mt. Tambora eruption, while for the 1809 eruption, *High* and *Best* show a distinct cooling in 1809 and *nNHP* in 1811. In contrast to the East Pacific, variability in the West Pacific is rather small (Fig. 7b). In all four experiments a clear volcanic signal is visible in the simulated ensemble mean SST anomaly after the 1809 eruption and the Mt. Tambora eruption, whereas only a weak signal appears for both eruptions in the reconstruction. Interestingly, in the West Atlantic, two distinct positive SST anomalies appear in the reconstructions in the aftermath of the unidentified 1809 and the Mt. Tambora eruption, while the MPI-ESM simulations shows cooling (Fig. 7c). Reasons for the



**Figure 7.** Coral-SST comparison. Comparison of reconstructed tropical annual mean SST (Tierney et al., 2015) with the MPI-ESM experiments over: a) eastern Pacific ( $10^{\circ}\text{N}$ – $10^{\circ}\text{S}$ ,  $175^{\circ}\text{E}$ – $85^{\circ}\text{W}$ ), b) western Pacific ( $25^{\circ}\text{N}$ – $25^{\circ}\text{S}$ ,  $110$ – $155^{\circ}\text{E}$ ), c) western Atlantic ( $15$ – $30^{\circ}\text{N}$ ,  $60$ – $90^{\circ}\text{W}$ ) and d) the Indian ( $20^{\circ}\text{N}$ – $15^{\circ}\text{S}$ ,  $40$ – $100^{\circ}\text{E}$ ) oceans. Black solid line: SST reconstruction; colored lines: ensemble means of the model simulations. Anomalies are taken with respect to the years 1800 to 1808. The squares on the bottom of each panel indicate years where the observation lies outside the simulated ensemble range (color code as for the ensemble mean).

anticorrelated behaviour are not obvious per se and may be related to changes in either ocean circulation or other climate factors than SST that influence the coral record, such as salinity and precipitation. In the reconstruction, the Indian Ocean is the only region that displays a peak cold anomaly after the Mt. Tambora eruption, but the magnitude of this cooling is comparable to an apparent cooling in 1807. A clear reference to the Mt. Tambora eruption is therefore difficult to establish. No large cooling is found in the coral data after 1809 over the Indian Ocean (Fig. 7d).

Instrumental measurements from the tropical region are sparse and no continuous temperature record covering the early 19th century exists. Fig. 8 shows a comparison between the model simulations with ship-based surface air temperature measurements from the tropical Atlantic and Indian Oceans. For each ocean basin, the model output is sampled at the locations and times of the ship measurements. For the Indian Ocean, observed temperature anomalies after 1809 are within the model ensemble spread of all the model ensembles. The model response in the Indian Ocean is quite variable for the *Low* forcing experiment, with some members showing no apparent cooling, and others with cooling of up to  $0.9^{\circ}\text{C}$ . Overall, observed Indian Ocean temperature anomalies are on the lower edge of the *Low* ensemble. For the *Best*, *High* and *nNHP* experiments, the simulated cooling over the Indian Ocean is more consistent across individual simulations, with the ensemble spread enveloping the ob-




**Figure 8.** Annual mean surface-air temperature anomalies from shipbourne measurements of the English East India Company (EEIC) (Brohan et al., 2012) over the tropical Indian and Atlantic oceans (black line) compared to similarly sampled model simulations from the *Low*, *Best*, *High* and *nNHP* forcing ensembles as labeled. Anomalies are taken with respect to the years 1800 to 1808.

415 served temperature time series. While *Low* forcing is not inconsistent with the observed Indian ocean temperatures, *Best*, *High* or *nNHP* appear more likely scenarios. In the tropical Atlantic, the observed cooling after the 1809 eruption is slightly stronger than for Mt. Tambora, and slightly stronger than that in the Indian Ocean. The maximum observed cooling in 1809 is roughly within the spread of all the model ensembles. However, while observed tropical Atlantic temperature anomalies are largest in 1809, the simulated cooling usually peaks in 1810. In 1810, the observed cooling is less than simulated in the *Best* ensemble, 420 and smaller than all but one of the individual simulations in the *High* ensemble. Looking at the years after the Mt. Tambora eruption, simulations and observations agree relatively well in the Indian Ocean, while in the Atlantic, the model simulations overestimate the post-Tambora cooling. Since satellite observations of the aerosol cloud from the 1991 eruption of Pinatubo show that aerosol quickly spreads uniformly across the tropics, it is unlikely that aerosol forcing from the 1809 eruption would be significantly different between the tropical Indian and Atlantic ocean basins. Therefore, differences in temperature response 425 in the model between the two regions seems more likely to be related to model sensitivity, which might particularly be linked to differences in ocean circulation and/or mixed layer depth.


#### 4.2.2 Northern hemisphere extratropics

Fig. 9 shows a comparison between the model experiments with four NH summer land near surface temperature reconstructions from tree-ring records, including the nested N-TREND (N) (Wilson et al., 2016) and the spatial N-TREND (S) reconstruction 430 (Anchukaitis et al., 2017). To ensure comparability between the reconstructions and the model results, the data are expressed as anomalies w.r.t. 1800-1808. The *High* and *Best* experiments show significantly larger cooling than the reconstructions, and are outside the 95% confidence interval of the N-TREND (N) reconstruction. Simulated SAT anomalies in *nNHP* are generally smaller than the reconstructions between 1809 and 1815. Best agreement between the ESM simulations and the data after the



**Figure 9.** Comparison of NH extratropical summer land temperatures. a) Comparison of simulated NH extratropical (40-75°N) summer land temperature anomalies (seasonal and spatial averaged) with four different NH tree-ring based temperature reconstructions (Wilson et al., 2016 (N-TREND (N)), Anchukaitis et al., 2017 ((N-TREND (S)), Guillet et al., 2017 (NVOLC), Schneider et al., 2015 (SCH15)). Anomalies are taken with respect to the years 1800-1808. The black lines represent the tree-ring records, the colored ones the ensemble mean of the four MPI-ESM model experiments. The shaded grey area indicates the 2 sigma uncertainty range for N-TREND (N). b) Comparison for the reconstructed and simulated anomalies for the year 1810. Uncertainty ranges for all reconstructions are based on the 2-sigma of the N-TREND (N) reconstruction. Simulated anomalies are shown as individual realizations.

1809 eruption is found for *Low*. In NH summer 1810 and 1811, *Low* matches the reconstructed temperature anomalies from the  
 435 NVOLC (Guillet et al., 2017) and N-TREND (S) records quite well. Interestingly, the devil really is in the detail. Despite the data richness of this period, the temporal evolution (trend) differs substantially between the different tree-ring reconstructions. In N-TREND (N) the evolution is a step-like temperature decrease with a 1st step in 1809, followed by a second one in 1812 and persistent low values until 1816. Distinct peak cooling appears in NVOLC and N-TREND (S) in NH 1810 followed by a short recovery phase in 1811 and a drop in 1812, but while summer SAT anomalies stay constant in the NVOLC reconstruction,  
 440 for N-TREND (S), they start to recover again after 1812. Schneider et al. (2015) show only a small cooling trend between 1809 and 1815. In their reconstruction, temperatures after the 1809 event did not return to their climatological mean after the initial drop, but remained low until the Mt. Tambora eruption in 1815. Compared to the reconstructions, the ESM simulations (*High*, *Best*, *Low*) show a very different temporal evolution with a relatively fast recovery after the 1809 eruption to near background conditions, followed by a second cooling peak for the Mt. Tambora eruption starting in 1816. In the MPI-ESM  
 445 model simulations no cooling peak appears in the ensemble mean for the summer of 1812 in contrast to the tree-ring records. *nNHP* is the only experiment which shows only a slight cooling trend between 1810 and 1815, appearing closer to Schneider




**Figure 10.** Spatial comparison of NH extratropical land temperatures for summer 1810. Statistical comparison of reconstructed surface summer temperatures from N-TREND (S) (Anchukaitis et al., 2017) with the ensemble distributions of the four MPI-ESM ensembles (10 members). Anomalies are for the year 1810 wrt to 1800-1808 mean. The shading shows, for each grid point, the percentile range of the ensemble simulation where the reconstructed temperature falls. Green patches indicate that the reconstruction lies in the interquartile range of the simulations, hence is in good agreement with the ensemble. Bluish patches indicate that the reconstruction lies in the higher range of the ensemble, i.e., the majority of simulations are colder than the reconstructions. Reddish patches indicate that the reconstruction lies in the lower range of the ensemble, i.e., the majority of simulations are warmer than the reconstructions. White dots indicate where the reconstruction is an outlier with respect to the distribution of the simulation ensembles, i.e., where the absolute difference between reconstruction and simulation ensemble mean is greater than three times the median absolute deviation of the simulation ensemble.

et al. (2015). Between 1813 and 1815, *nNHP* reveals similar temperature anomalies to *Best* and *High*, while *Low* shows less cooling than all other experiments which even disappears before the onset of the Tambora eruption

In Fig. 10, we analyse the spatial patterns of the percentiles of the model ensemble into which the reconstruction falls. If the reconstruction lies in the upper range of the distribution of ensemble members, the reconstructed temperature anomalies are warmer than most simulations, i.e. the majority of simulations are colder than the reconstruction. The *High* ensemble (Fig.

10a) is in many locations colder than the reconstructions, but the reconstruction from Central to Northern Europe lies mostly within the interquartile, i.e. the 25th-75th percentile range of the simulations. This behaviour results from the comparison of the variable local cooling in the individual simulations with highly heterogeneous temperature anomalies in the reconstruction (Fig. 2). Only in a few regions (Central Europe, West Russia, Alaska) are the simulated temperature anomalies much warmer and the reconstructed one below the 25th percentile. The *Best* experiment (Fig. 10b) indicates a similar behaviour as *High* albeit with more regions where the reconstruction lies within the interquartile range of the simulations, e.g., along the west coast of North America. *Low* (Fig. 10c) and *nNHP* (Fig. 10d) are the experiments with the best agreement between the simulated and the reconstructed surface temperature anomalies. *nNHP* is the experiment where, compared to the other model experiments, the reconstruction is in most locations within the interquartile range of the simulations, and which has the least number of locations for which the reconstruction is considered an outlier compared to the simulation ensemble. Overall, the N-TREND (S) reconstruction is colder over Eastern Europe and West Russia compared to simulated surface temperature anomaly distribution in all four experiments, and warmer over the Eastern part of North America.




**Figure 11.** Scatterplots of Root Mean Square Error (RMSE) versus spatial correlation between simulated NH summer temperature and the N-TREND (S) reconstruction (Anchukaitis et al., 2017) for summer 1810 and different regions: a) whole Northern Hemisphere, b) North America (180-60°W), c) Europe (60°W-60°E), d) Asia (60-180°E). Individual model realizations are indicated by squares, the ensemble mean with a full dot; small grey dots are for 1000 random samples from the control period (1800 to 1808). Analysis is restricted to grid-points where proxy data are available (number of data used for each region reported in the respective panel).

Another method to compare reconstructed and simulated spatial patterns of temperatures anomalies in summer 1810 is shown in Fig. 11, which illustrates the spatial correlation and root-mean-square-error (RMSE) between the reconstruction and individual ensemble members of each experiment for the whole Northern Hemisphere and three equal-area sections of

it. Similar metrics are calculated and shown for individual summers of an unforced control run, to illustrate the potential for the model to produce similar spatial structures as a result of natural variability. A perfect agreement between simulated and reconstructed data corresponds to spatial correlation of 1 and RMSE of 0, hence the best simulated representations of the reconstructed anomalies are found close to the top-left corner of each panel. A perfect correlation could result from a simulation which had a bias compared to the reconstruction, while RMSE results from absolute differences due to spatial differences and biases. Over the entire NH (Fig. 11a) the scatter of all ensembles largely overlap with each other and with the control run, reflecting the effects of the relatively large internal climate variability. The *High* and *Best* ensembles yield ensemble-mean metrics that are at the edge of the control run, with some realizations of the former ensemble being outside the spread of the control run for the NH and Asian regions. Both ensemble simulations are colder than the reconstruction (Fig. 10) for any year of the control run. The *Low* and *nNHP* ensemble compare best with the N-TREND (S) reconstruction according to this analysis, especially as they yield the smallest RMSE values both regarding individual realizations and the ensemble-mean. The best correlations for the NH, in both the control and the forced experiments, are only 0.5 which reveals that the model does not produce such a spatial pattern as we see in the reconstruction.

Model performance in terms of spatial correlation is especially interesting over North America (Fig. 11b), where a cold-warm zonal dipole is a major characteristic of the N-TREND (S) reconstruction but where also the proxy data quality is not optimal (Anchukaitis et al., 2017). The *Low*, *nNHP* and *Best* ensembles, as well as the control run, include realizations that yield high spatial correlations over North America. This suggests that such a continental anomaly is consistent with the variability produced naturally by the model. The spread of correlation values over North America is wide, with many realizations showing strongly negative correlation with the reconstruction, and correlations of the ensemble means are small. There is therefore no evidence from the model that the spatial pattern of temperature anomalies over North America is a specific response to the volcanic aerosol forcing. The results for Europe (Fig. 11c) are similar to those of North America, with a handful of simulations from the *Low* and *nNHP* experiments showing highest correlation with the reconstruction, but no clear improvement of the forced simulations in general compared to the control run. The range of correlations for the ensembles and the control run is smaller over Asia (Fig. 11d) than in the other considered regions, i.e., no realizations show strong positive (or negative) correlations with the reconstruction, as ensembles, except *nNHP*, yield too cold anomalies especially over Eastern Siberia that contrast the weak anomalies reconstructed there (Fig. 10). This is most likely related to a substantial data quality issue as the tree-ring data, especially for central Asia, is solely based on TRW data (Wilson et al., 2016). However, we can also not rule out the possibility of a model biases as the climatological mean state of near-surface air temperature in the MPI-ESM-LR over Central Asia deviates from ERA-Interim about a few degrees (Müller et al., 2018).

A comparison of simulated ensemble mean and observed near-surface air temperature anomalies over different European regions and New England is shown for NH summer and winter in Figs. 12,13, respectively, and with the individual ensemble members, which show a comparable variability to the observations in the supplementary material (Fig. S7-S14). Note that these figures neither account for the error in the observations nor for the error (difference between a station and an areal average). For NH summer, the station data reflect the findings from the spatial comparison with the tree-ring records (Fig. 10), i.e., most European station records indicate some cooling in summer 1810 (Fig. 12a-e), while the New England data shows no




**Figure 12.** Seasonal mean near-surface NH summer temperature anomalies ( $^{\circ}\text{C}$ ) with station data averaged over different regions: a) Central Europe ( $46.1\text{-}52.5^{\circ}\text{N}$ ,  $6\text{-}17.8^{\circ}\text{E}$ ), b) Eastern Europe ( $47\text{-}57^{\circ}\text{N}$ ,  $18\text{-}32^{\circ}\text{E}$ ), c) Northern Europe ( $55\text{-}66^{\circ}\text{N}$ ,  $10\text{-}31^{\circ}\text{E}$ ), d) Southern Europe ( $38\text{-}46^{\circ}\text{N}$ ,  $7\text{-}13.5^{\circ}\text{E}$ ), e) Western Europe ( $48.5\text{-}56^{\circ}\text{N}$ ,  $6^{\circ}\text{W}\text{-}6^{\circ}\text{E}$ ) and f) New England ( $41\text{-}44^{\circ}\text{N}$ ,  $73\text{-}69^{\circ}\text{W}$ ). The black line represents measurements, the colored lines the ensemble mean of the model simulations. The squares on the bottom of each panel indicate years where the measurement lies outside the simulation ensemble range (color code as for the ensemble mean). Anomalies are taken with respect to the years 1806-1820.

evidence of cooling in this year (Fig. 12f). Further, all of the stations show cooling in either 1812 or 1813, with many showing consecutive cold summers until the Tambora eruption of 1815. In boreal summer 1810, the *nNHP* and *Low* simulations and station data are inconsistent over Northern Europe (Fig. 13c), where the observed cooling is larger than in the simulations whereas for *High* and *Best* the station data lie within the ensemble range. Surface air temperature over Western and Central Europe seems to be not much affected by the 1809 eruption, neither in all model experiments nor in the station data (Fig. 12a,e). The simulated post volcanic cooling in summer 1810 is consistent with the station data over Southern and Eastern Europe in the

505





**Figure 13.** Same as Figure 12 but for seasonal mean near-surface NH winter temperature anomalies ( $^{\circ}\text{C}$ ). The year denotes the February month.

model experiments with symmetric volcanic forcing (*Best*, *High*, *Low*) but *nNHP* shows slightly warm anomalies for summer 1810 (Fig. 12d,b). In contrast, *nNHP* is the only experiment which shows a similar trend to the New England stations, whereas  
 510 the other experiments show stronger cooling there in 1810. Observed cooling after the Mt. Tambora eruption is matched quite well by the model, except for Central Europe and Western Europe where the observed anomalies in 1816 are larger. Excellent agreement between model simulations and station data is found for the “year without summer” for New England. An interesting feature is the observed warming peak of  $2^{\circ}\text{C}$  in summer 1811 over Central Europe, which is also found in one realization of the *Best* experiment (Fig. S7) suggesting the influence of internal variability. For NH winter, both model and station data show  
 515 higher variability than in NH summer (Fig. 13). Simulated and observed NH winter temperature anomalies agree quite well in the first three winters after the 1809 eruption. The only exception is New England (Fig. 13f) where, similar to NH summer (Fig. 12f), less agreement is found between the station data and the four experiments. The strong cooling signal of more than

-2 °C which is found at Northern, Western and Central European stations in winter 1813 /1814 is not reproduced by the model simulations. All experiments except *Low* produce positive post-volcanic winter temperature anomalies over Northern Europe  
520 with a warming signal in the first winter (1809/1810) after the 1809 eruption consistent with the station data.

## 5 Discussion

Significant surface cooling is found after the unidentified 1809 eruption in instrumental observations, proxy records and simulations, but the cooling is strongly spatially heterogeneous and is variable across the different data sources, especially across the NH extratropics.

525 Observed and simulated cooling in the tropics is considerable, with anomalies likely exceeding 0.5 °C. In general, the MPI-ESM simulations show stronger cooling over the tropical oceans than the reconstructed and observed temperatures (Fig. 6). Over the tropical Indian ocean, the simulated SST anomalies between 1809 and 1820 for the *Low*, *Best* and *nNHP* experiments agree quite well with the marine ship measurements (Fig. 8). Furthermore, the ensemble mean cooling in *Low* is close to the reconstruction of D'Arrigo et al. (2006) over the tropical IndoPacific in 1809 although the reconstruction does not pick up the  
530 1816 cooling expressed by all experiments (Fig. 6c). An even smaller SST variability range is also found for the Indian and the West Pacific SST reconstructions when compared to the models (Fig. 7). Uncertainties related to the proxy measurements likely explain some of the mismatch, as they depend on several factors such as mixed sea surface temperature and salinity influences on the  $\delta^{18}O$  measurements in coral archives as well as seasonal changes in the growth rate and other biological factors. The limited number of proxy records, which are not equally distributed in time and space, is also undoubtedly a  
535 relevant factor. Focusing on the tropical region, the comparison between simulations suggests that the most probable forcing would be somewhere between *Low*, *nNHP* and *Best*.

Simulated ensemble mean NH extratropical summer temperature anomalies are mostly for all forcing scenarios within the reconstructed range but the agreement is strongly dependent on the choice of the year, spatial coverage and the individual forcing scenarios and reconstruction used. Ensemble mean temperature differences between the different MPI-ESM experi-  
540 ments or the individual reconstructions could be in the order of the forced signal (Fig. 9). The NH mean summer temperature response to the 1809 eruption is smaller in tree-ring reconstructions than in the simulations (except for *nNHP*), consistent with previous comparisons of tree-ring reconstructions with climate models. Schneider et al. (2017), for example, compared tree-ring reconstructions with PMIP3 last millennium simulations from different models including the MPI-ESM. They also found that on average most model simulations feature a more pronounced volcanic signal than proxy reconstructions. The spread of  
545 simulated NH temperature arising from uncertainties in the volcanic forcing estimate (VSSI) for the 1809 eruption is generally comparable to the spread in the NH tree ring-based temperature reconstructions. It is therefore not possible to use one proxy result to strongly constrain the other, as we might if one distribution was much smaller than the other or if the overlap between the two was small. On the other hand, the disagreement between observed and simulated tropical temperatures in the *High* ensemble would suggest that VSSI estimates on the upper end of our estimated range are not likely and qualitatively, we might  
550 estimate that the most likely VSSI emissions for 1809 are between around 12 and 19 Tg S.

Instrumental records in Europe fit well with simulated temperatures using all forcing scenarios for both winter and summer seasons (Figs. 12,13). In boreal winter, observations from most stations (except New England and Eastern Europe) lie within the ensemble range of all four experiments within the 1st two post volcanic winters. Internal variability in NH winter is high and can overwhelm volcanic forcing uncertainties as pointed out by Zanchettin et al. (2019) for the 1st winter after the Mt. Tambora eruption. The spread of continental winter responses generated by the initial conditions for the unidentified 1809 eruption is expected to be larger than for the Mt. Tambora eruption, due to the lower signal to noise ratio. Ultimately, comparison with early instrumental data must be made cautiously. Although regular meteorological measurements have been performed across Europe since the 17th and 18th centuries (Brönnimann et al., 2019b), continuous temperature measurements in the 1810s are rare due to the Napoleonic wars and there is a significant lack of regular temperature time series from other parts of the world for this period.

Spatial correlations over the NH between the simulations and the NTREND (S) temperature reconstruction for NH summer 1810 are overall low. None of the anomalous patterns produced by the MPI-ESM, either under unperturbed conditions or volcanic-forcing scenarios, reproduces the amplitude and structure of the reconstructed hemispheric temperature anomaly pattern particularly well in summer 1810 suggesting that the mismatch is most likely not attributable to the volcanic forcing. This possibly stems from the heterogeneous reconstructed anomalies that lack a clear spatial structure and provide a “noisy” target to the simulations. The quite unusual temperature pattern in N-TREND (S), with cold temperatures over Europe and Alaska and warm temperatures over Eastern North America and Northern Siberia could be a potential cause for this. This heterogeneous spatial pattern in the tree-ring data is also reflected in the dampened cooling seen in the NH mean composite reconstructions (Fig. 9). Station observations seem to support this anomaly pattern by showing cooling only over Northern and Eastern Europe. Surface temperature reconstructions from an almost independent MXD network (Briffa et al., 2002b) indicate also positive temperature anomalies over East Canada, north of 55°N and East of 95°W but also regional cooling. Briffa et al. used MXD data only, which capture the short-time cooling after volcanic eruptions more reliably than RW based reconstructions which are influenced by biological persistence (e.g., Anchukaitis et al., 2012; Esper et al., 2015; Lücke et al., 2019). In N-TREND (S) a combination of MXD, RW and BI records are used and the prediction skill is spatially very variable, particularly over North America (Anchukaitis et al., 2017). The distinct East-West dipole pattern in the reconstruction could therefore be partly caused by a data quality issue as is certainly the case for Central Asia which is driven predominantly by tree ring-width data. One possibility is that 1810 was a dynamically unusual but not unlikely year (Fig. 11) in terms of NH atmospheric circulation that led to warming over Eastern North America which is not reflected in the simulated ensemble mean. While a few individual simulations produced a dipole temperature anomaly pattern over North America similar to that of the NTREND (S) reconstruction when forced with volcanic aerosol, many other simulations did not, and the ensemble mean correlation was close to zero for all forced ensembles. We can only conclude that in these simulations, the volcanic aerosol forcing did not lead to or increase the likelihood of the dipole North American pattern.

Overall, our model-data intercomparison suggests that the mid-to-low range of the sulfate estimates for the unidentified 1809 eruption as the most realistic. However, the range of uncertainty is large, as observed and reconstructed data are sparse and cover a wide post-volcanic temperature anomaly range. Uncertainties in volcanic forcing are often discussed as one of the main

causes for the discrepancy between simulated and reconstructed temperature response due to biases in magnitude and structure (e.g., Timmreck et al., 2009; Yang et al., 2019; Zhu et al., 2020). As discussed above, the spread of the different forcing ensembles is in the same range as the different tree ring reconstructions, hence it is difficult to reduce the forcing estimates. Nevertheless, our work does suggest that the response of large-scale NH climate indexes (Fig. S1) is sensitive not only to the magnitude of the volcanic forcing, but also its meridional structure, with, e.g., significant differences in the summer NPI response for the *Best* and *nNHP* ensembles. This would support the idea that the structure of forcing is important in influencing circulation changes (e.g., Toohey et al., 2019). An asymmetric volcanic forcing scenario with less or no NH extratropical forcing, as in *nNHP*, provides one way in which tropical temperatures could respond strongly to a tropical eruption, while the NH temperature response could be more muted and prolonged, as suggested by some tree-ring reconstructions for 1809. This scenario is however at odds with the ice-core records, which show comparable deposition to Antarctica and Greenland suggesting a global aerosol spread. Without a viable explanation for a mismatch between ice-core records and hemispheric radiative forcing, extreme hemispheric asymmetry (as in the *nNHP* scenario) must be concluded to be an unlikely reason for the muted NH mean temperature response to the 1809 eruption.

A factor which might have an influence on the mismatch between model results and proxy data is the season of the eruption, which has an impact on the climate response to volcanic eruptions (e.g. Stevenson et al., 2017; Predybaylo et al., 2020). We have not discussed this uncertainty for the 1809 eruption here as we chose January 1809 as the starting date following most studies that suggest that the eruption probably happened in boreal winter months across 1808 and 1809. Nevertheless the timing of the eruption remains uncertain. Chenoweth (2001) for example dated the 1809 eruption back to March-June 1808 based on a sudden cooling in Malaysian land surface temperature data and a peak cooling of marine air temperature in 1809. To take into account the eruption's season may be important to fully characterize the range of climate response compatible with the 1809 eruption and should be addressed in further studies.

One final caveat to our simulation-reconstruction comparison stems from the idealized nature of our experiments: our simulations are branched from a multi-centennial preindustrial control run and use only volcanic aerosol forcing; they are not transient simulations with realistic full forcing (including also variations in solar and land-use forcing, etc.). Neglecting the reduced solar irradiance in the early 19th century could, for example, be one of the factors why our model simulations do not show the slight cooling trend in summer surface temperature between 1810 and 1815 indicated by the NH tree-ring data. While this more idealized experimental set up has the advantage of better isolating the influence of volcanic forcing strength and structure alone, it complicates the comparison with observations and proxy reconstructions. Early 19th century ensemble simulations starting from a climate state with a more realistic past natural forcing, as planned for the CMIP6 VolMIP/PMIP “volc-cluster-mill” experiments (Zanchettin et al., 2016; Jungclaus et al., 2017), will allow for a more direct comparison with early instrumental data and paleoclimate reconstructions. Furthermore, they will also help to disentangle the processes which are specific to the MPI-ESM model and which need discussion in a broader context.

## 6 Conclusions

We used the MPI Earth system model to study the climatic impacts of the unidentified 1809 eruption, the 6th largest volcanic eruption of the last millennium. Our aim was to address the question of whether or not the short-term climate response to the 1809 eruption noted in observations and proxy-based climate reconstructions is compatible with a range of volcanic forcing estimates constructed based on estimated uncertainties in volcanic stratospheric injection from ice-core records. We demonstrated that the model simulations of the climate impacts of the 1809 eruption show generally good agreement with many of the available large-scale temperature reconstructions and early instrumental records. Assuming the model climate sensitivity is correct, this result implies that the eVolv2k VSSI estimate for the 1809 eruption is accurate within its reported uncertainty. Model-data intercomparison of tropical temperature anomalies, where the impact of direct radiative forcing is highlighted, suggests that the most likely forcing is somewhere between the *Low* and the *Best* scenarios with an estimated sulfur emission between 12 -19 Tg S. However, the limited data availability for this time period for the tropical region poses a caveat to this conclusion. Simulated NH-average summer temperature anomalies are for all forcing scenarios within the observational and reconstructed range. The long-lasting cooling trend detected in observed and reconstructed NH-average summer temperatures between 1810 and 1815, i.e., until the Mt. Tambora eruption, is not detected in the model simulations except, partly, in *nNHP*. The reason(s) for such a cooling trend remains thus unclear and needs to be further explored. The spatial correlation between simulated and reconstructed near-surface air temperatures over the NH in summer 1810 is low. Tree-ring based gridded reconstructions of NH extratropical summer SAT show a highly heterogeneous anomaly pattern in 1810 with extensive regions of cold and warm anomalies, with an east-west warm-cold dipole over North America, and variable temperature anomalies over Asia, which leads to a dampened NH-average cooling for this year (Fig. 10). The pattern over North America is consistent with the limited available station observations from New England, an almost independent MXD based summer temperature reconstruction (Briffa et al., 2002b), and is reproduced by some forced simulations and years from the control run, suggesting it is a plausible result of natural variability. In contrast, the reconstructed temperature pattern over Asia is not produced by any model simulation, forced or control. Possible explanations for this model-data disagreement include deficiencies and uncertainties regarding both tools. In particular, the dendrochronological network remains weakly constrained over the central Asian region, especially with respect to available MXD or density-like parameters, and instrumental observations are sparse, especially prior to the 1950s making calibration and validation difficult (Cook et al., 2013). Anchukaitis et al. (2017) clearly detailed poor reconstruction validation for many regions across North America as well as Northeast and central Asia. These regions must be targeted in ongoing proxy network development to update and develop new MXD or density-like parameter data-sets which are proven to be superior proxies of volcanically forced summer cooling (e.g., Anchukaitis et al., 2012; D'Arrigo et al., 2013; Esper et al., 2015; Lücke et al., 2019). Concerning the simulations, beyond model deficiencies that may bias the response, our study demonstrates that choices regarding both volcanic forcing strength and spatial structure can similarly affect reconstruction-simulation comparisons. Specifically, it was shown that for the muted response to the 1809 eruption in the NH extratropics, agreement between reconstructions and simulations improves by weakening the magnitude of the eruption and, alternatively, by preventing the volcanic aerosol to spread over the Northern Hemisphere. The forcing estimate of 1809, which

is based on ice core data, can only be improved by modelling experiments to narrow down the uncertainty range. This will be facilitated if further information on location and eruption season is identified. Recent advances in ice-core analysis can help to identify the source volcano of large historic eruptions by combining proximal tephra fallout with the geochemical analysis of ash shards in ice-cores as done for example for the 1257 Samalas eruption (Lavigne et al., 2013), the Changbaishan eruption in 946 (Oppenheimer et al., 2017), the 431 CE Tierra Blanca Joven eruption of Ilopango (Smith et al., 2020), or the 1477 CE eruption of the Icelandic Veiðivötn–Bárðarbunga volcanic system ((Abbott et al., 2021, this issue)

Finally, it is clear that improving temperature reconstruction accuracy by adding additional tree-ring density records in poorly replicated regions, including also the southern hemisphere, is needed. In summary, analysis of multiple climate records shows that the eruption of 1809 was certainly a “cold case” in the sense that it produced large-scale cooling, and that cooling is consistent with the range of sulfur emission and radiative forcing estimates deduced from ice-core records. In terms of attributing the eruption to a source volcano, or explaining the spatially heterogeneous and temporally delayed cooling in the NH suggested by tree ring networks, the investigation remains open.

## 1 Appendix Station data

**Table A1.** Temperature observations from land stations used in this study. n = number of available value in the 1806-1820 period. Source: Dove = digitized from the collection of Friedrich Wilhelm Dove (Dove, 1838, 1839, 1842,1845), NOAA = digitized from images held at the NOAA archive, DWD = German Weather Service, GHCN = Global Historical Climatology Network (Lawrimore et al., 2011), ISTI = International Surface Temperature Initiative (Rennie et al., 2014).

Source	Station	Start year	End year	lat	lon	N
Dove	London	1787	1838	51.51	-0.13	180
Dove	Arnhem	1790	1818	51.99	5.90	156
Dove	Penzance	1807	1827	50.14	-5.49	168
Dove	Dumfernline	1805	1824	56.08	-3.43	180
Dove	Carlisle	1801	1824	52.13	-0.50	180
NOAA	Brunswick	1807	1820	43.92	-69.94	142
DWD	Berlin	1719	2018	52.45	13.30	180
DWD	Hohenpeissenberg	1781	2018	47.80	11.01	156
DWD	München	1801	1953	48.14	11.60	180
DWD	Stuttgart	1792	1984	48.77	9.18	180
DWD	Trier	1788	2018	49.73	6.61	132
GHCN	Mikolai	1808	1875	47.03	31.95	96
GHCN	New Haven	1781	1970	41.30	-72.90	180
GHCN	Boston	1743	2010	42.37	-71.03	120
GHCN	Kremsmuenster	1767	2010	48.05	14.13	180
GHCN	Innsbruck	1777	1991	47.30	11.40	180

Source	Station	Start year	End year	lat	lon	N
GHCN	Praha	1771	1981	50.10	14.25	180
GHCN	Leobschutz	1805	1849	50.20	17.80	180
GHCN	Kobenhavn	1768	1980	55.68	12.55	180
GHCN	Torneo	1801	1832	66.40	23.80	180
GHCN	Woro	1800	1824	63.20	22.00	180
GHCN	Montdidier	1784	1869	49.70	2.60	180
GHCN	Paris	1757	1980	48.80	2.50	180
GHCN	Chalons	1806	1848	48.90	4.40	180
GHCN	Strasbourg	1801	1991	48.55	7.63	180
GHCN	Nice	1806	1991	43.65	7.20	180
GHCN	Karlsruhe	1779	1930	49.03	8.37	180
GHCN	Regensburg	1773	2010	49.05	12.10	180
GHCN	Budapest	1780	1991	47.52	19.03	180
GHCN	Udine	1803	1991	46.00	13.10	180
GHCN	Torino Casell	1753	1981	45.22	7.65	180
GHCN	Milano	1763	1987	45.43	9.28	180
GHCN	Verona	1788	1991	45.38	10.87	144
GHCN	Padova	1780	1827	45.40	11.85	167
GHCN	Bologna	1808	1981	44.53	11.30	156
GHCN	Palermo	1791	1868	38.10	13.40	179
GHCN	Riga	1795	1989	56.97	24.05	102
GHCN	Vilnius	1777	1981	54.63	25.10	180
GHCN	Trondheim	1761	1981	63.40	10.50	180
GHCN	Gdansk	1807	1984	54.40	18.60	168
GHCN	Warszawa	1779	1991	52.17	20.97	180
GHCN	Wroclaw	1792	1991	51.10	16.88	180
GHCN	St.Petersburg	1743	1991	59.97	30.30	180
GHCN	Stockholm	1756	1980	59.33	18.05	180
GHCN	Geneve	1753	1991	46.25	6.13	180
GHCN	Gordon Castle	1781	1975	57.60	-3.10	180
GHCN	Edinburgh	1764	1960	55.90	-3.20	180
GHCN	Manchester	1794	1991	53.35	-2.27	180
GHCN	Greenwich	1763	1969	51.50	0.00	180
ISTI	Torino	1760	2009	45.07	7.67	180
ISTI	Basel	1760	2010	47.54	7.58	180
ISTI	Bern	1777	2010	46.99	7.46	180
KNMI	DeBilt	1706	2018	52.10	5.19	180



**Table A2.** Overview of the different stations which are used in the compilation of the regional and seasonal station mean

Region	Latitude/ longitude	Number of stations	Stations
Central Europe	46.1-52.5°N,6-17.8°E	17	Geneve, Trier, Bern, Basel, Strasbourg, Karlsruhe, Stuttgart, Hohenpeißenberg, Innsbruck, München, Regensburg, Berlin, Kremsmuenster, Praha, Wroclaw, Leobschutz, Kobenhavn
Eastern Europe	47°N-57°N, 18-32°E	6	Gdansk, Budapest, Warszawa, Riga, Vilnius, Mikolai
Northern Europe	55-66°N, 10 -31°E	5	Trondheim, Stockholm, Woro, Torneo, St. Petersburg
Southern Europe	38-46°N, 7- 13.5°E	9	Nice, Torino-Casell, Torino, Milano, Verona, Bologna, Padova, Palermo, Udine
Western Europe	48.-56°N, 6 °W-6°O	13	Penzance, Dunfermline, Edinburgh, Gordon Castle, Carlisle, Manchester, London, Greenwich, Paris, Montdidier, Chalons, De Bilt, Arnhem
New England	41-44°N, 73-69°W	3	New Haven, Boston, Brunswick

665 *Code and data availability.* Primary data and scripts used in the analysis and other supplementary information that may be useful in reproducing the author's work are archived by the Max Planck Institute for Meteorology and can be obtained by contacting [publications@mpimet.mpg.de](mailto:publications@mpimet.mpg.de). Model results will be available under [cera-www.dkrz.de](http://cera-www.dkrz.de) soon. Proxy data can be obtained from NOAA/World Data Service for Paleoclimatology archives <https://www.ncdc.noaa.gov/paleo-search>.

*Author contributions.* CT, MT and DZ designed the study, analysed the results and wrote the paper. CT performed the simulations and coordinated the writing process. SB and RW provided data and contributed to the discussion and the writing process. EL compiled a new set of the station data.

*Competing interests.* The authors declare that they have no conflict of interest

*Acknowledgements.* The authors thank Stephan Lorenz who gave valuable comments on an earlier version of the paper. Computations were performed on the computer of the Deutsches Klima Rechenzentrum (DKRZ). The research has been supported by the Deutsche Forschungsgemeinschaft Research Unit VolImpact (FOR2820, grant no 398006378) within the project VolClim (TI 344/2-1, CT). SB acknowledges funding from the Swiss National Science Foundation project WeaR (188701) and the European Commission through H2020 (ERC Grant PALAEO-RA 787574). This work benefited from the participation of the authors in the Volcanic Impacts on Climate and Society (VICS) working group of the Past Global Changes (PAGES) project.

## References

- 680 Abbott, P. M., Plunkett, G., Corona, C., Chellman, N. J., McConnell, J. R., Pilcher, J. R., Stoffel, M., and Sigl, M.: Cryptotephra from the Icelandic Veiðivötn 1477 CE eruption in a Greenland ice core: confirming the dating of volcanic events in the 1450s CE and assessing the eruption's climatic impact, *Climate of the Past*, 17, 565–585, 2021.
- Anchukaitis, K. J., Breitenmoser, P., Briffa, K. R., Buchwal, A., Büntgen, U., Cook, E. R., D'Arrigo, R. D., Esper, J., Evans, M. N., Frank, D., et al.: Tree rings and volcanic cooling, *Nature Geoscience*, 5, 836–837, 2012.
- 685 Anchukaitis, K. J., Wilson, R., Briffa, K. R., Büntgen, U., Cook, E. R., D'Arrigo, R., Davi, N., Esper, J., Frank, D., Gunnarson, B. E., et al.: Last millennium Northern Hemisphere summer temperatures from tree rings: Part II, spatially resolved reconstructions, *Quaternary Science Reviews*, 163, 1–22, 2017.
- Anet, J., Muthers, S., Rozanov, E., Raible, C., Stenke, A., Shapiro, A., Brönnimann, S., Arfeuille, F. X., Brugnara, Y., Beer, J., et al.: Impact of solar versus volcanic activity variations on tropospheric temperatures and precipitation during the Dalton Minimum, *Climate of the*
- 690 *Past*, 10, 921–938, 2014.
- Briffa, K. R., Jones, P., and Schweingruber, F.: Tree-ring density reconstructions of summer temperature patterns across western North America since 1600, *Journal of Climate*, 5, 735–754, 1992.
- Briffa, K. R., Jones, P., and Schweingruber, F.: Summer temperatures across northern North America: Regional reconstructions from 1760 using tree-ring densities, *Journal of Geophysical Research: Atmospheres*, 99, 25 835–25 844, 1994.
- 695 Briffa, K. R., Schweingruber, F. H., Jones, P., Osborn, T., Shiyatov, S., and Vaganov, E.: Reduced sensitivity of recent tree-growth to temperature at high northern latitudes, *Nature*, 391, 678–682, 1998.
- Briffa, K. R., Osborn, T. J., Schweingruber, F. H., Jones, P. D., Shiyatov, S. G., and Vaganov, E. A.: Tree-ring width and density data around the Northern Hemisphere: Part 2, spatio-temporal variability and associated climate patterns, *The Holocene*, 12, 759–789, 2002b.
- Brohan, P., ALLAN, R., FREEMAN, J. E., Wheeler, D., WILKINSON, C., and WILLIAMSON, F.: Constraining the temperature history of
- 700 the past millennium using early instrumental observations, *Climate of the Past*, 8, 1551, 2012.
- Brönnimann, S. and Krämer, D.: Tambora and the "Year Without a Summer" of 1816. A Perspective on Earth and Human Systems Science, vol. 90, *Geographica Bernensia*, 2016.
- Brönnimann, S., Franke, J., Nussbaumer, S. U., Zumbühl, H. J., Steiner, D., Trachsel, M., Hegerl, G. C., Schurer, A., Worni, M., Malik, A., et al.: Last phase of the Little Ice Age forced by volcanic eruptions, *Nature geoscience*, 12, 650–656, 2019a.
- 705 Brönnimann, S., Allan, R., Ashcroft, L., Baer, S., Barriendos, M., Brázdil, R., Brugnara, Y., Brunet, M., Brunetti, M., Chimani, B., et al.: Unlocking pre-1850 instrumental meteorological records: A global inventory, *Bulletin of the American Meteorological Society*, 100, ES389–ES413, 2019b.
- Chenoweth, M.: Two major volcanic cooling episodes derived from global marine air temperature, AD 1807–1827, *Geophysical Research Letters*, 28, 2963–2966, 2001.
- 710 Clyne, M., Lamarque, J.-F., Mills, M. J., Khodri, M., Ball, W., Bekki, S., Dhomse, S. S., Lebas, N., Mann, G., Marshall, L., Niemeier, U., Poulain, V., Robock, A., Rozanov, E., Schmidt, A., Stenke, A., Sukhodolov, T., Timmreck, C., Toohey, M., Tummon, F., Zanchettin, D., Zhu, Y., and Toon, O. B.: Model physics and chemistry causing intermodel disagreement within the VolMIP-Tambora Interactive Stratospheric Aerosol ensemble, *Atmospheric Chemistry and Physics*, 21, 3317–3343, <https://doi.org/10.5194/acp-21-3317-2021>, <https://acp.copernicus.org/articles/21/3317/2021/>, 2021.
- 715 Cole-Dai, J.: Volcanoes and climate, *Wiley Interdisciplinary Reviews: Climate Change*, 1, 824–839, 2010.

- Cole-Dai, J., Ferris, D., Lanciki, A., Savarino, J., Baroni, M., and Thiemens, M. H.: Cold decade (AD 1810–1819) caused by Tambora (1815) and another (1809) stratospheric volcanic eruption, *Geophysical Research Letters*, 36, 2009.
- Cook, E. R., Briffa, K. R., and Jones, P. D.: Spatial regression methods in dendroclimatology: a review and comparison of two techniques, *International Journal of Climatology*, 14, 379–402, 1994.
- 720 Cook, E. R., Krusic, P. J., Anchukaitis, K. J., Buckley, B. M., Nakatsuka, T., Sano, M., et al.: Tree-ring reconstructed summer temperature anomalies for temperate East Asia since 800 CE, *Climate Dynamics*, 41, 2957–2972, 2013.
- Crowley, T. J. and Unterman, M. B.: Technical details concerning development of a 1200 yr proxy index for global volcanism, *Earth System Science Data*, 5, 187, 2013.
- Dai, J., Mosley-Thompson, E., and Thompson, L. G.: Ice core evidence for an explosive tropical volcanic eruption 6 years preceding Tambora, 725 *Journal of Geophysical Research: Atmospheres*, 96, 17 361–17 366, 1991.
- DallaSanta, K., Gerber, E. P., and Toohey, M.: The circulation response to volcanic eruptions: The key roles of stratospheric warming and eddy interactions, *Journal of Climate*, 32, 1101–1120, 2019.
- D’Arrigo, R., Wilson, R., Palmer, J., Krusic, P., Curtis, A., Sakulich, J., Bijaksana, S., Zulaikah, S., Ngkoimani, L. O., and Tudhope, A.: The reconstructed Indonesian warm pool sea surface temperatures from tree rings and corals: Linkages to Asian monsoon drought and El 730 Nino–Southern Oscillation, *Paleoceanography*, 21, 2006.
- D’Arrigo, R., Wilson, R., and Anchukaitis, K. J.: Volcanic cooling signal in tree ring temperature records for the past millennium, *Journal of Geophysical Research: Atmospheres*, 118, 9000–9010, 2013.
- Davi, N. K., Jacoby, G. C., and Wiles, G. C.: Boreal temperature variability inferred from maximum latewood density and tree-ring width data, Wrangell Mountain region, Alaska, *Quaternary Research*, 60, 252–262, 2003.
- 735 Dove, H. W.: Über die nicht periodischen Änderungen der Temperaturvertheilung auf der Oberfläche der Erde. 1. Teil, Reimer, Berlin, 1838.
- Dove, H. W.: Über die nicht periodischen Änderungen der Temperaturvertheilung auf der Oberfläche der Erde. 2. Teil, Reimer, Berlin, 1839.
- Dove, H. W.: Über die nicht periodischen Änderungen der Temperaturvertheilung auf der Oberfläche der Erde. 3. Teil, Reimer, Berlin, 1842.
- Dove, H. W.: Über die nicht periodischen Änderungen der Temperaturvertheilung auf der Oberfläche der Erde. 1 Teil, Reimer, Berlin, 1845.
- D’Arrigo, R., Wilson, R., and Tudhope, A.: The impact of volcanic forcing on tropical temperatures during the past four centuries, *Nature* 740 *Geoscience*, 2, 51–56, 2009.
- Esper, J., Schneider, L., Smerdon, J. E., Schöne, B. R., and Büntgen, U.: Signals and memory in tree-ring width and density data, *Dendrochronologia*, 35, 62–70, 2015.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, 2016.
- 745 Fischer, E. M., Luterbacher, J., Zorita, E., Tett, S., Casty, C., and Wanner, H.: European climate response to tropical volcanic eruptions over the last half millennium, *Geophysical research letters*, 34, 2007.
- Franke, J., Frank, D., Raible, C. C., Esper, J., and Brönnimann, S.: Spectral biases in tree-ring climate proxies, *Nature Climate Change*, 3, 360–364, 2013.
- Gao, C., Robock, A., and Ammann, C.: Volcanic forcing of climate over the past 1500 years: An improved ice core-based index for climate 750 models, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Gennaretti, F., Boucher, E., Nicault, A., Gea-Izquierdo, G., Arseneault, D., Berninger, F., Savard, M. M., Bégin, C., and Guiot, J.: Underestimation of the Tambora effects in North American taiga ecosystems, *Environmental Research Letters*, 13, 034 017, 2018.

- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., et al.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572–597, 2013.
- 755 Guevara-Murua, A., Williams, C., Hendy, E., Rust, A., and Cashman, K.: Observations of a stratospheric aerosol veil from a tropical volcanic eruption in December 1808: is this the Unknown 1809 eruption?, *Climate of the Past*, 10, 1707–1722, 2014.
- Guillet, S., Corona, C., Stoffel, M., Khodri, M., Lavigne, F., Ortega, P., Eckert, N., Sielenou, P. D., Daux, V., Churakova, O. v., et al.: Climate response to the Samalas volcanic eruption in 1257 revealed by proxy records, *Nature geoscience*, 10, 123–128, 2017.
- 760 Hakim, G. J., Emile-Geay, J., Steig, E. J., Noone, D., Anderson, D. M., Tardif, R., Steiger, N., and Perkins, W. A.: The last millennium climate reanalysis project: Framework and first results, *Journal of Geophysical Research: Atmospheres*, 121, 6745–6764, 2016.
- Hartl-Meier, C., Büntgen, U., Smerdon, J. E., Zorita, E., Krusic, P. J., Ljungqvist, F. C., Schneider, L., and Esper, J.: Temperature covariance in tree ring reconstructions and model simulations over the past millennium, *Geophysical Research Letters*, 44, 9458–9469, 2017.
- Haywood, J. M., Jones, A., Bellouin, N., and Stephenson, D.: Asymmetric forcing from stratospheric aerosols impacts Sahelian rainfall, *Nature Climate Change*, 3, 660–665, <https://doi.org/10.1038/NCLIMATE1857>, 2013.
- 765 Hegerl, G. C., Crowley, T. J., Baum, S. K., Kim, K.-Y., and Hyde, W. T.: Detection of volcanic, solar and greenhouse gas signals in paleo-reconstructions of Northern Hemispheric temperature, *Geophysical Research Letters*, 30, 2003.
- Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, *Journal of Advances in Modeling Earth Systems*, 5, 287–315, 2013.
- 770 Jungclaus, J., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and Von Storch, J.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *Journal of Advances in Modeling Earth Systems*, 5, 422–446, 2013.
- Jungclaus, J. H., Lohmann, K., and Zanchettin, D.: Enhanced 20th century heat transfer to the Arctic simulated in context of climate variations over last millennium, *Climate of the Past*, 10, 2201–2213, 2014.
- 775 Jungclaus, J. H., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L. P., Egorova, T., Evans, M., González-Rouco, J. F., Goosse, H., Hurtt, G. C., Joos, F., Kaplan, J. O., Khodri, M., Klein Goldewijk, K., Krivova, N., LeGrande, A. N., Lorenz, S. J., Luterbacher, J., Man, W., Maycock, A. C., Meinshausen, M., Moberg, A., Muscheler, R., Nehrbass-Ahles, C., Otto-Bliesner, B. I., Phipps, S. J., Pongratz, J., Rozanov, E., Schmidt, G. A., Schmidt, H., Schmutz, W., Schurer, A., Shapiro, A. I., Sigl, M., Smerdon, J. E., Solanki, S. K., Timmreck, C., Toohey, M., Usoskin, I. G., Wagner, S., Wu, C.-J., Yeo, K. L., Zanchettin, D., Zhang, Q., and Zorita, E.: The PMIP4 contribution to CMIP6 – Part 3: The last millennium, scientific objective, and experimental design for the PMIP4 *past1000* simulations, *Geoscientific Model Development*, 10, 4005–4033, <https://doi.org/10.5194/gmd-10-4005-2017>, <https://gmd.copernicus.org/articles/10/4005/2017/>, 2017.
- 780 Kent, E. C., Rayner, N. A., Berry, D. I., Saunby, M., Moat, B. I., Kennedy, J. J., and Parker, D. E.: Global analysis of night marine air temperature and its uncertainty since 1880: The HadNMAT2 data set, *Journal of Geophysical Research: Atmospheres*, 118, 1281–1298, 2013.
- Khodri, M., Izumo, T., Vialard, J., Janicot, S., Cassou, C., Lengaigne, M., Mignot, J., Gastineau, G., Guilyardi, E., Lebas, N., et al.: Tropical explosive volcanic eruptions can trigger El Niño by cooling tropical Africa, *Nature communications*, 8, 1–13, 2017.

- Lavigne, F., Degeai, J.-P., Komorowski, J.-C., Guillet, S., Robert, V., Lahitte, P., Oppenheimer, C., Stoffel, M., Vidal, C. M., Prato, I.,  
790 et al.: Source of the great AD 1257 mystery eruption unveiled, Samalas volcano, Rinjani Volcanic Complex, Indonesia, *Proceedings of the National Academy of Sciences*, 110, 16 742–16 747, 2013.
- Lawrimore, J. H., Menne, M. J., Gleason, B. E., Williams, C. N., Wuertz, D. B., Vose, R. S., and Rennie, J.: An overview of the Global Historical Climatology Network monthly mean temperature data set, version 3, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- 795 Li, J., Xie, S.-P., Cook, E. R., Morales, M. S., Christie, D. A., Johnson, N. C., Chen, F., D'Arrigo, R., Fowler, A. M., Gou, X., et al.: El Niño modulations over the past seven centuries, *Nature Climate Change*, 3, 822–826, 2013.
- Lücke, L. J., Hegerl, G. C., Schurer, A. P., and Wilson, R.: Effects of memory biases on variability of temperature reconstructions, *Journal of Climate*, 32, 8713–8731, 2019.
- Marshall, L., Johnson, J. S., Mann, G. W., Lee, L., Dhomse, S. S., Regayre, L., Yoshioka, M., Carslaw, K. S., and Schmidt, A.: Exploring how  
800 eruption source parameters affect volcanic radiative forcing using statistical emulation, *Journal of Geophysical Research: Atmospheres*, 124, 964–985, 2019.
- Marshall, L. R., Smith, C. J., Forster, P. M., Aubry, T. J., Andrews, T., and Schmidt, A.: Large variations in volcanic aerosol forcing efficiency due to eruption source parameters and rapid adjustments, *Geophysical Research Letters*, 47, e2020GL090 241, 2020.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., et al.: Devel-  
805 opments in the MPI-M Earth System Model version 1.2 (MPI-ESM1. 2) and its response to increasing CO<sub>2</sub>, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, 2019.
- McGregor, S., Timmermann, A., and Timm, O.: A unified proxy for ENSO and PDO variability since 1650., *Climate of the Past*, 6, 2010.
- Moreno-Chamarro, E., Zanchettin, D., Lohmann, K., and Jungclauss, J. H.: An abrupt weakening of the subpolar gyre as trigger of Little Ice Age-type episodes, *Climate Dynamics*, 48, 727–744, 2017.
- 810 Müller, W. A., Jungclauss, J. H., Mauritsen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., et al.: A Higher-resolution Version of the Max Planck Institute Earth System Model (MPI-ESM1. 2-HR), *Journal of Advances in Modeling Earth Systems*, 10, 1383–1413, 2018.
- Oppenheimer, C.: Climatic, environmental and human consequences of the largest known historic eruption: Tambora volcano (Indonesia) 1815, *Progress in physical geography*, 27, 230–259, 2003.
- 815 Oppenheimer, C., Wacker, L., Xu, J., Galván, J. D., Stoffel, M., Guillet, S., Corona, C., Sigl, M., Di Cosmo, N., Hajdas, I., et al.: Multi-proxy dating the ‘Millennium Eruption’ of Changbaishan to late 946 CE, *Quaternary Science Reviews*, 158, 164–171, 2017.
- PAGES 2k Consortium, .: Consistent multi-decadal variability in global temperature reconstructions and simulations over the Common Era, *Nature geoscience*, 12, 643, 2019.
- Pausata, F. S., Zanchettin, D., Karamperidou, C., Caballero, R., and Battisti, D. S.: ITCZ shift and extratropical teleconnections drive ENSO  
820 response to volcanic eruptions, *Science Advances*, 6, eaaz5006, 2020.
- Plummer, C. T., Curran, M. A., Ommen, T. D. v., Rasmussen, S. O., Moy, A. D., Vance, T. R., Clausen, H. B., Vinther, B. M., and Mayewski, P. A.: An independently dated 2000-yr volcanic record from Law Dome, East Antarctica, including a new perspective on the dating of the 1450s CE eruption of Kuwae, Vanuatu, *Climate of the Past*, 8, 1929–1940, 2012.
- Preddybaylo, E., Stenchikov, G., Wittenberg, A. T., and Osipov, S.: El Niño/Southern Oscillation response to low-latitude volcanic eruptions  
825 depends on ocean pre-conditions and eruption timing, *Communications Earth & Environment*, 1, 1–13, 2020.

- Raible, C. C., Brönnimann, S., Auchmann, R., Brohan, P., Frölicher, T. L., Graf, H.-F., Jones, P., Luterbacher, J., Muthers, S., Neukom, R., et al.: Tambora 1815 as a test case for high impact volcanic eruptions: Earth system effects, *Wiley Interdisciplinary Reviews: Climate Change*, 7, 569–589, 2016.
- 830 Rao, M. P., Cook, B. I., Cook, E. R., D'Arrigo, R. D., Krusic, P. J., Anchukaitis, K. J., LeGrande, A. N., Buckley, B. M., Davi, N. K., Leland, C., et al.: European and Mediterranean hydroclimate responses to tropical volcanic forcing over the last millennium, *Geophysical research letters*, 44, 5104–5112, 2017.
- Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 5, 459–482, <https://doi.org/10.1002/jame.20022>, 2013.
- 835 Rennie, J. J., Lawrimore, J. H., Gleason, B. E., Thorne, P., Morice, C., Menne, M., Williams, C., de Almeida, W. G., Christy, J., Flannery, M., et al.: The international surface temperature initiative global land surface databank: Monthly temperature data release description and methods, *Geoscience Data Journal*, 1, 75–102, 2014.
- Robock, A.: Volcanic eruptions and climate, *Reviews of Geophysics*, 38, 191–219, 2000.
- Schneider, L., Smerdon, J. E., Büntgen, U., Wilson, R. J., Myglan, V. S., Kirilyanov, A. V., and Esper, J.: Revising midlatitude summer temperatures back to AD 600 based on a wood density network, *Geophysical Research Letters*, 42, 4556–4562, 2015.
- 840 Schneider, L., Smerdon, J. E., Pretis, F., Hartl-Meier, C., and Esper, J.: A new archive of large volcanic events over the past millennium derived from reconstructed summer temperatures, *Environmental Research Letters*, 12, 094 005, 2017.
- Schurer, A. P., Hegerl, G. C., Luterbacher, J., Brönnimann, S., Cowan, T., Tett, S. F., Zanchettin, D., and Timmreck, C.: Disentangling the causes of the 1816 European year without a summer, *Environmental Research Letters*, 14, 094 019, 2019.
- Schweingruber, F., Briffa, K., and Jones, P.: Yearly maps of summer temperatures in Western Europe from AD 1750 to 1975 and Western 845 North America from 1600 to 1982, *Vegetatio*, 92, 5–71, 1991.
- Shindell, D. T., Schmidt, G. A., Mann, M. E., and Faluvegi, G.: Dynamic winter climate response to large tropical volcanic eruptions since 1600, *Journal of Geophysical Research: Atmospheres*, 109, 2004.
- Sigl, M., McConnell, J. R., Layman, L., Maselli, O., McGwire, K., Pasteris, D., Dahl-Jensen, D., Steffensen, J. P., Vinther, B., Edwards, R., et al.: A new bipolar ice core record of volcanism from WAIS Divide and NEEM and implications for climate forcing of the last 2000 850 years, *Journal of Geophysical Research: Atmospheres*, 118, 1151–1169, 2013.
- Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., et al.: Timing and climate forcing of volcanic eruptions for the past 2,500 years, *Nature*, 523, 543–549, 2015.
- Smith, V. C., Costa, A., Aguirre-Díaz, G., Pedrazzi, D., Scifo, A., Plunkett, G., Poret, M., Tournigand, P.-Y., Miles, D., Dee, M. W., et al.: The magnitude and impact of the 431 CE Tierra Blanca Joven eruption of Ilopango, El Salvador, *Proceedings of the National Academy of 855 Sciences*, 117, 26 061–26 068, 2020.
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., et al.: Atmospheric component of the MPI-M Earth system model: ECHAM6, *Journal of Advances in Modeling Earth Systems*, 5, 146–172, 2013.
- Stevenson, S., Fasullo, J. T., Otto-Bliesner, B. L., Tomas, R. A., and Gao, C.: Role of eruption season in reconciling model and proxy responses to tropical volcanism, *Proceedings of the National Academy of Sciences*, 114, 1822–1826, 2017.
- 860 Stoffel, M., Khodri, M., Corona, C., Guillet, S., Poulain, V., Bekki, S., Guiot, J., Luckman, B. H., Oppenheimer, C., Lebas, N., et al.: Estimates of volcanic-induced cooling in the Northern Hemisphere over the past 1,500 years, *Nature Geoscience*, 8, 784–788, 2015.
- Tierney, J. E., Abram, N. J., Anchukaitis, K. J., Evans, M. N., Giry, C., Kilbourne, K. H., Saenger, C. P., Wu, H. C., and Zinke, J.: Tropical sea surface temperatures for the past four centuries reconstructed from coral archives, *Paleoceanography*, 30, 226–252, 2015.

- Timmreck, C.: Modeling the climatic effects of large explosive volcanic eruptions, *Wiley Interdisciplinary Reviews: Climate Change*, 3, 865 545–564, 2012.
- Timmreck, C., Lorenz, S. J., Crowley, T. J., Kinne, S., Raddatz, T. J., Thomas, M. A., and Jungclaus, J. H.: Limited temperature response to the very large AD 1258 volcanic eruption, *Geophysical Research Letters*, 36, 2009.
- Toohey, M. and Sigl, M.: Volcanic stratospheric sulphur injections and aerosol optical depth from 500 BCE to 1900 CE, *Earth System Science Data*, 9, 809–831, 2017.
- 870 Toohey, M., Krüger, K., Niemeier, U., and Timmreck, C.: The influence of eruption season on the global aerosol evolution and radiative impact of tropical volcanic eruptions, *Atmospheric Chemistry and Physics*, 11, 12 351–12 367, <https://doi.org/10.5194/acp-11-12351-2011>, 2011.
- Toohey, M., Stevens, B., Schmidt, H., and Timmreck, C.: Easy Volcanic Aerosol (EVA v1. 0): an idealized forcing generator for climate simulations, *Geoscientific Model Development*, 9, 4049–4070, 2016.
- 875 Toohey, M., Krüger, K., Schmidt, H., Timmreck, C., Sigl, M., Stoffel, M., and Wilson, R.: Disproportionately strong climate forcing from extratropical explosive volcanic eruptions, *Nature Geoscience*, 12, 100, <https://doi.org/10.1038/s41561-018-0286-2>, <https://www.nature.com/articles/s41561-018-0286-2.pdf>, 2019.
- Wegmann, M., Brönnimann, S., Bhend, J., Franke, J., Folini, D., Wild, M., and Luterbacher, J.: Volcanic influence on European summer precipitation through monsoons: possible cause for “Years without Summer”, *Journal of climate*, 27, 3683–3691, 2014.
- 880 Wilson, R., Anchukaitis, K., Briffa, K. R., Büntgen, U., Cook, E., D’arrigo, R., Davi, N., Esper, J., Frank, D., Gunnarson, B., et al.: Last millennium northern hemisphere summer temperatures from tree rings: Part I: The long term context, *Quaternary Science Reviews*, 134, 1–18, 2016.
- Wilson, R., Anchukaitis, K., Andreu-Hayles, L., Cook, E., D’Arrigo, R., Davi, N., Haberbauer, L., Krusic, P., Luckman, B., Morimoto, D., et al.: Improved dendroclimatic calibration using blue intensity in the southern Yukon, *The Holocene*, 29, 1817–1830, 2019.
- 885 Winter, A., Zanchettin, D., Miller, T., Kushnir, Y., Black, D., Lohmann, G., Burnett, A., Haug, G. H., Estrella-Martínez, J., Breitenbach, S. F., et al.: Persistent drying in the tropics linked to natural forcing, *Nature Communications*, 6, 1–8, 2015.
- Wunderlich, F. and Mitchell, D. M.: Revisiting the observed surface climate response to large volcanic eruptions, *Atmospheric Chemistry and Physics*, 17, 485–499, <https://doi.org/10.5194/acp-17-485-2017>, 2017.
- Yang, W., Vecchi, G. A., Fueglistaler, S., Horowitz, L. W., Luet, D. J., Muñoz, Á. G., Paynter, D., and Underwood, S.: Climate Impacts From Large Volcanic Eruptions in a High-Resolution Climate Model: The Importance of Forcing Structure, *Geophysical Research Letters*, 46, 890 7690–7699, 2019.
- Zanchettin, D., Bothe, O., Graf, H. F., Lorenz, S. J., Luterbacher, J., Timmreck, C., and Jungclaus, J. H.: Background conditions influence the decadal climate response to strong volcanic eruptions, *Journal of Geophysical Research: Atmospheres*, 118, 4090–4106, 2013.
- Zanchettin, D., Bothe, O., Lehner, F., Ortega, P., Raible, C., and Swingedouw, D.: Reconciling reconstructed and simulated features of the 895 winter Pacific/North American pattern in the early 19th century, *Climate of the Past*, 11, 939–958, 2015.
- Zanchettin, D., Khodri, M., Timmreck, C., Toohey, M., Schmidt, A., Gerber, E. P., Hegerl, G., Robock, A., Pausata, F. S., Ball, W. T., et al.: The Model Intercomparison Project on the climatic response to Volcanic forcing (VolMIP): experimental design and forcing input data for CMIP6, *Geoscientific Model Development*, 9, 2701–2719, 2016.
- Zanchettin, D., Timmreck, C., Toohey, M., Jungclaus, J. H., Bittner, M., Lorenz, S. J., and Rubino, A.: Clarifying the relative role of forcing 900 uncertainties and initial-condition unknowns in spreading the climate response to volcanic eruptions, *Geophysical Research Letters*, 46, 1602–1611, 2019.



Zhu, F., Emile-Geay, J., Hakim, G. J., King, J., and Anchukaitis, K. J.: Resolving the differences in the simulated and reconstructed temperature response to volcanism, *Geophysical Research Letters*, 47, e2019GL086908, 2020.