

1 **Simulation of the mid-Pliocene Warm Period using HadGEM3:**
2 **Experimental design and results from model-model and model-data**
3 **comparison**

4
5 **Charles J. R. Williams^{1,6}, Alistair A. Sellar², Xin Ren¹, Alan M. Haywood³, Peter**
6 **Hopcroft⁴, Stephen J. Hunter³, William H. G. Roberts⁵, Robin S. Smith⁶, Emma J.**
7 **Stone¹, Julia C. Tindall³, Daniel J. Lunt¹**

8
9 ¹School of Geographical Sciences, University of Bristol, UK

10 ²Met Office Hadley Centre, UK

11 ³School of Earth and Environment, University of Leeds, UK

12 ⁴School of Geography, Earth and Environmental Sciences, University of Birmingham, UK

13 ⁵Department of Geography and Environmental Sciences, Northumbria University, UK

14 ⁶NCAS, Department of Meteorology, University of Reading, UK

15
16 **SUPPLEMENTARY MATERIAL**

17
18 **TEXT**

19 **Model description**

20 ***HadGEM2-AO***

21 The immediate predecessor to HadGEM3 is the family of HadGEM2 models, all of which vary in
22 terms of level of complexity but all of which have a common computational framework (Tindall and
23 Haywood 2020, Martin *et al.* 2011), which HadGEM3 also shares. The most complex full Earth
24 system version of the family, HadGEM2-ES, was included in the previous IPCC Assessment Report,
25 AR5. Tindall and Haywood (2020) conducted a Pliocene simulation for PlioMIP1 using the fully
26 coupled version of this model, HadGEM2-AO (hereafter referred to as HadGEM2). This model has
27 the same atmospheric spatial resolution as HadGEM3, but only 38 atmospheric vertical levels; for full
28 details on HadGEM2, see Collins *et al.* (2011), Martin *et al.* (2011) and Tindall and Haywood (2020).
29 In contrast to the HadGEM3 *mPWP* simulation, the HadGEM2 Pliocene simulation uses dynamic
30 vegetation from TRIFFID (Top-down Representation of Interactive Foliage and Flora Including
31 Dynamics, see Cox 2001), and a previous iteration of the PRISM boundary conditions, PRISM3 (see
32 Dowsett *et al.* 2007 and Dowsett *et al.* 2010). It should be noted that, whilst PRISM3 was mostly
33 implemented in this model, this does not include the orography, which was the same as pre-industrial
34 away from ice sheet regions. It should also be noted that the LSM used in this model differs slightly

35 from both PRISM3 and HadGEM3 simulations, in that the Bering Sea, Canadian Archipelago and
36 Hudson Bay gateways are all open (Tindall and Haywood 2020).

37

38 ***HadCM3***

39 The original fully-coupled atmosphere-ocean version of the UK's physical climate model is HadCM3
40 (Gordon *et al.* 2000), and over the years this has been used extensively for paleoclimate simulations
41 and has been updated/optimised according to the simulation in question. Although no longer
42 considered a state-of-the-art model, its fast speed and relatively cheap computational cost still makes
43 it appropriate for paleoclimate simulations (Hunter *et al.* 2019) and, thanks to this, it has been
44 included in every phase of CMIP to date. Both of the older HadCM3 simulations used here
45 (HadCM3-PRISM2 and HadCM3-PlioMIP1) have an atmospheric resolution of 3.75° longitude by
46 2.5° latitude with 19 vertical levels, and an ocean resolution of 1.25 ° longitude/latitude with 20
47 vertical levels; for full details, see Gordon *et al.* (2000). For a land surface scheme, both of these
48 simulations use the 1st generation Met Office Surface Exchange Scheme (MOSES1, see Cox *et al.*
49 1999), and both use dynamic vegetation. Concerning boundary conditions, HadCM3-PRISM2
50 predates PlioMIP1 and thus uses PRISM2, whereas HadCM3-PlioMIP1 was included in PlioMIP1
51 and, similar to HadGEM2, uses PRISM3 boundary conditions.

52

53 ***PlioMIP2 models***

54 The same 16 models as those in H16 are included here as a comparison to HadGEM3. These models,
55 along with their spatial resolutions, are listed in Table 4; see Table 1 in H16 for full information
56 (including boundary conditions, equilibrium climate sensitivity values and references) on each model.
57 It should be noted that one of these models is HadCM3 but is slightly different to the earlier versions
58 discussed here; the model, HadCM3-PlioMIP2, was run by Hunter *et al.* (2019) and is equivalent
59 (concerning updates) to the version developed by Valdes *et al.* (2017), HadCM3B-M2.1 (which
60 includes an updated land surface scheme, MOSES2, see Essery *et al.* 2001). In short, whereas
61 MOSES1 treats each model grid point as a homogeneous surface and calculates energy and moisture
62 fluxes using effective parameters, MOSES2 has subgrid heterogeneity and an improved representation
63 of surface and plant processes (Hunter *et al.* 2019); see Valdes *et al.* (2017) for a complete
64 comparison of MOSES1 and MOSES2. All of the models included in PlioMIP2 use PRISM4
65 boundary conditions.

66

67

68 TABLES

Value	Mega biome	BLT	NLT	C3 Grass	C4 Grass	Shrubs	Urban	Lakes	Bare soil	Land ice
1	Tropical forest	0.92	0	0	0.02	0.01	0	0	0.05	0
2	Warm-temperate forest	0.75	0	0.07	0.03	0.1	0	0	0.05	0
3	Savanna and dry woodland	0.18	0	0	0.67	0.05	0	0	0.1	0
4	Grassland and dry shrubland	0.05	0	0	0.55	0.3	0	0	0.1	0
5	Desert	0	0	0	0.02	0.13	0	0	0.85	0
6	Temperate forest	0	0.75	0.1	0	0.1	0	0	0.05	0
7	Boreal forest	0	0.7	0.2	0	0.025	0	0	0.075	0
8	Tundra	0	0	0	0	0.4	0	0	0.6	0
9	Dry tundra	0	0	0	0	0.4	0	0	0.6	0
28	Land ice	0	0	0	0	0	0	0	0	1

69


70 Table S1 - Lookup table to translate mega biomes from PRISM3 into HadGEM3 PFTs. Values in
71 first column correspond to those in Figure 2

72

73

74 **FIGURES**


75



76

77 Figure S1 - Land sea mask used in HadGEM3 *mPWP* and *piControl* simulations, with colours
78 showing fractional coverage of coastal grid points


79



80

81 Figure S2 – Canopy height used in HadGEM3, for each PFT. Dashed lines show global mean from
 82 *piControl* simulation, solid lines show latitudinally varying function of this global mean, used in
 83 *mPWP* simulation. a) broadleaf trees; b) needle-leaved trees; c) temperate C3 grass; d) tropical C4
 84 grass; e) shrubs


85



86

87 Figure S3 - Soil parameters used in HadGEM3. Left-hand column: *piControl* simulation, right-hand
 88 column: *mPWP* simulation. a) Volume fraction of condensed water in soil at wilting point, b)
 89 Volume fraction of condensed water in soil at critical point, c) Volume fraction of condensed water in
 90 soil at saturation point, d) Clapp-Hornberger "B" coefficient, e) Thermal conductivity, f) Hydraulic
 91 conductivity at saturation, g) Thermal capacity, h) Saturated soil water suction, i) Snow-free albedo of
 92 soil, j) Soil carbon content, k) Soil bulk density


93



94


95 Figure S4 - Soil dust properties used in HadGEM3. Left-hand column: *piControl* simulation, right-
 96 hand column: *mPWP* simulation. a) Dust parent soil clay fraction, b) Dust parent silt clay fraction, c)
 97 Dust parent soil sand fraction, d) Dust soil mass fraction (Division 1), e) Dust soil mass fraction
 98 (Division 2), f) Dust soil mass fraction (Division 3), g) Dust soil mass fraction (Division 4), h) Dust
 99 soil mass fraction (Division 5), i) Dust soil mass fraction (Division 6)

100




101
102

103 Figure S5 – Annual global mean net top of atmosphere (TOA) radiation from the HadGEM3 *mPWP* spin-up
 104 phase and production run, as well as the last 100 years from the CMIP6 *piControl* and the *piControl_mod*. See
 105 Williams *et al.* (2020) for the *piControl* spin-up phase that preceded this simulation
 106




107
108
109
110
111

Figure S6 - Statistically significant (as calculated by a Mann-Kendall test, using the 99% level) centennial trends in 1.5m temperature from the HadGEM3 Pliocene *mPWP* simulation




112

113 Figure S7 – Annual global mean measures of climate equilibrium from the HadGEM3 *mPWP* spin-up phase and
 114 production run, as well as the last 100 years from the CMIP6 *piControl* and the *piControl_mod*. See Williams *et*
 115 *al.* (2020) for the *piControl* spin-up phase that preceded this simulation: a) Full depth ocean temperature, b) Full
 116 depth ocean salinity
 117



118
 119 Figure S8 – Sea ice fraction climatology from HadGEM3. Left-hand column: *piControl_mod*
 120 simulation, right-hand column: *mPWP* simulation. a) Annual, b) DJF, c) JJA
 121



122

123

124 Figure S9 – 1.5 m air temperature climatology differences (Pliocene - PI) from HadGEM3 *mPWP*
 125 simulation and all other models in PlioMIP2, as well as multi-model ensemble mean (MME)