

Holocene vegetation transitions and their climatic drivers in MPI-ESM1.2

Anne Dallmeyer¹, Martin Claussen^{1,2}, Stephan J. Lorenz¹, Michael Sigl³, Matthew Toohey⁴, Ulrike Herzschuh^{5,6,7}

5 ¹Max Planck Institute for Meteorology, Bundesstrasse 53, 20146 Hamburg, Germany

²Centrum für Erdsystemforschung und Nachhaltigkeit (CEN), Universität Hamburg, Bundesstrasse 55, 20146 Hamburg, Germany

³Climate and Environmental Physics and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

10 ⁴ Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, Canada

⁵Alfred Wegner Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, Germany.

⁶Institute of Environmental Sciences and Geography, University of Potsdam, Germany

⁷Institute of Biochemistry and Biology, University of Potsdam, Germany

Correspondence to: Anne Dallmeyer (anne.dallmeyer@mpimet.mpg.de)

15 **Abstract**

We present a transient simulation of global vegetation and climate patterns of the mid- and late-Holocene using the MPI-ESM (Max Planck Institute for Meteorology Earth System Model) at T63 resolution. The simulated vegetation trend is discussed in the context of the simulated Holocene climate change. Our model captures the main trends found in reconstructions. Most prominent are the southward retreat of the northern treeline that is combined with the strong decrease of forest in the high northern latitudes during the Holocene and the vast increase of the Saharan desert, embedded in a general decrease in precipitation and vegetation in the northern hemispheric monsoon margin regions. The southern hemisphere experiences weaker changes in total vegetation cover during the last 8000 years. However, the monsoon-related increase in precipitation and the insolation-induced cooling of the winter climate lead to shifts in the vegetation composition, mainly between the woody plant functional types (PFTs).

The large-scale global patterns of vegetation almost linearly follow the subtle, approximately linear, orbital forcing. In some regions, however, non-linear, more rapid changes in vegetation are found in the simulation. The most striking region is the Sahel-Sahara domain with rapid vegetation transitions to a rather desertic state, despite a gradual insolation forcing. Rapid shifts in the simulated vegetation also occur in the high northern latitudes, in South Asia and in the monsoon margins of the southern hemisphere. These rapid changes are mainly triggered by changes in the winter temperatures, which go into, or move out of, the bioclimatic tolerance range of individual PFTs (Plant Functional Types). The dynamics of the transitions are determined by dynamics of the Net Primary Production (NPP) and the competition between PFTs. These changes mainly occur on timescales of centuries. More rapid changes in PFTs that occur within a few decades are mainly associated with the time scales of mortality and the bioclimatic thresholds implicit in the dynamic vegetation model, which have to be interpreted with caution.

Most of the simulated Holocene vegetation changes outside the high northern latitudes are associated with modifications in the intensity of the global summer monsoon dynamics that also affect the circulation in the

extra tropics via teleconnections. Based on our simulations, we thus identify the global monsoons as the key
40 player in Holocene climate and vegetation change.

1 Introduction

The Holocene (the last ~11,500 years) has been a period of strong global environmental changes. These were
mainly forced by the retreat of the continental ice masses, variations in the seasonal insolation associated
45 with changes in the Earth orbital constellation, and solar variability (Mayewski et al., 2004).

Due to the precession of the equinoxes and the earth axis, the timing of perihelion (the point closest to the
sun on the earth orbit) has been shifted continuously from August at 9ka BP (i.e. 9000 years before present),
to January at present day (Ruddiman, 2008). This contributed mainly to the gradual changes in the seasonal
energy input from the sun. For instance, the northern hemisphere received approximately 10% more summer
50 insolation at 9ka, whereas winter insolation was reduced by approximately 10% (Berger, 1978). Hence, the
seasonality in the northern hemisphere has weakened over the Holocene, while it has strengthen in the
southern hemisphere (e.g. Fischer and Jungclaus, 2011). As a consequence, the atmospheric and oceanic
circulations have continuously adjusted to the energy change, affecting in turn the precipitation and
temperature distribution and thus also the global vegetation patterns.

55 Winter climate was colder and summer climate was warmer on the northern hemispheric continents during
the mid-Holocene compared to present-day (e.g. Zhang et al., 2018, Viau et al., 2006, Davis et al., 2003).

Fewer studies have focused on Holocene temperature evolution on the southern hemisphere. Model
experiments reveal a cooling trend in austral winter and a warming trend in austral summer over the course
of the Holocene (Lorenz et al., 2006), in line with the orbital forcing.

60 The decrease in boreal summer insolation induced a southward movement of the Intertropical Convergence
Zone (ITCZ, e.g., Haug et al., 2001) and a southward retreat and weakening of the summer monsoons in the
northern hemisphere (e.g. Kutzbach 1981, Liu et al. 2003, Dallmeyer et al., 2015, D'Agostino et al., 2019),
implying a drying trend in subtropical Asia, Africa and Central America. In contrast, the summer monsoon
systems in the southern hemisphere strengthen during the Holocene (e.g. Zhao and Harrison, 2012, Jiang et
65 al., 2015, Prado et al. 2013a) and result in a moistening trend roughly north of 40°S (e.g. Royas and Moreno,
2009).

In the extra-tropics, reconstructions reveal a moistening trend in large parts of North America (Viau and
Gajewski, 2001, Williams et al., 2010) and South America since the mid-Holocene (Lamy et al, 2001, Prado
et al., 2013b). As possible explanations for this in-phase climate change in the Americas, Holocene
70 weakening of the northern hemispheric subtropical anticyclones and an enhancement of the north-southward
migration of the ITCZ have been proposed (Grimm et al., 2001). Furthermore, the southern hemispheric
westerly winds were probably shifted poleward and were intensified during the mid-Holocene, but
reconstructions reveal contradictory signals (Fletcher and Moreno and references therein, 2012). As another
explanation, the reduced El Niño Southern Oscillation activity and a more La Niña like mean state of the
75 Tropical Pacific during the mid-Holocene are proposed (e.g. Clement et al., 2004; Overpeck and Webb,

2000), leading to drier conditions on the extratropical American continents (Barr et al., 2019) and wetter conditions particularly in Australia. A detailed summary of the mid- to late-Holocene climate change is given in Wanner et al. (2008).

Pollen-based reconstructions reveal substantial vegetation changes as response to the Holocene climate signal. The forests in the northern mid- to high-latitudes were expanded during mid-Holocene compared to present day distributions, replacing tundra areas (MacDonald et al., 2000, Prentice et al., 2000) and coinciding a northward shifted northern treeline (Bigelow et al., 2003).

In the Afro-Asian monsoon region, the forest and steppe belts penetrated further northward onto the continents and the northern hemispheric desert regions were substantially reduced during mid-Holocene compared to present day (Zhao et al., 2009; Feng et al., 2006; Yu et al., 2000; Jolly et al., 1998, Metcalfe et al., 2015). At least for North Africa, reconstructions suggest a rapid increase of the Saharan desert over the course of the Holocene (e.g., deMenocal et al., 2000; Shanahan et al. 2015).

In North America, plant populations have expanded northwards over the Holocene (Williams et al., 2004). Biome reconstructions show rather stable conditions, but also reveal a northward expansion of the temperate forest and evergreen taiga (Cao et al., 2019a). Furthermore, forest taxa have re-invaded into the prairie (eastern Great Plains), reflecting the increase in precipitation over the mid- and late Holocene (Grimm et al., 2001). Most of the American equatorial savanna regions were more widespread during the mid-Holocene (Behling and Hooghiemstra, 2000).

In the Amazonian region, Holocene vegetation changes were minor, with the exception of southeast Brazil and southwest Amazonia indicating an increase in forests over the Holocene. Southern hemispheric grasslands have revealed a transition to communities requiring more moisture reflecting the increase in moisture levels over the course of the Holocene (Grimm et al., 2001).

Biome reconstructions for Australia show only little and locally varying vegetation changes since the mid-Holocene. Some records from the south-eastern part reveal moisture-stressed vegetation types at 6ka BP (i.e. before present), implying drier conditions there. Other records, e.g. from the Great Diving Range indicate increased vegetation cover and more moisture demanding plants during the mid-Holocene. The biome changes in the tropics have been minor, rather showing transitions between different forest types (Pickett et al., 2004).

Reconstructions for Southern Africa are rare, often incomplete, and show contradictory signals. During the early Holocene, desert pollen concentrations in sediments off the western South African coast were increased, followed by a period of increasing moisture during the mid-Holocene. Vegetation was more open on the southern continent (Scott and Lee-Thorp, 2004). Numerous records indicate an aridification during the mid- and late-Holocene (Burrough and Thomas, 2013). In tropical south east Africa, more humid vegetation types have occurred during the mid-Holocene (e.g. Marchant et al., 2018).

These assessments indicate that reconstructions have generally provided a comprehensive overview of the vegetation changes during the Holocene. But so far, most studies deal with the vegetation dynamics at individual sites or regional comparisons. Several continental-scale syntheses are available (e.g. Cao et al., 2019a; Williams, 2003; Williams et al. 2011; Trondman et al., 2015, Marquer et al., 2017), but no global

overview exists, showing the Holocene vegetation trend in its global context. This is partly related to the fact that the spatial coverage of records is low outside Europe and North America, limiting the possibilities of reconstructions. Even more systematic investigations of vegetation-climate relationships are not possible at all, because terrestrial climate proxy-data independent from vegetation are scarce. The climatic background of the vegetation changes can only be assessed. Therefore, we still lack substantial understanding of the climatic drivers associated with the Holocene vegetation dynamics, not only at the global scale, but also at the regional to continental scale. Earth System Modelling can provide insights into these processes.

So far, global simulations of Holocene vegetation change mostly focused on the comparison of vegetation composition between time-slices, while the dynamics were seldom taken into consideration (e.g. Brovkin et al., 2002; Crucifix et al., 2002). Recently, Braconnot et al. (2019) discussed the advantages and challenges of transient vegetation simulations in comparison to time-slice simulations by using the Earth System Model IPSL. In their simulation, the vegetation reduction in Northern Africa occurs and ends later than the forest decline in the northern hemisphere. In addition, the model indicates large variations in the forest composition throughout the Holocene. However, the vegetation-climate relationships in time and space were hitherto not investigated systematically in transient vegetation simulations conducted with Earth system models.

Here, we use a transient simulation of the mid- to late Holocene period performed with the comprehensive Earth System Model MPI-ESM1.2 to analyse the Holocene vegetation change in the context of the simulated global climate change. The aim of the study is to provide a detailed overview on the global and regional pattern of the Holocene vegetation changes and their main climatic drivers in the model. We focus on the long-term, multi-centennial vegetation trends.

After the description of the model and the transient simulation, the simulated vegetation change is evaluated against global biome-reconstructions for the mid-Holocene and present-day time-slices.

In section 4, we provide an overview of the general trend in the global vegetation pattern to specify the main regions of strong vegetation transitions. By performing a cluster analysis the coincidence of the regional vegetation trends are figured out. A redundancy analysis is undertaken to assess the main climatic drivers (precipitation vs temperature) of the vegetation change. In section 5, we zoom into the global Holocene vegetation clusters to discuss the regional vegetation transitions and interpret them in the context of the changes in climate and the large-scale atmospheric circulation systems (section 5). As some reconstructions reveal abrupt local transitions in the vegetation composition over the Holocene, we furthermore test and discuss the simulated vegetation changes with respect to their rapidity compared to the long-term change and orbital forcing (section 6). We summarize the results and conclude our study in section 7.

145

2 Methods

2.1 The Earth System Model MPI-ESM1.2

The model MPI-ESM1.2 (Mauritsen et al., 2019), developed at the Max-Planck-Institute for Meteorology, consists of the general circulation model of the ocean MPIOM (Jungclaus et al., 2013), including the ocean

150 biogeochemistry model HAMOCC (Ilyina et al., 2013), coupled to the atmospheric general circulation model
ECHAM6.3 (Stevens et al., 2013). The terrestrial carbon cycle and vegetation dynamics are calculated by the
land-surface scheme JSBACH3 (Reick et al., 2013) that incorporates the dynamic vegetation module
developed by Brovkin et al. (2009). Through the inclusion of the subsystem models JSBACH and
HAMOCC, the full carbon cycle is considered in MPI-ESM. Nevertheless, the carbon cycle is not fully
155 interactive in this simulation as the atmospheric CO₂ concentration is prescribed. ECHAM6 and the land
model was configured with a spectral resolution of T63 (corresponding to a horizontal resolution of
approximately 200km on a Gaussian grid) with 47 levels in the vertical atmosphere. For the ocean model the
horizontal resolution GR15 (i.e. 256x220 on a bipolar grid, corresponding to 12 to 180km) and 64 vertical
levels have been used.

160

2.2 The dynamic vegetation module

In the dynamic vegetation module (see Brovkin et al., 2009, Reick et al., 2013), natural vegetation is
represented by eight different plant functional types (PFT). Trees are classified into either tropical or
extratropical trees and their phenology can be evergreen or deciduous. Shrubs are distinguished as raingreen
165 (moisture limited) and cold-resistant (temperature limited) shrubs. Grass is differentiated between C3 and C4
grass. JSBACH uses a tiling approach, thus all PFTs can coexist in each grid-cell of the land-surface in
principle and are assigned certain cover fractions, ranging from 0 (not present) to 1 (covering the full grid-
cell).

The occurrence of each PFT is limited by temperature constraints that reflect the bioclimatic tolerance of the
170 individual PFTs. These environmental thresholds represent the chilling requirements of the plants by the
maximum mean temperature of the coldest month ($T_{c_{max}}$) and the cold resistance of the plants by the
minimum mean temperature of the coldest month ($T_{c_{min}}$). Furthermore, the presence of the PFTs is regulated
by their heat requirements during their growth phase. This limit is realized by temperature sums over days on
which mean temperature exceeds different thresholds (0°C and 5°C), called growing degree days (GDD0 and
175 GDD5, respectively). Cold shrubs are also not able to survive in too warm climates and are limited by the
maximum mean temperature of the warmest month ($T_{w_{max}}$). The different bioclimatic limits for the
individual PFTs are listed in Table 1.

In addition to these limits, the dynamics of changes in the fractional coverage of PFTs in response to climatic
change are governed by the dynamics of the Net Primary Production (NPP) of the competing PFTs. The time
180 scales of NPP dynamics include time scales for allocation, mortality and perturbation. The times scales of
allocation and mortality are the same for each individual PFT, but differ between PFTs. They range from 50
years for extratropical trees to 1 year for grass (see Table 1). If climate changes and if different PFTs can
exist within the range of climate variability, then the PFT with the strongest NPP dominates, and the time
scale of the transition mainly depends on the difference in the NPP allocation of the competing PFTs. If this
185 difference is small, the transition is slow. The transition can be faster, if one of the PFTs has much shorter
time scales of allocation and mortality. The fastest changes occur, if the climate crosses a bioclimatic

threshold (see Table 1) or if a large-scale perturbation, such as windbreak or fire, occurs. Then, the PFT affected decays exponentially within a few decades or less.

190 The fractional PFT coverage of a grid cell can also change, if the bare soil fraction (BSF), which represents the seasonal and permanently non-vegetated ground, varies. The BSF is determined by the relation of the carbon actually stored in the carbon pool for living tissues via the NPP of the individual plants to the maximum yearly carbon storage in this pool. This maximum is defined by the carbon costs needed by the plants to build up the leaves, stems etc. completely and depends on the ratio of the maximum leaf area index to the specific leaf area of the PFTs.

195

2.3 Transient simulation

To run the model in quasi-equilibrium between **early** mid-Holocene boundary conditions, climate and the carbon cycle, a **spin-up** simulation has been performed with all forcings fixed to the values of the year 6000 BCE. This simulation was initialised from the pre-industrial climate and vegetation state and was run for
200 more than 1000 years. The transient simulation then started from this equilibrium state and continued until pre-industrial times (i.e. 1850 CE). The transient simulation used in this study has been performed nearly identical to the simulation described in Bader et al. (2020), Brovkin et al. (2019), and Dallmeyer et al. (2020). The following forcings were prescribed:

205 a) orbital-induced insolation changes (Berger, 1978), updated every decade. These mainly impact the seasonal cycle of e.g. temperature and precipitation. The seasonality increases in the southern hemisphere and reduces in the northern hemisphere during the Holocene.

b) Greenhouse gas concentration (methane, carbon dioxide and nitrous oxide) inferred from ice core records (F. Joos, personal communication; see Brovkin et al., 2019 and Köhler, 2019), updated every decade. The difference in CO₂ between start and end of the simulations amounts to approximately 20 ppm.

210 c) Stratospheric sulfate aerosol injections imitating volcanic eruptions, prescribed from the Easy Volcanic Aerosol (EVA) forcing generator (Toohey et al., 2016), read annually, but calculated daily by linear interpolation. This forcing has been reconstructed based on ice-cores from both hemispheres to better identify volcanic eruptions with global impact on the climate. From Greenland, we employed sulfate data from the Greenland Ice Sheet Project Two (GISP2; 72.97°N, 38.80°W) ice core (Mayewski et al., 1997).
215 From Antarctica, we used sulfate data from the EPICA Dronning Maud Land (EDML; 75.00°S, 00.07°E) and Dome Concordia (EDC; 75.10°S, 123.35°E) ice cores (Severi et al., 2007), as well as sulfur and sulfate data from the WAIS (West Antarctic Ice Sheet) Divide ice core project (WD; 79.48°S, 112.11°W) (Cole-Dai et al., 2021). These four ice cores were synchronized to the WD2014 chronology (Sigl et al., 2016) and volcanic sulfate mass deposition rates over Greenland and Antarctica were estimated using established methods (Sigl et al., 2014; Sigl et al., 2015). Using the methodology developed (Gao et al., 2007) and applied to a similar bipolar network of ice cores over the past 2,500 years (Toohey and Sigl 2017) we estimated stratospheric sulfur injection (SSI) for in total 528 volcanic eruptions. Conversion of SSI into the aerosol properties is performed with the Easy Volcanic Aerosol module (Toohey et al. 2016). Aerosol extinction is assumed to be
220

linearly proportional to mass for eruptions smaller than the 1815 Tambora eruption, and follows a $2/3$ power-law scaling for larger eruptions (Crowley and Unterman, 2013). The previous volcanic forcing reconstruction used in Bader et al. (2020) was based on a Greenland ice-core (GISP2, Zielinski et al. 1996) only and therefore overestimated the effect of Icelandic and other extratropical volcanic eruptions on the global climate.

d) Spectral Solar Irradiance forcing, includes extrapolated 11-years solar cycle based on sun-spot observations data-sets of far infrared, near infrared and visible radiation (Krivova et al., 2011), read annually, but calculated daily by linear interpolation.

e) LUH2 land-use forcing by Hurtt et al. (2020) conform to the CMIP6 simulations, read annually, but calculated daily by linear interpolation. This forcing begins 850 CE with a quasi linear transition period (1000 years) starting 150 BCE to slowly build up the land-use. Bader et al. 2020 used a preliminary (unpublished) version of the LUH2 dataset.

In the analysis of this study, the period 6000BCE to 150BC is considered only, to exclude the effect of land use on the vegetation distribution **in the model**. For easier nomenclature, we define the **early** mid-Holocene time slice (further referred to 8ka before the year 2000, i.e. b2k) by the climatological mean of the first 100 years of this transient simulations (i.e. year 6000-5901 BCE) and the late-Holocene reference period (further referred to 2.15ka b2k) by the climatological mean of the last 100 years of this simulations without land-use (i.e. year 250-150 BCE). Accordingly, we define the 6ka b2k time-slice by the climatological mean of the years 4000-3901 BCE.

2.4 Analysis methods

To summarise Holocene vegetation change, we choose the fuzzy c-means clustering technique (Bezdek,1981). This method calculates a set of independent cluster centres (here: time-series of vegetation development), such that the sum of the square distances between the items assigned to the cluster and the cluster centre is minimized. This means, that grid-cells showing similar vegetation changes over the course of the Holocene are grouped into the same cluster. We here use the c-means clustering function in R (Meyer et al., 2017) and calculate the cluster on the basis of 100yr. running mean time-series for the PFT-groups "forest", "shrubs", "grass" and "bare soil". The number of independent cluster has been optimized by the Xie-Beni method (Xie and Beni, 1991).

To test the most important climatic driver in the different regions, we performed a constrained ordination via the redundancy analysis function "rda" implemented in the "vegan" package of R (Oksanen et al, 2018).

The redundancy analysis (RDA) is an extended multiple linear regression method and measures the linear relationship between a set of response variables (here the PFT-groups "forest", "grass", and "shrubs") and explanatory variables (here precipitation, temperature of the warmest month and temperature of the coldest month). It ordines the vegetation on axes that are constructed in such a way that their relationship to linear combinations of the climate variables is maximised. Therefore, the RDA technique can also be seen as a constrained version of the principal component analysis. In this study, it is taken as a measure of the variation in vegetation that can be explained by the climate variables. The temperature of the coldest month

explains only a small part of the variance in an initial RDA and was thus excluded in the final RDA shown here.

To assess the change in monsoon strength during the Holocene, we calculated the global monsoon area and the global monsoon precipitation following Zhou et al. (2008) with the modifications of Liu et al. (2009). In this definition, monsoon areas are characterized by a precipitation maximum (>55% of the annual total) in the summer season and a large annual range in rainfall. The summer season includes the months May to September for the northern hemisphere and the months November to March for the southern hemisphere. The reverse is applied for the respective winter season. As ‘annual range’ the difference between summer and winter precipitation is taken. The monsoon precipitation is given by the sum of total summer rainfall in the monsoon area, taking the latitudinal changes in the grid-cell area into account.

For some analysis, seasonal means are considered. We are aware of the fact, that the length of the season changes during the Holocene due to the precessional cycle (c.f. Bartlein and Shafer, 2018). However, this does not change the response of the atmospheric physics in the model to the insolation change, so we assume that the results are not affected by a fixed calendar. But please keep in mind, that we define the seasons according to the modern calendar implemented in the model, i.e. any statements regarding seasonal changes only refer to the simulation and might not necessarily be transferable to the real Holocene seasonal changes.

3 Evaluation of the simulated vegetation distribution against biome reconstructions

So far, palaeo-vegetation reconstructions have mostly been synthesized by biome distributions for individual time-slices (e.g. Harrison, 2017; Cao et al., 2019a). Quantitative vegetation reconstructions are only available for few regions (e.g. Trondman et al., 2015; Cao et al., 2019b, Li et al., 2020). Therefore, we decided to evaluate the simulated vegetation against pollen-based biome reconstructions for the mid-Holocene (6ka). The simulated PFT cover fractions were converted into mega-biome distributions using the approach by Dallmeyer et al., (2019). The agreement was tested by using the Best Neighbour Score (BNS, Tab.2, cf. Dallmeyer et al., 2019) that considers also the agreement in the surrounding grid-cells of the record sites. In principle, the vegetation in the model is well in line with the reconstructions (BNS=0.73, Fig. 1). All major biome belts are reproduced. In particular, the extratropical forest biomes are well represented showing a BNS of 0.79 (boreal forest) and 0.88 (temperate forest). The simulated deserts (BNS=0.39) and savannas (BNS=0.12) show little correspondence with the reconstructions. This is related to the fact that also in this simulation the northward spread of vegetation into the Sahara during the mid-Holocene is underestimated compared to the reconstructions (cf. Dallmeyer et al., 2020). Furthermore, the extent of the desert area in Central Asia is too small. The reconstructed desert in the southwest of North America (mainly around the Mojave and Sonoran desert) is not reflected in the model. Very few reconstructions exist for the savanna regions, probably biasing the comparison. However, the method for biomizing PFTs has shortcomings in mapping savannas as this biome can hardly be delimited by climatic constraints and PFT compositions (cf. Dallmeyer et al., 2019). Tropical savannas can only be distinguished from forests by e.g. their fire and shade

tolerance and unique functional ecology (Ratman et al., 2011). To analyse the main change in biomes towards PI, the biomes were further grouped into the main categories forests, savannas, grassland/tundra, and deserts. Fig.1c-d shows qualitatively the simulated and reconstructed difference for these biome categories between 6ka and PI on the basis of the ecological development. Positive development designates the transition from a more open landscape to a tree-dominated landscape (e.g. desert in 6ka to grassland, grasslands to savannas or to forests in PI). Similarly, negative development refers to the transition from tree-dominated biomes to a more open landscape (e.g. forest in 6ka to grassland in PI).

Since not all reconstructions mapped for 6ka reach into pre-industrial times, we also considered PI records in the grid-cells surrounding those of sites mapped in 6ka. This method does not lead to a significant improvement of the record density in the difference plot. Therefore, we retained the original selection, i.e. showing only records with a counterpart at PI. Both, the model and the reconstructions, show only little change in the area covered by these main biomes. Most remarkable is the decrease in forest and grassland in the northern hemispheric monsoon regions and the decrease of forest along the Taiga-Tundra boundary in the high northern latitudes indicated by the model. The reconstructions in principle also suggest this development, even if the record density is too low to show a detailed picture, particularly in North Africa.

The decrease in openness in central North America and central Asia is visible in the model and in the reconstructions. The records furthermore suggest a greening at the Labrador Sea coast, which is not shown by the model. This may be related to the migration of plants after the retreat of the Laurentine ice-sheet. The changes in land ice coverage are not considered in the model so that this process cannot be represented. The model indicates a positive ecological development in the South African and South American extratropics. This is not seen in the records showing rather no change in main biome category, but only few records exist that are located directly in the regions in which the model shows this positive signal. Hence our prediction, or hindcast, awaits further evaluation.

4 Global view of the simulated Holocene vegetation and climate trend

The simulated Holocene vegetation change from 8ka b2k to 2.15ka b2k has been clustered on the basis of the 100yr-running mean time-series. The desert and forest change can be grouped in four, the shrubs and grass change in five different cluster (Fig. 2). Figure 2a,c,e,h mark the regions that can be assigned to the respective cluster. Figure 2b,d,f,g show the trends in the respective cluster centre. Overall, the Holocene vegetation changes in this global view are quite linear, overlaid by weak multi-centennial variability.

However, non-linear changes in PFTs may occur on local or regional level. The most striking vegetation shifts seen in Fig. 2 are:

- a) a strong decrease in forest and increase in grass and shrubs in the high northern latitudes reflecting the southward retreat of the northern treeline and the shift from Taiga forest to Tundra
- b) a (strong) increase in the subtropical northern hemispheric deserts coinciding with a decrease and equator-ward retreat of all vegetation types in the monsoon regions
- c) an increase in the vegetated area in extratropical North America (mainly 30°-60°N, 90-120°W) related to an increase in forest and partly grass

- d) an increase in the vegetation in extratropical South America and South Africa due to an increase in forest, grass and shrubs (S Africa), including the southern hemispheric monsoon regions
- 330 e) a bipolar vegetation change in Australia, mostly driven by an increase in grass on the northern and a decrease of grass on the southern part of the continent.

Fig. 3 shows a similarly calculated cluster analysis for the 100yr running means of annual mean precipitation, and the temperature of the warmest and coldest month, respectively, assuming that these variables are the main climatic factors affecting the vegetation change. In line with the increasing wintertime insolation during the Holocene, the temperature of the coldest month (T_{cold}) uniformly increases in the northern hemisphere, in particular on the continents by up to 1.6K (cluster center). In the southern hemisphere, T_{cold} does only change in a few grid-boxes in South Africa and in the northern part of Australia, showing an overall decreasing trend with strong millennial variability.

The temperature of the warmest month (T_{warm}) shows the opposite trend, in line with the summertime insolation change. T_{warm} decreases in the northern hemisphere, particularly in the continental mid-latitudes (up to 3.2K). In the southern hemisphere, T_{warm} increases moderately during the Holocene by 0.7K. The signal is reversed in the tropical monsoon areas, probably due to a drop in evaporative cooling in the northern and an increase in evaporative cooling in the southern hemispheric tropical monsoon regions. Both, T_{warm} and T_{cold} are intercepted by relatively strong ‘events’ leading to a drop in temperature in all cluster centres by up to 0.7 for several hundred years. These events may at least partly be associated with the volcanic forcing prescribed to the model. The Pearson correlation coefficient between the detrended T_{warm} cluster centres and the volcanic forcing range from 0.6 (for the increasing cluster centre) to 0.8 (for the decreasing cluster centre). The Pearson correlation coefficient between the detrended T_{cold} cluster centres and the volcanic forcing range from 0.27 (for the decreasing cluster centre) to 0.74 (for the increasing cluster centre). These events have hardly any effect on the general vegetation trends.

In contrast to the hemispheric uniform temperature trends, the annual mean precipitation (P_{ann}) varies at regional level. P_{ann} increases in large parts of North America and in central northern Asia (ca 70-110°E, 45-60°N). In the northern hemispheric monsoon regions, precipitation decreases, particularly in the north African monsoon domain and the southern rim of the Himalayas.

355 In the southern hemisphere, P_{ann} increases in the monsoon region and the extratropics with the exception of Australia showing a decline in precipitation at the eastern coast and no change in the central and western parts (south of 22°S). The signal in the continental monsoon domains is well in line with the orbital monsoon hypothesis (Kutzbach et al. 1981) that directly links an increase in summer insolation with a strengthening of the summer monsoons and vice versa. P_{ann} also increases in northern Brasil and along the western coasts of the South American continent and South Africa. Precipitation does not change much north of 60°N, which implies that the shifts in vegetation distribution are rather controlled by temperature changes in this region.

To test the most important climatic driver in the different regions, we performed a constrained ordination via the redundancy analysis function “rda” implemented in the “vegan” package of R (Oksanen et al, 2018).

Fig. 4 summarizes the results, qualitatively showing the main contributors (T_{warm} vs. P_{ann}) explaining the variance in the vegetation change. The temperature of the coldest month explains only a small part of the variance in an initial RDA and was thus excluded in the final RDA.

The RDA result confirms that temperature is the most important variable constraining the vegetation change in the high northern latitudes, with the exception of East Siberia where the decrease in forest can only be explained by the combined effect of P_{ann} and T_{warm} . The shared variance of both variables is higher than the unique contribution by T_{warm} . Also, in most other regions, both variables jointly explain the vegetation dynamics during the Holocene (not shown), but in most regions, precipitation contributes most (Fig.4). Variations in T_{warm} are only important in the maritime areas of the northern hemisphere, where changes in vegetation are generally very small (i.e. cluster center shows no change), in Western Canada, Central-West Brasil, Patagonia, equatorial Africa and around the Tibetan Plateau.

This pattern of climate drivers resembles the findings by Seddon et al. (2016) analysing the effect of cloudiness, temperature and precipitation on recent vegetation productivity based on satellite data. In their study, temperature is also the most influential factor in the high northern latitudes, whereas elsewhere, precipitation explains most of the vegetation variability. In North Africa and the northern part of South America, the climatic driver is not as pronounced as in our RDA analysis, but the results can only be compared to a limited extent, as the modern data are influenced by land use that is excluded in our analysis.

Although the shared variance accounts for the largest part, precipitation in all monsoon regions explains much more of the vegetation change than T_{warm} . Given the uniform hemispheric trend in temperatures, precipitation is the factor that imprints the regional variations on the vegetation signal. We therefore focus on analysing the precipitation change as the main driver of the global vegetation trend rather than disentangling the temperature signal whose sign is directly forced by the Holocene insolation change. The simulated vegetation distributions for 2.15ka b2k and 8ka b2k are displayed in the Appendix A. Reference maps for the 2.15ka b2k climate states can be found in the Appendix B.

5. Regional insights into Holocene vegetation changes and the controlling factors of the precipitation change

5.1 The northern hemispheric monsoon domain

In both hemispheres, warm season precipitation is the most important driver of the Holocene precipitation signal (Fig.5). For the global monsoon domains, this is well in line with the orbital monsoon hypothesis (Kutzbach, 1981). The rise in northern hemispheric summertime insolation during the early Holocene leads to a strengthening of the thermal gradient between the continents and the oceans, enhancing the monsoon circulation and increasing the summertime precipitation over land. Climate model simulations and reconstructions indicate an increase in moisture availability in the monsoon margin regions and an inland penetration of the vegetation belts (Jolly et al., 1998; Hély et al. 2014; Feng et al., 2006; Zhao et al., 2009, Metcalfe et al., 2015). In the southern hemisphere, the decrease in insolation generally leads to a weakening of the monsoon circulation over land and the opposite effect on precipitation and vegetation (Wang et al. 2014 and references therein).

To assess the change in monsoon strength, we calculated the global monsoon area and the global monsoon precipitation following Zhou et al. (2008) with the modifications of Liu et al. (2009) (cf. methods).

According to our model, the northern hemispheric monsoon region is **expanded** by more than 25% on the continents at 8ka compared to 2.15ka b2k (Tab.3, Fig. 6). The monsoon precipitation is increased by 40%
405 over land. With a northward expansion of about 600 km (up to 22°N), the North African monsoon area is the most enlarged monsoon domain during 8ka b2k. This coincides with a vegetation expansion into the Sahara at 8ka b2k. In the extended monsoon area, raingreen shrubs are increased by up to 30% (Fig. 7e) and the grass fraction by up to 60%, reflected by the cluster centres with strong decreasing trend over the Holocene (cf. Fig 2). Raingreen shrubs populate the entire western Sahara and reach a coverage of about 10% in large
410 parts, going up to 20% in the coastal regions at 8ka b2k (Fig.7e, Fig A2 in the Appendix A). In the Eastern Sahara they are spread out **substantially** to only about 24°N. At the northwestern monsoon boundary, the forest cover fraction is decreased during 8ka b2k. This is related to the colder northern hemispheric winter climate during the mid-Holocene leading to unfavourable growth conditions for tropical trees in these grid-cells. The temperature of the coldest month is around 15.5°C which is a fixed bioclimatic limit for tropical
415 trees in the model. With the continuous boreal winter warming towards PI, cold season temperature more and more exceeds this limit leading to an establishment of tropical deciduous trees. The overall non-vegetated area north of 15°N is diminished by 22% in the mean at 8ka b2k, ranging up to 82% in the Central Sahel.

Thus, the vegetation changes in our model are significantly stronger and more expanded into the Sahara than in other recent global climate model simulations (e.g. Rachmayani et al. 2015, Hopcraft et al., 2017).

420 In the core monsoon domain (0-15°N), total vegetation cover is not increased during **early** mid-Holocene, but tropical evergreen trees are much more widespread (Fig. 7a). In the course of the Holocene these forested areas are replaced by grassland. Tropical Eastern Africa, experienced an increased total vegetation cover during **early** mid-Holocene that is determined by an enhanced monsoon related moisture flux convergence in spring **in the model** (not shown) leading to an expansion of grass covered areas. **This is in line with general
425 higher early- and mid-Holocene moisture levels in large parts of Tropical Eastern Africa as shown e.g. in lake level reconstructions (see overview by Tierney et al., 2011).**

In the other northern hemispheric monsoon domains, the vegetation change is more regionally confined. The East Asian monsoon area is expanded to the northwest into Western Mongolia by up to 800km at 8ka compared to 2.15ka b2k (Fig.6). This leads to an increased in total vegetation cover by up to 27% per grid-
430 cell, mainly due to an expansion of the grass cover (Fig.7g). The cover fraction of the extratropical deciduous and evergreen trees is also increased in total, although in several grid boxes it is only one tree PFT being replaced by the other. Pollen based-reconstructions reveal a regionally strongly increased tree cover fraction during the mid-Holocene, particularly in the middle and lower reaches of the Yellow River (Ren, 2007, Tian et al., 2016). This is also the area in which our model indicates the largest changes in tree cover.

435 Over the course of the Holocene, the vegetation belts retreat back to their modern position, coincident with the monsoon retreat, so that the grass fraction at the eastern rim (ca. 117-120°E, 41-46°N) increases towards 2.15ka b2k at the expense of the tree cover fraction (Fig. 7c,d).

The South Asian monsoon is expanded at its northwestern rim by about 1-2 grid-cells (ca. 200km) into western South Asia and onto the Tibetan Plateau (Fig.6), leading to a bare soil fraction decreased by up to 440 50% at 8ka b2k. The enhanced monsoon coincides with a widespread increase in area covered by raingreen shrubs, extratropical evergreen trees and grassland, in particular on the Tibetan Plateau and along its eastern margin, along the Gulf of Kutch, and in the Thar desert at **early** mid-Holocene (Fig.7). Tropical evergreen forest are decreased along the western coastal region of South Asia due to temperature limitations in the cold season. Our model reveals less vegetation in Central South Asia (17.5-21.5N, 75-82.5E) during **early** mid-445 Holocene, driven by a tropical evergreen forest cover decreased by up to 50 % and a C4 grass cover reduced by up to 35%. Partly, these PFTs are replaced by raingreen shrubs and C3 grass at 8ka b2k, reflecting a cooler winter climate. **This region is close to sites at which reconstructions reveal a less-humid climate and more open landscape during early mid-Holocene compared to late-Holocene (Chauhan et al. 2013).** Our model shows rather decreased precipitation **levels (less significant)** at 8ka summers, caused by a high 450 pressure anomaly in the lower atmosphere above South Asia and Indochina (Fig.8d). The anticyclonic atmospheric flow around this anomaly exhibits low level easterly wind above southern India (**Fig. 8c**), inhibiting the moisture flux to Central South Asia. Furthermore, the high pressure anomaly is associated with strong subsidence (Fig.8a). The monsoon flow stretches more to the north at 8ka compared to 2.15ka b2k. All these factors lead to a decreased moisture availability in this region, worsening the climatic conditions 455 for the establishment of vegetation in the model at **early** mid-Holocene.

The reduction in the South Asian monsoon strength during the Holocene also affect the vegetation on and around the Tibetan Plateau. At 8ka b2k, extratropical evergreen forests are more widespread along the northeastern and southwestern edge (Fig.7c). On the Tibetan Plateau, grass cover is strongly increased compared to 2.15ka b2k (up to 40%), owing to the higher summer precipitation (Fig. 8e). **This pattern of 460 changes is in line with pollen-based vegetation reconstructions (e.g. Herzschuh et al., 2010, vanCampo et al., 1996).**

The North American monsoon area is expanded south- and westward by **approximately** 200km at **early** mid-Holocene (Fig.6). Summer **precipitation** is substantially increased above the ocean, in Central America and the north western part of the South American continent, **similar as seen by reconstructions (Metcalf et al., 465 2015).** **The increase in precipitation leads** to a larger area covered by tropical evergreen forests in the model. In the course of the Holocene, tree cover reduces and is replaced by grass (Fig.7).

5.2 The southern hemispheric monsoon domain

In total, the southern hemispheric **monsoon** domain is enlarged by 8% and receives 7% more precipitation at 470 8ka compared to 2.15ka b2k. However, the continental monsoon area is smaller by 5% and precipitation level only reach 89% of the 2.15ka values (Tab.3). Thus the changes in the strength of the monsoon during the Holocene are much smaller in the southern hemisphere than in the northern hemisphere. For example, the monsoon area in South America is reduced by only a few grid cells, mainly in the Western part (Fig.6) at **early** mid-Holocene. Nevertheless, annual precipitation in nearly the entire continental monsoon area is

475 lower at 8ka than at 2.15ka b2k (Fig.3), due to a generally northward displaced Intertropical Convergence Zone and diminished uplift of moist air over the central continent during the monsoon season (cf. Fig. 9). Both is also mirrored in the slightly increased precipitation on the north-easternmost continent (i.e. Northeastern Brazil and the Caribbean region). This pattern is supported by reconstructions and other simulations (Maksic et al. 2018; Prado et al. 2013a and b, Cruz et al. 2009) for the mid-Holocene.

480 The reduced precipitation has a minor effect on the total vegetation cover that is only slightly decreased along the foot of the Bolivian Andes. The vegetation in the Amazon rainforest region is relatively stable, but forest is shrunk at its southern edge during **early** mid-Holocene, confirming the picture revealed by reconstructions (e.g. Mayle et al., 2000, Mayle and Power, 2008, Rossetti et al, 2017). Outside of this region, it is rather the vegetation composition that changes. The cover fraction of tropical evergreen trees is
485 increased by up to 86% in a broad band in southeastern Brazil (Fig.7a), which is overcompensated by a diminished fraction of extratropical evergreen trees and partly also extratropical deciduous trees. Thus, the total tree cover in this region is decreased and replaced by grassland at 8ka compared to 2.15ka b2k, indicating a decreased Atlantic rainforest and more open vegetation similar as proposed in reconstructions (e.g. Ledru et al., 2009). In the model, this is the effect of a climate characterised by warmer winters, but that
490 is still not moisture-limited in the monsoon domain.

On the central continent (ca. 15-30°S, along 60°W), extratropical and tropical evergreen tree covers are diminished by up to 22% leading to less forested area during **early** mid-Holocene. In this region, grass cover is slightly enhanced during 8ka b2k. This pattern of forest transitions fits well to a recent palaeo-record synthesis revealing the most significant Holocene transitions in vegetation for southwestern Amazonia and
495 southeastern Brazil, induced by a less intense South American summer monsoon (cf. Smith and Mayle, 2018).

The precipitation on the South African continent is diminished in the entire monsoon domain during **early** mid-Holocene austral summer (Fig. 9e) as well as the annual mean (Fig.3). This can mostly be attributed to the less powerful monsoon circulation at 8ka compared to 2.15ka b2k. The Angola low is weakened and the
500 South Atlantic high is shifted northwards (Fig. 9d), reducing the moisture flux to the continent and thereby decreasing precipitation (cf. Vigaud et al. 2009). The relatively weaker South Indian high induces an anticyclonic circulation with a northerly wind anomaly along the coast and landward low-level winds (Fig. 9c). This leads to increased precipitation over the ocean and decreased precipitation over land at 8ka b2k. The ITCZ is weakened, coinciding with reduced uplift (Fig. 9a) and inhibited convection.

505 The South African monsoon area is reduced by about 200km in its northwestern part (ca. 3.5-5.5°S, 14-24°E) at 8ka b2k (Fig.6). This is related to the fact that the West African monsoon dynamics also imprint on the Tropics in the southern hemisphere (up to ca. 10°S), probably due to easterly moisture-bearing trade winds off the Indian Ocean (e.g. Burrough and Thomas, 2013; Vincens et al., 2005). As a result, the tropical region not only experiences an increase in summer precipitation due to the intensification of the South African
510 monsoon over the course of the Holocene, but also a decrease in winter precipitation due to the weakening of the West African monsoon and easterly flux from the Indian Ocean (Fig.8). Therefore the annual range of

precipitation in this region is reduced during **early** mid-Holocene, not fulfilling the criteria of monsoon areas used in this study.

The overall result of the change in precipitation is an increased cover fraction of tropical evergreen trees (up to 16%) in parts of Central (12-20°S, 20-29°E) and Eastern (12-14S, 31-36°E) Africa at 8ka b2k (Fig.7a). Over the course of the Holocene, tree cover reduces and is replaced by grassland, in line with the tropical regions on the north African continent. South of 15°S, the vegetation change is more complex. The total vegetation is decreased in three north-south orientated stripes, mainly due to a decreased grass cover at 8ka b2k. East of 30°E, extratropical trees are reduced during **early** mid-Holocene and partly replaced by grassland. West of 30°E tropical evergreen trees strongly recede over the course of the Holocene and give way to raingreen shrubs and C4 grass. This is probably also related to the bioclimatic limit for the temperature of the coldest month needed for the establishment of tropical trees in the model (Tab.1), that is at some point in time no longer met in the cooling winter climate in the southern hemisphere during the Holocene. Southern Africa experiences rather an increase in C3 grass and extratropical trees from 8ka to 2.15ka b2k, while raingreen shrubs slightly decrease (Fig.7e).

Reconstructions show a diverse picture, but rather show more humid vegetation types at mid-Holocene and aridification afterwards (Olago, 2001; Burrough and Thomas, 2013 and references therein). This may at least partly be related to the fact that the vegetation distribution in Southern Africa is strongly controlled by fires, maintaining grassland and savannas despite sufficient rainfall for the establishment of forest (Bond et al., 2003, Bond, 2008). JSBACH includes a fire model, but the effect of fires may be underestimated in the Southern African region. At least, Southern African vegetation seems to be highly correlated with climate in our simulation.

The Australian monsoon region is shrunk at its south-eastern rim by **approximately** 200km at 8ka b2k (Fig.6). The high pressure anomaly above the continent leads to an anticyclonic flow with low-level westerly winds south of the monsoon domain (ca. 22°S) and easterly winds in the monsoon region (Fig.9c). This results in a weakened monsoon-related moisture flux and less precipitation during austral summer at **early** mid-Holocene **in the model (Fig.9e). Other climate model simulations reveal contradictory results with some even showing a stronger monsoon and increased precipitation at mid-Holocene (eg. Liu et al., 2004). In MPI.ESM, the total vegetation is decreased by up to 19% in the North Australian monsoon region during early** mid-Holocene, mainly due to a decreased C4 grass cover and less tropical evergreen trees. On the southern edge of the monsoon domain, the tropical evergreen tree cover is raised by up to 20% at the expense of extratropical evergreen trees and raingreen shrubs (Fig.7). This switch between different woody PFTs is mainly determined by the warmer southern hemispheric winter climate during mid-Holocene.

5.3 Extratropical North America

According to the model, the total vegetation cover in large parts of North America increases from 8ka to 2.15ka b2k (Fig.2). Most prominent is the vast expansion of trees, ranging from the southern Great Plains to the Canadian interior plains and the northeastern Rockies that is also seen in reconstructions (Grimm et al.,

2001). In the model, this is manifested by a rise in the extratropical evergreen tree fraction by up to 24%
550 west of 100°W and a rise in extratropical deciduous tree fraction by up to 45% east of 100°W (Fig.7). At
early mid-Holocene, most of this region is covered by grassland or bare soil. In the southern Great Plains,
raingreen shrubs are more widespread at 8ka, these are replaced by grass during the Holocene.

The precipitation response to the orbital forcing in the model is a complex mixture of several interacting
processes. In total, the Great Plains receive less precipitation during **early** mid-Holocene (Fig.3a). Fig. 5
555 shows that the summer months contribute most to the annual mean precipitation change. These are also the
months in which the northern hemispheric monsoon dynamics are most vigorous and the orbital forcing has
the strongest imprint on the monsoon circulation. Summer monsoon precipitation coincides with a strong
condensational latent heat release, affecting also the tropospheric circulation in the extratropics (e.g. Gaetani
et al., 2008; Nakanishi et al., 2021). The subtropical anticyclones are stated to be related to Kelvin and
560 Rossby wave responses to the heating in the monsoon rainband (Matsuno 1966, Gill 1980, Rodwell and
Hoskins, 2001, Trenberth et al., 2000). These anticyclones cover about 40% of the Earth surface (Rodwell
and Hoskins, 2001) and are therefore a major player in the global circulation of the atmosphere and oceans
and the atmospheric teleconnection pattern. This suggests that the strong changes in the northern hemispheric
monsoon systems also play a decisive role in modifying the extratropical vegetation change over the course
565 of the Holocene. Fig. 8d displays the mean change in the atmospheric standing waves during June, July and
August (JJA). The Bermuda high is strengthened in the core at 8ka b2k leading to a rerouting of the moisture
transport from the Gulf of Mexico along the Atlantic coast, thereby enhancing the precipitation along the
Appalachian mountain range (cf. also Williams et al., 2010). In this region of raised precipitation levels, the
extratropical deciduous tree fraction and with it the total forest cover is even higher during the mid-Holocene
570 than at 2.15ka b2k.

The moisture flux is more divergent in the Great Plains and the atmosphere is generally drier, inhibiting
convection and rainfall. The intensity change of the Bermuda high has so far been seen as one possible
explanation for the change in Holocene vegetation pattern seen in pollen-based reconstructions (e.g. Grimm
et al., 2001 and references therein). In addition, the model shows a rainfall-reducing subsidence anomaly
575 over the Great Plains south of 45°N at 8ka b2k (Fig.8a) that could at least partly be attributed to a
Rodwell&Hoskins-like Rossby wave response to the enhanced updrafts in the North American summer
monsoon (Harrison et al., 2003). The northward shifted upper level westerly Jet during the mid-Holocene
(Fig.8b) coincides with a northward replacement of the storm tracks, which probably results in less transient
eddy transport to the northern Great Plains at 8ka b2k (Seager et al., 2014). The model also reveals less
580 precipitation during spring and therefore limited evaporation during mid-Holocene summer (not shown),
reducing water recycling and the northward transport of moisture in the Great Plain low level Jet. The
influence of evaporation on vegetation change may also explain the relatively large proportion of vegetation
variance explained by the temperature of the warmest month (cf. Fig.4).

The more La Nina like sea surface temperature pattern (Fig. B1, Appendix B) may also contribute to the
585 drier surface conditions in the Great Plains during mid-Holocene, although the pattern is probably less
pronounced in the model than indicated by reconstructions (Shin et al, 2006). Which of these mechanism is

actually the main driver in the Holocene precipitation change can not be disentangled in the model without performing a complex set of sensitivity experiments. **Teleconnections with the northern hemispheric monsoons could strongly have contributed to the vegetation changes in the North American extratropics.**

590 The reduced precipitation above the Canadian interior Plains and northeastern Rockies (ca. 120-90°W, 45-60°N) during the mid-Holocene, accompanying the decreased extratropical evergreen tree fraction there (Fig.7c), is probably related to the strengthened and westward shifted North Pacific subtropical high (Fig.8d). This leads to enhanced northerly winds transporting rather dry continental air inland. The subsidence is increased in large parts during mid-Holocene summers (Fig.8a), further limiting the
595 convection. At least partly, the differences in subsidence can be contributed to the change in dynamics, coinciding with the northward displaced westerly jet at 8ka b2k. This also affects the region along the northern Rockies and Alaska (45-60°N, east of 120°W). Here, the westward shift in the north Pacific High reduce the moisture flux to the area, resulting in less moisture convergence in the region. Therefore precipitation is decreased at 8ka b2k in the annual mean that leads to less total vegetation, mainly due to a
600 reduced grass cover (up to 35%) at mid-Holocene. The fraction of extratropical evergreen trees decreases over the course of the Holocene and is replaced by cold-resistant shrubs and grassland (Fig.7), probably reflecting the typical Holocene signal in the Taiga-Tundra margin.

5.4 Taiga-Tundra dynamics in the high northern latitudes

605 The taiga-tundra region and the dependence of the vegetation signal on temperature changes has been widely discussed before (e.g. Bigelow et al., 2013, MacDonald et al. 2000, Wanner et al., 2008). Our model is in line with previous results and shows no new insights regarding the long-term trend. Therefore, this region will not be explored in detail in this study. The main signal is the decrease of tundra along the Arctic coast and the strong decrease of boreal forest further south, both reflecting the southward shift of the boreal vegetation
610 zones over the course of the Holocene (Fig.2 and Fig.7), also found in previous transient vegetation simulations (e.g. Braconnot et al., 2019). The main forcing is the cooling of the summer climate and the warming of the winter climate during the Holocene (Fig.3) that is induced by the seasonal changes in insolation. In North America, the boreal forest area is reduced by 25.5% and the mean northern treeline (here ad hoc defined as isoline of a zonal mean tree cover fraction of 10%) moves **approximately** 2° of latitudes to
615 the south over the course of the Holocene (Fig.7). Europe experiences less change. The boreal forest area shrinks by 5.9% and the mean northern treeline shifts southward by **approximately** 0.5° of latitudes. In Asia, the boreal tree cover fraction reduces by nearly one third from 8ka to 2.15ka b2k and the northern treeline shifts southward by **approximately** 3° of latitudes. The boreal trees are mainly replaced by C3 grass and cold-resistant shrubs.

620 However, two regions stand out due to an increase in the bare soil fraction during the Holocene. These are an area in mid-Siberia (about 100-140°E, north of 58°N) and a region in northern Canada (about 100-120°W, north of 60°N). The increase may be a model specific response. During the mid-Holocene, modelled winter temperatures are too cold to allow the establishment of extratropical evergreen trees in these regions (limited

by a mean temperature of the coldest month below -32.5°C). With the increase in wintertime insolation and a
625 successive winter warming over the Holocene, this limit no longer applies, and extratropical evergreen trees
replace the extratropical deciduous trees (Fig.7). However, the net primary productivity (NPP) of the
evergreen trees is too low to build up their living tissues completely. This, by definition, increases the
simulated bare soil fraction.

In another inland region in Siberia (ca. $70\text{-}110^{\circ}$, $48\text{-}58^{\circ}\text{N}$), the increase in forest cover south of the modern
630 treeline is disproportionately higher than in the other regions. This is in line with a substantial increase in
precipitation in this region (Fig.3) over the course of the Holocene. At mid-Holocene, the westerly wind jet is
shifted to the north and is weaker (Fig.8b). The coincident subsidence results in decreased summer
precipitation at 8ka b2k, similar to the changes in Canada.

In line with the results by Braconnot et al. (2019), the worldwide strongest changes in tree cover fraction
635 occur north of 60°N , while tree cover changes in the rest of Eurasia are rather small.

5.5 Extramonsoonal Australia and South America

The northern part of Australia (north of 22°S) is characterized by an increase in total vegetation during the
Holocene related to the strengthening of the Australian monsoon (cf. 4.2.2). South of 22°S vegetation
640 decreases during the Holocene (Fig.2a). At 8ka b2k, extratropical evergreen trees are more widespread.
Along the eastern coast, where proxy records and our model reveal wetter mid-Holocene conditions (e.g. Mc
Glone et al., 1992), the cover fraction of extratropical forests is increased by up to 16% (Fig.7). In central
Australia, forest fraction is raised by up to 6%. The grass cover is enhanced in large parts of the (today)
rather dry continental interior. Over the course of the Holocene, these grasslands and forested area are partly
645 replaced by raingreen shrubs.

These vegetation changes are related to a slightly wetter climate during the mid-Holocene. In large parts of
the continent, the precipitation difference between 8ka and 2.15ka b2k are largest during November
(continental interior, Fig.5), October (eastern coast) and June (parts of the southern coast). During these
months, the subtropical ridge is weaker at 8ka b2k (also seen in Fig. 10d). This leads to less subsidence (Fig.
650 10a) and favours convection and precipitation (Fig.10c). In addition, the moisture influx from the western
Pacific is increased. The enhanced low pressure system over northwestern Australia during austral spring
causes monsoon-like conditions and suggest an earlier onset of the Australian summer monsoon at 8ka b2k,
coinciding with an enhanced precipitation level in entire Australia. The warmer sea surface temperatures in
the Indonesian ocean may additionally favour increased winter precipitation during mid-Holocene (Verdon
655 and Franks, 2005). Furthermore, a more La Nina sea surface pattern in the tropical Pacific is known to
increase moisture level in arid Australia (Nicholls, 1992; Quigley et al., 2010, Barr et al., 2019).

The Gran Chaco and Pampas regions east of the South American Andes experience an increase in vegetation
during the Holocene, mainly due to an increase in grass cover **in the model** (Fig.7). On the Southern
continent, extratropical evergreen trees cover up to 27% less area at 8ka compared to 2.15ka b2k. These
660 vegetation changes are probably related to the increase in both, austral wintertime and summertime

precipitation during the Holocene. Large parts of the region get precipitation mainly due to the moisture influx in the north-easterly branch of the Subtropical Atlantic High. During 8ka b2k austral winters, this anticyclone is shifted equatorwards leading to diverging easterly wind anomalies in the lower level and a decrease in moisture flux convergence (not shown). The upper tropospheric westerly jet is squeezed and intensified along the core (ca.30°S), a pattern that is underlined by Holocene reconstructions (Lamy et al., 2010). The changes in Westerlies probably enhance the subsidence in the lee of the Andes. The subsidence is furthermore enhanced by the updraft of the Afro-Asian monsoon that leads to a strong divergence anomaly in the upper tropospheric velocity potential (Fig. 11) and accordingly to a strong convergence anomaly above South America during austral winter, triggering subsidence. Thus, the change in monsoons are also a key-driver of the South American vegetation changes.

Fig.5 indicates, that the precipitation difference between 8ka and 2.15ka b2k in this region is largest during the month November and December. During austral summer, a deep continental low evolves above the southern American continent that additionally deflects the easterly winds to the south so that the wind is channeled between the Brazilian Plateau and the Andes. This low level jet transports huge amounts of moisture southwards at present (Marengo et al., 2004; Garreaud et al., 2009). During mid-Holocene Nov. and Dec., this low is weakened, in line with the weakened South American monsoon (Fig.9), whose outflow touches this region at 2.15ka but not at 8ka b2k. The model indicate a southwind anomaly in the lower troposphere (i.e. a reduced low level jet), diminishing the inflow from the monsoon area (not shown).

Overall, the model reveals that vegetation changes south of 60°N during the Holocene occur predominantly in monsoon regions, their margins and in extratropical regions that are connected with the monsoons via teleconnections. Thus, our model results point to a strong effect of the monsoon-related changes in atmospheric circulation on the global Holocene vegetation dynamics outside the high northern latitudes. These results confirm earlier findings of monsoon-extratropical teleconnections (e.g. Rodwell and Hoskins, 2001, Gaetani et al., 2008; Nakanishi et al., 2021) and stress the importance of the monsoon system for the global climate and ecosystems. At least during the Holocene, the monsoon systems seemed to be the key player in the climate and vegetation dynamics.

6. Are the Holocene vegetation changes as linear as they appear in our global analysis?

The overview of the patterns of clustered vegetation trends reveals rather linear transitions between different PFT groups during the Holocene (Fig.2). This appears to be consistent with the assumption that large-scale vegetation change and climate change directly follow the subtle, approximately linear orbital forcing over the last 8000 years. In addition, it agrees with the analysis by Mottl et al. (2021) that reveals a linear, steady change in the vegetation composition between 8 ka b2k and 4.2 ka b2k before early land use sets in (which is excluded in our simulations in the millennia before 2 ka b2k). However, reconstructions also report of rapid, or abrupt, vegetation changes in different regions of the world. The most prominent example is the expansion of the Sahara from a rather 'green Sahara' to its current extent some 5500 years ago. Indicators of regionally (mainly in the western part) rapid changes include the rapid increase in dust deposition found in marine

sediments core off the Western Sahara coast (e.g. deMenocal et al, 2000). Abrupt regional vegetation changes, at least for individual taxa, are also suggested to exist in Central Asia (Zhao et al., 2017), the high northern latitudes (MacDonald et al. 2000), parts of Northern America (e.g. Marsicek et al., 2013; Shuman et al., 2009; Foster et al, 2006) and Europe (Giesecke et al., 2011, Seddon et al., 2015). How rapid these changes were and whether these changes only reflect local phenomena is still a matter of discussion. Mayewski et al. (2004) identified several phases with rapid global climate changes during the Holocene which might have forced changes in vegetation composition. Ecosystems can respond quite slowly to changes in the local environmental conditions, but can also change rapidly when the conditions approach critical thresholds (Scheffer et al., 2001). In addition, initially slow changes can be amplified by feedbacks in the climate system, giving rise to non-linear responses to the orbital forcing (Williams et al., 2002).

The cluster method used to group the globally-wide trend in main PFTs in this study is not designed to properly identify rapid changes. The aggregation of the PFTs in main PFT groups complicates the occurrence of rapid changes as these changes would have to be very extensive and accompanied by very strong climatic changes for an entire PFT group to collapse. Furthermore, the global view masks regionally appearing transitions. To detect non-linearity in the Holocene vegetation change, we suggest evaluating the relative change (R) in the individual PFT fractions (see Appendix C). The change in the individual PFT fraction between two consecutive time-steps is compared to the maximum possible change during the period of interest. The noise time-series are filtered by a Butterworth-Filter with a cut-off frequency $1/F = 500$ years. As measure for rapid changes, the maximum R_{max} of the absolute value of R is taken.

Fig.12 shows the global patterns of R_{max} for the individual PFTs. In total, six regions can be identified that indicate rapid changes in PFT cover fractions, i.e. shifts that exceed the linear trend by a factor of up to 5, given the low pass filter of 500 years. These regions include areas in the high northern latitudes and the monsoon margins in each hemisphere. The areas coincide with regions where the cluster method also shows relatively strong changes, except for the region of South America in which the rapid changes occur within the forest PFTs.

In northern Canada and eastern Siberia (region 1 and 2), rapid shifts mainly occur between the extratropical deciduous trees and extratropical evergreen trees, and, to a weaker extent, in the bare soil fraction. The main processes for the transition are the same as discussed above in section 5.4. A typical example of such a transition in region 1 and 2 is depicted in Fig 13a for a grid box in eastern Siberia. During the first millennium, extratropical evergreen trees cannot exist, because the temperature of the coldest month is too low. This bioclimatic limit mimics the dependence of extratropical trees on moisture in the high latitudes, because there are no moisture-related bioclimatic thresholds defined in JSBACH. In a very cold, and thus dry, climate, the deciduous trees have an advantage over evergreen trees. Extratropical evergreen trees and extratropical deciduous trees can coexist after about 6 ka b2k. The time scale of the transition from around 6 ka b2k to some 3.5 ka b2k is governed by the small difference in NPP between the tree PFTs. During the last millennium, rapid fluctuations within a few decades between extratropical evergreen and deciduous trees are found in the simulations. These are induced by fluctuations in the temperature of the coldest month, which may fall below the bioclimatic limit at which extratropical evergreen trees can exist in some years. The

extratropical evergreen trees decay within a few decades, and this decay is governed by the time scale of mortality of 50 years prescribed in JSBACH. The extratropical evergreen trees eventually recover at a slower pace which follows the difference in NPP development and which is not explicitly determined by the time scales prescribed in JSBACH.

740 In the Sahel-Sahara transition zone (region 3), the model reveals rapid changes which are also found in some, but not all, pollen records (e.g. Kröpelin et al., 2008, Shanahan et al., 2015) or in Saharan dust transport (e.g. deMenocal et al., 2000). The latter change can directly be linked to rapid landscape changes mainly in the western part of the Sahara (Egerer et al., 2016). **The simulated bare soil fraction rapidly increases at the expense of grass cover and, to a much lesser extent, of raingreen shrubs (see Fig. 12 and Fig. 14, which**
745 **provides a zoom into the region 19.59N-25.18N and 15.0W-5.63W). Bioclimatic limits affect the vegetation change only in the appearance of tropical tree types. Because early mid-Holocene insolation in the tropics is lower than today during winter, the simulated temperature of the coldest month falls below 15.5 °C at the beginning of the simulation. This temperature threshold represents the bioclimatic limit for frost-free climate, which is the prerequisite for the existence of tropical trees. During the Holocene the winter temperature**
750 **increases, and tropical trees emerge as seen in the in the grid cells centred at the latitude of 25.18 N (the first row of grid cells in Fig. 14) and east of 13.13 W (the second grid cell from the left in the first row of grid cells). The rapid transition between simulated tropical evergreen trees and tropical deciduous trees seen in some grid cells (e.g. Fig. 13b representing the grid cell centred at 19.59 N, 7.5 W) is not triggered by crossing a bioclimatic limit. The plant functional types ‘tropical evergreen tree’ and ‘tropical deciduous tree’**
755 **have the same temperature requirement in our model. They differ, however, in the parameter values implicit in the rate of carbon fixation computed for the photosynthesis. Therefore, it is the difference in NPP, which governs the transition between these PFTs. This transition can be interpreted as a non-linear response to the steady decline in moisture level caused by the retreating summer monsoon during the Holocene, and it generally takes several centuries in our simulations. Similar is valid for the transition between grass and bare**
760 **soil fraction. Also in this case, it is the decrease in NPP which leads to an increase in bare soil fraction. The simulated shift between tropical tree PFTs appears to be more rapid than the shift between grass and bare soil. The amplitude of the shift in tropical tree cover is, however much smaller than the decrease in grass and increase in bare soil fraction. A more detailed discussion of simulated and reconstructed vegetation changes in the Sahara-Sahel region is in preparation for a subsequent study.**

765 In of South Asia and in parts of Southeast Asia (region 4), various combinations of rapid changes are found in the simulation. In South Asia, raingreen shrubs decrease rapidly and are replaced by tropical evergreen trees and grasses. As an example, the vegetation dynamics of one grid cell from this region are depicted in Fig. 13b. The transition between the different PFTs is discussed in section 5.1. As in the high northern latitudes, the pace of the transition at the mid-Holocene is governed by the differences in NPP of the PFTs
770 involved. The time scales of allocation and mortality of the competing PFTs differ. They range from 30 years for tropical trees to 12 years for raingreen shrubs. The bioclimatic limit which restricts the occurrence of tropical trees is imposed by the temperature of the coldest month (see section 5.1). In the time period from around 7 ka b2k to 5.5 ka b2k this limit is frequently crossed. Because the time scales of allocation and

mortality are shorter than that of the PFTs at high northern latitudes, the change between PFTs appears to be more rapid, and the amplitude is larger.

The climatic changes that drive the rapid swings in PFTs in the South American region (region 5) and in the South African region (region 6) are discussed in section 5.2. The governing time scales of PFT dynamics and the bioclimatic limits involved are the same as for tropical Asia (region 4).

780 7 Summary and Conclusion

Using a cluster analysis technique on a Holocene simulation for the period 8ka to 2.15ka b2k with MPI-ESM, the main global natural vegetation trends during the Holocene are identified and discussed in the context of the climate change over the Holocene. A redundancy analysis reveals that precipitation is the main climatic driver leading to regional differences in the vegetation change outside the high northern latitudes.

785 The model is well in line with the overall trends reported by pollen-based vegetation reconstructions. These main trends and their atmospheric drivers in the model are summarized in Fig.15. They include:

a) The southward retreat of the tundra and northern treeline by up to 3° of latitudes, expressed by a bipolar change pattern for all PFT groups with a decrease on the northern site and an increase further south. Asia faces the major change in the model by a reduction of the boreal tree cover by nearly one third since the early mid-Holocene. This transition is mainly related to the seasonal temperature trend driven by the change in insolation during the Holocene. Boreal winter temperatures rise, summer temperatures decline, reducing the growing season and stressing the vegetation in the high northern latitudes. In the polar regions, climate becomes even too harsh for vegetation to survive.

b) A decrease and equator-ward retreat of all vegetation types in the northern hemispheric monsoon regions, leading to an increase in the deserts, particularly to a vast expansion of the Sahara. The mean non-vegetated area in North Africa north of 15°N increases by 22% during the Holocene, ranging up to 82% in southern Niger. The reduction in vegetation coincides with the equator-ward retreat and the weakening of the northern-hemispheric monsoon belts. Our model indicates a decrease of the continental northern hemispheric monsoon area by approximately 25% since 8ka b2k. The precipitation in the monsoon regions decreases by approximately 40% over land, leading to limited moisture availability in large regions.

c) An increase in the forest and partly the grass fraction in extratropical North America, leading to an overall increase in vegetated area over the Holocene, forced by a rising moisture level. From the model simulation we find several possible mechanisms triggering the Holocene increase in precipitation in North America. The North Pacific high is shifted westward and the westerly jet and the related storm tracks are shifted northward at mid-Holocene, leading to less moisture transport to Canada and the northern Great Plains and enhanced subsidence in large parts of the region. The Bermuda high is slightly strengthened during 8ka summer, rerouting the moisture transport along the coast and reducing the moisture flux convergence over the Great Plains. Additionally, the strengthening of the ascent in the North American summer monsoon is probably responsible for enhanced subsidence in the Great Plains at 8ka b2k.

810 d) A small increase in vegetation in the southern hemispheric monsoon regions, induced by an increase in forest, grass and shrubs (S Africa). Due to the decreased austral summer insolation during the mid-Holocene, the southern hemispheric monsoon systems are weaker at 8ka compared to 2.15ka b2k. The continental monsoon domain is reduced by 5% and the precipitation is decreased by 11%. Hence, the change in the monsoon strength is weaker than for the northern hemisphere. This is also reflected in the relatively minor
815 changes in total vegetation cover. Instead, the southern hemispheric monsoon regions experiences (strong) shifts in the woody PFT composition. The warmer winter climate during the mid-Holocene allows for the establishment of tropical evergreen trees in the model. In the course of the gradual cooling during the Holocene, winter temperatures at some point fall outside the tolerance range of tropical trees, which give way for other woody PFTs. Under the relatively humid conditions in South America, extratropical trees win
820 this competition. In South Africa, where the moisture is more limited, evergreen trees are replaced by raingreen shrubs. Australia experiences both, an increase in extratropical trees (mainly at the east coast) and in raingreen shrubs.

e) An increase in the bare soil fraction in extra-monsoonal Australia and an increase in grass and total vegetation in extra-monsoonal South America. Mid-Holocene Australian climate (south of 22°S) is slightly
825 wetter than at 2.15ka b2k due to a weaker subtropical ridge, less subsidence and a stronger moisture influx from the Pacific Ocean. During the Holocene, precipitation declines, leading to a decrease of the grass cover and extratropical trees that can not be overcompensated by the establishing raingreen shrubs. In South America, austral wintertime precipitation is decreased at the mid-Holocene due to increased subsidence and a decreased moisture flux convergence caused by the poleward shifted South Atlantic subtropical high. At least
830 partly, the stronger subsidence is related to a remote response to the enhanced North African summer monsoon. In addition, austral summer precipitation is decreased, probably due to a weaker influence of the South American monsoon, upstream of the main lower tropospheric wind field. The increase in rainfall during the Holocene leads to an increase in grass cover (Gran Chaco and Pampas region) and extratropical evergreen trees (Eastern Argentina).

835 At large scales, the global shifts in vegetation appear to follow the slow, to first order linear, orbital forcing. In some regions, however, the model simulates rapid, strongly nonlinear changes in PFTs. These regions include two domains in the high northern latitudes (Canada and mid-Siberia), and four regions in the monsoon margin area on both hemispheres (Sahel-Sahara, India & South East Asia, South America and South Africa). The rapid change in bare soil fraction in the Sahara region can be attributed to the nonlinear
840 response of the monsoon precipitation and vegetation to the orbital forcing (cf. Dallmeyer et al., 2020).

Transitions between individual PFTs triggered by changes in climate appear at a large spectrum of time scales. The transitions are governed by the difference in NPP of the competing PFTs. If the PFTs have the same time scales of allocation and mortality, then transitions can take place over many centuries. More rapid transitions in PFTs can occur within a few decades or less, if the climate moves in or out the bioclimatic
845 tolerance so that some temperature thresholds in the model are suddenly met or not fulfilled any more. These rapid changes in simulated vegetation pattern have to be interpreted with care. Bioclimatic limits are known to exist for plant species. Whether the concept of bioclimatic limits can be applied to the temporal dynamics

of PFTs has to be critically re-assessed. PFTs encompass a variety of different and diverse plant species. Because of this plant diversity within a PFT, PFTs presumably show a larger resilience to climate changes
850 than individual species do. Hence our model results should be taken as an indicator of possible rapid changes in the vegetation, rather than a precise prediction of rapid changes themselves.

This study explores the Holocene vegetation changes around the world and interprets them in terms of the Holocene climate change. The model results reveal that most of the Holocene vegetation trends seen outside
855 the high northern latitudes can be attributed to modifications in the intensity of the global summer monsoons. Due to the seasonal changes in insolation, the monsoon systems in the northern hemisphere weaken, while the monsoon systems in the southern hemisphere intensify during the Holocene. Due to teleconnections (Rodwell-Hoskins-response to the diabatic heating), the intensity change of the monsoons affect the subtropical anticyclones, whose modification in turn leads to an adjustment of the wind field over the
860 extratropical continents. In addition, the magnitude of the updrafts in the monsoon belts determines the strength of the subsidence in the extra-monsoonal areas, such that a weakening of the monsoon rainfall coinciding ascent also weakens the descent abroad. The model results thus identify the global monsoon system as the key player in Holocene climate and vegetation history and point to a far greater importance of the monsoon systems on the extra-monsoonal **vegetation dynamics** than previously assumed. The model
865 results indicate that changes in the global monsoon dynamic should always be considered as an actor in past and future climate changes, even in regions not primarily influenced by monsoons.

Appendix A: Simulated vegetation distributions for the 2.15ka b2k and the 8ka b2ka time-slices

Appendix B: simulated climate at 2.15ka

870 Appendix C: Definition of the relative change R

We evaluate the temporal change dV/dt of the simulated vegetation fraction V of a PFT relative to an overall amplitude ΔV and over the entire period ΔT considered, in our case, $\Delta T = 6000$ y. We define the relative change R by

$$875 \quad R = \left(\frac{dV}{dt} \right) \left(\frac{\Delta V}{\Delta T} \right)^{-1}.$$

We define the overall amplitude ΔV as the maximum possible change that the vegetation fraction can reach, i.e. $\Delta V = 1$, to focus on large and rapid vegetation changes. The simulated vegetation cover V varies on all time scales. Some of these variations can be attributed to fluctuations in climate which affect the net primary productivity (NPP). Variations in V also occur, when the bioclimatic limit at which a PFT can exist is
880 crossed. Then V decreases exponentially at times of decades and shorter, or it increases abruptly, if colonisable space is available, or the PFT grows at the expense of other PFTs. Examples are given in Fig. 13 a and c. This type of abrupt change, which is known to happen at species level, is likely to be a model artefact at biome level. Therefore, we use a Butterworth filter of the order 5 to separate very brief fluctuations in simulated vegetation fraction from fluctuations beyond the decadal time scale.

885 Figure C1 demonstrates the effect of filtering. In Figure C1, changes in extratropical deciduous trees in a grid box in East Siberia (the same grid box as shown in Fig. 13a) are depicted. The unfiltered time series reveals small, rapid changes at decadal time scale in the first millennia of the simulation, a large transition around 4000 ka b2k, and larger and rapid decadal fluctuations during the last millennia. When applying a Butterworth filter of order 5 with a cut-off frequency $1/F > 1/200$ y, then only the large transition around 4000 ka
890 b2k is identified as the most rapid change.

Figure 12 shows the maxima of the absolute value of R using $1/F = 1/500$ y,

$$R_{max} = \left| \frac{dV}{dt} \right|_{max} \left(\frac{|\Delta V|}{\Delta T} \right)^{-1},$$

hence we do not differentiate cases of strong increase or decrease in vegetation fraction. Using smaller or larger values of F would result in different values of R_{max} , as indicated in the example Fig.C1, but the overall
895 global patterns, specifically the regions of strongest non-linear change shown in Fig.12, would be more or less the same.

Code and Data availability

900 The primary data, i.e. the model code for MPI-ESM, are freely available to the scientific community and can be accessed with a license. In addition, secondary data and scripts that may be useful in reproducing the authors' work are archived by the Max Planck Institute for Meteorology and are accessible without any restrictions (link will be provided at the final submission).

The Biome6000 pollen-based biome reconstructions (Harrison, 2017) that we used for the evaluation of the
905 model can be downloaded from: <https://doi.org/10.17864/1947.99>.

The volcanic forcing data will be made available at <https://doi.pangaea.de/10.1594/PANGAEA.928646>.

Author contribution

AD and MC wrote the manuscript, SL ran the simulation, MS and MT provided the volcanic forcing data. MC, UH and AD planned the study and were involved in the analysis. All authors discussed the analysis and
910 the manuscript.

Competing interests

The authors declare that they have no conflict of interests.

Acknowledgements

This work contributes to the project PalMod, funded by the German Federal Ministry of Education and Research
915 (BMBF), Research for Sustainability initiative (FONA, www.fona.de). AD was financed by PalMod (Grant number: 01LP1920A). MS acknowledges funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 820047). We thank T. Kleinen (MPI-M) for his helpful comments on an earlier version of this manuscript. We acknowledge J. McConnell, J. Cole-Dai and M. Severi for providing ice-core data. We thank the referee Qiong Zhang and the anonymous referee for their constructive
920 suggestions for improving our manuscript.

References

- Bader, J., Jungclaus, J., Krivova, N., Lorenz, S., Maycock, A., Raddatz, T., Schmidt, H., Toohey, M., Wu, C.-J. & Claussen, M.: Global temperature modes shed light on the Holocene temperature conundrum. *Nature Communications*, 11: 4726. [doi:10.1038/s41467-020-18478-6](https://doi.org/10.1038/s41467-020-18478-6), 2020.
- Barr, C., Tibby, J., Leng, M.J., Tyler, J. J., Henderson, A. C. G., Overpeck, J. T., Simpson, G. L., Cole, J. E., Phipps, S. J., Marshall, J. C., Mcgregor, G. B., Hua, Q., Mcrobie, F. H.: Holocene El Niño–Southern Oscillation variability reflected in subtropical Australian precipitation. *Sci Rep* 9, 1627, <https://doi.org/10.1038/s41598-019-38626-3>, 2019.
- Bartlein, P. J. and Shafer, S. L.: Paleo calendar-effect adjustments in time-slice and transient climate-model simulations (PaleoCalAdjust v1.0): impact and strategies for data analysis, *Geosci. Model Dev.*, 12, 3889–3913, <https://doi.org/10.5194/gmd-12-3889-2019>, 2019.
- Behling, H., and H. Hooghiemstra: Neotropical savanna environments in space and time: Late Quaternary interhemispheric comparisons. In Markgraf, V. (ed.), *Interhemispheric Climate Linkages*. San Diego: Academic Press, Chapter 18, 2001.
- Berger, A. L.: Long-term variations of daily insolation and quater-nary climatic changes, *J. Atmos. Sci.*, 35, 2362–2367, 1978.
- Bezdek, J.C.: *Pattern Recognition with Fuzzy Objective Function Al-gorithms*. Plenum, New York, 1981.
- Bigelow, N.H., Brubaker, L.B., Edwards, M.E., Harrison, S.P., Prentice, I.C., Anderson, P.M., Andreev, A.A., Bartlein, P.J., Christensen, T.R., Cramer, W., Kaplan, J.O., Lozhkin, A.V., Matveyeva, N.V., Murray, D.F., McGuire, A.D., Razzhivin, V.Y., Ritchie, J.C., Smith, B., Walker, D.A., Gajewski, K., Wolf, V., Holmqvist, B.H., Igarashi, Y., Kremenetskii, K., Paus, A., Pisaric, M.F.J., Volkova, V.S.: Climate change and Arctic ecosystems: 1. Vegetation changes north of 55 degrees N between the last glacial maximum, mid-Holocene, and present. *Journal of Geophysical Research Atmospheres* 108 (D19), 8170, [doi:10.1029/2002JD002558](https://doi.org/10.1029/2002JD002558), 2003.
- Bond, W.J.: What limits trees in C₄ grasslands and savannas? *39*, 641, [doi:10.1146/annurev.ecolsys.39.110707.173411](https://doi.org/10.1146/annurev.ecolsys.39.110707.173411), 2008.
- Bond, W.J., Midgley, G.F., and Woodward, F.I.: What controls South African vegetation-climate or fire? *S Afr J Bot*, 69:79–91, 2003.
- Braconnot, P., Zhu, D., Marti, O., and Servonnat, J.: Strengths and challenges for transient Mid- to Late Holocene simulations with dynamical vegetation, *Clim. Past*, 15, 997–1024, <https://doi.org/10.5194/cp-15-997-2019>, 2019.
- Brovkin, V., Bendtsen, J., Claussen, M., Ganopolski, A., Kubatzki, C., Petoukhov, V., and Andreev, A.: Carbon cycle, vegetation, and climate dynamics in the Holocene: Experiments with the CLIMBER-2 model, *Global Biogeochem. Cy.*, 16, 1139, <https://doi.org/10.1029/2001GB001662>, 2002.
- Brovkin, V., Raddatz, T., Reick, C.H., Claussen, M., and Gayler, V.: Global biogeophysical interactions between forest and climate. *Geophys. Res. Lett.*, 36, L07405, [doi:10.1029/2009GL037543](https://doi.org/10.1029/2009GL037543), 2009.

- 960 Brovkin, V., Lorenz, S., Raddatz, T., Ilyina, T., Stemmler, I., Toohey, M., and Claussen, M.: What was the source of the atmospheric CO₂ increase during the Holocene?, *Biogeosciences*, 16, 2543–2555, <https://doi.org/10.5194/bg-16-2543-2019>, 2019.
- Burrough, S.L., and Thomas, D.S.G.: Central southern Africa at the time of the African Humid Period: a new analysis of Holocene palaeoenvironmental and palaeoclimate data, *Quaternary Science Reviews*, Volume 80, 2013, Pages 29–46, ISSN 0277-3791, <https://doi.org/10.1016/j.quascirev.2013.08.001>, 2013.
- 965 Cao .X., Tian, F., Dallmeyer, A., and Herzschuh, U.: Northern Hemisphere biome changes (>30°N) since 40 cal ka BP and their driving factors inferred from model-data comparisons, *Quaternary Science Reviews*, Volume 220, 291–309, ISSN 0277-3791, <https://doi.org/10.1016/j.quascirev.2019.07.034>, 2019a.
- Cao, X., Tian, F., Li, F., Gaillard, M.-J., Rudaya, N., Xu, Q., and Herzschuh, U.: Pollen-based quantitative land-cover reconstruction for northern Asia covering the last 40 ka cal BP, *Clim. Past*, 15, 1503–1536, <https://doi.org/10.5194/cp-15-1503-2019>, 2019b.
- 970 Chauhan, M.S., Sharma, A., Phartiyal, B., and Kumar, K.: Holocene vegetation and climatic variations in Central India: A study based on multiproxy evidences, *Journal of Asian Earth Sciences*, 77, 45–58, <https://doi.org/10.1016/j.jseaes.2013.08.005>, 2013.
- Clement, A.C., Hall, A., Broccoli, A.: The importance of precessional signals in the tropical climate. *Climate Dynamics* 22, 327–341, 2004.
- 975 Cole-Dai, J., Ferris, D. G., Kennedy, J. A., Sigl, M., McConnell, J. R., Fudge, T. J., Geng, L., Maselli, O. J., Taylor, K. C., and Souney, J. M.: Comprehensive record of volcanic eruptions in the Holocene (11,000 years) from the WAIS Divide, Antarctica ice core, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD032855, 2021.
- Crowley, T. J. and Unterman, M. B.: Technical details concerning development of a 1200-yr proxy index of global volcanism, *Earth System Science Data*, 5 187–197, 2013.
- 980 Cruz, F.W., Vuille, M., Burns, S.J.: Orbitally driven east–west antiphasing of South American precipitation. *Nature Geoscience* 2(3): 210–214, 2009.
- Crucifix, M., Loutre, M.F, Tulkens, P., Fichet, T., Berger, A: Climate evolution during the Holocene: A study with an Earth system model of intermediate complexity, *Clim. Dyn.*, 19, 43–60, 10.1007/S00382 001 0208 6, 2002.
- 985 D'Agostino, R., Bader, J., Bordoni, S., Ferreira, D., and Jungclaus, J: Northern hemisphere monsoon response to mid-holocene orbital forcing and greenhouse gas-induced global warming. *Geophysical Research Letters*, 46, 1591–1601., 2019.
- Dallmeyer, A., Claussen, M., Fischer, N., Haberkorn, K., Wagner, S., Pfeiffer, M., Jin, L., Khon, V., Wang, Y., and Herzschuh, U.: The evolution of sub-monsoon systems in the Afro-Asian monsoon region during the Holocene–comparison of different transient climate model simulations, *Clim. Past*, 11, 305–326, <https://doi.org/10.5194/cp-11-305-2015>, 2015.
- 990 Dallmeyer, A., Claussen, M., and Brovkin, V.: Harmonising plant functional type distributions for evaluating Earth System Models. *Climate of the Past*, 15, 335–366. [doi:10.5194/cp-2018-41](https://doi.org/10.5194/cp-2018-41), 2019.
- Dallmeyer, A., Claussen, M., Lorenz, S. & Shanahan, T.: The end of the African humid period as seen by transient comprehensive Earth system model simulation of the last 8000 years. *Climate of the Past*, 16, 117–140. [doi:10.5194/cp-16-117-2020](https://doi.org/10.5194/cp-16-117-2020), 2020.

- Davis, B.A.S., Brewer, S., Stevenson, A.C., Guiot, J., 2003. The temperature of Europe during the Holocene reconstructed from pollen data. *Quaternary Science Reviews* 22, 1701–1716.
- 1000 deMenocal, P. B., Ortiz, J., Guilderson, T., Adkins, J., Santhein, M., Baker, L., and Yarusinsky, M.: Abrupt onset and termination of the African Humid Period: Rapid climate responses to gradual insolation forcing. *Quaternary Science Reviews* 19 (1-5), 347-361, 2000.
- Egerer, S., Claussen, M., Reick, C., and Stanelle, T.: The link between marine sediment records and changes in Holocene Saharan landscape: simulating the dust cycle. *Clim. Past*, 12, 1009-1027, 2016
- 1005 Feng, Z.-D., An, C.B., and Wang, H.B.: Holocene climatic and environmental changes in the arid and semi-arid areas of China: a review. *The Holocene* 16(1), 119-130, 2006.
- Fischer, N., and Jungclauss, J.H.: Evolution of the seasonal temperature cycle in a transient Holocene simulation: orbital forcing and sea-ice. *Clim. Past*, 7, 1139-1148, 2011.
- 1010 Fletcher, M.S., and Moreno, P.I.: Have the Southern Westerlies changed in a zonally symmetric manner over the last 14,000 years? A hemisphere-wide take on a controversial problem, *Quaternary International*, Volume 253, Pages 32-46, ISSN 1040-6182, <https://doi.org/10.1016/j.quaint.2011.04.042>, 2012.
- Foster, D.R., Oswald, W.W., Faison, E.K., Doughty, E.D., Hansen, B.C.S.: A climatic driver for abrupt mid-Holocene vegetation dynamics and the hemlock decline in New England. *Ecology* 87(12): 2959–2966, 2006.
- Gaetani M, Pohl B, Douville H, and Fontaine B: West African Monsoon influence on the summer Euro-Atlantic circulation. *Geophys Res Lett* 38:38–42. <https://doi.org/10.1029/2011GL047150>, 2011.
- 1015 Gao, C. C., Oman, L., Robock, A., and Stenchikov, G. L.: Atmospheric volcanic loading derived from bipolar ice cores: Accounting for the spatial distribution of volcanic deposition, *J Geophys Res-Atmos*, 112, 2007.
- Garreaud, R.D., Vuille, M., Compagnucci, R.; Marengo, J.: Present-day South American climate. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 281(3-4):180-195, 2009.
- 1020 Giesecke, T., Bennett, K. D., Birks, H. J. B., Bjune, A. E., Bozilova, E., Feurdean, A., Finsinger, W., Froyd, C., Pokorny, P., Rosch, M., Seppa, H., Tonkov, S., Valsecchi, V., and Wolters, S.: The pace of Holocene vegetation change - testing for synchronous developments. *Quaternary Science Reviews*, 30(19-20), 2805–2814. <https://doi.org/10.1016/j.quascirev.2011.06.014>, 2011.
- Gill, A. E.: Some simple solutions for heat-induced tropical circulation. *Quart. J. Roy. Meteor. Soc.*, 106, 447–462, 1980.
- 1025 Grimm, E.C.: Lozano-García, S., Behling, H., Markgraf, V.: Holocene Vegetation and Climate Variability in the Americas, *Interhemispheric Climate Linkages*, Academic Press, Pages 325-370, Ch.19, ISBN 9780124726703, <https://doi.org/10.1016/B978-012472670-3/50022-7>, 2001.
- 1030 Harrison, S.: BIOME 6000 DB classified plotfile version 1. University of Reading. Dataset. <http://dx.doi.org/10.17864/1947.99>, 2017.
- Harrison, S. P., Kutzbach, J.E., Liu, Z., Bartlein, P.J., Otto-Bliesner, Muhs, D., Prentice, I.C., and Thompson, R.S.: Mid-Holocene climates of the Americas: A dynamical response to changed seasonality. *Clim. Dyn.* 20, 663–688, 2003.

- Haug, G.H., Hughen, K.A., Sigman, D.M., Peterson, L.C., Rfhl, U.: Southward migration of the Intertropical Convergence Zone through theHolocene. *Science* 293, 1304–1308, 2001.
- 1035 Hély, C., Lézine, A.-M., and contributors, A.: Holocene changes in African vegetation: tradeoff between climate and water availability, *Clim. Past*, 10, 681-686, <https://doi.org/10.5194/cp-10-681-2014>, 2014.
- Herzschuh, U., Birks, H. J. B., Ni, J., Zhao, Y., Liu, H., Liu, X., and Grosse, G.: Holocene land-cover changes on the Tibetan Plateau, *The Holocene*, 20, 91–104, 2010a.
- 1040 Hopcroft, P., Valdes, P., Harper, A., & Beerling, D.: Multi vegetation model evaluation of the Green Sahara climate regime. *Geophysical Research Letters*, 44, 6804– 6813. <https://doi.org/10.1002/2017GL073740>, 2017.
- Hurtt, G. C., Chini, L., Sahajpal, R., Frolking, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori, S., Goldewijk, K. K., Hasegawa, T., Havlik, P., Heinimann, A., Humpenöder, F., Jungclaus, J., Kaplan, J., Kennedy, J., Kristzin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X.: Harmonization of Global Land-Use Change and Management for the Period 850–2100 (LUH2) for CMIP6, *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2019-360>, in review, 2020.
- 1045 Ilyina, T., Six, K., Segsneider, J., Maier Reimer, E., Li, H., and Núñez Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI Earth System Model in different CMIP5 experimental realizations, *J. Adv. Model. Earth Syst.*, doi:10.1029/2012MS000178, 2013.
- 1050 Jiang, D., Tian, Z., and Lang, X.: Mid-Holocene global monsoon area and precipitation from PMIP simulations. *Clim Dyn* 44, 2493–2512, <https://doi-org.e-bis.mpimet.mpg.de/10.1007/s00382-014-2175-8>. 2015.
- Jolly, D., Prentice, I. C., Bonneille, R., Ballouche, A., Bengo, M., Brenac, P., Buchet, G., Burney, D., Cazet, J.-P., Cheddadi, R., Ectorh, T., Elenga, H., Elmoutaki, S., Guiot, J., Laarif, F., Lamb, H., Lezine, A.-M., Maley, J., Mbenza, M., Peyron, O., Reille, M., Reynaud-Farrera, I., Riollet, G., Ritchie, J. C., Roche, E., Scott, L., Ssemmanda, I., Straka, H., Umer, M., Van Campo, E., Vilimumbalo, S., Vincens, A., and Waller, M.: Biome reconstruction from pollen and plant macrofossil data for Africa and the Arabian peninsula at 0 and 6000 years. *Journal of Biogeography*, 25(6):1007-1027, 1998.
- 1055 1060 Jungclaus, J. H., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L. P., Egorova, T., Evans, M., González-Rouco, J. F., Goosse, H., Hurtt, G. C., Joos, F., Kaplan, J. O., Khodri, M., Klein Goldewijk, K., Krivova, N., LeGrande, A. N., Lorenz, S. J., Luterbacher, J., Man, W., Maycock, A. C., Meinshausen, M., Moberg, A., Muscheler, R., Nehrbass-Ahles, C., Otto-Bliesner, B. I., Phipps, S. J., Pongratz, J., Rozanov, E., Schmidt, G. A., Schmidt, H., Schmutz, W., Schurer, A., Shapiro, A. I., Sigl, M., Smerdon, J. E., Solanki, S. K., Timmreck, C., Toohey, M., Usoskin, I. G., Wagner, S., Wu, C. J., Yeo, K. L., Zanchettin, D., Zhang, Q., and Zorita, E.: The PMIP4 contribution to CMIP6 – Part 3: The last millennium, scientific objective, and experimental design for the PMIP4 past1000 simulations, *Geosci. Model Dev.*, 10, 4005-4033, 2017.
- 1065 Jungclaus, J. H., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and von Storch, J. S.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *J. Adv. Model. Earth Sy.*, 5, 422–446, <https://doi.org/10.1002/jame.20023>, 2013.
- 1070 Köhler: Interactive comment on “What was the source of the atmospheric CO₂ increase during the Holocene?” by Victor Brovkin et al., *Biogeosciences Discuss.*, <https://doi.org/10.5194/bg-2019-64-SC1>, 2019.
- Krivova, N. A., Solanki, S. K., and Unruh, Y. C.: Towards a long-term record of solar total and spectral irradiance, *Journal of Atmospheric and Solar-Terrestrial Physics*, 73, 223-234, 10.1016/j.jastp.2009.11.013, 2011.

- 1075 Kröpelin, S., Verschuren, D., Lézine, A.-M., Eggermont, H., Cocquyt, C., Francus, P., Cazet, J.-P., Fagot, M., Rumes, B., Russell, J.M., Darius, F., Conley, D.J., Schuster, M., von Suchodoletz, H., and Engstrom, D.R.: Climate-driven ecosystem succession in the Sahara: the past 6000 years, *Science*, 320, 765-768, 2008
- Kutzbach, J. E.: Monsoon climate of the early Holocene – climate experiment with the earths orbital parameters for 9000 years ago, *Science*, 214, 59–61, 1981.
- 1080 Lamy, F., Kilian, R., Arz, H. W., Francois, J.-P., Kaiser, J., Prange, M. and Steinke, T.: Holocene changes in the position and intensity of the southern westerly wind belt, *Nat. Geosci.*, 3(10), 695–699, doi:10.1038/ngeo959, 2010.
- Ledru, M.P., Mourguiart, P., Riccomini, C.: Related changes in biodiversity, insolation and climate in the Atlantic rainforest since the last interglacial. *Palaeogeography, Palaeoclimatology, Palaeoecology* 271(1): 140–152, 2009.
- 1085 Li, Furong, Gaillard, M.-J., Cao, X., , U. Herzschuh et al.: Toward quantification of Holocene anthropogenic land-cover change in temperate China: a review in the light of pollen-based REVEALS reconstructions of regional plant cover. *Earth-Science Reviews*, 103119. <https://doi.org/10.1016/j.earscirev.2020.103119>, 2020.
- Liu, Z., B. Otto-Bliesner, J. Kutzbach, L. Li, and C. Shields: Coupled Climate Simulation of the Evolution of Global Monsoons in the Holocene. *J. Climate*, 16, 2472–2490, [https://doi.org/10.1175/1520-0442\(2003\)016<2472:CCSOTE>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<2472:CCSOTE>2.0.CO;2), 2003.
- 1090 Liu, Z., Harrison, S.P., Kutzbach, J., and Otto-Bliesner, B.: Global monsoons in the mid-Holocene and oceanic feedback. *Climate Dynamics* 22, 157–182., <https://doi-org.e-bis.mpimet.mpg.de/10.1007/s00382-003-0372-y>, 2004.
- Liu, J., Wang, B., Ding, Q., Kuang, X., Soon, W., and Zorita, E.: Centennial variations of the global monsoon precipitation in the last millennium: Results from ECHO-G model, *J. Clim.*, 22, 2356–2371, doi:10.1175/2008JCLI2353.1, 2009.
- 1095 Lorenz, S. J., Kim, J.-H., Rimbu, N., Schneider, R. R., and G. Lohmann: Orbitally driven insolation forcing on Holocene climate trends: Evidence from alkenone data and climate modeling, *Paleoceanography*, 21, PA1002, doi:10.1029/2005PA001152, 2006.
- Maksic, J., Shimizu, M.H., Sampaio, G., Venancio, I.M., Cardoso, M., Ferreira, F.A.: Simulation of the holocene climate over South America and impacts on the vegetation. *Holocene* 29(2):287–299, 2019.
- 1100 Marchant, R., Richer, S., Boles, O., Capitani, C., Courtney-Mustaphi, C. J., Lane, P., Prendergast, M. E., Stump, D., De Cort, G., Kaplan, J. O., Phelps, L., Kay, A., Olago, D., Petek, N., Platts, P. J., Punwong, P., Widgren, M., Wynne-Jones, S., Ferro-Vázquez, C., Benard, J., Boivin, N., Crowther, A., Cuní-Sánchez, A., Deere, N. J., Ekblom, A., Farmer, J., Finch, J., Fuller, D., Gaillard-Lemdahl, M.-J., Gillson, L., Githumbi, E., Kabora, T., Kariuki, R., Kinyanjui, R., Kyazike, E., Lang, C., Lejju, J., Morrison, K. D., Muiruri, V., Mumbi, C., Muthoni, R., Muzuka, A., Ndiema, E., Kabonyi Nzabandora, C., Onjala, I., Schrijver, A. P., Rucina, S., Shoemaker, A., Thornton-Barnett, S., 1105 van der Plas, G., Watson, E. E., Williamson, D. and Wright, D.: Drivers and trajectories of land cover change in East Africa: Human and environmental interactions from 6000 years ago to present, *Earth-Science Rev.*, 178, 322–378, doi:<https://doi.org/10.1016/j.earscirev.2017.12.010>, 2018.
- 1110 Marengo, J. A., Soares, W.R., Saulo, C., and Nicolini, M.: Climatology of the Low-Level Jet East of the Andes as Derived from the NCEP–NCAR Reanalyses: Characteristics and Temporal Variability. *J. Climate*, 17, 2261–2280, [https://doi.org/10.1175/1520-0442\(2004\)017<2261:COTLJE>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2261:COTLJE>2.0.CO;2), 2004.
- Marquer, L., Gaillard, M.-J., Sugita, S., Poska, A., Trondman, A.-K., Mazier, F., Nielsen, A.B., Fyfe, R.M., Jönsson, A.M., Smith, B., Kaplan, J.O., Alenius, T., Birks, H.J.B., Bjune, A.E., Christiansen, J., Dodson, J., Edwards, K.J., Giesecke, T., Herzschuh, U., Kangur, M., Koff, T., Latalowa, M., Lechterbeck, J., Olofsson, J., Seppä, H.:

- 1115 Quantifying the effects of land use and climate on Holocene vegetation in Europe. *Quaternary Science Reviews* 171, 20-37, 2017.
- Marsicek, J.P., Shuman, B., Brewer, S., Foster, D.R., and Oswald, W.W.: Moisture and temperature changes associated with the mid-Holocene *Tsuga* decline in the northeastern United States. *Quaternary Science Reviews*, Volume 80, Pages 129-142, <https://doi.org/10.1016/j.quascirev.2013.09.001>, 2013.
- Matsuno, T.: Quasi-geostrophic motions in the equatorial area., *J. Meteor. Soc. Japan*, 44, 25–43, 1966.
- 1120 Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Popke, D., Gayler, V., Giorgetta, M., Goll, D., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez Cuesta de la Otero, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornblueh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W., Nabel, J., Nam, C., Notz, D., Nyawira, S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T., Rast, S., Redler, R., Reick, C., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stein, L., Stemmler, I., Stevens, B., von Storch, J., Tian, F., Voigt, A., de Vrese, P., Wieners, K.-H., Wilkenskield, S., Roeckner, E. & Winkler, A.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and its response to increasing CO₂. *Journal of Advances in Modeling Earth Systems*, 11, 998-1038. doi:10.1029/2018MS001400, 2019.
- 1125 Mayewski, P. A., Meeker, L. D., Twickler, M. S., Whitlow, S., Yang, Q. Z., Lyons, W. B., and Prentice, M.: Major features and forcing of high-latitude northern hemisphere atmospheric circulation using a 110,000-year-long glaciochemical series, *J Geophys Res-Oceans*, 102, 26345-26366, 1997. doi: 10.1029/96JC03365
- Mayewski, P. A., Rohling, E. E., Stager, J. C., Karlén, W., Maasch, K. A., Meeker, L. D., Meyerson, E. A., Gasse, F., Van Kreveld, S., Holmgren, K., Lee-Thorp, J., Rosqvist, G.; Rack, F., Staubwasser, M., Schneider, R., Steig, E.:
1135 Holocene climate variability. *Quat. Res.* 62, 243–255. doi: 10.1016/j.yqres.2004.07.001, 2004.
- Mayle, F.E, Power, M.J: Impact of a drier early-mid-Holocene climate upon Amazonian forests. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 363(1498): 1829–1838, 2008.
- Mayle, F.E., Burbridge, R., Killeen, T.J.: Millennial-scale dynamics of southern Amazonian rain forests. *Science* 290(5500): 2291–2294. DOI: [10.1126/science.290.5500.2291](https://doi.org/10.1126/science.290.5500.2291), 2000.
- 1140 MacDonald, G.M., Velichko, A.A., Kremenetski, C.V., Borisova, O.K., Goleva, A.A., Andreev, A.A., Cwynar, L.C., Riding, R.T., Forman, S.L., Edwards, T.W.D., Aravena, R., Hammarlund, D., Szeicz, J.M., and Gattaulin, V.N.: Holocene treeline history and climate change across northern Eurasia. *Quat Res*, 53, 302–311, 2000.
- McGlone, M.S., Kershaw, A.P., and Markgraf, V.: El Niño /Southern Oscillation climatic variability in Australasian and South American paleoenvironmental records. In: Diaz HF, Markgraf V (eds) *El Niño: Historical and Paleoclimatic Aspects of the Southern Oscillation*. Cambridge: Cambridge University Press, 436–462, 1992.
1145
- Metcalf, S. E., Barron, J.E., and Davies, S.J.: The holocene history of the North American Monsoon: Known knowns and known unknowns in understanding its spatial and temporal complexity, *Quat. Sci. Rev.*, 120, 1– 27, doi:[10.1016/j.quascirev.2015.04.004](https://doi.org/10.1016/j.quascirev.2015.04.004), 2015.
- 1150 Meyer, D, Dimitriadou, E., Hornik, K., Weingessel, A., and Leisch, F.: e1071: Misc Functions of the Department of Statistics, Probability Theory Group (Formerly: E1071), TU Wien. R package version 1.6-8. <https://CRAN.R-project.org/package=e1071>, last access: 14th September, 2020., 2017.
- Nakanishi, T., Tachibana, Y. and Ando, Y.: Possible semi-circumglobal teleconnection across Eurasia driven by deep convection over the Sahel. *Clim Dyn.* <https://doi.org/10.1007/s00382-021-05804-x>, 2021.

- Nicholls, N.: Historical El Niño/Southern Oscillation variability in the Australasian region. In: Diaz HF, Makrkrgraf V (eds) *El Niño: Historical and Paleoclimatic Aspects of the Southern Oscillation*. Cambridge: Cambridge University Press, 151–173, 1992.
- 1155
- Olago, D.: Vegetation changes over palaeo-time scales in Africa., *Climate Research*, 17: 105-121,2001.
- Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E, Wagner, H.: *vegan: Community Ecology Package*. R package version 2.4-6. <https://CRAN.R-project.org/package=vegan>, last access: 14th September, 2020, 2018.
- 1160
- Overpeck, J. T., and Webb, R.S.: Non-glacial rapid climate events: Past and future. *Proc. Natl. Acad. Sci. USA*,97,1335–1338, 2000.
- Pickett, E.J, Harrison, S.P, Hope, G. Harle, K., Dodson, J.R., Kershaw, A.P., Prentice, I.C., Backhouse, J., Colhoun, E.A., D'Costa, D., Flenley, J., Grindrod, J., Haberle, S., Hassell, C., Kenyon, C., Macphail, M., Martin, H., Martin, A.H., McKenzie, M., Newsome, J.C., Penny, D., Powell, J., Raine, J.I., Southern, W., Stevenson, J., Sutra, J.P., Thomas, I., van der Kaars, S., Ward, J.: Pollen-based reconstructions of biome distributions for Australia, Southeast Asia and the Pacific (SEAPAC region) at 0, 6000 and 18,000 ¹⁴C yr B.P. *Journal of Biogeography* 31: 1381-1444. (DOI: 10.1111/j.1365-2699.2004.01001.x), 2004.
- 1165
- Prado, LF, Wainer, I, Chiessi, C.M.: Mid-Holocene PMIP3/CMIP5 model results: Intercomparison for the South American Monsoon System. *The Holocene* 23(12): 1915–1920, 2013a.
- 1170
- Prado, LF, Wainer, I, Chiessi, C.: A mid-Holocene climate reconstruction for eastern South America. *Climate of the Past* 9(5): 2117–2133. <https://doi.org/10.5194/cp-9-2117-2013>, 2013b.
- Prentice, I.C., Jolly, D., and BIOME 6000 participants: Mid-Holocene and glacial-maximum vegetation geography of the northern continents and Africa *J. Biogeogr.*, 27, pp. 507-519, 2000.
- 1175
- Quigley, M.C, Horton, T., Hellstrom, J.C., Cupper, M.L., Sandiford, M.: Holocene climate change in arid Australia from speleothem and alluvial records. *The Holocene*, 20(7):1093-1104. doi:[10.1177/0959683610369508](https://doi.org/10.1177/0959683610369508), 2010.
- Rachmayani, R., Prange, M., and Schulz, M.: North African vegetation–precipitation feedback in early and mid-Holocene climate simulations with CCSM3-DGVM, *Clim. Past*, 11, 175–185, <https://doi.org/10.5194/cp-11-175-2015>, 2015.
- 1180
- Ratnam, J., Bond, W. J., Fensham, R. J., Hoffmann, W. A., Archibald, S., Lehmann, C. E. R., Anderson, M. T., Higgins, S. I., and Sankaran, M.: When is a “forest” a savanna, and why does it matter?, *Glob. Ecol. Biogeogr.*, 20, 653–660, <https://doi.org/10.1111/j.1466-8238.2010.00634.x>, 2011.
- 1185
- Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: The representation of natural and anthropogenic land cover change in MPI ESM, *J. Adv. Model. Earth Syst.*, 5, 1–24, doi:[10.1002/jame.20022](https://doi.org/10.1002/jame.20022), 2013.
- 1190
- Ren, G.: Changes in forest cover in China during the Holocene. *Veget. Hist. Archaeobot.*, 16, 119-126, 2007.
- Rodwell, M. J. and Hoskins, B. J.: Subtropical anticyclones and summer monsoons, *J. Clim.*, 14(15), 3192–3211, doi:[10.1175/1520-0442\(2001\)014<3192:SAASM>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<3192:SAASM>2.0.CO;2), 2001.

- Rossetti, D.F., Cohen, M.C.L, Pessenda, L.C.R.: Vegetation change in Southwestern Amazonia (Brazil) and relationship to the late Pleistocene and Holocene climate. *Radiocarbon* 59(1): 69–89, 2017.
- 1195 Rojas, M., Moreno, P., Kageyama, M., Crucifix, M., Hewitt, C., Abe, A., Ohgaito, R., Brady, E., and Hope, P.: The Last Glacial Maximum in the Southern Hemisphere in PMIP2 simulations, *Clim. Dynam.*, 32, 525–548, 2009.
- Ruddiman, W. F.: *Earth's Climate: Past and Future*, 2nd revised edition, W.H. Freeman and Company, p. 388, 2008.
- Scheffer, M, Carpenter,S., Foley, J.A.,Folke, C, Walker, B.: Catastrophic shifts in ecosystems. *Nature*, 413:591–596, 2001.
- 1200 Scott, L. and Tee-Thorp, J. A.: *Holocene climatic trends and rhythms in southern Africa, Past Climate Variability through Europe and Africa*, Springer, Dordrecht, The Netherlands, 69–91, 2004
- Seager, R., Neelin, D., Simpson, I., Liu, H., Henderson, N., Shaw, T., Kushnir, Y., Ting, M. and Cook, B.: Dynamical and Thermodynamical Causes of Large-Scale Changes in the Hydrological Cycle over North America in Response to Global Warming*, *J. Clim.*, 27(20), 7921–7948, doi:10.1175/JCLI-D-14-00153.1, 2014.
- 1205 Seddon, A.W., Macias-Fauria, M., and Willis, K.J.: Climate and abrupt vegetation change in Northern Europe since the last deglaciation. *The Holocene*. 2015; 25(1):25-36. doi:10.1177/0959683614556383, 2015.
- Seddon, A., Macias-Fauria, M., Long, P., Benz, D., Willis, K.J.: Sensitivity of global terrestrial ecosystems to climate variability. *Nature* 531, 229–232.. <https://doi-org.e-bis.mpimet.mpg.de/10.1038/nature16986>, 2016.
- 1210 Severi, M., Becagli, S., Castellano, E., Morganti, A., Traversi, R., Udisti, R., Ruth, U., Fischer, H., Huybrechts, P., Wolff, E., Parrenin, F., Kaufmann, P., Lambert, F., and Steffensen, J. P.: Synchronisation of the EDML and EDC ice cores for the last 52 kyr by volcanic signature matching, *Clim Past*, 3, 367-374, 2007. doi: 10.5194/cpd-3-409-2007
- Shanahan, T.M., McKay, N.P., Hughen, K.A., Overpeck, J.T., Otto-Bliesner, B., Heil, C.W., King, J., Scholz, C.A., and Peck, J., The time-transgressive termination of the African Humid Period, *Nature Geoscience*, 8 (2), 140-144, 2015.
- 1215 Shin, S.-J.K., Sardeshmukh, P.-D., Webb, R.S., Oglesby, R.J., and Barsugli, J.J.: Understanding the Mid-Holocene Climate *J. Climate*, 19 (12): 2801–2817, 2006.
- Shuman, B.N., Newby, P. and Donnelly, J.P: Abrupt climate change as an important agent of ecological change in the Northeast U.S. throughout the past 15,000 years. *Quaternary Science Reviews* 28(17–18): 1693–1709, 2009.
- 1220 Sigl, M., Fudge, T. J., Winstrup, M., Cole-Dai, J., Ferris, D., McConnell, J. R., Taylor, K. C., Welten, K. C., Woodruff, T. E., Adolphi, F., Bisiaux, M., Brook, E. J., Buizert, C., Caffee, M. W., Dunbar, N. W., Edwards, R., Geng, L., Iverson, N., Koffman, B., Layman, L., Maselli, O. J., McGwire, K., Muscheler, R., Nishiizumi, K., Pasteris, D. R., Rhodes, R. H., and Sowers, T. A.: The WAIS Divide deep ice core WD2014 chronology - Part 2: Annual-layer counting (0-31 ka BP), *Clim Past*, 12, 769-786, 2016. doi: 10.5194/cp-12-769-2016
- 1225 Sigl, M., McConnell, J. R., Toohey, M., Curran, M., Das, S. B., Edwards, R., Isaksson, E., Kawamura, K., Kipfstuhl, S., Krüger, K., Layman, L., Maselli, O. J., Motizuki, Y., Motoyama, H., Pasteris, D. R., and Severi, M.: Insights from Antarctica on volcanic forcing during the Common Era, *Nat Clim Change*, 4, 693-697, 2014. doi: 10.1038/nclimate2293
- 1230 Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O. J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D. R., Pilcher, J. R., Salzer, M., Schüpbach, S., Steffensen, J. P., Vinther, B. M., and Woodruff, T. E.:

- Timing and climate forcing of volcanic eruptions for the past 2,500 years, *Nature*, 523, 543-549, 2015. doi: 10.1038/nature14565
- 1235 Smith, R. J., and Mayle, F. E.: Impact of mid-to late Holocene precipitation changes on vegetation across lowland tropical South America: a paleo-data synthesis. *Quaternary Research* 89, 134–155, 2018.
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornblueh, L., Lohmann, U., Pincus, R., Reichler, T., and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, *J. Adv. Model. Earth Sy.*, 5, 146–172, <https://doi.org/10.1002/jame.20015>, 2013.
- 1240 Tian, F., Cao, X., Dallmeyer, A., Ni, J., Zhao, Y., Wang, Y., Herzschuh, U., 2016. Quantitative woody cover reconstructions from eastern continental Asia of the last 22 kyr reveal strong regional peculiarities. *Quaternary Science Reviews* 137, 33-44. <https://doi.org/10.1016/j.quascirev.2016.02.001>.
- 1245 Tierney, J.E., Lewis, S.C., Cook, B.I., LeGrande, A.N., and Schmidt, G.A., Model, proxy and isotopic perspectives on the East African Humid Period, *Earth and Planetary Science Letters*, Volume 307, Issues 1–2, Pages 103-112, <https://doi.org/10.1016/j.epsl.2011.04.038>, 2011.
- Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500 BCE to 1900 CE, *Earth System Science Data*, 9, 809-831, 2017. doi: 10.5194/essd-9-809-2017
- Toohey, M., Stevens, B., Schmidt, H., and Timmreck, C.: Easy Volcanic Aerosol (EVA v1.0): an idealized forcing generator for climate simulations, *Geosci. Model Dev.*, 9, 4049-4070, 2016.
- 1250 Trenberth, K. E., Stepaniak, D. P., and Caron, J. M.: The global monsoon as seen through the divergent atmospheric circulation, *J. Climate*, 13, 3969–3993, 2000
- 1255 Trondman, A.K., Gaillard, M.-J., Mazier, F., Sugita, S., Fyfe, R., Nielsen, A.B., Twiddle, C., Barratt, P., Birks, H.J.B., Bjune, A.E., Björkman, L., Broström, A., Caseldine, C., David, R., Dodson, J., Dörfler, W., Fischer, E., van Geel, B., Giesecke, T., Hultberg, T., Kalnina, L., Kangur, M., van der Knaap, P., Koff, T., Kuneš, P., Lagerås, P., Latalowa, M., Lechterbeck, J., Leroyer, C., Leydet, M., Lindbladh, M., Marquer, L., Mitchell, F.J.G., Odgaard, B.V., Peglar, S.M., Persson, T., Poska, A., Rösch, M., Seppä, H., Veski, S., and Wick, L.: Pollen-based quantitative reconstructions of Holocene regional vegetation cover (plant-functional types and land-cover types) in Europe suitable for climate modelling, *Global Change Biology*, 21(2), 676-697, DOI:10.1111/gcb.12737, 2015.
- 1260 Van Campo, E., Cour, P., and Sixuan, H.: Holocene environmental changes in Bangong Co basin (Western Tibet). Part 2: The pollen record. *Palaeogeogr. Palaeoclimatol.*, 120(1–2), 49–63, 1996.
- Verdon, D. C. and Franks, S. W.: Indian Ocean sea surface temperature variability and winter rainfall: Eastern Australia, *Water Resour. Res.*, 41, 1–10, <https://doi.org/10.1029/2004WR003845>, 2005.
- Viau, A. E., and K. Gajewski: Holocene variations in the global hydrological cycle quantified by objective gridding of lake databases. *J. Geophys. Res.*, 106, 31 703–31 716. 2001.
- 1265 Viau, A.E., Gajewski, K., Sawada, M.C., Fines, P.: Millennial-scale temperature variations in North America during the Holocene. *Journal of Geophysical Research* 111 (D9), D09102, 2006.
- Vincens, A., Buchet, G., Williamson, D., and Taieb, M.: A 23,000 yr pollen record from Lake Rukwa (8°S, SW Tanzania): new data on vegetation dynamics and climate in central eastern Africa, *Rev. Palaeobot. Palynol.*, 137, pp. 147-162, 2005.

- 1270 Vigaud, N., Richard, Y., Rouault, M. and Fauchereau, N.: Moisture transport between the South Atlantic Ocean and southern Africa: relationships with summer rainfall and associated dynamics, *Clim. Dyn.*, 32(1), 113–123, doi:10.1007/s00382-008-0377-7, 2009.
- Wanner, H., Beer, J., Bütikofer, J., Crowley, T.J., Cubasch, U., Flückiger, J., Goosse, H., Grosjean, M., Joos, F., Kaplan, J.O., Küttel, M., Müller, S.A., Prentice, I.C., Solomina, O., Stocker, T.F., Tarasov, P., Wagner, M., Widmann, M.:
1275 Mid- to Late Holocene climate change: an overview. *Quat. Sci. Rev.*, 27, 1791–1828. doi: 10.1016/j.quascirev.2008.06.013, 2008.
- Wang, P. X., Wang, B., Cheng, H., Fasullo, J., Guo, Z. T., Kiefer, T., and Liu, Z. Y.: The global monsoon across timescales: coherent variability of regional monsoons, *Clim. Past*, 10, 2007–2052, <https://doi.org/10.5194/cp-10-2007-2014>, 2014.
- 1280 Williams, J.W., Post, D.M., Cwynar, L. C., Lotter, A.F., and Levesque, A.J.: Rapid and widespread vegetation responses to past climate change in the North Atlantic region. *Geology*, 30: 971-974, 2002.
- Williams, J.W.: Variations in tree cover in North America since the Last Glacial Maximum. *Global and Planetary Change* 35, 1-23, 2003.
- Williams, J.W., Shuman, B.N., Webb, T. III, Bartlein, P.J, and Leduc, P.L: Late Quaternary vegetation dynamics in
1285 North America: scaling from taxa to biomes. *Ecological Monographs*, 74, 309– 334, 2004.
- Williams, J.W., Shuman, B., Bartlein, P.J., Diffenbaugh, N.S., Webb, III,T.: Rapid, time-transgressive, and variable responses to early Holocene midcontinental drying in North America. *Geology* 38:135–138. <https://doi.org/10.1130/G30413.1>. 2010.
- 1290 Williams, J.W., Tarasov, P.A., Brewer, S., Notaro, M.: Late-Quaternary variations in tree cover at the northern forest-tundra ecotone. *Journal of Geophysical Research - Biogeosciences* 116, G01017, doi: 01010.01029/02010JG001458, 2011.
- Xie, L.X., and Beni, G.; Validity measure for fuzzy clustering, *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol.3, n. 8, p. 841-847, 1991.
- 1295 Yu, G., Chen, X., Ni, J., Cheddadi, R., Guiot, J., Han, H., Harrison, S.P., Huang, C., Ke, M., Kong, Z., Li, S., Li, W., Liew, P., Liu, G., Liu, J., Liu, Q., Liu, K.-B., Prentice, I.C., Qui, W., Ren, G., Song, C., Sugita, S., Sun, X., Tang, L., Van Campo, E., Xia, Y., Xu, Q., Yan, S., Yang, X., Zhao, J., and Zheng, Z.: Palaeovegetation of China: a pollen date-based synthesis for the mid-Holocene and last glacial maximum. *J. Biogeogr.*, 27, 635–664, 2000.
- Zhang, Y, Renssen, H, Seppä, H, and Valdes, P.: Holocene temperature trends in the extratropical northern hemisphere based on inter-model comparisons. *Journal of Quaternary Science* 33(4): 464–476, 2018.
- 1300 Zhao, Y., Yu, Z., and Chen. F.: Spatial and temporal patterns of Holocene vegetation and climate changes in arid and semi-arid China. *Qut. Int.*, 194,6-18, 2009.
- Zhao, Y., and Harrison, S.: Mid-Holocene monsoons: A multi-model analysis of the inter-hemispheric differences in the responses to orbital forcing and ocean feedbacks. *Climate Dynamics*, 39(6), 1457–1487, 2012.
- 1305 Zhao, Y., Liu, Y. L., Guo, Z. T., Fang, K. Y., Li, Q., Cao, X. Y: Abrupt vegetation shifts caused by gradual climate changes in central Asia during the Holocene. *Sci China Earth Sci*, 60: 1317–1327, 2017.
- Zhou, T., Zhang, L., and Li, H.: Changes in global land monsoon area and total rainfall accumulation over the last half century, *Geophys. Res. Lett.*, 35, 1–6, <https://doi.org/10.1029/2008GL034881>, 2008.

No.	landcover classification	phenology type	$T_{C_{min}}$ [°C]	$T_{C_{max}}$ [°C]	$T_{W_{max}}$ [°C]	GDD5 [°C]	time scale[y]
1	tropical evergreen trees	raingreen	15.5	-	-	0	30
2	tropical deciduous trees	raingreen	15.5	-	-	0	30
3	extratrop. evergreen trees	evergreen	-32.5	18.5	-	400	60
4	extratrop. deciduous trees	summergreen	-	18.5	-	400	60
5	raingreen shrubs	raingreen	2.0	-	-	900	12
6	cold shrubs	summergreen	-	-2	18	350	24
7	C3 grass	grasses	-	15	-	0	1
8	C4 grass	grasses	10	-	-	0	1

Table.1: Bioclimatic limits and time-scale of allocation and mortality for the eight natural plant functional types (PFTs) used in the dynamic vegetation module of the model JSBACH. Listed are phenology type, PFT-specific minimum ($T_{C_{min}}$) and maximum ($T_{C_{max}}$) of the mean temperature of the coldest month, PFT-specific maximum mean temperature ($T_{W_{max}}$) of the warmest month, and growing degree days, i.e. temperature sum of days with temperatures exceeding 5°C (GDD5). All temperature values are given in °C.

1310

Biome	BNS
tropical forest	0.58
warm-temperate mixed forest	0.51
temperate forest	0.88
boreal forest	0.79
savanna and dry woodland	0.12
grassland and dry shrubland	0.63
desert	0.39
tundra	0.53
total	0.73

Table 2: Best neighbour score (BNS) showing the agreement between pollen-based reconstructions and the mega-biomes converted from the simulated plant functional type cover fractions. Values between 0.2 and 0.5 means ‘fair to good’ agreement, values above 0.5 ‘good to very good’ agreement, and above 0.8 reveal an ‘excellent’ agreement. For details on the metric see Dallmeyer et al. (2019).

Region		Monsoon precipitation	Monsoon area
NH	land	+40 %	+ 25 %
	total	+ 6 %	+ 4 %
SH	land	- 11 %	- 5 %
	total	+ 7 %	+ 8 %

1315

Table 3: Simulated change in northern hemispheric (NH) and southern hemispheric (SH) monsoon area and monsoon precipitation for the mid-Holocene (8ka b2k) compared to the late-Holocene (2.15ka b2k). The changes have been calculated following the method of Zhou et al. (2008) with the modifications of Liu et al. (2009).




Figure 1: Simulated (a) and reconstructed (b) biomes for the 6ka time-slice and the simulated (c) and reconstructed (d) change in vegetation between 6ka and pre-industrial (PI), expressed in terms of **landscape openness**. To assess the biome change, the biomes has been further grouped into the main categories forest, savannas, grassland/tundra and desert. Decrease in openness describes the transition from biomes indicating more open landscape to biomes indicating less open landscape (e.g. desert in 6ka to grasslands or to savannas or to forests in PI). **Increase in openness** Negative development describe the opposite transition (e.g. from grassland in 6ka to desert in PI).




Figure 2: Simulated Holocene vegetation cluster, derived by c-means clustering method (R, Meyer et al. 2017). The left panel displays the cluster pattern for (a) the bare soil cover, which is 1 minus the total vegetation in the model, (c) forest cover, (e) shrub cover, and (g) grass cover. The right panel shows the respective trends in the cluster centres, i.e. for (b) bare soil fraction, (d) forest fraction, (f) shrub fraction, and (h) grass fraction. For the analysis, 100 year running means of the anomaly time-series (fraction at certain time-step – 8ka b2k climatological mean) have been taken. The cover fraction change of the cluster centres can range from -1 to 1 in the model, i.e. decrease of 100% or increase of 100%, respectively.




Figure 3: Simulated Holocene climate cluster, derived by c-means clustering method (R, Meyer et al., 2017). The left panel displays the cluster pattern for (a) annual mean precipitation, (c) temperature of the warmest month (T_{warm}), and (e) temperature of the coldest month (T_{cold}). The right panel shows the respective trends in the cluster centres, i.e. for (b) annual mean precipitation [mm/d], (d) T_{warm} [K], and (f) T_{cold} [K]. For the analysis, 100 year running means of the anomaly time-series (value at certain time-step – 8ka b2k climatological mean) have been taken.




Figure 4: Results of the redundancy analyses performed by the VEGAN routine in R (Oksanen et al, 2018) for the vegetation groups forest, shrubs, and grass, with precipitation (precip) and temperature of the warmest month (Twarm) as explanatory variables. This figure is based on the ratio of the variance explained by Twarm (Tw) and by precipitation (P). $P > 2 \cdot Tw$ means that precipitation explains more than 2 times more of the variance than Twarm, vice versa for $Tw > 2 \cdot P$.




Figure 5: Season which contributes most to the annual mean precipitation change between 8ka b2k and 2.15ka b2k in the model, i.e. September to November (SON), June - August (JJA), March to May (MAM) and December to February (DJF). Please notice, that the seasons are calculated on the fixed model calendar and are not adjusted to the astronomical forcing (c.f. Bartlein and Shafer, 2018).




Figure 6: Monsoon region at 8ka and 2.15ka b2k, derived by the method by Zhou et al. (2008) with the modifications by Liu et al. (2009). The figure shows the monsoon area that is indicated for both time period, i.e. does not change from mid-to late Holocene (light blue), area additionally assigned to the monsoon domain at 8ka b2k, but not at 2.15 ka b2k (dark blue) and area additionally assigned to the monsoon domain at 2.15 ka b2k, but not at 8ka b2k.




Figure 7: Simulated changes in plant function types (PFT) for the **early** mid-Holocene time-slice (8ka b2k) compared to the late-Holocene (2.15ka b2k), i.e. yellowish colour indicate a decreased in PFT cover fraction at 8ka b2k, greenish colour an increased PFT cover fraction at 8ka b2k compared to 2.15ka b2k. Values of '1.0' mean that the grid-box is fully covered by the PFT during 8ka b2k, but does not occur at 2.15ka b2k in this grid-cell, and vice versa for values of '-1.0'. Only significant changes are shown (**t-test, 99% confidence interval**).





Figure 8: Simulated difference in June to August (JJA) mean climate in the model for the time-slice 8ka b2k compared to 2.15ka b2k, i.e. a) vertical velocity (ω) in 500hPa [10^{-2} m/s] with blue colour showing increased uplift during 8ka b2k; b) upper-tropospheric zonal wind (u) in 250hPa [m/s] with blue colours indicating enhanced easterly wind; c) low-level zonal flow (u) in 850hPa [m/s]; d) zonal anomaly in lower tropospheric streamfunction (ψ^* , in 10^7 m²/s) reflecting the change in standing waves, with blue colours indicating increased cyclonic circulation at 8ka b2k compared to 2.15ka b2k; and e) mean seasonal precipitation [mm/d] with blue colours showing enhanced precipitation during 8ka b2k. **Only significant changes are shown (t-test, 99% confidence interval).**

Figure 9:



Simulated difference in December to February (DJF, austral summer) mean climate in the model for the time-slice 8ka b2k compared to 2.15ka b2k, i.e. a) vertical velocity (omega) in 500hPa [10^{-2} m/s] with blue colour showing increased uplift during 8ka b2k; b) upper-tropospheric zonal wind (u) in 250hPa [m/s] with blue colours indicating enhanced easterly wind; c) low-level zonal flow (u) in 850hPa [m/s]; d) mean sea level pressure [hPa] with low pressure anomaly given in blue colours; and e) mean seasonal precipitation [mm/d] with blue colours showing enhanced precipitation during 8ka b2k. **Only significant changes are shown (t-test, 99% confidence interval).**




Figure 10: Simulated difference in September to November (SON) mean climate in the model for the time-slice 8ka b2k compared to 2.15ka b2k, i.e. a) vertical velocity (ω) in 500hPa [10^{-2} m/s] with blue colour showing increased uplift during 8ka b2k; b) upper-tropospheric zonal wind (u) in 250hPa [m/s] with blue colours indicating enhanced easterly wind; c) low-level zonal flow (u) in 850hPa [m/s]; d) zonal anomaly in lower tropospheric streamfunction (ψ^* , in 10^7 m²/s) reflecting the change in standing waves, with blue colours indicating increased cyclonic circulation at 8ka b2k compared to 2.15ka b2k; and e) mean seasonal precipitation [mm/d] with blue colours showing enhanced precipitation during 8ka b2k. **Only significant changes are shown (t-test, 99% confidence interval).**




Figure 11: Simulated difference (shaded) in June to September mean upper tropospheric (250hPa) velocity potential [km^2/s] between the time-slice 8ka b2k and 2.15ka b2k. Bluish colour indicate enhanced divergence, coinciding vertical ascent in the troposphere below, redish color indicate stronger convergence, coinciding subsidence below. The contours plotted on top show the respective velocity potential at 2.15ka b2k. Negative values display divergence, positive values convergence.




Figure 12: maximum relative change in PFTs (R), indicating the rapidness of the PFT change. The more yellowish the more rapid is the transition. The model reveals six different regions with rapid changes (region 1-6).




Figure 13: Examples of rapid transitions between simulated PFTs in a grid cell in a) East Siberia (at the grid cell centred at 62.49 N, 120 E in region 2 in Fig. 12), b) today's Sahara / Sahel region (at 19.59 N, 7.5 W in region 3), and c) South Asia (at 19.59 N, 78.75 E in region 4), shown as fractional coverage $V(t)$ as function of time t , given in ky b2k. The colors refer to different PFTs as indicated in the title line. EET: extratropical evergreen trees, EDT: extratropical deciduous trees, TET: tropical evergreen trees, TDT: tropical deciduous trees, SCR: shrubs, GRA: grass, BSF: bare soil fraction. The thin lines are annual values of fractional vegetation coverage, and the thick lines are filtered time series using a Butterworth filter of order 5 with a lowpass filter of 500 years.




Figure 14.: Simulated changes in PFTs in the today Sahara / Sahel region (part of region 3 in Fig. 12). The box in the upper left corner is centred at 25.18 N, 15 W, and box at the lower right corner, at 19.59 N, 5.63 W. Like in Fig. 13, the fractional coverages V of PFTs in each grid cell are depicted as function of time, given in ky b2k. The colors refer to bare soil fraction (yellow), grass (light green), tropical evergreen tree (dark green), tropical deciduous tree (green), shrubs (blue).




Figure 15: Summarising sketch of the main vegetation, climate and circulation changes from 8ka to 2.15ka b2k. South of 60°N, the vegetation trend positively correlates with the precipitation trend, i.e. regions with an increasing vegetation and/or forest fraction during the Holocene experienced increasing precipitation and vice versa for regions with decreasing trend.




Figure A1: Simulated plant functional type distributions at 2.15ka b2k.




Figure A2: Simulated plant functional type distributions at 8ka b2k.




Figure B1: Simulated annual mean sea surface temperatures (SST) for a) the 2.15ka b2k time-slice and b) the difference between 8ka and 2.15ka b2k.




Figure B2: Simulated JJA mean climate at 2.15ka b2k, i.e. a) vertical velocity (omega) in 500hPa [10^{-2} m/s] with blue colour showing uplift; b) upper-tropospheric zonal wind (u) in 250hPa [m/s] with blue colours indicating easterly wind; c) low-level zonal flow (u) in 850hPa [m/s]; d) zonal anomaly in lower tropospheric streamfunction (psi*, in 10^7 m²/s) reflecting the standing waves, with blue colours indicating cyclonic circulation; and e) mean seasonal precipitation [mm/d]




Figure B3: Simulated SON mean climate at 2.15ka b2k, i.e. a) vertical velocity (ω) in 500hPa [10^{-2} m/s] with blue colour showing uplift; b) upper-tropospheric zonal wind (u) in 250hPa [m/s] with blue colours indicating easterly wind; c) low-level zonal flow (u) in 850hPa [m/s]; d) mean sea level pressure anomaly [hPa] to standard atmospheric mean pressure (here 1013.25hPa), with blue colours indicating low-pressure systems; and e) mean seasonal precipitation [mm/d].




Figure B4: Simulated DJF mean climate at 2.15ka b2k, i.e. a) vertical velocity (omega) in 500hPa [10^{-2} m/s] with blue colour showing uplift; b) upper-tropospheric zonal wind (u) in 250hPa [m/s] with blue colours indicating easterly wind; c) low-level zonal flow (u) in 850hPa [m/s]; d) mean sea level pressure anomaly [hPa] to standard atmospheric mean pressure (here 1013.25hPa), with blue colours indicating low-pressure systems; and e) mean seasonal precipitation [mm/d].




Fig. C1: Black lines: Fractional coverage of extratropical deciduous tree in a grid box in East Siberia centred at 62.48 N, 120 E in region 2, see Fig. 12, and Fig 13 a) as function of time (in ky b2k). The thin lines are annual values of fractional vegetation coverage $V(t)$, and the thick lines are filtered time series of $V(t)$ using a Butterworth filter of order 5 with a low-pass filter of a) $F=100$ years, b) $F=250$ years, c) $F=500$ years, d) $F=750$ years. The absolute value of the relative change R is shown as red-dotted line. The titles of each subfigure indicate the strength of the low-pass filter F , the maximum of the absolute value of the relative change R_{\max} , and the time t at which R_{\max} occurs.