

Early Holocene cold snaps and their expression in the moraine record of the Eastern European Alps

Sandra M. Braumann^{1,2}, Joerg M. Schaefer², Stephanie M. Neuhuber¹, Christopher Lüthgens¹, Alan J. Hidy³, Markus Fiebig¹

5 ¹University of Natural Resources and Life Sciences (BOKU), Peter Jordan-Str. 82, 1190 Vienna, Austria

²Lamont-Doherty Earth Observatory of Columbia University, Division of Geochemistry, Palisades, NY 10964, USA

³Center for Accelerator Mass Spectrometry, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

Correspondence to: Sandra M. Braumann (sandra.braumann@boku.ac.at)

10 **Abstract.** Glaciers preserve climate variations in their geological and geomorphological records, which makes them prime candidates for climate reconstructions. Investigating the glacier-climate system over the past millennia is particularly relevant because, first, the amplitude and frequency of natural climate variability during the Holocene provides the climatic context against which modern, human-induced climate change must be assessed. Second, the transition from the last glacial to the current interglacial promises important insights into the climate system during warming, which is of particular interest with
15 respect to ongoing climate change.

Evidence of stable ice-margin positions that record cooling during the past 12 ka are preserved in two glaciated valleys of the Silvretta Massif in the Eastern European Alps, the Jamtal (JAM) and the Laraintal (LAR). We mapped and dated moraines in these catchments including historical ridges using Beryllium-10 Surface Exposure Dating (¹⁰Be SED) techniques, and correlate resulting moraine formation intervals with climate proxy records to evaluate the spatial and temporal scale of these cold phases.

20 The new geochronologies indicate the formation of moraines during the Early Holocene (EH), c. 11.0 ±0.7 ka (n=19). Boulder ages along historical moraines (n=6) suggest at least two glacier advances during the Little Ice Age (LIA; c. 1250-1850 CE), around 1300 CE and in the second half of the 18th century. An earlier advance to the same position may have occurred around 500 CE.

The Jamtal and Laraintal moraine chronologies provide evidence that millennial-scale EH warming was superimposed by
25 centennial-scale cooling. The timing of EH moraine formation coincides with brief temperature drops identified in local and regional paleoproxy records, most prominently with the Preboreal Oscillation (PBO), and is consistent with moraine deposition in other catchments in the European Alps, and in the Arctic region. This consistency points to cooling beyond the local scale and therefore a regional or even hemispheric climate driver. Freshwater input sourced from the Laurentide Ice Sheet (LIS), which changed circulation patterns in the North Atlantic, is a plausible explanation for EH cooling and moraine formation in
30 the Nordic region and in Europe.

1 Introduction

The transition from the Younger Dryas (YD; 12.9–11.7 ka; e.g., Alley, 2000) to the Holocene (c. 11.7 ka to present, e.g., Walker et al., 2008) is an important period for studying the climate system, its forcings and its feedbacks. Climatic conditions shifted from glacial to full interglacial conditions within approximately two millennia, between 12 and 10 ka (e.g., Cheng et al., 2020; Marcott et al., 2013; Rasmussen et al., 2006). This general warming trend was interrupted by abrupt centennial-scale cooling that appears linked to freshwater input into the North Atlantic (e.g., Bjorck et al., 1997; Hald and Hagen, 1998; Nesje et al., 2004; Thornalley et al., 2010). The climatic shift from the YD to the Early Holocene (EH) was accompanied by a multitude of major environmental changes that are interconnected, including the melting of ice caps and glaciers in both hemispheres, changes in the atmospheric composition and in circulation patterns, and the reorganization of ocean currents (e.g., Clark et al., 2012; Denton et al., 2021; Shakun et al., 2015). Human-induced warming since the beginning of the industrial era is on a trajectory to lead to changes of similar magnitude in our environmental system, yet at an even faster pace (Beniston et al., 2018; Gobiet et al., 2014). By investigating the timing of YD-EH warming and its perturbations, we can broaden our knowledge on natural drivers and physical mechanisms, which modulated the climate system at that time. Information on climate oscillations obtained from this major natural transition – from glacial to interglacial conditions – provides a valuable foundation for disentangling natural and anthropogenic forcings and their respective relevance. New knowledge in this field is particularly useful in the light of the ongoing transition from an interglacial to an industrialized world.

Glaciers respond to climate fluctuations sensitively and are important elements for understanding the climate of the past (Huston et al., 2021; Roe et al., 2017). Reconstructing former ice margins allows deciphering glacier fluctuations across time and space and informs us on climate variations that drove these changes. Mountain glaciers in alpine, melt-dominated regimes are most sensitive to changes in summer temperature, to a lesser extent to changes in precipitation (e.g., Oerlemans, 2005; Rupper and Roe, 2008; Steiner et al., 2008). At the end of the Late Glacial (LG), YD cooling resulted in glacier stabilization or readvance in the European Alps and lead to the deposition of moraine sets, whose estimated Equilibrium Line Altitudes (ELA) are approximately 250 to 350 m below ELAs of glaciers during the LIA (e.g., Ivy-Ochs, 2015). These moraines are termed ‘Egesen’ moraines and have been subject of numerous cosmogenic nuclide studies, which have advanced our understanding of glacier responses to cooling during the LG (e.g., Cossart et al., 2012; Federici et al., 2008; Ivy-Ochs et al., 2009; Ivy-Ochs et al., 2006; Kelly et al., 2004; Kerschner and Ivy-Ochs, 2008). In parallel, first attempts had been made to produce direct ages of younger moraines that were identified inboard the presumable Egesen moraines, but outboard historical LIA margins (Ivy-Ochs et al., 2006; Kerschner et al., 2006). Based on their morphostratigraphy, these moraine ridges were postulated as type localities for preboreal glacier advance, for instance the Kartell moraines in the Verwall area and the Kromer moraines in the Silvretta Massif, both in the Eastern Alps (e.g., Faedrich, 1979; Gross et al., 1978). Kartell moraines are today placed into the latest YD (Egesen-III). Recalculated ^{10}Be ages of Kromer moraines suggest moraine deposition during the EH, around 10 ka (Ivy-Ochs et al., 2006; Kerschner et al., 2006; Moran et al., 2016b). Dating efforts that address presumable EH

moraines continued toward the Central and Western Alps and have produced chronologies, which substantiate moraine
65 formation between 12 and 10 ka, although not necessarily synchronously (Baroni et al., 2017; Boxleitner et al., 2019a;
Boxleitner et al., 2019b; Cossart et al., 2012; Hofmann et al., 2019; Moran et al., 2016a; Moran et al., 2017a; Moran et al.,
2016b; Protin et al., 2021; Protin et al., 2019; Schimmelpfennig et al., 2012; Schimmelpfennig et al., 2014; Schindelwig et al.,
2012). In a few recent studies this pattern of moraine deposition has been confirmed in the Eastern Alpine region (Bichler et
70 al., 2016; Moran et al., 2016a; Moran et al., 2017a; Moran et al., 2017b). The youngest dated EH moraine is located in the
Ochsental, a valley adjacent to the sites discussed in this study (Braumann et al., 2020). The relevance of this chronology lies
in the finding that glaciers in the valley had melted back to historical sizes around 10 ka, and that they have remained within
these limits throughout the Holocene, which is consistent with complementary paleoproxy records from the Eastern Alps (e.g.,
Dietre et al., 2014; Nicolussi and Patzelt, 2000; Patzelt, 2016).

75 To intensify our knowledge on EH glacier configurations in the Eastern Alps, where directly dated moraine ages remain sparser
compared to the Western and Central Alps, we conducted a geochronological study in two glaciated catchments in the Silvretta
region in the Eastern Alps. We applied state-of-the-art cosmogenic nuclide techniques to date moraines inboard presumable
LG ice margins to constrain the timing of Holocene cold phases recorded in the moraine record. We chose this region for two
reasons: First, moraine sets which postdate the LG phase are well-preserved in the Silvretta Massif and show multi-ridge
80 structures. These geomorphological features promise insights into repeated Holocene cooling at times when glaciers were
larger than during the LIA. Second, in addition to comparable cosmogenic nuclide records in the region (Braumann et al.,
2020; Ivy-Ochs et al., 2006; Moran et al., 2016b), high-quality paleoenvironmental, archeological and historical information
on Holocene climate, which complements the moraine record, is available (Dietre et al., 2014; Kasper, 2015, 2013; Nicolussi,
2010; Patzelt, 2019).

85 The primary objective of this study is to generate more detailed and robust moraine chronologies in the Eastern Alpine region,
which contribute to our understanding of the spatial and temporal pattern of glacier advances throughout the Holocene with
an emphasis on the EH. We correlate the new moraine chronologies with moraine records and climate proxy data from the
Alpine region and from other glaciated regions in the Northern Hemisphere, and identify climate signals that are coherent with
Holocene glacier and climate evolution in the Silvretta Massif. Finally, we discuss possible links between climatic trigger
90 events and EH cold snaps, which manifest in the moraine record of the Northern Hemisphere.

2 Setting

The study sites are located at the north-facing side of the Silvretta Massif, a mountain range in the transition zone between the
Eastern and Western European Alps, at the border of Austria and Switzerland (Fig. 1a). Moraine sets from two adjacent valleys,
the Jamtal (JAM) and the Laraintal (LAR), were investigated and used for glacier reconstructions. Both valleys are north-south
95 oriented, drain northwards into the Danube catchment and are at present glaciated only in their highest sections (>2400 m

a.s.l.; Fig. 1b). The main and most prominent glacier of the Jamtal is the Jamtalferner with an area of approximately 2.8 km² (DEM of 2018). Smaller glaciers such as the Totenfeld, the Getschnerferner and the Augustenferner have retreated to cirque positions and are not connected to the valley glacier today (Fig. 1c). The situation is different in the neighboring Laraintal, where the Larainferner, which covers about 1 km² (DEM 2018), is the only glacier still present in the valley (Fig. 1d).

100




Figure 1: Location and lithology of investigated glaciated catchments. (a) Central Europe with the Alpine mountain range highlighted in dark grey and Austria outlined with black line. The Silvretta Massif (star symbol) is located in the westernmost part of the Eastern Alps. (b) North-facing section of the Silvretta Massif. Blue shading illustrates glacier extents in the year of 2012 (Fischer et al., 2015). Jamtal (JAM) and Laraintal (LAR) are subject of this study. Moraine chronologies of Kromertal (KR), Klostersal (KL) and Ochental (OcG-GrK) will be discussed later in this article (Braumann et al., 2020; Moran et al., 2016b). Red circles mark locations of subfossil tree findings. 1 – Bielerhöhe (Patzelt, 2019), 2 – Klostersal (Nicolussi, 2010), 3 – Futschöltal (F) (Patzelt, 2019), 4 – Las Gondas (Dietre et al., 2014; Nicolussi, 2010) (c) Updated geological and geomorphological maps of Jamtal

105

110 and (d) Laraintal, modified from Fuchs and Oberhauser (1990). Viewpoints from which photos in Figs. 3 and 5 were taken are denoted with white-blue symbols. AUT – Austria; CH – Switzerland. Digital Elevation Model (DEM) provided by © Land Tirol (resolution 1 m).

The closest meteorological station recorded a mean annual atmospheric temperature of 3.1° C averaged over the period 1981–2010 (station number 101949; 1587 m a.s.l.; BMNT, 2016). The mean annual precipitation measured at the same station amounts to 1087 mm/yr for the same reference period, with snow cover present on 175 days/yr on average. Precipitation is also measured in proximity to the Jamtalferner tongue (2400 m a.s.l.) and yields an annual mean of 1507 mm/yr (reference period 1989–2017; Fischer et al., 2019). Since the end of the LIA, all Silvretta glaciers have retreated in response to almost continuous warming and have lost about two thirds of their areas relative to the LIA maximum (Fischer et al., 2021; WGMS, 2018). Geodetic mass balance estimates of Silvretta glaciers document increased losses within recent decades (Fischer et al., 2021). While geodetic mass balance averaged across all Silvretta glaciers amounted to -0.2 ± 0.1 m w.e./yr in the reference period from 1969–2002, this value increased to -0.8 ± 0.2 m w.e./yr between 2006–2018. For Jamtalferner and Larainferner, mass losses from 2006–2018 are quantified to -1.0 ± 0.2 m w.e./yr and to -0.8 ± 0.2 m w.e./yr, respectively.

The Silvretta Massif contains some of the oldest rocks of the Eastern Alps with a presumed depositional age in the Precambrian followed by several metamorphic events (Bertle, 1973; Maggetti and Flisch, 1993). Lithology in the region consists of crystalline rocks, which are part of the Upper Eastern Alpine tectonic unit, more precisely the Silvretta-Seckau nappe (Fuchs and Oberhauser, 1990; Schuster, 2015). Rocks at the Jamtal and Laraintal formed during the Permian, experienced repeated faulting prior to and throughout the Alpine orogeny and are thus highly metamorphic (Friebe, 2007, and references therein). Predominant rock types in the study area are different gneiss variations and amphibolites (Fig. 1c and 1d). Rock samples that were collected from moraines had quartz yields – the target mineral for the applied cosmogenic nuclide method – ranging from 0.3 % to 26.1 % with a median of 3.9 % (Table S1).

130 3 Methods

3.1 Principle of ^{10}Be Surface Exposure Dating (SED)

Glaciers erode into bedrock and transport rock material to their margins. When glaciers are stationary for several years (or longer), linear landforms – moraines – that consist of glacial debris, accumulate at their ice margins. Dating these moraines unravels the timing of glacier stabilization or rather the beginning of glacier retreat, and allows the reconstruction of glacier configurations of the past. For ^{10}Be SED – the cosmogenic nuclide approach used in this study – rock samples were extracted from boulder surfaces deposited along moraines. Sub-rounded to rounded boulders were prioritized for sampling. Compared to angular boulders, (sub-)rounded ones are more likely to have been carved out of bedrock by glacial flow. When these englacially or subglacially transported boulders melted out of glacial ice, their surfaces were for the first time exposed to high-energetic cosmic radiation. Secondary cosmic rays interact with Si and O in quartz and produce radionuclides, among others

140 ^{10}Be (Lal, 1988). The annual production rate of ^{10}Be is well constrained today and the accumulation of the radionuclide is used to determine the duration of exposure by quantifying ^{10}Be in rock surfaces of moraine boulders (Gosse and Phillips, 2001).

3.2 Geomorphological mapping and rock sample collection

We build upon geological and geomorphological maps, which were produced in previous studies and which were the basis for further detailed field investigations in the years of 2018 to 2020 (Fischer et al., 2019; Fischer et al., 2015; Fuchs and Oberhauser, 1990; Hertl, 2001). In the course of a general survey of the Jamtal and the Laraintal area, we updated preexisting maps according to our own mapping. We then focused on the mapping of glacial features and placed particular emphasis on the fine structure of moraines that were presumably deposited during the LIA, and ridges that were identified outboard these moraines. The dating of these structures promises to shed light onto the timing of climate perturbations, which favored moraine formation when glaciers were still relatively large compared to their present-day configurations. In order to ensure robust landform age calculations, we took three or more rock samples from each selected ridge, provided that they fulfilled our sample selection criteria described in detail in Braumann et al. (2020; Appendix S-Table 1). In total, 27 samples were extracted from boulders using hammer and chisel, and an electric saw. Geographic coordinates of sampled boulders were measured using a hand-held GPS device. Strike and dip of sampled surfaces were quantified with a geological compass. Sample elevations were taken from the DEM of 2018 (x-y-z resolution 1 m, © Land Tirol). Shielding was calculated using the ArcGIS ‘Skyline’ Toolbox.

3.3 Sample preparation and age calculation

All samples were processed at the Cosmogenic Isotope Laboratory of the Lamont-Doherty Earth Observatory (LDEO) following the geochemical standard protocol for quartz preparation and the extraction of ^{10}Be (LDEO, 2012a, b; Schaefer et al., 2009). Prior to quartz digestion using concentrated hydrofluoric acid, approximately 180 μg of the LDEO ^9Be carrier made of deep-mine Beryl was added to the samples (carrier concentration c. 1000 ppm). Samples LAR-19-14 and LAR-19-16 with extremely low quartz yields of 0.3–0.4 %, equivalent to c. 2.6–2.7 g of purified quartz per sample, were treated differently. Due to their small sample sizes combined with our EH age estimates, we expected low total numbers of cosmogenic ^{10}Be atoms in the samples. For these samples we adopted a sample preparation procedure where ^9Be carrier is reduced and replaced with Fe carrier. Only c. 100 μg ^9Be carrier was added during digestion and then c. 100 μg of Fe (concentration 1000 mg/L) was added to the samples prior to hydroxide precipitation (as $\text{Be}(\text{OH})_2 + \text{Fe}(\text{OH})_2$), subsequent to the cation columns step. The Fe addition allowed us to maintain a manageable sample volume, which facilitates the handling of the samples. This procedure was recommended for exceptionally low-level samples based on unpublished experimental data that suggests it optimizes ^{10}Be counting efficiency at the Center for Accelerator Mass Spectrometry (CAMS) facility, Lawrence Livermore National Laboratory (LLNL) (pers. comm. A. J. Hidy). We proceeded with subsequent steps of sample preparation according to the LDEO protocol (LDEO, 2012a). Isotope ratios ($^{10}\text{Be}/^9\text{Be}$) in samples were measured at CAMS-LLNL using the 07KNDSTD3110 standard with a $^{10}\text{Be}/^9\text{Be}$ ratio of 2.85×10^{-12} (Nishiizumi et al., 2007).

Exposure ages were calculated using the online calculator formerly known as the CRONUS-Earth online calculator (v3; Balco et al., 2008). We applied the regional Swiss production rate (Claude et al., 2014), and ‘Lm’ scaling to account for site specific nuclide production. All ^{10}Be boulder ages are based on the arithmetic mean of 3–5 replicate AMS measurements and are presented with 1σ analytical uncertainties, including a 1 % uncertainty on the carrier concentration. Moraine ages represent arithmetic means of exposure ages of three or more boulder ages. Uncertainties reported with moraine ages include the production rate uncertainty (for the Swiss production rate c. 6.3 %) in addition to the analytical and carrier uncertainty, and were propagated in quadrature. Identification of potential outliers was accomplished following the χ^2 statistics implemented in the online calculator.

Corrections of exposure ages for seasonal snow cover were not applied. First, samples were primarily taken from boulder tops or their upper sections, preferably located at windswept locations to minimize potential snow cover (see Supplement, sect. 5). Second, if exposure ages were significantly influenced by snow effects, age dispersion would be expected among boulders embedded in the same moraine, but whose shapes and exposures vary. Our data do not show a significant bias of this type; therefore, snow cover effects appear to be insignificant at our study sites. However, if a snow correction was applied to a boulder assuming a 1 m thick snowpack that is preserved over four months and has a snow density of c. 0.3 g/cm^3 (estimates based on modern values; BMLRT, 2021), its exposure age would become around 5–6 % older (Gosse and Phillips, 2001). The preservation of striations on rock surfaces and the general condition of boulder surfaces in the valleys suggest that erosion has not significantly impacted their surfaces since deposition. Therefore, all ages that are presented and discussed in the following represent values without any erosion correction applied. However, in some studies addressing the Holocene time scale, an erosion rate of 1 mm/ka is used (André, 2002). To test the sensitivity of our ages to this erosion rate, we recalculated our data using this value and find that ages become at most 1 % older (median 0.8 %; Table S3), an age shift that is not significant on the 1σ level and that does not change our interpretation of the data.

4 Results

4.1 Geomorphology

In both valleys, distinct moraine sets, which mark Holocene paleo-ice margins, are preserved. We numbered the moraines from the youngest J0 to the oldest J5 at the Jamtal, and moraines at the Laraintal from L1 to L5 in analogy.

4.1.1 Jamtal

At the Jamtal, the innermost moraine we address in this article, is **J0**. The moraine was deposited at the left-lateral valley flank, inboard the presumable LIA moraine (**J1**; Fig. 2). J0’s age of deposition falls into the period between the end of the LIA and the turn of the 20th century according to Fischer et al. (2019) and a historical map that was composed in the years between 1870–1877 (K. u. k. Militärgeographisches Institut, 1870-1887). J0 and J1 are punctuated by an approximately 100 m wide drainage channel, which evolved along the flowline of a former tributary glacier (Totenfeld, Fig. 3a and A1c). With numerous

bedrock outcrops along the valley flank and a slope of $>35^\circ$ the terrain is steep and impedes the accumulation moraines higher up (Fig. 3b). On the right-lateral side, in turn, slope angles are lower ($5\text{-}35^\circ$) and allowed the formation and preservation of multi-ridge moraine complexes (J1 to J4; Fig. 2, A1d and A1f). J1 consists of fresh, blocky debris with little to no lichen colonialization. Pioneer plants grow in voids in between blocks, whereas segments with more fine sediment are covered with a thin vegetation layer (Fig. A1a–c). In some sections, J1 is several tens of meters wide, which contrasts with the adjacent J2 with a maximum width of about 8 m. J2 is located in a depression between J1 and a till covered slope, on top of which J3 and J4 were deposited (Fig. A1e–f). J2 is rich in fine sediment but does not feature boulders that meet our ^{10}Be sample selection criteria.





Figure 2: Holocene moraine chronology of Jamtal. J0 (red) has been deposited after the LIA but prior to the 20th century (Fischer et al., 2019; K. u. k. Militärgeographisches Institut, 1870-1887). J1, J3R and J4R were dated in this study. Ages along the J1 moraine (pink) indicate that Jamtalferner reached its historical maximum during the second half of the 18th century, and during the Neoglacial (NG), c. 500 CE. Earlier phases of glacier stabilization that exceeded subsequent Holocene culminations, are evidenced by moraines J3R and J4R, which both date to the EH. DEM provided by © Land Tirol (resolution 1 m).

215



220 **Figure 3: Photographs of Jamtal. (a) View toward Jamtalferner and Totenfeld. ^{10}Be sample locations are marked with colored circles (pink – LIA; black – Neoglaical, cyan and blue – EH). (b) Downvalley view depicting sample locations JAM-18-06 (pink) and JAM-18-07 (black). (c) Valley flank below Chalausferner and Augustenferner with sampled boulders along the innermost (pink), the middle (cyan) and the outer (blue) right lateral moraine of former Chalausferner. For the broader context of the individual sites, see Figure 1c.**

225 **J3** and **J4**, two parallel, curved moraines about 20 to 30 m further uphill relative to J2, are right-lateral moraines of Chalausferner, and evidence the convergence of this tributary glacier and the Jamtalferner (Fig. 3c). Boulder surfaces embedded in these moraines are populated with black and green lichens and show signs of weathering, for instance cracks and exfoliation. On the valley floor, a moraine with a frontal position at an altitude of about 2120 m a.s.l. is preserved. The ridge consists of weakly weathered material and is in this respect as well as with respect to geometry the terminal equivalent of the


230 lateral **J1** moraine (Fig. 2 and A2c). This correlation is in accordance with glacier outlines of the Austrian Glacier Inventory

(AGI; Fischer et al., 2015), with a geomorphological map compiled by Hertl (2001:226) and with results from a recent study on vegetation dynamics at the Jamtal, which includes ice-margin reconstructions since the end of the LIA (Fischer et al., 2019). North of the terminal section of J1 is an area covered with angular and subangular blocks, whose surfaces are significantly more weathered compared to J1, and which exhibit extensive lichen population (Fig. A2c–e). Many blocks have cracks and are fractured, which may indicate impacts associated with gravitational movement. The morphology outboard J1 is convex – unusual for in situ rockfall deposits, which typically form lobate structures with large boulders in frontal positions. We therefore hypothesize that these deposits stem from rock failures along one (or both) valley flank(s) farther uphill. The material collapsed onto the formerly larger glacier and was transported downstream through glacial flow. As the glacier retreated, these rockfall deposits melted out and accumulated on the valley floor. An additional argument supporting this scenario is the provenance area of a potential rockfall event, which could not clearly be identified along the surrounding walls and peaks. The blockfield was in part overprinted by one (or multiple) glacier advances, as evidenced by the position of moraine J1. A set of ridges in the right latero-frontal section outboard the J1 moraine appears to be somewhat displaced (Fig. 2 and A2f). A scarp above this moraine set and a stabilized sliding mass below caused an offset of formerly connected crests. Together with rockfall deposits from bedrock outcrops in higher up sections, these moraines are not considered as prime candidates for ¹⁰Be sampling. We also avoided structures close to the Jamtal hut. Even though we identified several ridges in its vicinity, which were presumably deposited during the Holocene, land surfaces in this area have been anthropogenically altered. For instance, the road leading up to the hut and the hut itself are built on moraines (Hertl, 2001:77). The same argument applies to a ridge at an elevation of approximately 2045 m a.s.l. denoted as **J5** in (Fig. 2 and A2a–b). Although the structure may be interpreted as a Holocene terminal moraine it was rejected for sample collection as it is in part anthropogenically overprinted and may comprise boulders disintegrated from the right-lateral slope above. Evidence of older, LG terminal positions farther downstream is scarce. Hertl (2001:77) describes a lineament that dips to the valley floor about 2.5 km downstream of J5, at an elevation of 1900–1920 m a.s.l. The author tentatively interprets the structure as a latero-frontal moraine deposited towards the YD termination. Around 8 km downvalley from J5, a tripartite moraine set (Gaffelar settlement) is attributed to an earlier YD phase, yet not the YD maximum. Lateral LG moraines are absent in the main valley, but are preserved in the Futschöl tributary valley, which joins the Jamtal from the East in the area of the Jamtal hut (Fig. 1b).

4.1.2 Laraintal


At the Laraintal, we focused on valley sections outlined in Figure 1d and detailed in Figure 4. Texture, relative positions, and structure of moraines in this valley resemble moraine sets at the Jamtal. L1 – the presumable LIA ridge – consists of fresh, sparsely vegetated debris and is traceable along both valley flanks. On the eastern side, we identified a fine-structured set of moraines which we refer to as L2, L3R and L4R in Figure 4. Similar to J2, L2 with a width of approximately 8–10 m is less prominent compared to L1 (>20 m; Fig. A3b). Also, L2 has a high fine sediment content with no large boulders on the crest. The fine-grained texture of L2 contrasts the subsequent outer LR3 and LR4 ridges, which are both blockier. The distinct nature

of L2 (and also J2) in comparison with the outer moraines allows speculations about different ice dynamics that lead to their
 265 formation, for instance glacier advance (push moraine) versus glaciers in equilibrium (dump moraines). LR3 has a broader,
 but less pronounced crest compared to LR4, the outermost moraine in this valley section (Figure 5c). Boulders of both
 moraines, L3R and L4R, were sampled for ^{10}Be extraction. Vegetation cover has developed on the surfaces of both ridges and
 soil formation processes are advanced on the glacier-distal side of L4R.



270

Figure 4: Holocene moraine chronology of Laraintal with moraine ages displayed. The ^{10}Be sample collected from the L1 suggests an early LIA advance around 1300 CE. Lateral and terminal moraines outboard L1 yield EH ages, which agree well with the Jamtal moraine record. DEM provided by © Land Tirol (resolution 1 m).



275 **Figure 5: Photographs of Laraintal. (a) Terminal moraine section with sample locations marked with circles. (b) Left lateral valley flank with LAR-19-22 (cyan), LAR-19-23 (pink) and LAR-19-22 (cyan) from left to right. (c) Right lateral moraine set with LAR-19-15 in blue in the foreground, and LAR-19-13 and LAR-19-14 in the background (cyan). For the broader context of the individual sites, see Figure 1d.**

280 Analogous to the Jamtal, the terrain at the western valley side is generally steeper compared to the eastern side, with slope angles of c. $>30^\circ$. One exception is a riegel (bedrock bar) consisting of amphibolite that forms the basis for a relatively flat area (Fig. 5b and A3d). Geomorphology in this section has been influenced and shaped by debris flows, slope erosion and the

thawing of permafrost, in addition to glacial processes. Two ridges were deposited directly below the headwall, amid a mix of scree, till and rockfall deposits. According to Hertl (2001:69), these ridges may mark late LG ice margins (Fig. 4) with a terminal equivalent around 4 km downstream at an elevation of c. 1870 m a.s.l. In contrast to this complex section, ridges **L1** and **L3L** farther away from the headwall and closer to the western edge of the riegel are well preserved and continue northwards and below the riegel. **L3L** can be traced along the valley flank until it is cut by a debris cone. **L1** dips towards the valley floor, where its frontal segment splits up in at least two ridges. The area outboard the **L1** terminus is covered with glaciofluvial sediments, and is delimited by an arc-shaped blocky structure traversing the valley downstream, at an altitude of c. 2180 m a.s.l. (Fig. 5a, **L3T**). On the structure's glacier-proximal side, fine sediment has accumulated and gives this landform an almost terrace-like character with the blocky ridge acting as a barrier. The ridge is dissected by a creek (Larainbach); its western end partly overburdened by the same debris cone, which cuts into **L3L**. Overall, the sedimentary composition, shape, and orientation of the ridge indicate that **L3T** is a moraine. An almost identical landform (**L4T**) replicates outside **L3T**, at a horizontal distance of 50–60 m and with its crest at an elevation of about 2170 m a.s.l. (Fig. 5a). Both moraines, **L3T** and **L4T**, evidence former terminal positions of Larainferner and are correlated with lateral moraines **L3-4L** at the left-lateral side and with **L3R** and **L4R** at the right-lateral side.


Approximately 200–250 m further downstream, at an elevation of 2130 m a.s.l., is another set of ridges, on one of which a small hut (“Zollhütte”) was built (Fig. 4). This Zollhütte ridge (**L5**) is framed by a block field consisting of a blend of angular and rounded boulders. A massive debris cone west of Zollhütte, which has a layer of coarse and medium-sized, fresh material on top, points to continuous sediment supply from the left-lateral wall. To the east, a scarp with a concave surface below indicates former (and possibly ongoing) sliding processes directed towards Zollhütte. Moreover, rockfall events with material disintegrating from the wall below the “Hoher Kogel” peak have been witnessed during field work in the year of 2019, with boulder volumes of multiple cubic meters that have crashed on the valley floor (Fig. 1d and A4a–b). Such events have probably also occurred in the past as the wall exhibits multiple lighter sections, which are indicative of removed material and thus of previous rock failures. Due to the manifold processes, which impacted the Zollhütte area, the **J5** ridge is scientifically risky to tackle with SED of boulders, even though it probably delimits a terminal glacier position.

4.2 ¹⁰Be results

¹⁰Be analytical data of all 27 boulder samples and corresponding age information are listed in Table 1 (Jamtal) and Table 2 (Laraintal). Kernel plots of moraine ages are displayed in Figure 6 (LIA) and Figure 7 (EH). Ages are reported for each valley individually and are discussed according to their landform number in ascending order, from **J1** to **J4** and from **L1** to **L4**. All exposure ages fall into three periods of high(er) glacier activity within the past 12 ka: the LIA, the first millennium common era, and the EH.

4.2.1 Jamtal

315 Five samples were collected from boulders along moraine **J1** (Fig. 2, Table 1). Three of them (JAM-18-06, JAM-18-17, JAM-18-18) were deposited during the second half of the 18th century and yield a rounded mean age of **260 ±20 yrs** (Fig. 6). The other two samples, JAM-18-07 (1500 ±50 yrs) and JAM-18-16 (1070 ±20 yrs), both produce neoglacial ages. Since J2 lacks suitable boulders for ¹⁰Be sampling, the subsequent dated ridge is J3R. Based on three boulder ages of moraine **J3R** (JAM-18-01: 11,020 ±200 yrs, JAM-18-02: 11,280 ±180 yrs, JAM-18-03: 11,850 ±220 yrs), we calculate a landform age of 11,380 ±830
320 yrs, rounded to **11.4 ±0.8 ka**. **J4R** outside J3R gives a moraine age of 10,750 ±690 yrs (**10.8 ±0.7 ka**) derived from boulder ages of samples JAM-18-04 (10,680 ±200 yrs), JAM-19-21 (10,920 ±210 yrs) and JAM-19-22 (10,660 ±130 yrs) (Fig. 7a–b).



325 **Figure 6: Kernel plot of LIA ages produced from boulders embedded in the historical moraine at Jamtal. The gray-shaded bar illustrates the 1 σ Standard Deviation (SD) of the landform age calculated from analytical uncertainties of individual samples. Red dashed lines add the production rate uncertainty and the uncertainty of the carrier concentration to the 1 σ SD and indicate the total uncertainty. Green dotted lines show the Standard Error (SE), which describes the dispersion of different sample means from the population mean.**

330 Table 1: ^{10}Be analytical data and corresponding exposure ages of Jamtal samples. Samples were analyzed at the CAMS-LLNL. All
 335 samples were measured against the 07KNSTD3110 standard with a ratio of 2.85×10^{-12} (Nishiizumi et al., 2007). Two procedural
 blanks were processed with each batch of samples with ratios ranging from 2.8 to 9.1×10^{-16} (Supplement, Table S2). The ^{10}Be
 background contaminations measured in the blanks were subtracted from the samples. Exposure ages were calculated with the
 calculator formerly known as CRONUS-Earth online calculator (v3; Balco et al., 2008), using the Swiss ^{10}Be production rate (Claude
 et al., 2014) and choosing the ‘Lm’ scaling scheme. Ages are calculated relative to the sampling year denoted by the first number in
 the sample ID and are rounded to the nearest 10 years. Uncertainties of boulder ages include the 1σ analytical error and a 1 %
 uncertainty on the carrier concentration.

	Sample ID	Latitude	Longitude	Elev.	Av. Thickness	Shielding factor	Quartz mass	^{9}Be added	$^{10}\text{Be}/^{9}\text{Be}$ ratio $\pm 1\sigma$ analytical unc. (10^{-14})	^{10}Be atoms $\pm 1\sigma$ analytical unc.	^{10}Be conc. $\pm 1\sigma$ analytical unc.	^{10}Be exposure age and carrier unc.
	[DD]	[DD]	[m a.s.l.]	[cm]	[g]	[μg]			[atoms]	[atoms/g qtz]	[yrs]	
J1	JAM-18-06	46.8823	10.1684	2223	1.5	0.9317	13.3800	186	0.70 ± 0.09 (12.6%)	86597 ± 10881	6141 ± 772	270 ± 30
	JAM-18-17	46.8774	10.1723	2297	1.1	0.9760	10.4745	186	0.55 ± 0.07 (12.5%)	67819 ± 8457	6052 ± 755	240 ± 30
	JAM-18-18	46.8776	10.1725	2289	2.9	0.9748	23.4243	186	1.30 ± 0.07 (5.4%)	161887 ± 8777	6528 ± 354	270 ± 20
NG	JAM-18-07	46.8820	10.1682	2231	1.6	0.9044	17.4383	187	4.59 ± 0.15 (3.3%)	572357 ± 18941	32307 ± 1069	1500 ± 50
	JAM-18-16	46.8766	10.1718	2316	2.4	0.9770	31.2824	186	6.51 ± 0.14 (2.1%)	804960 ± 16997	25445 ± 537	1070 ± 20
J3R	JAM-18-01	46.8766	10.1736	2350	1.6	0.9557	31.1628	185	67.00 ± 1.22 (1.8%)	8250915 ± 149649	264480 ± 4797	11020 ± 200
	JAM-18-02	46.8767	10.1735	2339	1.7	0.9752	11.4783	186	25.38 ± 0.41 (1.6%)	3151014 ± 51230	274134 ± 4457	11280 ± 180
	JAM-18-03	46.8770	10.1734	2328	4.0	0.9759	7.2439	186	16.50 ± 0.31 (1.9%)	2045059 ± 38161	281704 ± 5257	11850 ± 220
J4R	JAM-18-04	46.8771	10.1739	2335	2.6	0.9748	20.2670	183	42.55 ± 0.79 (1.9%)	5196242 ± 96265	255946 ± 4742	10680 ± 200
	JAM-19-21	46.8776	10.1738	2318	3.2	0.9756	10.4519	185	21.79 ± 0.41 (1.9%)	2703364 ± 50631	258051 ± 4833	10920 ± 210
	JAM-19-22	46.8773	10.1738	2329	1.6	0.9729	10.6823	185	22.09 ± 0.28 (1.2%)	2739711 ± 34113	255888 ± 3186	10660 ± 130


340 4.2.2 Laraintal

Sample LAR-19-23 (700 ± 20 yrs) stems from **L1** and might capture a LIA maximum early in the 14th century (Fig. 4, Table
 2). The deposition of the left-lateral **L3-4L** moraine is constrained by LAR-19-22 with a boulder age of $11,210 \pm 210$ yrs and
 by LAR-19-24 dated to $10,930 \pm 210$ yrs. These ages are statistically indistinguishable and cannot clearly be assigned to either
 the inner moraine (L3) or the outer one (L4). Therefore, they are included in landform age calculations of both ridges. The
 345 right-lateral **L3R** ridge yields boulder ages of $11,120 \pm 210$ yrs (LAR-19-13) and $11,200 \pm 280$ yrs (LAR-19-14). The age of
 the terminal segment of L3-4L – L3T – is derived from samples LAR-18-09 ($11,540 \pm 200$ yrs), LAR-19-17 ($11,070 \pm 210$ yrs),
 LAR-19-18 ($11,330 \pm 290$ yrs), LAR-19-19 ($10,730 \pm 200$ yrs), LAR-19-20 ($11,480 \pm 210$ yrs) and results in a landform age of
 $11,230 \pm 780$ yrs. Based on our mapping and dating results, we are confident that moraine segments L3R and L3T, and
 potentially L3-4L, can be attributed to the same glacier advance or stabilization. Therefore, we aggregate all nine boulder ages
 350 and compute a moraine age of $11\,180 \pm 750$ yrs (L3: **11.2 ± 0.8 ka**; Fig. 7c). Based on the same reasoning, we combine L3-4L,
L4R (LAR-19-12: $10,890 \pm 180$ yrs, LAR-19-15: $10,860 \pm 200$ yrs, LAR-19-16: $11,060 \pm 220$ yrs – landform age $10\,940 \pm 690$
 yrs) and **L4T** (LAR-18-03: $10,160 \pm 190$ yrs, LAR-18-10: $10,880 \pm 210$ yrs, LAR-19-21: $10,660 \pm 200$ yrs – landform age
 $10\,570 \pm 760$ yrs) and suggest a moraine age of $10,830 \pm 750$ yrs (L4: **10.8 ± 0.8 ka**; Fig. 7d).

355 Analytical results from samples, which were spiked with Fe, show that ages calculated from both samples are consistent with
boulder ages obtained for the same landforms, but processed according to the standard protocol (L3R: LAR-19-13, L4R: LAR-
19-12 and LAR-19-15). Analytical uncertainties of corresponding samples amount to 2.0 % (LAR-19-16) and 2.5 % (LAR-
19-14) and are within the expected range of ^{10}Be AMS measurement uncertainties at LLNL-CAMS (Rood et al., 2013). By
replacing a portion of the ^9Be carrier with Fe, we achieved similar analytical precision as with routinely processed samples,
360 but with only a quarter of the sample mass used. Our results suggest that the substitution of a fraction of ^9Be carrier using Fe
is a viable and promising advancement in the sample preparation protocol that extends the application field of the ^{10}Be SED
method to younger samples and more challenging lithologies.

365 **Table 2: ^{10}Be analytical data and corresponding exposure ages of Laraintal samples. For details on analytics, processing or age
calculation see captions of Table 1 and Supplement, Tables S1 and S2. Samples LAR-19-14 and LAR-19-16 were spiked with Fe in
addition to (a reduced amount of) ^9Be carrier.**

	Sample ID	Latitude	Longitude	Elev.	Av. Thickness	Shieldin g factor	Quartz mass	^9Be added	Fe added	$^{10}\text{Be}/^9\text{Be}$ ratio $\pm 1\sigma$ analytical unc. (10^{-14})	^{10}Be atoms $\pm 1\sigma$ analytical unc.	^{10}Be conc. $\pm 1\sigma$ analytical unc.	^{10}Be exposure age $\pm 1\sigma$ analytical unc. and carrier unc.
		[DD]	[DD]	[m a.s.l.]	[cm]		[g]	[μg]	[μg]		[atoms]	[atoms/g qtz]	[yrs]
L1	LAR-19-23	46.9234	10.2216	2292	1.5	0.9433	30.2276	181		4.20 ± 0.10 (2.3%)	507590 ± 11631	16570 ± 380	700 ± 20
L3	LAR-18-09	46.9272	10.2225	2177	2.0	0.8899	18.9009	186		35.05 ± 0.62 (1.8%)	4340818 ± 76283	229187 ± 4028	11540 ± 200
	LAR-19-17	46.9267	10.2223	2179	1.5	0.9424	10.1141	182		19.47 ± 0.36 (1.9%)	2366107 ± 43845	233453 ± 4326	11070 ± 210
	LAR-19-18	46.9268	10.2223	2178	2.2	0.9427	10.5868	182		20.76 ± 0.54 (2.6%)	2520432 ± 65068	237607 ± 6134	11330 ± 290
	LAR-19-19	46.9267	10.2226	2178	1.6	0.9413	10.2762	185		18.72 ± 0.35 (1.9%)	2319925 ± 42983	225277 ± 4174	10730 ± 200
	LAR-19-20	46.9267	10.2224	2180	2.5	0.9434	10.1923	182		20.22 ± 0.38 (1.9%)	2460269 ± 45624	240901 ± 4467	11480 ± 210
	LAR-19-13	46.9224	10.2258	2303	1.3	0.9604	10.2457	180		22.20 ± 0.41 (1.9%)	2676391 ± 49575	260739 ± 4830	11120 ± 210
	LAR-19-14	46.9225	10.2258	2303	2.0	0.9584	2.6078	102	99	10.02 ± 0.25 (2.5%)	686210 ± 17191	260711 ± 6531	11200 ± 280
	LAR-19-22	46.9245	10.2213	2256	1.9	0.8546	10.3313	181		19.30 ± 0.36 (1.9%)	2335406 ± 43280	225574 ± 4180	11210 ± 210
L4	LAR-19-24	46.9225	10.2201	2357	1.7	0.9581	10.4515	185		22.38 ± 0.42 (1.9%)	2767560 ± 51798	264203 ± 4945	10930 ± 210
	LAR-18-03	46.9273	10.2220	2162	1.5	0.9245	24.1911	186		40.30 ± 0.75 (1.9%)	5004558 ± 92711	206505 ± 3826	10160 ± 190
	LAR-18-10	46.9272	10.2225	2161	2.8	0.9401	6.2606	187		11.25 ± 0.21 (1.9%)	1404082 ± 26340	223566 ± 4194	10880 ± 210
	LAR-19-21	46.9273	10.2222	2160	2.3	0.9383	10.3970	180		18.92 ± 0.35 (1.9%)	2281226 ± 42291	218937 ± 4059	10660 ± 200
	LAR-19-12	46.9231	10.2261	2295	1.4	0.9585	10.0087	182		20.82 ± 0.33 (1.6%)	2536338 ± 40705	252920 ± 4059	10890 ± 180
	LAR-19-15	46.9240	10.2259	2275	3.5	0.9563	10.4801	182		21.05 ± 0.39 (1.9%)	2561640 ± 47473	243958 ± 4521	10860 ± 200
LAR-19-16	46.9238	10.2260	2280	3.2	0.9453	2.5099	104	99	9.01 ± 0.18 (2.0%)	627353 ± 12552	247430 ± 4950	11060 ± 220	



370 Figure 7: Kernel plots of Holocene moraine ages. Left column (a) and (c): inner dated EH moraines at Jamtal and Laraintal. Right
 375 column (b) and (d): outermost dated EH moraines at Jamtal and Laraintal. (e) Synthesis of moraine ages across both valleys grouped
 to Moraine Formation Intervals (MFI) MFI 3-4 yielding an age of $11,030 \pm 740$ yrs rounded to 11.0 ± 0.7 ka ($n=19$; outliers: JAM-18-
 03 and LAR-18-03). Gray-shaded bars illustrate 1σ Standard Deviations (SD) of landform age uncertainties calculated from
 analytical uncertainties of individual samples. Red dashed lines add the production rate uncertainty (c. 6.3 %) and the uncertainty
 of the carrier concentration to the 1σ SD and indicate total uncertainties, which results in conservative uncertainty estimates. Green
 dotted lines indicate Standard Errors (SE).


5 Discussion

5.1 The moraine record of the past two millennia

The classical LIA moraines of both valleys (J1 and L1) feature boulders deposited within the expected time interval, i.e. between 1250 and 1850 CE (Fig. 8f; e.g., Grove, 2004; PAGES 2k Consortium, 2013). An early LIA advance of Larainferner to L1's position is suggested by LAR-19-23 and may have occurred at the beginning of 14th century. Three consistent boulder ages from J1 are aggregated to a mean age of 260 ± 25 yrs and indicate an advance of Jamtalferner between c. 1735 and 1790 CE. A recent geochronological study in the adjacent Ochsental comes to remarkably similar results with boulder ages from the LIA moraine yielding a mean age of 260 ± 30 yrs (Fig. 8e; Braumann et al., 2020). Glacier advances during this period are also documented in the Eastern Alps, for instance at the Zillertal and at the Ötztal (Nicolussi, 2013; Pindur and Heuberger, 2010), in the Central Alps at the Lower Grindelwald glacier (Zumbühl and Nussbaumer, 2018) and in the Western Alps at the Mer de Glace (Nussbaumer et al., 2007). High glacial activity during the second half of the 18th century with termini coming close to, or reaching their LIA maximum, is congruent with a phase of decreased summer temperatures detected in proxy records in the vicinity of our study site (Fig. 8a–b; Fohlmeister et al., 2013; Ilyashuk et al., 2019; Larocque-Tobler et al., 2010a; Vollweiler et al., 2006), and with reconstructed summer and mean annual temperatures from Greenland ice cores (Fig. 8d; Buizert et al., 2018; Kobashi et al., 2017).

Besides LIA-aged boulders along the LIA moraine we sampled two blocks of J1, which were deposited during the first millennium of the common era. The younger boulder, JAM-18-16, dates to the beginning of the Medieval Warm Period (MWP). By that time, glaciers in the region were likely smaller relative to their LIA maxima (e.g., Solomina et al., 2016). The boulder's position and its bedding were re-evaluated in the field after age calculation, and we cannot exclude that the boulder has tilted (Fig. S7). Therefore, we interpret the exposure age as a minimum age. The older neoglacial boulder, JAM-18-07, was exposed 1500 ± 50 yrs ago, which is again in good agreement with the neighboring Ochsental chronology, where a block in an identical setting (embedded in the classical LIA moraine) – was dated to 1500 ± 40 yrs (Fig. 8e; Braumann et al., 2020). There is a possibility of pre-exposure of both samples that could produce erroneous Neoglacial ages. However, evidence for a period of glacier advance in the Eastern Alps during the 5th and 6th century CE was found beyond the Silvretta region, for instance in sediment profiles and peat cores in the forefield of Fernaufener, Mittelbergferner and Simonykees (Patzelt, 2016; Patzelt and Bortenschlager, 1973). This timing coincides with prominent episodes of glacier advance in the Western Alps, most notably at the Aletsch glacier (Holzhauser et al., 2005), and at the Miage and Mer de Glace, both in the Mont Blanc massif (Deline and Orombelli, 2005; Le Roy et al., 2015). Concurrent glacier advances have also been reported from Alaska, Iceland, Scandinavia and from Greenland (Barclay et al., 2009; Biette et al., 2020; Solomina et al., 2016). Glacier advance during this period is consistent with decreasing summer temperatures (Fig. 8c–d) and higher precipitation rates in Europe (e.g., Büntgen et al., 2011). This regional climate perturbation, which is often referred to as Dark Ages Cold Period (DACP) and which occurred in tandem with the migration period in Europe, began around 400 CE and lasted into the 8th century CE in the region (e.g., Helama et al., 2017). During that time, the Atlantic Meridional Overturning Circulation (AMOC), which transports

410 heat from the South Atlantic towards the North was weakened (Thornalley et al., 2018), which lead to cooling in the Nordic region. Potential volcanic eruption(s) in the Northern Hemisphere in the year of 536 CE may have amplified cooling across Europe and define the onset of the recently postulated Late Antique Little Ice Age (LALIA; 536 to c. 660 CE; Büntgen et al., 2016). Consistent with the timing of the regional DACP, Helama et al. (2021) suggests centennial-scale phases in Northern Europe during the Holocene, which resemble the LIA climatic regime, one among them beginning around 540 CE.



415

420 **Figure 8:** Youngest part of the ^{10}Be chronology from Jamtal and Laraintal correlated with climate proxy data from the Alps (local records in dark blue), Europe (green) and the Greenland Ice Sheet (GrIS in red) covering the past c. 2000 yrs. Proxy records indicate cooler climate conditions during the Dark Ages Cold Period (DACP) and during the Little Ice Age (LIA), synchronous with periods of moraine formation in the Silvretta region. Chironomid-inferred (CI) July temperatures from (a) Mutterbergersee in the Stubai Alps, Austria (Ilyashuk et al., 2019) approximately 70 km E of study site (north of the Alpine drainage divide) and (b) lake Silvaplana around 60 km SW of study site (Engadin, Switzerland; south of the Alpine drainage divide) (Larocque-Tobler et al., 2010a). (c) European summer temperature anomalies (reference period 1961-1990, 60-year low-pass filter) identified in tree-ring chronologies (Büntgen et al., 2011). (d) Mean summer temperature reconstructions (JJA) derived from nine Greenland ice cores (Buizert et al., 2018). (e) ^{10}Be ages of boulders sampled from the presumable LIA (Holocene composite moraine) from Ochsental (OcG) (Braumann et al., 2020) and from (f) Jamtal and Laraintal (this study). RWP – Roman Warm Period; MWP – Medieval Warm Period.

425

In summary, samples collected along the classical LIA moraine at the Jamtal (this study) and at the adjacent Ochsental (Braumann et al., 2020) yield ages that fall into the regional DACP and the LIA. These results are consistent with the timing of glacier advances across the Alps and in other places of the Northern Hemisphere. The advance of Silvretta glaciers coincides

with cooling trends captured in local, regional and hemispheric proxy data. Moraines J1 and L1 are probably composite
430 moraines that have formed ice margins at least once prior to the LIA. J1 and L1, in the following sections termed ‘Holocene
composite moraines’, mark the maximum of glacier advances and corresponding temperature minima since the end of the YD-
EH transition.

5.2 The moraine record of the Early Holocene

5.2.1 Local correlation

435 Moraine records at Jamtal and at Laraintal are remarkably similar and point to synchronous glacier dynamics throughout the
Holocene, particularly during its onset. In both valleys, we identified up to three lateral ridges just outboard the J1 and L1
composite moraines, and their terminal equivalents, albeit in varying states of preservation. The outermost ridges in both
valleys, J3R and L3, and J4R and L4, respectively, yield statistically identical landform ages (Fig. 7a–d). Interestingly, they
are chronostratigraphically inverse, i.e., JR3 and L3 are systematically several centuries older than JR4 and L4. An explanation
440 for this age pattern may be decadal- to centennial-scale pre-exposure of J3R and L3 boulders, which would lead to an
overestimation of ages. However, if pre-exposure was a problem in the data set, we would expect greater scatter in boulder
ages inferred for the same moraine. Another explanation for age inversion is post-depositional displacement such as sacking
or tilting of J4R and L4 boulders, for instance through the thawing of permafrost, which would lead to underestimation of
corresponding ages. Yet again boulder ages along JR4 and L4 are in good agreement, making this explanation unlikely. A
445 process, which may have affected JR4 and L4 boulder surfaces and could cause a systematic shift towards younger ages, is
katabatic winds, when the glacier abandoned moraines JR4 and L4 and halted for a few centuries at the positions of or close
to JR3 and L3. Consolidation of the age difference between moraines JR3/L3 and J4R/L4 would necessitate the removal of
approximately 2.5 mm of rock from boulder surfaces along the outermost moraines (J4R and L4) within 500 years.

The age pattern of EH moraines in the Jamtal and Laraintal is noteworthy, but we emphasize that moraine ages of JR3, L3,
450 JR4 and L4 overlap within 1σ uncertainties and that age inversion is non-existent from a statistical point of view. As we
attribute the landforms to a climatic state we group them across both valleys and refer to them as **Moraine Formation Interval
(MFI) 3-4**, equivalent to $11,030 \pm 740$ yrs, rounded to **11.0 \pm 0.7 ka** (Fig. 7f).

We correlate EH Jamtal and Laraintal moraine chronologies with moraine chronologies and glacier proxy records at the local
455 scale and propose a concept of YD-EH deglaciation. MFI 3-4 falls well into the EH and is different from advances during the
LG. Presumable YD moraines are identified at considerable distance downstream and outboard of landforms addressed in this
study (Hertl, 2001, and references therein), which implicates that glaciers shrank from their LG ice margin to a position close
to the LIA maximum within a few centuries. Rapid deglaciation is a direct response of glaciers to an increase of summer
temperatures by several degrees during the YD-EH transition, in the Eastern Alpine region and across the Alps (Fig. 9c–f; e.g.,
460 Affolter et al., 2019; Heiri et al., 2014; Ilyashuk et al., 2009; Larocque-Tobler et al., 2010b; Samartin et al., 2012). This

warming trend was interrupted by brief cold spells, which manifest in moraine records in the Silvretta Massif and in the adjacent Verwall mountains to the NE. Based on these moraine chronologies, the following local YD-EH glacier history emerges (Figure 10):

- 465 i. **Latest YD — Kartell moraines (Verwall):** Moraines that formed within this period in the region have been dated at the Kartell site, around 15–20 km NE of Jamtal and Laraintal (Ivy-Ochs et al., 2006). Because of the previously higher ^{10}Be production rate estimate, Kartell moraines were initially placed into the EH. Recalculations using the updated production rate yield boulder ages ranging between 11.8 ± 0.6 ka and 12.5 ± 0.9 ka with the production rate uncertainty excluded. The age update now assigns Kartell moraines to the late(st) YD (Boxleitner et al., 2019b; Ivy-Ochs, 2015). A local lowering of the ELA of approximately 110–120 m relative to the historical moraine that was calculated using the Accumulation Area Ratio (AAR) method (Gross et al., 1978), was reported by Sailer and Kerschner (1999) and (Ivy-Ochs et al., 2006).
- 470
- ii. **YD-EH transition/Preboreal — MFI 3--4:** The shift from glacial to interglacial conditions is captured at the Jamtal and Laraintal. Corresponding moraine chronologies indicate ice margins at terminal positions some hundreds of meters outboard of the LIA maximum, equivalent to an estimated ELA depression of approximately 70 m (AAR method; Hertl, 2001:80). These chronologies evidence abrupt cold snaps, which punctuated the general warming trend during the EH and which caused decadal- to centennial-scale glacier oscillations. The mean age of MFI 3-4 (11.0 ± 0.7 ka) follows **the** **Preboreal Oscillation (PBO)** as defined based on paleoenvironmental records from Europe ($11.30\text{--}11.15$ ka; e.g., Björck et al., 1997; Joannin et al., 2013; Magny et al., 2007; Schwander et al., 2000). Evidence of subsequent summer cooling between 10,700–10,500 cal BP is detected in lake sediments in the Swiss and Austrian Alps (Fig. 9c–d; Heiri et al., 2003; Lauterbach et al., 2011).
- 475
- 480

Kromer moraines identified in valleys 10–15 km further towards the West (Fig. 1b) were originally placed into the Preboreal (Gross et al., 1978). Morphologically, these moraines resemble the blocky, multi-ridge structures of J3/L3 and J4/L4 at the Jamtal and Laraintal and yield similar snowline depression estimates of 70–90 m. Updated ^{10}Be moraine ages of 9.9 ± 0.7 ka and 10.2 ± 0.7 ka fall within a somewhat younger age spectrum compared to the Jamtal and Laraintal moraines (Kerschner et al., 2006; Moran et al., 2016b). However, the age discrepancy between Kromer moraines and Jamtal and Laraintal moraines could be reconciled considering age uncertainties.


485

- iii. **Interglacial/Holocene mode — Ochsental–Grüne Kuppe (GrK):** Deglaciation patterns in the Ochsental to the West suggest ice-margin configurations similar to the LIA around 10 ka. The timing of moraine formation adjacent to the lateral Holocene composite moraine (equivalent to J1 and L1 in this study) was constrained to 9.9 ± 0.7 ka (Grüne Kuppe site, $n=4$) (Braumann et al., 2020). In addition, two boulders of the same age were found in latero-frontal sections of the Holocene composite moraine and point to similar ice margins around 10 ka and during the LIA. Although no exposure
- 490

ages are available for J2 and L2, and although these moraines could have been deposited within any cold phase between MFI 3-4 and the LIA, we note that J2 and L2 potentially may correlate with the Grüne Kuppe moraine, particularly as the ridges are in a morphostratigraphically similar position. The timing of the Grüne Kuppe moraine stabilization aligns with a climate anomaly detected in some proxy records of the Alps around 10.5 ka which persisted for several centuries, sometimes termed **Central European cold phase 1** (CE-1; e.g., Boch et al., 2009; Haas et al., 1998; Schmidt et al., 2006). This phase was followed by deglaciation and glaciers in the Alps receded to sizes smaller than their historical maximum (Patzelt, 2019; Solomina et al., 2015). Glacier retreat led to a vegetation change with trees spreading to high(er) elevations. At the Las Gondas bog in the adjacent Fimbatal (Fig. 1b), subfossil wood and tree logs were found up to an elevation of 2370 m a.s.l., with the oldest sample dated to 8620-8480 cal BP at 2355 m.a.s.l. (Nicolussi, 2010). In close vicinity to our study sites (Futschöltal), evidence of *Pinus cembra* populations growing at an elevation of c. 2290 m a.s.l. within the period between c. 5580 and 4970 calBP was found (Patzelt, 2019). Consistent results were reported from other valleys in the region, for instance from the Klostertal and the Bielerhöhe sites (Nicolussi, 2010), and from Kaunertal (Nicolussi et al., 2005) (Fig. 1b).


The timing of moraine formation at the Kartell site overlaps with MFI 3–4, just as MFI 3–4 overlaps with moraine formation at GrK in the Ochsental. Instead of indicating individual phases of glacier advance or stabilization, differences in the landforms' central ages could also be owed to uncertainties of the dating method, which we cannot exclude, or to catchment-specific effects such as shading or bedrock topography. We note that catchments where the evaluated ¹⁰Be moraine chronologies were generated are comparable in terms of elevation, glacier size, exposure, geographical location (North of the Alpine drainage divide), and orientation (Fig. B1a). Furthermore, summer temperature time series of meteorological stations at St. Anton in the Verwall region and at Galtür in the Silvretta region show similar trends in the period from 1957–2001 (Fig. B1b). We assume that glaciers in the region respond(ed) to the same temperature forcing, and probably at a similar level of sensitivity not only in recent decades, but also before. Hence, the multiphase age structure of EH moraine formation in the Verwall and Silvretta regions points to distinct episodes of glacier advance or stabilization.

515



520 Figure 9: Proxy records capturing the YD-EH transition. (a) Detrital Carbonate Peaks (DCP) identified in marine sediment cores from the Labrador sea (Jennings et al., 2015); TC – Total Carbonate. (b) Mean summer temperature reconstructions (JJA) derived from nine Greenland ice cores (Buizert et al., 2018). (c) Ostracods record extracted from lake sediments of Mondsee (Austrian Alps) (Lauterbach et al., 2011). (d) Chironomid-inferred (CI) atmospheric July temperatures from lake sediments of Hinterburgsee in the Swiss Alps. (e) Stacked CI July temperatures in the European Alps (Heiri et al., 2014). (f) Mean Annual Temperatures (MAT) in Central Europe reconstructed based on speleothems from Milandre Cave, Switzerland (Affolter et al., 2019). (g) Arctic moraine record: GrIS – Greenland Ice Sheet moraines, LIS – Laurentide Ice Sheet (LIS) moraines, NA Alpine – North America Alpine mountain glacier moraines (Young et al., 2020). (h) Moraine Formation Interval (MFI) identified in this study. Purple bars highlight

 525 Holocene cold events detected in Greenland ice cores (Rasmussen et al., 2007).



530 Figure 10: Glacier retreat during the transition from glacial to interglacial conditions evidenced in moraine chronologies from the Silvretta and Verwall regions. KAR – Kartell moraines (Ivy-Ochs, 2015; Ivy-Ochs et al., 2006), site is located approximately 15-20 km NE of Jamtal and Laraintal. MFI 3-4 described in this study. GrK – Grüne Kuppe moraine identified in the adjacent Ochsental suggesting LIA-like glacier extents around 10,000 years (Braumann et al., 2020).

5.2.2 Alpine-wide and hemispheric correlation

535 Moraine formation during the transition from glacial to interglacial climatic conditions that is presented in Figure 10, builds on glacier records in the Silvretta Massif and in the Verwall mountains in the Eastern Alps. This model is consistent with results of previous studies that have addressed glacier evolution during LG and EH based on moraine chronologies (e.g., Baroni et al., 2017; Hofmann et al., 2019; Moran et al., 2016a; Moran et al., 2017a; Protin et al., 2021; Protin et al., 2019; Schimmelpfennig et al., 2012; Schimmelpfennig et al., 2014; Schindelwig et al., 2012). Investigated moraine sets may differ with respect to their structure, state of preservation or distance relative to the LIA maximum, but they share their position
 540 (outboard the LIA maximum but inboard the presumable LG ice margin), and their age of deposition between c. 12 and 10 ka before present and imply substantial large-scale cooling during this period. This pattern of EH moraine stabilization is not

limited to the Alpine realm, but has been observed in other places in the North Atlantic and Arctic region, for instance along the Fennoscandian ice sheet (e.g., Briner et al., 2014; Nesje, 2009), the Icelandic ice sheet (e.g., Sigfusdottir and Benediktsson, 2020), at Svalbard (e.g., Farnsworth et al., 2020), along the eastern part of the LIS (e.g., Corbett et al., 2016; Ullman et al., 2016; Young et al., 2020) and along the Greenland ice sheet (e.g., Biette et al., 2020; Levy et al., 2016; Young et al., 2020) (Fig. 9g). Atmospheric temperatures were certainly different in these regions during the EH (Fig. 9b–f), and glaciers in the European Alps retreated much earlier to positions inboard their subsequent historical margin compared to Arctic glaciers which continued to deposit moraines outboard their LIA at least two more millennia. However, concurrent moraine stabilization during the EH raises the question what caused this synchronicity in climatic cooling during the first millennia of the Holocene.

5.2.3 Climatic drivers of EH moraine formation in the Northern Hemisphere

Freshwater influx into the North Atlantic and into the Arctic Ocean is known as a driver for climate of the Northern Hemisphere and acts as a plausible cause for abrupt centennial-scale cold snaps during the LG and the EH (e.g., Bjorck et al., 1997; Fisher et al., 2002; Hald and Hagen, 1998; Nesje et al., 2004). Prominent examples are repeated outbursts of the north American proglacial Agassiz lake, whose final drainage caused a sharp temperature drop in the Northern Hemisphere, the 8.2 ka event detected in Greenland ice cores (Alley and Agustsdottir, 2005; Clarke et al., 2009; Hillaire-Marcel et al., 2007; Teller et al., 2002; Thornalley et al., 2010). Besides abrupt high-volume releases of freshwater to the North Atlantic or Arctic Ocean via major lake drainages or iceberg armadas, there is evidence of more subdued glacial discharge during the EH that results in a deceleration of the thermohaline circulation (Bamberg et al., 2010; Renssen et al., 2010; Thornalley et al., 2009). Weakening of the AMOC leads to less heat transported to the North Atlantic region, which can prompt brief, decadal- to centennial-scale cold snaps in the North, and also at lower latitudes. During the PBO and subsequent centennial-scale cold phases, harsher climate conditions are reported in the North Atlantic region (e.g., Bos et al., 2007; Knudsen et al., 2008; Paus et al., 2015; Timms et al., 2021) with cooling extending towards Western and Central Europe (Fig. 9c–f). Glaciers in North Atlantic regions and in the Alps responded to these climate perturbations with stabilization or advance, hence moraine deposition.

To test the linkage between EH moraine formation and freshwater discharge of the LIS, we review corresponding markers in marine sediment cores, temperature proxy records and moraine records in these regions. We begin with the YD termination, when icebergs and meltwater plumes were released into the North Atlantic, evidenced by ice-rafted debris and layers of ‘foreign’ sediment enriched with detrital carbonates in marine sediments. These layers, often referred to as Heinrich-0 (H0) and characterized by a Detrital Carbonate Peak (DCP) date to the earliest Holocene and were identified in Baffin Bay (Simon et al., 2014), in the Labrador Sea (Andrews et al., 1995; Rashid et al., 2011), at the coast of Newfoundland (Pearce et al., 2015), including the Flemish cap (Li and Piper, 2015). Jennings et al. (2015) found multiple subsequent detrital carbonate peaks (DCP 1–7 in Fig. 9a) between 11,500 and 8,000 yrs BP in a core from the Labrador shelf, which is attributed to freshwater sourced from Hudson Strait. **DCP1** detected around 11,500 yrs BP coincides with the end of H0 and with the onset of 11.4 ka event (c. 11,450 to 11,350 cal BP; Rasmussen et al., 2007). During the subsequent PBO captured in Nordic records, mean

575 annual and summer temperatures in the European Alps declined (Fig. 8e–f; e.g., Affolter et al., 2019; Heiri et al., 2014; Lauterbach et al., 2011). The propagation of this cooling trend towards Western and Central Europe is supported by cooler and more humid climate conditions in these regions c. 11,300–11,150 cal BP (Magny et al., 2007). In parallel, ¹⁰Be concentration in Greenland ice cores, a proxy for solar activity, decreases towards a minimum (Adolphi et al., 2014; Finkel and Nishiizumi, 1997; Mekhaldi et al., 2020). Low solar activity may have amplified the cooling imposed by freshening of the Atlantic Ocean, or vice versa. Glaciers in the North Atlantic region and in the European Alps advanced or stabilized repeatedly during the first millennia of the Holocene and deposited moraines, as evidenced by moraines dated in this study (Fig. 9g–h). A similar chain of events may have occurred some centuries later. The deposition of the **DCP2** c. 10,600 cal BP was preceded by the so-called Gold Cove advance of the LIS’s Labrador sector across Hudson Strait and its subsequent retreat (Jennings et al., 2015; Kaufman et al., 1993; Rashid et al., 2014). The resulting freshwater input may have weakened the AMOC, which in turn led to a drop in mean annual temperatures in Greenland and moraine formation in the Arctic (e.g., Biette et al., 2020; Young et al., 2020). Temperatures in the Alps decreased or stagnated around that time (Fig. 9c–d, f). Moraine formation between 10,700 yrs BP and 10,500 yrs BP, concurrent with DCP2, is observed across the European Alps (e.g., Moran et al., 2017a; Protin et al., 2019; Schimmelpfennig et al., 2012; Schimmelpfennig et al., 2014).

The linkage between DCPs, freshwater input into the Atlantic and Arctic Ocean and subsequent EH glacier advances has been put forward before in the context of the North American and Arctic region (e.g., Andrews et al., 2014; Nesje, 2009; Young et al., 2020). The authors of a recent geochronological study carried out in the Western Alps go a step further and propose that freshwater forcing in the North Atlantic region acted as a driver for moraine formation in the Mont Blanc Massif (Protin et al., 2021). They suggest that a decrease in AMOC strength led to extended sea-ice periods during winter in the North Atlantic, which in turn caused a southwards shift of the westerlies. Cold air was then transported to Europe and led to moraine deposition in the European Alps. With our new moraine chronologies from the Silvretta region, we complement the glacier record from the Western Alps with robust evidence for EH moraine formation in the Eastern Alps and corroborate the hypothesis that centennial-scale cold phases occurred at a regional (hemispheric) scale between 12 and 10 ka.


6 Summary and Conclusions

- Glaciers at both study sites, the Jamtal and the Laraintal, stabilized or advanced at least twice during the EH, which is evidenced by moraines deposited outboard the historical moraine (LIA maximum) and inboard the presumable LG ice margin. The timing of moraine formation is consistent across both valleys and is constrained with ¹⁰Be SED, yielding a combined age of 11.0 ± 0.7 ka (MFI 3–4, n=19). EH moraines in the Silvretta region indicate repeated punctuation of the general postglacial warming trend by short cold episodes in Europe. These cold snaps, most prominently the PBO, appear to have their origin in the North Atlantic region. Layers of ice-rafted debris and DCPs in marine sediment cores along the eastern margin of the LIS point to glacial discharge during the earliest Holocene. Resulting freshwater released into the North Atlantic probably caused a drop in salinity and led to a slowdown of the AMOC. Reduced heat transport northwards

caused cooling in the North Atlantic region, which propagated toward Europe. Glaciers and ice sheets in the Northern Hemisphere responded to this cooling via moraine deposition.

- 610 ■ Based on Holocene moraine chronologies of the Silvretta Massif and the adjacent Verwall mountains, we propose the following local model that describes alternating phases of glacier retreat and stabilization between 12 and 10 ka: the YD termination (Kartell; Ivy-Ochs, 2015; Ivy-Ochs et al., 2006), the YD-EH transition (MFI 3–4, this study), and the Holocene mode (GrK; Braumann et al., 2020). The proposed concept confirms the hypothesis formulated by Patzelt and Bortenschlager (1973) almost 50 years ago, that glaciers in the Eastern Alps deposited moraines during the Preboreal, and had retreated to their subsequent historical ice margins by c. 10 to 9.5 ka. During the rest of the Holocene, the magnitude
615 of cooling was most likely too small in the Eastern Alps to force advances which exceeded dimensions that glaciers had around 10 ka.
- 620 ■ Our data suggests that glaciers in the Silvretta region advanced to a position close or equivalent to their LIA maximum around 500 CE. The timing of this advance is concurrent with the migration period in Europe, which is often associated with regional climate deterioration. In the Silvretta Massif, this hypothesis rests on a few ¹⁰Be exposure ages of the Jamtal (this study) and the Ochsental (Braumann et al., 2020), but there is growing evidence that many glaciers in the Alps, North America and the Nordic region advanced at that time and reached their historical margin (e.g., Barclay et al., 2009; Holzhauser et al., 2005; Le Roy et al., 2015; Patzelt, 2016).
- 625 ■ Silvretta glaciers may have advanced to their LIA maximum as early as c. 1300 CE. A subsequent advance to the same position took place in the second half of the 18th century, documented by three boulders along the historic moraine yielding a mean age of c. 260 ±25 yrs. This result agrees well with an advance of glaciers in the adjacent Ochsental (Braumann et al., 2020). Contemporaneous advances have also been reported for glaciated areas beyond the Silvretta region, e.g., Lower Grindelwald glacier, Mer de Glace and for glaciers in the Ötztal (Nicolussi, 2013; Nussbaumer et al., 2007; Pindur and Heuberger, 2010; Zumbühl and Nussbaumer, 2018).
- 630 ■ The classical LIA moraine, traditionally referred to as the ‘1850 moraine’ in the European Alps, marks the LIA maximum glacier extent that was reached multiple times during the LIA, but perhaps also earlier during the Holocene, most likely around 500 CE. As the moraine comprises glacial sediments deposited during several glacier advances during the past millennia, we propose that these landforms should rather be viewed as ‘Holocene composite moraines’ instead of ‘1850 moraines’ or ‘LIA moraines’.
- 635 ■ Our observations document sensitive glacier response to the natural warming in the Early Holocene. Accelerating anthropogenic warming has been driving the rapid retreat of highly temperature-sensitive glaciers over the last century (e.g., Oerlemans, 2005; Zemp et al., 2019), and will lead to the disappearance of most glaciers in the European Alps unless greenhouse gas forcing is substantially reduced in the very near future (Zekollari et al., 2019).


Appendix A: Complementary photo documentation of landforms



- 640 A1: Jamtal. (a) J1 moraine with Jamtal hut in the background. Boulder surfaces are fresh and are not colonized by lichens; pioneer
 plants growing on the moraine. (b) J1 multi-ridge structure with view towards with Jamtalferner in the back. (c) Left-lateral side of
 645 Jamtal below Totenfeld glacier. (d) View towards Chalausferner (in the background); J1 in the foreground, J3R and J4R outboard
 J1 are curved lateral moraines of former Chalausferner. (e) J2 moraine (undated) at the center with till covered slope to the left and
 J1 to the right. (f) Right-lateral moraine sets. J4R and J3R date to the Early Holocene, J1 comprises debris that was deposited during
 the LIA and during at least one earlier glacier advance around 500 CE.




A2: Jamtal. (a) J5 ridge; view from right-lateral valley side towards W, and (b) view from downvalley position southwards. The structure was reworked in the course of trail construction and maintenance. (c) Transition zone between J1 terminal moraine and blockfield. We speculate that the blockfield originates from rock failures along valley flanks farther upvalley. Corresponding debris was transported downstream by the glacier. During glacier retreat, the blocks melted out and covered the valley floor. The blockfield (at least its uppermost section) was then partly overprinted by subsequent glacier advances. (d) Blockfield consisting of coarse, angular to subangular components; view towards SE. (e) Transition zone between fresh J1 moraine deposits and the blockfield, whose blocks are populated by lichens and are partly overgrown with vegetation. (f) Displaced moraines along the right-lateral valley side outboard J1; Futschöltal (tributary valley) in the back.




A3: Laraintal. (a) Terminal moraines L3T and L4T dissected by a creek (Larainbach). (b) Right-lateral moraine set: L1, L2 (undated) and L3R. (c) Larainbach, modern river plane with L1 double-ridge structure identified on both sides of the creek. (d) Left-lateral flatter valley section, where moraines L3L and L1, and presumable Late Glacial moraines accumulated. (e) Left-lateral glacier side. Person standing on L1 ridge with L3L to the left. Hoher Kogel peak in the background. (f) Provenance area of rockfall event in 2019, Hoher Kogel peak.

660



665 **A4: Laraintal.** (a) Fresh rockfall deposits (2019) in the Zollhütte area. Note person at the center of the photograph for scale. (b) Zollhütte area (L5); unweathered scree on the left side of the photograph; mass movement slab to the right of the hiking trail; fresh rockfall path behind the slab.

Appendix B: Observations of summer temperature (JJA) in the Silvretta and Verwall regions



670 **B1: Overview of Silvretta and Verwall regions** addressed in local correlation of moraine records (section 5.2.2). (a) ^{10}Be moraine records generated in the following valleys: KAR – Kartell (Ivy-Ochs et al., 2006), KL – Kloistertal and KR – Kromertal (Moran et al., 2016b), OcG-GrK – Ochsental/Grüne Kuppe (Braumann et al., 2020), JAM – Jamtal and LAR – Laraintal (this study). Locations of meteorological stations in the Verwall region (St. Anton, station #14300) and in the Silvretta region (Galtür; homogenized HISTALP data), (Auer et al., 2007). Distance between the two stations amounts to approximately 20 km. Orthophoto accessed via <https://maps.tirol.gv.at>, © Land Tirol. (b) Summer (JJA) temperature time series (1957–2001) from the two stations that exhibit similar trends in the period between 1957–2001 (data provided by Austrian Weather Service ZAMG).

675

Data availability

All analytical information associated with cosmogenic nuclide measurements is listed in the tables in the Supplement and will be made available via the ICE-D Alpine database (<http://alpine.ice-d.org/>).

Author contribution

680 SMB and JMS designed the study. SMB, MF and JMS carried out field work. SMB was responsible for cosmogenic nuclide sample preparation. AJH performed sample measurements and developed the strategy for low-level ^{10}Be sample processing. All authors were involved in data interpretation. SMB wrote the manuscript, which was revised and edited by all authors.

Competing interests

The authors declare that they have no conflict of interest.

685 **Disclaimer**

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

Acknowledgements

SMB is grateful to the LDEO cosmogenic nuclide laboratory staff Roseanne Schwartz and Jean Hanley for help and advice during sample processing. Furthermore, SMB thanks Günther Gross for helpful discussions on the Holocene moraine record in the Silvretta Massif, Dominik Blasenbauer for assistance during field work, and Herbert Formayer for providing meteorological data. The authors express their gratitude to their hosts at the Jamtal hut, at Larainalpe and at Pension Türtscher, who have facilitated the field campaigns. Finally, the authors thank two anonymous reviewers, whose comments greatly improved the manuscript, and Alberto Reyes, who was the editor of the manuscript. This work was performed in part under 695 the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344; this is LLNL-JRNL-821951.

Financial support

This study was sponsored by inatura Museum GmbH. SMB is a recipient of a DOC Fellowship of the Austrian Academy of Sciences (OeAW) at the Institute of Applied Geology, University of Natural Resources and Life Sciences (BOKU) Vienna. 700 For research visits at the LDEO, SMB received a Marietta Blau scholarship sponsored by OeAD GmbH, a Marshall Plan

scholarship provided by the Austrian Marshall Plan Foundation and financial support from BOKU's 'Transitions to Sustainability' (T2S) doctoral school. This material is based upon work supported by the National Science Foundation under Grant No. NSF-1853881.

705 **References**

- Adolphi, F., Muscheler, R., Svensson, A., Aldahan, A., Possnert, G., Beer, J., Sjolte, J., Bjorck, S., Matthes, K., and Thieblemont, R.: Persistent link between solar activity and Greenland climate during the Last Glacial Maximum, *Nat Geosci*, 7, 662-666, 2014.
- Affolter, S., Hauselmann, A., Fleitmann, D., Edwards, R. L., Cheng, H., and Leuenberger, M.: Central Europe temperature constrained by speleothem fluid inclusion water isotopes over the past 14,000 years, *Sci Adv*, 5, 2019.
- 710 Alley, R. B.: The Younger Dryas cold interval as viewed from central Greenland, *Quaternary Sci Rev*, 19, 213-226, 2000.
- Alley, R. B. and Agustsdottir, A. M.: The 8k event: cause and consequences of a major Holocene abrupt climate change, *Quaternary Sci Rev*, 24, 1123-1149, 2005.
- André, M. F.: Rates of postglacial rock weathering on glacially scoured outcrops (Abisko-Riksgransen area, 68 degrees N), *Geogr Ann A*, 84A, 139-150, 2002.
- 715 Andrews, J. T., Gibb, O. T., Jennings, A. E., and Simon, Q.: Variations in the provenance of sediment from ice sheets surrounding Baffin Bay during MIS 2 and 3 and export to the Labrador Shelf Sea: site HU 2008029-0008 Davis Strait, *J Quaternary Sci*, 29, 3-13, 2014.
- Andrews, J. T., Jennings, A. E., Kerwin, M., Kirby, M., Manley, W., Miller, G. H., Bond, G., and Maclean, B.: A Heinrich-Like Event, H-0 (Dc-0) - Source(S) for Detrital Carbonate in the North-Atlantic during the Younger Dryas Chronozone, *Paleoceanography*, 10, 943-952, 1995.
- 720 Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C., Briffa, K., Jones, P., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin, J. M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S., Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., and Niepova, E.: HISTALP - historical instrumental climatological surface time series of the Greater Alpine Region, *Int J Climatol*, 27, 17-46, 2007.
- 725 Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J.: A complete and easily accessible means of calculating surface exposure ages or erosion rates from Be-10 and Al-26 measurements, *Quat Geochronol*, 3, 174-195, 2008.
- Bamberg, A., Rosenthal, Y., Paul, A., Heslop, D., Mulitza, S., Ruhlemann, C., and Schulz, M.: Reduced North Atlantic Central Water formation in response to early Holocene ice-sheet melting, *Geophys Res Lett*, 37, 2010.
- Barclay, D. J., Wiles, G. C., and Calkin, P. E.: Holocene glacier fluctuations in Alaska, *Quaternary Sci Rev*, 28, 2034-2048, 2009.
- 730 Baroni, C., Casale, S., Salvatore, M. C., Ivy-Ochs, S., Christl, M., Carturan, L., Seppi, R., and Carton, A.: Double response of glaciers in the Upper Peio Valley (Rhaetian Alps, Italy) to the Younger Dryas climatic deterioration, *Boreas*, 46, 783-798, 2017.
- Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J. I., Magnusson, J., Marty, C., Morán-Tejeda, E., Morin, S., Naaim, M., Provenzale, A., Rabatel, A., Six, D., Stötter, J., Strasser, U., Terzago, S., and Vincent, C.: The European mountain cryosphere: a review of its current state, trends, and future challenges, *The Cryosphere*, 12, 759-794, 2018.
- 735 Bertle, H.: Zur Geologie des Fensters von Gargellen (Vorarlberg) und seines Kristallinen Rahmens - Österreich, *Mitt. Ges. Geol. Berbaustud.*, 22, 1-60, 1973.
- Bichler, M. G., Reindl, M., Reitner, J. M., Drescher-Schneider, R., Wirsig, C., Christl, M., Hajdas, I., and Ivy-Ochs, S.: Landslide deposits as stratigraphical markers for a sequence-based glacial stratigraphy: a case study of a Younger Dryas system in the Eastern Alps, *Boreas*, 45, 537-551, 2016.
- 740 Biette, M., Jomelli, V., Chenet, M., Braucher, R., Rinterknecht, V., Lane, T., and Team, A.: Mountain glacier fluctuations during the Lateglacial and Holocene on Clavering Island (northeastern Greenland) from ¹⁰Be moraine dating, *Boreas*, 49, 873-885, 2020.
- Bjorck, S., Rundgren, M., Ingolfsson, O., and Funder, S.: The Preboreal oscillation around the Nordic Seas: terrestrial and lacustrine responses, *J Quaternary Sci*, 12, 455-465, 1997.
- 745 BMLRT: eHYD - Hydrographisches Jahrbuch. BMNT, Abteilung I/3 – Wasserhaushalt (HZB), Vienna, 2021.
- BMNT: Hydrographisches Jahrbuch von Österreich. BMNT (Federal Ministry for Sustainability and Tourism), Vienna, 2016.
- Boch, R., Spötl, C., and Kramers, J.: High-resolution isotope records of early Holocene rapid climate change from two coeval stalagmites of Katerloch Cave, Austria, *Quaternary Sci Rev*, 28, 2527-2538, 2009.
- Bos, J. A. A., van Geel, B., van der Plicht, J., and Bohncke, S. J. P.: Preboreal climate oscillations in Europe: Wiggle-match dating and synthesis of Dutch high-resolution multi-proxy records, *Quaternary Sci Rev*, 26, 1927-1950, 2007.
- 750 Boxleitner, M., Ivy-Ochs, S., Egli, M., Brandova, D., Christl, M., Dahms, D., and Maisch, M.: The ¹⁰Be deglaciation chronology of the Göschenental, central Swiss Alps, and new insights into the Göschenen Cold Phases, *Boreas*, 48, 867-878, 2019a.
- Boxleitner, M., Ivy-Ochs, S., Egli, M., Brandova, D., Christl, M., and Maisch, M.: Lateglacial and Early Holocene glacier stages – New dating evidence from the Meiental in central Switzerland, *Geomorphology*, 340, 15-31, 2019b.
- 755 Braumann, S. M., Schaefer, J. M., Neuhuber, S. M., Reitner, J. M., Lüthgens, C., and Fiebig, M.: Holocene glacier change in the Silvretta Massif (Austrian Alps) constrained by a new ¹⁰Be chronology, historical records and modern observations, *Quaternary Sci Rev*, 245, 1-21, 2020.
- Briner, J. P., Svendsen, J. I., Mangerud, J., Lohne, O. S., and Young, N. E.: A Be-10 chronology of south-western Scandinavian Ice Sheet history during the Lateglacial period, *J Quaternary Sci*, 29, 370-380, 2014.

- 760 Buizert, C., Keisling, B. A., Box, J. E., He, F., Carlson, A. E., Sinclair, G., and DeConto, R. M.: Greenland-Wide Seasonal Temperatures During the Last Deglaciation, *Geophys Res Lett*, 45, 1905-1914, 2018.
- Büntgen, U., Myglan, V. S., Ljungqvist, F. C., McCormick, M., Di Cosmo, N., Sigl, M., Jungclauss, J., Wagner, S., Krusic, P. J., Esper, J., Kaplan, J. O., de Vaan, M. A. C., Luterbacher, J., Wacker, L., Tegel, W., and Kirilyanov, A. V.: Cooling and societal change during the Late Antique Little Ice Age from 536 to around 660 AD, *Nat Geosci*, 9, 231-236, 2016.
- 765 Büntgen, U., Tegel, W., Nicolussi, K., McCormick, M., Frank, D., Trouet, V., Kaplan, J. O., Herzig, F., Heussner, K. U., Wanner, H., Luterbacher, J., and Esper, J.: 2500 Years of European Climate Variability and Human Susceptibility, *Science*, 331, 578-582, 2011.
- Cheng, H., Zhang, H. W., Spotl, C., Baker, J., Sinha, A., Li, H. Y., Bartolome, M., Moreno, A., Kathayat, G., Zhao, J. Y., Dong, X. Y., Li, Y. W., Ning, Y. F., Jia, X., Zong, B. Y., Brahim, Y. A., Perez-Mejias, C., Cai, Y. J., Novello, V. F., Cruz, F. W., Severinghaus, J. P., An, Z. S., and Edwards, R. L.: Timing and structure of the Younger Dryas event and its underlying climate dynamics, *P Natl Acad Sci USA*, 117, 23408-23417, 2020.
- 770 Clark, P. U., Shakun, J. D., Baker, P. A., Bartlein, P. J., Brewer, S., Brook, E., Carlson, A. E., Cheng, H., Kaufman, D. S., Liu, Z. Y., Marchitto, T. M., Mix, A. C., Morrill, C., Otto-Bliesner, B. L., Pahnke, K., Russell, J. M., Whitlock, C., Adkins, J. F., Blois, J. L., Clark, J., Colman, S. M., Curry, W. B., Flower, B. P., He, F., Johnson, T. C., Lynch-Stieglitz, J., Markgraf, V., McManus, J., Mitrovica, J. X., Moreno, P. I., and Williams, J. W.: Global climate evolution during the last deglaciation, *P Natl Acad Sci USA*, 109, E1134-E1142, 2012.
- 775 Clarke, G. K. C., Bush, A. B. G., and Bush, J. W. M.: Freshwater Discharge, Sediment Transport, and Modeled Climate Impacts of the Final Drainage of Glacial Lake Agassiz, *J Climate*, 22, 2161-2180, 2009.
- Claude, A., Ivy-Ochs, S., Kober, F., Antognini, M., Salcher, B., and Kubik, P. W.: The Chironico landslide (Valle Leventina, southern Swiss Alps): age and evolution, *Swiss J Geosci*, 107, 273-291, 2014.
- Corbett, L. B., Bierman, P. R., and Davis, P. T.: Glacial history and landscape evolution of southern Cumberland Peninsula, Baffin Island, Canada, constrained by cosmogenic Be-10 and Al-26, *Geol Soc Am Bull*, 128, 1173-1192, 2016.
- 780 Cossart, E., Fort, M., Bourles, D., Braucher, R., Perrier, R., and Sime, L.: Deglaciation pattern during the Lateglacial/Holocene transition in the southern French Alps. Chronological data and geographical reconstruction from the Claree Valley (upper Durance catchment, southeastern France), *Palaeogeogr Palaeoclimatol*, 315, 109-123, 2012.
- Deline, P. and Orombelli, G.: Glacier fluctuations in the western Alps during the Neoglacial, as indicated by the Miage morainic amphitheatre (Mont Blanc massif, Italy), *Boreas*, 34, 456-467, 2005.
- 785 Denton, G. H., Putnam, A. E., Russell, J. L., Barrell, D. J. A., Schaefer, J. M., Kaplan, M. R., and Strand, P. D.: The Zealandia Switch: Ice age climate shifts viewed from Southern Hemisphere moraines, *Quaternary Sci Rev*, 257, 2021.
- Dietre, B., Walser, C., Lambers, K., Reitmaier, T., Hajdas, I., and Haas, J. N.: Palaeoecological evidence for Mesolithic to Medieval climatic change and anthropogenic impact on the Alpine flora and vegetation of the Silvretta Massif (Switzerland/Austria), *Quatern Int*, 353, 3-16, 2014.
- 790 Faedrich, R.: Spät- und postglaziale Gletscherschwankungen in der Ferwallgruppe (Tirol/Vorarlberg), *Dusseldorfer Geographische Schriften*, 12, 1-161, 1979.
- Farnsworth, W. R., Allaart, L., Ingolfsson, O., Alexanderson, H., Forwick, M., Noormets, R., Retelle, M., and Schomacker, A.: Holocene glacial history of Svalbard: Status, perspectives and challenges, *Earth-Sci Rev*, 208, 2020.
- 795 Federici, P. R., Granger, D. E., Pappalardo, M., Ribolini, A., Spagnolo, M., and Cyr, A. J.: Exposure age dating and Equilibrium Line Altitude reconstruction of an Egesen moraine in the Maritime Alps, Italy, *Boreas*, 37, 245-253, 2008.
- Finkel, R. C. and Nishiizumi, K.: Beryllium 10 concentrations in the Greenland Ice Sheet Project 2 ice core from 3-40 ka, *J Geophys Res-Oceans*, 102, 26699-26706, 1997.
- Fischer, A., Fickert, T., Schweizer, G., Patzelt, G., and Gross, G.: Vegetation dynamics in Alpine glacier forelands tackled from space, *Sci Rep-Uk*, 9, 2019.
- 800 Fischer, A., Seiser, B., Helfricht, K., and Stocker-Waldhuber, M.: High-resolution inventory to capture glacier disintegration in the Austrian Silvretta, *The Cryosphere Discuss.*, 2021, 1-31, 2021.
- Fischer, A., Seiser, B., Waldhuber, M. S., Mitterer, C., and Abermann, J.: Tracing glacier changes in Austria from the Little Ice Age to the present using a lidar-based high-resolution glacier inventory in Austria, *Cryosphere*, 9, 753-766, 2015.
- 805 Fisher, T. G., Smith, D. G., and Andrews, J. T.: Preboreal oscillation caused by a glacial Lake Agassiz flood, *Quaternary Sci Rev*, 21, 873-878, 2002.
- Fohlmeister, J., Vollweiler, N., Spotl, C., and Mangini, A.: COMNISPA II: Update of a mid-European isotope climate record, 11 ka to present, *Holocene*, 23, 749-754, 2013.
- Friebe, G.: *Geologie der österreichischen Bundesländer - Vorarlberg*, Verlag der Geologischen Bundesanstalt (GBA), Wien, 2007.
- 810 Fuchs, G. and Oberhauser, R.: *170 Galtür*, Geologische Bundesanstalt, Vienna, 1990.
- Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., and Stoffel, M.: 21st century climate change in the European Alps - A review, *Sci Total Environ*, 493, 1138-1151, 2014.
- Gosse, J. C. and Phillips, F. M.: Terrestrial in situ cosmogenic nuclides: theory and application, *Quaternary Sci Rev*, 20, 1475-1560, 2001.
- 815 Gross, G., Kerschner, H., and Patzelt, G.: Methodische Untersuchungen über die Schneegrenze in den alpinen Gletschergebieten, *Zeitschrift für Gletscherkunde und Glaziologie*, 12, 223-251, 1978.

- Grove, J. M.: Little ice ages : ancient and modern, Routledge, London ; New York, 2004.
- Haas, J. N., Richoz, I., Tinner, W., and Wick, L.: Synchronous Holocene climatic oscillations recorded on the Swiss Plateau and at timberline in the Alps, *Holocene*, 8, 301-309, 1998.
- Hald, M. and Hagen, S.: Early preboreal cooling in the Nordic seas region triggered by meltwater, *Geology*, 26, 615-618, 1998.
- 820 Heiri, O., Koinig, K. A., Spotl, C., Barrett, S., Brauer, A., Drescher-Schneider, R., Gaar, D., Ivy-Ochs, S., Kerschner, H., Luetscher, M., Moran, A., Nicolussi, K., Preusser, F., Schmidt, R., Schoeneich, P., Schworer, C., Sprafke, T., Terhorst, B., and Tinner, W.: Palaeoclimate records 60-8 ka in the Austrian and Swiss Alps and their forelands, *Quaternary Sci Rev*, 106, 186-205, 2014.
- Heiri, O., Wick, L., van Leeuwen, J. F. N., van der Knaap, W. O., and Lotter, A. F.: Holocene tree immigration and the chironomid fauna of a small Swiss subalpine lake (Hinterburgsee, 1515 m asl), *Palaeogeogr Palaeocl*, 189, 35-53, 2003.
- 825 Helama, S., Jones, P. D., and Briffa, K. R.: Dark Ages Cold Period: A literature review and directions for future research, *Holocene*, 27, 1600-1606, 2017.
- Helama, S., Stoffel, M., Hall, R. J., Jones, P. D., Arppe, L., Matskovsky, V. V., Timonen, M., Nojd, P., Mielikainen, K., and Oinonen, M.: Recurrent transitions to Little Ice Age-like climatic regimes over the Holocene, *Clim Dynam*, doi: 10.1007/s00382-021-05669-0, 2021. 2021.
- 830 Hertl, A.: Untersuchungen zur spätglazialen Gletscher- und Klimageschichte der Österreichischen Silvretta-Gruppe, 2001. Doctoral Thesis, 716, Leopold-Franzens-Universität Innsbruck, Innsbruck, 265 pp., 2001.
- Hillaire-Marcel, C., de Vernal, A., and Piper, D. J. W.: Lake Agassiz final drainage event in the northwest North Atlantic, *Geophys Res Lett*, 34, 2007.
- Hofmann, F. M., Alexanderson, H., Schoeneich, P., Mertes, J. R., Léanni, L., and Team, A.: Post-Last Glacial Maximum glacier fluctuations in the southern Écrins massif (westernmost Alps): insights from ¹⁰Be cosmic ray exposure dating *Boreas*, 48, 1019-1041, 2019.
- 835 Holzhauser, H., Magny, M., and Zumbühl, H. J.: Glacier and lake-level variations in west-central Europe over the last 3500 years, *Holocene*, 15, 789-801, 2005.
- Huston, A., Siler, N., Roe, G. H., Pettit, E., and Steiger, N. J.: Understanding drivers of glacier-length variability over the last millennium, *The Cryosphere*, 15, 1645-1662, 2021.
- 840 Ilyashuk, B., Gobet, E., Heiri, O., Lotter, A. F., van Leeuwen, J. F. N., van der Knaap, W. O., Ilyashuk, E., Oberli, F., and Ammann, B.: Lateglacial environmental and climatic changes at the Maloja Pass, Central Swiss Alps, as recorded by chironomids and pollen, *Quaternary Sci Rev*, 28, 1340-1353, 2009.
- Ilyashuk, E. A., Heiri, O., Ilyashuk, B. P., Koinig, K. A., and Psenner, R.: The Little Ice Age signature in a 700-year high-resolution chironomid record of summer temperatures in the Central Eastern Alps, *Clim Dynam*, 52, 6953-6967, 2019.
- 845 Ivy-Ochs, S.: Glacier Variations in the European Alps at the End of the Last Glaciation, *Cuad Investig Geogr*, 41, 295-315, 2015.
- Ivy-Ochs, S., Kerschner, H., Maisch, M., Christl, M., Kubik, P. W., and Schlüchter, C.: Latest Pleistocene and Holocene glacier variations in the European Alps, *Quaternary Sci Rev*, 28, 2137-2149, 2009.
- Ivy-Ochs, S., Kerschner, H., Reuther, A., Maisch, M., Sailer, R., Schaefer, J., Kubik, P. W., Synal, H.-A., and Schlüchter, C.: The timing of glacier advances in the northern European Alps based on surface exposure dating with cosmogenic ¹⁰Be, ²⁶Al, ³⁶Cl, and ²¹Ne. In: *In Situ-Produced Cosmogenic Nuclides and Quantification of Geological Processes*, Siame, L. L., Bourlès, D. L., and Brown, E. T. (Eds.), Geological Society of America, 2006.
- 850 Jennings, A., Andrews, J., Pearce, C., Wilson, L., and Olfasdottir, S.: Detrital carbonate peaks on the Labrador shelf, a 13-7 ka template for freshwater forcing from the Hudson Strait outlet of the Laurentide Ice Sheet into the subpolar gyre, *Quaternary Sci Rev*, 107, 62-80, 2015.
- Joannin, S., Vanniere, B., Galop, D., Peyron, O., Haas, J. N., Gilli, A., Chapron, E., Wirth, S. B., Anselmetti, F., Desmet, M., and Magny, M.: Climate and vegetation changes during the Lateglacial and early-middle Holocene at Lake Ledro (southern Alps, Italy), *Clim Past*, 9, 913-933, 2013.
- 855 K. u. k. Militärgeographisches Institut: Dritte Reambulierte Landesaufnahme (Franzisco-Josephinische Landesaufnahme) - Blatt 5244, K. u. k. Militärgeographisches Institut, Vienna, 1870-1887.
- Kasper, M.: *Mythos Piz Buin - Kulturgeschichte eines Berges*, 2015.
- 860 Kasper, M.: *Silvretta historica - Zeitreise durch die Silvretta*, Heimatschutzverein Montafon, Schruns, 2013.
- Kaufman, D. S., Miller, G. H., Stravers, J. A., Manley, W. F., and Duvall, M. L.: Late-Glacial Ice Margins and Deglacial Chronology for Southeastern Baffin-Island and Hudson Strait, Eastern Canadian Arctic - Reply, *Can J Earth Sci*, 30, 1753-1758, 1993.
- Kelly, M. A., Kubik, P. W., Von Blanckenburg, F., and Schlüchter, C.: Surface exposure dating of the Great Aletsch Glacier Egesen moraine system, western Swiss Alps, using the cosmogenic nuclide Be-10, *J Quaternary Sci*, 19, 431-441, 2004.
- 865 Kerschner, H., Hertl, A., Gross, G., Ivy-Ochs, S., and Kubik, P. W.: Surface exposure dating of moraines in the Kromer valley (Silvretta Mountains, Austria) - evidence for glacial response to the 8.2 ka event in the Eastern Alps?, *Holocene*, 16, 7-15, 2006.
- Kerschner, H. and Ivy-Ochs, S.: Palaeoclimate from glaciers: Examples from the Eastern Alps during the Alpine Lateglacial and early Holocene, *Global Planet Change*, 60, 58-71, 2008.
- Knudsen, K. L., Stabell, B., Seidenkrantz, M. S., Eriksson, J., and Blake, W.: Deglacial and Holocene conditions in northernmost Baffin Bay: sediments, foraminifera, diatoms and stable isotopes, *Boreas*, 37, 346-376, 2008.
- 870

- Kobashi, T., Menviel, L., Jeltsch-Thommes, A., Vinther, B. M., Box, J. E., Muscheler, R., Nakaegawa, T., Pfister, P. L., Doring, M., Leuenberger, M., Wanner, H., and Ohmura, A.: Volcanic influence on centennial to millennial Holocene Greenland temperature change, *Sci Rep-Uk*, 7, 2017.
- Lal, D.: In situ-produced Cosmogenic Isotopes in Terrestrial Rocks, *Annu Rev Earth Pl Sc*, 16, 355-388, 1988.
- 875 Larocque-Tobler, I., Grosjean, M., Heiri, O., Trachsel, M., and Kamenik, C.: Thousand years of climate change reconstructed from chironomid subfossils preserved in varved lake Silvaplana, Engadine, Switzerland, *Quaternary Sci Rev*, 29, 1940-1949, 2010a.
- Larocque-Tobler, I., Heiri, O., and Wehrli, M.: Late Glacial and Holocene temperature changes at Egelsee, Switzerland, reconstructed using subfossil chironomids, *J Paleolimnol*, 43, 649-666, 2010b.
- Lauterbach, S., Brauer, A., Andersen, N., Danielopol, D. L., Dulski, P., Huls, M., Milecka, K., Namiotko, T., Obremaska, M., Von Grafenstein, U., and Participants, D.: Environmental responses to Lateglacial climatic fluctuations recorded in the sediments of pre-Alpine Lake Mondsee (northeastern Alps), *J Quaternary Sci*, 26, 253-267, 2011.
- LDEO: Extraction of Beryllium from Quartz. LDEO Cosmogenic Isotope Lab, Palisades, NY, 2012a.
- LDEO: Separation and Purification of Quartz from Whole Rock. LDEO Cosmogenic Isotope Lab, Palisades, NY, 2012b.
- 885 Le Roy, M., Nicolussi, K., Deline, P., Astrade, L., Edouard, J. L., Miramont, C., and Arnaud, F.: Calendar-dated glacier variations in the western European Alps during the Neoglacial: the Mer de Glace record, Mont Blanc massif, *Quaternary Sci Rev*, 108, 1-22, 2015.
- Levy, L. B., Kelly, M. A., Lowell, T. V., Hall, B. L., Howley, J. A., and Smith, C. A.: Coeval fluctuations of the Greenland ice sheet and a local glacier, central East Greenland, during late glacial and early Holocene time, *Geophys Res Lett*, 43, 1623-1631, 2016.
- Li, G. and Piper, D. J. W.: The influence of meltwater on the Labrador Current in Heinrich event 1 and the Younger Dryas, *Quaternary Sci Rev*, 107, 129-137, 2015.
- 890 Maggetti, M. and Flisch, M.: Evolution of the Silvretta Nappe. In: Pre-Mesozoic Geology in the Alps, von Raumer, J. F. and Neubauer, F. (Eds.), Springer Berlin Heidelberg, Berlin, Heidelberg, 1993.
- Magny, M., Vanniere, B., de Beaulieu, J. L., Begeot, C., Heiri, O., Millet, L., Peyron, O., and Walter-Simonnet, A. V.: Early-Holocene climatic oscillations recorded by lake-level fluctuations in west-central Europe and in central Italy, *Quaternary Sci Rev*, 26, 1951-1964, 2007.
- 895 Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A Reconstruction of Regional and Global Temperature for the Past 11,300 Years, *Science*, 339, 1198-1201, 2013.
- Mekhaldi, F., Czymzik, M., Adolphi, F., Sjolte, J., Bjorck, S., Aldahan, A., Brauer, A., Martin-Puertas, C., Possnert, G., and Muscheler, R.: Radionuclide wiggle matching reveals a nonsynchronous early Holocene climate oscillation in Greenland and western Europe around a grand solar minimum, *Clim Past*, 16, 1145-1157, 2020.
- 900 Moran, A. P., Ivy-Ochs, S., Schuh, M., Christl, M., and Kerschner, H.: Evidence of central Alpine glacier advances during the Younger Dryas-early Holocene transition period, *Boreas*, 45, 398-410, 2016a.
- Moran, A. P., Ivy-Ochs, S., Vockenhuber, C., and Kerschner, H.: First ³⁶Cl exposure ages from a moraine in the Northern Calcareous Alps, *E&G Quaternary Sci. J.*, 65, 145-155, 2017a.
- Moran, A. P., Ivy Ochs, S., Christl, M., and Kerschner, H.: Exposure dating of a pronounced glacier advance at the onset of the late-Holocene in the central Tyrolean Alps, *The Holocene*, 27, 1350-1358, 2017b.
- 905 Moran, A. P., Kerschner, H., and Ivy-Ochs, S.: Redating the moraines in the Kromer Valley (Silvretta Mountains) - New evidence for an early Holocene glacier advance, *Holocene*, 26, 655-664, 2016b.
- Nesje, A.: Latest Pleistocene and Holocene alpine glacier fluctuations in Scandinavia, *Quaternary Sci Rev*, 28, 2119-2136, 2009.
- Nesje, A., Dahl, S. O., and Bakke, J.: Were abrupt Lateglacial and early-Holocene climatic changes in northwest Europe linked to freshwater outbursts to the North Atlantic and Arctic Oceans?, *Holocene*, 14, 299-310, 2004.
- 910 Nicolussi, K.: Die historischen Vorstöße und Hochstände des Vernagtferners 1600-1850 AD, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 45/46, 9-23, 2013.
- Nicolussi, K.: Jahrringdaten zur nacheiszeitlichen Waldverbreitung in der Silvretta. In: Letzte Jäger, erste Hirten - Hochalpine Archäologie in der Silvretta - Begleitheft zur Ausstellung, Reitmaier, T. (Ed.), Abt. Ur- und Frühgeschichte der Universität Zürich, Zürich, 2010.
- 915 Nicolussi, K., Kaufmann, M., Patzelt, G., van der Plicht, J., and Thurner, A.: Holocene tree-line variability in the Kauner Valley, Central Eastern Alps, indicated by dendrochronological analysis of living trees and subfossil logs, *Veg Hist Archaeobot*, 14, 221-234, 2005.
- Nicolussi, K. and Patzelt, G.: Discovery of early-Holocene wood and peat on the forefield of the Pasterze Glacier, Eastern Alps, Austria, *Holocene*, 10, 191-199, 2000.
- Nishiizumi, K., Imamura, M., Caffee, M. W., Southon, J. R., Finkel, R. C., and McAninch, J.: Absolute calibration of Be-10 AMS standards, *Nucl Instrum Meth B*, 258, 403-413, 2007.
- 920 Nussbaumer, S. U., Zumbühl, H. J., and Steiner, D.: Fluctuations of the "Mer de Glace" (Mont Blanc area, France) AD 1500-2050: an interdisciplinary approach using new historical data and neural network simulations, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 40, 5-175, 2007.
- Oerlemans, J.: Extracting a climate signal from 169 glacier records, *Science*, 308, 675-677, 2005.
- 925 PAGES 2k Consortium: Continental-scale temperature variability during the past two millennia, *Nat Geosci*, 6, 339-346, 2013.

- Patzelt, G.: Das Bunte Moor in der Oberfernau (Stubai Alpen, Tirol) – Eine neu bearbeitete Schlüsselstelle für die Kenntnis der nacheiszeitlichen Gletscherschwankungen der Ostalpen, *Jahrbuch der Geologischen Bundesanstalt*, Band 156, 97-107, 2016.
- Patzelt, G.: *Gletscher : Klimazeugen von der Eiszeit bis zur Gegenwart*, Hatje Cantz, Berlin, 2019.
- 930 Patzelt, G. and Bortenschlager, S.: Die postglazialen Gletscher- und Klimaschwankungen in der Venedigergruppe (Hohe Tauern, Ostalpen), *Z. Geomorph. Suppl.*, 16, 25-72, 1973.
- Paus, A., Boessenkool, S., Brochmann, C., Epp, L. S., Fabel, D., Hafliðason, H., and Linge, H.: Lake Store Finnsjøen - a key for understanding Lateglacial/early Holocene vegetation and ice sheet dynamics in the central Scandes Mountains, *Quaternary Sci Rev*, 121, 36-51, 2015.
- 935 Pearce, C., Andrews, J. T., Bouloubassi, I., Hillaire-Marcel, C., Jennings, A. E., Olsen, J., Kuijpers, A., and Seidenkrantz, M. S.: Heinrich 0 on the east Canadian margin: Source, distribution, and timing, *Paleoceanography*, 30, 1613-1624, 2015.
- Pindur, P. and Heuberger, H.: Zur holozänen Gletschergeschichte im Zemmgrund in den Zillertaler Alpen, Tirol/Österreich (Ostalpen), *Zeitschrift für Gletscherkunde und Glazialgeologie*, 42, 2010.
- Protin, M., Schimmelpfennig, I., Mugnier, J.-L., Buoncristiani, J.-F., Le Roy, M., Pohl, B., Moreau, L., and Team, A.: Millennial-scale deglaciation across the European Alps at the transition between the Younger Dryas and the Early Holocene – evidence from a new cosmogenic nuclide chronology, *Boreas*, doi: <https://doi.org/10.1111/bor.12519>, 2021. 2021.
- 940 Protin, M., Schimmelpfennig, I., Mugnier, J.-L., Ravanel, L., Le Roy, M., Deline, P., Favier, V., Buoncristiani, J.-F., Aumaître, G., Bourlès, D. L., and Keddadouche, K.: Climatic reconstruction for the Younger Dryas/Early Holocene transition and the Little Ice Age based on paleo-extents of Argentière glacier (French Alps), *Quaternary Sci Rev*, 221, 105863, 2019.
- Rashid, H., Piper, D. J. W., and Flower, B. P.: The role of Hudson Strait outlet in Younger Dryas sedimentation in the Labrador Sea, *Geophys. Monogr. Ser.*, 193, 93-110, 2011.
- 945 Rashid, H., Piper, D. J. W., Mansfield, C., Saint-Ange, F., and Polyak, L.: Signature of the Gold Cove event (10.2 ka) in the Labrador Sea, *Quatern Int*, 352, 212-221, 2014.
- Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new
- 950 Greenland ice core chronology for the last glacial termination, *J Geophys Res-Atmos*, 111, 2006.
- Rasmussen, S. O., Vinther, B. M., Clausen, H. B., and Andersen, K. K.: Early Holocene climate oscillations recorded in three Greenland ice cores, *Quaternary Sci Rev*, 26, 1907-1914, 2007.
- Renssen, H., Goosse, H., Crosta, X., and Roche, D. M.: Early Holocene Laurentide Ice Sheet deglaciation causes cooling in the high-latitude Southern Hemisphere through oceanic teleconnection, *Paleoceanography*, 25, 2010.
- 955 Roe, G. H., Baker, M. B., and Herla, F.: Centennial glacier retreat as categorical evidence of regional climate change, *Nat Geosci*, 10, 95+, 2017.
- Rood, D. H., Brown, T. A., Finkel, R. C., and Guilderson, T. P.: Poisson and non-Poisson uncertainty estimations of Be-10/Be-9 measurements at LLNL-CAMS, *Nucl Instrum Meth B*, 294, 426-429, 2013.
- Rupper, S. and Roe, G.: Glacier Changes and Regional Climate: A Mass and Energy Balance Approach, *J Climate*, 21, 5384-5401, 2008.
- 960 Sailer, R. and Kerschner, H.: Equilibrium-line altitudes and rock glaciers during the Younger Dryas cooling event, Ferwall group, western Tyrol, Austria, *Annals of Glaciology*, Vol 28, 28, 141-145, 1999.
- Samartin, S., Heiri, O., Vescovi, E., Brooks, S. J., and Tinner, W.: Lateglacial and early Holocene summer temperatures in the southern Swiss Alps reconstructed using fossil chironomids, *J Quaternary Sci*, 27, 279-289, 2012.
- Schaefer, J. M., Denton, G. H., Kaplan, M. R., Putnam, A., Finkel, R. C., Barrell, D. J. A., Andersen, B. G., Schwartz, R., Mackintosh, A.,
- 965 Chinn, T., and Schlüchter, C.: High-Frequency Holocene Glacier Fluctuations in New Zealand Differ from the Northern Signature, *Science*, 324, 622-625, 2009.
- Schimmelpfennig, I., Schaefer, J. M., Akcar, N., Ivy-Ochs, S., Finkel, R. C., and Schlüchter, C.: Holocene glacier culminations in the Western Alps and their hemispheric relevance, *Geology*, 40, 891-894, 2012.
- 970 Schimmelpfennig, I., Schaefer, J. M., Akcar, N., Koffman, T., Ivy-Ochs, S., Schwartz, R., Finkel, R. C., Zimmerman, S., and Schlüchter, C.: A chronology of Holocene and Little Ice Age glacier culminations of the Steingletscher, Central Alps, Switzerland, based on high-sensitivity beryllium-10 moraine dating, *Earth Planet Sc Lett*, 393, 220-230, 2014.
- Schindelwig, I., Akcar, N., Kubik, P. W., and Schlüchter, C.: Lateglacial and early Holocene dynamics of adjacent valley glaciers in the Western Swiss Alps, *J Quaternary Sci*, 27, 114-124, 2012.
- Schmidt, R., Kamenik, C., Tessadri, R., and Koinig, K. A.: Climatic changes from 12,000 to 4,000 years ago in the Austrian Central Alps tracked by sedimentological and biological proxies of a lake sediment core, *J Paleolimnol*, 35, 491-505, 2006.
- 975 Schuster, R.: *Zur Geologie der Ostalpen*, *Abh. Geol. B.-A.*, 64, 143-165, 2015.
- Schwander, J., Eicher, U., and Ammann, B.: Oxygen isotopes of lake marl at Gerzensee and Leysin (Switzerland), covering the Younger Dryas and two minor oscillations, and their correlation to the GRIP ice core, *Palaeogeogr Palaeoclimatol*, 159, 203-214, 2000.
- 980 Shakun, J. D., Clark, P. U., He, F., Lifton, N. A., Liu, Z. Y., and Otto-Bliesner, B. L.: Regional and global forcing of glacier retreat during the last deglaciation, *Nat Commun*, 6, 2015.

- Sigfusdottir, T. and Benediktsson, I. O.: Refining the history of Younger Dryas and Early Holocene glacier oscillations in the Borgarfjorour region, western Iceland, *Boreas*, 49, 296-314, 2020.
- Simon, Q., Hillaire-Marcel, C., St-Onge, G., and Andrews, J. T.: North-eastern Laurentide, western Greenland and southern Innuitian ice stream dynamics during the last glacial cycle, *J Quaternary Sci*, 29, 14-26, 2014.
- 985 Solomina, O. N., Bradley, R. S., Hodgson, D. A., Ivy-Ochs, S., Jomelli, V., Mackintosh, A. N., Nesje, A., Owen, L. A., Wanner, H., Wiles, G. C., and Young, N. E.: Holocene glacier fluctuations, *Quaternary Sci Rev*, 111, 9-34, 2015.
- Solomina, O. N., Bradley, R. S., Jomelli, V., Geirsdottir, A., Kaufman, D. S., Koch, J., McKay, N. P., Masiokas, M., Miller, G., Nesje, A., Nicolussi, K., Owen, L. A., Putnam, A. E., Wanner, H., Wiles, G., and Yang, B.: Glacier fluctuations during the past 2000 years, *Quaternary Sci Rev*, 149, 61-90, 2016.
- 990 Steiner, D., Pauling, A., Nussbaumer, S. U., Nesje, A., Luterbacher, J., Wanner, H., and Zumbühl, H. J.: Sensitivity of European glaciers to precipitation and temperature - two case studies, *Climatic Change*, 90, 413-441, 2008.
- Teller, J. T., Leverington, D. W., and Mann, J. D.: Freshwater outbursts to the oceans from glacial Lake Agassiz and their role in climate change during the last deglaciation, *Quaternary Sci Rev*, 21, 879-887, 2002.
- Thornalley, D. J. R., Elderfield, H., and McCave, I. N.: Holocene oscillations in temperature and salinity of the surface subpolar North Atlantic, *Nature*, 457, 711-714, 2009.
- 995 Thornalley, D. J. R., McCave, I. N., and Elderfield, H.: Freshwater input and abrupt deglacial climate change in the North Atlantic, *Paleoceanography*, 25, 2010.
- Thornalley, D. J. R., Oppo, D. W., Ortega, P., Robson, J. I., Brierley, C. M., Davis, R., Hall, I. R., Moffa-Sanchez, P., Rose, N. L., Spooner, P. T., Yashayaev, I., and Keigwin, L. D.: Anomalously weak Labrador Sea convection and Atlantic overturning during the past 150 years, *Nature*, 556, 227-+, 2018.
- 1000 Timms, R. G. O., Abrook, A. M., Matthews, I. P., Francis, C. P., Mroczkowska, A., Candy, I., Brooks, S. J., Milner, A. M., and Palmer, A. P.: Evidence for centennial-scale Lateglacial and early Holocene climatic complexity from Quoyloo Meadow, Orkney, Scotland, *J Quaternary Sci*, 2021. 2021.
- Ullman, D. J., Carlson, A. E., Hostetler, S. W., Clark, P. U., Cuzzone, J., Milne, G. A., Winsor, K., and Caffee, M.: Final Laurentide ice-sheet deglaciation and Holocene climate-sea level change, *Quaternary Sci Rev*, 152, 49-59, 2016.
- 1005 Vollweiler, N., Scholz, D., Muhlinghaus, C., Mangini, A., and Spotl, C.: A precisely dated climate record for the last 9 kyr from three high alpine stalagmites, Spannagel Cave, Austria, *Geophys Res Lett*, 33, 2006.
- Walker, M., Johnsen, S., Rasmussen, S. O., Steffensen, J. P., Popp, T., Gibbard, P., Hoek, W., Lowe, J., Andrews, J., Björck, S., Cwynar, L., Hughen, K., Kershaw, P., Kromer, B., Litt, T., Lowe, D. J., Nakagawa, T., Newnham, R., and Schwander, J.: The Global Stratotype Section and Point (GSSP) for the base of the Holocene Series/Epoch (Quaternary System/Period) in the NGRIP ice core, *Episodes*, 31, 264-267, 2008.
- WGMS, W. G. M. S.: *Fluctuations of Glaciers Database*. World Glacier Monitoring Service, Zürich, 2018.
- Young, N. E., Briner, J. P., Miller, G. H., Lesnek, A. J., Crump, S. E., Thomas, E. K., Pendleton, S. L., Cuzzone, J., Lamp, J., Zimmerman, S., Caffee, M., and Schaefer, J. M.: Deglaciation of the Greenland and Laurentide ice sheets interrupted by glacier advance during abrupt coolings, *Quaternary Sci Rev*, 229, 2020.
- 1015 Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-CORDEX RCM ensemble, *Cryosphere*, 13, 1125-1146, 2019.
- Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer, S. U., Gartner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., and Cogley, J. G.: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568, 382-+, 2019.
- 1020 Zumbühl, H. J. and Nussbaumer, S. U.: Little Ice Age glacier history of the Central and Western Alps from pictorial documents, 2018, 44, 22, 2018.