

Holocene palaeoceanography of the Northeast Greenland shelf

Teodora Pados-Dibattista¹, Christof Pearce¹, Henrieka Detlef¹, Jørgen Brendtsen², Marit-Solveig Seidenkrantz¹

¹Paleoceanography and Paleoclimate Group, Arctic Research Centre, and iClimate Centre, Department of Geoscience, Aarhus University, Aarhus, 8000, Denmark

²ClimateLab, Brønshøj, 2700, Denmark

Correspondence to: Teodora Pados-Dibattista (pados.theo@gmx.at) and Marit-Solveig Seidenkrantz (mss@geo.au.dk)

Abstract. The Northeast Greenland shelf is highly sensitive to climate and ocean variability because it is swept by the East Greenland Current, which, through the western Fram Strait, forms the main pathway of export of sea ice and cold water masses from the Arctic Ocean into the North Atlantic Ocean. In order to reconstruct the variability of the East Greenland Current and general palaeoceanographic conditions in the area during the Holocene, we carried out benthic foraminiferal assemblage, stable isotope, and sedimentological analyses of a marine sediment core retrieved from the Northeast Greenland shelf (core DA17-NG-ST07-73G). The results reveal significant variations in the water masses and thus, in the strength of the East Greenland Current over the last ca. 9.4 ka BP. Between 9.4 and 8.2 ka BP the water column off Northeast Greenland was highly stratified, with cold, sea ice-loaded surface waters and strong influx of warm Atlantic Water in the subsurface. At ~8.4 ka BP a short-lived peak in terrestrial elements may be linked to influx of iceberg transported sediments and thus, to the so-called 8.2 ka event. Holocene Thermal Maximum like conditions prevailed from 8.2 to 6.2 ka BP, with a strong influence of the Return Atlantic Current and a weakened transport of Polar Water in the upper East Greenland Current. After 6.2 ka BP we recorded a return to a more stratified water column with sea-ice loaded surface waters and still Atlantic-sourced subsurface waters. After 4.2 ka BP increased Polar Water at the surface of the East Greenland Current and reduction of the Return Atlantic Water at subsurface levels led to freshening and reduced stratification of the water column and (near) perennial sea-ice cover. The Neoglaciation started at 3.2 ka BP at our location, characterised by a strengthened East Greenland Current. Cold subsurface water conditions with possible sea-ice cover and minimum surface water productivity persisted here throughout the last ~3 kyr.

1 Introduction

The acceleration of climatic changes in the Arctic and Subarctic regions is particularly marked by the drastic reduction of summer sea-ice cover. According to model simulations, the Arctic Ocean may become seasonally ice-free as early as 2040-2050 (Stroeve et al., 2012). This sea-ice reduction has extreme societal and environmental relevance, as it may open new opportunities for shipping and societal development of Arctic regions and Greenland. Furthermore, the sea-ice decline has been shown to alter Arctic ecosystems (Ardyna et al., 2014). However, despite their importance, Holocene marine environments and corresponding natural sea-ice states around Greenland are still not well understood. The Northeast (NE) Greenland continental shelf is a particularly important region for studying and understanding the mechanisms that control sea-ice formation, melt and drift. It is the broadest shelf along the Greenland margin, extending more than 300 km from the coastline. To the west it reaches the East Greenland coastline with its major marine-terminating outlets of the Greenland Ice Sheet, while to the east it is bounded by the Fram Strait, which is the only deep passage between the Arctic Ocean and the rest of the world oceans. This shelf region is highly sensitive to climate and ocean variability because it underlies the East Greenland Current (EGC), which, through the western Fram Strait, is the main pathway of export of sea ice and cold water masses from the Arctic Ocean into the North Atlantic Ocean (Rudels and Quadfasel, 1991). The outflow of meltwater/sea ice is a key factor that determines the North Atlantic freshwater budget and

stratification, and it influences the deep-water formation. Moreover, waters on the NE Greenland continental shelf play a major role in the stability of the glacier outlets (Nioghalvfjordsfjorden Glacier, Zachariæ Isstrøm, Storstrømmen) of the Northeast Greenland Ice Stream (Wilson and Straneo, 2015; Schaffer et al, 2017).




Figure 1(a): Close-up of the Northeast Greenland Shelf showing its characteristic features, the location of the studied sediment core (black circle) and the schematic subsurface circulation of the continental shelf (orange arrows). Arctic Atlantic Water (AAW), Return Atlantic Water (RAW), Nioghalvfjordsfjorden Glacier (79NG), Zachariæ Isstrøm (ZI), Storstrømmen (SS), Northeast Greenland Ice Stream (NEGIS, grey arrows), Norske Øer Ice Barrier (NØIB; white dashed line). (b): Overview map of the study area with the studied sediment core DA17-NG-ST07-73G (black circle; ‘Rumohr core DA17-NG-ST07-72R was taken from the same site), other core sites mentioned in the text (open circles) and schematic illustration of the major currents. East Greenland Current (EGC), West Spitsbergen Current (WSC), Return Atlantic Current (RAC), Irminger Current (IC), West Greenland Current (WGC). Red arrows: warm surface currents, blue arrows: cold surface currents, orange arrow: Atlantic originated subsurface water mass, Arctic Atlantic Water (AAW). Core sites: PS93/025 (Syring et al., 2020a; Zehnich et al., 2020), PS100/270 (Syring et al., 2020b), JM10-330 (Consolaro et al., 2018), MSM5/5-712 (Müller et al., 2012; Werner et al., 2013), JM02-440 (Ślubowska-Woldengen et al., 2007), PS1878 (Telesinski et al., 2014a-b), PS2641-4 (Müller et al., 2012; Perner et al., 2015; Kolling et al., 2017), PS1243 (Bauch et al., 2001), DA00-02 (Seidenkrantz et al., 2008), MD99-2317 (Jennings et al., 2011), JM96-1206 (Jennings et al., 2002; Perner et al., 2016), MD99-2322 (Jennings et al., 2011), MD99-2275 (Ran et al., 2006), MD95-2011 (Giraudeau et al., 2010), MD99-2269 (Giraudeau et al., 2004; Giraudeau et al., 2010), MD99-2266 (Moossen et al., 2015), 248260 (Seidenkrantz et al., 2007), DA12-11/2 (Van Nieuwenhové et al., 2018; Orme et al., 2018). Black line: location of the hydrographic transect pictured on Fig. 2b. Black dashed line: median sea-ice extent in September (1981-2010). The ocean bathymetry data are derived from GEBCO (Weatherall et al., 2015).

At the surface, the EGC carries cold and fresh polar waters southward. Below this relatively low-saline layer, the water masses originate from submerged and recirculated Atlantic-sourced waters from the Arctic (Arctic Atlantic Water), and partly from the returning branch of the West Spitsbergen Current (Return Atlantic Current, RAC; Quadfasel et al., 1987) (Fig. 1). The EGC transports these water masses from the north and along the East Greenland shelf towards the North Atlantic Ocean.

The freshwater budget here is one of the major components that affects the strength of the Atlantic Meridional Overturning Circulation (AMOC; Rahmstorf, 1995; Clark et al., 2002). The AMOC is a main mechanism for the heat redistribution on our planet and with impact on the global climate. It transports warm and salty surface waters to high latitudes in the northern North Atlantic, where they cool, sink and return southwards at depth. Increased freshwater flux from the EGC to the northern North Atlantic may prevent deep convection, thus, reducing meridional heat transport, which causes cooling of the high latitudes (Clark et al., 2001). At present, due to the anthropogenic global warming, the AMOC may be in its weakest state in the last 1000 years (Caesar et al., 2021), demanding an improved understanding of this complex system and its components.

The amount of polar water transported southward in the EGC is related to dynamics in the Arctic Ocean, the regional wind stress above the Nordic Seas and Greenland shelf area and, thereby, also depends on the North Atlantic Oscillation (NAO) and its Arctic counterpart, the Arctic Oscillation (AO). The NAO/AO is one of the most prominent and recurrent patterns of atmospheric circulation variability. The NAO/AO complex refers to a redistribution of air masses between the Arctic and the subtropical Atlantic, and its shifts from positive to negative phase produce large changes in the weather of the middle and high altitudes of the Northern Hemisphere (Hurrell et al., 2003). In case of positive NAO/AO, stronger westerlies carry moist air over Europe and Siberia, Atlantic Water inflow to the Arctic Ocean through the Fram Strait increases, and temperatures in the Arctic increase, which lead to a reduction in sea ice formation. During intervals of a negative NAO/AO these phenomena occur to be reversed (Kwok, 2000; Hurrell and Deser, 2010).

Several studies suggest that the East Greenland shelf has been subjected to a series of oceanographic and palaeoclimatic changes during the Holocene, induced by changes in the strength of the EGC linked to fluctuations in Atlantic Water entrainment. Most of these studies focus on the shelf region of Middle and South East Greenland (e.g., Jennings et al., 2002; Jennings et al., 2011; Müller et al., 2012; Perner et al., 2015; Perner et al., 2016; Kolling et al., 2017) and only few investigated the palaeoceanographic evolution of the EGC in the northern part of the East Greenland shelf (Bauch et al., 2001; Syring et al., 2020a, 2020b; Zehnich et al., 2020). In this study, we aim to reconstruct the dynamics and variability of the EGC off NE Greenland at centennial resolution throughout most of the Holocene by micropalaeontological, sedimentological and geochemical analysis of a sediment core retrieved from the central NE Greenland shelf. Faunal assemblage analysis of benthic and planktic foraminifera, stable isotope analysis, radiocarbon datings and X-ray fluorescence data allow us to infer changes in sea surface productivity, subsurface temperatures, sea-ice conditions and Greenland Ice Sheet melting over the last ca. 9.4 ka BP. The reconstruction provides new insights into the not yet resolved palaeoceanographic evolution of this region, with comparisons to published records from the polar/subpolar North Atlantic.

2 Regional setting

The bathymetry of the NE Greenland continental shelf is characterised by a trough system, consisting of five prominent cross-shelf troughs, which are deeper than the surrounding banks (Arndt et al., 2015). Nioghalvfjærdsfjorden Glacier (or 79 North Glacier), Zachariæ Isstrøm and Storstrømmen (Fig. 1) are the three major marine-terminating glaciers of the Northeast Greenland Ice Stream that drain in the area. Our study site is located approximately in the middle of the shelf in an inter-trough area between the Westwind Trough and Norske Trough, facing Nioghalvfjærdsfjorden Glacier and Zachariæ Isstrøm (Fig. 1). The location lies directly on the flow path of the EGC, as it flows southward along the Greenland shelf break (Johannessen, 1986), and over the middle to outer shelf (Bourke et al., 1987). Below a layer of the relatively low-saline Polar Surface Water, mainly originating from the Arctic Ocean and with contribution from local melt, the upper part of the EGC carries cold, low-saline (T: 0-1 °C, S<32) Polar Water

(PW) from the Arctic Ocean in the upper ca. 200 m of the water column. At subsurface levels below the PW, the water mass has mainly its origin from Atlantic sources, to varying extent from the Return Atlantic Water (RAW; transported by the RAC) and from Arctic Atlantic Water (AAW; also sometimes named “Arctic Intermediate Water”) (Rudels et al., 2005) (Fig. 2). The RAC is the western branch of the West Spitsbergen Current that transports relatively warm and saline water masses ($T < 2$ °C, $S: 34-35$) from the North Atlantic towards NE Greenland and then, by joining the EGC, southwards along the Greenland coast. In contrast, the eastern branch of the West Spitsbergen Current transports Atlantic Water (AW) into the Arctic Ocean, where it circulates at subsurface levels and cools. The AAW ($T \geq 0$ °C, $S: 34-35$) is formed by this recirculated cool AW, which again exits the Arctic Ocean and, together with the RAC, it forms the subsurface section of the EGC, as it flows southward along the East Greenland shelf (e.g., Quadfasel et al, 1987; Rudels, 2012). Along this southward path, the bathymetry on the NE Greenland shelf steers parts of the AW into the prominent cross-shelf troughs, toward the marine terminating glaciers of the Northeast Greenland Ice Stream, and thereby modulating the glaciers basal melt rates (Arndt et al., 2015; Schaffer et al., 2017).

The Polar Front in the Greenland Sea represents the eastern limit of perennial sea-ice cover. The location of the sea-ice edge in the summer depends on the extent of the sea-ice export from the Arctic Ocean (Vinje, 1977). Today the sea-ice margin and the location of the Polar Front lies east of our study site (Danish Meteorological Institute (DMI) and NSIDC, 2012). However, the Northeast Water Polynya is in fairly close proximity to our core location. This is a seasonally ice-free or only loosely ice-covered area south of Nordøstrundingen (Fig. 1) on the eastern Greenland coast. Two ice barriers support the seasonal formation of the Northeast Water Polynya. The shelf ice of the Ob Bank (Fig. 1a) pushes drift ice eastward, while the Norske Øer Ice Barrier (Fig. 1a) blocks the northward flowing sea ice, which is entrained in the NE Greenland coastal current (Schneider and Budéus, 1994). It starts to open around May/June, then gradually increases in size and closes in September (Pedersen et al., 1993). The polynya’s maximum extent to the north can reach high latitudes up to 83°N, beyond the NE Greenland shelf. To the east, the polynya can occupy the entire NE Greenland shelf with only little sea ice left in the area (Schneider and Budéus, 1997).




Figure 2(a): Temperature and salinity profile of the water column at the coring station (79.073° N 11.918° W) obtained during the NorthGreen17 expedition, showing the main water masses. (b): Annual average temperature of the water column in the upper 1000 m along a transect at 78.876° N, showing the main water masses at the study site. Polar Surface Water (PSW), Polar Water (PW), combined Arctic Atlantic Water (AAW) and Return Atlantic Water (RAW), as well as upper Polar Deep Water (uPDW). The yellow star indicates the location of the core site at 385 m water depth. Temperature data from World Ocean Atlas 2018 (Boyer et al., 2018). The location of this transects is marked as a black line on Fig. 1b.

130 **3 Material and methods**

Gravity core DA17-NG-ST07-73G (79.068° N 11.903° W; hereafter 73G) was collected during the NorthGreen Expedition in September 2017 at 385 m water depth in an isolated bathymetrical depression on the NE Greenland shelf (Fig. 1). In addition, we collected Rumohr core DA17-NG-ST07-72R (79.072° N 11.888° W, 384.6 m water depth; hereon 72G) with an intact sediment-water interface from the same station, which provided information for the age model of the gravity core. Temperature and salinity of the water column were measured in situ by a Seabird CTD (Conductivity-Temperature-Depth) profiler (Seabird SBE911 plus system equipped with two temperature and conductivity sensors) before the deployment of the gravity corer, with the core site being identified based on Innomar® subbottom profiler data obtained prior to coring.

140 The 410 cm long gravity core (73G) and the 88 cm long Rumohr core (72R) were stored at ca 3 °C before they were split lengthwise. The archive halves were scanned with an Itrax X-ray fluorescence core scanner (Cox Analytical) at the Department of Geoscience, Aarhus University (AU), Denmark. The scan was conducted with a Molybdenum tube (other settings: step size: 200 µm; exposure time: 10 s; voltage: 30 kV, current: 30 mA). The core scanning provided a line scan image, a 2-cm-wide radiographic image of the centre of the core (voltage: 60 kV; current: 45 mA; exposure time: 2000 ms) and high-resolution analysis of bulk geochemical composition. From the results of the X-ray fluorescence scanning, we show the elements Si, K, Ti and the Ca/Fe ratio on Fig. 3.

145 The age-depth model of core 73G is based on six Accelerator Mass Spectrometry radiocarbon (AMS ¹⁴C) dates, of which five are of mixed benthic foraminifera and one is of a worm tube lining. AMS ¹⁴C dating was performed at the Laboratory of Ion Beam Physics, ETH Zürich, Switzerland. Radiocarbon ages were calibrated using the Marine20 calibration dataset (Heaton et al., 2020) and a local marine reservoir age correction (ΔR) of 0 ± 50 years (Reimer and Reimer, 2001). We are aware that Heaton et al. (2020) state that the Marine20 is not suitable for polar regions. However, the same was true, just not explicitly stated, for the earlier marine radiocarbon age calibration curve Marine13 (Reimer et al., 2013), which is commonly used in arctic palaeoceanography (e.g., Perner et al., 2015; Seidenkrantz et al., 2019; Syring et al., 2020a). The presence of sea ice in polar regions impacts the local reservoir age, and therefore there is added uncertainty; this issue is not resolved by using an older calibration dataset. Since its publication, the Marine20 has been widely used in the Arctic realm (e.g., Farmer et al., 2021; Altuna et al., 2021). Moreover, for this specific Holocene reconstruction, the differences between using Marine13 and Marine20 are much smaller than the associated uncertainties (see Table A1 in the appendices).

160 The age-depth model was made with the OxCal v4.4 software (Ramsey, 2008), using a P-sequence depositional model. Beyond the lowest radiocarbon date, the model is extrapolated toward the bottom of the core. The age of the upper part of the core is constrained by correlation to Rumohr core 72R by selected XRF elements (appendices, Fig. A1), to determine the amount of top sediment loss during gravity coring.

165 Samples for foraminiferal, ice-rafted debris (IRD) and stable isotope analyses were taken at 5 cm intervals at the Department of Geoscience, AU. The samples were weighed, wet-sieved with distilled water through sieves of 1000, 100 and 63 µm, dried at 40 °C for 24 hours and weighed again. For the foraminiferal assemblage analyses both planktic and benthic (calcareous and agglutinated) species were counted and identified to species level under a stereomicroscope for the size fractions of 63-100 µm and >100 µm; counts were subsequently combined prior to percentage calculations. When the samples contained excessive material, prior to counting they were split into equal parts containing >300 specimens, using a microsplitter. In these cases, foraminiferal tests of one part were counted and identified, and the total number of foraminifera was calculated. A minimum of 300 benthic specimens were identified for each sample, except for four samples (10-11 cm, 15-16 cm, 30-31 cm, 195-196 cm).

170 These four samples contained only between 242-296 specimens. Absolute foraminiferal concentrations were calculated as
individuals/g wet sediment. Relative abundances of both benthic agglutinated and calcareous species were calculated as a
percentage of the total benthic (calcareous and agglutinated) foraminiferal fauna. However, diagrams of the separate percentage
calculations of the two benthic foraminiferal groups (agglutinated species of total agglutinated assemblage and calcareous species
of total calcareous assemblage) are shown in the appendices (Figs. A2 and A3), as are diagrams of the same species as number of
175 individuals/g wet sediment.

After counting, we placed selected benthic calcareous foraminifera species into groups according to their main environmental
preferences; for details see the discussion (chapter 5.2) and Table A3 in the appendices. The Atlantic Water group includes *C.*
neoteretis and *P. bulloides*. The chilled Atlantic Water group includes *I. norcrossi* and *M. barleeaanum*. The Arctic Water group
includes *S. horvathi* and *E. arctica*. The sea-ice edge group includes *S. feylingi*.

180 For stable isotope analysis 5-200 specimens of the benthic foraminiferal species *Elphidium clavatum* were picked from every
sample. The number of picked tests was restricted by the number of available, clearly identifiable tests without any corrosion or
non-typical shapes. The specimens belonged mainly to the 100-1000 μm fraction, supplemented with few specimens from the 63-
100 μm fraction, when necessary. The oxygen and carbon isotope analysis of foraminiferal calcite was performed at the Leibniz
185 Laboratory for Radiometric Dating and Stable Isotope Research at the Christian-Albrechts-University of Kiel using a MAT253
(Thermo Scientific) mass spectrometer system and a Kiel IV carbonate preparation device, with an analytical accuracy of <0.08
‰ for $\delta^{18}\text{O}$ and <0.05 ‰ for $\delta^{13}\text{C}$. Results are expressed in δ notation referring to the PDB (Pee Dee Belmnite) standard, while
using NBS-19 and IAEA-603 carbonate standards.

190 The content of IRD was calculated at 5 cm sample intervals using the dry weight of the size fraction >1000 μm divided by the wet
weight of the bulk sample.

Grain size distribution was measured on 42 samples (taken at 10-cm intervals) using a laser diffraction particle sizer (Sympatec
Helos) at the Department of Geoscience, AU, and grouped into three fractions: sand (>60 μm), silt (2-60 μm) and clay (<2 μm),
195 according to the available particle size intervals.

4 Results

4.1 Core description

Core DA17-NG-ST07-73G consist entirely of olive grey (5Y 4/1) marine silt with some clay, except the lowermost part of the
core. The sediment of the lowest 40 cm of the core is much coarser, containing up to 51 % sand; it is also darker in colour (Fig. 3).
200 In the rest of the core the sand fraction is on average 1.5 % and the clay : silt : sand ratio does not show any significant changes.




Figure 3: Results of the sedimentological and XRF core scanning analyses of core DA17-NG-ST07-73G. From left to right: photograph of the core with the amount of IRD (black line), radiograph, grain size analysis and XRF counts of selected elements. Black dashed lines in the XRF curves represent the mean of the values measured throughout the core. Black arrows next to the left y-axis (age) mark radiocarbon dates. The bottom 65 cm of the sediment core is marked with rose-pink colour. The photograph and radiograph of the core were horizontally stretched.

4.2 Chronology

The age model was developed using six AMS ^{14}C measurements on mixed benthic foraminifera and on a worm tube lining from core 73G (Table 1). In order to evaluate potential loss of core top sediment during the coring process, we compared the XRF data of gravity core 73G and Rumohr core 72R (Fig. A1 in appendices), which indicated that 12 cm sediment was missing from the top of core 73G; this information was used when constructing the age model of core 73G. This age model suggest that the 410 cm long sediment core covers the last ~11.2 ka BP (Fig. 4). However, the results of the grain size, IRD and X-ray fluorescence analysis indicate that beyond the lowest radiocarbon-dated sample (at 345 cm depth, 9.4 ka BP) the sediments change significantly (Fig. 4). In the bottom 65 cm of the core, preliminary analysis of foraminiferal content revealed the presence of species indicating a Pliocene or early Pleistocene age (Feyling-Hanssen, 1976; Feyling-Hanssen, 1980; Feyling-Hanssen et al., 1983), in addition to the normal in situ foraminiferal assemblage (see chapter 4.4.1). We suggest that these Plio-/Pleistocene foraminifera are reworked into a glaciomarine setting. Due to this very different environment and the lack of ^{14}C dates in this interval, the age model of the bottom 65 cm of core 73G is currently uncertain. Therefore, the main focus of this paper is on the last 9.4 ka.

Table 1: List of radiocarbon dates and modelled ages in core DA17-NG-ST07-73G. All dates were calibrated using the Marine20 calibration curve and $\Delta R=0 \pm 50$ years.

Lab. ID	Depth (cm)	Material	Radiocarbon age (yr BP)	Error (yr)	Calibrated 2-sigma (BP)			Modelled 2-sigma (cal yr BP)		
					from	to	median	from	to	median

ETH-95387	70.5	Mixed benthic foraminifera	2475	60	2200	1705	1952	2161	1817	1989
ETH-95388	110.5	Mixed benthic foraminifera	3275	70	3190	2715	2935	3165	2783	2972
ETH-110893	226	Worm tube lining	5645	30	6029	5622	5835	6031	5678	5857
ETH-95389	250.5	Mixed benthic foraminifera	6015	70	6468	5994	6241	6643	6275	6458
ETH-95390	300.5	Mixed benthic foraminifera	7595	70	8097	7654	7870	8151	7761	7942
ETH-95391	345.5	Mixed benthic foraminifera	9015	70	9823	9295	9536	9602	9125	9383




Figure 4: Age-depth model of core DA17-NG-ST07-73G based on six AMS 14C dates and on the comparison with Rumohr core DA17-NG-ST7-72R. The shading in pink illustrates the 2-sigma uncertainty range of the model. The hatched box in the bottom of the core indicates an interval containing reworked microfossils and is therefore of uncertain age (see pink shading on Fig. 3).

4.3 X-ray fluorescence

Throughout the whole core, Si, K and Ti follow the same pattern. After a drop in values in the lowermost section of yet uncertain age, all three records show a relatively constant values until ~318 cm depth (ca. 8.5 ka BP). At 316 cm (ca. 8.4 ka BP) all three elements displays peak counts, followed by a decrease until 161 cm (ca. 4.2 ka BP). There is a short-term drop in all three element counts at 161 cm depth (ca. 4.2 ka BP), followed by a steady increase until 125 cm (ca. 3.2 ka BP) and relatively constant values until the top of the core (Fig. 3). The Ca/Fe ratio follows a steadily decreasing trend throughout the core, with slightly stronger fluctuations from 345 cm until ~270 cm (ca. 7 ka BP) and in the top 55 cm (last ~1.5 ka BP). The record shows a pronounced peak between 335 and 338 cm (ca. 9.1 ka BP), and two minor peaks at ~213 cm (ca. 5.5 ka BP) and at ~34 cm (ca. 0.9 ka BP) (Fig. 3).

4.4 Foraminiferal analysis

4.4.1 410-345 cm core depth

240 The bottom 65 cm of the core are characterised by a mixture of two assemblages. An assemblage of relatively small, well-preserved specimens of Arctic species include *Cassidulina reniforme*, *Elphidium clavatum*, *Islandiella norcrossi* and *Melonis barleeanum*. This assemblage is mixed with large specimens of *Cibicidoides grossus*, *Cibicides scaldicensis*, *Islandiella inflata*, *Cassidulina teretis* and *Elphidium funderi*, which are extinct since the early or mid Pleistocene (Feyling-Hanssen, 1976; Feyling-Hanssen, 1980; Feyling-Hanssen et al., 1983; Seidenkrantz, 1995) and by species that are only present in temperate to tropical regions (*Bulimina marginata*, *Faujasina* sp.). The assemblage also contain some unusually large specimens of *E. clavatum*, *Haynesina orbiculare* and *Islandiella islandica*.

4.4.2 345-0 cm core depth

We counted in total 70 samples, and identified 65 benthic (44 calcareous and 21 agglutinated) and three planktic foraminiferal species (Table A2 in the appendices) in the top 345 cm of core 73G. Both benthic and planktic specimens were well-preserved and showed minor or no signs of post-mortem dissolution of the tests.

250 The foraminiferal content is relatively low throughout the core, the benthic foraminiferal concentration varies between 17 and 158 individuals (ind.) g⁻¹ wet sediment (sed.) (calcareous 2-128 ind. g⁻¹ sed., agglutinated 10-55 ind./g sed.) (Fig. 5).

The concentration of planktic foraminifera is extremely low throughout the core, it lies between 0 and 13 ind. g⁻¹ sed. Planktic foraminifera are represented primarily by the polar species *Neogloboquadrina pachyderma*, occasionally accompanied in the interval between 340-130 cm (ca. 9.4-3.4 ka BP) by specimens of the subpolar species *Turboratalita quinqueloba* and *Neogloboquadrina incompta*. The overall occurrence of planktic foraminifera is averaging 2 % of the total foraminiferal fauna.

255 On average, benthic agglutinated foraminifera account for 65 % of the total foraminiferal assemblage. From the end of the record until 210 cm (ca. 5.4 ka BP) the calcareous species periodically outnumber the agglutinated ones. However, from 210 cm until the top of the record, agglutinated foraminifera continuously dominate the benthic assemblage (Fig. 5). Throughout the core, the most abundant benthic agglutinated species are *Portatrochammina bipolaris*, followed by *Ammoglobogerina globigeriniformis*, representing on average 42 % and 16 % (respectively) of the benthic agglutinated assemblage, and 27.5 % and 10.5 % of the total benthic assemblage. They are both continuously present throughout the core and their relative abundances do not show strong fluctuations (Fig. 6).

260 Benthic calcareous foraminifera represent on average 33 % of the total foraminiferal assemblage. The most abundant benthic calcareous species are *C. reniforme*, *Cassidulina neoteretis*, *E. clavatum* and *Stetsonia horvathi* (representing on average 8.7%, 6.4%, 5.9% and 4.4% of the whole benthic assemblage, respectively). From these four species the abundances of *E. clavatum* shows the least variability throughout the record (Fig. 5).




Figure 5: Foraminiferal concentrations and relative abundances of nine selected benthic calcareous species (expressed as a percentage of total benthic foraminiferal content) versus calibrated age along sediment core DA17-NG-ST7-73G. The depicted species were chosen in order to show changes in the water masses. Red colour represents species that indicate warm Atlantic Water inflow; orange represents species that indicate in the Arctic recirculated, chilled Atlantic Water influence; dark blue represents Arctic Water species; light blue represents a species that indicates sea ice. Ecozones (I-V) are shown on the right side of the figure. Black arrows next to the left primary y-axis (Age) mark radiocarbon dates. Note that the x-axes have different scaling.


In the following sections, we describe foraminiferal assemblage zones (ecozones) in the dated section of the core (345-0 cm core depth). These zones were defined by visual evaluation and interpretation of the species abundances. Here it should be mentioned that also species with relatively low abundances can be significant for the environmental interpretations. Boundaries were placed where major changes occurred in the relative abundances of the most important benthic calcareous and agglutinated species, indicating changes in the environment (Figs. 5 and 6). Unless otherwise specified, all relative frequencies are provided as average values for the interval.

4.4.3 Ecozone I (345-310 cm; ca. 9.4-8.2 ka BP)

This ecozone is characterised by the highest content of total foraminifera (171 ind. g⁻¹ sed.), including the highest concentrations of planktic (13 ind. g⁻¹ sed.) and benthic calcareous (128 ind. g⁻¹ sed.) foraminifera throughout the core. These three categories show two peaks in this interval: the first and bigger at ca. 9 ka BP and the second, smaller one at ca. 8.7 ka BP. We find in this interval the lowest agglutinated/calcareous ratio (unshown), as the calcareous specimens almost continuously outnumber the agglutinated ones. The average concentration of agglutinated foraminifera is 29 ind. g⁻¹ sed. in this zone, and it does not vary significantly throughout the other intervals either (27-29 ind. g⁻¹ sed.). The benthic calcareous assemblage is dominated by *C. reniforme* (relative abundance on average 14 %), followed by *S. horvathi* (12 %), *E. clavatum* (11 %) and *C. neoteretis* (9 %), as important accessory species.

4.4.4 Ecozone II (310-245 cm; ca. 8.2-6.2 ka BP)

290 In this ecozone, we can recognise a distinctive decrease in planktic and benthic calcareous foraminiferal concentrations, and as a consequence, a higher benthic agglutinated/calcareous species ratio. The relative abundance of *C. neoteretis* promptly increases, and it becomes the dominating species of the benthic calcareous fauna in this interval (highest relative abundance 27 %, on average 13 %), followed by *C. reniforme* (10 %). *Pullenia bulloides* suddenly appears in the record becoming an important accessory species, and, at the same time, the relative abundance *S. horvathi* drastically decreases and it remains around 1 % during the whole period. Even though the relative abundances of *E. clavatum* decrease compared to the previous interval, this species shows its highest relative abundance throughout the core in this ecozone at 255 cm (17 %; ca. 6.5 ka BP). The relative abundance of agglutinated species *Adercotryma glomerata* increases from 310 cm on and remains on the same level throughout this zone.



300 **Figure 6: Relative abundances of the most abundant (>5% in at least one sample) benthic agglutinated foraminiferal species (expressed as a percentage of total benthic foraminiferal content) versus calibrated age along sediment core DA17-NG-ST07-73G. Ecozones (I-V) are shown on the right side of the figure. Black arrows next to the left primary y-axis (Age) mark radiocarbon dates. Note that the x-axes have different scaling.**

4.4.5 Ecozone III (245-165 cm; ca. 6.2-4.2 ka BP)

305 The concentrations of planktic and benthic calcareous foraminifera continue to decrease in this ecozone. From 245 cm, the relative abundance of *C. neoteretis* drops back to an average of 7 %, while the relative abundance of *S. horvathi* rises (from 1 % to 7 %). Although still only found in low numbers, *Epistominella arctica* also increases (from 0.4 % (previous interval) to 0.8 %), just like *Textularia torquata*, while the relative abundances of *A. glomerata* decreases.

4.4.6 Ecozone IV (165-125 cm; ca. 4.2-3.2 ka BP)

310 The base of this ecozone is defined by the drastic decrease in benthic calcareous foraminifera concentrations (from 15.7 to 8.4 ind. g⁻¹ sed.), and increased relative abundances of *E. arctica* and *Stainforthia feylingi*. *C. neoteretis* drops dramatically (to 3 %), just like *C. reniforme* that reaches its lowest relative abundance in this zone (5 %). On the other hand, agglutinated species *A. glomerata* and *Reophax fusiformis* shows a drastic peak at the beginning of this ecozone, and the relative abundances of *T. torquata*, *Textularia* cf. *earlandi* and *Spiroplectammina biformis* starts to increase as compared to the previous interval.

4.4.7 Ecozone V (125-0 cm; ca. 3.2-0.3 ka BP)

315 The concentrations of planktic foraminifera and the relative abundances of *C. neoteretis* further decrease and reach their lowest level throughout the core, while *C. reniforme* shows an increase compared to ecozone IV, just like *S. feylingi*. The relative abundances of agglutinated species *T. torquata*, *T. cf. earlandi* and *S. biformis* continue to increase, while *Saccamina difflugiformis* shows a steep rise unique to this interval.

4.5 Stable isotopes

320 From the 70 analysed samples the results of three samples were not accepted: one (180 cm) due to technical problems, and two gave too low signal due to small sample size (0 cm and 130 cm). The sample at 140 cm gave a comparatively very low $\delta^{18}\text{O}$ (2.25 ‰) and a very high $\delta^{13}\text{C}$ value (-1.07 ‰), but these outliers were nonetheless accepted.

325 Except for the one outlier, the variation of $\delta^{18}\text{O}$ values is rather small throughout the core (between 2.85 and 3.69 ‰, with a mean of 3.29 ‰), and the values remain relatively constant, albeit with a slight increase towards the top of the core. The $\delta^{13}\text{C}$ values show a constant increase from the bottom to the top of the core, varying between -1.07 and -2.61 ‰ (Fig. 7).

5 Discussion

5.1 Origin of Atlantic-sourced water at the core site

330 Although the study site is located on the central flow route of the EGC, the path of the influx of the Atlantic-sourced water, which is clearly identified as the bottom layer in the CTD profiles (Fig. 2), is less clear. The CTD profile with bottom water temperatures below 1 °C and salinities of 34.8 suggest that the Atlantic-sourced water is closer in character to the water found in the Westwind Trough than the Atlantic-sourced water presently seen in the Norske Trough (c.f., Budéus et al., 1997). This may indicate that cool Arctic Atlantic Water is the primary source of the Atlantic subsurface waters, rather than the warmer Return Atlantic Water (cf. Rudels et al., 2005; Schaffer et al., 2017). The Atlantic-sourced water may thus reach our study site via the Westwind Trough, but may also be present due to wind-driven upwelling. This process leads to considerable modification of local water masses and shelf break exchanges (e.g., Estrade et al., 2008; Kirillov et al., 2016).

5.2 Environmental significance of foraminiferal assemblages

340 In order to be able to describe the changes in water masses over time on the NE Greenland shelf, we place selected benthic calcareous foraminifera species into groups that are based on environmental preferences of the species (Table A3 in appendices). The Atlantic Water group includes *C. neoteretis* and *P. bulloides*. These species indicate warm and saline AW inflow underneath cold and low salinity surface waters (e.g., Mackensen and Hald, 1988; Seidenkrantz 1995; Rytter et al., 2002; Jennings et al., 2004; Jennings et al., 2011; Cage et al., 2021). The chilled Atlantic Water group includes *I. norcrossi* and *M. barleeaanum*. These species have been previously linked to cool AW (e.g., Slubowska-Woldengen et al., 2007; Perner et al., 2011; Perner et al., 2015; Cage et

al., 2021), and we use them to represent the relative contribution of chilled AW recirculated in the Arctic Ocean to the EGC. The Arctic Water group includes *S. horvathi* and *E. arctica*. Both species live at present beneath perennial sea ice, in the deeper part of the Arctic Ocean, thus indicating cold, Arctic originated deep-waters (e.g., Green, 1960; Lagoe, 1979; Wollenburg and Mackensen, 1998; Jennings et al, 2020). *Cassidulina reniforme* is also an Arctic species, although it requires stable conditions (e.g., Hald and Korsun, 1997; Consolaro et al., 2018), and in some studies it has been considered linked to highly chilled Atlantic Water (Ślubowska et al., 2005; Hansen et al., 2020). We consider *S. feylingi* as a sea-ice edge indicator species that tolerates unstable conditions (Knudsen and Seidenkrantz, 1994; Seidenkrantz, 2013); its increase may refer to the location of a sea-ice margin at the study site. Moreover, we use in our interpretation the abundances of the agglutinated species *A. glomerata*, *T. earlandi*, *T. torquata*, *S. biformis* and *S. difflugiformis*. *A. glomerata* is often correlated with Atlantic sourced waters in high latitudes (e.g., Hald and Korsun, 1997, Lloyd 2006; Perner et al., 2012). Many studies find *T. earlandi*, *T. torquata* and *S. biformis* in areas influenced by cold, low-salinity Arctic water, often in glaciomarine environments (e.g., Jennings and Helgadottir, 1994; Korsun and Hald, 2000; Perner et al, 2012; Perner et al., 2015, Wangner et al., 2018). *S. difflugiformis* is described as closely associated with Polar Water on the East Greenland shelf and in fjords (Jennings and Helgadottir, 1994).




Figure 7: From the bottom to the top: $\delta^{13}\text{C}$ (a) and $\delta^{18}\text{O}$ (b) results and relative abundances of the four benthic calcareous foraminifera groupings (c-f) (based on environmental preferences of the contained species, see Fig. 5 and Appendix Table A3) from this study (core 73G), compared to other proxy records from the NE Greenland shelf and western Svalbard. IP_{25} (g) and HBI III (h) from core PS93/025 (Syring et al., 2020a). Relative abundances of planktic polar species *Neogloboquadrina pachyderma* (sin.) and subpolar species *Turborotalita quinqueloba* (i) from core JM10-330 (Consolaro et al., 2018). HTM: Holocene Thermal Maximum. Black arrows on the x-axis mark radiocarbon dates.

5.3 Palaeoenvironmental interpretation

5.3.1 Bottom 65 cm of the sediment core

The combined presence of an assemblage of well-preserved small tests of Arctic foraminiferal species and many extinct Pliocene to early-Pleistocene specimens, today found in Plio-Pleistocene assemblages from Baffin Island and Peary Land (Feyling-Hanssen, 1976; Feyling-Hanssen, 1980; Feyling-Hanssen et al., 1983; Funder et al., 2001), suggest that these Plio-Pleistocene foraminifera have been reworked from older deposits into a glaciomarine environment with *in situ* Arctic foraminifera. The closest known Plio-Pleistocene deposit is the Kap København Formation from Peary Land, northern Greenland (Funder et al., 2001; see Fig. 1b), but

370 it is unknown if any such deposits may be found outcropping to the sea floor off NE Greenland. Reworking of older sediments during glacier ice retreat has previously been observed (e.g., Seidenkrantz et al., 2019). Therefore, we here suggest that the breakup and significant retreat of a nearby glacier caused reworking of older sediments and thus, the lowermost section of sediment from before 9.4 ka was deposited during a period of glacier retreat. However, further studies are required to resolve this issue.

5.3.2 Early Holocene (ca. 9.4-8.2 ka BP; ecozone I)

375 This interval was characterised by high percentages of *S. horvathi*, which has previously been considered linked to high sea-ice concentrations (e.g., Green, 1960; Lagoe, 1979; Wollenburg and Mackensen, 1988; Jennings et al., 2020) combined with the presence of Atlantic Water indicator species such as *C. neoteretis*, *I. norcrossi* and *M. barleeaanum*. This combination of Atlantic-sourced bottom water and sea ice suggest a highly stratified water column in the early Holocene (Fig. 5). Cold, heavily sea ice-loaded surface waters entrained in PW of the upper EGC characterised the upper water column, while at the same time the influx of warm Atlantic-sourced water from either the RAC or the AAW (or a combination of both) was also strong. When compared to the data from the top of the core, the water column stratification between the cold and warmer waters was likely more pronounced than today. Relatively high counts of the terrestrially-derived elements Si, K and Ti, (Ren et al., 2009) together with relatively low $\delta^{18}\text{O}$ values may indicate increased melt water influence from the Greenland Ice Sheet (Figs. 3 and 7). However, the low $\delta^{18}\text{O}$ values seen here may be also a consequence of the warmer bottom waters.

385 The interval was also likely characterised by high surface water productivity, as shown by high concentrations of planktic and benthic calcareous foraminifera and the presence of *E. arctica*, which thrives in high-productivity environments (Wollenburg and Kuhnt, 2000). It is also supported by a pronounced peak in Ca/Fe ratio, which is considered as a general proxy for palaeo-productivity (e.g., Vare et al., 2009) and low and slowly rising benthic $\delta^{13}\text{C}$ values that can also be related to high level of bioproductivity, due to decomposition of isotopically light organic carbon in the lower part of the water column. High surface water productivity may sound contradictory to heavy sea-ice cover, however, the planktic foraminifera may have been transported to the site in the subsurface water layer from a nearby more open-ocean site, suggesting increased surface water-bioproductivity in the general region. We propose that high productivity combined with sea ice indicates that our site was located west of the Polar Front, but that the Polar Front was situated in close proximity to our location in this time interval. It should be noted that in the XRF data, we recognise a short-lived, but clear peak in terrestrially-derived sediments around 8.5-8.4 ka BP (Fig. 3). This may be potentially linked to increased iceberg release during the so-called 8.2 ka event (i.a., Alley et al., 1997; Barber et al., 1999), however, this signal seems to be only reflected in the results of the sedimentological analyses. It is not recognizable in the foraminiferal assemblage changes/stable isotope results, although the lower temporal resolution of the latter may also not allow us to identify any changes.

400 Our suggestions of cold water and sea ice at the surface combined with Atlantic-sourced bottom waters is in accordance with a recent reconstructions from the NE Greenland shelf (Syring et al., 2020a; Zehnich et al., 2020; core PS93/025). The study of Syring et al. (2020a) reveals a cold interval with increased sea-ice cover between 9.3 and 7.9 ka BP, shown by lower concentrations of marginal sea-ice proxy HBI III and IP₂₅, interpreted as low sea-ice algal productivity caused by an expansion of the sea-ice cover (Fig. 7). The stable isotope results of Zehnich et al. (2020) point to a relatively thick subsurface Atlantic water mass and high levels of productivity in the upper water column. Foraminiferal records from the Southeast Greenland shelf (Jennings et al., 2011; cores MD99-2322 and MD992317) suggests cold and unstable sea-surface conditions between 9.4 and 8.1 ka BP. Moreover, cool deep-water conditions in the eastern Fram Strait during this period (Consolaro et al., 2018; core JM10-330) point to a weakened RAC, suggesting that the strong water column stratification on the NE Greenland shelf was due to increased AAW influx.

5.3.3 Holocene Thermal Maximum (ca. 8.2-6.2 ka BP; ecozone II)

410 The interval from 8.2 to 6.2 ka BP was characterised by the warmest bottom-water conditions of the Holocene on the NE Greenland shelf, as well as reduced sea-ice cover, as indicated by the almost complete absence of sea-ice indicator species and maximum abundances of Atlantic Water species and other species indicating stable bottom waters. *Cassidulina neoteretis* and *A. glomerata* had their highest relative abundances during this interval (except for an extreme peak of *A. glomerata* at ~4 ka BP), and *P. bulloides* appeared around 8 ka BP in the record after a long absence (Figs. 5 and 6), suggesting highly stable bottom waters (Rytter et al., 415 2002). A concurrent significant decrease in planktic and benthic calcareous foraminiferal concentrations indicates a transition towards lower surface and subsurface water productivity, also shown in decreasing Ca/Fe element ratios. We suggest that this may be explained by a strengthening of the RAC, transporting RAW to our site and causing higher-than-present entrainment of warm Atlantic-sourced water into the EGC. ~~As a consequence of likely vertical expansion of the Atlantic-sourced water layer, surface waters likely also warmed, and the Polar Front moved from our location to further north or further inland during this period, which represents the warmest of our record, and which we here refer to as the Holocene Thermal Maximum (HTM).~~

In the eastern Fram Strait several studies (Slubowska-Woldengen et al., 2007; core JM02-440; Werner et al., 2013; core MSM5/5-712; Werner et al., 2016; core MSM5/5-723; Consolaro et al., 2011; core JM10-330, see Fig. 1) recorded strong AW flow and warm sea-surface conditions until 7-6.8 ka BP (Fig. 7), which would have resulted in a stronger RAC. In line with our findings, 425 Müller et al. (2012) (core PS2641-4) and Jennings et al. (2011) (cores MD99-2322 and MD99-2317) reconstructed relatively warm conditions during the first part of the mid Holocene on the middle East and Southeast Greenland shelf in surface and subsurface waters, respectively. Also further south in the Nordic Seas, thermal optimum-like conditions with warm sea surface temperatures prevailed until ca. 6 ka BP (Bauch et al., 2001; core PS1243).

5.3.4 Cooling after the HTM (ca. 6.2-4.2 ka BP; ecozone III)

430 From 6.2-4.2 ka BP benthic foraminiferal assemblages again start to resemble those of the period prior to 8.2 ka BP, with increased percentage of *S. horvathi* and a decrease in the Atlantic water indicator *C. neoteretis* (Fig. 5). This suggests return to a more stratified water column with sea-ice loaded surface water and Atlantic-sourced subsurface waters. However, in contrast to Ecozone I, neither the benthic and planktic foraminiferal concentrations nor stable isotope data suggest increased bioproductivity.

435 Along West Spitsbergen the inflow of subsurface AW, and thus the RAC weakened after 7 ka BP (Consolaro et al., 2018; core JM10-330;), and a gradual cooling of surface waters started around 7-6.6 ka BP (Müller et al., 2012; core MSM5/5-712), accompanied by increased sea-ice cover (Slubowska-Woldengen et al., 2007; core JM02-440). A sediment core retrieved north from our location (Zehlich et al., 2020; Syring et al., 2020a; core PS93/025) shows after 7.2 ka BP stronger vertical stratification (derived from benthic-planktic $\delta^{18}\text{O}$ difference) and after 5.5 ka BP seasonally more extended sea-ice cover (derived from decreasing HBI III and increasing IP₂₅ concentrations; Fig. 7) and decreasing primary production. Accordingly, at the same time, 440 Perner et al. (2015) (core PS641-4) recorded the southward relocation of Polar Front to close to their site on the middle East Greenland shelf. Giraudeau et al. (2004) (core MD99-2269) and Ran et al. (2006) (core MD99-2275) recorded increased influence of cool, low salinity polar waters of the EGC on the North Iceland Shelf between 6.5-3.5 ka BP and 6.8-5.6 ka BP, respectively. In the Iceland Basin a long- term cooling trend started at between 6.8 and 6.1 ka BP (Orme et al., 2018; Van Nieuwenhove et al., 445 2018; core DA12-11/2).

5.3.5 Further cooling in the mid Holocene (ca. 4.2-3.2 ka BP; ecozone IV)

Our record shows clear evidence for significant changes occurring at 4.2 ka BP. The period starts at our location with a sudden rise in the relative abundances of calcareous sea-ice indicator species *S. feylingi* and in the relative abundances of agglutinated species *T. earlandi*, *T. torquata* and *S. biformis* (Figs. 5 and 6), which are often connected to cold, low-salinity PW (Jennings and Helgadottir, 1994; Korsun and Hald, 2000; Perner et al., 2012; Perner et al., 2015; Wangner et al., 2018). Furthermore, the interval is marked by lowest abundances of warm Atlantic Water species and very low concentrations (<1 ind./g) of planktic foraminifera. The benthic agglutinated/calcareous foraminiferal ratio increases and the terrestrially derived elements Si, K and Ti show minimal values (Fig. 3), indicating very low meltwater influence in the EGC. Since the foraminiferal assemblage points to low bottom water temperatures, we interpret the slightly higher benthic $\delta^{18}\text{O}$ values and increasing $\delta^{13}\text{C}$ values after 4.2 ka BP (Fig. 7) as indicative of general cooling and weak influence of AW. Accordingly, the light $\delta^{18}\text{O}$ spike found in the middle of this interval cannot be attributed to sudden bottom water warming on the basis of the foraminiferal fauna. It is more readily explained by local brine rejection during sea ice formation, which can carry the light isotopic signal from the surface to the ocean bottom (e.g., Mackensen and Schmiedl, 2016). It should here be noted that apart from a single sample with increased relative frequencies of the warmer-water agglutinated species *A. glomerata* (Figs. 6 and A3), there is no evidence of a shorter-lived climate excursion that might be linked to the so-called 4.2 ka BP event (Weiss, 2017), only a general change in environment at this time.

We suggest that increased PW at the surface of the EGC and reduction of the RAC at subsurface levels led to freshening and reduced stratification of the water column at our site; it may also have experienced (near) perennial sea-ice cover. The reduced strength of the RAC was likely caused by an overall decrease in AW transport of the North Atlantic and West Spitsbergen currents to the eastern Fram Strait, accompanied by surface-water cooling and increased sea-ice coverage after 5.5 ka BP (Hald et al., 2007; core MSM5/5-712; Werner et al., 2013; core PS1878; Telesinski et al., 2014b; core PS1878; Consolaro et al., 2018; core JM10-330; Fig. 7). $\delta^{18}\text{O}$ results of Zehnich et al. (2020; core PS93/025) point to colder conditions and weak AW advection after 5 ka BP north from our location. Harsh conditions and strengthened EGC with permanent sea-ice cover dominated the central Greenland shelf as well (Perner et al., 2015; Kolling et al., 2017; core PS641-4). Further south on the shelf Jennings et al. (2002) and Perner et al. (2016) (core JM96-1206) described increased PW influence in the EGC, glacier advance and iceberg rafting during this period.

5.3.6 Neoglaciation (ca. 3.2-0.3 ka BP; ecozone V)

The period from ca. 3.2 to 0.3 ka BP is characterised by a further increase in the relative abundances of sea-ice edge indicator species *S. feylingi* and the agglutinated species *S. difflugiformis*. Atlantic Water indicator *C. neoteretis* is only found in relatively low numbers, although there is some increase in species linked to chilled Atlantic Water (Figs. 5, 6 and 7). Continuously increasing benthic $\delta^{13}\text{C}$ values indicate strong ventilation of the water column (Fig. 7). The results point to cold and unstable conditions with minimum surface water productivity and increasing sea-ice cover at our location. During this period, the EGC was likely strengthened compared to previous times, with a thick layer of cold and fresh PW on the surface and recirculated AAW inflow from the Arctic at subsurface levels, as seen in the chilled Atlantic Water group.

Previous studies have indicated that during the late Holocene, the West Spitsbergen Current transported less and less heat (Slubowska-Woldengen et al., 2007; core JM02-440), causing less warm water to reach the EGC through the RAC. From West Spitsbergen (Müller et al., 2012; core MSM5/5-712; Consolaro et al., 2018; core JM10-330) and from several other locations in the Greenland Sea (e.g., Telesinski et al., 2014b, core PS1878) and West Greenland (e.g., Seidenkrantz et al., 2007; core 248260;

485 Seidenkrantz et al., 2008; cores DA00-02P and DA00-03P) a general cooling trend has been reported, with more severe sea-ice conditions. On the Iceland shelf strongly reduced Irminger surface water and low coccolith carbonate sedimentation indicate extreme advection of polar waters and extended sea-ice development (Giraudeau et al., 2004; core MD99-2269).

5.4 Development of Holocene oceanographic conditions on the NE Greenland shelf

490 Early Holocene warming of the circum-Arctic region (Kaufmann et al., 2004) led to enhanced Greenland Ice Sheet and glacier melting around the Arctic. During the Last Glacial Maximum (~26-19 ka BP) the Greenland Ice Sheet extended onto the shelf offshore NE Greenland, perhaps even reaching the shelf break (Bennike and Weidick, 2001; Arndt et al., 2017). However, as a consequence of rising air temperatures due to the orbitally-forced Northern Hemisphere summer insolation maximum (reported at approximately 10 ka BP; Andersen et al., 2004; Jansen et al., 2008), the deglaciation of the outer coast and retreat of the ice margin to its present location occurred already between 11.7 and 9.3 ka BP (Larsen et al., 2018); the ice shelf of 79NG rapidly retreated through the fjord between 9.6 and 7.5 ka BP (Syring et al., 2020b). The drastic ice recession of the early Holocene produced an extended meltwater surface layer in the Greenland region prior to 8.6 ka BP (Seidenkrantz et al., 2013). The extensive melting of the Greenland Ice Sheet was strong enough to act as negative feedback to the early Holocene warming, and delayed the HTM with 2 kyr at our location compared to the eastern parts of the Nordic Seas (Blaschek and Renssen, 2013). This thus explains the cold, heavily sea ice-loaded surface waters and strongly stratified water column that characterised our study site on the NE Greenland shelf between 9.4 and 8.2 ka BP.

The strong freshwater discharge might have also weakened the recording of the 8.2 ka cooling event at our location, although a short-term increase in terrestrial deposits suggests transport of sediments from land via meltwater or icebergs during this event. This 8.2 ka cold event has been described in a number of palaeoceanographic archives from Greenland and the northern North Atlantic (e.g., Risebrobakken et al., 2003; Ellison et al., 2006; Rasmussen et al., 2007), and it is believed to be connected to the collapse of the Laurentide Ice Sheet and drainage of Lake Agassiz (e.g., Alley et al., 1997; Barber et al., 1999; Hillaire-Marcel et al., 2007; Hoffman et al., 2012). However, in many other records from areas that are mainly influenced by the EGC, WGC or the Baffin-Labrador Current system (Keigwin et al., 2005; Sachs 2007; Seidenkrantz et al., 2013), the 8.2 ka cooling event cannot be recognised; these sites were all characterised by the permanent presence of a low-salinity, cold surface layer at that time, which may also explain the weak signal of this event at our site.

After ca. 8.2 ka BP, our data suggest that increased entrainment of warm Atlantic-sourced water within the RAC changed the water column properties on the NE Greenland shelf. During the subsequent Holocene Thermal Maximum, both subsurface and surface water temperatures increased, and consequently, open water conditions prevailed in the next ca. 2 kyr close to our location. Due to the warming, the floating ice margin of 79NG decreased (Bennike and Weidick, 2001), but on the other hand, the fresh, less dense basal melt water of the glacier isolated the landfast sea-ice cover from the warm subsurface waters, and thus, stabilised the Norske Øer Ice Barrier (Mayer et al., 2000; Syring et al., 2020b). Further south, parallel to the warming of the EGC (Jennings et al., 2011; Müller et al., 2012), the Irminger Current increased its strength, as recorded on the North Icelandic Shelf (Ran et al., 2006; Cabedo-Sanz et al., 2016).

520 After the Thermal Maximum, cooling started on the NE Greenland shelf between 6.2 and 4.2 ka BP (e.g., this study; Andrews et al., 1997; Jennings et al., 2002; Zehnich et al., 2020), with decreased, but still persistent inflow of subsurface AW from the RAC. The EGC became stronger, with sea-ice loaded surface waters and relatively warm Atlantic-sourced subsurface waters. Coincident

with the expansion of the EGC, several studies from the Nordic Seas (e.g., Bauch et al., 2001; Hall et al., 2004; Hald et al., 2007) infer a weakening of the AMOC, increased water column stratification and less ventilated subsurface during this period. In line with a decreased flux of recirculating AW onto the NE Greenland shelf, the Northeast Greenland Ice Margin started to advance from its mid Holocene minimum around 6 ka BP (Larsen et al., 2018). Reduced amounts of warm AW on the inner continental shelf most probably reduced the basal melting within the 79NG fjord and may have contributed to the re-advance of the ice shelf of 79NG seen from around 4.5 ka BP on (Bennike and Weidick, 2001; Syring et al., 2020b).

The Neoglacial cold interval started on the East Greenland shelf approx. 3.5-3.2 ka BP, with increased freshwater forcing from the Arctic Ocean and advance of the Greenland Ice Sheet (this study; Andersen et al., 2004; Jennings et al., 2011). According to model simulations of Renssen et al. (2006), the expansion of sea ice may be associated with a cooling triggered by a negative solar irradiance anomaly, which was amplified through a positive oceanic feedback mechanism. The cooling caused temporary relocation of deep-water formation sites in the Nordic Seas, which was accompanied by a distinct reduction in AMOC strength (Hall et al., 2004). The increase in sea-ice extent stratified the water column and hampered the deep-water formation, leading to additional cooling and more sea ice (Renssen et al., 2006).

5.5 Variations of Atlantic and Polar Water entrainment in the EGC during the Holocene and their implications for the general climatic trends in the Nordic Seas

Distinct variability in subsurface water mass properties on the NE Greenland shelf through the Holocene point to broad-scale changes in the proportion of RAW/AAW and PW in the EGC, and thus, to a strong covariance with the northward heat transport in the Nordic Seas. The increased entrainment of warm Atlantic-sourced water into the EGC that we document during the HTM, coincides with increased advection of warm AW in the West Spitsbergen Current, driven by increased wind force and/or by stronger thermohaline circulation (Sarthein et al., 2003; Slubowska-Woldengen et al., 2007; Knudsen et al., 2011; Werner et al., 2013; Werner et al., 2016; Consolaro et al., 2018).

During the mid Holocene, the summer insolation declined at the northern latitudes due to the changes in the Earth's orbital parameter (Berger, 1978). From 7 ka BP, the summer sea surface temperatures of the Nordic Seas started to decrease (Koç et al., 1993; Andersen et al., 2004). This cooled AW progressed northward carrying less and less heat to the North Atlantic and West Spitsbergen currents (Sarthein et al., 2003; Slubowska-Woldengen et al., 2007; Risebrobakken et al., 2011; Consolaro et al., 2018; Hald et al., 2007), also due to a weakened AMOC (e.g., Bauch et al., 2001; Hall et al., 2004; Hald et al., 2007; Knudsen et al., 2011). We suggest that, in response to the reduced warm water transport through the RAC, the proportion of cooled AAW and PW in the EGC increased, which ended the ameliorated conditions on the East Greenland shelf seen at our study site.

The northward flow of AW with the North Atlantic and West Spitsbergen currents to the eastern Fram Strait, and consequently the RAW reaching the EGC, arrived to a minimum during the late Holocene, after 4.5 ka BP (Slubowska-Woldengen et al., 2007). Consequently, the EGC strengthened, carrying a thick layer of fresh PW southward, as also seen after ca. 4.2 ka BP (and even more so after 3.2 ka BP) in DA17-NG-ST07-73G. A similar pronounced colder period after 4 ka BP is recognised in some (but not all) reconstructions north and south of Iceland, in the Greenland Sea and to the west of Svalbard (Moossen et al., 2015; Orme et al., 2018; Telesinski et al., 2014b; Werner et al., 2016; respectively); in particular at locations influenced by the EGC. This gradual cooling trend in the northern North Atlantic is associated therefore with a continuous transition from a more prevailing positive phase towards a period when the prevailing NAO and AO phase was commonly negative (e.g., Andersen et al., 2004; Orme et al., 2018). During intervals of a negative NAO/AO high pressure over Greenland and consequently strengthened northerly

winds over East Greenland can lead to strengthened outflow of PW from the Arctic Ocean through the Fram Strait; this water will be carried southward by the EGC (e.g., Ionita et al., 2016). The late-Holocene trend towards a stronger and fresher EGC is likely to have increasingly impacted the Subpolar Gyre dynamics. Increased freshwater input to the Labrador Sea may have weakened the Subpolar Gyre circulation by preventing deep convection (Hillaire-Marcel et al., 2001), and consequently reduced the amount of water entrained into the North Atlantic Current (Moros et al., 2012). This, on the other hand, may have forced enhanced northward heat transport via the North Atlantic Current, and may have helped to restart (Thornalley et al., 2009) the previously weakened AMOC (e.g., Cronin et al., 2003; Oppo et al., 2003).

6 Conclusions

The presented multiproxy study, based on benthic foraminiferal assemblage-, geochemical- and sedimentological analyses of sediment core DA17-NG-ST07-73G allowed us to reconstruct changes in sea surface productivity, subsurface water temperatures, sea-ice conditions, Greenland Ice Sheet melting and thus, in the strength of the EGC over the last ca. 9.4 ka BP on the NE Greenland shelf.

- Between 9.4 and 8.2 ka BP the water column was highly stratified, with cold, heavily sea ice-loaded surface waters and strong influx of warm waters in the subsurface. High surface water productivity suggests that the polar front was close to our location.
- A short-lived peak in terrestrially-derived elements suggesting transport of sediments from land via meltwater or icebergs may be linked to the so-called 8.2 ka BP event.
- The interval from 8.2 to 6.2 ka BP was characterised by the warmest bottom-water conditions of the Holocene on the NE Greenland shelf, with low surface water productivity and strong AW influence (RAC) in the weakened EGC.
- After the HTM the water column started to resemble that of the period prior to 8.2 ka BP. The EGC became stronger, with sea-ice loaded surface waters and relatively warm Atlantic-sourced subsurface waters. The subsurface inflow of warm AW from the RAC decreased, but remained persistent.
- After 4.2 ka BP increased PW at the surface of the EGC and reduction of the RAC at subsurface levels led to freshening and reduced stratification of the water column and to a (near) perennial sea-ice cover.
- The period from ca. 3.2 until 0.3 ka BP is characterised by cold and unstable conditions, with minimum surface water productivity, and possibly, sea-ice cover at our location. During this period, the EGC was likely strengthened, with a thick layer of cold and fresh PW on the surface and strong recirculated AAW inflow from the Arctic at subsurface levels.
- The proportion of PW and AW in the EGC shows a strong covariance with the northward heat transport to the northern North Atlantic. Thus, the cooling trend that characterises the Holocene after the HTM led to strengthened EGC and increased PW transport, probably due to transition to a more negative NAO/AO scenario, causing more PW to exit the Arctic Ocean via the Fram Strait. This stronger and fresher EGC in the late-Holocene is likely to have impacted the Subpolar Gyre circulation and helped to strengthen the AMOC.

Appendices

Table A1: Comparison of radiocarbon dates calibrated with the Marine20 (Heaton et al., 2020) and the Marine13 (Reimer et al., 2013) calibration curve in core DA17-NG-ST07-73G. For the dates calibrated with the Marine20 we used a local marine reservoir age correction (ΔR) of 0 ± 50 years; for the dates calibrated with the Marine13 we used a ΔR of 150 ± 50 years, as this is the approximate offset between the two curves (Heaton et al., 2020).

Lab. ID	Depth (cm)	Radiocarbon age (yr BP)	Error (yr)	Median cal. age (yr BP) Marine20, $\Delta R=0\pm 50$ yr	Median cal. age (yr BP) Marine13, $\Delta R=150\pm 50$ yr
ETH-95387	70.5	2475	60	1952	1946
ETH-95388	110.5	3275	70	2935	2915
ETH-110893	226	5645	30	5835	5872
ETH-95389	250.5	6015	70	6241	6284
ETH-95390	300.5	7595	70	7870	7902
ETH-95391	345.5	9015	70	9536	9528

Table A2: Foraminiferal taxa identified in this study (core depth 345-0 cm), with their original reference. Benthic calcareous, agglutinated and planktic species are separated and listed alphabetically.

Benthic agglutinated

Adercotryma glomerata (Brady, 1878)
Ammodiscus sp.
Ammoglobogerina globigeriniformis (Parker and Jones, 1865)
Cibrostomoides kosterensis (Höglund, 1947)
Deuterammia grahami Brönnimann and Whittaker, 1988
Deuterammia montagui Brönnimann and Whittaker, 1988
Hormosinella sp.
Portatrochammina bipolaris Brönnimann and Whittaker, 1980
Recurvoides turbinatus (Brady, 1881)
Reophax fusiformis (Williamson, 1858)
Reophax guttifer/rostrata (Brady, 1881)/Höglund, 1947
Reophax sp.
Reophax subfusiformis Earland, 1933
Rhabdammina sp.
Saccammina difflugiformis (Brady, 1879)
Saccammina sphaerica Brady, 1871
Saccorhiza ramosa (Brady, 1879)
Spiroplectammina biformis (Parker and Jones, 1865)
Textularia cf. *earlandi* Parker, 1952
Textularia torquata Parker, 1952
Textularia kattagatensis Höglund, 1948

Benthic calcareous

Islandiella helanae Feyling-Hanssen and Buzas, 1976
Ammonia sp.
Astrononion gallowayi Loeblich and Tappan, 1953
Bolivina aff. *albatrossi* Cushman, 1922
Buccella frigida (Cushman, 1922)
Buliminella elegantissima (d'Orbigny, 1839)
Cassidulina neoteretis Seidenkrantz, 1995
Cassidulina reniforme Nørvang, 1945
Ceratobulimina arctica Green, 1959
Cibicides lobatulus (Walker and Jacob, 1798)
Dentalina pauperata d'Orbigny, 1846
Elphidium albiumbilicatum (Weiss, 1964)

Elphidium clavatum Cushman, 1930
Elphidium frigidum Cushman, 1933
Elphidium hallandense Brotzen, 1943
Eoponidella pulchella (Parker, 1952)
Epistominella arctica Green, 1959
Epistominella vitrea Parker, 1953
Fisurina sp.
Florius sp.
Glabratella arctica Scott and Vilks, 1991
Globulina oculus Jennings, Seidenkrantz and Knudsen, 2020
Guttulina glacialis (Cushman and Ozawa, 1930)
Haynesina nivea (Lafrenz, 1963)
Islandiella norcrossi (Cushman, 1933)
Lagena sp./*Procerolagena*
Melonis barleeaanum (Williamson, 1858)
Miliolinella subrotunda (Montague, 1803)
Nonionella iridea Heron-Allen and Earland, 1932
Nonionella labradorica (Dawson, 1860)
Polymorphina sp.
Pullenia bulloides (d'Orbigny, 1846)
Pyrgo williamsoni (Silvestri, 1923)
Quinqueloculina seminulum (Linnaeus, 1758)
Quinqueloculina stalker Loeblich and Tappan, 1953
Robertina arctica d'Orbigny, 1846
Sagrina sp.
Stainforthia concava (Höglund, 1947)
Stainforthia feylingi Knudsen and Seidenkrantz, 1994
Stainforthia fusiformis (Williamson, 1858)
Stetsonia horvathi Green, 1959
Trifarina fluens (Todd in Cushman and McCulloch, 1948)
Triloculina tricarinata d'Orbigny, 1846
Triloculina trihedra Loeblich and Tappan, 1953

Planktic

Neogloboquadrina incompta (Cifelli, 1961)
Neogloboquadrina pachyderma (Ehrenberg, 1861)
Turborotalita quinqueloba (Natland, 1938)

605 Table A3: List of benthic foraminiferal key species used for palaeoenvironmental reconstruction.


Species	Environmental preferences	References
Calcareous		
<i>Cassidulina neoteretis</i>	warm, saline Atlantic Water	Mackensen and Hald, 1988; Seidenkrantz, 1995; Rytter et al., 2002; Jennings et al., 2004; Jennigns et al., 2011, Perner et al., 2015; Cage et al., 2021
<i>Pullenia bulloides</i>	warm, saline Atlantic Water	Rytter et al., 2002; Jennings et al., 2011

<i>Cassidulina reniforme</i>	chilled Atlantic Water, stable conditions	Hald and Korsun, 1997; Slubowska et al., 2005; Slubowska-Woldengen et al., 2007; Perner et al., 2011; Consolaro et al., 2018
<i>Islandiella norcrossi</i>	chilled Atlantic Water	Slubowska-Woldengen et al., 2007; Perner et al., 2011; Perner et al., 2015; Cage et al., 2021
<i>Melonis barleeanum</i>	productivity, cool Atlantic Water	Perner et al., 2015
<i>Elphidium clavatum</i>	tolerates unstable conditions, in glaciomarine environments	Slubowska-Woldengen et al., 2007; Jennings et al., 2011; Perner et al., 2011
<i>Stetsonia horvathi</i>	cold, Arctic deep water	Green, 1960; Lagoe, 1979; Wollenburg and Mackensen, 1988; Jennings et al., 2020
<i>Epistominella arctica</i>	cold, Arctic deep water, productivity	Green, 1960; Lagoe, 1979; Wollenburg and Mackensen, 1988; Jennings et al., 2020
<i>Stainforthia feylingi</i>	tolerates unstable conditions, high productivity, often in sea-ice edge regions	Knudsen and Seidenkrantz, 1994
Agglutinated		
<i>Adercotryma glomerata</i>	Atlantic sourced waters	Hald and Korsun, 1997, Lloyd 2006; Perner et al., 2012; Wangner et al., 2018
<i>Textularia earlandi</i>	cold, low-salinity Arctic waters	Jennings and Helgadottir, 1994; Korsun and Hald, 2000; Wangner et al., 2018
<i>Textularia torquata</i>	cold, low-salinity Arctic waters	Perner et al., 2012; Perner et al., 2015; Wangner et al., 2018
<i>Spiroplectammina biformis</i>	cold, low-salinity Arctic waters	Jennings and Helgadottir, 1994; Korsun and Hald, 2000; Perner et al., 2012; Wangner et al., 2018
<i>Saccammina difflugiformis</i>	Polar Waters on the EG shelf	Jennings and Helgadottir, 1994

Table A4: List of abbreviations used in the manuscript.

NE Greenland	North East Greenland
EGC	East Greenland Current
RAC	Return Atlantic Current
AMOC	Atlantic Meridional Overturning Circulation
NAO	North Atlantic Oscillation
AO	Arctic Oscillation
PW	Polar Water
RAW	Return Atlantic Water
AAW	Arctic Atlantic Water
AW	Atlantic Water
HTM	Holocene Thermal Maximum

DA17-NG-ST07-72R



DA17-NG-ST07-73G

610

Figure A1: Comparison of XRF data from gravity core DA17-NG-ST07-73G (black) and Rumohr core DA17-NG-ST07-72R (red).






Figure A2: **Foraminiferal** concentrations and relative abundances of nine selected benthic calcareous species (expressed as a percentage of total benthic calcareous foraminiferal content) versus calibrated age along sediment core **DA17-NG-ST07-73G**. The depicted species were chosen in order to show changes in the environment. Red colour represents species that indicate warm Atlantic Water inflow; orange represents species that indicate in the Arctic recirculated, chilled Atlantic Water influence; dark blue represents Arctic Water species; light blue represents a species that indicates sea ice. Ecozones (I-V) are shown on the right side of the figure. Black arrows next to **the left primary y-axis (Age)** mark radiocarbon dates. Note that the x-axes have different scaling.

615



620 **Figure A3: Relative abundances of the most abundant (>5% in at least one sample) benthic agglutinated foraminifera species (expressed as a percentage of total benthic agglutinated foraminiferal content) versus calibrated age along sediment core DA17-NG-ST07-73G. Ecozones (I-V) are shown on the right side of the figure. Black arrows next to the left primary y-axis (Age) mark radiocarbon dates. Note that the x-axes have different scaling.**



625 **Figure A4: Foraminiferal concentrations showing the concentrations of nine selected benthic calcareous species versus calibrated age**
along sediment core DA17-NG-ST07-73G. The depicted species were chosen in order to show changes in the environment. Red colour
represents species that indicate warm Atlantic Water inflow; orange represents species that indicate in the Arctic recirculated, chilled
Atlantic Water influence; dark blue represents Arctic Water species; light blue represents a species that indicates sea ice. Ecozones (I-V)
are shown on the right side of the figure. Black arrows next to the left primary y-axis (Age) mark radiocarbon dates. Note that the x-
 630 **axes have different scaling.**




Figure A5: Foraminiferal concentrations of the most abundant (>5% in at least one sample) benthic agglutinated foraminifera species versus calibrated age along sediment core DA17-NG-ST07-73G. Ecozones (I-V) are shown on the right side of the figure. Black arrows next to the left primary y-axis (Age) mark radiocarbon dates. Note that the x-axes have different scaling.

635 Data availability

Data presented in this manuscript have been uploaded to PANGAEA (<https://doi.pangaea.de/10.1594/PANGAEA.934100>) and will be publicly accessible right after the publication of the manuscript.

Author contribution

640 Teodora Pados-Dibattista and Marit-Solveig Seidenkrantz developed the research idea. Teodora Pados-Dibattista carried out the sampling, data collection and the data analysis, with the help of Christof Pearce, Henrieka Detlef, Jørgen Brendtsen and Marit-Solveig Seidenkrantz. Christof Pearce performed the age modelling of the core. Teodora Pados-Dibattista prepared the manuscript with contributions from all co-authors.

Competing interests

Author Marit-Solveig Seidenkrantz is co-editor-in-chief of the journal.

645 **Acknowledgements**

We would like to thank the captain and crew as well as the shipboard scientific party onboard RV *Dana*. We also wish to thank Marianne Lyngholm Nielsen for carrying out the X-Ray Fluorescence core scanning and the magnetic susceptibility measurements and Nils Andersen, Leibniz Laboratory for Radiometric Dating and Stable Isotope Research at the Christian-Albrechts-University of Kiel, for the stable isotope measurements.

650 **Financial support**

The NorthGreen17 expedition was funded by the Danish Centre for Marine Research and the Natural Science and Engineering Research Council of Canada. The research was funded by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 792639, with further support from the Danish Council for Independent Research (grants no. 7014-00113B (G-Ice) and 0135-00165B (GreenShelf) to MSS) and the European Union's Horizon 2020 research and innovation program under grant agreement No. 869383 (ECOTIP).

References

Alley, R. B., Mayewski, P. A., Sowers, T., Stuiver, M., Taylor, K. C., and Clark, P. U.: Holocene climatic instability: A prominent, widespread event 8200 yr ago, *Geology*, 25(6), 483-486, 1997.

660 Andersen, C., Koç, N., Jennings, A. E., and Andrews, J. T.: Nonuniform response of the major surface currents in the Nordic Seas to insolation forcing: Implications for the Holocene climate variability, *Paleoceanography*, 19, PA2003, <https://doi:10.1029/2002PA000873>, 2004.

Andrews, J. T., Smith, L. M., Preston, R., Cooper, T., and Jennings, A. E.: Spatial and temporal patterns of iceberg rafting (IRD) along the East Greenland margin, ca. 68 N, over the last 14 cal. Ka, *J. Quaternary Sci.*: Published for the Quaternary Research Association, 12(1), 1-13, 1997.

665 **Ardyna, M., Babin, M., Gosselin, M., Devred, E., Rainville, L., and Tremblay, J.-É.: Recent Arctic Ocean sea ice loss triggers novel fall phytoplankton blooms, *Geophys. Res. Lett.*, 41, 6207–6212, doi:10.1002/2014GL061047, 2014.**

Arndt, J. E., Jokat, W., Dorschel, B., Myklebust, R., Dowdeswell, J. A., and Evans, J.: A new bathymetry of the Northeast Greenland continental shelf: Constraints on glacial and other processes, *Geochem. Geophys. Geosy.*, 16(10), 3733-3753, 2015.

670 Arndt, J. E., Jokat, W., and Dorschel, B.: The last glaciation and deglaciation of the Northeast Greenland continental shelf revealed by hydro-acoustic data, *Quaternary Sci. Rev.*, 160, 45–56, 2017.

Barber, D. C., Dyke, A., Hillaire-Marcel, C., Jennings, A. E., Andrews, J. T., Kerwin, M. W., Bilodeau, G., McNeely, R., Southon, J., Morehead, M. D., and Gagnon, J.-M.: Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes, *Nature*, 400(6742), 344-348, 1999.

675 Bauch, H. A., Erlenkeuser, H., Spielhagen, R. F., Struck, U., Matthiessen, J., Thiede, J., and Heinemeier, J.: A multiproxy reconstruction of the evolution of deep and surface waters in the subarctic Nordic seas over the last 30,000 yr., *Quaternary Sci. Rev.*, 20(4), 659-678, 2001.

Bennike, O. L. E. and Weidick, A.: Late Quaternary history around Nioghalvfjærdsfjorden and Jøkelbugten, North-East Greenland, *Boreas*, 30(3), 205-227, 2001.

680 Berger, A.: Long-term variations of caloric insolation resulting from the Earth's orbital elements, *Quaternary Res.*, 9(2), 139-167, 1978.

- Blaschek, M. and Renssen, H.: The Holocene thermal maximum in the Nordic Seas: the impact of Greenland Ice Sheet melt and other forcings in a coupled atmosphere–sea-ice–ocean model, *Clim. Past*, 9(4), 1629–1643, 2013.
- Bourke, R. H., Newton, J. L., Paquette, R. G., and Tunncliffe, M. D.: Circulation and water masses of the East Greenland shelf, *J. Geophys. Res.*, 92(C7), 6729–6740, <https://doi.org/10.1029/JC092iC07p06729>, 1987.
- 685 Boyer, T. P., Garcia, H. E., Locarnini, R. A., Ricardo, A., Zweng, M. M., Mishonov, A. V., Reagan, J. R., Weathers, K. A., Baranova, O. K., Seidov, D., and Smolyar, I. V.: World Ocean Atlas 2018, NOAA National Centers for Environmental Information, dataset, <https://accession.nodc.noaa.gov/NCEI-WOA18>, 2018.
- Budéus, G., Schneider, W., and Kattner, G.: Distribution and exchange of water masses in the Northeast Water polynya (Greenland Sea), *J. Mar. Syst.*, 10, 123–138, doi:10.1016/S0924-7963(96)00074-7, 1997.**
- 690 Cabedo-Sanz, P., Belt, S. T., Jennings, A. E., Andrews, J. T., and Geirsdóttir, Á.: Variability in drift ice export from the Arctic Ocean to the North Icelandic Shelf over the last 8000 years: a multi-proxy evaluation, *Quaternary Sci. Rev.*, 146, 99–115, 2016.
- Caesar, L., McCarthy, G. D., Thornalley, D. J. R., Cahill, N., and Rahmstorf, S.: Current Atlantic Meridional Overturning Circulation weakest in last millennium, *Nat. Geosci.*, 1–3, <https://doi.org/10.1038/s41561-021-00699-z>, 2021.
- Cage, A.G., Pieńkowski, A.J., Jennings, A., Knudsen, K.L., and Seidenkrantz, M.-S.: Comparative analysis of six common foraminiferal species of the genera *Cassidulina*, *Paracassidulina* and *Islandiella* from the Arctic-North Atlantic domain, *J. Micropalaeontol.*, in press, 2021.
- 695 Clark, P. U., Marshall, S. J., Clarke, G. K., Hostetler, S. W., Licciardi, J. M., and Teller, J. T.: Freshwater forcing of abrupt climate change during the last glaciation, *Science*, 293(5528), 283–287, 2001.
- Clark, P. U., Pisias, N. G., Stocker, T. F., and Weaver, A. J.: The role of the thermohaline circulation in abrupt climate change, *Nature*, 415(6874), 863–869, 2002.
- 700 Consolaro, C., Rasmussen, T. L., and Panieri, G.: Palaeoceanographic and environmental changes in the eastern Fram Strait during the last 14,000 years based on benthic and planktonic foraminifera, *Mar. Micropalaeontol.*, 139, 84–101, 2018.
- Cronin, T. M., Dwyer, G. S., Kamiya, T., Schwede, S., and Willard, D. A.: Medieval warm period, little ice age and 20th century temperature variability from Chesapeake Bay, *Global Planet. Change*, 36(1–2), 17–29, 2003.
- 705 Danish Meteorological Institute (DMI) and NSIDC: Arctic Sea Ice Charts from the Danish Meteorological Institute, 1893–1995, edited by: Underhill, V., Fetterer, F., National Snow and Ice Data Center, Boulder, Colorado USA, 2012.
- El bani Altuna, N., Rasmussen, T. L., Ezat, M. M., Vadakkepuliambatta, S., Groeneveld, J., and Greaves, M.: Deglacial bottom water warming intensified Arctic methane seepage in the NW Barents Sea, *Communications Earth & Environment*, 2, 188, <https://doi.org/10.1038/s43247-021-00264-x>, 2021.**
- 710 Ellison, C. R. W., Chapman, M. R., and Hall, I. R.: Surface and deep ocean interactions during the cold climate event 8200 years ago, *Science*, 312(5782), 1929–1932, <https://doi.org/10.1126/science.1127213>, 2006.
- Estrade, P., Marchesiello, P., Colin De Verdiere, A., and Roy, C.: Cross-shelf structure of coastal upwelling: A two-dimensional extension of Ekman’s theory and a mechanism for inner shelf upwelling shut down, *J. Mar. Res.*, 66, 589–616, <https://doi.org/10.1357/002224008787536790>, 2008.
- 715 **Farmer, J. R., Sigman, D. M., Granger, J., Underwood, O. M., Fripiat, F., Cronin, T. M., Martínez-García, A., and Haug, G. H.: Arctic Ocean stratification set by sea level and freshwater inputs since the last ice age, *Nat. Geosci.*, 14, 684–689, <https://doi.org/10.1038/s41561-021-00789-y>, 2021.**
- Feyling-Hanssen, R. W.: The stratigraphy of the Quaternary Clyde Foreland Formation, Baffin Island, illustrated by the distribution of benthic foraminifera, *Boreas*, 5, 77–94, 1976.

- 720 Feyling-Hanssen, R. W.: Microbiostratigraphy of young Cenozoic marine deposits of Qivituq Peninsula, Baffin Island, Mar. Micropaleontol., 5, 153–184, 1980.
- Feyling-Hanssen, R., Funder, S., and Petersen, K.S.: The Lodin Elv Formation; a Plio-Pleistocene occurrence in Greenland, B. Geol. Soc. Den., 31, 81–106, 1983.
- Funder, S., Bennike, O., Böcher, J., Israelson, C., Strand Petersen, K., and Símonarson, L. A.: Late Pliocene Greenland–The Kap
725 Kopenhagen Formation in North Greenland, B. Geol. Soc. Den., 48, 117-134, 2001.
- Giraudeau, J., Jennings, A. E., and Andrews, J. T.: Timing and mechanisms of surface and intermediate water circulation changes in the Nordic Seas over the last 10,000 cal years: a view from the North Iceland shelf, Quaternary Sci. Rev., 23(20-22), 2127-2139, 2004.
- Giraudeau, J., Grelaud, M., Solignac, S., Andrews, J. T., Moros, M., and Jansen, E.: Millennial-scale variability in Atlantic water advection to the Nordic Seas derived from Holocene coccolith concentration records, Quaternary Sci. Rev., 29(9-10), 1276-1287,
730 2010.
- Green, K. E.: Ecology of some Arctic foraminifera, Micropaleontol., 6, 57–78, 1960.
- Hald, M. and Korsun, S.: Distribution of modern benthic foraminifera from fjords of Svalbard, European Arctic, J. Foramin. Res., 27(2), 101-122, 1997.
- 735 Hald, M., Andersson, C., Ebbesen, H., Jansen, E., Klitgaard-Kristensen, D., Risebrobakken, B., Salomonsen, G. R., Sarnthein, M., Sejrup, H. P., and Telford, R. J.: Variations in temperature and extent of Atlantic Water in the northern North Atlantic during the Holocene, Quaternary Sci. Rev., 26(25-28), 3423-3440, 2007.
- Hall, I. R., Bianchi, G. G., and Evans, J. R.: Centennial to millennial scale Holocene climate-deep water linkage in the North Atlantic. Quaternary Sci. Rev., 23(14-15), 1529-1536, 2004.
- 740 Hansen, K. E., Giraudeau, J., Wacker, L., Pearce, C., and Seidenkrantz, M.-S: Reconstruction of Holocene oceanographic conditions in the Eastern Baffin Bay, Clim. Past, 16, 1075–1095, <https://doi.org/10.5194/cp-16-1075-2020>, 2020.
- Heaton, T. J., Köhler, P., Butzin, M., Bard, E., Reimer, R. W., Austin, W. E. N., Bronk Ramsey, C., Grootes, P. M., Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J., and Skinner, L. C.: Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 Cal BP), Radiocarbon, 62(4), 779–820, <https://doi.org/10.1017/RDC.2020.68>,
745 2020.
- Hillaire-Marcel, C., De Vernal, A., Bilodeau, G., and Weaver, A. J.: Absence of deep-water formation in the Labrador Sea during the last interglacial period, Nature, 410(6832), 1073-1077, 2001.
- Hillaire-Marcel, C., De Vernal, A., and Piper, D. J.: Lake Agassiz final drainage event in the northwest North Atlantic, Geophys. Res. Lett., 34(15), 2007.
- 750 Hoffman, J. S., Carlson, A. E., Winsor, K., Klinkhammer, G. P., LeGrande, A. N., Andrews, J. T., and Strasser, J. C.: Linking the 8.2 ka event and its freshwater forcing in the Labrador Sea, Geophys. Res. Lett., 39(18), 2012.
- Hurrell, J. W. and Deser, C.: North Atlantic climate variability: the role of the North Atlantic Oscillation, J. Marine Syst., 79(3-4), 231-244, 2010.
- Hurrell, J. W., Kushnir, Y., Ottersen, G., and Visbeck, M.: An overview of the North Atlantic oscillation. Geophysical Monograph-American Geophysical Union, 134, 1-36, 2003.
- 755 Ionita, M., Scholz, P., Lohmann, G., Dima, M., and Prange, M.: Linkages between atmospheric blocking, sea ice export through Fram Strait and the Atlantic meridional overturning circulation, Sci. Rep.-UK, 6, 32881, <https://doi.org/10.1038/srep32881>, 2016.

- Jansen, E., Andersson, C., Moros, M., Nisancioglu, K. H., Nyland, B. F., and Telford, R. J.: The Early to Mid-Holocene Thermal Optimum in the North Atlantic, in: Natural Climate Variability and Global Warming, edited by: Battarbee, R. W., Binney, H. A., Wiley Blackwell, 123–137, <https://doi.org/10.1002/9781444300932.ch5>, 2008.
- Jennings, A. E. and Helgadottir, G.: Foraminiferal assemblages from the fjords and shelf of eastern Greenland. *J. Foramin. Res.*, 24(2), 123-144, 1994.
- Jennings, A. E., Knudsen, K. L., Hald, M., Hansen, C. V., and Andrews, J. T.: A mid-Holocene shift in Arctic sea-ice variability on the East Greenland Shelf, *The Holocene*, 12(1), 49-58, 2002.
- Jennings, A. E., Weiner, N. J., Helgadottir, G., and Andrews, J. T.: Modern foraminiferal faunas of the southwestern to northern Iceland shelf: oceanographic and environmental controls, *J. Foramin. Res.*, 34(3), 180-207, 2004.
- Jennings, A., Andrews, J., and Wilson, L.: Holocene environmental evolution of the SE Greenland Shelf North and South of the Denmark Strait: Irminger and East Greenland current interactions, *Quaternary Sci. Rev.*, 30(7-8), 980-998, 2011.
- Jennings, A., Andrews, J., Reilly, B., Walczak, M., Jakobsson, M., Mix, A., Stoner, J., Nicholls, K. W., and Cheseby, M.: Modern foraminiferal assemblages in northern Nares Strait, Petermann Fjord, and beneath Petermann ice tongue, NW Greenland, *Arct. Antarct. Alp. Res.*, 52(1), 491-511, 2020.
- Johannessen, O. M.: Brief overview of the physical oceanography, in: The Nordic Seas, edited by: Hurdle, B.G., Springer-Verlag, New York, https://doi.org/10.1007/978-1-4615-8035-5_4, 1986.
- Kaufmann, D. S., Ager, T. A., Anderson, N. J., Anderson, P. M., Andrews, J. T., Bartlein, P. J., Brubaker, L. B., Coats, L. L., Cwynar, L. C., Duvall, M. L., Dyke, A. S., Edwards, M. E., Eisner, W. R., Gajewski, K., Geirsdóttir, A., Hu, F. S., Jennings, A. E., Kaplan, M. R., Kerwin, M. W., Lozhkin, A. V., MacDonald, G. M., Miller, G. H., Mock, C. J., Oswald, W. W., Otto-Bliesner, B. L., Porinchu, D. F., Rühland, K., Smol, J. P., Steig, E. J., and Wolfe, B. B.: Holocene thermal maximum in the western Arctic (0-180°W), *Quaternary Sci. Rev.*, 23, 529–560, 2004.
- Keigwin, L. D., Sachs, J. P., Rosenthal, Y., and Boyle, E. A.: The 8200 year B.P. event in the slope water system, western subpolar North Atlantic, *Paleoceanography*, 20, <https://dx.doi.org/10.1029/2004PA001074>, 2005.
- Kirillov, S., Dmitrenko, I., Tremblay, B., Gratton, Y., Barber, D., and Rysgaard, S.: Upwelling of Atlantic Water along the Canadian Beaufort Sea continental slope: Favorable atmospheric conditions and seasonal and interannual variations, *J. Clim.*, 29(12), 4509-4523, <https://doi.org/10.1175/JCLI-D-15-0804.1>, 2016.
- Knudsen, K. L. and Seidenkrantz, M. S.: *Stainforthia feylingi* new species from arctic to subarctic environments, previously recorded as *Stainforthia schreibersiana* (Czjzek). *Cushman Foundation for Foraminiferal Research Special Publication*, 32, 5-13, 1994.
- Knudsen, M. F., Seidenkrantz, M. S., Jacobsen, B. H., and Kuijpers, A.: Tracking the Atlantic Multidecadal Oscillation through the last 8,000 years, *Nat. Commun.*, 2(1), 1-8, 2011.
- Koç, N., Jansen, E., and Haflidason, H.: Paleoceanographic reconstructions of surface ocean conditions in the Greenland, Iceland and Norwegian seas through the last 14 ka based on diatoms, *Quaternary Sci. Rev.*, 12, 115–40, 1993.
- Kolling, H. M., Stein, R., Fahl, K., Perner, K., and Moros, M.: Short-term variability in late Holocene sea ice cover on the East Greenland Shelf and its driving mechanisms, *Palaeogeogr. Palaeoclimatol.*, 485, 336-350, 2017.
- Korsun, S. and Hald, M.: Seasonal dynamics of benthic foraminifera in a glacially fed fjord of Svalbard, European Arctic, *J. Foramin. Res.*, 30(4), 251-271, 2000.
- Kwok, R.: Recent changes in Arctic Ocean sea ice motion associated with the North Atlantic Oscillation, *Geophys. Res. Lett.*, 27(6), 775-778, 2000.
- Lagoe, M. B.: Recent Benthic foraminifera from the Central Arctic Ocean, *J. Foramin. Res.*, 7, 106–129, 1979.

- Larsen, N. K., Levy, L. B., Carlson, A. E., Buizert, C., Olsen, J., Strunk, A., Bjørk, A. A., and Skov, D. S.: Instability of the Northeast Greenland Ice Stream over the last 45,000 years, *Nat. Commun.*, 9(1), 1-8, 2018.
- 800 Lloyd, J. M.: Modern distribution of benthic foraminifera from Disko Bugt, West Greenland, *J. Foramin. Res.*, 36(4), 315-331, 2006.
- Mackensen, A. and Hald, M.: *Cassidulina teretis* Tappan and *C. laevigata* D'Orbigny: their modern and Late Quaternary distribution in Northern seas, *J. Foramin. Res.*, 18(1), 16–24, 1988.
- Mackensen, A. and Schmiedl, G.: Brine formation recorded by stable isotopes of recent benthic foraminifera in Storfjorden, Svalbard: palaeoceanographical implications, *Boreas*, 45, 552– 556, 10.1111/bor.12174. ISSN 0300-9483, 2016.**
- 805 Mayer, C., Reeh, N., Jung-Rothenhäusler, F., Huybrechts, P., and Oerter, H.: The subglacial cavity and implied dynamics under Nioghalvfjærdsfjorden Glacier, NE-Greenland, *Geophys. Res. Lett.*, 27(15), 2289-2292, 2000.
- Moossen, H., Bendle, J., Seki, O., Quillmann, U., and Kawamura, K.: North Atlantic Holocene climate evolution recorded by high-resolution terrestrial and marine biomarker records, *Quaternary Sci. Rev.*, 129, 111e127, 2015.
- 810 Moros, M., Jansen, E., Oppo, D. W., Giraudeau, J., and Kuijpers, A.: Reconstruction of the late-Holocene changes in the Sub-Arctic Front position at the Reykjanes Ridge, north Atlantic, *The Holocene*, 22(8), 877-886, 2012.
- Müller, J., Werner, K., Stein, R., Fahl, K., Moros, M., and Jansen, E.: Holocene cooling culminates in sea ice oscillations in Fram Strait, *Quaternary Sci. Rev.*, 47, 1-14, 2012.
- Oppo, D. W., McManus, J. F., and Cullen, J. L.: Deepwater variability in the Holocene epoch, *Nature*, 422(6929), 277-277, 2003.
- 815 Orme, L. C., Miettinen, A., Divine, D., Husum, K., Pearce, C., Van Nieuwenhove, N., Born, A., Mohan, R., and Seidenkrantz, M. S.: Subpolar North Atlantic sea surface temperature since 6 ka BP: Indications of anomalous ocean-atmosphere interactions at 4-2 ka BP, *Quaternary Sci. Rev.*, 194, 128-142, 2018.
- Pedersen, L.T., Gudmandsen, P., and Skriver, H.: North-East Water - A remote sensing study, Report 545, Electromagnetic Institute, Technical University of Denmark, 1993.
- 820 Perner, K., Moros, M., Lloyd, J. M., Kuijpers, A., Telford, R. J., and Harff, J.: Centennial scale benthic foraminiferal record of late Holocene oceanographic variability in Disko Bugt, West Greenland, *Quaternary Sci. Rev.*, 30(19-20), 2815-2826, 2011.
- Perner, K., Moros, M., Jennings, A., Lloyd, J. M., and Knudsen, K. L.: Holocene palaeoceanographic evolution off West Greenland, *The Holocene*, 23(3), 374-387, 2012.
- Perner, K., Moros, M., Lloyd, J. M., Jansen, E., and Stein, R.: Mid to late Holocene strengthening of the East Greenland Current linked to warm subsurface Atlantic water, *Quaternary Sci. Rev.*, 129, 296-307, 2015.
- 825 Perner, K., Jennings, A. E., Moros, M., Andrews, J. T., and Wacker, L.: Interaction between warm Atlantic-sourced waters and the East Greenland Current in northern Denmark Strait (68 N) during the last 10 600 cal a BP, *J. Quaternary Sci.*, 31(5), 472-483, 2016.
- Quadfasel, D., Gascard, J. C., and Koltermann, K. P.: Large-scale oceanography in Fram Strait during the 1984 marginal ice zone experiment, *J. Geophys. Res.-Oceans*, 92, 6719-6728, 1987.
- 830 Rahmstorf, S.: Bifurcations of the Atlantic thermohaline circulation in response to changes in the hydrological cycle, *Nature*, 378(6553), 145-149, 1995.
- Ramsey, C. B.: Deposition models for chronological records. *Quaternary Sc. Rev.*, 27(1-2), 42-60, 2008.
- Ran, L., Jiang, H., Knudsen, K. L., Eiríksson, J., and Gu, Z.: Diatom response to the Holocene climatic optimum on the North Icelandic shelf, *Mar. Micropaleontol.*, 60(3), 226-241, 2006.
- 835 Rasmussen, S. O., Vinther, B. M., Clausen, H. B., and Andersen, K. K.: Early Holocene climate oscillations recorded in three Greenland ice cores, *Quaternary Sci. Rev.*, 26, 1907–1914, <https://doi.org/10.1016/j.quascirev.2007.06.015>, 2007.

- Reimer, P. J. and Reimer, R. W.: A marine reservoir correction database and on-line interface, *Radiocarbon*, 43(2A), 461-463, 2001.
- 840 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Backwell, P. G., Ramsey, C. B., Buck, C. E., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R. A., Turney, C. S. M., and van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0-50,000 Years cal BP, *Radiocarbon*, 55(4), 1869-1887, https://doi.org/10.2458/azu_js_rc.55.16947, 2013.
- 845 Ren, J., H. Jiang, M. S. Seidenkrantz, and Kuijpers, A.: A Diatom-Based Reconstruction of Early Holocene Hydrographic and Climatic Change in a Southwest Greenland Fjord, *Mar. Micropaleontol.*, 70, 166-76, <https://doi.org/10.1016/j.marmicro.2008.12.003>, 2009.
- Renssen, H., Goosse, H., and Muscheler, R.: Coupled climate model simulation of Holocene cooling events: oceanic feedback amplifies solar forcing, *Clim. Past*, 2(2), 79-90, 2006.
- 850 Risebrobakken, B., Jansen, E., Andersson, C., Mjelde, E., and Hevrøy, K.: A high-resolution study of Holocene paleoclimatic and paleoceanographic changes in the Nordic Seas, *Paleoceanography*, 18(1), 2003.
- Risebrobakken, B., Dokken, T., Smedsrud, L. H., Andersson, C., Jansen, E., Moros, M., and Ivanova, E. V.: Early Holocene temperature variability in the Nordic Seas: The role of oceanic heat advection versus changes in orbital forcing, *Paleoceanography*, 26(4), PA4206, <https://doi:10.1029/2011PA002117>, 2011.
- 855 Rudels, B.: Arctic Ocean circulation and variability—advection and external forcing encounter constraints and local processes, *Ocean Sci.*, 8(2), 261-286, 2012.
- Rudels, B. and Quadfasel, D.: Convection and deep water formation in the Arctic Ocean-Greenland Sea system, *J. Marine Syst.*, 2(3-4), 435-450, 1991.
- Rudels, B., Björk, G., Nilsson, J., Winsor, P., Lake, I., and Nohr, C.: The interaction between waters from the Arctic Ocean and the Nordic Seas north of Fram Strait and along the East Greenland Current: results from the Arctic Ocean-02 Oden expedition, *J. Marine Syst.*, 55(1-2), 1-30, 2005.
- 860 Rytter, F., Knudsen, K. L., Seidenkrantz, M. S., and Eiríksson, J.: Modern distribution of benthic foraminifera on the North Icelandic shelf and slope, *J. Foramin. Res.*, 32(3), 217-244, 2002.
- Sachs, J.P.: Cooling of Northwest Atlantic slope waters during the Holocene, *Geophys. Res. Lett.*, 34, L03609, <http://dx.doi.org/10.1029/2006GL028495>, 2007.
- 865 Sarnthein, M., Van Kreveland, S., Erlenkeuser, H., Grootes, P. M., Kucera, M., Pflaumann, U., and Schulz, M.: Centennial-to-millennial-scale periodicities of Holocene climate and sediment injections off the western Barents shelf, 75 N, *Boreas*, 32(3), 447-461, 2003.
- Schaffer, J., von Appen, W. J., Dodd, P. A., Hofstede, C., Mayer, C., de Steur, L., and Kanzow, T.: Warm water pathways toward Nioghalvfjærdsfjorden Glacier, Northeast Greenland. *J. Geophys. Res.-Oceans*, 122(5), 4004-4020, 2017.
- 870 Schneider, W. and Budéus, G.: The northeast water polynya (Greenland Sea), *Polar Biol.*, 14(1), 1-9, 1994.
- Schneider, W. and Budéus, G.: Summary of the Northeast Water polynya formation and development (Greenland Sea), *J. Marine Syst.*, 10(1-4), 107-122, 1997.
- Seidenkrantz, M. S.: *Cassidulina teretis* Tappan and *Cassidulina neoteretis* new species (Foraminifera): stratigraphic markers for deep sea and outer shelf areas, *J. Micropalaeontol.*, 14(2), 145-157, 1995.
- 875 Seidenkrantz, M. S.: Benthic foraminifera as palaeo sea-ice indicators in the subarctic realm – examples from the Labrador Sea – Baffin Bay region, *Quaternary Sci. Rev.*, 79, 135-144; <https://doi:10.1016/j.quascirev.2013.03.014>, 2013.

- Seidenkrantz, M. S., Aagaard-Sørensen, S., Sulsbrück, H., Kuijpers, A., Jensen, K. G., and Kunzendorf, H.: Hydrography and climate of the last 4400 years in a SW Greenland fjord: implications for Labrador Sea palaeoceanography, *The Holocene*, 17(3), 387-401, 2007.
- 880 Seidenkrantz, M. S., Roncaglia, L., Fischel, A., Heilmann-Clausen, C., Kuijpers, A., and Moros, M.: Variable North Atlantic climate seesaw patterns documented by a late Holocene marine record from Disko Bugt, West Greenland, *Mar. Micropaleontol.*, 68(1-2), 66-83, 2008.
- Seidenkrantz, M. S., Ebbesen, H., Aagaard-Sørensen, S., Moros, M., Lloyd, J. M., Olsen, J., Knudsen, M. F. and Kuijpers, A.:
- 885 Early Holocene large-scale meltwater discharge from Greenland documented by foraminifera and sediment parameters, *Palaeogeogr. Palaeoclimatol.*, 391, 71-81, 2013.
- Seidenkrantz, M.-S., Kuijpers, A., Olsen, J., Pearce, C., Lindblom, S., Ploug, J., Przybyło, P., and Snowball, I.: Southwest Greenland shelf glaciation during MIS 4 more extensive than during the Last Glacial Maximum, *Sci. Rep.-UK*, 9, 15617; [https://doi: 10.1038/s41598-019-51983-3](https://doi.org/10.1038/s41598-019-51983-3), 2019.
- 890 Ślubowska, M. A., Koç, N., Rasmussen, T. L., and Klitgaard-Kristensen, D.: Changes in the flow of Atlantic water into the Arctic Ocean since the last deglaciation: Evidence from the northern Svalbard continental margin, 80°N, *Paleoceanography*, 20, <https://doi.org/10.1029/2005PA001141>, 2005.
- Ślubowska-Woldengen, M., Rasmussen, T. L., Koc, N., Klitgaard-Kristensen, D., Nilsen, F., and Solheim, A.: Advection of Atlantic Water to the western and northern Svalbard shelf since 17,500 cal yr BP. *Quaternary Sci. Rev.*, 26(3-4), 463-478, 2007.
- 895 Stroeve, J. C., Serreze, M. C., Holland, M. M., Kay, J. E., Malanik, J., and Barrett, A. P.: The Arctic's rapidly shrinking sea ice cover: a research synthesis, *Climatic Change*, 110(3), 1005-1027, 2012.
- Syring, N., Stein, R., Fahl, K., Vahlenkamp, M., Zehnick, M., Spielhagen, R. F., and Niessen, F.: Holocene changes in sea-ice cover and polynya formation along the eastern North Greenland shelf: New insights from biomarker records, *Quaternary Sci. Rev.*, 231, 106173, 2020a.
- 900 Syring, N., Lloyd, J. M., Stein, R., Fahl, K., Roberts, D. H., Callard, L., and O'Cofaigh, C.: Holocene Interactions Between Glacier Retreat, Sea Ice Formation, and Atlantic Water Advection at the Inner Northeast Greenland Continental Shelf, *Paleoceanography and Paleoclimatology*, 35(11), e2020PA004019, 2020b.
- Telesiński, M. M., Spielhagen, R. F., and Bauch, H. A.: Water mass evolution of the Greenland Sea since late glacial times, *Clim. Past*, 10(1), 123-136, 2014a.
- 905 Telesiński, M. M., Spielhagen, R. F., and Lind, E. M.: A high-resolution Lateglacial and Holocene palaeoceanographic record from the Greenland Sea, *Boreas*, 43(2), 273-285, 2014b.
- Thornalley, D. J., Elderfield, H., and McCave, I. N.: Holocene oscillations in temperature and salinity of the surface subpolar North Atlantic, *Nature*, 457(7230), 711-714, 2009.
- Van Nieuwenhove, N., Pearce, C., Knudsen, M. F., Røy, H., and Seidenkrantz, M. S.: Meltwater and seasonality influence on Subpolar Gyre circulation during the Holocene, *Palaeogeogr. Palaeoclimatol.*, 502, 104-118, 2018.
- 910 Vare, L. L., Masse, G., Gregory, T. R., Smart, C. W., and Belt, S. T.: Sea ice variations in the central Canadian Arctic Archipelago during the Holocene, *Quaternary Sci. Rev.*, 28, 1354-1366, 2009.
- Vinje, T. E.: Sea Ice Conditions in the European Sector of the Marginal Seas of the Arctic 1966-75, 1975, Norwegian Polar Institut, 163-174, 1977.
- 915 Wangner, D. J., Jennings, A. E., Vermassen, F., Dyke, L. M., Hogan, K. A., Schmidt, S., Kjær, K. H., Knudsen, M. F., and Andresen, C. S.: A 2000-year record of ocean influence on Jakobshavn Isbræ calving activity, based on marine sediment cores, *The Holocene*, 28(11), 1731-1744, 2018.

- Weatherall, P., Marks, K. M., Jakobsson, M., Schmitt, T., Tani, S., Arndt, J. E., Rovere, M., Chayes, D., Ferrini, V. and Wigley, R.: A new digital bathymetric model of the world's oceans, *Earth and Space Science*, 2(8), 331-345, 2015.
- 920 Weiss, H.: 4.2 ka BP megadrought and the Akkadian collapse, in: *Megadrought and Collapse: from early agriculture to Angkor*, edited by: Weiss, H., Oxford University Press, Oxford, 93–160, 2017.
- Werner, K., Spielhagen, R. F., Bauch, D., Hass, H. C., and Kandiano, E.: Atlantic Water advection versus sea-ice advances in the eastern Fram Strait during the last 9 ka: Multiproxy evidence for a two-phase Holocene, *Paleoceanography*, 28(2), 283-295, 2013.
- 925 Werner, K., Müller, J., Husum, K., Spielhagen, R. F., Kandiano, E. S., and Polyak, L.: Holocene sea subsurface and surface water masses in the Fram Strait—Comparisons of temperature and sea-ice reconstructions, *Quaternary Sci. Rev.*, 147, 194-209, 2016.
- Wilson, N. J. and Straneo, F.: Water exchange between the continental shelf and the cavity beneath Nioghalvfjærdsbræ (79 North Glacier), *Geophys. Res. Lett.*, 42(18), 7648-7654, 2015.
- Wollenburg, J. E. and Mackensen, A.: Living benthic foraminifera from the central Arctic Ocean: Faunal composition, standing stock and diversity, *Mar. Micropaleontol.*, 34, 153–185, 1998.
- 930 Wollenburg, J. E. and Kuhnt, W.: The response of benthic foraminifers to carbon flux and primary production in the Arctic Ocean, *Mar. Micropaleontol.*, 40(3), 189-231, 2000.
- Zehnich, M., Spielhagen, R. F., Bauch, H. A., Forwick, M., Hass, H. C., Palme, T., Stein, R., and Syring, N.: Environmental variability off NE Greenland (western Fram Strait) during the past 10,600 years, *The Holocene*, 30(12), 1752-1766, 2020.