

Pliocene evolution of the tropical Atlantic thermocline depth

5 Carolien M. H. van der Weijst¹, Josse Winkelhorst¹, Wesley de Nooijer¹, Anna von der Heydt², Gert-Jan Reichart^{1,3}, Francesca Sangiorgi¹, Appy Sluijs¹

¹Department of Earth Sciences, Utrecht University, 3584 CB Utrecht, the Netherlands

²Department of Physics, Utrecht University, 3584CC Utrecht, the Netherlands

³NIOZ Royal Netherlands Institute for Sea Research, 1797 SZ 't Horntje, the Netherlands

Correspondence to: C.M.H. van der Weijst (c.m.h.vanderweijst@uu.nl)

10

Abstract.

It has been hypothesized that global temperature trends are tightly linked to tropical thermocline depth, and that thermocline shoaling played a crucial role in the intensification of late Pliocene northern hemisphere glaciation. The Pliocene thermocline evolution in the Pacific Ocean is well documented and supports this hypothesis, but thermocline records from the tropical Atlantic Ocean are limited. We present new planktonic foraminiferal Mg/Ca, $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ records from the late Pliocene interval at Ocean Drilling Program Site 959 in the eastern equatorial Atlantic (EEA), which we use to reconstruct ocean temperatures and relative changes in salinity and thermocline depth. Data were generated using surface-dwelling *Globigerinoides ruber* and subsurface-dwelling *Neogloboquadrina dutertrei*. Reduced gradients between the surface and subsurface records indicate deepening of the EEA thermocline at the end of the mid-Piacenzian Warm Period (mPWP; ~3.3-3.0 Ma). We connect our late Pliocene records to previously published early Pliocene $\delta^{18}\text{O}$ data from Site 959 and compare these to the Site 1000 in the Caribbean Sea. Over the course of the Pliocene, thermocline changes in the EEA and Caribbean Sea follow similar patterns, with prominent step-wise thermocline deepening between ~5.5 and 4.0 Ma, gradual shoaling up to the mPWP, followed by minor deepening at the end of the mPWP. The tropical thermocline depth evolution of the tropical Atlantic differs from the Pacific, which is characterized by gradual basin-wide shoaling across the Pliocene. These results potentially challenge the hypothesized link between tropical thermocline depth and global climate. The mechanisms behind the periodically divergent Pacific and Atlantic thermocline movements remain speculative. We suggest that they are related to basin geometry and heterogenous temperature evolutions in regions from where thermocline waters are sourced. A positive feedback loop between source region temperature and tropical cyclone activity may have amplified tropical thermocline adjustments.

30

1 Introduction

The Pliocene (5.33-2.58 Ma) is the most recent geological epoch with substantially higher greenhouse gas concentrations (de la Vega et al., 2020; Martínez-Botí et al., 2015; Stap et al., 2016) and elevated global surface temperatures (Brierley et al., 2009; Dowsett et al., 2016; McClymont et al., 2020) compared to preindustrial times. This makes it an interesting interval to study for its potential analogies with our future climate (Burke et al., 2018). Typical features of the warm Pliocene ocean are the poleward expansion of the tropical warm pools and a reduction of the zonal sea surface temperature (SST) gradient in the Pacific (Brierley et al., 2009; Dekens et al., 2008; Fedorov et al., 2013). Another key feature of the Pliocene ocean is the deep tropical Pacific thermocline that gradually shoaled towards the end of the Pliocene (Dekens et al., 2007; Ford et al., 2012, 2015; Steph et al., 2010), while the meridional and zonal SST gradient steepened (Fedorov et al., 2015). It has been suggested that this shoaling reached a critical threshold around 3 Ma and played an important role in the onset of northern hemisphere glaciation (Fedorov et al., 2006; Philander and Fedorov, 2003; Steph et al., 2010). Moreover, because there is general coherence between Pliocene changes in meridional, zonal, and vertical (i.e. thermocline depth) temperature gradients in the Pacific, Fedorov et al. (2015) suggested that these temperature gradients are somehow mechanistically linked.

Interestingly, while the thermocline shoaled across the tropical Pacific, $\delta^{18}\text{O}$ and Mg/Ca records of planktic foraminifera show that the Caribbean thermocline deepened during the early Pliocene (Steph et al., 2010). This was interpreted by Steph et al. (2010) as a local phenomenon. However, early Pliocene $\delta^{18}\text{O}$ records from the Eastern Equatorial Atlantic (EEA) also show a thermocline deepening (Norris, 1998). This suggests that the entire tropical Atlantic may have had a different early Pliocene thermocline evolution than the Pacific. If so, the link between tropical thermocline depth and Pliocene global temperature trends may need to be reconsidered. Late Pliocene thermocline records from the EEA are currently unavailable, yet necessary for a complete evaluation of Pliocene tropical thermocline depth evolution in a context of global climate changes.

We present the first records of late Pliocene thermocline depth variability in the EEA. We generated $\delta^{18}\text{O}$, $\delta^{13}\text{C}$ and Mg/Ca records on surface-dwelling *Globigerinoides ruber* and subsurface-dwelling *Neogloboquadrina dutertrei* from Ocean Drilling Program (ODP) Site 959 sediments. We use the offsets between these records to infer changes in thermocline/nutricline depth and relative salinity changes. We evaluate our new records against previously published thermocline depth records from the Atlantic and Pacific to evaluate global patterns of thermocline depth changes across the Pliocene.

2 Methods

2.1 Site and modern oceanographic setting

We used sediments recovered at ODP Site 959 during Leg 159 in the Gulf of Guinea at ~160 km offshore Ghana and Ivory Coast (3.62°N, 2.73°W; 2090 m depth; Mascle et al., 1996, Figure 1). At present, this region is characterized by a shallow

thermocline compared to the west Atlantic (Figure 2). Surface ocean circulation at Site 959 is characterized by the eastward flowing geostrophic Guinea Current that originates from the North Equatorial Counter Current and the Canary Current (Figure 1). The thermal structure of the local water column varies seasonally in response to monsoon-induced changes in upwelling intensity (Figure 1b). In West Africa, the intertropical convergence zone (ITCZ) shifts from 5-10°N in February to 15-20°N in August, which forces strong southwesterly monsoonal winds over the Gulf of Guinea in boreal summer (Gu and Adler, 2004). A major upwelling event occurs along the northern coast in summer, which is likely forced by a combination of local wind-stress, wind-induced eastward propagating Kelvin waves and intensification of the Guinea Current (Djakouré et al., 2017; Verstraete, 1992), which together raise the thermocline. A shorter and weaker coastal upwelling event typically occurs in winter (Wiafe and Nyadjro, 2015). South of Site 959, upwelling occurs at the equator in response to equatorial divergence and along the African west coast due to persistent Ekman-pumping (Bakun and Nelson, 1991; Figure 1).

2.2 Thermocline depth reconstructions

The thermocline is typically defined as the depth at which the vertical temperature change is at its steepest. This depth can be approximated with the 20°C isotherm in the modern ocean, which is typically done in modelling studies, but different isotherms may be more appropriate in warmer climates (e.g. Von der Heydt and Dijkstra, 2011). A common approach for qualitative reconstructions of past thermocline depth variability is by paired tracing of surface ocean and shallow subsurface ocean conditions. We trace surface ocean conditions with *Globigerinoides ruber* and subsurface conditions with *Neogloboquadrina dutertrei*, which typically reside in the upper ocean and at thermocline depths respectively (Anand et al., 2003; Rebotim et al., 2017; Steph et al., 2009). Figure 3 conceptually shows how thermocline depth determines tracer offsets between surface- and subsurface-dwellers ($\Delta_{\text{surface-subsurface}}$). We use foraminiferal $\Delta_{\text{surface-subsurface}}$ $\delta^{18}\text{O}$, Mg/Ca, and $\delta^{13}\text{C}$ (hereafter $\Delta\delta^{18}\text{O}$, $\Delta\text{Mg/Ca}$ and $\Delta\delta^{13}\text{C}$) to trace changes in thermocline and nutricline depth, and paired $\Delta\delta^{18}\text{O}$ - $\Delta\text{Mg/Ca}$ to trace relative salinity changes.

2.3 Stable isotope analysis

Samples were taken at 5-20 cm intervals from cores 959C-5H and 959C-6H between 35.77 and 48.73 revised Meters Composite Depth (rMCD; splice by Vallé et al., 2016). The studied interval of nannofossil/foraminifer ooze (Masclé et al., 1996) was dated using oxygen isotope stratigraphy (van der Weijst et al., 2020) and spans 3.5-2.8 Ma. The preservation of planktonic foraminifera in the Pliocene of 959 is generally very good; a large proportion of the tests have a “glassy” appearance, and *G. ruber* often still has spines around its aperture (Figure S1). These features suggest that diagenetic calcite overgrowth, dissolution and recrystallisation are minimal (Edgar et al., 2015).

For each measurement, 50-60 foraminiferal tests (*G. ruber* white s.s. and *N. dutertrei*) were picked from the 250-355 μm size fraction. Cleaning of *G. ruber* was performed at Utrecht University and that of *N. dutertrei* at the Royal Netherlands Institute for Sea Research (NIOZ) following slightly different protocols to remove clay and coccoliths. After gently crushing the tests between two glass plates, *G. ruber* was ultrasonically cleaned with ethanol and rinsed with distilled water. Crushed tests of *N. dutertrei* were ultrasonically cleaned three times with methanol, two times with hot (95°C) 1% NaOH/H₂O₂ solution and rinsed with Milli-Q in between each step. Stable isotope analyses were performed at Utrecht University on a Thermo Finnigan GasBench-II carbonate preparation device coupled to a Thermo Finnigan Delta-V mass spectrometer. The international IAEA-CO-1 standard was measured to calibrate to the Vienna Pee Dee Belemnite (VPDB) and the inhouse NAXOS standard to correct for drift and track analytical precision, which was better than 0.1‰ for both carbon and oxygen isotopes.

2.4 Mg/Ca analysis and calibration

Sediment samples were wet-sieved with tap water and dried overnight at a low temperature. Per sample, between 20 and 60 specimens of the surface-dwelling *G. ruber* (white, s.s.) and the thermocline-dwelling *N. dutertrei* were picked from the 250-355 μm size fraction and cleaned following the protocol of Barker et al. (2003). Tests were gently crushed between two glass plates and ultrasonically treated with methanol (3x), hot 1% NaOH/H₂O₂ solution at 95°C (2x), 0.001M HNO₃ (1x) and rinsed with Milli-Q in between each step. Trace element analyses were performed at the Royal Netherlands Institute for Sea Research (NIOZ) on an Element-2 ICP-MS. Note that the four M2 samples (identified by peak benthic $\delta^{18}\text{O}$; Van der Weijst et al., 2020) were also used in a separate analysis for which a large quantity of carbonate was needed. For this reason, specimens of *G. ruber* from samples 959C/6H-2/102-104, 959C/6H-2/107-109, 959C/6H-2/112-114 and 959C/6H-2/117-119 were combined to a “M2 bulk sample”, and only 959C/6H-2/102-104 could also be analyzed separately. We calculate ocean temperatures using Mg/Ca values (mmol/mol).

Ambient seawater temperature is typically the dominant control on the ratio of magnesium to calcium in foraminiferal shells (e.g. Hönisch et al., 2013; Tierney and Malevich, 2019). At present, there is no standard approach to calibrate Mg/Ca to temperature, which complicates site-to-site comparisons (McClymont et al., 2020). However, there is general consensus that calibrating Mg/Ca from fossil foraminifera to paleotemperature requires corrections for preferential removal of Mg over Ca during dissolution of calcite in the water column and sediment (Dekens et al., 2002; Regenberg et al., 2006) and the effect of the changing Mg/Ca of seawater (Mg/Ca_{sw}) on geological timescales in relation to rates of oceanic crust production (Hardie, 1996). Moreover, most calibrations assume a constant partitioning coefficient (D_{Mg}) at a given temperature, i.e., a linear relationship between Mg/Ca_{sw} and calcite Mg/Ca (Mg/Ca_{calcite}), but experiments suggest that D_{Mg} changes in response to Mg/Ca_{sw} (De Nooijer et al., 2017; Evans et al., 2016; Evans and Müller, 2012; Ries, 2004). To improve calibrations, species-specific D_{Mg} sensitivities and past Mg/Ca_{sw} need better constraints.

125

Here, we apply a constant Pliocene Mg/Ca_{sw} correction to the species-specific core top calibrations from Dekens et al. (2002), in which a constant partitioning coefficient is assumed, and a dissolution correction is embedded based on species, ocean basin and core depth:

$$130 \quad \text{Mg/Ca (calcite)} = B \cdot \frac{\text{Past Mg/Ca}_{\text{sw}}}{\text{Present Mg/Ca}_{\text{sw}}} \cdot e^{A(T - C \cdot \text{core depth})} \quad (1).$$

The slope (A) is 0.09 for *G. ruber* and 0.08 for *N. dutertrei*, the intersect (B) is 0.38 for *G. ruber* and 0.60 for *N. dutertrei*, and the dissolution parameter (C) is 0.61 for *G. ruber* and 2.8 for *N. dutertrei* (Dekens et al., 2002). The present-day Mg/Ca_{sw} is 5.2 mmol/mol and we use the Pliocene estimate of 4.3 mmol/mol (Evans et al., 2016). The core depth at Site 959 is 2.09 km.

135 Note that *G. ruber* is calibrated to SST and *N. dutertrei* to 50 m depth subsurface temperature respectively (Dekens et al., 2002).

3 Results

Late Pliocene Mg/Ca SST estimates from surface dwelling *G. ruber* vary between 26.5°C and 29.0°C and Mg/Ca subsurface
140 temperatures from *N. dutertrei* between 21.0°C and 25.4°C (Figure 4). Surface δ¹⁸O varies largely between -2.18‰ and -1.31‰ and subsurface δ¹⁸O between -1.19‰ and -0.14‰. Surface δ¹⁸O and Mg/Ca barely show a long-term trend, although both record distinct cooling during the M2 glacial and peak temperatures during the KM3 interglacial. Subsurface δ¹⁸O and Mg/Ca both show a warming trend following the M2 glacial, although it is more gradual in Mg/Ca-SubST than in δ¹⁸O-SubST. Surface and subsurface δ¹³C values vary between 0.91-2.13‰ and 0.56-1.45‰ respectively. Both show an increasing trend in
145 the older part of the interval, and abruptly drop around KM3. In the younger part of the interval, surface δ¹³C remains stable, while subsurface δ¹³C increases.

The Δ(surface-subsurface) Mg/Ca, δ¹⁸O and δ¹³C records (Figure 4) follow a similar trend towards smaller offsets. Both ΔMg/Ca and Δδ¹⁸O signal a vertical temperature gradient decrease from ~5.0 to 3.5°C. The Δδ¹⁸O record decreases after the
150 M2 glacial, thereby leading the ΔMg/Ca and Δδ¹³C decrease around the KM3 interglacial by ~150 kyr. Moreover, Δδ¹⁸O stabilizes in the younger part of the interval, while ΔMg/Ca and Δδ¹³C continue to decrease.

4 Discussion

4.1 Late Pliocene thermocline, nutricline and salinity changes in the Eastern Equatorial Atlantic

155 The decreasing surface-subsurface Mg/Ca and $\delta^{18}\text{O}$ gradients between 3.3 Ma and 2.8 Ma at Site 959 reflect thermocline
deepening in the eastern equatorial Atlantic (Figure 4). These decreasing vertical gradients are driven by a gradual warming
trend in the subsurface ocean: *G. ruber* surface ocean Mg/Ca and $\delta^{18}\text{O}$ records remain stable across the interval, whereas the
N. dutertrei subsurface records show an increase in Mg/Ca and decrease in $\delta^{18}\text{O}$ following the M2 glacial. Crude $\delta^{18}\text{O}$ -based
temperature estimates calculated with the linear Shackleton (1974) calibration have similar values and amplitude of variation
as the Mg/Ca estimates, but the decreasing $\Delta\delta^{18}\text{O}$ trend leads decreasing $\Delta\text{Mg/Ca}$ (Figure 4).

160

Besides analytical and calibration errors, several confounding factors limit the use of $\Delta\delta^{18}\text{O}$ and $\Delta\text{Mg/Ca}$ as quantitative
proxies for thermocline depth, including: vertical migration of foraminifera in the water column (e.g. Steph et al., 2009),
variability in the mass and/or preservation of the high- $\delta^{18}\text{O}$ and low-Mg/Ca secondary calcite crust that *N. dutertrei* precipitates
at the end of its lifecycle (Jonkers et al., 2012; Steinhardt et al., 2015), seawater carbon chemistry (Spero et al., 1997) and light
165 intensity (Spero and Michael, 1987). Moreover, a discrepancy is observed in depth habitat ranges of planktonic species based
on tows versus those reconstructed based on their $\delta^{18}\text{O}$ values in core tops (Rebotim et al., 2017). Crucially, however, $\Delta\delta^{18}\text{O}$
and $\Delta\text{Mg/Ca}$ perform well in predicting larger thermocline patterns in the tropical Atlantic (Anand et al., 2003; Steph et al.,
2009).

170 In addition to ambient temperature, calcite $\delta^{18}\text{O}$ is determined by local seawater $\delta^{18}\text{O}$ (δw), which is in turn tightly linked to
salinity (Bigg and Rohling, 2000). Recently, Dämmer et al. (2020) advised against using coupled Mg/Ca- $\delta^{18}\text{O}$ data for salinity
reconstructions, because cumulative uncertainties from aforementioned confounding factors are too large to reliably
reconstruct measured δw . We are therefore reluctant to calculate absolute salinities with our Pliocene data. However, these
uncertainties are mostly relevant when applying this method quantitatively to discrete samples. To reduce these uncertainties,
175 we use a semi-quantitative approach to assess regional long-term salinity trends by calculating the relative vertical salinity
gradient ($\Delta\text{salinity}$) from the discrepancy between $\Delta\text{Mg/Ca}$ and $\Delta\delta^{18}\text{O}$ (Figure 5). We assume that $\Delta\text{Mg/Ca}$ best approaches
 $\Delta\text{temperature}$, despite the salinity effect on Mg/Ca (Gray et al., 2018; Hönisch et al., 2013), and that the residual $\Delta\delta^{18}\text{O}$ is
explained by salinity changes. Note that the intersects of the $\Delta\text{Mg/Ca}$ and $\Delta\delta^{18}\text{O}$ temperature reconstructions in Figure 5 may
not correspond to a vertical salinity gradient of 0 due to uncertainties in both the $\delta^{18}\text{O}$ -paleotemperature and the Mg/Ca
180 calibrations. Rather, the intercepts indicate an unknown baseline $\Delta\text{salinity}$ to which the rest of the record can be compared.

The magnitude of variability in the $\Delta\text{salinity}$ record is strongly affected by the assumption of the δw -salinity slope (Figure 5).
In the modern tropical Pacific, the δw -S slope varies with a factor >3 (0.09-0.32‰/salinity unit) between across the equatorial
Pacific, because the precipitation-forced isotopic “amount effect” (lower δw with higher tropical precipitation rates), is stronger

185 in the western than eastern Pacific (Conroy et al., 2017). In the tropical Atlantic, however, slopes of 0.11‰/salinity unit (Craig
and Gordon, 1965) and 0.15‰/salinity unit (LeGrande and Schmidt, 2006) have been reported for the tropical Atlantic region,
but the zonal variability of the δw -S relationship is poorly documented. Using the slope provided by Craig and Gordon (1965;
0.11‰/salinity unit), we calculate that Δ salinity was reduced by ~ 2 salinity units during the mid-Piacenzian Warm Period
(mPWP; 3.264-3.025 Ma) compared to the bordering intervals (Figure 5). At present, the vertical salinity gradient is ~ 1 in the
190 Gulf of Guinea (Figure 1b), suggesting that the Pliocene baseline salinity gradient was stronger than at present and/or that the
 δw -S slope was steeper. A doubling of the assumed tropical Atlantic δw -S slope (0.22‰/salinity unit) is needed to solve the
 Δ salinity peak to the modern gradient of ~ 1 . It falls within range of modern tropical Pacific slopes and may be realistic for the
interglacial Pliocene Atlantic, which was characterized by intensified monsoonal precipitation over West Africa (Haywood et
al., 2020).

195

In the absence of major circulation changes, the $\Delta\delta^{13}\text{C}$ record mainly reflects changes in the biological carbon pump and
nutricline depth, which in the modern tropical ocean is tightly coupled to thermocline depth (Wilson and Adamec, 2002). We
here consider coupled thermocline and nutricline deepening to be the most likely explanation for the observed synchronous
changes in $\Delta\text{Mg}/\text{Ca}$ and $\Delta\delta^{13}\text{C}$ around KM3. The relative lead of the $\Delta\delta^{18}\text{O}$ decrease may explained by subsurface freshening
200 (i.e., deepening of the halocline, Figure 2b) in response to increased precipitation rates during the mPWP. This implies that
 $\Delta\text{Mg}/\text{Ca}$ may outperform $\Delta\delta^{18}\text{O}$ as a proxy for thermocline depth, but salinity changes would have to be strong in order to
fully obscure the thermocline depth signal in the $\Delta\delta^{18}\text{O}$ record. Despite temporarily increased precipitation during the mPWP,
the net Δ temperature estimates from $\Delta\delta^{18}\text{O}$ and $\Delta\text{Mg}/\text{Ca}$ are equal between 3.5 and 2.8 Ma.

205 **4.2 Dissimilar evolutions of the tropical Atlantic and Pacific thermocline**

Our results show that the eastern equatorial Atlantic thermocline and nutricline deepened following the warmest mPWP
interglacials, while SST and deep ocean records show global cooling (Herbert et al., 2016; Lisiecki and Raymo, 2005). Late
Pliocene thermocline and nutricline changes are similar at Site 959 in the eastern equatorial Atlantic and Site 1000 in the
western tropical Atlantic (Figure 5), indicating that tropical thermocline deepening occurred across the entire basin. To further
210 explore global patterns of Pliocene tropical thermocline movements, we connect our new Site 959 $\delta^{18}\text{O}$ data to the early
Pliocene records of Norris (1998) in Figure 6. Across the Pliocene, thermocline depth evolution was similar in the eastern
equatorial Atlantic and Caribbean Sea (Figure 6). The late Pliocene $\sim 0.4\%$ $\Delta\delta^{18}\text{O}$ decrease at Site 959 and Site 1000 is
relatively small compared to the fluctuations in the latest Miocene and early Pliocene (Figure 6). Both sites show a net $\sim 1.5\%$
 $\Delta\delta^{18}\text{O}$ decrease between 5.5 and 4.0 Ma, indicating significant thermocline deepening. This is followed by an interval of
215 moderate thermocline shoaling, before deepening again in the late Pliocene. With the exception of the late Pliocene interval at
Site 1241 (Figure 6), tropical Pacific proxy records reflect overall Pliocene thermocline shoaling (Ford et al., 2012, 2015;
LaRiviere et al., 2012; Steph et al., 2006b, 2010). Although changes in $\Delta\delta^{18}\text{O}$ may not always be proportional to vertical

movement of the thermocline (see Section 4.1), the early Pliocene fluctuations in the Atlantic records are appreciably larger than the long-term shifts towards lower $\Delta\delta^{18}\text{O}$ values in the Pacific (Figure 6). This suggests that the tropical Atlantic thermocline underwent major changes during the early Pliocene, which were not paralleled in magnitude by late Pliocene changes in the Atlantic or by any Pliocene thermocline change in the Pacific.

It was suggested by Fedorov et al. (2015) that global temperature trends are tightly linked to tropical thermocline depth and that thermocline shoaling played a crucial role in the onset of Northern Hemisphere glaciations around 3 Ma Fedorov et al., 2006). Heat that is gained in the tropics must be balanced by heat loss at high latitudes, and because more heat can be gained by an ocean with a shallow thermocline, tropical thermocline shoaling should lead to surface ocean cooling at high latitudes (Boccaletti et al., 2004; Fedorov et al., 2006; Philander and Fedorov, 2003). While this theory is supported by the majority of the Pacific thermocline records (Ford et al., 2012, 2015; LaRiviere et al., 2012; Steph et al., 2006b, 2010), it appears to be inconsistent with basin-wide deepening of the tropical Atlantic thermocline (Figure 6). The link between ocean SST gradients, tropical thermocline depth, and ocean heat transport may be different in the Atlantic and Pacific oceans, because the asymmetric geometry of the Atlantic basin leads to a different pattern of tropical thermocline ventilation (Harper, 2000). Also, Atlantic Meridional Overturning Circulation (AMOC) leads to northward ocean heat transport in both hemispheres, in contrast to the Pacific Ocean (Forget and Ferreira, 2019). However, because the tropical Pacific is larger than the tropical Atlantic, Pacific thermocline shoaling may have been sufficient to balance heat loss at high latitudes, even if partly counteracted by Atlantic thermocline deepening.

4.3 Controls on Pliocene thermocline changes

4.3.1 Central American Seaway closure

It has been hypothesized that Central American Seaway (CAS) closure played a major role in the global tropical thermocline depth (Steph et al., 2010; Zhang et al., 2012). In this scenario, salt transport to the North Atlantic increased as a consequence of reduced inflow of relatively fresh Pacific surface waters, which in model simulations promotes the production of NADW (Lunt et al., 2007; Sepulchre et al., 2014; Steph et al., 2010; Zhang et al., 2012). Steph et al. (2010) reasoned that an increased volume of NADW was associated with a greater volume of the “cold water sphere”, which raised the tropical thermocline everywhere except for the Caribbean region. There, reduced inflow of cold Pacific waters caused the thermocline to deepen locally. However, our study shows that tropical thermocline deepening occurred across the basin (Figure 6). Even if CAS closure had promoted NADW production and/or AMOC strength, it is not clear how this, in turn, would have affected tropical thermocline depth, as both negative (Lopes dos Santos et al., 2010) and positive (Venancio et al., 2018) relationships between thermocline depth and AMOC strength have been inferred from proxy data. Moreover, it is unclear if CAS closure could have had an opposite effect on tropical Atlantic and Pacific thermocline depths, as this conflicts with modeling studies (Steph et al.,

250 2010). A dominant role of CAS closure in shaping the tropical thermocline evolution in the Atlantic and Pacific basins is not consistently supported by data and models. Future work could be aimed at modelling the spatial extent of the region where the inflow of cold Pacific waters is reduced in response to CAS closure. Moreover, it is vital to constrain the chronology of CAS closure (Montes et al., 2015; O’Dea et al., 2016) and determine during which intervals this process might have had extensive and long-lasting oceanographic effects (Molnar, 2008; Sepulchre et al., 2014).

255 **4.3.2 Source region SSTs and tropical cyclones**

Tropical thermocline waters are sourced from mid-latitude surface waters (Harper, 2000). It has therefore been suggested that extratropical cooling contributed to the Pliocene shoaling of the tropical Pacific thermocline (Ford et al., 2012, 2015). During the early Pliocene, midlatitude SSTs were indeed generally higher than preindustrial (Brierley et al., 2009). However, Pliocene midlatitude temperature evolutions were regionally variable (Herbert et al., 2016; Karas et al., 2017), with a notable asymmetry
260 between the northern and southern hemisphere (Pontes et al., 2020). At present, Pacific thermocline waters are sourced from mid-latitude surface waters in both the northern and southern hemisphere, whereas in the Atlantic, the thermocline is predominantly sourced from the southern hemisphere as a consequence of the asymmetric basin geometry (Harper, 2000). Mid-latitude SST records from potential Pacific tropical thermocline source regions register predominantly cooling between 5.5 and 2.8 Ma. In contrast, South Atlantic Site 1088 registers warming during the latest Miocene/early Pliocene, and during
265 the mPWP (Figure 7), in tandem with tropical Atlantic thermocline deepening. However, South Atlantic cooling between ~4.5 and 3.5 Ma (Figure 7) is not mirrored by thermocline shoaling, suggesting that source region SST was not likely the only driver of tropical Atlantic thermocline depth.

Tropical thermocline depth is also affected by cyclone activity, which, in turn, is linked to the latitudinal SST gradient in a
270 positive feedback mechanism (Fedorov et al., 2010). In other words, source region SSTs are potentially relevant to tropical thermocline dynamics in terms of both stand-alone trends, and in the context of distant SST trends. Tropical cyclones force vertical mixing in the upper ocean, which deepens the tropical thermocline (Buetti et al., 2014; Jansen et al., 2010). Tropical cyclone activity was higher in the early Pliocene than at present, especially in the Pacific (Fedorov et al., 2010). Gradually increasing zonal and meridional SST gradients (Fedorov et al., 2015) would have promoted stronger Walker and Hadley
275 circulation, thereby reducing tropical cyclone activity and raising the tropical thermocline (Brierley et al., 2009; Fedorov et al., 2010). The Pacific underwent a stronger reduction in tropical cyclone activity than the Atlantic (Fedorov et al., 2010), which may be linked to the contrasting evolution of the Pacific and Atlantic tropical thermocline depth. This feedback mechanism between midlatitude temperatures and tropical cyclone activity could further be explored in relationship to tropical thermocline depth. The hypothesized link should be tested in model simulations, and could additionally be supported or refuted
280 by better proxy records from mid-latitudes.

5 Conclusions

Our new Mg/Ca, $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ records from Site 959 indicate late Pliocene thermocline and nutricline deepening in the eastern equatorial Atlantic, starting at the end of the mPWP. A temporal discrepancy between $\Delta\delta^{18}\text{O}$ and $\Delta\text{Mg/Ca}$ can be explained
285 by a transient reduction of the vertical salinity gradient in response to intensification of west African monsoonal precipitation during the mPWP. Tropical thermocline deepening occurred across the Atlantic basin, as is indicated by a nearly identical $\Delta\delta^{18}\text{O}$ record at Site 1000 in the Caribbean Sea. This seems inconsistent with the hypothesis that Pliocene global cooling was linked to tropical thermocline shoaling, as observed in the Pacific, although Atlantic thermocline adjustments may have had less effect on the global heat budget because the Atlantic basin is smaller. The mechanisms driving Pliocene thermocline
290 changes are currently speculative. In climate models, Central American Seaway closure leads to global thermocline shoaling, but during most of the early and late Pliocene, opposite changes occurred in the tropical Atlantic and Pacific thermoclines. We therefore suggest to further explore alternative mechanisms, such as source water temperature forcing in the Southern Ocean in combination with tropical cyclone activity. Due to basin morphology, tropical Atlantic thermocline waters are sourced from the southern hemisphere, whereas both hemispheres feed the tropical thermocline waters in the Pacific. Asymmetric northern
295 and southern hemisphere temperature trends may therefore have contributed to the differences in Atlantic and Pacific thermocline depth changes. A positive feedback loop between meridional SST gradients and tropical cyclones may have amplified vertical thermocline movements.

Data availability

New Site 959 data are available as a supplement to this paper and will be uploaded to the PANGAEA online data repository
300 upon publication.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We thank the International Ocean Discovery Program and the predecessors for samples and data, Arnold van Dijk (UU), Cindy
305 Remijnse-Schrader (UU), Wim Boer (NIOZ) and Geert-Jan Brummer (NIOZ) for technical support and advice. We are also grateful to editor Luc Beaufort (Cerege) and two anonymous reviewers for constructive criticism on an earlier draft of this work. This work was carried out under the program of the Netherlands Earth System Science Centre (NESSC), financially supported by the Ministry of Education, Culture and Science (OCW). AS thanks the European Research Council for Consolidator grant 771497.

310 References

- Anand, P., Elderfield, H. and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series, *Paleoceanography*, 18(2), doi:10.1029/2002PA000846, 2003.
- Bakun, A. and Nelson, C. S.: The Seasonal Cycle of Wind-Stress Curl in Subtropical Eastern Boundary Current Regions, *J. Phys. Oceanogr.*, 21(12), 1815–1834, doi:10.1175/1520-0485(1991)021<1815:TSCOWS>2.0.CO;2, 1991.
- 315 Barker, S., Greaves, M. and Elderfield, H.: A study of cleaning procedures used for foraminiferal Mg/Ca paleothermometry, *Geochemistry, Geophys. Geosystems*, 4(9), 1–20, doi:10.1029/2003GC000559, 2003.
- Bigg, G. R. and Rohling, E. J.: An oxygen isotope data set for marine waters, *J. Geophys. Res. Ocean.*, 105(C4), 8527–8535, doi:10.1029/2000JC900005, 2000.
- Boccaletti, G., Pacanowski, R. C., Philander, S. G. H. and Fedorov, A. V.: The thermal structure of the Upper Ocean, *J. Phys. Oceanogr.*, 34(4), 888–902, doi:10.1175/1520-0485(2004)034<0888:TTSOTU>2.0.CO;2, 2004.
- 320 Brierley, C. M., Fedorov, A. V., Liu, Z., Herbert, T. D., Lawrence, K. T. and Lariviere, J. P.: Greatly expanded tropical warm pool and weakened Hadley circulation in the early Pliocene., *Science*, 323(5922), 1714–1718, doi:10.1126/science.1167625, 2009.
- Bueti, M. R., Ginis, I., Rothstein, L. M. and Griffies, S. M.: Tropical cyclone-induced thermocline warming and its regional and global impacts, *J. Clim.*, 27(18), 6978–6999, doi:10.1175/JCLI-D-14-00152.1, 2014.
- Burke, K. D., Williams, J. W., Chandler, M. A., Haywood, A. M., Lunt, D. J. and Otto-Bliesner, B. L.: Pliocene and Eocene provide best analogs for near-future climates, *Proc. Natl. Acad. Sci. U. S. A.*, 115(52), 13288–13293, doi:10.1073/pnas.1809600115, 2018.
- Conroy, J. L., Thompson, D. M., Cobb, K. M., Noone, D., Rea, S. and LeGrande, A. N.: Spatiotemporal variability in the $\delta^{18}\text{O}$ -salinity relationship of seawater across the tropical Pacific Ocean, *Paleoceanography*, doi:10.1002/2016PA003073, 2017.
- 330 Craig, H. and Gordon, L. I.: Deuterium and oxygen-18 variations in the ocean and marine atmosphere, in *Stable isotopes in oceanographic studies and paleotemperatures*, edited by E. Tongiorgi, pp. 9–130, Proceedings, Spoleto, Italy, Consiglio Nazionale delle Ricerche, Lab. de Geologia Nucleare, Pisa, Italy., 1965.
- 335 Dämmer, L. K., de Nooijer, L., van Sebille, E., Haak, J. G. and Reichert, G.-J.: Evaluation of oxygen isotopes and trace elements in planktonic foraminifera from the Mediterranean Sea as recorders of seawater oxygen isotopes and salinity, *Clim. Past*, 16(6), 2401–2414, doi:10.5194/cp-16-2401-2020, 2020.
- Dekens, P. S., Lea, D. W., Pak, D. K. and Spero, H. J.: Core top calibration of Mg/Ca in tropical foraminifera: Refining paleotemperature estimation, *Geochemistry, Geophys. Geosystems*, 3(4), 1–29, doi:10.1029/2001GC000200, 2002.
- 340 Dekens, P. S., Ravelo, A. C. and McCarthy, M. D.: Warm upwelling regions in the Pliocene warm period, *Paleoceanography*, 22(August), 1–12, doi:10.1029/2006PA001394, 2007.
- Dekens, P. S., Ravelo, A. C., McCarthy, M. D. and Edwards, C. A.: A 5 million year comparison of Mg/Ca and alkenone paleothermometers, *Geochemistry, Geophys. Geosystems*, 9(10), doi:10.1029/2007GC001931, 2008.
- 345 De Nooijer, L. J., Van Dijk, I., Toyofuku, T., & Reichert, G. J.: The impacts of seawater Mg/Ca and temperature on element incorporation in benthic foraminiferal calcite, *Geochemistry, Geophys. Geosystems*, 18(10), 3617–3630, doi:10.1002/2017GC007183


- Djakouré, S., Penven, P., Bourlès, B., Koné, V. and Veitch, J.: Respective roles of the Guinea current and local winds on the coastal upwelling in the northern Gulf of Guinea, *J. Phys. Oceanogr.*, 47(6), 1367–1387, doi:10.1175/JPO-D-16-0126.1, 2017.
- 350 Dowsett, H. J., Dolan, A. M., Rowley, D., Pound, M., Salzmann, U., Robinson, M. M., Chandler, M., Foley, K. and Haywood, A. M.: The PRISM4 (mid-Piacenzian) palaeoenvironmental reconstruction, *Clim. Past Discuss.*, 4(March), 1–39, doi:10.5194/cp-2016-33, 2016.
- Edgar, K. M., Anagnostou, E., Pearson, P. N. and Foster, G. L.: Assessing the impact of diagenesis on $\delta^{11}\text{B}$, $\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Sr/Ca and B/Ca values in fossil planktic foraminiferal calcite, *Geochim. Cosmochim. Acta*, 166, 189–209, doi:10.1016/j.gca.2015.06.018, 2015.
- 355 Evans, D., & Müller, W.: Deep time foraminifera Mg/Ca paleothermometry: Nonlinear correction for secular change in seawater Mg/Ca. *Paleoceanography*, 27(4), doi:10.1029/2012PA002315, 2012.
- Evans, D., Brierley, C. M., Raymo, M. E., Erez, J. and Müller, W.: Planktic foraminifera shell chemistry response to seawater chemistry: Pliocene–Pleistocene seawater Mg/Ca, temperature and sea level change, *Earth Planet. Sci. Lett.*, 438, 360 139–148, doi:10.1016/j.epsl.2016.01.013, 2016.
- Fedorov, A. V., Dekens, P. S., McCarthy, M., Ravelo, A. C., DeMenocal, P. B., Barreiro, M., Pacanowski, R. C. and Philander, S. G.: The Pliocene Paradox (Mechanisms for a Permanent El Niño), *Science* (80-.), 312(5779), 1485–1489, doi:10.1126/science.1122666, 2006.
- Fedorov, A. V., Brierley, C. M. and Emanuel, K.: Tropical cyclones and permanent El Niño in the early Pliocene epoch., 365 *Nature*, 463(7284), 1066–1070, doi:10.1038/nature08831, 2010.
- Fedorov, A. V., Brierley, C. M., Lawrence, K. T., Liu, Z., Dekens, P. S. and Ravelo, A. C.: Patterns and mechanisms of early Pliocene warmth., *Nature*, 496(7443), 43–9, doi:10.1038/nature12003, 2013.
- Fedorov, A. V., Burls, N. J., Lawrence, K. T. and Peterson, L. C.: Tightly linked zonal and meridional sea surface temperature gradients over the past five million years, *Nat. Geosci.*, (November), doi:10.1038/ngeo2577, 2015.
- 370 Ford, H. L., Ravelo, A. C. and Hovan, S.: A deep Eastern Equatorial Pacific thermocline during the early Pliocene warm period, *Earth Planet. Sci. Lett.*, 355–356, 152–161, doi:10.1016/j.epsl.2012.08.027, 2012.
- Ford, H. L., Ravelo, A. C., Dekens, P. S., Lariviere, J. P. and Wara, M. W.: The evolution of the equatorial thermocline and the early Pliocene El Padre mean state, *Geophys. Res. Lett.*, 42, 4878–4887, doi:10.1002/2015GL064215, 2015.
- Forget, G. and Ferreira, D.: Global ocean heat transport dominated by heat export from the tropical Pacific, *Nat. Geosci.*, 375 12(5), 351–354, doi:10.1038/s41561-019-0333-7, 2019.
- Gray, W. R., Rae, J. W. B., Wills, R. C. J., Shevenell, A. E., Taylor, B. J., Burke, A., Foster, G.L., and Lear, C. H.: Deglacial planktic foraminiferal boron isotope and Mg/Ca data from sediment core MD01-2416 in the western North Pacific Ocean, *Nat. Geosci.*, 11, 340–344, doi:10.1594/PANGAEA.887381. 2018.
- Gu, G. and Adler, R. F.: Seasonal evolution and variability associated with the West African monsoon system, *J. Clim.*, 380 17(17), 3364–3377, doi:10.1175/1520-0442(2004)017<3364:SEAVAW>2.0.CO;2, 2004.
- Hardie, L. A.: Secular variation in seawater chemistry: An explanation for the coupled secular variation in the mineralogies of marine limestones and potash evaporites over the past 600 my. *Geology*, 24(3), 279–283, doi: 10.1130/0091-7613(1996)024<0279:SVISCA>2.3.CO;2, 1996.

- Harper, S.: Thermocline ventilation and pathways of tropical-subtropical water mass exchange, *Tellus, Ser. A Dyn. Meteorol. Oceanogr.*, 52(3), 330–345, doi:10.3402/tellusa.v52i3.12269, 2000.
- Haywood, A. M., Tindall, J. C., Dowsett, H. J., Dolan, A. M., Foley, K. M., Hunter, S. J., Hill, D. J., Chan, W. Le, Abe-Ouchi, A., Stepanek, C., Lohmann, G., Chandan, D., Richard Peltier, W., Tan, N., Contoux, C., Ramstein, G., Li, X., Zhang, Z., Guo, C., Nisancioglu, K. H., Zhang, Q., Li, Q., Kamae, Y., Chandler, M. A., Sohl, L. E., Otto-Bliesner, B. L., Feng, R., Brady, E. C., Von Der Heydt, A. S., Baatsen, M. L. J. and Lunt, D. J.: The Pliocene Model Intercomparison Project Phase 2: Large-scale climate features and climate sensitivity, *Clim. Past*, 16(6), 2095–2123, doi:10.5194/cp-16-2095-2020, 2020.
- Herbert, T. D., Lawrence, K. T., Tzanova, A., Peterson, L. C., Caballero-Gill, R. and Kelly, C. S.: Late Miocene global cooling and the rise of modern ecosystems, *Nat. Geosci.*, 9(September), doi:10.1038/ngeo2813, 2016.
- von der Heydt, A. S. and Dijkstra, H. A.: The impact of ocean gateways on ENSO variability in the Miocene, *Geol. Soc. Spec. Publ.*, 355, 305–318, doi:10.1144/SP355.15, 2011.
- Hammer, Ø., Harper, D. A., and Ryan, P. D. PAST: Paleontological statistics software package for education and data analysis, *Palaeontologia electronica*, 4(1), 9, 2001.
- Hönisch, B., Allen, K. A., Lea, D. W., Spero, H. J., Eggins, S. M., Arbuszewski, J., DeMenocal, P., Rosenthal, Y., Russell, A. D. and Elderfield, H.: The influence of salinity on Mg/Ca in planktic foraminifers – Evidence from cultures, core-top sediments and complementary $\delta^{18}\text{O}$, *Geochim. Cosmochim. Acta*, 121, 196–213, doi:10.1016/j.gca.2013.07.028, 2013.
- Jansen, M. F., Ferrari, R. and Mooring, T. A.: Seasonal versus permanent thermocline warming by tropical cyclones, *Geophys. Res. Lett.*, 37(3), doi:10.1029/2009gl041808, 2010.
- Jonkers, L., De Nooijer, L. J., Reichart, G. J., Zahn, R. and Brummer, G. J. A.: Encrustation and trace element composition of *Neogloboquadrina dutertrei* assessed from single chamber analyses implications for paleotemperature estimates, *Biogeosciences*, doi:10.5194/bg-9-4851-2012, 2012.
- Karas, C., Nürnberg, D., Bahr, A., Groeneveld, J., Herrle, J. O., Tiedemann, R. and DeMenocal, P. B.: Pliocene oceanic seaways and global climate, *Sci. Rep.*, 7, 39842, doi:10.1038/srep39842, 2017.
- de la Vega, E., Chalk, T. B., Wilson, P. A., Bysani, R. P. and Foster, G. L.: Atmospheric CO₂ during the Mid-Piacenzian Warm Period and the M2 glaciation, *Sci. Rep.*, 10(1), 11002, doi:10.1038/s41598-020-67154-8, 2020.
- LeGrande, A. N. and Schmidt, G. A.: Global gridded data set of the oxygen isotopic composition in seawater, *Geophys. Res. Lett.*, 33(12), 1–5, doi:10.1029/2006GL026011, 2006.
- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20(1), doi:10.1029/2004PA001071, 2005.
- Liu, C., Liang, X., Ponte, R. M., Vinogradova, N. and Wang, O.: Vertical redistribution of salt and layered changes in global ocean salinity, *Nat. Commun.*, 10(1), doi:10.1038/s41467-019-11436-x, 2019.
- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: World Ocean Atlas 2013, Volume 1: Temperature, edited by: Levitus, S., NOAA Atlas NESDIS, 73, 40 pp., 2013.
- Lopes dos Santos, R. A., Prange, M., Castañeda, I. S., Schefuß, E., Mulitza, S., Schulz, M., Niedermeyer, E. M., Sinninghe Damsté, J. S. and Schouten, S.: Glacial-interglacial variability in Atlantic meridional overturning circulation and thermocline adjustments in the tropical North Atlantic, *Earth Planet. Sci. Lett.*, 300(3–4), 407–414, doi:10.1016/j.epsl.2010.10.030, 2010.



- Lunt, D. J., Valdes, P. J., Haywood, A. M. and Rutt, I. C.: Closure of the Panama Seaway during the Pliocene: implications for climate and Northern Hemisphere glaciation, *Clim. Dyn.*, 30(1), 1–18, doi:10.1007/s00382-007-0265-6, 2007.
- 425 Martínez-Botí, M. a., Foster, G. L., Chalk, T. B., Rohling, E. J., Sexton, P. F., Lunt, D. J., Pancost, R. D., Badger, M. P. S. and Schmidt, D. N.: Plio-Pleistocene climate sensitivity evaluated using high-resolution CO₂ records, *Nature*, 518(7537), 49–54, doi:10.1038/nature14145, 2015.
- Masclé, J., Lohmann, G., Clift, P. D. and the shipboard scientific party: Proceedings of the Ocean Drilling Program, Initial Reports 159, Proc. Ocean Drill. Program, Initial Reports, 159(College Station, TX, Ocean Drilling Program), 1996.
- 430 McClymont, E. L., Ford, H. L., Ho, S. L., Tindall, J. C., Haywood, A. M., Alonso-Garcia, M., Bailey, I., Berke, M. A., Littler, K., Patterson, M. O., Petrick, B., Peterse, F., Ravelo, A. C., Risebrobakken, B., De Schepper, S., Swann, G. E. A., Thirumalai, K., Tierney, J. E., van der Weijst, C., White, S., Abe-Ouchi, A., Baatsen, M. L. J., Brady, E. C., Chan, W.-L., Chandan, D., Feng, R., Guo, C., von der Heydt, A. S., Hunter, S., Li, X., Lohmann, G., Nisancioglu, K. H., Otto-Bliesner, B. L., Peltier, W. R., Stepanek, C. and Zhang, Z.: Lessons from a high-CO₂ world: an ocean view from ~ 3 million years ago, *Clim. Past*, 16(4), 1599–1615, doi:10.5194/cp-16-1599-2020, 2020.
- 435 Molnar, P.: Closing of the Central American Seaway and the ice age: A critical review, *Paleoceanography*, 23(2), 1–15, doi:10.1029/2007PA001574, 2008.
- Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J. C., Valencia, V., Ayala, C., Pérez-Angel, L. C., Rodriguez-Parra, L. A., Ramirez, V. and Niño, H.: Middle Miocene closure of the Central American Seaway, *Science* (80-.), 348(6231), 226–229, doi:10.1126/science.aaa2815, 2015.
- 440 Norris, R.D.: Miocene-Pliocene surface-water hydrography of the eastern equatorial Atlantic. In: Masclé, J., Lohmann, G.P., Moullade, M. (eds.), Proceedings of the Ocean Drilling Program, Scientific Results Vol. 159, College Station, TX (Ocean Drilling Program), 539-555, 1998.
- 445 O’Dea, A., Lessios, H. A., Coates, A. G., Eytan, R. I., Restrepo-Moreno, S. A., Cione, A. L., Collins, L. S., De Queiroz, A., Farris, D. W., Norris, R. D., Stallard, R. F., Woodburne, M. O., Aguilera, O., Aubry, M. P., Berggren, W. A., Budd, A. F., Cozzuol, M. A., Coppard, S. E., Duque-Caro, H., Finnegan, S., Gasparini, G. M., Grossman, E. L., Johnson, K. G., Keigwin, L. D., Knowlton, N., Leigh, E. G., Leonard-Pingel, J. S., Marko, P. B., Pyenson, N. D., Rachello-Dolmen, P. G., Soibelzon, E., Soibelzon, L., Todd, J. A., Vermeij, G. J. and Jackson, J. B. C.: Formation of the Isthmus of Panama, *Sci. Adv.*, 2(8), 1–12, doi:10.1126/sciadv.1600883, 2016.
- Philander, S. G. and Fedorov, A. V.: Role of tropics in changing the response to Milankovich forcing some three million years ago, *Paleoceanography*, 18(2), doi:10.1029/2002PA000837, 2003.
- 450 Pontes, G. M., Wainer, I., Taschetto, A. S., Sen Gupta, A., Abe-Ouchi, A., Brady, E. C., Chan, W. Le, Chandan, D., Contoux, C., Feng, R., Hunter, S. J., Kame, Y., Lohmann, G., Otto-Bliesner, B. L., Peltier, W. R., Stepanek, C., Tindall, J., Tan, N., Zhang, Q. and Zhang, Z.: Drier tropical and subtropical Southern Hemisphere in the mid-Pliocene Warm Period, *Sci. Rep.*, 10(1), 1–11, doi:10.1038/s41598-020-68884-5, 2020.
- 455 Rebotim, A., Voelker, A. H. L., Jonkers, L., Waniek, J. J., Meggers, H., Schiebel, R., Fraile, I., Schulz, M. and Kucera, M.: Factors controlling the depth habitat of planktonic foraminifera in the subtropical eastern North Atlantic, *Biogeosciences*, 14(4), 827-859, doi: 10.5194/bg-14-827-2017, 2017.
- Regenberg, M., Nürnberg, D., Steph, S., Groeneveld, J., Garbe-Schönberg, D., Tiedemann, R. and Dullo, W. C.: Assessing the effect of dissolution on planktonic foraminiferal Mg/Ca ratios: Evidence from Caribbean core tops, *Geochemistry, Geophys. Geosystems*, 7(7), doi:10.1029/2005GC001019, 2006.

- 460 Ries, J. B.: Effect of ambient Mg/Ca ratio on Mg fractionation in calcareous marine invertebrates: A record of the oceanic Mg/Ca ratio over the Phanerozoic, *Geology*, 32(11), 981–984, doi:10.1130/G20851.1, 2004.
- Schlitzer, R.: Ocean Data View 5.2.0, <https://odv.awi.de>, 2019
- Sepulchre, P., Arsouze, T., Donnadiou, Y., Dutay, J. C., Jaramillo, C., Le Bras, J., Martin, E., Montes, C. and Waite, A. J.: Consequences of shoaling of the Central American Seaway determined from modeling Nd isotopes, *Paleoceanography*, 29(3), 176–189, doi:10.1002/2013PA002501, 2014.
- 465 Shackleton, N. J.: Attainment of isotopic equilibrium between ocean water and the benthonic foraminifera genus *Uvigerina*: isotopic changes in the ocean during the last glacial, *Colloq. Int. du C.N.R.S.*, 219, 203–209, 1974.
- Spero, H. J. and Michael, D.: The Influence of Symbiont Photosynthesis on the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ Values of Planktonic Foraminiferal Shell Calcite, , 4, 213–228, 1987.
- 470 Spero, H. J., Bijma, J., Lea, D. W. and Bemis, B. E.: Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes, *Nature*, 390(6659), 497–500, doi:10.1038/37333, 1997.
- Stap, L. B., de Boer, B., Ziegler, M., Bintanja, R., Lourens, L. J. and van de Wal, R. S. W.: CO₂ over the past 5 million years: Continuous simulation and new $\delta^{11}\text{B}$ -based proxy data, *Earth Planet. Sci. Lett.*, 439, 1–10, doi:10.1016/j.epsl.2016.01.022, 2016.
- 475 Steinhardt, J., De Nooijer, L. L. J., Brummer, G. J. and Reichart, G. J.: Profiling planktonic foraminiferal crust formation, *Geochemistry, Geophys. Geosystems*, 16(7), 2409–2430, doi:10.1002/2015GC005752, 2015.
- Steph, S.: Oxygen and carbon isotope ratios of foraminifers of ODP Site 165-1000. PANGAEA, doi:10.1594/PANGAEA.272449, 2005.
- Steph, S., Tiedemann, R., Prange, M., Groeneveld, J., Nürnberg, D., Reuning, L., Schulz, M. and Haug, G. H.: Changes in Caribbean surface hydrography during the Pliocene shoaling of the Central American Seaway, *Paleoceanography*, 21(4), 1–25, doi:10.1029/2004PA001092, 2006a.
- 480 Steph, S., Tiedemann, R., Groeneveld, J., Sturm, A. and Nürnberg, D.: Pliocene Changes in Tropical East Pacific Upper Ocean Stratification: Response to Tropical Gateways?, *Proc. Ocean Drill. Program, 202 Sci. Results*, 202(October), doi:10.2973/odp.proc.sr.202.211.2006, 2006b.
- 485 Steph, S., Regenberg, M., Tiedemann, R., Mulitza, S. and Nürnberg, D.: Stable isotopes of planktonic foraminifera from tropical Atlantic/Caribbean core-tops: Implications for reconstructing upper ocean stratification, *Mar. Micropaleontol.*, 71(1–2), 1–19, doi:10.1016/j.marmicro.2008.12.004, 2009.
- Steph, S., Tiedemann, R., Prange, M., Groeneveld, J., Schulz, M., Timmermann, A., Nürnberg, D., Rühlemann, C., Saukel, C. and Haug, G. H.: Early Pliocene increase in thermohaline overturning: A precondition for the development of the modern equatorial Pacific cold tongue, *Paleoceanography*, 25(2), 1–17, doi:10.1029/2008PA001645, 2010.
- 490 Tierney, J. E. and Malevich, S. B.: Bayesian Calibration of the Mg/Ca Paleothermometer in Planktic Foraminifera, *Paleoceanography and Paleoclimatology*, 34(12), 2005–2030., doi:10.1029/2019PA003744, 2019.
- Venancio, I. M., Mulitza, S., Govin, A., Santos, T. P., Lessa, D. O., Albuquerque, A. L. S., Chiessi, C. M., Tiedemann, R., Vahlenkamp, M., Bickert, T. and Schulz, M.: Millennial- to Orbital-Scale Responses of Western Equatorial Atlantic Thermocline Depth to Changes in the Trade Wind System Since the Last Interglacial, *Paleoceanogr. Paleoclimatology*, 33(12), 1490–1507, doi:10.1029/2018PA003437, 2018.

- Verstraete, J. M.: The seasonal upwellings in the Gulf of Guinea, *Prog. Oceanogr.*, 29(1), 1–60, doi:10.1016/0079-6611(92)90002-H, 1992.
- 500 Wagner, T.: Pliocene-Pleistocene deposition of carbonate and organic carbon at Site 959: Paleoenvironmental implications for the eastern equatorial Atlantic off the Ivory Coast/Ghana. In: Mascle, J., Lohmann, G.P., Moullade, M. (eds.), *Proceedings of the Ocean Drilling Program, Scientific Results Vol. 159*, College Station, TX (Ocean Drilling Program), 539-555, 1998.
- 505 van der Weijst, C. M. H., Winkelhorst, J., Lourens, L., Raymo, M. E., Sangiorgi, F. and Sluijs, A.: A Ternary Mixing Model Approach Using Benthic Foraminifer $\delta^{13}\text{C}$ - $\delta^{18}\text{O}$ Data to Reconstruct Late Pliocene Deep Atlantic Water Mass Mixing, *Paleoceanogr. Paleoclimatology*, 35(12), doi:10.1029/2019PA003804, 2020.
- Wiafe, G. and Nyadjro, E. S.: Satellite observations of upwelling in the gulf of Guinea, *IEEE Geosci. Remote Sens. Lett.*, 12(5), 1066–1070, doi:10.1109/LGRS.2014.2379474, 2015.
- Wilson, C. and Adamec, D.: A global view of bio-physical coupling from SeaWiFS and TOPEX satellite data, 1997-2001, *Geophys. Res. Lett.*, 29(8), 1–4, doi:10.1029/2001GL014063, 2002.
- 510 Zhang, X., Prange, M., Steph, S., Butzin, M., Krebs, U., Lunt, D. J., Nisancioglu, K. H., Park, W., Schmittner, A., Schneider, B. and Schulz, M.: Changes in equatorial Pacific thermocline depth in response to Panamanian seaway closure: Insights from a multi-model study, *Earth Planet. Sci. Lett.*, 317–318, 76–84, doi:10.1016/j.epsl.2011.11.028, 2012.
- 515 Zweng, M. M., Reagan, J. R., Antonov, J. I., Locarnini, R. A., Mishonov, A. V., Boyer, T. P., Garcia, H. E., Baranova, O. K., Johnson, D. R., Seidov, D., and Biddle, M. M.: *World Ocean Atlas 2013, Volume 2: Salinity*, edited by: Levitus, S., NOAA Atlas NESDIS, 74, 39 pp., 2013.




520 **Figure 1. General surface currents and upwelling areas in the eastern equatorial Atlantic. Blue shading: coastal upwelling in boreal summer; pink shading: equatorial upwelling; orange shading: permanent coastal upwelling. NEC: North Equatorial Current; NECC: North Equatorial Counter Current; CC: Canary Current; GC: Guinea Current; SEC: South Equatorial Current; EUC: Equatorial Undercurrent; SECC: South Equatorial Counter Current; BCC: Benguela Coastal Current. Green stippled lines mark the annual range of the Intertropical Convergence Zone (ITCZ). Figure after Norris, (1998); Wagner, (1998); Wiafe and Nyadjro (2015). Map generated with Ocean Data View (Schlitzer, 2020).**




525

530

Figure 2. (a): Upper ocean temperature (shaded contours; Locarnini et al., 2013) and salinity (black contours; Zweng et al., 2013) profiles across the black stippled line (see map) between Site 1000 and Site 959. The white stippled line shows the 20°C isotherm depth. Cross section generated with Ocean Data View (Schlitzer, 2020). (b): Seasonal vertical temperature and salinity profiles in the Gulf of Guinea (2.5°N 3.5°W; Locarnini et al., 2013; Zweng et al., 2013). Symbols (left panel) mark the 20°C isotherm depth, which varies seasonally between ~40-60 meters.




535 **Figure 3.** Conceptual manifestation of the thermocline depth as a $\Delta\text{Mg}/\text{Ca}$ $\Delta\delta^{18}\text{O}$ signal. Depth differentiation of the *G. ruber* and *N. dutertrei* habitats (stippled arrows) on a vertical temperature gradient (black line) sets a $\delta^{18}\text{O}$ and Mg/Ca offset in exported foraminiferal tests (solid arrows). The magnitude of this offset changes in response to thermocline shoaling and deepening.




540 **Figure 4.** (a): Surface (*G. ruber*; blue) and subsurface (*N. dutertrei*; orange) $\delta^{18}\text{O}$ and Mg/Ca temperature (b): Surface (*G. ruber*; green) and subsurface (*N. dutertrei*; pink) $\delta^{13}\text{C}$ records. Bold lines in (a) and (b) were smoothed with a LOESS moving regression model using a span of 0.5 in PAST software (Hammer et al., 2001). (c): Δ (Surface-subsurface) offsets from (a) and (b). Site 959 benthic $\delta^{18}\text{O}$ record (van der Weijst et al., 2020) as grey silhouette for reference. Vertical bands indicate the M2 glacial and KM3 interglacial stage. Note that planktic $\delta^{18}\text{O}$ and temperature axes are evenly scaled according to the $\delta^{18}\text{O}$ -T relationship of Shackleton (1974).


545



550 **Figure 5. (a):** Smoothed Mg/Ca and $\delta^{18}\text{O}$ records from Site 959 **(b):** Relative magnitude of the salinity gradient between surface and subsurface, calculated by deconvolving $\Delta(\text{surface-subsurface } \delta^{18}\text{O} \text{ and Mg/Ca})$, and assuming a modern tropical Atlantic $\delta^{18}\text{O}$ -S slope ($0.11\text{‰}/\text{salinity unit}$; solid line) and a doubled slope (dashed line). **(c):** LOESS smoothed (span of 0.5 in PAST software, Hammer et al., 2001) thermocline and nutricline tracers from Site 959 (as in Figure 4c) and Site 1000 in the Caribbean Sea (stippled lines, data from Steph, 2005). Site 1000 $\Delta\delta^{13}\text{C}$ was steeper than at Site 959, and was corrected with -0.4‰ to aid visual comparison.



555 **Figure 6.** Compilation of tropical $\Delta\delta^{18}\text{O}$ data. (a): Eastern Equatorial Atlantic (EEA) Site 959, this study and Norris (1998). (b): Western Tropical Atlantic (WTA) Site 1000, Steph, (2005) and Steph et al. (2006a). (c): Eastern Equatorial Pacific (EEP) Site 1241, Steph et al. (2006b). (d): Western Equatorial Pacific (WEP) Site 806, LaRiviere et al. (2012). (e) LOESS smoothed records (span of 0.5 in PAST software, Hammer et al., 2001). Surface *Trilobatus sacculifer* records were corrected with +0.33‰ to *G. ruber* based on their offset in the modern ocean (Steph et al., 2009). Blue square marks late Pliocene interval covered in Figures 4 and 5.



560

Figure 7.

565

Regionally sorted mid-latitude temperature records. (a): North Atlantic; (b): South Atlantic; (c): North Pacific; (d): South Atlantic. The records from Site 590 and 516 are Mg/Ca-based (*T. sacculifer*) from Karas et al. (2017), all other records are $U^{k_{37}}$ temperature records from the Herbert et al. (2016) compilation.