

Secular and orbital-scale variability of equatorial Indian Ocean summer monsoon winds during the late Miocene

5 Clara T. Bolton¹, Emmeline Gray^{1,2}, Wolfgang Kuhnt³, Ann Holbourn³, Julia Lübbers³,
Katharine Grant⁴, Kazuyo Tachikawa¹, Gianluca Marino^{4,5}, Eelco J. Rohling^{4,6}, Anta-Clarisse
Sarr¹, Nils Andersen⁷

1. Aix Marseille Univ, CNRS, IRD, INRAE, Coll France, CEREGE, Aix-en-Provence, France

2. Now at: School of Environment, Earth and Ecosystem Sciences, The Open University, Milton Keynes, UK

10 3. Institute of Geosciences, University of Kiel, D-24118 Kiel, Germany

4. Research School of Earth Sciences, Australian National University, Canberra ACT 2601, Australia

5. Centro de Investigación Mariña, Universidade de Vigo, GEOMA, Palaeoclimatology Lab, Vigo, 36310,
Spain

6. Ocean and Earth Science, University of Southampton, Southampton SO14 3ZH, UK

15 7. Leibniz Laboratory for Radiometric Dating and Stable Isotope Research, University of Kiel, D-24118 Kiel,
Germany

Correspondence to: Clara T. Bolton (bolton@cerege.fr)

Abstract

20 In the modern northern Indian Ocean, biological productivity is intimately linked to near-
surface oceanographic dynamics forced by the South Asian, or Indian, monsoon. In the late
Pleistocene, this strong seasonal signal is transferred to the sedimentary record in the form of
strong variance in the precession band (19-23 kyr) because precession dominates low-latitude
insolation variations and drives seasonal contrast in oceanographic conditions. In addition,
25 internal climate system feedbacks (e.g., ice-sheet albedo, carbon cycle, topography) play a
key role in monsoon variability. Little is known about orbital-scale monsoon variability in the
pre-Pleistocene, when atmospheric CO₂ levels and global temperatures were higher. In
addition, many questions remain open regarding the timing of the initiation and
intensification of the South Asian monsoon during the Miocene, an interval of significant
30 global climate change that culminated in bipolar glaciation. Here, we present new high-
resolution (< 1 kyr) records of export productivity and sediment accumulation from
International Ocean Discovery Program Site U1443 in the southernmost Bay of Bengal
spanning the late Miocene (9 to 5 million years ago). Underpinned by a new orbitally tuned
benthic isotope stratigraphy, we use X-Ray Fluorescence-derived biogenic barium variations
35 to discern productivity trends and rhythms. Results show strong eccentricity-modulated
precession-band productivity variations throughout the late Miocene, interpreted to reflect
insolation forcing of summer monsoon wind strength in the equatorial Indian Ocean. On long
timescales, our data support the interpretation that South Asian monsoon winds were already

established by 9 Ma in the equatorial sector of the Indian Ocean, with no apparent
40 intensification over the latest Miocene.

1 Introduction

The Asian monsoon is a major hydrological phenomenon that is driven by atmospheric
pressure gradients created by asymmetric heating between the equatorial Indian and western
45 Pacific Oceans and the Indo-Asian landmass, creating strong seasonally reversing winds and
ocean currents and heavy boreal summer precipitation over the Bay of Bengal and Indian
subcontinent (Webster, 1987a, b; Schott and McCreary Jr, 2001; Gadgil, 2003; Schott et al.,
2009). Surface winds over the northern Indian Ocean (Arabian Sea and Bay of Bengal) and
South China Sea are strong indicators of the strength of the South Asian and East Asian
50 monsoon subsystems, respectively, and precipitation amount and seasonality can also be
diagnostic of monsoon strength (e.g., Webster and Yang, 1992; Goswami et al., 1999). Thus,
past monsoon dynamics can be reconstructed using wind, runoff, and precipitation indicators
recorded in marine sediments from these core convective regions in the Bay of Bengal and
South China Sea. The Asian monsoon is known to have varied substantially over short
55 (interannual to suborbital) and long (orbital to geological) timescales in response to forcing
factors both external and internal to Earth's climate system (e.g., Wang et al., 2005; Clemens
and Prell, 2003; Farnsworth et al., 2019; Kathayat et al., 2016).

There is great uncertainty surrounding the timing of Asian monsoon initiation and
60 intensification, and the degree of coupling between regional monsoon subsystems.
Discrepancies in part stems from the fact that records come from the South Asian or East
Asian monsoon subsystems, which are sensitive to different aspects of regional topography
(Molnar et al., 2010; Clift et al., 2008; Clift and Webb, 2019; Boos and Kuang, 2010; Acosta
and Huber, 2020). Further, differences in monsoon expression occur even within the core
65 convective region of the South Asian monsoon (e.g., dominance of summer monsoon winds
in the southern Bay of Bengal *versus* monsoonal rainfall/runoff in the northern and eastern
parts). Meanwhile, proxies generally record singular aspects of monsoonal climate that are
not necessarily coupled on all timescales (e.g., winter or summer monsoon winds,
precipitation total amount or degree of seasonality). Evidence for strong monsoonal climates
70 (i.e. with strong seasonality of precipitation) exists across Asia during the Paleogene (Spicer
et al., 2017; Licht et al., 2014), yet many terrestrial records from southeast Asia suggest an

onset of the monsoon near the Oligocene-Miocene boundary (~24-22 Ma) (Guo et al., 2002; Sun and Wang, 2005; Wang et al., 2005). Marine records of drift sedimentation near the Maldives archipelago as well as upwelling and oxygenation indicators from the Arabian Sea, both influenced by wind and surface ocean circulation, suggest an onset of strong seasonally-reversing South Asian (monsoon) winds and Arabian Sea upwelling during the late middle Miocene (~13-10 Ma) (Zhuang et al., 2017; Gupta et al., 2015; Betzler et al., 2016; Betzler et al., 2018; Bialik et al., 2020; Nigrini, 1991).

In contrast, a time interval in which records from different regions and proxies converge somewhat is the late Miocene. Magnetic records from the Chinese Loess Plateau are interpreted to show a long-term intensification of the East Asian summer monsoon (EASM) from ~8.2 to 2.6 Ma (Ao et al., 2016). A late Miocene strengthening of Asian winter monsoons is inferred from South China Sea (Holbourn et al., 2018; Jia et al., 2003; Wan et al., 2007) and Andaman Sea (Lee et al., 2020) records. In the Arabian Sea, an intensification of upwelling and productivity at ~8 Ma is interpreted to reflect a strengthening of the South Asian Summer Monsoon (SASM) (Kroon et al., 1991; Singh and Gupta, 2014; Gupta et al., 2004), although other studies find evidence contrary to this (Tripathi et al., 2017; Huang et al., 2007). Proposed monsoon intensifications during the late Miocene roughly coincide with strong global cooling (Herbert et al., 2016), and a number of studies have implicated cooling and the ramp-up of Antarctic glaciation in monsoon strengthening (Ao et al., 2016; Holbourn et al., 2018; Gupta et al., 2004). Until now, a lack of continuous, well-preserved marine sequences from the South Asian monsoon region has stalled our understanding of its complex evolution during the Miocene.

The SASM is thought to have varied strongly on orbital timescales because monsoon strength responds, both directly and via internal feedback mechanisms, to insolation forcing. Model simulations predict a stronger South Asian monsoon during summer insolation maxima at both precession minima and (to a lesser degree) obliquity maxima (Bosmans et al., 2018; Prell and Kutzbach, 1992, 1987; Kutzbach, 1981; Jalihal et al., 2019; Tabor et al., 2018; Ding et al., 2021). Precession is the dominant control on insolation and its seasonal distribution near the equator, and proxy-based Pleistocene SASM records show strong precession-band (19-23 kyr) variability (Kathayat et al., 2016; Prell and Kutzbach, 1987; Clemens et al., 1991; Zhisheng et al., 2011; Bolton et al., 2013; Caley et al., 2011; Gebregiorgis et al., 2018; Wang et al., 2005; Clemens and Prell, 1990; Rostek et al., 1997). The influence of global boundary

conditions related to global ice volume and greenhouse gas concentrations on SASM winds and precipitation/runoff is also demonstrated by strong obliquity- and eccentricity-band variance in Plio-Pleistocene records (e.g., Clemens and Prell, 2003; Gebregiorgis et al., 2018; Clemens et al., 2021; An et al., 2011). Our current knowledge of orbital-resolution past productivity fluctuations in the South Asian monsoon region and their relationship with local (insolation) or remote (global ice volume, greenhouse gases) forcing mechanisms comes almost entirely from Pleistocene Arabian Sea sedimentary records (Clemens and Prell, 2003; Caley et al., 2011; Singh et al., 2011; Shimmield and Mowbray, 1991; Rogalla and Andruleit, 2005; Clemens et al., 1991; Ziegler et al., 2010). These records suggest that summer monsoon proxies significantly lag northern hemisphere summer insolation maxima in the precession and obliquity bands due to climate feedbacks. Orbital control on past SASM strength in the pre-Pleistocene, when boundary conditions were different, has so far only been investigated in the Andaman Sea over the latest Miocene-early Pliocene, where seawater oxygen isotope data suggest high-amplitude precession and obliquity forcing of monsoon rainfall with significant phase lags (Jöhnck et al., 2020).

In this paper, we investigate sediment accumulation and export productivity dynamics at millennial resolution in late Miocene sediments from southern Bay of Bengal (BOB) International Ocean Discovery Program (IODP) Site U1443 (5°23'N, 90°22'E, Fig. 1). The late Miocene (11.6-5.3 Ma) is an interval of major global climate change, with long-term cooling between ~7.5 and 5.5 Ma (Herbert et al., 2016) culminating in major high-latitude cooling events (Holbourn et al., 2018), and important carbon cycle shifts recorded in the marine and terrestrial realms potentially linked to atmospheric CO₂ decline (Tauxe and Feakins, 2020; Steinthorsdottir et al., 2020; Rae et al., 2021). The region of Site U1443 is strongly influenced by seasonally reversing monsoon winds today, and primary productivity is tightly coupled to the annual monsoon cycle (Fig. 1a-b, Fig. 2). During the SASM, strong moisture-laden winds blow inland, driving surface circulation changes and increased mixed layer depth (Fig. 1a-b) (Webster, 1987a, b; Schott and McCreary Jr, 2001; Gadgil, 2003; Schott et al., 2009). Strong Ekman pumping mixes nutrients into the surface layer during the South Asian summer monsoon and, to a lesser extent, the winter monsoon, stimulating biological productivity (Fig. 2) (Lévy et al., 2007; McCreary et al., 2009; Koné et al., 2009; Behrenfeld et al., 2005; Longhurst, 1995). This strong seasonal signal is transferred to the sedimentary record in the form of strong variance at orbital periods because insolation variations drive seasonal contrast. Here, we generate a new orbitally tuned age model based

140 on benthic foraminiferal stable isotopes spanning ~9 to 5 Ma. Core-scanning X-Ray
Fluorescence (XRF) data are then used to reconstruct bulk, carbonate, and biogenic barium
content and mass accumulation rates (MARs), shedding light on secular and orbital-scale
export productivity and sedimentation changes over the late Miocene.

145 **2. Background**

In the modern southern BOB waters overlying Site U1443, both primary and export
productivity are strongly controlled by seasonally reversing winds associated with the South
Asian monsoon. Figure 2 shows the annual cycle of wind stress, mixed layer depth (MLD),
net primary productivity, and biogenic particle export based on recent oceanographic,
150 satellite, and sediment trap data (see methods for details). In boreal summer (JJA) strong
southwest winds mix the upper water column, deepening the MLD to ~60 m and entraining
nutrients into the photic zone, leading to enhanced primary productivity and biogenic particle
export (with a lag of ~3-4 weeks) (Fig. 1a-b, Fig. 2). During boreal winter, northeast winds
deepen the MLD to a lesser extent, resulting in a second smaller peak in productivity during
155 the winter monsoon (DJF) (Fig. 2). During the inter-monsoon seasons, lowest wind stress is
recorded leading to a shallow MLD, higher Sea Surface Temperatures (SSTs), and a more
stratified upper water column, resulting in increased oligotrophy and reduced biological
productivity. The biannual productivity maxima observed in the surface ocean above Site
U1443 is characteristic of monsoon-dominated tropical regions (Longhurst, 1995; Lévy et al.,
160 2007).

Sediment trap data suggest that primary productivity is the dominant control on organic
carbon export at a location west of the northern end of the Ninetyeast Ridge (SBBT site; 5°N,
87°E, Fig. 1c). However, lithogenic mineral ballasting at this location is not negligible
165 (average ~13% and 15% lithogenic particles in shallow and deep traps respectively) (Rixen et
al., 2019; Unger et al., 2003) and could in part explain the bias towards the late summer peak
seen in biogenic fluxes compared to net primary productivity, as maximum concentrations of
lithogenic particles at SBBT occur during the summer monsoon. While wind forcing is
identified as the dominant factor controlling biogenic particle fluxes at the SBBT site,
170 advection of nutrient- and chlorophyll-rich waters originating from the eastern Arabian Sea
via the Southwest Monsoon Current may further contribute to the summer productivity peak
in this region (Unger et al., 2003). During the summer monsoon, the relatively salty and

nutrient-rich Southwest Monsoon Current flows eastwards south of Sri Lanka then turns northwards into the BOB (Fig. 1c) (Schott et al., 2009; Jensen, 2003). The Southwest Monsoon Current and associated eddies have been shown to increase chlorophyll concentrations and average phytoplankton size along their paths as far east as 88-90°E, with the current's influence generally restricted to north of 6°N at this longitude (Jyothibabu et al., 2015; Vinayachandran et al., 2004; Webber et al., 2018). While river runoff and resultant salinity stratification during the summer monsoon suppress primary productivity further north in the BOB (Prasanna Kumar et al., 2002), seasonal surface salinity variations are very small (<0.2 psu) at 5°N (Zweng et al., 2013). Accordingly, monsoon impacts on nutrients and productivity in our study area are limited to those driven by surface currents and wind mixing, and biogenic export fluxes during the SASM are similar (particulate organic carbon) or higher (CaCO₃ and biogenic SiO₂) than at sites further north in the BOB (Unger et al., 2003). Thus, modern data give us confidence that export productivity at our site is likely a reflection of South Asian (primarily summer) monsoon wind strength, via its control on MLD and nutrient entrainment into the mixed layer and surface ocean currents.

3 Materials & Methods

3.1 Site and sampling

Samples used in this study are from Site U1443 (Latitude 5°23.01'N, Longitude 90°21.71'E, water depth 2924 m), drilled during IODP Expedition 353 in the southernmost BOB on the crest of Ninetyeast Ridge (NER) (Clemens et al., 2016) (Fig. 1). During the late Miocene, Site U1443 is estimated to have migrated northeastwards from 1°71'N, 88°06'E at 9 Ma to 3°27'N, 89°04'E at 5 Ma (Fig. 1; paleo-location estimates from <http://portal.gplates.org>), although its position relative to peninsular India remained stable. Site U1443 is located ~100 m southeast of Ocean Drilling Program (ODP) Site 758 and is a re-drill of this legacy site (Shipboard Scientific Party, 1989). At Site U1443, the use of Advanced Piston Coring (APC) and half-length APC drilling techniques down to >200 m CCSF (core depth below sea floor) in three holes allowed recovery of a complete, spliced Neogene sedimentary section spanning 0-195 m CCSF (core composite depth below sea floor). Late Miocene records cover the interval 70.06 m CCSF (U1443B 7H 5W 75-76 cm) to 122.76 m CCSF (U1443C 15H 4W 148-149 cm), following the revised shipboard splice (Table S1), spanning the interval ~9.5 to 5 Ma based on initial bio-magneto-stratigraphy. Samples come from lithologic Units Ib and IIa, and sediments mainly consist of light grey to pale yellow nannofossil ooze with clay and

foraminifers, and occasional volcanic ash (Clemens et al., 2016). Cores were sampled (1 cm half rounds) at a depth resolution of 4 cm in the upper part of the late Miocene interval (70.06-114.18 m) and 2 cm in the lower part (114.18-122.76 m) where sedimentation rate estimates were lower. U-channels for XRF scanning were sampled from archive halves of sediment cores at Kochi Core Centre (Japan) during the post-cruise sampling party for 39 sections included in the splice between U1443C 9H 2A (69.95 m CCSF) and U1443C 13H 5A (113.58 m CCSF).

3.2 Modern Oceanography

Modern oceanographic conditions over the seasonal cycle above Site U1443 were assessed using recent datasets to investigate the regional relationship between monsoon winds and biological productivity (Figs. 1a-b, 2). Monthly data for wind (Wind Stress, Metop-A ASCAT, 0.25°, Global, Near Real Time, 2009-present) (Fig. 2a), MLD (1969-2010) (Keerthi et al., 2013) (Fig. 2b) and depth-integrated net primary productivity estimated from satellite-derived surface chlorophyll concentrations (Primary Productivity, Aqua MODIS, NPP, Global, 2003-present, EXPERIMENTAL (Monthly Composite) calculated using method of Behrenfeld and Falkowski (1997); (Erd, 2020)) (Fig. 2c) were extracted for a box between ~4.5-5.5°N latitude and 89-91°E longitude (depending on grid resolution) and binned by month. Scatter thus reflects a combination of spatial variability within our 1° by 2° box and interannual variability; monthly means over each time series are also shown. Particulate organic carbon, biogenic SiO₂, and CaCO₃ fluxes (Fig. 2d, e) are from SBBT sediment trap samples (~5°N, 87°E, Fig. 1c) (Unger et al., 2003). In Fig. 2 we show data from the deep traps (~3000 m, ~21 day sampling intervals) (Rixen et al., 2019), but seasonal patterns of biogenic particle flux are very similar in the shallow (~1000 m) SBBT traps (Unger and Jennerjahn, 2009; Vidya et al., 2013). Data points show fluxes recorded in individual years (1987-1997, plotted against mid-time for the trap deployment), with monthly averages also shown. Monthly wind and net primary productivity data were downloaded from the ERDDAP server (<https://coastwatch.pfeg.noaa.gov/erddap/index.html>) and Indian Ocean MLD data (Keerthi et al., 2013) from http://www.ifremer.fr/cerweb/deboyer/mlD/Surface_Mixed_Layer_Depth.php.

3.3 Late Miocene benthic foraminiferal stable isotope data

Bulk sediment samples were oven-dried at 50°C, weighed, and washed over a 63 µm sieve in tap water at CEREGE (Centre Européen de Recherche et d'Enseignement des Géosciences de l'Environnement). The >63 µm fraction was oven-dried at 50°C on a filter paper and weighed to determine percentage coarse fraction. The <63 µm fraction was centrifuged and dried at 50°C. Depth resolution for the benthic isotope record is 8 cm (70.06 m to 112.87 m CCSF), 4 cm (112.86 m and 114.18 m CCSF) or 2cm (114.18-122.76 m CCSF), except near splice tie-points where sampling included overlap between cores increasing resolution. Six to twelve specimens of the epibenthic foraminiferal species *Cibicidoides wuellerstorfi* were picked from the >212µm fraction, with 6-8 well-preserved specimens selected for analysis. Tests were broken into fragments, cleaned in ethanol in an ultrasonic bath, and oven dried at 40°C. Stable carbon and oxygen isotopes were measured on a Thermo Scientific MAT 253 dual-inlet isotope ratio mass spectrometer (DI-IRMS) coupled to Kiel IV carbonate preparation device at the Leibniz Laboratory, University of Kiel. Based on long-term analysis of international and internal carbonate standards, precision (1σ) is better than ±0.08‰ for δ¹⁸O and 0.05‰ for δ¹³C. Results were calibrated using the National Institute of Standard and Technology (NIST) carbonate isotope standard NBS (National Bureau of Standard) 19, and are reported on the Vienna PeeDee Belemnite (VPDB) scale. *C. wuellerstorfi* isotope data below 112.86 m CCSF were originally published in Lübbers et al. (2019), and a low-resolution version of the long-term δ¹³C record in *Bretschneider et al.* [submitted].

3.4 Age model

Seven calcareous nannofossil bio-events dated between 9.53 Ma and 5.04 Ma were identified at ~0.5 to 1 m resolution in Site U1443 splice samples (Table S2) to increase the resolution of shipboard biostratigraphy (Robinson et al., 2016). To check for orbital periodicities prior to tuning, wavelet analyses were performed on benthic isotope records in the depth domain and on the revised nannofossil-based age model (using a 4th order polynomial fit, Fig. S1) in R using the biwavelet package (Gouhier et al., 2016; Grinsted et al., 2004) (Fig. S2a-d). All time-series were first interpolated to constant depth or age resolution, such that the maximum resolution present was preserved (2 cm and 2 kyr for benthic isotope records in the depth and age domain, respectively). Records were then detrended to remove signals with periods longer than one third of the length of the dataset using the “bandpass” function in the R package Astrochron (Meyers, 2014). Wavelets indicate that obliquity-driven cycles are

270 present throughout (~0.53 m, Fig S2 a-b; 41 kyr and 53 kyr, Fig. S2 c-d), confirming that the
U1443 record is suitable for orbital tuning. Using revised nannofossil datums (Table S2, Fig.
S1) and shipboard magnetostratigraphy (Clemens et al., 2016) as preliminary age-depth tie-
points (Fig. 3), an astronomical age model was constructed by tuning our monospecific
benthic $\delta^{18}\text{O}$ record to an eccentricity plus tilt (ET) target curve (Laskar et al., 2004) (Table
275 S3, Fig. 4). We did not include precession in our tuning target so as not to introduce
assumptions related to which hemisphere was controlling climate at our site, and because the
temporal resolution of our benthic record does not permit accurate resolution of precession
cycles in some intervals. We used a minimal tuning approach, tying ET maxima to benthic
 $\delta^{18}\text{O}$ minima, with at most one tie-point per ~100 kyr and often one every 200-300 kyr (Fig.
280 4, Fig. S1), so as not to artificially introduce frequency modulations (Zeeden et al., 2015).

3.5 XRF Scanning and Calibration

U-channels were scanned at 1 cm intervals at The Australian National University (ANU) on a
third generation Avaatech XRF core scanner. All cores were scanned sequentially and
285 standards measured daily were consistent across all runs. Core sections were covered with 4
micron-thick Ultralene film and measured at 10 kv with a 500 μA current and no filter, then
at 30 kv with a 200 μA current and Pd thin filter, and finally at 50 kv with a 50 μA current
and Cu filter. A 30 s count time was used for all runs. Late Miocene XRF data generated at
ANU (72.75 m to 113.56 m CCSF) were spliced with data from Lübbers et al. (2019)
290 (112.80m to 122.76m CCSF), also scanned on an Avaatech XRF core scanner but with
different machine settings. To splice the records, we rescaled the raw Lübbers et al. (2019)
elemental count data so that absolute values and variance matched our data, based on an
overlapping interval between 112.80 and 113.56m CCSF.

295 Quantitative chemical compositions of a subset of discrete bulk sediment samples were
determined at CEREGE after total digestion by Inductively Coupled Plasma Mass
Spectrometer (ICP-MS Agilent 7500 ce). Twenty samples from the scanned late Miocene
interval, selected to cover the range of values in the raw XRF count data for elements of
interest, were analysed and concentrations of Al, K, Ca, Ti, Mn, Fe, Rb, Sr, and Ba were
300 determined. Prior to analysis, samples were dried and homogenised in a pestle and mortar.
About 30 mg of sediment was completely dissolved by acid digestion using a 2:1 mixture of
ultrapure acids (15 M HNO_3 and 22 M HF with HClO_4) on a hot plate. Blank contribution

was estimated to be negligible. The accuracy of measurements was evaluated using analysis of geostandards MAG-1 (marine mud) and BE-N (basalt). The typical analytical uncertainty was better than 5%. XRF-derived element counts were converted into element concentrations by direct linear calibration. This allowed us to reduce uncertainties related to the variable matrix effect and physical properties such as moisture content that typically change downcore. Linear calibration between element counts and concentrations in discrete samples over the interval ~8.15 to 5 Ma showed significant coefficients of determination, with R^2 values ranging from 0.68 (Al) to 0.87 (Fe) (Fig. S3). For Ca and Sr, R^2 values were lower (0.39 and 0.42 respectively, Fig S3) due to the consistently high Ca and Sr contents and small variability in the selected calibration samples. To estimate percent CaCO_3 , we therefore used a Ca/Fe ratio calibration rather than a direct linear calibration. We first used the linear relationship between Ca/Fe counts and Ca/Fe as determined by ICP-MS (Fig. S3, $R^2=0.93$). Then % CaCO_3 was calculated assuming that all Ca was contained in CaCO_3 – a reasonable assumption at Site U1443 given the relatively low clay content in lithological subunits Ib and IIa (Clemens et al., 2016). Additionally, we used calibrated XRF data to calculate biogenic barium concentrations and “carbonate-free basis” (cfb) elemental concentrations, permitting evaluation of the extent to which dilution by the dominant sediment constituent (here CaCO_3) is driving trends and variability of more minor constituents in our records. To represent the relative contributions of CaCO_3 versus terrigenous sediment components, we use the log count ratio of $\text{Ca}/(\sum(\text{Al, K, Ti, Fe, Rb}))$, termed $\log(\text{Ca}/\text{Terr})$.

3.6 Ba-based export productivity proxies

The accumulation of biogenic barium in sediments is a reliable proxy for export production in certain environments (Paytan and Griffith, 2007). Micron-sized barite (BaSO_4) crystals are the main carriers of particulate barium in the ocean, with a maximum in concentration occurring just below the euphotic zone (Bishop, 1988; Dehairs et al., 1980). Although the exact mechanisms governing the precipitation of barite in the water column are only now coming to light (Martínez-Ruiz et al., 2019), its formation is thought to be associated with decaying organic matter. Depth profiles of dissolved Ba suggest that passive adsorption of barite onto mainly biogenic particles as they sink through the water column, combined with vertical mixing of dissolved Ba from the deep ocean and riverine input, can best explain Ba’s nutrient-like water column distribution (Dehairs et al., 1980; Cao et al., 2016). Goldberg and Arrhenius (1958) first hypothesised that an increase in Ba accumulation rate in sediments

underlying the equatorial Pacific divergence zone was directly linked to overlying high productivity, followed by similar observations in equatorial Indian Ocean sediments (Schmitz, 1987). Subsequently, evidence for strong correlations between fluxes of Ba and organic carbon in Atlantic and Pacific sediment traps led to algorithms relating new productivity to particulate Ba flux (Dymond et al., 1992; Francois et al., 1995). A further study focusing on the accumulation of barite (Ba_{barite}) extracted from core-top and late Pleistocene sediments refined its use as a proxy for export productivity (Paytan et al., 1996). Although significant Ba regeneration occurs in the uppermost few millimetres of sediment (Paytan and Kastner, 1996), barite dissolution is thought to cease after burial due to supersaturation in interstitial waters (Gingele and Dahmke, 1994; Dymond et al., 1992) and barite is not subject to burial diagenesis in oxic sediments (Paytan et al., 1993). Ocean sedimentary Ba has both a biogenic (Ba_{bio}) and a terrigenous (Ba_{detrital}) component, so estimates of past export productivity using barium must distinguish between these sources. This can either be done by chemical leaching of bulk sediment (assuming that all barite is Ba_{bio}) e.g. (Paytan et al., 1996), or by determination of total barium (Ba_{total}) and subtraction of Ba_{detrital} using Al content and the terrigenous Ba/Al ratio, resulting in a record of Ba_{xs} ($Ba_{\text{total}} - Ba_{\text{detrital}}$), see equation 1 (Dymond et al., 1992).

$$[Ba_{\text{xs}}]_{\text{ppm}} = [Ba_{\text{total}}]_{\text{ppm}} - (Ba/Al_{\text{terrigenous}}) * [Al_{\text{total}}]_{\text{ppm}} \quad (1)$$

Direct comparisons of measurements of Ba_{barite} and Ba_{xs} suggest that non-barite phases of barium may be included in the calculation of Ba_{xs} ; nevertheless Ba_{xs} is most representative of Ba_{barite} and therefore export productivity in oxic carbonate-rich sediments with low terrigenous, biogenic silica, and organic carbon contents (Eagle et al., 2003; Averyt and Paytan, 2004; Gonnee and Paytan, 2006). The use of bulk Ba/Ti, Ba/Al, and Ba/Fe ratios is another approach to evaluate relative changes in export productivity (i.e., normalisation to an element presumed to be predominantly of terrigenous origin), but without precisely predefining the Ba/terrigenous ratio that could vary over time, and also removing the effect of dilution by a dominant sedimentary component such as $CaCO_3$ (Murray et al., 2000).

Here, we reconstruct changes in export productivity at Site U1443 over the late Miocene using XRF-derived Ba data and compare elemental count ratios of $\log(Ba/Fe)$, $\log(Ba/Ti)$, and $\log(Ba/Al)$, with $[Ba]_{\text{cfb}}$ and $[Ba]_{\text{xs}}$ calculated following equation (1), using a $Ba/Al_{\text{terrigenous}}$ value of 0.0075 g/g following Dymond et al. (1992). To verify consistency of

370 trends, we also calculate $[Ba]_{xs}$ using $[Ti]$ to represent $Ba_{detrital}$, applying a $Ba/Ti_{terrigenous}$ ratio
of 0.183 g/g (Mclennan, 2001), and carbonate-free $[Ba]_{xs}$.

3.7 Mass Accumulation rates

MARs of bulk sediment, $CaCO_3$, $[Ba]_{xs}$, and summed terrigenous elements (Al, K, Ti, Fe,
375 and Rb) were calculated by multiplying concentrations by linear sedimentation rates (in
cm/kyr) derived from our new age model and dry bulk densities (in g/cm^3). Dry bulk density
values were estimated from high-resolution shipboard Gamma Ray Attenuation bulk density
scanning data, and the linear relationship between all shipboard U1443 wet bulk density and
dry bulk density measurements ($n=164$, $R^2 > 0.99$) (Clemens et al., 2016). Units are $g/cm^2/kyr$
380 for bulk and $CaCO_3$ MAR, $\mu g/cm^2/kyr$ for $[Ba]_{xs}$ MAR and $mg/cm^2/kyr$ for terrigenous
MAR.

3.8 Time series analysis

Spectral analyses of benthic isotope and XRF data against tuned age were performed on
385 detrended records (bandpass filtered to remove signals with periods longer than one third of
the length of each dataset) with a constant time step of 0.5 kyr for XRF records and 2 kyr for
isotope records. Multi-taper method (MTM) spectral analyses using a robust red-noise model
were performed using Acycle (Li et al., 2019). Blackman-Tukey cross-spectral analyses were
performed in Arand to assess phase and coherence (Howell et al., 2006). Wavelet analyses
390 were performed in R using the biwavelet package (Gouhier et al., 2016; Grinsted et al.,
2004). To illustrate precession-band variance and amplitude modulation, certain records (with
identified significant precession variance) were filtered using a Tanner-Hilbert filter centred
on 46.5 cycles/Myr with bandwidth ± 8.5 (designed to include all precession terms with
periods between 18 and 26 kyr) in Acycle (Li et al., 2019).

395 4. Results

4.1 Age model and benthic foraminiferal isotope data

U1443 benthic $\delta^{18}O$ and $\delta^{13}C$ data between 70.06 m and 122.76 m CCSF are shown in the
depth domain in Figure 3 alongside calcareous nannofossil datums (revised herein, Table S2)
and magnetochron boundaries. Our tuned benthic $\delta^{18}O$ and $\delta^{13}C$ records, shown in Figure 4,
400 span the interval 8.99 to 4.96 Ma and our age model shows excellent agreement with revised
biostratigraphic and shipboard magnetostratigraphic datums (Fig. S1). Sedimentation rates

generally vary between 1 and 1.7 cm/kyr, with a minimum of ~0.5-0.7 cm/kyr in the oldest part of the record (9 to 8.6 Ma) and a maximum of ~1.9 cm/kyr at 8 to 7.8 Ma (Fig. 4d). Between 112.86 m and 121 m CCSF (8.7-8.1 Ma), our age model differs by up to 60 kyr
405 from that of Lübbers et al. (2019), which is based on correlation of Site U1443 benthic $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ data to the orbitally tuned ODP Site 1146 $\delta^{13}\text{C}$ record (Holbourn et al., 2018) using a limited number of tie points (Site locations in Fig. 1c). Both wavelet analyses (Fig. 4 f, g) and spectral analyses (Fig. S2e, f) of the tuned benthic records reveal significant (>99%) orbital periodicities of ~405 kyr and 41 kyr for $\delta^{18}\text{O}$ and ~405 kyr, 125 kyr, 95 kyr, 53 kyr
410 and 41 kyr for $\delta^{13}\text{C}$, and filtered isotope records show good correspondence with filtered ET (Fig. 4h). Cross-spectral analysis between $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ reveals >95% coherency in the 41 kyr and 405 kyr bands (Fig. S4a). Our age model is supported by close agreement between Site U1443 benthic $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ data and independent orbitally tuned benthic isotope records from the South China Sea (ODP 1146) (Holbourn et al., 2018; Holbourn et al., 2021),
415 equatorial Pacific (IODP Sites U1338 and 1337) (Drury et al., 2016; Drury et al., 2018; Drury et al., 2017), and equatorial Atlantic (ODP Sites 926 and 999) (Bickert et al., 2004; Shackleton and Hall, 1997; Drury et al., 2017; Zeeden et al., 2013) (Figs S5, S6).

Mean temporal resolution of the Site U1443 benthic isotope record is 4.2 kyr. Between 9 and
420 7.6 Ma, mean benthic $\delta^{18}\text{O}$ values vary between 2.5 and 2.8 ‰, with an overall decreasing trend culminating in minimum values averaging ~2.4 ‰ between 7.6 and 7 Ma (Fig. 4c). Between 7 and 6.5 Ma, mean $\delta^{18}\text{O}$ values increase by ~0.25 ‰, and between 6.5 and 5 Ma, mean values vary between 2.55 and 2.8 ‰. Between 6 and 5 Ma, a number of prominent benthic $\delta^{18}\text{O}$ maxima are identified in the U1443 $\delta^{18}\text{O}$ record, namely TG2, TG12, TG14,
425 TG20 and TG22 (following nomenclature of Shackleton et al. (1995)) (Fig. 4c). Between 7.7 and 6.9 Ma, strong obliquity modulation of the U1443 $\delta^{18}\text{O}$ record is seen (Fig. 4f), as also noted at Sites U1337 (Drury et al., 2017) and 1146 (Holbourn et al., 2018) (Fig. S5). Long-term trends are similar to those recorded at Pacific sites, with benthic $\delta^{18}\text{O}$ values at Indian Ocean Site U1443 ~0.1‰ heavier than at Pacific Sites U1337/U1338 and ~0.2-0.3‰ heavier
430 than at South China Sea Site 1146 (Fig. 5a).

Mean benthic $\delta^{13}\text{C}$ values at Site U1443 vary between 0.7 and 1.1‰ from 9 to 7.6 Ma, then decrease from ~1 to -0.2‰ between 7.6 and 6.7 Ma, reflecting the globally recognised Late Miocene Carbon Isotope Shift (LMCIS) (Keigwin, 1979; Keigwin and Shackleton, 1980)
435 (Fig. 4e). From 6.7 to 5 Ma, mean values vary between -0.2 and 0.4 ‰. The timing of the

LMCIS at Site U1443 (~7.6 to ~6.7 Ma) is synchronous with the event in independent orbitally tuned high-resolution records (Drury et al., 2018; Drury et al., 2017; Holbourn et al., 2018; Drury et al., 2016) (Fig. 5b, Fig. S6), and its magnitude (~1‰ decrease in $\delta^{13}\text{C}$ in smoothed record) is similar to that recorded in Pacific Ocean sediments from Sites U1338, U1337 and 1146. The Site U1443 $\delta^{13}\text{C}$ record shows a consistent positive offset of 0.15-0.25 ‰ relative to South China Sea Site 1146 over the 9 to 5 Ma interval (Fig. 5b).

4.2 XRF data

Scanning XRF results are shown in Figure 6. Raw and calibrated elemental data show consistent trends and amplitude variability (Fig. 6). For Ti, Ba, Al, and Mn, the re-scaled counts/s values in the 113.37 to 122.76 m CCSF interval (Lübbbers et al., 2019) fell outside of our calibration range, thus data below 113.6 m (~8.15 Ma) were not converted to concentrations (Fig. 6). In brief, Al, Si, Ti, Fe, Rb, and K show similar trends, with a long-term small increase in concentrations from 8.15 to 5 Ma and spikes (particularly pronounced in Rb and K) corresponding in some cases to described ash layers (Clemens et al., 2016). Ca and Ba show a minor long-term decrease over the study interval, while Sr and Mn increase from ~8.15 to 6 Ma, then stabilise or decrease slightly. All elements show high-frequency variability throughout. Confidence in our method of calculating CaCO_3 content is provided by very good agreement with independent % CaCO_3 estimates for the middle and early late Miocene interval of Site U1443 based on calibration of XRF-derived counts of $(\text{Ca}/\sum(\text{Ca}, \text{Al}, \text{Si}, \text{K}, \text{Ti}, \text{Mn}, \text{Fe}, \text{S}))$ to discrete CaCO_3 measurements (Lübbbers et al., 2019), including an overlapping interval based on an alternate splice from 112.8 and 113.6 m CCSF (Fig. 7b).

4.3 CaCO_3 content, sediment accumulation patterns and Ba proxies

Late Miocene estimated % CaCO_3 varies between ~60 and 90% with a slight long-term decrease over the 9 to 5 Ma interval (Fig. 7b). This long-term trend is also visible in the $\log(\text{Ca}/\text{Terr})$ record (Fig. 7c), implying a small increase in the contribution of terrigenous material relative to CaCO_3 in Site U1443 sediments over time. Three % CaCO_3 and $\log(\text{Ca}/\text{Terr})$ minima between 6 and 5 Ma occur in identified ash layers. $\log(\text{Ba}/\text{Al})$, $\log(\text{Ba}/\text{Fe})$ and $\log(\text{Ba}/\text{Ti})$ show identical long-term and orbital-scale trends (Fig. S7), therefore we only discuss $\log(\text{Ba}/\text{Fe})$ in the main text. $\log(\text{Ba}/\text{Fe})$ shows a long-term decrease between 9 and 5.3 Ma, and a smaller increase from 5.3 to 5 Ma (Fig. 7d). $[\text{Ba}]_{\text{xs}}$ shows identical variability whether calculated using Al or Ti (Fig. 7e), and generally shows

similar patterns to $\log(\text{Ba}/\text{Fe})$ where records overlap (8.15 to 5 Ma). Values of $[\text{Ba}]_{\text{xs}}$ generally vary between 400 and 800 ppm (representing on average 83% of total $[\text{Ba}]$), and carbonate-free $[\text{Ba}]_{\text{xs}}$ concentrations are typically 1000 to 4000 ppm (Fig. S7). A peak in $\log(\text{Ba}/\text{Fe})$ between 7.6 and 7.3 Ma is less pronounced in the $[\text{Ba}]_{\text{xs}}$ record, but is prominent in the carbonate-free $[\text{Ba}]_{\text{xs}}$ record (Fig. S7, grey shading), suggesting that this peak is suppressed in the $[\text{Ba}]_{\text{xs}}$ record as a result of dilution by carbonate. The trough between 7.9 and 7.6 Ma seen in $\log(\text{Ba}/\text{Fe})$, $[\text{Ba}]_{\text{xs}}$ and to a lesser extent in % CaCO_3 , $\log(\text{Ca}/\text{Terr})$, and carbonate-free $[\text{Ba}]_{\text{xs}}$ appears not to be an artefact of dilution.

Bulk sediment MARs vary between 0.5 and 2.1 $\text{g}/\text{cm}^2/\text{kyr}$ with a step increase from ~ 0.5 to 1.5 $\text{g}/\text{cm}^2/\text{kyr}$ occurring at 8.66 Ma (Fig. 7g), concurrent with a major sedimentation rate increase (Fig. 4d). The stepwise nature of MAR records results from age model-imposed stepped changes in sedimentation rate. CaCO_3 MARs range from 0.4 to 2 $\text{g}/\text{cm}^2/\text{kyr}$ and covary with bulk sediment MARs, with the increasing difference between the two records reflecting a small long-term increase in non- CaCO_3 components (Fig. 7g). This small increase is reflected in terrigenous element MARs, which vary between ~ 20 and 80 $\text{mg}/\text{cm}^2/\text{kyr}$ (excluding volcanic ash layers) (Fig. 7i). We note that absolute values of terrigenous MAR should be interpreted with caution, because this calculation does not include Si as this element was not quantified in discrete samples. Nevertheless, a significant correlation between Al and Si counts ($R^2 = 0.8$, Fig. 6) suggests that Si is primarily of terrigenous origin, therefore trends in $\log(\text{Ca}/\text{Terr})$ and terrigenous MAR in Fig. 7 are likely robust despite the exclusion of Si. From 8.3 to 5 Ma, $[\text{Ba}]_{\text{xs}}$ MAR shows similar patterns to CaCO_3 MAR, with no clear long-term trend and maximum values driven by higher sedimentation rates in the intervals 5.2 to 5 Ma, 6.3 to 6.1 Ma, 7.7 to 7.5 Ma, and 8 to 7.8 Ma (Fig. 7f).

Spectral analyses reveal significant orbital periods in all late Miocene XRF records (Fig. 8). The ~ 405 kyr period is $>99\%$ significant in $[\text{Al}]$, $[\text{Ba}]$, $[\text{Ba}]_{\text{cfb}}$, $[\text{Ba}]_{\text{xs}}$, and $\log(\text{Ba}/\text{Fe})$, whereas the ~ 125 kyr period is significant ($>95\%$ or $>99\%$) in $[\text{Ba}]$, $[\text{Ba}]_{\text{cfb}}$, $[\text{Ba}]_{\text{xs}}$, $\log(\text{Ba}/\text{Fe})$, and % CaCO_3 records. At higher frequencies, the spectral signatures of $[\text{Fe}]$, $[\text{Al}]$, $[\text{Ti}]$, $[\text{Ba}]$, are dominated by significant peaks at 24 kyr ($>99\%$) and 41 kyr ($>90\%$), with $[\text{Ba}]$ additionally showing peaks at 22.5 kyr ($>99\%$) and at 26 and 30 kyr ($>95\%$). $\log(\text{Ba}/\text{Fe})$, $[\text{Ba}]_{\text{cfb}}$, and $[\text{Ba}]_{\text{xs}}$ show dominant ($>99\%$ significant) 22.5 kyr variability, with additional $>95\%$ significant peaks at 24 kyr and 30 kyr (for $[\text{Ba}]_{\text{xs}}$ only). $\log(\text{Ca}/\text{Terr})$ shows significant

peaks at 24 and 22.5 kyr (both >99%), and also at 68 kyr (>95%). Percent CaCO₃ contains significant (>95%) variability at 22.5 kyr, with additional peaks at 68 kyr and 37 kyr. In
505 summary, all records show highly significant variability in the precession band (22-24 kyr),
with variability at the 22.5 kyr period and the ~125 kyr period most strongly associated with
the biogenic component of Ba and with CaCO₃. Wavelet analyses of log(Ba/Fe) and [Ba]_{xs}
confirm significant precession-scale variability in these records throughout the 9 to 5 Ma
interval (Fig. 9).

510

5. Discussion

5.1 Late Miocene sedimentation patterns in the southern Bay of Bengal

We first examine the drivers of changes in sediment MAR identified in our record, and their
possible link to regional and global productivity trends. The three-fold increase in CaCO₃
515 MAR at 8.66 Ma at Site U1443, originally described in Lübbers et al. (2019), could result
from improved preservation and/or increased carbonate export by pelagic calcifiers
(coccolithophores and/or foraminifera). Based on CaCO₃ percentages, MARs, and benthic to
planktic foraminiferal ratios, Lübbers et al. (2019) identified the mid to late Miocene
“carbonate crash” in Site U1443 sediments between ~12.2 and 10 Ma, with a slow recovery
520 from ~10 to 8.7 Ma, favouring an interpretation that the increase in CaCO₃ MAR at 8.66 Ma
reflects improved preservation. A record of planktic foraminiferal fragmentation between 9
and 8 Ma generated in the present study, interpreted to reflect a decrease in carbonate
dissolution (Le and Shackleton, 1992), supports this interpretation (Fig. 7h). We see no
change in log(Ba/Fe) concurrent with the CaCO₃ MAR increase at 8.66 Ma, which suggests
525 that total export productivity at Site U1443 remained stable over this transition. However, our
data suggest that an increase in coccolithophore production may have occurred. The
contribution of foraminifera to total CaCO₃ over our study interval is low (see >63 μm MAR
in Fig. 7g), leading us to infer that higher CaCO₃ MARs between 8.66 and 5 Ma are primarily
driven by coccoliths. A 3-fold increase in sediment accumulation rate at ~8.6 Ma with no
530 change in CaCO₃ content (%), implying a large increase in CaCO₃ MARs, is also seen at
shallower (2247 m) Deep Sea Drilling Project (DSDP) Site 216 on the NER near the equator
(Fig. 1c) (Bukry, 1974; McNeill et al., 2017; Pimm, 1974). This suggests that production
played a role in driving regional carbonate MAR increases as well as improved preservation
at deeper sites. A recent study decoupling coccolith and foraminiferal MARs in relatively
535 shallow, globally-distributed sites (minimally affected by dissolution) records a late Miocene

pulse in coccolith MARs beginning at ~9 to 8 Ma and persisting into the Pliocene at ~4 to 3 Ma, which is interpreted to reflect high coccolithophore productivity and calcification driven by weathering alkalinity inputs and regional nutrient changes (Si and Rosenthal, 2019).

540 Interestingly, the increase in bulk MARs at 8.66 Ma is driven by both CaCO_3 and to a lesser extent non- CaCO_3 components (clays), implicating another mechanism as well as improved carbonate preservation and increased coccolith export productivity affecting sedimentation at Site U1443. Miocene Site U1443 clays are thought to be primarily supplied by the major rivers, with limited contribution of fine-grained mineral dust originating from the deserts
545 bordering the Arabian Sea (Bretschneider et al., 2021), suggesting that an increase in dust delivery at this time is an unlikely candidate for driving the ~50% increase in clay content. We speculate that an increase in coccolith CaCO_3 flux to the seafloor could have led to increased scavenging by sinking biogenic aggregates of fine clays. Fine clays are present in the southern BOB water column as a direct result of riverine flux (Rixen et al., 2019;
550 Ramaswamy, 1993), and in nepheloid layers above the NER where high clay concentrations occur due to proximity to the sedimentary fan systems to the east (Nicobar Fan) and west (Bengal Fan) (Stow et al., 1990). Recent studies of sedimentation patterns on the Bengal and Nicobar Fans, separated by the NER, interpret a large increase in sediment accumulation rate both on the NER and the Nicobar Fan at ~10 to 8 Ma to reflect increased lithogenic sediment
555 flux to the eastern Indian Ocean (McNeill et al., 2017; Pickering et al., 2020b). Our data from the NER show that >75% of the 3-fold increase in sediment accumulation rate at 8.66 Ma is driven by biogenic CaCO_3 , thus we caution against using sediment accumulation rate at Site U1443/758 as representative of changes in sediment flux to the Bengal-Nicobar Fan system. Data from Site U1443, as well as from nearby DSDP Site 216 (Bukry, 1974; McNeill et al.,
560 2017; Pimm, 1974) (Fig. 1c), suggest that increases in biogenic carbonate accumulation on the NER are decoupled, both temporally and mechanistically, from the increase in sediment delivery to the Nicobar Fan system. The gradual increase in terrigenous element and non- CaCO_3 MARs over the 9 to 5 Ma interval seen at Site U1443 (Fig. 7g,i) is part of a longer-term trend of increasing mineral flux in this region of the NER from the Miocene to the
565 Pleistocene, beginning at ~12 Ma at ODP Site 758, that is thought to reflect increased Himalayan erosion (Ali et al., 2021; Hovan and Rea, 1992).

Increases in the MAR of biogenic components (CaCO_3 , opal, organic carbon, phosphorus) between ~9 and 4 Ma have been measured in sediments from the Pacific, Indian and Atlantic

570 Oceans (Farrell et al., 1995; Lyle and Baldauf, 2015; Van Andel et al., 1975; Grant and
Dickens, 2002; Delaney and Filippelli, 1994; Hermoyian and Owen, 2001; Dickens and
Owen, 1999; Drury et al., 2020). This period of increased biogenic sedimentation, supported
by independent paleoproductivity proxies (e.g., Diester Haass et al., 2005), is thought to
reflect higher biological productivity and was dubbed the “biogenic bloom” by Farrell et al.
575 (1995). A low-resolution CaCO₃ MAR record from Site 758 shows higher values between 8
and 4 Ma, which in the absence of evidence for an increase in carbonate dissolution at 4 Ma,
could suggest an end to the biogenic bloom at this site in the early Pliocene (Dickens and
Owen, 1999; Pierce et al., 1989; Si and Rosenthal, 2019), although improved age control for
the Pliocene interval and independent paleoproductivity reconstructions are needed to verify
580 this. Hypotheses to explain the biogenic bloom invoke a change in global nutrient cycling;
i.e., a global increase in nutrient input, and/or redistribution of nutrients between basins
(Grant and Dickens, 2002), although the asynchronous timing of the biogenic bloom between
regions, its variable expression, and its differentiation from the carbonate crash recovery
complicate its interpretation. Diester Haass et al. (2006) hypothesised that changes in
585 reconstructed productivity were correlated to the LMCIS at four Indo-Pacific sites, and
tentatively proposed a link to a strengthened wind regime at this time. At Site U1443, we find
no clear link between export productivity or carbonate sedimentation and the LMCIS (Fig. 7).
In the northern Indian Ocean, the influence of possible concurrent changes in monsoon
strength on paleoproductivity and biogenic MARs must also be considered, and these are
590 discussed in Section 5.3.

5.2 Orbital forcing of late Miocene South Asian summer monsoon winds

On orbital timescales, time series analyses reveal dominant precession-band (22-24 kyr)
variance in late Miocene export productivity records (Fig. 8, 9). Spectral analyses of
595 individual calibrated timeseries of [Ba], [Al], [Fe], [Ti], [Ba]_{cfb} and %CaCO₃ allow us to
tease apart the effects of sediment dilution and the competing influence of Ba_{terr} and Ba_{bio} on
our Ba proxies, [Ba]_{xs} and log(Ba/Fe). The 41 kyr obliquity period is most significant (>90%)
in the terrigenous element records (Al, Fe, Ti, Ba), and absent or less significant in [Ba]_{xs},
log(Ba/Fe), log(Ca/Terr), and %CaCO₃ (Fig 8). The 24 kyr period stands out as highly
600 significant in all records (>99%, except for %CaCO₃ and [Ba]_{cfb} where >95%). In contrast,
the 125-kyr and 22.5 kyr peaks that dominates the [Ba]_{xs} and log(Ba/Fe) spectra (>99%
significant), are also highly significant only in the [Ba], [Ba]_{cfb}, %CaCO₃, and log(Ca/Terr)

records. This suggests that strong variability at the 125-kyr (eccentricity) and 22.5 kyr (precession) periods is related to biological productivity (i.e. Ba_{bio} and not Ba_{detrital} , as well as biogenic CaCO_3). The 30-kyr peak in $[Ba]_{\text{xs}}$ is also seen in $[Ba]$ but not in $[Fe]$, $[Al]$, or $[Ti]$, so we similarly interpret this period as being related to biological productivity. The 23.6-kyr and 22.3-kyr periods (highlighted together in Fig. 8 as one grey band spanning 22-24 kyr) are primary periods of Earth's precession, whereas the 53-kyr and 41-kyr periods are related to Earth's obliquity (Laskar et al., 2004).

610

Although our export productivity records are of sufficient resolution to detect it, half-precession (~ 11 kyr) cycles were not identified in spectral analyses. The presence of half-precession cycles might be expected because the bi-annual primary productivity peak observed today in southern BOB waters (Fig. 2) could fuel high productivity during both precession minima (strong summer winds) and precession maxima (strong winter winds). The lack of a half-precession signal could be explained by the distinct particle export seasonality, such that the fraction of net primary productivity exported from the photic zone and accumulating in underlying sediments is strongly biased towards the late summer months (Fig. 2d,e), perhaps as a result of increased ballasting by terrigenous particles carried into the BOB by summer monsoon runoff. Thus, the winter productivity maximum appears not to be efficiently transferred to the fossil record.

620

Significant variability at obliquity and precession periods has been identified in high-resolution late Miocene-early Pliocene records of precipitation/runoff based on planktic foraminiferal $\delta^{18}\text{O}$ and seawater $\delta^{18}\text{O}$ in the nearby Andaman Sea (Jöhnck et al., 2020). These authors suggest that, prior to a distinct switch to obliquity-driven variability around 5.55 Ma, their records reflect strong precession (insolation) control on South Asian monsoon rainfall from 6.2-5.55 Ma, with significant phase lags between proxy variations and precession. Wavelet analyses of $\log(\text{Ba}/\text{Fe})$ and $[Ba]_{\text{xs}}$ show that precession-band (22-24 kyr) variability dominated throughout our 9 to 5 Ma study interval at Site U1443 (Fig. 9). Although phase relationships with insolation should be interpreted with caution because of errors inherent to our late Miocene age model, export productivity appears to be coherent and in phase (within error) with the Summer Inter-Tropical Insolation Gradient, SITIG (the insolation gradient between 23°N and 23°S on June 21st) (Fig. S4d). SITIG has been proposed as a primary control on the strength of SASM winds, because a stronger SITIG increases the pressure gradient between the two limbs of the winter hemisphere Hadley cell, which drives monsoon

630

635

winds into the summer hemisphere (Bosmans et al., 2015). Our new Ba-based export productivity records corroborate the hypothesis that insolation played a dominant role in driving late Miocene South Asian summer monsoon wind variability in the equatorial sector of the SAM region, as predicted by general circulation models (Bosmans et al., 2018), with internal climate processes such as ice volume playing a more minor role than in the Late Pleistocene when large glacial-interglacial cycles and related feedbacks drove variability in the Asian monsoon on 100-kyr timescales (Clemens et al., 2018; Clemens et al., 2021) and SASM wind proxies from the Arabian Sea and southern BOB record up to ~9 kyr phase lags relative to precession (Bolton et al., 2013; Caley et al., 2011; Clemens and Prell, 2003; Clemens et al., 1991). A half-precession cycle related to equatorial insolation has been previously identified in Miocene and Pleistocene climate proxy records (Fox et al., 2017; Bolton et al., 2013; Sun and Huang, 2006). The lack of a half-precession signal in our records corroborates the idea that the SITIG, rather than local insolation (which contains a significant half-precession component between the equator and 5° latitude), was the primary driver of export productivity variations at our site.

In addition to the periods discussed above, a number of non-primary orbital periods termed heterodynes, which result from non-linear interactions between variables operating at Earth's primary orbital periods (Rial and Anaclerio, 2000; Thomas et al., 2016; Clemens et al., 2010), stand out in our late Miocene records (24, 26, 30, 37, 49, and 68 kyr periods; Fig. 8). For example, the 1/24 kyr heterodyne, prominent in all our records, could result from the interference between eccentricity and precession, and the 1/30 kyr heterodyne seen in Ba records from an interaction between obliquity and precession. Several of these heterodynes have been previously identified in spectra of seawater $\delta^{18}\text{O}$ that reflect Asian monsoon precipitation and runoff, both in the Andaman Sea (30 and 130 kyr during the Pleistocene, 27 and 30 kyr in the latest Miocene) (Gebregiorgis et al., 2018; Jöhnck et al., 2020) and in the East China Sea (29 and 69 kyr during the Pleistocene) (Clemens et al., 2018), suggesting high sensitivity of the monsoon to orbital forcing. We favour the interpretation that the prominent 24-kyr variability in our records reflects a primary period of precession, because precession filters of $[\text{Ba}]_{\text{xs}}$ and $\log(\text{Ba}/\text{Fe})$ spanning 18 to 26 kyr show strong amplitude modulation of the precession signal at a period of ~405 kyr, which results from the interaction of the 23.6-kyr and 22.3-kyr periods ($1/[(1/22.3) - (1/23.6)] = 404.8$ kyr) (Fig. 10). Amplitude modulation of precession-scale variability in our productivity records broadly follows that of

670 SITIG (Fig. 10), suggesting a direct response of SASM winds to cross-equatorial insolation gradients during the late Miocene.

Cross-spectral analysis of our $[Ba]_{xs}$ productivity record with the Site U1448 seawater $\delta^{18}O$ record (Jöhnck et al., 2020) over the interval 6.19 to 4.95 Ma (where records overlap) shows
675 >80% coherency and an in-phase relationship at the 30-kyr period, suggesting that monsoon winds and precipitation/runoff in the BOB were to some degree coupled on orbital timescales during this time (Fig. S4c), as is the case in the late Pleistocene (Clemens et al., 2021).
Nevertheless, our $[Ba]_{xs}$ record over this interval contains stronger primary precession (22-24 kyr) and obliquity (41 kyr) signals than the Site U1448 seawater $\delta^{18}O$ record, which cannot
680 be explained by differences in resolution, highlighting that different climatic processes and feedbacks operating on orbital timescales must contribute to the two records (interpreted to reflect runoff/precipitation and wind, respectively) to different extents. In the late Pleistocene, strong obliquity-band and precession-band variance is found in Andaman Sea proxy records of monsoon precipitation/runoff (Gebregiorgis et al., 2018), whereas records of upper ocean stratification controlled by South Asian monsoon wind mixing at Site 758 (~100 m from Site
685 U1443) show only precession-band variance (Bolton et al., 2013). The significant 41-kyr variability seen in late Miocene terrigenous elements at Site U1443 (Fig. 8a-c) could also suggest obliquity control on monsoon runoff into the BOB at this time. Clemens et al. (2021) show that 100 kyr and 41 kyr variability are at least as important as precession in Pleistocene
690 proxy records of monsoon precipitation/runoff in the Bay of Bengal, and suggest that summer monsoon precipitation is strongly influenced by global boundary conditions related to ice-volume and greenhouse gas feedbacks. On the other hand, obliquity forcing of tropical climate has been shown to occur independently of high-latitude ice-sheet growth and decay as a result of interhemispheric insolation gradients (Bosmans et al., 2015), **consistent with**
695 studies showing strong obliquity control on African monsoon runoff prior to the establishment of large northern hemisphere ice sheets (Zeeden et al., 2014; Lourens et al., 2001; Lourens et al., 1996). It is important to note that whilst SAM expression above Site U1443 in the southern BOB is dominated by summer monsoon winds that drive surface ocean currents and deeper mixing, oceanographic conditions in the northern and eastern BOB
700 (e.g., Sites U1447 and U1448) are instead primarily controlled by summer monsoon freshwater inputs (Jöhnck et al., 2020; Kuhnt et al., 2020). Runoff and direct precipitation during the SASM lead to strong salinity stratification in the northern parts of the BOB in the late summer and autumn, which prevents upper ocean mixing (e.g., Sengupta et al., 2016).

705 These regional differences in the manifestation of the monsoon must be considered when interpreting records from the heterogeneous BOB, and records from multiple locations and proxies are needed to achieve a comprehensive picture of the SAM subsystem.

710 The 405 kyr eccentricity modulation of precession-scale export productivity variability broadly coincides with 405-kyr cycles in benthic $\delta^{13}\text{C}$ at Site U1443, with higher export productivity during benthic $\delta^{13}\text{C}$ minima and eccentricity maxima on these timescales (Fig. 10). Cross-spectral analysis indicates that $\log(\text{Ba}/\text{Fe})$ and benthic $\delta^{13}\text{C}$ are > 95% coherent at the ~405 kyr and 22-24 kyr periods, with an in-phase relationship in the precession band ($-13^\circ \pm 25$) and a near antiphase relationship on 405-kyr timescales ($151^\circ \pm 27$) (Fig. S4b). 405-kyr modulation of the ocean carbon cycle, primarily recorded in carbonate content and
715 benthic $\delta^{13}\text{C}$ records (Herbert, 1997; Drury et al., 2020; De Vleeschouwer et al., 2020; Westerhold et al., 2020; Pälike et al., 2012; Paillard, 2017; Holbourn et al., 2007) but also in productivity and monsoon-related dust records (Rickaby et al., 2007; Wang et al., 2010), has been observed throughout the Cenozoic and Mesozoic sedimentary record. In middle
720 Miocene records, poor carbonate preservation noted during eccentricity maxima is interpreted in terms of transient shoaling of the carbonate saturation horizon (Holbourn et al., 2007; Flower and Kennett, 1994). Here, we see a broad positive correlation between $\log(\text{Ba}/\text{Fe})$ and $\log(\text{Ca}/\text{Terr})$ records on 405-kyr timescales (Figs. 9a,b), suggesting that carbonate content fluctuations at Site U1443 in the late Miocene were more strongly related to biogenic production than to dissolution on long eccentricity timescales. The coincidence of late
725 Miocene eccentricity maxima with productivity maxima and benthic $\delta^{13}\text{C}$ minima at Site U1443 is compatible with the hypothesis that a strengthened monsoon induced 405-kyr cycles in the marine carbon cycle via increased weathering and nutrient inputs during eccentricity maxima, leading to enhanced marine biological productivity and deep-ocean organic carbon burial (Ma et al., 2011).

730

5.3 Late Miocene monsoon evolution

On long timescales, our Site U1443 biogenic Ba records show relatively stable (9 to 6.5 Ma) or slightly decreasing (6.5-5.3 Ma) export productivity between 9 and 5 Ma (Fig. 7d-f, Fig. S7). Based on sediment colour properties and XRF-derived Ba/Ti ratios in the preceding
735 interval (~13.5-8.3 Ma), Lübbers et al. (2019) suggested that a shift towards a higher productivity regime occurred at ~11.2 Ma at Site U1443, 2.5 Ma before the rise in CaCO_3

MAR at Site U1443 and also significantly earlier than the onset of the biogenic bloom at other sites (Farrell et al., 1995; Grant and Dickens, 2002; Dickens and Owen, 1999; Diester Haass et al., 2005). Those authors also suggested that this shift was potentially linked to an intensification of the South Asian monsoon. An increase in export productivity at ~11 Ma is consistent with long-term changes in benthic foraminiferal assemblages at Site 758 (Gupta et al., 2004; Nomura, 1995) and at sites in the western tropical Indian Ocean (Smart et al., 2007), as well as with opal records from the more remote equatorial Pacific (Lyle and Baldauf, 2015). Reconstructions of Arabian Sea upwelling, export productivity, and deoxygenation (Bialik et al., 2020; Gupta et al., 2015; Huang et al., 2007; Zhuang et al., 2017), as well as the abrupt appearance of drift sediments in the Maldives Archipelago at ~13 Ma (Betzler et al., 2016), point towards an intensification of seasonally reversing South Asian monsoon winds between 13 and 11 Ma, consistent with Site U1443 export productivity records. Our new data suggest that similar levels of export productivity to those seen from 11.2 to 9 Ma persisted until at least 5 Ma at Site U1443.

Compiled Asian monsoon proxy records spanning the 9 to 5 Ma interval show relatively stable long-term SASM strength (Fig. 11, Fig. 1c for site map). Similar to records from the Maldives and Arabian Sea (Tripathi et al., 2017; Huang et al., 2007; Betzler et al., 2016; Zhuang et al., 2017), Site U1443 records do not corroborate the hypothesis that SASM winds intensified at ~8 to 7 Ma as suggested by some studies (Kroon et al., 1991; Singh and Gupta, 2014; Gupta et al., 2015; An et al., 2001).

The long-term trend in our record shows broad agreement with a low-resolution clay mineralogy record from Site U1447 in the Andaman Sea showing gradual long-term decrease in smectite/(illite and chlorite) over the late Miocene (Fig. 11g), indicating strengthened physical weathering and/or weakened chemical weathering that can be attributed to the South Asian winter and summer monsoons respectively (Lee et al., 2020). Also at Site U1447, records of potassium content (%K, Fig. 11x) are interpreted to show a shift in sediment provenance and/or an increase in physical weathering and erosion in the sediment source region between ~7 and ~6 Ma (Fig. 11h), which may be linked to an increase in monsoon rainfall intensity and global cooling (Kuhnt et al., 2020). Between 6.2 and 5 Ma, our equatorial Indian Ocean wind records show good long-term agreement with a seawater $\delta^{18}\text{O}$ record from the Andaman Sea (Jöhneck et al., 2020), with a minimum in export productivity at ~5.3 Ma at Site U1443 coinciding with a maximum in seawater $\delta^{18}\text{O}$ at Site U1448 (Fig. 11i,

j). One interpretation of this could be a coupled reduction in both SASM wind intensity and runoff/precipitation over this interval, although Jöhnck et al. (2020) invoke an increase in local evaporation and/or a change in precipitation source to explain the decreasing seawater $\delta^{18}\text{O}$ values between 5.6 and 5.2 Ma. High-resolution records of precipitation and runoff from the SASM region further back in time are needed to verify to what extent monsoon winds and precipitation/runoff are coupled on long timescales.

In contrast to the relatively stable South Asian summer monsoon between 9 and 5 Ma, evidence for a step strengthening of the East Asian winter monsoon during the late Miocene (~7 Ma) comes from the South China Sea (Holbourn et al., 2018; Wan et al., 2007) (Fig. 11a), whereas records from a site on the Chinese Loess Plateau suggest a more gradual intensification of the East Asian summer monsoon from 8.2 to 2.6 Ma (Ao et al., 2016) (Fig. 11b). The step change in South China Sea surface water geochemistry has been interpreted to reflect drying and cooling of the Asian interior and a related southward shift of the ITCZ leading to an intensified dry winter monsoon over southeast Asia (Holbourn et al., 2018). An intensification of the East Asian winter monsoon around 7 Ma is consistent with an increase in aeolian dust delivery to the South China Sea at this time (Wan et al., 2007). The long-term increase in East Asian summer monsoon strength inferred from magnetic records in Chinese loess sequences is attributed to progressive Antarctic glaciation that drives an increased pressure gradient between the Australian High and Asian Low pressure cells, a mechanism supported by numerical simulations (Ao et al., 2016). However, a late Miocene intensification of the East Asian summer monsoon is not corroborated by studies from the South China Sea (Wan et al., 2007; Holbourn et al., 2021; Holbourn et al., 2018). The apparent insensitivity of equatorial SASM wind intensity to global late Miocene sea surface cooling, which began at ~7.5 Ma and culminated in an SST minimum at ~6 Ma (Fig. 11k) (Herbert et al., 2016), is consistent with climate modelling studies that show limited impact of different $p\text{CO}_2$ scenarios on SASM wind patterns and strength (Kitoh et al., 1997). The South Asian summer monsoon has been described as a thermally direct circulation driven by the thermal contrast between the Indian subcontinent and the equatorial Indian Ocean that develops in summer, with abrupt transitions between summer and winter monsoon regimes suggested to be related to feedbacks between extratropical eddies and tropical circulation (Privé and Plumb, 2007a, b; Lutsko et al., 2019; Bordoni and Schneider, 2008; Geen et al., 2018). Changes in the cross-equatorial temperature gradient have been shown to substantially impact the strength of onshore SASM monsoon flow (Lutsko et al., 2019; Acosta and Huber,

805 2020). The lack of a long-term trend in wind and surface circulation proxies over the 9 to 5
Ma interval (Fig. 11) suggests a relatively constant land-sea temperature gradient despite
global cooling. Thus, although single site records may not be representative of the East or
South Asian monsoon subsystems as a whole our data add to the body of evidence suggesting
decoupling between the East Asian and South Asian monsoons on long timescales.

810

6. Summary & Conclusions

We present new equatorial Indian Ocean benthic $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ records and an age model
spanning the interval between 9 and 5 Ma (late Miocene-earliest Pliocene), and analyse
sedimentation and productivity trends and cyclicity using XRF-derived records and MARs.
815 Biogenic sediment MARs reveal a modest imprint of the late Miocene biogenic bloom at Site
U1443 lasting until at least 5 Ma, primarily driven by fine-fraction (coccolith) CaCO_3
accumulation, as noted at other sites (Si and Rosenthal, 2019). Nevertheless, the carbonate
MAR record of Site U1443 is clearly influenced by carbonate preservation as well as
production over the late Miocene, so independent productivity proxies must be considered
820 when defining the duration of the biogenic bloom. Our results indicate that step increases in
sediment accumulation rate recorded on the Ninetyeast Ridge at ~9-8 Ma primarily reflect an
increase in CaCO_3 accumulation, and that this is likely independent from the increase in
lithogenic sediment flux recorded in nearby Nicobar Fan sites, itself related to sediment re-
routing within the Nicobar-Bengal Fan system around the same time (McNeill et al., 2017;
825 Pickering et al., 2020a). Our data show no long-term increase in export productivity between
9 and 5 Ma (and by inference no intensification of SASM winds), therefore these data
support existing evidence for an early late Miocene (~13-10 Ma) establishment of strong
seasonally reversing South Asian monsoon winds and Arabian Sea upwelling, with relatively
stable or slightly weakening SASM winds over the remainder of the late Miocene and earliest
830 Pliocene between 9 and 5 Ma. Spectral and cross-spectral analyses of XRF-based biogenic
barium records reveal that export productivity in waters overlying Site U1443 was
consistently paced by precession, with amplitude modulation of the precession signal on
~405 kyr timescales and no significant variability at glacial-interglacial (obliquity)
timescales. Coeval late Miocene productivity maxima and benthic $\delta^{13}\text{C}$ minima during
835 eccentricity maxima at Site U1443 provides support for the hypothesis that the monsoon may
have paced changes in the carbon cycle on ~405 kyr timescales (Ma et al., 2011). Significant
coherence and an in-phase relationship at the precession band between biogenic barium and

the SITIG suggests direct forcing of South Asian monsoon winds by insolation gradients over the late Miocene, relatively unaffected by glacial boundary conditions and long-term global cooling trends (Herbert et al., 2016). In contrast, East Asian summer and winter monsoons appear to have intensified during the late Miocene in response to global cooling and Antarctic ice sheet growth and related feedbacks (Ao et al., 2016; Holbourn et al., 2018), although more continuous records over the late Miocene are needed to understand regional trends due to the heterogeneous nature of Asian monsoon expression.

845

References

Data Availability

All data is available on the PANGAEA database at
850 <https://doi.org/10.1594/PANGAEA.935166>

Author contributions

C.T.B. designed the study. Sample processing and picking of benthic foraminifera was carried out by E.G and C.T.B. Picked foraminifera samples were verified and cleaned by A.H., W.K., and J.L., and stable isotopes measurements were performed by N.A., A.H.,
855 W.K., and J.L. XRF scanning was carried out by C.T.B. in collaboration with K.G., G.M., and E.J.R. XRF calibration was carried out by E.G. and K.T. The manuscript was written by C.T.B with feedback from all authors.

Acknowledgements

860 This research used samples and data provided by the International Ocean Discovery Program (IODP). We thank the science party, technical staff and crew of IODP Expedition 353. Funding for this research was provided by French ANR project iMonsoon ANR-16-CE01-0004-01, IODP France, and Deutsche Forschungsgemeinschaft (grant Ku649/36-1). KMG is supported by Australian Research Council grant DE190100042. CTB thanks Anna Joy Drury
865 and Tim Herbert for age model feedback, and Luc Beaufort, Baptiste Suchéras-Marx, and Ian Bailey for discussions that helped improve the manuscript. Marta Garcia Molina, Jean-Charles Mazur, and Christine Pailles are thanked for technical laboratory support at CEREGE.

870 **Figures**






Figure 1: Seasonal contrast (February vs August) in wind stress (a) and mixed layer depth (b). Yellow dots indicate the modern position of IODP Site U1443, and yellow stars its paleo-position at 9 Ma (1.71°N, 88.06°E) and 5 Ma (3.27°N, 89.04°E) calculated using G-Plates: http://portal.gplates.org/service/reconstruct_points). Maps were created on the ERDDAP website using the datasets *Wind Stress, Metop-A ASCAT, 0.25°, Global, Near Real Time, 2009-present (Monthly)* and *Ocean Climatology Ocean Mixed Layer Depth MLD T02 kriging* (see methods for details). c: Regional bathymetric map showing modern locations of marine and terrestrial sites discussed in this study (white squares) and the position of Site U1443, a redrill of Site 758, in the modern ocean (yellow square) and its paleo-position at 9 Ma and 5 Ma (yellow stars). The red circle shows the location of the SBBT sediment trap. Green arrows show surface ocean circulation during the summer monsoon (July/August) and the eastward flow of waters from the Arabian Sea into the BOB via the Southwest Monsoon Current (SMC), after Schott et al. (2009).



890

Figure 2: Modern, seasonal oceanographic variability above Site U1443 in the southern BOB. **a:** wind stress, **b:** mixed layer depth (MLD), **c:** net primary productivity (NPP), **d:** particulate organic carbon (POC) flux, **e:** biogenic silica (bSi) and calcium carbonate (CaCO₃) flux. See Section 3.1 for details of individual datasets and sources. Points represent individual months, diamonds and triangles with lines represent monthly mean values over entire time series. Months (x-axis) run from 1 (1st January) to 12 (1st December). JJA = June, July August, DJF = December, January, February.

895



900

Figure 3: U1443 benthic foraminiferal (*Cibicoides wuellerstorfi*) $\delta^{18}\text{O}$ (bottom) and $\delta^{13}\text{C}$ (top) data on the composite depth scale. Blue squares show refined depth ranges for calcareous nannofossil datums (see Table S2), and shipboard magnetostratigraphy is also shown for the interval over which it could be reliably determined. Between ~90 and 128 m CCSF, sediments in cores from all holes showed scattered directional signals during pass-through magnetic remanence measurement, which hindered any determination of polarity patterns in this interval across the whole site (Clemens et al., 2016). Black and white zones = normal and reversed polarity, respectively; grey zones = magnetic polarity not clearly determined. All numbers are age assignments for boundaries/nannofossil events in Ma (Gradstein et al., 2012).

905

910






Figure 4: Astronomical (minimal) tuning of Site U1443 benthic $\delta^{18}\text{O}$ record to ET target. **a:** benthic $\delta^{18}\text{O}$ record on depth scale with nannofossil-based age constraints, **b:** ET tuning target (1:1 weighting, normalised). ETP (1:0.5:-0.4 weighting, normalised) is also shown in grey. Astronomical time series from (Laskar et al., 2004). **c:** tuned benthic $\delta^{18}\text{O}$ vs age, **d:** sedimentation rates, **e:** tuned benthic $\delta^{13}\text{C}$ vs age. Tie-points between **a** and **b** are shown in orange (Table S3). An age-depth plot showing ET tie-points and good agreement with biostratigraphic age control is shown in Fig S1. **f** and **g:** wavelet analyses of tuned isotope records; white shaded area shows cone of influence and contours show 95% significance level. Main orbital periods are shown on the right in kyr. **h:** Filtered tuned benthic isotope records compared to filtered ET (as in **b**). Top: 100-kyr filtered benthic $\delta^{13}\text{C}$ (blue; Gaussian filter centred on 100 kyr with bandwidth ± 25 kyr to include 95 and 125 kyr peaks) compared with filtered ET (grey, identical filter design). Bottom: 41-kyr filtered benthic $\delta^{18}\text{O}$ (blue, Gaussian filter centred on 41.5 kyr with bandwidth ± 1.5 kyr) compared with filtered ET (grey, identical filter design).




930 **Figure 5:** Late Miocene evolution of low-latitude deep-ocean inter-basin benthic $\delta^{13}\text{C}$ and
 $\delta^{18}\text{O}$ gradients. South China Sea ODP Site 1146 (Holbourn et al., 2018) with age model
revised in Holbourn et al. (2021), equatorial Pacific IODP Sites U1338 (Drury et al., 2018;
935 Drury et al., 2016) and U1337 (Drury et al., 2017), and ODP Site 926 (Shackleton and Hall,
1997; Drury et al., 2017; Zeeden et al., 2013). All records are shown on their latest
independent orbitally tuned chronologies. We have excluded Caribbean ODP Site 999 from
this figure because it is bathed in intermediate water masses due to basin geometry and sill
depths (Bickert et al., 2004). Deep South Atlantic Site 704 (Müller et al., 1991) data are not
plotted due to clear age model discrepancies when compared to orbitally tuned records. All
 $\delta^{18}\text{O}$ records are based on *Cibicidoides wuellerstorfi* or *C. mundulus* therefore no corrections
940 are applied, following (Jöhneck et al., 2021). The Site 926 record includes $\delta^{13}\text{C}$ corrections for
some samples due to the multispecific nature of the record (Drury et al., 2017). No correction
was applied to *C. wuellerstorfi* or *C. mundulus* $\delta^{13}\text{C}$ values.



945

Figure 6: Raw counts/sec (grey lines) and calibrated concentration (blue lines) scanning XRF elemental data over the late Miocene interval. Pink circles show samples used for calibration (See Fig. S3).



950

Figure 7: XRF-derived late Miocene CaCO_3 and export productivity records from Site U1443. **a:** benthic $\delta^{13}\text{C}$, **b:** percent CaCO_3 , **c:** $\log(\text{Ca}/\text{Terr})$, **d:** $\log(\text{Ba}/\text{Fe})$, **e:** Ba_{XS} calculated with both $[\text{Al}]$ (light green) and $[\text{Ti}]$ (dark green), **f:** $[\text{Ba}]_{\text{XS}}$ MAR, **g:** bulk (black), CaCO_3 (blue), and $>63\mu\text{m}$ fraction (light blue) MAR, **h:** foraminiferal fragmentation index (Le and Shackleton, 1992), **i:** Terrigenous ($\text{Al}+\text{Fe}+\text{Ti}+\text{K}+\text{Rb}$) MAR. For CaCO_3 records, lighter blue lines are based on % CaCO_3 from Lübbers et al. (2019) and darker blue lines are based on % CaCO_3 estimates in this study. The late Miocene carbon isotope excursion and the main interval of late Miocene SST cooling are shown as grey bars. Shaded intervals are identified ash layers.

960





Figure 8: Spectral analyses for over the 9 to 5 Ma interval (log ratios and % CaCO₃) or 8.15 to 5 Ma interval (calibrated element concentrations and [Ba]_{xs}). **a:** [Ti], **b:** [Fe], **c:** [Al], **d:** [Ba], **e:** [Ba]_{cfb} (carbonate free basis), **f:** [Ba]_{xs}, **g:** log(Ba/Fe), **h:** log(Ca/Terr), **i:** % CaCO₃. Grey bands denote primary orbital periods based on the La04 astronomical solution. The 22-24 kyr band covers two peaks centred at 23.5 and 22.3 kyr.



970

Figure 9: Orbital-scale variability of productivity and CaCO_3 proxies at Site U1443. All records shown here are bandpassed as described in the methods. (a) benthic $\delta^{18}\text{O}$, (b) benthic $\delta^{13}\text{C}$, (c) $\log(\text{Ca}/\text{Terr})$, (d) $\log(\text{Ba}/\text{Fe})$, (e) $[\text{Ba}]_{\text{xs}}$, (f) ETP. For c-e, wavelet analyses are shown, illustrating dominant precession-scale (22-24 kyr) variability in Ba proxies and both precession and obliquity (41-kyr) variability in $\log(\text{Ca}/\text{Terr})$.

975





Figure 10: Detrended and filtered late Miocene U1443 records to illustrate precession-band variance and amplitude modulation. **a:** eccentricity (Laskar et al., 2004), **b:** benthic $\delta^{13}\text{C}$ (bandpassed as in Fig. 9) and its 405-kyr filter (above) (note reversed y-axes). **c:** Lowess-detrended $\log(\text{Ba}/\text{Fe})$ (window = 0.1 Ma) and 18 to 26 kyr filter (above), **d:** Lowess-detrended Ba_{xs} (window = 0.1 Ma) and its 18 to 26 kyr filter (above), **e:** The summer inter-tropical insolation gradient (SITIG, calculated as the insolation difference between 23°N and 23°S on 21st June using orbital solution of Laskar et al. (2004)) and its 18 to 26 kyr filter and amplitude modulation (above); For panels **c-e**, raw datasets were filtered using a Tanner-Hilbert filter centred on 46.5 cycles/Myr with bandwidth ± 8 (designed to include all precession terms with periods between 18 and 26 kyr). For panel **b**, a Tanner-Hilbert filter centred on 2.47 cycles/myr with bandwidth ± 0.8 was applied.



990

Figure 11: Compilation of late Miocene (9-5 Ma) Asian monsoon reconstructions, showing representative records from different regions and their published interpretations (bars on left).

995

a: Seawater $\delta^{18}\text{O}$ showing an increase in East Asian winter monsoon strength at ~ 7.4 Ma (Holbourn et al., 2018; Holbourn et al., 2021), **b:** Stacked magnetic records of the East Asian summer monsoon from the Chinese Loess Plateau (Ao et al., 2016), **c:** Mn/Ca ratios used to trace oxygen minimum zone (OMZ) variations from Maldives Site IODP U1471 (Betzler et al., 2016), **d:** $\delta^{15}\text{N}$ record from Arabian Sea site IODP U1456 showing OMZ intensity (Tripathi et al., 2017), **e:** total organic carbon (TOC) % from Arabian Sea sites IODP U1456 and ODP Site 722 (Huang et al., 2007; Tripathi et al., 2017), **f:** % *Gobigerinoides bulloides*, a planktic foraminiferal upwelling indicator, at ODP Site 722 (Huang et al., 2007; Kroon et al., 1991), *G. bulloides* was counted in the $>150 \mu\text{m}$ fraction in Huang et al. (white diamonds) and the $>125 \mu\text{m}$ fraction in Kroon et al. (grey circles), **g:** clay mineralogy (smectite/(illite + chlorite)) at IODP Site U1447 in the Andaman Sea (Lee et al., 2020), **h:** Percentage

1000

Potassium (K%) at IODP Site U1447 derived from spectral natural gamma ray measurements (Kuhnt et al., 2020), **i**: Andaman Sea IODP Site U1448 ice-volume-corrected seawater $\delta^{18}\text{O}$ record (Jöhnck et al., 2020), and **j**: Export productivity records from Site U1443 (this study). **k**: Global sea surface temperature trends (expressed as anomalies relative to the present), stacked by latitude band (Herbert et al., 2016). All records are on their original age models, and Loess smooths are shown for high-resolution records.

1010

References

- 1015 Acosta, R. and Huber, M.: Competing topographic mechanisms for the summer Indo-Asian monsoon, *Geophysical Research Letters*, 47, e2019GL085112, 2020.
- Ali, S., Hathorne, E., and Frank, M.: Persistent provenance of South Asian Monsoon induced silicate weathering over the past 27 million years, *Paleoceanography and Paleoclimatology*, accepted, 2021.
- 1020 An, Z., Kutzbach, J. E., Prell, W. L., and Porter, S. C.: Evolution of Asian monsoons and phased uplift of the Himalaya–Tibetan plateau since Late Miocene times, *nature*, 411, 62-66, 2001.
- An, Z., Clemens, S. C., Shen, J., Qiang, X., Jin, Z., Sun, Y., Prell, W. L., Luo, J., Wang, S., and Xu, H.: Glacial-interglacial Indian summer monsoon dynamics, *science*, 333, 719-723, 1025 2011.
- Ao, H., Roberts, A. P., Dekkers, M. J., Liu, X., Rohling, E. J., Shi, Z., An, Z., and Zhao, X.: Late Miocene–Pliocene Asian monsoon intensification linked to Antarctic ice-sheet growth, *Earth and Planetary Science Letters*, 444, 75-87, 2016.
- Averyt, K. B. and Paytan, A.: A comparison of multiple proxies for export production in the equatorial Pacific, *Paleoceanography*, 19, 2004. 1030
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnology and oceanography*, 42, 1-20, 1997.
- Behrenfeld, M. J., Boss, E., Siegel, D. A., and Shea, D. M.: Carbon-based ocean productivity and phytoplankton physiology from space, *Global biogeochemical cycles*, 19, 2005.
- 1035 Betzler, C., Eberli, G., Lüdmann, T., Reolid, J., Kroon, D., Reijmer, J., Swart, P., Wright, J., Young, J., and Alvarez-Zarikian, C.: Refinement of Miocene sea level and monsoon events from the sedimentary archive of the Maldives (Indian Ocean), *Progress in Earth and Planetary Science*, 5, 1-18, 2018.

- Betzler, C., Eberli, G. P., Kroon, D., Wright, J. D., Swart, P. K., Nath, B. N., Alvarez-Zarikian, C. A., Alonso-García, M., Bialik, O. M., and Blättler, C. L.: The abrupt onset of the modern South Asian Monsoon winds, *Scientific reports*, 6, 29838, 2016.
- Bialik, O. M., Auer, G., Ogawa, N. O., Kroon, D., Waldmann, N. D., and Ohkouchi, N.: Monsoons, upwelling, and the deoxygenation of the northwestern Indian Ocean in response to middle to late Miocene global climatic shifts, *Paleoceanography and Paleoclimatology*, 35, 2020.
- Bickert, T., Haug, G., and Tiedemann, R.: Late Neogene benthic stable isotope record of ODP Site 999: Implications for Caribbean paleoceanography, organic carbon burial and the Messinian Salinity Crisis, *Paleoceanography*, 19, PA1023, 2004.
- Bishop, J. K.: The barite-opal-organic carbon association in oceanic particulate matter, *Nature*, 332, 341-343, 1988.
- Bolton, C. T., Chang, L., Clemens, S. C., Kodama, K., Ikehara, M., Medina-Elizalde, M., Paterson, G. A., Roberts, A. P., Rohling, E. J., and Yamamoto, Y.: A 500,000 year record of Indian summer monsoon dynamics recorded by eastern equatorial Indian Ocean upper water-column structure, *Quaternary Science Reviews*, 77, 167-180, 10.1016/j.quascirev.2013.07.031, 2013.
- Boos, W. R. and Kuang, Z.: Dominant control of the South Asian monsoon by orographic insulation versus plateau heating, *Nature*, 463, 218-222, 2010.
- Bordoni, S. and Schneider, T.: Monsoons as eddy-mediated regime transitions of the tropical overturning circulation, *Nature Geoscience*, 1, 515-519, 2008.
- Bosmans, J., Hilgen, F., Tuenter, E., and Lourens, L.: Obliquity forcing of low-latitude climate, *Climate of the Past*, 11, 1335-1346, 2015.
- Bosmans, J., Erb, M., Dolan, A., Drijfhout, S., Tuenter, E., Hilgen, F., Edge, D., Pope, J. O., and Lourens, L.: Response of the Asian summer monsoons to idealized precession and obliquity forcing in a set of GCMs, *Quaternary Science Reviews*, 188, 121-135, 2018.
- Bretschneider, L., Hathorne, E. C., Bolton, C. T., Gebregiorgis, D., Giosan, L., Gray, E., Huang, H., Holbourn, A., Kuhnt, W., and Frank, M.: Enhanced late Miocene chemical weathering and altered precipitation patterns in the watersheds of the Bay of Bengal recorded by detrital clay radiogenic isotopes, *Paleoceanography and Paleoclimatology*, e2021PA004252, 2021.
- Bukry, D.: Coccolith and silicoflagellate stratigraphy, eastern Indian Ocean, Deep Sea Drilling Project Leg 22 in: Initial Reports of the Deep Sea Drilling Project, edited by: von der Borch, C. C., Sclater, J. G., and al., e., Washington, 10.2973/dsdp.proc.22.127.1974, 1974.

- Caley, T., Malaizé, B., Zaragosi, S., Rossignol, L., Bourget, J., Eynaud, F., Martinez, P., Giraudeau, J., Charlier, K., and Ellouz-Zimmermann, N.: New Arabian Sea records help
1075 decipher orbital timing of Indo-Asian monsoon, *Earth and Planetary Science Letters*, 308, 433-444, 10.1016/j.epsl.2011.06.019, 2011.
- Cao, Z., Siebert, C., Hathorne, E. C., Dai, M., and Frank, M.: Constraining the oceanic barium cycle with stable barium isotopes, *Earth and Planetary Science Letters*, 434, 1-9, 2016.
- 1080 Clemens, S. C. and Prell, W. L.: Late Pleistocene variability of Arabian Sea summer monsoon winds and continental aridity: Eolian records from the lithogenic component of deep-sea sediments, *Paleoceanography*, 5, 109-145, 1990.
- Clemens, S. C. and Prell, W. L.: A 350,000 year summer-monsoon multi-proxy stack from the Owen Ridge, Northern Arabian Sea, *Marine Geology*, 201, 35-51, 10.1016/S0025-
1085 3227(03)00207-X, 2003.
- Clemens, S. C., Prell, W. L., and Sun, Y.: Orbital-scale timing and mechanisms driving Late Pleistocene Indo-Asian summer monsoons: Reinterpreting cave speleothem $\delta^{18}\text{O}$, *Paleoceanography*, 25, 2010.
- Clemens, S. C., Prell, W., Murray, D., Shimmield, G., and Weedon, G.: Forcing mechanisms
1090 of the Indian Ocean monsoon, *Nature*, 353, 720-725, DOI 10.1038/353720a0, 1991.
- Clemens, S. C., Holbourn, A., Kubota, Y., Lee, K., Liu, Z., Chen, G., Nelson, A., and Fox-Kemper, B.: Precession-band variance missing from East Asian monsoon runoff, *Nature communications*, 9, 1-12, 2018.
- Clemens, S. C., Yamamoto, M., Thirumalai, K., Giosan, L., Richey, J. N., Nilsson-Kerr, K.,
1095 Rosenthal, Y., Anand, P., and McGrath, S. M.: Remote and local drivers of Pleistocene South Asian summer monsoon precipitation: A test for future predictions, *Science Advances*, 7, eabg3848, 2021.
- Clemens, S. C., Kuhnt, W., LeVay, L. J., Anand, P., Ando, T., Bartol, M., Bolton, C. T., Ding, X., Gariboldi, K., Giosan, L., Hathorne, E., Huang, Y., Jaiswal, P., Kim, S.,
1100 Kirkpatrick, J. B., Littler, K., Marino, G., Martinez, P., Naik, D., Peketi, A., Phillips, S. C., Robinson, M. M., Romero, O. E., Sagar, N., Taladay, K., Taylor, S. N., Thirumalai, K., Uramoto, G., Y. Usui, Wang, J., Yamamoto, M., and Zhou, L.: Site U1443, in: *Indian Monsoon Rainfall. Proceedings of the International Ocean Discovery Program*, edited by: Clemens, S. C., Kuhnt, W., LeVay, L.J., and the Expedition 353 Scientists, College Station,
1105 TX (International Ocean Discovery Program),
<http://dx.doi.org/10.14379/iodp.proc.353.103.2016>, 2016.

- Clift, P. D. and Webb, A. A. G.: A history of the Asian monsoon and its interactions with solid Earth tectonics in Cenozoic South Asia, Geological Society, London, Special Publications, 483, 631-652, 2019.
- 1110 Clift, P. D., Hodges, K. V., Heslop, D., Hannigan, R., Van Long, H., and Calves, G.: Correlation of Himalayan exhumation rates and Asian monsoon intensity, *Nature Geoscience*, 1, 875-880, 2008.
- De Vleeschouwer, D., Drury, A. J., Vahlenkamp, M., Rochholz, F., Liebrand, D., and Pälike, H.: High-latitude biomes and rock weathering mediate climate–carbon cycle feedbacks on eccentricity timescales, *Nature communications*, 11, 1-10, 2020.
- 1115 Dehairs, F., Chesselet, R., and Jedwab, J.: Discrete suspended particles of barite and the barium cycle in the open ocean, *Earth and Planetary Science Letters*, 49, 528-550, 1980.
- Delaney, M. L. and Filippelli, G. M.: An apparent contradiction in the role of phosphorus in Cenozoic chemical mass balances for the world ocean, *Paleoceanography*, 9, 513-527, 1994.
- 1120 Dickens, G. R. and Owen, R. M.: The latest Miocene–early Pliocene biogenic bloom: a revised Indian Ocean perspective, *Marine Geology*, 161, 75-91, 1999.
- Diester Haass, L., Billups, K., and Emeis, K. C.: In search of the late Miocene–early Pliocene “biogenic bloom” in the Atlantic Ocean (Ocean Drilling Program Sites 982, 925, and 1088), *Paleoceanography*, 20, 2005.
- 1125 Diester Haass, L., Billups, K., and Emeis, K. C.: Late Miocene carbon isotope records and marine biological productivity: Was there a (dusty) link?, *Paleoceanography*, 21, 2006.
- Ding, Z., Huang, G., Liu, F., Wu, R., and Wang, P.: Responses of global monsoon and seasonal cycle of precipitation to precession and obliquity forcing, *Climate Dynamics*, 56, 3733-3747, 2021.
- 1130 Drury, A. J., John, C. M., and Shevenell, A. E.: Evaluating climatic response to external radiative forcing during the late Miocene to early Pliocene: New perspectives from eastern equatorial Pacific (IODP U1338) and North Atlantic (ODP 982) locations, *Paleoceanography*, 31, 167-184, 2016.
- Drury, A. J., Lee, G., Gray, W., Lyle, M., Westerhold, T., Shevenell, A. E., and John, C.:
- 1135 Deciphering the state of the late Miocene to early Pliocene equatorial Pacific, *Paleoceanography and Paleoclimatology*, 33, 246-263, 2018.
- Drury, A. J., Liebrand, D., Westerhold, T., Beddow, H. M., Hodell, D. A., Rohlfs, N., Wilkens, R. H., Lyle, M., Bell, D. B., and Kroon, D.: Climate, cryosphere and carbon cycle controls on Southeast Atlantic orbital-scale carbonate deposition since the Oligocene (30–0
- 1140 Ma), *Climate of the Past Discussions*, 1-35, 2020.

- Drury, A. J., Westerhold, T., Frederichs, T., Tian, J., Wilkens, R., Channell, J. E., Evans, H., John, C. M., Lyle, M., and Röhl, U.: Late Miocene climate and time scale reconciliation: Accurate orbital calibration from a deep-sea perspective, *Earth and Planetary Science Letters*, 475, 254-266, 2017.
- 1145 Dymond, J., Suess, E., and Lyle, M.: Barium in deep-sea sediment: A geochemical proxy for paleoproductivity, *Paleoceanography*, 7, 163-181, 1992.
- Eagle, M., Paytan, A., Arrigo, K. R., van Dijken, G., and Murray, R. W.: A comparison between excess barium and barite as indicators of carbon export, *Paleoceanography*, 18, 2003.
- 1150 ERD, N. N. S.: Primary Productivity, Aqua MODIS, NPP, Global, 2003-present, EXPERIMENTAL (Monthly Composite), erdMH1ppmday, 2020.
- Farnsworth, A., Lunt, D. J., Robinson, S. A., Valdes, P. J., Roberts, W. H., Clift, P. D., Markwick, P., Su, T., Wrobel, N., and Bragg, F.: Past East Asian monsoon evolution controlled by paleogeography, not CO₂, *Science advances*, 5, eaax1697, 2019.
- 1155 Farrell, J. W., Raffi, I., Janecek, T. R., Murray, D. W., Levitan, M., Dadey, K. A., Emeis, K.-C., Lyle, M., Flores, J.-A., and Hovan, S.: Late Neogene sedimentation patterns in the eastern equatorial Pacific Ocean, in: *Proceedings of the Ocean Drilling Program Scientific Results*, edited by: Pisias, N. G., Mayer, L. A., Janecek, T. R., Palmer-Julson, A., and van Andel, T. H., 1995.
- 1160 Flower, B. P. and Kennett, J. P.: The middle Miocene climatic transition: East Antarctic ice sheet development, deep ocean circulation and global carbon cycling, *Palaeogeography, palaeoclimatology, palaeoecology*, 108, 537-555, 1994.
- Fox, B. R., D'Andrea, W., Wilson, G., Lee, D., and Wartho, J.-A.: Interaction of polar and tropical influences in the mid-latitudes of the Southern Hemisphere during the Mi-1 deglaciation, *Global and Planetary Change*, 155, 109-120, 2017.
- 1165 Francois, R., Honjo, S., Manganini, S. J., and Ravizza, G. E.: Biogenic barium fluxes to the deep sea: Implications for paleoproductivity reconstruction, *Global Biogeochemical Cycles*, 9, 289-303, 1995.
- Gadgil, S.: The Indian monsoon and its variability, *Annual Review of Earth and Planetary Sciences*, 31, 429-467, 2003.
- 1170 Gebregiorgis, D., Hathorne, E. C., Giosan, L., Clemens, S., Nürnberg, D., and Frank, M.: Southern Hemisphere forcing of South Asian monsoon precipitation over the past ~1 million years, *Nature Communications*, 9, 4702, 10.1038/s41467-018-07076-2, 2018.

- 1175 Geen, R., Lambert, F., and Vallis, G.: Regime change behavior during Asian monsoon onset, *Journal of Climate*, 31, 3327-3348, 2018.
- Gingele, F. and Dahmke, A.: Discrete barite particles and barium as tracers of paleoproductivity in South Atlantic sediments, *Paleoceanography*, 9, 151-168, 1994.
- Goldberg, E. D. and Arrhenius, G.: Chemistry of Pacific pelagic sediments, *Geochimica et cosmochimica acta*, 13, 153-212, 1958.
- 1180 Gonneea, M. E. and Paytan, A.: Phase associations of barium in marine sediments, *Marine Chemistry*, 100, 124-135, 2006.
- Goswami, B., Krishnamurthy, V., and Annmalai, H.: A broad-scale circulation index for the interannual variability of the Indian summer monsoon, *Quarterly Journal of the Royal Meteorological Society*, 125, 611-633, 1999.
- 1185 Gouhier, T. C., Grinsted, A., and Simko, V.: Package 'biwavelet' [code], 2016.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. B., and Ogg, G. M.: *The geologic time scale 2012*, elsevier2012.
- Grant, K. M. and Dickens, G. R.: Coupled productivity and carbon isotope records in the southwest Pacific Ocean during the late Miocene–early Pliocene biogenic bloom, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 187, 61-82, 2002.
- 1190 Grinsted, A., Moore, J. C., and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, 2004.
- Guo, Z., Ruddiman, W. F., Hao, Q., Wu, H., Qiao, Y., Zhu, R. X., Peng, S., Wei, J., Yuan, B., and Liu, T.: Onset of Asian desertification by 22 Myr ago inferred from loess deposits in China, *Nature*, 416, 159-163, 2002.
- 1195 Gupta, A. K., Singh, R. K., Joseph, S., and Thomas, E.: Indian Ocean high-productivity event (10–8 Ma): Linked to global cooling or to the initiation of the Indian monsoons?, *Geology*, 32, 753-756, 2004.
- Gupta, A. K., Yuvaraja, A., Prakasam, M., Clemens, S. C., and Velu, A.: Evolution of the South Asian monsoon wind system since the late Middle Miocene, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 438, 160-167, 2015.
- 1200 Herbert, T. D.: A long marine history of carbon cycle modulation by orbital-climatic changes, *Proceedings of the National Academy of Sciences*, 94, 8362-8369, 1997.
- Herbert, T. D., Lawrence, K. T., Tzanova, A., Peterson, L. C., Caballero-Gill, R., and Kelly, C. S.: Late Miocene global cooling and the rise of modern ecosystems, *Nature Geoscience*, 9, 843, 10.1038/Ngeo2813, 2016.
- 1205

- Hermoyian, C. S. and Owen, R. M.: Late Miocene-early Pliocene biogenic bloom: Evidence from low-productivity regions of the Indian and Atlantic Oceans, *Paleoceanography*, 16, 95-100, 2001.
- 1210 Holbourn, A., Kuhnt, W., Clemens, S. C., and Heslop, D.: A ~12 Myr Miocene record of East Asian Monsoon variability from the South China Sea, *Paleoceanography and Paleoclimatology*, 36, e2021PA004267, <https://doi.org/10.1029/2021PA004267>, 2021.
- Holbourn, A., Kuhnt, W., Schulz, M., Flores, J.-A., and Andersen, N.: Orbitally-paced climate evolution during the middle Miocene “Monterey” carbon-isotope excursion, *Earth and Planetary Science Letters*, 261, 534-550, 2007.
- 1215 Holbourn, A., Kuhnt, W., Clemens, S. C., Kochhann, K. G., Jöhnck, J., Lübbers, J., and Andersen, N.: Late Miocene climate cooling and intensification of southeast Asian winter monsoon, *Nature communications*, 9, 1584, 2018.
- Hovan, S. A. and Rea, D. K.: The Cenozoic record of continental mineral deposition on Broken and Ninetyeast Ridges, Indian Ocean: southern African aridity and sediment delivery from the Himalayas, *Paleoceanography*, 7, 833-860, 1992.
- 1220 Howell, P., Piasias, N., J. Ballance, Baughman, J., and Ochs, L.: ARAND Time-Series Analysis Software [code], 2006.
- Huang, Y., Clemens, S. C., Liu, W., Wang, Y., and Prell, W. L.: Large-scale hydrological change drove the late Miocene C₄ plant expansion in the Himalayan foreland and Arabian Peninsula, *Geology*, 35, 531–534, doi: 10.1130/G23666A, 2007.
- 1225 Jalihal, C., Bosmans, J. H. C., Srinivasan, J., and Chakraborty, A.: The response of tropical precipitation to Earth's precession: the role of energy fluxes and vertical stability, 2019.
- Jensen, T. G.: Cross-equatorial pathways of salt and tracers from the northern Indian Ocean: Modelling results, *Deep Sea Research Part II: Topical Studies in Oceanography*, 50, 2111-2127, 2003.
- 1230 Jia, G., Peng, P. a., Zhao, Q., and Jian, Z.: Changes in terrestrial ecosystem since 30 Ma in East Asia: Stable isotope evidence from black carbon in the South China Sea, *Geology*, 31, 1093-1096, 2003.
- 1235 Jöhnck, J., Holbourn, A. E., Kuhnt, W., and Andersen, N.: Oxygen isotope offsets in deep-water benthic foraminifera, *Journal of Faoraminiferal Research*, 51, 225-244, 2021.
- Jöhnck, J., Kuhnt, W., Holbourn, A., and Andersen, N.: Variability of the Indian Monsoon in the Andaman Sea across the Miocene-Pliocene transition, *Paleoceanography and Paleoclimatology*, e2020PA003923, 2020.

- 1240 Jyothibabu, R., Vinayachandran, P., Madhu, N., Robin, R., Karnan, C., Jagadeesan, L., and Anjusha, A.: Phytoplankton size structure in the southern Bay of Bengal modified by the Summer Monsoon Current and associated eddies: Implications on the vertical biogenic flux, *Journal of Marine Systems*, 143, 98-119, 2015.
- Kathayat, G., Cheng, H., Sinha, A., Spötl, C., Edwards, R. L., Zhang, H., Li, X., Yi, L., Ning, Y., and Cai, Y.: Indian monsoon variability on millennial-orbital timescales, *Scientific reports*, 6, 24374, 2016.
- 1245 Keerthi, M., Lengaigne, M., Vialard, J., de Boyer Montégut, C., and Muraleedharan, P.: Interannual variability of the Tropical Indian Ocean mixed layer depth, *Climate dynamics*, 40, 743-759, 2013.
- 1250 Keigwin, L.: Late Cenozoic stable isotope stratigraphy and paleoceanography of DSDP sites from the east equatorial and central North Pacific Ocean, *Earth and Planetary Science Letters*, 45, 361-382, 1979.
- Keigwin, L. and Shackleton, N.: Uppermost Miocene carbon isotope stratigraphy of a piston core in the equatorial Pacific, *Nature*, 284, 613-614, 1980.
- 1255 Kitoh, A., Yukimoto, S., Noda, A., and Motoi, T.: Simulated changes in the Asian summer monsoon at times of increased atmospheric CO₂, *Journal of the Meteorological Society of Japan. Ser. II*, 75, 1019-1031, 1997.
- Koné, V., Aumont, O., Lévy, M., and Resplandy, L.: Physical and biogeochemical controls of the phytoplankton seasonal cycle in the Indian Ocean: A modeling study, *Indian Ocean Biogeochemical Processes and Ecological Variability*, 185, 350, 2009.
- 1260 Kroon, D., Steens, T., and Troelstra, S. R.: 13. ONSET OF MONSOONAL RELATED UPWELLING IN THE WESTERN ARABIAN SEA AS REVEALED BY PLANKTONIC FORAMINIFERS1, *Proceedings of the ocean drilling program, scientific results*, Kuhnt, W., Holbourn, A. E., Jöhnck, J., and Lübbers, J.: Miocene to Pleistocene
- 1265 Palaeoceanography of the Andaman Region: Evolution of the Indian Monsoon on a Warmer-Than-Present Earth, in: *The Andaman Islands and Adjoining Offshore: Geology, Tectonics and Palaeoclimate*, Springer, 261-288, 2020.
- Kutzbach, J. E.: Monsoon climate of the early Holocene: climate experiment with the earth's orbital parameters for 9000 years ago, *Science*, 214, 59-61, 1981.
- 1270 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astronomy & Astrophysics*, 428, 261-285, 2004.

- Le, J. and Shackleton, N. J.: Carbonate dissolution fluctuations in the western equatorial Pacific during the late Quaternary, *Paleoceanography*, 7, 21-42, 1992.
- 1275 Lee, J., Kim, S., Lee, J. I., Cho, H. G., Phillips, S. C., and Khim, B.-K.: Monsoon-influenced variation of clay mineral compositions and detrital Nd-Sr isotopes in the western Andaman Sea (IODP Site U1447) since the late Miocene, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 538, 109339, 2020.
- Lévy, M., Shankar, D., André, J. M., Shenoi, S., Durand, F., and de Boyer Montégut, C.:
1280 Basin-wide seasonal evolution of the Indian Ocean's phytoplankton blooms, *Journal of Geophysical Research: Oceans*, 112, 2007.
- Li, M., Hinnov, L., and Kump, L.: Acycle: Time-series analysis software for paleoclimate research and education, *Computers & Geosciences*, 127, 12-22, 10.1016/j.cageo.2019.02.011, 2019.
- 1285 Licht, A., Van Cappelle, M., Abels, H., Ladant, J.-B., Trabucho-Alexandre, J., France-Lanord, C., Donnadieu, Y., Vandenberghe, J., Rigaudier, T., and Lécuyer, C.: Asian monsoons in a late Eocene greenhouse world, *Nature*, 513, 501-506, 2014.
- Longhurst, A.: Seasonal cycles of pelagic production and consumption, *Progress in oceanography*, 36, 77-167, 1995.
- 1290 Lourens, L. J., Wehausen, R., and Brumsack, H. J.: Geological constraints on tidal dissipation and dynamical ellipticity of the Earth over the past three million years, *Nature*, 409, 1029-1033, 2001.
- Lourens, L. J., Antonarakou, A., Hilgen, F., Van Hoof, A. A. M., Vergnaud-Grazzini, C., and Zachariasse, W. J.: Evaluation of the Plio-Pleistocene astronomical timescale,
1295 *Paleoceanography*, 11, 391-413, Doi 10.1029/96pa01125, 1996.
- Lübbbers, J., Kuhnt, W., Holbourn, A. E., Bolton, C. T., Gray, E., Usui, Y., Kochhann, K. G., Beil, S., and Andersen, N.: The middle to late Miocene “Carbonate Crash” in the equatorial Indian Ocean, *Paleoceanography and Paleoclimatology*, 34, 813-832, 2019.
- Lutsko, N. J., Marshall, J., and Green, B.: Modulation of monsoon circulations by cross-
1300 equatorial ocean heat transport, *Journal of Climate*, 32, 3471-3485, 2019.
- Lyle, M. and Baldauf, J.: Biogenic sediment regimes in the Neogene equatorial Pacific, IODP Site U1338: Burial, production, and diatom community, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 433, 106-128, 2015.
- Ma, W., Tian, J., Li, Q., and Wang, P.: Simulation of long eccentricity (400-kyr) cycle in
1305 ocean carbon reservoir during Miocene Climate Optimum: Weathering and nutrient response to orbital change, *Geophysical Research Letters*, 38, 2011.

- Martínez-Ruiz, F., Paytan, A., Gonzalez-Muñoz, M., Jroundi, F., Abad, M. d. M., Lam, P. J., Bishop, J., Horner, T., Morton, P. L., and Kastner, M.: Barite formation in the ocean: Origin of amorphous and crystalline precipitates, *Chemical Geology*, 511, 441-451, 2019.
- 1310 McCreary, J., Murtugudde, R., Vialard, J., Vinayachandran, P., Wiggert, J. D., Hood, R. R., Shankar, D., and Shetye, S.: Biophysical processes in the Indian Ocean, *Indian Ocean biogeochemical processes and ecological variability*, 185, 9-32, 2009.
- McLennan, S. M.: Relationships between the trace element composition of sedimentary rocks and upper continental crust, *Geochemistry, Geophysics, Geosystems*, 2, 2001.
- 1315 McNeill, L. C., Dugan, B., Backman, J., Pickering, K. T., Poudroux, H. F., Henstock, T. J., Petronotis, K. E., Carter, A., Chemale Jr, F., and Milliken, K. L.: Understanding Himalayan erosion and the significance of the Nicobar Fan, *Earth and Planetary Science Letters*, 475, 134-142, 2017.
- Meyers, S. R.: *Astrochron: An R Package for Astrochronology [code]*, 2014.
- 1320 Molnar, P., Boos, W. R., and Battisti, D. S.: Orographic controls on climate and paleoclimate of Asia: thermal and mechanical roles for the Tibetan Plateau, *Annual Review of Earth and Planetary Sciences*, 38, 2010.
- Müller, D. W., Hodell, D. A., and Ciesielski, P. F.: 25. Late Miocene to earliest Pliocene (9.8-4.5 Ma) paleoceanography of the subantarctic southeast atlantic: stable isotopic, sedimentologic, and microfossil evidence, in: *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by: Ciesielski, P. F., Kristoffersen, Y., and al., e., 1991.
- 1325 Murray, R., Knowlton, C., Leinen, M., Mix, A. C., and Polsky, C.: Export production and carbonate dissolution in the central equatorial Pacific Ocean over the past 1 Myr, *Paleoceanography*, 15, 570-592, 2000.
- 1330 Nigrini, C.: Composition and biostratigraphy of radiolarian assemblages from an area of upwelling (northwestern Arabian Sea, Leg 117), in: *Proceedings of the Ocean Drilling Program, Scientific Results*, edited by: Prell, W., Niitsuma, N., and al, e., 89-126, 1991.
- Nomura, R.: Paleogene to Neogene deep-sea paleoceanography in the eastern Indian Ocean: benthic foraminifera from ODP Sites 747, 757 and 758, *Micropaleontology*, 251-290, 1995.
- 1335 Paillard, D.: The Plio-Pleistocene climatic evolution as a consequence of orbital forcing on the carbon cycle, *Climate of the Past*, 13, 1259-1267, 2017.
- Pälike, H., Lyle, M. W., Nishi, H., Raffi, I., Ridgwell, A., Gamage, K., Klaus, A., Acton, G., Anderson, L., and Backman, J.: A Cenozoic record of the equatorial Pacific carbonate compensation depth, *Nature*, 488, 609-614, 2012.

- 1340 Paytan, A. and Griffith, E. M.: Marine barite: Recorder of variations in ocean export productivity, *Deep Sea Research Part II: Topical Studies in Oceanography*, 54, 687-705, 2007.
- Paytan, A. and Kastner, M.: Benthic Ba fluxes in the central Equatorial Pacific, implications for the oceanic Ba cycle, *Earth and Planetary Science Letters*, 142, 439-450, 1996.
- 1345 Paytan, A., Kastner, M., and Chavez, F.: Glacial to interglacial fluctuations in productivity in the equatorial Pacific as indicated by marine barite, *Science*, 274, 1355-1357, 1996.
- Paytan, A., Kastner, M., Martin, E., Macdougall, J., and Herbert, T.: Marine barite as a monitor of seawater strontium isotope composition, *Nature*, 366, 445-449, 1993.
- Pickering, K. T., Carter, A., Andò, S., Garzanti, E., Limonta, M., Vezzoli, G., and Milliken, K. L.: Deciphering relationships between the Nicobar and Bengal submarine fans, *Indian Ocean*, *Earth and Planetary Science Letters*, 544, 116329, 2020a.
- 1350 Pickering, K. T., Poudoux, H., McNeill, L. C., Backman, J., Chemale, F., Kutterolf, S., Milliken, K. L., Mukoyoshi, H., Henstock, T. J., and Stevens, D. E.: Sedimentology, stratigraphy and architecture of the Nicobar Fan (Bengal–Nicobar Fan System), *Indian Ocean: Results from International Ocean Discovery Program Expedition 362*, *Sedimentology*, 67, 2248-2281, 2020b.
- 1355 Pierce, J., Weissel, J., and al., e.: *Proceedings of the Ocean Drilling Program, Initial Reports 121*, 1000 pp. pp.1989.
- Pimm, A.: Sedimentology and history of the northeastern Indian Ocean from late Cretaceous to Recent, in: *Initial Reports of the Deep Sea Drilling Project*, edited by: von der Borch, C. C., Sclater, J. G., and al., e., U.S. Gov. Print. Office, Washington, 717-803, 10.2973/dsdp.proc.22.139.1974, 1974.
- 1360 Prasanna Kumar, S., Muraleedharan, P., Prasad, T., Gauns, M., Ramaiah, N., De Souza, S., Sardesai, S., and Madhupratap, M.: Why is the Bay of Bengal less productive during summer monsoon compared to the Arabian Sea?, *Geophysical Research Letters*, 29, 88-81-88-84, 2002.
- 1365 Prell, W. L. and Kutzbach, J. E.: Monsoon variability over the past 150,000 years, *Journal of Geophysical Research: Atmospheres*, 92, 8411-8425, 1987.
- Prell, W. L. and Kutzbach, J. E.: Sensitivity of the Indian monsoon to forcing parameters and implications for its evolution, *Nature*, 360, 17, 1992.
- 1370 Privé, N. C. and Plumb, R. A.: Monsoon dynamics with interactive forcing. Part I: Axisymmetric studies, *Journal of the atmospheric sciences*, 64, 1417-1430, 2007a.

- Privé, N. C. and Plumb, R. A.: Monsoon dynamics with interactive forcing. Part II: Impact of eddies and asymmetric geometries, *Journal of the atmospheric sciences*, 64, 1431-1442, 2007b.
- 1375
- Rae, J. W., Zhang, Y. G., Liu, X., Foster, G. L., Stoll, H. M., and Whiteford, R. D.: Atmospheric CO₂ over the Past 66 Million Years from Marine Archives, *Annual Review of Earth and Planetary Sciences*, 49, 2021.
- Ramaswamy, V.: Lithogenic Fluxes to the Northern Indian Ocean, *Monsoon biogeochemistry*, 97, 1993.
- 1380
- Rial, J. A. and Anaclerio, C.: Understanding nonlinear responses of the climate system to orbital forcing, *Quaternary Science Reviews*, 19, 1709-1722, 2000.
- Rickaby, R., Bard, E., Sonzogni, C., Rostek, F., Beaufort, L., Barker, S., Rees, G., and Schrag, D.: Coccolith chemistry reveals secular variations in the global ocean carbon cycle?, *Earth and Planetary Science Letters*, 253, 83-95, 2007.
- 1385
- Rixen, T., Gaye, B., Emeis, K.-C., and Ramaswamy, V.: The ballast effect of lithogenic matter and its influences on the carbon fluxes in the Indian Ocean, *Biogeosciences*, 16, 485-503, 2019.
- Robinson, M. M., Bartol, M., Bolton, C. T., Ding, X., Gariboldi, K., Romero, O. E., and Scientists, E.: Biostratigraphic Summary, in: *Proceedings of the International Ocean Discovery Program*, 2016.
- 1390
- Rogalla, U. and Andruleit, H.: Precessional forcing of coccolithophore assemblages in the northern Arabian Sea: Implications for monsoonal dynamics during the last 200,000 years, *Marine Geology*, 217, 31-48, 2005.
- 1395
- Rostek, F., Bard, E., Beaufort, L., Sonzogni, C., and Ganssen, G.: Sea surface temperature and productivity records for the past 240 kyr in the Arabian Sea, *Deep Sea Research Part II: Topical Studies in Oceanography*, 44, 1461-1480, 1997.
- Schmitz, B.: Barium, equatorial high productivity, and the northward wandering of the Indian continent, *Paleoceanography*, 2, 63-77, 1987.
- 1400
- Schott, F. A. and McCreary Jr, J. P.: The monsoon circulation of the Indian Ocean, *Progress in Oceanography*, 51, 1-123, 2001.
- Schott, F. A., Xie, S. P., and McCreary Jr, J. P.: Indian Ocean circulation and climate variability, *Reviews of Geophysics*, 47, 2009.
- Sengupta, D., Bharath Raj, G., Ravichandran, M., Sree Lekha, J., and Papa, F.: Near-surface salinity and stratification in the north Bay of Bengal from moored observations, *Geophysical Research Letters*, 43, 4448-4456, 2016.
- 1405

- Shackleton, N. J. and Hall, M. A.: The Late Miocene stable isotope record, Site 926, in: Proc. ODP Sci. Res. 154, edited by: Shackleton, N. J., Curry, W. B., Richter, C., and Bralower, T. J., 367-373, 1997.
- 1410 Shackleton, N. J., Hall, M., and Pate, D.: 15. Pliocene stable isotope stratigraphy of Site 846, Proc. Ocean Drill. Program Sci. Results, 337-355,
Shimmiel, G. B. and Mowbray, S. R.: The inorganic geochemical record of the northwest Arabian Sea: a history of productivity variation over the last 400 ky from sites 722 and 724, 1991.
- 1415 Shipboard Scientific Party: Site 758, in: Proceedings of the ODP Initial Reports 121, edited by: J. Peirce, J. W., et al., ODP: College Station, TX, 359-453, 1989.
Si, W. and Rosenthal, Y.: Reduced continental weathering and marine calcification linked to late Neogene decline in atmospheric CO₂, *Nature Geoscience*, 12, 833-838, 2019.
- 1420 Singh, A. D., Jung, S. J., Darling, K., Ganeshram, R., Ivanochko, T., and Kroon, D.:
Productivity collapses in the Arabian Sea during glacial cold phases, *Paleoceanography*, 26, 2011.
Singh, R. K. and Gupta, A.: Miocene history of Indian monsoon: A review of marine records, *Indian Miocene: A Geodynamic and chronologic framework for palaeobiota, sedimentary environments and palaeoclimates* (Ed: Tiwari RP) Special Publication of the Palaeontological Society of India, 5, 101-109, 2014.
- 1425 Smart, C. W., Thomas, E., and Ramsay, A. T.: Middle-late Miocene benthic foraminifera in a western equatorial Indian Ocean depth transect: paleoceanographic implications, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 247, 402-420, 2007.
Spicer, R., Yang, J., Herman, A., Kodrul, T., Aleksandrova, G., Maslova, N., Spicer, T.,
- 1430 Ding, L., Xu, Q., and Shukla, A.: Paleogene monsoons across India and South China: Drivers of biotic change, *Gondwana Research*, 49, 350-363, 2017.
Steinthorsdottir, M., Coxall, H., de Boer, A., Huber, M., Barbolini, N., Bradshaw, C., Burls, N., Feakins, S., Gasson, E., and Henderiks, J.: The Miocene: the Future of the Past, *Paleoceanography and Paleoclimatology*, e2020PA004037, 2020.
- 1435 Stow, D. A., Amano, K., Batson, B., Brass, G. W., Corrigan, J., Raman, C., Tiercelin, J.-J., Townsend, M., Wijayananda, N., and Cochran, J.: Sediment facies and processes on the distal Bengal Fan, Leg 116, in: Proceedings of the Ocean Drilling Project, Scientific Results, Leg 116, edited by: Cochran, J. R., Stow, D. A. V., and al., e., 377-396, 1990.

- 1440 Sun, J. and Huang, X.: Half-precessional cycles recorded in Chinese loess: response to low-latitude insolation forcing during the Last Interglaciation, *Quaternary Science Reviews*, 25, 1065-1072, 2006.
- Sun, X. and Wang, P.: How old is the Asian monsoon system?—Palaeobotanical records from China, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 222, 181-222, 2005.
- 1445 Tabor, C. R., Otto-Bliesner, B. L., Brady, E. C., Nusbaumer, J., Zhu, J., Erb, M. P., Wong, T. E., Liu, Z., and Noone, D.: Interpreting precession-driven $\delta^{18}\text{O}$ variability in the South Asian monsoon region, *Journal of Geophysical Research: Atmospheres*, 123, 5927-5946, 2018.
- Tauxe, L. and Feakins, S.: A re-assessment of the chronostratigraphy of late Miocene C3-C4 transitions, *Paleoceanography and Paleoclimatology*, e2020PA003857, 2020.
- 1450 Thomas, E. K., Clemens, S. C., Sun, Y., Prell, W. L., Huang, Y., Gao, L., Loomis, S., Chen, G., and Liu, Z.: Heterodynes dominate precipitation isotopes in the East Asian monsoon region, reflecting interaction of multiple climate factors, *Earth and Planetary Science Letters*, 455, 196-206, 2016.
- Tripathi, S., Tiwari, M., Lee, J., Khim, B.-K., Expedition, I., Pandey, D. K., Clift, P. D., Kulhanek, D. K., Andò, S., and Bendle, J. A.: First evidence of denitrification vis-à-vis monsoon in the Arabian Sea since Late Miocene, *Scientific reports*, 7, 43056, 2017.
- 1455 Unger, D. and Jennerjahn, T.: Impact of regional Indian Ocean characteristics on the biogeochemical variability of settling particles, *GMS*, 185, 257-280, 2009.
- Unger, D., Ittekkot, V., Schäfer, P., Tiemann, J., and Reschke, S.: Seasonality and interannual variability of particle fluxes to the deep Bay of Bengal: influence of riverine input and oceanographic processes, *Deep Sea Research Part II: Topical Studies in Oceanography*, 50, 897-923, 2003.
- 1460 Van Andel, T. H., Heath, G. R., and T.C. Moore, J.: Cenozoic History and Paleooceanography of the Central Equatorial Pacific Ocean: A Regional Synthesis of Deep Sea Drilling Project Data, in: *The Geological Society of America*, <https://doi.org/10.1130/MEM143-p1>, 1975.
- 1465 Vidya, P., Prasanna Kumar, S., Gauns, M., Verenkar, A., Unger, D., and Ramaswamy, V.: Influence of physical and biological processes on the seasonal cycle of biogenic flux in the equatorial Indian Ocean, *Biogeosciences Discussions*, 10, 2889-2936, 2013.
- Vinayachandran, P., Chauhan, P., Mohan, M., and Nayak, S.: Biological response of the sea around Sri Lanka to summer monsoon, *Geophysical Research Letters*, 31, 2004.
- 1470 Wan, S., Li, A., Clift, P. D., and Stuut, J.-B. W.: Development of the East Asian monsoon: mineralogical and sedimentologic records in the northern South China Sea since 20 Ma, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 254, 561-582, 2007.

- Wang, P., Tian, J., and Lourens, L. J.: Obscuring of long eccentricity cyclicity in Pleistocene oceanic carbon isotope records, *Earth and Planetary Science Letters*, 290, 319-330, 10.1016/j.epsl.2009.12.028, 2010.
- 1475
- Wang, P., Clemens, S., Beaufort, L., Braconnot, P., Ganssen, G., Jian, Z., Kershaw, P., and Sarnthein, M.: Evolution and variability of the Asian monsoon system: state of the art and outstanding issues, *Quaternary Science Reviews*, 24, 595-629, 2005.
- Webber, B. G., Matthews, A. J., Vinayachandran, P., Neema, C., Sanchez-Franks, A., Vijith, V., Amol, P., and Baranowski, D. B.: The dynamics of the Southwest Monsoon current in 2016 from high-resolution in situ observations and models, *Journal of Physical Oceanography*, 48, 2259-2282, 2018.
- 1480
- Webster, P. J.: *The elementary monsoon*, 1987a.
- Webster, P. J.: *The variable and interactive monsoon*, *Monsoons*. Wiley, New York, 269, 1987b.
- 1485
- Webster, P. J. and Yang, S.: Monsoon and ENSO: Selectively interactive systems, *Quarterly Journal of the Royal Meteorological Society*, 118, 877-926, 1992.
- Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnett, J. S., Bohaty, S. M., De Vleeschouwer, D., and Florindo, F.: An astronomically dated record of Earth's climate and its predictability over the last 66 million years, *Science*, 369, 1383-1387, 2020.
- 1490
- Zeeden, C., Hilgen, F. J., Hüsing, S. K., and Lourens, L. L.: The Miocene astronomical time scale 9–12 Ma: New constraints on tidal dissipation and their implications for paleoclimatic investigations, *Paleoceanography*, 29, 296-307, 2014.
- 1495
- Zeeden, C., Meyers, S. R., Lourens, L. J., and Hilgen, F. J.: Testing astronomically tuned age models, *Paleoceanography*, 30, 369-383, 2015.
- Zeeden, C., Hilgen, F., Westerhold, T., Lourens, L., Röhl, U., and Bickert, T.: Revised Miocene splice, astronomical tuning and calcareous plankton biochronology of ODP Site 926 between 5 and 14.4 Ma, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 369, 430-451, 2013.
- 1500
- Zhisheng, A., Clemens, S. C., Shen, J., Qiang, X., Jin, Z., Sun, Y., Prell, W. L., Luo, J., Wang, S., and Xu, H.: Glacial-interglacial Indian summer monsoon dynamics, *science*, 333, 719-723, 2011.
- Zhuang, G., Pagani, M., and Zhang, Y. G.: Monsoonal upwelling in the western Arabian Sea since the middle Miocene, *Geology*, 45, 655-658, 2017.
- 1505

Ziegler, M., Lourens, L. J., Tuenter, E., Hilgen, F., Reichert, G. J., and Weber, N.: Precession phasing offset between Indian summer monsoon and Arabian Sea productivity linked to changes in Atlantic overturning circulation, *Paleoceanography*, 25, 2010.

1510 Zweng, M., Reagan, J., Antonov, J., Locarnini, R., Mishonov, A., Boyer, T., Garcia, H., Baranova, O., Johnson, D., and Seidov, D.: World Ocean Atlas 2013, Volume 2: Salinity, NOAA Atlas NESDIS, 74, 39, 2013.