

Climate Change Detection and Attribution using observed and simulated Tree-Ring Width

Jörg Franke^{1,2,*}, Michael N. Evans^{2,3,*}, Andrew Schurer⁴, Gabriele C. Hegerl⁴

¹Institute of Geography, University of Bern, Switzerland.

5 ²Oeschger Centre for Climate Change Research, University of Bern, Switzerland.

³Department of Geology and Earth System Science Interdisciplinary Center, University of Maryland, College Park, Maryland 20742, USA

⁴School of GeoSciences, The University of Edinburgh, Edinburgh, United Kingdom

* Both authors contributed equally to this study

10 *Corresponding author(s): Jörg Franke (franke@giub.unibe.ch) and Michael N. Evans (mnevans@umd.edu)*

Abstract. The detection and attribution (D&A) of paleoclimatic change to external radiative forcing relies on regression of statistical reconstructions on simulations. However, this procedure may be biased by assumptions of stationarity and univariate linear response of the underlying paleoclimatic observations. Here we perform a D&A study, modeling paleoclimate data observations as a function of paleoclimatic data simulations. Specifically, we model tree-ring width (TRW) observations as a linear function of TRW simulations which are themselves forward modeled from realistic singly-forced and cumulatively forced climate simulations for the period 1401-2000. Temperature and moisture-sensitive TRW simulations detect distinct patterns in time and space. Temperature-sensitive TRW observations and simulations are significantly correlated for northern hemisphere averages, and their variation is attributed to volcanic forcing. In decadal smoothed temporal fingerprints, we find the observed responses to be significantly larger and/or more persistent than the simulated responses. The pattern of simulated TRW of moisture-limited trees is consistent with the observed anomalies in the two years following major volcanic eruptions. We can for the first time attribute this spatiotemporal fingerprint in moisture limited tree-ring records to volcanic forcing. These results suggest that use of nonlinear and multivariate proxy system models in paleoclimatic detection and attribution studies may permit more realistic, spatially resolved and multivariate fingerprint detection studies, and evaluation of the climate sensitivity to external radiative forcing, than has previously been possible.

30 1 Introduction

One of the crucial questions in climate change research is to determine how external radiative forcings bring about climate variation and change, if the forced response may be distinguished from the internal, unforced variability, and between different forcings, and if the model simulated change is consistent with that detected in observed or reconstructed climate. Major contributions to answer this question come from so-called “detection and attribution” (D&A) studies (see review by Hegerl and Zwiers, 2011) for methods, and Gillett et al. (2021) for a recent update. Methods are generally based on matching observed changes with patterns derived from climate model simulations, which were driven by single and multiple external forcings, including solar variability, volcanic aerosols, the well-mixed greenhouse gases, orbital variations, and land use change. The idea initiated in early work by Hasselmann (1979). After methodological refinements and advances in climate modeling in the early 1990s (e.g. Hasselmann, 1993; Santer et al., 1993) there was growing evidence that the external greenhouse gas signal can be differentiated from climate variability generated within Earths’ climate system (Hegerl et al., 1996). Detection and attribution studies have been an important part of the Assessment Reports of Working Group I of the Intergovernmental Panel on Climate Change, from the calling for better detection of the role of human activities in climate forcing in the first Assessment report (1990), to formal detection and attribution studies comparing observed and simulated climate change in all Assessment reports since, with increasingly confident assessments of the detection of human influences and estimates of the human contribution derived from attribution results.

Typically, D&A analyses have been limited to periods when instrumental observations of physically measurable variables and derived diagnostics are available, with global observation networks becoming dense enough for such studies about 100 to 150 years before present. This period allowed for attribution of trends in many thermodynamic and dynamic characteristics of the climate system, including global and regional temperature, temperature extremes, ocean heat content, tropopause height, specific humidity, zonal mean precipitation, air pressure fields to potential forcings (e.g. Hegerl et al., 1996; Santer, 2003; Polson et al., 2013a; Bindoff et al., 2014; Eyring et al., 2021; Gillett et al., 2021). While


instrumental observations cover the period of a major increase in greenhouse gases and other human influences, studying the climate system responses to non-anthropogenic external radiative forcings, such as solar variability or volcanic eruption, benefits from studying longer periods over which more realizations and/or longer-term processes are evident, and where the anthropogenic influence is less dominant. For instance, very few climatically important volcanic eruptions occurred in the past 150 years, but more than a dozen occurred over the past 600 years (Sigl et al., 2015) at nonuniform frequency in time, possibly creating long-term forcing of the climate system (McGregor et al., 2015; PAGES 2k Consortium, 2019; Brönnimann et al., 2019). Such longer-term studies would integrate longer-term responses of the climate system to external radiative forcing, enabling a more complete picture of the equilibrium and transient response, and ultimately of the climate sensitivity to external radiative forcing. Paleoclimatology allows extension of the observational record into the past using indirect measurements of climatic conditions, which can be used to reconstruct past climate. Previous studies have detected a role of external forcing in the climate of the last millennium using annual mean surface temperature anomaly reconstructions on both a hemispheric scale (Hegerl et al., 2003; Schurer et al., 2013, 2014) and regionally (PAGES 2k-PMIP3 group, 2015). These analyses have found that volcanic forcing is detected with a smaller contribution from greenhouse gases that is detectable by 1900, and a contribution from solar forcing that was not detectable against climate variability. However, the reconstruction process itself introduces additional assumptions into detection and attribution studies that arise from the nature of the reconstructions, but which may not be justified. Many of these are demonstrated in pseudoproxy experiments (Smerdon et al., 2011) and through study of the extensive network of tree-ring width observations. These include assumed univariate, normally distributed and linear response of the paleoclimatic indicators to the target reconstruction variable (Evans et al., 2014; Wang et al., 2014); stationarity of patterns of regional and global scale climate variability (Wilson et al., 2010); seasonal and spatial representation (St. George, 2014; Smerdon et al., 2011); and autoregression characteristics in observations and target variables (Cook et al., 1999). Limited adherence to assumptions in observations and statistical modeling has been found to introduce biases into reconstructed variables, even in large scale averages (PAGES2k Consortium, 2017) and may lead to the underestimation of errors in D&A studies that are necessary to separate the forced and unforced responses (Neukom et al., 2019). In particular, autocorrelation due to memory in TRW affects the response to volcanism which, if not accounted for, biases D&A results (Lücke et al., 2019).

Progress in process understanding of paleoclimatic observations has led to the development of proxy system models (Evans et al., 2013), which may be used to identify systematic uncertainties and evaluate the extent of biases introduced by the reconstruction process into the D&A problem. One recent example is the Vaganov–Shashkin Lite (VSL) sensor model, which simulates standardized tree ring width (TRW) chronology variations based on monthly mean temperature, precipitation, and latitude. These inputs are used to estimate nondimensional growth arising from temperature and soil moisture conditions (G_T , G_M) either of which may stoichiometrically limit growth at each monthly time step: a multivariate and nonlinear mimic of the processes by which forests sense and filter climatic variability and imprint those results in observable tree ring width variations (Tolwinski-Ward et al., 2011a, b). VSL has been widely tested for parameter estimation and global applicability.

Here we leverage VSL, historical gridded climate data products (Harris et al., 2014), singly and multiply forced climate simulations for the period 1401 to 2000 C.E. (Schurer et al., 2013), and the nearly 3000 consistently detrended TRW observations (B14, Breitenmoser et al., 2014) to perform an extratropical northern hemisphere D&A exercise directly using observed and simulated TRW data (Fig. 1, Eq. 1):

$$\alpha = \beta_0 + \beta_1 \hat{\alpha} \quad (1)$$

With α representing the paleoclimatic observations (TRW in this case), and $\hat{\alpha}$ representing the sensor modeled TRW simulations, themselves employing as input the output of a realistically forced climate model. Coefficients β_0 and β_1 represent, respectively, the unforced and forced amplitudes of variability (for a more detailed introduction, see Section 2.4 below). This approach stands in contrast to prior studies, which perform the D&A analysis in the space of reconstructed northern hemisphere mean surface temperature at annual resolution (Schurer et al., 2013, 2014). It has the potential advantages of circumventing assumptions required in the reconstruction process, and exploiting the “several-to-one” mapping that might reinforce environmental signatures in TRW data, such as spatially and temporally correlated patterns of moisture and temperature variability that mimic drought indices (Cook et al., 1999, 2004, 2010; Meko et al., 1995). Conversely, we may also identify key uncertainties in the sensor modeling, and the potential for the several-to-one mapping to obfuscate the detection and attribution of a forced response in the TRW observations.



115 **Figure 1: Schematic overview of the performed analysis. General steps are indicated in bold, study-specific procedures in normal text. B14 refers to the Breitenmoser et al. (2014) data set, S/N stands for signal to noise ratio, T for temperature and PREC for precipitation.**

120 The remainder of this paper is organized as follows. First, we estimate and evaluate parameters for VSL, using gridded instrumental temperature (T) and precipitation (PREC) estimates and contemporaneous TRW observations (Section 2.2) in the period 1901-1970. Then VSL is used to build ensembles of simulated TRW series, in response to singly and cumulatively forced simulations of T and PREC, given uncertainty propagated through the parameter estimation process, and with bias corrections for simulated T and PREC ensembles (Section 2.3). We then estimate the D&A coefficients and their propagated uncertainty (Eq. 2; Section 2.4). The results are analyzed locally, regionally, and globally for detection and attribution of a forced climate response, vis a vis the simulated and actual TRW observations (Section 3). We discuss the results and the potential to extend the approach in Section 4; conclusions are drawn in Section 5.

130 2 Data and Methods

The inputs into and process of this detection and attribution study is illustrated in Fig. 1 and described below.

135 2.1 Tree-ring width measurements

We use the tree-ring width (TRW) collection described by and employed in (Breitenmoser et al., 2014) as the observational basis for the development and validation of VSL parameters, and as the D&A predictand (Eq 1). B14 consists of 2918 uniformly detrended and standardized tree-ring width chronologies from six continents and 163 species that have been upload to the International Tree-Ring Data Base (ITRDB, Zhao et al., 2019) until 2014. These series have been quality controlled for metadata errors, repetitive measurements, incorrect units, decimal point errors and misplaced positions (Tab. S1 in Breitenmoser et al., 2014). Detrending for biological age trends and stand dynamics and standardization to dimensionless growth indices was done in a hierarchical approach. If possible, negative exponential curves and linear regression curves of any slope were fitted. In case both methods failed, “a smoothing spline was fit with a 50% cut-off frequency at 75 % of each series length” (ARSTAN, Cook, 1985; Breitenmoser et al., 2014). Multiple measurements at the same site have been combined into robust means (Cook and Kairiukstis, 1990), which are variance adjusted for changing sample size through time (Osborn et al., 1997). For every point in time, which is explicitly resolved as one value per growing season each year, a chronology is based on at least 8 samples. We use the autoregressive-standardized (Osborn et al., 1997; Frank et al., 2007) version of the available chronologies from B14. We require that the chronologies have at least 40 years observed within the period 1901 – 1970. (See section 2.2 below)

150 We restrict subsequent analysis of simulations and the D&A exercise to the extratropical northern hemisphere continental areas, where the vast majority of TRW observations are located, with high concentrations in the North America, Europe and northern Asia (Fig. 2). Record length varies from 100-

600 years (Fig. 2). Series availability is generally greatest between the mid-19th century and the late 20th century (Fig. 3), and the longest records are equally distributed in longitude across the north hemisphere boreal terrestrial latitudes (Fig. 2).

155 2.2 VSL parameter estimation

For the purpose of VSL parameter estimation, we use the global, gridded instrumental temperature and precipitation data sets CRU TS 3.23 (Harris et al., 2014), regridded to 64 longitude x 32 latitude (~5.6°) using a distance-weighted average of the four nearest neighbor values. To correct for mean temperature biases, we applied an adiabatic (-6 K/km) T correction to the regridded CRU product, based on differences between elevations of grid points and elevations of observed TRW chronologies (Evans et al., 2006). Parameters T_1 , T_2 , M_1 and M_2 describe the onset of growth (1) and point above which climate is no longer a limiting factor (2) for temperature (T) and moisture (M), respectively (Tolwinski-Ward et al., 2011a, 2013). We conditioned and validated all four parameters simultaneously using contemporaneous observations and VSL simulations within the period 1901-1970. The growth period is defined as a 16-month interval. To integrate monthly incremental growth arising from pre-season and growing season, the growth integration period starts in September of the previous year and ends in December of the current year in the northern hemisphere (previous March to current June for the southern hemisphere), the same period as in Tolwinski-Ward et al. (2011a) and Breitenmoser et al. (2014). Other VSL parameters are not calibrated, but taken from other studies (Evans et al., 2006; Fan, 2004; Huang et al., 1996; Tolwinski-Ward et al., 2011a, 2013; Vaganov et al., 2006; van den Dool, 2003). Within the chosen parameter estimation time window 1901-1970, with available $N \geq 40$, half of the years for which observed TRW data were available were chosen at random for parameter estimation (“calibration” Tolwinski-Ward et al., 2013). The other half were reserved for validation of the estimated parameters, via simulation using the estimated parameters, T and PREC not used to estimate the parameters, and comparison with the TRW observations withheld from the calibration process. The process was then repeated, using now the second half of data for parameter estimation (calibration), and the first half for validation of this parameter set. Up to 200 parameter sets were stored as valid, if all four calibration and validation correlations between simulated and observed TRW were all independently significant at the $p < 0.1$ level; all others failing this validation test were discarded.

180 2.3 VSL simulations, 1401-2000

Temperature and precipitation input data for VSL are derived from climate model simulations. We use the set of simulations described in Schurer et al. (2014), which have been conducted with HADCM3, interpolated to the same 64 x 32 grid as described in Section 2.2, to produce TRW simulations driven by singly and cumulatively forced climate simulations (Table 1). Because simulated T and PREC are spatially and seasonally biased relative to historical gridded T and PREC, we first bias-correct the HadCM3 T and PREC fields by computing T and PREC anomaly fields and adding them to (scaling them by) the CRU TS3.23 T climatology (PREC variability) for the overlapping period 1901-2000 C.E. This step also ensures that systematic differences in mean simulated T and PREC will not systematically bias VSL simulations based on parameter estimates conditioned on the historical CRU TS3.23 T and PREC products. We then identified the primary limiting factor for simulated growth (at $p < 0.05$, assuming a binomial distribution) and divided the simulated chronologies into primarily temperature, moisture (M), both, or neither limited TRW simulations. The median, over parameter estimate realizations, of T and M TRW simulations, were then separately weighted by inverse distance between observed and simulated grid point, observed expressed population signal (EPS, Wigley et al., 1984), and observed mean correlation between increment series within a chronology (RBAR, Cook and Kairiukstis, 1990) and averaged. Observed TRW were gridded and averaged in the same way as described above for subsequent D&A analysis (see Fig. 1 for a schematic overview of the entire process chain). Because centennial-scale climate variability may not be consistently preserved in the TRW records (Cook et al., 1995; Franke et al., 2013), and these timescales are poorly sampled in the 600 year period available for study, we removed low frequency variability by applying a 71-year high pass LOESS filter to both observed and simulated gridded TRW and focus our analyses on this residual interannual to multidecadal variance at annual resolution. The choice of annual resolution reflects the observational resolution (Section 2.1), the time-integrating nature of the sensor model (Section 2.2), the uncertainty of implementing radiative forcing estimates in the climate simulations (Gao et al., 2008), and the response timescale of the climate system to the large scale forcing. We call the results, on which we base the detection and attribution analyses, climate-sensor simulations. This nomenclature reflects modeling of both the climate in response to external radiative forcing(s), and the tree ring width observation that is basis for the comparison with actual TRW observations.

210 **Table 1: All forcing and single forcing HADCM3 simulations as well as control runs used in this study (V: volcanic, S: solar, G: greenhouse gases, L: land-use, A: tropospheric aerosols).**

Number of simulations	Forcings	Period
2	NO forcing	1401– 2000
4	V, S, G, L, A	1401 – 2000
3	V	1401 – 2000
4	S	1401 – 2000

2.4 Detection and Attribution

215 To solve for the D&A coefficients in (Eq. 1), we use the total least squares (TLS) D&A technique to account for errors in both dependent and independent variables within the regression procedure (Allen and Stott, 2003) to account for internal variability in both observations and model simulations. We follow the analysis used in Allen and Stott (2003), Polson et al., (2013a), and Schurer et al. (2013), which estimates a best fit regression coefficient (β) given by the equation:

$$\alpha = \beta (\hat{\alpha} - v) + v \quad (2)$$

220 In this study, $\hat{\alpha}$ are the simulated tree-ring widths and α are the observed tree-ring widths, either at each grid box or spatially and/or temporally aggregated to increase the signal to noise ratio. v are realizations of internal variability. Confidence intervals are obtained with the bootstrap method described in DelSole et al. (2019). They are calculated by randomly sampling, with replacement, pairs of values from the arrays of observed and simulated tree-ring widths to form new arrays the same length as the originals. A new scaling factor is then calculated by regressing the resampled model onto the resampled observations. This process is repeated 10000 times and a 5%–95% confidence interval is estimated from the distribution. If the distribution of beta values is significantly greater than 0 ($p < 0.05$) then the effect of the response to the forcing is considered to have been detected. If the distribution of β -values is significantly less than unity, the response in the climate-sensor simulations is too large; the response in climate-sensor simulations is significantly greater than observed, and the simulated climate sensitivity is smaller than observed. Conversely, if the scaling range is significantly greater than unity, the simulated climate-sensor response is significantly smaller than observed in TRW, and the climate sensitivity of the model may be inferred to be larger than observed. The estimate of the unforced variability as the residual of the D&A regression model provides another important result that needs to be compared with unforced variability of climate simulations (control runs) as a check of variability (PAGES 2k Consortium, 2019).

3 Results

3.1 Parameter estimation, TRW simulations and TRW observations

240 1664 of 2761 TRW chronologies in the B14 compilation were climate sensitive and therefore successfully simulated and retained for further analysis. With small differences between climate simulations, and using the methodology as described in (Tolwinski-Ward et al., 2013), their Fig 8, 21% of the successfully simulated chronologies are temperature sensitive, ca. 57% are moisture sensitive, ca. 11% are both moisture and temperature sensitive, and ca. 11% are not climate sensitive, i.e. neither moisture nor temperature sensitive (Fig 2). Distributions of temperature, moisture, both temperature and moisture, and neither temperature nor moisture sensitivity overlap in space. There are many moisture-sensitive TRW chronologies found in North America, the Mediterranean and other arid regions (Fig. 2, top right panel). However, there are also temperature-sensitive chronologies (upper left panel) and mixed responders (lower left panel) which are collocated in arid regions (upper left panel). Chronologies found to be neither temperature nor moisture sensitive (Fig 2, lower right panel) tend to be found at the highest latitudes, but not exclusively so.

250




Figure 2: Limitations determined for all TRW chronologies with valid parameter sets, separated into temperature sensitive (top left), moisture sensitive (top right), both T and M sensitive (bottom left) and neither T nor M sensitive (bottom right). (Maps created with Matlab mapping package M_map (Pawlowicz, 2022))

255

We found that bootstrapped VSL parameter estimates were in many cases distinctly non-normal in distribution for some or all of the four parameters, and for some TRW simulations. Distributions were sometimes uniformly distributed across the prior expected parameter ranges, unimodal non-normal, and even bimodal. Because there were not necessarily well-defined means or medians across parameter sets and simulations, we used all valid parameter sets to produce TRW simulations. Hence, we propagate uncertainty arising from stochastic variation in the climate simulations through parameter and structural uncertainty in the ring width sensor model.

260

Because the fingerprint of external radiative forcing may or may not be distinct and unique in temperature and moisture, we use the fit of VSL diagnostic variables G_T and G_M (estimate of nondimensional growth arising from temperature and soil moisture conditions, respectively) to binomial distributions to determine whether each simulation is primarily controlled by temperature, moisture, both or neither control at the $p < 0.05$ significance level (Tolwinski-Ward et al., 2013). We perform a similar analysis to determine the same primary growth controls in the TRW observations, using the same diagnostics from the parameter estimation exercise. We then average TRW observations to the simulation grid resolution for temperature and moisture-limited simulations separately. Where there are multiple observed TRW chronologies available within a particular grid box, we construct a weighted average using inverse grid-point distance and intra-chronology mean incremental growth series correlation as weighting factors.


265

270

TRW simulations (Sections 2.2, 2.3) are developed for all locations, where TRW observations exist and the parameter estimation has been successful, i.e. for most of the extratropical northern hemisphere (Fig. 3). We exclude the southern hemisphere in this because only 5 temperature-sensitive chronologies and 1 moisture-sensitive chronology are located there. The record length of the simulations is constrained by TRW observations (see 2.1 and Fig. 3). The longest records are equally distributed in longitude across the north hemisphere boreal terrestrial latitudes (Fig. 3). Thus, statistics assessed across the simulations and observations are best described as representing the northern hemisphere temperate and subpolar terrestrial regions. Furthermore, we note that the locations of temperature and moisture-sensitive chronologies overlap (Fig 2) but are generally not coincident (Fig. 3): for either the observed or simulated sets, only about 1/3 are both coincident in space and significantly correlated with each other (Fig. 2). Hence, for the remainder of the analysis presented here, we develop and discuss the temperature and moisture-sensitive results separately.

275

280



285


Figure 3: Numbers of years (color scale) available for comparison between gridded, observed and climate-sensor simulated TRW chronologies, within the period 1401-2000. Top row: observed (left) and simulated (right) temperature-sensitive chronologies. Simulated chronologies are masked by observational availability, and in some cases (e.g., subpolar Eurasia) are neither moisture nor temperature-sensitive or else are both temperature and moisture sensitive, and do not appear in either simulation map. Lower panels: as for top panels, except for moisture-sensitive chronologies. (Maps created with Matlab mapping package M_map (Pawlowicz, 2022))

290

295


300

Figure 4 shows the growth functions GT and GM for the subsets of temperature and moisture-sensitive identified TRW chronologies (section 3.1). Although VSL has well-known limitations, for instance the lack of a soil moisture model allowing for snow, and despite the potential for an unrealistic and coarsely resolved annual cycle in the HadCM3 simulations, the results suggest plausible seasonality of the growth response of the TRW simulations. In particular, GT for T sensitive chronologies is maximum but limiting in June-October with a median response (black line) maximum for July-September. GM for the T sensitive subset of chronologies is not limiting through the same period. Similarly, for M sensitive chronologies, GM is limiting between July-December. GE (the scaling associated with insolation (energy) as a function of latitude) is limiting ($GE < 0.7$) after September for latitudes poleward of 20N (results not shown), and GT is not limiting through the warm months.



305

Figure 4: Simulated intra-year partial growth response functions G_T (left column) and G_M (right column) for T sensitive (top row) and M sensitive (bottom row) simulations using ALL climate simulations, with parameters conditioned and validated using observed TRW data within the period 1901-1970. Heavy solid lines are median values across all simulations.



310

Figure 5. Top: mean series for observed (black) and ALL-forcing simulated temperature-sensitive chronologies (red). Annually resolved and a 11-year Hanning window filtered time series are shown with thin and bold lines, respectively. Labels quantify the Pearson correlation (r), effective degrees of freedom (edf, Hu et al., 2017) and the p-value. x-axis units are years CE. Bottom: the same above but for moisture-sensitive observations (black) and simulations (blue). Note, this plot shows the global means of standardized TRW at all grid boxes with data before high-pass filtering

315

and variance adjustment. There are small differences in numbers of observed and simulated chronologies that arise from both the observational masking and from the simulation parameter validation procedure (Section 2).

320 **3.2. Comparison of observed and sensor-simulated TRW**

To detect an external-forcing signal in noisy, local observations, the signal-to-noise ratio has to be enhanced. This is commonly achieved by averaging in space and/or time (Sections 1, 2.4). We begin with analysis of global mean TRW variability at all locations where tree growth is either temperature or moisture limited, for comparisons between TRW observations and climate-sensor simulations driven with ALL forcings (Table 1). The variance of the average over all grid boxes increases back in time because of the decreasing numbers of records (Fig. 5), likely reflecting increasing uncertainty; the variance in the beginning of the 15th century is twice as large as that observed at the end of the 20th century. To reduce sensitivity of the detection and attribution analysis to observational uncertainty, we homogenize the variance through time by multiplication of a time dependent scaling factor that is estimated by linear regression of the observed variance on the variance of TRW climate-sensor simulations from the control simulations.


325 Results suggest limited but significant correlation between global mean observed and simulated TRW temperature-sensitive simulations, for both annual and decadal-filtered series (Fig. 5). Nonsignificant correlations are found for moisture-sensitive observations vs simulations at both annual and decadal timescales (Fig. 5). We find similar results for correlations between VOLC-forced simulations and temperature and moisture-sensitive observations (T sensitive: $r_1=0.22$, $edf=201$, $p=0.001$; $r_{11}=0.48$, $edf=19$, $p=0.02$; M sensitive: $r_1=0.01$, $edf=380$, $p=0.40$; $r_{11}=0.00$, $edf=54$, $p=0.51$). Correlations are not significant for comparisons between observed and SOLAR-forced or unforced TRW simulations (results not shown).

330 Based on these results, we test for detection of patterns in the TRW following volcanic eruptions in temperature and moisture-sensitive TRW chronologies, using a composite analysis across the 7 largest (above 95th quantile) volcanic forcing event responses based on Crowley and Unterman (2013) between 1670 and 1970 (Fig. 6). We do not use the full period back to 1401 because only few locations have data reaching that far back in time. However, including up to 12 eruptions where available leads to a very similar pattern (not shown). The event size is measured in stratospheric Atmospheric Optical Depth (AOD, Table 2) Consistent with the results for averaged temperature-sensitive chronologies, we find a reduction in simulated tree growth in the first two years after the eruption in nearly all locations worldwide (Fig. 6, top right). Observed tree growth at the temperature sensitive sites is reduced in most locations, but not as homogeneously as in the simulations (Fig. 6, top left). Possible reasons may be related to the small sample size, uncertainties in the reconstruction of the volcanic forcing (Sigl et al., 2015), a low climate signal-to-noise ratio in ring width, and an enhanced signal-to-noise ratio in the simulations, which are represented by their 4-member ensemble mean (Tab. 1). Additionally, moisture influences may not be perfectly removed from the temperature-sensitive observations, because some of the positive growth anomalies appear in locations for which tree growth tends to be generally moisture limited (in southwestern North America, north European lowlands and the eastern Mediterranean (St. George and Ault, 2014). Thus, the composite observed temperature-sensitive response may in part also reflect increased moisture in dry regions following volcanic eruptions (Iles and Hegerl, 2015).

335 **Table 2: Largest volcanic eruptions used in the composite analysis based on atmospheric optical depth.**

Crowley and Unterman, 2013											
1442	1456	1594	1600	1641	1673	1694	1809	1815	1832	1884	1903
Sigl et al., 2015											
1453	1458	1595	1601	1641	1695	1783	1809	1815	1832	1836	1884


360 For our moisture-sensitive comparison, we do not find a global volcanic response of the same sign, but rather regions with uniform responses (Fig. 6, bottom row). The simulated composite (Fig 6, bottom row, rightmost panel) produces anomalously lower (higher) growth at high (low) latitudes. We show the composite for observations based on two forcings, Crowley and Unterman (2013) that was used to force the climate simulations (Fig 1; Table 2) and the more recent and probably more realistic inferred Global Volcanic aerosol Forcing (GVF, in W/m^2) by Sigl et al. (2015). For the composite based on Crowley and Unterman (2013), we find positive growth anomalies around the Mediterranean and in southwestern North America, and negative anomalies in Eurasia, with more prominent composite positive than negative regions. The observed composite based on the Toohey and Sigl (2017) chronology produces a small negative composite response in eastern North America and small positive response in western North America, which is weaker but consistent with the pattern in the simulations (Fig. 6, bottom row, middle and right panels).



375 **Figure 6: Composite average ring width anomaly (standardized units) in temperature-sensitive TRW chronologies in the first two years after volcanic eruptions in observations and VOLC-forcing simulations (top). Because relatively few TRW records are available for 1400-1700 (Fig 2), the composite includes the 7 strongest eruptions between 1670 and 1970 based on the eruption chronologies of Crowley et al. (2013) (left and right column) and Sigl et al. (2015 middle column). Bottom row: as for top row, except for moisture-sensitive TRW observations and simulations.**

380 3.3 Detection and Attribution analysis

We detect and attribute a response to volcanic forcing in both, the spatial mean temperature timeseries and the spatio-temporal pattern of moisture limited tree-ring records. As large volcanic eruptions disturb the climate system for a few years, we show results 3-year and 11-year moving averages. For the 3-year smoothed temperature-sensitive TRW averaged over all grid boxes, we find a significantly detectable scaling factor β not significantly different from one; in other words, observed and simulated temperature sensitive chronologies agree within uncertainty (Fig. 7, left panel). This is true for both, the all-forcing and the volcanic-forcing based TRW simulations (β_{ALL} and β_{VOLC}). For decadal averages (Fig. 7, middle panel), both scaling factors are likely greater than 1, and indicate that the observed responses are larger and/or more persistent than the simulated responses. This is consistent with superposed epoch analysis results for volcanic eruptions using tree-ring data (see Lücke et al., 2019).



385 **Figure 7: Left: Beta values and uncertainties for 3-year moving averages (following DelSole et al., 2019) in the TLS D&A analysis for temperature-sensitive TRW (Fig 2, top panels). ALL and VOLC indicate regression on the all-forcing and volcanic-only based TRW simulations, respectively. Middle: as in left panel, except for 11-year running means; uncertainties adjusted for serial autocorrelation. Right: as in left panel, except for moisture-sensitive TRW (Fig 2, bottom panels) but with the aggregate mean response grouped by the two regions based on the positive or negative response of the TRW simulations (Fig 6, bottom right). Uncertainty ranges are based on 90% confidence intervals of scaling factors.**

395 As described above, moisture sensitive trees show positive growth anomalies in some regions and negative anomalies in other regions. We define a two-region spatio-temporal pattern identified in the moisture sensitive TRW simulations (Fig. 6, bottom right panel). Again β -values not significantly

different from one for both ALL and VOLC forced simulations (Fig. 7, right panel) allow us to attribute moisture changes in response to volcanism.

4 Discussion

400 Detection and attribution studies using paleoclimatic data have previously focused on regression of reconstructed climate variables on realistically forced climate simulations (PAGES 2k Consortium, 2019; Schurer et al., 2014). In this study, we have attached a validated, realistically multivariate and nonlinear, intermediate complexity proxy-sensor model (Evans et al., 2013; Tolwinski-Ward et al., 2013, 2015) to enable the D&A framework within the space of the paleoclimatic observation – in this study, tree ring
405 width chronologies. Because this particular sensor model is a scaled and time-integrated transformation of temperature and precipitation variations into a single diagnostic, which is commonly observed across the terrestrial landscape, the potential for fingerprinting either distinct univariate or integrated plant-stress-like signatures of the different radiative forcings becomes possible. The approach also substitutes structural and parametric uncertainty in the sensor model for the uncertainty arising from inversion of
410 multivariate paleoclimatic observations for univariate climatic reconstruction, and so provides a complementary assessment of the uncertainty that propagates into the D&A results.

We find that the global mean forced response in temperature-sensitive TRW chronologies is consistent with observations within the 1401-2000 period, a result that supports the prior work using global mean surface temperature reconstructions as predictand (Hegerl et al., 2006 and references therein), and
415 implicitly the use of temperature-sensitive TRW chronologies for producing those results. However, we also find that moisture and temperature sensitive chronologies (Fig. 3) form distinct subgroups in space (Fig. 2) and in temporal average (Fig. 4). The fingerprint of climate forcing, as determined by comparison between all-series averaged temperature and moisture sensitive observations and simulations is statistically significant in temperature, but not in moisture, for both ALL and VOLC forced simulations
420 (Fig. 7).

For the attribution analysis targeting volcanic forcing (Figs. 6, 7), we find disagreement in the amplitude of the temperature-sensitive forcing as a function of timescale, with the observed annual/3-year/decadal timescale variance being smaller/equal/greater than the simulated variance (Fig. 7, left and middle panels). One explanation would be that the simulated temperature response to volcanic forcing is
425 unrealistically large. This has been observed for the HadCM3 climate model simulation in a previous study (Schurer et al., 2013). Volcanic forcings used to produce the climate simulations may also be oversimplified in time and/or space relative to actual forcing (Stevenson et al., 2017), or its timing may be incorrect (yielding suppressed amplitude in reconstructions). For many eruptions with an unknown date, the eruption was set to January 1 and the AOD is entered into the model in four equal latitude bands
430 only, proportional to the amount of Sulphur in the Antarctic and Greenland ice cores (Crowley and Unterman, 2013). Because TRW simulations are a simplified representation of actual TRW variation, they neglect the observational uncertainty and the potential for superimposed and competing influences, such that the simulated TRW response to forcing may be relatively large. This is indeed the case; for either VOLC or ALL forcing, simulated variance is about one-third larger than observed variance (results not shown).
435

A further explanation could be that autocorrelations in observed and simulated TRW are different. We find observed mean TRW autocorrelation to be about two-thirds larger than that of VOLC forced simulations (results not shown). Consequently, we find the observed TRW variance at decadal resolution to be significantly greater than simulated TRW variance. This result suggests that i) the observed
440 response contains decadal timescale non-climatic variation not adequately removed by observational signal processing (Cook and Kairiukstis, 1990), ii) mechanisms represented in the climate simulations are inadequate to represent slower response timescales of volcanic forcing (Miller et al., 2012), iii) mechanisms of forest response to volcanic forcing via soil moisture, air temperature or insolation variations, as represented in VSL, are insufficient to represent the observed lower frequency response
445 (Esper et al., 2015; Lücke et al., 2019) or a combination of all three factors. Previous studies found scaling factors to increase as more smoothing is applied (Schurer et al., 2013). However, they did not reach the point of a significantly larger response in observations than simulations.

Previous studies based on historical observations found that volcanic eruptions produced positive precipitation and streamflow anomalies in the Mediterranean and the Southwest of the United States,
450 whereas negative anomalies were observed at high latitudes, and in western North America, the Indian to South-East-Asian region and the tropics (Iles et al., 2013; Iles and Hegerl, 2015). This was in agreement with the CMIP5 simulated precipitation response (Fig. 1a in Iles and Hegerl, 2015), although the pattern in observed precipitation was very noisy and not clearly observed. In contrast, the response was identifiable in observed streamflow data which covers a longer period and integrates the precipitation
455 response. Reasons that the precipitation response couldn't be detected are likely to include the small number of eruptions in the instrumental period over which a composite was formed, combined with low

signal-to-noise ratio for precipitation (Fig. 1a in Iles and Hegerl, 2015), and the complex precipitation response pattern with regions of increases and decreases which is more difficult to detect (see also Polson et al., 2013b). We would obtain similarly non-detection and non-attribution were we to define regions manually (e.g. Northern Europe vs Mediterranean or Western vs Eastern North America), or for a smaller integration over the years following an eruption. This finding is in agreement with (Rao et al., 2017), who see the effect in tree-ring reconstructed PDSI only in a very small region of north western Europe, southern Spain and northern Morocco, and with Fischer et al. (2007), who found increased precipitation in the Mediterranean and Scandinavia, and decreased precipitation in Northwestern and Central Europe following volcanic eruptions, although not statistically significant in many locations. For such small-scale regions, our TRW network is too sparse, our simulation grid too coarse and the time span of the TRW series is too limited to calculate robust composites. The present study sidesteps or respects some of these challenges by extending the analysis several centuries into the past (Table 1), integrating the forced response over time and space (Fig 7), and forming the attribution model using the native observed variable rather than a reconstructed climatic variable (Section 1; Fig. 1). We find a similar pattern in moisture sensitive TRW (Fig 6). Simulations are most consistent with the expected pattern if the composite is based on the same forcing chronology as that used to drive the underlying HadCM3 simulations (Crowley and Unterman, 2013). The pattern in TRW observations agrees better with the more recent volcanic forcing chronology of Sigl (2015). This suggests the latter forcing series reconstruction may be more consistent with the response as observed in TRW. However, the two forcing chronologies are similar enough, that the two-region detection and attribution analysis (Fig 7, right panel) produces the significant detection of both the ALL and VOLC forced TRW signals, within uncertainty of unity, lending support to the conclusions of (Iles and Hegerl, 2015).

5 Conclusion

We have performed a detection and attribution study using observed and modeled tree-ring width data directly for the exercise. We found that temperature and moisture sensitive TRW contain different signatures of the forced climate response over the past six centuries. Specifically, we find that the signature of the ALL- and VOLC-forcing response is most evident across the mean of all temperature-sensitive chronologies, but not across the mean of all moisture sensitive chronologies. The amplitude of the temperature-sensitive forced response is larger than expected from the model simulations in decadal filtered results, suggesting inaccuracies in the representation of forcing and/or response on those timescales in observations, simulations, or both sources of information. Additionally, we detect and attribute a previously identified spatial pattern in moisture-sensitive response to volcanic forcing at annual timescales, with a dipole drying/moistening pattern similar to the one previously identified by others within the historical time period and with direct moisture observations. In this study we demonstrate for the first time that climate change D&A can be conducted directly on paleoclimatic observations and their multivariate, non-linear proxy system simulations, allowing for a much more reliable model evaluation than possible if using reconstructed climate variables. The results may realistically diverge from those obtained by D&A studies using univariate surface temperatures reconstructed from similar datasets, because the underlying observations may in reality be multivariate, nonlinear responders. Further studies could improve upon this proof of concept by incorporating stable isotopic observations in combination with isotope enabled climate model simulations, and by accessing a longer time interval for developing composite analyses, additional data types, and a larger ensemble of realistically forced climate simulations.

Data availability

The HADCM3 simulations are available at the Center for Environmental Data Analysis: <https://catalogue.ceda.ac.uk/uuid/9b51eb5d44524e4c9aebcbbc53d79a27>. Gridded instrumental data is available at the Climate Research Unit: <https://crudata.uea.ac.uk/cru/data/hrg/>. The B14 TRW data collection is available at the World Data Center for Climate (WDCC) at DKRZ, where it was used as input data for a data assimilation based climate reconstruction called EKF400 (Franke et al., 2017): http://cera-www.dkrz.de/WDCC/ui/Compact.jsp?acronym=EKF400_Input_Data_v1.1. TRW simulations are available upon request from the MNE.

Author contribution

JF and MNE designed the study with contributions of GCH. JF prepared the inputs for the TRW simulation conducted by MNE. JF performed the D&A analysis with support of AS. JF and MNE wrote the paper with help of AS and GCH.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

520 This study originated in a sabbatical stay of MNE at the University of Bern supported by the Oeschger
Center for Climate Change, the Sigrist Foundation and the University of Bern, Department of Geography.
JF was funded by the Swiss National Science Foundation (project 162668) and by the European Union
(H2020/ERC grant number 787574 PALAEO-RA). A.S. and G.H. were funded by the UK Natural
Environmental Research Council via the grant Vol-Clim (NE/S000887/1) and under the Belmont forum,
525 Grant PacMedy (NE/P006752/1). We thank the British Atmospheric Data Centre (BADC) for access to
the HadCM3 climate simulation (<http://badc.nerc.ac.uk/browse/badc/euroclim500>). MNE is especially
grateful to Martin Grosjean, Raphael Neukom, Christoph Dätwyler, Jörg Franke, Stefan Brönnimann and
their research groups for stimulating conversations, chocolate, coffee and emotional support through a
particularly difficult life transition; and to Anupma Gupta (1971-2015), and Aditi and Maya for their
ongoing teachings and wisdom.

530 References

- Allen, M. R. and Stott, P. A.: Estimating signal amplitudes in optimal fingerprinting, part I: theory, *Climate Dynamics*, 21, 477–491, <https://doi.org/10.1007/s00382-003-0313-9>, 2003.
- 535 Bindoff, N. L., Stott, P. A., AchutaRao, K. M., Allen, M. R., Gillett, N., Gutzler, D., Hansingo, K.,
Hegerl, G., Yu, H., Jain, S., Mokhov, I. I., Overland, J., Perlwitz, J., Sebbari, R., and Zhang, X.: Detec-
tion and Attribution of Climate Change: from Global to Regional, in: *Climate Change 2013 - The Phys-
ical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergov-
ernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and
New York, NY, USA, 867–952, <https://doi.org/10.1017/CBO9781107415324.022>, 2014.
- 540 Breitenmoser, P., Brönnimann, S., and Frank, D.: Forward modelling of tree-ring width and compari-
son with a global network of tree-ring chronologies, *Clim. Past*, 10, 437–449,
<https://doi.org/10.5194/cp-10-437-2014>, 2014.
- 545 Brönnimann, S., Franke, J., Nussbaumer, S. U., Zumbühl, H. J., Steiner, D., Trachsel, M., Hegerl, G.
C., Schurer, A., Worni, M., Malik, A., Flückiger, J., and Raible, C. C.: Last phase of the Little Ice Age
forced by volcanic eruptions, *Nat. Geosci.*, 12, 650–656, <https://doi.org/10.1038/s41561-019-0402-y>,
2019.
- Cook, E. R.: *A Time Series Analysis Approach to Tree-Ring Standardization*, The University of Ari-
zona, 1985.
- Cook, E. R. and Kairiukstis, L. A. (Eds.): *Methods of Dendrochronology*, Springer Netherlands, Dor-
drecht, <https://doi.org/10.1007/978-94-015-7879-0>, 1990.
- 550 Cook, E. R., Briffa, K. R., Meko, D. M., Graybill, D. A., and Funkhouser, G.: The “segment length
curse” in long tree-ring chronology development for palaeoclimatic studies, *The Holocene*, 5, 229–237,
<https://doi.org/10.1177/095968369500500211>, 1995.
- Cook, E. R., Meko, D. M., Stahle, D. W., and Cleaveland, M. K.: Drought Reconstructions for the
Continental United States, 12, 18, 1999.
- 555 Cook, E. R., Woodhouse, C. A., Eakin, C. M., Meko, D. M., and Stahle, D. W.: Long-Term Aridity
Changes in the Western United States, *Science*, 306, 1015–1018, [https://doi.org/10.1126/sci-
ence.1102586](https://doi.org/10.1126/sci-
ence.1102586), 2004.
- Cook, E. R., Anchukaitis, K. J., Buckley, B. M., D’Arrigo, R. D., Jacoby, G. C., and Wright, W. E.:
560 Asian Monsoon Failure and Megadrought During the Last Millennium, *Science*, 328, 486–489,
<https://doi.org/10.1126/science.1185188>, 2010.
- Crowley, T. J. and Unterman, M. B.: Technical details concerning development of a 1200 yr proxy in-
dex for global volcanism, *Earth Syst. Sci. Data*, 5, 187–197, <https://doi.org/10.5194/essd-5-187-2013>,
2013.
- 565 DelSole, T., Trenary, L., Yan, X., and Tippett, M. K.: Confidence intervals in optimal fingerprinting,
Clim Dyn, 52, 4111–4126, <https://doi.org/10.1007/s00382-018-4356-3>, 2019.

- van den Dool, H.: Performance and analysis of the constructed analogue method applied to U.S. soil moisture over 1981–2001, *J. Geophys. Res.*, 108, 8617, <https://doi.org/10.1029/2002JD003114>, 2003.
- 570 Esper, J., Schneider, L., Smerdon, J. E., Schöne, B. R., and Büntgen, U.: Signals and memory in tree-ring width and density data, *Dendrochronologia*, 35, 62–70, <https://doi.org/10.1016/j.dendro.2015.07.001>, 2015.
- Evans, M. N., Reichert, B. K., Kaplan, A., Anchukaitis, K. J., Vaganov, E. A., Hughes, M. K., and Cane, M. A.: A forward modeling approach to paleoclimatic interpretation of tree-ring data, *J. Geophys. Res.*, 111, G03008, <https://doi.org/10.1029/2006JG000166>, 2006.
- 575 Evans, M. N., Tolwinski-Ward, S. E., Thompson, D. M., and Anchukaitis, K. J.: Applications of proxy system modeling in high resolution paleoclimatology, *Quaternary Science Reviews*, 76, 16–28, <https://doi.org/10.1016/j.quascirev.2013.05.024>, 2013.
- Evans, M. N., Smerdon, J. E., Kaplan, A., Tolwinski-Ward, S. E., and González-Rouco, J. F.: Climate field reconstruction uncertainty arising from multivariate and nonlinear properties of predictors, *Geophys. Res. Lett.*, 41, 9127–9134, <https://doi.org/10.1002/2014GL062063>, 2014.
- 580 Eyring, V., Gillett, N. P., Achuta Rao, K. M., Barimalala, R., Barreiro Parrillo, M., Bellouin, N., Casou, C., Durack, P. J., Kosaka, Y., McGregor, H., Min, S., Morgenstern, O., and Sun, Y.: Chapter 3: Human Influence on the Climate System, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 423–552, <https://doi.org/10.1017/9781009157896.005>, 2021.
- 585 Fan, Y.: Climate Prediction Center global monthly soil moisture data set at 0.5° resolution for 1948 to present, *J. Geophys. Res.*, 109, D10102, <https://doi.org/10.1029/2003JD004345>, 2004.
- Fischer, E. M., Luterbacher, J., Zorita, E., Tett, S. F. B., Casty, C., and Wanner, H.: European climate response to tropical volcanic eruptions over the last half millennium, *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2006GL027992>, 2007.
- 590 Frank, D., Büntgen, U., Böhm, R., Maugeri, M., and Esper, J.: Warmer early instrumental measurements versus colder reconstructed temperatures: shooting at a moving target, *Quaternary Science Reviews*, 26, 3298–3310, <https://doi.org/10.1016/j.quascirev.2007.08.002>, 2007.
- 595 Franke, J., Frank, D., Raible, C. C., Esper, J., and Brönnimann, S.: Spectral biases in tree-ring climate proxies, *Nature Clim Change*, 3, 360–364, <https://doi.org/10.1038/nclimate1816>, 2013.
- Franke, J., Brönnimann, S., Bhend, J., and Brugnara, Y.: A monthly global paleo-reanalysis of the atmosphere from 1600 to 2005 for studying past climatic variations, *Sci Data*, 4, 170076, <https://doi.org/10.1038/sdata.2017.76>, 2017.
- 600 Gao, C., Robock, A., and Ammann, C.: Volcanic forcing of climate over the past 1500 years: An improved ice core-based index for climate models, *J. Geophys. Res.*, 113, D23111, <https://doi.org/10.1029/2008JD010239>, 2008.
- 605 Gillett, N. P., Kirchmeier-Young, M., Ribes, A., Shiogama, H., Hegerl, G. C., Knutti, R., Gastineau, G., John, J. G., Li, L., Nazarenko, L., Rosenbloom, N., Seland, Ø., Wu, T., Yukimoto, S., and Ziehn, T.: Constraining human contributions to observed warming since the pre-industrial period, *Nat. Clim. Chang.*, 11, 207–212, <https://doi.org/10.1038/s41558-020-00965-9>, 2021.
- Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset: updated high-resolution grids of monthly climatic observations, *Int. J. Climatol.*, 34, 623–642, <https://doi.org/10.1002/joc.3711>, 2014.
- 610 Hasselmann, K.: On the signal-to-noise problem in atmospheric response studies, in: *Meteorology of Tropical Oceans*, Royal Meteorological Society, London, UK, 251–259, 1979.

- Hasselmann, K.: Optimal Fingerprints for the Detection of Time-dependent Climate Change, *J. Climate*, 6, 1957–1971, [https://doi.org/10.1175/1520-0442\(1993\)006<1957:OFFTDO>2.0.CO;2](https://doi.org/10.1175/1520-0442(1993)006<1957:OFFTDO>2.0.CO;2), 1993.
- Hegerl, G. and Zwiers, F.: Use of models in detection and attribution of climate change, *WIREs Clim Change*, 2, 570–591, <https://doi.org/10.1002/wcc.121>, 2011.
- 615 Hegerl, G. C., Storch, H. von, Hasselmann, K., Santer, B. D., Cubasch, U., and Jones, P. D.: Detecting Greenhouse-Gas-Induced Climate Change with an Optimal Fingerprint Method, 9, 2281–2306, [https://doi.org/10.1175/1520-0442\(1996\)009<2281:DGGICC>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<2281:DGGICC>2.0.CO;2), 1996.
- Hegerl, G. C., Crowley, T. J., Baum, S. K., Kim, K.-Y., and Hyde, W. T.: Detection of volcanic, solar and greenhouse gas signals in paleo-reconstructions of Northern Hemispheric temperature, *Geophys. Res. Lett.*, 30, <https://doi.org/10.1029/2002GL016635>, 2003.
- 620 Hegerl, G. C., Crowley, T. J., Hyde, W. T., and Frame, D. J.: Climate sensitivity constrained by temperature reconstructions over the past seven centuries, *Nature*, 440, 1029–1032, <https://doi.org/10.1038/nature04679>, 2006.
- Hu, J., Emile-Geay, J., and Partin, J.: Correlation-based interpretations of paleoclimate data – where statistics meet past climates, *Earth and Planetary Science Letters*, 459, 362–371, <https://doi.org/10.1016/j.epsl.2016.11.048>, 2017.
- 625 Huang, J., van den Dool, H., M., and Georgarakos, K. P.: Analysis of Model-Calculated Soil Moisture over the United States (1931–1993) and Applications to Long-Range Temperature Forecasts, 9, 1350–1362, [https://doi.org/10.1175/1520-0442\(1996\)009<1350:AOMCSM>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<1350:AOMCSM>2.0.CO;2), 1996.
- 630 Iles, C. E. and Hegerl, G. C.: Systematic change in global patterns of streamflow following volcanic eruptions, *Nature Geosci*, 8, 838–842, <https://doi.org/10.1038/ngeo2545>, 2015.
- Iles, C. E., Hegerl, G. C., Schurer, A. P., and Zhang, X.: The effect of volcanic eruptions on global precipitation, *J. Geophys. Res. Atmos.*, 118, 8770–8786, <https://doi.org/10.1002/jgrd.50678>, 2013.
- Lücke, L. J., Hegerl, G. C., Schurer, A. P., and Wilson, R.: Effects of Memory Biases on Variability of Temperature Reconstructions, 32, 8713–8731, <https://doi.org/10.1175/JCLI-D-19-0184.1>, 2019.
- 635 McGregor, H. V., Evans, M. N., Goosse, H., Leduc, G., Martrat, B., Addison, J. A., Mortyn, P. G., Oppo, D. W., Seidenkrantz, M.-S., Sicre, M.-A., Phipps, S. J., Selvaraj, K., Thirumalai, K., Filipsson, H. L., and Ersek, V.: Robust global ocean cooling trend for the pre-industrial Common Era, *Nature Geosci*, 8, 671–677, <https://doi.org/10.1038/ngeo2510>, 2015.
- 640 Meko, D., Stockton, C. W., and Boggess, W. R.: The Tree-Ring Record of severe sustained Drought, *J Am Water Resources Assoc*, 31, 789–801, <https://doi.org/10.1111/j.1752-1688.1995.tb03401.x>, 1995.
- Miller, G. H., Geirsdóttir, Á., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson, C., Björnsson, H., and Thordarson, T.: Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks, *Geophys. Res. Lett.*, 39, n/a-n/a, <https://doi.org/10.1029/2011GL050168>, 2012.
- 645 Neukom, R., Steiger, N., Gómez-Navarro, J. J., Wang, J., and Werner, J. P.: No evidence for globally coherent warm and cold periods over the preindustrial Common Era, *Nature*, 571, 550–554, <https://doi.org/10.1038/s41586-019-1401-2>, 2019.
- Osborn, T. J., Briffa, K. R., and Jones, P. D.: Adjusting variance for sample-size in tree-ring chronologies and other regional mean series, *Dendrochronologia*, 15, 89–99, 1997.
- 650 PAGES 2k Consortium: Consistent multidecadal variability in global temperature reconstructions and simulations over the Common Era, *Nat. Geosci.*, 12, 643–649, <https://doi.org/10.1038/s41561-019-0400-0>, 2019.

- 655 PAGES 2k-PMIP3 group: Continental-scale temperature variability in PMIP3 simulations and PAGES 2k regional temperature reconstructions over the past millennium, *Clim. Past*, 11, 1673–1699, <https://doi.org/10.5194/cp-11-1673-2015>, 2015.
- PAGES2k Consortium: A global multiproxy database for temperature reconstructions of the Common Era, *Sci Data*, 4, 170088, <https://doi.org/10.1038/sdata.2017.88>, 2017.
- Pawlowicz, R.: *M_Map* A Mapping package for Matlab, 2022.
- 660 Polson, D., Hegerl, G. C., Zhang, X., and Osborn, T. J.: Causes of Robust Seasonal Land Precipitation Changes*, 26, 6679–6697, <https://doi.org/10.1175/JCLI-D-12-00474.1>, 2013a.
- Polson, D., Hegerl, G. C., Allan, R. P., and Sarojini, B. B.: Have greenhouse gases intensified the contrast between wet and dry regions?, *Geophys. Res. Lett.*, 40, 4783–4787, <https://doi.org/10.1002/grl.50923>, 2013b.
- 665 Rao, M. P., Cook, B. I., Cook, E. R., D'Arrigo, R. D., Krusic, P. J., Anchukaitis, K. J., LeGrande, A. N., Buckley, B. M., Davi, N. K., Leland, C., and Griffin, K. L.: European and Mediterranean hydroclimate responses to tropical volcanic forcing over the last millennium, *Geophys. Res. Lett.*, 44, 5104–5112, <https://doi.org/10.1002/2017GL073057>, 2017.
- 670 Santer, B. D.: Influence of Satellite Data Uncertainties on the Detection of Externally Forced Climate Change, 300, 1280–1284, <https://doi.org/10.1126/science.1082393>, 2003.
- Santer, B. D., Wigley, T. M. L., and Jones, P. D.: Correlation methods in fingerprint detection studies, *Climate Dynamics*, 8, 265–276, <https://doi.org/10.1007/BF00209666>, 1993.
- Schurer, A. P., Hegerl, G. C., Mann, M. E., Tett, S. F. B., and Phipps, S. J.: Separating Forced from Chaotic Climate Variability over the Past Millennium, 26, 6954–6973, <https://doi.org/10.1175/JCLI-D-12-00826.1>, 2013.
- 675 Schurer, A. P., Tett, S. F. B., and Hegerl, G. C.: Small influence of solar variability on climate over the past millennium, *Nature Geosci*, 7, 104–108, <https://doi.org/10.1038/ngeo2040>, 2014.
- 680 Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O. J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D. R., Pilcher, J. R., Salzer, M., Schüpbach, S., Steffensen, J. P., Vinther, B. M., and Woodruff, T. E.: Timing and climate forcing of volcanic eruptions for the past 2,500 years, *Nature*, 523, 543–549, <https://doi.org/10.1038/nature14565>, 2015.
- 685 Smerdon, J. E., Kaplan, A., Zorita, E., González-Rouco, J. F., and Evans, M. N.: Spatial performance of four climate field reconstruction methods targeting the Common Era, *Geophys. Res. Lett.*, 38, n/a–n/a, <https://doi.org/10.1029/2011GL047372>, 2011.
- St. George, S.: An overview of tree-ring width records across the Northern Hemisphere, *Quaternary Science Reviews*, 95, 132–150, <https://doi.org/10.1016/j.quascirev.2014.04.029>, 2014.
- St. George, S. and Ault, T. R.: The imprint of climate within Northern Hemisphere trees, *Quaternary Science Reviews*, 89, 1–4, <https://doi.org/10.1016/j.quascirev.2014.01.007>, 2014.
- 690 Stevenson, S., Fasullo, J. T., Otto-Bliesner, B. L., Tomas, R. A., and Gao, C.: Role of eruption season in reconciling model and proxy responses to tropical volcanism, *Proc. Natl. Acad. Sci. U.S.A.*, 114, 1822–1826, <https://doi.org/10.1073/pnas.1612505114>, 2017.
- 695 Tolwinski-Ward, S. E., Evans, M. N., Hughes, M. K., and Anchukaitis, K. J.: An efficient forward model of the climate controls on interannual variation in tree-ring width, *Clim Dyn*, 36, 2419–2439, <https://doi.org/10.1007/s00382-010-0945-5>, 2011a.
- Tolwinski-Ward, S. E., Evans, M. N., Hughes, M. K., and Anchukaitis, K. J.: Erratum to: An efficient forward model of the climate controls on interannual variation in tree-ring width, *Clim Dyn*, 36, 2441–2445, <https://doi.org/10.1007/s00382-011-1062-9>, 2011b.

- 700 Tolwinski-Ward, S. E., Anchukaitis, K. J., and Evans, M. N.: Bayesian parameter estimation and interpretation for an intermediate model of tree-ring width, *Clim. Past*, 9, 1481–1493, <https://doi.org/10.5194/cp-9-1481-2013>, 2013.
- Tolwinski-Ward, S. E., Tingley, M. P., Evans, M. N., Hughes, M. K., and Nychka, D. W.: Probabilistic reconstructions of local temperature and soil moisture from tree-ring data with potentially time-varying climatic response, *Clim Dyn*, 44, 791–806, <https://doi.org/10.1007/s00382-014-2139-z>, 2015.
- 705 Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500 BCE to 1900 CE, *Earth Syst. Sci. Data*, 9, 809–831, <https://doi.org/10.5194/essd-9-809-2017>, 2017.
- Vaganov, E. A., Hughes, M. K., and Šaškin, A. V.: Growth dynamics of conifer tree rings: images of past and future environments, Springer, Berlin, 354 pp., 2006.
- 710 Wang, J., Emile-Geay, J., Guillot, D., Smerdon, J. E., and Rajaratnam, B.: Evaluating climate field reconstruction techniques using improved emulations of real-world conditions, *Clim. Past*, 10, 1–19, <https://doi.org/10.5194/cp-10-1-2014>, 2014.
- Wigley, T. M. L., Briffa, K. R., and Jones, P. D.: On the Average Value of Correlated Time Series, with Applications in Dendroclimatology and Hydrometeorology, 23, 201–213, [https://doi.org/10.1175/1520-0450\(1984\)023<0201:OTAVOC>2.0.CO;2](https://doi.org/10.1175/1520-0450(1984)023<0201:OTAVOC>2.0.CO;2), 1984.
- 715 Wilson, R., Cook, E., D’Arrigo, R., Riedwyl, N., Evans, M. N., Tudhope, A., and Allan, R.: Reconstructing ENSO: the influence of method, proxy data, climate forcing and teleconnections, *J. Quaternary Sci.*, 25, 62–78, <https://doi.org/10.1002/jqs.1297>, 2010.
- 720 Zhao, S., Pederson, N., D’Orangeville, L., HilleRisLambers, J., Boose, E., Penone, C., Bauer, B., Jiang, Y., and Manzanedo, R. D.: The International Tree-Ring Data Bank (ITRDB) revisited: Data availability and global ecological representativity, *J Biogeogr*, 46, 355–368, <https://doi.org/10.1111/jbi.13488>, 2019.