

Variations in export production, lithogenic sediment transport and iron fertilization in the Pacific sector of the Drake Passage over the past 400 ka

5 María H. Toyos^{1,2,3}, Gisela Winckler^{4,5}, Helge W. Arz⁶, Lester Lembke-Jene³, Carina B. Lange^{2,7,8},
Gerhard Kuhn³, and Frank Lamy³

¹ Programa de Postgrados en Oceanografía, Departamento de Oceanografía, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Concepción, Chile

² Centro de Investigación Dinámica de Ecosistemas Marinos de Altas Latitudes (IDEAL), Universidad Austral de Chile, Valdivia, Chile

10 ³ Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar und Meeresforschung, Bremerhaven, Germany

⁴ Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964

⁵ Department of Earth and Environmental Sciences, Columbia University, New York, NY 10027

⁶ Leibniz-Institut für Ostseeforschung Warnemünde (IOW), Rostock-Warnemünde, Germany

⁷ Departamento de Oceanografía and Centro Oceanográfico COPAS Sur-Austral, Universidad de Concepción, Chile

15 ⁸ Scripps Institution of Oceanography, La Jolla, California 92037

Correspondence to: Maria H Toyos (mtoyos@udec.cl)

Abstract. Changes in Southern Ocean export production have broad biogeochemical and climatic implications. Specifically, iron fertilization likely increased subantarctic nutrient utilization and enhanced the efficiency of the biological pump during
20 glacial. However, past export production in the subantarctic Southeast Pacific is poorly documented, and its connection to Fe fertilization, potentially related to Patagonian Ice Sheet dynamics is unknown. We report on biological productivity changes over the past 400 ka, based on a combination of ²³⁰Th_{xs}-normalized and stratigraphy-based mass accumulation rates of biogenic barium, organic carbon, biogenic opal, and calcium carbonate as indicators of paleo-export production in a sediment core upstream of the Drake Passage (57.5° S; 70.3° W). In addition, we use fluxes of iron and lithogenic material as proxies for
25 terrigenous **matter input**, and thus potential micronutrient supply. Stratigraphy-based mass accumulation rates are strongly influenced by bottom-current dynamics, which result in variable sediment focussing or winnowing at our site. Carbonate is virtually absent in the core, except during peak interglacial intervals of the Holocene, and Marine Isotope Stages (MIS) 5 and 11, likely caused by transient decreases in carbonate dissolution. All other proxies suggest that export production increased during most glacial periods, coinciding with high iron fluxes. Such augmented glacial iron fluxes at the core site were most
30 likely derived from glaciogenic input from the Patagonian Ice Sheet promoting the growth of phytoplankton. Additionally, glacial export production peaks are also consistent with northward shifts of the Subantarctic and Polar Fronts, which positioned our site south of the Subantarctic Front and closer to silicic acid-rich waters of the Polar Frontal Zone, ~~as well as a with a decrease in the diatom utilization of Si relative to nitrate under Fe replete conditions~~. However, glacial export production near the Drake Passage was lower than in the Atlantic and Indian sectors of the Southern Ocean, which may relate to complete
35 consumption of silicic acid in the study area. Our results underline the importance of micro-nutrient fertilization through lateral terrigenous input from South America rather than aeolian transport, and exemplify the role of frontal shifts and nutrient limitation for past productivity changes in the Pacific entrance to the Drake Passage.

1 Introduction

The Southern Ocean (SO) plays an essential role in modulating glacial-interglacial variations of atmospheric pCO₂ (Sigman
40 et al., 2010). Increased biological export production, fueled by enhanced iron (Fe) fertilization in a more stratified glacial SO, is thought to have been a key driver of increased deep marine carbon storage (Jaccard et al., 2013). In the SO, the wind-driven,

eastward flowing Antarctic Circumpolar Current (ACC) enhances the air-sea exchange of CO₂ and the upwelling of nutrient- and CO₂-rich subsurface water masses (e.g., Marshall and Speer, 2012). Its flow is concentrated along several fronts, which are the Subantarctic Front (SAF), Polar Front (PF), and Southern ACC Front (SACCF; Orsi et al., 1995). These fronts act as barriers, inhibiting the exchange of the upwelled waters and their associated nutrients with neighboring fronts, and therefore also represent the limits between geochemical provinces (Chapman et al., 2020; Paparazzo, 2016). **Nevertheless, in some regions of the SO with weaker jets the mixing barrier effect is lower, allowing some degree of meridional exchange of nutrients and upwelled waters by eddy fields (Naveira Garabato et al., 2011).** Furthermore, these SO fronts are not stationary and their positions have been shown to change on seasonal to orbital timescales (e.g. Gille, 2014; Kemp et al., 2010). ~~The major~~
50 ~~constriction for the ACC flow and SO fronts is~~ The Drake Passage (DP), located between the southern tip of South America and the Antarctic Peninsula, **is a major constriction for the ACC flow and SO fronts.** At the DP entrance, a consistent pattern of glacial reduction of the ACC throughflow has been previously linked to a northward shift of the SAF (Lamy et al., 2015; Toyos et al., 2020).

Today, the SO represents the major region in the world ocean where the efficiency of the biological carbon pump, **the processes**
55 **by which marine organisms consume inorganic nutrients and CO₂ during photosynthesis and transform these elements into organic material into the deep sea,** is low. Incomplete nutrient utilization arises from a combination of high overturning rates, yielding fast replenishment of most macronutrients to the photic zone, and Fe limitation, restricting phytoplankton growth (Boyd et al., 2012; Moore et al., 2013). Therefore, mean chlorophyll-*a* concentrations above 2 mg m⁻³ are only found within 50 km off a continental or island coastline (Fig 1, Graham et al., 2015). Specifically, in the Subantarctic Zone (SAZ; the area
60 between the SAF and the Subtropical Front), waters feature excess nitrate relative to silicate (Dugdale et al., 1995) and Fe limits phytoplankton growth (Boyd et al., 1999). Thus, diatom production is relatively low, coccolithophores control primary production (Rigual-Hernández et al., 2015), and the biological pump is predominantly driven by carbonate producing organisms (Honjo, 2004). In contrast, the region between the SAF and the PF, known as the Polar Frontal Zone (the PFZ), is characterized by high abundances of large diatoms (Kopczynska et al., 2001), and by relatively low mean chlorophyll-*a*
65 concentrations (Fig 1, Graham et al., 2015).

The modern low export production in the SAZ contrasts with increased biological activity during glacials, predominantly fueled by enhanced Fe fertilization (e.g., Anderson et al., 2014; Jaccard et al., 2013; Kohfeld et al., 2005; Lamy et al., 2014; Martínez-García et al., 2009; Thöle et al., 2019). However, other mechanisms such as a more efficient diatom growth, shifts in the dominant plankton types, and increased nutrient utilization due to the lack of Fe limitation may also explain the glacial
70 increase in export production, which may lead to an increase in the biological carbon pump's efficiency in the SAZ (François et al., 1997; Galbraith and Skinner, 2020; Matsumoto et al., 2014). Of the proposed mechanisms, only Fe fertilization contributed significantly to the lowering of atmospheric CO₂, explaining ~~the last~~ 30 to 50 ppm of atmospheric CO₂ drawdown during the last glacial period (Kohfeld et al. 2005, Martin, 1990). It is widely believed that the Fe driving this fertilization of the glacial subantarctic SO is primarily delivered via aeolian dust (e.g., Martínez-García et al., 2009, 2014). This idea has been
75 corroborated by reconstructions of past glacial-interglacial variability in the southern westerly wind belt, which indicate a glacial strengthening and an equatorward migration and/or extension (Ho et al., 2012; Kohfeld et al., 2013; Lamy et al., 2014). In addition to dust input, especially in the vicinity of continents, Fe may also be brought to the surface ocean via continental runoff, iceberg transport or meltwater, and coastal upwelling (De Baar and De Jong, 2001). An increasing number of studies have recently shown that icebergs could provide a comparable or larger amount of bioavailable Fe than dust to the SO (e.g.,
80 Hopwood et al., 2019), and therefore significantly influence primary productivity (Wu and Hou, 2017). For instance, Fe associated with subglacial meltwater and icebergs stimulates and enhances marine primary productivity in ecosystems around the Antarctic Ice Sheet (Laufkötter et al., 2018; Vernet et al., 2011), and in the SAZ during the Last Glacial Maximum (LGM), where a combination of high ice discharge and slower iceberg melting due to colder sea surface temperatures increased the supply of Fe-rich terrigenous material (Wadham et al., 2019). It is well known that during glacial stages, erosion increased

85 significantly, enhancing physical weathering and, in turn, the sediment supply to the deep ocean. However, despite comprising a non-trivial portion of the SO, past export production in the Southeast Pacific, and its connection to direct Fe fertilization via meltwater or icebergs, potentially related to Patagonian Ice sheet (PIS) dynamics, remains unexplored. The subantarctic Pacific in the vicinity of the entrance to the DP is proximal to Patagonia, but likely does not receive substantial dust from this region, given the prevailing wind direction of the southern westerlies. In fact, Patagonian dust is predominantly transported eastward
90 to the SO's Atlantic sector (Li et al., 2008, 2010). Therefore, the DP entrance location might provide a unique opportunity to explore the PIS's potential as a direct source of Fe for fertilization of the southeastern Pacific during glacial intervals.


In this study, we reconstruct and characterize export production changes off southernmost Patagonia in the subantarctic Southeast Pacific over the past 400 ka and investigate ~~its~~ their link to Fe fertilization and SO frontal shifts. We use a combination of $^{230}\text{Th}_{\text{xs}}$ -normalized and stratigraphy-based mass accumulation rates of the lithogenic fraction, Fe, excess
95 Barium (Ba_{xs}), i.e. the fraction of Ba that is not supplied by terrigenous material, total organic carbon (TOC), biogenic opal, and carbonate (CaCO_3) from sediment core PS97/093-2 located near the SAF at the DP entrance (Fig 1). While Ba_{xs} and TOC ~~may~~ reflect the integrated total export production, CaCO_3 and biogenic opal indicate changes in the export production related to specific organisms. These are mainly coccolithophores and foraminifera for CaCO_3 and diatoms and radiolarians for opal (Dymond et al., 1992; Paytan, 2009). We show that variations in export production were closely linked to terrigenous sediment
100 and Fe delivery from the Patagonian hinterland, glacier dynamics and fundamental climate/ocean changes in the region. Lastly, we discuss potential underlying mechanisms, and evaluate our results with respect to published SAZ paleoproductivity reconstructions from other SO sectors.

2 Study area

Our site PS97/093 is located in the subantarctic Southeast Pacific, at the western entrance to the DP ($57^\circ 29.94' \text{ S}$; $70^\circ 16.48' \text{ W}$, 3781 m water depth). The DP, located between South America and the Antarctic Peninsula, is 850 km wide and represents the major geographical constriction for ACC transport into the Atlantic (Gordon et al., 1978). At present, the geostrophic transport of the ACC in the DP is associated with the SAF and PF, where strong surface and bottom velocities have been observed (Meredith et al., 2011; Renault et al., 2011). Furthermore, glacial-interglacial variability in bottom current circulation of *ca.* 16% during the last 1.3 Ma. has been reported at our coring site (Toyos et al., 2020). At present, Site PS97/093 is bathed
110 in Circumpolar Deep Water (CDW), which consists of a mixture of aged North Atlantic Deep Water that enters the SO through the Atlantic, recirculated Pacific and Indian Deep waters, and dense bottom waters from Antarctica (e.g., Carter et al., 2008). The carbonate ion concentration of CDW is nearly constant at $85 \pm 5 \mu\text{mol/kg}$ (Broecker and Clark, 2001), and it is slightly undersaturated with respect to calcium (Key et al., 2004). In the DP, the seafloor features, the position of the core southern westerly wind belt, and nutrient concentrations define the location of the ACC fronts (e.g., Ferrari et al., 2014; Meredith et al.,
115 2011; Paparazzo et al., 2016). Here, the PF is associated with the northern expression of the silicate front, indicating the geographical boundary between the silicate-poor waters to the north of it and silicate-rich waters to the south of it (Freeman et al., 2018, 2019). Our sediment core was retrieved *ca.* 40 km NW of the present-day position of the SAF, and *ca.* 350 km NW of the PF, within the main flow of the ACC (Fig 1b).

The DP is the region of the SO with the largest vertical velocities associated with topographically induced upwelling (Graham et al., 2015). Nevertheless, the DP is not considered an area of intensive phytoplankton growth, as illustrated by low chlorophyll-*a* concentrations (Fig 1; Demidov et al., 2011). Primary production is characterized by phytoplankton blooms in austral spring (Demidov et al., 2011), when Fe availability is the critical factor for the occurrence of the blooms (De Baar et al., 1995). Additionally, a lack of dissolved silica compared to phosphorus may limit the growth of siliceous phytoplankton species north of the PF (mainly diatoms, Demidov et al., 2011; Freeman et al., 2019). An analysis of the composition of the

125 surface sediments shows that in the DP a gradual change occurs from low organic carbon and high CaCO₃ content in the SAZ to opal-rich sediments at the Polar Front zone and in the Permanently Open Ocean Zone (Cárdenas et al., 2019). In the northern part of the DP, the provenance of terrigenous materials is restricted to proximal source regions, specifically southern Patagonia (Wu et al., 2019). Furthermore, analyses of surface sediments from the eastern South Pacific and DP have excluded both a westward dust transport pathway from southern South America dust sources to this region (Wengler et al., 130 2019), as well as and a substantial modern dust contribution from either Patagonia or Australia to the DP (Wu et al., 2019). After the Great Patagonian Glaciation (*ca.* 1 Ma), in at least five major glaciations, the Patagonian Andes were covered by a continuous mountain ice sheet, extending from 37°S to the Cape Horn and onto the Pacific Patagonian Shelf, with most ice likely calving into the Pacific Ocean on the western side, south of Chiloe island (Fig. 1; Davies et al., 2020; Gowan et al., 2021; Rabassa, 2008; Rabassa et al., 2011). As a result, at the southernmost Chilean continental margin, higher ice-rafted debris (IRD) contents occurred during cold intervals, interpreted as advances of the PIS (Caniupán et al., 2011). The distance 135 between core PS97/093-2 and the PIS during the last glacial (situated at that reached ~56°S at that time, Glasser et al., 2008) was *ca.* 180 km (Fig 1b).



140 Figure 1: (a) Map of the Southern Ocean mean chlorophyll-*a* concentrations for the years 2009–2019 with core locations. Yellow diamond indicates the location of core PS97/093-2 (this study), and white diamonds the location of published records discussed in the text: PS75/076 and PS75/059 (Lamy et al. 2014); PS2498-1, TN057-21, and TN057-06 (Anderson et al., 2014); ODP1090 (Martínez-García et al. 2014); PS2082-1 (Frank et al. 1999; Nürnberg, 1997); MD11-3357 (Thöle et al., 2019), and EPICA Dome C ice Core (Lambert et al., 2008). (b) Map of the Drake Passage mean chlorophyll-*a* concentrations for the years 2009–2019 showing the location of PS97/093-2 (this study), MD07-3128 (Caniupán et al., 2011), PS75/034-2 (Ho et al., 2012), GeoB3325-5 (Tapia et al., 2021), and ODP Site 1233 (Lamy et al., 2010; yellow diamonds); cyan area indicates the extension of the Patagonian Ice Sheet at the Last Glacial Maximum based on Glasser et al. (2008), white arrows show trajectories of the Antarctic Circumpolar Current (ACC), South Pacific Current (SPC), Cape Horn Current (CHC), Humboldt Current (Strub et al., 2019), and Antarctic Coastal Current (AACC, Deacon, 1984). Black lines mark ACC modern fronts (Orsi et al., 1995). SAF, Subantarctic Front; PF, Polar Front; SACCF, southern Antarctic Circumpolar Current Front, and SB, southern boundary of the ACC. We used the MODIS-Aqua Level-3 145 150

3 Material and methods

155 Piston core PS97/093-2 was retrieved from the Pacific entrance of the Drake Passage (57° 29.94' S; 70° 16.48' W; 3781 m
water depth; 16.45 m length; Fig. 1) during expedition PS97 "Paleo Drake" with R/V Polarstern (Lamy, 2016). Lithologically,
different types of hemipelagic sediments occur at this location that vary in composition on orbital timescales (Lamy, 2016).
Sediments from the interglacials MIS 11, MIS 5 and the Holocene are primarily composed of calcareous oozes (nannofossil
or foraminifera-nannofossil oozes) with minor concentrations of diatoms. In contrast, glacials and sediments from MIS 9, 7
160 and 3 are clayey silt with rare biogenic components, intercalated with some layers of diatomaceous fine-grained clayey silts.

3.1 Age model

The age model for core PS97/093-2 was developed by ~~taken from~~ Toyos et al. (2020) using the AnalySeries software (Paillard
et al., 1996), and is based ~~in~~ on a two-step approach: 1) establishment of a preliminary age model based on [shipboard physical
property data](#) and biostratigraphic time markers from calcareous nannofossils and diatoms, and 2) fine-tuning of the high-
165 resolution XRF-derived elemental Fe and Ca counts and CaCO₃ contents to the LR04 benthic δ¹⁸O stack (Lisiecki and Raymo,
2005). [The iron content is generally representative of the sediment's siliciclastic fraction, which is most likely controlled by a
combination of factors, including dilution of biogenic material \(primarily CaCO₃\) together with varying eolian and/or
glaciogenic sediment input from South America. CaCO₃ content usually is lower during glacial stages when enhanced detrital
fluxes suppress the relative amount of carbonate \(e.g., Diekmann, 2007\).](#) For tuning, we assumed that low Fe contents
170 characterize interglacial periods, whereas high contents represent glacials. Additionally, XRF Ca counts and CaCO₃ contents
were used for additional tuning in the intervals where they are present.

3.2 Bulk sediment parameters and geochemistry

Total carbon and nitrogen (TC, TN) were quantified using a CNS analyzer (Elementar Varia EL III) at the Alfred-Wegener-
Institute, Bremerhaven (AWI) using 100 mg of freeze-dried and homogenized sediment. Total organic carbon (TOC) contents
175 were measured with a carbon-sulfur determinator (CS-2000, ELTRA) after the removal of inorganic carbon (total inorganic
carbon) by adding 37% (vol/vol) of hydrochloric acid. CaCO₃ was calculated employing the standard equation Eq. (1):

$$CaCO_3[wt\%] = (TC[wt\%] - TOC[wt\%]) * 8.333 \quad (1)$$

Biogenic opal was determined at the Laboratory of Paleoceanography, University of Concepción (UdeC), Chile. The alkaline
180 extraction was conducted following the procedure described by Mortlock and Froelich, (1989), but using NaOH as a digestion
solution (Müller and Schneider, 1993). Between 50 and 70 milligrams of freeze-dried sediment were first treated with 10%
H₂O₂ and 1N HCl, and then extracted with 1M NaOH (40 mL; pH~13) at 85 °C for five hours. The analysis was carried out
by molybdate-blue spectrophotometry. Values are expressed as biogenic opal [percent](#) by multiplying the Si (%) by 2.4
(Mortlock and Froelich, 1989). We did not correct for the release of extractable Si from coexisting clay minerals, and thus
185 biogenic opal values could be overestimated (Schlüter and Rickert, 1998). Biogenic opal was also measured at AWI
Bremerhaven, though [at significantly lower temporal resolution and with smaller sample sizes \(30 mg\)](#), using the sequential
leaching method of Müller and Schneider (1993), and [differences](#) between the overlapping data [points](#) were observed. For
terrigenous contents exceeding 70%, opal concentrations measured at UdeC [were](#) consistently 3–5% higher than those
measured at AWI. When the lithogenic content was below 40%, the inter-lab difference was less than 1%. [The discrepancy](#)

190 between both datasets is most probably due to leaching of clay minerals like smectite (Cárdenas et al., 2019; Wu et al., 2019)
and a higher pH base employed at UdeC. Despite the difference in values, both records show a similar pattern of variability.
Given the importance of high-resolution data, we here use the opal results from UdeC. Dry bulk densities were quantified on
a total of 162 samples with a gas pycnometer (Micromeritics AccPyc II 1340) at AWI Bremerhaven, using the density
measurements of freeze-dried and homogenized bulk sediment samples and calculated by incorporating the water content of
195 the samples.

The archive half of core PS97/093-2 was measured with an AVAATECH X-Ray Fluorescence Core Scanner (XRF-CS) at
AWI Bremerhaven for high-resolution semi-quantitative element intensities of Ca, Fe, Ba and Ti at 0.5 cm resolution (0.5 x
1.2 cm measurement area, slit size down- and across-core). Three consecutive runs were performed with tube voltages of
10 kV (no filter), 30 kV (Pd-thick filter) and 50 kV (Cu filter), a current of 0.15, 0.175 and 1 mA, acquisition times of 10 s,
200 15 s and 20 s, respectively. Raw data were processed using Canberra *Eurisys'* iterative least squares software (WIN AXIL)
package. To obtain a high-resolution CaCO₃ record we calibrated the XRF Ca intensities with the bulk sediment CaCO₃
measurements ($r^2=0.92$, $n=157$, $P<0.0001$, Fig S1).

3.3 Elements and U/Th isotope analysis

Concentrations of Fe, Ti and Ba, along with U/Th isotopes, were determined at Lamont-Doherty Earth Observatory (LDEO).
205 Freeze dried samples (100 mg) were spiked with a ²³⁶U – ²²⁹Th solution, followed by complete acid digestion (Fleisher and
Anderson, 2003). Digests were taken up in 10 ml of 0.5 M HNO₃, and subsequently split in two aliquots. First, 0.4 ml was
diluted again in 0.5 M HNO₃ to get a final dilution of 2000x that was used for the determination of Al, Fe, Ti and Ba
concentrations. For the determination of U/Th isotopes, the remainder of the initial dilution (~ 9.6 ml) was utilized, with U/Th
purification achieved via Fe-coprecipitation and anion exchange chromatography following the methodology of Fleisher and
210 Anderson, (2003). To check for reproducibility and for quality control purposes, an internal sediment standard (VOICE Internal
MegaStandard (VIMS)) was run in each batch. U/Th isotopes were measured on a Thermo Scientific Element 2 ICP-MS, and
absolute elemental concentrations of Fe, Ti, and Ba were determined using an ICP-OES. Finally, to obtain a high-resolution
Fe record, we calibrated the 3-point smoothed Fe XRF intensities with bulk sediment (wt%) Fe from our discrete samples
($r^2=0.90$, $n=132$, $P<0.0001$) (Fig S1).

215 3.4 Excess barium

Ba_{xs} was calculated as:

$$Ba_{xs} = Ba_{total} - \left(Ti_{total} * \left[\frac{Ba}{Ti} \right]_{detrital} \right) \quad \text{Eq. (2):} \quad (2)$$

where Ba_{total} is the total measured Ba, Ti_{total} is the total measured Ti, and [Ba/Ti]_{detrital} is the ratio of Ba and Ti in crustal material
(assumed here to be 0.126 after Turekian and Wedepohl, 1961). This methodology assumes that 1) the major source of
220 elemental Ba to deep-sea sediments is marine barite, and 2) terrigenous material has a known and constant Ba/Ti ratio (e.g.,
Winckler et al., 2016). Our Ba_{xs} record was obtained calibrating the 3-point smoothed Ba and Ti XRF intensities with bulk
sediment (wt%) Ba ($r^2=0.76$, $n=132$, $P<0.0001$) and Ti ($r^2=0.79$, $n=132$, $P<0.0001$) (Fig S1).

3.5 Mass accumulation rates

Mass accumulation rates of individual sediment components (Fe, Ba_{xs}, TOC, CaCO₃ and biogenic opal) were calculated by
225 using the ²³⁰Th_{xs} normalization method (Bacon, 1984; Francois et al., 2004). ²³⁰Th is produced in the water column by decay
of ²³⁴U, and has a short residence time, settling quickly to the underlying sediments by proximal scavenging. As such, the flux

of ^{230}Th scavenged from the water column is considered to be nearly equal to its production rate. Therefore, the $^{230}\text{Th}_{\text{xs}}$ -normalized mass accumulation rate (MAR) for a given sample can be determined by Eq (3):

$$\text{MAR} = \frac{\beta \times z}{[^{230}\text{Th}_{\text{xs}}^0]}, \quad (3)$$

230 where $\beta \times z$ is the integrated ^{230}Th production in the overlying water column that depends on the water depth (z), and $^{230}\text{Th}_{\text{xs}}^0$ is the measured ^{230}Th activity after corrections for i) ^{230}Th supported by ^{238}U in detrital sediments, ii) ^{230}Th supported by authigenic ^{238}U from the seawater, and iii) radioactive decay of ^{230}Th since deposition. **Uncertainties of $^{230}\text{Th}_{\text{xs}}^0$ were propagated by considering analytical errors, uncertainties in the lithogenic and autigenic corrections, and decay corrections. In this study, if a sample has $^{230}\text{Th}_{\text{xs}} \leq 30\%$ of the total ^{230}Th concentration and a propagated uncertainty $\geq 60\%$, it has been excluded. We determine MAR back to *ca.* 400 ka, with the relative uncertainties of the $^{230}\text{Th}_{\text{xs}}$ constrained MAR increasing with the age of the sediments ~~due to uncertainties in the lithogenic and authigenic corrections.~~** $^{230}\text{Th}_{\text{xs}}$ -normalized mass accumulation rates of individual components were obtained by multiplying their respective concentrations by the ^{230}Th -derived MAR.

240 Additionally, stratigraphy-based bulk mass accumulation rates (BMAR) of individual components were obtained by multiplying the concentration of the respective component by the linear sedimentation rate and the dry bulk density.

3.6 Calculation of focusing factor

The degree of sediment focusing (Ψ) was calculated following the approach of Suman and Bacon, (1989) Eq (4):

$$\Psi = \left(\int_{r_1}^{r_2} r^2 {}^{230}\text{Th}_{\text{xs}}^0 \rho_r dr \right) / \beta_z (t_2 - t_1), \quad (4)$$

245 where ρ_r is the dry bulk density (g/cm^3), $^{230}\text{Th}_{\text{xs}}^0$ is the concentration of excess ^{230}Th in the sediment corrected for decay since deposition, t_1 and t_2 are the corresponding ages (kyr) of sediment depths r_1 and r_2 (cm), and β_z ($\text{dpm}/\text{cm}^2/\text{kyr}$) is the integrated ^{230}Th production in the overlying water column from ^{234}U decay. $\Psi > 1$ indicates sediment focusing, whereas $\Psi < 1$ denotes sediment winnowing. Values of $\Psi = 1$ indicate that the amount of ^{230}Th buried in the sediment is equivalent to the amount of ^{230}Th produced in the water column.

3.7 Lithogenic content

250 The concentrations of lithogenic material were calculated using ^{232}Th (e.g., Anderson et al., 2014; Winckler et al., 2008) assuming that i) ^{232}Th is exclusively of detrital origin (Francois et al., 2004), and ii) the ^{232}Th concentration of terrigenous material is relatively constant. ^{232}Th -derived lithogenic MAR were calculated by dividing the **lithogenic material concentration** by average ^{232}Th concentration of lithogenic material from Patagonia ($9 \mu\text{g}/\text{g}$, McGee et al., 2016).

255 Additionally, the lithogenic content was also determined by subtraction of the biogenic sediment components from the total bulk, Eq (5):

$$\text{lithogenic (wt\%)} = 100 - [\text{biogenic opal (wt\%)} + \text{CaCO}_3 \text{ (wt\%)} + (2\text{TOC (wt\%)})), \quad (5)$$


4 Results

260 CaCO_3 content is almost zero throughout most of the record, peaking only during MIS 11 ($\sim 80\%$, which coincides with a prominent layer of nannofossil ooze, Lamy, 2016), MIS 5 ($\sim 40\%$), and the Holocene ($\sim 40\%$, Fig 2). In contrast, **percentages of lithogenic material content** (^{232}Th -derived, and those obtained by subtracting biogenic sediment components from the total bulk) and Fe (wt%) show minima during these interglacials, and higher values during the rest of the record. Overall, the ^{232}Th -

based lithogenic fraction is 10% lower than the lithogenic fraction obtained by subtraction (Fig 2b). The Ba_{xs} concentrations show glacial-interglacial variability with broadly elevated higher values during interglacials. Specifically, in our Ba_{xs} record, interglacial peaks are more pronounced in MIS 5, 7 and 9 than in MIS 1 and MIS 11 (Fig 2d). whereas The percentage of TOC does not display glacial-interglacial variability, and values increase gradually from MIS 5e to the Holocene. The biogenic opal percentage shows strong similarities with the lithogenic and Fe records with pronounced minima of less than four percent during MIS11, MIS 5 and the Holocene, and relatively high values during the rest of the record (~ 7.5–10 %).

$CaCO_3$ and Ba_{xs} BMAR records vary in parallel with percentages of $CaCO_3$ and Ba_{xs} , respectively (Fig 2d, g), except for MIS 5e compared to MIS 5a–d, the latter caused by an increase in the sedimentation rates from ca. 0.5 to 2 cm/ka during MIS 5e (Figs 2d,g and 4). In contrast to the $CaCO_3$ and Ba_{xs} BMAR records, with all other BMAR of individual components that show no strong similarities with the corresponding percentages (Fig 2b, c, e, f). From MIS 11 to MIS 6/5 transition all other BMAR records generally co-vary, showing two small peaks during MIS 9 and MIS 7, and a larger peak during Termination II that continues through the beginning of MIS 5. However, from MIS 4 to the Holocene there are some differences between all records; the lithogenic, Fe and opal BMAR records increase gradually from MIS 4 to MIS 2, and decrease during the Holocene (Fig 2b, c, f), whereas TOC BMAR increases gradually from MIS 4 to the Holocene (Fig 2e).

$^{230}Th_{xs}$ -normalized mass accumulation rates (MAR) of individual components show that lithogenics, Fe, Ba_{xs} , TOC and opal co-vary and display certain glacial-interglacial variability, whereas $CaCO_3$ peaks when almost all other fluxes are low (Fig 3). From MIS 7, Lithogenic and Fe MARs show a pattern of higher values during glacials than in preceding interglacial stages and reach maxima during the MIS 6/5 transition. In contrast, the lowest lithogenic and Fe MARs are recorded in the Holocene, MIS 5, and possibly during MIS10-11 (Fig 3b, c). Opal MAR follows the same pattern with three strongest minima during MIS 11, MIS 5 and the Holocene. From mid MIS 9 to the end of MIS 6, the Opal MAR gradually increases until the end of MIS 6, reaching the highest values at the MIS 6/5 transition (Fig. 3g). Glacial-interglacial variability is not recorded between MIS 4 and MIS 2 in the opal record despite the relatively high values. TOC and Ba_{xs} MAR records are in good agreement with opal MAR except for the glacial peaks in the interval MIS 4 to MIS 2, and MIS11 (Fig. 3e, f). $CaCO_3$ MAR displays strong increases during MIS 11, MIS 5 and the Holocene only (Fig 3h).



290 Figure 2: Southeast Pacific (PS97/093-2) stratigraphy-based mass accumulation rates (BMAR) and percentages of individual components compared to dust flux in the EPICA Dome C (EDC) ice core (Lambert et al. 2008). (a) EDC dust flux; (b) Lithogenic content (wt. %; calculated from: $100 - [\text{opal wt. \%} + \text{CaCO}_3 \text{ wt. \%} + 2 \times \text{TOC wt. \%}]$, grey line) and lithogenic content based on ^{232}Th (grey stars); lithogenic BMAR (purple line); (c) Fe content (grey line) and Fe BMAR (green line); (d) Ba_{ss} content (grey line) and Ba_{ss} BMAR (pink line); (e) TOC content (light grey line) and TOC BMAR (light blue line); (f) Opal content (wt. %; grey line) and opal BMAR (blue line); (g) CaCO_3 content (grey line) and CaCO_3 BMAR (orange line). Numbers in the lower part of the figure indicate Marine Isotope Stages (MIS). Vertical grey bars mark glacial stages according to Lisiecki and Raymo (2005).
 295





Figure 3: Southeast Pacific (PS97/093-2) Comparison of export production and terrigenous fluxes $^{230}\text{Th}_{\text{xs}}$ -normalized mass accumulation rates (MAR) from Southeast Pacific core PS97/093-2 compared to dust flux in the EDC ice core (Lambert et al. 2008) and bottom current strengths of core PS97/093-2 (Toyos et al., 2020). (a) EDC dust flux, (b) ^{232}Th based lithogenic MAR; (c) Fe MAR; (d) changes in bottom current strength as indicated by the sortable silt record of core PS97/093-2 (note that Y-axis is reversed); (e) Ba_{xs} MAR; (f) TOC MAR; (g) biogenic opal MAR; and (h) CaCO_3 MAR. Shaded areas indicate associated errors. Numbers in the lower part of the figure indicate Marine Isotope Stages (MIS). Grey bars denote glacial stages according to Lisiecki & Raymo (2005). Dashed lines during MIS 6 denote the interval of extreme winnowing where MAR might be overestimated, and for MIS 10 they indicate uncertainty due to the lack of data points.

Focusing factors range from significant winnowing ($\Psi=0.08$) to considerable focusing ($\Psi=3.55$) throughout the past 400 ka. At the coring site, winnowing dominates, and we only have brief periods of focusing during the peak interglacials MIS 9, MIS 5e and the Holocene. The intervals from MIS 11 to MIS 9 and from mid-MIS 8 to MIS 6 show net winnowing, whereas the interval between mid-MIS 9 to mid-MIS 8 was net neutral, and from MIS 5e to MIS 2 was nearly neutral with moderate winnowing (Fig 4b). Furthermore, the focusing factor record broadly agrees with the sedimentation rates, with winnowing in the intervals of low sedimentation rates and focusing during periods of relatively high sedimentation rates (Fig 4b, c).

A comparison between the MAR and the BMAR (Figs 4d, e) shows discrepancies between both records. For most of the record, MAR is higher than BMAR, except for the focusing intervals (MIS 9, MIS 5e and the Holocene), where an opposite trend is observed. The largest divergence between both records happens during MIS 6, which is characterized by significant

315 winnowing (Fig 4). As a result, during MIS 9, MIS 6, MIS 5e and the Holocene, the BMAR and MAR of the individual components (lithogenic, Fe, Ba_{xc}, TOC, opal, and CaCO₃) diverge from each other to varying degrees (Table 1), showing only similar values from mid-MIS 5 to MIS 2 and in MIS 9/8 (Figs 2 and 3).



320 **Figure 4: Comparison of sediment mass accumulation rates, focusing factors, bottom current strength and sedimentation rates of core PS97/093-2 for the last 400 ka. (a) Changes in bottom current strength based on the Sortable Silt grain size record (lilac line, Toyos et al., 2020) compared to (b) focusing factors (red line, black dashed line marks focusing factor=1); (c) sedimentation rates; (d) ²³⁰Th_{xs}- normalized mass accumulation rate (MAR, black line and dots) and associated errors (2σ; pink shadow), which grow with the age of the sample, dashed lines indicate intervals with uncertainties due to extreme winnowing (Marine Isotope Stage 6) or lack of data (Marine Isotope Stage 10); (e) Stratigraphy-based bulk mass accumulation rates (BMAR) obtained by multiplying sedimentation rates by dry bulk densities. Numbers in the lower part of the figure indicate Marine Isotope Stages (MIS). Grey bars denote glacial stages according to Lisiecki & Raymo (2005).**

325

330

335 **Table 1: Comparison between the average MAR and BMARs for the intervals with high (0–15 kyr, 324–334 kyr) and low (212–243 kyr, 243–279 kyr) focusing factors.**

average intervals	Accumulation rate of Fe (mg/cm ² /kyr)	Accumulation rate of Ba _{xs} (mg/cm ² /kyr)	Accumulation rate of TOC (mg/cm ² /kyr)	Accumulation rate of opal(mg/cm ² /kyr)	Accumulation rate of CaCO ₃ (mg/cm ² /kyr)	
0–15 kyr ($\Psi=2.07$)	30.06	0.53	1.07	53.04	185	MAR
	40.65	0.94	1.82	70.05	411	BMARs
	1.35	1.77	1.70	1.32	2.21	overestimation factor
324–334 kyr ($\Psi=3.55$)	55.44	0.46	0.59	125.46	11.50	MAR
	72.99	1.32	0.82	184.97	12.53	BMARs
	1.31	2.84	1.38	1.47	1.08	overestimation factor
212–243 kyr ($\Psi=0.40$)	77.49	0.52	1.17	153.67	4.52	MAR
	34.21	0.30	0.50	69.14	1.40	BMARs
	2.26	1.71	2.35	2.22	3.21	underestimation factor
243–279 kyr ($\Psi=0.24$)	47.11	0.29	0.66	86.05	6.62	MAR
	12.03	0.07	0.23	22.99	2.07	BMARs
	3.91	3.75	2.83	3.74	3.00	underestimation factor

5 Discussion

5.1 Influence of current dynamics on sediment redistribution and its effect on proxy record interpretations

340 The study area at the Pacific entrance of the DP is strongly influenced by the ACC, as documented by highly variable current strengths on millennial and glacial-interglacial timescales (Lamy et al., 2015; Toyos et al., 2020; Wu et al., 2021). Our results indicate that this variability in current strength influenced changes in sedimentation patterns (Fig. 4). Specifically, ~~our~~ the focusing factors ~~indicate~~ imply that sediment redistribution is highly variable in our core, encompassing significant winnowing during MIS 6 (Fig 4b). Under these extreme conditions winnowing may bias the ²³⁰Th normalization reconstructions because of a preferential removal of the fine ²³⁰Th-rich grains (Marcantonio et al., 2014). Costa and McManus (2017) documented a potential overestimation of MAR when winnowing is severe (focusing factors of 0.158–0.20). With moderate winnowing $\Psi > 0.24$, this effect is negligible and ²³⁰Th_{xs}-normalized accumulation rates are robust. Therefore, the high MAR during MIS 6 in core PS97/093-2 may in part stem from the extreme winnowing ($\Psi=0.08$) at that time, whereas in all other intervals Ψ exceeds 0.24 and thus estimated MARs are reliable (Fig 4).

350 In our core, we observe winnowing during glacial periods in intervals with relatively low current strengths, and increased focusing factors contemporaneous with increased sortable silt values in the same core, reflecting strengthened bottom currents during MIS 9, MIS 5e and the Holocene (Toyos et al., 2020; Fig. 4a-c). The consistent pattern of winnowing (focusing) in intervals with relatively low (high) current strength might at first sight seem counterintuitive. We suggest that ~~Since~~ under strong bottom current conditions, the fine sediment fraction (<10 μ m) potentially behaves similarly to the >63 μ m ~~sediment~~ fraction because of cohesive effects and flocculation of the fine fraction during transport (McCave and Hall, 2006), ~~but~~ whereas 355 under slower current conditions a loss of such effects in the fine sediment fraction might trigger the winnowing of ~~the fine~~ such sediment fraction (Marcantonio et al., 2014). We propose that winnowing at our site, in a generally fine-grained sediment setting, is due to current velocities that occasionally ~~are overruling~~ overrule the cohesive forces, resulting in the loss of a small amount of the fine fraction. Prior studies have pointed out that in the deep sea, current velocities between *ca.* 6.5 and 10.5 cm/s are enough to trigger surface sediment erosion of the aggregated particulate matter that compose the surface sediments (Peine 360 et al., 2009; Turnewitsch et al., 2008). However, our results suggest that even lower flow speeds of ~5.5–6 cm/s during glacials (Fig 4, Toyos et al., 2020, calibrated following Wu et al., 2019) reached the threshold of intermittent erosion and resuspension

of the fine sediment fraction. During interglacial periods MIS 9, MIS 5 and the Holocene, strong currents plausibly caused a gradual mobilization of more and more coarse-grained material that was laterally accumulated in certain areas, resulting in a coarser-grained focusing at our site. A similar pattern, characterized by strengthened bottom currents inferred from grain size analysis, and increased sediment focusing has also been observed for interglacials on the Weddell Sea continental margin (Frank et al., 1996).

The dynamic bottom water circulation [appears to have](#) led to frequent syndepositional redistribution. For core PS97/093-2, when comparing MAR and BMAR of Fe, Ba_{xs}, TOC, opal and CaCO₃ in the intervals with high and low focusing factors (excluding MIS 6, where we cannot discard an overestimation of the ²³⁰Th_{xs,0}-constrained fluxes because of extreme winnowing), large discrepancies are seen (Table 1, Figs 2 and 3). BMAR of the individual components are up to three times higher than MAR in intervals with highest focusing, and ~~ea-~~ [about](#) four times smaller in intervals with pronounced winnowing (Table 1). These differences confirm that the PS97/093-2 BMAR are strongly biased by sediment redistribution due to current dynamics. Given that the PS97/93 site is affected by substantial lateral redistribution of sediment particles, we therefore base our paleoproductivity reconstruction on the MAR, which allows for the quantification of lateral sediment redistribution and accurate vertical rain rates (e.g., Costa et al., 2020; Suman and Bacon, 1989).

5.2 Southeast Pacific lithogenic material, sources and iron fertilization potential

A comparison of lithogenic fluxes in core PS97/093-02 with [eolian dust](#) records from open ocean sites in the SAZ (Anderson et al., 2014, Lamy et al., 2014; Martínez-García et al., 2009; Thöle et al., 2019) and in Antarctic ice cores (Lambert et al., 2008), shows ~~a-~~ [generally](#) similar patterns (Fig. 5). However, for MIS 10 and MIS 8, some characteristic features are not observed in our sediment core due to a lower temporal resolution that prevents an accurate evaluation during such intervals (Fig 5). Our glacial lithogenic MAR peaks were one order of magnitude higher than those ~~reconstructions from observed in~~ the open Pacific, Atlantic and Indian SAZ (Fig 5). Furthermore, studies of modern surface sediments [do no indicate](#) ~~discard~~ a substantial contribution of dust to the DP region and the Southeast Pacific north of the SAF (Wengler et al., 2019; Wu et al., 2019). Thus, accounting [for these very high lithogenic fluxes in the proximity of the DP requires](#) an additional non-eolian source of terrigenous material ~~is required~~.




Figure 5: Changes in ^{230}Th -normalized lithogenic MARs in the subantarctic Southern Ocean (see Fig 1 for location of sites discussed). From top to bottom: Cores PS75/076 and PS75/059 (Lamy et al., 2014); Core PS97/093-2 (this study); Core PS2498 (Anderson et al., 2014); Cores TN057-21 and TN056-06 (Anderson et al., 2014), and ODP1090 (Martínez-García et al., 2009); Core MD11-3357 (Thöle et al., 2019), and dust MAR in the EPICA Dome C ice core (Lambert et al., 2008). Marine Isotope Stage (MIS); grey bars denote glacial stages. Continental masses are located in their relative longitudinal position to the marine sediment cores and represent the primary source of terrigenous inputs to the Subantarctic Zone.

We suggest that the comparably high lithogenic MARs observed in our core are due to the relative proximity of our site to southern South America as a substantial source of runoff-derived and glaciogenic sediments, particularly during glacial stages when large parts of the southern Andes were covered by the PIS (Fig 1b). South America is considered a substantial source of runoff-derived and glaciogenic sediments, particularly during glacial stages when large parts of the southern Andes were covered by the PIS (Fig 1b). Therefore, we suggest that the PS97/093-2's high lithogenic MARs are related to the relative proximity of our site to southern South America. From MIS 6 onwards, PS97/093-2 lithogenic MARs reveal significant increases during glacial stages. Specifically, the most intense growth occurred during MIS 6, due to a combination of the penultimate local glaciation (Rabassa, 2008) and a possible winnowing bias, followed by a smaller rise during MIS 4, and a gradual increase to higher values in the LGM. This pattern closely corresponds to reported Patagonian ice expansions (e.g., Gowan et al., 2021; Lowell et al., 1995; Rabassa, 2008; Rabassa and Clapperton, 1990; Fig S2). Furthermore, an IRD record from Core MD07-3128,

retrieved from the Pacific entrance to the Strait of Magellan (*ca.* 620 km NNW of core PS97/093-2, Fig 1b), shows a consistent
405 pattern of higher IRD contents during cold intervals over the past ~60 kyr, which have been interpreted as advances of the PIS
(Caniupán et al., 2011). Moreover, enhanced Fe concentrations linked to PIS advances during colder periods were reported at
ODP Site 1233 from the Chilean continental margin (Fig 1b, Kaiser and Lamy, 2010; Lamy et al., 2004), and Antarctic dust
~~peaks maxima in Antarctica~~ and ~~have been mechanistically linked to the presence of active outwash plains in Patagonia~~
~~advances of the PIS~~ (Sugden et al. (2009). ~~This mechanism explains enhanced dust mobilisation and uptake from these outwash~~
410 ~~for transport to Antarctica.~~

It has been suggested that Fe bio-availability depends on the mineralogy of the detrital fraction (Schroth et al., 2009).
Therefore, it is unclear which amount of the Fe that reached our site was bioavailable, and how this availability changed
through time. Previous studies suggest that physical weathering increases the labile primary Fe (II) content of sediments that
is associated with phytoplankton fertilization (Shoenfelt et al., 2017, 2018), indicating that a stronger contribution of
415 glaciogenic sediments ~~might~~ increase the proportion of bioavailable Fe in a given sediment flux (Shoenfelt et al., 2019). ~~As~~
~~Nevertheless, since~~ the study site is proximal to the continental sources (tip of South America), which are dominated by
physical weathering during both glacial and interglacial intervals, we hypothesize that temporal changes in the proportion of
bioavailable Fe in the Southeast Pacific are less pronounced than in more open ocean locations. We thus expect that the Fe
supply delivered to our site might have always been relatively enriched in bio-available Fe compared to other oceanographic
420 sectors. In summary, because dust does not seem to be the main source of lithogenic material at site PS97/093, we propose
that advances of the PIS and iceberg discharge during glacial intervals (e.g., Kaiser et al., 2007; Rabassa, 2008) could have
increased lithogenic fluxes and potentially bio-available Fe reaching the subantarctic Pacific entrance to the DP and thus
promoting biological productivity.

In the present-day Atlantic SAZ, chlorophyll-*a* blooms are supported by an advective supply of Fe from the South American
425 coastline along the western boundary current first and the Subtropical Front later, rather than by continental dust sources
(Graham et al., 2015). Other studies also invoke this mechanism for the LGM, where increased supply of glaciogenic debris
by expanded Patagonian and Antarctic Peninsula ice sheets travel long distances via the ACC at depth (Noble et al., 2012).
However, in the subantarctic region of our core location, the glacial northward shift of the SAF likely decreased the DP
throughflow of the ACC, weakening the cold water route into the Atlantic (Lamy et al., 2015; Toyos et al., 2020), which would
430 have hindered the transport of Fe released from the PIS to the Southeast Pacific into the Atlantic. It has been suggested that
the reduced supply to the Atlantic basin through the DP during glacials might have been compensated by a stronger
recirculation within the South Pacific, causing a stronger South Pacific Gyre recirculation (Lamy et al., 2015). Therefore, the
bioavailable Fe might have been transported in the South Pacific gyre, enhancing productivity in this region, which is currently
limited by Fe availability (Bonnet et al., 2008). To evaluate the reach of Fe fertilization in the Southeast Pacific linked to PIS
435 dynamics, further reconstructions along the Pacific SAZ and Cape Horn-Humboldt current systems ~~and iron speciation data~~
are needed. Currently, the knowledge is restricted to one recently published record (core GeoB3327-5, located at 43°S, *ca.* 400
km off the Chilean coast) that reportedly featured increases in primary production during MIS 2–4, linked to increased supply
of micronutrients via continental runoff, primarily controlled by PIS variability or precipitation (Tapia et al., 2021).

5.3 Orbital-scale variations and drivers of export production in the Southeast Pacific SO

440 5.3.1 ~~Assessing the proxies' response to~~ Export production fluxes

Our ²³⁰Th-normalized results show that the proxies for integrated export production (Ba_{xs} MAR and TOC MAR) generally co-
vary and correlate with biogenic opal MAR; all being higher during MIS 6 and MIS 4–2, and lower during ~~full interglacials~~
~~(the Holocene, and MIS 5 and MIS 11; PAGES, 2016) and~~ (Table S1, Fig 3).

Usually, variable dissolution in the water column does not significantly affect opal burial in the SO (Chase et al., 2015),
445 suggesting that opal MAR might be only marginally affected by preservation changes, thus providing valid information on

past diatom productivity changes (e.g., Bradtmiller et al., 2009; Sprenk et al., 2013). Since the preservation of organic carbon in sediments is globally scarce (only about 1% of the TOC produced in the water column gets preserved in the sediments, Berger et al., 1989), the good correspondence and positive correlation between opal and TOC MARs ($r^2=0.56$, Table S1, Fig 3) might imply that the TOC preserved in the sediments is related to carbon locked in diatom frustules. Opal MAR also correlates with Ba_{xs} MAR, ~~which has a preservation rate of ca. 30% in sediments under oxic conditions (Dymond, 1992)~~, suggesting that most of the glacial export production was fueled by diatoms. An exception to the strong correlation between the export production proxies occurred from MIS 4 to MIS 2, when TOC and Ba_{xs} MARs show relatively high values with glacial/interglacial variability, whereas the opal record is rather flat within this interval. Several studies proposed that diatom growth rate was ~~inferred from opal burial during the LGM may have been~~ more effective during the LGM compared to ~~than~~ today (resulting in higher opal burial during the LGM), as ~~because~~ diatoms can reduce their Si/C uptake ratio under Fe-replete conditions (Anderson et al., 2002; Chase et al., 2015). Thus, a glacial increase of Fe input may have created conditions under which relatively more TOC and biogenic barium were exported per unit of opal buried (Fig. 3).

Unlike our other MAR records, high $CaCO_3$ MAR values are observed only during the Holocene, MIS 5 and MIS 11 (Fig. 3). To explain the different pattern, we have to consider changes in carbonate chemistry and variations in the depth of the lysocline affecting our site's carbonate contents through time. At present, the lysocline ~~occurs~~ is located at ca. 4000 m (Sulpis et al., 2018), which is only slightly below the depth of our core (3781 m water depth). Site PS97/093 is bathed in Circumpolar Deep Water, which is characterized by relatively low $[CO_3^{2-}]$, being undersaturated with respect to $CaCO_3$, and therefore promotes carbonate dissolution (Key et al., 2004). In the SO, the lysocline depth shoaled at least 0.5 km during the glacial stages of the past 500 kyr, triggering enhanced carbonate dissolution and concomitant reduced carbonate MAR (Howard and Prell, 1994). Moreover, poor carbonate preservation has also been observed in another Southeast Pacific core with an alkenone-based SST record (PS75/034-2, 54° 22'S, 80° 05'W, 4425 m water depth; Ho et al., 2012, Fig 1b). Nevertheless, since carbonate preservation is favored in intervals with high sedimentation rates (Gottschalk et al., 2018), in our core the relatively high sedimentation rates during MIS1–3 or MIS 5e (Fig 4c) might allow the burial of carbonates in the sediment before being dissolved during such intervals. On the other hand, we suggest that the absence of carbonates between MIS 10 to MIS 6, and in MIS 4 indicates carbonate dissolution, rather than changes in export production of coccolithophores and foraminifera (Fig. 3). As a result, $CaCO_3$ MAR in our record are only present in the globally strong interglacials MIS 11, MIS 5 and the Holocene (PAGES, 2016), while during weaker interglacials, such as MIS 7 and MIS 9, carbonate productivity did not offset carbonate dissolution.

5.3.2 Drivers of local export production patterns

In the DP, the oceanographic fronts may act as barriers to mixing in the ocean, preventing cross-frontal movement and meridional exchange of water masses and their associated nutrients (Naveira Garabato et al., 2011; Paparazzo, 2016). On longer timescales, studies of sediment cores located upstream and downstream of the northern and central DP suggest a northward shift of the SO frontal system during glacial times, lower ACC flow speed and reduced transport through the DP (Lamy et al., 2015; Roberts et al., 2017; Toyos et al., 2020; Wu et al., 2021). Particularly at site PS97/093, a comparison between the \overline{SS} record and the opal and TOC MAR records shows that decreases in current strengths correspond with increases in opal and TOC (Fig 3). The inverse correspondence between current strength and opal export supports the idea that the glacial decrease in current vigor, linked to the northward shift of the frontal systems, would locate our coring site south of the strongest ACC flow in the vicinity of the SAF (Toyos et al., 2020) and closer to the Si-rich southern waters. Furthermore, it is likely that the waters between the PF and SAF received an additional supply of dissolved $Si(OH)_4$ during glacial intervals, as inferred from the displacement of the opal belt north of the present day PF position (e.g., Diekmann, 2007). This pattern is explained by upwelling of dissolved $Si(OH)_4$ south of the PF as it does today, but due to expanded ice cover during glacials, very little silica was used by diatoms south of the PF, and the upwelled $Si(OH)_4$ was transported northwards from the PF

(Chase et al., 2003; Freeman et al., 2018). At the same time, we discard a substantial arrival at our core site of either hydrothermally sourced Fe from the East Pacific Rise (Fitzsimmons et al., 2014) or potentially upwelled Fe south of the PF. In ~~the hypothetical~~ case of upwelling of Fe south of the PF, its strong particle reactivity might have caused efficient scavenging of Fe beneath the ice (Chase et al., 2003). ~~Therefore, enhanced nutrient utilization and opal production can only be achieved through an additional Fe source.~~ In our core, the coeval increase in MAR of lithogenics, Fe, biogenic opal, TOC and Ba_{xs} suggests that sufficient Fe was transported to our site to consume all the supplied Si(OH)₄ during glacial intervals. Furthermore, under Fe-replete conditions, diatoms take up substantially less silicic acid relative to nitrate (Matsumoto et al., 2014). Modern observations and incubation experiments show that the Si:N uptake ratio lowers from ~ 4:1 to ~ 1:1 (Brzezinski et al., 2002; Franck et al., 2000). Therefore, since our core received enough glacial Fe to relieve the limitation, the enhanced diatom production, indicated by higher biogenic opal MAR, may ~~be the result of respond to~~ a reduced diatom Si:N uptake ratio, combined with an increase of the amount of Si(OH)₄ that reached the core site.

In contrast with this glacial scenario, increases in bottom current strength, usually during interglacials, indicate that the SAF was located south of our site, resulting in a production regime similar to present-day conditions. This area is characterized by low average chlorophyll-*a* concentrations (Fig 1b), where the lack of dissolved silica may limit the growth of diatoms (Demidov et al., 2011; Freeman et al., 2019), Fig 6a and 6b) and might cause coccolithophores and other phytoplankters to become the dominant group (Rigual Hernández et al., 2020; Saavedra-Pellitero et al., 2019). However, the absence of peaks in the TOC and Ba_{xs} MAR during the Holocene and MIS 5 suggests that the increased CaCO₃ export production did not ~~impact~~ significantly ~~influence in~~ the integrated proxies of export production, implying ~~no~~ noticeable impact ~~by calcareous organisms~~ on the total export during such intervals (Fig 3).

Unlike other interglacials, during MIS 11, a prominent nannofossil ooze, primarily composed of *Geophyrocapsa* coccolithophore and foraminifera has been reported in the area (Lamy, 2016; Toyos et al., 2020) that is mirrored at our site by CaCO₃ MARs one order of magnitude higher than in the Holocene and in MIS 5, and by increases in the Ba_{xs} and TOC MARs (Fig 3). A similar rise in CaCO₃ accumulation has also been recognized at other SAZ locations in the Atlantic (Hodell et al., 2000) and the Pacific (Gersonde, 2011; Saavedra-Pellitero et al., 2017) sectors of the SO, suggesting that the increase in CaCO₃ MAR during MIS 11 in core PS97/093-2 is part of a general pattern comprising at least the Pacific and Atlantic SAZ, likely caused by an exceptional southward migration of the ACC frontal systems in the Pacific sector of the SO (Saavedra-Pellitero et al., 2017).

5.3.3 Comparison of PS97/093-2 export fluxes/MARs to other Southern Ocean SAZ sites

In order to assess the ~~spatially complex~~ zonal dynamics of productivity and lithogenic flux variations along the Southern Ocean's SAZ, we compare our MAR reconstructions with previously published records from open ocean areas of the SAZ (Anderson et al., 2014; Frank et al., 1999; Lamy et al., 2014; Martínez-García et al., 2014; Nürnberg et al., 1997; Thöle et al., 2019). We use a circum-Antarctic subantarctic transect of MAR of lithogenics, Fe, Ba_{xs}, TOC, opal and CaCO₃ for the Holocene (0–10 ka), the LGM (19–27 ka; Clark et al., 2009), the last interglacial MIS 5e (119–124 ka), and the end of MIS 6 during full glacial conditions, just before the beginning of the warming in the SO at 132–131 ka (Bianchi and Gersonde, 2002; Fig 6). Overall, our productivity indicators agree with subantarctic records from other locations, displaying higher CaCO₃ during interglacials and higher opal, TOC and Ba_{xs} during glacials, the latter pattern most likely enabled by an increase of lithogenic Fe fluxes (Martin, 1990; Fig 6). But, as it has previously been reported, it is likely that other factors than Fe ~~played~~ ~~a role~~ in regulating the biological production in the SO (Anderson et al., 2014; Kohfeld et al., 2005).




Figure 6: Comparison of MAR across the Subantarctic Southern Ocean, during the Holocene (0–10 ka, red), the Last Glacial Maximum (19–27 ka, blue), Marine Isotope Stage 5e (119–124 ka, yellow), and late Marine Isotope Stage 6 (132–150 ka, green): Maps of surface water nitrate (a) and silicate (b) concentrations (data from WOCE, Global Hydrographic Climatology, Gouretski and Koltermann, 2004); black lines indicate the modern location of the Subantarctic Front (SAF), and Polar Front (PF, Orsi et al., 1995); white diamonds refer to core locations (PS75/076 and PS75/059, Lamy et al. 2014; PS97/093-2, this study; PS2498-1, TN057-21 and TN057-06, Anderson et al., 2014; ODP1090, Martínez-García et al. 2014; PS2082-1, Nürnberg, 1997 and Frank, 2002; MD11-3357, Thöle et al., 2019); (c) lithogenic flux; (d) Fe flux; (e) Ba_s flux; (f) total organic carbon flux; (g) opal flux; (h) CaCO₃ flux. Grey bars indicate the projected core positions.

At our core location, we observe that the LGM lithogenic (Fe) MARs were about eightfold (ninefold) larger than during the Holocene. Furthermore, our site records the highest glacial lithogenic and Fe fluxes of the subantarctic SO; the LGM Indian–Atlantic and Pacific lithogenic fluxes are four to eleven times lower than at the PS97/093 site (Fig 6). However, the corresponding glacial increases in our productivity MARs are not as high as expected when considering the relative increase in the lithogenic MAR. In the Atlantic and Indian sectors, the LGM rise in export production was larger than the increase in lithogenic fluxes, whereas the opposite pattern is observed in our core (Fig 6). Furthermore, the glacial and interglacial export production fluxes are higher in the Atlantic and Indian sectors than at the DP entrance and in the central Pacific (Fig 6). Whereas in the central Pacific, the relatively low export production is explained by weaker Fe fertilization caused by lower

local glacial dust fluxes (Lamy et al., 2014), the results at our site indicate that export production did not respond exclusively
545 to glacial Fe fertilization ~~as efficiently as in the other sectors of the SAZ~~ (Fig 6). Since it takes very little bioavailable iron to
get out of the Fe-limited conditions (*ca.* 3nM, Boyd et al., 2000), and our core site is proximal to Fe sources, which might
facilitate a faster transition from a Fe-deplete to Fe-replete conditions. ~~Therefore~~, we suggest that under bioavailable Fe-replete
conditions, another mechanism than Fe fertilization might ultimately regulate export production at our site.

The region is currently nitrate and phosphate-rich (Fig 6a; phosphate shown in Fig S3), and previous works north of the PF
550 have shown that productivity increased during the LGM, whereas nitrate utilization decreased only slightly, implying a glacial
intensification in the supply of nitrate to surface waters (François et al., 1997; Martínez-García et al., 2014). Since nitrate is
not the macronutrient that was limiting export production during glacials, we propose Si(OH)₄ as a suitable candidate.

Under high Fe conditions, a glacial northward migration of the SO frontal system, together with the transport of silicic acid
northwards of the PF (Chase et al., 2003), and a reduced Si:N uptake ratio by diatoms (Brzezinski et al., 2002) should have
555 contributed to the relief of silicate limitation in the SAZ, resulting in an increase in export production (predominantly diatom
production). Given that LGM (and the possibly overestimated MIS 6) opal fluxes at our site were over fivefold larger than
during the Holocene (Fig 6g), to sustain such a diatom productivity implicated by the opal flux would require a supply of
Si(OH)₄ much greater than exists today (Anderson et al., 2014). Thus, we propose that if all the Si(OH)₄ brought to our site
was consumed entirely, silica (diatom) production would have been inhibited by Si(OH)₄ limitation, as it is happening today
560 in some areas of the Polar Front Zone, where nearly depleted levels of silicic acid have been reported (Mengelt et al., 2001,
Fig 6b). This hypothesis might be supported by modern experiments that show that Si(OH)₄ limitation significantly restrict
the response of diatom production to Fe in the SAZ (Brzezinski et al., 2005). Consequently, the growth of other phytoplankton
groups (e.g., small flagellates) or small, weakly-silicified diatom species would have been favored, which are strongly affected
by remineralization, thus lowering export production (Buesseler, 1998). Alternatively, since increases in productivity at the
565 SO fronts are explained by horizontal advection rather than upwelling (Graham et al., 2015), it might be possible that during
past glacials only a portion of the Si(OH)₄ supplied via upwelling at the PF travelled northwards across the PFZ reaching the
core location. The other portion of the upwelled silica might have been advected laterally into the Atlantic via the ACC
(Graham et al., 2015), causing a more intensive phytoplankton growth there instead of in the DP. Although the latter scenario
is somewhat speculative, both assumptions (a moderate glacial Si(OH)₄ limitation and dominance of horizontal advection
570 rather than upwelling) are not mutually exclusive; hence the two scenarios would explain the weaker productivity response to
Fe at the PS97/093 site compared to other subantarctic records.

6 Conclusions

We present a multi-proxy approach for reconstructing paleo-export production in the SAZ at the Pacific entrance to the DP
covering the past 400 ka. As our core is located in an area with relatively high bottom current speeds, we show that our site is
575 strongly affected by the lateral redistribution of sediments, with high focusing (winnowing) factors associated with high (low)
bottom current speeds. This counterintuitive pattern is most likely due to a dominance of the coarse (fine)-grained fraction
together with a preservation (loss) of the flocculation and cohesive effects of the fine sediment fraction under relatively high
(slow) bottom currents. As a result, focusing (winnowing) leads to an overestimation (underestimation) of the BMAR by a
factor of *ca.* 3 (4) during such intervals in our core. The frequent syndepositional redistribution emphasizes the importance of
580 the utilization of the ²³⁰Th_{xs}-normalization method to reconstruct fluxes in the subantarctic SO.

Whereas our lithogenic record follows the temporal pattern of other lithogenic records at open ocean sites in the SAZ and in
the Antarctic ice core (EDC dust flux), we present evidence that dust was not the main source of lithogenic material. Instead,
advances of the PIS during glacial intervals increased the portion of glaciogenic sediments and therefore released lithogenics.

585 ~~Since the core site is relatively proximal to continental sources, which are dominated by physical weathering and likely enriched in bioavailable Fe, we suggest that the Southeast Pacific (Drake Passage) received enough glacial bioavailable Fe to not limit phytoplankton growth.~~

Our results show orbital-scale variations from a predominantly high opal-rich export production for most glacials to lower levels of export productivity during interglacials dominated by carbonates. We suggest that such fluctuations are responding to glacial-interglacial frontal shifts coupled with nutrient limitation. We hypothesize that during interglacials the SAF was located slightly south of our core location, so that a reduction in Fe and dissolved silica supply may have limited the growth of siliceous phytoplankton (diatoms), leading to calcareous plankton (i.e. coccolithophores) becoming the dominant group. Yet, the reduced total export production during such intervals suggests that export production by calcareous organisms was too small to have a noticeable impact on the total export, except for MIS 11, where the impact was perceptible, as demonstrated by the occurrence of ~~due to~~ a prominent calcareous ooze. Conversely, the glacial northward migration of the SO frontal system would have positioned the core south of the SAF and closer to Si-rich waters in an environment characterized by extremely high lithogenic and Fe fluxes from a more extensive PIS as mentioned above. The coeval increase in lithogenic, Fe, opal, TOC, and Ba_{xs} suggests that our core received sufficient Fe and Si(OH)₄ to fuel higher export production during glacial intervals.

A spatial comparison with previously published records from different locations in the SAZ shows that our core displays the highest glacial lithogenic and Fe MARs. However, the corresponding glacial increases in productivity fluxes are not as high as we expected given the relative increase in lithogenics, suggesting that export production did not respond to glacial Fe fertilization as efficiently as in other sectors of the SAZ. We hypothesize that under these Fe-repleted glacial conditions, a depletion in the silicic acid that was transported northwards of the PF, could explain the glacial export production pattern at our core location.

Data availability

605 All data are available in PANGAEA repository: <https://doi.pangaea.de/10.1594/PANGAEA.934588>

Supplement

The supplement related to this article is available online at: XXXX

Author contribution

This study is part of MHT's thesis under the supervision of CBL and FL. MHT carried out most of the measurements. MHT, GW and HWA made the mass accumulation rate calculations. MHT, CBL, FL, HWA, LLJ and GW analysed and interpreted the data. MHT wrote the first draft and produced the figures for the manuscript with substantial contributions of GW, HWA, LLJ, CBL and FL. All authors interpreted, edited, and reviewed the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

615 Acknowledgements

This work was funded by the Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung through its research programs “PACES-II” and “Changing Earth – Sustaining our Future”. Additional [financial](#) support was provided by the Chilean oceanographic centers FONDAP-IDEAL (project number 1500003) and COPAS Sur-Austral (AFB170006) to CBL and MHT, and Lamont-Doherty Earth Observatory. MHT acknowledges support from scholarship CONICYT-
620 PCHA/Doctorado Nacional/2016-21160454, Postgraduate Office of Universidad de Concepción, Red Clima Red (project number LPR163), and Doctorado MaReA. [We acknowledge the comments and suggestions by Dr. Giuseppe Cortese and Dr. Louisa Bradtmiller, which helped us to improve the revised version of this manuscript.](#) We thank the captain, crew and scientific party of R/V Polarstern for a successful PS97 cruise. We acknowledge A. Ávila for technical support at the Paleooceanography lab of Universidad de Concepción, R. Schwarz, M. Fleisher and J. Abell for assistance at Lamont-Doherty
625 Earth Observatory, and S. Wiebe, R. Fröhlking and V. Schumacher for technical support at AWI.

References

- Anderson, R. F., Chase, Z., Fleisher, M. Q. and Sachs, J.: The Southern Ocean’s biological pump during the Last Glacial Maximum, *Deep. Res. Part II Top. Stud. Oceanogr.*, 49(9–10), 1909–1938, doi:10.1016/S0967-0645(02)00018-8, 2002.
- Anderson, R. F., Barker, S., Fleisher, M., Gersonde, R., Goldstein, S. L., Kuhn, G., Mortyn, P. G., Pahnke, K. and Sachs, J.
630 P.: Biological response to millennial variability of dust and nutrient supply in the Subantarctic South Atlantic Ocean, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 372(2019), doi:10.1098/rsta.2013.0054, 2014.
- de Baar, H. J. W. and De Jong, J. T. M.: Distributions, sources and sinks of iron in seawater, in *The Biogeochemistry of Iron in Seawater*, edited by D. R. . Turner and K. A. Hunter, pp. 123–253, John Wiley & Sons Limited., 2001.
- de Baar, H. J. W., Bathmann, U., Smetacek, V., Löscher, B. M. and Veth, C.: Importance of iron for plankton blooms and
635 carbon dioxide drawdown in the Southern Ocean, *Nature*, 373(6513), 412–415, doi:10.1038/373412a0, 1995.
- Bacon, M. P.: Glacial to interglacial changes in carbonate and clay sedimentation in the Atlantic Ocean estimated from 230Th measurements., *Isot. Geosci.*, 2(2), 97–111, 1984.
- Berger, W. H., Smetacek, V. and Wefer, G.: Ocean Productivity and Paleoproductivity - An Overview, in *Productivity of the Ocean: Present and Past*, edited by W. . Berger, V. Smetacek, and G. Wefer, pp. 1–34, John Wiley & Sons Limited., 1989.
- 640 Bianchi, C. and Gersonde, R.: The Southern Ocean surface between Marine Isotope Stages 6 and 5d: Shape and timing of climate changes, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 187(1–2), 151–177, doi:10.1016/S0031-0182(02)00516-3, 2002.
- Bonnet, S., Guieu, C., Bruyant, F., Prášil, O., Van Wambeke, F., Raimbault, P., Moutin, T., Grob, C., Gorbunov, M. Y., Zehr, J. P., Masquelier, S. M., Garczarek, L. and Claustre, H.: Nutrient limitation of primary productivity in the Southeast Pacific (BIO-SOPE cruise), *Biogeosciences*, 5(1), 215–225, doi:10.5194/bg-5-215-2008, 2008.
- 645 Boyd, P., LaRoche, J., Gall, M., Frew, R. and McKay, R. M. L.: Role of iron, light, and silicate in controlling algal biomass in subantarctic waters SE of New Zealand, *J. Geophys. Res. Ocean.*, 104(C6), 13395–13408, doi:10.1029/1999JC900009, 1999.

- Boyd, P. W., Watson, a J., Law, C. S., Abraham, E. R., Trull, T., Murdoch, R., Bakker, D. C., Bowie, a R., Buesseler, K. O., Chang, H., Charette, M., Croot, P., Downing, K., Frew, R., Gall, M., Hadfield, M., Hall, J., Harvey, M., Jameson, G., LaRoche, J., Liddicoat, M., Ling, R., Maldonado, M. T., McKay, R. M., Nodder, S., Pickmere, S., Pridmore, R., Rintoul, S., Safi, K., Sutton, P., Strzepek, R., Tanneberger, K., Turner, S., Waite, A. and Zeldis, J.: A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization, *Nature*, 407(6805), 695–702, doi:10.1038/35037500, 2000.
- Boyd, P. W., Arrigo, K. R., Strzepek, R. and Van Dijken, G. L.: Mapping phytoplankton iron utilization: Insights into Southern Ocean supply mechanisms, *J. Geophys. Res. Ocean.*, 117(6), 1–18, doi:10.1029/2011JC007726, 2012.
- 655 Bradtmiller, L. I., Anderson, R. F., Fleisher, M. Q. and Burekle, L. H.: Comparing glacial and Holocene opal fluxes in the Pacific sector of the Southern Ocean, *Paleoceanography*, 24(2), 1–20, doi:10.1029/2008PA001693, 2009.
- Broecker, W. S. and Clark, E.: Glacial-to-Holocene Redistribution of Carbonate Ion in the Deep Sea, *Science*, 294(5549), 2152–2155, doi:10.1126/science.1064171, 2001.
- Brzezinski, M. A., Pride, C. J., Franck, V. M., Sigman, D. M., Sarmiento, J. L., Matsumoto, K., Gruber, N., Rau, G. H. and Coale, K. H.: A switch from Si(OH) 4 to NO 3 – depletion in the glacial Southern Ocean, *Geophys. Res. Lett.*, 29(12), 1564, doi:10.1029/2001GL014349, 2002.
- 660 Brzezinski, M. A., Jones, J. L. and Demarest, M. S.: Control of silica production by iron and silicic acid during the Southern Ocean Iron Experiment (SOFeX), *Limnol. Oceanogr.*, 50(3), 810–824, doi:10.4319/lo.2005.50.3.0810, 2005.
- Buesseler, K. O.: The decoupling of production and particulate export in the surface ocean, *Global Biogeochem. Cycles*, 12(2), 297–310, doi:10.1029/97GB03366, 1998.
- 665 Caniupán, M., Lamy, F., Lange, C. B., Kaiser, J., Arz, H., Kilian, R., Baeza Urrea, O., Aracena, C., Hebbeln, D., Kissel, C., Laj, C., Mollenhauer, G. and Tiedemann, R.: Millennial-scale sea surface temperature and Patagonian Ice Sheet changes off southernmost Chile (53 °s) over the past ~60 kyr, *Paleoceanography*, 26(3), doi:10.1029/2010PA002049, 2011.
- Cárdenas, P., Lange, C. B., Vernet, M., Esper, O., Srain, B., Vorrath, M.-E., Ehrhardt, S., Müller, J., Kuhn, G., Arz, H. W., Lembke-Jene, L. and Lamy, F.: Biogeochemical proxies and diatoms in surface sediments across the Drake Passage reflect oceanic domains and frontal systems in the region, *Prog. Oceanogr.*, 174, 72–88, doi:10.1016/j.pcean.2018.10.004, 2019.
- 670 Carter, L., McCave, I. N. and Williams, M. J. M.: Chapter 4 Circulation and Water Masses of the Southern Ocean: A Review, in *Developments in Earth and Environmental Sciences*, vol. 8, edited by F. Florindo and M. Soegert, pp. 85–114, Elsevier B.V., Amsterdam., 2008.
- 675 Chapman, C. C., Lea, M.-A., Meyer, A., Sallée, J.-B. and Hindell, M.: Defining Southern Ocean fronts and their influence on biological and physical processes in a changing climate, *Nat. Clim. Chang.*, 10(3), 209–219, doi:10.1038/s41558-020-0705-4, 2020.
- Chase, Z., Anderson, R. F., Fleisher, M. Q. and Kubik, P. W.: Accumulation of biogenic and lithogenic material in the Pacific sector of the Southern Ocean during the past 40,000 years, *Deep. Res. Part II Top. Stud. Oceanogr.*, 50(3–4), 799–832, doi:10.1016/S0967-0645(02)00595-7, 2003.
- 680

- Chase, Z., Kohfeld, K. E. and Matsumoto, K.: Controls on biogenic silica burial in the Southern Ocean, *Global Biogeochem. Cycles*, 29(10), 1599–1616, doi:10.1002/2015GB005186, 2015.
- Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J. X., Hostetler, S. W. and McCabe, A. M.: The Last Glacial Maximum, *Science*, 325(5941), 710–714, doi:10.1126/science.1172873, 2009.
- 685 Costa, K. and McManus, J.: Efficacy of ²³⁰Th normalization in sediments from the Juan de Fuca Ridge, northeast Pacific Ocean, *Geochim. Cosmochim. Acta*, 197, 215–225, doi:10.1016/j.gca.2016.10.034, 2017.
- Costa, K. M., Hayes, C. T., Anderson, R. F., Pavia, F. J., Bausch, A., Deng, F., Dutay, J. C., Geibert, W., Heinze, C., Henderson, G., Hillaire-Marcel, C., Hoffmann, S., Jaccard, S. L., Jacobel, A. W., Kienast, S. S., Kipp, L., Lerner, P., Lippold, J., Lund, D., Marcantonio, F., McGee, D., McManus, J. F., Mekik, F., Middleton, J. L., Missiaen, L., Not, C., Pichat, S.,
- 690 Robinson, L. F., Rowland, G. H., Roy-Barman, M., Tagliabue, A., Torfstein, A., Winckler, G. and Zhou, Y.: ²³⁰Th Normalization: New Insights on an Essential Tool for Quantifying Sedimentary Fluxes in the Modern and Quaternary Ocean, *Paleoceanogr. Paleoclimatology*, 35(2), 1–36, doi:10.1029/2019PA003820, 2020.
- Davies, B. J., Darvill, C. M., Lovell, H., Bendle, J. M., Dowdeswell, J. A., Fabel, D., García, J.-L., Geiger, A., Glasser, N. F., Gheorghiu, D. M., Harrison, S., Hein, A. S., Kaplan, M. R., Martin, J. R. V., Mendelova, M., Palmer, A., Pelto, M., Rodés, Á., Sagredo, E. A., Smedley, R. K., Smellie, J. L. and Thorndycraft, V. R.: The evolution of the Patagonian Ice Sheet from 35
- 695 ka to the present day (PATICE), *Earth-Science Rev.*, 204, 103152, doi:10.1016/j.earscirev.2020.103152, 2020.
- Deacon, G.: The Antarctic Circumpolar Ocean. *Studies in Polar Research Series*, viii, in *Geological Magazine*, vol. 122, edited by Cambridge University Press., pp. 306–306, Cambridge, London, New York, New Rochelle, Melbourne, Sydney., 1984.
- Demidov, A. B., Mosharov, S. A., Gagarin, V. I. and Romanova, N. D.: Spatial variability of the primary production and
- 700 chlorophyll-*a* concentration in the drake passage in the austral spring, *Oceanology*, 51(2), 281–294, doi:10.1134/S0001437011020056, 2011.
- Diekmann, B.: Sedimentary patterns in the late Quaternary Southern Ocean, *Deep. Res. Part II Top. Stud. Oceanogr.*, 54(21–22), 2350–2366, doi:10.1016/j.dsr2.2007.07.025, 2007.
- Dugdale, R. C., Wilkerson, F. P. and Minas, H. J.: The role of a silicate pump in driving new production, *Deep Sea Res. Part*
- 705 *I Oceanogr. Res. Pap.*, 42(5), 697–719, doi:10.1016/0967-0637(95)00015-X, 1995.
- Dymond, J.: Barium in deep sea sediments: A geo-chemical proxy for paleoproductivity: *Paleoceanography*, *Paleoceanography*, 7, 163–181, doi:0883-8305/92/92PA-00181, 1992.
- Ferrari, R., Jansen, M. F., Adkins, J. F., Burke, A., Stewart, A. L. and Thompson, A. F.: Antarctic sea ice control on ocean circulation in present and glacial climates, *Proc. Natl. Acad. Sci.*, 111(24), 8753–8758, doi:10.1073/pnas.1323922111, 2014.
- 710 Fitzsimmons, J. N., Boyle, E. A. and Jenkins, W. J.: Distal transport of dissolved hydrothermal iron in the deep South Pacific Ocean, *Proc. Natl. Acad. Sci.*, 111(47), 16654–16661, doi:10.1073/pnas.1418778111, 2014.
- Fleisher, M. Q. and Anderson, R. F.: Assessing the collection efficiency of Ross Sea sediment traps using ²³⁰Th and ²³¹Pa, *Deep. Res.*, 50, 693–712, doi:10.1016/S0967-0645(02)00591-X, 2003.

- 715 Franck, V. M., Brzezinski, M. A., Coale, K. H. and Nelson, D. M.: Iron and silicic acid concentrations regulate Si uptake north and south of the Polar Frontal Zone in the Pacific Sector of the Southern Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 47(15–16), 3315–3338, doi:10.1016/S0967-0645(00)00070-9, 2000.
- Francois, R., Frank, M., Rutgers van der Loeff, M. M. and Bacon, M. P.: 230 Th normalization: An essential tool for interpreting sedimentary fluxes during the late Quaternary, *Paleoceanography*, 19(1), PA1018, doi:10.1029/2003PA000939, 2004.
- 720 François, R., Altabet, M. A., Yu, E.-F., Sigman, D. M., Bacon, M. P., Frank, M., Bohrmann, G., Bareille, G. and Labeyrie, L. D.: Contribution of Southern Ocean surface-water stratification to low atmospheric CO₂ concentrations during the last glacial period, *Nature*, 389(6654), 929–935, doi:10.1038/40073, 1997.
- Frank, M., Mangini, A., Gersonde, R., Rutgers van der Loeff, M. and Kuhn, G.: Late Quaternary sediment dating and quantification of lateral sediment redistribution applying 230Th_{ex}: a study from the eastern Atlantic sector of the Southern Ocean, *Geol. Rundschau*, 85(3), 554–566, doi:10.1007/BF02369010, 1996.
- 725 Frank, M., Gersonde, R. and Mangini, A.: Sediment Redistribution, 230Th_{ex}- Normalization and Implications for the Reconstruction of Particle Flux and Export Paleoproductivity, in *Use of Proxies in Paleoceanography*, pp. 409–426, Springer Berlin Heidelberg, Berlin, Heidelberg., 1999.
- Frank, M.: Accumulation rate and vertical rain rate of sediment core PS2082-1. *PANGAEA*,
730 <https://doi.org/10.1594/PANGAEA.81101>, 2002.
- Freeman, N. M., Lovenduski, N. S., Munro, D. R., Krumhardt, K. M., Lindsay, K., Long, M. C., and MacLennan, M.: The Variable and Changing Southern Ocean Silicate Front: Insights from the CESM Large Ensemble. *Global Biogeochemical Cycles*, 32(5), 752–768. <https://doi.org/10.1029/2017GB005816>, 2018.
- Freeman, N. M., Munro, D. R., Sprintall, J., Mazloff, M. R., Purkey, S., Rosso, I., DeRanek, C. A. and Sweeney, C.: The
735 Observed Seasonal Cycle of Macronutrients in Drake Passage: Relationship to Fronts and Utility as a Model Metric, *J. Geophys. Res. Ocean.*, 124(7), 4763–4783, doi:10.1029/2019JC015052, 2019.
- Gersonde, R.: The expedition of the research vessel “Polarstern” to the polar South Pacific in 2009/2010 (ANT-XXVI/2 - BIPOMAC), Gersonde, edited by H. Bornemann; and B. Chiaventone, *Berichte zur Polar- und Meeresforschung (Reports on Polar and Marine Research)*, Bremerhaven, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven., 2011.
- 740 Gille, S. T.: Meridional displacement of the Antarctic Circumpolar Current, *Philos. Trans. R. Soc.*, 372, 20130273, doi:<http://dx.doi.org/10.1098/rsta.2013.0273>, 2014.
- Glasser, N. F., Jansson, K. N., Harrison, S. and Kleman, J.: The glacial geomorphology and Pleistocene history of South America between 38°S and 56°S, *Quat. Sci. Rev.*, 27(3–4), 365–390, doi:10.1016/j.quascirev.2007.11.011, 2008.
- Gordon, A. L., Molinelli, E. and Baker, T.: Large-scale relative dynamic topography of the Southern Ocean, *J. Geophys. Res.*,
745 83(C6), 3023, doi:10.1029/jc083ic06p03023, 1978.

- Gottschalk, J., Hodell, D. A., Skinner, L. C., Crowhurst, S. J., Jaccard, S. L. and Charles, C.: Past Carbonate Preservation Events in the Deep Southeast Atlantic Ocean (Cape Basin) and Their Implications for Atlantic Overturning Dynamics and Marine Carbon Cycling, *Paleoceanogr. Paleoclimatology*, 33(6), 643–663, doi:10.1029/2018PA003353, 2018.
- 750 Gouretski, V. V. and Koltermann, K. P.: The World Ocean Circulation Experiment (WOCE) Global Hydrographic Climatology, 35th ed., edited by B. BHS, Hamburg. [online] Available from: <http://rda.ucar.edu/datasets/ds285.4/>, 2004.
- Gowan, E. J., Zhang, X., Khosravi, S., Rovere, A., Stocchi, P., Hughes, A. L. C., Gyllencreutz, R., Mangerud, J., Svendsen, J.-I. and Lohmann, G.: A new global ice sheet reconstruction for the past 80 000 years, *Nat. Commun.*, 12(1), 1199, doi:10.1038/s41467-021-21469-w, 2021.
- 755 Graham, R. M., De Boer, A. M., van Sebille, E., Kohfeld, K. E. and Schlosser, C.: Inferring source regions and supply mechanisms of iron in the Southern Ocean from satellite chlorophyll data, *Deep. Res. Part I Oceanogr. Res. Pap.*, 104, 9–25, doi:10.1016/j.dsr.2015.05.007, 2015.
- Ho, S. L., Mollenhauer, G., Lamy, F., Martínez-García, A., Mohtadi, M., Gersonde, R., Hebbeln, D., Nunez-Ricardo, S., Rosell-Melé, A. and Tiedemann, R.: Sea surface temperature variability in the Pacific sector of the Southern Ocean over the past 700 kyr, *Paleoceanography*, 27(4), PA4202, doi:10.1029/2012PA002317, 2012.
- 760 Hodell, D. A., Charles, C. D. and Ninnemann, U. S.: Comparison of interglacial stages in the South Atlantic sector of the Southern Ocean for the past 450 kyr: Implications for Marine Isotope Stage (MIS) 11, *Glob. Planet. Change*, 24(1), 7–26, doi:10.1016/S0921-8181(99)00069-7, 2000.
- Honjo, S.: Particle export and the biological pump in the Southern Ocean, *Antarct. Sci.*, 16(4), 501–516, doi:10.1017/S0954102004002287, 2004.
- 765 Hopwood, M. J., Carroll, D., Höfer, J., Achterberg, E. P., Meire, L., Le Moigne, F. A. C., Bach, L. T., Eich, C., Sutherland, D. A. and González, H. E.: Highly variable iron content modulates iceberg-ocean fertilisation and potential carbon export, *Nat. Commun.*, 10(1), doi:10.1038/s41467-019-13231-0, 2019.
- Howard, W. R. and Prell, W. L.: Late Quaternary CaCO₃ production and preservation in the Southern Ocean: Implications for oceanic and atmospheric carbon cycling, *Paleoceanography*, 9(3), 453–482, doi:10.1029/93PA03524, 1994.
- 770 Jaccard, S. L., Hayes, C. T., Martínez-García, A., Hodell, D. A., Anderson, R. F., Sigman, D. M. and Haug, G. H.: Supplementary materials: Two modes of change in Southern Ocean productivity over the past million years, *Science*, 339(6126), 1419–1423, doi:10.1126/science.1227545, 2013.
- Kaiser, J. and Lamy, F.: Links between Patagonian Ice Sheet fluctuations and Antarctic dust variability during the last glacial period (MIS 4-2), *Quat. Sci. Rev.*, 29(11–12), 1464–1471, doi:10.1016/j.quascirev.2010.03.005, 2010.
- 775 Kaiser, J., Lamy, F., Arz, H. W. and Hebbeln, D.: Dynamics of the millennial-scale sea surface temperature and Patagonian Ice Sheet fluctuations in southern Chile during the last 70 kyr (ODP Site 1233), *Quat. Int.*, 161(1), 77–89, doi:10.1016/j.quaint.2006.10.024, 2007.

- 780 Kemp, A. E. S., Grigorov, I., Pearce, R. B. and Naveira Garabato, A. C.: Migration of the Antarctic Polar Front through the mid-Pleistocene transition: evidence and climatic implications, *Quat. Sci. Rev.*, 29(17–18), 1993–2009, doi:10.1016/j.quascirev.2010.04.027, 2010.
- Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C. and Peng, T. H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cycles*, 18(4), 1–23, doi:10.1029/2004GB002247, 2004.
- 785 Kohfeld, K. E., Le Quéré, C., Harrison, S. P. and Anderson, R. F.: Role of marine biology in glacial-interglacial CO₂ cycles, *Science*, 308(5718), 74–78, doi:10.1126/science.1105375, 2005.
- Kohfeld, K. E., Graham, R. M., de Boer, A. M., Sime, L. C., Wolff, E. W., Le Quéré, C. and Bopp, L.: Southern Hemisphere westerly wind changes during the Last Glacial Maximum: Paleo-data synthesis, *Quat. Sci. Rev.*, 68(April 2013), 76–95, doi:10.1016/j.quascirev.2013.01.017, 2013.
- 790 Kocczynska, E. E., Dehairs, F., Elskens, M. and Wright, S.: Phytoplankton and microzooplankton variability between the Subtropical and Polar Fronts south of Australia: Thriving under regenerative and new production in late summer, *J. Geophys. Res. Ocean.*, 106(C12), 31597–31609, doi:10.1029/2000JC000278, 2001.
- Lambert, F., Delmonte, B., Petit, J. R., Bigler, M., Kaufmann, P. R., Hutterli, M. A., Stocker, T. F., Ruth, U., Steffensen, J. P. and Maggi, V.: Dust - Climate couplings over the past 800,000 years from the EPICA Dome C ice core, *Nature*, 452(7187), 616–619, doi:10.1038/nature06763, 2008.
- 795 Lamy, F.: The expedition PS97 of the research vessel POLARSTERN to the Drake Passage in 2016, Bremerhaven, Germany., 2016, [Berichte zur Polar- und Meeresforschung = Reports on Polar and Marine Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany. 701, 571 p., doi:10.2312/BzPM_0701_2016, 2016.](#)
- Lamy, F., Kaiser, J., Ninnemann, U., Hebbeln, D., Arz, H. W. and Stoner, J.: Antarctic Timing of Surface Water Changes off Chile and Patagonian Ice Sheet Response, *Science*, 304(5679), 1959–1962, doi:10.1126/science.1097863, 2004.
- 800 Lamy, F., Gersonde, R., Winckler, G., Esper, O., Jaeschke, A., Kuhn, G., Ullermann, J., Martinez-Garcia, A., Lambert, F. and Kilian, R.: Increased Dust Deposition in the Pacific Southern Ocean During Glacial Periods, *Science*, 343(6169), 403–407, doi:10.1126/science.1245424, 2014.
- Lamy, F., Arz, H. W., Kilian, R., Lange, C. B., Lembke-Jene, L., Wengler, M., Kaiser, J., Baeza-Urrea, O., Hall, I. R., Harada, N. and Tiedemann, R.: Glacial reduction and millennial-scale variations in Drake Passage throughflow., *Proc. Natl. Acad. Sci. U. S. A.*, 112(44), 13496–501, doi:10.1073/pnas.1509203112, 2015.
- 805 Laufkötter, C., Stern, A. A., John, J. G., Stock, C. A. and Dunne, J. P.: Glacial Iron Sources Stimulate the Southern Ocean Carbon Cycle, *Geophys. Res. Lett.*, 45(24), 13,377–13,385, doi:10.1029/2018GL079797, 2018.
- Li, F., Ginoux, P. and Ramaswamy, V.: Distribution, transport, and deposition of mineral dust in the Southern Ocean and Antarctica: Contribution of major sources, *J. Geophys. Res.*, 113(D10), D10207, doi:10.1029/2007JD009190, 2008.

- 810 Li, F., Ginoux, P. and Ramaswamy, V.: Transport of Patagonian dust to Antarctica, *J. Geophys. Res. Atmos.*, 115(18), 1–9, doi:10.1029/2009JD012356, 2010.
- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20(1), PA1003, doi:10.1029/2004PA001071, 2005.
- Lowell, T. V., Heusser, C. J., Andersen, B. G., Moreno, P. I., Hauser, A., Heusser, L. E., Schluchter, C., Marchant, D. R. and
815 Denton, G. H.: Interhemispheric Correlation of Late Pleistocene Glacial Events, *Science*, 269(5230), 1541–1549, doi:10.1126/science.269.5230.1541, 1995.
- Marcantonio, F., Lyle, M. and Ibrahim, R.: Particle sorting during sediment redistribution processes and the effect on ^{230}Th -normalized mass accumulation rates, *Geophys. Res. Lett.*, 41(15), 5547–5554, doi:10.1002/2014GL060477, 2014.
- Marshall, J. and Speer, K.: Closure of the meridional overturning circulation through Southern Ocean upwelling, *Nat. Geosci.*,
820 5(3), 171–180, doi:10.1038/ngeo1391, 2012.
- Martin, J. H.: Glacial-interglacial CO_2 change: The Iron Hypothesis, *Paleoceanography*, 5(1), 1–13, doi:10.1029/PA005i001p00001, 1990.
- Martínez-García, A., Rosell-Melé, A., Geibert, W., Gersonde, R., Masqué, P., Gaspari, V. and Barbante, C.: Links between iron supply, marine productivity, sea surface temperature, and CO_2 over the last 1.1 Ma, *Paleoceanography*, 24(1), PA1207, 825 doi:10.1029/2008PA001657, 2009.
- Martínez-García, A., Sigman, D. M., Ren, H., Anderson, R. F., Straub, M., Hodell, D. A., Jaccard, S. L., Eglinton, T. I. and Haug, G. H.: Iron Fertilization of the Subantarctic Ocean During the Last Ice Age, *Science*, 343(6177), 1347–1350, doi:10.1126/science.1246848, 2014.
- Matsumoto, K., Chase, Z. and Kohfeld, K.: Different mechanisms of silicic acid leakage and their biogeochemical
830 consequences, *Paleoceanography*, 29(3), 238–254, doi:10.1002/2013PA002588, 2014.
- McCave, I. N. and Hall, I. R.: Size sorting in marine muds: Processes, pitfalls, and prospects for paleoflow-speed proxies, *Geochemistry, Geophys. Geosystems*, 7(10), Q10N05, doi:10.1029/2006GC001284, 2006.
- McGee, D., Winckler, G., Borunda, A., Serno, S., Anderson, R. F., Recasens, C., Bory, A., Gaiero, D., Jaccard, S. L., Kaplan, M., McManus, J. F., Revel, M. and Sun, Y.: Tracking eolian dust with helium and thorium: Impacts of grain size and
835 provenance, *Geochim. Cosmochim. Acta*, 175, 47–67, doi:10.1016/j.gca.2015.11.023, 2016.
- Mengelt, C., Abbott, M. R., Barth, J. A., Letelier, R. M., Measures, C. I. and Vink, S.: Phytoplankton pigment distribution in relation to silicic acid, iron and the physical structure across the Antarctic Polar Front, 170°W , during austral summer, *Deep. Res. Part II Top. Stud. Oceanogr.*, 48(19–20), 4081–4100, doi:10.1016/S0967-0645(01)00081-9, 2001.
- Meredith, M. P., Woodworth, P. L., Chereskin, T. K., Marshall, D. P., Allison, L. C., Bigg, G. R., Donohue, K., Heywood, K.
840 J., Hughes, C. W., Hibbert, A., Hogg, A. M., Johnson, H. L., Jullion, L., King, B. A., Leach, H., Lenn, Y. D., Maqueda, M. A. M., Munday, D. R., Garabato, A. C. N., Provost, C., Sallée, J. B. and Sprintall, J.: SUSTAINED MONITORING OF THE

- SOUTHERN OCEAN AT DRAKE PASSAGE : PAST ACHIEVEMENTS AND FUTURE PRIORITIES, *Rev. Geophys.*, 49, 8755–1209, doi:10.1029/2010RG000348.1., 2011.
- Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., Saito, M. A., Thingstad, T. F., Tsuda, A. and Ulloa, O.: Processes and patterns of oceanic nutrient limitation, *Nat. Geosci.*, 6(9), 701–710, doi:10.1038/ngeo1765, 2013.
- Mortlock, R. A. and Froelich, P. N.: A simple method for the rapid determination of biogenic opal in pelagic marine sediments, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 36(9), 1415–1426, doi:10.1016/0198-0149(89)90092-7, 1989.
- 850 Müller, P. J. and Schneider, R.: An automated leaching method for the determination of opal in sediments and particulate matter, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 40(3), 425–444, doi:10.1016/0967-0637(93)90140-X, 1993.
- Naveira Garabato, A. C., Ferrari, R. and Polzin, K. L.: Eddy stirring in the Southern Ocean, *J. Geophys. Res.*, 116(C9), C09019, doi:10.1029/2010JC006818, 2011.
- Noble, T. L., Piotrowski, A. M., Robinson, L. F., McManus, J. F., Hillenbrand, C. D. and Bory, A. J. M.: Greater supply of Patagonian-sourced detritus and transport by the ACC to the Atlantic sector of the Southern Ocean during the last glacial period, *Earth Planet. Sci. Lett.*, 317–318, 374–385, doi:10.1016/j.epsl.2011.10.007, 2012.
- 855 Nürnberg, C. C., Bohrmann, G., Schlüter, M. and Frank, M.: Barium accumulation in the Atlantic sector of the Southern Ocean: Results From 190,000-year records, *Paleoceanography*, 12(4), 594–603, doi:10.1029/97PA01130, 1997.
- Orsi, H., Whitworth, T. and Jr, W. D. N.: On the meridional extent and fronts of the Antarctic Circumpolar Current, *Deep Sea Res.*, 42(5), 641–673, doi:10.1016/0967-0637(95)00021-W, 1995.
- 860 PAGES, Past Interglacials Working Group of PAGES: Interglacials of the last 800,000years, *Rev. Geophys.*, 54, 162–219, doi:10.1002/2015RG000482.Received, 2016.
- Paillard, D., Labeyrie, L. and Yiou, P.: Macintosh Program performs time-series analysis, *Eos, Trans. Am. Geophys. Union*, 77(39), 379–379, doi:10.1029/96EO00259, 1996.
- 865 Papparazzo, F. E.: Tendencias espaciales y temporales en la distribución de macronutrientes en aguas superficiales del Pasaje Drake, *Ecol. Austral*, 26(1), 027–039, doi:10.25260/EA.16.26.1.0.142, 2016.
- Paytan, A.: Ocean Paleoproductivity, in *Encyclopedia of Paleoclimatology and Ancient Environments. Encyclopedia of Earth Sciences Series.*, edited by V. Gornitz, pp. 643–651, Springer Netherlands, Dordrecht, 2009.
- Peine, F., Turnewitsch, R., Mohn, C., Reichelt, T., Springer, B. and Kaufmann, M.: The importance of tides for sediment dynamics in the deep sea—Evidence from the particulate-matter tracer ^{234}Th in deep-sea environments with different tidal forcing, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 56(7), 1182–1202, doi:10.1016/j.dsr.2009.03.009, 2009.
- 870 Rabassa, J.: Late Cenozoic Glaciations in Patagonia and Tierra del Fuego, *Dev. Quat. Sci.*, 11, 151–204, doi:10.1016/S1571-0866(07)10008-7, 2008.

- Rabassa, J. and Clapperton, C. M.: Quaternary glaciations in the southern hemisphere: An overview, *Quat. Sci. Rev.*, 9(2–3), 299–304, doi:10.1016/0277-3791(90)90024-5, 1990.
- Rabassa, J., Coronato, A. and Martínez, O.: Late Cenozoic glaciations in Patagonia and Tierra del Fuego: An updated review, *Biol. J. Linn. Soc.*, 103(2), 316–335, doi:10.1111/j.1095-8312.2011.01681.x, 2011.
- Renault, A., Provost, C., Sennéchaël, N., Barré, N. and Kartavtseff, A.: Two full-depth velocity sections in the Drake Passage in 2006—Transport estimates, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 58(25–26), 2572–2591, doi:10.1016/j.dsr2.2011.01.004, 2011.
- Rigual-Hernández, A. S., Trull, T. W., Bray, S. G., Cortina, A. and Armand, L. K.: Latitudinal and temporal distributions of diatom populations in the pelagic waters of the Subantarctic and Polar Frontal zones of the Southern Ocean and their role in the biological pump, *Biogeosciences*, 12(18), 5309–5337, doi:10.5194/bg-12-5309-2015, 2015.
- Rigual Hernández, A. S., Trull, T. W., Nodder, S. D., Flores, J. A., Bostock, H., Abrantes, F., Eriksen, R. S., Sierro, F. J., Davies, D. M., Ballegeer, A.-M., Fuertes, M. A. and Northcote, L. C.: Coccolithophore biodiversity controls carbonate export in the Southern Ocean, *Biogeosciences*, 17(1), 245–263, doi:10.5194/bg-17-245-2020, 2020.
- Roberts, J., McCave, I. N., McClymont, E. L., Kender, S., Hillenbrand, C.-D., Matano, R., Hodell, D. A. and Peck, V. L.: Deglacial changes in flow and frontal structure through the Drake Passage, *Earth Planet. Sci. Lett.*, 474, 397–408, doi:10.1016/j.epsl.2017.07.004, 2017.
- Saavedra-Pellitero, M., Baumann, K. H., Lamy, F. and Köhler, P.: Coccolithophore variability across Marine Isotope Stage 11 in the Pacific sector of the Southern Ocean and its potential impact on the carbon cycle, *Paleoceanography*, 32(8), 864–880, doi:10.1002/2017PA003156, 2017.
- Saavedra-Pellitero, M., Baumann, K. H., Fuertes, M. Á., Schulz, H., Marcon, Y., Manon Vollmar, N., Flores, J. A. and Lamy, F.: Calcification and latitudinal distribution of extant coccolithophores across the Drake Passage during late austral summer 2016, *Biogeosciences*, 16(19), 3679–3702, doi:10.5194/bg-16-3679-2019, 2019.
- Schlitzer, Reiner, Ocean Data View, <https://odv.awi.de>, 2021
- Schlüter, M. and Rickert, D.: Effect of pH on the measurement of biogenic silica, *Mar. Chem.*, 63(1–2), 81–92, doi:10.1016/S0304-4203(98)00052-8, 1998.
- Schroth, A. W., Crusius, J., Sholkovitz, E. R. and Bostick, B. C.: Iron solubility driven by speciation in dust sources to the ocean, *Nat. Geosci.*, 2(5), 337–340, doi:10.1038/ngeo501, 2009.
- Shoenfelt, E. M., Sun, J., Winckler, G., Kaplan, M. R., Borunda, A. L., Farrell, K. R., Moreno, P. I., Gaiero, D. M., Recasens, C., Sambrotto, R. N. and Bostick, B. C.: High particulate iron(II) content in glacially sourced dusts enhances productivity of a model diatom, *Sci. Adv.*, 3(6), e1700314, doi:10.1126/sciadv.1700314, 2017.
- Shoenfelt, E. M., Winckler, G., Lamy, F., Anderson, R. F. and Bostick, B. C.: Highly bioavailable dust-borne iron delivered to the Southern Ocean during glacial periods, *Proc. Natl. Acad. Sci. U. S. A.*, 115(44), 11180–11185, doi:10.1073/pnas.1809755115, 2018.

- Shoenfelt, E. M., Winckler, G., Annett, A. L., Hendry, K. R. and Bostick, B. C.: Physical Weathering Intensity Controls Bioavailable Primary Iron(II) Silicate Content in Major Global Dust Sources, *Geophys. Res. Lett.*, 46(19), 10854–10864, doi:10.1029/2019GL084180, 2019.
- 910 Sigman, D. M., Hain, M. P. and Haug, G. H.: The polar ocean and glacial cycles in atmospheric CO₂ concentration, *Nature*, 466(7302), 47–55, doi:10.1038/nature09149, 2010.
- Sprenk, D., Weber, E. M., Kuhn, G., Rosén, P., Frank, M., Molina-Kescher, M., Liebetrau, V. and Röhling, H. G.: Southern Ocean bioproductivity during the last glacial cycle - New detection method and decadal-scale insight from the Scotia Sea, *Geol. Soc. Spec. Publ.*, 381(1), 245–261, doi:10.1144/SP381.17, 2013.
- 915 Strub, P. T., James, C., Montecino, V., Rutllant, J. A. and Blanco, J. L.: Ocean circulation along the southern Chile transition region (38°–46°S): Mean, seasonal and interannual variability, with a focus on 2014–2016, *Prog. Oceanogr.*, 172(May 2018), 159–198, doi:10.1016/j.pocean.2019.01.004, 2019.
- Sugden, D. E., McCulloch, R. D., Bory, A. J.-M. and Hein, A. S.: Influence of Patagonian glaciers on Antarctic dust deposition during the last glacial period, *Nat. Geosci.*, 2(4), 281–285, doi:10.1038/ngeo474, 2009.
- 920 Sulpis, O., Boudreau, B. P., Mucci, A., Jenkins, C., Trossman, D. S., Arbic, B. K. and Key, R. M.: Current CaCO₃ dissolution at the seafloor caused by anthropogenic CO₂, *Proc. Natl. Acad. Sci.*, 115(46), 11700–11705, doi:10.1073/pnas.1804250115, 2018.
- Suman, D. O. and Bacon, M. P.: Variations in Holocene sedimentation in the North American Basin determined from ²³⁰Th measurements, *Deep Sea Res. Part A, Oceanogr. Res. Pap.*, 36(6), 869–878, doi:10.1016/0198-0149(89)90033-2, 1989.
- 925 Tapia, R., Ho, S. L., Núñez-Ricardo, S., Marchant, M., Lamy, F. and Hebbeln, D.: Increased marine productivity in the southern Humboldt Current System during MIS 2–4 and 10–11, *Paleoceanogr. Paleoclimatology*, 33(0), 2–31, doi:10.1029/2020PA004066, 2021.
- Thöle, L. M., Amsler, H. E., Moretti, S., Auderset, A., Gilgannon, J., Lippold, J., Vogel, H., Crosta, X., Mazaud, A., Michel, E., Martínez-García, A. and Jaccard, S. L.: Glacial-interglacial dust and export production records from the Southern Indian Ocean, *Earth Planet. Sci. Lett.*, 525, [115716](#), doi:10.1016/j.epsl.2019.115716, 2019.
- 930 Toyos, M. H., Lamy, F., Lange, C. B., Lembke-Jene, L., Saavedra-Pellitero, M., Esper, O. and Arz, H. W.: Antarctic Circumpolar Current Dynamics at the Pacific Entrance to the Drake Passage Over the Past 1.3 Million Years, *Paleoceanogr. Paleoclimatology*, 35(7), 1–20, doi:10.1029/2019PA003773, 2020.
- Turekian, K. K. and Wedepohl, K. H.: Distribution of the Elements in Some Major Units of the Earth's Crust, *GSA Bull.*, 935 72(2), 175–192, doi:10.1130/0016-7606(1961)72[175:DOTEIS]2.0.CO;2, 1961.
- Turnewitsch, R., Reyss, J.-L., Nycander, J., Waniek, J. J. and Lampitt, R. S.: Internal tides and sediment dynamics in the deep sea—Evidence from radioactive ²³⁴Th/²³⁸U disequilibria, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 55(12), 1727–1747, doi:10.1016/j.dsr.2008.07.008, 2008.

- Vernet, M., Sines, K., Chakos, D., Cefarelli, A. O. and Ekern, L.: Impacts on phytoplankton dynamics by free-drifting icebergs in the NW Weddell Sea, *Deep. Res. Part II Top. Stud. Oceanogr.*, 58(11–12), 1422–1435, doi:10.1016/j.dsr2.2010.11.022, 2011.
- 940
- Wadham, J. L., Hawkings, J. R., Tarasov, L., Gregoire, L. J., Spencer, R. G. M., Gutjahr, M., Ridgwell, A. and Kohfeld, K. E.: Ice sheets matter for the global carbon cycle, *Nat. Commun.*, 10(1), 3567, doi:10.1038/s41467-019-11394-4, 2019.
- Wengler, M., Lamy, F., Struve, T., Borunda, A., Böning, P., Geibert, W., Kuhn, G., Pahnke, K., Roberts, J., Tiedemann, R. and Winckler, G.: A geochemical approach to reconstruct modern dust fluxes and sources to the South Pacific, *Geochim. Cosmochim. Acta*, 264, 205–223, doi:10.1016/j.gca.2019.08.024, 2019.
- 945
- Winckler, G., Anderson, R. F., Fleisher, M. Q., McGee, D. and Mahowald, N.: Covariant glacial-interglacial dust fluxes in the equatorial Pacific and Antarctica., *Science*, 320(5872), 93–96, doi:10.1126/science.1150595, 2008.
- Winckler, G., Anderson, R. F., Jaccard, S. L. and Marcantonio, F.: Ocean dynamics, not dust, have controlled equatorial Pacific productivity over the past 500,000 years, *Proc. Natl. Acad. Sci. U. S. A.*, 113(22), 6119–6124, doi:10.1073/pnas.1600616113, 2016.
- 950
- Wu, S.Y. and Hou, S.: Impact of icebergs on net primary productivity in the Southern Ocean, *Cryosph.*, 11(2), 707–722, doi:10.5194/tc-11-707-2017, 2017.
- Wu, S., Kuhn, G., Diekmann, B., Lembke-Jene, L., Tiedemann, R., Zheng, X., Ehrhardt, S., Arz, H. W. and Lamy, F.: Surface sediment characteristics related to provenance and ocean circulation in the Drake Passage sector of the Southern Ocean, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 154(December), 103135, doi:10.1016/j.dsr.2019.103135, 2019.
- 955
- Wu, S., Lembke-Jene, L., Lamy, F., Arz, H. W., Nowaczyk, N., Xiao, W., Zhang, X., Hass, H. C., Titschack, J., Zheng, X., Liu, J., Dumm, L., Diekmann, B., Nürnberg, D., Tiedemann, R. and Kuhn, G.: Orbital- and millennial-scale Antarctic Circumpolar Current variability in Drake Passage over the past 140,000 years, *Nat. Commun.*, 12(1), 3948, doi:10.1038/s41467-021-24264-9, 2021.
- 960