

Investigating hydroclimatic impacts of the 168-158 BCE volcanic quartet and their relevance to the Nile River basin and Egyptian history

5 Ram Singh^{1,2}, Kostas Tsigaridis^{1,2}, Allegra N. LeGrande^{2,1}, Francis Ludlow³, Joseph G Manning⁴

¹ Center for Climate Systems Research, Columbia University, New York

² NASA Goddard Institute for Space Studies, New York, NY-10025

³ Department of History, School of Histories and Humanities, Trinity College, Dublin 2, Ireland

10 ⁴ Departments of History and Classics, Yale University, New Haven, CT 06520, USA

Correspondence to: Ram Singh (rs4068@columbia.edu)

Abstract.

The Ptolemaic era (305-30BCE) is an important period of Ancient Egyptian history known for its material and scientific advances, but also ongoing episodes of political and social unrest in the form of (sometimes
15 widespread) revolts against the Ptolemaic elites. While the role of environmental pressures has long been overlooked in this period of Egyptian history, ice-core-based volcanic histories have identified the period as experiencing multiple notable eruptions, and a repeated temporal association between explosive volcanism and revolt has recently been noted. Here we analyze the global and regional (Nile River Basin) climate response to a unique historical case of 4 consecutive and closely timed large, strato-volcanic
20 eruptions (first a tropical one, closely followed by 3 extratropical northern hemispheric events) between 168 and 158 BCE, a particularly troubled period in Ptolemaic history for which we now provide a more detailed hydroclimatic context. The NASA GISS ModelE2.1 Earth system model simulates a strong radiative response with a radiative forcing (top of atmosphere) of -7.5 W/m² (following the first eruption) and -2.5 w/m² (after each of the 3 remaining eruptions) at a global scale. Associated with this, we observe
25 a global surface cooling of the order of 1.5°C following the first (tropical) eruption, with the following three extratropical eruptions extending the cooling period for more than 15 years. Consequently, this series of eruptions constrained the northward migration of the inter-tropical convergence zone (ITCZ) during the northern hemisphere summer monsoon season, and major monsoon zones (African, South

Asian, and East Asian) experienced a suppression of rainfall of >1 mm/day during the monsoon (JJAS) season averaged for 2 years after each eruption. A substantial suppression of the Indian and north African summer monsoon (over the Nile River headwater region) strongly affected the modelled river flow in the catchment and river discharge at river mouth. River mass flow over the basin was observed to consecutively decrease by up to more than 30% relative to an unperturbed (non-volcanic) annual mean flow for 2 years after the tropical eruption. A moderate decrease of up to 10-20% was observed after each of the remaining eruptions. These results indicate, in sum, that the first eruption likely produced a strong hydroclimate response, with the following 3 eruptions prolonging this condition. These results also support the recent hypothesized association between ice-core-based signals of explosive volcanism and hydroclimatic variability during the Ptolemaic era, including the suppression of the agriculturally critical Nile summer flooding.

Key Words: Volcanic eruption, hydroclimate impacts, Inter-tropical Convergence Zone (ITCZ), Monsoon, Nile River basin

1. Introduction

Explosive volcanic eruptions that result in high altitude sulfate aerosol distribution across one or both hemispheres can diminish insolation, with global and regional climatic impacts (e.g., Robock, 2000; Toohey et al., 2019). The sulfate aerosols resulting from (for example) the 1991 Mt. Pinatubo eruption of ~18 Tg SO₂ increased the atmospheric optical depth from ~0.6 to ~0.75 and was associated with surface cooling of 0.5 °C (Robock and Mao 1995; Parker et al., 1996). Volcanically induced cooling can also reduce net evaporation and hence precipitation over large areas (Lui et al., 2016; Iles et al., 2013), while also potentially producing a near global dynamical suppression of the northward migration of the inter-tropical convergence zone (ITCZ) during the boreal summer, as the convergence follows the surface area of maximum temperature (Pettersen et al., 2000; Chiang and Bitz, 2005; Broccoli et al., 2006; Colose et al., 2016). These changes can impact river outflow (Oman et al., 2006; Sabzevari et al., 2015; Kostić et al., 2016) with implications for civilizations from antiquity to the present-day. The Nile River, upon which

- Deleted: Large explosive
- Deleted: , leading to both
- Deleted: on climate and society
- Deleted: of the atmosphere
- Deleted: The
- Deleted: caused by such events
- Deleted: leading to
- Deleted: -scale
- Deleted: in precipitation
- Deleted: , moreover,
- Deleted:), which has
- Deleted: , not least by affecting food security where agriculture is dependent upon this outflow.

Egyptian agriculture was heavily dependent, is a key example. With ice-core-based volcanic histories
 70 now identifying several hundred potentially climatically effective eruptions over the past 2.5 millennia
 (Sigl et al., 2015), Egyptian civilization provides a test-case for the study of human vulnerability to abrupt
 environmental changes in having experienced repeated volcanically induced “hydroclimatic shocks”
 (e.g., Mikhail, 2015; Manning et al., 2017, 2021).

Explosive volcanic eruptions represent the major natural source of forced variability in the climate system
 75 at yearly to decadal time scales (Schmidt et al., 2011; Colose et al., 2016; Swingeduow et al., 2017; Khodri
 et al. 2017). Powerful explosive eruptions can inject sulfur-rich gases into the stratosphere, where they
 oxidize to form sulfate aerosols that can persist for months to years, impacting climate on regional to
 global scales. Volcanic stratospheric aerosols can cause troposphere cooling by scattering incoming
 shortwave radiation, while also heating the stratosphere (Robock and Mao, 1992). Unequal north-south
 80 stratospheric heating due to volcanic aerosols concentrated in lower latitudes after tropical eruptions can
 influence major modes of circulation and surface climate variability such as the Arctic Oscillation/North
 Annular Mode (AO/NAM) and North Atlantic Oscillation (NAO), driving an enhanced westerly airflow
 (Shindell et al., 2004; Zanchettin et al., 2021). The post-volcanic surface temperature response can affect
 the El Niño/La Niña phase and Southern Oscillation, and have a long-term impact on Atlantic Meridional
 85 Overturning Circulation (AMOC) strength (Khodri et al., 2017; Wahl et al., 2014; Robock and Mao,
 1995; Pausata et al., 2015). Extratropical eruptions are usually thought to have a weaker impact than
 tropical eruptions. This arises in part because of the background Brewer-Dobson circulation upwelling in
 the tropics and downwelling at higher latitudes, which affects the stratospheric lifetime of volcanic
 aerosols (Kirtman et al., 2013; Myhre et al., 2013; Schneider et al., 2009). Recent studies have, however,
 90 illustrated the potential for a dynamically induced and disproportionally strong climate forcing from such
 eruptions (Toohey et al. 2019).

Volcanic injection of sulfur-containing compounds can, too, influence stratospheric chemistry, yielding
 further complex atmospheric and climatic responses upon interacting with water and halogens (LeGrande
 et al., 2016; Brenna et al., 2020; Staunton-Sykes et al., 2021). Paleoclimate records and climate modeling
 95 suggest that the dynamical response to volcanic aerosol causes a net (but regionally variable) drying and
 impacts global rainfall patterns (PagesHydro2k/Smerdon et al., 2017; Colose et al., 2016; Liu et al., 2016;

- Deleted: (and hence civilization)
- Deleted: forcing
- Deleted: several
- Deleted: valuable
- Deleted: and resilience
- Deleted: change
- Deleted: potentially experiencing
- Deleted: induced by these events
- Deleted: have become increasingly recognized as
- Deleted: stratovolcanic
- Deleted: high
- Deleted: aerosol particles
- Deleted: in turn
- Deleted: the
- Deleted: in the stratosphere
- Deleted: cooling in the
- Deleted: aerosol presence
- Deleted: atmospheric
- Deleted: in effect by
- Deleted: also
- Deleted: , as well as having
- Deleted: comparatively
- Deleted: climate
- Deleted: directly
- Deleted: extratropical eruptions to
- Deleted: include a
- Deleted: efforts
- Deleted: of
- Deleted: effect
- Deleted: significantly
- Deleted: the
- Deleted: pattern

Iles and Hegerl, 2014). Trenberth and Dai (2007) thus analyzed the impact of the Pinatubo (1991) eruption on terrestrial precipitation and river streamflow and found an increase in associated drought conditions in 1992. Joseph and Zeng (2011) suggested that volcanically induced rainfall anomalies over land and ocean can seasonally modulate tropical drought. Hemispheric biases in volcanic aerosol distribution can, moreover, impact the movement of ITCZ by constraining its summertime migration into the energetically deficit hemisphere (Colose et al., 2016; Xian and Miller, 2008). Effectively, the ITCZ shifts “away” from the hemisphere with the greatest aerosol burden. For tropical eruptions, even those producing roughly even hemispheric burdens, this movement is typically more southward owing to the larger amount of land in the Northern Hemisphere and relative the greater ocean area (and higher thermal capacity) of the Southern Hemisphere.

For Africa, eruptions producing asymmetrical latitudinal aerosol burdens (e.g., Katmai in 1912, El Chichón in 1982) may have enhanced 20th century Sahelian droughts by influencing the strength and position of Hadley cells (Haywood et al., 2013). Of the Nile, monsoon rainfall over the Ethiopian highlands contributes (via the Blue Nile and Atbara River) ~85% to the summer flood in Egypt and is a strong control on its interannual variability (Melesse et al., 2011). Ancient Egypt was famed for the productivity of its flood recession agriculture, but the Nile summer flood was also famously variable, with insufficient flooding impacting Egyptian society (e.g., Bell, 1975; Butzer, 1976, 1984; Said, 1993; Hassan, 1997a, b; Hassan, 2007; McCormick, 2013). Some of this variability was driven by explosive volcanism. The Laki fissure eruption (1783/84) injected approximately 122 Mt of SO₂ into the atmosphere over eight months, producing a strong cooling and suppressing the African monsoon (Oman et al., 2006; D’Arrigo et al., 2011). The resulting diminished Nile flooding (Oman et al., 2005, 2006; Mikhail, 2015) is known colloquially as “Nile failure”. Similar impacts were simulated and can be observed for the Katmai (1912) eruption (Vorosmarty et al., 1998; Thordarson and Self, 2003; Oman et al., 2005, 2006; Manning, 2018).

The most richly documented period of ancient Egyptian history is the Ptolemaic era, 305-30 BCE, with Egypt ruled by Greeks in a lineage beginning with Ptolemy I Soter (d. 283 BCE), previously one of Alexander the Great’s key generals and instrumental in Egypt’s conquest. The period distinguishes itself for mixing Greek and Egyptian traditions, its material, cultural and scientific achievement (e.g., the

Deleted: For example, ...renberth and Dai (2007) thus analyzed the impact of the Pinatubo (1991) eruption on land...errestrial precipitation and river streamflow and found an increase in associated drought conditions after the eruption ...n 1992. Joseph and Zeng (2011) suggested that varying responses to ...olcanically induced rainfall anomalies over land and ocean can seasonally modulate drought conditions in the tropics. In addition, hemispherical asymmetrical radiative forcing due to ...ropical drought. Hemispheric biases in the ...olcanic aerosol distribution of volcanic aerosols creates a radiative imbalance across the hemisphere, impacts ...an, moreover, impact the movement of ITCZ,...by constraining the extent of (... [1]

Deleted: amount of volcanic ...erosol burden. For tropical eruptions, even those producing a ...oughly even hemispheric aerosol burden... (... [2]

Deleted: volcanic ...ruptions resulting in an ...roducing asymmetrical latitudinal aerosol burden...urdens (e.g., the ...atmai eruption ...n 1912 and (... [3]

Deleted: are thus thought to ...ay have enhanced 20th century Sahelian drought conditions by shifting the surface temperature maxima and...roughs by influencing the strength and position of Hadley cells (Haywood et al., 2013). Of the Nile, monsoon rainfall over the Ethiopian highlands contributes (mainly ...ia the Blue Nile and Atbara River) ~85% to the summer flood over the Egyptian plains ...n Egypt and is a strong control over the associated ...n its interannual variability of the flood (... [4]

Deleted: Years of diminished African monsoon rainfall were historically associated with insufficient river water to extensively practice the ...ncient Egypt was famed for the productivity of its flood recession agriculture that ordinarily delivered such high agriculturally productivity in the Nile valley, and for which ancient Egypt was famed. But ... but the Nile summer flood was also famously mercurial...ariable, with insufficient flooding often to adverse societal impacts (... [5]

Deleted: likely

Deleted: series ...1783/84) thus ...njected approximately 122 Mt of SO₂ into the atmosphere over eight months and produced... producing a strong cooling and suppression of (... [6]

Deleted:),... The resulting in reduced...iminished Nile River flooding (Oman et al., 2005, 2006; Mikhail, 2015), or what... is known colloquially as “Nile failure”. Similar impacts have been...re simulated over the African region (... [7]

Deleted: One of the...he most richly documented periods...eriod of ancient Egyptian history is the Ptolemaic era, 305-30 BCE, during which time...ith Egypt was (... [8]

Deleted: who had been...reviously one of Alexander the Great’s key generals and instrumental in the...gypt’s conquest of Egypt... The period distinguishes itself through its...or mixing of ...reek and Egyptian traditions and... its great ...aterial, cultural and scientific achievement (not least in (... [9]

265 founding of Alexandria with its Great Library), but also its chronic political instability (McGing, 1997; Ludlow and Manning, 2016; 2021). External environmental influences have been little considered in this, despite the dependence of Egyptian agriculture on the summer flood. However, recent work has revealed a repeated close coincidence in the timing of many (if certainly not all) internal revolts and ice-core-based dates of inferred-tropical and NH extratropical eruptions that appears statistically significance (i.e., non-random) (Ludlow and Manning, 2016, 2021; Manning et al., 2017). One example is the “Great Theban Revolt” starting in c.207 BCE, shortly after a 209 BCE tropical eruption (Sigl et al., 2015), with extensive territories lost to two native Pharaohs (Ludlow and Manning, 2016; Ludlow et al., 2023). That the revolt and eruption dates under study derive from independent chronologies (documentary and ice-core) helps exclude potential biases in estimating this statistical significance. For example, inflated correlations may result between events known from the same sources (e.g., between extreme weather and societal stresses such as famine, if those instances of extreme weather that contributed to such stresses were more likely to have been documented than those that didn’t (White and Pei, 2020)). It is a truism that correlation does not establish causation. Genuine causality is, however, implied where significance testing suggests an observed correlation is unlikely to have arisen randomly, though this does not determine the direction or character of causality (Izdebski et al., 2023). Statistical significance may, moreover, be sensitive to many factors. These include here (1) the choice of statistical test, (2) the choice of revolt and eruption dates (if uncertainties exist), (3) judgements as to what constitutes “revolt” (vs. phenomena like food riots motivated more by desperation than politics), and (4) judgements concerning which eruptions to include in testing (e.g., seeking those with a meaningful impact vs. non-climatically effective eruptions introducing “noise” into the analysis), assessed by estimated volumes and heights of atmospheric SO₂ injections, eruption locations, and more. Notably, thus, testing by Ludlow and Manning (2016) was followed by Manning et al. (2017) who also observed a statistically significant coincidence between eruptions and Ptolemaic-era revolts using different methods and variant dates. Logic dictates that the direction of any genuine causality must flow here from eruption to revolt (Izdebski et al., 2023). Further confidence in its reality arises from the existence of plausible mechanisms connecting volcanic hydroclimatic variability with revolt (i.e., via reduction of the Nile summer flood and consequent societal impacts). Much work remains to further characterize this causality, how direct or

- Deleted: the city of
- Deleted: on the Mediterranean coast
- Deleted: famed
- Deleted: and Lighthouse
- Deleted: through
- Deleted: Little consideration has been given to external
- Deleted: history
- Deleted: great
- Deleted: Nile
- Deleted: correlation between
- Deleted: and frequency
- Deleted: the
- Deleted: explosive
- Deleted: Just one
- Deleted: of
- Deleted: occurring
- Deleted: notable
- Deleted: , when the Ptolemies
- Deleted: control of large areas of the Nile Valley to a sequence of
- Deleted: apparently
- Deleted: Egyptian
- Deleted: Sigl et al., 2015;
- Deleted: 2022).
- Deleted: The temporal correspondence between internal revolts and explosive volcanism certainly appears recurrent and non-random in the Ptolemaic Egypt (Ludlow and Manning, 2016, 2021; Manning et al., 2017; Izdebski et al., 2022).
- Deleted: revolts
- Deleted: volcanic eruptions
- Deleted: are known
- Deleted: different archives with
- Deleted: historical
- Deleted: has also helped to
- Deleted: positive
- Deleted: when
- Deleted: are
- Deleted: or disease
- Deleted: While the results of Ludlow and Manning (2016, 2021) and Manning et al. (2017) thus implies a causal linkage between explosive eruptions and Ptolemaic-era revolts, much
- Deleted: determine its underlying character, including

indirect it may have been, and whether this changed meaningfully through time (and between revolts that
 335 varied in geography and scale) according to (or in interaction with) other coincident potential causes (from
longer term developments promoting chronic vulnerabilities, to more acute political and socioeconomic
stresses). White and Pei (2020) argue that such questions represent a key challenge for climate historians
and related scholars, recommending a framework wherein potential causes are assessed using a
framework of necessary and sufficient conditions (put simply; see also Ludlow et al. (2023)). Gao et al.
 340 (2021) employ a framework wherein the role of volcanically induced hydroclimatic “shocks” in the
collapse of Chinese dynasties is characterized along a spectrum from “ultimate” to “proximate” causation
(see also Villmoare (2022) for this framework). Here, smaller hydroclimatic shocks could act as the
ultimate cause of collapse when enabled by high pre-existing stress, while larger shocks could act with
greater independence as proximate causes without substantial pre-existing stress.
 345 An alternative framing in many climate-conflict studies (not incompatible with that proposed by White
and Pei (2020) or employed by Gao et al. (2021)) is to delineate multiple identifiable “pathways” that
may enable or lead (through material (economic), political, cultural or psychological channels) to links
between hydroclimatic variability and various forms of conflict (see Hsiang and Burke (2014) and Ide
(2017) for reviews). Success here requires that statistical findings are interpreted with due reference to
 350 the relevant historical, political, geographical contexts. For Ptolemaic Egypt, one hypothesized pathway
involves the societal impacts and responses to sudden hydroclimatic variability that can follow explosive
volcanism, and influence (alongside other regional factors) the intensity of the African monsoon. When
this causes a “failure” of the Nile summer flood, many adverse societal impacts may occur. These include
harvest failure (seen also in other periods of Egyptian history (e.g., Hassan, 1997a,b; Mikhail, 2015)),
 355 possibly prompting subsistence-driven migration to urban areas, with inability to meet state taxation
demands, (payable in grain) also potentially necessitating the sale of hereditary familial lands (Manning,
2003; Manning et al., 2017). These stressors might work in tandem with the psychological, and religious
significance of a “failed” flood, something widely feared, and which could be interpreted (even
propagandized to foment revolt) as signaling divine displeasure at the Pharaoh (Ludlow and Manning,
 360 2021; Ludlow et al., 2023). In the Ptolemaic context, when some native Egyptian elites were likely

- Deleted: (which
- Deleted: date,
- Deleted:), and (relatedly) what pathways were in effect
- Deleted: “operationalize” any such linkage. Answering
- Deleted: is now deemed
- Deleted: (White and Pei, 2020). Taken alone, such a correlation does not establish (nor necessarily even imply) causation. Causality is, however, at least implied in cases where analyses are conducted alongside statistical significance testing, with the resulting correlations considered unlikely to have arisen purely by chance, and when such results
- Deleted: context, allowing causal “pathways” to be credibly hypothesized (Izdebski et al., 2022).⁴
- Deleted: the fundamental
- Deleted: linkage
- Deleted: pronounced changes in hydrological cycle wrought by the widespread northern hemispheric cooling that can follow major explosive volcanic eruptions, and which can act (as noted at (... [10])
- Deleted: , alongside other regional factors.
- Deleted: leads to
- Deleted: agriculturally critical
- Deleted: a range of
- Deleted: most obviously
- Deleted: agricultural and economic impacts (as can be
- Deleted: many
- Deleted: with reduced food security for families who may (... [11])
- Deleted: ,
- Deleted: their
- Deleted: and prompting migration from rural areas to large (... [12])
- Deleted: This would likely compound
- Deleted: ,
- Deleted: and, ultimately, political
- Deleted: Nile
- Deleted: with such an event being
- Deleted: among the general populace and with the potential to
- Deleted: and
- Deleted: a reflection of
- Deleted: 2022
- Deleted: context of the
- Deleted: period
- Deleted: parts of the populace, including at least
- Deleted: of the older
- Deleted: and priesthood,

resentful of Greek rule, [with](#) taxation and other advantages given to those of Greek backgrounds (McGing, 1997; Ludlow et al., [2023](#)), a Nile failure may have held [particular](#) political potency. Huhtamaa et al. (2022) have called for [case studies of the hydroclimatic and socioeconomic impacts of specific eruptions to advance understandings of human-environmental causalities](#). For Ptolemaic Egypt, the 160s BCE are [particularly relevant](#) in experiencing considerable internal revolt and instability. Indeed, the Ptolemaic dynasty might have fallen here [without](#) self-interested Roman intervention against the Seleukid empire ([rivals to the Ptolemies](#)) after their invasion (170-168 BCE) of Egypt under [King](#) Antiochus IV (Grainger, 2010; Blouin, 2014; Manning et al., 2017). This [period](#) is also remarkable for three notable volcanic eruptions (168, 164, 161 BCE), with [another](#) in 158 BCE (Sigl et al., 2015), [which we term the “eruption quartet”](#). Substantial sulfate across both poles identifies the first (168 BCE) as the largest and likely [tropical](#), followed by three moderate eruptions in the northern hemisphere, [extratropics](#) (Sigl et al., 2015). While high-resolution [paleoenvironmental](#) proxies for Egypt are effectively absent [for](#) this period, the hydroclimatic impacts of [these](#) eruptions can be [explored](#) by climate modeling. Few studies have examined the climatic societal effects of eruption clusters. These include the cluster between 1108 and 1110 CE (Guillet et al., 2020), the “double event” of the 6th century in 536 and 540 CE (Toohey et al., 2016), and the cluster from 1637 to 1641 (Stoffel et al., 2022; Huhtamaa et al., 2022). These studies variously employed [paleoclimatic](#) data, written evidence and/or climate [modelling](#) to reveal strong negative [post-eruption](#) temperature anomalies [for](#) the Northern hemisphere, thereby suggesting the potential for [diminished](#) crop yields and providing a climatic context to better understand the human history of these periods. [Here, we intend](#) to advance our understanding of the likely hydroclimatic impact of [the 168-158 BCE eruption quartet as a foundation for ongoing efforts to more securely establish and qualify](#) the causality underlying the observed association between [eruptions, Ptolemaic-era revolts, and other political and socioeconomic phenomena and developments](#).

We thus use a computationally expensive but more sophisticated version of the National Aeronautics and Space Administration (NASA), Goddard Institute for Space Studies (GISS) Earth system model, GISS ModelE2.1-MATRIX (Bauer et al., 2008; Bauer et al., 2020), to simulate the [168-158 BCE eruption quartet](#) and regional hydroclimate [responses](#) over the Nile River basin. [Section 2 presents](#) model details and, experiment [methodology](#). [Estimation of](#) background climate [for](#) the 2.5k [period](#) (orbital and

Deleted: and the

Deleted: 2022...023), a Nile failure may have held additional...articular political potency, offering further pathways by which volcanically induced hydroclimatic variability might motivate revolt onset. (... [13])

Deleted: detailed ...ase studies of both ...he hydroclimatic and socioeconomic impacts of specific volcanic ...ruptions to advance our ...nderstandings of human-environmental causalities. For Ptolemaic Egypt, the 160s BCE are thus of particular relevance, ...articularly relevant in experiencing considerable internal revolt and instability. Indeed, the Ptolemaic dynasty might well have fallen here if not for ...ithout self-interested Roman intervention against the Seleukid empire (great ...ivals to the Ptolemies),... after their successful ...nvasion (170-168 BCE) of Egypt under the command of...ing Antiochus IV (Grainger, 2010; Blouin, 2014; Manning et al., 2017; see section S1.1 for additional historical context).... This period is also a decade...remarkable for three notable volcanic eruptions (168, 164 and... 161 BCE), with a further event...nother in 158 BCE (Sigl et al., 2015). The of... which we term the “eruption quartet”. Substantial sulfate across both poles (Sigl et al., 2015) ...dentifies the first eruption (168 BCE) as the largest and likely occurring in the tropics...ropical, tropics...ropical, followed by three equally separated and comparably ...oderate-sized extratropical...ruptions in the northern hemisphere. ... extratropics (Sigl et al., 2015). While high-resolution paleoenvironmental...aleoenvironmental proxies for Egypt are effectively absent in...or this early ...eriod, our understanding of...the hydroclimatic impacts of the sequence of...hese eruptions between 168 and 158 BCE ...an be advanced...explored by climate modeling. No previous study has specifically explored such a set of four closely consecutive eruptions or their impacts on the regional hydroclimate of a major ancient-era civilization. The few previous studies that have thus far (... [14])

Deleted: and ...societal effects of eruption clusters. These include an exploration of ...he volcanic event...cluster of the early 12th century (...etween 1108 and 1110 CE)...(Guillet et al., 2020), the “double event” of the 6th century in 536 and 540 CE (Toohey et al., 2016), perhaps better seen as a triple event, in view of the additional, additional, if much smaller, eruption in 546 (Sigl et al., 2015), and the eruption...016), and the cluster from 1637 to 1641 (Stoffel et al., 2022; Huhtamaa et al., 2022). These studies have ...ariously employed paleoclimatic...aleoclimatic data, written evidence climate model simulation ...odelling to reveal the ...trong negative post-eruption temperature anomalies over...or the Northern hemisphere following these eruptions, ... thereby suggesting the potential for adverse effects on...inished crop yields and providing a climatic context by which ...o better understand the human history of these periods (Guillet et al. 2020; Toohey et al. 2016; Stoffel et al., 2022). In this study, our main intent is... Here, we intend to advance our understanding of the likely hydroclimatic impact of his ...he 168-158 BCE eruption quartet as a foundation for further work aimed at establishing the nature of...ngoing efforts to more securely (... [15])

Deleted: ..., 2020), to simulate the series of eruptions from ...68 to ...158 BCE eruption quartet and analyze the impacts on ...egional hydroclimate responses over the Nile River basin. Details of the...ection 2 presents model details and methodology employed to conduct the...experiment and analysis are discussed in section 2. estimation of the...ethodology. Estimation of background cl (... [16])

greenhouse gases (GHGs) changes), alongside impacts due to PMIP4 (Paleoclimate Model Intercomparison Project, phase 4) vegetation cover estimates for the period are considered in section 3. Particular subsections evaluate the GISS ModelE for its capability to resolve the microphysical properties of volcanic aerosols during this period and to analyze the radiative impacts of the aerosols from these eruptions, which control their radiative and hydroclimatic impacts (Timmreck et al., 2009; Timmreck et al., 2010; Schmidt et al., 2010). The discussion and conclusion (section 4) summarize our results and considers how they can advance understandings of the period's fraught history in Egypt.

2. Methodology & Experiment design

2.1 Model Description

We used the NINT (Non-INTERactive) version (Kelley et al., 2020) of GISS ModelE2.1 to simulate background climate conditions corresponding to 2500 years before present (2.5ka, kilo-years BP), similar to protocols for the mid-Holocene (6ka) coordinated experiment (Kageyama et al, 2017), but adjusting trace gases and orbital forcing for 2.5ka. The term “non-interactive” means that atmospheric composition and climate are decoupled, so any changes in composition are handled by external input only. Once our model attained an equilibrium climate state, we enabled atmospheric composition-climate interactions for our experiments, described below.

GISS ModelE2.1 is a state-of-the-art Earth System Model contributing to the Climate Model Intercomparison Project (CMIP) phase 6 (Eyring et al., 2016). The model's atmospheric component simulates on a horizontal resolution of 2° latitude by 2.5° longitude with 40 vertical layers and a model top at 0.1 hPa. It is coupled to the GISS Ocean v1 model at a horizontal resolution of 1° latitude by 1.25° longitude with 40 layers. The Demographic Global Vegetation Model (DGVM) is the Ent Terrestrial Biosphere Model (TBM) (Kiang, 2012; Kim et al., 2015) and was used to implement climate-influencing vegetation properties, including satellite-driven (MODIS) plant functional types (PFTs) and the monthly varying leaf area index (LAI) (Gao et al., 2008; Myneni et al., 2002). Tree heights come from Simard et al. (2011) and include an interactive carbon cycle (Ito et al., 2020). The MATRIX (Multiconfiguration Aerosol TRacker of mIXing state) aerosol microphysics module (Bauer et al., 2008; 2020) was used in

- Deleted: only), together with the
- Deleted: vegetations
- Deleted: under
- Deleted: Further
- Deleted: here
- Deleted: NASA
- Deleted: modelling
- Deleted: in resolving
- Deleted: arising
- Deleted: this eruption
- Deleted: the
- Deleted: the
- Deleted: of volcanic eruptions
- Deleted: .
- Deleted: Finally, the
- Deleted: summarizes
- Deleted: our understanding
- Deleted: human history in Egypt. This case of 4 closely consecutive eruptions presents a unique case to study the role of multiple eruptions on regional climate over such a critical region (Nile Basin).
- Deleted: the
- Deleted: the period
- Deleted: the
- Deleted: developed
- Deleted: except with
- Deleted: adjusted
- Deleted: the
- Deleted: performed here, as
- Deleted: of GISS ModelE2.1
- Deleted: is
- Deleted: the
- Deleted: controlling
- Deleted: the
- Deleted: Bauer et al.,
- Deleted: is

our coupled composition-climate runs, to simulate the active volcanism and corresponding climate conditions. MATRIX is an aerosol microphysics scheme using the quadrature method of moments, representing new particle formation (Vehkamäki et al., 2002), aerosol-phase chemistry, condensational growth, coagulation and the mixing state of aerosols (Bauer et al., 2013). MATRIX tracks 16 mixing states with 51 aerosol tracers and resolves mixtures of sulfate, nitrate, ammonium, aerosol water, black carbon, organic carbon, sea salt and mineral dust (Bauer et al., 2008). MATRIX includes the direct effect and the first indirect effect of aerosols on climate.

Approximate eruption locations are crucial to estimate climatic impacts (Toohey et al., 2016, Aquila et al., 2018 [https://acd-ext.gsfc.nasa.gov/Documents/NASA_reports/Docs/VolcanoWorkshopReport_v12.pdf]). Locations for the 168-158 BCE eruptions were first chosen following the bi-polar multi-ice-core sulphate deposition data of Sigl et al. (2015), which allow a discrimination between likely tropical (low-latitude) eruptions and those likely in the extratropics of either hemisphere. Without additional data (e.g., ice-core tephra) providing more precise locations, the ultimate model location must be selected more arbitrarily. Our chosen locations are shown in Fig S3, and we note that eruption longitude (versus latitude) is not expected to play a major role as an uncertainty factor, in the modelled climatic response. The forcing potential of each eruption in terms of atmospheric SO₂ injection was also estimated using the Sigl et al. (2015) multi-ice-core record of sulfate deposition over Greenland and Antarctica, linearly scaled corresponding to Pinatubo (1991) eruption estimates of 18.5 Tg SO₂ (Wolfe and Hoblitt, 1996). Injection height was selected to match Pinatubo, absent further information.

Deleted: the
Deleted: described here

Deleted: An approximate
Deleted: location is
Deleted: accurately
Deleted: volcanic
Deleted: The broad hemispheric position of
Deleted: to
Deleted: volcanic quartet is thus
Deleted: (to begin) with reference to
Deleted: allows
Deleted: occurring
Deleted: any firm
Deleted: or direct geologic evidence) indicative of a
Deleted: location, however
Deleted: The
Deleted: of all eruptions
Deleted: . We
Deleted: , however,
Deleted: the
Deleted: of each eruption
Deleted: .
Deleted: these four eruptions
Deleted: is
Deleted: The injection
Deleted: is
Deleted: that of
Deleted: in the absence of any
Deleted: is
Deleted: PMIP4 (Paleoclimate Model Intercomparison Project)
Deleted: These include
Deleted: as well as the
Deleted: are
Deleted: change
Deleted: the
Deleted: a
Deleted: amount of
Deleted: change. The orbital

2.2 Experiment Design

A control simulation for the 2.5ka period was performed using the PMIP phase 4 protocols for the mid-Holocene (6ka) experiment, altered for conditions appropriate to 2.5ka. This included altering the orbital forcing, greenhouse gases (CO₂: 279 ppm, N₂O: 266 ppb, and CH₄: 610 ppb), plus vegetation in Africa and high boreal Eurasia and North America (Otto-Bliesner et al., 2017). Ozone and aerosols were prescribed to non-anthropogenic conditions only – this is distinct from pre-industrial simulations that do include small, anthropogenic changes and attendant aerosol and atmospheric chemistry changes. Orbital

and greenhouse gas forcings for the 2.5ka period are expected to play a vital role in producing the correct equilibrium climate. We ran a control run with the NINT configuration for 1000 years to get the model in equilibrium, then extended this for 100 years by adding the MATRIX version of ModelE2.1 to again achieve an equilibrium state for a 2.5ka period with composition-climate interactions enabled. Vegetation cover, LAI and vegetation height were prescribed corresponding to the piControl period climate. A lack of exact vegetation cover information prevents GCMs without dynamic vegetation model from reproducing mid-Holocene warm Northern hemisphere summer and enhanced NH monsoons conditions (Tierney et al., 2017; Larrasoana et al., 2013). However, the vegetation cover used here as defined by the PMIP4 protocol vegetation sensitivity experiment (Otto-Bliesner et al., 2017) for the mid-Holocene period shows an intense impact on North African rainfall and explains the difference between simulated and reconstructed climate conditions (Braconnot et al., 1999; Pausata et al., 2016). To address this, we created a modified mid-Holocene boundary condition sensitivity vegetation map by linearly interpolating between pre-industrial vegetation and the Mid-Holocene vegetation sensitivity experiment (i.e., northern hemisphere high latitude tundra during the pre-industrial was replaced by boreal forests, while African vegetation was altered with evergreen shrubs replacing all vegetation up to 25N and grasslands up through the Mediterranean Coast in 6ka (Otto-Bliesner et al., 2017)).

Fig S1 (Supplementary Information) shows the major vegetation plant function type (PFT) cover changes under the PMIP4 sensitivity vegetation protocols after linearly interpolating for the 2.5ka period. The 2.5k equilibrated simulation with MATRIX was then extended for 70 more years with a corrected dust tuning (a typical process when equilibrating the model on a new climate state), and a further 130 years with the linearly interpolated PMIP4 vegetation described above (see table TS1 for details of control runs and annual global mean time series of surface air temperature and precipitation in Fig. S2). This run equilibrated very quickly and no further tuning was needed. We thus used the last 100 of the total 130 years of that equilibrated run as the base climate for our analysis. An ensemble of 10 members with active volcanic eruptions was simulated using a restart file every 10 years during the last 100 years of the control simulation corresponding to 2.5ka period as summarized in table TS1, following the same approach as performed for the CMIP6 ensemble simulations (Kelley et al., 2020).

Deleted: and

Deleted: are

Deleted: The

Deleted: for the relevant period

Deleted: the

Deleted: that lack

Deleted: ,

Deleted: ,

Deleted: -

Deleted: Pre-Industrial

Deleted: preindustrial is

Deleted: and

Deleted: ,

Deleted:).

Deleted: information

Deleted: ¶

Deleted: is

Deleted: ,

Deleted: ,

Deleted: refer to

Moved down [1]: The starting timepoint for each ensemble member is shown by blue vertical lines in fig S2.

Deleted: Each ensemble member started on January 1st of the year 169 BCE and ran for 16 years, with each eruption happening on the 15th of June of the 2nd, 6th, 9th and 12th years modelled. Because the exact date of an eruption cannot be directly determined based upon ice-core sulphate deposition data, both because of possible uncertainties in the ice-core chronologies and because of variable time lags between eruptions and the atmospheric circulation of the resulting sulphate and its deposition in the polar ice, we selected a summer eruption date to investigate the impact on northern hemisphere monsoon and wintertime atmospheric circulation. We also note that the accuracy of our modelling will depend in part upon the accuracy of the ice-core-based volcanic forcing reconstruction being employed. Uncertainties in reconstructed forcing can arise, for example, because of variation in the deposition of sulphate across the polar regions for any given eruption. In this respect, it is important to note that the Sigl et al. (2015) volcanic forcing reconstruction employs several ice-cores from Antarctica and Greenland, but our results can be revisited as reconstructions become more reliable by incorporating larger numbers of ice-cores.

Formatted: Font: Times New Roman

The starting timepoint for each ensemble member is shown by blue vertical lines in fig S2. Each member started January 1st of the year 169 BCE and ran for 16 years, with each eruption happening on the 15th of June of the 2nd, 6th, 9th and 12th years modelled. Because exact eruption dates cannot be determined from ice-core sulphate deposition data, due to ice-core chronological uncertainties and lags between eruptions and the deposition of sulphate in the ice, we selected a summer eruption date to investigate impacts on northern hemisphere monsoon and wintertime atmospheric circulation. We also note that modelling accuracy will depend partly upon the accuracy of the ice-core-based forcing reconstruction employed. Uncertainties can arise, for example, because of variation in sulphate deposition across the polar regions for any given eruption. It is thus notable Sigl et al. (2015) employ several Antarctic and Greenland ice-cores in their reconstruction, helping to average out regional variability in deposition, but our results can be revisited as reconstructions incorporate more ice-cores.

Moved (insertion) [1]

Table 1. Details of eruptions applied in this experiment, with each eruption happening on the 15th of June of the 2nd, 6th, 9th and 12th model years.

Eruption	Year (BCE)	Position	Eruption injection (SO ₂)	Injection Height (km)
E1	168	Pinatubo (15.13, 120.35) (Tropical)	22.5 Tg	22-26
E2	164	Mt Laki, Iceland (64.03, -18.13 W) (NH)	6.5 Tg	22-26
E3	161	Mt Katmai, Alaska Peninsula (58.28, -154.95 W) (NH)	7.2 Tg	22-26
E4	158	Shiveluch, Kamchatka, Russia (56.39, 161.21) (NH)	7.5 Tg	22-26

3. Results

810 3.1. 2.5ka control runs

We first evaluated the 2.5ka control climate run for providing a precise background climate to investigate the impacts of the 168-158 BCE volcanic quartet.

3.1.1 2.5Ka GHG+ORB climate

We compared the 2.5ka equilibrium climate with only GHG, orbital, and non-anthropogenic forcing changes against a preindustrial (year 1850) control run to evaluate the impact (alone) of orbital and greenhouse gas changes on our base climate state. Surface air temperatures showed globally minimal differences with a warming of northern hemisphere high latitudes due to the different orbital forcing for all seasons (Fig 1). The implications of changes in orbital forcing for 2.5k are thus evident in the surface temperature but the northern hemisphere monsoon season (JJAS) and winter season (DJF) rainfall slightly was reduced along the northern equatorial belt. This points to the limitation of the GISS model in not having an interactively dynamic vegetation component to reproduce the known mid-Holocene wet African landcover conditions (Harrison et al., 2015; Tiwari et al., 2022). Numerous studies have demonstrated that including bio-geophysical feedbacks and atmospheric dynamics helps to successfully model the wet African conditions for mid-Holocene (Kutzbach et al., 1996; Claussen et al., 2003; Kutzbach and Liu, 1997; Hewitt and Mitchell, 1998). Using PMIP4 vegetation over northern hemispheric regions is known to provide a solution to the same long-standing issue with CMIP3/CMIP5 models that fail to reproduce these wet African conditions for mid-Holocene (Harrison et al., 2015).

- Deleted: for investigating
- Deleted: hydroclimatic
- Deleted: from 168-158 BCE
- Deleted: alone
- Deleted: show
- Deleted: the
- Deleted: decreases
- Deleted: land cover
- Deleted: ,
- Deleted: both biogeophysical feedback processes
- Deleted: in achieving
- Deleted: ..
- Deleted: the
- Deleted: the
- Deleted: hemisphere
- Deleted: has been shown
- Deleted: a




Fig 1. Seasonal mean (Annual, DJF & JJAS) surface air temperature (top row) for 2.5k period equilibrium run, differences from the preindustrial period (2.5ka-preindustrial) for all three seasons (2nd row from top) and seasonal (Annual, DJF & JJAS) mean precipitation (3rd row from top) and the difference (bottom row) from preindustrial period (2.5ka-preindustrial). The equilibrium run for the 2.5k period includes the orbital and GHG concentration changes for the 2.5k period (referred to as OG), the preindustrial period (as PI), and their difference (OG-PI) as simulated by GISS ModelE2.1.

Deleted: means

Deleted: of

Deleted: include

3.1.2 2.5Ka ORB+GHG+VEG climate

The comparison of mean climate for the 2.5ka period for inclusion of PMIP4 vegetation is shown in fig 2 for the mean surface air temperature and precipitation for the Annual, DJF and northern hemisphere monsoon (JJAS) seasons.




Fig 2. Mean surface air temperature for Annual, DJF and JJAS seasons (top row) and seasonal mean precipitation (3rd row from top) for the equilibrium runs with the PMIP4 vegetation for the 2.5k period and surface temperature difference (2nd row from top) plus the seasonal precipitation differences (bottom row) for the 2.5k period as simulated by GISS ModelE2.1. We have used a short initial notation for our forcings to denote the difference (ORB+GHG+VEG = OGV and ORB+GHG= OG)

GISS ModelE2.1 simulated a global a mean surface air temperature (SAT) of 14.4 °C, 12.8 °C and 15.7 °C for Annual, DJF and JJAS seasons respectively for the 2.5ka (ORB+GHG+VEG) simulation, which is

- Deleted: as well as
- Deleted: forcing
- Deleted: simulates

0.11 C (Annual), 0.11 °C (DJF) and 0.08 °C (JJAS) higher than the 2.5ka (ORB+GHG) simulation without including these vegetation changes. A strong increase in surface air temperature of greater than 2°C was calculated for northern hemisphere high latitude land regions, particularly where landcover (tundra) was replaced by boreal forest, decreasing ground surface albedo during snowy winter months. A moderate rise of 0.5°C was also simulated for Africa, coinciding with regions of vegetation changes (described in section 2.2). The regional pattern of difference in rainfall in the northern hemisphere monsoon season (JJAS) was observed mostly over the North African and Asian regions. The observed increase of 0.4 mm/day or greater over the North African and Southwest Asian monsoon regions indicates a northward movement of the ITCZ during the monsoon season, consistent with expectations given the modified vegetation for this period and in agreement with current understanding of mid-Holocene rainfall regimes (Tierney et al., 2017; Tiwari et al., 2022). These results also acknowledge the sensitivity of the hydroclimatic impacts to regional landcover and hydrological conditions (Singh et al., 2020).

We also analyzed zonal changes of longwave and shortwave radiation at the top of the atmosphere with our altered surface albedo (Fig 3). Vegetation-albedo feedback from the inclusion of woody forest over higher latitudes and shrubs and steppes over northern Africa is important in the additional monsoon season rainfall seen for North Africa. Greater vegetation cover for the Sahara and at higher Northern Hemispheric latitudes alters surface albedo by $\geq 10\%$ regionally as well as altering absorption of incoming solar radiations across northern hemisphere higher latitudes (Fig 3). Consequently, the pole-equator temperature gradient increases, pulling the ITCZ northwards (see Fig 2). We thus concluded that the control climate generated using PMIP4 vegetation scaled from the mid-Holocene to the 2.5k period provides more precise control conditions to investigate the hydroclimatic impact of volcanic forcing perturbations. Vegetation boundary conditions implemented according to the PMIP4 sensitivity experiments with orbital and greenhouse gas forcing thus helped to produce a precise equilibrium climate, for this historically and climatically critical period 2.5ka years ago.

Deleted: the ...urface air temperature of greater than 2°C is...as calculated over the...or northern hemisphere high latitude land regions, particularly in areas ...here land cover...andcover (tundra) is...as replaced by boreal forest, decreasing ground surface albedo during snowy winter months. A moderate rise of 0.5°C over Africa is...as also simulated, which coincides...for Africa, coinciding with the ...egions of vegetation changes as ...described in section 2.2.... The regional pattern of difference in rainfall in the northern hemisphere monsoon season (JJAS) is...as observed mostly over the North African and Asian regions. This...he observed increase of 0.4 mm/day or greater over the North African and Southwest Asian monsoon regions indicates a northward movement of the ITCZ during the monsoon season that is... consistent with expectations given the modified vegetation for this period and is ...n agreement with our ...urrent understanding of mid-Holocene rainfall regimes (Tierney et al., 2017; Tiwari et al., ..., 2022). These results also acknowledge the sensitivity of the hydroclimatic impacts due ...o volcanic eruption to the ...egional land cover...andcover and hydrology alteration

... [17]

Deleted: the ...onal changes of longwave and shortwave radiation at the top of the atmosphere with our altered ground...urface albedo, as shown in ... (Fig 3). The vegetation.... Vegetation-albedo feedback due to...rom the inclusion of woody forest on...ver higher latitudes and shrubs and steppes over northern Africa plays a crucial role...s important in the additional monsoon season rainfall over the...een for North African region....frica. Greater vegetation cover for the Sahara and at higher Northern Hemispheric latitudes in the Northern hemisphere ...lters the ground...urface albedo by more than ...10% regionally as well as altering the ...bsorption of incoming solar radiations across the ...orthern hemisphere higher latitudes (Fig 3). Consequently, it increases ...he pole-equator temperature gradient and pulls...ncreases, pulling the ITCZ northwards as shown in...see Fig 2. It was.... We thus concluded that the control climate using the ...MIP4 vegetation scaled from the mid-Holocene to the 2.5k period provides more precise and suitable ...ontrol conditions to investigate the climatic...ydroclimatic impact of volcanic forcing perturbations due to volcanic eruptions... Vegetation boundary conditions implemented according to the PMIP4 sensitivity experiments with orbital and greenhouse gas forcing...thus helped to produce a precise equilibrium climate condition

... [18]




Fig 3. Annual zonal mean of ground albedo (A), longwave & shortwave radiation at the top of the atmosphere (ToA; B and C) for 2.5ka climates with ORB+GHG and ORB+GHG+VEG (red and green lines), respectively. The blue line shows the difference between them. The blue line in panel B & C is **exaggerated** 10-fold (x10) to **show the difference more clearly** on the same vertical axis.

Deleted: the original difference values in order to clearly

3.2. Radiative forcing and climate response to volcanic aerosols

We **simulated** a series of four eruptions occurring mid-June during the 2nd, 6th, 9th and 12th years of **our** runs (as **per** section 2.2 and Table 1). Explosively injected SO₂ oxidizes to form **stratospheric sulphate** aerosols that can alter the radiative balance at the top of the atmosphere by scattering incoming solar radiation and absorbing and re-emitting longwave radiation. Fig 4 shows the different components of the radiative budget on a monthly scale, with the annual cycle climatology removed for the entire period covering all four eruptions. The relative impacts of scattering **shortwave (SW)** and absorbing **longwave (LW)** radiation is proportional to the sulfate aerosol size (Lacis, 1992). The model simulated a lifetime for volcanically injected SO₂ as 31.4±0.72 days for eruption E1 and 24.4±0.44, 25.02±0.40 and 25.5±0.36 days for eruptions E2, E3 and E4, respectively. Other studies have reported a comparable average lifetime of 33 days (Read et al., 1993), 25±5 days (Guo et al., 2004), 35 and 25 days (Bluth et al., 1992; Schnetzler et al., 1995) for SO₂ injected from **the** 1991 Pinatubo eruption using various satellite retrievals.

Deleted: simulate

Deleted: all

Deleted: ,

Deleted: the simulation,

Deleted: described in

Deleted: .

Deleted: in the stratosphere

Deleted: then

Deleted: the

Deleted: the




Fig 4: Monthly mean top of the atmosphere radiative balance perturbation due to volcanic aerosols for the entire simulation length. Orange/red shows the longwave radiative response, light/dark blue represents the shortwave, and grey/black represents the net (ToA) radiative change averaged at the global scale. The light-colored solid lines represent individual ensemble members, and the dark-colored lines show the ensemble mean. The green vertical dashed lines show when the eruptions happened.

With 22.5 Tg of SO₂ injected, the first, tropical eruption was larger than Pinatubo (~30%) and altered the longwave radiation budget by a mean of ~3 W/m² for almost a year, while the other three eruptions were approximately 1/3rd of Pinatubo and produced a perturbation of the longwave radiative budget by ~1 W/m². The model [simulated](#) a strong impact on the shortwave radiation budget up to a mean of ~10 W/m² for a few months after the first eruption and of ~4 W/m² for a few months after each of the subsequent eruptions. A mean imbalance of up to -7.5 W/m² after the first eruption and -2.5 to -3 W/m² after the other eruptions in the top of atmosphere net radiative forcing suggests a strong corresponding surface cooling.

Deleted: after the eruption

Deleted: simulates

Note that the bumps in the various radiative forcing trajectories (Fig. 5) in the year after each eruption reflect the northern hemispheric seasonal cycle. The presence of volcanic aerosols in the atmosphere impacted climate in several important ways, described below.

Deleted: apparent in
Deleted: in the northern hemisphere.
Deleted: these
Deleted: keyways, as




Fig 5. Globally averaged changes in MSU TLS (top panel), surface air temperature (middle panel) and total atmospheric column AOD at 550 nm for each month for the entire simulation period. The light-colored solid lines represent individual ensemble members, and the solid dark colors show the ensemble means. The green vertical dashed lines show when the eruptions happened.

The top panel in Fig. 5 shows the monthly change in microwave sounding unit (MSU) temperature for the lower stratosphere (TLS) as calculated by the model, which is a typical metric for present-day evaluation of modeled stratospheric temperatures against satellite data. It covers the lower stratosphere, where volcanic aerosols mostly reside, and represents the local atmospheric response of longwave absorption by them. After Pinatubo (1991), a lower stratospheric warming on the order of 2-3°C for a year was estimated using multiple reanalysis products (Labitzke and McCormick, 1992; Fujiwara et al., 2015). This is comparable to the ~25% larger eruption simulated here, E1, in which volcanic aerosols spread over a larger region (in both the northern and southern hemispheres) and absorbed a significant portion of longwave radiation, warming the lower stratosphere by up to 3°C for the first two years post-eruption. This effect intensified during the second year, before steadily declining in Years 3 and 4 with the scavenging of volcanic aerosols. The three subsequent extratropical eruptions warmed the lower stratosphere by up to 0.5°C only, because these were weaker and only affected the northern hemisphere for a shorter period (~18 months).

The lower panel in Fig 5 presents the aerosol optical depth (AOD), a measure of atmospheric opacity to incoming radiation calculated as the extinction (sum of scattering and absorption) of shortwave radiation at 550nm. The model simulated an AOD anomaly of around 0.21 for the first 18 months after E1, decreasing as aerosols were progressively removed. The subsequent eruptions produced an AOD of the order of ~0.1 that similarly decreased with time. For comparison, the AOD estimation for Pinatubo (1991) is 0.15 for approximately 12 months over a background optical depth of ~0.6 (Russell et al., 1996; English et al., 2013).

In the upper troposphere and lower stratosphere, the impact of each eruption was distinct with a near-complete recovery to background AOD levels observed after each (i.e., before the next) event; however, at the surface, a lag in recovery time was evident (middle panel, Fig 5). The net impact of radiative flux perturbations following the eruptions is summarized using the global surface air temperature change for the entire period. The model produced a mean cooling of ~1.5°C in the second year after E1 and ~1.0°C after all other eruptions. Although AOD recovered a few years after each eruption, the surface temperature response was more prolonged. This lag in response can be mainly ascribed to the thermal inertia of the oceans, requiring a greater recovery time. This is borne out by the sea surface temperature

- Deleted: lie
- Deleted: the Mount
- Deleted:) eruption,
- Deleted: of
- Deleted: has been
- Deleted: after the
- Deleted: was seen to intensify
- Deleted: starting to
- Deleted: decline
- Deleted: other
- Deleted: both
- Deleted: extratropical eruptions that
- Deleted: the
- Deleted: simulates
- Deleted: the first eruption, which decreases
- Deleted: are
- Deleted: produce
- Deleted: which
- Deleted: decreases
- Deleted: the
- Deleted: eruption
- Deleted: of these eruptions is
- Deleted: and
- Deleted: is
- Deleted: in
- Deleted: in the form of
- Deleted: over
- Deleted: produces
- Deleted: robust
- Deleted: (or proceeding year)
- Deleted: the first eruption, and although
- Deleted: recovers
- Deleted: is
- Deleted: global mean surface air temperature
- Deleted: which need more
- Deleted: to return to the normal.

(SST) response shown in fig S4, with a slower post-eruption recovery and remanent cooling effect. The smaller extratropical eruptions (E2-E4) that followed the large tropical E1, were then observed to hinder surface temperature recovery, maintaining a surface cooling of around 1.0°C for the entire simulation period. Comparatively, Pinatubo (1991) created an ~0.5°C (peak) cooling over 1-2 years post-eruption (Hansen et al., 1996).

3.3 Volcanic aerosol properties

Radiative forcing from volcanic aerosols is tightly controlled by aerosol size (Lacis et al., 1992; Hansen et al., 1980). The aerosol effective radius, R_{eff} , is a key metric in linking aerosol microphysical properties with their SW and LW impacts. The vertical profile of aerosol size as represented by R_{eff} was calculated for each month (Fig 6). After the tropical eruption (E1), new aerosols nucleated and grew rapidly via coagulation and, while SO₂ was still available, by condensation, and attained a maximum R_{eff} of $\geq 0.5 \mu\text{m}$ approximately for 2 years. In comparison, R_{eff} after Pinatubo (1991) increased to $0.6 \mu\text{m}$ and sustained that size for approximately 2 years (Russell et al., 1996). Sulfate aerosol sizes for the three subsequent extratropical eruptions (E2 to E4) grew up to $0.3 \mu\text{m}$.

- Deleted: is
- Deleted: (as shown in the supplementary information (Fig S4)).
- Deleted: one (
- Deleted:) are
- Deleted: the
- Deleted: and maintain
- Deleted: during
- Deleted: period of
- Deleted: the 1991 Mt.
- Deleted: eruption
- Deleted: after the
- Deleted: The effect of
- Deleted: on radiative forcing
- Deleted: is
- Deleted: and is shown in
- Deleted: .
- Deleted: greater than
- Deleted: the
- Deleted: eruption went up
- Deleted: three smaller




Fig 6. Timeseries of global ensemble mean vertical profile of sulfate aerosol R_{eff} for the entire simulation period. The vertical black line with a circled cross mark on the horizontal axis shows the eruption timings.

The aerosol extinction vertical profile (Fig S5A) shows that the radiative impact of the E1 tropical eruption in the lower stratosphere was prolonged as compared to the later extratropical eruptions. Lower stratospheric heating affects the dynamics of the stratosphere; after tropical eruptions enhanced tropical upwelling and extratropical downwelling with the phase of Brewer-Dobson circulation impact upon the transportation of trace species such as Ozone (O_3) and NO_2 (Aquila et al., 2013; Trepte et al., 1992; Pitari et al., 2016; Pitari and Mancini, 2002). Fig S5B shows a strong positive (≥ 10 ppbv) anomaly of CH_4 in the upper stratosphere and negative (≤ 10 ppbv) anomalies in the lower stratosphere, especially after the tropical eruption (E1). Changes in the mean concentration of upper and lower stratospheric methane (CH_4) suggest a strong vertical transport (Kilian et al., 2020).

3.4. Latitudinal temperature response to volcanic aerosol forcing

The Hovmöller diagram (Fig 7A, 7B) shows the differences between the zonally averaged AOD at 550nm and surface air temperature response for the ensemble means of the volcanic eruption simulations compared to the mean climatology of the control simulation. The statistical significance level is estimated using the 2-tail student t-test after Deser et al. (2012) and following the assertion that 10 ensembles are sufficient for reasonable estimation of internal variability at a regional scale (Singh and AchutaRao, 2019). The pattern of total AOD after the first eruption (E1) showed a strong cross-equatorial transport of stratospheric aerosols into the southern hemisphere, with a similar pattern in the northern hemisphere. This is consistent with the hypothesis that an enhanced Brewer-Dobson circulation in the southern hemisphere during the austral winter can lead to the southward transportation of volcanic aerosols after a Pinatubo type (tropical) eruption (Aquila et al., 2012). The initial dispersal of aerosols from eruption E1 was strongly influenced by its timing and exhibited a seasonal dependence (consistent with Toohey et al., 2011). However, the other three eruptions (E2, E3 and E4) in the high latitude Northern Hemisphere only yielded an increased AOD in that hemisphere.

A lag of more than 12 months in peak surface temperature response (Fig 5, Middle panel) after E1 correlates well with the modelled aerosol distribution, consistent with reporting for similar events (e.g.,

- Deleted: the
- Deleted: timing of the eruptions
- Deleted: Heating of the lower stratosphere
- Deleted: have an
- Deleted: on
- Deleted: and
- Deleted: as
- Deleted: ,
- Deleted: shows
- Deleted: transportation
- Deleted: the
- Deleted: season
- Deleted: .
- Deleted: extratropic
- Deleted: the northern
- Deleted: the
- Deleted: the first eruption
- Deleted: of aerosols
- Deleted: findings reported in the literature

190 Jungclauss et al., 2010; Klocke, 2011). The peak global mean surface temperature response thus appeared when aerosols from the tropical eruption (E1) had extended across the northern hemisphere extratropics and polar regions. It should be noted that northern extratropical land surfaces responded quickly to the attenuated post-eruption shortwave radiative flux compared to the tropics. The zonally averaged surface temperature response (fig 7B) showed that a strong cooling of 1.0-1.5°C lasted over the tropical north and

195 partially over the tropical southern hemisphere for more than 30 months after E1. Further, the largest anomalies of >2.0 °C cooling mostly appeared six months after E1, with the subsequent extratropical eruptions helping to maintain the northern hemispheric cooling.

The seasonality of surface temperature response revealed a more substantial cooling during the boreal summer season for all four eruptions and for E1 also revealed the expected post-tropical-eruption winter warming pattern over Europe. Fig S6 shows the spatial pattern of the surface temperature response to volcanic aerosols over the four seasons directly following E1 (JJA & SON for the eruption year and DJF & MAM for first year following). The response for the first two seasons was confined to the tropics but moved to higher latitudes thereafter. As evident in fig S3, the winter (DJF) post-eruption warming over

200 Europe and an observed cooling over Northern America may result from the same fundamental

205 atmospheric dynamics noticed after Pinatubo (1991) (Robock, 2000; Robock and Mao, 1992).

- Deleted: peak
- Deleted: appears
- Deleted: the volcanic
- Deleted: have
- Deleted: the
- Deleted: the land surface over the
- Deleted: extratropics is observed to respond
- Deleted: shows
- Deleted: the first eruption.
- Deleted: greater
- Deleted: the first eruption
- Deleted: reveals
- Deleted: reveals
- Deleted: Supplementary
- Deleted: the first eruption (
- Deleted:)
- Deleted: of eruption
- Deleted: the next
- Deleted: surface temperature
- Deleted: is
- Deleted: and moves
- Deleted: after six months.
- Deleted: anomalous
- Deleted: pattern after the eruption
- Deleted: be a product of
- Deleted: as
- Deleted: eruption




Fig 7. Hovmöller diagram showing the zonally averaged temporal dispersion of volcanic aerosols in terms of AOD change at 550nm (A), surface temperature response (B), and lower stratospheric temperature response (C). Anomalies were calculated with respect to a climatological annual cycle calculated from the control simulation. The gray color is painted over regions where changes are not statistically significant at the 95% confidence level. Circled cross marks show the modeled spatial and temporal position of the eruptions.

The global lower stratospheric temperature response in terms of MSU TLS data was discussed in section 3.2. Interestingly, fig 7C shows that the latitudinal anomaly of the lower stratosphere warming was broadly limited to the equatorial lower stratosphere. E1 induced lower stratosphere warming on the order of >3 °C, with a weaker warming of up to 1-2 °C after E2, E3 and E4. Lower stratosphere warming is also known to affect the polar vortex strength in the northern hemisphere and atmospheric circulations

Deleted: the

Deleted: has been

Deleted: is

Deleted: Eruption

Deleted: induces

Deleted: the three extratropical eruptions (

Deleted:).

Deleted: affects

into the troposphere, with repercussions for surface climate and variability patterns (e.g., Graf et al., 1993, 2007; Shindell et al., 2004).

- Deleted: substantial
- Deleted: as suggested in previous research

3.5 Latitudinal precipitation response to volcanic aerosols

We used a coarser resolution earth system model having a simplified parameterization and successful in simulating the large-scale patterns of rainfall change (Kelley et al., 2020). Studies of observational records plus modeling efforts have demonstrated that the cascading impact of an altered radiative balance at the top of the atmosphere due to volcanic eruptions is reflected in the hydrological cycle in regional patterns of seasonal rainfall change (e.g., Robock and Liu, 1994; Robock, 2000; Trenberth & Dai, 2006; Schneider et al., 2009; Iles et al., 2012; Iles and Hegerl, 2014; Timmreck, 2012). We investigated the hydrological cycle response to the 168-158 BCE eruption quartet at both a global and regional scale, paying particular attention to the northern hemispherical monsoon season (JJAS) for the first 2 years following each eruption. Any individual ensemble member might best represent the historical reality, but it is impossible to select the most accurate member absent supporting observational data from the period. Also, added noise due to natural variability can be greater at the regional scale, even to the extent of altering the sign of observed changes among the individual ensemble runs. Thus, we mainly focused upon the mean from across the ensemble when examining the response to the eruptions for the various climate variables considered. Fig 8 shows the Hovmöller diagram of the zonal mean precipitation anomaly relative to the annual cycle climatology of the 100-year-long control simulation.

- Deleted: Numerous studies using the
- Deleted: record in addition to
- Deleted: terms of
- Deleted: Societies conducting agriculture in arid and semi-arid regions before the advent of modern reservoirs and irrigation, such as in ancient Egypt, are perhaps most impacted by such changes.
- Deleted: thus investigate
- Deleted: this set of eruptions
- Deleted: for the period under study. It can be argued that any
- Deleted: period
- Deleted: in the absence of
- Deleted: of interest
- Deleted: the
- Deleted: alter the sign of change
- Deleted: spatial
- Deleted: ensembles.
- Deleted: selected
- Deleted: and its statistical inference to show
- Deleted: volcanic
- Deleted: with robustness
- Deleted: specific
- Deleted: under consideration




Fig 8. Hovmöller diagram showing the zonally averaged rainfall anomaly for the entire period as the spatiotemporal response of global rainfall to our series of volcanic eruptions. Circled cross marks show the locations and timing of the eruptions. Black dots point out regions where changes are not statistically significant at the 95% confidence level.


The ensemble zonal mean post-eruption rainfall change showed a substantial negative trend in the northern hemisphere due to the volcanic aerosol-induced cooling. A robust negative anomaly on the order of 0.3-0.4 mm/day in the northern hemisphere rain belt (ITCZ) region appeared shortly following E1 and persisting for several years (Fig 8). A pattern of strong drying in the equator also coincided with the northern hemisphere monsoon season (JJAS) for 2 to 3 years after E1. However, because the northern hemispheric extratropical rainfall response strongly correlates with the surface temperature response, it thus emerged here 12 months after E1, with the model calculating a moderate to high decrease on the order of 0.1-0.2 mm/day, persisting throughout the year for three post eruption years. A shift in the northern hemisphere rainfall pattern was also evident for the region around 30°N, with slight increases in rainfall for 2 years after E1. This response was statistically significant over only a few spots, however. Fig 8 demonstrates that a drying pattern also prevailed after the later extratropical eruptions (E2-E4).

- Deleted: the
- Deleted: after the eruptions
- Deleted: as a result of cooling induced by
- Deleted: aerosols.
- Deleted: of
- Deleted: the first eruption (E1) and persisted during the following couple of
- Deleted: the eruption (
- Deleted:).
- Deleted: in the northern hemisphere extratropics
- Deleted: was
- Deleted: seen to emerge
- Deleted: the tropical eruption (
- Deleted:),
- Deleted: a
- Deleted: in rainfall, a response that persisted
- Deleted: the tropical eruption (
- Deleted:).
- Deleted: after the first eruption
- Deleted: clearly

1335 Thus, the northern hemisphere experienced a sustained net (albeit temporally varying) precipitation decline for the entire modeled period, with a distinct seasonal character.

Deleted: Because of this

Deleted: and



1340 Fig 9. Mean change (mm/day) in northern hemisphere monsoon season (JJAS) rainfall averaged for three consecutive years after eruption E1 (including the eruption year) and two years each after E2, E3 and E4 (including the eruption years) (left to right and top to bottom). The caption over each panel shows the eruption characteristics. A gray color is painted over the grid boxes for which rainfall change is not significant at the 95% confidence level. Years indicated in parentheses follow the order of the eruptions in our simulation period, i.e., E1 occurs in the 2nd simulation year, and E2-E4 occur in the 6th, 9th and 12th years, respectively.

Deleted: each of

Deleted: change in

We further evaluated the spatial patterns of change in mean rainfall during the northern hemisphere monsoon season (JJAS) (fig 9). We averaged the three monsoon seasons (eruption year and next 2 years) after the more potent tropical eruption (E1) and two monsoon seasons (eruption year and next year) after

Deleted: as shown in Fig

Deleted: .

each of the extratropical eruptions (E2, E3 and E4), focusing principally on statistically significant responses. After E1, summer monsoon rainfall appeared strongly suppressed over many major northern hemisphere monsoon regions. Importantly for our focus on Egypt, African monsoon rainfall showed a notable decrease of 0.5-1.0 mm/day during the three-year post-eruption JJAS season average (i.e., derived from the eruption year and first two post-eruption years). This decrease covered a large area in Africa from (approximately) the equator to (approximately) 17°N. The South and East Asian monsoon regions also exhibited a robust negative rainfall anomaly of >1.0 mm/day over the Indian subcontinent and (more variably) several regions of China, though with some isolated increase over the eastern Vietnamese landmass. Similar patterns of decrease appeared over the western (particularly northwestern) Pacific and northern hemispheric high latitude regions more broadly. The model also simulated a (statistically significant) band of enhanced JJAS rainfall stretching from Central Asia westward through the Middle East and into the Mediterranean (touching parts of northern Africa), Western European and parts of the North Atlantic (roughly between a latitudinal band of 30°N to 50°N). A contiguous band of increased rainfall was observed further south and west in the Atlantic, stretching into parts of the northern Caribbean, southeastern Gulf of Mexico and Mesoamerica (fig 9).

Similar patterns of suppressed boreal monsoon season rainfall were observed following the extratropical eruptions (E2-E4), but a particularly notable east-west band over land and ocean (broadly confined between slightly north of the equator and 30°S) shows a positive rainfall anomaly (most clearly statistically significant between (approximately) 5°N and 10°S (fig 9)). This is largely consistent with observations and modeling of volcanic climatic impacts under a range of scenarios and periods (e.g., Robock, 2000; Robock and Liu, 1994; Iles et al., 2012; Liu et al., 2016; Haywood et al., 2013; Schneider et al., 2009; Trenberth and Dai, 2007; Joseph and Zeng, 2011; Gu and Adler, 2011). Of mechanisms, for many northern hemisphere landmasses, these eruptions induced a surface cooling that altered the meridional (equator-to-pole) surface temperature gradient (fig 7). Given this energetic deficit, we posit that the northern hemisphere experienced a post-eruption alteration of large-scale circulation patterns and moisture convergence, producing a constrained northward ITCZ migration during the boreal summer, diminishing rainfall over many northern hemispheric monsoon regions and (relatedly) promoting increased rainfall in the above-described band from the equator southward (Liu et al., 2016; Oman et al.,

Deleted: remaining
Deleted: and focused
Deleted: identifying
Deleted: Hence, after the tropical eruption, the
Deleted: historical

Deleted: were
Deleted: shown to experience
Deleted: as well as
Deleted: decreased rainfall also
Deleted: and

Deleted: Near
Deleted: on

Deleted: also

Deleted: each of
Deleted: both
Deleted: being
Deleted: pattern
Deleted: past literature employing

Deleted: In terms of
Deleted: clearly
Deleted: northern hemisphere
Deleted: may
Deleted: (NH)
Deleted: resulting in
Deleted: of the ITCZ
Deleted: suppressing large scale
Deleted: patterns
Deleted: as a related consequence

2006; Graf, 1992; Dogar, 2018). This is consistent with Colose et al. (2016). They demonstrated that a hemispherically asymmetric volcanic forcing creates energetically deficient conditions in the hemisphere of the greatest forcing that “pushes” the ITCZ away from it. Palaeoclimatic data also shows that tropical and northern hemispheric eruptions can create a dipole resulting in wetter summer conditions over extensive parts of the Mediterranean, with correspondingly drier conditions over northern Europe (Rao et al., 2017). This is also largely consistent with our model output (fig 9).

Deleted: analyses of Colose et al. (2016), who have demonstrated that a hemispherically asymmetric volcanic forcing creates energetically deficient conditions in the hemisphere of the greatest forcing and which “pushes” the ITCZ away from it. It has also been shown with palaeoclimatic data also shows that tropical and northern hemisphere hemispheric eruptions can create a dipole that results

[19]

3.6 African monsoon and Nile River response

Our modeling suggests that all four eruptions, 168-158 BCE, are likely to have influenced rainfall over different monsoon regions in the northern hemisphere for 2-3 years after each eruption, in combination producing a sustained deficit for more than a decade. Focusing on the North African monsoon region, fig 10 shows three consecutive years of JJAS rainfall over equatorial and northern Africa (encompassing the Nile basin) after each eruption. The African monsoon exhibited notably reduced rainfall of ≥ 1 mm/day following the (mid-June) E1 tropical eruption, starting in the eruption year itself (Year 0, fig 10), affecting both the White Nile watershed in the south of the basin and the Blue Nile and Atbara River watersheds further north and east in the Ethiopian Highlands. Reduced precipitation was also observed following each (also mid-June) extratropical eruption (E2-E4) during the eruption years but was more spatially constrained (and particularly for E2, less severe). This is perhaps unsurprising as the estimated SO_2 output of E2-E4 is approximately 1/3 of E1. Nonetheless, in each case the Blue Nile and Atbara River headwaters experienced a statistically significant decrease, with implications for the summer flood in Egypt, which depends for approximately 80% of its floodwater on rainfall there (Melesse et al., 2011). For E2-E4, this response intensified in the first full post-eruption year (Year 1, fig 10), persisting into the second full post-eruption year (Year 2, fig 10), while for E1 the response contracted geographically in Years 1-2, resembling that seen after E2-E4 (fig 9, fig 10). This suppression of the African monsoon following tropical and northern hemispheric extratropical eruptions is consistent with previous studies (e.g., Colose et al., 2016; Oman et al., 2006; Haywood et al., 2013; Jacobson et al., 2020; Manning et al., 2017).

Deleted: during the period... 168-158 BCE, are likely to have strongly influenced the rainfall pattern over different monsoon rainfall regions in the northern hemisphere consecutively for 2-3 years after each eruption, and in combination producing a sustained deficit in monsoon rainfall (on average) for more than a decade. We now focus on the North African monsoon region, which strongly affects the Nile River summer flooding. Fig. 10 shows the three consecutive years of monsoon season (JJAS) rainfall over equatorial and northern Africa (encompassing the Nile River basin) after each eruption. The African monsoon exhibited a strong response with notably reduced rainfall of more than 1 mm/day following the (mid-June) E1 tropical eruption, starting in the year of the eruption year itself (Year 0, fig 10), affecting both the White Nile watershed in the south of the basin and the Blue Nile and Atbara River watersheds further north and east in the Ethiopian Highlands. Reduced precipitation was also observed following each of the also mid-June extratropical eruptions (E2-E4) during the eruption years was more spatially constrained (and particularly for E2, less severe). This result is perhaps unsurprising as the estimated SO_2 output of E2-E4 is only approximately 1/3 that of the tropical eruption, ...1. Nonetheless, in each case the Blue Nile and Atbara River headwaters in the Ethiopian highlands were observed to experience a statistically significant decrease, with important implications for the summer flood in Egypt, which depends for approximately 80% of its floodwater on rainfall here (Melesse et al., 2011). For E2-E4, this response was seen to intensify in the first full post-eruption year (Year 1, fig 10), persisting into the second full post-eruption year (Year 2, fig 10), while for E1 the response contracted geographically to resemble the response in Years 1-2, resembling that seen after E2-E4 (fig 9, fig 10). These results are indicative of an effective suppression of the African monsoon following tropical and northern hemispheric extratropical volcanic eruptions, a finding consistent with previous studies (e.g., Colose et al., 2016; Oman et al., 2006; Haywood et al., 2013; Jacobson et al., 2020; Manning et al., 2017).

[20]




Fig 10. Ensemble mean rainfall difference from the climatological control for each of the first three monsoon seasons (JJAS; rows) after each eruption (columns) over equatorial and North Africa. The blue boundary line shows the present-day Nile River basin, broadly similar to the river extent approximately 2.5ka years ago. The red stippling indicates regions over which change in rainfall is not statistically significant at a 95% confidence level.

Spatial patterns in total cloud cover (Fig S7) for the three consecutive post-eruption monsoon seasons showed a decrease of up to 10% over East Africa and the adjacent Indian Ocean region. This is consistent with the above-reported negative rainfall anomalies (>1 mm/day) over North African land regions, especially over the watershed of the Nile River basin, again suggesting a notable weakening of the summer monsoon (Graf, 1992, Oman et al., 2005). Positive total cloud cover anomalies also coincided with regions having a positive rainfall response (e.g., Mediterranean, Middle East).

We also analyzed the mass of total annual water flow averaged over the Nile River basin (blue line, Fig 11) as representative of Nile flooding and discharge at the river's mouth to summarize the volcanic impacts on Nile flooding. Table 2 presents the percentage annual deficit (-) or excess (+) of water flow in

- Deleted: which is
- Deleted: the
- Deleted: of
- Deleted: Fig
- Deleted: show
- Deleted: cloud cover
- Deleted: These spatial patterns are
- Deleted: and
- Deleted: suggest
- Deleted: strong
- Deleted: anomalies of
- Deleted: coincide
- Deleted: observed as
- Deleted: and
- Deleted: being
- Deleted: river
- Deleted: at an annual scale
- Deleted: thus

the Nile River basin, after each eruption along with the variability (one standard deviation) observed across the model ensemble members, relative to the 100 years climatological mean (base climate). E1 had a strong impact (>30% deficit) on water mass over the Nile catchment during the eruption year and first full post-eruption year (Years 0 and 1, table 2), with a more moderate decrease of ~13% during the second full post-eruption year (Year 2, table 2). The first extratropical eruption (E1) showed a minor decrease in the eruption year (Year 0, table 2) that was not statistically significant. The following two full post-eruption years reversed this pattern to exhibit a modest (though not statistically significant) increase. Extratropical eruptions E3 and E4 showed a more consistent decrease. This was on the order of ~-5% in the eruption year (Year 0, table 2) for both, becoming notably greater in the first full post-eruption year (Year 1, table 2), being ~-18% for E3 and ~-12% for E4. This decrease persisted into the second full post-eruption year (Year 2, table 2) for E4 (~-12%), but fell back in line with the 100 years climatological mean for E3 (though this mean exhibited the highest variance among ensemble members (table 2)). Several individual ensemble members have simulated the change in river flow at the 95% confidence levels ($1.95 \cdot \sigma_{\text{ctrl}}$; σ denotes standard deviation) for a few years when compared against the variability for the control period.

Table 2. Annual mean change (%) and standard deviation in water mass flow over the Nile River catchment for 3 consecutive years after each eruption. Control run variability (interannual standard deviation about the decadal mean, σ_{ctrl}) for Nile basin river flow is 25.2%.

	E1(Tr, 22.5 Tg) Change /Std	E2(NHE, 6.5 Tg) Change /Std	E3(NHE, 7.2 Tg) Change /Std	E4(NHE, 7.5 Tg) Change /Std
Year 0 (eruption year, mid-June)	-28.7±39.9	-3.02±22.5	-4.9±35.2	-4.7±29.6
Year 1	-37.8±22.5	2.5±36.7	-18.1±28.9	-11.7±29.9
Year 2	-13.4±32.2	10.7±39.9	0.9±47.8	-12.1±28.0

- Deleted: on an annual basis
- Deleted: The tropical eruption (
- Deleted:)
- Deleted: the annual
- Deleted: i.e.,
- Deleted: i.e.,
- Deleted: but this
- Deleted: deemed
- Deleted: increase, but this was also
- Deleted: deemed
- Deleted: . The next two extratropical
- Deleted: (
- Deleted:) instead
- Deleted: response in the form of a
- Deleted: For E3 and E4 this decrease
- Deleted: ~
- Deleted: i.e.,
- Deleted: and was
- Deleted: i.e.,
- Deleted:) for both E3 and E4 (
- Deleted: %, respectively). The
- Deleted: i.e.,
- Deleted: (--
- Deleted: %)
- Deleted: effectively
- Deleted: although
- Deleted: annual whole-basin
- Deleted: change does exhibit
- Deleted: observed
- Formatted: German

610 The spatial patterning of response across a basin as complex as the Nile is a critical consideration (fig 11). After E1, the above-described rainfall suppression is associated with a drastic reduction in annual river flow observed over effectively the entire river basin, with a simulated decrease of approximately 30, 40 and 15 km³ relative to the 100 years climatological mean (~104 km³) for Years 0 to 2, respectively. After E2, total annual river flow in Year 0 slightly increased (table 2), although this response was not statistically significant, and in Year 1 exhibited a marked contrast between (particularly) the southern (greater flow) and northern (lesser flow) parts of the basin, before a more consistent increased flow was observed in Year 2. The contrast between a reduced flow over (broadly) the northern part of the basin versus increased flow over the southern part was then observed consistently for all post eruption years for E3 and E4 (fig 11). This contrast may arise in part as a function of the size and complexity of the Nile basin (and the markedly different geographical location of rainfall supplying the White Nile to the south and Blue Nile and Atbara River to the northeast), combined with the asymmetrical (northern hemispherically biased) aerosol loading after extratropical eruptions, suppressing the northward boreal summer migration of the ITCZ and associated rain-bearing monsoon winds (as discussed earlier). As these winds are the primary driver of summer rainfall over the Ethiopian highlands, summer flooding of the Blue Nile and Atbara River into the north of the basin and Egypt would be diminished, while water flow down the White Nile (fed by rainfall over the equatorial lakes) would be potentially enhanced by the failure of the ITCZ to migrate northward beyond this region.

- Deleted: further
- Deleted: Fig
- Deleted: the tropical eruption (
- Deleted:),
- Deleted: can be
- Deleted: per year
- Deleted: the second eruption (
- Deleted:), the
- Deleted: was observed to
- Deleted: increase
- Deleted: shown in fig 11
- Deleted: both eruptions
- Deleted: . We can hypothesize that this
- Deleted: arises
- Deleted: large
- Deleted: loading of sulfate aerosols in the higher latitudes of the
- Deleted: hemisphere
- Deleted: . This may lead to a post-eruption scenario in which
- Deleted: were suppressed
- Deleted: Given that
- Deleted: the





Fig 11. Annual river flow anomaly (km^3/year) relative to the control climatology for 3 consecutive years after each eruption (columns) over North Africa. Other details as per fig 10.

To summarize the hydroclimatic impact of this volcanic quartet on the Nile River basin, fig 12 (top panel) shows that the northern hemisphere experienced a substantial cooling of $\sim 2.5^\circ \text{C}$ (1.0°C greater than the global average response) with a lower spread among ensembles after the first eruption (E1). The subsequent eruptions (E2, E3, and E4), reoccurring at equal temporal intervals, maintained a cooling of $\sim 1.75^\circ \text{C}$ for at least a decade. The monthly mean rainfall anomaly over the Nile basin was observed as a considerable decrease varying between ~ 1.0 and 1.5 mm/day during the monsoon seasons (JJAS) for each post-eruption year (fig 12, middle panel). The impact of decreased rainfall over this region is strongly evident after the tropical eruption E1 in discharge at the Nile mouth (grid box centered at 29.0N , 31.25E) in the Nile delta region of Egypt (fig 12, bottom panel). Our modelling showed here a mean deficit beginning in the eruption year (Year 0) and peaking at a reduction of more than 50% during the first full post-eruption year (Year 1), effectively requiring 2 further years to recover. There was no persistent negative discharge anomaly evident after E2, although individual ensemble variability around the mean

Deleted: the
Deleted: African continent.
Deleted: are same
Deleted: Fig
Deleted: these four sequential
Deleted: eruptions
Deleted: Fig
Deleted: then
Deleted: anomaly of
Deleted: the years following
Deleted: (Figure
Deleted: terms of Nile River
Deleted: river
Deleted: Fig
Deleted: shows
Deleted: that begins
Deleted: year of the
Deleted: (i.e.,
Deleted: peaks
Deleted: of water discharge
Deleted: i.e.,
Deleted:)
Deleted: is
Deleted: the second eruption (
Deleted:),


690 is quite high. By contrast, a deficit began in the year of E3 (Year 0) that persisted throughout the first full post-eruption year (Year 1) and into the start of the second full post-eruption year (Year 2). This deficit peaked in Year 2 at just less than 50%. A similar response was observed after E4, with a persistent negative anomaly starting in the first full post-eruption year (Year 1), continuing throughout Year 2 and into the start of Year 3. This deficit also peaked in Year 1 (at approximately 30%).

Spatially Averaged Anomaly for ModelE diagnostics



695
1700 Fig 12: Monthly time series of individual ensemble and mean of surface temperature response (°C) averaged over northern hemisphere (NH) (top panel), rainfall change (mm/day) for the model's spatial box representing the Nile River watershed (Latitude: (5N, 18N), Longitude: (30E, 42E)) (middle panel) and Nile River discharge anomaly (%) at the delta region (grid box centered at 29.0N, 31.25E). For each panel, the darker solid (thick) line shows the multi-ensemble mean, individual member (thin line), and

Deleted: here
Deleted: is observed to begin
Deleted: the third eruption (
Deleted: , i.e.,
Deleted: then persists
Deleted: i.e.,
Deleted: i.e.,
Deleted: peaks
Deleted: is
Deleted: the fourth (
Deleted:) eruption
Deleted: i.e.,
Deleted: peaks



Deleted:

the color envelope shows the associated variability ($\pm\sigma$; Standard deviation). The annual cycle of climate variability of the control run is shown as $2\sigma_{\text{ctrl}}$ lines (red solid line) along the x-axis for all three variables. The vertical dotted green line shows when each eruption happened.

720 It is evident that the mean surface temperature response in the northern hemisphere is significant at the control period's $2\sigma_{\text{ctrl}}$ level (95% significance). However, while rainfall and river discharge responses are not significant at the $2\sigma_{\text{ctrl}}$ levels, several individual members do show significance at $2\sigma_{\text{ctrl}}$ as well. This signifies the important influence of the model's internal variability in representing the regional hydrological response to volcanic eruptions. However, the statistical significance of the rainfall and

725 discharge response may be sensitive to the relatively coarse resolution of the GISS ModelE, as well as the boundaries chosen to model the Nile basin and its headwaters. We thus investigated the post-volcanic change in river flow for the southern (White Nile-dominated) and northern (Blue Nile and Atbara river-dominated) parts of the basin by dividing it at 10° N (Fig 13). Annual mean river flow change for the south (blue lines) and north (red lines) of the Nile basin were in broad agreement with a negative flow

1730 anomaly after eruption E1. This was most notable in the eruption year and the first year following, with the 95th percentile envelopes (dotted lines) deemed significant at the 95% confidence level for both these years (i.e., crossing the dashed lines parallel to the x-axis (fig 13)). In contrast, the mean north and south responses disagreed, including in the sign of the observed changes, after the extratropical eruptions (E2, E3 & E4). More specifically, while the mean flow anomalies in the year of E2 were unremarkable, with

735 little north-south contrast, a more notable divergence was observed in the first full year following, with a positive flow anomaly in the south and negative in the north. In the year of E3, flow in the south showed no notable anomaly, while flow in the north was marginally negative. This distinction became more marked in the first full year following, mainly due to a larger negative anomaly in the north. In the year of E4, a negative anomaly was again observed in the north, persisting for three post-eruption years, and

1740 contrasting with positive or unremarkable anomalies in the south.

These results are consistent with our earlier-described results (e.g., spatial rainfall variability over the Nile River basin, as per figs 10 and 11) and proposed mechanisms, alongside expectations from the literature (e.g., Manning et al., 2017). Thus, tropical eruptions (like E1) may produce a more consistent (negative) north-south flow response due to their more even interhemispheric aerosol burden and

Deleted: $1\sigma_{\text{ctrl}}$ (red dashed line) and

Formatted: Line spacing: Multiple 1.15 li

Formatted: Font color: Text 1, English (UK)

Deleted: $1\sigma_{\text{ctrl}}$ and $2\sigma_{\text{ctrl}}$ levels.

Formatted: Font color: Text 1, English (UK)

Formatted: Font color: Text 1

Deleted: $1\sigma_{\text{ctrl}}$ level, they fall within the $2\sigma_{\text{ctrl}}$

Deleted: although a few

Deleted: $2\sigma_{\text{ctrl}}$

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1, English (US)

Deleted: dearth in the modeling Nile River basin at a

Deleted: In particular, given the complexity of the Nile's hydrology and disparate sources of discharge for the White and Blue Niles.

Deleted: Fig

Deleted:).

Deleted: and showed

Deleted: Figs.

Deleted: result in

associated radiative impact. Extratropical NH eruptions (like E2-E4) that can result in a more asymmetric hemispheric aerosol burden may, by contrast, introduce contrasting flow anomalies by suppressing the northward migration of the ITCZ, negatively impacting flow in the Blue Nile and Atbara rivers by diminishing monsoon rainfall in the Ethiopian highlands, while potentially enhancing flow in the White Nile, fed by rainfall over the equatorial lakes.




Fig 13. Annual Nile River flow changes averaged over the northern (red) and southern (blue) parts of the basin (divided at 10° N) for the entire simulation period. The solid lines represent the ensemble mean for each part of the basin; the dotted lines are $\pm 1.95\sigma$, where σ is derived from across all the ensembles, and the horizontal dashed lines parallel to x-axis are the $\pm 1.95\sigma_{ctrl}$ where σ_{ctrl} is the standard deviation across the 100-year control run. Red and blue lines correspond to the northern and southern parts of the Nile basin, respectively.

4. Discussion and Conclusions

Recent years have seen increasing interest in the role of hydroclimatic variability in human history, including by interdisciplinary teams combining evidence and methods across disciplinary divides (e.g., McCormick, 2011, 2019; Manning et al., 2017; Ludlow and Travis, 2019; van Bavel et al., 2019; Degroot et al., 2021; Ljungqvist et al., 2021; [Travis et al., 2022](#); Izdebski et al., [2023](#)). For the pre-modern era, when systematic observations of hydroclimate become scarce, this effort depends increasingly upon natural archives (palaeoclimatic proxies) that track variability at spatial and temporal resolutions sufficiently high to convincingly identify associations with societal phenomena (e.g., subsistence crises, migration, conflict), economic and demographic processes, and major historical events (e.g., “collapse” of empires). Work such as that by PAGES2k Network members offering paleoclimatic reconstructions and data collections (e.g., PAGES 2k Consortium, 2013; PAGES 2k Consortium, 2017) are thus crucial, although here the exclusive focus on the past 2k years (for some proxies an artificial horizon and for others an aspiration), excludes many foundational periods and events in human civilization. This includes the development of advanced ancient societies in Asia, the Near East and Mediterranean that are richly documented and offer considerable potential for the study of socioecological systems.

Important work has still been possible using speleothems, sedimentary and other archives (e.g., Drake, 2012; Schneider and Adali, 2014; Knapp and Manning, 2016; Sołtysiak, 2016), but there is often little direct temporal and/or geographical overlap between these early ancient world regions of rich human documentation and proxies (e.g., tree-ring based) with precision and accuracy at annual-or-better resolutions. A major advance has been the publication of a chronologically precise and accurate bipolar ice-core-based volcanic forcing reconstruction for the past 2,500 years (Sigl et al., 2015; Toohey and Sigl, 2017). The potentially global hydroclimatic impacts of major explosive eruptions makes this record widely geographically relevant, while the repeated incidence of major eruptions that can be identified through sulphate deposition in the polar ice sheets has allowed their use as “tests” of societal vulnerability and response to sudden hydroclimatic shocks in both a statistical manner (e.g., Manning et al., 2017; Gao et al., 2021; Ludlow et al., [2023](#)) and in a complementary qualitative manner as “revelatory crises” (Solway, 1994; Dove, 2014), in which tensions and vulnerabilities in political and economic systems are potentially exposed under pressure from sudden environmental variability (e.g., Ludlow and Crampsie, 2019; Huhtamaa et al., [2022](#); Ludlow et al., [2023](#)).

Deleted: from

Deleted: traditional

Deleted: 2022, Travis et al., 2022

Deleted: , and boasting sufficiently accurate dating,

Deleted: the

Deleted: kingdoms and

Deleted: curated

Deleted: as temporal coverage and sampling depth improves), presently...

Deleted: some of the most

Deleted: history

Deleted: hence

Deleted: recent

Deleted: , however,

Deleted: now be seen in ever greater detail

Deleted: in-press, 2022

Deleted: , inequalities

Deleted: given

Deleted: the

Deleted: of

Deleted: Ludlow et al., in-press, 2022;

Deleted: 2022). The scrutiny that such exposure can bring to prevailing human systems (social, economic, ideological) can spur their potentially rapid transformation, while the related suspension of cultural norms and business-as-usual practices under the “state of exception” that may prevail during “natural” disasters and other crises can facilitate this, though such conditions can also be exploited to further entrench existing power bases (e.g., Dove, 2014; McConnell et al., 2020

1830 For historical eruptions to act as tests or be studied as potential “revelatory” crises, knowledge of their
dating alone is insufficient, particularly given the regional and seasonal variability of volcanic
hydroclimatic impacts, and the sensitivity of these impacts to multiple variables such as the location,
season, chemical composition, and height attained by volcanic ejecta (Robock, 2000; Cole-Dai, 2010;
Ludlow et al., 2013). Even where instrumental or natural archives are available, but especially where
1835 these are thin or absent, climate modelling can provide insights into the expected impacts for particular
regions, seasons and related physical (e.g., riverine) systems (e.g., Toohey et al., 2016; McConnell et al.,
2020; Mackay et al., 2022). This is true for modelling of idealized eruptions, but potentially even more
so for models that produce “historical realizations” based upon actual forcing reconstructions (e.g., Tardif
et al., 2019).

1840 In this context we have presented a modeling effort that explores the impacts of a unique eruption quartet
during the (historically tumultuous) decade 168-158 BCE, with a focus on the Nile River basin. These
target years are intermediate between the mid-Holocene and end of the preindustrial periods, and
representative background climate conditions are necessary to investigate the climatic impact of such a
short-term forcing (Zanchettin et al., 2013). PMIP4 vegetation distributions (linearly interpolated for the
1845 2.5ka period from the mid-Holocene (Otto-Bliesner et al., 2017) to the end of the preindustrial (taken as
1850 1850) for the GISS ModelE2.1 (MATRIX) version (Kelley et al., 2020; Bauer et al., 2020)) were therefore
used to improve GCM simulations without a fully dynamic vegetation implementation (Harrison et al.,
2015). Vegetation-albedo feedbacks due to greater prevalence of arid shrubs/steppe over Africa and of
boreal forests over high latitudes were observed to induce a northward movement of the ITCZ over Africa
(Sahara region) promoting a simulated rainfall increase of the order of 0.5-1.0 mm/day in the region (a
response consistent with theoretical expectations and other estimates (e.g., Otterman, 1975; Charney,
1975; Claussen, 2009; Rachmayini et al., 2015; Pausata et al., 2016)).

The GISS ModelE2.1 simulated a strong shortwave and longwave global radiative forcing of -10 and +3.0
W/m², respectively, following the tropical eruption (E1) and a roughly equal forcing of -3.5 and +1.0
1855 W/m², respectively, for each of the 3 extratropical eruptions (E2-E4). The peak net radiative volcanic
forcing was calculated at -7.5 W/m² and -2.5 W/m² for the tropical and extratropical eruptions,
respectively. The model calculated a global AOD at 550 nm of 0.22 and 0.1 after the tropical and

Deleted: climatic

Deleted: .

Deleted: particular

Deleted:)

Deleted: a

Deleted: thus

Deleted: such

Deleted: is

Deleted: Pausata et al., 2016;

Deleted:).

extratropical eruptions, respectively, and estimated a peak cooling of ~1.5 °C almost 12-months after the first eruption (E1), with the three consecutive eruptions then sustaining a surface cooling of about 1.0 °C for almost all of the 15 years modelled. The first eruption (E1) was 30% larger than Pinatubo and the GISS ModelE2.1 simulated proportionally stronger radiative impacts as compared to Pinatubo (for which see: Hansen et al., 1992; Robock and Mao, 1994; Parker et al., 1996; McCormick et al., 1995; Stenchikov, 2015). A detailed analysis of the impacts of volcanic aerosols on the chemical composition of the stratosphere was not part of this study.

The global hydrological cycle responded vigorously to the volcanically induced surface cooling in the GISS ModelE2.1, with a >1.0 mm/day decrease in rainfall observed over African, Indian, and Chinese regions during the summer monsoon season consecutively for 3 years after E1 (tropical) and for 2 years after each of the eruptions E2-E4 (extratropical northern hemispheric). Statistically significant decreases in rainfall over the major tropical northern hemisphere rain belt was also calculated by the model, as well as more broadly over higher latitudes for this hemisphere. Some smaller regions of positive rainfall anomalies were, however, simulated over the northern hemisphere mid-latitudes (land and ocean) around 30° N. These patterns of hydrological response are consistent with studies reporting changes in rainfall and large-scale atmospheric circulations (such as Hadley cell weakening) (e.g., Robock and Liu, 1994; Gillett et al., 2004; Trenberth and Dai, 2007; Crowley et al., 2008; Fischer et al., 2008; Joseph and Zeng, 2011; Timmreck, 2012; Iles et al., 2012; Haywood et al., 2013; Liu et al., 2016). However, we note that particularly on smaller spatial scales, as examined here, the variability in the modelled response as observed across our individual ensemble members may reduce the representativeness of the mean. The notable variability on display across our individual ensemble members, even to the quite substantial forcing represented by E1, also suggests that hydroclimatic responses on local to regional scales may depart from broader regional or hemispheric averages even after quite large volcanic forcings.

For the equatorial and northern African landmass specifically, the GISS ModelE2.1 produced a notable suppression of monsoon (JJAS) rainfall for all eruptions, E1-E4. The onset of this response can be observed in the JJAS season beginning with each eruption year itself, though the timing of the peak intensity and/or greatest spatial extent of this suppression varied (e.g., for E1 the greatest extent and peak intensity occurred for JJAS in Year 0, while for E2-E3 the peak intensity and greatest extent occurred in

Deleted: of simulations.

Deleted: details of

Deleted: ,

Deleted: responds

Deleted: greater than

Deleted: observed

Deleted: the

Deleted: eruption

Deleted: both

Deleted: cycle

Deleted: previous

Deleted: 2016).

Deleted: between eruptions

Year 1, and for E4 in Year 0). The suppression centered (for all eruptions and each plotted post-eruption
1910 JJAS season, fig 10) around latitudes 10-15°N, where it ran in an east-west band that in some years was
effectively contiguous across the continent (approx. 16°W to 52°E). There was, however, a tendency for
this response to be more marked and long-lived (into JJAS of Year 2, fig 10) in the central and eastern
portions of this range, where it was statistically significant and could surpass 1 mm/day (up to 30-40% of
climatology for control period).

1915 Importantly, the regions of the most rapid onset, greatest persistence and intensity of response included
Lake Tana (12°0'N 37°15'E) and the Ethiopian highlands that comprise the headwaters of the Blue Nile
and Atbara rivers and which supply most summer floodwater in Egypt (Melesse et al., 2011). This result
is broadly consistent with CMIP5 model runs forced with large twentieth century eruptions (e.g., Iles and
Hegerl, 2014; Manning et al., 2017). Annual river flow for the Nile River basin (fig 11) closely followed
1920 the patterns of decreased JJAS rainfall over the headwater region. Simulated river flow showed a deficit
in the range of 15-40 km³/year up to 3 years following the modelled extratropical northern hemisphere
eruptions. Simulated variability in river discharge also increased 3/4-fold following the extratropical
eruptions, but there is no way to tell which ensemble member best describes the historical conditions that
actually occurred following the eruptions between 168 and 158 BCE. The large variability between
1925 ensembles and the statistical significance of the drying implies that Nile summer flooding might have
been considerably lower than the simulated mean anomaly.

What is certain is that the scale and persistence of the hydroclimatic impacts implied by our modelling
for the 168-158 BCE eruption quartet supports, to begin, inferences of poor Nile flooding in 166 and 161
BCE from scattered references in surviving written sources (Bonneau, 1971). These also identify 169
1930 BCE as potentially experiencing poor flooding. This suggests (assuming sufficiently accurate ice-core
dating (Sigl et al., 2015)) that the eruption quartet (the impacts of which are now better supported and
characterised by our modelling) may have compounded the stresses arising from this initial shock,
contributing to what is long recognized as a tumultuous decade in Egyptian history. During the 6th Syrian
War, Antiochus IV and his Seleukid army invaded Egypt twice. The first invasion occurred in 170 BCE
1935 and the second, more serious occupation, in 168 BCE. This takeover might indeed have re-shaped
Mediterranean history had it not been averted by Roman diplomatic intervention (Höbl, 2001). Internal

Deleted: is

Deleted: can

Deleted: the vast majority of

Deleted: apparent

Deleted: was

Deleted: seen to increase

Deleted: because of the spatial variability in the rainfall response across ensemble members. There

Deleted: best

Deleted: happened

Deleted: , but the

Deleted: tells us

Deleted: may

Deleted: nonetheless

Deleted: , which

Deleted: may have compounded any societal impacts already arising from this. Indeed, our modelling supports the contention that there is a largely overlooked but significant environmental context to what has long been recognized as a tumultuous decade in Egyptian history. Acknowledging the historical context evolving over the preceding decades, and the resulting political, military, economic and cultural setting through which any hydroclimatic shock will have propagated, is of course also key to achieving a fuller understanding of the human-environmental entanglements in the 160s, as indeed it is for any period or region (White and Pei, 2020). This now more clearly includes the role of explosive volcanism and (relatedly) where along the spectrum of proximate to ultimate causality (as per Gao et al., 2021) any resulting hydroclimatic shocks lay in contributing to the revolts and other societal stressors in evidence. Thus, the increasing dominance of Rome in the eastern Mediterranean, and the growing internal political weakness of the Ptolemaic kingdom and their great rivals the Seleukid empire are writ large in the historical narrative of the second century BCE Mediterranean world, but the extent to which the instability of the major eastern Mediterranean powers was an outcome of the rising power of Rome has been heavily debated. A consensus view is now that these developments were directly coupled, and that eastward Roman expansion was driven not by the exceptional aggressiveness of Rome so much as by a "power transition crisis" in the eastern states around 207-200 BCE that drew Rome in (Eckstein, 2008). ¶ The Ptolemies also began to face notable internal dynastic disputes and broader internal unrest among (at least certain sections of) the populace in the years after the Battle of Raphia in 217 BCE. Despite their success in that battle against the Seleukids at the close of the so-called Fourth Syrian War (Grainger, 2010), this was marked as a turning point for longer-term Ptolemaic fortunes by the Greek historian Polybius (V.107.1-3). The high cost of this war and subsequent continued conflict with the Seleukids (that also ... [21])

Deleted: occurred

Deleted: would

Deleted: commonly referred to as "the Day of Eleusis"

Deleted: , pp.147-8).¶ ... [22]

turmoil [continued](#) in Egypt in the 160s and 150s BCE, affecting both the capital Alexandria and the countryside. Surviving sources refer, for example, to “bad times and [having](#) been driven to every extremity owing to the price of wheat” in 168 BCE (*UPZ* 1 59; Bagnall and Derow, 2004, pp. 281-82), and it is known that by the middle of the decade an Egypt-wide agricultural crisis, described as a “calamity” was underway, [droving](#) Ptolemaic officials to near panic (*UPZ* 1 110, 165-164 BCE). Manning et al. (2017) have identified dates of probable revolt onset in Ptolemaic history, with such onset dates identified in 168 BCE and 156 BCE, both coinciding closely with the dates of our eruption quartet. A study of the longevity and geography of these revolts is now of considerable interest. The surviving texts do not tell a complete story but scattered written references that [imply](#) a long persistence of revolt throughout the decade (Veisse 2004) are now rendered more explicable given the modelled persistence of reduced temperatures and suppressed Nile summer flooding for more than a decade following the 168 BCE tropical eruption and the three following extratropical eruptions. [More precisely delineating the political, military, economic and cultural pathways through which any volcanically induced hydroclimatic shock will have propagated is the subject of ongoing efforts to achieve a fuller understanding of the human-environmental entanglements of the 160s. Relatedly, open questions remain as to where along the spectrum from proximate to ultimate causality \(as per Gao et al., 2021\) or necessary and sufficient conditions \(as per White and Pei, 2020\) hydroclimatic shocks lay in contributing to the revolts and other societal stresses that feature so prominently in Ptolemaic history.](#)

Deleted: that drove

Deleted:)

Deleted: .

Deleted: already

Deleted: also

Deleted: yet

Deleted: suggest

Deleted: , including for the years 168-157 BCE

Deleted: , pp. 78-79),

Deleted: credible and

Deleted: NH eruptions

Deleted: ¶

Code/Data availability

Details to support the results in the manuscript is available as supplementary information is provided with the manuscript. Raw data and codes are available on request to author.

Acknowledgements

RS, KT, FL and JM acknowledge support by the National Science Foundation under Grant No. ICER-1824770. ANL acknowledges institutional support from NASA GISS. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Center for Climate Simulation (NCCS) at Goddard Space Flight Center. The authors thank for their input through multiple

discussions the project members and collaborators of the ICER-1824770 project, ‘Volcanism, Hydrology
2080 and Social Conflict: Lessons from Hellenistic and Roman-Era Egypt and Mesopotamia’. FL
acknowledges support from the Trinity Center for Environmental Humanities. This paper benefited from
discussion facilitated by the ‘Volcanic Impacts on Climate and Society’ (VICS) Working Group of
PAGES.

2085 **Author’s contributions**

FL and JM identified the study period in consultation with the other authors. RS, KT and ANL designed
the model simulations. RS performed the simulations, created the figures in close collaboration with KT,
ANL, FL and JM. RS wrote the first draft of the manuscript and [RS, and FL](#) led the writing of subsequent
drafts. All authors contributed to the interpretation of results and the drafting of the text.

2090

Competing interests

The authors declare no competing interests.

Short Summary

2095 This study is a modelling effort to investigate hydroclimate impacts for the Nile River basin induced by
a volcanic “quartet” of four closely spaced eruptions in ice-core volcanic chronology for the decade 168-
158 BCE in a context to ancient Egyptian history. The NASA GISS ModelE simulated a strong response
in sustained temperature reduction and suppressed monsoon rainfall over East Africa following these
eruptions, leading to a deficit in Egypt's agriculturally critical Nile summer flooding.

2100 **References**

Aquila, V., Oman, L. D., Stolarski, R. S., Colarco, P. R., and Newman, P. A.: Dispersion of the volcanic
sulfate cloud from a Mount Pinatubo-like eruption, 117, <https://doi.org/10.1029/2011JD016968>, 2012.
Aquila, V., Oman, L. D., Stolarski, R., Douglass, A. R., and Newman, P. A.: The Response of Ozone and
Nitrogen Dioxide to the Eruption of Mt. Pinatubo at Southern and Northern Midlatitudes, 70, 894–900,
2105 <https://doi.org/10.1175/JAS-D-12-0143.1>, 2013.

- Bauer, S. E., Wright, D. L., Koch, D., Lewis, E. R., and McGraw, R.: MATRIX (Multiconfiguration Aerosol TRacker of mIXing state): an aerosol microphysical module for global atmospheric models, 33, 2008.
- Bauer, S. E., Ault, A., and Prather, K. A.: Evaluation of aerosol mixing state classes in the GISS modelE-
 2110 MATRIX climate model using single-particle mass spectrometry measurements, 118, 9834–9844,
<https://doi.org/10.1002/jgrd.50700>, 2013.
- Bauer, S. E., Tsigaridis, K., Faluvegi, G., Kelley, M., Lo, K. K., Miller, R. L., Nazarenko, L., Schmidt,
 G. A., and Wu, J.: Historical (1850–2014) Aerosol Evolution and Role on Climate Forcing Using the
 GISS ModelE2.1 Contribution to CMIP6, 12, e2019MS001978, <https://doi.org/10.1029/2019MS001978>,
 2115 2020.
- Bell, B. 1975. ‘Climate and the History of Egypt: The Middle Kingdom’, *American Journal of
 Archaeology* 79(3): 223-69.
- Berhane, F., Zaitchik, B., and Dezfuli, A.: Subseasonal Analysis of Precipitation Variability in the Blue
 Nile River Basin, 27, 325–344, <https://doi.org/10.1175/JCLI-D-13-00094.1>, 2014.
- 2120 Blouin, J.: Defining and measuring tax planning aggressiveness, 67, 875–899,
<https://doi.org/10.17310/ntj.2014.4.06>, 2014.
- Bluth, G. J. S., Doiron, S. D., Schnetzler, C. C., Krueger, A. J., and Walter, L. S.: Global tracking of the
 SO₂ clouds from the June, 1991 Mount Pinatubo eruptions, 19, 151–154,
<https://doi.org/10.1029/91GL02792>, 1992.
- 2125 Braconnot, P., Joussaume, S., Marti, O., and de Noblet, N.: Synergistic feedbacks from ocean and
 vegetation on the African Monsoon response to Mid-Holocene insolation, 26, 2481–2484,
<https://doi.org/10.1029/1999GL006047>, 1999.
- Brenna, H., Kutterolf, S., Mills, M. J., and Krüger, K.: The potential impacts of a sulfur- and halogen-
 rich supereruption such as Los Chocoyos on the atmosphere and climate, 20, 6521–6539,
 2130 <https://doi.org/10.5194/acp-20-6521-2020>, 2020.
- Broccoli, A. J., Dahl, K. A., and Stouffer, R. J.: Response of the ITCZ to Northern Hemisphere cooling,
 33, <https://doi.org/10.1029/2005GL024546>, 2006.

- Butzer, K. W.: Early hydraulic civilization in Egypt: a study in cultural ecology, The University of Chicago Press, Chicago London, 134 pp., 1976.
- 2135 Butzer, K.W. 1984. 'Long-term Nile flood variation and political discontinuities in pharaonic Egypt', in: J. Desmond Clark and S.A. Brandt (eds.), *From Hunters to Farmers: The Causes and Consequences of Food Production in Africa*, Berkeley, 102-12.
- Charney, J. G.: Dynamics of deserts and drought in the Sahel, 101, 193–202, <https://doi.org/10.1002/qj.49710142802>, 1975.
- 2140 Christiansen, B.: Volcanic Eruptions, Large-Scale Modes in the Northern Hemisphere, and the El Niño–Southern Oscillation, 21, 910–922, <https://doi.org/10.1175/2007JCLI11657.1>, 2008.
- Chiang, J. C. H. and Bitz, C. M.: Influence of high latitude ice cover on the marine Intertropical Convergence Zone, *Climate Dynamics*, 25, 477–496, <https://doi.org/10.1007/s00382-005-0040-5>, 2005.
- Campbell, B. M. S. and Ludlow, F. (2020) "Climate, Disease and Society in Late-Medieval Ireland", *Proceedings of the Royal Irish Academy*, 120C, 159-252.
- 2145 Claussen*, M.: Late Quaternary vegetation-climate feedbacks, 5, 203–216, <https://doi.org/10.5194/cp-5-203-2009>, 2009.
- Claussen, M., Brovkin, V., Ganopolski, A., Kubatzki, C., and Petoukhov, V.: Climate Change in Northern Africa: The Past is Not the Future, *Climatic Change*, 57, 99–118, <https://doi.org/10.1023/A:1022115604225>, 2003.
- 2150 Cole- Dai J., 'Volcanoes and Climate', *WIREs Climate Change*, 1, 824– 39, 2010.
- Colose, C. M., LeGrande, A. N., and Vuille, M.: Hemispherically asymmetric volcanic forcing of tropical hydroclimate during the last millennium, *Earth Syst. Dynam.*, 7, 681–696, <https://doi.org/10.5194/esd-7-681-2016>, 2016.
- 2155 Crowley, T., GA, Z., Vinther, B., Udisti, R., Kreutzs, K., Cole-Dai, J., and Castellano, E.: Volcanism and the Little Ice Age, *PAGES Newsl.*, 16, 22–23, <https://doi.org/10.1029/2002GL0166335>, 2008.
- D'Arrigo, R., Seager, R., Smerdon, J. E., LeGrande, A. N., and Cook, E. R.: The anomalous winter of 1783–1784: Was the Laki eruption or an analog of the 2009–2010 winter to blame?, 38, <https://doi.org/10.1029/2011GL046696>, 2011.

- 2160 Dogar, M. M.: Impact of Tropical Volcanic Eruptions on Hadley Circulation Using a High-Resolution AGCM, 114, 1284, <https://doi.org/10.18520/cs/v114/i06/1284-1294>, 2018.
- Dove M. R.: *Anthropology of Climate Change* (Chichester, Wiley & Sons, 2014).
- Degroot, D., Anchukaitis, K., Bauch, M., Burnham, J., Carnegie, F., Cui, J., de Luna, K., Guzowski, P., Hambrecht, G., Huhtamaa, H., Izdebski, A., Kleemann, K., Moesswilde, E., Neupane, N., Newfield, T.,
- 2165 Pei, Q., Xoplaki, E., and Zappia, N.: Towards a rigorous understanding of societal responses to climate change. *Nature*, 591, 539–550, doi:10.1038/s41586-021-03190-2, 2021.
- Drake, B.L.: ‘The Influence of Climatic Change on the Late Bronze Age Collapse and the Greek Dark Ages’, *Journal of Archaeological Science* 39, pp.1862-1870, 2012
- A. Eckstein, Rome enters the Greek East: From Anarchy to Hierarchy in the Hellenistic Mediterranean, 230-170 BC. Blackwell, 2008.
- 2170 English, J. M., Toon, O. B., and Mills, M. J.: Microphysical simulations of large volcanic eruptions: Pinatubo and Toba, 118, 1880–1895, <https://doi.org/10.1002/jgrd.50196>, 2013.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 2175 Fischer, E. M., Luterbacher, J., Zorita, E., Tett, S. F. B., Casty, C., and Wanner, H.: European climate response to tropical volcanic eruptions over the last half millennium, 34, <https://doi.org/10.1029/2006GL027992>, 2007.
- Fujiwara, M., Hibino, T., Mehta, S. K., Gray, L., Mitchell, D., and Anstey, J.: Global temperature response to the major volcanic eruptions in multiple reanalysis data sets, *Atmos. Chem. Phys.*, 15, 13507–13518, <https://doi.org/10.5194/acp-15-13507-2015>, 2015.
- 2180 Gao, F., Morissette, J. T., Wolfe, R. E., Ederer, G., Pedelty, J., Masuoka, E., Myneni, R., Tan, B., and Nightingale, J.: An Algorithm to Produce Temporally and Spatially Continuous MODIS-LAI Time Series, 5, 60–64, <https://doi.org/10.1109/LGRS.2007.907971>, 2008.
- 2185 Gao, C., Ludlow, F., Matthews, A., Stine, A. R., Robock, A., Pan, Y., Breen, R. and Sigl. M. (2021) “Volcanic Climate Impacts Can Act as Ultimate and Proximate Causes of Chinese Dynastic Collapse”, *Communications Earth & Environment*, 2, Article Number: 234. DOI: 10.1038/s43247-021-00284-7

- Gillett, N. P., Weaver, A. J., Zwiers, F. W., and Wehner, M. F.: Detection of volcanic influence on global precipitation, 31, <https://doi.org/10.1029/2004GL020044>, 2004.
- 2190 Guillet, S., Corona, C., Ludlow, F., Oppenheimer, C., and Stoffel, M.: Climatic and societal impacts of a “forgotten” cluster of volcanic eruptions in 1108-1110 CE, *Sci Rep*, 10, 6715, <https://doi.org/10.1038/s41598-020-63339-3>, 2020.
- Guo, S., Bluth, G. J. S., Rose, W. I., Watson, I. M., and Prata, A. J.: Re-evaluation of SO₂ release of the 15 June 1991 Pinatubo eruption using ultraviolet and infrared satellite sensors, 5, 2195 <https://doi.org/10.1029/2003GC000654>, 2004.
- Gu, G. and Adler, R. F.: Precipitation and Temperature Variations on the Interannual Time Scale: Assessing the Impact of ENSO and Volcanic Eruptions, 24, 2258–2270, <https://doi.org/10.1175/2010JCLI3727.1>, 2011.
- Graf, H.-F.: Arctic radiation deficit and climate variability, *Climate Dynamics*, 7, 19–28, 2200 <https://doi.org/10.1007/BF00204818>, 1992.
- Graft, H.-F., Kirchner, I., Robock, A., and Schult, I.: Pinatubo eruption winter climate effects: model versus observations, *Climate Dynamics*, 9, 81–93, <https://doi.org/10.1007/BF00210011>, 1993.
- Graf, H.-F., Li, Q., and Giorgetta, M. A.: Volcanic effects on climate: revisiting the mechanisms, 7, 4503–4511, <https://doi.org/10.5194/acp-7-4503-2007>, 2007.
- 2205 Grainger, J. *The Syrian Wars* (Brill, 2010).
- Hansen, J. E., Lacis, A. A., Lee, P., and Wang, W.-C.: Climatic Effects of Atmospheric Aerosols, 338, 575–587, <https://doi.org/10.1111/j.1749-6632.1980.tb17151.x>, 1980.
- Hansen, J., Lacis, A., Ruedy, R., and Sato, M.: Potential climate impact of Mount Pinatubo eruption, 19, 215–218, <https://doi.org/10.1029/91GL02788>, 1992.
- 2210 Hassan, F. A.: Nile Floods and Political Disorder in Early Egypt, in: *Third Millennium BC Climate Change and Old World Collapse*, Berlin, Heidelberg, 1–23, https://doi.org/10.1007/978-3-642-60616-8_1, 1997b.
- Hassan, F.A. 1997a. ‘The Dynamics of a Riverine Civilization: A Geoarchaeological Perspective on the Nile Valley, Egypt’, *World Archaeology* 29(1): 51-74.

- 2215 Hassan, F. A.: Extreme Nile floods and famines in Medieval Egypt (AD 930–1500) and their climatic implications, *Quaternary International*, 173–174, 101–112, <https://doi.org/10.1016/j.quaint.2007.06.001>, 2007.
- Harrison, S. P., Bartlein, P. J., Izumi, K., Li, G., Annan, J., Hargreaves, J., Braconnot, P., and Kageyama, M.: Evaluation of CMIP5 palaeo-simulations to improve climate projections, *Nature Clim Change*, 5, 735–743, <https://doi.org/10.1038/nclimate2649>, 2015.
- 2220 Haywood, J. M., Jones, A., Bellouin, N., and Stephenson, D.: Asymmetric forcing from stratospheric aerosols impacts Sahelian rainfall, *Nature Clim Change*, 3, 660–665, <https://doi.org/10.1038/nclimate1857>, 2013.
- Hewitt, C. D. and Mitchell, J. F. B.: A fully coupled GCM simulation of the climate of the mid-Holocene, 25, 361–364, <https://doi.org/10.1029/97GL03721>, 1998.
- 2225 Höbl, G. A History of the Ptolemaic Empire (Routledge, 2001).
- [Hsiang, S.M., Burke, M. \(2014\) Climate, conflict, and social stability: what does the evidence say? Climatic Change, 123, 39–55.](#)
- Huhtamaa, H., Stoffel, M. and Corona, C. (2022). Recession or resilience? Long-range socioeconomic consequences of the 17th-century volcanic eruptions in northern Fennoscandia, *Climate of the Past Discussions*. <https://doi.org/10.5194/cp-2021-147>.
- 2230 [Ide, T. \(2017\). Research methods for exploring the links between climate change and conflict. WIREs Clim. Change, 8: e456. https://doi.org/10.1002/wcc.456.](#)
- Iles, C. E., Hegerl, G. C., Schurer, A. P., and Zhang, X.: The effect of volcanic eruptions on global precipitation, 118, 8770–8786, <https://doi.org/10.1002/jgrd.50678>, 2013.
- 2235 Iles, C. E. and Hegerl, G. C.: The global precipitation response to volcanic eruptions in the CMIP5 models, *Environ. Res. Lett.*, 9, 104012, <https://doi.org/10.1088/1748-9326/9/10/104012>, 2014.
- Ito, G., Romanou, A., Kiang, N. Y., Faluvegi, G., Aleinov, I., Ruedy, R., Russell, G., Lerner, P., Kelley, M., and Lo, K.: Global Carbon Cycle and Climate Feedbacks in the NASA GISS ModelE2.1, 12, e2019MS002030, <https://doi.org/10.1029/2019MS002030>, 2020.
- 2240 Izdebski, A., Bloomfield, K., Eastwood, W. J., Fernandes, R., Fleitmann, D., Guzowski, P., Haldon, J., Ludlow, F., Luterbacher, J., Manning, J. G., Masi, A., Mordechai, L., Newfield, T., Stine, A. R., Senkul,

- C. and Xoplaki, E. (In Press, [2023](#)), “The Emergence of Interdisciplinary Environmental History: Bridging the Gap between the Humanistic and Scientific Approaches to the Late Holocene,” *Annales*, 77 (2), 1-48.
- Jacobson, T. W. P., Yang, W., Vecchi, G. A., and Horowitz, L. W.: Impact of volcanic aerosol hemispheric symmetry on Sahel rainfall, *Clim Dyn*, 55, 1733–1758, <https://doi.org/10.1007/s00382-020-05347-7>, 2020.
- Joseph, R. and Zeng, N.: Seasonally Modulated Tropical Drought Induced by Volcanic Aerosol, 24, 2045–2060, <https://doi.org/10.1175/2009JCLI3170.1>, 2011.
- Jungclauss, J. H., Lorenz, S. J., Timmreck, C., Reick, C. H., Brovkin, V., Six, K., Segschneider, J., Giorgetta, M. A., Crowley, T. J., Pongratz, J., Krivova, N. A., Vieira, L. E., Solanki, S. K., Klocke, D., Botzet, M., Esch, M., Gayler, V., Haak, H., Raddatz, T. J., Roeckner, E., Schnur, R., Widmann, H., Claussen, M., Stevens, B., and Marotzke, J.: Climate and carbon-cycle variability over the last millennium, *Clim. Past*, 6, 723–737, <https://doi.org/10.5194/cp-6-723-2010>, 2010.
- Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Genio, A. D. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezuman, K., Miller, R. L., Murray, L. T., Oinas, V., Orbe, C., García-Pando, C. P., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: GISS-E2.1: Configurations and Climatology, 12, e2019MS002025, <https://doi.org/10.1029/2019MS002025>, 2020.
- Kiang, N. Y.: Description of the NASA GISS vegetation dynamics model Tech. rep. NASA, 2012
- Khodri, M., Izumo, T., Vialard, J., Janicot, S., Cassou, C., Lengaigne, M., Mignot, J., Gastineau, G., Guilyardi, E., Lebas, N., Robock, A., and McPhaden, M. J.: Tropical explosive volcanic eruptions can trigger El Niño by cooling tropical Africa, *Nat Commun*, 8, 778, <https://doi.org/10.1038/s41467-017-00755-6>, 2017.
- Kilian, M., Brinkop, S., and Jöckel, P.: Impact of the eruption of Mt Pinatubo on the chemical composition of the stratosphere, 20, 11697–11715, <https://doi.org/10.5194/acp-20-11697-2020>, 2020.

- Kim, Y., Moorcroft, P. R., Aleinov, I., Puma, M. J., and Kiang, N. Y.: Variability of phenology and fluxes of water and carbon with observed and simulated soil moisture in the Ent Terrestrial Biosphere Model (Ent TBM version 1.0.1.0.0), Biogeosciences, <https://doi.org/10.5194/gmdd-8-5809-2015>, 2015.
- 2275 Kirtman, B., Power, S. B., Adedoyin, A. J., Boer, G. J., Bojariu, R., Camilloni, I., Doblas-Reyes, F., Fiore, A. M., Kimoto, M., Meehl, G., Prather, M., Sarr, A., Schar, C., Sutton, R., van Oldenborgh, G. J., Vecchi, G., and Wang, H.-J.: Chapter 11 - Near-term climate change: Projections and predictability, edited by: IPCC, Cambridge University Press, Cambridge, 2013.
- Klock D: Assessing the uncertainty in climate sensitivity, Report on Earth System Science 95, Max Plank
 2280 Institute of Meteorology ISSN 1614-1199, 87pp, Phd Thesis 2011.
- Knapp, A.B.and Manning, S.W. :‘Crisis in Context: The End of the Late Bronze Age in the Eastern Mediterranean’, *American Journal of Archaeology* 120 pp.99-149, 2016.
- Kostić, S., Stojković, M., Prohaska, S., and Vasović, N.: Modeling of river flow rate as a function of rainfall and temperature using response surface methodology based on historical time series, *Journal of*
 2285 *Hydroinformatics*, 18, 651–665, <https://doi.org/10.2166/hydro.2016.153>, 2016.
- Kutzbach, J., Bonan, G., Foley, J., and Harrison, S. P.: Vegetation and soil feedbacks on the response of the African monsoon to orbital forcing in the early to middle Holocene, *Nature*, 384, 623–626, <https://doi.org/10.1038/384623a0>, 1996.
- Kutzbach, J. E. and Liu, Z.: Response of the African Monsoon to Orbital Forcing and Ocean Feedbacks
 2290 in the Middle Holocene, 278, 440–443, <https://doi.org/10.1126/science.278.5337.440>, 1997.
- Labitzke, K. and McCormick, M. P.: Stratospheric temperature increases due to Pinatubo aerosols, 19, 207–210, <https://doi.org/10.1029/91GL02940>, 1992.
- Lacis, A., Hansen, J., and Sato, M.: Climate forcing by stratospheric aerosols, *Geophys. Res. Lett.*, 19, 1607–1610, <https://doi.org/10.1029/92GL01620>, 1992.
- 2295 Larrasoana, J. C., Roberts, A. P., and Rohling, E. J.: Dynamics of Green Sahara Periods and Their Role in Hominin Evolution, *PLOS ONE*, 8, e76514, <https://doi.org/10.1371/journal.pone.0076514>, 2013.
- LeGrande, A. N., Tsigaridis, K., and Bauer, S. E.: Role of atmospheric chemistry in the climate impacts of stratospheric volcanic injections, *Nature Geosci*, 9, 652–655, <https://doi.org/10.1038/ngeo2771>, 2016.

- Liu, F., Chai, J., Wang, B., Liu, J., Zhang, X., and Wang, Z.: Global monsoon precipitation responses to large volcanic eruptions, *Sci Rep*, 6, 24331, <https://doi.org/10.1038/srep24331>, 2016.
- Ludlow, F., Stine, A. R., Leahy, P., Murphy, E., Mayewski, P., Taylor, D., Killen, J., Baillie, M., Hennessy, M. and Kiely, G. “Medieval Irish Chronicles Reveal Persistent Volcanic Forcing of Severe Winter Cold Events, 431-1649 CE”, *Environmental Research Letters*, 8 (2), L024035, doi:10.1088/1748-9326/8/2/024035, 2013.
- Ludlow, F. & Manning, J. G. in *Climate Change and Ancient Societies in Europe and the Near East: Diversity in Collapse and Resilience* (eds Erdkamp, P., Manning, J. G. and Verboven K.) 301-320 (Palgrave Macmillan, 2021).
- Ludlow, F. & Manning, J. G. in *Revolt and resistance in the Ancient Classical World and the Near East: The crucible of empire* (eds Collins, J. J. & Manning, J. G.) 154–171 (Brill, 2016)
- Ludlow, F. and Travis, C. (2019) “STEAM Approaches to Climate Change, Extreme Weather and Social-Political Conflict”, In: de la Garza, A. & Travis, C. (eds.), *The STEAM Revolution: Transdisciplinary Approaches to Science, Technology, Engineering, Arts, Humanities and Mathematics*. New York: Springer, 33-65.
- Ludlow, F., Kostick, C. and Morris, C. in *The Cambridge World History of Genocide* (eds Ben Kiernan, Tracy Maria Lemos and Tristan Taylor) (Cambridge University Press, In Press, 2023).
- Ludlow, F. and Crampsie, A.: “Climate, Debt and Conflict: Environmental History as a New Direction in Understanding Early Modern Ireland”, In: Sarah Covington, Vincent Carey and Valerie McGowan-Doyle (eds.), *Early Modern Ireland: New Sources, Methods, and Directions*. London: Routledge, 269-300, 2019.
- Ljungqvist, F. C., Seim, A. and Huhtamaa, H.: Climate and Society in European History. *WIRE’s Climate Change*, 12, e691. <https://doi.org/10.1002/wcc.691>.
- Mackay, H., Plunkett, G., Jensen, B., Aubry, T., Corona, C., Mi Kim, W., Toohey, M., Sigl, M., Stoffel, M., Anchukaitis, K., Raible, C., Bolton, M., Manning, J., Newfield, T., di Cosmo, N., Ludlow, F., Kostick, C., Yang, Z., Coyle McClung, L., Amesbury, M., Monteath, A., Hughes, P., Langdon, P., Charman, D., Booth, R., Davies, K., Blundell, A. and Swindles, G. (2022), “The 852/3 CE Mount Churchill Eruption: Examining the Potential Climatic and Societal Impacts and the Timing of the Medieval Climate Anomaly

Deleted: 2022

Formatted: Font color: Auto, English (UK), Pattern: Clear

in the North Atlantic Region”, *Climate of the Past*, 18, 1475-1508. <https://doi.org/10.5194/cp-18-1475-2022>.

2330 Manning, J. G. *Land and Power in Ptolemaic Egypt: The Structure of Land Tenure* (Cambridge University Press, 2003).

Manning, J. G., Ludlow, F., Stine, A. R., Boos, W. R., Sigl, M., and Marlon, J. R.: Volcanic suppression of Nile summer flooding triggers revolt and constrains interstate conflict in ancient Egypt, *Nat Commun*, 8, 900, <https://doi.org/10.1038/s41467-017-00957-y>, 2017.

2335 J. G. Manning, *The Open Sea: The Economic Life of the Ancient Mediterranean World from the Iron Age to the Rise of Rome* (Princeton: Princeton University Press, 2018)

McCormick, M. P., Thomason, L. W., and Trepte, C. R.: Atmospheric effects of the Mt Pinatubo eruption, *Nature*, 373, 399–404, <https://doi.org/10.1038/373399a0>, 1995.

2340 McCormick, M. History's changing climate: Climate science, genomics and the emerging consilient approach to interdisciplinary history,” *Journal of Interdisciplinary History* 42 (2011): 252-73.

McCormick, M. 2013. ‘What climate science, Ausonius, Nile floods, rye, and thatch tell us about the environmental history of the Roman Empire’, in 61-88.

McCormick, M. *Climates of History, Histories of Climate: From History to Archaeoscience*, *Journal of Interdisciplinary History*, 50 (2019): 3–30.

2345 McConnell, J. R., Sigl, M., Plunkett, G., Burke, A., Kim, W. M., Raible, C. C., Wilson, A. I., Manning, J. G., Ludlow, F., Chellman, N. J., Innes, H. M., Yang, Z., Larsen, J. F., Schaefer, J. R., Kipfstuhl, S., Mojtabavi, S., Wilhelms, F., Opel, T., Meyer, H., and Steffensen, J. P.: Extreme climate after massive eruption of Alaska’s Okmok volcano in 43 BCE and effects on the late Roman Republic and Ptolemaic Kingdom, *Proc Natl Acad Sci USA*, 117, 15443–15449, <https://doi.org/10.1073/pnas.2002722117>, 2020.

2350 McGing, B. C. Revolt Egyptian style: Internal opposition to Ptolemaic rule. *Arch. Papyrusforschung* 43, 274–277 (1997).

Mikhail, A. 2015. ‘Ottoman Iceland: A climate history’, *Environmental History* 20: 262–284.

Melesse, A. M. (Ed.): *Nile River Basin*, Springer Netherlands, Dordrecht, <https://doi.org/10.1007/978-94-007-0689-7>, 2011.

Formatted: Font: 12 pt

- 2355 Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestad, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura, and H. Zhang.: Anthropogenic and natural radiative forcing. In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Doschung, A. Nauels, Y. Xia, V. Bex, and P.M.
- 2360 Midgley, Eds. Cambridge University Press, pp. 659-740, doi:10.1017/CBO9781107415324.018, 2013
- Myneni, R. B., Hoffman, S., Knyazikhin, Y., Privette, J. L., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G. R., Lotsch, A., Friedl, M., Morisette, J. T., Votava, P., Nemani, R. R., and Running, S. W.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote Sensing of Environment*, 83, 214–231, [https://doi.org/10.1016/S0034-4257\(02\)00074-3](https://doi.org/10.1016/S0034-4257(02)00074-3), 2002.
- 2365 Oman, L., Robock, A., Stenchikov, G., Schmidt, G. A., and Ruedy, R.: Climatic response to high-latitude volcanic eruptions, 110, <https://doi.org/10.1029/2004JD005487>, 2005.
- Oman, L., Robock, A., Stenchikov, G. L., Thordarson, T., Koch, D., Shindell, D. T., and Gao, C.: Modeling the distribution of the volcanic aerosol cloud from the 1783–1784 Laki eruption, *J. Geophys. Res.*, 111, D12209, <https://doi.org/10.1029/2005JD006899>, 2006.
- 2370 Otterman, J.: Baring High-Albedo Soils by Overgrazing: A Hypothesized Desertification Mechanism, 186, 531–533, <https://doi.org/10.1126/science.186.4163.531>, 1974.
- Otto-Bliesner, B. L., Braconnot, P., Harrison, S. P., Lunt, D. J., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Capron, E., Carlson, A. E., Dutton, A., Fischer, H., Goelzer, H., Govin, A., Haywood, A., Joos, F., LeGrande, A. N., Lipscomb, W. H., Lohmann, G., Mahowald, N., Nehrbass-Ahles, C., Pausata, F. S. R.,
- 2375 Peterschmitt, J.-Y., Phipps, S. J., Renssen, H., and Zhang, Q.: The PMIP4 contribution to CMIP6 – Part 2: Two interglacials, scientific objective and experimental design for Holocene and Last Interglacial simulations, 10, 3979–4003, <https://doi.org/10.5194/gmd-10-3979-2017>, 2017.
- PAGES 2k Consortium, Continental-scale temperature variability during the past two millennia, *Nature Geoscience*, 6, 339-346 2013.
- 2380 PAGES 2k Consortium, A global database for temperature reconstructions of the Common Era, *Scientific Data*, 4:170088 | DOI: 10.1038/sdata.2017.88

PAGES Hydro2k Consortium: Comparing proxy and model estimates of hydroclimate variability and change over the Common Era, 13, 1851–1900, <https://doi.org/10.5194/cp-13-1851-2017>, 2017.

Parker, D. E., Wilson, H., Jones, P. D., Christy, J. R., and Folland, C. K.: The Impact of Mount Pinatubo
 2385 on World-Wide Temperatures, 16, 487–497, [https://doi.org/10.1002/\(SICI\)1097-0088\(199605\)16:5<487::AID-JOC39>3.0.CO;2-J](https://doi.org/10.1002/(SICI)1097-0088(199605)16:5<487::AID-JOC39>3.0.CO;2-J), 1996.

Pausata, F. S. R., Chafik, L., Caballero, R., and Battisti, D. S.: Impacts of high-latitude volcanic eruptions on ENSO and AMOC, *Proc Natl Acad Sci USA*, 112, 13784–13788, <https://doi.org/10.1073/pnas.1509153112>, 2015.

2390 Pausata, F. S. R., Messori, G., and Zhang, Q.: Impacts of dust reduction on the northward expansion of the African monsoon during the Green Sahara period, *Earth and Planetary Science Letters*, 434, 298–307, <https://doi.org/10.1016/j.epsl.2015.11.049>, 2016.

Peterson, L. C., Haug, G. H., Hughen, K. A., and Röhl, U.: Rapid Changes in the Hydrologic Cycle of the Tropical Atlantic During the Last Glacial, <https://doi.org/10.1126/science.290.5498.1947>, 2000.

2395 Pitari, G. and Mancini, E.: Short-term climatic impact of the 1991 volcanic eruption of Mt. Pinatubo and effects on atmospheric tracers, *Nat. Hazards Earth Syst. Sci.*, 2, 91–108, <https://doi.org/10.5194/nhess-2-91-2002>, 2002.

Pitari, G., Cionni, I., Di Genova, G., Visioni, D., Gandolfi, I., and Mancini, E.: Impact of Stratospheric Volcanic Aerosols on Age-of-Air and Transport of Long-Lived Species, 7, 149, <https://doi.org/10.3390/atmos7110149>, 2016.

2400 Rachmayani, R., Prange, M., and Schulz, M.: North African vegetation–precipitation feedback in early and mid-Holocene climate simulations with CCSM3-DGVM, *Clim. Past*, 11, 175–185, <https://doi.org/10.5194/cp-11-175-2015>, 2015. <https://doi.org/10.5194/cp-11-175-2015>, 2015.

Rao, M. P., Cook, B. I., Cook, E. R., D’Arrigo, R. D., Krusic, P. J., Anchukaitis, K. J., LeGrande, A. N.,
 2405 Buckley, B. M., Davi, N. K., Leland, C., and Griffin, K. L.: European and Mediterranean hydroclimate responses to tropical volcanic forcing over the last millennium, 44, 5104–5112, <https://doi.org/10.1002/2017GL073057>, 2017.

Read, W. G., Froidevaux, L., and Waters, J. W.: Microwave limb sounder measurement of stratospheric SO₂ from the Mt. Pinatubo Volcano, 20, 1299–1302, <https://doi.org/10.1029/93GL00831>, 1993.

- Robock, A.: Volcanic eruptions and climate, 38, 191–219, <https://doi.org/10.1029/1998RG000054>, 2000.
- Robock, A. and Mao, J.: Winter warming from large volcanic eruptions, *Geophys. Res. Lett.*, 19, 2405–2408, <https://doi.org/10.1029/92GL02627>, 1992.
- Robock, A. and Liu, Y.: The Volcanic Signal in Goddard Institute for Space Studies Three-Dimensional Model Simulations, 7, 44–55, [https://doi.org/10.1175/1520-0442\(1994\)007<0044:TVSIGI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1994)007<0044:TVSIGI>2.0.CO;2), 1994.
- Robock, A. and Mao, J.: The Volcanic Signal in Surface Temperature Observations, 8, 1086–1103, [https://doi.org/10.1175/1520-0442\(1995\)008<1086:TVSIST>2.0.CO;2](https://doi.org/10.1175/1520-0442(1995)008<1086:TVSIST>2.0.CO;2), 1995.
- Roger S. Bagnall and Peter Derow, *Historical Sources in Translation. The Hellenistic Period*. New Edition, Blackwell, 2004.
- Roller, D. W. *Cleaopatra: A Biography* (Oxford University Press, 2010).
- Sabzevari, A. A., Zarenistanak, M., Tabari, H., and Moghimi, S.: Evaluation of precipitation and river discharge variations over southwestern Iran during recent decades, *J Earth Syst Sci*, 124, 335–352, <https://doi.org/10.1007/s12040-015-0549-x>, 2015.
- Russell, P. B., Livingston, J. M., Pueschel, R. F., Bauman, J. J., Pollack, J. B., Brooks, S. L., Hamill, P., Thomason, L. W., Stowe, L. L., Deshler, T., Dutton, E. G., and Bergstrom, R. W.: Global to microscale evolution of the Pinatubo volcanic aerosol derived from diverse measurements and analyses, 101, 18745–18763, <https://doi.org/10.1029/96JD01162>, 1996.
- Schmidt, A., Carslaw, K. S., Mann, G. W., Wilson, M., Breider, T. J., Pickering, S. J., and Thordarson, T.: The impact of the 1783–1784 AD Laki eruption on global aerosol formation processes and cloud condensation nuclei, 10, 6025–6041, <https://doi.org/10.5194/acp-10-6025-2010>, 2010.
- Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y., Del Genio, A., Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., Lerner, J., Lo, K. K., Miller, R. L., Nazarenko, L., Oinas, V., Perlwitz, J., Perlwitz, J., Rind, D., Romanou, A., Russell, G. L., Sato, M., Shindell, D. T., Stone, P. H., Sun, S., Tausnev, N., Thresher, D., and Yao, M.-S.: Present-Day Atmospheric Simulations Using GISS ModelE: Comparison to In Situ, Satellite, and Reanalysis Data, 19, 153–192, <https://doi.org/10.1175/JCLI3612.1>, 2006.

Schmidt, G. A., Jungclauss, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J., Delaygue, G., Joos, F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J., Shindell, D. T., Solanki, S. K., Steinhilber, F., and Vieira, L. E. A.: Climate forcing reconstructions for use in PMIP simulations of the last millennium (v1.0), *Geosci. Model Dev.*, 4, 33–45, <https://doi.org/10.5194/gmd-4-33-2011>, 2011.

Schmidt, G. A., Kelley, M., Nazarenko, L., Ruedy, R., Russell, G. L., Aleinov, I., Bauer, M., Bauer, S. E., Bhat, M. K., Bleck, R., Canuto, V., Chen, Y.-H., Cheng, Y., Clune, T. L., Del Genio, A., de Fainchtein, R., Faluvegi, G., Hansen, J. E., Healy, R. J., Kiang, N. Y., Koch, D., Lacis, A. A., LeGrande, A. N., Lerner, J., Lo, K. K., Matthews, E. E., Menon, S., Miller, R. L., Oinas, V., Oloso, A. O., Perlwitz, J. P., Puma, M. J., Putman, W. M., Rind, D., Romanou, A., Sato, M., Shindell, D. T., Sun, S., Syed, R. A., Tausnev, N., Tsigaridis, K., Unger, N., Voulgarakis, A., Yao, M.-S., and Zhang, J.: Configuration and assessment of the GISS ModelE2 contributions to the CMIP5 archive: GISS MODEL-E2 CMIP5 SIMULATIONS, *J. Adv. Model. Earth Syst.*, 6, 141–184, <https://doi.org/10.1002/2013MS000265>, 2014.

Said, R. 1993. *The River Nile: Geology, Hydrology and Utilization*. Oxford.

Schneider, D. P., Ammann, C. M., Otto-Bliesner, B. L., and Kaufman, D. S.: Climate response to large, high-latitude and low-latitude volcanic eruptions in the Community Climate System Model, 114, <https://doi.org/10.1029/2008JD011222>, 2009.

Schneider A. W. and Adali, S.F. : ‘‘No Harvest was Reaped’’: Demographic and Climatic Factors in the Decline of the Neo-Assyrian Empire’, *Climatic Change* 127, pp.435-446, 2014.

Schnetzler, C. C., Krueger, A. J., Bluth, G. S., Sprod, I. E., and Walter, L. S.: Comment on the Paper ‘‘The atmospheric SO₂ budget for Pinatubo derived from NOAA-11 SBUV/2 spectral data’’ by R. D. McPeters, 22, 315–316, <https://doi.org/10.1029/94GL02406>, 1995.

Segele, Z. T., Lamb, P. J., and Leslie, L. M.: Large-scale atmospheric circulation and global sea surface temperature associations with Horn of Africa June–September rainfall, 29, 1075–1100, <https://doi.org/10.1002/joc.1751>, 2009.

Shindell, D. T.: Dynamic winter climate response to large tropical volcanic eruptions since 1600, *J. Geophys. Res.*, 109, D05104, <https://doi.org/10.1029/2003JD004151>, 2004.

Shindell, D. T., Faluvegi, G., Unger, N., Aguilar, E., Schmidt, G. A., Koch, D. M., Bauer, S. E., and Miller, R. L.: Simulations of preindustrial, present-day, and 2100 conditions in the NASA GISS

composition and climate model G-PUCCINI, 6, 4427–4459, <https://doi.org/10.5194/acp-6-4427-2006>, 2006.

Sigl, M., Winstrup, M., McConnell, J. R., Welten, K. C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O. J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D. R., Pilcher, J. R., Salzer, M., Schüpbach, S., Steffensen, J. P., Vinther, B. M., and Woodruff, T. E.: Timing and climate forcing of volcanic eruptions for the past 2,500 years, *Nature*, 523, 543–549, <https://doi.org/10.1038/nature14565>, 2015.

Simard, M., Pinto, N., Fisher, J. B., and Baccini, A.: Mapping forest canopy height globally with spaceborne lidar, 116, <https://doi.org/10.1029/2011JG001708>, 2011.

Singh, R. and AchutaRao, K.: Quantifying uncertainty in twenty-first century climate change over India, *Clim Dyn*, 52, 3905–3928, <https://doi.org/10.1007/s00382-018-4361-6>, 2019.

Singh, R., LeGrande, A. N., and Tsigaridis, K.: Influence of regional anthropogenic changes over Nile region on the climate system during the late Holocene (~2500 years before present), EGU General Assembly 2020, Online, 4–8 May 2020, EGU2020-12338, <https://doi.org/10.5194/egusphere-egu2020-12338>, 2020.

Staunton-Sykes, J., Aubry, T. J., Shin, Y. M., Weber, J., Marshall, L. R., Luke Abraham, N., Archibald, A., and Schmidt, A.: Co-emission of volcanic sulfur and halogens amplifies volcanic effective radiative forcing, 21, 9009–9029, <https://doi.org/10.5194/acp-21-9009-2021>, 2021.

Solway, J. S. : ‘Drought as a Revelatory Crisis: An Exploration of Shifting Entitlements and Hierarchies in the Kalahari, Botswana’, *Development and Change* 25, pp.471-495, 1994.

Stenchikov, G.: Chapter 26 - The Role of Volcanic Activity in Climate and Global Change, in: *Climate Change (Second Edition)*, edited by: Letcher, T. M., Elsevier, Boston, 419–447, <https://doi.org/10.1016/B978-0-444-63524-2.00026-9>, 2016.

Stoffel, M., Corona, C., Ludlow, F., Sigl, M., Huhtamaa, H., Garnier, E., Helama, S., Guillet, S., Crampsie, A., Kleemann, K., Camenisch, C., McConnell, J. and Gao, C. (In Review, 2022), “Climatic, Weather and Socio-Economic Conditions Corresponding with the mid-17th Century Eruption Cluster”, *Climate of the Past*.

- Swingedouw, D., Mignot, J., Ortega, P., Khodri, M., Menegoz, M., Cassou, C., and Hanquiez, V.: Impact of explosive volcanic eruptions on the main climate variability modes, *Global and Planetary Change*, 150, 24–45, <https://doi.org/10.1016/j.gloplacha.2017.01.006>, 2017.
- 2495 Sołtysiak, A. : ‘Drought and the Fall of Assyria: Quite Another Story’, *Climatic Change* 136, 2016, pp.389-394, 2016.
- Tardif, R., Hakim, G.J., Perkins, W.A., Horlick, K.A., Erb, M.P., Emile-Geay, J., Anderson, D.M., Steig, E.J. & Noone, D. (2019). Last Millennium Reanalysis with an Expanded Proxy Database and Seasonal Proxy Modeling, *Climate of the Past*, 15, pp. 1251-1273.
- 2500 Thordarson, T.: Atmospheric and environmental effects of the 1783–1784 Laki eruption: A review and reassessment, *J. Geophys. Res.*, 108, 4011, <https://doi.org/10.1029/2001JD002042>, 2003.
- Tierney, J. E., Pausata, F. S. R., and deMenocal, P. B.: Rainfall regimes of the Green Sahara, <https://doi.org/10.1126/sciadv.1601503>, 2017.
- Timmreck, C., Lorenz, S. J., Crowley, T. J., Kinne, S., Raddatz, T. J., Thomas, M. A., and Jungclaus, J.
- 2505 H.: Limited temperature response to the very large AD 1258 volcanic eruption, *Geophys. Res. Lett.*, 36, L21708, <https://doi.org/10.1029/2009GL040083>, 2009.
- Timmreck, C., Graf, H.-F., Lorenz, S. J., Niemeier, U., Zanchettin, D., Matei, D., Jungclaus, J. H., and Crowley, T. J.: Aerosol size confines climate response to volcanic super-eruptions, 37, <https://doi.org/10.1029/2010GL045464>, 2010.
- 2510 Timmreck, C.: Modeling the climatic effects of large explosive volcanic eruptions, 3, 545–564, <https://doi.org/10.1002/wcc.192>, 2012.
- Tiwari, S., Ramos, R., Pausata, F. S. R., LeGrande, A. N., Griffiths, M. L., Beltrami, H., Chandan, D., de Vernal, A., Litchmore, D., Peltier, R., and Tabor, C. R.: Model performance in simulating the mid-Holocene Green Sahara, EGU General Assembly 2022, Vienna, Austria, 23–27 May 2022, EGU22-3233, <https://doi.org/10.5194/egusphere-egu22-3233>, 2022
- 2515 Toohey, M., Krüger, K., Niemeier, U., and Timmreck, C.: The influence of eruption season on the global aerosol evolution and radiative impact of tropical volcanic eruptions, *Atmos. Chem. Phys.*, 11, 12351–12367, <https://doi.org/10.5194/acp-11-12351-2011>, 2011.

Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500 BCE
 2520 to 1900 CE, 9, 809–831, <https://doi.org/10.5194/essd-9-809-2017>, 2017.

Toohey, M., Krüger, K., Sigl, M., Stordal, F., and Svensen, H.: Climatic and societal impacts of a volcanic
 double event at the dawn of the Middle Ages, *Climatic Change*, 136, 401–412,
<https://doi.org/10.1007/s10584-016-1648-7>, 2016.

Toohey, M., Krüger, K., Schmidt, H., Timmreck, C., Sigl, M., Stoffel, M., and Wilson, R.:
 2525 Disproportionately strong climate forcing from extratropical explosive volcanic eruptions, 12, 100–107,
<https://doi.org/10.1038/s41561-018-0286-2>, 2019.

Trepte, C. R. and Hitchman, M. H.: Tropical stratospheric circulation deduced from satellite aerosol data,
Nature, 355, 626–628, <https://doi.org/10.1038/355626a0>, 1992.

Travis, C., Holm, P., Ludlow, F., Kostick, C., McGovern, R. and Nicholls, J. (2022) “Cowboys, Cod,
 2530 Climate and Conflict: Navigations in the Digital Environmental Humanities”, In: Travis, C., Legg, R.,
 Bergmann, L., Crampsie, A. and Dixon, D. (eds.), *Routledge Handbook of the Digital Environmental
 Humanities*. London: Routledge, 17-39.

Trenberth, K. E. and Dai, A.: Effects of Mount Pinatubo volcanic eruption on the hydrological cycle as
 an analog of geoengineering: PINATUBO AND THE HYDROLOGICAL CYCLE, *Geophys. Res. Lett.*,
 2535 34, <https://doi.org/10.1029/2007GL030524>, 2007. Vashisht, A., Zaitchik, B., and Gnanadesikan, A.:
 ENSO Teleconnection to Eastern African Summer Rainfall in Global Climate Models: Role of the
 Tropical Easterly Jet, 34, 293–312, <https://doi.org/10.1175/JCLI-D-20-0222.1>, 2021.

Vashisht, A., Zaitchik, B., and Gnanadesikan, A.: ENSO Teleconnection to Eastern African Summer
 Rainfall in Global Climate Models: Role of the Tropical Easterly Jet, 34, 293–312,
 2540 <https://doi.org/10.1175/JCLI-D-20-0222.1>, 2021.

van Bavel, B.J.P., Curtis, D.R., Hannaford, M.J., Moatsos, M., Roosen, J., and Soens, T.: Climate and
 society in long-term perspective: Opportunities and pitfalls in the use of historical datasets, *Wires Clim.*
Change, 10(6), e611, doi: 10.1002/wcc.611, 2019.

Vehkamäki, H., Kulmala, M., Napari, I., Lehtinen, K. E. J., Timmreck, C., Noppel, M., and Laaksonen,
 2545 A.: An improved parameterization for sulfuric acid–water nucleation rates for tropospheric and
 stratospheric conditions, 107, AAC 3-1-AAC 3-10, <https://doi.org/10.1029/2002JD002184>, 2002.

- Veisse, A.-E.: Les révoltes égyptiennes: recherches sur les troubles intérieurs en Égypte du règne de Ptolémée III à la conquête romaine, Peeters, Leuven; Paris; Dudley, MA, 2004.
- 2550 [Villmoare B., *The Evolution of Everything: The Patterns and Causes of Big History* \(Cambridge University Press, 2022\)](#)
- Vörösmarty, CJ; Fekete, BM; Tucker, BA (1998): Global River Discharge, 1807-1991, V. 1.1 (RivDIS). Data set. Available on-line [<http://www.daac.ornl.gov>] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
- 2555 Wahl, E. R., Diaz, H. F., Smerdon, J. E., and Ammann, C. M.: Late winter temperature response to large tropical volcanic eruptions in temperate western North America: Relationship to ENSO phases, *Global and Planetary Change*, 122, 238–250, <https://doi.org/10.1016/j.gloplacha.2014.08.005>, 2014.
- Wolfe, E. W., and Hoblitt R. P.: Overview of the eruptions, in *Fire and Mud: Eruptions and lahars of the Moun Pinatubo, Philippines*, edited by Newhall C.G. and Punongbayan, pp. 415-433, Univ of Wash. Press, Seattle.
- 2560 White, S. and Pei, Q. (2020). Attribution of historical societal impacts and adaptations to climate and extreme events: Integrating quantitative and qualitative perspectives. *Past Global Changes Magazine*, 28(2), 44-45.
- Xian, P. and Miller, R. L.: Abrupt Seasonal Migration of the ITCZ into the Summer Hemisphere, 65, 1878–1895, <https://doi.org/10.1175/2007JAS2367.1>, 2008.
- 2565 Zanchettin, D., Bothe, O., Graf, H. F., Lorenz, S. J., Luterbacher, J., Timmreck, C., and Jungclaus, J. H.: Background conditions influence the decadal climate response to strong volcanic eruptions, 118, 4090–4106, <https://doi.org/10.1002/jgrd.50229>, 2013.
- Zanchettin, D., Timmreck, C., Khodri, M., Schmidt, A., Toohey, M., Abe, M., Bekki, S., Cole, J., Fang, S.-W., Feng, W., Hegerl, G., Johnson, B., Lebas, N., LeGrande, A. N., Mann, G. W., Marshall, L., Rieger, 2570 L., Robock, A., Rubinetti, S., Tsigaridis, K., and Weierbach, H.: Effects of forcing differences and initial conditions on inter-model agreement in the VolMIP volc-pinatubo-full experiment, 1–39, <https://doi.org/10.5194/gmd-2021-372>, 2021.

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [1] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [2] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [2] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [2] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [3] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [3] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [3] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [3] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [3] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [4] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [4] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [4] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [4] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [4] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [4] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [5] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [5] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [5] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [5] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [6] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [6] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [6] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [6] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [7] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [7] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [7] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [7] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [7] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 4: [8] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [8] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [8] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [8] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
3:07:00 AM Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 4: [9] Deleted
Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

Page 6: [10] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 6: [11] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 6: [12] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [13] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [13] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [13] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [14] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [14] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

Page 7: [15] Deleted
3:07:00 AM

Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]

11/22/22

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [15] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [15] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [15] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [15] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [15] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [15] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [15] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [16] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [16] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [16] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [16] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [16] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 7: [16] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 7: [16] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [17] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [17] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 15: [18] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 15: [18] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [19] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [19] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [19] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [19] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [19] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted **Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]** **11/22/22**
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22
3:07:00 AM

▼
▲
Page 28: [20] Deleted Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY] 11/22/22

▼

▲

Page 28: [20] Deleted 3:07:00 AM	Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]	11/22/22
-------------------------------------	---	----------

Page 39: [21] Deleted 3:07:00 AM	Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]	11/22/22
-------------------------------------	---	----------

Page 39: [22] Deleted 3:07:00 AM	Singh, Ram (GISS-6110)[TRUSTEES OF COLUMBIA UNIVERSITY]	11/22/22
-------------------------------------	---	----------

Investigating hydroclimatic impacts of the 168-158 BCE volcanic quartet and their relevance to the Nile River basin and Egyptian history

Ram Singh^{1,2}, Kostas Tsigaridis^{1,2}, Allegra N. LeGrande^{2,1}, Francis Ludlow³, Joseph G Manning⁴

¹ Center for Climate Systems Research, Columbia University, New York

² NASA Goddard Institute for Space Studies, New York, NY-10025

³ Department of History, School of Histories and Humanities, Trinity College, Dublin 2, Ireland

⁴ Departments of History and Classics, Yale University, New Haven, CT 06520, USA

Correspondence to: Ram Singh (rs4068@columbia.edu)

Supplementary information

Deleted: S1.1 Introduction (Historical Context)

Important questions remain, however, in particular on the role of hydroclimatic shocks in the longer-term declining stability of the state and its ability to project power across the eastern Mediterranean. Repeated revolt in the third to first centuries BCE speaks to persistent vulnerability, yet despite experiencing the hydroclimatic effects of multiple eruptions (including those with a greater climate forcing potential than has been experienced in the twentieth and twenty-first centuries (Sigl et al., 2015) and multiple such revolts, the dynasty persisted for almost three centuries, simultaneously suggesting a considerable level of resilience. It is conspicuous, however, that the dynasty ultimately ended (with Cleopatra's defeat by Rome at the naval battle of Actium in 31 BCE and her suicide in 30 BCE) in a decade that followed one of the largest explosive eruptions of the last 2,500 years in terms of climate forcing potential (based upon polar ice-core sulfate deposition levels), that of Okmok (Alaska) in early 43 BCE (McConnell et al., 2020). This itself followed a smaller but notable (likely extratropical NH eruption) in 46 BCE (McConnell et al., 2020). Egypt in the 40s BCE had, perhaps unsurprisingly therefore, experienced repeated Nile failure, famine, plague, inflation, administrative corruption, rural depopulation, migration, and land abandonment (Höbl, 2001; Röller, 2010). It is notable though that there is no convincingly documented revolt, perhaps owing to Cleopatra's abilities as a leader and interventions in grain distribution to prevent starvation of the population. The stresses of the 40s BCE may still, however, be credibly posited as weakening Egypt's hand against Rome as it became entangled in the complex political and military developments of this major moment in world history, as Rome transitioned from its republic to imperial form (Manning et al., 2017; McConnell et al., 2020; Ludlow and Manning, 2021). ¶ Here clearly, as at any other time, the societal impact of a given hydroclimatic shock will be mediated by the prevailing historical context, but this does not mean all hydroclimatic shocks are of equal potential impact. Thus, regarding the association between explosive volcanism and Chinese dynastic collapse over the Common Era, the societal efficacy of volcanic climate forcing was observed to depend not only on levels of pre-existing or contemporaneous societal stress or instability (i.e., the historical context), but also on the magnitude of the climate forcing itself (Gao et al., 2021). Ice-core data suggest that the sequence of approximately 24 tropical and extratropical NH eruptions experienced by Ptolemaic Egypt varied in their climate forcing potential and were not distributed evenly in time (Sigl et al., 2015; Manning et al., 2017). Clusters of historical eruptions have at other times been examined for their potentially severe climatic and societal impacts (e.g., Toohey et al., 2016; Guillet et al., 2020; Campbell and Ludlow, 2020; Stoffel et al., 2022) and such an investigation can make a meaningful contribution to our understanding of the role of explosive volcanism in the history of Ptolemaic Egypt

Table TS1: Table showing the order and specification of the 2.5ka control run. The latest 100 years of the 2.5ka (GHG+ORB+VEG) run are used to represent the base climate for the 2.5ka period. NINT: Non-INTERactive version, MATRIX: Multiconfiguration Aerosol TRacker of mIXing state.


order of control runs and volcanic experiments				
				
ModelE (NINT) 2.5ka control run (1000 years)	ModelE (MATRIX) 2.5ka control run (100 years)	ModelE (MATRIX) 2.5ka control run (70 years) 2.5ka (GHG+ORB)	ModelE (MATRIX) 2.5ka control run (130 years) 2.5ka (GHG+ORB+VEG)	ModelE (MATRIX) 10 ensemble members (Initialized using restart files at 10 years gap) 2.5ka (GHG+ORB+VEG)
Orbital forcing (2.5ka), Greenhouse gases (2.5ka), Vegetations (PI)	Orbital forcing (2.5ka), Greenhouse gases (2.5ka), Vegetations (PI)	Orbital forcing (2.5ka), Greenhouse gases (2.5ka), Vegetations (PI) Retuned Dust Parameters	Orbital forcing (2.5ka), Greenhouse gases (2.5ka), PMIP4 Vegetations (Interpolated for 2.5ka) Retuned Dust Parameters	Orbital forcing (2.5ka), Greenhouse gases (2.5ka), PMIP4 Vegetations (Interpolated for 2.5ka) Retuned Dust Parameters Volcanic SO ₂ injection for 4 eruptions during 158-168 BCE (Sigl et al., 2015)

Fig S1: Spatial representation of changes in major vegetation plant function types (PFTs) under the PMIP4 sensitivity protocol (Otto-Bliesner et al., 2017) after interpolating for the 2.5ka period. Top row shows arid shrub and boreal forest changes, bottom row show C3 and C4 grasses.




Fig S2. Timeseries of annual global mean surface air temperature (top) and precipitation (bottom) for the 2.5ka control run with (green) and without (red) the altered vegetation (see text). Vertical blue lines show the starting points of all 10 ensemble members.




Fig S3. Geographical positions of all 4 volcanic eruptions during the study period are distinguished by different colours. First eruption (E1, Red) erupted during 168 BCE, second (E2, Blue) during 164 BCE, third (E3, Magenta) during 161 BCE and fourth (E4, Orange) erupted during 158 BCE. Spatial box (with caption Nile H) shows the spatial extent of the region considered as the Nile River watershed (discussed in Fig12, main text).




Fig S4: Globally averaged change in sea surface temperature (top panel), surface air temperature over land (middle panel) and over ocean (bottom panel) for each month for the entire simulation period. The light-colored solid lines represent individual ensemble members, and the solid dark colors show the ensemble means. The green vertical dashed lines show the dating of the four eruptions (E1-E4).




Fig S5. Vertical profile of extinction (A) and methane (CH_4 ; B) change following the volcanic eruptions. Vertical black line represents the timing of each eruption.




Fig S6. Spatial pattern of seasonal mean surface temperature response ($^{\circ}\text{C}$) for the four seasons following the first eruption E1. Column 1 show the JJA (June, July August) and SON (September, October, November) for the year of eruption. Column 2 shows the DJF (December, January, February) of the eruption year and first following year, and the MAM (March, April, May) of the first following year. Stippling marks the regions where differences are not significant.




Fig S7. Total cloud cover change during the monsoon season (JJAS) for 3 consecutive years after each eruption (columns) over the North African continent. The blue line demarks the present-day Nile River basin boundary, which is broadly similar to the river extent approximately 2.5ka years ago. The red stippling indicates regions over which change in rainfall is not significant at the 95% confidence level.

