

A cosmogenic nuclide-derived chronology of pre-Last Glacial Cycle glaciations during MIS 8 and MIS 6 in northern Patagonia

Tancrède P. M. Leger¹, Andrew S. Hein¹, Ángel Rodés², Robert G. Bingham¹, Irene Schimmelpfennig³,
5 Derek Fabel², Pablo T. Gonzalez⁴, ASTER Team⁺

¹ School of GeoSciences, University of Edinburgh, Drummond Street, Edinburgh, EH8 9XP, UK

² Scottish Universities Environmental Research Centre, Scottish Enterprise Technology Park, East Kilbride, G75 0QF, Glasgow, UK

³ Aix-Marseille Université, CNRS, Coll. France, IRD, INRAE, CEREGE, Aix en Provence, 13545, France

⁴ Instituto de Geografía, Facultad de Historia, Geografía y Ciencia Política, Pontificia Universidad Católica de Chile, Santiago, Chile

⁺ A full list of authors appears at the end of the paper.

15 *Correspondence:* Tancrède P. M. Leger (Tancrede.leger@ed.ac.uk)

Abstract. The precise environmental mechanisms controlling Quaternary glacial cycles remain ambiguous. To address this problem, it is critical to better comprehend the drivers of spatio-temporal variability in ice-sheet evolution by establishing reliable chronologies of former outlet-glacier advances. When spanning multiple glacial cycles, such chronologies have the capacity to ~~resolve conundrums on~~ contribute to knowledge on the topic of interhemispheric phasing of glaciations and climate events. In southern Argentina, reconstructions of this kind are achievable, as Quaternary expansions of the Patagonian Ice Sheet have emplaced a well-preserved geomorphological record covering several glacial cycles. Moreover, robust ice-sheet reconstructions from Patagonia are powerful barometers of former climate change, as Patagonian glaciers are influenced by the Southern Westerly Winds and its coupled Antarctic Circumpolar Current. Former shifts in these circulation mechanisms are essential to better constrain, as they may have played a critical role in pacing regional and possibly global Quaternary climate change. Here, we present a new set of cosmogenic ¹⁰Be and ²⁶Al exposure ages from pre-Last Glacial Cycle moraine boulder, glaciofluvial outwash cobble and bedrock samples. This dataset constitutes the first direct chronology dating pre-LGM glacier advances in northern Patagonia, and completes our effort to date the entire preserved moraine record of the Río Corcovado valley system (43°S, 71°W). We find the outermost margins of the study site depict at least three distinct pre-Last Glacial Cycle glaciations occurring at ~~284 ± 7~~ around 290-270 ka, ~~257 ± 7~~ 270-245 ka, and ~~447 ± 4~~ 130-150 ka. Combined with the local LGM chronology, we discover that a minimum of four distinct Pleistocene glaciations occurred during Marine Isotope Stages eight, six, and two in northern Patagonia. Evidence for ~~stage~~ Marine Isotope Stage four and three deposits were not found at the study site. ~~which, This~~ illustrates former longitudinal and latitudinal asynchronies in Patagonian Ice Sheet mass

Formatted: French (France)

Formatted: Spanish (Spain, Traditional Sort)

balance during these [stages: Marine Isotope Stages](#). We find the most extensive middle-to-late Pleistocene expansions of the Patagonian Ice Sheet appear to be out-of-phase with local summer insolation intensity, but synchronous with orbitally-controlled periods of longer and colder winters. Our findings thus enable to [explore/investigate](#) the potential roles of seasonality and seasonal duration in driving southern mid-latitude ice-sheet mass balance and facilitate novel glacio-geomorphological interpretations for the study region. They also provide empirical constraints of former ice-sheet extent and dynamics that are essential to calibrating numerical ice-sheet and glacial isostatic adjustment models.

40

45

50

55

60

65

1 Introduction

Understanding the environmental mechanisms driving the Quaternary build-up and collapse cycles of Earth's major ice sheets remains a focus of ongoing research. Key to this research is the establishment of detailed chronologies offering direct dating of major ice-sheet expansion and recession events (Darvill *et al.*, 2016). Numerical glacier chronologies are most informative when presenting unambiguous dating of ice-sheet margins spanning the entirety of the last glacial cycle (LGC: ~125-15 ka) and ideally earlier Quaternary glacial cycles (Kaplan *et al.*, 2010; Schaefer *et al.*, 2015). To the east of the Patagonian Andes, the Argentinian foreland contains one of the most complete and well-preserved sequences of Quaternary glacial deposits in the world (Clapperton, 1993). This unique geomorphological record provides the opportunity to reconstruct and date glacier fluctuations of the formerly 2500 km-long (from north to south) Patagonian Ice Sheet (PIS) over several glacial cycles (Mercer, 1976; Hein *et al.*, 2011). Furthermore, the PIS occupied a key mid-latitude position in the ocean-dominated Southern Hemisphere (SH) and was the only ice mass to fully intersect the precipitation-bearing southern westerly wind (SWW) belt, a fundamental feature of the southern mid-latitude climate system. Hence, reconstructing the timing of PIS expansion and recession events during several glacial cycles can provide rare insight into southern mid-latitude Quaternary climate evolution (Davies *et al.*, 2020).

Palaeo-climate proxy records such as those retrieved from ice- and ocean-sediment cores indicate that some of the most potent Quaternary cooling events occurred during the Middle Pleistocene (Parrenin *et al.*, 2013; Shakun *et al.*, 2015), a period (~800 - 130 ka; Hughes *et al.*, 2020) characterized by 100-ka pacing of glacial-interglacial cycles. Paradoxically, there is a dearth of directly-dated terrestrial glacial records for these glacial events (Hughes *et al.*, 2020). In Patagonia, bracketing ages for pre-LGC glaciations have been produced in some places (*e.g.* near Lago Argentino; 50.5°S) with $^{40}\text{Ar}/^{39}\text{Ar}$ dating, *K-Ar* dating, and palaeo-magnetic measurements on lava flows interbedded between Pliocene/Pleistocene glacial-till units (Mercer, 1976; Sylwan *et al.*, 1991; Meglioli, 1992; Singer *et al.*, 2004). For instance, Mercer (1976) used such methods to constrain the likely timing of the Greatest Patagonian Glaciation to between 1.2 Ma and 1.0 Ma in the Río Gallegos (52°S) and Lago Buenos Aires (46.5°S) regions. While such studies have been instrumental in establishing the broad chronological framework for Patagonian glaciations (Rabassa and Coronato, 2009), direct numerical dating of individual outlet-glacier advances is now required to develop detailed knowledge of pre-LGM and pre-LGC ice-sheet activity in Patagonia.

In Patagonia, direct numerical dating of pre-LGC glacial advances, achievable using terrestrial cosmogenic nuclide (TCN) surface exposure dating, has been limited to two valleys of eastern central Patagonia (Kaplan *et al.*, 2005; Hein *et al.*, 2009; 2011; 2017; Cogež *et al.*, 2018). Throughout the rest of the southern mid-latitudes (23 - 66 °S), terrestrial glacial records yielding TCN exposure ages that pre-date the LGC have been reported using moraine boulders in New Zealand (Putnam *et al.*, 2013), the southern central Andes (Terrizzano *et al.*, 2017) and Tasmania (Barrows *et al.*, 2002; Kiernan *et al.* 2004; 2010;

2014; 2017; Colhoun and Barrows, 2011, Augustinus *et al.*, 2017). Because pre-LGC moraines and boulders are often poorly
100 preserved, the majority of these investigations have identified the approximate timing of individual pre-LGC glacial advances
with small ($n = 5$ per moraine) and scattered sets of ages. Consequently, robust and direct numerical chronologies of pre-
LGC glacial expansion events remain particularly scarce in the southern mid-latitudes. Patagonian investigations have shown
that TCN exposure dating can yield direct ages for the deposition of pre-LGC moraine-outwash complexes preserved in the
Argentinian steppe (*e.g.* Hein *et al.*, 2017). While moraine boulders are well-suited to dating glacier advances of the LGC,
105 Hein *et al.* (2009) revealed that targeting outwash terraces in eastern Patagonia can be more appropriate for dating pre-LGC
glacial margins, because their low-gradient surfaces are less prone to degradation via gravity-driven diffusion when compared
to steep-sided and unconsolidated moraines. On pre-LGC moraines, such degradation more frequently causes moraine surface
lowering and boulder exhumation post-deposition, leading to significant exposure-age scatter and underestimations of moraine
formation age. In contrast, surface cobbles from well-preserved outwash plains tend to display comparatively less post-
110 depositional scatter and are thought to date the glacial event more closely (Hein *et al.*, 2009). This approach has been used
successfully to constrain the timing of glacial advances in several locations of central and southern Patagonia (Darvill *et al.*,
2015; Hein *et al.*, 2009; 2011; 2017; Mendelová *et al.*, 2020). However, to date, no pre-LGC glacial events have been dated
using this technique in northern Patagonia. Consequently, here we adopt this approach and target both moraine boulders and
outwash surface cobbles to constrain the timing of pre-LGC glacial advances in northeastern Patagonia.


115
We focus our investigation on the Río Corcovado (RC) valley system (43°S; 71°W), formerly host to a major outlet glacier of
the PIS in northeastern Patagonia (Caldenius, 1932). We present a multi-glacial-cycle ^{10}Be and ^{26}Al TCN exposure-age
chronology from the three stratigraphically oldest glacial moraine/outwash complexes deposited by the RC outlet glacier. This
study builds on our earlier work (Leger *et al.*, 2020, 2021a,b) to compile a geochronological reconstruction that spans the
120 entire preserved moraine-outwash record of the RC valley system. We use our cosmogenic nuclide-derived chronology to test
whether major middle Pleistocene advances previously dated in central Patagonia (Hein *et al.*, 2009; 2017) coincide with
moraines found in its northeastern sector (this study), and thus whether such advances depict strong regional climate signals
expressed across much of the palaeo-ice front. We then compare the different magnitudes of middle Pleistocene Patagonian
glaciations relative to other formerly glaciated regions of the world, and thereby assess the presence of former interhemispheric
125 and inter-regional *a*/synchronies in ice-sheet behaviour. Finally, we evaluate the synchronicity of PIS expansion events with
other global and regional palaeo-climate proxy records, and explore the potential role of various insolation parameters on
controlling middle-to-late Pleistocene climate variations in Patagonia and the southern mid-latitudes.

130 2 Setting

Our study site focuses on the ice-marginal environment of the former PIS during Quaternary ice-age maxima, at the latitude of its major eastward-flowing Palena outlet glacier (43°S; Leger *et al.*, 2020; 2021a). This up to 100 km-long outlet glacier diverged towards the eastern mountain front into three glaciers occupying the Río Frío valley to the northeast, the Río Huemul (RH) valley to the east, and the RC valley to the south (Fig. 1; Leger *et al.*, 2021a). The local valleys are characterised by a westward-sloping bed topography such that the former outlet glaciers flowed up along reverse-sloping beds. The surface geology of the study area is characterised by well-preserved and distinct sequences of ice-contact glaciogenic deposits interspersed among their associated glaciolacustrine and glaciofluvial sediment-landform assemblages (Leger *et al.*, 2020). Terminal moraine complexes indicate that the RC and RH outlet glaciers re-converged to deposit moraine sequences located approximately 60 km to the east of the Argentinian town of Corcovado (43°54'S; 71°46'W), into an environment presently climatically defined as the semi-arid Patagonian steppe (Fig. 1; Garreaud *et al.*, 2013). Local aridity (~570 mm a⁻¹; Fick and Hijmans, 2017; Fig. 1a) results from the Patagonian Andes acting as a powerful orographic barrier to the dominant ~~SW~~**Southern Westerly Winds** (Fig. 1a), causing a potent west-east rain-shadow effect (Garreaud *et al.*, 2013; Fig. 1a). Along the flowlines of the RC outlet glacier, ice flow eroded bedrock mainly sourced from the Mesozoic north Patagonian batholith towards the west (Hervé *et al.*, 2017) along with formations of Jurassic/Cretaceous volcanic and sedimentary units located ~~farther~~**farther** east (Haller *et al.*, 2003). Moraine-outwash complexes studied here thus contain a mixture of rock types including quartz-bearing leuco-monzonite, granite and white granodiorite that are well-suited to TCN exposure dating using *in situ* ¹⁰Be and ²⁶Al radionuclides.

150 Ice sheet-wide geomorphological mapping of glacial sediment-landform assemblages has been conducted by Caldenius (1932), Glasser and Jansson (2008), and Davies *et al.* (2020) while the first detailed glacial geomorphological map specific to the RC and RH valleys was generated by Leger *et al.* (2020). These maps enabled the identification of at least eight stratigraphically-distinct moraine-outwash complexes deposited by the RC outlet glacier (Fig. 1c). This study focuses on dating results from the three stratigraphically oldest glacial margins: *i.e.* the RC 0 (outermost), RC I and RC II (Fig. 1c) moraine-outwash complexes. Directly inboard of these margins, the RC outlet glacier deposited another series of at least five younger moraine complexes (RC III-VII moraines), previously dated to between 26.4 ± 1.4 ka and 19.9 ± 1.1 ka using TCN exposure dating and Bayesian age modelling (Leger *et al.*, 2021a). The latter moraine belts reflect the local Last Glacial Maximum (LGM) expansions of the PIS.

160



165

170

175

180

185

190

195

Figure 1. (A): 1-km² spatial resolution total annual precipitation (mm a⁻¹) and mean annual windspeed (m s⁻¹) land data (1970-2000 mean) from the WorldClim version 2.1 dataset (Fick and Hijmans, 2017) overlaying maps of southernmost South America with location of study area highlighted by black box. (B): Adapted from Leger *et al.* (2020; 2021 a,b): Approximate former extent of the PIS during the LGM, modified from Glasser and Jansson (2008). Locations of major PIS paleo-outlet glaciers are designated: BI: Bahía Inútil, EM: Estrecho de Magallanes, SS: Seno Skyring, BV: Bella Vista, TDP & UE: Torres del Paine & Última Esperanza, LA: Lago Argentino, LV: Lago Viedma, LC/P: Lago Cochrane/Pueyrredón, LBA: Lago Buenos Aires, LP/LGV: Lago Palena/General Vintter. NPI, SPI and CDI stand for North, South, and Cordillera Darwin Patagonian ice fields, respectively. A - 125 m (brown-coloured) topographic contour roughly simulating former coastline locations at the LGM is displayed (after Lambeck *et al.*, 2014). Bathymetric data were acquired from the General Bathymetric Chart of the Oceans (GEBCO). (C): Glacial geomorphology of the study area indicating stratigraphic relationship of moraine-outwash complexes preserved, sample site location, and former ice-flow direction. *Other TCN dating samples: published in Leger *et al.* (2021 a,b).

210 3 Methods

Preliminary identification and mapping of major glacial sediment-landform assemblages studied here was carried out using remotely-sensed imagery and topographic data. These geomorphological interpretations were then ground-checked during a 5-week field season during the 2020 austral summer. Details of the geomorphological mapping methodology and criteria, and the complete map of the study area, are given by Leger *et al.* (2020). To date the RC 0 - II glacier advances, we sampled both outwash surface cobbles and moraine boulders. Specifically, we measured ¹⁰Be in five cobbles from the outermost proglacial outwash surface sampled, here termed the “Tecka outwash”, which we correlate stratigraphically with the RC 0 advance (Fig. 1c). We sampled five and six cobbles respectively from the RC I and RC II outwash terraces, six moraine boulders from the RC II moraine complex, and one ice-moulded bedrock surface located between the RC I and II margins (Figs. 1-3). Additionally, we measured ²⁶Al in five of these 23 samples to determine whether they present simple or complex exposure/burial histories. The chosen moraine boulders are large (90-200 cm high) and rounded to sub-rounded granodiorite and granite erratics embedded in the RC II moraine crest (Fig. 3). TheWhere found, the top 2-5 cm of boulder surfaces exhibiting glacial polish smooth rock fragments protruding from more eroded surrounding surfaces were sampled using hammer, chisel and angle grinder.

To date the formation of proglacial outwash plains, quartz-bearing and fluvially-rounded cobbles (4-10 cm diameter; crushed whole) were sampled from well-preserved terraces characterised by low surface gradients (~0.5°-1°), following the procedure established by Hein *et al.* (2009; 2011) (Fig. 2a-d). Surface ventifacts and desert varnish, indicative of prolonged cobble exposure, were considered a required sampling criterion. Sample preparation and cosmogenic isotope ratio measurements were conducted at three different laboratories: the cosmogenic isotope analysis and AMS Accelerator Mass Spectrometer (AMS)

facilities of the Scottish Universities Environmental Research Centre (SUERC) (East Kilbride, UK), the French National Cosmogenic Nuclides Laboratory (LN2C) and the ASTER AMS facility of the European Centre for Research and Teaching in Environmental Geosciences (CEREGE, Aix-en-Provence, France), and the University of Edinburgh's Cosmogenic Nuclide Laboratory (Edinburgh, UK). Sample details and nuclide concentrations are displayed in Table 1. Details relevant to sample

235 location, photographs, preparation, and wet chemistry are provided in the supplementary materials.

^{10}Be and ^{26}Al exposure ages were calculated using the online calculator formerly known as the CRONUS-Earth online calculator version 3 (Balco *et al.*, 2008). Rock density is assumed to be 2.65 g cm^{-3} and elevation flag is STD. For ^{10}Be exposure-age calculations, we used the *NIST_27900* standardisation (equivalent to 07KNSTD within rounding error) and the central Patagonia production rate (Kaplan *et al.*, 2011) obtained and re-calculated from the ICE-D online database (<http://calibration.ice-d.org/>). Details regarding specific AMS standards, nominal $^{10}\text{Be}/^9\text{Be}$ ratios and ^{10}Be half-life employed are included in the supplementary materials. ^{26}Al exposure ages were calculated using the Purdue Z92-0222 AMS standardisation (equivalent to KNSTD within rounding error) and the default CRONUS-Earth online calculator's global production rate of Borchers *et al.* (2016), as the central Patagonia production rate does not feature $^{26}\text{Al}/^{27}\text{Al}$ measurements. We

240 however consider it reasonable to use both the central Patagonia and global production rates when comparing ^{10}Be and ^{26}Al ages, as their respective ^{10}Be calibrations agree within error, with reported LSDn scaling- fitting parameter values of 0.823 ± 0.067 and 0.846 ± 0.040 , respectively (calibration: ICE-D online database). All exposure ages are displayed in Table 2. Below, we discuss exposure ages calculated using the time-dependent LSDn scaling scheme of Lifton *et al.* (2014) with 1σ analytical uncertainties. Exposure ages are between 6-9% and 0.6-1% older when using the *St* and *Lm* scaling schemes of Lal (1991) and

250 Stone (2000), respectively. The ^{10}Be New Zealand production rate (Putnam *et al.*, 2010), often used for Patagonian TCN exposure-age datasets due to its southern mid-latitude location (44°S), decreases our exposure ages by between 1.7% and 1.5% using LSDn scaling, which is less than 1σ analytical uncertainties (3.5-1.6%).

255

260




Figure 2. Field photographs of: (A-D) proglacial outwash plains and outwash surface cobbles sampled for TCN exposure dating and; (E, F) the ice-moulded bedrock surface sampled (RC20-01). Ages displayed are ¹⁰Be exposure ages ± 1σ analytical uncertainties. Further sample coordinates and characteristics are presented in Table 1.

300

305

310

315

320




Figure 3. Field photographs of moraine granite and granodiorite boulders sampled for TCN exposure dating. Ages displayed are ^{10}Be exposure ages $\pm 1\sigma$ analytical uncertainties.

325

330

4 Results

335 4.1 Geomorphology

In the RC valley, the three most distal preserved moraine complexes (RC 0-II) are morphologically distinct from the younger LGM moraines (RC III-VII). They display greater relief (70-90 m) than the RC III – VII moraines (10-40 m; Leger *et al.*, 2021a), are more subdued, and have broader-crested ridges. [TheAlong our sampling transect, the](#) three moraine complexes are 14 km (RC 0), 12 km (RC I) and 5 km (RC II) more distal than the outermost-dated LGM moraine (RC III). The sampled RC II moraine is a ~1 km-wide multi-ridge complex presenting a flat-topped, subdued crest surface elevated up to 45 m above its ice-distal outwash plain. Ice-distal slope gradient is variable but does not exceed 5°, a relatively low grade compared to the slopes of the younger, better preserved LGM moraines which feature slope gradients of 9-19° (Leger *et al.*, 2021a). The RC I moraine complex features slope gradients (>5°) and surface geomorphologies for individual moraine ridges comparable to the RC II moraine belt, but is much wider in places (up to 4.6 km) and exhibits a greater concentration of distinct hummocky ridges and moraines. The subdued moraine geomorphologies characteristic of the RC 0-II margins are suggestive of progressive lateral slope downwasting and moraine surface erosion of fine material post-deposition (Clapperton, 1993; Glasser and Jansson, 2008; Hein *et al.*, 2017). Geomorphological observations thus suggest the RC II and older moraine complexes were deposited significantly prior to the local LGM. In the adjacent RH valley, the outermost glacial margin consists of a ~2 km-wide belt of kettle-kame topography displaying a chaotic hummocks-and-hollows configuration, and was formerly termed the “Tecka drift” (Haller *et al.*, 2003; Leger *et al.*, 2020). The sampled Tecka outwash surface marks the distal limit of this kettle-kame [glaciogenic](#) deposit (Fig. 1c). Based on geomorphological mapping and moraine-complex traceability across the RC and RH valleys, the Tecka limit (RH valley) is assumed here to correspond stratigraphically to the RC 0 advance, for which no well-preserved outwash plain surface could be found in the RC valley (Fig. 1c).

355 The outwash surfaces sampled in this study all present long-term preservation properties (Fig. 2). They display low-gradient surfaces (0.5-1°), desert pavements, sparse vegetation cover, and preserved braided palaeo-meltwater channels suggestive of limited surface erosion/deflation (Fig. 2a,d; Leger *et al.*, 2020). These glaciofluvial terraces are composed of sorted and fluvially-rounded sands and gravel deposits. Near the locations of outwash cobble sampling, the outwash surfaces display no detectable signs of post-deposition activation and/or incision by post-glacial hydrology. Where checked, modern soil thickness varied between 0 and 20 cm. This is comparable to observations made further south in the Lago Pueyrredón (47.3°S) and Lago Belgrano (47.5°S) valleys (Hein *et al.*, 2009; Mendelová *et al.*, 2020). At the locations of sampling, these terraces occur adjacent to moraine complexes and mark their distal limits (Fig. 1c), thus enabling the unambiguous stratigraphic association of an outwash deposit with a specific glacial advance.

365

Table 1. TCN exposure dating sample details and nuclide concentrations.

Sample ID	Latitude (DD)	Longitude (DD)	Elevation (m a.s.l.)	Boulder height (m)	Thickness (cm)	Shielding correction	SiO ₂ mass dissolved (g)	Nuclide	Total ²⁷ Al mass (mg)	±1σ (mg)	¹⁰ Be or ²⁶ Al/ ²⁷ Al ratio	±1σ ratio	Blank correction	¹⁰ Be (or ²⁶ Al) atoms g ⁻¹ (SiO ₂)	±1σ atoms g ⁻¹ (SiO ₂)	AMS Carboide code	²⁶ Al/ ¹⁰ Be
RH Teika outwash surface cobbles																	
19RHS09 ^Y	-43.5975	-70.88756	828	n/a	9.0	0.9993	21.4250	¹⁰ Be	0.24843	0.0005	2.6145E-12	6.2270E-14	0.27%	2.0277E+06	6.3160E+04	b11920	n/a
19RHS10 [*]	-43.5975	-70.88756	828	n/a	9.0	0.9993	21.4477	¹⁰ Be	0.24980	0.0005	2.4737E-12	6.2290E-14	0.28%	1.9201E+06	6.2099E+04	b11921	n/a
19RHS12 [*]	-43.5975	-70.88756	828	n/a	9.0	0.9993	20.3129	¹⁰ Be	0.24824	0.0005	2.7806E-12	3.5790E-14	0.25%	2.2683E+06	6.4118E+04	b11922	n/a
19RHS13 [*]	-43.5975	-70.88756	828	n/a	10.0	0.9993	20.5896	¹⁰ Be	0.24882	0.0005	2.5540E-12	2.7600E-14	0.28%	2.0579E+06	4.7039E+04	b11923	n/a
19RHS15 [*]	-43.5975	-70.88756	828	n/a	7.5	0.9993	20.6152	¹⁰ Be	0.24833	0.0005	2.3316E-12	2.7170E-14	0.30%	1.8715E+06	4.3560E+04	b11924	n/a
RC I outwash surface cobbles																	
RC20-12 [*]	-43.86717	-71.09769	989	n/a	5.0	0.9998	1.9101	¹⁰ Be	0.45617	0.0005	1.3234E-13	4.6198E-15	1.46%	2.0812E+06	7.3876E+04	HDL	n/a
RC20-13 [*]	-43.86717	-71.09769	989	n/a	4.0	0.9998	19.9300	¹⁰ Be	0.22898	0.0005	2.4409E-12	5.0220E-14	0.82%	1.8577E+06	3.8833E+04	b12115	n/a
RC20-14 [*]	-43.87017	-71.10005	993	n/a	9.0	0.9998	3.3260	¹⁰ Be	2.54209 [†]	0.1086	4.1826E-12	5.810E-14	0.16%	1.1803E+07	5.3809E+05	a3380	6.4 ± 0.3
RC20-15 [*]	-43.87017	-71.10005	993	n/a	5.0	0.9998	19.6520	¹⁰ Be	0.44201	0.0005	1.9378E-13	5.8984E-15	1.02%	1.7071E+06	5.2408E+04	HDM	n/a
RC20-16 [*]	-43.87132	-71.09883	991	n/a	4.5	0.9998	4.5739	¹⁰ Be	0.22940	0.0005	2.1267E-12	3.1670E-14	0.94%	1.6425E+06	2.5061E+04	b12118	6.0 ± 0.6
RC20-01 [*]	-43.7036	-71.10018	1026	n/a	1.5	0.9990	7.3150	¹⁰ Be	2.26658 [†]	0.2340	3.8320E-12	5.8150E-14	0.19%	9.8440E+06	1.0466E+06	a3381	n/a
RC20-01 [*]	-43.7036	-71.10018	1026	n/a	1.5	0.9990	7.3150	²⁶ Al	0.44368	0.0005	3.5153E-13	8.3475E-15	0.56%	2.2687E+06	5.4143E+04	HDK	n/a
RC I/II ice-moulded bedrock surface																	
RC20-01 [*]	-43.7036	-71.10018	1026	n/a	1.5	0.9990	7.3150	¹⁰ Be	0.24810	0.0005	8.1307E-13	1.9970E-14	2.35%	1.7405E+06	4.4314E+04	b12105	6.8 ± 0.3
RC20-01 [*]	-43.7036	-71.10018	1026	n/a	1.5	0.9990	7.3150	²⁶ Al	2.13039 [†]	0.0748	1.8242E-12	4.4450E-14	0.43%	1.1803E+07	5.0785E+05	a3375	n/a
RC II outwash surface cobbles																	
RC20-06 [*]	-43.79607	-71.11590	974	n/a	7.0	1.00000	13.9770	¹⁰ Be	0.24171	0.0005	9.5491E-13	1.8140E-14	1.99%	1.0800E+06	2.1289E+04	b12111	n/a
RC20-07 [*]	-43.79607	-71.11590	974	n/a	7.5	1.00000	20.5390	¹⁰ Be	0.24078	0.0005	1.3495E-12	2.1480E-14	1.42%	1.0412E+06	1.7088E+04	b12112	n/a
RC20-10 [*]	-43.80491	-71.12183	976	n/a	6.0	1.00000	19.8810	¹⁰ Be	0.22016	0.0005	1.4072E-12	2.6350E-14	1.42%	1.0724E+06	2.0627E+04	b12119	n/a
RC20-20 [*]	-43.80491	-71.12183	976	n/a	4.0	1.00000	4.5249	¹⁰ Be	0.43875	0.0005	2.0274E-13	5.7110E-15	0.89%	1.3006E+06	2.7075E+04	HDV	n/a
RC20-21 [*]	-43.80383	-71.12228	977	n/a	5.5	1.00000	6.4173	¹⁰ Be	0.44192	0.0005	2.8334E-13	7.4642E-15	0.70%	1.2555E+06	3.3338E+04	HDQ	n/a
RC20-22 [*]	-43.80383	-71.12228	977	n/a	4.0	1.00000	20.7920	¹⁰ Be	0.23398	0.0005	1.5691E-12	2.4620E-14	1.25%	1.1643E+06	1.8792E+04	b12120	n/a
RC II moraine boulders																	
RC20-02 [*]	-43.79239	-71.12655	984	1.56	2.0	0.9999	15.0190	¹⁰ Be	0.24273	0.0005	1.0789E-12	2.7230E-14	1.76%	1.1433E+06	2.9631E+04	b12106	n/a
RC20-03 [*]	-43.79345	-71.12502	986	0.86	1.4	0.9999	20.0790	¹⁰ Be	0.24451	0.0005	1.2374E-12	1.7440E-14	1.52%	9.9050E+05	1.4486E+04	b12107	n/a
RC20-04 [*]	-43.79600	-71.13014	983	2.00	2.3	0.9999	19.7420	¹⁰ Be	0.23424	0.0005	1.8605E-12	3.8960E-14	1.05%	1.4566E+06	3.1076E+04	b12108	6.3 ± 0.3
RC20-05 [*]	-43.81415	-71.15841	1020	1.15	2.5	0.9999	20.0200	¹⁰ Be	2.86499 [†]	0.0895	2.8283E-12	4.3000E-14	0.21%	9.1319E+06	3.1778E+05	a3376	6.5 ± 0.3
RC20-10 [*]	-43.81509	-71.15717	1029	1.66	1.7	0.9999	10.5800	¹⁰ Be	0.23186	0.0005	1.8099E-12	3.4030E-14	1.10%	1.3845E+06	2.6868E+04	b12109	n/a
RC20-11 [*]	-43.80433	-71.13924	1002	1.55	1.2	1.00000	19.8670	¹⁰ Be	0.20353 [†]	0.0661	3.9912E-12	6.4770E-14	0.21%	9.0374E+06	3.2871E+05	a3377	n/a
RC20-11 [*]	-43.80433	-71.13924	1002	1.55	1.2	1.00000	19.8670	¹⁰ Be	0.23772	0.0005	5.1964E-13	9.4730E-15	3.73%	7.4934E+05	1.4825E+04	b12113	n/a
RC20-11 [*]	-43.80433	-71.13924	1002	1.55	1.2	1.00000	19.8670	²⁶ Al	0.23441	0.0005	1.2858E-12	2.4140E-14	1.53%	9.9741E+05	1.9268E+04	b12114	n/a
Blanks																	
RC20-BL	Processed with	n/a	n/a	n/a	n/a	n/a	n/a	¹⁰ Be	0.44728	0.0005	1.9671E-15	3.0249E-16	n/a	n/a	n/a	HDP	n/a
CB190421	Δ samples	n/a	n/a	n/a	n/a	n/a	n/a	¹⁰ Be	0.22855	0.0005	2.1105E-14	2.6500E-15	n/a	n/a	n/a	b12121-bb1	n/a
EUTL9	Y samples	n/a	n/a	n/a	n/a	n/a	n/a	²⁶ Al	2.97988 [†]	0.1120	6.1162E-15	1.7660E-15	n/a	n/a	n/a	a3382	n/a
EUTL9	Y samples	n/a	n/a	n/a	n/a	n/a	n/a	¹⁰ Be	0.24843	0.0005	7.0592E-15	1.5050E-15	n/a	n/a	n/a	b11930	n/a
EUTL9	Y samples	n/a	n/a	n/a	n/a	n/a	n/a	¹⁰ Be	0.24843	0.0005	7.0592E-15	1.5050E-15	n/a	n/a	n/a	b11930	n/a

385

390

395

400

Table footnotes. All cobble samples were crushed whole, without prior cutting. Sample preparation and wet chemistry was conducted at three different laboratories: Y samples: The University of Edinburgh's Cosmogenic Nuclide Laboratory (Edinburgh, UK), * samples: The Scottish Universities Environmental Research Centre (SUERC) (East Kilbride, UK), Δ samples: The French National Cosmogenic Nuclides Laboratory (LN2C) of the European Centre for Research and Teaching in Environmental Geosciences (CEREGE, Aix-en-Provence, France). Nuclide ratios in Y and * samples were measured at the SUERC AMS facility (East Kilbride, UK), while Nuclide ratios in Δ samples were measured at the ASTER AMS facility (CEREGE, Aix-en-Provence, France). SiO₂ is the chemical formula of silica (quartz). # Total ²⁷Al mass from both carrier (²⁷Al concentration: 982 ± 0.1 μg g⁻¹) and sample, determined by ICP-OES. Be Carrier solution used at LN2C has a ⁹Be concentration of 3025 ± 9 μg g⁻¹ while carrier solutions used at SUERC and the University of Edinburgh's Cosmogenic Nuclide Laboratory have ⁹Be concentrations of 849 ± 12 μg g⁻¹ and 1000 μg g⁻¹, respectively. Conversions of isotope ratios to ¹⁰Be and ²⁶Al concentrations were conducted following standard equations, as described by Balco (2006).

4.2 TCN exposure-age chronology

The exposure-age results for the 23 samples are presented in Table 2. Four of the five cobbles sampled on the outermost Tecka/RC 0 outwash surface (Fig. 1c) range from 252.0 ± 6.3 ka to 284.1 ± 7.0 ka and afford a mean ^{10}Be exposure age of 268.7 ± 14.4 ka (arithmetic mean $\pm 1\sigma$ standard deviation). One older age (19RHS12: 311.5 ± 8.1 ka) lies outside the remaining population's 95% (2σ) confidence level and is rejected as a statistical outlier. The remaining exposure-age population is relatively well-clustered given the landform age, but still displays a >1 mean square weighted deviation (MSWD) value of 4.38 ($>k = 2.63$). Such a MSWD value is indicative of greater exposure-age scatter than can be predicted by 1σ analytical uncertainties alone (Jones *et al.*, 2019).

The five ^{10}Be ages from the RC I outwash surface cobbles range from 185.0 ± 3.0 ka to 257.4 ± 6.6 ka and afford a mean exposure age of 214.0 ± 29.5 ka. Although the age population presents no obvious statistical or stratigraphical outliers, the exposure ages are poorly clustered, and present a high MSWD value of 33.6 ($>k = 2.41$), indicating significantly more exposure-age scatter than solely predicted by analytical uncertainties. From the ^{26}Al measured in two samples (RC20-13 and RC20-15) from this population, the resulting $^{26}\text{Al}/^{10}\text{Be}$ concentration ratios are 6.4 ± 0.3 and 6.0 ± 0.6 , respectively, and are consistent with a single, continuous exposure history ([within uncertainty](#)) post erosion (Granger and Muzikar, 2001; Balco and Rovey, 2008; [supplementary materials figure 1](#)).

The ice-moulded bedrock surface (RC20-01), located inboard of the RC I moraine and outboard of the RC II moraine, yields a ^{10}Be exposure age of 185.7 ± 5.0 ka, an ^{26}Al exposure age of 181.1 ± 8.5 ka, and an $^{26}\text{Al}/^{10}\text{Be}$ concentration ratio of 6.8 ± 0.3 , which is consistent with continuous exposure post erosion of the surface.

The six ^{10}Be ages from the RC II outwash surface cobbles range from 120.6 ± 2.0 ka to 146.6 ± 4.3 ka with a mean exposure age of 131.3 ± 11.1 ka. The age population is tightly clustered considering analytical uncertainties associated with TCN exposure dating of pre-LGC landforms, and thus features no obvious statistical or stratigraphical outliers. However, the population still yields a MSWD value of 11.4 ($>k = 2.26$), diagnostic of some non-analytical exposure-age scatter in the ^{10}Be age distribution.

Three of the six ^{10}Be ages from the RC II moraine boulders range from 125.5 ± 3.4 ka to 161.9 ± 3.6 ka and afford a mean exposure age of 145.3 ± 18.4 ka. Younger ages from the remaining three boulders (RC20-03, 10, 11) plot outside the remaining boulder population's 95% confidence level and the 2σ envelope associated with the youngest RC II outwash cobble (RC20-07). We consider ^{10}Be inheritance an unlikely source of exposure-age scatter compared to the high potential for boulder exhumation causing young outliers. Moreover, the RC II outwash and RC II moraine belt are geomorphologically likely to

represent the same glacier expansion event. Hence, these three younger boulder ages were rejected as stratigraphical and statistical outliers (Table 2). The three remaining moraine boulder ages still exhibit a high MSWD value of 28.7 ($>k = 3.0$), indicating that the remaining dataset displays substantially more exposure-age scatter than can be predicted solely by analytical uncertainties. The two oldest sampled boulders, RC20-04 (161.9 ± 3.6 ka) and RC20-05 (148.6 ± 3.0 ka), yielded ^{26}Al exposure ages of 143.6 ± 5.4 ka and 137.5 ± 5.4 ka, corresponding to $^{26}\text{Al}/^{10}\text{Be}$ concentration ratios of 6.3 ± 0.3 and 6.5 ± 0.3 , respectively. Such ratios are consistent with continuous exposure post erosion.

435

440

445

450

455

460

465

Table 2. TCN exposure ages and summary statistics.

Sample ID	Nuclide	LSDn			St			Lm			Outlier
		Age Years BP	Internal	External	Age Years BP	Internal	External	Age Years BP	Internal	External	
Tecka drift outwash surface cobbles											
19RHS09	¹⁰ Be	276461	9267	25853	299616	10102	28347	278319	9333	26176	
19RHS10	¹⁰ Be	262261	9057	24545	283596	9847	26839	263892	9117	24837	
19RHS12	¹⁰ Be	311517	8053	28598	339128	8829	31545	313965	8121	29001	Yes
19RHS13	¹⁰ Be	284138	6979	25816	308209	7617	28348	286092	7030	26152	
19RHS15	¹⁰ Be	251988	6251	22738	272373	6791	24856	253561	6292	23017	
Mean (n = 5): 277.27 ka; 1σ std: 22.83 ka											
Mean (n = 4): 268.71 ka; 1σ std: 14.37 ka; 1σ internal: 15.99 ka; 1σ internal + PR%: 28.35 ka											
Oldest cobble (n=4): 284.14 ka; 1σ internal: 6.98 ka											
Uncertainty weighted mean (n = 4): 268.06 ka; 1σ std: 13.11 ka											
MSWD: 4.38 > k: 2.63 (n=4); Peak age (n=4): 281.26 ka											
RC I outwash surface cobbles											
RC20-12	¹⁰ Be	236643	8917	22310	258263	9785	24618	240146	9057	22775	
RC20-13	¹⁰ Be	208105	4584	18433	226933	5023	20322	211056	4653	18815	
	²⁶ Al	191679	9487	20471	203478	10131	25523	189094	9347	21655	
RC20-14	¹⁰ Be	198062	6403	18118	215770	7007	19941	200857	6498	18484	
	²⁶ Al	184989	2957	16053	200289	3214	17559	187105	2993	16340	
RC20-15	¹⁰ Be	157741	18142	23333	166261	19204	26863	155152	17821	23787	
	²⁶ Al	257360	6563	23302	281123	7213	25764	261214	6668	23806	
RC20-16	¹⁰ Be	257360	6563	23302	281123	7213	25764	261214	6668	23806	
Mean (n = 5): 214.03 ka; 1σ std: 29.48 ka; 1σ internal: 13.90 ka; 1σ internal + PR%: 23.27 ka											
Oldest cobble (n=2): 257.36 ka; 1σ internal: 6.56 ka;											
Uncertainty weighted mean (n = 5): 209.46 ka; 1σ std: 25.91 ka											
MSWD: 33.57 > k: 2.41 (n=5); Peak age (n=5): 185.20 ka											
RC I-II ice-moulded bedrock surface											
RC20-01	¹⁰ Be	185674	4954	16596	201425	5395	18184	187977	5018	16904	
	²⁶ Al	181079	8514	19053	191447	9049	23703	178385	8376	20151	
RC II outwash surface cobbles											
RC20-06	¹⁰ Be	124477	2532	10758	133895	2730	11671	125552	2554	10915	
RC20-07	¹⁰ Be	120592	2040	10323	129458	2195	11177	121543	2057	10467	
RC20-19	¹⁰ Be	122414	2428	10560	131578	2616	11448	123440	2449	10712	
RC20-20	¹⁰ Be	146616	4334	13121	158063	4686	14269	148078	4379	13328	
RC20-21	¹⁰ Be	142948	3935	12689	154172	4256	13805	144388	3976	12891	
RC20-22	¹⁰ Be	130620	2179	11202	140785	2354	12182	131858	2200	11378	
Mean (n = 6): 131.28 ka; 1σ std: 11.05 ka; 1σ internal: 7.45 ka; 1σ internal + PR%: 13.32 ka											
Oldest cobble (n=6): 146.62 ka; 1σ internal: 4.33 ka;											
Uncertainty weighted mean (n = 6): 128.88 ka; 1σ std: 9.03 ka											
MSWD: 11.36 > k: 2.26 (n=6); Peak age (n=6): 121.96 ka											
RC II moraine boulders											
RC20-02	¹⁰ Be	125486	3356	11064	135098	3622	12012	126630	3388	11230	
RC20-03	¹⁰ Be	108441	1630	9217	115730	1742	9917	109171	1641	9335	Yes
RC20-04	¹⁰ Be	161918	3593	14192	174602	3887	15448	163599	3632	14426	
	²⁶ Al	143604	5367	14309	151277	5676	17887	141129	5268	15174	
RC20-05	¹⁰ Be	148589	2962	12902	160552	3210	14070	150344	2998	13134	
	²⁶ Al	137464	5353	13743	145113	5673	17185	135316	5264	14587	
RC20-10	¹⁰ Be	79902	1613	6830	84106	1699	7241	80260	1620	6900	Yes
RC20-11	¹⁰ Be	107548	2134	9245	114817	2283	9949	108339	2151	9368	Yes
Mean (n = 6): 121.98 ka; 1σ std: 29.92 ka											
Mean (n = 3): 145.33 ka; 1σ std: 18.43 ka; 1σ internal: 5.74 ka; 1σ internal + PR%: 13.55 ka											
Uncertainty weighted mean (n = 3): 145.12 ka; 1σ std: 14.70 ka											
MSWD: 28.73 > k: 3.0 (n=3); Peak age (n=3): 148.61 ka											
MSWD: 28.15 > k: 3.0 (n=3); Peak age (n=3): 148.61 ka											

470

475

480

485

490

495

500

505 Table footnotes. ^{10}Be and ^{26}Al exposure ages were calculated using “the online calculator formerly known as the CRONUS-Earth
online calculator version 3” (Balco *et al.*, 2008). Rock density is assumed to be 2.65 g cm^{-3} and elevation flag is STD. All samples were
collected in 2020, thus considered the year of reference for “before present” (BP). ^{10}Be ages reported here are calculated using the
central Patagonia production rate (Kaplan *et al.*, 2011) while ^{26}Al ages are here calculated using the global ^{26}Al production rate of
Borchers *et al.* (2016). AMS standardizations employed for calculator input data are NIST_27900 (^{10}Be) and Purdue Z92-0222 (^{26}Al).
510 Scaling schemes: St is the time-independent version of Lal (1991) and Stone (2000), Lm is the time-dependent version of Lal (1991)
and Stone (2000), and LSDn is the time-dependent scheme of Lifton *et al.* (2014). Ages are reported with 1σ analytical and external
uncertainties, the latter including production rate and scaling uncertainties. Summary statistics were calculated for each dated
landform using only ^{10}Be exposure ages and LSDn scaling. This includes arithmetic means with 1σ standard deviations (std), 1σ
propagated (from individual ages) internal uncertainties, and propagated 1σ internal plus production rate uncertainty (PR%).
515 Summary statistics for sets of outwash surface cobble ages also display the oldest cobble exposure age; here considered a better
minimum-age estimate of outwash deposit formation. Summary statistics also include uncertainty-weighted means and 1σ standard
deviations, MSWDs and Peak ages; calculated using standard equations as described by Jones *et al.* (2019). Ages in bold are here
interpreted as the most appropriate summary ages per landform and are the ones used throughout the paper for discussion.

520

525

530

535


540

4.3 TCN exposure-age scatter sources

We consider post-depositional shielding of TCN exposure dating samples by vegetation and/or snow to be negligible given the semi-arid and windy conditions (annual 1970-2000 mean wind speed of $\sim 5.3 \text{ m s}^{-1}$; WorldClim 2.1 data; Fick and Hijmans, 2017) that characterise the study site and more generally the Argentinian steppe foreland (Garreaud *et al.*, 2013; Hein *et al.*, 2009; 2017). Modern daily winter (June-August) precipitation is estimated at 2.2 mm near the sampling site (Fick and Hijmans, 2017), which would represent between 20 and 50 cm thick snowfalls given various snow densities. Snow covers of between 75-150 cm thick persisting for four months of the year are typically required to reduce nuclide production by $\sim 5\%$ (snow density range: $0.16\text{-}0.33 \text{ g cm}^{-3}$; Dunai, 2010). Snow accumulation at the study site is thus considered too low to impact exposure ages beyond analytical uncertainties on 10^5 yr timescales. Cosmogenic nuclide inheritance from previous exposure has been shown to be negligible in other eastern Patagonian valleys (*e.g.* Douglass *et al.*, 2007; Hein *et al.*, 2009) and is considered **minimal/unlikely** for our samples based on the long ($>80 \text{ km}$) distance separating glacial deposits and source regions. Such transport distance should have enabled efficient glacial erosion of transported clasts. For moraine boulder samples, we expect the sources of exposure-age scatter to originate mainly from rock surface erosion and boulder exhumation through moraine surface deflation and lateral creep (Putkonen and Swanson, 2003; Briner *et al.*, 2005; Hein *et al.*, 2017).

For all outwash surface cobbles sampled, total rock-surface erosion is considered negligible due to same reasons as described for moraine boulder samples, but also due to the fluvially-rounded and polished nature of target cobbles. Such interpretation is further supported by the analysis of ^{10}Be concentrations in a proglacial outwash depth-profile of MIS 8 - old sediments deposited more than 65 km east of the closest bedrock source region, in an eastern Patagonian setting similar to our study site (Hein *et al.*, 2009), the fluvially-rounded nature of target samples. Results from this analysis indicate that nuclide inheritance is negligible in outwash deposits of the Río Blanco and Hatcher units, in the Lago Pueyrredón valley (47.5°S). We do, however, expect exposure-age scatter to reflect cobble exhumation via a combination of outwash surface deflation (Hein *et al.*, 2009; Darvill *et al.*, 2015) and near surface turbation (*e.g.* cryoturbation) caused by possible local development of soil permafrost during cold intervals (Trombotto, 2008). We thus assume post-depositional disturbance to predominantly cause cobble exhumation and young apparent exposure ages (Phillips *et al.*, 1990; Hein *et al.*, 2011), and thus consider the oldest cobble exposure age a better minimum-age estimate of outwash-plain stabilisation following glacier-front retreat, with the exception of obvious statistical outliers (Table 2, Fig. 4).

570



610 Figure 4. (A): Kernel density plots, adapted from IceTEA tools for exposure dating outputs (Jones *et al.*, 2019) and summary statistics for dated landforms. Thick red curve represent the summed probability distribution for the exposure-age population, excluding outliers, while thin red curves depict Gaussian curves for individual samples. Outliers are denoted by grey dashed curves. Vertical black and purple dashed lines denote the population's arithmetic mean and peak ages, respectively. (B): Same kernel density plots as (A) but visualised on single time scale to enable better comparison of different landform ages and relative exposure-age scatter between surface outwash cobble and moraine boulder samples. On both panels, white and blue vertical bands differentiate Marine Isotope Stages (after Lisiecki and Raymo, 2005), Panel B also features the global LR04 stack of benthic $\delta^{18}\text{O}$ records of Lisiecki and Raymo (2005); commonly used to define the timing of Marine Isotope Stages, here also denoted by alternating white and blue vertical bands.

620 5 Discussion

5.1 TCN exposure age interpretations

5.1.1 The Tecka/RC 0 outwash cobble exposure ages

625 Surface cobbles from the Tecka/RC 0 outwash (Fig. 1c) yield a mean exposure age of 268.7 ± 14.4 ka (n=4), excluding one statistical outlier with an age of 311.5 ± 8.1 ka, and display a discernible amount of exposure-age scatter (MSWD: $4.38 > k$). Following the rationale outlined in Sect. 4.3., we consider the outwash's oldest cobble (284.1 ± 7.0 ka) as the closest minimum-age estimate for the RC 0 advance (Figs. 4, 5). The relatively tight cluster of remaining exposure ages ($284 \pm 7 - 252 \pm 6$ ka) supports this interpretation (Fig. 4).

630 HereAt the sampling sites, the outwash terrace sampled-featuresurface displays preserved braided meltwater channels that demonstratesuggest minimal outwash surface deflation post deposition. We therefore assess the possible impact of cryoturbation alone on exposure-age underestimation, and estimate broadly the potential magnitude of former cobble exhumation through soil. To do so, we model the constant exhumation of all sampled cobbles (excluding outliers) through a given soil horizon (density: 1.3 g cm^{-3}) and calculate the resulting exposure-age bias caused by cosmic-ray attenuation with depth (Gosse and Phillips, 2001; see supplementary materials). We acknowledge that cobble upfreezing is unlikely to have been continuous and of similar magnitude for all cobbles. We assume that some of this variability is considered by reporting a population mean exposure age post-simulation that matches the original age of the oldest cobble, although this assumption undoubtedly yields uncertainties. Using such simulations, we calculated that a soil thickness of ~ 12 cm would have been
635
640 required for the Tecka/ RC 0 cobbles to yield a mean ^{10}Be exposure age (283.1 ± 15.0 ka: +5.7%) that approximately matches

that of the oldest original surface cobble. This estimated magnitude of cobble upfreezing is plausible given our observations of modern soil thicknesses (0-20 cm) near to the sampling locations. Therefore, despite local semi-arid conditions, cobble exhumation due to cryoturbation through a realistic soil thickness can potentially explain the observed exposure-age scatter (Fig. 5). Such results may also support the hypothesis that outwash surface erosion and deflation was minimal at this location.

645


In summary, our interpretations suggest that the exposure ages from the Tecka outwash place the timing of the RC 0 advance at around 290-270 ka, thus during the Marine Isotope Stage (MIS) 8 (300-243 ka; Lisiecki and Raymo, 2005) interval (Figs. 4,6).

650

655

660

665



670 **Figure 5. Illustration of the impact of combined outwash-plain surface deflation and near-surface turbation on exposure-age scatter for outwash surface cobbles sampled. The relationship between depth and nuclide concentration is illustrated by the decay curve (purely schematic) on the bottom left-hand corner of the diagram, which demonstrates that radionuclide production in rock minerals is reduced with depth below surface due to cosmic-ray attenuation. The attenuation effect causes cobbles that have experienced some exhumation post-deposition, due to the effect of upfreezing (dashed vertical arrows; scenario 1), to display exposure-age scatter causing landform deposition age underestimation. Cobbles that remained near the outwash surface since deposition (scenario 2) will display less/no scatter and better estimate the true minimum outwash deposition age. Cobble upfreezing is considered to mainly**

675

680 occur within the soil horizon (brown-coloured) due to the impact of potential permafrost formation. Soil thickness is locally estimated to reach a maximum of ~20 cm. The outwash surface photograph focuses on one of the surface cobble sampled (yellow circle), and enables the assessment of surface cobble density, roundness, size, vegetation type and cover density that are characteristic of outwash plain surfaces sampled within the context of this investigation.

5.1.2 The RC I outwash cobbles and timing of the RC I outermost advance

685 Surface cobbles from the RC I outwash are poorly clustered (MSWD: $33.6 > k$) and afford a mean exposure age of 214.0 ± 29.5 ka and an oldest cobble age of 257.4 ± 6.6 ka (Table 2, Fig. 4). As previously justified (Sect. 4.3.), we consider the population's oldest cobble (257.4 ± 6.6 ka) as the closest minimum-age estimate for the RC I outwash deposition. Simulations of constant cobble upfreezing through soil (Sect. 5.1.1.) demonstrate that a mean exposure age (257.2 ± 35.3 ka: +20%) similar to that of the oldest original cobble is obtained when simulating constant exhumation through a ~39 cm-thick soil horizon. Although reasonable, this average soil thickness is greater than modern observations (0-20 cm). For these cobbles, three distinct outwash surface locations were sampled. They display near identical surface morphologies, vegetation cover and apparent modern soil thicknesses (Fig. 2d). It is thus challenging to assess whether spatially variable cryoturbation could have contributed to the observed scatter. This may indicate that, additionally to cobble upfreezing, outwash surface deflation played, in this instance, an important role in causing significant exhumation of certain RC I cobbles post deposition. This hypothesis is supported by the observation that RC I cobble exposure ages display three distinct age groups that correspond with the three different sampling locations (Table 1, SM figure 3). The first group (sample RC20-16) yields an exposure age of 257.4 ± 6.6 ka, while the second (RC20-12 and 13) and third (RC20-14 and 15) groups display mean exposure ages of 222.4 ± 20.2 ka and 191.5 ± 9.3 ka, respectively. We note that the 1σ standard deviations associated with these age groups do not overlap. Thus, it seems plausible that outwash surface deflation was greater at the locations of the second and third sample groups and, combined with clast upfreezing, could therefore explain the substantial exposure-age scatter observed in the ^{10}Be age distribution of the RC I outwash cobbles.

700 An alternative hypothesis to explain the large RC I exposure-age scatter would be to argue that the RC I outwash could be a composite glaciofluvial deposit formed by meltwater runoff associated with numerous distinct RC outlet-glacier advances. While we commonly expect younger glaciofluvial sedimentation to bury previous deposits, certain cobbles associated with the oldest advances could have been remobilised by either meltwater or ice during younger advances and re-deposited towards the outwash surface. The overlap between the youngest RC I outwash cobble age (185.0 ± 3.0 ka) and the deglacial ^{10}Be age (185.7 ± 5.0 ka) obtained from the ice-moulded bedrock surface sampled directly inboard of the RC I limit could support this hypothesis. Moreover, the RC I moraine belt is a wider (~4 km) complex constituted of more ice-contact hummocks and hummocky ridges suggestive of a greater glaciogenic sediment volume than the RC 0 and RC II moraine complexes (Fig. 6a). This geomorphological distinction could signal that a higher number of individual outlet-glacier advances have reached the RC I margin.

Establishing which of the previous hypotheses are more likely to be valid would require a more detailed chronological and sedimentological analysis through a depth profile of the RC I outwash deposit. However, regardless of which mechanism is responsible for the observed exposure-age scatter, we argue the oldest exposure age from the RC I outwash suggests a separate extensive advance of the RC outlet glacier occurred at around 270-245 ka, thus during the latest stage of the MIS 8 interval (Figs. 4,6).

5.1.3 The RC I H-sampled bedrock surface and timing of the innermost RC I advances

The sampled ice-moulded bedrock's position and elevation (1026 m *a.s.l.*) suggests it was exposed after deglaciation from the RC I moraine-outwash complex, and prior to the RC II advance (Fig. 1c, 6a). Its ^{10}Be age of 185.7 ± 5.0 ka is thus interpreted as a minimum-age estimate of ice-front retreat following the innermost RC outlet-glacier advance reaching the RC I moraine complex. The older RC I outwash-cobble mean exposure age (214.0 ± 29.5 ka) brings further evidence for this stratigraphic relationship. The $^{26}\text{Al}/^{10}\text{Be}$ ratio (6.8 ± 0.3) from this surface bedrock sample does not indicate a complex exposure-burial history, (within uncertainty), thus suggesting the RC glacier did not advance beyond and bury the bedrock for ~~any~~ a prolonged and >100 ka period (SM Fig. 1). This ratio also demonstrates efficient subglacial erosion of the bedrock surface, which created a fresh surface free of inherited nuclides when last ice-covered.

Unlike other samples, we noticed signs of surface erosion on the bedrock outcrop (Fig. 2e). The surface demonstrated a lack of fluvio-glacial polish preservation and showed signs of homogeneous granular disintegration of a depth that proved challenging to quantify in the field. We thus suspect a certain degree of exposure-age underestimation from this bedrock sample. (RC20-01). Applying the rock-surface erosion rate of 0.2 mm ka^{-1} estimated for semi-arid central Patagonia (Douglass *et al.*, 2007; Hein *et al.*, 2017) increases its age by 3% to 191.3 ± 5.3 ka, which is within the 1σ analytical uncertainty of the exposure age. Subsequently, this minimum deglacial exposure age suggests the youngest outlet-glacier advance to have reached the RC I moraine complex had to occur prior to ~190 ka. This interpretation is coherent with RC I outwash cobble exposure ages, and together indicate the RC I complex was most likely deposited prior to the MIS 6 (191-130 ka; Lisiecki and Raymo, 2005) cold interval. Because palaeo-climate proxy records such as Antarctic atmospheric temperature and global SST suggest a return to warmer, near-interglacial conditions between the MIS 7d interstadial (~220-230 ka) and ~190 ka (Fig. 8; Parrenin *et al.*, 2013; Shakun *et al.*, 2015), it is likely possible that the innermost RC I advance occurred during, or prior to the MIS 7d interstadial. In fact, the exposure age of the bedrock surface would date to peak MIS-7d Antarctic cooling (*i.e.* 225 ka; Fig. 8) if a surface erosion rate of 1.07 mm ka^{-1} was applied (LSDn age: 225.1 ± 7.5 ka), which lies within reasonable bounds for rock erosion rates in eastern Patagonia (maximum rate: 1.4 mm ka^{-1} ; Kaplan *et al.*, 2005).

745

To summarise, outwash cobble and bedrock ages suggest that outermost RC I glaciogenic deposits were likely deposited by a MIS 8 glacier advance (~270-245 ka), as portrayed by the oldest outwash cobble age (Figs. 6b,7). On the other hand, the innermost and youngest outlet-glacier advance to have reached the multi-ridge, 4 km-wide RC I moraine complex (Fig. 1c, 6a) may have been significantly younger, and perhaps occurred during the MIS 7d interstadial. However, the latter part of this interpretation remains unclear, due to the lack of chronological data for the innermost RC I deposits combined with the analytical uncertainties and the large scatter displayed by outwash cobble exposure ages from this margin. More chronological constraints are thus required to test this hypothesis.

755

5.1.4 The RC II ~~outwash-cobble and moraine boulder~~ exposure ages

As for the older RC I and Tecka outwash cobble samples, we argue the observed scatter (MSWD: $11.4 > k$) in the RC II outwash cobbles exposure ages (mean: 131.3 ± 11.1 ka) likely originates from cobble exhumation and minor outwash-surface deflation (Fig. 5). This interpretation is reinforced by cobble ages displaying a younger mean age and a tighter cluster than moraine boulders from the stratigraphically-equivalent glacial margin (RC II; Table 2, Fig. 4). Consequently, the oldest surface cobble (146.7 ± 4.3 ka) is deemed a better minimum-age estimate for the RC II glacial advance. We find that a mean exposure age (145.5 ± 12.6 ka: + 11%) similar to that of the oldest cobble can be obtained when simulating constant exhumation of all cobbles (see Sect. 5.1.1.) through a 24 cm-thick soil horizon. This estimate, although highly uncertain, is consistent with modern soil thickness observations near sampling locations (0-20 cm). This simulation suggests that, despite local semi-arid conditions, cobble upfreezing through a realistic soil thickness can cause the observed exposure-age scatter leading to the age underestimation of outwash plain formation. In summary, the RC II outwash cobble ages seem to suggest a minimum-age estimate of ~145 ka. Therefore, we argue the RC II limit reflects an extensive expansion of the local PIS outlet glacier occurring towards the latest stage of ~~the MIS 6 cold interval~~ (Figs. 4,6,7).


770

Following outlier removal ($n = 3$), the RC II moraine boulder mean exposure age ($n = 3$; 145.3 ± 18.4 ka) overlaps the RC II mean outwash cobble age within 1σ analytical uncertainties. The boulder population also displays two ages that are consistent with the set of outwash cobbles (125.5 ± 3.4 ka and 148.6 ± 3.0 ka). These results support our stratigraphic interpretation that these two landforms were deposited by the same outlet-glacier advance. Because we sampled rounded boulders and specifically targeted raised rock fragments presenting ~~fluvio-glacial polish~~ polished surfaces, we here consider boulder surface erosion to

775

be negligible, in contrast to the sampled ice-moulded bedrock surface, where granular disintegration was ubiquitous. The population's oldest boulder (RC20-04: 161.9 ± 3.6 ka) displays a ~15 ka older exposure age than the oldest RC II outwash surface cobble. Cosmogenic nuclide inheritance from pre-glacial-transport exposure is considered unlikely given the >80 km distance to source region and the sub-rounded morphology of the boulders sampled indicative of subglacial clast erosion (Fig. 3b,c). We instead propose this older boulder may reflect at least one earlier glacial advance to the same terminal position, or may have been re-deposited during the RC II advance following exposure during previous glacier advances reaching proximal inboard ice-front positions. The $^{26}\text{Al}/^{10}\text{Be}$ concentration ratio from this sample (6.3 ± 0.3) does not suggest a prolonged period of boulder burial (SM Fig. 1). We therefore hypothesise previous PIS advances might have locally occurred during earlier MIS 6 cold intervals (e.g. MIS 6c,e), while the youngest penultimate MIS 6 glacial maximum is more accurately dated by the RC II surface outwash cobbles, as the glaciofluvial terraces likely remained active until the end of the glaciation.

Overall, our results from the RC II limit ($n = 12$ ages) indicate that outwash surface cobble ages display less post-depositional scatter than moraine boulder ages, with MSWD values of ~11 vs ~30, respectively. This comparison thus agrees with results of Hein *et al.* (2009, 2017) suggesting that well-rounded cobbles sampled from the surface of former proglacial outwash plains are better-suited to numerically date pre-LGC glaciations in semi-arid, undisturbed environments such as the Argentinian steppe of Patagonia. However, even if semi-arid conditions usually limit soil thickness to <20 cm on outwash plains of the Argentinian steppe, our simulations show that potential cobble exhumation due to cryoturbation alone can cause meaningful landform-age underestimations for pre-LGC glacial deposits. This source of post-depositional exposure-age scatter is thus important to consider when interpreting outwash surface cobble exposure ages. Another advantage of targeting outwash cobbles is their abundance in formerly active glaciofluvial environments, while moraine boulders on pre-LGC margins are often sparse and where found, are highly eroded. This enables rigorous selection of cobbles that are most suitable to TCN exposure dating. On the other hand, obtaining exposure ages from both moraine boulders and outwash cobbles can provide additional insight on former glacier activity that may not be evident without the combined approach.



845

Figure 6. (A): Glacial geomorphological map focused on the RC 0 - RC II moraine-outwash sequence around the location of TCN exposure dating samples, highlighted by white dots (with the exception of Tecka/RC 0 outwash samples (Fig. 1c)). Exposure ages displayed are ^{10}Be ages $\pm 1\sigma$ analytical uncertainties, calibrated with LSDn scaling and central Patagonia production rate (Kapan *et al.*, 2011). Ages in red were interpreted as outliers. "Bed" stands for "bedrock" and depicts the ~~RC I~~—RC II ice moulded bedrock surface sampled. Elevation data is provided by the 30-m resolution DEM from the ALOS World 3D missions (AW3D30, version 2.2; JAXA; <https://www.Eorc.jaxa.jp/ALOS/en/aw3d30/>) with a shaded relief background and elevation contour lines at 15-m intervals. (B): Simplified and smoothed elevation profile graph (DEM data: AW3D30) drawn along the black line denoted in panel (A), with TCN exposure dating results (^{10}Be) from the RC 0 - III moraine-outwash dataset after interpretation of the best minimum-age estimate for each dated landform. Palaeo-ice surface elevation and slope gradients are hypothetical and purely schematic. *: Stratigraphic order of events (1: oldest, 5: youngest).

850

855

860 5.2 Comparisons with palaeo-climate and global cryosphere contexts

5.2.1 The MIS 8 glaciations

Our chronology suggests that two of the most extensive PIS advances preserved at the study site (RC 0 and RC I outermost) occurred during MIS 8, at around 290-270 ka, and 270-245 ka, respectively (Figs. 6,7). These expansion events are coeval, within dating uncertainties, with glacial advances recorded by the two largest outlet glaciers of the former PIS in central Patagonia. There, the penultimate glacial advance in both the Lago Buenos Aires valley (46.5°S; Hein *et al.*, 2017; Fig. 1) and the Lago Pueyrredón valley (47.5°S; Hein *et al.*, 2009), are thought to have occurred during MIS 8. Altogether, these three pre-LGC Patagonian TCN chronologies indicate that the MIS 8 glacial events were some of the PIS's most extensive Pleistocene glaciations. These advances likely affected the former PIS across a large latitudinal range, as they can now be traced across multiple valleys from central to northern Patagonia. The geographical ubiquity of these events is diagnostic of a strong climatic forcing, while their timing coincides with some of the lowest middle Pleistocene δD values recorded over Antarctica at around 280-270 ka, associated with maximum MIS 8 Antarctic cooling of -8.5°C relative to pre-industrial modern temperatures (Parrenin *et al.*, 2013; Fig. 8). The RC 0 advance may also be coeval with a severe peak in dust flux recorded in the EDC Antarctic ice core at ~ 270 ka (Lambert *et al.*, 2008; Fig. 8). Isotopic composition analyses have demonstrated that Antarctic ice-core dust concentrations are positively correlated to outwash-plain activity in southern Patagonia, and thus may directly reflect PIS outlet-glacier expansion events (Sugden *et al.*, 2009; Koffman *et al.*, 2021).

870

875

Formatted: English (United Kingdom)

880 During MIS 8, Northern Hemisphere (NH) mid-latitude summer insolation intensity reached two distinct minima at 280 ka and 255 ka (Berger and Loutre, 1991; Fig. 8). These troughs are synchronous with peaks in the SH mid-latitude seasonality curve indicative of colder winters and warmer summers in Patagonia at those times (Fig. 8d; Darvill *et al.*, 2016). Importantly, the signal of NH mid-latitude summer insolation intensity also mirrors orbitally-controlled SH seasonal duration (Fig. 8c; Denton *et al.*, 2021). Hence, Patagonia was experiencing longer, colder winters and shorter, warmer summers around 280 ka and 255 ka. In contrast, SH mid-latitude summer insolation intensity reached maximum values at 280 ka and 255 ka, but 885 minimum values at 292 ka and 268 ka, which are within exposure-age scatter and analytical uncertainties associated with dating of the RC 0 and outermost RC I advances (this study), and of the MIS 8 advances of the Lago Buenos Aires and Lago Pueyrredón outlet glaciers also (Hein *et al.*, 2009; 2017). Therefore, due to current TCN exposure dating uncertainties, it remains a challenge to determine whether the MIS 8 Patagonian glaciation was coeval with a SH, or a NH mid-latitude summer insolation intensity minimum. However, more accurate dating of younger PIS outlet-glacier advances occurring during the 890 global LGM (MIS 2; *e.g.* Peltier *et al.*, 2021), including the RC outlet glacier (RC III-VII; Leger *et al.*, 2021a), has shown that these later SH glacier advances, and also SSTs and Antarctic atmospheric temperature depressions, were anti-phased with SH summer insolation intensity, and were instead coeval with high SH seasonality and winter duration. Consequently, it is plausible that the PIS advanced at ~280 ka (RC 0) and ~255 ka (RC I), when SH summer insolation intensity was high, but when SH winters were cold and long. The proposed timing for the RC 0 advance is consistent with other SH palaeo-climate proxy records which suggest that peak MIS 8 cooling occurred relatively early during the glacial cycle (*i.e.* 280-270 ka; Fig. 8), in contrast with the later MIS 6 and MIS 5d to MIS 2 glaciations, which are characterised by maximum global cooling occurring just prior to sudden glacial terminations (Hughes *et al.*, 2020).

Paradoxically, MIS 8 displays a relatively weak global cooling signal in several records such as benthic $\delta^{18}\text{O}$ (Hughes *et al.*, 900 2020), and is thought to have been characterised by the lowest sea-level depression relative to other middle Pleistocene 100 ka glacial intervals (-93 m; Spratt and Lisiecki, 2016). Moreover, glacial terrestrial records from different regions demonstrate contrasting ice-sheet and glacier behaviours during MIS 8. Indeed, although stratigraphic evidence for MIS 8 glaciations is often elusive, some has been reported for the Fennoscandian and British-Irish ice sheets (Beets *et al.*, 2005; Davies *et al.*, 2012; White *et al.*, 2010; 2017), former ice sheets and glaciers overprinting Poland (Krznanian glaciations; Lindner and Marks; 905 1999), Siberia and Russia (Astakhov *et al.*, 2011; 2016), Central and southern Europe (*e.g.* Preusser *et al.*, 2011), and the West Coast Range of Tasmania (Colhoun and Barrows, 2011; Kiernan *et al.*, 2010). In North America, chronological constraints for stratigraphic evidence of pre-MIS 6 glacial deposits are, in most cases, too imprecise to determine the ~~marine isotope stage~~ [Marine Isotope Stage](#) to which they relate (Hughes *et al.*, 2020). The review by Hughes *et al.* (2020) concludes that robust chronological evidence of MIS 8 glaciations is rare, and, where it is found, is often characterised by margins located inboard 910 of MIS 6 limits. In general, direct chronological evidence of MIS 8 glaciations more extensive than MIS 6-2 advances has only been found in Russia (east of the Urals; Astakhov *et al.*, 2016) and Patagonia. Therefore, the extensive MIS 8 glaciation of Patagonia implies a strong regional cooling signal (Hein *et al.*, 2017).

It has been noted in previous work that the major eastern Patagonian valleys, including the RC valley (Figs. 6,7), feature records of progressively decreasing glacier extents as the glacier advances get younger (Singer *et al.*, 2004; Kaplan *et al.*, 2009; Coge *et al.*, 2018). This pattern may not be entirely climate-related. Instead, it might be caused by a combination of long-term erosion of the accumulation area and focused subglacial erosion along former glacial valleys that resulted in glacial overdeepening (Kaplan *et al.*, 2009). During Quaternary glacial cycles, selective subglacial erosion, shown to be most effective along the flowlines of warm-based PIS outlet glaciers and near the centre of the ice sheet where ice was thickest (Clapperton, 1983), has caused rapid and zonally asymmetric overdeepening of main Patagonian valleys (Rabassa and Clapperton, 1990). In the past ~1.2 Ma, this erosive process has been estimated to cause >1000 m of bedrock elevation lowering in certain Patagonian valleys (Singer *et al.*, 2004). Kaplan *et al.* (2009) argued that such overdeepening would likely have overcome the effects of tectonic- and glacial isostatic adjustment-related uplift. For a given location and identical climate forcing, erosion-driven bed lowering may have led to progressively more negative glacier mass balances during each major PIS expansion event, causing less extensive glacier advances with time. Moreover, if more effective towards the former ice divide, subglacial erosion would also have increased retrograde basal slope gradients and thus augmented basal shear stress during subsequent glacier advances. These processes could therefore partly explain the observed pattern of decreasing outlet-glacier extent through time in eastern Patagonia (Fig. 6). Consequently, although the geographically widespread MIS 8 advances of the PIS were most likely the result of strong climate forcing, their outboard positions relative to MIS 6-2 deposits do not necessarily indicate colder regional conditions during MIS 8 than during the younger advances.

935

940

945




Figure 7. Palaeo-glacial reconstructions for the RC, RH, RF and Lago Palena/General Vintter (LP/LGV) valleys, for each advances/stillstands and deglacial events interpreted in this investigation (A-C). Panel D displays the ice extent associated with the RC III moraine complex, described in detail in companion publication (Leger *et al.*, 2021a). The latter reconstruction includes the formation of glaciolacustrine phase one (here termed PL1; Leger *et al.*, 2021a). Ice-sheet and mountain glacier models were digitised manually in ArcGIS. Apart from when delineated by confidently-mapped moraine limits (*e.g.* the RC, RH and LP/LGV moraine sequences), the geographical location of ice margins are inferred from topography. Topographic data is from the AW3D30 DEM. “OC” and “MB” denote the type of rock samples used for TCN exposure dating of the represented event, and stand for “outwash cobbles”, and “moraine boulders”, respectively. Yellow stars indicate the approximate sample locations for TCN exposure dating of each represented event, while each panel features the number of ^{10}Be exposure ages produced ($n = x$) on the bottom left-hand corner (this excludes ^{26}Al ages). There are no previously published glacier chronologies in the RH, RF and LP/LGV valleys. Relative ice extents in those neighbouring valleys are thus inferred from our RC chronology and cross-valley comparisons of moraine set numbers, preservation and morphostratigraphy. Hence, these inferences yield some uncertainties. Proglacial lake volume estimates were computed from DEM data (AW3D30).

985

5.2.2 The MIS 7₆ glaciation

990

Our chronology provides ineptive evidence that the innermost and youngest advance to have reached the RC I moraine complex may have occurred during MIS 7 (243–191 ka; Lisiecki and Raymo, 2005). Although the resolution of our chronology is not sufficient to assert the following with confidence, we argue that this potential event might relate to interstadial MIS 7d (~235–225 ka), described in detail by Ruddiman and McIntyre (1982, p1273) as an event that “bisected “interglacial” isotopic stage 7, lasted no more than 23,000 yr from inception to end, and resulted in a very large ice volume build-up and a full-glacial climatic response of the North Atlantic Ocean”. Direct terrestrial evidence and robust dating of this event are currently non-existent in the southern mid-latitudes, and hence more empirical data from several locations are required to assess the geographical ubiquity of this glacial expansion event.

995

1000

During the relatively brief MIS 7d interstadial, global atmospheric CO_2 concentrations are estimated to have diminished by ~75 ppm (Bereiter *et al.*, 2015), and the atmospheric temperature depression over Antarctica was of similar magnitude to that recorded during the MIS 8 and MIS 6 stadials (Fig. 8). These records indicate that a 20 ka long Antarctic atmospheric cooling phase, with a peak cooling of approximately $-7.5^\circ C$ relative to pre-industrial modern temperatures, occurred at ~223 ka, 7 ka after NH mid-latitude summer insolation intensity reached the lowest values of the past 1 Ma, at 230 ka (Fig. 8). Therefore, while interstadial MIS 7d was relatively short-lived, it was characterised by record-high winter duration and a peak in local seasonality (colder winters and warmer summers) at the SH mid-latitudes. Antarctic dust concentration records also indicate a significant maximum at 225–220 ka (Lambert *et al.*, 2008; 2012). This implies the MIS 7d interstadial was likely associated with an expansion of the PIS inducing outwash plain activity in southern Patagonia (Sugden *et al.*, 2009).

1005

5.2.3 The penultimate MIS 6 glaciation

1010

Our interpretation of TCN exposure ages from the penultimate RC II margin suggests deposit suggest that a major expansion of the PIS occurred during MIS 6, and towards the latter part of the glacial interval, at around 140–130–150 ka (Figs. 4,6,7). Our chronology suggests that maximum PIS expansion occurred just prior to the rapid global warming leading to MIS 5e, one of the warmer middle-to-late Pleistocene interglacial periods (Fig. 8). This is consistent with global sea-level reconstructions suggesting that the second greatest global ice volume of the past 800 ka, associated with a sea-level minimum of -125 m, was reached towards the end of MIS 6, at ~140 – 135 ka (Spratt and Lisiecki, 2016). At that time, mean atmospheric temperatures

1015


over Antarctica are estimated to have reached their second lowest values of the past 800 ka (-9.3 °C relative to pre-industrial period; Fig. 8). Global cooling during MIS 6 was more persistent than during other middle Pleistocene glacial cycles. Indeed, this cycle is associated with four major minima in Antarctic atmospheric temperatures spread over ~ 50 ka, and occurring at around ~ 185 , ~ 165 , ~ 155 and ~ 140 ka (Parrenin *et al.*, 2013; Fig. 8). As observed for the MIS 8 interval, these peaks in atmospheric cooling occurred a few ka after NH summer insolation intensity minima, associated with SH seasonality and winter duration maxima. Global SSTs and atmospheric CO_2 concentrations remained consistently low for the same 50 ka time-window, between ~ 190 and ~ 140 ka (Bereiter *et al.*, 2015; Shakun *et al.*, 2015). The persistence of the cold phase throughout MIS 6 may have facilitated multiple expansions of the PIS at this time. In our ^{10}Be chronology, this may be reflected by the potential recycling and re-deposition of one RC II moraine boulder sample, yielding a ^{10}Be exposure ages of 162 ± 4 ka, during the younger penultimate advance dated by the RC II outwash cobbles (~ 130 -~~140~~150 ka).

Evidence for several distinct MIS 6 glaciations has been reported from other regions of the world. For the European Ice Sheet, for instance, stratigraphic investigations have led to the dating, and naming, of two glaciations within MIS 6, respectively the Late Saalian Drenthe glaciation at 175-155 ka, and the Warthe Stadial between 150 ka and 135 ka (Toucanne *et al.*, 2009; Margari *et al.*, 2014). More generally, MIS 6 glacial deposits are ubiquitous near former glacial margins and are documented by a wider body of evidence than other pre-LGC glaciations (*e.g.* Putnam *et al.*, 2013; Kiernan *et al.*, 2014; Evans *et al.*, 2019; Fernandes *et al.*, 2021), suggesting that the greatest global ice volume of the middle Pleistocene was likely reached during that interval (Hughes *et al.*, 2020). In Patagonia, however, our chronology is amongst the first published datasets, ~~along with one other ongoing investigation (abstract: Peltier *et al.*, 2017)~~, offering direct and robust dating of a MIS 6 glacial expansion event. Indeed, the well-studied Lago Pueyrredón and Lago Buenos Aires pre-LGC moraine-outwash records do not provide clear chronological evidence for MIS 6 advances (Hein *et al.*, 2009; 2017). However, this may just imply that the MIS 2 advances of the Lago Pueyrredón and Lago Buenos Aires outlet glaciers were more extensive than any MIS 6 glacial events. Therefore, as for the MIS 8 glaciation, MIS 6 expansions of outlet glaciers may have been ubiquitous through the former PIS. However, the current lack of robust pre-LGC glacier chronologies in Patagonia prevents this hypothesis from being tested adequately.

Formatted: List Paragraph

5.2.43 Synthesis: The timing of Patagonian glaciations

Along with companion publications focusing on the LGC (Leger *et al.*, 2020; 2021a), our findings illustrate that the preserved sequence of moraine-outwash complexes in the RC valley system records a minimum of four major PIS expansion events occurring during MIS 8, 6, and 2; ~~and possibly also during MIS 7.~~ In Antarctic atmospheric paleo-temperature proxy records, global SST records, and global atmospheric CO_2 records, the strongest minima between today and 330 ka are all recorded by the RC moraine-outwash complexes, with the exception of the 70-60 ka (MIS 4) cold interval, for which no preserved moraine-outwash complex could be found at the study site. However, we note that a major advance of the PIS near this time did occur on the western side of the Andes at 42°S (García *et al.*, 2021; Gómez *et al.*, 2022). There, Garcia *et al.* (2021) dated glacial sediments on the Isle of Chiloe to 57.8 ± 4.7 ka, or early MIS 3, with a somewhat smaller advance occurring during MIS 2 at 26.0 ± 2.9 ka. The lack of a corresponding MIS 3/4 moraine east of the mountains suggests that the PIS behaved asynchronously between its western and eastern margins at this time. In southeastern Patagonia, more-extensive-than MIS 2 advances were also dated to MIS 4 and 3 using TCN exposure dating (*e.g.* Darvill *et al.*, 2016; García *et al.*, 2018; Peltier *et al.*, 2021). We argue that while the RC outlet glacier most likely advanced during the MIS 4 and 3 cooling events, these intervals were relatively short-lived, impeding the PIS from becoming locally thick enough to allow the RC outlet glacier to advance along its retrograde bed slope and generate an extensive advance. Very likely, in addition, all glaciogenic deposits associated with MIS 3/4 advances may have been eroded during the younger MIS 2 advances. This may suggest that steady cooling over periods >15-20 ka is an important pre-requisite for northeastern PIS outlet glaciers to reach a mass balance positive enough to enable highly extensive advances.



1085

1090

1095

1100

1105

1110

1115

1120 Figure 8. Vertical plot comparing this investigation's TCN ^{10}Be chronology to other Patagonian outlet-glacier TCN chronologies, to
insolation intensity variations, and to palaeo-climate proxy records over the past 350 ka. (A) Stable isotope δ Deuterium record
(orange curve) from the EPICA Dome C (EDC) ice core, 75°S, east-Antarctica and Antarctic atmospheric temperature (overlying
dashed black curve) from a stack of isotopic temperature reconstructions from five different ice cores (EDC, Vostok, Dome Fuji,
1125 TALDICE, and EDML) (Parrenin *et al.*, 2013), with most potent cooling events of the past 350 ka highlighted by black circles. TCN
exposure ages from the RC valley interpreted as best minimum-age estimate for each reconstructed glacial event are displayed (red
dots) with analytical uncertainty (black horizontal bars). (B): Global stack (n = 49 cores) of paired planktonic $\delta^{18}\text{O}$ -derived Sea
Surface Temperature (SST) data with Marine Isotope Stage numbers indicated (Shakun *et al.*, 2015). (C): Dark green line: summer
(mid-June) insolation intensity for 60°N (Berger and Loutre, 1991). Dashed black curve: seasonal duration at 60°S (insolation
1130 threshold = 300 W m⁻²) after Huybers and Denton (2008) and Denton *et al.* (2021). (D): Southern Hemisphere seasonality at 30°C,
40°S, and 60°S calculated by subtracting June from December insolation intensity values (Berger and Loutre, 1991), such that
increasing seasonality indicates colder winters and warmer summers (after Darvill *et al.*, 2016). (E): Atmospheric CO₂ concentrations
composite record from several Antarctic ice cores including EDC, WAIS, Vostock, Siple Dome, TALDICE, EDML, and LAW Dome
(Bereiter *et al.*, 2015). (F): Dust flux record from the EDC ice core (Lambert *et al.*, 2008). (G-I): Kernel density distribution curves
(1 σ) for TCN exposure ages from the RC O – RC II moraine-outwash records (this study: G), the Hatcher moraine-outwash complex,
1135 in the Lago Cochrane/Pueyrredón valley (Hein *et al.*, 2009; H), and the Moreno 2/3 moraine-outwash limits, in the Lago General
Carerra / Buenos Aires valley (Hein *et al.*, 2017; I).

1140

5.3 Climatic and orbital drivers of SH glaciations - hypotheses

Our chronology implies that the most extensive middle Pleistocene PIS expansion events preserved match the strongest minima
in Antarctic atmospheric temperatures (Parrenin *et al.*, 2013), global oceanic SSTs (Shakun *et al.*, 2015), and global
1145 atmospheric CO₂ concentrations (Bereiter *et al.*, 2015) (Fig. 8). They also often appear to occur a few ka after around the timing
of minima in NH summer insolation intensity (60°N) and maxima in SH seasonality, while being out-of-phase with mid-
latitude SH summer insolation intensity (Fig. 8e)-8c). However, one must note that this statement can only be advanced with
confidence for the local MIS 2 expansions of the PIS. For the local MIS 8 and MIS 6 glaciations, this observation is based on
current knowledge of ^{10}Be production rates and the assumptions made in this paper, and does not take into account the full
1150 exposure-age range covered by dating analytical uncertainties. The first observations of the asynchrony between SH mid-
latitude glacier advances/recessions and local summer insolation intensity (*e.g.* Mercer, 1976; Denton *et al.*, 1999) encouraged
the hypothesis that major glacial-termination-inducing warming signals were primarily controlled by NH summer insolation
intensity driving NH glacial mass balance, and were propagated from the northern to the southern hemisphere through oceanic
and atmospheric circulation shifts (*e.g.* Broecker, 1998; Denton *et al.*, 2010). Some of these interhemispheric climate-
1155 transmission mechanisms, such as the impact of the abrupt weakening of the Atlantic meridional overturning circulation in the

north Atlantic following the last deglaciation, have been well documented (Barker *et al.*, 2009; Denton *et al.*, 2010). However, there is still a great deal of debate regarding the dominant trigger mechanisms responsible for global ice-sheet expansion/retreat cycles (Putnam *et al.*, 2013). Indeed, the above hypothesis entails the dilemma that while NH summer insolation intensity would be a key driver of NH ice-sheet mass balance, it would instead have little impact on mass fluctuations of SH mid-latitude ice sheets and glaciers (Denton *et al.*, 2021). This problem invites the consideration of alternative hypotheses, such as the potential roles of SH seasonality and seasonal duration on controlling global ice-sheet mass fluctuations.

García *et al.* (2018) noted that the extensive advances of the Torres del Paine (51°S) and the Última Esperanza (51.5°S) outlet glaciers at 48 ka (MIS 3) coincide with a minimum in obliquity-modulated high-latitude winter insolation intensity. The authors therefore hypothesise that colder SH winters during periods of high obliquity may favour SH mid-latitude glaciation by promoting local atmospheric cooling, Southern Ocean stratification and expanded Antarctic sea ice. Since we observe that middle-to-late Pleistocene PIS expansion events appear to occur during periods of longer, colder winters and shorter, warmer summers, we suggest that other PIS chronologies (including this study) may support García *et al.* (2018)'s hypothesis.

Alternatively, the “Zealandia Switch” hypothesis (Denton *et al.*, 2021) focuses on the SH seasonal-duration aspect of orbitally-controlled insolation as a potential driver of late-Quaternary glaciation. By pacing the seasonal cycle of weakened and northward-migrated [SWW Southern Westerly Winds](#) during winter versus strengthened and southward-migrated [SWW Southern Westerly Winds](#) during summer, this hypothesis implies that SH seasonal duration might be a key driver of global climate change during middle-to-late Pleistocene glacial/interglacial cycles. Using global climate model simulations, the authors show that a reduced SH summer duration and increased SH winter duration could cause millennial-scale equatorward migrations of the [SWW Southern Westerly Winds](#) and its coupled Antarctic Circumpolar Current. Such zonal wind configuration would induce a northward migration of the subtropical front and a latitudinally-restricted subtropical gyre system in the Indo-Pacific Ocean basins (*ibid*). The bathymetry of the Australia/Zealandia continent is such that a restricted Indo-Pacific subtropical gyre could decrease Indonesian Throughflow and restrict westward Agulhas Leakage into the South Atlantic Ocean (Lorrey and Bostock, 2017). A reduced Agulhas Leakage has implications for the Atlantic salt input resulting in a potential weakening of Atlantic meridional overturning circulation (Caley *et al.*, 2012); [De Boer et al., 2013](#)). The authors thus argue that such a southern oceanic and atmospheric circulation configuration could promote a global glacial mode (Denton *et al.*, 2021) that would greatly affect SH and thus Patagonian glacier mass balance.

Moreover, variations in the intensity and latitudinal position of the [SWW Southern Westerly Winds](#) influence Southern Ocean upwelling of deep-stored CO_2 and nutrients (Anderson *et al.*, 2009; Ai *et al.*, 2020). This mechanism is thought to represent a major contribution to variations in global atmospheric CO_2 concentrations over millennial timescales (Sarmiento and Toggweiler, 1984). Former equatorward shifts and weakening of the [SWW Southern Westerly Winds](#) could have decreased Southern Ocean upwelling and enhanced CO_2 storage in the deep ocean, thus causing delayed global cooling amplification

1190 (Sime *et al.*, 2013). Conversely, a poleward migration and strengthening of the SWWSouthern Westerly Winds is thought to
amplify global warming via increasing Southern Ocean upwelling, CO_2 outgassing and increasing CO_2 atmospheric
concentrations. This hypothetical CO_2 capacitor storage-and-release mechanism may explain the highly amplified warming
observed during major glacial terminations (Anderson *et al.*, 2009; Denton *et al.*, 2021). This positive feedback mechanism
may partly justify why middle-to-late Pleistocene global oceanic and atmospheric temperature variations often reached their
1195 most extreme values with some notable delay (a few ka) relative to NH mid-latitude summer insolation intensity.

To summarise, empirical data including this study suggest that major middle-to-late Pleistocene cycles of SH mid-latitude
glaciers and ice sheets appear to be coeval with NH glaciers and ice sheets, but out-of-phase with local mid-latitude summer
insolation intensity (Putnam *et al.*, 2013; Peltier *et al.*, 2021), traditionally thought to be the primary driver of middle-to-late
1200 Pleistocene glacial expansion/demise cycles (Mercer, 1984). However, recent hypotheses suggest that this paradox might be
explained by the potentially more dominant climate-forcing effect of the seasonality and seasonal-duration expressions of SH
insolation, through their impact on atmospheric temperature, the position of the SWWSouthern Westerly Winds belt and
Subtropical Front, Antarctic sea ice extent, and Southern Ocean stratification and CO_2 storage-and-release feedback
mechanisms. Testing the above hypotheses, and determining which of seasonality versus seasonal duration played a primary
1205 role in driving SH climate and glacial variations during the middle-to-late Pleistocene, represent key avenuesremains a major
challenge and represents a key avenue for future research.

1210

1215

1220

6 Conclusions

- 1225 • We provide geomorphological and direct chronological evidence for at least three extensive, stratigraphically distinct pre-Last Glacial Cycle advances of the Río Corcovado glacier, a major outlet of the former northern Patagonian Ice Sheet. TCN surface exposure dating of outwash surface cobbles yields best ~~minimum~~-age ~~range~~ estimates for the RC 0, RC I and RC II advances of ~~284 ± 7290-270~~ ka, ~~257 ± 7270-245~~ ka, and ~~147 ± 4130-150~~ ka, respectively.
- 1230 • Together with evidence we have reported in Leger *et al.* (2021a, b), our complete chronology reveals a minimum of four distinct expansion events of northeastern Patagonian Ice Sheet outlet glaciers that occurred during the MIS 8, MIS 6, and MIS 2 intervals. ~~Further results gathered from bedrock surface and moraine boulder exposure ages also provide ineptive evidence for another MIS 7 expansion event and at least another MIS 6 advance previously reaching a close extent to that of the penultimate RC II advance.~~
- 1235 • Our dataset presents the first robust direct dating evidence of MIS 6 glaciations in Patagonia.
- 1240 • We did not find evidence for MIS 4 and 3 deposits in the RC valley geomorphological record. However, extensive glaciations did occur at these times on the former western margin of the ~~Jee-Sheet~~ice sheet, and in other, more southern Patagonian regions. This highlights longitudinal and latitudinal asynchronies in former Patagonian Ice Sheet mass balance during these stages. We suggest that MIS 4 and 3 outlet-glacier advances likely occurred in northeastern Patagonia, but were of relatively shorter extent, and their associated deposits were likely eroded by the younger, more extensive MIS 2 glaciation.
- 1245 • We interpret that major middle-to-late Pleistocene Patagonian Ice Sheet expansions ~~may have~~ occurred in synchrony with the greatest cooling events recorded in global SST and Antarctic atmospheric temperatures, with the most prominent peaks in Antarctic ice-core dust concentrations, and with major minima in global CO_2 atmospheric concentrations. We observe a particularly good correlation between Antarctic atmospheric temperature signals and southern mid-latitude ice-sheet volume fluctuations, over several glacial cycles.
- 1250 • Our results suggest that Patagonian glaciations were more extensive during MIS 8 than during MIS 6-2, while global ice volumes and sea level depression were not particularly high during MIS 8. This implies ~~that there was an~~ interhemispheric asynchrony in the relative volume and extent of Earth's major ice sheets during MIS 8. However, we acknowledge that a larger MIS 8 eastern Patagonian Ice Sheet is not necessarily indicative of a more potent local climate-forcing event than
- 1255

during younger glaciations (MIS 6-2) as glacial erosion, which greatly influences the local topography, may play a role in controlling ice-sheet extent.

- Major middle-to-late Pleistocene advances of the Patagonian Ice Sheet appear to have occurred out-of-phase with local summer insolation intensity but in synchrony with orbitally-controlled periods of longer and colder winters. We hypothesise that seasonality and seasonal duration may exert more control over southern mid-latitude ice-sheet mass balance than mid-latitude summer insolation intensity over millennial timescales. This may be a consequence of the possible effect of seasonal duration variability on the strength and migration of the [SWW Southern Westerly Winds](#), and the impact of these migrations on global climate and Southern Ocean circulation and CO_2 upwelling, as proposed by the “*Zealandia Switch*” hypothesis.
- Our findings corroborate work by Hein *et al.* (2009; 2017) and indicate that, in eastern Patagonia, pre-LGC outwash surface cobbles yield exposure ages displaying less post-depositional scatter than moraine boulders from the same margins, and hence facilitate more accurate estimations of the approximate timing of pre-LGC glacial advances.

1260

1265

1270

1275

1280

1285

Data availability.

All data associated with the production of new ^{10}Be and ^{26}Al TCN exposure ages, including sample characteristics and preparation protocols, are provided in the manuscript tables, figures, and the Supplementary Materials. Further inquiries can be directed to the corresponding author.

1290

Data online repository.

The data needed to re-calculate the ^{10}Be and ^{26}Al TCN exposure ages presented in this paper using the "online calculator formerly known as the CRONUS-Earth online calculators" version 3 is freely available online, as long as original publication is cited when used and/or referred to. This includes a textfile with data blocks in the correct input format and arranged by landforms studied in this investigation. The data are located in an open-access Mendeley Data online repository, accessible using this DOI: <https://doi.org/10.17632/gg4b3sh9k2.1>.

1295

Team list.

ASTER Team: Georges Aumaître (Aix-Marseille Université, CNRS, Coll. France, IRD, INRAE, CEREGE, Aix en Provence, 13545, France), Fawzi Zaidi (Aix-Marseille Université, CNRS, Coll. France, IRD, INRAE, CEREGE, Aix en Provence, 13545, France) and Karim Keddadouche (Aix-Marseille Université, CNRS, Coll. France, IRD, INRAE, CEREGE, Aix en Provence, 13545, France).

1300

1305

Author contributions.

T.P.M.L co-concieved the study with A.S.H. raised the finances for fieldwork, conducted fieldwork with help from P.T.G, wrote NERC NEIF grant application together with A.S.H, conducted result analyses and interpretations, wrote the manuscript and supplementary materials with feedback from A.S.H. and R.G.B primarily, and A.R., I.S, and D.F. in a second time, and generated maps and figures. TCN exposure-age analyses including wet chemistry and AMS measurements were conducted by A.R and D.F at SUERC and T.P.M.L, I.S and the ASTER Team at CEREGE and the ASTER AMS.

1310

Competing interest.

We (the authors) hereby declare that this scientific investigation presents no known competing financial interests or personal conflicts influencing research output.

1315

Disclaimer.

Formatted: French (France)

1320

Acknowledgements.

We express our gratitude towards all individuals who contributed to the <https://crowd.science/> crowdfunding campaign that provided us with the means to conduct a field expedition to the study site in January and February 2020. We thank all local landowners who authorized access and sampling on their properties, in particular, the Estancia Tecka (Chubut province, Argentina) for enabling access to their land and private roads. We thank Elaine McDougall and Allan Davidson for their invaluable laboratory work enabling efficient quartz isolation and purification from our rock samples. [Finally, we are thankful to the two anonymous reviewers for providing valuable feedback and comments which greatly improved the manuscript.](#)

1325

1330

Financial support.

This study was conducted within the context of a University of Edinburgh E³ Doctoral Training Partnership Ph.D. studentship (award code: NE/L002558/1) awarded by the Natural Environment Research Council (NERC) and the University of Edinburgh School of GeoSciences to T.P.M.L. Fieldwork (01/02 2020) was partly supported by a crowd-funding campaign through the Crowd.Science fundraising platform (<https://crowd.science>) and a British Society for Geomorphology Postgraduate Research Grant award (BSG-2019-04) awarded to T.P.M.L. TCN exposure dating laboratory analysis and AMS measurements were funded by a NERC National Environmental Isotope Facility (NEIF) grant (2245-0320) awarded to A.S.H. and T.P.M.L. in July 2020.

1335

1340

Review statement.

1345

1350

References.

1355 [Ai, X. E., Studer, A. S., Sigman, D. M., Martínez-García, A., Fripiat, F., Thöle, L. M., Michel, E., Gottschalk, J., Arnold, L., Moretti, S., Schmitt, M., Oleynik, S., Jaccard, S. L., and Haug, G. H., 2020., Southern Ocean upwelling, Earth's obliquity, and glacial-interglacial atmospheric CO₂ change: Science, v. 370\(6522\), p. 1348–1352, doi: <https://doi.org/10.1126/science.abd2115>, 2020.](#)

Formatted

1360 [Anderson, R. F., Ali, S., Bradtmiller, L. I., Nielsen, S. H. H., Fleisher, M. Q., Anderson, B. E., and Burckle, L. H., 2009., Wind-driven upwelling in the Southern Ocean and the deglacial rise in atmospheric CO₂: Science, v. 323\(5920\), p. 1443–1448, doi: <https://doi.org/10.1126/science.1167441>, 2009.](#)

Formatted: Font: Not Bold, Font color: Black

Formatted

Formatted: Bibliography

Formatted: Font color: Black

1365 [Astakhov, V., 2011. Ice margins of northern Russia revisited. In: Ehlers, J., Gibbard, P.L., Hughes, P.D. \(Eds.\), Quaternary Glaciations - Extent and Chronology: A Closer Look. Developments in Quaternary Sciences 15. Elsevier, Amsterdam, pp. 1–14, 2011.](#)

Formatted: Font: Not Bold

Formatted: Left

[Astakhov, V., Shkatova, V., Zastrozhnov, A., and Chuyko, M., 2016., Glaciomorphological map of the Russian Federation: Quaternary International, v. Quat. Int., 420, p. 4–14, doi: <https://doi.org/10.1016/j.quaint.2015.09.024>, 2016.](#)

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

1370 [Augustinus, P., Fink, D., Fletcher, M. S., and Thomas, I., 2017., Re-assessment of the mid to late Quaternary glacial and environmental history of the Boco Plain, western Tasmania: Quaternary Science Reviews, v. Quat. Sci. Rev., 160, p. 31–44, doi: <https://doi.org/10.1016/j.quascirev.2017.01.015>, 2017.](#)

Formatted: Font color: Black

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1375 [Balco, G., 2006., Converting Al and Be isotope ratio measurements to nuclide concentrations in quartz, Documentation—Be-10-Be/26-Al exposure age calculator, 2006.](#)

Formatted: Font: Not Bold

[Balco, G., and Rovey, C. W., 2008., An isochron method for cosmogenic-nuclide dating of buried soils and sediments, Am. J., American Journal of Science, v. Sci., 308\(10\), p. 1083–1114, doi: <https://doi.org/10.2475/10.2008.02>, 2008.](#)

Formatted

Formatted

1380 [Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J., 2008., A complete and easily accessible means of calculating surface exposure ages or erosion rates from 10Be and 26Al measurements: Quaternary geochronology, v. Quat. Geochronol., 3\(3\), p. 174–195, doi: <https://doi.org/10.1016/j.quageo.2007.12.001>, 2008.](#)

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1385

Barker, S., Diz, P., Vautravers, M. J., Pike, J., Knorr, G., Hall, I. R., and Broecker, W. S., 2009. Interhemispheric Atlantic seesaw response during the last deglaciation. *Nature*, v. 457(7233), p. 1097–1102, doi: <https://doi.org/10.1038/nature07770>, 2009.

Formatted

1390

Barrows, T. T., Stone, J. O., Fifield, L. K., and Cresswell, R. G., 2002. The timing of the last glacial maximum in Australia. *Quaternary Science Reviews*, v. 21(4-3), p. 159–173, doi: [https://doi.org/10.1016/S0277-3791\(01\)00109-3](https://doi.org/10.1016/S0277-3791(01)00109-3), 2002.

Formatted: Font: Not Bold, Font color: Black

Formatted

Formatted: Bibliography

Formatted: Font color: Black

Beets, D. J., Meijer, T., Beets, C. J., Cleveringa, P., Laban, C., and Van der Spek, A. J. F., 2005. Evidence for a Middle Pleistocene glaciation of MIS 8 age in the southern North Sea. *Quaternary International*, v. 133, p. 7–19, doi: <https://doi.org/10.1016/j.quaint.2004.10.002>, 2005.

Formatted

1395

Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H., Kipfstuhl, S., and Chappellaz, J., 2015. Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present: *Geophysical Research Letters*, v. 42(2), p. 542–549, doi: <https://doi.org/10.1002/2014GL061957>, 2015.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1400

Berger, A., and Loutre, M. F., 1991. Insolation values for the climate of the last 10 million years. *Quaternary Science Reviews*, v. 10(4), p. 297–317, doi: [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.

Formatted

1405

Borchers, B., Marrero, S., Balco, G., Caffee, M., Goehring, B., Lifton, N., Nishiizumi, K., Phillips, F., Schaefer, J., and Stone, J., 2016. Geological calibration of spallation production rates in the CRONUS-Earth project. *Quaternary Geochronology*, v. 31, p. 188–198, doi: <https://doi.org/10.1016/j.quageo.2015.01.009>, 2016.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1410

Briner, J. P., Kaufman, D. S., Manley, W. F., Finkel, R. C., and Caffee, M. W., 2005. Cosmogenic exposure dating of late Pleistocene moraine stabilization in Alaska. *Geological Society of America Bulletin*, v. 117(7-8), p. 1108–1120, doi: <https://doi.org/10.1130/B25649.1>, 2005.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1415

Broecker, W. S., 1998. Paleocan circulation during the last deglaciation: a Last Deglaciation: A bipolar seesaw? *Paleoceanography*, v. 13(2), p. 119–121, doi: <https://doi.org/10.1029/97PA03707>, 1998.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Caldenius, C. C. Z., 1932. Las glaciaciones cuaternarias en la patagonia y tierra del fuego. *Geografiska Annaler*, v. 14, p. 1–164, doi: <https://doi.org/10.1080/20014422.1932.11880545>, 1932.

Formatted: Font color: Black

Formatted

1420 Caley, T., Giraudeau, J., Malaizé, B., Rossignol, L., and Pierre, C., 2012. Agulhas leakage as a key process in the modes of Quaternary climate changes: *Proceedings of the National Academy of Sciences*, v. *Proc. Natl. Acad. Sci.*, 109(18), p. 6835-6839, doi: <https://doi.org/10.1073/pnas.1115545109>, 2012.

Formatted: Font: Not Bold, Font color: Black

Formatted

Formatted: Bibliography

Formatted: Font color: Black

Clapperton, C.M., 1993. Quaternary geology and geomorphology of South America: Amsterdam, Elsevier Science Publishers B.V., 779 p., 1993.

Formatted: Font: Not Bold

1425 Cogež, A., Herman, F., Pelt, É., Reuschlé, T., Morvan, G., Darvill, C. M., Norton, K. P., Christl, M., Märki, L., and Chabaux, F., 2018. U-Th and 10Be constraints on sediment recycling in proglacial settings, Lago Buenos Aires, Patagonia: *Earth Surface Dynamics*, v. *Surf. Dyn.*, 6(1), p. 121-140, doi: <https://doi.org/10.5194/esurf-6-121-2018>, 2018.

Formatted: Font: Not Bold, Font color: Black

Formatted

Formatted: Bibliography

Formatted: Font color: Black

1430 Colhoun, E.A., Barrows, T.T., 2011. The glaciation of Australia. In: Ehlers, J., Gibbard, P.L., Hughes, P.D. (Eds.), Quaternary Glaciations - Extent and Chronology: A Closer Look. *Developments in Quaternary Sciences* 15. Elsevier, Amsterdam, pp. 1037-1045, 2011.

Formatted: Font: Not Bold

1435 Darvill, C. M., Bentley, M. J., Stokes, C. R., Hein, A. S., and Rodés, Á., 2015. Extensive MIS 3 glaciation in southernmost Patagonia revealed by cosmogenic nuclide dating of outwash sediments: *Earth and Planetary Science Letters*, v. *Planet. Sci. Lett.*, 429, p. 157-169, doi: <https://doi.org/10.1016/j.epsl.2015.07.030>, 2015.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1440 Darvill, C. M., Bentley, M. J., Stokes, C. R., and Shulmeister, J., 2016. The timing and cause of glacial advances in the southern mid-latitudes during the last glacial cycle based on a synthesis of exposure ages from Patagonia and New Zealand: *Quaternary Science Reviews*, v. *Quat. Sci. Rev.*, 149, p. 200-214, doi: <https://doi.org/10.1016/j.quascirev.2016.07.024>, 2016.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1445 Davies, B. J., Roberts, D. H., Bridgland, D. R., Ó Cofaigh, C. Ó., Riding, J. B., Demarchi, B., Penkman, K. E. H., and Pawley, S. M., 2012. Timing and depositional environments of a Middle Pleistocene glaciation of northeast England: New evidence from Warren House Gill, County Durham: *Quaternary Science Reviews*, v. *Quat. Sci. Rev.*, 44, p. 180-212, <https://doi.org/10.1016/j.quascirev.2010.02.003>, 2012.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Davies, B. J., Darvill, C. M., Lovell, H., Bendle, J. M., Dowdeswell, J. A., Fabel, D., ~~García~~ García, J.-L., Geiger, A., Glasser, N. F., Gheorghiu, D. M., Harrison, S., Hein, A. S., Kaplan, M. R., Martin, J. R. V., Mendelova, M., Palmer, A., Pelto, M., ~~Rodés~~ Rodés, Á., Sagredo, E. A., Smedley, R. K., Smellie, J. L., and Thorndyraft, V. R., 2020. The evolution of the

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

1450 Patagonian Ice Sheet from 35 ka to the ~~Present Day~~present day (PATICE)), ~~Earth-Science Reviews~~Sci. Rev., 204, 103152, doi: <https://doi.org/10.1016/j.earscirev.2020.103152>, 2020.

~~De Boer, A. M., Graham, R. M., Thomas, M. D., and Kohfeld, K. E.: The control of the Southern Hemisphere Westerlies on the position of the Subtropical Front, J. Geophys. Res. Oceans, 118(10), 5669-5675, <https://doi.org/10.1002/jgrc.20407>, 2013.~~

1455 Denton, G. H., Lowell, T. V., Heusser, C. J., ~~Schlichter~~Schluchter, C., Andersen, B. G., Heusser, L. E., Moreno, P. J., and Marchant, D. R., 1999.: Geomorphology, ~~stratigraphy~~Stratigraphy, and ~~radio-carbon chronology~~Radiocarbon Chronology of ~~Hanquihue drift~~Llanquihue Drift in the ~~area~~Area of the ~~southern lake district~~Southern Lake District, ~~Seno Reloncavi~~Reloncavi, and Isla ~~grande~~Grande de ~~Chiloé~~Chiloe, Chile: ~~Geografiska Annaler, Series A, Physical Geography, v., Geogr. Ann. Ser. Phys. Geogr.~~ 81 (2), p. 167-229, doi: <https://doi.org/10.1111/j.0435-3676.1999.00057.x>, 1999.

~~Denton, G. H., Anderson, R. F., Toggweiler, J. R., Edwards, R. L., Schaefer, J. M., and Putnam, A. E., 2010.: The last glacial termination: Last Glacial Termination, Science, v., 328, p., 1652-1656, doi: <https://doi.org/10.1126/science.1184119>, 2010.~~

1465 Denton, G. H., Putnam, A. E., Russell, J. L., Barrell, D. J. A., Schaefer, J. M., Kaplan, M. R., and Strand, P. D., 2021.: The Zealandia Switch: Ice age climate shifts viewed from Southern Hemisphere moraines: ~~Quaternary Science Reviews, v., Quat. Sci. Rev.~~ 257, 106771, doi: <https://doi.org/10.1016/j.quascirev.2020.106771>, 2021.

1470 Douglass, D.C., Singer, B.S., Ackert, R.P., Kaplan, M.R., and Caffee, M.W., 2007.: Constraining Boulder Erosion Rates and Ages of Mid-Pleistocene Moraines. Lago Buenos Aires, Argentina. Geological Society of America Abstracts and Programs, Northeastern Section, 42nd Annual Meeting, 2007.

1475 Evans, D. J. A., Roberts, D. H., Bateman, M. D., Ely, J., Medialdea, A., Burke, M. J., Chiverrell, R. C., Clark, C. D., and Fabel, D., 2019.: A chronology for North Sea Lobe advance and recession on the Lincolnshire and Norfolk coasts during MIS 2 and 6: ~~Proceedings of the Geologists' Association, v., Proc. Geol. Assoc.~~, 130, p. 523-540, doi: <https://doi.org/10.1016/j.pgeola.2018.10.004>, 2019.

1480 Fernandes, M., Oliva, M., Vieira, G., Palacios, D., Fernández-Fernández, J. M., Delmas, M., García-Oteyza, J., Schimmelpennig, I., Ventura, J., ASTER Team., 2021., ~~Aumaitre, G., and Keddadouche, K.:~~ Maximum glacier extent of the Penultimate Glacial Cycle in the Upper Garonne Basin (Pyrenees): new chronological evidence: ~~Environmental, Environ. Earth Sciences, v., Sci.~~ 80(24), p. 1-20, doi: 796, <https://doi.org/10.1007/s12665-021-10022-z>, 2021.

Formatted

Formatted: Font color: Black

Formatted: Font color: Auto, Pattern: Clear

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1485
1490
1495
1500
1505
1510
1515

Fick, S. E., and Hijmans, R. J., 2017., WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas: *International journal of climatology*, v., *Int. J. Climatol.*, 37(42), p., 4302–4315, doi: <https://doi.org/10.1002/joc.5086>, 2017.

García, J., L., Hein, A. S., Binnie, S. A., Gómez, G. A., González, M. A., and Dunai, T. J., 2018., The MIS 3 maximum of the Torres del Paine and Última Esperanza ice lobes in Patagonia and the pacing of southern mountain glaciation: *Quaternary Science Reviews*, v., *Quat. Sci. Rev.*, 185, p., 9–26, doi: <https://doi.org/10.1016/j.quascirev.2018.01.013>, 2018.

García, J., L., Lüthgens, C., Vega, R. M., Rodés, Á., Hein, A. S., and Binnie, S., 2021., A composite 10Be, IR-50 and 14C chronology of the pre-Last Glacial Maximum (LGM) full ice extent of the western Patagonian Ice Sheet in the Isla de Chiloé, south Chile (42° S): *E&G Quaternary Science Journal*, v., *EampG Quat. Sci. J.*, 70, p., 105–128, doi: <https://doi.org/10.5194/egqsj-70-105-2021>, 2021.

Garreaud, R., Lopez, P., Minvielle, M., and Rojas, M., 2013., Large-scale control on the Patagonian climate: *Journal of Climate*, v., *J. Clim.*, 26(4), p., 215–230, doi: <https://doi.org/10.1175/JCLI-D-12-00001.1>, 2013.

Glasser, N., and Jansson, K., 2008., The glacial map of southern South America: *Journal of Maps*, v., 4(1), p., 175–196, doi: <https://doi.org/10.4113/jom.2008.1020>, 2008.

Gómez, G. A., García, J., L., Villagrán, C., Lüthgens, C., and Abarzúa, A. M., 2022., Vegetation, glacier, and climate changes before the global last glacial maximum in the Isla Grande de Chiloé, southern Chile (42° S): *Quaternary Science Reviews*, v., *Quat. Sci. Rev.*, 276, 107301, doi: <https://doi.org/10.1016/j.quascirev.2021.107301>, 2022.

Gosse, J. C., and Phillips, F. M., 2001., Terrestrial in situ cosmogenic nuclides: theory and application: *Quaternary Science Reviews*, v., *Quat. Sci. Rev.*, 20(14), p., 1475–1560, doi: [https://doi.org/10.1016/S0277-3791\(00\)00171-2](https://doi.org/10.1016/S0277-3791(00)00171-2), 2001.

Granger, D. E., and Muzikar, P. F., 2001., Dating sediment burial with in situ-produced cosmogenic nuclides: theory, techniques, and limitations: *Earth and Planetary Science Letters*, v., *Planct. Sci. Lett.*, 188(1–2), p., 269–281, doi: [https://doi.org/10.1016/S0012-821X\(01\)00309-0](https://doi.org/10.1016/S0012-821X(01)00309-0), 2001.

Haller, M., Lech, R.R., Martinez, O.A., Meister, C.M., and Page, S.M., 2003., Hoja Geologica 4373IV/III, Trevelin, Provincia del Chubut. Programa Nacional de Cartas Geologicas de la Republica Argentina: Servicio Geologico Nacional, Buenos Aires, scale 1:250.000, 2003.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

Formatted

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted

1520 Hein, A. S., Hulton, N. R. J., Dunai, T. J., Schnabel, C., Kaplan, M. R., Naylor, M., and Xu, S., 2009.; Middle Pleistocene glaciation in Patagonia dated by cosmogenic-nuclide measurements on outwash gravels: *Earth and Planetary Science Letters*, v. *Planet. Sci. Lett.*, 286(1-2), p. 184–197, doi: <https://doi.org/10.1016/j.epsl.2009.06.026>, 2009.

Formatted: Font: Not Bold, Font color: Black

Formatted

Formatted: Bibliography

Formatted: Font color: Black

1520 Hein, A. S., Dunai, T. J., Hulton, N. R. J., and Xu, S., 2011.; Exposure dating outwash gravels to determine the age of the greatest Patagonian glaciations: *Geology*, v. 39(2), p. 103–106, doi: <https://doi.org/10.1130/G31215.1>, 2011.

Formatted: Font: Not Bold, Font color: Black

Formatted: Bibliography

Formatted

Formatted: Font color: Black

1525 Hein, A. S., Coge, A., Darvill, C. M., Mendelová Mendelova, M., Kaplan, M. R., Herman, F., Dunai, T. J., Norton, K., Xu, S., Christl, M., and Rodés, Á., 2017.; Regional mid-Pleistocene glaciation in central Patagonia: *Quaternary Science Reviews*, v. *Quat. Sci. Rev.*, 164, p. 77–94, doi: <https://doi.org/10.1016/j.quascirev.2017.03.023>, 2017.

Formatted

Formatted: Font color: Black

Formatted

1530 Hervé, F., Fuentes, F. J., Calderón, M., Fanning, M., Quezada, P., Pankhurst, R., and Rapela, C., 2017.; Ultramafic rocks in the North Patagonian Andes: is their emplacement associated with the Neogene tectonics of the Liquiñe–Ofqui Fault Zone? *Andean Geology*, v. *Geol.*, 44(1), p. 1–16, doi: <https://doi.org/10.5027/andgeoV44n1-a01>, 2017.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1535 Hughes, P. D., Gibbard, P. L., and Ehlers, J., 2020.; The “missing glaciations” of the Middle Pleistocene: *Quaternary Research*, v. *Quat. Res.*, 96, p. 161–183, doi: <https://doi.org/10.1017/qua.2019.76>, 2020.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted

Formatted: Bibliography

Formatted: Font color: Black, Pattern: Clear

1535 Huybers, P., and Denton, G., 2008.; Antarctic temperature at orbital timescales controlled by local summer duration: *Nature Geoscience*, v. *Nat. Geosci.*, 1, p. 787–792, doi: <https://doi.org/10.1038/ngeo311>, 2008.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

1540 Jones, R. S., Small, D., Cahill, N., Bentley, M. J., and Whitehouse, P. L., 2019.; iceTEA: Tools for plotting and analysing cosmogenic-nuclide surface-exposure data from former ice margins: *Quaternary Geochronology*, v. *Quat. Geochronol.*, 51, p. 72–86, doi: <https://doi.org/10.1016/j.quageo.2019.01.001>, 2019.

Formatted: Font color: Black, Pattern: Clear

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

1540 Kaplan, M. R., Douglass, D. C., Singer, B. S., Ackert, R. P., and Caffee, M. W., 2005.; Cosmogenic nuclide chronology of pre-last glacial maximum moraines at Lago Buenos Aires, 46 degrees S, Argentina: *Quaternary Research*, v. *Quat. Res.*, 63, p. 301–315, doi: <https://doi.org/10.1016/j.yqres.2004.12.003>, 2005.

Formatted: Font color: Black, Pattern: Clear

Formatted

Formatted: Pattern: Clear

1545 Kaplan, M. R., Hein, A. S., Hubbard, A., and Lax, S. M., 2009.; Can glacial erosion limit the extent of glaciation? *Geomorphology*, v. 103(2), p. 172–179, doi: <https://doi.org/10.1016/j.geomorph.2008.04.020>, 2009.

Formatted: Font color: Auto, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

550 Kaplan, M. R., Schaefer, J. M., Denton, G. H., Barrell, D. J. A., Chinn, T. J. H., Putnam, A. E., Andersen, B. G., Finkel, R. C., Schwartz, R., and Doughty, A. M., 2010. Glacier retreat in New Zealand during the Younger Dryas stadial. *Nature*, *V*, 467(7312), p. 194–197, doi: <https://doi.org/10.1038/nature09313>, 2010.

555 Kaplan, M. R., Strelin, J. A., Schaefer, J. M., Denton, G. H., Finkel, R. C., Schwartz, R., Putnam, A. E., Vandergoes, M. J., Goehring, B. M., and Travis, S. G., 2011. In-situ cosmogenic ¹⁰Be production rate at Lago Argentino, Patagonia: implications for late-glacial climate chronology. *Earth and Planetary Science Letters*, *v. Planet. Sci. Lett.*, 309(1–2), p. 21–32, doi: <https://doi.org/10.1016/j.epsl.2011.06.018>, 2011.

560 Kiernan, K., Fifield, L. K., & Chappell, J., 2004. Cosmogenic nuclide ages for last glacial maximum Last Glacial Maximum moraine at Schnells Ridge, southwest Tasmania. *Quaternary Research*, *v. Quat. Res.*, 61(3), p. 335–338, doi: <https://doi.org/10.1016/j.yqres.2004.02.004>, 2004.

Kiernan, K., Fink, D., Greig, D., & Mifud, C., 2010. Cosmogenic radionuclide chronology of pre-last glacial cycle moraines in the Western Arthur range, Southwest Tasmania. *Quaternary Science Reviews*, *v. Quat. Sci. Rev.*, 29(23–24), p. 3286–3297, doi: <https://doi.org/10.1016/j.quascirev.2010.07.023>, 2010.

565 Kiernan, K., McMinn, M. S., and Fink, D., 2014. Topographic and microclimatic impacts on glaciation of the Denison Range, southwest Tasmania. *Quaternary Science Reviews*, *v. Quat. Sci. Rev.*, 97, p. 136–147, doi: <https://doi.org/10.1016/j.quascirev.2014.05.008>, 2014.

570 Kiernan, K., Fink, D., & McConnell, A., 2017. Cosmogenic ¹⁰Be and ²⁶Al exposure ages of glaciations in the Frankland Range, southwest Tasmania reveal a limited MIS-2 ice advance. *Quaternary Science Reviews*, *v. Quat. Sci. Rev.*, 157, p. 141–151, doi: <https://doi.org/10.1016/j.quascirev.2016.12.008>, 2017.

575 Koffman, B. G., Goldstein, S. L., Winckler, G., Borunda, A., Kaplan, M. R., Bolge, L., Cai, Y., Recasens, C., Koffman, T. N. B., and Vallelonga, P., 2021. New Zealand as a source of mineral dust to the atmosphere and ocean. *Quaternary Science Reviews*, *v. Quat. Sci. Rev.*, 251, 106659, doi: <https://doi.org/10.1016/j.quascirev.2020.106659>, 2021.

Lal, D., 1991. Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models. *Earth and Planetary Science Letters*, *v. Planet. Sci. Lett.*, 104, p. 424–439, doi: [https://doi.org/10.1016/0012-821X\(91\)90220-C](https://doi.org/10.1016/0012-821X(91)90220-C), 1991.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

1580 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M., 2014., Sea level and global ice volumes from the Last Glacial Maximum to the Holocene: *Proceedings of the National Academy of Sciences*, v., *Proc. Natl. Acad. Sci.*, 111(43), p., 15296–15303., <https://doi.org/10.1073/pnas.1411762111>, 2014.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

1585 Lambert, F., Delmonte, B., Petit, J. R., Bigler, M., Kaufmann, P. R., Hutterli, M. A., Stocker, T. F., Ruth, U., Steffensen, J. P., and Maggi, V., 2008., Dust-climate couplings over the past 800,000 years from the EPICA Dome C ice core., *Nature*, v., 452(7187), p., 616–619, doi:<https://doi.org/10.1038/nature06763>, 2008.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

1590 Lambert, F., Bigler, M., Steffensen, J. P., Hutterli, M., and Fischer, H., 2012., Centennial mineral dust variability in high-resolution ice core data from Dome C, Antarctica: *Climate of the Past*, v., 8, p., 609–623, doi:<https://doi.org/10.5194/cp-8-609-2012>, 2012.

Formatted

Formatted: Bibliography

Formatted: Font color: Black, Pattern: Clear

1595 Leger, T. P. M., Hein, A. S., Bingham, R. G., Martini, M. A., Rodrigo Soteres, R. L. S., Sagredo, E. A., and Martínez, O. A., 2020., The glacial geomorphology of the Río Corcovado, Río Huemul and Lago Palena/General Vinter valleys, northeastern Patagonia (43°S, 71°W): *Journal of Maps*, v., 16(2), p., 651–668, doi:<https://doi.org/10.1080/17445647.2020.1794990>, 2020.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1600 Leger, T. P. M., Hein, A. S., Bingham, R. G., Rodés, Á., Fabel, D., and Smedley, R., 2021a., K., Geomorphology and ¹⁰Be chronology of the Last Glacial Maximum and deglaciation in northeastern Patagonia, 43°S–71°W: *Quaternary Science Reviews*, v., *Quat. Sci. Rev.*, 272, doi:107194, <https://doi.org/10.1016/j.quascirev.2021.107194>, 2021a.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted

Formatted: Bibliography

Formatted: Font color: Black, Pattern: Clear

1605 Leger, T. P. M., Hein, A. S., Goldberg, D., Schimmelfennig, I., Van Wyk de Vries, M. S., Bingham, R. G., and ASTER Team., 2021b., Northeastern Patagonian Glacier Advances (43°S) Reflect Northward Migration of the Southern Westerlies Towards the End of the Last Glaciation: *Frontiers in Earth Science*, v., *Sci.*, 9, 751987, doi:<https://doi.org/10.3389/feart.2021.751987>, 2021b.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1610 Lifton, N., Sato, T., and Dunai, T. J., 2014., Scaling in situ cosmogenic nuclide production rates using analytical approximations to atmospheric cosmic-ray fluxes: *Earth and Planetary Science Letters*, v., *Planet. Sci. Lett.*, 386, p., 149–160, doi:<https://doi.org/10.1016/j.epsl.2013.10.052>, 2014.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Lindner, L., and Marks, L., 1999., New approach to stratigraphy of palaeolake and glacial sediments of the younger Middle Pleistocene in mid-eastern Poland: *Geological Quarterly*, v. 43(1), p. 1–8, 1999.

Formatted: Font: Not Bold

1615 [Lisiecki, L. E., and Raymo, M. E., 2005., A Pliocene–Pleistocene stack of 57 globally distributed benthic \$\delta^{18}O\$ records., *PLIOCENE-PLEISTOCENE BENTHIC STACK*, *Paleoceanography*, v. 20\(1\), doi: n/a-n/a, <https://doi.org/10.1029/2004PA001071>, 2005.](#)

[Lorrey, A.M., Bostock, H., 2017., The Quaternary climate of New Zealand. *Advances in Quaternary Science - the New Zealand Landscape*. Springer-Verlag, pp. 67-139., 2017](#)

1620 [Margari, V., Skinner, L. C., Hodell, D. A., Martrat, B., Toucanne, S., Grimalt, J. O., Gibbard, P. L., Lunkka, J. P., and Tzedakis, P. C., 2014., Land-ocean changes on orbital and millennial time scales and the penultimate glaciation., *Geology*, v. 42, p. 183–186, doi: <https://doi.org/10.1130/G35070.1>, 2014.](#)

1625 [Martin, L. C. P., Blard, P., H., Balco, G., Lavé, J., Delunel, R., Lifton, N., and Laurent, V., 2017., The CREp program and the ICE-D production rate calibration database: A fully parameterizable and updated online tool to compute cosmic-ray exposure ages., *Quaternary geochronology*, v. Quat. Geochronol., 38, p. 25–49, doi: <https://doi.org/10.1016/j.quageo.2016.11.006>, 2017.](#)

1630 [Meglioli, A., 1992., Glacial geology and chronology of southernmost Patagonia and Tierra del Fuego, Argentina and Chile \[Ph.D. thesis\]: Bethlehem, Pennsylvania, Lehigh University, 216 p., 1992.](#)

1635 [Mendelová, M., Hein, A. S., Rodés, Á., & Xu, S., 2020., Extensive mountain glaciation in central Patagonia during Marine Isotope Stage 5., *Quaternary Science Reviews*, v. Quat. Sci. Rev., 227, 105996, doi: <https://doi.org/10.1016/j.quascirev.2019.105996>, 2020.](#)

[Mercer, J. H., 1976., Glacial history of southernmost Southernmost South America., *Quaternary Research*, v. Quat. Res., 6, p. 125–166, doi: \[https://doi.org/10.1016/0033-5894\\(76\\)90047-8\]\(https://doi.org/10.1016/0033-5894\(76\)90047-8\), 1976.](#)

1640 [Mercer, J. H., 1984., Simultaneous climatic change in both hemispheres and similar bipolar interglacial warming: Evidence and implications, in: *Geophysical Monograph Series*, vol. 29, edited by: Hansen, J. E., *Climate processes and climate sensitivity*, v. 29, p. 307–313, doi: and Takahashi, T., American Geophysical Union, Washington, D. C., 307–313, <https://doi.org/10.1029/GM029p0307>, 1984.](#)

1645 [Parrenin, F., Masson-Delmotte, V., Köhler, P., Raynaud, D., Paillard, D., Schwander, J., Barbante, C., Landais, A., Wegner, A., and Jouzel, J., 2013., Synchronous changeChange of atmosphericAtmospheric, CO₂ and Antarctic temperature](#)

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted: Font: Not Bold

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Formatted: Font color: Black, Pattern: Clear

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Formatted: Bibliography

Formatted

during Temperature During the last deglacial warming: Last Deglacial Warming, *Science*, v. 339(6123), p. 1060–1063, doi: <https://doi.org/10.1126/science.1226368>, 2013.

Formatted

Formatted: Font color: Black, Pattern: Clear

Peltier, C., Kaplan, M. R., Schaefer, J. M., Araos, J., Sagredo, E. A., and Schwartz, R., 2017 (December), Glacier Climate Change in Southern South America over Multiple Glacial Cycles and Implications for Glacial to Interglacial Shifts in the Westerly Winds: In 2017 AGU Fall Meeting, AGU, Bibecode: 2017AGUFMPP43B1349P

Formatted: Font: Not Bold, Font color: Black, Swedish (Sweden), Pattern: Clear

Formatted: Font color: Black, Swedish (Sweden), Pattern: Clear

Peltier, C., Kaplan, M. R., Birkel, S. D., Soteres, R. L., Sagredo, E. A., Aravena, J. C., Araos, J., Moreno, P. J., Schwartz, R., and Schaefer, J. M., 2021, A.: The large MIS 4 and long MIS 2 glacier maximum maxima on the southern tip of South America: *Quaternary Science Reviews*, v. *Quat. Sci. Rev.*, 262, 106858, doi: <https://doi.org/10.1016/j.quascirev.2021.106858>, 2021.

Formatted: Font color: Black, Swedish (Sweden), Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Phillips, F. M., Zreda, M. G., Smith, S. S., Elmore, D., Kubik, P. W., and Sharma, P., 1990.: Cosmogenic chlorine Chlorine-36 chronology Chronology for glacial deposits Glacial Deposits at Bloody Canyon, eastern Eastern Sierra Nevada, *Science*, v. 248, p. 1529–1532, doi: <https://doi.org/10.1126/science.248.4962.1529>, 1990.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Preusser, F., Graf, H. R., Keller, O., Krayss, E., and Schlüchter, C., 2011.: Quaternary glaciation history of northern Switzerland: *E&G Quaternary Science Journal*, v. *EampG Quat. Sci. J.*, 60(2/3), p. 282–305, doi: <https://doi.org/10.3285/eg.60.2-3.06>, 2011.

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Putkonen, J., and Swanson, T., 2003.: Accuracy of cosmogenic ages for moraines: *Quaternary Research*, v. *Quat. Res.*, 59(2), p. 255–261, doi: [https://doi.org/10.1016/S0033-5894\(03\)00006-1](https://doi.org/10.1016/S0033-5894(03)00006-1), 2003.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Putnam, A. E., Schaefer, J. M., Barrell, D. J. A., Vandergoes, M., Denton, G. H., Kaplan, M. R., Finkel, R. C., Schwartz, R., Goehring, B. M., and Kelley, S. E., 2010.: In situ cosmogenic ¹⁰Be production-rate calibration from the Southern Alps, New Zealand: *Quaternary Geochronology*, v. *Quat. Geochronol.*, 5(4), p. 392–409, doi: <https://doi.org/10.1016/j.quageo.2009.12.001>, 2010.

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Putnam, A. E., Schaefer, J. M., Denton, G. H., Barrell, D. J. A., Birkel, S. D., Andersen, B. G., Kaplan, M. R., Finkel, R. C., Schwartz, R., and Doughty, A. M., 2013.: The Last Glacial Maximum at 44°S documented by a ¹⁰Be moraine chronology at Lake Ohau, Southern Alps of New Zealand. *Quaternary Science Reviews*, v. 62, 114e141, doi: <https://doi.org/10.1016/j.quascirev.2012.10.034>, 2013.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Rabassa, J., and Clapperton, C. M., 1990.: Quaternary glaciations of the southern Andes: *Quaternary Science Reviews*, v. *Quat. Sci. Rev.*, 9(2-3), p. 153–174, doi: [https://doi.org/10.1016/0277-3791\(90\)90016-4](https://doi.org/10.1016/0277-3791(90)90016-4), 1990.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1680 Rabassa, J., and Coronato, A., 2009., Glaciations in Patagonia and Tierra del Fuego during the Ensenadan stage/age (earlyStage/Age (Early Pleistocene–earliest middleMiddle Pleistocene): Quaternary International, v., Quat. Int., 210(1–2), p. 18–36, doi: <https://doi.org/10.1016/j.quaint.2009.06.019>, 2009.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

1685 Sarmiento, J. L., and Toggweiler, J. R., 1984., A new model for the role of the oceans in determining atmospheric pP CO₂. Nature, v., 308(5960), p. 621–624, doi: <https://doi.org/10.1038/308621a0>, 1984.

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1690 Schaefer, J. M., Putnam, A. E., Denton, G. H., Kaplan, M. R., Birkel, S., Doughty, A. M., Kelley, S. E., Barrell, D. J. A., Finkel, R. C., Winckler, G., Anderson, R. F., Ninneman, U. S., Barker, S., Schwartz, R., Andersen, B. J. G., and Schluechter, C., 2015., The southern glacial maximum Southern Glacial Maximum 65,000 years ago and its unfinished termination: Quaternary Science Reviews, v., Unfinished Termination, Quat. Sci. Rev., 114, p. 52–60, doi: <https://doi.org/10.1016/j.quascirev.2015.02.009>, 2015.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1695 Shakun, J. D., Lea, D. W., Lisiecki, L. E., and Raymo, M. E., 2015., An 800-kyr record of global surface ocean δ¹⁸O₁₈O and implications for ice volume-temperature coupling: Earth and Planetary Science Letters, v., Planet. Sci. Lett., 426, p. 58–68, doi: <https://doi.org/10.1016/j.epsl.2015.05.042>, 2015.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

1700 Sime, L. C., Kohfeld, K. E., Le Quéré, C., Wolff, E. W., de Boer, A. M., Graham, R. M., and Bopp, L., 2013., Southern Hemisphere westerly wind changes during the Last Glacial Maximum: model-data comparison: Quaternary Science Reviews, v., Quat. Sci. Rev., 64, p. 104–120, doi: <https://doi.org/10.1016/j.quascirev.2012.12.008>, 2013.

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1705 Singer, B. S., Brown, L. L., Rabassa, J. O., and Guillou, H., 2004., ⁴⁰Ar/³⁹Ar chronology of Late Pliocene and Early Pleistocene geomagnetic and glacial events in southern Argentina: In Timescales Of The Paleomagnetic Field (eds J., in: Geophysical Monograph Series, edited by: Channell, D. J. E. T., Kent, W. D. V., Lowrie, W., and J. Meert), doi: , J. G., American Geophysical Union, Washington, D. C., 175–190, <https://doi.org/10.1029/145GM13>, 2013.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1710 Spratt, R. M., and Lisiecki, L. E., 2016., A Late Pleistocene sea level stack: Climate of the Clim. Past, v., 12(4), p. 1079–1092, doi: <https://doi.org/10.5194/cp-12-1079-2016>, 2016.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

Stone, J. O., 2000., Air pressure and cosmogenic isotope production: Journal of Geophysical Research: J. Geophys. Res. Solid Earth, v., 105(B10), p. 23753–23759, doi: <https://doi.org/10.1029/2000jb900181>, <https://doi.org/10.1029/2000JB900181>, 2000.

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1715 Sugden, D. E., McCulloch, R. D., Bory, A. J. M., and Hein, A. S., 2009. Influence of Patagonian glaciers on Antarctic dust deposition during the last glacial period: *Nature Geoscience*, v. *Nat. Geosci.*, 2(4), p. 281–285, doi: <https://doi.org/10.1038/ngeo474>, 2009.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

Sylwan, C., Beraza, L., and Caselli, A., 1991. Magnetostratigrafía de la secuencia morenica en el valle del Lago Pueyrredon, provincia de Santa Cruz: *Revista de la Asociación Geológica Argentina*, v. 46, p. 235–238, 1991.

Formatted: Font: Not Bold

1720 Terrizzano, C. M., García Morabito, E. G., Christl, M., Likerman, J., Tobal, J., Yamin, M., & Zech, R., 2017. Climatic and Tectonic forcing on alluvial fans in the Southern Central Andes: *Quaternary science reviews*, v. *Quat. Sci. Rev.*, 172, p. 131–141, doi: <https://doi.org/10.1016/j.quascirev.2017.08.002>, 2017.

Formatted

Formatted: Pattern: Clear

Formatted: Font color: Auto, Pattern: Clear

1725 Toucanne, S., Zaragosi, S., Bourillet, J. F., Cremer, M., Eynaud, F., Van Vliet-Lanoë, B., Penaud, A., Fontanier, C., Turon, J. L., and Cortijo, E., and Gibbard, P. L., 2009. Timing of massive “Fleuve Manche” discharges over the last 350 kyr: insights into the European ice-sheet oscillations and the European drainage network from MIS 10 to 2: *Quaternary Science Reviews*, v. *Quat. Sci. Rev.*, 28, p. 1238–1256, doi: <https://doi.org/10.1016/j.quascirev.2009.01.006>, 2009.

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1730 Trombotto, D., 2008. Geocryology of southern South America, in Rabassa, J., ed., *The late Cenozoic of Patagonia and Tierra del Fuego: Developments in Quaternary Sciences Volume, vol. 11*: Amsterdam, Elsevier, p. 255–268, doi: [https://doi.org/10.1016/S1571-0866\(07\)10012-9](https://doi.org/10.1016/S1571-0866(07)10012-9), 2008.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1735 White, T. S., Bridgland, D. R., Howard, A. J., Westaway, R., Howard, A. J., and White, M. J., 2010. Evidence from the Trent terrace archive, Lincolnshire, UK, for lowland glaciation of Britain during the Middle and Late Pleistocene: *Proceedings of the Geologists' Association*, v. *Proc. Geol. Assoc.*, 121, p. 141–153, doi: <https://doi.org/10.1016/j.pgeola.2010.05.001>, 2010.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1740 White, T. S., Bridgland, D. R., Westaway, R., and Straw, A., 2017. Evidence for late Middle Pleistocene glaciation of the British margin of the southern North Sea: *Journal of Quaternary Science*, v. *LATE MIDDLE PLEISTOCENE GLACIATION OF THE BRITISH NORTH SEA MARGIN*, *J. Quat. Sci.*, 32(2), p. 261–275, doi: <https://doi.org/10.1002/jqs.2826>, 2017.

Formatted: Font: Not Bold, Font color: Black, Pattern: Clear

Formatted: Bibliography

Formatted

Formatted: Font color: Black, Pattern: Clear

1745

